

Pierre-Antoine Graham

HOMEWORK 1

Aldo Riello
Classical Physics

Perimeter Institute for Theoretical Physics
September 12, 2023

Contents

1	Bead on a rotating wire	2
A	Lagrangian	2
B	Time Translation Symmetry Conserved Charge	2
C	Effective Potential	3
D	Discussion	3
2	Brachistochrone	4
A	Travel time fonctionnal	4
B	Conserved quantity	4
3	A sneaky recipe for the Noether charge	5
A	Alternate Disformation	5
B	Variation of the Action	5
C	Conserved quantity	5
D	Application to time translations	6

1 Bead on a rotating wire

A point particle of mass m is constrained to move on a parabola rotating with angular velocity ω around a z axis aligned with the opposite of a gravitationnal field of intensity g . The parabola is described in cylindrical coordinates ρ, ϕ, z by the relations $z = \alpha\rho^2$ and $\phi = \omega t$.

A Lagrangian

Following the constraints imposed on the particle, its cartesian position can be written as the following

$$x = \rho \cos(\omega t), \quad y = \rho \sin(\omega t), \quad \text{and} \quad z = \alpha\rho^2$$

leaving ρ as the only free parameter describing the motion of the particle. The cartesian velocity of the particle is then given by the time derivative of its cartesian coordinates. We have

$$\dot{x} = \dot{\rho} \cos(\omega t) - \omega\rho \sin(\omega t), \quad \dot{y} = \dot{\rho} \sin(\omega t) + \omega\rho \cos(\omega t), \quad \text{and} \quad \dot{z} = 2\alpha\rho\dot{\rho}.$$

The square magnitude of the cartesian velocity of the particle can be written in cylindrical coordinates as

$$\begin{aligned} \dot{x}^2 + \dot{y}^2 + \dot{z}^2 &= (\dot{\rho} \cos(\omega t) - \omega\rho \sin(\omega t))^2 + (\dot{\rho} \sin(\omega t) + \omega\rho \cos(\omega t))^2 + (2\alpha\rho\dot{\rho})^2 + (2\alpha\rho\dot{\rho})^2 \\ &= \dot{\rho}^2 \cos^2(\omega t) + \omega^2 \rho^2 \sin^2(\omega t) + \dot{\rho}^2 \sin^2(\omega t) + \omega^2 \rho^2 \cos^2(\omega t) \\ &\quad + 2\dot{\rho}\rho\omega \cos(\omega t) \sin(\omega t) - 2\dot{\rho}\rho\omega \cos(\omega t) \sin(\omega t) + 4\alpha^2 \rho^2 \dot{\rho}^2 \\ &= \dot{\rho}^2 + \omega^2 \rho^2 + 4\alpha^2 \rho^2 \dot{\rho}^2. \end{aligned}$$

Following from this expression, the kinetic energy reads $T = \frac{1}{2}m(\dot{\rho}^2 + \omega^2 \rho^2 + 4\alpha^2 \rho^2 \dot{\rho}^2)$. The potential energy du to gravitationnal interaction is given by $V = mgz = mg\alpha\rho^2$. We can finally write the lagrangian

$$L = T - V = \frac{1}{2}m((1 + 4\alpha^2 \rho^2)\dot{\rho}^2 + \omega^2 \rho^2) - mg\alpha\rho^2.$$

B Time Translation Symmetry Conserved Charge

Since $\frac{\partial L}{\partial t} = 0$, Noether's theorem ensures that

$$\begin{aligned} E &= \frac{\partial L}{\partial \dot{\rho}} \dot{\rho} - L = m(1 + 4\alpha^2)\dot{\rho}^2 - \frac{1}{2}m((1 + 4\alpha^2 \rho^2)\dot{\rho}^2 + \omega^2 \rho^2) + mg\alpha\rho^2 \\ &= \frac{1}{2}m(1 + 4\alpha^2 \rho^2)\dot{\rho}^2 + \underbrace{m\rho^2 \left(g\alpha - \frac{1}{2}\omega^2 \right)}_{U_{\text{eff}}} \end{aligned}$$


is a conserved charge on physical trajectories which corresponds to an effective energy as it is associated to time translation symmetry of L . The potential-like term in E can be written as $U_{\text{eff}} = \frac{1}{2}m\rho^2(\omega_0^2 - \omega^2)$ where $\omega_0 = \sqrt{2g\alpha}$.

C

Effective Potential

The qualitative dynamics of the system can be described with U_{eff} by considering the three following regimes:

- $\omega_0 > \omega$ Gravity is stronger than rotation and the particle trapped in an harmonic potential. Note that when it oscillates through " $\rho = 0$ " it doesn't stop. Indeed, " $\rho = 0$ " is not included in the coordinate chart provided by spherical coordinates. In reality, nothing special happens at the bottom of the parabola and the particle continues to the other side;
- $\omega_0 = \omega$ In this situation gravity balances the centripetal and parabolic normal forces resulting in a net force accounting only for uniform rotation motion at fixed z . This is an unstable equilibrium (the two other regimes bring the particle far from this equilibrium);
- $\omega_0 < \omega$ This situation corresponds to an unbounded growth of z . The effective potential represents the effect of the centrifugal effect pushing the particle away from the origin more than gravity is pulling it.



In each case the arrows indicate the direction of the force which is constant in each regime.

D

Discussion

The total energy of the particle is given by

$$E_0 = T + V = \frac{1}{2}m((1 + 4\alpha^2\rho^2)\dot{\rho}^2 + \omega^2\rho^2) + mg\alpha\rho^2 = E + m\rho^2\omega^2.$$

The conservation of E implies $\dot{E}_0 = \dot{E} + 2m\rho\dot{\rho}\omega^2 = 2m\rho\dot{\rho}\omega^2$. From this, we see that E_0 is conserved on shell iff $\dot{\rho} = 0$ or $\rho = 0$. In the former case, ρ is a constant and this can only be realised if $\omega_0 = \omega$ or $\rho = 0$ (no effective force). In the latter case, $\rho = 0$ which is a subcase of $\dot{\rho} = 0$. If $\omega_0 < \omega$, the parabola is rotating fast enough to go beyond the gravitational pull of the particle towards $\rho = 0$ and the velocity of the particle grows without bounds (the motor rotating the parabola is transferring energy to the particle and E_0 is not conserved).

2 Brachistochrone

Consider a particle subject to a uniform gravitational field of strength g oriented in the negative y direction of a planar Oxy cartesian coordinate system. The particle starts at point A of coordinates x_A, y_A and ends at point B of coordinates x_B, y_B . We are interested in the curve linking A and B minimizing the travel time of the particle.

A Travel time fonctionnal

The first step in constructing the travel time fonctionnal is the parametrisation of its curve argument. Suppose the trajectory is parametrised by y so that it takes the form of a function $x(y)$. The length element at x is $dl = \sqrt{\dot{x}(y)^2 + \dot{y}^2} dy = \sqrt{1 + \dot{x}(y)^2} dy$ with $\dot{() = \frac{d}{dy}}$. By a mass independant analogue of the conservation of energy $0 = \frac{1}{2}v^2 - gy$ (with "potential" energy zero at y_B) where v is the velocity, we have $v = \sqrt{2gy}$. For motion along an element dl of the curve, the time elapsed is $dt = dl / \sqrt{2gy}$ leading to the following travel time fonctionnal

$$T[x(y)] = \int_{y_A}^{y_B} dy \frac{\sqrt{1 + \dot{x}(y)^2}}{\sqrt{2gy}}.$$

B Conserved quantity

Since the integrand (effective lagrangian $L = \sqrt{1 + \dot{x}(y)^2} / \sqrt{2gy}$) of the previous fonctionnal is cyclic in x , the Euler-Lagrange equations imply

$$\frac{d}{dy} \left(\frac{\partial L}{\partial \dot{x}} \right) = 0 \implies Q = \frac{\partial L}{\partial \dot{x}} = \frac{2\dot{x}(y)}{\sqrt{1 + \dot{x}(y)^2} \sqrt{2gy}} \text{ is conserved on Brachistochrone.}$$

We can now write

$$\begin{aligned} gyQ^2(1 + \dot{x}(y)^2) &= 2\dot{x}(y)^2 \implies \dot{x}(y) = \sqrt{\frac{gyQ^2}{2 - gyQ^2}} \\ \implies x(y) &= \int dy \sqrt{\frac{gyQ^2}{2 - gyQ^2}} = \int dy \sqrt{\frac{gyQ^2}{2 - gyQ^2}} \end{aligned}$$

3 A sneaky recipe for the Noether charge

A Alternate Disformation

Consider a particle described at time t by the generalized coordinate $q(t)$ evolving with respect to a lagrangian L . A disformation of the particle's history is given by $q(t) \mapsto q' = q(t) + \varepsilon \tilde{\delta}_s q(t)$ where $\delta \varepsilon$ is an infinitesimal constant. This disformation contributes to the action S with a boundary term if there exists a function R_s such that $\tilde{\delta}_s L = \frac{d}{dt} R_s$ (first order variation of L in ε).

B Variation of the Action

An alternative disformation of the history of the particle reads $q(t) \mapsto q'(t) = q(t) + \varepsilon f(t) \tilde{\delta}_s q(t)$ where f is a function vanishing at the start ($t = t_0$) and end ($t = t_1$) of the history. The action following from this disformation is

$$\begin{aligned}
 & S(q(t) + \varepsilon f(t) \tilde{\delta}_s q(t)) \\
 &= \int_{t_0}^{t_1} dt \, L(q(t) + \varepsilon f(t) \tilde{\delta}_s q(t), \dot{q}(t) + \varepsilon \frac{d}{dt}(f(t) \tilde{\delta}_s q(t)), t) \\
 &= S(q(t)) + \varepsilon \int_{t_0}^{t_1} dt \left[\frac{\partial L}{\partial q} f(t) \tilde{\delta}_s q(t) + \frac{\partial L}{\partial \dot{q}} (\dot{f}(t) \tilde{\delta}_s q(t) + f(t) \tilde{\delta}_s \dot{q}(t)) \right] + O(\varepsilon^2) \\
 &= S(q(t)) + \varepsilon \int_{t_0}^{t_1} dt \left[\frac{\partial L}{\partial q} f(t) \tilde{\delta}_s q(t) - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}} \tilde{\delta}_s q(t) \right) f(t) + \frac{\partial L}{\partial \dot{q}} f(t) \tilde{\delta}_s \dot{q}(t) \right] + \left[\frac{\partial L}{\partial \dot{q}} f(t) \tilde{\delta}_s q(t) \right]_{t_0}^{t_1} + O(\varepsilon^2) \\
 &= S(q(t)) + \varepsilon \int_{t_0}^{t_1} dt \left[\left(\frac{\partial L}{\partial q} \tilde{\delta}_s q(t) + \frac{\partial L}{\partial \dot{q}} \tilde{\delta}_s \dot{q}(t) \right) - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}} \tilde{\delta}_s q(t) \right) \right] f(t) + O(\varepsilon^2) \\
 &= S(q(t)) + \varepsilon \int_{t_0}^{t_1} dt \left[\frac{d}{dt} R_s - \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}} \tilde{\delta}_s q(t) \right) \right] f(t) + O(\varepsilon^2)
 \end{aligned}$$

and its first order variation reads $\delta_s^f S = - \int_{t_0}^{t_1} dt \dot{Q}_s f$ with $Q_s = \frac{\partial L}{\partial \dot{q}} \tilde{\delta}_s q(t) - R_s$. Using integration by parts we get

$$\delta_s^f S = \int_{t_0}^{t_1} dt \, Q_s \dot{f} + [\dot{Q}_s f]_{t_0}^{t_1} = \int_{t_0}^{t_1} dt \, Q_s \dot{f}.$$

C Conserved quantity

Since $f(t) \tilde{\delta}_s q(t)$ vanishes on the end points of the history, it can be interpreted the usual $\tilde{\delta}_s q(t)$ used in the action principle leading to on-shell histories. This means that the first order variation of S vanishes independently of f on-shell. We have $0 = \delta_s^f S = - \int_{t_0}^{t_1} dt \dot{Q}_s f \implies \dot{Q}_s = 0$. Note that since \dot{f} excludes constant functions except $f = 0$, $\int_{t_0}^{t_1} dt \, Q_s \dot{f}$ can't be used to conclude $Q_s = 0$ (\dot{f} is not arbitrary).

D

Application to time translations

Consider the Lagrangian $L = m\dot{q}^2/2 - V(q)$ describing a particle of mass m in a potential V . The infinitesimal disformation of an history due to time translation is given by $q'(t) = q(t + \varepsilon) = q(t) + \varepsilon \dot{q}$ is associated to $\tilde{\delta}_s q = \dot{q}$. Here, we consider an alternate disformation given by $q'(t) = q(t + \varepsilon) = q(t) + f(t)\varepsilon \dot{q}$ leading to the following variation of the action:

$$\begin{aligned} & S(q(t) + \varepsilon f(t)\tilde{\delta}_s q(t)) \\ &= \int_{t_0}^{t_1} dt \, L(q(t) + \varepsilon f(t)\tilde{\delta}_s q(t), \dot{q}(t) + \varepsilon \frac{d}{dt}(f(t)\tilde{\delta}_s q(t)), t) \end{aligned}$$

References

- [1] Aldo Riello. *Fourteen Lectures in CLASSICAL PHYSICS*. 2023.