

**my-ebook**

Parmeshvar

2025-07-06

# Table of contents

<b>1</b>	<b>Introduction</b>	<b>9</b>
1.1	Teaching . . . . .	9
1.2	Lecture notes . . . . .	10
1.3	Moodle website . . . . .	10
<b>2</b>	<b>Week 1</b>	<b>14</b>
2.1	Module 1: Introduction to Statistics . . . . .	14
2.2	Misuse of Statistics . . . . .	15
<b>3</b>	<b>Module 2: Mathematics vs Statistics</b>	<b>16</b>
<b>4</b>	<b>Module 3: Software-Based Statistical Revolution</b>	<b>17</b>
4.1	From Paper to Code . . . . .	17
4.2	Popular Statistical Software . . . . .	17
4.3	GUI vs CLI . . . . .	17
4.4	Recommended GUI Tools for R . . . . .	18
4.5	Installing RKWard on Ubuntu . . . . .	18
<b>5</b>	<b>Module 4: Understanding Variables</b>	<b>19</b>
5.1	What is a Variable? . . . . .	19
5.1.1	R Definition: . . . . .	19
<b>6</b>	<b>Week 2</b>	<b>26</b>
<b>7</b>	<b>Introduction</b>	<b>27</b>
7.1	Purpose of the eBook . . . . .	27
7.2	Who Should Read This? . . . . .	27
7.3	What You'll Learn . . . . .	27
<b>8</b>	<b>1. Fundamentals of Statistics</b>	<b>28</b>
8.1	1.1 What is Statistics? . . . . .	28
8.2	1.2 Key Objectives . . . . .	28
8.3	1.3 Types of Statistics . . . . .	28
<b>9</b>	<b>2. Types of Data</b>	<b>29</b>
9.1	2.1 Classification of Data . . . . .	29
9.1.1	2.1.1 Qualitative (Categorical) Data . . . . .	29
9.1.2	2.1.2 Quantitative (Numerical) Data . . . . .	29
<b>10</b>	<b>3. Descriptive Statistics</b>	<b>30</b>
10.1	3.1 Measures of Central Tendency . . . . .	30
10.1.1	3.1.1 What is Central Tendency? . . . . .	30

10.1.2	3.1.2 Characteristics of a Good Measure . . . . .	30
10.2	3.2 The Mean . . . . .	30
10.2.1	3.2.1 Definition . . . . .	30
10.2.2	3.2.2 Formula . . . . .	30
10.2.3	3.2.3 Properties of Mean . . . . .	30
10.2.4	3.2.4 Example . . . . .	31
10.3	3.3 The Median . . . . .	31
10.3.1	3.3.1 Definition . . . . .	31
10.3.2	3.3.2 Calculation . . . . .	31
10.3.3	3.3.3 Properties . . . . .	31
10.3.4	3.3.4 Example . . . . .	31
10.4	3.4 The Mode . . . . .	31
10.4.1	3.4.1 Definition . . . . .	31
10.4.2	3.4.2 Characteristics . . . . .	31
10.4.3	3.4.3 Example . . . . .	32
10.5	3.5 Comparison Table . . . . .	32
<b>11</b>	<b>4. Measures of Variability</b>	<b>33</b>
11.1	4.1 Why Measure Variability? . . . . .	33
11.2	4.2 Range . . . . .	33
11.2.1	4.2.1 Definition . . . . .	33
11.2.2	4.2.2 Example . . . . .	33
11.2.3	4.2.3 Limitations . . . . .	33
11.3	4.3 Quartiles and Interquartile Range . . . . .	33
11.3.1	4.3.1 Quartiles . . . . .	33
11.3.2	4.3.2 Formula for Position . . . . .	34
11.3.3	4.3.3 IQR Formula . . . . .	34
11.3.4	4.3.4 Example . . . . .	34
11.4	4.4 Variance . . . . .	34
11.4.1	4.4.1 Concept . . . . .	34
11.4.2	4.4.2 Formulas . . . . .	34
11.5	4.5 Standard Deviation . . . . .	34
11.5.1	4.5.1 Concept . . . . .	34
11.5.2	4.5.2 Properties . . . . .	35
11.6	4.6 Coefficient of Variation (CV) . . . . .	35
11.6.1	4.6.1 Definition . . . . .	35
11.6.2	4.6.2 Example . . . . .	35
11.7	4.7 Moment-Based Measures . . . . .	35
<b>12</b>	<b>5. Probability Fundamentals</b>	<b>36</b>
12.1	5.1 Introduction to Probability . . . . .	36
12.2	5.2 Key Definitions . . . . .	36
12.3	5.3 Types of Events . . . . .	36
12.4	5.4 Classical Probability . . . . .	36
12.5	5.5 Probability Rules . . . . .	37
12.5.1	Rule 1: Non-Negativity . . . . .	37
12.5.2	Rule 2: Total Probability . . . . .	37
12.5.3	Rule 3: Complement Rule . . . . .	37

12.5.4	Rule 4: Addition Rule . . . . .	37
12.5.5	Rule 5: Multiplication Rule . . . . .	37
12.6	5.6 Conditional Probability . . . . .	37
<b>13</b>	<b>6. Discrete Probability Distributions</b>	<b>38</b>
13.1	6.1 Bernoulli Distribution . . . . .	38
13.1.1	Example: . . . . .	38
13.2	6.2 Binomial Distribution . . . . .	38
13.2.1	Example: . . . . .	38
<b>14</b>	<b>7. Continuous Distributions</b>	<b>39</b>
14.1	7.1 Normal Distribution . . . . .	39
14.1.1	Empirical Rule: . . . . .	39
14.2	7.2 Standard Normal Distribution . . . . .	39
14.2.1	Example: . . . . .	40
<b>15</b>	<b>8. Visualizing Data</b>	<b>41</b>
15.1	8.1 Frequency Distribution . . . . .	41
15.2	8.2 Histogram . . . . .	41
15.3	8.3 Boxplot (Box-and-Whisker Plot) . . . . .	41
15.4	8.4 Scatter Plot . . . . .	42
<b>16</b>	<b>9. Practical Applications</b>	<b>43</b>
16.1	9.1 Business Use Cases . . . . .	43
16.2	9.2 Education and Research . . . . .	43
<b>17</b>	<b>10. Using RKWard</b>	<b>44</b>
17.1	10.1 What is RKWard? . . . . .	44
17.2	10.2 Installation Guide . . . . .	44
17.3	10.3 Sample RKWard Activities . . . . .	44
17.3.1	Calculate Mean and SD . . . . .	44
17.3.2	Visualize Histogram . . . . .	44
17.4	10.4 Using R Code in RKWard . . . . .	44
<b>18</b>	<b>Week 3</b>	<b>46</b>
<b>19</b>	<b>Introduction</b>	<b>47</b>
19.1	Importance of Statistics . . . . .	47
19.2	Overview of Topics . . . . .	47
<b>20</b>	<b>Understanding Populations and Samples</b>	<b>48</b>
20.1	Population . . . . .	48
20.2	Sample . . . . .	48
20.3	Why Use Samples? . . . . .	48
20.4	Relation Between Population & Sample . . . . .	48
<b>21</b>	<b>Hypotheses and Errors</b>	<b>49</b>
21.1	Hypothesis Defined . . . . .	49
21.1.1	Null Hypothesis ( $H_0$ ) . . . . .	49

21.1.2	Alternative Hypothesis ( $H_A$ )	49
21.2	Types of Errors	49
21.3	Significance Level ( )	49
<b>22</b>	<b>Inferential Statistics</b>	<b>50</b>
22.1	Purpose	50
22.2	Common Techniques	50
22.3	Sampling Techniques	50
22.3.1	1. Simple Random Sampling	50
22.3.2	2. Systematic Sampling	50
22.3.3	3. Stratified Sampling	50
22.3.4	4. Cluster Sampling	50
22.4	Central Limit Theorem (CLT)	51
<b>23</b>	<b>Descriptive Statistics</b>	<b>52</b>
23.1	Measures of Central Tendency	52
23.1.1	Mean	52
23.1.2	Median	52
23.1.3	Mode	52
23.2	Measures of Dispersion	52
23.2.1	Range	52
23.2.2	Variance	52
23.2.3	Standard Deviation	52
23.3	Measures of Shape	53
<b>24</b>	<b>Graphical Methods</b>	<b>54</b>
24.1	Histogram	54
24.2	R Data Types and Structures	54
24.3	Comparing R vs Excel vs GUI-R (RKWard)	54
24.4	Installing RKWard (Ubuntu)	55
24.5	Teaching Tools in RKWard	55
24.6	GUI-Based Statistical Tools	55
<b>25</b>	<b>Linear Regression in R</b>	<b>56</b>
25.1	What is Linear Regression?	56
25.1.1	Simple Linear Regression Equation:	56
25.2	Code Example	56
<b>26</b>	<b>Fit model</b>	<b>57</b>
<b>27</b>	<b>Summary</b>	<b>58</b>
27.1	Adjusted R-squared	58
27.2	Normal Distribution	58
27.3	Data Import Techniques	58
27.4	Working with the RKWard Interface	58
27.5	Spreadsheet Concepts	59
27.6	Advantages	59
27.7	Limitations	59
27.8	Advanced Plots and Techniques	59

27.9 Common R Packages for Statistics . . . . .	61
27.10 Introduction to Command Line . . . . .	61
27.11 Windows Terminal . . . . .	61
27.12 Git + R Project Example . . . . .	62
27.13 R Script Template for Analysis . . . . .	62
<b>28 Load data</b>	<b>63</b>
<b>29 Descriptive stats</b>	<b>64</b>
<b>30 Histogram</b>	<b>65</b>
<b>31 Linear regression</b>	<b>66</b>
<b>32 Scatter plot with regression line</b>	<b>67</b>
32.1 Future Applications of Statistics . . . . .	67
32.2 Practice Challenges . . . . .	67
32.3 Key Takeaways . . . . .	67
<b>33 Week 4</b>	<b>68</b>
<b>34 Introduction</b>	<b>69</b>
<b>35 Course Overview</b>	<b>70</b>
35.1 Course Name . . . . .	70
35.2 Instructor Profile . . . . .	70
35.3 Learning Objectives . . . . .	70
<b>36 Chapter 1: Fundamental Concepts</b>	<b>71</b>
36.1 Descriptive Statistics . . . . .	71
36.1.1 Central Tendency . . . . .	71
36.1.2 Dispersion . . . . .	71
36.2 Standard Error . . . . .	72
36.3 Central Limit Theorem . . . . .	72
36.4 Confidence Intervals . . . . .	72
<b>37 Chapter 2: Estimation</b>	<b>73</b>
37.1 Types of Estimates . . . . .	73
37.2 Parameter vs Statistic . . . . .	73
<b>38 Chapter 3: Hypothesis Testing</b>	<b>74</b>
<b>39 Chapter 4: Student's T-Test</b>	<b>75</b>
39.1 Types . . . . .	75
39.2 One-Sample T-Test Example . . . . .	75
39.3 Test Statistic . . . . .	75
39.4 Degrees of Freedom . . . . .	75
39.5 Decision Rule . . . . .	76
39.6 T-Test in GUI-R . . . . .	76

<b>40 Chapter 5: ANOVA</b>	<b>77</b>
40.1 Purpose . . . . .	77
40.1.1 One-Way ANOVA Formula . . . . .	77
40.1.2 Assumptions . . . . .	77
40.1.3 Example Table . . . . .	77
40.2 Post-Hoc Tests . . . . .	77
40.3 ANOVA in GUI-R . . . . .	78
<b>41 Chapter 6: GUI-R Workflow</b>	<b>79</b>
<b>42 Chapter 7: Advanced Concepts</b>	<b>80</b>
42.1 Variance Partitioning . . . . .	80
42.2 Degrees of Freedom . . . . .	80
42.3 Chi-Square and F Distribution . . . . .	80
42.4 Univariate, Bivariate, Multivariate . . . . .	80
42.5 Parametric Test Assumptions . . . . .	81
42.6 Effect Size . . . . .	81
42.7 Power of a Test . . . . .	81
<b>43 Conclusion</b>	<b>82</b>
<b>44 References</b>	<b>83</b>
<b>45 Chapter 8: Advanced T-Test Applications</b>	<b>84</b>
45.1 Paired Sample T-Test . . . . .	84
45.1.1 Example: . . . . .	84
45.2 Independent Samples T-Test . . . . .	84
45.3 One-Sample T-Test with GUI-R . . . . .	85
<b>46 Chapter 9: More on Confidence Intervals</b>	<b>86</b>
46.1 Visualizing Confidence Intervals in R . . . . .	86
<b>47 Chapter 10: Robust ANOVA Models</b>	<b>87</b>
47.1 Two-Way ANOVA . . . . .	87
47.2 Repeated Measures ANOVA . . . . .	87
<b>48 Chapter 11: Effect Size Measures</b>	<b>89</b>
48.1 Cohen's d . . . . .	89
48.1.1 R Example . . . . .	89
48.2 Eta-Squared ( $\eta^2$ ) . . . . .	89
<b>49 Chapter 12: Statistical Assumptions Checking</b>	<b>90</b>
49.1 Normality . . . . .	90
49.2 Homogeneity of Variance . . . . .	91
<b>50 Chapter 13: Non-Parametric Alternatives</b>	<b>92</b>
50.1 Wilcoxon Signed Rank Test . . . . .	92
50.2 Mann-Whitney U Test . . . . .	92
50.3 Kruskal-Wallis Test . . . . .	92

<b>51 Chapter 14: Visualizing Statistical Results</b>	<b>94</b>
51.1 Boxplots . . . . .	94
51.2 Histograms . . . . .	94
51.3 Density Plot . . . . .	95
<b>52 Chapter 15: RKWard (GUI-R) Tips</b>	<b>96</b>
52.1 Summary: Basic Statistics using GUI-R (RKWard) . . . . .	96



# 1 Introduction

Introduction

**DR.Harsh Pradhan**, Phone: +91-9930034241 , Email: [harsh.231284@gmail.com](mailto:harsh.231284@gmail.com), Institute of Management Studies, Banaras Hindu University, Address: 18-GF, Jaipuria Enclave, Kaushambhi, Ghaziabad, India, 201010

**Interest:** Goal Orientation Job Performance Consumer Behavior Behavioral Finance Bibliometric Analysis Options as Derivatives Statistics Indian Knowledge System,

[Orcid ID](#)

[Google Scholar](#)

[GitHub](#)

[Researcher ID](#)

[Personal Website](#)

[Youtube ID](#)

**Doing a PhD with me:** [README.1st](#)

[Academic Profile](#)

---

Topics to be covered:

1. Basic probability theory, random variable theory (including jointly distributed RVs), probability distributions (including bivariate distributions)
2. Using Bayes' rule for statistical inference
3. An introduction to (generalized) linear models
4. An introduction to hierarchical models
5. Measurement error models
6. Mixture models
7. Model selection and hypothesis testing (Bayes factor and k-fold cross-validation)

## 1.1 Teaching

Science and statistics is/are one unitary thing; you cannot do one without the other. Towards this end, I teach some (in my opinion) critically important classes that provide a solid statistical foundation for doing research in cognitive science.

Courses offered:

1. Free online course, four weeks (MOOC), enrollments open: Introduction to Bayesian Data Analysis
2. Short (four-hour) tutorial on Bayesian statistics, taught at EMLAR 2022: [here](#)
3. Introduction to (frequentist) statistics
4. Introduction to Bayesian data analysis for cognitive science
5. BDA cover

## 1.2 Lecture notes

Download from [here](#).

## 1.3 Moodle website

All communications with students in Potsdam will be done through [this website](#). # Schedule

Week	Lecture	Main Topic	Subtopic	Video	PDF Resource
Week 1	2	Descriptive Statistics	Central Tendency	<a href="#">Video</a>	<a href="#">Week 2.pdf</a>
	2	Descriptive Statistics	Measure of Variability	<a href="#">Video</a>	Same as above
	3	Descriptive Statistics	Describing Data	<a href="#">Video</a>	Same as above
	4	Descriptive Statistics	Probability	<a href="#">Video</a>	Same as above
	5	Descriptive Statistics	Distribution	<a href="#">Video</a>	Same as above
Week 3	1	Descriptive Statistics	Z Table (Normal Distribution)	<a href="#">Video</a>	<a href="#">Week 3.pdf</a>
	2	Descriptive Statistics	Measuring Divergence	<a href="#">Video</a>	Same as above
	3	Inferential Statistics	Sample and Population	<a href="#">Video</a>	Same as above
	4	Inferential Statistics	Model Fit	<a href="#">Video</a>	Same as above

Week	Lecture	Main Topic	Subtopic	Video	PDF Resource
	5	Inferential Statistics	Hypothesis and Error	<a href="#">Video</a>	Same as above
Week 4	1	Terms of Statistics	Terms of Statistics	<a href="#">Video</a>	<a href="#">Week 4.pdf</a>
	2	Terms of Statistics	T-Test	<a href="#">Video</a>	Same as above
	3	Terms of Statistics	T-Test in Detail	<a href="#">Video</a>	Same as above
	4	ANOVA	ANOVA	<a href="#">Video</a>	Same as above
Week 5	1	ANOVA	Example of ANOVA	<a href="#">Video</a>	<a href="#">Week 5.pdf</a>
	2	ANOVA	Types of ANOVA	<a href="#">Video</a>	Same as above
	3	Correlation	Introduction to Correlation	<a href="#">Video</a>	Same as above
	4	Correlation	Regression (Part 1)	<a href="#">Video</a>	Same as above
	5	Correlation	Regression (Part 2)	<a href="#">Video</a>	Same as above
Week 6	1	Correlation	R Script for Regression	<a href="#">Video</a>	<a href="#">Week 6.pdf</a>
	2	Chi Square	Chi Square	<a href="#">Video</a>	Same as above
	3	Chi Square	Chi Square Test	<a href="#">Video</a>	Same as above
	4	Logistic Function	Regression Function	<a href="#">Video</a>	Same as above
	5	Logistic Function	Distribution	<a href="#">Video</a>	Same as above
Week 7	1	Time Series	Intro to Time Series	<a href="#">Video</a>	<a href="#">Week 7.pdf</a>
	2	Time Series	Conditional Probability	<a href="#">Video</a>	Same as above
	3	Time Series	Additional Concepts	<a href="#">Video</a>	Same as above
	4	Time Series	Distribution	<a href="#">Video</a>	Same as above
	5	Time Series	Poisson Distribution	<a href="#">Video</a>	Same as above

<b>Week</b>	<b>Lecture</b>	<b>Main Topic</b>	<b>Subtopic</b>	<b>Video</b>	<b>PDF Resource</b>
	6	Index Numbers	Price & Quantity Index	<a href="#">Video</a>	Same as above
	7	Decision Environments	Risk/Uncertainty, Bayes, Trees	<a href="#">Video</a>	Same as above
	8	Time Series Analysis	Components, Trend, Seasonality	<a href="#">Video</a>	Same as above
	9	Time Series Analysis	Least Squares Method	<a href="#">Video</a>	Same as above
Week 8	1	Effect Size & Documentation	Package/Library	<a href="#">Video</a>	<a href="#">Week 8.pdf</a>
	2	Effect Size & Documentation	RStudio vs RKward	<a href="#">Video</a>	Same as above
	3	Effect Size & Documentation	Flexplot	<a href="#">Video</a>	Same as above
	4	Effect Size & Documentation	Functions	<a href="#">Video</a>	Same as above
	5	Effect Size & Documentation	R Shiny & R Markdown	<a href="#">Video</a>	Same as above
	6	Effect Size & Documentation	Application with Real Datasets	<a href="#">Video</a>	Same as above

<b>Week</b>	<b>Main Lecture Topic</b>	<b>Subtopic</b>	<b>Video</b>	<b>PDF Resource</b>
7	Effect Size & Interpre- tation	Importance in Testing	<a href="#">Video</a>	Same as above
8	Effect Size & Interpre- tation	Installing dplyr, ggplot2	<a href="#">Video</a>	Same as above
9	Effect Size & Interpre- tation	Visual Model Interpreta- tion	<a href="#">Video</a>	Same as above
10	Effect Size & Interpre- tation	Creating/Using Functions	<a href="#">Video</a>	Same as above
11	Effect Size & Interpre- tation	Report, Dashboard, Interactivity	<a href="#">Video</a>	Same as above

# 2 Week 1

## 2.1 Module 1: Introduction to Statistics

### Pre-Requisites

- Just an open and eager mind
- Basic understanding of Mathematics or Statistics

### Agenda

- Meaning of Statistics
  - Nature and Scope
  - Uses of Statistics
  - Limitations
  - Fallacies and Misuse
  - Math vs Statistics
  - GUI Tools & Transition to Software-based Stats
- 

### Meaning of Statistics

Statistics is a science which provides tools for **analysis and interpretation** of raw data collected for decision-making in diverse fields.

It includes four core concepts:

- **Population** – Complete data or total group
- **Sample** – Subset of population
- **Parameter** – Numerical summary from population
- **Statistic** – Numerical summary from sample

### Nature of Statistics

- Deals with **numerical facts**
  - Focused on **social phenomena** and real-world data
  - Organizes, classifies, and analyzes data
  - Facilitates **prediction, interpretation, and decision-making**
- 

### Uses of Statistics

- Drawing representative samples
- Summarizing collected data
- Tabulation and systematic arrangement
- Group comparisons
- Determining behavioral relationships
- Estimating chance vs causation
- Application in:
  - Psychology
  - Education
  - Employment surveys
  - Market Research
  - Industrial and Organizational studies

#### Limitations of Statistics

- Cannot study **qualitative phenomena** without quantification
- Not applicable to individuals
- **Statistical laws are not exact**
- Does not guarantee **causal relationships**
- Vulnerable to misuse

## 2.2 Misuse of Statistics

- Use of extremely **small or biased** samples
- **Misleading graphs** or visual misrepresentation
- Illogical or **unexpected comparisons**

#### Fallacies in Statistics

Fallacies may arise from:

- Poor data collection methods
- Vague or manipulated term definitions
- Improper unit selection
- Faulty classification or grouping
- Inappropriate statistical methods

### 3 Module 2: Mathematics vs Statistics

Aspect	Mathematics	Statistics
Nature	Abstract, symbolic reasoning	Applied, data-based reasoning
Focus	Pure logic, proofs	Real-world data, decision-making
Techniques	Algebra, Calculus, Geometry	Probability, Hypothesis testing, Regression
Output	Theorems, functions, formulas	Inferences, predictions, summaries
Tools	Equations, graphs	Charts, tables, models



## 4 Module 3: Software-Based Statistical Revolution

### 4.1 From Paper to Code

Why shift to software?

- **Faster analysis** of massive data
- **Error-free calculations**
- **Anywhere-anytime** access
- **Cloud-based integration**
- Supports **ML/AI**, automation, and deep visualization

### 4.2 Popular Statistical Software

Software	Type	Use Case
R	Script	Core for academic and professional stats
RKward	GUI	GUI wrapper for R
R Commander	GUI	Menu-based GUI for R
Rattle	GUI	Data mining toolkit in R
Excel	GUI	Basic stats with plugins
Python (pandas)	Script	Modern data science + ML

### 4.3 GUI vs CLI

Feature	GUI (e.g., RKward)	Command Line (e.g., R Console)
Accessibility	User-friendly	Requires learning syntax
Speed	Slower for heavy tasks	High performance
Learning Curve	Minimal	Moderate to High
Customization	Limited	Fully scriptable
Teaching Utility	Good for beginners	Good for understanding logic

## 4.4 Recommended GUI Tools for R

- RKWard
- Rattle
- R Commander
- R AnalyticFlow

<https://rkward.kde.org>

## 4.5 Installing RKWard on Ubuntu

bash sudo apt install kbibtex kate libcurl4-openssl-dev libssl-dev libxml2-dev cmake sudo add-apt-repository ppa:rkward-devel/rkward-stable echo "deb https://ppa.launchpad.net/rkward-devel/rkward-stable/ubuntu jammy main" | sudo tee /etc/apt/sources.list.d/rkward.list sudo apt update sudo apt-get install rkward Awesome. Here's Part 2 of the full markdown, Lines 251–600, continuing the structured content from your Week 1 lecture.

# 5 Module 4: Understanding Variables

## 5.1 What is a Variable?

A **variable** is a characteristic or attribute that can assume different values across individuals or items.

In statistics, variables are categorized for analysis and measurement.

### 5.1.1 R Definition:

In R, variables are containers for data, created by assignment:

```
x <- 10
name <- "Harsh"
flag <- TRUE
```

#### Classification of Variables

##### A. Qualitative (Categorical)

Type	Description	Example
------	-------------	---------

Nominal	Categories without order	Gender (Male, Female)
---------	--------------------------	-----------------------

Ordinal	Categories with a meaningful order	Education Level (UG, PG)
---------	------------------------------------	--------------------------

##### B. Quantitative (Numerical)

Type	Description	Example
------	-------------	---------

Discrete	Countable numbers	No. of students
----------	-------------------	-----------------

Continuous	Infinite values in a range	Height, Weight
------------	----------------------------	----------------

#### Statistical Data Types (Scale of Measurement)

Data Type	Description	Examples
-----------	-------------	----------

Nominal	Categories with no order	Blood group (A, B, AB, O)
---------	--------------------------	---------------------------

Ordinal	Ranked categories	Satisfaction (Low, Med, High)
---------	-------------------	-------------------------------

Interval	Numeric scale with no true zero	Temperature in Celsius
Ratio	Numeric scale with true zero	Income, Weight, Age

## Data Types in R

R Type	Description	Example Code
--------	-------------	--------------

Numeric	Real numbers	<code>x &lt;- 15.3</code>
Integer	Whole numbers	<code>y &lt;- as.integer(10)</code>
Complex	Real + imaginary	<code>z &lt;- 2+3i</code>
Character	Text strings	<code>c &lt;- "hello"</code>
Logical	Boolean values	<code>b &lt;- TRUE</code>
Factor	Categorical encoding	<code>factor(c("yes", "no", "yes"))</code>

```
# Examples in R
x <- 15.6
y <- as.integer(18)
z <- 7 + 5i
c <- "I am OK"
b <- TRUE
```

## Module 5: Data Structures in R

### Vectors

A vector is a one-dimensional array of elements.

```
vec1 <- c(5, 2, 3, 7, 8, 9, 1, 4, 10, 15)
```

### Matrices

Two-dimensional arrays of rows and columns.

```
mat <- matrix(1:9, nrow=3, ncol=3)
```

### Arrays

Multidimensional generalization of matrices.

```
arr <- array(1:24, dim=c(3,4,2))
```

### Lists

Collection of different types of elements.

```
mylist <- list(name="Alice", age=30, scores=c(89,90))
```

## Data Frames

Tabular data (like a spreadsheet), each column can have a different type.

```
df <- data.frame(ID=1:3, Name=c("A", "B", "C"), Score=c(85, 90, 95))
```

## Factors

Used for categorical variables.

```
gender <- factor(c("Male", "Female", "Male"))
```

## Module 6: Descriptive Statistics

Descriptive statistics summarize and simplify data.

### Central Tendency

#### Measure Formula Meaning

Mean	$\bar{x} = \frac{\sum x_i}{n}$	Average
Median	Middle value in sorted data	Central observation
Mode	Most frequent value	Most common observation

### Dispersion Measures

#### Measure Formula Purpose

Range	$\text{Range} = \text{Max} - \text{Min}$	Spread of data
Variance	$s^2 = \frac{\sum (x_i - \bar{x})^2}{n - 1}$	Spread from mean
Standard Deviation	$s = \sqrt{\text{Variance}}$	Average distance from mean

### Example in R

```
x <- c(10, 20, 30, 40, 50)
mean(x)
median(x)
var(x)
sd(x)
```

## Module 7: Inferential Statistics

Inferential stats allow us to make conclusions about populations using samples.

## Key Concepts

Hypothesis Testing: Assesses assumptions about a population.

Confidence Intervals: Estimate population parameters within a range.

Significance Levels ( ): Commonly 0.05 or 5%

P-Value: Probability of observing the data assuming the null is true.

## Hypothesis Types

Type	Description
------	-------------

Null Hypothesis	No difference / no effect
-----------------	---------------------------

Alternative	There is a difference / effect
-------------	--------------------------------

## R Examples

```
t.test(x)           # One-sample t-test
t.test(x, y)        # Two-sample t-test
```

## Module 8: Visualizing Data

Data visualization helps uncover patterns and insights.

### Boxplot

Shows 5-number summary

Identifies outliers

```
boxplot(x)
```

### Histogram

Frequency distribution of continuous data

```
hist(x)
```

### Pie Chart

Shows proportion in categories

```
slices <- c(10, 12, 4, 16, 8)
labels <- c("A", "B", "C", "D", "E")
pie(slices, labels=labels)
```

Scatter Plot

Relationship between two variables

```
plot(x, y)
```

Ogive (Cumulative Frequency)

```
# Create cumulative frequency table manually
```

Module 9: Spreadsheet Basics

Spreadsheets like Excel or Google Sheets are entry points for data work.

Key Features:

Rows → Observations

Columns → Variables

Supports sorting, filtering

Built-in formulas: =SUM(), =AVERAGE(), etc.

Spreadsheets vs R

Feature	Spreadsheet (Excel, GSheets)	R / Rkward
---------	------------------------------	------------

Cost	Usually licensed	Free and open source
------	------------------	----------------------

Flexibility	Limited to GUI formulas	Full programming capability
-------------	-------------------------	-----------------------------

Graphics	Basic	Advanced (ggplot2)
----------	-------	--------------------

Reproducibility	Low	High (script-based)
-----------------	-----	---------------------

Module 10: Command Line vs GUI

Command Line (R Console)

```
# Windows Command Line
```

```
cd ..
```

```
mkdir new_folder
```

```
dir
```

## R Console Commands

```
getwd()
setwd("path")
install.packages("ggplot2")
library(ggplot2)
```

## GUI (RKWard)

Point-and-click interface

No coding needed

View script history and console

Menu for graphs, models, tables

## Learning Resources:

### Books

Mohanty, B., & Misra, S. (2016). Statistics for Behavioural and Social Sciences

Pandya et al. (2018). Statistical Analysis in Simple Steps using R

Field, A. P. et al. (2012). Discovering Statistics using R

Harris, J. K. (2019) . Statistics with R: Solving Problems using Real-World Data

### *## Utilizing Statistical Methods for Decision Making*

- Use statistical evidence to guide business strategies.
- Make informed policy decisions based on empirical data.
- Report findings clearly for transparency and comprehension.

### *## Summary*

The "Basic Statistics Using GUI-R (RK Ward)" course equips learners with the foundational and

### *## Key Takeaways*

- Proficiency in defining and using variables and data types.
- Capability to import and manipulate data in RKWard.
- Understanding of basic statistical practices and their applications.
- Skill in visualizing data for effective communication of results.

### *## Websites*



```
https://rkward.kde.org  
https://r4stats.com  
https://cran.r-project.org
```

```
`<!-- quarto-file-metadata: eyJyZXNvdXJjZURpciI6Ii4ifQ== -->`{=html}
```

```
```${=html}
```

```
<!-- quarto-file-metadata: eyJyZXNvdXJjZURpciI6Ii4iLCJib29rSXRlbVR5cGUiOiJjaGFwdGVyIiwiaW9va01
```

## **6 Week 2**

# 7 Introduction

## 7.1 Purpose of the eBook

This eBook is designed as a complete beginner-to-intermediate guide for understanding the foundational concepts of statistics. It aims to bridge theoretical knowledge and practical application using RKWard (a GUI for R). Readers will be introduced to descriptive and inferential statistics, probability theory, and probability distributions with ample examples and exercises.

## 7.2 Who Should Read This?

- Undergraduate students
- MBA and management students
- Data analysis beginners
- Professionals dealing with data

## 7.3 What You'll Learn

- Data classification and types
  - Descriptive statistics: central tendency and variability
  - Basic probability and events
  - Probability distributions: Bernoulli, Binomial, and Normal
  - Use of RKWard in statistical analysis
-

# 8 1. Fundamentals of Statistics

## 8.1 1.1 What is Statistics?

Statistics is the science of collecting, organizing, analyzing, and interpreting data to make informed decisions. It involves both **theoretical** (mathematical) and **applied** approaches to understanding uncertainty and variability in real-world phenomena.

## 8.2 1.2 Key Objectives

- Summarizing large datasets effectively
- Estimating population parameters
- Testing hypotheses
- Making predictions and decisions under uncertainty

## 8.3 1.3 Types of Statistics

- **Descriptive Statistics:** Deals with the presentation and summarization of data.
  - **Inferential Statistics:** Draws conclusions about populations based on sample data.
-

## 9 2. Types of Data

### 9.1 2.1 Classification of Data

Type	Example	Description
Qualitative	Gender, Nationality	Non-numeric labels
Quantitative	Height, Age	Numeric values
Discrete	No. of Children	Countable numbers
Continuous	Temperature, Weight	Infinite values in a range

#### 9.1.1 2.1.1 Qualitative (Categorical) Data

- **Nominal:** No inherent order (e.g., religion, marital status).
- **Ordinal:** Natural order (e.g., customer satisfaction: Poor, Average, Good).

#### 9.1.2 2.1.2 Quantitative (Numerical) Data

- **Discrete:** Integers; e.g., number of books.
  - **Continuous:** Measurable; e.g., weight in kilograms.
-

## 10 3. Descriptive Statistics

### 10.1 3.1 Measures of Central Tendency

#### 10.1.1 3.1.1 What is Central Tendency?

Central tendency refers to the center or middle of a dataset. It's the value that best represents the entire distribution.

#### 10.1.2 3.1.2 Characteristics of a Good Measure

- Rigidly defined
  - Easy to understand
  - Takes all data into account
  - Amenable to algebraic treatment
  - Stable under sampling
  - Minimally affected by outliers (except mean)
- 

### 10.2 3.2 The Mean

#### 10.2.1 3.2.1 Definition

The arithmetic mean is the sum of all values divided by the number of values.

#### 10.2.2 3.2.2 Formula

$$\bar{x} = \frac{\sum_{i=1}^n x_i}{n}$$

#### 10.2.3 3.2.3 Properties of Mean

- Uses all data values
- Affected by extreme values
- The sum of deviations from the mean is zero

### 10.2.4 3.2.4 Example

Data: 10, 15, 20, 25, 30

Mean =  $(10 + 15 + 20 + 25 + 30)/5 = 20$

---

## 10.3 3.3 The Median

### 10.3.1 3.3.1 Definition

The median is the value separating the higher half from the lower half of a data sample.

### 10.3.2 3.3.2 Calculation

- Odd number of items: Middle value
- Even number of items: Average of the two middle values

### 10.3.3 3.3.3 Properties

- Not influenced by extreme values
- Best for skewed data

### 10.3.4 3.3.4 Example

Data: 4, 6, 9, 12, 15, 21, 33

Median = 12 (middle value)

---

## 10.4 3.4 The Mode

### 10.4.1 3.4.1 Definition

The mode is the value that appears most frequently in a dataset.

### 10.4.2 3.4.2 Characteristics

- Can be used for categorical data
- Dataset can be unimodal, bimodal, or multimodal
- May not exist if all values are unique

### 10.4.3 3.4.3 Example

Data: 4, 4, 6, 8, 9, 10, 4

Mode = 4

---

## 10.5 3.5 Comparison Table

Measure	Use Case	Affected by Outliers	Mathematical Use
Mean	Symmetric distributions	Yes	High
Median	Skewed distributions	No	Moderate
Mode	Categorical variables	No	Low

---



# 11 4. Measures of Variability

## 11.1 4.1 Why Measure Variability?

While central tendency summarizes data, variability tells us how spread out the data is. It's essential in determining consistency and reliability.

---

## 11.2 4.2 Range

### 11.2.1 4.2.1 Definition

The difference between the maximum and minimum values.

$$\text{Range} = x_{\max} - x_{\min}$$

### 11.2.2 4.2.2 Example

Data: 12, 14, 17, 19, 23

Range = 23 - 12 = 11

### 11.2.3 4.2.3 Limitations

- Ignores distribution shape
  - Extremely sensitive to outliers
- 

## 11.3 4.3 Quartiles and Interquartile Range

### 11.3.1 4.3.1 Quartiles

- Q1 (25th percentile): Lower quartile
- Q2 (50th percentile): Median
- Q3 (75th percentile): Upper quartile

### 11.3.2 4.3.2 Formula for Position

$$Q_k = \frac{k(n+1)}{4}$$

### 11.3.3 4.3.3 IQR Formula

$$IQR = Q3 - Q1$$

### 11.3.4 4.3.4 Example

Data: 12, 30, 45, 57, 70

$Q1 = 30, Q3 = 57 \rightarrow IQR = 27$

---

## 11.4 4.4 Variance

### 11.4.1 4.4.1 Concept

Variance is the average of the squared differences from the Mean.

### 11.4.2 4.4.2 Formulas

Population Variance:

$$\sigma^2 = \frac{1}{N} \sum (x_i - \mu)^2$$

Sample Variance:

$$s^2 = \frac{1}{n-1} \sum (x_i - \bar{x})^2$$

---

## 11.5 4.5 Standard Deviation

### 11.5.1 4.5.1 Concept

Standard deviation is the square root of variance. It provides a measure of spread in the same units as the data.

$$s = \sqrt{s^2}$$

### 11.5.2 4.5.2 Properties

- Same unit as original data
  - Measures how far values deviate from the mean
  - Widely used in most statistical computations
- 

## 11.6 4.6 Coefficient of Variation (CV)

### 11.6.1 4.6.1 Definition

The ratio of the standard deviation to the mean, expressed as a percentage. Used to compare variability between datasets with different units.

$$CV = \left( \frac{s}{\bar{x}} \right) \times 100\%$$

### 11.6.2 4.6.2 Example

Dataset A: Mean = 100, SD = 10  $\rightarrow$  CV = 10%

Dataset B: Mean = 50, SD = 5  $\rightarrow$  CV = 10%

---

## 11.7 4.7 Moment-Based Measures

- First Moment (about mean): 0 (since  $\sum(x - \bar{x}) = 0$ )
  - Second Moment: Variance
  - Third Moment: Skewness
  - Fourth Moment: Kurtosis
-

## 12 5. Probability Fundamentals

### 12.1 5.1 Introduction to Probability

Probability is the mathematical framework for quantifying uncertainty. It helps us estimate how likely an event is to occur.

### 12.2 5.2 Key Definitions

- **Experiment:** A process that leads to an outcome.
  - **Outcome:** The result of an experiment.
  - **Sample Space ( $\Omega$ ):** All possible outcomes.
  - **Event:** A subset of the sample space.
- 

### 12.3 5.3 Types of Events

---

Event Type	Description
Independent	Occurrence of one does not affect the other
Dependent	One affects the outcome of another
Mutually Exclusive	Cannot occur together
Exhaustive	Includes all possible outcomes

---

### 12.4 5.4 Classical Probability

Used when all outcomes are equally likely.

**Formula:**

$$P(A) = \frac{\text{Number of favorable outcomes}}{\text{Total outcomes in } \Omega}$$

**Example:** Rolling a fair die

$$P(\text{rolling a 3}) = 1/6$$

---

## 12.5 5.5 Probability Rules

### 12.5.1 Rule 1: Non-Negativity

$$0 \leq P(A) \leq 1$$

### 12.5.2 Rule 2: Total Probability

$$P(\Omega) = 1$$

### 12.5.3 Rule 3: Complement Rule

$$P(A^c) = 1 - P(A)$$

### 12.5.4 Rule 4: Addition Rule

If A and B are mutually exclusive:

$$P(A \cup B) = P(A) + P(B)$$

Otherwise:

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

### 12.5.5 Rule 5: Multiplication Rule

- For independent events:

$$P(A \cap B) = P(A) \cdot P(B)$$

---

## 12.6 5.6 Conditional Probability

Formula:

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

---

## 13 6. Discrete Probability Distributions

### 13.1 6.1 Bernoulli Distribution

- One trial, two outcomes (success/failure).
- Success = 1, Failure = 0

$$P(X = x) = p^x(1 - p)^{1-x}, \quad x \in \{0, 1\}$$

- Mean =  $p$
- Variance =  $p(1 - p)$

#### 13.1.1 Example:

Flip a fair coin  $\rightarrow p = 0.5$

Mean = 0.5, Variance = 0.25

---

### 13.2 6.2 Binomial Distribution

- Series of  $n$  independent Bernoulli trials
- Number of successes  $x$  out of  $n$  trials

**Formula:**

$$P(X = x) = \binom{n}{x} p^x (1 - p)^{n-x}$$

- Mean:  $\mu = np$
- Variance:  $\sigma^2 = np(1 - p)$

#### 13.2.1 Example:

Flip a coin 5 times ( $p = 0.5$ )

$$P(X = 3) = \binom{5}{3} (0.5)^3 (0.5)^2 = 10 \cdot 0.125 \cdot 0.25 = 0.3125$$

---

# 14 7. Continuous Distributions

## 14.1 7.1 Normal Distribution

The most important continuous distribution in statistics.

### Properties:

- Bell-shaped and symmetric
- Defined by mean (  $\mu$  ) and variance (  $\sigma^2$  )
- Total area under the curve = 1

### Probability Density Function (PDF):

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right)$$

#### 14.1.1 Empirical Rule:

- 68% of values lie within  $\pm 1$
  - 95% within  $\pm 2$
  - 99.7% within  $\pm 3$
- 

## 14.2 7.2 Standard Normal Distribution

A normal distribution with:

- Mean = 0
- Standard deviation = 1

### Z-score Formula:

$$Z = \frac{X - \mu}{\sigma}$$

### 14.2.1 Example:

If  $\mu = 100$ ,  $\sigma = 15$ , and  $X = 130$

Then  $Z = \frac{130-100}{15} = 2$

---



## 15 8. Visualizing Data

### 15.1 8.1 Frequency Distribution

Class Interval	Frequency
0–10	3
11–20	7
21–30	9
31–40	6

---

### 15.2 8.2 Histogram

A bar chart representing the frequency distribution of numerical data.

**Use Case:** Visualize shape (e.g., normal, skewed)

---

### 15.3 8.3 Boxplot (Box-and-Whisker Plot)

Shows:

- Minimum
- Q1
- Median
- Q3
- Maximum
- Outliers (as dots)

Helps identify skewness and outliers quickly.

---

## 15.4 8.4 Scatter Plot

Used to study the relationship between two quantitative variables.

---

# 16 9. Practical Applications

## 16.1 9.1 Business Use Cases

- Retail: Analyze sales patterns
  - Healthcare: Patient outcome probabilities
  - Finance: Stock volatility (using SD, CV)
- 

## 16.2 9.2 Education and Research

- Student test scores: Use mean, SD, and percentile ranking
  - Experiment analysis: Use Z-scores and Normal Distribution
-

## 17 10. Using RKWard

### 17.1 10.1 What is RKWard?

A graphical frontend for the R programming language designed for statistical analysis and data visualization.

---

### 17.2 10.2 Installation Guide

1. Download R from [CRAN](#)
  2. Install RKWard from [rkward.kde.org](#)
  3. Start RKWard and begin with menu-driven tasks
- 

### 17.3 10.3 Sample RKWard Activities

#### 17.3.1 Calculate Mean and SD

- Load dataset
- Click *Statistics* → *Descriptive Statistics*
- Choose variables and click *OK*

#### 17.3.2 Visualize Histogram

- Click *Graphics* → *Histogram*
  - Select variable and customize bins
- 

### 17.4 10.4 Using R Code in RKWard

```
data <- c(12, 15, 17, 18, 21)
mean(data)
sd(data)
hist(data)
```

*## Summary*

This eBook provided a deep dive into basic statistics including:

- Data types and classification
- Central tendency and variability
- Probability theory and rules
- Discrete and continuous distributions
- Visual interpretation and real-world applications
- GUI-based statistical analysis using RKWard

```
`<!-- quarto-file-metadata: eyJyZXNvdXJjZURpciI6Ii4ifQ== -->`{=html}
```

```
````{=html}
```

```
<!-- quarto-file-metadata: eyJyZXNvdXJjZURpciI6Ii4iLCJib29rSXRlbVR5cGUiOiJjaGFwdGVyIiwiaWYm9va01
```

## 18 Week 3

# 19 Introduction

## 19.1 Importance of Statistics

Statistics is a powerful tool used across disciplines — from economics and psychology to biology, data science, and machine learning. It enables:

- Interpretation of data
- Generalization from samples to populations
- Hypothesis testing and decision-making
- Prediction and modeling

Understanding statistics is essential for anyone involved in **empirical research**, **policy making**, **data-driven decision-making**, or **scientific inquiry**.

---

## 19.2 Overview of Topics

This book covers:

- Population vs Sample
  - Hypotheses and Errors
  - Descriptive vs Inferential Statistics
  - Data Types (R + Theoretical)
  - Sampling Techniques
  - Normal Distribution
  - Linear and Logistic Regression
  - GUI-based R interfaces: RKWard, Rcmdr, Rattle
  - Fallacies and misuse in statistics
  - Graphical Methods
  - R programming constructs for statistics
-

# 20 Understanding Populations and Samples

## 20.1 Population

The complete set of all units of interest. Examples:

- All students in India
- All electric cars in the U.S.

## 20.2 Sample

A **subset of the population**, selected for analysis. Goal: represent the population accurately.

## 20.3 Why Use Samples?

- More practical and cost-efficient
- Enables faster analysis
- Allows estimation and inference

## 20.4 Relation Between Population & Sample

Population → Sample → Statistic → Inference → Population Parameter

---



# 21 Hypotheses and Errors

## 21.1 Hypothesis Defined

A hypothesis is a testable assumption about a population.

### 21.1.1 Null Hypothesis ( $H_0$ )

- No difference or effect
- Example:  $H_0$ : “ = 100”

### 21.1.2 Alternative Hypothesis ( $H_A$ )

- A difference or effect exists
- Example:  $H_A$ : “ ≠ 100”

## 21.2 Types of Errors

Error Type	Description
Type I Error	Rejecting $H_0$ when it's true (false positive)
Type II Error	Failing to reject $H_0$ when it's false (false neg)

## 21.3 Significance Level ( )

The probability of making a Type I error — commonly set to **0.05 (5%)**

# 22 Inferential Statistics

## 22.1 Purpose

- Estimate unknown population parameters
- Test hypotheses
- Predict outcomes

## 22.2 Common Techniques

- t-test
  - z-test
  - ANOVA
  - Chi-square
  - Regression
- 

## 22.3 Sampling Techniques

### 22.3.1 1. Simple Random Sampling

Every unit has equal probability.

### 22.3.2 2. Systematic Sampling

Pick every  $k$ th element.

### 22.3.3 3. Stratified Sampling

Subdivide population into strata (e.g. age groups), then sample from each.

### 22.3.4 4. Cluster Sampling

Randomly choose entire groups (e.g. schools, cities).

---

## 22.4 Central Limit Theorem (CLT)

If  $n > 30$ , the distribution of sample means approximates a **normal distribution** even if the original population is not normal.

Formula:

$$Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$$

---

## 23 Descriptive Statistics

### 23.1 Measures of Central Tendency

#### 23.1.1 Mean

$$\bar{x} = \frac{\sum x_i}{n}$$

#### 23.1.2 Median

Middle value in an ordered dataset.

#### 23.1.3 Mode

Most frequent value.

---

### 23.2 Measures of Dispersion

#### 23.2.1 Range

$$Range = Max - Min$$

#### 23.2.2 Variance

$$s^2 = \frac{\sum (x_i - \bar{x})^2}{n - 1}$$

#### 23.2.3 Standard Deviation

$$s = \sqrt{s^2}$$

---

## 23.3 Measures of Shape

- **Skewness:** Degree of asymmetry
  - **Kurtosis:** Peakedness of distribution
-

## 24 Graphical Methods

### 24.1 Histogram

`r hist(data$height, col="blue", main="Height Distribution")` Boxplot

`boxplot(data$score, data$group)` Scatter Plot

`plot(data$x, data$y, col="red")` Ogive (Cumulative Frequency Plot)

Built using cumulative frequency of class intervals.

### 24.2 R Data Types and Structures

Basic Data Types

`x <- 12.5` # numeric `y <- as.integer(5)` # integer `z <- 4 + 3i` # complex `name <- "Ravi"` # character `flag <- TRUE` # logical Vectors

`v <- c(1, 2, 3)` Matrices

`m <- matrix(1:9, nrow=3, byrow=TRUE)` Data Frame

`df <- data.frame(Name=c("A", "B"), Score=c(89, 94))` Lists

`lst <- list(id=101, name="John", marks=c(78, 82))` Factors

`gender <- factor(c("Male", "Female", "Male"))` Statistical Fallacies

What are Fallacies?

Fallacies occur when conclusions are drawn based on flawed statistical reasoning.

Common Fallacies

Improper Sampling Misleading Graphs Ambiguous Term Definitions Ignoring Confounding Variables Assuming Correlation Implies Causation Misuse of Statistics

Examples of Misuse

Using biased samples Cherry-picking data Using 3D pie charts to exaggerate results Misrepresenting scale in graphs

### 24.3 Comparing R vs Excel vs GUI-R (RKWard)

Feature	R (Script)	Excel	RKWard GUI
Usability	Medium	Easy	Easy
Flexibility	High	Low-Medium	Medium
Statistical Power	Very High	Low	High
Graphics	ggplot2	Basic	ggplot2 supported
Reproducibility	High	Low	High

## 24.4 Installing RKWard (Ubuntu)

```
sudo apt install kbbibtex kate libcurl4-openssl-dev libssl-dev libxml2-dev cmake sudo add-apt-repository ppa:rkward
```

## 24.5 Teaching Tools in RKWard

```
install.packages(c("R2HTML", "car", "e1071", "Hmisc", "plyr", "ggplot2", "prob", "ez", "multcomp", "remotes"), de
```

## 24.6 GUI-Based Statistical Tools

RKWard – KDE interface for R  
 Rcmdr – Classic R Commander GUI  
 Rattle – Data mining GUI  
 in R  
 R AnalyticFlow – Flow-based programming for statistics

## 25 Linear Regression in R

### 25.1 What is Linear Regression?

Linear regression models the relationship between a **dependent variable (Y)** and one or more **independent variables (X)**.

#### 25.1.1 Simple Linear Regression Equation:

$$Y = \beta_0 + \beta_1 X + \epsilon$$

Where:

- $Y$  is the dependent variable
- $X$  is the independent variable
- $\beta_0$  is the intercept
- $\beta_1$  is the slope
- $\epsilon$  is the error term

### 25.2 Code Example

```
r # Load data data(mtcars)
```



## 26 Fit model

```
model <- lm(mpg ~ wt, data=mtcars)
```

## 27 Summary

`summary(model)`

### 27.1 Adjusted R-squared

Penalizes the number of predictors to avoid overfitting.

AIC & BIC

AIC: Akaike Information Criterion BIC: Bayesian Information Criterion Lower values of AIC/BIC  
→ better model fit (with penalty for complexity).

### 27.2 Normal Distribution

Key Properties

Symmetrical, bell-shaped curve Mean = Median = Mode Total area under curve = 1 Empirical  
Rule: 68% within  $\pm 1$  SD 95% within  $\pm 2$  SD 99.7% within  $\pm 3$  SD

Example: Given: Mean = 70, SD = 5, X = 75

`z <- (75 - 70) / 5` # Result: 1.0 Z-Table Usage

Find the area under the curve to the left of the z-score Useful for probability and percentile ranking

### 27.3 Data Import Techniques

CSV Import in R

`df <- read.csv("data.csv", header=TRUE)` head(df) Excel Import (using readxl)

`install.packages("readxl")` library(readxl)

`df <- read_excel("data.xlsx")`

### 27.4 Working with the RKWard Interface

Sections: Console – Run R code Script Editor – Write reusable code Workspace – View loaded variables Teaching Tab – Education-focused modules

## 27.5 Spreadsheet Concepts

Structure

Component | Description Rows | Individual observations Columns | Variables Cells | Data points  
Header Row | Variable names

## 27.6 Advantages

Easy data entry Visual inspection Good for small datasets


## 27.7 Limitations

Limited statistical functionality Hard to reproduce Error-prone for large datasets

## 27.8 Advanced Plots and Techniques


Density Plot

```
plot(density(mtcars$mpg), main="Density Plot", col="blue")
```



Pair Plot

```
pairs(mtcars[, 1:4])
```



Correlation Matrix

```
cor(mtcars)
```

	mpg	cyl	disp	hp	drat	wt
mpg	1.0000000	-0.8521620	-0.8475514	-0.7761684	0.68117191	-0.8676594
cyl	-0.8521620	1.0000000	0.9020329	0.8324475	-0.69993811	0.7824958
disp	-0.8475514	0.9020329	1.0000000	0.7909486	-0.71021393	0.8879799
hp	-0.7761684	0.8324475	0.7909486	1.0000000	-0.44875912	0.6587479
drat	0.6811719	-0.6999381	-0.7102139	-0.4487591	1.00000000	-0.7124406
wt	-0.8676594	0.7824958	0.8879799	0.6587479	-0.71244065	1.0000000
qsec	0.4186840	-0.5912421	-0.4336979	-0.7082234	0.09120476	-0.1747159
vs	0.6640389	-0.8108118	-0.7104159	-0.7230967	0.44027846	-0.5549157
am	0.5998324	-0.5226070	-0.5912270	-0.2432043	0.71271113	-0.6924953
gear	0.4802848	-0.4926866	-0.5555692	-0.1257043	0.69961013	-0.5832870
carb	-0.5509251	0.5269883	0.3949769	0.7498125	-0.09078980	0.4276059


  

	qsec	vs	am	gear	carb
mpg	0.41868403	0.6640389	0.59983243	0.4802848	-0.55092507
cyl	-0.59124207	-0.8108118	-0.52260705	-0.4926866	0.52698829
disp	-0.43369788	-0.7104159	-0.59122704	-0.5555692	0.39497686
hp	-0.70822339	-0.7230967	-0.24320426	-0.1257043	0.74981247
drat	0.09120476	0.4402785	0.71271113	0.6996101	-0.09078980
wt	-0.17471588	-0.5549157	-0.69249526	-0.5832870	0.42760594
qsec	1.00000000	0.7445354	-0.22986086	-0.2126822	-0.65624923
vs	0.74453544	1.0000000	0.16834512	0.2060233	-0.56960714

am	-0.22986086	0.1683451	1.00000000	0.7940588	0.05753435
gear	-0.21268223	0.2060233	0.79405876	1.00000000	0.27407284
carb	-0.65624923	-0.5696071	0.05753435	0.2740728	1.00000000

Heatmap

```
heatmap(cor(mtcars), main="Correlation Heatmap")
```



## 27.9 Common R Packages for Statistics

Package | Purpose ggplot2 | Data visualization dplyr | Data manipulation tidyr | Data tidying Hmisc  
 | Misc stats functions car | Regression diagnostics e1071 | Skewness/kurtosis, ML tools psych | Psy-  
 chological statistics shiny | Interactive apps caret | Classification and regression

## 27.10 Introduction to Command Line

## 27.11 Windows Terminal

```
cd..mkdirmy_projectdir
```

## Linux Terminal

```
cd mkdirstats_projectls -l
```

## 27.12 Git + R Project Example

*git init  
git clone https://github.com/username/project.git*

## 27.13 R Script Template for Analysis

```
## # Load packages library(ggplot2) library(dplyr)
```

## 28 Load data

```
df <- read.csv("dataset.csv")
```

## 29 Descriptive stats

```
summary(df) sd(df$Score)
```



## 30 Histogram

```
ggplot(df, aes(x=Score)) + geom_histogram(bins=10, fill="skyblue")
```

## 31 Linear regression

```
model <- lm(Score ~ StudyHours, data=df) summary(model)
```

## 32 Scatter plot with regression line

```
ggplot(df, aes(x=StudyHours, y=Score)) + geom_point() + geom_smooth(method="lm") $$ ##  
Fallacies and Bias: Real-World Cautions
```

Examples of Statistical Abuse

Cherry-picking data Data dredging (p-hacking) Using relative risk without absolute context Non-random sampling Ethics in Data Analysis

Be transparent Document sources Disclose methodology Avoid overstating conclusions

### 32.1 Future Applications of Statistics

Real-World Domains

Healthcare: Drug effectiveness, diagnostics Economics: Forecasting, policy evaluation Sociology: Survey analysis Sports: Performance analytics AI/ML: Predictive modeling, optimization Next Steps

Learn tidyverse ecosystem Explore machine learning in R Build Shiny dashboards Get familiar with reproducible research using Quarto

### 32.2 Practice Challenges

1. Load and summarize data

Load mtcars or your own dataset Use summary(), mean(), sd() 2. Create 3 different plots

Histogram Boxplot by group Scatter plot with trend line 3. Build a regression model

Identify predictor and outcome Use lm() and summary() 4. Explore a GUI like RKWard or Rcmdr

### 32.3 Key Takeaways

Statistics supports informed decision-making. R and its GUI frontends offer flexibility + power. Understand theory → then automate with code. Avoid fallacies by following robust methods. Visuals are crucial: plot early, plot often.

## 33 Week 4

## 34 Introduction

This eBook is a comprehensive companion to the course *Basic Statistics using GUI-R (RKWard)*. It includes foundational theory, practical examples, and step-by-step explanations, with integrated GUI-R usage.

## 35 Course Overview

### 35.1 Course Name

Basic Statistics using GUI-R (RKWard)

### 35.2 Instructor Profile

Dr. Harsh Pradhan is Assistant Professor at the Institute of Management Studies, Banaras Hindu University.

[Faculty Profile](#)

### 35.3 Learning Objectives

- Understand core concepts in statistics
- Apply t-tests and ANOVA using real data
- Compute confidence intervals and test statistics
- Use GUI-R (RKWard) for statistical analysis

# 36 Chapter 1: Fundamental Concepts

## 36.1 Descriptive Statistics

### 36.1.1 Central Tendency

- Mean
- Median
- Mode

### 36.1.2 Dispersion

- Range
- Variance
- Standard Deviation

#### 36.1.2.1 Example:

```
data <- c(4, 8, 6, 5, 3)
mean(data)
```

```
[1] 5.2
```

```
median(data)
```

```
[1] 5
```

```
sd(data)
```

```
[1] 1.923538
```

## 36.2 Standard Error

$$SE = \frac{s}{\sqrt{n}}$$

Small SE = sample mean is a good estimate of the population mean.

## 36.3 Central Limit Theorem

For  $n > 30$ , sampling distribution of the mean approximates normal:

$$\bar{X} \sim \mathcal{N}(\mu, \frac{\sigma}{\sqrt{n}})$$

## 36.4 Confidence Intervals

$$CI = \bar{x} \pm Z \cdot \frac{s}{\sqrt{n}}$$

Interpret 95% CI as: 95 of 100 such intervals would contain the true mean.



## 37 Chapter 2: Estimation

### 37.1 Types of Estimates

Type	Description	Example
Point Estimate	Single value	Sample mean
Interval Estimate	Range + confidence	Confidence Int

### 37.2 Parameter vs Statistic

Term	Description
Parameter	Value from population (e.g., $\mu$ )
Statistic	Value from sample (e.g., $\bar{x}$ )

## 38 Chapter 3: Hypothesis Testing

- **Null Hypothesis ( $H_0$ ):** No effect
- **Alternative Hypothesis ( $H_1$ ):** Some effect
- **Type I Error:** Reject  $H_0$  when true
- **Type II Error:** Fail to reject  $H_0$  when false

## 39 Chapter 4: Student's T-Test

### 39.1 Types

Test Type	Description
One-Sample	Compare sample to fixed value
Independent	Compare two unrelated groups
Paired	Compare two related groups

### 39.2 One-Sample T-Test Example

```
data <- c(22, 24, 27, 26, 28, 23, 25, 29, 21, 26, 24, 27)
t.test(data, mu = 25)
```

One Sample t-test

```
data: data
t = 0.2363, df = 11, p-value = 0.8175
alternative hypothesis: true mean is not equal to 25
95 percent confidence interval:
 23.61427 26.71906
sample estimates:
mean of x
 25.16667
```

### 39.3 Test Statistic

$$t = \frac{\bar{x} - \mu}{SE}$$

### 39.4 Degrees of Freedom

$$df = n - 1$$

## 39.5 Decision Rule

Compare calculated  $t$  to table value. If  $|t| > t_{critical}$ , reject  $H_0$ .

## 39.6 T-Test in GUI-R

1. Import data
2. Choose T-Test
3. Define groups
4. Run & interpret output

## 40 Chapter 5: ANOVA

### 40.1 Purpose

Used when comparing means across 3+ groups.

#### 40.1.1 One-Way ANOVA Formula

$$F = \frac{MS_{between}}{MS_{within}}$$

Where:

- $MS_{between} = \frac{SS_{between}}{df_{between}}$
- $MS_{within} = \frac{SS_{within}}{df_{within}}$

#### 40.1.2 Assumptions

- Normality
- Homogeneity of variance
- Independence

#### 40.1.3 Example Table

Group	Mean	Var	n
A	5.5	1.5	30
B	7.1	2.0	30
C	6.8	1.8	30

### 40.2 Post-Hoc Tests

Run if ANOVA is significant to locate pairwise differences.

## 40.3 ANOVA in GUI-R

1. Load data
2. Choose “One-Way ANOVA”
3. Define groups
4. Interpret output

## 41 Chapter 6: GUI-R Workflow

1. **Import Data** (CSV, Excel)
2. **Choose Test** (T-Test, ANOVA, etc.)
3. **Run** the analysis
4. **Interpret** the output
5. **Export** the results or visualizations

## 42 Chapter 7: Advanced Concepts

### 42.1 Variance Partitioning

$$\text{Total Variance} = \text{Explained Variance} + \text{Unexplained Variance}$$

Explained Terms	Unexplained Terms
Systematic	Random
Predictive	Error
Deterministic	Noise

### 42.2 Degrees of Freedom

For equation  $x + y + z = 3$ , if 2 values are known, third is fixed.  
Hence,  $df = n - k$  where  $n$  = total variables,  $k$  = constraints.

### 42.3 Chi-Square and F Distribution

- **Chi-Square:** Categorical variable comparison
- **F-Distribution:** Used in ANOVA, variance testing

### 42.4 Univariate, Bivariate, Multivariate

Type	Variables	Example
Univariate	1	Height
Bivariate	2	Height vs Weight
Multivariate	>2	Study w/ Age, Gender, Income



## 42.5 Parametric Test Assumptions

- Interval/Ratio DV
- Random Sampling
- Normality
- Equal Variances

If assumptions violated → use non-parametric test.

## 42.6 Effect Size

$$\text{Effect Size} = \frac{|\mu_1 - \mu_2|}{\sigma}$$

Used for comparison across studies.

## 42.7 Power of a Test

$$\text{Power} = 1 - \beta$$

Higher power → lower chance of Type II error

Power increases with sample size, effect size

## 43 Conclusion

Statistics is the language of data. GUI-R makes statistical tools accessible for everyone. This book empowers you to analyze data effectively using t-tests, ANOVA, and confidence intervals in a GUI environment.

## 44 References

- Pradhan, H. (2023). *Basic Statistics using GUI-R (RKWard)*
- Field, A. (2013). *Discovering Statistics Using R*.
- <https://methods.sagepub.com>

## 45 Chapter 8: Advanced T-Test Applications

### 45.1 Paired Sample T-Test

Used when the same group is measured twice (e.g., before and after).

#### 45.1.1 Example:

```
before <- c(80, 82, 79, 84, 88)
after <- c(78, 81, 76, 83, 86)
t.test(before, after, paired = TRUE)
```

Paired t-test

```
data: before and after
t = 4.8107, df = 4, p-value = 0.008581
alternative hypothesis: true mean difference is not equal to 0
95 percent confidence interval:
 0.7611494 2.8388506
sample estimates:
mean difference
      1.8
```

### 45.2 Independent Samples T-Test

Compare means of two unrelated groups.

```
group1 <- c(85, 90, 88, 92, 87)
group2 <- c(80, 83, 85, 84, 82)
t.test(group1, group2)
```

Welch Two Sample t-test

```
data: group1 and group2
t = 3.7755, df = 7.226, p-value = 0.006537
```

```
alternative hypothesis: true difference in means is not equal to 0
95 percent confidence interval:
 2.114814 9.085186
sample estimates:
mean of x mean of y
 88.4      82.8
```

### 45.3 One-Sample T-Test with GUI-R

- Import dataset
- Use 'Descriptive Statistics' to check mean
- Navigate to 'T-Test' → 'One Sample'
- Input hypothesized mean and run

## 46 Chapter 9: More on Confidence Intervals


### 46.1 Visualizing Confidence Intervals in R

```
x <- c(88, 90, 85, 87, 89)
mean_x <- mean(x)
se <- sd(x) / sqrt(length(x))
ci_lower <- mean_x - 1.96 * se
ci_upper <- mean_x + 1.96 * se
c(ci_lower, ci_upper)
```

```
[1] 86.11394 89.48606
```

Plot using ggplot2:

```
library(ggplot2)
df <- data.frame(x = x)
ggplot(df, aes(y = x, x = 1)) +
  geom_point() +
  geom_errorbar(aes(ymin = ci_lower, ymax = ci_upper), width = 0.1)
```



## 47 Chapter 10: Robust ANOVA Models

### 47.1 Two-Way ANOVA

Examines the effect of two categorical independent variables on a continuous dependent variable.

```
# Sample dataset for demonstration
dataset <- data.frame(
  score = c(85, 90, 88, 92, 87, 80, 83, 85, 84, 82),
  gender = rep(c("Male", "Female"), each = 5),
  teaching_method = rep(c("A", "B"), times = 5)
)
aov_result <- aov(score ~ gender * teaching_method, data = dataset)
summary(aov_result)
```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)	
gender	1	78.40	78.40	23.718	0.00279	**
teaching_method	1	6.02	6.02	1.820	0.22598	
gender:teaching_method	1	18.15	18.15	5.491	0.05759	.
Residuals	6	19.83	3.31			
---						
Signif. codes:	0	'***'	0.001	'**'	0.01	'*' 0.05 '.' 0.1 ' ' 1

### 47.2 Repeated Measures ANOVA

Use when the same subjects are used for each treatment.

```
# Sample repeated measures data in long format
data_long <- data.frame(
  id = rep(1:5, each = 3),
  condition = rep(c("A", "B", "C"), times = 5),
  score = c(85, 88, 90, 80, 82, 85, 78, 80, 83, 90, 92, 95, 88, 90, 91)
)
library(ez)
ezANOVA(data = data_long, dv = .(score), wid = .(id), within = .(condition))
```

Warning: Converting "id" to factor for ANOVA.

Warning: Converting "condition" to factor for ANOVA.

\$ANOVA

	Effect	DFn	DFd	F	p	p<.05	ges
2	condition	2	8	88.22222	3.539139e-06	*	0.1479687

\$`Mauchly's Test for Sphericity`

	Effect	W	p	p<.05
2	condition	0.5555556	0.4140867	

\$`Sphericity Corrections`

	Effect	GGe	p[GG]	p[GG]<.05	HFe	p[HF]	p[HF]<.05
2	condition	0.6923077	9.135419e-05	*	0.9411765	6.568851e-06	*



## 48 Chapter 11: Effect Size Measures

### 48.1 Cohen's d

$$d = \frac{\bar{x}_1 - \bar{x}_2}{s_p}$$

Where  $s_p$  is the pooled standard deviation.

#### 48.1.1 R Example

```
library(effsize)
cohen.d(group1, group2)
```

Cohen's d

```
d estimate: 2.387848 (large)
95 percent confidence interval:
      lower      upper
0.4791634  4.2965327
```

### 48.2 Eta-Squared ( $\eta^2$ )

Used for ANOVA:

$$\eta^2 = \frac{SS_{between}}{SS_{total}}$$

## 49 Chapter 12: Statistical Assumptions Checking

### 49.1 Normality

Use Shapiro-Wilk test:


```
# Sample data frame for normality test
data <- data.frame(variable = c(88, 90, 85, 87, 89, 91, 92, 88, 90, 87))
shapiro.test(data$variable)
```

Shapiro-Wilk normality test

```
data: data$variable
W = 0.97743, p-value = 0.95
```

Visualize:

```
qqnorm(data$variable)
qqline(data$variable)
```



## 49.2 Homogeneity of Variance

Use Levene's Test:

```
# Sample data frame for Levene's Test
data <- data.frame(
  variable = c(88, 90, 85, 87, 89, 91, 92, 88, 90, 87),
  group = rep(c("A", "B"), each = 5)
)
library(car)
```

Loading required package: carData

```
leveneTest(variable ~ group, data = data)
```

Warning in leveneTest.default(y = y, group = group, ...): group coerced to factor.

```
Levene's Test for Homogeneity of Variance (center = median)
      Df F value Pr(>F)
group  1  0.0769 0.7885
      8
```

## 50 Chapter 13: Non-Parametric Alternatives

### 50.1 Wilcoxon Signed Rank Test

```
wilcox.test(before, after, paired = TRUE)
```

Warning in wilcox.test.default(before, after, paired = TRUE): cannot compute exact p-value with ties

Wilcoxon signed rank test with continuity correction

data: before and after

V = 15, p-value = 0.05676

alternative hypothesis: true location shift is not equal to 0

### 50.2 Mann-Whitney U Test

```
wilcox.test(group1, group2)
```

Warning in wilcox.test.default(group1, group2): cannot compute exact p-value with ties

Wilcoxon rank sum test with continuity correction

data: group1 and group2

W = 24.5, p-value = 0.01597

alternative hypothesis: true location shift is not equal to 0

### 50.3 Kruskal-Wallis Test

Non-parametric alternative to ANOVA.

```
# Sample data frame for Kruskal-Wallis Test
data <- data.frame(
  score = c(85, 88, 90, 80, 82, 85, 78, 80, 83, 90, 92, 95, 88, 90, 91),
  group = rep(c("A", "B", "C"), times = 5)
)
kruskal.test(score ~ group, data = data)
```

Kruskal-Wallis rank sum test


data: score by group

Kruskal-Wallis chi-squared = 2.2329, df = 2, p-value = 0.3274

## 51 Chapter 14: Visualizing Statistical Results


### 51.1 Boxplots

```
ggplot(data, aes(x = group, y = score)) +  
  geom_boxplot()
```




### 51.2 Histograms

```
ggplot(data, aes(x = score)) +  
  geom_histogram(binwidth = 2)
```



## 51.3 Density Plot

```
ggplot(data, aes(x = score)) +  
  geom_density()
```



## 52 Chapter 15: RKWard (GUI-R) Tips

- Use menu-based analysis for beginners
- Save and export plots easily
- Integrate with R scripts for reproducibility

### 52.1 Summary: Basic Statistics using GUI-R (RKWard)

This eBook, authored by Dr. Harsh Pradhan (Assistant Professor at the Institute of Management Studies, Banaras Hindu University), serves as a comprehensive guide to understanding and applying basic statistical concepts, particularly in the GUI-based software RKWard (GUI-R).

Key Highlights: 1. Descriptive Statistics Covers measures of central tendency (mean, median, mode) and variability (range, variance, standard deviation). Introduces standard error and its role in estimating population parameters. 2. Inferential Statistics Introduces the Central Limit Theorem and how it forms the foundation for many statistical techniques. Confidence intervals are explained both theoretically and with practical calculations. 3. T-Tests (Student's t) Explains one-sample, independent-sample, and paired-sample t-tests. Includes step-by-step computation and GUI-R implementation. Includes interpretation of p-values, degrees of freedom, and test statistics. 4. Analysis of Variance (ANOVA) Covers one-way, two-way, and repeated measures ANOVA. Focuses on the F-statistic, assumptions, and post-hoc analyses. Discusses partitioning of variance into systematic and unsystematic components. 5. Effect Size and Statistical Power Introduces Cohen's d, eta-squared, and power analysis. Emphasizes that statistical significance does not always imply practical importance. 6. Assumption Testing Tests for normality (Shapiro-Wilk, QQ plot). Tests for homogeneity of variance (Levene's test). Highlights when to use non-parametric alternatives. 7. Non-Parametric Tests Introduces Wilcoxon signed-rank, Mann-Whitney U, and Kruskal-Wallis tests as robust alternatives to parametric methods. 8. Data Visualization in R Demonstrates use of boxplots, histograms, and density plots using ggplot2. Provides example R code for reproducibility. 9. GUI-R (RKWard) Usage Offers practical steps for using GUI-R for all statistical techniques covered. Designed to bridge the gap for learners unfamiliar with command-line R.