Rappel de cours:

• 
$$\int_a^b f(u(x))u'(x) dx = \int_{x(a)}^{u(b)} u du$$

## Exercice 1

### Exercice 1.a

$$\int tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx$$

Soit u(x) = cos(x), u'(x) = -sin(x) et  $f(u) = -\frac{1}{u}$  donc

$$\int \frac{\sin(x)}{\cos(x)} dx = \int f(\cos(x))\sin(x) dx = \int -\frac{1}{u} du = -\ln(u) = -\ln(\cos(x))$$

### Exercice 1.b

$$\int \frac{\ln(x+1)}{x+1} \, dx$$

Soit u(x) = ln(x+1),  $u'(x) = \frac{1}{x+1}$  et f(u) = u donc

$$\int \frac{\ln(x+1)}{x+1} \, dx = \int f(\ln(x+1)) \cdot \frac{1}{x+1} \, dx = \int u \, du = \frac{u^2}{2} = \frac{\ln^2(x+1)}{2}$$

### Exercice 1.c

$$\int \frac{x}{\sqrt{2x^2+3}}$$

Soit  $u(x) = 2x^2 + 3$ , u'(x) = 4x et  $f(u) = \frac{1}{4\sqrt{u}}$  donc

$$\int \frac{x}{\sqrt{2x^2 + 3}} \, dx = \int f(2x^2 + 3) \cdot 4x \, dx = \int \frac{1}{4\sqrt{u}} \, du = \frac{1}{4} \int \frac{1}{\sqrt{u}} \, du = \frac{\sqrt{u}}{2} = \frac{\sqrt{2x^2 + 3}}{2}$$

### Exercice 1.d

$$\int \frac{\cos(x)}{\sin^2(x)} \, dx$$

Soit u(x) = sin(x), u'(x) = cos(x) et  $f(u) = \frac{1}{u^2}$  donc

$$\int \frac{\cos(x)}{\sin^2(x)} \, dx = \int f(\sin(x)) \cdot \cos(x) \, dx = \int \frac{1}{u^2} \, du = -\frac{1}{u} = -\frac{1}{\sin(x)}$$

### Exercice 1.e

$$\int x\sqrt{x^2+1}\,dx$$

Soit  $u(x) = x^2 + 1$ , u'(x) = 2x et  $f(u) = \frac{\sqrt(u)}{2}$  donc

$$\int x\sqrt{x^2+1}\,dx = \int f(x^2+1).2x\,dx = \int \frac{\sqrt(u)}{2}\,du = \frac{1}{2}\int \sqrt(u)\,du = \frac{1}{2}\frac{2}{3}u^{\frac{3}{2}} = \frac{1}{3}u^{\frac{3}{2}} = \frac{(x^2+1)^{\frac{3}{2}}}{3}u^{\frac{3}{2}} = \frac{1}{3}u^{\frac{3}{2}} = \frac{1}{3}u^{\frac{3}{2$$

### Exercice 1.f

$$\int \frac{1}{x \ln(x)} \, dx$$

Soit u(x) = ln(x),  $u'(x) = \frac{1}{x}$  et  $f(u) = \frac{1}{u}$  donc

$$\int \frac{1}{x \ln(x)} \, dx = \int f(\ln(x)) \cdot \frac{1}{x} \, dx = \int \frac{1}{u} \, du = \ln(|u|) = \ln(|\ln(x)|)$$

### Exercice 1.g

$$\int \sin^2(x)\cos(x)\,dx$$

Soit u(x) = sin(x), u'(x) = cos(x) et  $f(u) = u^2$  donc

$$\int \sin^2(x)\cos(x) \, dx = \int f(\sin(x)).\cos(x) \, dx = \int u^2 \, du = \frac{u^3}{3} = \frac{\sin^3(x)}{3}$$

### Exercice 1.h

$$\int e^{\sin(x)}\cos(x)\,dx$$

Soit u(x) = sin(x), u'(x) = cos(x) et  $f(u) = e^u$  donc

$$\int e^{\sin(x)}\cos(x) dx = \int f(\sin(x)).\cos(x) dx = \int e^{u} du = e^{u} = e^{\sin(x)}$$

### Exercice 1.i

$$\int \frac{\ln^2(x)}{x} \, dx$$

Soit u(x) = ln(x),  $u'(x) = \frac{1}{x}$  et  $f(u) = u^2$  donc

$$\int \frac{\ln^2(x)}{x} \, dx = \int f(\ln(x)) \cdot \frac{1}{x} \, dx = \int u^2 \, du = \frac{u^3}{3} = \frac{\ln^3(x)}{3}$$

## Exercice 2

### Exercice 2.a

 $t=u(x)=tan(\frac{x}{2})$ donc $x=u^{-1}(t)=2arctan(t)$  et  $(u^{-1})'(t)=\frac{dx}{dt}=\frac{2}{1+t^2}.$  On a

$$sin(x) = \frac{2tan(\frac{x}{2})}{1 + tan^2(\frac{x}{2})}$$

$$1-\sin(x) = 1 - \frac{2tan(\frac{x}{2})}{1+tan^2(\frac{x}{2})} = \frac{tan^2(\frac{x}{2}) - 2tan^2(\frac{x}{2}) + 1}{1+tan^2(\frac{x}{2})} = \frac{(tan(\frac{x}{2}) - 1)^2}{1+tan^2(\frac{x}{2})}$$

Donc

$$\frac{1}{1-sin(x)} = \frac{1+tan^2(\frac{x}{2})}{(tan(\frac{x}{2})-1)^2}$$

 $\operatorname{Et}$ 

$$\int_0^{\pi/3} \frac{1}{1 - \sin(x)} dx = \int_0^{\pi/3} \frac{1 + \tan^2(\frac{x}{2})}{(\tan(\frac{x}{2}) - 1)^2} dx$$
$$\int_{tan(0)}^{tan(\pi/6)} \frac{1 + t^2}{(t - 1)^2} \cdot \frac{2}{1 + t^2} dt = \int_{tan(0)}^{tan(\pi/6)} \frac{2}{(t - 1)^2} dt$$

Soit 
$$u(x) = x - 1$$
,  $u'(x) = 1$  et  $f(u) = \frac{1}{u^2}$  donc

$$\int_{tan(0)}^{tan(\pi/6)} \frac{2}{(t-1)^2} \, dt = 2 \int_{tan(0)}^{tan(\pi/6)} f(t-1).1 \, dt = 2 \int_{tan(0)-1}^{tan(\pi/6)-1} \frac{1}{u^2} \, du = [\frac{2}{u}]_{tan(0)-1}^{tan(\pi/6)-1} = -2 - \frac{2}{tan(\pi/6)-1}$$

### Exercice 2.b

 $t = u(x) = tan(\frac{x}{2})$  donc  $x = u^{-1}(t) = 2arctan(t)$  et  $(u^{-1})'(t) = \frac{dx}{dt} = \frac{2}{1+t^2}$ . On a

$$cos(x) = \frac{1 - tan^2(\frac{x}{2})}{1 + tan^2(\frac{x}{2})}$$

$$5 - 3\cos(x) = 5 - 3 \cdot \frac{1 - \tan^2(\frac{x}{2})}{1 + \tan^2(\frac{x}{2})} = \frac{2 + 8\tan^2(\frac{x}{2})}{1 + \tan^2(\frac{x}{2})}$$

Donc

$$\frac{1}{5 - 3\cos(x)} = \frac{1 + \tan^2(\frac{x}{2})}{2(1 + 4\tan^2(\frac{x}{2}))}$$

 $\operatorname{Et}$ 

$$\int_0^{\pi/2} \frac{1}{5 - 3\cos(x)} \, dx = \int_0^{\pi/2} \frac{1 + \tan^2(\frac{x}{2})}{2(1 + 4\tan^2(\frac{x}{2}))} \, dx$$

$$\int_{4\pi\pi(0)}^{\tan(\pi/4)} \frac{1 + t^2}{2(1 + 4t^2)} \cdot \frac{2}{1 + t^2} \, dt = \int_{4\pi\pi(0)}^{\tan(\pi/4)} \frac{1}{1 + 4t^2} \, dt$$

Soit 
$$u(t)=2t,\,u'(t)=2$$
 et  $f(u)=\frac{1}{2(u^2+1)}$  donc

$$\int_{tan(0)}^{tan(\pi/4)} \frac{1}{1+4t^2} dt = \int_{tan(0)}^{tan(\pi/4)} f(2t) \cdot 2 dt = \int_{2tan(0)}^{2tan(\pi/4)} \frac{1}{2(u^2+1)} du = \frac{1}{2} \left[arctan(u)\right]_{2tan(0)}^{2tan(\pi/4)}$$

### Exercice 2.c

 $t=u(x)=tan(\frac{x}{2})$ donc $x=u^{-1}(t)=2arctan(t)$  et  $(u^{-1})'(t)=\frac{dx}{dt}=\frac{2}{1+t^2}.$  On a

$$cos(x) = \frac{1 - tan^2(\frac{x}{2})}{1 + tan^2(\frac{x}{2})}$$

$$1 + \cos(x) = 1 + \frac{1 - \tan^2(\frac{x}{2})}{1 + \tan^2(\frac{x}{2})} = \frac{2}{1 + \tan^2(\frac{x}{2})}$$

$$(1 + \cos(x))^2 = \frac{4}{(1 + \tan^2(\frac{x}{2}))^2}$$

Donc

$$\frac{1}{(1+\cos(x))^2} = \frac{(1+\tan^2(\frac{x}{2}))^2}{4}$$

 $\operatorname{Et}$ 

$$\int_0^{\pi/2} \frac{1}{(1+\cos(x))^2} dx = \int_0^{\pi/2} \frac{(1+\tan^2(\frac{x}{2}))^2}{4} dx$$

$$\int_{tan(0)}^{tan(\pi/4)} \frac{(1+t^2)^2}{4} \cdot \frac{2}{1+t^2} dt = \int_{tan(0)}^{tan(\pi/4)} \frac{1}{2} (1+t^2) dt = \frac{1}{2} [t+\frac{t^3}{3}]_{tan(0)}^{tan(\pi/4)}$$

#### Exercice 2.c

Trop complexe.

# Exercice 3

Exercice 3.a

$$\int arcsin(x) \, dx$$

Exercice 3.b

$$\int x^n ln(x) \, dx$$

Par partie avec

$$f(x) = ln(x)$$
  $f'(x) = \frac{1}{x}$   
 $g'(x) = x^n$   $g(x) = \frac{x^{n+1}}{n+1}$ 

Donc

$$\int x^n \ln(x) \, dx = \frac{x^{n+1} \ln(x)}{n+1} - \int \frac{1}{x} \cdot \frac{x^{n+1}}{n+1} \, dx = \frac{x^{n+1} \ln(x)}{n+1} - \frac{1}{n+1} \int x^n \, dx = \frac{x^{n+1} \ln(x)}{n+1} - \frac{x^{n+1} \ln(x)}{(n+1)^2} - \frac{x^{n+1}$$

Exercice 3.c

$$\int x \arcsin(x) \, dx$$

Exercice 3.d

$$\int \ln(x^2+1)\,dx$$

 $\operatorname{QED}$