

On the sensitivities of marine low cloud cover to the strength of the tropical inversion and sea surface temperature

Xin Qu (UCLA), Alex Hall (UCLA)



Steve Klein (LLNL)




Anthony DeAngelis (UCLA)



CFMIP Meeting on Cloud Processes and Feedbacks
June 8, 2015




A simple framework to understand and constrain tropical marine low-cloud cover (LCC) feedback



EIS: estimated inversion strength
SST: sea surface temperature


What type of climate variability contains the “DNA” of climate change?

The sensitivity of LCC to EIS, averaged over five subtropical low-cloud regions



- *LCC generally increases with increasing EIS.*
- *There is a high degree of time-scale invariance in the EIS sensitivity.*

The sensitivity of LCC to SST, averaged over five subtropical low-cloud regions




- *LCC generally decreases with increasing SST, except for the seasonal cycle.*
- *The SST sensitivity is similar on interannual and decadal time scales.*

What type of climate variability contains the “DNA” of climate change?

Cross-model correlation between the simple model-predicted and actual GCM LCC changes in Scenario A1B and RCP8.5 simulations

	CMIP3	CMIP5
Seasonal cycle	0.12	-0.09
Intermonthly	0.51	0.15
Interannual	0.81	0.46
Decadal	0.91	0.84



- *Decadal variations contain the information most relevant for climate change.*

What drives the SST sensitivity?: Lesson from a seven-variable framework

$$\Delta LCC \approx \frac{\partial LCC}{\partial EIS} \cdot \Delta EIS + \frac{\partial LCC}{\partial LHF} \cdot \Delta LHF + \frac{\partial LCC}{\partial \delta q} \cdot \Delta \delta q + \frac{\partial LCC}{\partial R} \cdot \Delta R + \frac{\partial LCC}{\partial \omega 700} \cdot \Delta \omega 700 + \frac{\partial LCC}{\partial U_s} \cdot \Delta U_s + \frac{\partial LCC}{\partial T_{adv}} \cdot \Delta T_{adv}$$

Sensitivities of LCC to various factors in current climate

Projected changes in those factors in the 21st century

Six other factors:

LHF: surface latent heat flux

$\delta q = q(\text{surface}) - q(700\text{hPa})$: vertical moisture gradient in the lower-troposphere


R: mean relative humidity in the free-troposphere

$\omega 700$: vertical velocity at 700 hPa.

U_s: surface wind speed


T_{adv}: temperature advection

Comparison of 7 terms: $\frac{\partial LCC}{\partial x} \cdot \Delta x$



- The dominant terms are the EIS, LHF and δq terms.
- The EIS term is generally positive, while the LHF and δq terms are generally negative.


Interpreting the SST contribution in the 2-variable model



➤ *The SST contribution in the 2-variable model can be interpreted by the contributions of two-warming induced changes:*


- an increase in vertical moisture gradient in the lower-troposphere
- an increase in surface latent heat flux

Agreement between the 2- and 7-variable models



➤ *Two simple models capture almost exactly the same portion of the intermodel variance in LCC changes.*


Observed constraints of EIS and SST sensitivities in the 2-variable model



- ISCCP (1984-2009)
- MISR (2000-2013)
- △ PATMOS-x (1982-2009)
- ✱ MODIS (2002-2014)

- *Observed EIS sensitivity ranges from 2.5 to 3.5 %/K, while observed SST sensitivity ranges from -3 to -1 %/K.*
- *Many GCMs underestimate the magnitudes of both EIS and SST sensitivities.*

Observed constraints of LCC changes



Using observed sensitivities and ensemble-mean EIS and SST changes

Using observed sensitivities and EIS and SST changes in each model

- *In almost all predictions constrained by observations, the large LCC increases in some GCMs are excluded.*
- Applying a similar methodology to abrupt4xCO2 simulations, we find that a strong negative LCC feedback is unlikely.

Take home message

- 1. LCC changes in GCMs can be interpreted as a linear combination of contributions by EIS and SST.*
- 2. The negative SST contribution (due to a negative LCC sensitivity to SST) originates primarily from two warming-induced changes: increases in surface latent heat flux and increases in vertical moisture gradient.*
- 3. The sensitivities of LCC to EIS and SST can be constrained by observations.*
- 4. If realistic EIS and SST sensitivities are simulated in GCMs, a strong negative LCC feedback is unlikely.*

Outstanding questions

1. *Why are EIS and SST sensitivities diagnosed from longer term climate variations better predictive of climate change than those from shorter term climate variations?*
2. *What are the causes of model bias in EIS and SST sensitivities?*
3. *Is there any way to reduce the uncertainties in observed EIS and SST sensitivities?*