mielke-geometrodynamics — formula report

1794 inline formulas (section omitted; all rendered), 1149 display equations, 5 tables, 18 unrecovered image regions.

Corrected (3)

<table>
<tr><th>Identifier</th><th>Page</th><th>Conf.</th><th>LaTeX source</th><th>Rendered</th><th>Scan image</th></tr>
<tr><td>mielke-geometrodynamics_
EQ0378 (was)</td><td>128</td><td>■ 0.136</td><td>D D \vartheta^{\alpha}=R_{\beta}{ }^{\alpha} \wedge \vartheta^{\beta}=0, \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha}=-\stackrel{(\pm)}{R}_{\alpha}{ }^{\beta} \wedge \stackrel{(\pm)}{\Pi}_{\beta}= \pm \frac{i}{2} \ell^{2}\left(e_{\alpha} \downharpoonleft \stackrel{(\pm)}{\Pi^{\beta}}\right) \wedge \Sigma_{\beta} \cong 0 .</td><td>$D D \vartheta^{\alpha}=R_{\beta}{ }^{\alpha} \wedge \vartheta^{\beta}=0, \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha}=-\stackrel{(\pm)}{R}_{\alpha}{ }^{\beta} \wedge \stackrel{(\pm)}{\Pi}_{\beta}= \pm \frac{i}{2} \ell^{2}\left(e_{\alpha} \downharpoonleft \stackrel{(\pm)}{\Pi^{\beta}}\right) \wedge \Sigma_{\beta} \cong 0 .$</td><td>—</td></tr>
<tr><td>mielke-geometrodynamics_
EQ0378 (now)</td><td>128</td><td>—</td><td>preserve the term Pontryagin index for the classification of real associated vector bundles (\textsc{Mayer} \& \textsc{Drechsler} 1977). Nevertheless, the denotation of \textsc{Belavin}</td><td>$preservethetermPontryaginindexfortheclassificationofrealassociatedvectorbundles($Mayer&Drechsler$1977).Nevertheless,thedenotationof$Belavin</td><td>—</td></tr>
<tr><td colspan="6">basis: inferred / ink</td></tr>
<tr><td>mielke-geometrodynamics_
EQ0402 (was)</td><td>134</td><td>■ 0.042</td><td>\begin{array}{ll} q: & \underline{*}^{\vartheta^{\beta}} \Psi_{\|}(\vartheta)=\underline{*}^{\prime}\underline{\vartheta}^{\beta} \Psi_{\|}(\vartheta), \\ \stackrel{ \pm}{p}: & \stackrel{*}{+}_{\underline{\Pi}} \Psi_{\|}(\vartheta)=-i \ell^{2} \frac{\delta}{\delta \underline{*}_{\alpha}} \Psi_{\|}(\vartheta), \end{array}</td><td>$\begin{array}{ll} q: & \underline{*}^{\vartheta^{\beta}} \Psi_{\|}(\vartheta)=\underline{*}^{\prime}\underline{\vartheta}^{\beta} \Psi_{\|}(\vartheta), \\ \stackrel{ \pm}{p}: & \stackrel{*}{+}_{\underline{\Pi}} \Psi_{\|}(\vartheta)=-i \ell^{2} \frac{\delta}{\delta \underline{*}_{\alpha}} \Psi_{\|}(\vartheta), \end{array}$</td><td>—</td></tr>
<tr><td>mielke-geometrodynamics_
EQ0402 (now)</td><td>134</td><td>—</td><td>\begin{array}{ll} q: & \stackrel{*}{\underline{\vartheta}}^{\beta} \Psi_{\|}(\vartheta) = \stackrel{*'}{\underline{\vartheta}}^{\beta} \Psi_{\|}(\vartheta), \\ \ \stackrel{\pm}{p}: & \stackrel{*^{(\pm)}}{\underline{\Pi}_{\alpha}} \Psi_{\|}(\vartheta) = -i\ell^{2} \dfrac{\delta}{\delta \stackrel{*}{\underline{\vartheta}}_{\alpha}} \Psi_{\|}(\vartheta), \end{array}</td><td>$\begin{array}{ll} q: & \stackrel{*}{\underline{\vartheta}}^{\beta} \Psi_{\|}(\vartheta) = \stackrel{*'}{\underline{\vartheta}}^{\beta} \Psi_{\|}(\vartheta), \\ \ \stackrel{\pm}{p}: & \stackrel{*^{(\pm)}}{\underline{\Pi}_{\alpha}} \Psi_{\|}(\vartheta) = -i\ell^{2} \dfrac{\delta}{\delta \stackrel{*}{\underline{\vartheta}}_{\alpha}} \Psi_{\|}(\vartheta), \end{array}$</td><td>—</td></tr>
<tr><td colspan="6">basis: inferred / ink</td></tr>
<tr><td>mielke-geometrodynamics_
EQ0972 (was)</td><td>314</td><td>■ 0.153</td><td>\stackrel{\text { (१) }}{D} \alpha:=\stackrel{\text { (०) }}{d} \alpha-\omega^{g} \wedge \alpha-\alpha \wedge \stackrel{\circ}{\circ}</td><td>$\stackrel{\text { (१) }}{D} \alpha:=\stackrel{\text { (०) }}{d} \alpha-\omega^{g} \wedge \alpha-\alpha \wedge \stackrel{\circ}{\circ}$</td><td>—</td></tr>
<tr><td>mielke-geometrodynamics_
EQ0972 (now)</td><td>314</td><td>—</td><td>\overset{(1)}{D}\alpha:=\overset{(0)}{d}\alpha-\omega^{g} \wedge\alpha-\alpha\wedge\omega^{g}</td><td>$\overset{(1)}{D}\alpha:=\overset{(0)}{d}\alpha-\omega^{g} \wedge\alpha-\alpha\wedge\omega^{g}$</td><td>—</td></tr>
<tr><td colspan="6">basis: inferred / ink</td></tr>
<tr><td colspan="6">Conf. MathPix confidence: ■ $\geq$0.9 ■ 0.5–0.9 ■ <0.5 — no value
Residual render vs scan (inkdrill): • C component • W weak • S stable • N noise • K clean • not measured</td></tr>
</table>

Unresolved (2)

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_EQ0472	155	—	`\stackrel{十}{\Omega}^{(t)}=0 \Longleftrightarrow R_{i j}^{\{ \}}=0 \Longleftrightarrow R_{i j}^{\{ \}}= \Lambda^{\prime} g_{i j} .`	*(not rendered)*	$\overset{+}{\Omega}^{\{\}} = 0 \Longleftrightarrow \not{R}_{ij}^{\{\}} = 0 \Longleftrightarrow R_{ij}^{\{\}} = \Lambda' g_{ij}.$
mielke-geometrodynamics_EQ0857	278	—	`\left.\mathbf{ヒ}_{n} \mathscr{G}^{A} \cong n\right \rfloor\left(D \underline{\tau}^{A}\right),`	*(not rendered)*	$Ł_n \mathscr{G}^A \cong n \rfloor (D \underline{\tau}^A)\,,$

Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value

Residual render vs scan (inkdrill): • C component • W weak • S stable • N noise • K clean • not measured

Flagged, not acted on: 176 rows, none above the reporting threshold.

Each differs from its scan by fewer than 20 components, which is the tail of the distribution rather than its bulk: 53 are component (`C`), 123 weak (`W`). By MathPix confidence: 105 at $\geq$0.9, 30 in 0.6–0.9, 41 below 0.6, 0 with none. Out/465 sampled twenty rows drawn from the $\geq$0.9 band corpus-wide and found eighteen purely typographic differences, one typographic plus a crop overrun, one borderline and *no* unambiguous content errors — so a high-confidence flag here is evidence of a rendering difference, not of a defect. Every row is in `report.ink.json`.