

cardona-qft-methods: residuals

equations measured 2026-09-15T14:37:04Z against the model of 2026-09-15

Corrected (2)

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
cardona-qft-methods_ E00048 (was)	35	■ 1.000	$c\left(d\,x^{\{j\}}\right)\left(x_0\right)=\gamma^j,\quad \sigma_1^D\left(x_0,\xi\right)=c\left(\xi\right)=\gamma^j\xi_j$		
cardona-qft-methods_ E00048 (now)	35	—	$c\left(d\,x^{\{j\}}\right)\left(x_0\right)=\gamma^j,\quad \sigma_1^{\mathcal{D}}\left(x_0,\xi\right)=c\left(\xi\right)=\gamma^j\xi_j$		
basis: census / census					
cardona-qft-methods_ E00025 (was)	313	■ 1.000	$F^{\{10\}}=\mathrm{d}\,C-H\wedge C+m\,e^B=\hat{F}-H\wedge C$	$F^{(10)}=\mathrm{d}C-H\wedge C+m\,e^B=\hat{F}-H\wedge C$	$F^{(10)}=\mathrm{d}C-H\wedge C+m\,e^B=\hat{F}-H\wedge C$
cardona-qft-methods_ E00025 (now)	313	—	$\cancel{F}^{\{10\}}=\mathrm{d}\,C-H\wedge C+m\,e^B=\hat{F}-H\wedge C$	$\cancel{F}^{(10)}=\mathrm{d}C-H\wedge C+m\,e^B=\hat{F}-H\wedge C$	$F^{(10)}=\mathrm{d}C-H\wedge C+m\,e^B=\hat{F}-H\wedge C$
basis: census / census					
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Unresolved

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E00654 (4)	261	■ 0.167 ●	$\phi_{\theta}(x)\phi_{\theta}(y)=e^{-\frac{1}{2}\frac{\partial}{\partial x^{\mu}}\frac{\partial}{\partial y^{\nu}}\theta^{\mu\nu}}\phi_0(x)\phi_0(y)e^{\frac{1}{2}\left(\frac{\overleftarrow{\partial}}{\partial x^{\mu}}+\frac{\overleftarrow{\partial}}{\partial y^{\mu}}\right)\theta^{\mu\nu}P_{\nu}}\neq\phi_0(x)\phi_0(y).$	(not rendered)	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: basis] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Flagged, not acted on

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0677	46	■ 0.150 ● C +44	\left.\begin{array}{ccccc} 1 & \rightarrow & \operatorname{Inn}(\mathcal{A}) & \rightarrow & \operatorname{Aut}(\mathcal{A}) & \rightarrow & \operatorname{Aut}(\mathcal{A})/\operatorname{Inn}(\mathcal{A}) & \rightarrow & 1, \\ 1 & \rightarrow & \mathcal{G} & \rightarrow & \mathcal{G} \rtimes \operatorname{Diff}(M) & \rightarrow & \operatorname{Diff}(M) & \rightarrow & 1. \end{array}\right\}	(not rendered)	
cardona-qft-methods_ EQ0748	284	■ 0.547 ● C -27	\begin{aligned} A d S_5 &: & -R^2 &= -Z_0^2 + Z_1^2 + Z_2^2 + Z_3^2 + Z_4^2 - Z_5^2, \\ S^5 &: & R^2 &= Y_1^2 + Y_2^2 + Y_3^2 + Y_4^2 + Y_5^2 + Y_6^2. \end{aligned}		
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

86 more flagged rows below the band, stated as a count: 18 component, 68 weak; confidence high 59, mid 14, low 13, none 0

Low confidence

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ9908 [conf 0.080]	350	0.080 U +5	$\begin{array}{rcl} P_g M : {}_1 \times {}_0 P_h M & \longrightarrow & P_g M \times P_h M \\ \downarrow & e_0 \circ \pi_1 & \downarrow \\ M & & \downarrow \times M \\ \downarrow & & \end{array}$	$\begin{array}{ccc} P_g M : {}_1 \times {}_0 P_h M & \longrightarrow & P_g M \times P_h M \\ \downarrow e_0 \circ \pi_1 & & \downarrow e_0 \times e_0 \\ M & \xrightarrow{\Delta_g} & M \times M \end{array}$	
Conf. MathPix confidence: ≥0.9 0.5-0.9 <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured refined: <i>basis</i> beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					