

gilmore-lie-groups: equation evidence

6 rows show a REFINEMENT below, not the OCR reading: 6 verified against the census. Each is marked [refined] beside its identifier. The model is unchanged — the original reading is still what ‘latex’ holds; this report chose the refinement for them. Rows: gilmore-lie-groups_EQ0012, gilmore-lie-groups_EQ0412, gilmore-lie-groups_EQ0413, gilmore-lie-groups_EQ0486, gilmore-lie-groups_EQ0556, gilmore-lie-groups_EQ0788

Display equations

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_EQ0584 (11.4) [conf 0.001]	188	0.001	$\left.\left.\left.\begin{bmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & \ddots & \\ & & & & 1 \\ & & & & & -1 \\ & & & & & & 1 \\ & & & & & & & -1 \\ & & & & & & & & \ddots \end{bmatrix} \begin{matrix} H_1 \\ H_2 \\ H_3 \\ \vdots \\ H_n \\ \frac{1}{\sqrt{2}}(E_\alpha + E_{-\alpha}) \\ \frac{1}{\sqrt{2}}(E_\alpha - E_{-\alpha}) \\ \frac{1}{\sqrt{2}}(E_\beta + E_{-\beta}) \\ \frac{1}{\sqrt{2}}(E_\beta - E_{-\beta}) \\ \vdots \end{matrix} \right.\right.\right.$	(not rendered)	
gilmore-lie-groups_EQ0686 (13.4) [conf 0.014]	220	0.014	$\begin{aligned} & \left[\frac{h}{[h',p']} = [h, \epsilon p] = \epsilon [h, p] \right. \\ & \left. \lim_{\epsilon \rightarrow 0} \frac{[p', p']}{\epsilon p} = \epsilon^2 [p, p] \right] \end{aligned}$	$\begin{aligned} & \lim_{\epsilon \rightarrow 0} \frac{[h', p']}{\epsilon p} = \epsilon^2 [p, p] \\ & \lim_{\epsilon \rightarrow 0} \frac{[p', p']}{\epsilon^2 (h + p)} = \epsilon^2 [p, p] \end{aligned}$	$\begin{aligned} & [h', h'] = [h, h] \subseteq h \\ & [h', p'] = [h, \epsilon p] = \epsilon [h, p] \lim_{\epsilon \rightarrow 0} \epsilon p \rightarrow p' \\ & [p', p'] = [\epsilon p, \epsilon p] = \epsilon^2 [p, p] \lim_{\epsilon \rightarrow 0} \epsilon^2 (h + p) \rightarrow 0 \end{aligned}$
gilmore-lie-groups_EQ0249 [conf 0.031]	84	0.031	$f_0(\theta) = \cos \theta \quad \text{or} \quad f_0(\theta) = \cos \theta \quad f_1(\theta) = \sin(\theta) / \theta \quad \theta f_1(\theta) = \sin(\theta) \quad f_2(\theta) = (1 - \cos(\theta)) / \theta^2 \quad \theta^2 f_2(\theta) = 1 - \cos(\theta),$	$f_0(\theta) = \cos \theta \quad \text{or} \quad f_0(\theta) = \cos \theta \quad f_1(\theta) = \sin(\theta) / \theta \quad \theta f_1(\theta) = \sin(\theta) \quad f_2(\theta) = (1 - \cos(\theta)) / \theta^2 \quad \theta^2 f_2(\theta) = 1 - \cos(\theta),$	$\begin{aligned} & f_0(\theta) = \cos \theta \quad \text{or} \quad f_0(\theta) = \cos \theta \\ & f_1(\theta) = \sin(\theta) / \theta \quad \theta f_1(\theta) = \sin(\theta) \\ & f_2(\theta) = (1 - \cos(\theta)) / \theta^2 \quad \theta^2 f_2(\theta) = 1 - \cos(\theta) \end{aligned}$
gilmore-lie-groups_EQ0368 (7.6) [conf 0.032]	114	0.032	$\operatorname{tr} e^{SXS^{-1}} \longrightarrow \left\{ \begin{array}{l} 2 \cosh \theta \geq 2 \\ 2 \cos \theta \geq -2 \end{array} \right. \text{ real eigenvalues}$	$\operatorname{tr} e^{SXS^{-1}} \longrightarrow \left\{ \begin{array}{l} 2 \cosh \theta \geq 2 \\ 2 \cos \theta \geq -2 \end{array} \right. \text{ imaginary eigenvalues}$	$\operatorname{tr} e^{SXS^{-1}} \longrightarrow \left\{ \begin{array}{l} 2 \cosh \theta \geq 2 \\ 2 \cos \theta \geq -2 \end{array} \right. \text{ real eigenvalues}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E09141 [conf 0.052]	56	■ 0.052	$\begin{array}{l} A^{t}=I_{3} \text { \& \& \text { quadratic } } \\ \text { constraints, } A \in O(3) \text { \& } C=0 \text { \& } B, D \text { \& } \\ \text { arbitrary linear constraints } \end{array}$	$\begin{array}{l} A A^{t}=I_{3} \\ C=0 \\ B, D \text { arbitrary } \end{array} \quad \begin{array}{l} \text { quadratic constraints, } A \in O(3) \\ \text { linear constraints} \end{array}$	$\begin{array}{l} A A^{t}=I_{3} \quad \text { quadratic constraints, } A \in O(3) \\ C=0 \quad \text { linear constraints} \\ B, D \text { arbitrary } \quad \text { no constraints} \end{array}$
gilmore-lie-groups_ E09142 (3.25) [conf 0.071]	56	■ 0.071	$\begin{array}{l} x^{\prime} \text { \& } y^{\prime} \text { \& } z^{\prime} \text { \& } \\ \text { \& } \end{array}$	$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} A & t_1 \\ t_2 & \\ t_3 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} A & x \\ y & y \\ z & z \end{bmatrix} + t_1 = +t_2.$	$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} A & t_1 \\ t_2 & \\ t_3 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} A & x \\ y & y \\ z & z \end{bmatrix} + t_1$
gilmore-lie-groups_ E09684 (13.2) [conf 0.084]	220	■ 0.084	$\begin{array}{l} \mathfrak{g}=\mathfrak{h}+\mathfrak{p} \\ \mathfrak{h}, \mathfrak{p} \subseteq \mathfrak{g} \\ \mathfrak{h}, \mathfrak{p} \subseteq \mathfrak{p} \end{array}$	$\begin{array}{l} \mathfrak{g}=\mathfrak{h}+\mathfrak{p} \\ \mathfrak{h}, \mathfrak{p} \subseteq \mathfrak{h} \\ \mathfrak{h}, \mathfrak{p} \subseteq \mathfrak{p} \end{array}$	$\begin{array}{l} \mathfrak{g}=\mathfrak{h}+\mathfrak{p} \\ \mathfrak{h}, \mathfrak{p} \subseteq \mathfrak{h} \\ \mathfrak{h}, \mathfrak{p} \subseteq \mathfrak{p} \end{array}$
gilmore-lie-groups_ E09923 (15.57)	286	■ 0.123	$\begin{array}{l} C_k k^0\rangle \\ \begin{array}{l} j \\ \mu \end{array} \end{array}$	$C_k k^0\rangle \left \begin{array}{l} j \\ \mu \end{array} \right\rangle = k\rangle \left \begin{array}{l} j \\ v \end{array} \right\rangle D_{vv';\mu\mu'}^{jj'}(C_k)$	$C_k k^0\rangle \left \begin{array}{l} j \\ \mu \end{array} \right\rangle = k\rangle \left \begin{array}{l} j \\ v \end{array} \right\rangle D_{vv';\mu\mu'}^{jj'}(C_k)$
gilmore-lie-groups_ E09591 (11.12)	191	■ 0.125	$\begin{array}{l} \mathfrak{g}=\mathfrak{h}+\mathfrak{p} \\ \mathfrak{h}, \mathfrak{p} \subseteq \mathfrak{g} \\ \mathfrak{h}, \mathfrak{p} \subseteq \mathfrak{p} \end{array}$	$\begin{array}{l} \mathfrak{g}=\mathfrak{h}+\mathfrak{p} \\ \mathfrak{h}, \mathfrak{p} \subseteq \mathfrak{h} \\ \mathfrak{h}, \mathfrak{p} \subseteq \mathfrak{p} \end{array}$	$\begin{array}{l} \mathfrak{g}=\mathfrak{h}+\mathfrak{p} \\ \mathfrak{h}, \mathfrak{p} \subseteq \mathfrak{h} \\ \mathfrak{h}, \mathfrak{p} \subseteq \mathfrak{p} \end{array}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00586 (11.6)	189	■ 0.128	$\left. \begin{array}{c} h^1 \\ \vdots \\ h^n \\ e^{+\alpha} \\ e^{-\alpha} \\ e^{+\beta} \\ e^{-\beta} \\ \vdots \end{array} \right\} \begin{array}{c} -1 \\ \\ -1 \\ \ddots \\ -1 \\ -1 \\ -1 \\ \ddots \end{array} \begin{array}{c} iH_1 \\ iH_2 \\ iH_3 \\ \vdots \\ iH_n \\ \frac{1}{\sqrt{2}}(E_\alpha + E_{-\alpha}) \\ i\frac{1}{\sqrt{2}}(E_\beta + E_{-\beta}) \\ \frac{1}{\sqrt{2}}(E_\beta - E_{-\beta}) \end{array}$	$\begin{bmatrix} h^1 & & & & & & \\ & h^2 & & & & & \\ & & h^3 & & & & \\ & & & \ddots & & & \\ & & & & -1 & & \\ & & & & & -1 & \\ & & & & & & -1 \\ & & & & & & & \ddots \end{bmatrix} \begin{array}{c} iH_1 \\ iH_2 \\ iH_3 \\ \vdots \\ iH_n \\ \frac{1}{\sqrt{2}}(E_\alpha + E_{-\alpha}) \\ i\frac{1}{\sqrt{2}}(E_\beta + E_{-\beta}) \\ \frac{1}{\sqrt{2}}(E_\beta - E_{-\beta}) \end{array}$	$\begin{bmatrix} h^1 & & & & & & \\ & h^2 & & & & & \\ & & h^3 & & & & \\ & & & \ddots & & & \\ & & & & -1 & & \\ & & & & & -1 & \\ & & & & & & -1 \\ & & & & & & & \ddots \end{bmatrix} \begin{array}{c} iH_1 \\ iH_2 \\ iH_3 \\ \vdots \\ iH_n \\ \frac{1}{\sqrt{2}}(E_\alpha + E_{-\alpha}) \\ i\frac{1}{\sqrt{2}}(E_\beta + E_{-\beta}) \\ \frac{1}{\sqrt{2}}(E_\beta - E_{-\beta}) \end{array}$
gilmore-lie-groups_ E00378 (7.16)	118	■ 0.146	$\begin{aligned} &\text{algebra} = \text{noncompact generators} + \text{compact generators} \\ &\text{EXP} \downarrow \\ &\text{EXP} \downarrow \\ &\text{group} = \text{coset representatives} \end{aligned}$	$\begin{aligned} &\text{algebra} = \text{noncompact generators} + \text{compact generators} \\ &\text{EXP} \downarrow \\ &\text{EXP} \downarrow \\ &\text{group} = \text{coset representatives} \end{aligned}$	$\begin{aligned} &\text{algebra} = \text{noncompact generators} + \text{compact generators} \\ &\text{EXP} \downarrow \quad \text{EXP} \downarrow \quad \downarrow \quad \downarrow \text{EXP} \\ &\text{group} = \text{coset representatives} \times \text{compact subgroup} \end{aligned}$
gilmore-lie-groups_ E00687 (13.5)	220	■ 0.160	$\begin{aligned} &\mathfrak{g} = \mathfrak{h} + \mathfrak{p} \quad \mathfrak{p} \rightarrow \mathfrak{p}' = \epsilon \mathfrak{p} \\ &[\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h} \\ &[\mathfrak{h}, \mathfrak{p}] \subseteq \mathfrak{p} \\ &[\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h} + \mathfrak{p} \end{aligned}$	$\begin{aligned} &\mathfrak{g} = \mathfrak{h} + \mathfrak{p} \quad \mathfrak{p} \rightarrow \mathfrak{p}' = \epsilon \mathfrak{p} \\ &[\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h} \\ &[\mathfrak{h}, \mathfrak{p}] \subseteq \mathfrak{p} \\ &[\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h} + \mathfrak{p} \end{aligned}$	$\begin{aligned} &\mathfrak{g}' = \mathfrak{h} + \mathfrak{p}' \\ &[\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h} \\ &[\mathfrak{h}, \mathfrak{p}] \subseteq \mathfrak{p} \\ &[\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h} + \mathfrak{p} \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0478 (8.10)	146	■ 0.182	$\begin{aligned} X &= \sum y_i Y_i + \sum z_j Z_j \\ \mathfrak{def}(X) &\rightarrow \begin{bmatrix} 0 & +(y_3 - z_3) & -(y_2 - z_2) & +(y_1 + z_1) \\ -(y_3 - z_3) & 0 & +(y_1 - z_1) & +(y_2 + z_2) \\ +(y_2 - z_2) & -(y_1 - z_1) & 0 & +(y_3 + z_3) \\ -(y_1 + z_1) & -(y_2 + z_2) & -(y_3 + z_3) & 0 \end{bmatrix} \\ \mathfrak{Reg}(X) &= \left[\begin{array}{ccc ccc} 0 & -y_3 & +y_2 & & & \\ +y_3 & 0 & -y_1 & & & \\ -y_2 & +y_1 & 0 & & & \\ \hline & & & -z_3 & 0 & +z_1 \\ & & & +z_2 & -z_1 & 0 \end{array} \right] \begin{matrix} Y_1 \\ Y_2 \\ Y_3 \\ Z_1 \\ Z_2 \\ Z_3 \end{matrix} \end{aligned}$	$X = \sum y_i Y_i + \sum z_j Z_j$ $\mathfrak{def}(X) \rightarrow \begin{bmatrix} 0 & +(y_3 - z_3) & -(y_2 - z_2) & +(y_1 + z_1) \\ -(y_3 - z_3) & 0 & +(y_1 - z_1) & +(y_2 + z_2) \\ +(y_2 - z_2) & -(y_1 - z_1) & 0 & +(y_3 + z_3) \\ -(y_1 + z_1) & -(y_2 + z_2) & -(y_3 + z_3) & 0 \end{bmatrix}$ $\mathfrak{Reg}(X) = \left[\begin{array}{ccc ccc} 0 & -y_3 & +y_2 & & & \\ +y_3 & 0 & -y_1 & & & \\ -y_2 & +y_1 & 0 & & & \\ \hline & & & -z_3 & 0 & +z_1 \\ & & & +z_2 & -z_1 & 0 \end{array} \right] \begin{matrix} Y_1 \\ Y_2 \\ Y_3 \\ Z_1 \\ Z_2 \\ Z_3 \end{matrix}$	
gilmore-lie-groups_ EQ0097	44	■ 0.191	$\left[\begin{array}{c} x' \\ y' \\ z' \end{array}\right] = \left[\begin{array}{c c} M_1 & a \\ b & b \\ a & \\ M_2 & \end{array}\right] \left[\begin{array}{c c} SO(2) & 0 \\ 0 & 0 \end{array}\right] \left[\begin{array}{c} x \\ y \\ z \end{array}\right]$	$\left[\begin{array}{c} x' \\ y' \\ z' \end{array}\right] = \left[\begin{array}{c c} M_1 & a \\ b & b \\ a & \\ M_2 & \end{array}\right] \left[\begin{array}{c c} SO(2) & 0 \\ 0 & 0 \end{array}\right] \left[\begin{array}{c} x \\ y \\ z \end{array}\right]$	
gilmore-lie-groups_ EQ0770 (14.12)	240	■ 0.195	$\begin{aligned} &\text{Klein-Gordon equation} \\ E(n, l) &= \text{Schrödinger equation} \\ \sqrt{1 + (\alpha/N')^2} W(n, l) &= -\frac{1}{2} m c^2 \alpha^2 \frac{1}{N^2} \quad N = n + l + 1 \\ N' &= n + \frac{1}{2} + \sqrt{\left(l + \frac{1}{2}\right)^2 - \alpha^2} \end{aligned}$	Klein–Gordon equation $E(n, l) = \frac{m c^2}{\sqrt{1 + (\alpha/N')^2}}$ Schrödinger equation $W(n, l) = -\frac{1}{2} m c^2 \alpha^2 \frac{1}{N^2}$ $N' = n + \frac{1}{2} + \sqrt{\left(l + \frac{1}{2}\right)^2 - \alpha^2} \quad N = n + l + 1$	

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0384 (7.21)	120	■ 0.217	$\begin{aligned} \mathfrak{so}(3) = & \left\{ \begin{bmatrix} 0 & 0 & -a_2 \\ 0 & 0 & a_1 \\ a_2 & -a_1 & 0 \end{bmatrix} \mid a_1, a_2 \in \mathbb{R} \right\} \\ & \oplus \left\{ \begin{bmatrix} \cos a_3 & \sin a_3 & 0 \\ -\sin a_3 & \cos a_3 & 0 \\ 0 & 0 & 1 \end{bmatrix} \mid a_3 \in \mathbb{R} \right\} \end{aligned}$	$\mathfrak{so}(3) = \left\{ \begin{bmatrix} 0 & 0 & -a_2 \\ 0 & 0 & a_1 \\ a_2 & -a_1 & 0 \end{bmatrix} \mid a_1, a_2 \in \mathbb{R} \right\} \oplus \left\{ \begin{bmatrix} \cos a_3 & \sin a_3 & 0 \\ -\sin a_3 & \cos a_3 & 0 \\ 0 & 0 & 1 \end{bmatrix} \mid a_3 \in \mathbb{R} \right\}$	$\mathfrak{so}(3) = \left\{ \begin{bmatrix} 0 & 0 & -a_2 \\ 0 & 0 & a_1 \\ a_2 & -a_1 & 0 \end{bmatrix} \mid a_1, a_2 \in \mathbb{R} \right\} \oplus \left\{ \begin{bmatrix} \cos a_3 & \sin a_3 & 0 \\ -\sin a_3 & \cos a_3 & 0 \\ 0 & 0 & 1 \end{bmatrix} \mid a_3 \in \mathbb{R} \right\}$
gilmore-lie-groups_ EQ0624 (1.20)	24	■ 0.222	$\left(\left(r_1 - r_2 \right) \right) \xrightarrow{I} (r_1 - r_2),$	$(r_1 - r_2) \xrightarrow{I} (r_1 - r_2),$	$(r_1 - r_2) \left\{ \begin{array}{l} \xrightarrow{I} (r_1 - r_2) \\ \xrightarrow{(12)} (r_2 - r_1) = -(r_1 - r_2) \end{array} \right.$
gilmore-lie-groups_ EQ0474 (8.6)	145	■ 0.233	$\begin{aligned} & \eta \left(\hat{n} + \frac{1}{2} \right) + R a^{\dagger 2} + L a^2 + r a^{\dagger} + l a + \delta I \\ & \mathfrak{d} \mathfrak{e} \mathfrak{f} = \begin{bmatrix} 0 & l & r & -2\delta \\ 0 & \eta & 2R & -r \\ 0 & -2L & -\eta & l \\ 0 & 0 & 0 & 0 \end{bmatrix} \\ & \mathfrak{R} \mathfrak{e} \mathfrak{g} = \left[\begin{array}{ccc ccc} 0 & -2R & 2L & -r & l & 0 \\ 4L & 2\eta & 0 & 2l & 0 & 0 \\ -4R & 0 & -2\eta & 0 & -2r & 0 \\ \hline & & & \eta & 2L & l \\ & & & -2R & -\eta & -r \\ & & & 0 & 0 & 0 \end{array} \right] \begin{matrix} \hat{n} + \frac{1}{2} \\ a^{\dagger 2} \\ a^2 \\ a^{\dagger} \\ a \\ I \end{matrix} \end{aligned}$	(not rendered)	$\eta \left(\hat{n} + \frac{1}{2} \right) + R a^{\dagger 2} + L a^2 + r a^{\dagger} + l a + \delta I$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0897 (15.31)	281	■ 0.247	$\begin{array}{lll} \text{ (i) } & k \cdot k > 0 & k^0 = (0, 0, 1, 0) \\ \text{ (ii) } & k \cdot k = 0 & k \neq 0 \quad k^0 = (0, 0, 1, +i) \quad \text{ (a)} \\ & & k^0 = (0, 0, 1, -i) \quad \text{ (b)} \\ \text{ (iii) } & k \cdot k < 0 & k^0 = (0, 0, 0, +i) \quad \text{ (a)} \\ & & k^0 = (0, 0, 0, -i) \quad \text{ (b)} \\ \text{ (iv) } & k \cdot k = 0 & k = 0 \quad k^0 = (0, 0, 0, 0) \end{array}$	$\begin{array}{ll} \text{ (i) } & k \cdot k > 0 \quad k^0 = (0, 0, 1, 0) \\ \text{ (ii) } & k \cdot k = 0 \quad k \neq 0 \\ & k^0 = (0, 0, 1, +i) \\ & k^0 = (0, 0, 1, -i) \\ \text{ (iii) } & k \cdot k < 0 \quad k^0 = (0, 0, 0, +i) \\ & k^0 = (0, 0, 0, -i) \\ \text{ (iv) } & k \cdot k = 0 \quad k = 0 \end{array}$	$\begin{array}{ll} \text{ (i) } & k \cdot k > 0 \quad k^0 = (0, 0, 1, 0) \\ \text{ (ii) } & k \cdot k = 0 \quad k \neq 0 \quad k^0 = (0, 0, 1, +i) \quad \text{ (a)} \\ & k^0 = (0, 0, 1, -i) \quad \text{ (b)} \\ \text{ (iii) } & k \cdot k < 0 \quad k^0 = (0, 0, 0, +i) \quad \text{ (a)} \\ & k^0 = (0, 0, 0, -i) \quad \text{ (b)} \\ \text{ (iv) } & k \cdot k = 0 \quad k = 0 \quad k^0 = (0, 0, 0, 0) \end{array}$
gilmore-lie-groups_ EQ0726 (13.44)	228	■ 0.260	$\begin{aligned} & a^{\dagger} a \left \begin{array}{c} \infty \\ n \end{array} \right\rangle = \lim_{c \rightarrow 0} \left[J_3 + \frac{1}{2c^2} \right] \left \begin{array}{c} J \\ M \end{array} \right\rangle \\ & = \lim_{c \rightarrow 0} \left[J + M + \left(\frac{1}{2c^2} - J \right) \right] \left \begin{array}{c} J \\ M = n - J \end{array} \right\rangle \rightarrow (n + 0) \left \begin{array}{c} \infty \\ n \end{array} \right\rangle \\ & a^{\dagger} \left \begin{array}{c} \infty \\ n \end{array} \right\rangle = \lim_{c \rightarrow 0} c J_+ \left \begin{array}{c} J \\ M \end{array} \right\rangle \\ & = \lim_{c \rightarrow 0} \left \begin{array}{c} J \\ M + 1 \end{array} \right\rangle \sqrt{(J - M)(J + M + 1)c^2} \rightarrow \sqrt{n + 1} \left \begin{array}{c} \infty \\ n + 1 \end{array} \right\rangle \\ & a \left \begin{array}{c} \infty \\ n \end{array} \right\rangle = \lim_{c \rightarrow 0} c J_- \left \begin{array}{c} J \\ M \end{array} \right\rangle \\ & = \lim_{c \rightarrow 0} \left \begin{array}{c} J \\ M - 1 \end{array} \right\rangle \sqrt{(J + M)(J - M + 1)c^2} \rightarrow \sqrt{n} \left \begin{array}{c} \infty \\ n - 1 \end{array} \right\rangle \end{aligned}$	$\begin{aligned} & a^{\dagger} a \left \begin{array}{c} \infty \\ n \end{array} \right\rangle = \lim_{c \rightarrow 0} \left(J_3 + \frac{1}{2c^2} \right) \left \begin{array}{c} J \\ M \end{array} \right\rangle \\ & = \lim_{c \rightarrow 0} \left[J + M + \left(\frac{1}{2c^2} - J \right) \right] \left \begin{array}{c} J \\ M = n - J \end{array} \right\rangle \rightarrow (n + 0) \left \begin{array}{c} \infty \\ n \end{array} \right\rangle \\ & a^{\dagger} \left \begin{array}{c} \infty \\ n \end{array} \right\rangle = \lim_{c \rightarrow 0} c J_+ \left \begin{array}{c} J \\ M \end{array} \right\rangle \\ & = \lim_{c \rightarrow 0} \left \begin{array}{c} J \\ M + 1 \end{array} \right\rangle \sqrt{(J - M)(J + M + 1)c^2} \rightarrow \sqrt{n + 1} \left \begin{array}{c} \infty \\ n + 1 \end{array} \right\rangle \\ & a \left \begin{array}{c} \infty \\ n \end{array} \right\rangle = \lim_{c \rightarrow 0} c J_- \left \begin{array}{c} J \\ M \end{array} \right\rangle \\ & = \lim_{c \rightarrow 0} \left \begin{array}{c} J \\ M - 1 \end{array} \right\rangle \sqrt{(J + M)(J - M + 1)c^2} \rightarrow \sqrt{n} \left \begin{array}{c} \infty \\ n - 1 \end{array} \right\rangle \end{aligned}$	$\begin{aligned} & a^{\dagger} a \left \begin{array}{c} \infty \\ n \end{array} \right\rangle = \lim_{c \rightarrow 0} \left(J_3 + \frac{1}{2c^2} \right) \left \begin{array}{c} J \\ M \end{array} \right\rangle \\ & = \lim_{c \rightarrow 0} \left[J + M + \left(\frac{1}{2c^2} - J \right) \right] \left \begin{array}{c} J \\ M = n - J \end{array} \right\rangle \rightarrow (n + 0) \left \begin{array}{c} \infty \\ n \end{array} \right\rangle \\ & a^{\dagger} \left \begin{array}{c} \infty \\ n \end{array} \right\rangle = \lim_{c \rightarrow 0} c J_+ \left \begin{array}{c} J \\ M \end{array} \right\rangle \\ & = \lim_{c \rightarrow 0} \left \begin{array}{c} J \\ M + 1 \end{array} \right\rangle \sqrt{(J - M)(J + M + 1)c^2} \rightarrow \sqrt{n + 1} \left \begin{array}{c} \infty \\ n + 1 \end{array} \right\rangle \\ & a \left \begin{array}{c} \infty \\ n \end{array} \right\rangle = \lim_{c \rightarrow 0} c J_- \left \begin{array}{c} J \\ M \end{array} \right\rangle \\ & = \lim_{c \rightarrow 0} \left \begin{array}{c} J \\ M - 1 \end{array} \right\rangle \sqrt{(J + M)(J - M + 1)c^2} \rightarrow \sqrt{n} \left \begin{array}{c} \infty \\ n - 1 \end{array} \right\rangle \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0778	243	■ 0.264	$\begin{array}{ll} \angle \theta \ \phi L_{-} \theta' \phi' \rangle \\ \theta^{\prime} \phi^{\prime} \rangle \\ \theta^{\prime} \phi^{\prime} \rangle \end{array}$ Matrix element of the angular momentum shift down operator in the coordinate representation: $\hbar(-\partial/\partial\theta + i(\cos\theta/\sin\theta)(\partial/\partial\phi))\delta(\cos\theta' - \cos\theta)\delta(\phi' - \phi)$. Matrix element of the angular momentum shift down operator in the algebraic representation: $\hbar\sqrt{(l'-m')(l+m)}$. Matrix element of the similarity transformation between the coordinate representation and algebraic representation. Also called spherical harmonic: $Y_m^l(\theta, \phi)$.	$\langle \theta \phi L_{-} \theta' \phi' \rangle$ $\langle \theta \phi L_{-} \theta' \phi' \rangle$ $\langle \theta \phi L_{-} \theta' \phi' \rangle$	Matrix element of the angular momentum shift down operator in the coordinate representation: $\hbar(-\partial/\partial\theta + i(\cos\theta/\sin\theta)(\partial/\partial\phi))\delta(\cos\theta' - \cos\theta)\delta(\phi' - \phi)$. Matrix element of the angular momentum shift down operator in the algebraic representation: $\hbar\sqrt{(l'-m')(l+m)}$. Matrix element of the similarity transformation between the coordinate representation and algebraic representation. Also called spherical harmonic: $Y_m^l(\theta, \phi)$.
gilmore-lie-groups_ EQ0712 (13.29)	225	■ 0.265	$\begin{array}{ll} \left[\begin{array}{c} h_{+} \\ h_{-} \\ h_3 \\ h_0 \end{array} \right] = \left[\begin{array}{ccc} c & & \\ & c & \\ & & 1 \end{array} \right] \left[\begin{array}{c} J_{+} \\ J_{-} \\ J_3 \\ J_0 \end{array} \right] \\ c \ \& \ \& \ 1 \ \& \ \frac{1}{2c^2} \end{array} \right] \left[\begin{array}{c} J_{+} \\ J_{-} \\ J_3 \\ J_0 \end{array} \right] \\ \right] \left[\begin{array}{c} J_{+} \\ J_{-} \\ J_3 \\ J_0 \end{array} \right] \end{array}$	$\left[\begin{array}{c} h_{+} \\ h_{-} \\ h_3 \\ h_0 \end{array} \right] = \left[\begin{array}{ccc} c & & \\ & c & \\ & & 1 \end{array} \right] \left[\begin{array}{c} J_{+} \\ J_{-} \\ J_3 \\ J_0 \end{array} \right]$	$\left[\begin{array}{c} h_{+} \\ h_{-} \\ h_3 \\ h_0 \end{array} \right] = \left[\begin{array}{ccc} c & & \\ & c & \\ & & 1 \end{array} \right] \left[\begin{array}{c} J_{+} \\ J_{-} \\ J_3 \\ J_0 \end{array} \right]$
gilmore-lie-groups_ EQ0753	234	■ 0.270	$\mathfrak{g} = \mathfrak{h} + \mathfrak{p} \quad [\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h}, \quad [\mathfrak{h}, \mathfrak{p}] = \mathfrak{p}, \quad [\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h}$	$\mathfrak{g} = \mathfrak{h} + \mathfrak{p} \quad [\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h}, \quad [\mathfrak{h}, \mathfrak{p}] = \mathfrak{p}, \quad [\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h}$	$\mathfrak{g} = \mathfrak{h} + \mathfrak{p} \quad [\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h}, \quad [\mathfrak{h}, \mathfrak{p}] = \mathfrak{p}, \quad [\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h}$
gilmore-lie-groups_ EQ0070 (1.65)	34	■ 0.272	$\begin{array}{ll} v_2^3 + v_3^3 = & 27 J_3' = & 27 \times 518400 = & 1399680 \\ J_3^{\prime} = & 27 \times 518400 = & 1399680 \\ v_2^3 v_3^3 = & -27 J_2^{\prime 3} = & -27 \times (-2736)^3 = & 552983334912 \end{array}$	$v_2^3 + v_3^3 = 27 J_3' = 27 \times 518400 = 1399680$ $v_2^3 v_3^3 = -27 J_2^{\prime 3} = -27 \times (-2736)^3 = 552983334912$	$v_2^3 + v_3^3 = 27 J_3' = 27 \times 518400 = 1399680$ $v_2^3 v_3^3 = -27 J_2^{\prime 3} = -27 \times (-2736)^3 = 552983334912$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0311	100	■ 0.278	$\begin{array}{l} \backslash\mathrm{begin}\{\mathrm{array}\}\{\mathrm{ll}\}\{\backslash\mathrm{mathfrak{h}},\backslash\mathrm{mathfrak{h}}\}\backslash\mathrm{subse} \\ \mathrm{teq}\backslash\mathrm{mathfrak{h}}\}\&(\backslash\mathrm{mathfrak{h}},\backslash\mathrm{mathfrak{h}})\backslash\mathrm{leq}\,0\backslash \\ \backslash\{\backslash\mathrm{left}[\backslash\mathrm{mathfrak{h}},\backslash\mathrm{mathfrak{p}}^{\backslash\mathrm{prime}}\backslash\mathrm{right}]\backslash \\ \mathrm{subse} \\ \mathrm{teq}\backslash\mathrm{mathfrak{p}}^{\backslash\mathrm{prime}}\}\backslash\backslash\{\backslash\mathrm{left}[\backslash\mathrm{mathfrak{p}}^{\backslash\mathrm{prime}}\backslash\mathrm{right}]\backslash\mathrm{subse} \\ \mathrm{teq}\backslash\mathrm{mathfrak{h}}\}\&\backslash\mathrm{left}(\backslash\mathrm{mathfrak{h}},\backslash\mathrm{mathfrak{p}}^{\backslash\mathrm{prime}}\backslash\mathrm{right})=0\backslash\backslash\&\backslash\mathrm{left}(\backslash\mathrm{mathfrak{p}}^{\backslash\mathrm{prime}}, \\ \backslash\mathrm{mathfrak{p}}^{\backslash\mathrm{prime}}\backslash\mathrm{right})\backslash\mathrm{leq}\,0\backslash\mathrm{end}\{\mathrm{array}\} \end{array}$	$\begin{array}{l} [\mathfrak{h},\mathfrak{h}]\subseteq\mathfrak{h}\qquad(\mathfrak{h},\mathfrak{h})\leq 0 \\ [\mathfrak{h},\mathfrak{p}']\subseteq\mathfrak{p}' \\ [\mathfrak{p}',\mathfrak{p}']\subseteq\mathfrak{h}\qquad(\mathfrak{h},\mathfrak{p}')=0 \\ \qquad\qquad\qquad(\mathfrak{p}',\mathfrak{p}')\leq 0 \end{array}$	$\begin{array}{l} [\mathfrak{h},\mathfrak{h}]\subseteq\mathfrak{h}\qquad(\mathfrak{h},\mathfrak{h})\leq 0 \\ [\mathfrak{h},\mathfrak{p}']\subseteq\mathfrak{p}'\quad\text{and}\quad(\mathfrak{h},\mathfrak{p}')=0 \\ [\mathfrak{p}',\mathfrak{p}']\subseteq\mathfrak{h}\qquad(\mathfrak{p}',\mathfrak{p}')\leq 0 \end{array}$
gilmore-lie-groups_ EQ0789 (14.31)	247	■ 0.300	$\begin{array}{l} \backslash\mathrm{begin}\{\mathrm{aligned}\}\&\mathrm{M}_{\{z\}^{\backslash\mathrm{prime}}}\backslash\mathrm{left} \mathrm{N}_{\{\backslash\mathrm{pm}\,l\}^{\backslash\mathrm{l}}}\backslash \\ \mathrm{right}\backslash\mathrm{rangle}=\mathrm{D}_{\{1\}}\backslash\mathrm{left} \backslash\mathrm{begin}\{\mathrm{array}\}\{\mathrm{c}\}\,l+1\backslash\backslash\backslash\mathrm{pm} \\ l\backslash\mathrm{end}\{\mathrm{array}\}\backslash\mathrm{right}\backslash\mathrm{rangle}\quad\mathrm{D}_{\{1\}}=\mathrm{frac}\{1\}\{\mathrm{N}\} \\ \backslash\mathrm{sqrt}\{\backslash\mathrm{frac}\{\mathrm{N}^{\{2\}}-(l+1)^{\{2\}}\}\{2\,l+3\}\}\backslash\backslash\&\mathrm{M}_{\{\backslash\mathrm{pm}\,l\}^{\backslash\mathrm{prime}}}\backslash\mathrm{left} \backslash\mathrm{begin}\{\mathrm{array}\}\{\mathrm{c}\}\,\mathrm{N}^{\{2\}}\backslash\backslash\backslash\mathrm{pm}\,l\backslash\mathrm{end}\{\mathrm{array}\} \\ \backslash\mathrm{right}\backslash\mathrm{rangle}=\backslash\mathrm{pm}\,\mathrm{D}_{\{2\}}\backslash\mathrm{left} \backslash\mathrm{begin}\{\mathrm{array}\}\{\mathrm{c}\}\,l+1\backslash\backslash \\ \backslash\mathrm{pm}(l+1)\backslash\mathrm{end}\{\mathrm{array}\}\backslash\mathrm{right}\backslash\mathrm{rangle}\quad\mathrm{D}_{\{2\}}=\mathrm{frac}\{1\}\{\mathrm{N}\} \\ \backslash\mathrm{sqrt}\{\backslash\mathrm{frac}\{2\,l+2\}\{2\,l+3\}\backslash\mathrm{left}[\mathrm{N}^{\{2\}}-(l+1)^{\{2\}}\backslash\mathrm{right}]\}\backslash \\ \mathrm{end}\{\mathrm{aligned}\} \end{array}$	$\begin{array}{l} M'_z\left N_{\pm l}^l\right\rangle = D_1\left \begin{array}{c} l+1 \\ \pm l \end{array}\right\rangle \quad D_1 = \frac{1}{N}\sqrt{\frac{N^2-(l+1)^2}{2l+3}} \\ M'_{\pm}\left \begin{array}{c} N^2 \\ \pm l \end{array}\right\rangle = \pm D_2\left \begin{array}{c} l+1 \\ \pm(l+1) \end{array}\right\rangle \quad D_2 = \frac{1}{N}\sqrt{\frac{2l+2}{2l+3}}[N^2-(l+1)^2] \end{array}$	$\begin{array}{l} M'_z\left N\begin{array}{c} l \\ \pm l \end{array}\right\rangle = D_1\left N\begin{array}{c} l+1 \\ \pm l \end{array}\right\rangle \quad D_1 = \frac{1}{N}\sqrt{\frac{N^2-(l+1)^2}{2l+3}} \\ M'_{\pm}\left N\begin{array}{c} l \\ \pm l \end{array}\right\rangle = \pm D_2\left N\begin{array}{c} l+1 \\ \pm(l+1) \end{array}\right\rangle \quad D_2 = \frac{1}{N}\sqrt{\frac{2l+2}{2l+3}}[N^2-(l+1)^2] \end{array}$
gilmore-lie-groups_ EQ0398 (7.34)	124	■ 0.317	$\begin{array}{l} \backslash\mathrm{begin}\{\mathrm{gathered}\}\backslash\mathrm{operatorname}\{\mathrm{EXP}\}\backslash\mathrm{left}[\\ \backslash\mathrm{begin}\{\mathrm{array}\}\{\mathrm{lll}\}\,0\&l\&\backslash\mathrm{delta}\backslash\backslash\,0\&0\&r\backslash\backslash\,0\& \\ 0\&0\backslash\mathrm{end}\{\mathrm{array}\}\backslash\mathrm{right}]=\mathrm{frac}\{\backslash\mathrm{left}[\backslash\mathrm{begin}\{\mathrm{array}\}\{\mathrm{ccc}\} \\ 1\&l\&\backslash\mathrm{delta}+\mathrm{frac}\{1\}\{2\}\,l\,r\backslash\backslash\,0\&1\&r\backslash\backslash\,0\&0 \\ \&1\backslash\mathrm{end}\{\mathrm{array}\}\backslash\mathrm{right}]\}\{\backslash \}\backslash\backslash\backslash \backslash\backslash\mathrm{e}^{\{r^{\backslash\mathrm{prime}}\} \\ \mathrm{a}^{\backslash\mathrm{dagger}}\}\}\mathrm{e}^{\{\backslash\mathrm{delta}^{\backslash\mathrm{prime}}\}\,I}\}\mathrm{e}^{\{l^{\backslash\mathrm{prime}}\}\,\mathrm{a}\} \\ \backslash\mathrm{rightarrow}\backslash\mathrm{left}[\backslash\mathrm{begin}\{\mathrm{array}\}\{\mathrm{ccc}\}\,1\&l^{\backslash\mathrm{prime}}\& \\ \backslash\mathrm{delta}^{\backslash\mathrm{prime}}\backslash\backslash\,0\&1\&r^{\backslash\mathrm{prime}}\backslash\backslash\,0\&0\&1 \\ \backslash\mathrm{end}\{\mathrm{array}\}\backslash\mathrm{right}]=\backslash\mathrm{left}[\backslash\mathrm{begin}\{\mathrm{array}\}\{\mathrm{ccc}\}\,1\&l^{\backslash\mathrm{prime}} \\ \backslash\mathrm{prime}\}\&\backslash\mathrm{delta}^{\backslash\mathrm{prime}}\backslash\mathrm{prime}\}+l^{\backslash\mathrm{prime}}\backslash\mathrm{prime}\}\,r^{\backslash\mathrm{prime}} \\ \backslash\mathrm{prime}\}\backslash\backslash\,0\&1\&r^{\backslash\mathrm{prime}}\backslash\mathrm{prime}\}\backslash\backslash\,0\&0\&1 \\ \backslash\mathrm{end}\{\mathrm{array}\}\backslash\mathrm{right}]\backslash\mathrm{leftarrow}\mathrm{e}^{\{l^{\backslash\mathrm{prime}}\}\backslash\mathrm{prime}\,\mathrm{prime}\}\,\mathrm{a}\} \\ \mathrm{e}^{\{\backslash\mathrm{delta}^{\backslash\mathrm{prime}}\}\backslash\mathrm{prime}\,\mathrm{prime}\}\,I}\}\mathrm{e}^{\{r^{\backslash\mathrm{prime}}\}\backslash\mathrm{prime}\,\mathrm{prime}\}\,\mathrm{a}^{\backslash\mathrm{dagger}}\}\}\backslash\mathrm{end}\{\mathrm{gathered}\} \end{array}$	$\begin{array}{l} \mathrm{EXP}\left[\begin{array}{ccc} 0 & l & \delta \\ 0 & 0 & r \\ 0 & 0 & 0 \end{array}\right] = \frac{\left[\begin{array}{ccc} 1 & l & \delta + \frac{1}{2}lr \\ 0 & 1 & r \\ 0 & 0 & 1 \end{array}\right]}{\parallel} \\ \parallel \\ e^{r'a^\dagger}e^{\delta'I}e^{l'a} \rightarrow \left[\begin{array}{ccc} 1 & l' & \delta' \\ 0 & 1 & r' \\ 0 & 0 & 1 \end{array}\right] = \left[\begin{array}{ccc} 1 & l'' & \delta'' + l''r'' \\ 0 & 1 & r'' \\ 0 & 0 & 1 \end{array}\right] \leftarrow e^{l''a}e^{\delta''I}e^{r''a^\dagger} \end{array}$	$\begin{array}{l} \mathrm{EXP}\left[\begin{array}{ccc} 0 & l & \delta \\ 0 & 0 & r \\ 0 & 0 & 0 \end{array}\right] = \left[\begin{array}{ccc} 1 & l & \delta + \frac{1}{2}lr \\ 0 & 1 & r \\ 0 & 0 & 1 \end{array}\right] \\ \parallel \\ e^{r'a^\dagger}e^{\delta'I}e^{l'a} \rightarrow \left[\begin{array}{ccc} 1 & l' & \delta' \\ 0 & 1 & r' \\ 0 & 0 & 1 \end{array}\right] = \left[\begin{array}{ccc} 1 & l'' & \delta'' + l''r'' \\ 0 & 1 & r'' \\ 0 & 0 & 1 \end{array}\right] \leftarrow e^{l''a}e^{\delta''I}e^{r''a^\dagger} \end{array}$
gilmore-lie-groups_ EQ0544	170	■ 0.319	$\frac{\mathrm{dim}(\mathrm{operatorname}\{\mathrm{dim}\}(\backslash\mathrm{mathfrak{g}}))}{\{\mathrm{rank}(\backslash\mathrm{mathfrak{g}})\}^2}=\left\{\begin{array}{lll} \text{ratio} & n\rightarrow\infty & \text{algebra} \\ 2-\frac{1}{n} & 2 & A_n \\ 2+\frac{1}{n} & 2 & B_n, C_n \end{array}\right.$	$\frac{\mathrm{dim}(\mathfrak{g})}{\{\mathrm{rank}(\mathfrak{g})\}^2}=\left\{\begin{array}{lll} \text{ratio} & n\rightarrow\infty & \text{algebra} \\ 2-\frac{1}{n} & 2 & A_n \\ 2+\frac{1}{n} & 2 & B_n, C_n \end{array}\right.$	$\frac{\mathrm{dim}(\mathfrak{g})}{\{\mathrm{rank}(\mathfrak{g})\}^2}=\begin{array}{ccc} \text{ratio} & n\rightarrow\infty & \text{algebra} \\ \left\{\begin{array}{l} 1+\frac{2}{n} \\ 2-\frac{1}{n} \\ 2+\frac{1}{n} \end{array}\right. & \begin{array}{l} 1 \\ 2 \\ 2 \end{array} & \begin{array}{l} A_n \\ D_n \\ B_n, C_n \end{array} \end{array}$


Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ8581 (9.10)	156	■ 0.333	$\begin{aligned} \mathfrak{d e f}(Y) &= \frac{1}{2} \left[\begin{array}{cc} i b_3 & b_1 - i b_2 \\ b_1 + i b_2 & -i b_3 \end{array} \right] \\ \mathfrak{R e g}(y) &= \left[\begin{array}{ccc} 0 & -b_3 & -b_2 \\ b_3 & 0 & b_1 \\ -b_2 & b_1 & 0 \end{array} \right] \end{aligned}$	$\mathfrak{d e f}(Y) = \frac{1}{2} \left[\begin{array}{cc} i b_3 & b_1 - i b_2 \\ b_1 + i b_2 & -i b_3 \end{array} \right]$ $\mathfrak{R e g}(y) = \left[\begin{array}{ccc} 0 & -b_3 & -b_2 \\ b_3 & 0 & b_1 \\ -b_2 & b_1 & 0 \end{array} \right]$	$\mathfrak{d e f}(Y) = \frac{1}{2} \left[\begin{array}{cc} i b_3 & b_1 - i b_2 \\ b_1 + i b_2 & -i b_3 \end{array} \right]$ $\mathfrak{R e g}(y) = \left[\begin{array}{ccc} 0 & -b_3 & -b_2 \\ b_3 & 0 & b_1 \\ -b_2 & b_1 & 0 \end{array} \right]$
gilmore-lie-groups_ EQ8377 (7.15)	118	■ 0.340	$\begin{aligned} &\left[\begin{array}{cc} a & b \\ b & -a \end{array} \right] = \left[\begin{array}{cc} a & b+c \\ b-c & -a \end{array} \right] - \left[\begin{array}{cc} 0 & c \\ -c & 0 \end{array} \right] \\ &\quad \text{EXP} \downarrow \text{EXP} \downarrow \\ &\left[\begin{array}{cc} z+y & x \\ x & z-y \end{array} \right] = \text{SL}(2; \mathbb{R}) \end{aligned}$	$\left[\begin{array}{cc} a & b \\ b & -a \end{array} \right] = \left[\begin{array}{cc} a & b+c \\ b-c & -a \end{array} \right] - \left[\begin{array}{cc} 0 & c \\ -c & 0 \end{array} \right]$ $\text{EXP} \downarrow \text{EXP} \downarrow$ $\left[\begin{array}{cc} z+y & x \\ x & z-y \end{array} \right] = \text{SL}(2; \mathbb{R})$	$\left[\begin{array}{cc} a & b \\ b & -a \end{array} \right] = \left[\begin{array}{cc} a & b+c \\ b-c & -a \end{array} \right] - \left[\begin{array}{cc} 0 & c \\ -c & 0 \end{array} \right]$ $\text{EXP} \downarrow \quad \text{EXP} \downarrow \quad \downarrow \text{EXP} \downarrow \text{EXP}$ $\left[\begin{array}{cc} z+y & x \\ x & z-y \end{array} \right] = \text{SL}(2; \mathbb{R}) \quad / \quad \text{SO}(2)$
gilmore-lie-groups_ EQ8479 (8.11)	147	■ 0.354	$\mathfrak{d e f} = \frac{i}{2} \left[\begin{array}{cc} a_3 & a_1 - i a_2 \\ a_1 + i a_2 & -a_3 \end{array} \right] \mathfrak{R e g} = \left[\begin{array}{ccc} 0 & -a_3 & +a_2 \\ +a_3 & 0 & -a_1 \\ -a_2 & +a_1 & 0 \end{array} \right] \begin{matrix} X_1 \\ X_2 \\ X_3 \end{matrix}$	$\mathfrak{d e f} = \frac{i}{2} \left[\begin{array}{cc} a_3 & a_1 - i a_2 \\ a_1 + i a_2 & -a_3 \end{array} \right] \mathfrak{R e g} = \left[\begin{array}{ccc} 0 & -a_3 & +a_2 \\ +a_3 & 0 & -a_1 \\ -a_2 & +a_1 & 0 \end{array} \right] \begin{matrix} X_1 \\ X_2 \\ X_3 \end{matrix}$	$\mathfrak{d e f} = \frac{i}{2} \left[\begin{array}{cc} a_3 & a_1 - i a_2 \\ a_1 + i a_2 & -a_3 \end{array} \right] \mathfrak{R e g} = \left[\begin{array}{ccc} 0 & -a_3 & +a_2 \\ +a_3 & 0 & -a_1 \\ -a_2 & +a_1 & 0 \end{array} \right] \begin{matrix} X_1 \\ X_2 \\ X_3 \end{matrix}$
gilmore-lie-groups_ EQ8684 (11.25)	193	■ 0.354	$\begin{aligned} &\mathfrak{s p}(n) = \mathfrak{u}(n) + \mathfrak{[sp(n) - u(n)]} \\ &\quad \downarrow \quad \downarrow \quad \downarrow \\ &\mathfrak{u s p}(2n) = \underbrace{\mathfrak{ou}(2n)}_{\text{real}} + \underbrace{\mathfrak{[usp(2n) - ou(2n)]}}_{\text{imaginary}} \\ &\quad \downarrow \quad \downarrow \quad \downarrow \mathfrak{p} \rightarrow i \mathfrak{p} \\ &\mathfrak{s p}(2n; R) = \underbrace{\mathfrak{ou}(2n)}_{\text{real}} + i \underbrace{\mathfrak{[usp(2n) - ou(2n)]}}_{\text{real}} \end{aligned}$	$\mathfrak{s p}(n) = \mathfrak{u}(n) + \underbrace{\mathfrak{[sp(n) - u(n)]}}_{\downarrow}$ $\mathfrak{u s p}(2n) = \underbrace{\mathfrak{ou}(2n)}_{\text{real}} + \underbrace{\mathfrak{[usp(2n) - ou(2n)]}}_{\text{imaginary}}$ $\mathfrak{s p}(2n; R) = \underbrace{\mathfrak{ou}(2n)}_{\text{real}} + i \underbrace{\mathfrak{[usp(2n) - ou(2n)]}}_{\text{real}}$ $\downarrow \mathfrak{p} \rightarrow i \mathfrak{p}$	$\mathfrak{s p}(n) = \mathfrak{u}(n) + \mathfrak{[sp(n) - u(n)]}$ $\downarrow \quad \downarrow \quad \downarrow$ $\mathfrak{u s p}(2n) = \underbrace{\mathfrak{ou}(2n)}_{\text{real}} + \underbrace{\mathfrak{[usp(2n) - ou(2n)]}}_{\text{imaginary}}$ $\downarrow \quad \downarrow \quad \downarrow \mathfrak{p} \rightarrow i \mathfrak{p}$ $\mathfrak{s p}(2n; R) = \underbrace{\mathfrak{ou}(2n)}_{\text{real}} + i \underbrace{\mathfrak{[usp(2n) - ou(2n)]}}_{\text{real}}$
gilmore-lie-groups_ EQ8683 (11.24)	193	■ 0.357	$\begin{aligned} &\mathfrak{u}(n) \longrightarrow \mathfrak{ou}(2n) \\ &\mathfrak{s p}(n) \longrightarrow \mathfrak{usp}(2n) \end{aligned}$	$\mathfrak{s p}(n) \longrightarrow \mathfrak{ou}(2n)$ $\mathfrak{u s p}(2n)$	$\mathfrak{u}(n) \longrightarrow \mathfrak{ou}(2n)$ $\mathfrak{s p}(n) \longrightarrow \mathfrak{usp}(2n)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0381 (7.19)	119	■ 0.448	$\begin{array}{c} \frac{1}{2} \left[\begin{array}{cc} 0 & ib_1 + b_2 \\ ib_1 - b_2 & 0 \end{array} \right] + \frac{1}{2} \left[\begin{array}{cc} b_3 & 0 \\ 0 & -b_3 \end{array} \right] \\ \text{EXP} \downarrow \quad \downarrow \quad \downarrow \text{EXP} \\ \left[\begin{array}{cc} e^{+ib_3/2} & 0 \\ 0 & e^{-ib_3/2} \end{array} \right] \end{array}$	$\frac{i}{2} \left[\begin{array}{cc} 0 & ib_1 + b_2 \\ ib_1 - b_2 & 0 \end{array} \right] + \frac{i}{2} \left[\begin{array}{cc} b_3 & 0 \\ 0 & -b_3 \end{array} \right]$ $\text{EXP} \downarrow \quad \downarrow \quad \downarrow \text{EXP}$ $[SU(1, 1)/U(1)] \quad \times \left[\begin{array}{cc} e^{+ib_3/2} & 0 \\ 0 & e^{-ib_3/2} \end{array} \right]$	
gilmore-lie-groups_ EQ0765 (14.11)	239	■ 0.452	$\alpha = \frac{e^2}{\hbar c} = \frac{1}{137.035999796(70)} = 0.0072973525313(38)$	$\alpha = \frac{e^2}{\hbar c} = \frac{1}{137.035999796(70)} = 0.0072973525313(38)$	$\alpha = \frac{e^2}{\hbar c} = \frac{1}{137.035999796(70)} = 0.0072973525313(38)$
gilmore-lie-groups_ EQ0590 (11.11)	190	■ 0.454	$\begin{array}{c} \frac{1}{2} \left[\begin{array}{cc} a_3 & a_1 - ia_2 \\ a_1 + ia_2 & -a_3 \end{array} \right] = \frac{1}{2} \left[\begin{array}{cc} a_3 & 0 \\ 0 & -a_3 \end{array} \right] + \frac{1}{2} \left[\begin{array}{cc} a_1 - ia_2 & a_1 + ia_2 \end{array} \right] \\ \text{u(1)} \end{array}$	$\frac{1}{2} \left[\begin{array}{cc} a_3 & a_1 - ia_2 \\ a_1 + ia_2 & -a_3 \end{array} \right] = \frac{1}{2} \left[\begin{array}{cc} a_3 & 0 \\ 0 & -a_3 \end{array} \right] + \frac{1}{2} \left[\begin{array}{cc} a_1 - ia_2 & a_1 + ia_2 \end{array} \right]$	$\frac{1}{2} \left[\begin{array}{cc} a_3 & a_1 - ia_2 \\ a_1 + ia_2 & -a_3 \end{array} \right] = \frac{1}{2} \left[\begin{array}{cc} a_3 & 0 \\ 0 & -a_3 \end{array} \right] + \frac{1}{2} \left[\begin{array}{cc} 0 & a_1 - ia_2 \\ a_1 + ia_2 & 0 \end{array} \right]$
gilmore-lie-groups_ EQ0890 (15.23)	279	■ 0.463	$\left\{ \Lambda, 0 \right\} \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle = \left \begin{array}{cc} j & j' \\ v & v' \end{array} \right\rangle D_{vv';\mu\mu'}^{jj'}(\Lambda)$	$\left\{ \Lambda, 0 \right\} \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle = \left \begin{array}{cc} j & j' \\ v & v' \end{array} \right\rangle D_{vv';\mu\mu'}^{jj'}(\Lambda)$	$\left\{ \Lambda, 0 \right\} \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle = \left \begin{array}{cc} j & j' \\ v & v' \end{array} \right\rangle D_{vv';\mu\mu'}^{jj'}(\Lambda)$
gilmore-lie-groups_ EQ0568 (10.9)	177	■ 0.466	$\begin{array}{c} E_8 \pm \mathbf{e}_i \pm \mathbf{e}_j \\ \frac{1}{2} (\pm \mathbf{e}_1 \pm \mathbf{e}_2 \pm \mathbf{e}_3 \pm \mathbf{e}_4 \pm \mathbf{e}_5 \pm \mathbf{e}_6 \pm \mathbf{e}_7 \pm \mathbf{e}_8) \\ \text{even number of + signs} \\ 1 \leq i \neq j \leq 8 \end{array}$	$E_8 \pm \mathbf{e}_i \pm \mathbf{e}_j$ $\frac{1}{2} (\pm \mathbf{e}_1 \pm \mathbf{e}_2 \pm \mathbf{e}_3 \pm \mathbf{e}_4 \pm \mathbf{e}_5 \pm \mathbf{e}_6 \pm \mathbf{e}_7 \pm \mathbf{e}_8)$ $\text{even number of + signs}$ $1 \leq i \neq j \leq 8$	$E_8 \pm \mathbf{e}_i \pm \mathbf{e}_j$ $\frac{1}{2} (\pm \mathbf{e}_1 \pm \mathbf{e}_2 \pm \mathbf{e}_3 \pm \mathbf{e}_4 \pm \mathbf{e}_5 \pm \mathbf{e}_6 \pm \mathbf{e}_7 \pm \mathbf{e}_8)$ $\text{even number of + signs}$ $1 \leq i \neq j \leq 8$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0916 (15.51)	285	■ 0.497	$\left\langle H_{k^0}, 0 \right \left k^0 \right\rangle \left\langle \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right = \left k^0 \right\rangle \left\langle \begin{array}{cc} j & j' \\ v & v' \end{array} \right D_{\nu v'; \mu \mu'}^{jj'}(H_{k^0})$		$\{H_{k^0}, 0\} k^0\rangle \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle = k^0\rangle \left \begin{array}{cc} j & j' \\ v & v' \end{array} \right\rangle D_{\nu v'; \mu \mu'}^{jj'}(H_{k^0})$
gilmore-lie-groups_ EQ0884 (15.17)	278	■ 0.500	$\Lambda \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle = \left \begin{array}{cc} j & j' \\ v & v' \end{array} \right\rangle D_{\nu v'; \mu \mu'}^{jj'}(\Lambda)$		$\Lambda \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle = \left \begin{array}{cc} j & j' \\ v & v' \end{array} \right\rangle D_{\nu v'; \mu \mu'}^{jj'}(\Lambda)$
gilmore-lie-groups_ EQ0308	99	■ 0.509	$\begin{aligned} & \mathfrak{sl}(n; \mathbb{C}) = \text{antihermitian} + \text{hermitian} \\ & \mathfrak{g} = \mathfrak{h} + \mathfrak{p} \end{aligned}$	$\mathfrak{sl}(n; \mathbb{C}) = \text{antihermitian} + \text{hermitian}$ $\mathfrak{g} = \mathfrak{h} + \mathfrak{p}$	$\mathfrak{sl}(n; \mathbb{C}) = \text{antihermitian} + \text{hermitian}$ $\mathfrak{g} = \mathfrak{h} + \mathfrak{p}$
gilmore-lie-groups_ EQ0605 (11.26)	194	■ 0.511	$\begin{aligned} & v_i^{\prime} G_{ij} v_j = (p_1, q_1, p_2, q_2, \dots, p_n, q_n)' \\ & = \sum_{i=1}^n (p_i' q_i - q_i' p_i) \end{aligned}$	$v_i' G_{ij} v_j = (p_1, q_1, p_2, q_2, \dots, p_n, q_n)'$ $= \sum_{i=1}^n (p_i' q_i - q_i' p_i)$	$v_i' G_{ij} v_j = (p_1, q_1, p_2, q_2, \dots, p_n, q_n)'$ $= \sum_{i=1}^n (p_i' q_i - q_i' p_i) \quad (11.26)$
gilmore-lie-groups_ EQ0829 (14.64)	259	■ 0.513	$\begin{aligned} & [rD, r] = +r \quad [a^\dagger a, a^\dagger] = +a^\dagger \\ & [rD, rD^2] = -rD^2 \quad [a^\dagger a, a^\dagger aa] = -a^\dagger aa \\ & [r, rD^2] = -2rD \quad [a^\dagger, a^\dagger aa] = -2a^\dagger a \end{aligned}$	$\begin{aligned} [rD, r] &= +r \quad [a^\dagger a, a^\dagger] = +a^\dagger \\ [rD, rD^2] &= -rD^2 \quad [a^\dagger a, a^\dagger aa] \\ [r, rD^2] &= -2rD \quad [a^\dagger, a^\dagger aa] \\ &= -2a^\dagger a \end{aligned}$	$\begin{aligned} [rD, r] &= +r \quad [a^\dagger a, a^\dagger] = +a^\dagger \\ [rD, rD^2] &= -rD^2 \quad [a^\dagger a, a^\dagger aa] = -a^\dagger aa \\ [r, rD^2] &= -2rD \quad [a^\dagger, a^\dagger aa] = -2a^\dagger a \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08283 (5.22)	93	0.520	$M \mathbb{G} + G \mathbb{M}^{\dagger} = 0 \quad \begin{array}{l} \begin{array}{l} \mathbb{R} \\ \mathbb{C} \\ \mathbb{Q} \end{array} \end{array} \quad \begin{array}{l} \mathfrak{o}(n; G) \\ \mathfrak{u}(n; G) \\ \mathfrak{sp}(n; G) \end{array}$	$MG + GM^{\dagger} = 0$	$MG + GM^{\dagger} = 0 \quad \begin{array}{l} \mathbb{R} \\ \mathbb{C} \\ \mathbb{Q} \end{array} \quad \begin{array}{l} \mathfrak{o}(n; G) \\ \mathfrak{u}(n; G) \\ \mathfrak{sp}(n; G) \end{array}$
gilmore-lie-groups_ E08558 (10.3)	174	0.523	$\sum_{\alpha \neq 0} \alpha \cdot \alpha = \text{rank} = l$	$\sum_{\alpha \neq 0} \alpha \cdot \alpha = \text{rank} = l$	$\sum_{\alpha \neq 0} \alpha \cdot \alpha = \text{rank} = l$
gilmore-lie-groups_ E08672	216	0.523	$z \rightarrow z' = \frac{az+b}{cz+d} \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL(2; \mathbb{R}), \quad ad - bc = 1$	$z \rightarrow z' = \frac{az+b}{cz+d} \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL(2; \mathbb{R}), \quad ad - bc = 1$	$z \rightarrow z' = \frac{az+b}{cz+d} \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL(2; \mathbb{R}), \quad ad - bc = 1$
gilmore-lie-groups_ E08588 (9.9)	155	0.530	$\begin{aligned} & \left(\text{Re } g(X) - \lambda I_3 \right) \cdot (-\lambda) = (-\lambda)^3 + (-\lambda)(+a_1^2 + a_2^2 + a_3^2) = 0 \\ & \left(+a_1^2 + a_2^2 + a_3^2 \right) \cdot (-\lambda) = (-\lambda)(\lambda^2 + \phi_2(\mathbf{a})) \\ & \phi_2(\mathbf{a}) = +a_1^2 + a_2^2 + a_3^2 \end{aligned}$	$\begin{aligned} \ \Re g(X) - \lambda I_3\ &= (-\lambda)^3 + (-\lambda)(+a_1^2 + a_2^2 + a_3^2) = 0 \\ &= (-\lambda)(\lambda^2 + \phi_2(\mathbf{a})) \\ \phi_2(\mathbf{a}) &= +a_1^2 + a_2^2 + a_3^2 \end{aligned}$	$\begin{aligned} \ \Re g(X) - \lambda I_3\ &= (-\lambda)^3 + (-\lambda)(+a_1^2 + a_2^2 + a_3^2) = 0 \\ &= (-\lambda)(\lambda^2 + \phi_2(\mathbf{a})) \\ \phi_2(\mathbf{a}) &= +a_1^2 + a_2^2 + a_3^2 \end{aligned}$
gilmore-lie-groups_ E08152	59	0.532	$\begin{array}{l} \text{F(n)=F(n-1)+F(n-2)} \\ \text{F(n)} \end{array}$	$\begin{array}{l} F(n) = F(n-1) + F(n-2) \\ \begin{array}{cccccccccc} n & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & \dots \\ F(n) & 0 & 1 & 1 & 2 & 3 & 5 & 8 & 13 & \dots \end{array} \end{array}$	$F(n) = F(n-1) + F(n-2)$
gilmore-lie-groups_ E08887 (14.47)	253	0.534	$\begin{array}{l} a_1^{\dagger} a_2 \quad (+1, -1, 0, 0) \\ a_2^{\dagger} a_1 \quad (-1, +1, 0, 0) \\ b_1^{\dagger} b_2 \quad (0, 0 + 1, -1) \\ b_2^{\dagger} b_1 \quad (0, 0, -1, +1) \end{array}$	$\begin{array}{l} a_1^{\dagger} a_2 \quad (+1, -1, 0, 0) \\ a_2^{\dagger} a_1 \quad (-1, +1, 0, 0) \\ b_1^{\dagger} b_2 \quad (0, 0 + 1, -1) \\ b_2^{\dagger} b_1 \quad (0, 0, -1, +1) \end{array}$	$\begin{array}{l} a_1^{\dagger} a_2 \quad (+1, -1, 0, 0) \\ a_2^{\dagger} a_1 \quad (-1, +1, 0, 0) \\ b_1^{\dagger} b_2 \quad (0, 0 + 1, -1) \\ b_2^{\dagger} b_1 \quad (0, 0, -1, +1) \end{array}$
gilmore-lie-groups_ E08658	214	0.536	$ds^2 = \frac{(dx_1 dx_2) \begin{bmatrix} 1 + x_2^2 - x_1 x_2 \\ -x_1 x_2 \\ 1 + x_1^2 \end{bmatrix} (dx_1)}{1 + x_1^2 + x_2^2}$	$ds^2 = \frac{(dx_1 dx_2) \begin{bmatrix} 1 + x_2^2 - x_1 x_2 \\ -x_1 x_2 \\ 1 + x_1^2 \end{bmatrix} (dx_1)}{1 + x_1^2 + x_2^2}$	$ds^2 = \frac{(dx_1 dx_2) \begin{bmatrix} 1 + x_2^2 - x_1 x_2 \\ -x_1 x_2 \\ 1 + x_1^2 \end{bmatrix} (dx_1)}{1 + x_1^2 + x_2^2}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00380 (7.18)	119	■0.541	$\begin{array}{c} \begin{array}{c} \left[\begin{array}{ccc} 0 & 0 & a_2 \\ 0 & 0 & a_1 \\ a_2 & a_1 & 0 \end{array}\right] \\ \text{EXP} \downarrow \end{array} \\ \downarrow \text{SO}(2,1)/\text{SO}(2) \times \begin{array}{c} \begin{array}{ccc} \cos a_3 & \sin a_3 & 0 \\ -\sin a_3 & \cos a_3 & 0 \\ 0 & 0 & 1 \end{array} \end{array} \end{array}$	$\begin{array}{c} \begin{array}{ccc} 0 & 0 & a_2 \\ 0 & 0 & a_1 \\ a_2 & a_1 & 0 \end{array} \\ \text{EXP} \downarrow \end{array} + c \begin{array}{c} \begin{array}{ccc} 0 & a_3 & 0 \\ -a_3 & 0 & 0 \\ 0 & 0 & 0 \end{array} \\ \downarrow \text{EXP} \end{array}$	$\begin{array}{c} \begin{array}{ccc} 0 & 0 & a_2 \\ 0 & 0 & a_1 \\ a_2 & a_1 & 0 \end{array} \\ \text{EXP} \downarrow \end{array} + \begin{array}{c} \begin{array}{ccc} 0 & a_3 & 0 \\ -a_3 & 0 & 0 \\ 0 & 0 & 0 \end{array} \\ \downarrow \text{EXP} \end{array}$ $[SO(2,1)/SO(2)] \times \begin{array}{ccc} \cos a_3 & \sin a_3 & 0 \\ -\sin a_3 & \cos a_3 & 0 \\ 0 & 0 & 1 \end{array}$
gilmore-lie-groups_ E00172	64	■0.546	$\{f,g\}=(Df)^tG(Dg) \quad \text{where} \quad G=\begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix} \quad \text{and} \quad (Dg)=\begin{bmatrix} \partial g/\partial q \\ \partial g/\partial p \end{bmatrix}$	$\{f,g\}=(Df)^tG(Dg) \quad \text{where} \quad G=\begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix} \quad \text{and} \quad (Dg)=\begin{bmatrix} \partial g/\partial q \\ \partial g/\partial p \end{bmatrix}$	$\{f,g\}=(Df)^tG(Dg) \quad \text{where} \quad G=\begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix} \quad \text{and} \quad (Dg)=\begin{bmatrix} \partial g/\partial q \\ \partial g/\partial p \end{bmatrix}$
gilmore-lie-groups_ E00100	45	■0.551	$X=H(x,y,z,ct)=\begin{bmatrix} ct+z & x-iy \\ x+iy & ct-z \end{bmatrix}=ctI_2+\sigma\cdot\mathbf{x}$	$X=H(x,y,z,ct)=\begin{bmatrix} ct+z & x-iy \\ x+iy & ct-z \end{bmatrix}=ctI_2+\sigma\cdot\mathbf{x}$	$X=H(x,y,z,ct)=\begin{bmatrix} ct+z & x-iy \\ x+iy & ct-z \end{bmatrix}=ctI_2+\sigma\cdot\mathbf{x}$
gilmore-lie-groups_ E00663	214	■0.554	$dV=\sqrt{\ g\ }r^{n-1}\sin^{n-2}\phi_2\sin^{n-3}\phi_3\cdots\sin^1\phi_{n-1}\sin^0\phi_n$ $dr\wedge d\phi_2\wedge d\phi_3\wedge\cdots\wedge d\phi_n$	$dV=\sqrt{\ g\ }r^{n-1}\sin^{n-2}\phi_2\sin^{n-3}\phi_3\cdots\sin^1\phi_{n-1}\sin^0\phi_n$ $dr\wedge d\phi_2\wedge d\phi_3\wedge\cdots\wedge d\phi_n$	$dV=\sqrt{\ g\ }r^{n-1}\sin^{n-2}\phi_2\sin^{n-3}\phi_3\cdots\sin^1\phi_{n-1}\sin^0\phi_n$ $dr\wedge d\phi_2\wedge d\phi_3\wedge\cdots\wedge d\phi_n$
gilmore-lie-groups_ E00227 (4.37)	79	■0.562	$(X_i,X_j)=\text{Tr}R(X_i)R(X_j)=\sum_r\sum_sR(X_i)_r^sR(X_j)_s^r=\sum_r\sum_sC_{ir}^sC_{js}^r$	$(X_i,X_j)=\text{Tr}R(X_i)R(X_j)=\sum_r\sum_sR(X_i)_r^sR(X_j)_s^r=\sum_r\sum_sC_{ir}^sC_{js}^r$	$(X_i,X_j)=\text{Tr}R(X_i)R(X_j)=\sum_r\sum_sR(X_i)_r^sR(X_j)_s^r=\sum_r\sum_sC_{ir}^sC_{js}^r$
gilmore-lie-groups_ E00724 (13.41)	228	■0.562	$\lim_{\substack{c\rightarrow 0 \\ J\rightarrow\infty}}h_3\left \begin{array}{c}J\\M\end{array}\right\rangle=n\left \begin{array}{c}\infty\\n\end{array}\right\rangle$	$\lim_{\substack{c\rightarrow 0 \\ J\rightarrow\infty}}h_3\left \begin{array}{c}J\\M\end{array}\right\rangle=n\left \begin{array}{c}\infty\\n\end{array}\right\rangle$	$\lim_{\substack{c\rightarrow 0 \\ J\rightarrow\infty}}h_3\left \begin{array}{c}J\\M\end{array}\right\rangle=n\left \begin{array}{c}\infty\\n\end{array}\right\rangle$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0638 (12.10)	208	0.570	$\begin{array}{cc} S_{0(2,1)} / S_{0(2)} & \& S_{0(3)} / S_{0(2)} \\ \left[\begin{array}{cc} W & X \\ +X^t & Y \end{array} \right] & \& \left[\begin{array}{cc} W & X \\ -X^t & Y \end{array} \right] \\ \& \& W^2 = I_2 + \binom{x}{y} \begin{pmatrix} x & y \end{pmatrix} & \& W^2 = I_2 - \binom{x}{y} \begin{pmatrix} x & y \end{pmatrix} \\ \& \& Y^2 = I_1 + \begin{pmatrix} x & y \end{pmatrix} \binom{x}{y} & \& Y^2 = I_1 - \begin{pmatrix} x & y \end{pmatrix} \binom{x}{y} \end{array}$	$SO(2,1)/SO(2) \qquad SO(3)/SO(2)$ $\begin{bmatrix} W & X \\ +X^t & Y \end{bmatrix} \qquad \begin{bmatrix} W & X \\ -X^t & Y \end{bmatrix}$ $W^2 = I_2 + \binom{x}{y} \begin{pmatrix} x & y \end{pmatrix} \qquad W^2 = I_2 - \binom{x}{y} \begin{pmatrix} x & y \end{pmatrix}$ $Y^2 = I_1 + \begin{pmatrix} x & y \end{pmatrix} \binom{x}{y} \qquad Y^2 = I_1 - \begin{pmatrix} x & y \end{pmatrix} \binom{x}{y}$	
gilmore-lie-groups_ EQ0705	224	0.574	$\begin{gathered} \mathcal{C}^2[S_{0(4,1)}] = \mathbf{L} \cdot \mathbf{L} - \mathbf{B} \cdot \mathbf{B} + \mathbf{T} \cdot \mathbf{T} - T_{45}^2 \\ \mathcal{C}^2[ISO(3,1)] = 0 - 0 + \nabla \cdot \nabla - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \end{gathered}$	$\mathcal{C}^2[SO(4,1)] = \mathbf{L} \cdot \mathbf{L} - \mathbf{B} \cdot \mathbf{B} + \mathbf{T} \cdot \mathbf{T} - T_{45}^2$ $\mathcal{C}^2[ISO(3,1)] = 0 - 0 + \nabla \cdot \nabla - \frac{1}{c^2} \frac{\partial^2}{\partial t^2}$	$\mathcal{C}^2[SO(4,1)] = \mathbf{L} \cdot \mathbf{L} - \mathbf{B} \cdot \mathbf{B} + \mathbf{T} \cdot \mathbf{T} - T_{45}^2$ $\mathcal{C}^2[ISO(3,1)] = 0 - 0 + \nabla \cdot \nabla - \frac{1}{c^2} \frac{\partial^2}{\partial t^2}$
gilmore-lie-groups_ EQ0390 (7.26)	123	0.578	$(x, y) \rightarrow \left[\begin{array}{cc} x & y \\ 0 & 1 \end{array} \right] = \left[\begin{array}{cc} e^w & (e^w - 1)z/w \\ 0 & 1 \end{array} \right] \leftarrow (w, z)$	$(x, y) \rightarrow \left[\begin{array}{cc} x & y \\ 0 & 1 \end{array} \right] = \left[\begin{array}{cc} e^w & (e^w - 1)z/w \\ 0 & 1 \end{array} \right] \leftarrow (w, z)$	$(x, y) \rightarrow \left[\begin{array}{cc} x & y \\ 0 & 1 \end{array} \right] = \left[\begin{array}{cc} e^w & (e^w - 1)z/w \\ 0 & 1 \end{array} \right] \leftarrow (w, z)$
gilmore-lie-groups_ EQ0798 (14.41)	251	0.582	$\left[\begin{array}{c} \lambda \mathbf{u} \\ \lambda \end{array} \right]' = \left[\begin{array}{cc} A & B \\ C & D \end{array} \right] \left[\begin{array}{c} \lambda \mathbf{u} \\ \lambda \end{array} \right]$	$\left[\begin{array}{c} \lambda \mathbf{u} \\ \lambda \end{array} \right]' = \left[\begin{array}{cc} A & B \\ C & D \end{array} \right] \left[\begin{array}{c} \lambda \mathbf{u} \\ \lambda \end{array} \right]$	$\left[\begin{array}{c} \lambda \mathbf{u} \\ \lambda \end{array} \right]' = \left[\begin{array}{cc} A & B \\ C & D \end{array} \right] \left[\begin{array}{c} \lambda \mathbf{u} \\ \lambda \end{array} \right]$
gilmore-lie-groups_ EQ1022	319	0.583	$\begin{array}{l} \text{displacements} \\ \text{rotations} \\ \text{boosts} \\ \text{dilations} \end{array} \quad \begin{array}{l} P_i \partial_x, \partial_y, \partial_t \\ L_z \quad x \partial_y - y \partial_x \\ B_i \quad x \partial_t + t \partial_x, y \partial_t + t \partial_y \\ D_i \quad x \partial_x + y \partial_y + t \partial_t, u \partial_u \end{array}$	$\begin{array}{l} \text{displacements} \\ \text{rotations} \\ \text{boosts} \\ \text{dilations} \end{array} \quad \begin{array}{l} P_i \partial_x, \partial_y, \partial_t \\ L_z \quad x \partial_y - y \partial_x \\ B_i \quad x \partial_t + t \partial_x, y \partial_t + t \partial_y \\ D_i \quad x \partial_x + y \partial_y + t \partial_t, u \partial_u \end{array}$	$\begin{array}{l} \text{displacements} \\ \text{rotations} \\ \text{boosts} \\ \text{dilations} \end{array} \quad \begin{array}{l} P_i \partial_x, \partial_y, \partial_t \\ L_z \quad x \partial_y - y \partial_x \\ B_i \quad x \partial_t + t \partial_x, y \partial_t + t \partial_y \\ D_i \quad x \partial_x + y \partial_y + t \partial_t, u \partial_u \end{array}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0091	42	■ 0.584	$\begin{aligned} & \backslash begin{aligned} g \backslash left(x_{\{1\}}, x_{\{2\}}, x_{\{3\}} \backslash right) \\ & \& \backslash circ g \backslash left(y_{\{1\}}, y_{\{2\}}, y_{\{3\}} \backslash right) \backslash \backslash \{ \\ & \backslash left[\backslash begin{array}{cc} x_{\{1\}} \& x_{\{2\}} \backslash \backslash x_{\{3\}} \& \\ & \backslash frac{1+x_{\{2\}} x_{\{3\}}}{x_{\{1\}}} \backslash end{array} \backslash right] \backslash times \\ & \backslash left[\backslash begin{array}{cc} y_{\{1\}} \& y_{\{2\}} \backslash \backslash y_{\{3\}} \& \\ & \backslash frac{1+y_{\{2\}} y_{\{3\}}}{y_{\{1\}}} \backslash end{array} \backslash right] \} \\ & \& = \backslash left[\backslash begin{array}{cc} z_{\{1\}} \& z_{\{2\}} \backslash \backslash z_{\{3\}} \& \\ & \backslash frac{1+z_{\{2\}} z_{\{3\}}}{z_{\{1\}}} \backslash end{array} \backslash right] \\ & \backslash end{aligned} \end{aligned}$	$g(x_1, x_2, x_3) \circ g(y_1, y_2, y_3)$ $\begin{bmatrix} x_1 & x_2 \\ x_3 & \frac{1+x_2x_3}{x_1} \end{bmatrix} \times \begin{bmatrix} y_1 & y_2 \\ y_3 & \frac{1+y_2y_3}{y_1} \end{bmatrix} = \begin{bmatrix} z_1 & z_2 \\ z_3 & \frac{1+z_2z_3}{z_1} \end{bmatrix}$	$g(x_1, x_2, x_3) \circ g(y_1, y_2, y_3) = g(z_1, z_2, z_3)$ $\begin{bmatrix} x_1 & x_2 \\ x_3 & \frac{1+x_2x_3}{x_1} \end{bmatrix} \times \begin{bmatrix} y_1 & y_2 \\ y_3 & \frac{1+y_2y_3}{y_1} \end{bmatrix} = \begin{bmatrix} z_1 & z_2 \\ z_3 & \frac{1+z_2z_3}{z_1} \end{bmatrix}$
gilmore-lie-groups_ EQ0946	294	■ 0.584	$\begin{aligned} & g_{\{\text{flat}\}} = \backslash left[\backslash begin{array}{c ccc} c^2 \\ & \& \& \backslash hline \& -1 \& \& \backslash \& -1 \& \backslash \& \& -1 \\ & \backslash end{array} \backslash right] \quad g_{\{\text{grav}\}} = \backslash left[\\ & \backslash begin{array}{c ccc} c^2 \backslash left(1 + \backslash frac{2 \Phi(x)}{c^2} \backslash right) \\ & \backslash right) \& \& \backslash \backslash \backslash hline \& -1 \& \& \backslash \& -1 \& \backslash \& \& -1 \\ & \backslash end{array} \backslash right] \end{aligned}$	$g_{\text{flat}} = \left[\begin{array}{c ccc} c^2 & & & \\ \hline & -1 & & \\ & & -1 & \\ & & & -1 \end{array} \right] \quad g_{\text{grav}} = \left[\begin{array}{c ccc} c^2 \left(1 + \frac{2\Phi(x)}{c^2} \right) & & & \\ \hline & -1 & & \\ & & -1 & \\ & & & -1 \end{array} \right]$	$g_{\text{flat}} = \left[\begin{array}{c ccc} c^2 & & & \\ \hline & -1 & & \\ & & -1 & \\ & & & -1 \end{array} \right] \quad g_{\text{grav}} = \left[\begin{array}{c ccc} c^2 \left(1 + \frac{2\Phi(x)}{c^2} \right) & & & \\ \hline & -1 & & \\ & & -1 & \\ & & & -1 \end{array} \right]$
gilmore-lie-groups_ EQ0639 (12.11)	209	■ 0.594	$\begin{aligned} & \backslash begin{gathered} S_{0(2,1)} / S_{0(2)} \quad \backslash quad \quad S_{0(3)} / S_{0(2)} \backslash \backslash g_{\{*,*\}} = W^{-2} = \backslash left[I_{\{2\}} + \backslash binom{x}{y} \backslash left(\\ & \backslash begin{array}{ll} x \& y \backslash end{array} \backslash right) \backslash right]^{-1} \\ & \quad \backslash quad \quad g_{\{*,*\}} = W^{-2} = \backslash left[I_{\{2\}} - \backslash binom{x}{y} \backslash left(\\ & \backslash begin{array}{ll} x \& y \backslash end{array} \backslash right) \backslash right]^{-1} \backslash \backslash \\ & g^{\{*,*\}} = W^{+2} = \backslash left[\backslash begin{array}{cc} 1+x^2 \& +x y \backslash \backslash \\ & +y x \& 1+y^2 \backslash end{array} \backslash right] \quad \backslash quad \quad g^{\{*,*\}} = W^{+2} = \\ & \backslash left[\backslash begin{array}{cc} 1-x^2 \& -x y \backslash \backslash \\ & -y x \& 1-y^2 \backslash end{array} \backslash right] \backslash end{gathered} \end{aligned}$	$\begin{aligned} & SO(2,1)/SO(2) \quad SO(3)/SO(2) \\ & g_{*,*} = W^{-2} = \left[I_2 + \begin{pmatrix} x \\ y \end{pmatrix} \begin{pmatrix} x & y \end{pmatrix} \right]^{-1} \quad g_{*,*} = W^{-2} = \left[I_2 - \begin{pmatrix} x \\ y \end{pmatrix} \begin{pmatrix} x & y \end{pmatrix} \right]^{-1} \\ & g^{*,*} = W^{+2} = \left[\begin{array}{cc} 1+x^2 & +xy \\ +yx & 1+y^2 \end{array} \right] \quad g^{*,*} = W^{+2} = \left[\begin{array}{cc} 1-x^2 & -xy \\ -yx & 1-y^2 \end{array} \right] \end{aligned}$	$\begin{aligned} & SO(2,1)/SO(2) \quad SO(3)/SO(2) \\ & g_{*,*} = W^{-2} = \left[I_2 + \begin{pmatrix} x \\ y \end{pmatrix} \begin{pmatrix} x & y \end{pmatrix} \right]^{-1} \quad g_{*,*} = W^{-2} = \left[I_2 - \begin{pmatrix} x \\ y \end{pmatrix} \begin{pmatrix} x & y \end{pmatrix} \right]^{-1} \\ & g^{*,*} = W^{+2} = \left[\begin{array}{cc} 1+x^2 & +xy \\ +yx & 1+y^2 \end{array} \right] \quad g^{*,*} = W^{+2} = \left[\begin{array}{cc} 1-x^2 & -xy \\ -yx & 1-y^2 \end{array} \right] \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00019 (1.15)	22	■ 0.596	$\begin{aligned} & \left(z-z_1\right) \left(z-z_2\right) \cdots \left(z-z_n\right)=z^n-I_1 z^{n-1}+I_2 z^{n-2}-\cdots+(-)^n I_n=0 \\ & I_1=\sum_{i=1}^n z_i=z_1+z_2+\cdots+z_n \\ & I_2=\sum_{i<j}^n z_i z_j=z_1 z_2+z_1 z_3+\cdots+z_1 z_n+z_2 z_3+\cdots+z_{n-1} z_n \\ & \vdots \\ & I_n=\sum_{i<j<\cdots<k}^n z_i z_j \cdots z_k=z_1 z_2 \cdots z_n \end{aligned}$	$\begin{aligned} & (z-z_1)(z-z_2)\cdots(z-z_n)=z^n-I_1z^{n-1}+I_2z^{n-2}-\cdots+(-)^nI_n=0 \\ & I_1=\sum_{i=1}^nz_i=z_1+z_2+\cdots+z_n \\ & I_2=\sum_{i<j}^nz_iz_j=z_1z_2+z_1z_3+\cdots+z_1z_n+z_2z_3+\cdots+z_{n-1}z_n \\ & \vdots \\ & I_n=\sum_{i<j<\cdots<k}^nz_iz_j\cdots z_k=z_1z_2\cdots z_n \end{aligned}$	$\begin{aligned} & (z-z_1)(z-z_2)\cdots(z-z_n)=z^n-I_1z^{n-1}+I_2z^{n-2}-\cdots+(-)^nI_n=0 \\ & I_1=\sum_{i=1}^nz_i=z_1+z_2+\cdots+z_n \\ & I_2=\sum_{i<j}^nz_iz_j=z_1z_2+z_1z_3+\cdots+z_1z_n+z_2z_3+\cdots+z_{n-1}z_n \\ & \vdots \\ & I_n=\sum_{i<j<\cdots<k}^nz_iz_j\cdots z_k=z_1z_2\cdots z_n \end{aligned} \quad (1)$
gilmore-lie-groups_ E00567	177	■ 0.603	$\begin{aligned} & E_7 \pm \mathbf{e}_i \pm \mathbf{e}_j \\ & \frac{1}{2} \underbrace{(\pm \mathbf{e}_1 \pm \mathbf{e}_2 \pm \mathbf{e}_3 \pm \mathbf{e}_4 \pm \mathbf{e}_5 \pm \mathbf{e}_6)}_{\text{even number of + signs}} \pm \frac{\sqrt{2}}{4} \mathbf{e}_7 \quad 1 \leq i \neq j \leq 6 \end{aligned}$	$E_7 \pm \mathbf{e}_i \pm \mathbf{e}_j$	$E_7 \pm \mathbf{e}_i \pm \mathbf{e}_j$
gilmore-lie-groups_ E00310	99	■ 0.610	$\begin{aligned} & [\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h} \quad (\mathfrak{h}, \mathfrak{h}) \leq 0 \\ & [\mathfrak{h}, \mathfrak{p}] \subseteq \mathfrak{p} \quad \text{and} \quad (\mathfrak{h}, \mathfrak{p}) = 0 \\ & [\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h} \quad \text{and} \quad (\mathfrak{p}, \mathfrak{p}) \geq 0 \end{aligned}$	$\begin{aligned} & [\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h} \quad (\mathfrak{h}, \mathfrak{h}) \leq 0 \\ & [\mathfrak{h}, \mathfrak{p}] \subseteq \mathfrak{p} \quad \text{and} \quad (\mathfrak{h}, \mathfrak{p}) = 0 \\ & [\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h} \quad (\mathfrak{p}, \mathfrak{p}) \geq 0 \end{aligned}$	$\begin{aligned} & [\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h} \quad (\mathfrak{h}, \mathfrak{h}) \leq 0 \\ & [\mathfrak{h}, \mathfrak{p}] \subseteq \mathfrak{p} \quad \text{and} \quad (\mathfrak{h}, \mathfrak{p}) = 0 \\ & [\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h} \quad (\mathfrak{p}, \mathfrak{p}) \geq 0 \end{aligned}$
gilmore-lie-groups_ E00961 (16.4)	300	■ 0.612	$\begin{aligned} & \text{surface equation} \quad F(x, y, p) = 0 \\ & \text{constraint equation} \quad p = dy/dx \quad \text{when } F(x, y, p) = 0 \end{aligned}$	$\begin{aligned} & \text{surface equation} \quad F(x, y, p) = 0 \\ & \text{constraint equation} \quad p = dy/dx \quad \text{when } F(x, y, p) = 0 \end{aligned}$	$\begin{aligned} & \text{surface equation} \quad F(x, y, p) = 0 \\ & \text{constraint equation} \quad p = dy/dx \quad \text{when } F(x, y, p) = 0 \end{aligned}$
gilmore-lie-groups_ E00635 (12.7)	208	■ 0.613	$\begin{aligned} & P' = \begin{bmatrix} W & X \\ X^\dagger & Y \end{bmatrix} \quad X = \begin{bmatrix} x^1 \\ x^2 \\ \vdots \\ x^n \end{bmatrix} \\ & W^2 = I_n + XX^\dagger \\ & Y^2 = 1 + X^\dagger X \end{aligned}$	$\begin{aligned} & P' = \begin{bmatrix} W & X \\ X^\dagger & Y \end{bmatrix} \quad X = \begin{bmatrix} x^1 \\ x^2 \\ \vdots \\ x^n \end{bmatrix} \\ & W^2 = I_n + XX^\dagger \\ & Y^2 = 1 + X^\dagger X \end{aligned}$	$\begin{aligned} & P' = \begin{bmatrix} W & X \\ X^\dagger & Y \end{bmatrix} \quad X = \begin{bmatrix} x^1 \\ x^2 \\ \vdots \\ x^n \end{bmatrix} \\ & W^2 = I_n + XX^\dagger \\ & Y^2 = 1 + X^\dagger X \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E06606 (11.21)	192	0.619	$\begin{aligned} \mathfrak{g} &= \mathfrak{h} + \mathfrak{p} \\ \mathfrak{su}(n) &= \mathfrak{so}(n) + [\mathfrak{su}(n) - \mathfrak{so}(n)] \\ \mathfrak{sp}(n) &= \mathfrak{u}(n) + [\mathfrak{sp}(n) - \mathfrak{u}(n)] \end{aligned}$	$\mathfrak{g} = \mathfrak{h} + \mathfrak{p} \longrightarrow \mathfrak{g}'$ $\mathfrak{su}(n) = \mathfrak{so}(n) + [\mathfrak{su}(n) - \mathfrak{so}(n)] \longrightarrow \mathfrak{sl}(n; \mathbb{R})$ $\mathfrak{sp}(n) = \mathfrak{u}(n) + [\mathfrak{sp}(n) - \mathfrak{u}(n)] \longrightarrow \mathfrak{sp}(2n; \mathbb{R})$	$\begin{aligned} \mathfrak{g} &= \mathfrak{h} + \mathfrak{p} \longrightarrow \mathfrak{g}' \\ \mathfrak{su}(n) &= \mathfrak{so}(n) + [\mathfrak{su}(n) - \mathfrak{so}(n)] \longrightarrow \mathfrak{sl}(n; \mathbb{R}) \\ \mathfrak{sp}(n) &= \mathfrak{u}(n) + [\mathfrak{sp}(n) - \mathfrak{u}(n)] \longrightarrow \mathfrak{sp}(2n; \mathbb{R}) \end{aligned}$
gilmore-lie-groups_ E06411 (7.47)	126	0.625	$\begin{aligned} &\operatorname{EXP}\left(i\theta_{+}^{\prime}\right) \quad \operatorname{EXP}\left(i\theta_{z}^{\prime}\right) \quad \operatorname{EXP}\left(i\theta_{-}^{\prime}\right) \\ &\downarrow \downarrow \\ &\left[\begin{array}{cc} 1 & i\theta_{+}^{\prime} \\ 0 & 1 \end{array}\right]\left[\begin{array}{cc} e^{i\theta_{z}^{\prime} / 2} & 0 \\ 0 & e^{-i\theta_{z}^{\prime} / 2} \end{array}\right]\left[\begin{array}{cc} 1 & 0 \\ i\theta_{-}^{\prime} & 1 \end{array}\right] \\ &= \left[\begin{array}{cc} e^{i\theta_{z}^{\prime} / 2}-\theta_{+}^{\prime} \theta_{-}^{\prime} e^{-i\theta_{z}^{\prime} / 2} & i\theta_{+}^{\prime} e^{-i\theta_{z}^{\prime} / 2} \\ i\theta_{-}^{\prime} e^{-i\theta_{z}^{\prime} / 2} & e^{-i\theta_{z}^{\prime} / 2} \end{array}\right] \end{aligned}$	$\begin{aligned} &\operatorname{EXP}\left(i\theta_{+}^{\prime} J_{+}\right) \quad \operatorname{EXP}\left(i\theta_{z}^{\prime} J_{z}\right) \quad \operatorname{EXP}\left(i\theta_{-}^{\prime} J_{-}\right) \\ &\downarrow \downarrow \\ &\left[\begin{array}{cc} 1 & i\theta_{+}^{\prime} \\ 0 & 1 \end{array}\right]\left[\begin{array}{cc} e^{i\theta_{z}^{\prime} / 2} & 0 \\ 0 & e^{-i\theta_{z}^{\prime} / 2} \end{array}\right]\left[\begin{array}{cc} 1 & 0 \\ i\theta_{-}^{\prime} & 1 \end{array}\right] \\ &= \left[\begin{array}{cc} e^{i\theta_{z}^{\prime} / 2}-\theta_{+}^{\prime} \theta_{-}^{\prime} e^{-i\theta_{z}^{\prime} / 2} & i\theta_{+}^{\prime} e^{-i\theta_{z}^{\prime} / 2} \\ i\theta_{-}^{\prime} e^{-i\theta_{z}^{\prime} / 2} & e^{-i\theta_{z}^{\prime} / 2} \end{array}\right] \end{aligned}$	$\begin{aligned} &\operatorname{EXP}\left(i\theta_{+}^{\prime} J_{+}\right) \quad \operatorname{EXP}\left(i\theta_{z}^{\prime} J_{z}\right) \quad \operatorname{EXP}\left(i\theta_{-}^{\prime} J_{-}\right) \\ &\downarrow \downarrow \\ &\left[\begin{array}{cc} 1 & i\theta_{+}^{\prime} \\ 0 & 1 \end{array}\right]\left[\begin{array}{cc} e^{i\theta_{z}^{\prime} / 2} & 0 \\ 0 & e^{-i\theta_{z}^{\prime} / 2} \end{array}\right]\left[\begin{array}{cc} 1 & 0 \\ i\theta_{-}^{\prime} & 1 \end{array}\right] \\ &= \left[\begin{array}{cc} e^{i\theta_{z}^{\prime} / 2}-\theta_{+}^{\prime} \theta_{-}^{\prime} e^{-i\theta_{z}^{\prime} / 2} & i\theta_{+}^{\prime} e^{-i\theta_{z}^{\prime} / 2} \\ i\theta_{-}^{\prime} e^{-i\theta_{z}^{\prime} / 2} & e^{-i\theta_{z}^{\prime} / 2} \end{array}\right] \end{aligned}$
gilmore-lie-groups_ E06343 (6.28)	106	0.637	$\begin{aligned} \mathcal{X}_a &= -x^{\prime} \partial / \partial x^{\prime} - y^{\prime} \partial / \partial y^{\prime} \\ \mathcal{X}_b &= -\partial / \partial y^{\prime} \end{aligned}$	$\begin{aligned} \mathcal{X}_a &= -x^{\prime} \partial / \partial x^{\prime} - y^{\prime} \partial / \partial y^{\prime} \\ \mathcal{X}_b &= -\partial / \partial y^{\prime} \end{aligned}$	$\begin{aligned} \mathcal{X}_a &= -x^{\prime} \partial / \partial x^{\prime} - y^{\prime} \partial / \partial y^{\prime} \\ \mathcal{X}_b &= -\partial / \partial y^{\prime} \end{aligned}$
gilmore-lie-groups_ E06690 (13.8)	221	0.639	$\begin{aligned} &\left\{ \left[L_3, P_1 \right] \right\} = -P_2 \\ &\left\{ \left[L_3, P_2 \right] \right\} = +P_1 \\ &\left\{ \left[P_1, P_2 \right] \right\} = 0 \end{aligned}$	$\begin{aligned} [L_3, P_1] &= -P_2 \\ [L_3, P_2] &= +P_1 \\ [P_1, P_2] &= 0 \end{aligned}$	$\begin{aligned} [L_3, P_1] &= -P_2 \\ [L_3, P_2] &= +P_1 \\ [P_1, P_2] &= 0 \end{aligned}$
gilmore-lie-groups_ E06736	231	0.640	$\begin{aligned} &\left\langle \begin{array}{c} c \\ m^{\prime} \end{array} \left a J_{\pm} \right \begin{array}{c} c \\ m \end{array} \right\rangle \xrightarrow{\lim} \left(\begin{array}{c} p \\ m^{\prime} \end{array} \left P_{\pm} \right \begin{array}{c} p \\ m \end{array} \right) \\ &a \sqrt{(l \mp m)(l \pm m + 1)} \delta_{m^{\prime}, m \pm 1} \xrightarrow{\lim} p \delta_{m^{\prime}, m \pm 1} \end{aligned}$	$\begin{aligned} &\left\langle \begin{array}{c} l \\ m^{\prime} \end{array} \left a J_{\pm} \right \begin{array}{c} l \\ m \end{array} \right\rangle \xrightarrow{\lim} \left\langle \begin{array}{c} p \\ m^{\prime} \end{array} \left P_{\pm} \right \begin{array}{c} p \\ m \end{array} \right\rangle \\ &a \sqrt{(l \mp m)(l \pm m + 1)} \delta_{m^{\prime}, m \pm 1} \xrightarrow{\lim} p \delta_{m^{\prime}, m \pm 1} \end{aligned}$	$\begin{aligned} &\left\langle \begin{array}{c} l \\ m^{\prime} \end{array} \left a J_{\pm} \right \begin{array}{c} l \\ m \end{array} \right\rangle \xrightarrow{\lim} \left\langle \begin{array}{c} p \\ m^{\prime} \end{array} \left P_{\pm} \right \begin{array}{c} p \\ m \end{array} \right\rangle \\ &a \sqrt{(l \mp m)(l \pm m + 1)} \delta_{m^{\prime}, m \pm 1} \xrightarrow{\lim} p \delta_{m^{\prime}, m \pm 1} \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00873 (15.6)	276	0.640	$\{\Lambda, a\} = \left[\begin{array}{ccc c} & & & x \\ & \Lambda & & y \\ & & & z \\ \hline 0 & 0 & 0 & 0 \end{array} \right] \begin{array}{c} \\ \\ \\ 1 \end{array}$	$\{\Lambda, a\} = \left[\begin{array}{ccc c} & & & x \\ & \Lambda & & y \\ & & & z \\ \hline 0 & 0 & 0 & 0 \end{array} \right] \begin{array}{c} \\ \\ \\ 1 \end{array}$	$\{\Lambda, a\} = \left[\begin{array}{ccc c} & & & x \\ & \Lambda & & y \\ & & & z \\ \hline 0 & 0 & 0 & 0 \end{array} \right] \begin{array}{c} \\ \\ \\ 1 \end{array}$
gilmore-lie-groups_ E00766	240	0.655	$\begin{aligned} & r \rightarrow 0 : R \simeq r^\gamma, \gamma(\gamma - 1) + A = 0 \\ & r \rightarrow \infty : R \simeq e^{\lambda r}, \lambda^2 + C = 0 \end{aligned}$	$r \rightarrow 0 : R \simeq r^\gamma, \gamma(\gamma - 1) + A = 0$ $r \rightarrow \infty : R \simeq e^{\lambda r}, \lambda^2 + C = 0$	$r \rightarrow 0 : R \simeq r^\gamma, \gamma(\gamma - 1) + A = 0$ $r \rightarrow \infty : R \simeq e^{\lambda r}, \lambda^2 + C = 0$
gilmore-lie-groups_ E00919	286	0.656	$\begin{aligned} & \psi\rangle = \sum_{k,\xi} k;\xi\rangle \langle k;\xi \psi\rangle \\ & \psi\rangle = \sum_{k,\mu\mu'} k\rangle \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle \left\langle k; \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right \psi \rangle \end{aligned}$	$ \psi\rangle = \sum_{k,\xi} k;\xi\rangle \langle k;\xi \psi\rangle$ $ \psi\rangle = \sum_{k,\mu\mu'} k\rangle \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle \left\langle k; \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right \psi \rangle$	$ \psi\rangle = \sum_{k,\xi} k;\xi\rangle \langle k;\xi \psi\rangle$ $ \psi\rangle = \sum_{k,\mu\mu'} k\rangle \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle \left\langle k; \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right \psi \rangle$
gilmore-lie-groups_ E00375 (7.13)	117	0.666	$\begin{aligned} & \begin{array}{ccc} \left[\begin{array}{cc} a & c \\ -c & -a \end{array} \right] & + & \left[\begin{array}{cc} 0 & b \\ b & 0 \end{array} \right] \\ \text{EXP} \downarrow & & \downarrow \text{EXP} \\ \left[\begin{array}{cc} z+y & x \\ -x & z-y \end{array} \right] \times \left[\begin{array}{cc} \cosh b & \sinh b \\ \sinh b & \cosh b \end{array} \right] & & \end{array} \end{aligned}$	$\begin{aligned} & \begin{array}{ccc} \left[\begin{array}{cc} a & c \\ -c & -a \end{array} \right] & + & \left[\begin{array}{cc} 0 & b \\ b & 0 \end{array} \right] \\ \text{EXP} \downarrow & & \downarrow \text{EXP} \\ \left[\begin{array}{cc} z+y & x \\ -x & z-y \end{array} \right] \times \left[\begin{array}{cc} \cosh b & \sinh b \\ \sinh b & \cosh b \end{array} \right] & & \end{array} \end{aligned}$	$\begin{aligned} & \begin{array}{ccc} \left[\begin{array}{cc} a & c \\ -c & -a \end{array} \right] & + & \left[\begin{array}{cc} 0 & b \\ b & 0 \end{array} \right] \\ \text{EXP} \downarrow & & \downarrow \text{EXP} \\ \left[\begin{array}{cc} z+y & x \\ -x & z-y \end{array} \right] \times \left[\begin{array}{cc} \cosh b & \sinh b \\ \sinh b & \cosh b \end{array} \right] & & \end{array} \end{aligned}$
gilmore-lie-groups_ E00149 (3.32)	58	0.672	$\operatorname{Sp}(n) \xrightarrow{\mathbb{Q} \rightarrow 2 \times 2} \mathbb{C} U \operatorname{Sp}(2n) \subset \operatorname{SU}(2n)$	$\operatorname{Sp}(n) \xrightarrow{\mathbb{Q} \rightarrow 2 \times 2} \mathbb{C} U \operatorname{Sp}(2n) \subset \operatorname{SU}(2n)$	$Sp(n) \xrightarrow{\mathbb{Q} \rightarrow 2 \times 2} \mathbb{C} U Sp(2n) \subset SU(2n)$
gilmore-lie-groups_ E01034	321	0.672	$\begin{aligned} & \int \frac{\dim \lambda}{\operatorname{Vol}(G)} \Gamma_{\mu\nu}^{\lambda'}(g) \Gamma_{\mu\nu}^\lambda(g) d\rho(g) = \delta^{\lambda'\lambda} \delta_{\mu'\mu} \delta_{\nu'\nu} \\ & \sum_\lambda \sum_\mu \sum_\nu \frac{\dim \lambda}{\operatorname{Vol}(G)} \Gamma_{\mu\nu}^{\lambda*}(g') \Gamma_{\mu\nu}^\lambda(g) = \delta(g', g) \end{aligned}$	$\int \frac{\dim \lambda}{\operatorname{Vol}(G)} \Gamma_{\mu\nu}^{\lambda'}(g) \Gamma_{\mu\nu}^\lambda(g) d\rho(g) = \delta^{\lambda'\lambda} \delta_{\mu'\mu} \delta_{\nu'\nu}$ $\sum_\lambda \sum_\mu \sum_\nu \frac{\dim \lambda}{\operatorname{Vol}(G)} \Gamma_{\mu\nu}^{\lambda*}(g') \Gamma_{\mu\nu}^\lambda(g) = \delta(g', g)$	$\int \frac{\dim \lambda}{\operatorname{Vol}(G)} \Gamma_{\mu\nu}^{\lambda'}(g) \Gamma_{\mu\nu}^\lambda(g) d\rho(g) = \delta^{\lambda'\lambda} \delta_{\mu'\mu} \delta_{\nu'\nu}$ $\sum_\lambda \sum_\mu \sum_\nu \frac{\dim \lambda}{\operatorname{Vol}(G)} \Gamma_{\mu\nu}^{\lambda*}(g') \Gamma_{\mu\nu}^\lambda(g) = \delta(g', g)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0817 (14.53)	256	0.675	$\begin{array}{lrl} M_{\{1\}}=\left[\begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array}\right] & M_{\{2\}}=\left[\begin{array}{cccc} 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \end{array}\right] \\ \text{\texttt{\end{array}\right]} & M_{\{1\}}^t=+M_{\{1\}} & G_{\{1\}}^t=-G_{\{1\}} & M_{\{2\}}^t=-M_{\{2\}} & G_{\{2\}}^t=+G_{\{2\}} \end{array}$	$M_1=\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad M_2=\begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{bmatrix}$ $M_1^t=+M_1 \quad G_1^t=-G_1 \qquad M_2^t=-M_2 \quad G_2^t=+G_2$	
gilmore-lie-groups_ EQ0335 (6.20)	105	0.676	$\left[\begin{array}{c} x^{\prime} \\ 1 \end{array}\right]=\left[\begin{array}{cc} 1+d a & d b \\ 0 & 1 \end{array}\right]\left[\begin{array}{c} x \\ 1 \end{array}\right]$	$\left[\begin{array}{c} x' \\ 1 \end{array}\right] = \left[\begin{array}{cc} 1+da & db \\ 0 & 1 \end{array}\right] \left[\begin{array}{c} x \\ 1 \end{array}\right]$	$\left[\begin{array}{c} x' \\ 1 \end{array}\right] = \left[\begin{array}{cc} 1+da & db \\ 0 & 1 \end{array}\right] \left[\begin{array}{c} x \\ 1 \end{array}\right]$
gilmore-lie-groups_ EQ0905 (15.39)	282	0.677	$\begin{array}{l} \text{\texttt{\begin{aligned} \{\left[Y_{\{3\}}, Y_{\{1\}}\right] \} \& =-Y_{\{2\}} \\ \{\left[Y_{\{3\}}, Y_{\{2\}}\right] \} \& =+Y_{\{1\}} \\ \left[Y_{\{1\}}, Y_{\{2\}}\right] \} \& =0 \end{aligned}} \quad \operatorname{ISO}(2)} \end{array}$	$\begin{array}{l} [Y_3, Y_1] = -Y_2 \\ [Y_3, Y_2] = +Y_1 \quad \operatorname{ISO}(2) \\ [Y_1, Y_2] = 0 \end{array}$	$\begin{array}{l} [Y_3, Y_1] = -Y_2 \\ [Y_3, Y_2] = +Y_1 \qquad \operatorname{ISO}(2) \\ [Y_1, Y_2] = 0 \end{array}$
gilmore-lie-groups_ EQ0459 (7.85)	138	0.683	$\left \begin{array}{c} j \\ \theta \end{array}\right\rangle=S U(2)\left \begin{array}{c} j \\ -j \end{array}\right\rangle$	$\left \begin{array}{c} j \\ \theta \end{array}\right\rangle = S U(2) \left \begin{array}{c} j \\ -j \end{array}\right\rangle$	$\left \begin{array}{c} j \\ \theta \end{array}\right\rangle = S U(2) \left \begin{array}{c} j \\ -j \end{array}\right\rangle$
gilmore-lie-groups_ EQ0212 (4.22)	76	0.685	$C_{rs}^{\prime \, t} = (A^{-1})_r^i (A^{-1})_s^j C_{ij}^k A_k^t$	$C_{rs}^{\prime \, t} = (A^{-1})_r^i (A^{-1})_s^j C_{ij}^k A_k^t$	$C_{rs}^{\prime \, t} = (A^{-1})_r^i (A^{-1})_s^j C_{ij}^k A_k^t$
gilmore-lie-groups_ EQ0026 (1.22)	25	0.689	$\left[\begin{array}{rr} 1 & 1 \\ 1 & -1 \end{array}\right]\left[\begin{array}{l} r_1 \\ r_2 \end{array}\right]=\left[\begin{array}{c} I_1 \\ \pm\left[I_1^2-4 I_2\right]^{1 / 2} \end{array}\right]$	$\left[\begin{array}{cc} 1 & 1 \\ 1 & -1 \end{array}\right] \left[\begin{array}{c} r_1 \\ r_2 \end{array}\right] = \left[\begin{array}{c} I_1 \\ \pm\left[I_1^2-4 I_2\right]^{1 / 2} \end{array}\right]$	$\left[\begin{array}{cc} 1 & 1 \\ 1 & -1 \end{array}\right] \left[\begin{array}{c} r_1 \\ r_2 \end{array}\right] = \left[\begin{array}{c} I_1 \\ \pm\left[I_1^2-4 I_2\right]^{1 / 2} \end{array}\right]$
gilmore-lie-groups_ EQ1037	321	0.697	$\begin{array}{l} \text{\texttt{\begin{aligned} g\rangle\langle g \& =\int g\rangle d\rho(g)\langle g \quad \text{\texttt{\text{in group space}}} \\ \left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right]\left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right] \& =\sum_{\lambda} \sum_{\mu} \sum_{\nu}\left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right]\left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right] \quad \text{\texttt{\text{in representation space}}} \\ \left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right] \& =\sum_{\lambda} \sum_{\mu} \sum_{\nu}\left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right] \quad \text{\texttt{\text{in representation space}}} \end{aligned}} \end{array}$	$\begin{array}{l} g\rangle\langle g = \int g\rangle d\rho(g)\langle g \quad \text{in group space} \\ \left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right] \left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right] = \sum_{\lambda} \sum_{\mu} \sum_{\nu} \left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right] \left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right] = I \text{ in representation space} \end{array}$	$\begin{array}{l} g\rangle\langle g \quad = \quad \int g\rangle d\rho(g)\langle g \quad = I \quad \text{in group space} \\ \left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right] \left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right] = \sum_{\lambda} \sum_{\mu} \sum_{\nu} \left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right] \left[\begin{array}{c} \lambda \\ \mu \nu \end{array}\right] = I \text{ in representation space} \end{array}$
gilmore-lie-groups_ EQ0272 (5.11)	91	0.699	$\left[\begin{array}{lll} 0 & l & \delta \\ 0 & 0 & r \\ 0 & 0 & 0 \end{array}\right]=l X_l+r X_r+\delta X_{\delta}$	$\left[\begin{array}{ccc} 0 & l & \delta \\ 0 & 0 & r \\ 0 & 0 & 0 \end{array}\right] = l X_l + r X_r + \delta X_{\delta}$	$\left[\begin{array}{ccc} 0 & l & \delta \\ 0 & 0 & r \\ 0 & 0 & 0 \end{array}\right] = l X_l + r X_r + \delta X_{\delta}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E09165	63	■ 0.704	$SO(3) \rightarrow \left[\begin{array}{c c} Z_2 & \begin{smallmatrix} x \\ y \end{smallmatrix} \\ \hline -x & -y \end{array} \middle \begin{array}{c} Z_1 \\ \hline 1 \end{array} \right] \times \left[\begin{array}{cc c} \pm\sqrt{1-z^2} & z & 0 \\ -z & \pm\sqrt{1-z^2} & 0 \\ \hline 0 & 0 & 1 \end{array} \right]$	$SO(3) \rightarrow \left[\begin{array}{c c} Z_2 & \begin{smallmatrix} x \\ y \end{smallmatrix} \\ \hline -x & -y \end{array} \middle \begin{array}{c} Z_1 \\ \hline 1 \end{array} \right] \times \left[\begin{array}{cc c} \pm\sqrt{1-z^2} & z & 0 \\ -z & \pm\sqrt{1-z^2} & 0 \\ \hline 0 & 0 & 1 \end{array} \right]$	$SO(3) \rightarrow \left[\begin{array}{c c} Z_2 & \begin{smallmatrix} x \\ y \end{smallmatrix} \\ \hline -x & -y \end{array} \middle \begin{array}{c} Z_1 \\ \hline 1 \end{array} \right] \times \left[\begin{array}{cc c} \pm\sqrt{1-z^2} & z & 0 \\ -z & \pm\sqrt{1-z^2} & 0 \\ \hline 0 & 0 & 1 \end{array} \right]$
gilmore-lie-groups_ E08889	253	■ 0.712	$\begin{array}{l} \begin{array}{c} a_1^{\dagger} a_2 \\ b_1^{\dagger} b_2 \\ a_i^{\dagger} b_j^{\dagger} \end{array} \quad \begin{array}{c} a_2^{\dagger} a_1 \\ b_2^{\dagger} b_1 \\ a_i b_j \end{array} \quad \begin{array}{c} \frac{1}{2} (a_1^{\dagger} a_1 - a_2^{\dagger} a_2) \\ \frac{1}{2} (b_1^{\dagger} b_1 - b_2^{\dagger} b_2) \\ \frac{1}{2} (a_i^{\dagger} a_i + b_j^{\dagger} b_j + 1) \end{array} \quad \begin{array}{c} \mathfrak{su}(2) \\ \mathfrak{su}(2) \\ \mathfrak{su}(1, 1) \end{array} \end{array}$	$\begin{array}{l} \begin{array}{c} a_1^{\dagger} a_2 \\ b_1^{\dagger} b_2 \\ a_i^{\dagger} b_j^{\dagger} \end{array} \quad \begin{array}{c} a_2^{\dagger} a_1 \\ b_2^{\dagger} b_1 \\ a_i b_j \end{array} \quad \begin{array}{c} \frac{1}{2} (a_1^{\dagger} a_1 - a_2^{\dagger} a_2) \\ \frac{1}{2} (b_1^{\dagger} b_1 - b_2^{\dagger} b_2) \\ \frac{1}{2} (a_i^{\dagger} a_i + b_j^{\dagger} b_j + 1) \end{array} \quad \begin{array}{c} \mathfrak{su}(2) \\ \mathfrak{su}(2) \\ \mathfrak{su}(1, 1) \end{array} \end{array}$	$\begin{array}{l} a_1^{\dagger} a_2 \quad a_2^{\dagger} a_1 \quad \frac{1}{2} (a_1^{\dagger} a_1 - a_2^{\dagger} a_2) \quad \mathfrak{su}(2) \\ b_1^{\dagger} b_2 \quad b_2^{\dagger} b_1 \quad \frac{1}{2} (b_1^{\dagger} b_1 - b_2^{\dagger} b_2) \quad \mathfrak{su}(2) \\ a_i^{\dagger} b_j^{\dagger} \quad a_i b_j \quad \frac{1}{2} (a_i^{\dagger} a_i + b_j^{\dagger} b_j + 1) \quad \mathfrak{su}(1, 1) \end{array}$
gilmore-lie-groups_ E09902 (15.36)	282	■ 0.712	$\begin{array}{l} \begin{array}{c} \theta \\ -\theta_3 \\ +\theta_2 \\ -ib_1 \end{array} \quad \begin{array}{c} +\theta_3 \\ 0 \\ -\theta_1 \\ -ib_2 \end{array} \quad \begin{array}{c} -\theta_2 \\ +\theta_1 \\ 0 \\ -ib_3 \end{array} \quad \begin{array}{c} ib_1 \\ ib_2 \\ ib_3 \\ 0 \end{array} \quad \begin{array}{c} 0 \\ 0 \\ 1 \\ i \end{array} \quad \begin{array}{c} -\theta_2 - b_1 \\ +\theta_1 - b_2 \\ -b_3 \\ -ib_3 \end{array} \quad \begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array} \end{array}$	$\begin{bmatrix} 0 & +\theta_3 & -\theta_2 & ib_1 \\ -\theta_3 & 0 & +\theta_1 & ib_2 \\ +\theta_2 & -\theta_1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 1 \\ i \end{bmatrix} = \begin{bmatrix} -\theta_2 - b_1 \\ +\theta_1 - b_2 \\ -b_3 \\ -ib_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$	$\begin{bmatrix} 0 & +\theta_3 & -\theta_2 & ib_1 \\ -\theta_3 & 0 & +\theta_1 & ib_2 \\ +\theta_2 & -\theta_1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 1 \\ i \end{bmatrix} = \begin{bmatrix} -\theta_2 - b_1 \\ +\theta_1 - b_2 \\ -b_3 \\ -ib_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$
gilmore-lie-groups_ E09470 (8.1)	143	■ 0.714	$\left[Z, X_i \right] = R(Z)_i^j X_j$	$[Z, X_i] = R(Z)_i^j X_j$	$[Z, X_i] = R(Z)_i^j X_j$
gilmore-lie-groups_ E09822 (14.58)	257	■ 0.715	$\begin{array}{l} \Gamma_i = r \mathbf{p}_i \\ \Gamma_4 = \frac{1}{2} (r \mathbf{p} \cdot \mathbf{p} - r) \\ \Gamma_5 = \frac{1}{2} (r \mathbf{p} \cdot \mathbf{p} + r) \end{array}$	$\begin{array}{l} \Gamma_i = r \mathbf{p}_i \\ \Gamma_4 = \frac{1}{2} (r \mathbf{p} \cdot \mathbf{p} - r) \\ \Gamma_5 = \frac{1}{2} (r \mathbf{p} \cdot \mathbf{p} + r) \end{array}$	$\begin{array}{l} \Gamma_i = r \mathbf{p}_i \\ \Gamma_4 = \frac{1}{2} (r \mathbf{p} \cdot \mathbf{p} - r) \\ \Gamma_5 = \frac{1}{2} (r \mathbf{p} \cdot \mathbf{p} + r) \end{array}$
gilmore-lie-groups_ E09578	185	■ 0.715	$L_{ij} = a_i^{\dagger} a_j - a_j^{\dagger} a_i = -L_{ji} \quad Q_{ij} = a_i^{\dagger} a_j + a_j^{\dagger} a_i = +Q_{ji}$	$L_{ij} = a_i^{\dagger} a_j - a_j^{\dagger} a_i = -L_{ji} \quad Q_{ij} = a_i^{\dagger} a_j + a_j^{\dagger} a_i = +Q_{ji}$	$\begin{array}{l} \mathbf{L} \\ L_{ij} = a_i^{\dagger} a_j - a_j^{\dagger} a_i = -L_{ji} \quad Q_{ij} = a_i^{\dagger} a_j + a_j^{\dagger} a_i = +Q_{ji} \end{array}$
gilmore-lie-groups_ E09564	176	■ 0.717	$G_2 + \mathbf{e}_i - \mathbf{e}_j \pm [(\mathbf{e}_i + \mathbf{e}_j) - 2\mathbf{e}_k] \quad 1 \leq i \neq j \neq k \leq 3$	$G_2 + \mathbf{e}_i - \mathbf{e}_j \pm [(\mathbf{e}_i + \mathbf{e}_j) - 2\mathbf{e}_k] \quad 1 \leq i \neq j \neq k \leq 3$	$G_2 + \mathbf{e}_i - \mathbf{e}_j \pm [(\mathbf{e}_i + \mathbf{e}_j) - 2\mathbf{e}_k] \quad 1 \leq i \neq j \neq k \leq 3$
gilmore-lie-groups_ E09699	197	■ 0.718	$w = \alpha + \beta \quad x = \gamma + \delta \quad \text{inner product} \rightarrow 8(\beta^2 + \gamma^2 - \delta^2)$	$w = \alpha + \beta \quad x = \gamma + \delta \quad \text{inner product} \rightarrow 8(\beta^2 + \gamma^2 - \delta^2)$	$w = \alpha + \beta \quad x = \gamma + \delta \quad \text{inner product} \rightarrow 8(\beta^2 + \gamma^2 - \delta^2)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00792 (14.34)	248	■ 0.724	$\begin{gathered} (x, y, z) \in R^3 \rightarrow (w, x, y, z) \in R^4 \\ (p_x, p_y, p_z) \in R^3 \rightarrow (p_w, p_x, p_y, p_z) \in R^4 \end{gathered}$	$(x, y, z) \in R^3 \rightarrow (w, x, y, z) \in R^4$ $(p_x, p_y, p_z) \in R^3 \rightarrow (p_w, p_x, p_y, p_z) \in R^4$	$(x, y, z) \in R^3 \rightarrow (w, x, y, z) \in R^4$ $(p_x, p_y, p_z) \in R^3 \rightarrow (p_w, p_x, p_y, p_z) \in R^4$
gilmore-lie-groups_ E00042 (1.39)	28	■ 0.725	$\begin{aligned} & \quad y^3 - 0y^2 + I_2' y - I_3' = 0 \\ & y - I_3'^{\prime} = 0 \quad \& \quad I_1'^{\prime} = s_1' + s_2' + s_3' = 0 \\ & I_2' = s_1' s_2' + s_1' s_3' + s_2' s_3' = I_2 - (1/3) I_1^2 \\ & I_3' = s_1' s_2' s_3' = I_3 - (1/3) I_2 I_1 + (2/27) I_1^3 \end{aligned}$	$y^3 - 0y^2 + I_2' y - I_3' = 0$ $I_1' = s_1' + s_2' + s_3' = 0$ $I_2' = s_1' s_2' + s_1' s_3' + s_2' s_3' = I_2 - (1/3) I_1^2$ $I_3' = s_1' s_2' s_3' = I_3 - (1/3) I_2 I_1 + (2/27) I_1^3$	$y^3 - 0y^2 + I_2' y - I_3' = 0$ $I_1' = s_1' + s_2' + s_3' = 0$ $I_2' = s_1' s_2' + s_1' s_3' + s_2' s_3' = I_2 - (1/3) I_1^2$ $I_3' = s_1' s_2' s_3' = I_3 - (1/3) I_2 I_1 + (2/27) I_1^3$
gilmore-lie-groups_ E00114 (3.4)	49	■ 0.726	$\begin{array}{c} c_1 = a_1 + i b_1 \in \mathbb{C} \\ c_2 = a_2 + i b_2 \in \mathbb{C} \\ i^2 = -1, \quad j^2 = -1, \quad ij + ji = 0 \\ (c_1, c_2) = c_1 + j c_2 \\ (c_1, c_2) \longrightarrow \begin{bmatrix} c_1 & c_2 \\ -c_2^* & c_1^* \end{bmatrix} \end{array}$	$c_1 = a_1 + i b_1 \in \mathbb{C}$ $c_2 = a_2 + i b_2 \in \mathbb{C}$ $i^2 = -1, \quad j^2 = -1, \quad ij + ji = 0$ $(c_1, c_2) = c_1 + j c_2$ $(c_1, c_2) \longrightarrow \begin{bmatrix} c_1 & c_2 \\ -c_2^* & c_1^* \end{bmatrix}$	$c_1 = a_1 + i b_1 \in \mathbb{C}$ $q = (c_1, c_2) = c_1 + j c_2 \quad c_2 = a_2 + i b_2 \in \mathbb{C}$ $i^2 = -1, \quad j^2 = -1, \quad ij + ji = 0$ $(c_1, c_2) \longrightarrow \begin{bmatrix} c_1 & c_2 \\ -c_2^* & c_1^* \end{bmatrix}$
gilmore-lie-groups_ E00616	198	■ 0.728	$\begin{aligned} & \mathfrak{su}(2) \quad J_3, J_{\pm} \quad [J_3, J_{\pm}] = \pm J_{\pm} \quad [J_+, J_-] = +2J_3 \\ & \mathfrak{su}(1, 1) K_3, K_{\pm} \quad [K_3, K_{\pm}] = \pm K_{\pm} \quad [K_+, K_-] = -2K_3 \end{aligned}$	$\mathfrak{su}(2) \quad J_3, J_{\pm} \quad [J_3, J_{\pm}] = \pm J_{\pm} \quad [J_+, J_-] = +2J_3$ $\mathfrak{su}(1, 1) K_3, K_{\pm} \quad [K_3, K_{\pm}] = \pm K_{\pm} \quad [K_+, K_-] = -2K_3$	$\mathfrak{su}(2) \quad J_3, J_{\pm} \quad [J_3, J_{\pm}] = \pm J_{\pm} \quad [J_+, J_-] = +2J_3$ $\mathfrak{su}(1, 1) K_3, K_{\pm} \quad [K_3, K_{\pm}] = \pm K_{\pm} \quad [K_+, K_-] = -2K_3$
gilmore-lie-groups_ E00612	197	■ 0.729	$\begin{aligned} & \operatorname{Spin}(3) = S U(2) \quad \& \quad \operatorname{Spin}(4) = S U(2) \otimes S U(2) \\ & \operatorname{Spin}(5) = U \operatorname{Sp}(4) \\ & \operatorname{Spin}(6) = S U(4) \end{aligned}$	$\operatorname{Spin}(3) = S U(2)$ $\operatorname{Spin}(4) = S U(2) \otimes S U(2)$ $\operatorname{Spin}(5) = U \operatorname{Sp}(4)$ $\operatorname{Spin}(6) = S U(4)$	$\operatorname{Spin}(3) = S U(2)$ $\operatorname{Spin}(4) = S U(2) \otimes S U(2)$ $\operatorname{Spin}(5) = U \operatorname{Sp}(4)$ $\operatorname{Spin}(6) = S U(4)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00788 (14.30) [re-fined: census]	247	0.732	$\begin{aligned} M_{+}^{\prime}\wedge Y_m^l &= \left\langle \theta \phi \middle \left(\begin{array}{cc} j/2 & j/2 \\ m_a+1 & m_b \end{array} \right) \begin{array}{c} j/2 \\ m_a \end{array} \begin{array}{c} j/2 \\ m_b \end{array} \middle m \right\rangle \times \sqrt{(j/2-m_a)(j/2+m_a+1)} \\ &\quad - \begin{array}{c} j/2 \\ m_a \end{array} \begin{array}{c} j/2 \\ m_b+1 \end{array} \middle \begin{array}{c} j/2 \\ m_a \end{array} \begin{array}{c} j/2 \\ m_b \end{array} \middle m \right\rangle \times \sqrt{(j/2-m_b)(j/2+m_b+1)} \Bigg) \\ &\quad - \left\langle \begin{array}{cc} j/2 & j/2 \\ m_a & m_b+1 \end{array} \right\rangle \begin{array}{c} j/2 \\ m_a \end{array} \begin{array}{c} j/2 \\ m_b \end{array} \middle m \right\rangle \times \sqrt{(j/2-m_b)(j/2+m_b+1)} \Bigg) \end{aligned}$	$M_{+}^{\prime}Y_m^l = \left\langle \theta \phi \middle \left(\begin{array}{cc} j/2 & j/2 \\ m_a+1 & m_b \end{array} \right) \begin{array}{c} j/2 \\ m_a \end{array} \begin{array}{c} j/2 \\ m_b \end{array} \middle m \right\rangle \times \sqrt{(j/2-m_a)(j/2+m_a+1)} - \begin{array}{c} j/2 \\ m_a \end{array} \begin{array}{c} j/2 \\ m_b+1 \end{array} \middle \begin{array}{c} j/2 \\ m_a \end{array} \begin{array}{c} j/2 \\ m_b \end{array} \middle m \right\rangle \times \sqrt{(j/2-m_b)(j/2+m_b+1)} \Bigg)$	$M_{+}^{\prime}Y_m^l = \langle \theta \phi \left(\begin{array}{cc} j/2 & j/2 \\ m_a+1 & m_b \end{array} \right) \begin{array}{c} j/2 \\ m_a \end{array} \begin{array}{c} j/2 \\ m_b \end{array} \middle m \right\rangle \times \sqrt{(j/2-m_a)(j/2+m_a+1)} - \begin{array}{c} j/2 \\ m_a \end{array} \begin{array}{c} j/2 \\ m_b+1 \end{array} \middle \begin{array}{c} j/2 \\ m_a \end{array} \begin{array}{c} j/2 \\ m_b \end{array} \middle m \right\rangle \times \sqrt{(j/2-m_b)(j/2+m_b+1)} \Bigg)$
gilmore-lie-groups_ E00938	290	0.733	$\begin{aligned} \mathbf{B} + i\mathbf{E} &\rightarrow \mathbf{B}' + i\mathbf{E}' = e^{i\phi}(\mathbf{B} + i\mathbf{E}) \\ \mathbf{J}_m + i\mathbf{J}_e &\rightarrow \mathbf{J}'_m + i\mathbf{J}'_e = e^{i\phi}(\mathbf{J}_m + i\mathbf{J}_e) \end{aligned}$	$\mathbf{B} + i\mathbf{E} \rightarrow \mathbf{B}' + i\mathbf{E}' = e^{i\phi}(\mathbf{B} + i\mathbf{E})$ $\mathbf{J}_m + i\mathbf{J}_e \rightarrow \mathbf{J}'_m + i\mathbf{J}'_e = e^{i\phi}(\mathbf{J}_m + i\mathbf{J}_e)$	$\mathbf{B} + i\mathbf{E} \rightarrow \mathbf{B}' + i\mathbf{E}' = e^{i\phi}(\mathbf{B} + i\mathbf{E})$ $\mathbf{J}_m + i\mathbf{J}_e \rightarrow \mathbf{J}'_m + i\mathbf{J}'_e = e^{i\phi}(\mathbf{J}_m + i\mathbf{J}_e)$
gilmore-lie-groups_ E00226 (4.36)	79	0.737	$\left(X_i,X_j\right)=\operatorname{tr}R\left(X_i\right)\left(X_j\right)=\sum_r\sum_sR\left(X_i\right)_r^{s*}R\left(X_j\right)_r^s=\sum_r\sum_sC_{ir}^{s*}C_{jr}^s$	$(X_i,X_j)=\operatorname{tr}R(X_i)^{\dagger}R(X_j)=\sum_r\sum_sR(X_i)_r^{s*}R(X_j)_r^s=\sum_r\sum_sC_{ir}^{s*}C_{jr}^s$	$(X_i,X_j)=\operatorname{tr}R(X_i)^{\dagger}R(X_j)=\sum_r\sum_sR(X_i)_r^{s*}R(X_j)_r^s=\sum_r\sum_sC_{ir}^{s*}C_{jr}^s$
gilmore-lie-groups_ E00723 (13.40)	227	0.737	$\lim_{c\rightarrow 0}(-J+\frac{1}{2c^2})=0 \text{ implies } 2Jc^2=1$	$\lim_{c\rightarrow 0}\left(-J+\frac{1}{2c^2}\right)=0 \text{ implies } 2Jc^2=1$	$\lim_{c\rightarrow 0}\left(-J+\frac{1}{2c^2}\right)=0 \text{ implies } 2Jc^2=1$
gilmore-lie-groups_ E00683 (13.1)	219	0.743	$C_{rs}^t(0)=\lim_{\epsilon\rightarrow 0}C_{rs}^t(\epsilon)$	$C_{rs}^t(0)=\lim_{\epsilon\rightarrow 0}C_{rs}^t(\epsilon)$	$C_{rs}^{\textcolor{teal}{t}}(0)=\lim_{\epsilon\rightarrow 0}C_{rs}^{\textcolor{teal}{t}}(\epsilon)$
gilmore-lie-groups_ E00685 (13.3)	220	0.743	$\left[\begin{array}{l} \mathbf{h}' \\ \mathbf{p}' \end{array}\right]=\left[\begin{array}{cc} I_{\dim(\mathbf{h})} & 0 \\ 0 & \epsilon I_{\dim(\mathbf{p})} \end{array}\right]\left[\begin{array}{l} \mathbf{h} \\ \mathbf{p} \end{array}\right]$	$\left[\begin{array}{l} \mathbf{h}' \\ \mathbf{p}' \end{array}\right]=\left[\begin{array}{cc} I_{\dim(\mathbf{h})} & 0 \\ 0 & \epsilon I_{\dim(\mathbf{p})} \end{array}\right]\left[\begin{array}{l} \mathbf{h} \\ \mathbf{p} \end{array}\right]$	$\left[\begin{array}{l} \mathbf{h}' \\ \mathbf{p}' \end{array}\right]=\left[\begin{array}{cc} I_{\dim(\mathbf{h})} & 0 \\ 0 & \epsilon I_{\dim(\mathbf{p})} \end{array}\right]\left[\begin{array}{l} \mathbf{h} \\ \mathbf{p} \end{array}\right]$
gilmore-lie-groups_ E00184	67	0.749	$S_{\mathrm{Tot}}=\left(\begin{array}{cc} 0 & s_{34} \\ s_{21} & 0 \end{array}\right)+\frac{1}{1-s_{12}s_{43}}\left[\begin{array}{cc} s_{33}s_{22} & s_{33}s_{112}s_{44} \\ s_{22}s_{43}s_{11} & s_{22}s_{44} \end{array}\right]$	$S_{\mathrm{Tot}}=\left[\begin{array}{cc} 0 & s_{34} \\ s_{21} & 0 \end{array}\right]+\frac{1}{1-s_{12}s_{43}}\left[\begin{array}{cc} s_{33}s_{22} & s_{33}s_{112}s_{44} \\ s_{22}s_{43}s_{11} & s_{22}s_{44} \end{array}\right]$	$S_{\mathrm{Tot}}=\left[\begin{array}{cc} 0 & s_{34} \\ s_{21} & 0 \end{array}\right]+\frac{1}{1-s_{12}s_{43}}\left[\begin{array}{cc} s_{33}s_{22} & s_{33}s_{112}s_{44} \\ s_{22}s_{43}s_{11} & s_{22}s_{44} \end{array}\right]$
gilmore-lie-groups_ E00618	199	0.749	$\begin{aligned} &\left J_{+} n_1,n_2\rangle\right =\sqrt{(n_1+1)n_2} \\ &\left J_{-} n_1,n_2\rangle\right =\sqrt{n_1(n_2+1)} \end{aligned}$	$\begin{aligned} \langle n_1+1,n_2-1 J_{+} n_1,n_2\rangle &=\sqrt{(n_1+1)n_2} \\ \langle n_1,n_2 J_{-} n_1+1,n_2-1\rangle^* &=\sqrt{n_1(n_2+1)}^* \end{aligned}$	$\begin{aligned} \langle n_1+1,n_2-1 J_{+} n_1,n_2\rangle &=\sqrt{(n_1+1)n_2} \\ \langle n_1,n_2 J_{-} n_1+1,n_2-1\rangle^* &=\sqrt{n_1(n_2+1)}^* \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08942 (15.72)	291	0.751	$\begin{array}{ll} \operatorname{Re} & +\frac{2}{c}\frac{\partial \mathbf{G}_{\mathbf{e}}}{\partial t}-\nabla \times \mathbf{G}_{\mathbf{m}}=0 \\ \operatorname{Im} & -\frac{2}{c}\frac{\partial \mathbf{G}_{\mathbf{m}}}{\partial t}-\nabla \times \mathbf{G}_{\mathbf{e}}=0 \end{array}$	$\begin{array}{ll} \operatorname{Re} & +\frac{2}{c}\frac{\partial \mathbf{G}_{\mathbf{e}}}{\partial t}-\nabla \times \mathbf{G}_{\mathbf{m}}=0 \\ \operatorname{Im} & -\frac{2}{c}\frac{\partial \mathbf{G}_{\mathbf{m}}}{\partial t}-\nabla \times \mathbf{G}_{\mathbf{e}}=0 \end{array}$	$\begin{array}{ll} \operatorname{Re} & +\frac{2}{c}\frac{\partial \mathbf{G}_{\mathbf{e}}}{\partial t}-\nabla \times \mathbf{G}_{\mathbf{m}}=0 \\ \operatorname{Im} & -\frac{2}{c}\frac{\partial \mathbf{G}_{\mathbf{m}}}{\partial t}-\nabla \times \mathbf{G}_{\mathbf{e}}=0 \end{array}$
gilmore-lie-groups_ E08741	232	0.752	$\begin{array}{l} \lim \epsilon \rightarrow \infty \quad \epsilon J_{\pm} \rightarrow h_{\pm} \quad 2j\epsilon^2 \rightarrow 1 \\ j \rightarrow +\infty, m \rightarrow -\infty \quad J_3 + \frac{1}{2\epsilon^2} \rightarrow h_3 \quad \left \begin{array}{c} j \\ m \end{array} \right\rangle \rightarrow \left \begin{array}{c} \infty \\ n \end{array} \right\rangle \\ j+m=n \text{ (finite)} \quad \theta \rightarrow \frac{\pi}{2} - \sqrt{2}\epsilon x \end{array}$	$\begin{array}{l} \lim \epsilon \rightarrow \infty \quad \epsilon J_{\pm} \rightarrow h_{\pm} \quad 2j\epsilon^2 \rightarrow 1 \\ j \rightarrow +\infty, m \rightarrow -\infty \quad J_3 + \frac{1}{2\epsilon^2} \rightarrow h_3 \quad \left \begin{array}{c} j \\ m \end{array} \right\rangle \rightarrow \\ j+m=n \text{ (finite)} \quad \theta \rightarrow \frac{\pi}{2} - \sqrt{2}\epsilon x \end{array}$	$\begin{array}{l} \lim \epsilon \rightarrow \infty \quad \epsilon J_{\pm} \rightarrow h_{\pm} \quad 2j\epsilon^2 \rightarrow 1 \\ j \rightarrow +\infty, m \rightarrow -\infty \quad J_3 + \frac{1}{2\epsilon^2} \rightarrow h_3 \quad \left \begin{array}{c} j \\ m \end{array} \right\rangle \rightarrow \left \begin{array}{c} \infty \\ n \end{array} \right\rangle \\ j+m=n \text{ (finite)} \quad \theta \rightarrow \frac{\pi}{2} - \sqrt{2}\epsilon x \end{array}$
gilmore-lie-groups_ E08884	252	0.753	$\begin{array}{l} \left[\begin{array}{cccc c} 0 & L_3 & -L_2 & M_1 & B_1 \\ -L_3 & 0 & L_1 & M_2 & B_2 \\ L_2 & -L_1 & 0 & M_3 & B_3 \\ -M_1 & -M_2 & -M_3 & 0 & B_4 \\ \hline B_1 & B_2 & B_3 & B_4 & 0 \end{array} \right] + \\ + \\ + \\ + \\ + \\ - \end{array}$	$\begin{array}{l} \left[\begin{array}{cccc c} 0 & L_3 & -L_2 & M_1 & B_1 \\ -L_3 & 0 & L_1 & M_2 & B_2 \\ L_2 & -L_1 & 0 & M_3 & B_3 \\ -M_1 & -M_2 & -M_3 & 0 & B_4 \\ \hline B_1 & B_2 & B_3 & B_4 & 0 \end{array} \right] + \\ + \\ + \\ + \\ + \\ - \end{array}$	$\begin{array}{l} \left[\begin{array}{cccc c} 0 & L_3 & -L_2 & M_1 & B_1 \\ -L_3 & 0 & L_1 & M_2 & B_2 \\ L_2 & -L_1 & 0 & M_3 & B_3 \\ -M_1 & -M_2 & -M_3 & 0 & B_4 \\ \hline B_1 & B_2 & B_3 & B_4 & 0 \end{array} \right] + \\ + \\ + \\ + \\ + \\ - \end{array}$
gilmore-lie-groups_ E08873 (1.68)	34	0.760	$\begin{array}{l} x_1' - x_2' = x_1 - x_2 \\ = \pm 2i \times 145152\sqrt{3} = \pm 2\sqrt{-K_2} = \pm 2i\sqrt{K_2} \\ = \pm 290304\sqrt{3} \end{array}$	$\begin{array}{l} x_1' - x_2' = x_1 - x_2 \\ = \pm 2i \times 145152\sqrt{3} = \pm 2\sqrt{-K_2} = \pm 2i\sqrt{K_2} \\ = \pm 290304\sqrt{3} \end{array}$	$\begin{array}{l} x_1' - x_2' = x_1 - x_2 \\ = \pm 2i \times 145152\sqrt{3} = \pm 2\sqrt{-K_2} = \pm 2i\sqrt{K_2} \\ = \pm 290304\sqrt{3} \end{array}$
gilmore-lie-groups_ E08735	231	0.761	$\begin{array}{l} \lim a \downarrow 0 \quad aJ_{\pm} \rightarrow P_{\pm} \quad a^2l(l+1) \rightarrow p^2 \text{ finite} \\ l \uparrow \infty \quad J_3 \rightarrow P_3 \quad \left \begin{array}{c} l \\ m \end{array} \right\rangle \rightarrow \left \begin{array}{c} p \\ m \end{array} \right\rangle \\ (p/a)\beta = l\beta = x \text{ finite} \end{array}$	$\begin{array}{l} \lim a \downarrow 0 \quad aJ_{\pm} \rightarrow P_{\pm} \quad a^2l(l+1) \rightarrow p^2 \text{ finite} \\ l \uparrow \infty \quad J_3 \rightarrow P_3 \quad \left \begin{array}{c} l \\ m \end{array} \right\rangle \rightarrow \left \begin{array}{c} p \\ m \end{array} \right\rangle \\ (p/a)\beta = l\beta = x \text{ finite} \end{array}$	$\begin{array}{l} \lim a \downarrow 0 \quad aJ_{\pm} \rightarrow P_{\pm} \quad a^2l(l+1) \rightarrow p^2 \text{ finite} \\ l \uparrow \infty \quad J_3 \rightarrow P_3 \quad \left \begin{array}{c} l \\ m \end{array} \right\rangle \rightarrow \left \begin{array}{c} p \\ m \end{array} \right\rangle \\ (p/a)\beta = l\beta = x \text{ finite} \end{array}$
gilmore-lie-groups_ E08397 (7.33)	124	0.762	$\begin{array}{l} e^{ra^{\dagger}+la+\delta I} \\ \parallel \\ e^{r'a^{\dagger}}e^{\delta'I}e^{l'a} = \parallel \\ e^{l''a}e^{\delta''I}e^{r''a^{\dagger}} \end{array}$	$\begin{array}{l} e^{ra^{\dagger}+la+\delta I} \\ \parallel \\ e^{r'a^{\dagger}}e^{\delta'I}e^{l'a} = \parallel \\ e^{l''a}e^{\delta''I}e^{r''a^{\dagger}} \end{array}$	$\begin{array}{l} e^{ra^{\dagger}+la+\delta I} \\ \parallel \\ e^{r'a^{\dagger}}e^{\delta'I}e^{l'a} = \parallel \\ e^{l''a}e^{\delta''I}e^{r''a^{\dagger}} \end{array}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08280 (4.13)	74	0.766	$\begin{aligned} & \operatorname{EXP}(\epsilon X) \operatorname{EXP}(\delta Y) \operatorname{EXP}(-\epsilon X) \operatorname{EXP}(-\delta Y) \\ &= I + \epsilon \delta (XY - YX) = I + \epsilon \delta [X, Y] \end{aligned}$	$\operatorname{EXP}(\epsilon X) \operatorname{EXP}(\delta Y) \operatorname{EXP}(-\epsilon X) \operatorname{EXP}(-\delta Y) = I + \epsilon \delta (XY - YX) = I + \epsilon \delta [X, Y]$	$\operatorname{EXP}(\epsilon X) \operatorname{EXP}(\delta Y) \operatorname{EXP}(-\epsilon X) \operatorname{EXP}(-\delta Y) = I + \epsilon \delta (XY - YX) = I + \epsilon \delta [X, Y]$
gilmore-lie-groups_ E08499 (9.8)	155	0.768	$\begin{aligned} & \operatorname{def}(X) = \frac{1}{2} \begin{bmatrix} ia_3 & i(a_1 - ia_2) \\ i(a_1 + ia_2) & -ia_3 \end{bmatrix} \\ & \operatorname{Reg}(X) = \begin{bmatrix} 0 & -a_3 & a_2 \\ a_3 & 0 & -a_1 \\ -a_2 & a_1 & 0 \end{bmatrix} \end{aligned}$	$\operatorname{def}(X) = \frac{1}{2} \begin{bmatrix} ia_3 & i(a_1 - ia_2) \\ i(a_1 + ia_2) & -ia_3 \end{bmatrix}$ $\operatorname{Reg}(X) = \begin{bmatrix} 0 & -a_3 & a_2 \\ a_3 & 0 & -a_1 \\ -a_2 & a_1 & 0 \end{bmatrix}$	$\operatorname{def}(X) = \frac{1}{2} \begin{bmatrix} ia_3 & i(a_1 - ia_2) \\ i(a_1 + ia_2) & -ia_3 \end{bmatrix}$ $\operatorname{Reg}(X) = \begin{bmatrix} 0 & -a_3 & a_2 \\ a_3 & 0 & -a_1 \\ -a_2 & a_1 & 0 \end{bmatrix}$
gilmore-lie-groups_ E0887 (15.20)	279	0.769	$ k\rangle \times \begin{bmatrix} j & j' \\ \mu & \mu' \end{bmatrix}$	$ k\rangle \times \begin{bmatrix} j & j' \\ \mu & \mu' \end{bmatrix}$	$ k\rangle \times \begin{bmatrix} j & j' \\ \mu & \mu' \end{bmatrix}$
gilmore-lie-groups_ E0848 (1.37)	27	0.772	$\begin{aligned} & J_1 = \sum_{i+2j+3k=3} A_{ijk} I_1^i I_2^j I_3^k \\ & J_2 = \sum_{i+2j+3k=6} B_{ijk} I_1^i I_2^j I_3^k \end{aligned}$	$J_1 = \sum_{i+2j+3k=3} A_{ijk} I_1^i I_2^j I_3^k$ $J_2 = \sum_{i+2j+3k=6} B_{ijk} I_1^i I_2^j I_3^k$	$J_1 = \sum_{i+2j+3k=3} A_{ijk} I_1^i I_2^j I_3^k$ $J_2 = \sum_{i+2j+3k=6} B_{ijk} I_1^i I_2^j I_3^k$
gilmore-lie-groups_ E0841 (6.26)	106	0.772	$\begin{aligned} & \begin{bmatrix} x' & y' \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1+da & db \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x & y \\ 0 & 1 \end{bmatrix} \\ & \begin{bmatrix} x' & y' \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1+da & db \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x & y \\ 0 & 1 \end{bmatrix} \end{aligned}$	$\begin{bmatrix} x' & y' \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1+da & db \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x & y \\ 0 & 1 \end{bmatrix}$	$\begin{bmatrix} x' & y' \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1+da & db \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x & y \\ 0 & 1 \end{bmatrix}$
gilmore-lie-groups_ E0928 (15.54)	286	0.772	$\begin{aligned} & \left\{ J_3^{(j)} k_3^0 - j k_4^0 I_{2j+1} \right\} \left\langle k^0; \begin{bmatrix} j & 0 \\ m & 0 \end{bmatrix} \middle \psi \right\rangle = 0 \\ & \left\{ J_3^{(j)} k_3^0 - j k_4^0 I_{2j+1} \right\} \left\langle k^0; \begin{bmatrix} j & 0 \\ m & 0 \end{bmatrix} \middle \psi \right\rangle = 0 \end{aligned}$	$\left\{ J_3^{(j)} k_3^0 - j k_4^0 I_{2j+1} \right\} \left\langle k^0; \begin{bmatrix} j & 0 \\ m & 0 \end{bmatrix} \middle \psi \right\rangle = 0$	$\left\{ J_3^{(j)} k_3^0 - j k_4^0 I_{2j+1} \right\} \left\langle k^0; \begin{bmatrix} j & 0 \\ m & 0 \end{bmatrix} \middle \psi \right\rangle = 0$
gilmore-lie-groups_ E0146 (3.29)	57	0.774	$\begin{aligned} & SO(n) = O(n) \cap SL(n; \mathbb{R}) \\ & SO(p, q) = O(p, q) \cap SL(p + q; \mathbb{R}) \\ & SU(n) = U(n) \cap SL(n; \mathbb{C}) \\ & SU(p, q) = U(p, q) \cap SL(p + q; \mathbb{C}) \end{aligned}$	$SO(n) = O(n) \cap SL(n; \mathbb{R})$ $SO(p, q) = O(p, q) \cap SL(p + q; \mathbb{R})$ $SU(n) = U(n) \cap SL(n; \mathbb{C})$ $SU(p, q) = U(p, q) \cap SL(p + q; \mathbb{C})$	$SO(n) = O(n) \cap SL(n; \mathbb{R})$ $SO(p, q) = O(p, q) \cap SL(p + q; \mathbb{R})$ $SU(n) = U(n) \cap SL(n; \mathbb{C})$ $SU(p, q) = U(p, q) \cap SL(p + q; \mathbb{C})$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08725 (13.42)	228	0.776	$\begin{aligned} M=-J+n\rangle &= \frac{(J_+)^n}{[(2J)!n!/(2J-n)!]^{1/2}} \begin{vmatrix} J \\ -J \end{vmatrix} \\ &= \lim_{J \rightarrow \infty} \frac{(cJ_+)^n}{[(2Jc^2)^n n!]^{1/2}} \begin{vmatrix} \infty \\ 0 \end{vmatrix} \\ &= \frac{(a^\dagger)^n}{\sqrt{n!}} \begin{vmatrix} \infty \\ 0 \end{vmatrix} \end{aligned}$	$ M=-J+n\rangle = \frac{(J_+)^n}{[(2J)!n!/(2J-n)!]^{1/2}} \begin{vmatrix} J \\ -J \end{vmatrix}$ $\begin{vmatrix} \infty \\ n \end{vmatrix} = \lim_{J \rightarrow \infty} \frac{(cJ_+)^n}{[(2Jc^2)^n n!]^{1/2}} \begin{vmatrix} \infty \\ 0 \end{vmatrix} = \frac{(a^\dagger)^n}{\sqrt{n!}} \begin{vmatrix} \infty \\ 0 \end{vmatrix}$	$\begin{vmatrix} J \\ M=-J+n \end{vmatrix} = \frac{(J_+)^n}{[(2J)!n!/(2J-n)!]^{1/2}} \begin{vmatrix} J \\ -J \end{vmatrix}$ $\begin{vmatrix} \infty \\ n \end{vmatrix} = \lim_{J \rightarrow \infty} \frac{(cJ_+)^n}{[(2Jc^2)^n n!]^{1/2}} \begin{vmatrix} \infty \\ 0 \end{vmatrix} = \frac{(a^\dagger)^n}{\sqrt{n!}} \begin{vmatrix} \infty \\ 0 \end{vmatrix}$
gilmore-lie-groups_ E08118 (3.6)	50	0.791	$\begin{array}{l} m_{i\alpha} = 0 \quad p+1 \leq i \leq p+q \\ \quad \quad \quad 1 \leq \alpha \leq p \end{array}$	$m_{i\alpha} = 0 \quad \begin{array}{l} p+1 \leq i \leq p+q \\ 1 \leq \alpha \leq p \end{array}$	$m_{i\alpha} = 0 \quad \begin{array}{l} p+1 \leq i \leq p+q \\ 1 \leq \alpha \leq p \end{array}$
gilmore-lie-groups_ E08565	176	0.791	$\begin{aligned} F_4 &\pm \mathbf{e}_i \pm \mathbf{e}_j \\ &\pm 2\mathbf{e}_i \\ &\pm \mathbf{e}_1 \pm \mathbf{e}_2 \pm \mathbf{e}_3 \pm \mathbf{e}_4 \\ 1 \leq i \neq j \leq 4 \end{aligned}$	$F_4 \pm \mathbf{e}_i \pm \mathbf{e}_j \pm 2\mathbf{e}_i \pm \mathbf{e}_1 \pm \mathbf{e}_2 \pm \mathbf{e}_3 \pm \mathbf{e}_4 \quad 1 \leq i \neq j \leq 4$	$F_4 \pm \mathbf{e}_i \pm \mathbf{e}_j \pm 2\mathbf{e}_i \pm \mathbf{e}_1 \pm \mathbf{e}_2 \pm \mathbf{e}_3 \pm \mathbf{e}_4 \quad 1 \leq i \neq j \leq 4$
gilmore-lie-groups_ E08658 (1.46)	29	0.796	$\begin{aligned} &\left(z^{\{\prime\}}-t_1\right)\left(z^{\{\prime\}}-t_2\right)\left(z^{\{\prime\}}-t_3\right)\left(z^{\{\prime\}}-t_4\right)=z^{\prime 4}-I_1^{\prime} z^{\prime 3}+I_2^{\prime} z^{\prime 2}-I_3^{\prime} z^{\prime}+I_4^{\prime}=0 \\ &I_1^{\prime}=0 \\ &I_2^{\prime}=I_2-\frac{3}{8} I_1^2 \\ &I_3^{\prime}=I_3-\frac{1}{2} I_2 I_1+\frac{1}{8} I_3^3 \\ &I_4^{\prime}=I_4-\frac{1}{4} I_3 I_1+\frac{1}{16} I_2 I_1^2-\frac{3}{4^4} I_1^4 \end{aligned}$	$\begin{aligned} (z'-t_1)(z'-t_2)(z'-t_3)(z'-t_4) &= z'^4 - I_1' z'^3 + I_2' z'^2 - I_3' z' + I_4' = 0 \\ I_1' &= 0 \\ I_2' &= I_2 - \frac{3}{8} I_1^2 \\ I_3' &= I_3 - \frac{1}{2} I_2 I_1 + \frac{1}{8} I_3^3 \\ I_4' &= I_4 - \frac{1}{4} I_3 I_1 + \frac{1}{16} I_2 I_1^2 - \frac{3}{4^4} I_1^4 \end{aligned}$	$\begin{aligned} (z'-t_1)(z'-t_2)(z'-t_3)(z'-t_4) &= z'^4 - I_1' z'^3 + I_2' z'^2 - I_3' z' + I_4' = 0 \\ I_1' &= 0 \\ I_2' &= I_2 - \frac{3}{8} I_1^2 \\ I_3' &= I_3 - \frac{1}{2} I_2 I_1 + \frac{1}{8} I_3^3 \\ I_4' &= I_4 - \frac{1}{4} I_3 I_1 + \frac{1}{16} I_2 I_1^2 - \frac{3}{4^4} I_1^4 \end{aligned}$
gilmore-lie-groups_ E08131 (3.17)	54	0.798	$S U(1 ; \mathbb{Q}) \sim S^3 \subset R^4$	$S U(1; \mathbb{Q}) \sim S^3 \subset R^4$	$S U(1; \mathbb{Q}) \sim S^3 \subset R^4$
gilmore-lie-groups_ E08695 (13.13)	222	0.798	$\begin{aligned} L_i &= \epsilon_{ijk} x_j \partial_k \\ V_i &= x_i \partial_4 - x_4 \partial_i \quad 1 \leq i \neq j \neq k \leq 3 \end{aligned}$	$L_i = \epsilon_{ijk} x_j \partial_k$ $V_i = x_i \partial_4 - x_4 \partial_i \quad 1 \leq i \neq j \neq k \leq 3$	$L_i = \epsilon_{ijk} x_j \partial_k$ $V_i = x_i \partial_4 - x_4 \partial_i \quad 1 \leq i \neq j \neq k \leq 3$
gilmore-lie-groups_ E08363 (7.1)	113	0.799	$\text{Lie groups} \underset{\text{EXP}}{\stackrel{\text{ln}}{\rightleftharpoons}} \text{Lie algebras}$	$\text{Lie groups} \underset{\text{EXP}}{\stackrel{\text{ln}}{\rightleftharpoons}} \text{Lie algebras}$	$\text{Lie groups} \underset{\text{EXP}}{\stackrel{\text{ln}}{\rightleftharpoons}} \text{Lie algebras}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E0655 (12.24)	212	0.802	$\begin{aligned} \mathfrak{g}^{\prime} &= \left(\mathfrak{g}_{+,+} + i \mathfrak{g}_{+,-} \right) + \left(i \mathfrak{g}_{-,+} + \mathfrak{g}_{-,-} \right) \\ &= \mathfrak{h}^{\prime} + \mathfrak{p}^{\prime} \end{aligned}$	$\mathfrak{g}'' = (\mathfrak{g}_{+,+} + i\mathfrak{g}_{+,-}) + (i\mathfrak{g}_{-,+} + \mathfrak{g}_{-,-})$ $= \mathfrak{h}'' + \mathfrak{p}''$	$\mathfrak{g}'' = (\mathfrak{g}_{+,+} + i\mathfrak{g}_{+,-}) + (i\mathfrak{g}_{-,+} + \mathfrak{g}_{-,-})$ $= \mathfrak{h}'' + \mathfrak{p}''$
gilmore-lie-groups_ E0300	98	0.803	$\begin{aligned} Z &= \left[\begin{array}{c c} A & B \\ \hline C & D \end{array} \right] = \left[\begin{array}{c c} A & 0 \\ \hline 0 & D \end{array} \right] + \left[\begin{array}{c c} 0 & B \\ \hline C & 0 \end{array} \right] \\ &= \mathfrak{h} + \mathfrak{p} \end{aligned}$	$Z = \left[\begin{array}{c c} A & B \\ \hline C & D \end{array} \right] = \left[\begin{array}{c c} A & 0 \\ \hline 0 & D \end{array} \right] + \left[\begin{array}{c c} 0 & B \\ \hline C & 0 \end{array} \right]$ $\mathfrak{g} = \mathfrak{h} + \mathfrak{p}$	$Z = \left[\begin{array}{c c} A & B \\ \hline C & D \end{array} \right] = \left[\begin{array}{c c} A & 0 \\ \hline 0 & D \end{array} \right] + \left[\begin{array}{c c} 0 & B \\ \hline C & 0 \end{array} \right]$ $\mathfrak{g} = \mathfrak{h} + \mathfrak{p}$
gilmore-lie-groups_ E0299	97	0.803	$R(A) = \left[\begin{array}{cc c} \theta & \theta & -b \\ \hline \theta & \theta & a-c \end{array} \right] \begin{array}{l} X_a \\ X_c \\ X_b \end{array}$	$R(A) = \left[\begin{array}{cc c} 0 & 0 & -b \\ \hline 0 & 0 & +b \\ 0 & 0 & a-c \end{array} \right] \begin{array}{l} X_a \\ X_c \\ X_b \end{array}$	$R(A) = \left[\begin{array}{cc c} 0 & 0 & -b \\ \hline 0 & 0 & +b \\ 0 & 0 & a-c \end{array} \right] \begin{array}{l} X_a \\ X_c \\ X_b \end{array}$
gilmore-lie-groups_ E0346	107	0.803	$\begin{array}{cccc} a_1^{\dagger} a_1 + a_2^{\dagger} a_2 & a_1^{\dagger} a_1 - a_2^{\dagger} a_2 & a_1^{\dagger} a_2 & a_2^{\dagger} a_1 \\ \downarrow & \downarrow & \downarrow & \downarrow \\ \left[\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array} \right] & \left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array} \right] & \left[\begin{array}{cc} 0 & 1 \\ 0 & 0 \end{array} \right] & \left[\begin{array}{cc} 0 & 0 \\ 1 & 0 \end{array} \right] \end{array}$	$\begin{array}{cccc} a_1^{\dagger} a_1 + a_2^{\dagger} a_2 & a_1^{\dagger} a_1 - a_2^{\dagger} a_2 & a_1^{\dagger} a_2 & a_2^{\dagger} a_1 \\ \downarrow & \downarrow & \downarrow & \downarrow \\ \left[\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array} \right] & \left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array} \right] & \left[\begin{array}{cc} 0 & 1 \\ 0 & 0 \end{array} \right] & \left[\begin{array}{cc} 0 & 0 \\ 1 & 0 \end{array} \right] \end{array}$	$\begin{array}{cccc} a_1^{\dagger} a_1 + a_2^{\dagger} a_2 & a_1^{\dagger} a_1 - a_2^{\dagger} a_2 & a_1^{\dagger} a_2 & a_2^{\dagger} a_1 \\ \downarrow & \downarrow & \downarrow & \downarrow \\ \left[\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array} \right] & \left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array} \right] & \left[\begin{array}{cc} 0 & 1 \\ 0 & 0 \end{array} \right] & \left[\begin{array}{cc} 0 & 0 \\ 1 & 0 \end{array} \right] \end{array}$
gilmore-lie-groups_ E0502 (9.11)	156	0.812	$\begin{aligned} \left\ \operatorname{Reg}(Y) - \lambda I_3 \right\ &= (-\lambda)^3 + (-\lambda)(-b_1^2 - b_2^2 + b_3^2) = 0 \\ &= (-\lambda)(\lambda^2 + \phi_2(\mathbf{b})) \\ \phi_2(\mathbf{b}) &= -b_1^2 - b_2^2 + b_3^2 \end{aligned}$	$\ \operatorname{Reg}(Y) - \lambda I_3 \ = (-\lambda)^3 + (-\lambda)(-b_1^2 - b_2^2 + b_3^2) = 0$ $= (-\lambda)(\lambda^2 + \phi_2(\mathbf{b}))$ $\phi_2(\mathbf{b}) = -b_1^2 - b_2^2 + b_3^2$	$\ \operatorname{Reg}(Y) - \lambda I_3 \ = (-\lambda)^3 + (-\lambda)(-b_1^2 - b_2^2 + b_3^2) = 0$ $= (-\lambda)(\lambda^2 + \phi_2(\mathbf{b}))$ $\phi_2(\mathbf{b}) = -b_1^2 - b_2^2 + b_3^2$
gilmore-lie-groups_ E0266 (5.4)	89	0.815	$\begin{array}{l} m_{ij} = 0 \quad p+1 \leq i \leq p+q \\ \quad \quad \quad 1 \leq j \leq p+q \end{array}$	$m_{ij} = 0 \quad p+1 \leq i \leq p+q$ $\quad \quad \quad 1 \leq j \leq p+q$	$m_{ij} = 0 \quad p+1 \leq i \leq p+q$ $\quad \quad \quad 1 \leq j \leq p+q$
gilmore-lie-groups_ E0388 (7.24)	123	0.817	$(x, y) \rightarrow \left[\begin{array}{cc} x & y \\ 0 & 1 \end{array} \right] \quad x > 0$	$(x, y) \rightarrow \left[\begin{array}{cc} x & y \\ 0 & 1 \end{array} \right] \quad x > 0$	$(x, y) \rightarrow \left[\begin{array}{cc} x & y \\ 0 & 1 \end{array} \right] \quad x > 0$
gilmore-lie-groups_ E0581 (11.1)	186	0.817	$\sum_{i=1}^{\text{rank}} h^i H_i + \sum_{\alpha \neq 0} e^{\alpha} E_{\alpha}$	$\sum_{i=1}^{\text{rank}} h^i H_i + \sum_{\alpha \neq 0} e^{\alpha} E_{\alpha}$	$\sum_{i=1}^{\text{rank}} h^i H_i + \sum_{\alpha \neq 0} e^{\alpha} E_{\alpha}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0022 (1.18)	24	■ 0.819	$\begin{array}{l rrc} & I & (12) & \text{Basis functions} \\ \hline \Gamma^1 & 1 & 1 & u_1 = r_1 + r_2 \\ \Gamma^2 & 1 & -1 & u_2 = r_1 - r_2 \end{array}$	$\begin{array}{l rrc} & I & (12) & \text{Basis functions} \\ \hline \Gamma^1 & 1 & 1 & u_1 = r_1 + r_2 \\ \Gamma^2 & 1 & -1 & u_2 = r_1 - r_2 \end{array}$	$\begin{array}{l rrc} & I & (12) & \text{Basis functions} \\ \hline \Gamma^1 & 1 & 1 & u_1 = r_1 + r_2 \\ \Gamma^2 & 1 & -1 & u_2 = r_1 - r_2 \end{array}$
gilmore-lie-groups_ EQ0208 (4.18)	75	■ 0.820	$\begin{aligned} & \left\{ \left[X_a, X_b \right] \right\} = 2 X_b \\ & \left\{ \left[X_a, X_c \right] \right\} = -2 X_c \\ & \left\{ \left[X_b, X_c \right] \right\} = X_a \end{aligned}$	$\begin{aligned} [X_a, X_b] &= 2X_b \\ [X_a, X_c] &= -2X_c \\ [X_b, X_c] &= X_a \end{aligned}$	$\begin{aligned} [X_a, X_b] &= 2X_b \\ [X_a, X_c] &= -2X_c \\ [X_b, X_c] &= X_a \end{aligned}$
gilmore-lie-groups_ EQ0334 (6.19)	105	■ 0.820	$\begin{gathered} f(p) \\\downarrow \\ f_S[x(p)] \stackrel{\perp}{=} f_{S'}[x'(p)] \end{gathered}$	$\begin{gathered} f(p) \\\downarrow \\ f_S[x(p)] \stackrel{\perp}{=} f_{S'}[x'(p)] \end{gathered}$	$\begin{array}{c} f(p) \\ \downarrow \quad \downarrow \\ f_S[x(p)] = f_{S'}[x'(p)] \end{array}$
gilmore-lie-groups_ EQ0660	214	■ 0.824	$\begin{aligned} & ds^2 = \frac{(dr \, d\theta)}{1+r^2} \left[\begin{array}{cc} \frac{1}{1+r^2} & 0 \\ 0 & r^2 \end{array} \right] \begin{pmatrix} dr \\ d\theta \end{pmatrix} \\ & d\mu = \frac{r \, dr \, d\theta}{\sqrt{1+r^2}} \end{aligned}$	$\begin{aligned} ds^2 &= \frac{(drd\theta) \left[\begin{array}{cc} \frac{1}{1+r^2} & 0 \\ 0 & r^2 \end{array} \right] \begin{pmatrix} dr \\ d\theta \end{pmatrix}}{1+r^2} \\ d\mu &= \frac{rdrd\theta}{\sqrt{1+r^2}} \end{aligned}$	$\begin{aligned} ds^2 &= \frac{(dr \, d\theta) \left[\begin{array}{cc} \frac{1}{1+r^2} & 0 \\ 0 & r^2 \end{array} \right] \begin{pmatrix} dr \\ d\theta \end{pmatrix}}{1+r^2} \\ d\mu &= \frac{r \, dr \, d\theta}{\sqrt{1+r^2}} \end{aligned}$
gilmore-lie-groups_ EQ0257 (4.61a)	86	■ 0.825	$\left[\begin{array}{ll} \alpha_R + i\alpha_I & \beta_R + i\beta_I \\ \beta_R - i\beta_I & \alpha_R - i\alpha_I \end{array} \right]$	$\left[\begin{array}{cc} \alpha_R + i\alpha_I & \beta_R + i\beta_I \\ \beta_R - i\beta_I & \alpha_R - i\alpha_I \end{array} \right]$	$\left[\begin{array}{cc} \alpha_R + i\alpha_I & \beta_R + i\beta_I \\ \beta_R - i\beta_I & \alpha_R - i\alpha_I \end{array} \right]$
gilmore-lie-groups_ EQ0891 (15.24)	279	■ 0.826	$ k\rangle \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right $	$ k\rangle \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right $	$ k\rangle \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right $

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0930 (15.65)	288	■ 0.826	$\begin{aligned} & \left[\frac{\partial}{\partial t} \frac{\partial}{\partial x} + \frac{\partial}{\partial y} \frac{\partial}{\partial x} + \frac{\partial}{\partial z} \frac{\partial}{\partial x} + \frac{\partial}{\partial t} \frac{\partial}{\partial y} + \frac{\partial}{\partial x} \frac{\partial}{\partial y} + \frac{\partial}{\partial z} \frac{\partial}{\partial y} + \frac{\partial}{\partial t} \frac{\partial}{\partial z} + \frac{\partial}{\partial x} \frac{\partial}{\partial z} + \frac{\partial}{\partial y} \frac{\partial}{\partial z} \right] \begin{bmatrix} B_1 + iE_1 \\ B_2 + iE_2 \\ B_3 + iE_3 \end{bmatrix} = 0 \\ & -\frac{i}{c} \frac{\partial}{\partial t} (B + iE)_1 + \partial_3 (B + iE)_2 - \partial_2 (B + iE)_3 = 0 \\ & -\partial_3 (B + iE)_1 - \frac{i}{c} \frac{\partial}{\partial t} (B + iE)_2 + \partial_1 (B + iE)_3 = 0 \\ & +\partial_2 (B + iE)_1 - \partial_1 (B + iE)_2 - \frac{i}{c} \frac{\partial}{\partial t} (B + iE)_3 = 0 \end{aligned}$	$\begin{bmatrix} -\frac{i}{c} \frac{\partial}{\partial t} & +\partial_3 & -\partial_2 \\ -\partial_3 & -\frac{i}{c} \frac{\partial}{\partial t} & +\partial_1 \\ +\partial_2 & -\partial_1 & -\frac{i}{c} \frac{\partial}{\partial t} \end{bmatrix} \begin{bmatrix} B_1 + iE_1 \\ B_2 + iE_2 \\ B_3 + iE_3 \end{bmatrix} = 0$ $-\frac{i}{c} \frac{\partial}{\partial t} (B + iE)_1 + \partial_3 (B + iE)_2 - \partial_2 (B + iE)_3 = 0$ $-\partial_3 (B + iE)_1 - \frac{i}{c} \frac{\partial}{\partial t} (B + iE)_2 + \partial_1 (B + iE)_3 = 0$ $+\partial_2 (B + iE)_1 - \partial_1 (B + iE)_2 - \frac{i}{c} \frac{\partial}{\partial t} (B + iE)_3 = 0$	$\begin{bmatrix} -\frac{i}{c} \frac{\partial}{\partial t} & +\partial_3 & -\partial_2 \\ -\partial_3 & -\frac{i}{c} \frac{\partial}{\partial t} & +\partial_1 \\ +\partial_2 & -\partial_1 & -\frac{i}{c} \frac{\partial}{\partial t} \end{bmatrix} \begin{bmatrix} B_1 + iE_1 \\ B_2 + iE_2 \\ B_3 + iE_3 \end{bmatrix} = 0$ $-\frac{i}{c} \frac{\partial}{\partial t} (B + iE)_1 + \partial_3 (B + iE)_2 - \partial_2 (B + iE)_3 = 0$ $-\partial_3 (B + iE)_1 - \frac{i}{c} \frac{\partial}{\partial t} (B + iE)_2 + \partial_1 (B + iE)_3 = 0$ $+\partial_2 (B + iE)_1 - \partial_1 (B + iE)_2 - \frac{i}{c} \frac{\partial}{\partial t} (B + iE)_3 = 0$
gilmore-lie-groups_ EQ0993 (16.36)	309	■ 0.831	$\operatorname{EXP} \left(\lambda \left(x \frac{\partial}{\partial x} - y \frac{\partial}{\partial y} - 2p \frac{\partial}{\partial p} \right) \right) \begin{bmatrix} x \\ y \\ p \end{bmatrix} = \begin{bmatrix} e^{\lambda x} x \\ e^{-\lambda y} y \\ e^{-2\lambda p} p \end{bmatrix}$	$\operatorname{EXP} \left(\lambda \left\{ x \frac{\partial}{\partial x} - y \frac{\partial}{\partial y} - 2p \frac{\partial}{\partial p} \right\} \right) \begin{bmatrix} x \\ y \\ p \end{bmatrix} = \begin{bmatrix} e^{\lambda x} x \\ e^{-\lambda y} y \\ e^{-2\lambda p} p \end{bmatrix}$	$\operatorname{EXP} \left(\lambda \left\{ x \frac{\partial}{\partial x} - y \frac{\partial}{\partial y} - 2p \frac{\partial}{\partial p} \right\} \right) \begin{bmatrix} x \\ y \\ p \end{bmatrix} = \begin{bmatrix} e^{\lambda x} x \\ e^{-\lambda y} y \\ e^{-2\lambda p} p \end{bmatrix}$
gilmore-lie-groups_ EQ0576 (10.18)	182	■ 0.837	$\begin{gathered} p \text{ arbitrary, } q=1, B_l, C_l \\ \quad \quad \quad p=2, q=2, F_4 \end{gathered}$	$p \text{ arbitrary, } q=1, B_l, C_l \quad l=p+1$ $p=2, q=2, F_4$	$p \text{ arbitrary, } q=1, B_l, C_l \quad l=p+1$ $p=2, q=2, F_4$
gilmore-lie-groups_ EQ0277 (5.16)	92	■ 0.839	$\begin{aligned} & \left[\begin{array}{cc} A^t & C^t \\ B^t & D \end{array} \right] \left[\begin{array}{cc} I_2 & 0 \\ 0 & -1 \end{array} \right] + \left[\begin{array}{cc} I_2 & 0 \\ 0 & -1 \end{array} \right] \left[\begin{array}{cc} A & B \\ C & D \end{array} \right] = 0 \\ & \left[\begin{array}{cc} A^t & C^t \\ B^t & D \end{array} \right] \left[\begin{array}{cc} I_2 & 0 \\ 0 & -1 \end{array} \right] + \left[\begin{array}{cc} I_2 & 0 \\ 0 & -1 \end{array} \right] \left[\begin{array}{cc} A & B \\ C & D \end{array} \right] = 0 \end{aligned}$	$\begin{bmatrix} A^t & C^t \\ B^t & D \end{bmatrix} \begin{bmatrix} I_2 & 0 \\ 0 & -1 \end{bmatrix} + \begin{bmatrix} I_2 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} = 0$	$\begin{bmatrix} A^t & C^t \\ B^t & D \end{bmatrix} \begin{bmatrix} I_2 & 0 \\ 0 & -1 \end{bmatrix} + \begin{bmatrix} I_2 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} = 0$
gilmore-lie-groups_ EQ0629 (12.2)	205	■ 0.847	$\begin{aligned} & G/H = \operatorname{EXP}(p) \quad ds^2 = g_{\mu, \nu} dx^\mu dx^\nu < 0 \\ & G'/H = \operatorname{EXP}(ip) \quad ds^2 = g_{\mu, \nu} dx^\mu dx^\nu > 0 \end{aligned}$	$G/H = \operatorname{EXP}(p) \quad ds^2 = g_{\mu, \nu} dx^\mu dx^\nu < 0$ $G'/H = \operatorname{EXP}(ip) \quad ds^2 = g_{\mu, \nu} dx^\mu dx^\nu > 0$	$G/H = \operatorname{EXP}(p) \quad ds^2 = g_{\mu, \nu} dx^\mu dx^\nu < 0$ $G'/H = \operatorname{EXP}(ip) \quad ds^2 = g_{\mu, \nu} dx^\mu dx^\nu > 0$
gilmore-lie-groups_ EQ0217 (4.27)	76	■ 0.848	$Z \stackrel{\text{regular}}{\longrightarrow} R(Z)$	$Z \stackrel{\text{regular}}{\longrightarrow} R(Z)$	$Z \stackrel{\text{regular}}{\longrightarrow} R(Z)$
gilmore-lie-groups_ EQ0543	169	■ 0.848	$\mathcal{C}^2(\mathbf{M}) = (\mathbf{M} + \mathbf{R}) \cdot (\mathbf{M} + \mathbf{R}) - (\mathbf{R}) \cdot (\mathbf{R}) = \mathbf{M} \cdot \mathbf{M} + \mathbf{M} \cdot 2\mathbf{R}$	$\mathcal{C}^2(\mathbf{M}) = (\mathbf{M} + \mathbf{R}) \cdot (\mathbf{M} + \mathbf{R}) - (\mathbf{R}) \cdot (\mathbf{R}) = \mathbf{M} \cdot \mathbf{M} + \mathbf{M} \cdot 2\mathbf{R}$	$\mathcal{C}^2(\mathbf{M}) = (\mathbf{M} + \mathbf{R}) \cdot (\mathbf{M} + \mathbf{R}) - (\mathbf{R}) \cdot (\mathbf{R}) = \mathbf{M} \cdot \mathbf{M} + \mathbf{M} \cdot 2\mathbf{R}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0693 (13.11)	221	■ 0.854	$\begin{aligned} & \text{\texttt{\textbackslash begin{aligned}}} & \text{\texttt{d_1}} = +R \text{\texttt{\textbackslash theta_2}} \text{\texttt{\textbackslash\ \&}} \text{\texttt{d_2}} = -R \\ & \text{\texttt{\textbackslash theta_1}} \text{\texttt{\textbackslash end{aligned}}}\end{aligned}$	$\begin{aligned} d_1 &= +R\theta_2 \\ d_2 &= -R\theta_1 \end{aligned}$	$\begin{aligned} d_1 &= +R\theta_2 \\ d_2 &= -R\theta_1 \end{aligned}$
gilmore-lie-groups_ EQ0009 (1.6)	19	■ 0.855	$\begin{array}{ccccccc} \text{\texttt{\textbackslash begin{array}{cccccc}}} & \text{\texttt{g_i}} & \text{\texttt{\& \textbackslash cdot \& g_j}} & \text{\texttt{\& = \& g_i}} \\ & \text{\texttt{\& \textbackslash cdot g_j}} & \text{\texttt{\& \textbackslash\ \downarrow \& \textbackslash downarrow \& \textbackslash downarrow \&}} \\ & \text{\texttt{\& \textbackslash downarrow \textbackslash \Gamma\textbackslash left(g_i\textbackslash right)}} & \text{\texttt{\& \textbackslash times \&}} \\ & \text{\texttt{\Gamma\textbackslash left(g_j\textbackslash right)}} & \text{\texttt{\& =\Gamma\textbackslash left(g_i} \text{\texttt{\& \textbackslash cdot g_j}} \\ & \text{\texttt{\textbackslash right)}}} & \text{\texttt{\textbackslash end{array}}}\end{array}$	$\begin{array}{ccccccc} g_i & \cdot & g_j & = & g_i \cdot g_j \\ \downarrow & \downarrow & \downarrow & & \downarrow \\ \Gamma(g_i) & \times & \Gamma(g_j) & = & \Gamma(g_i \cdot g_j) \end{array}$	$\begin{array}{ccccccc} g_i & \cdot & g_j & = & g_i \cdot g_j \\ \downarrow & \downarrow & \downarrow & & \downarrow \\ \Gamma(g_i) & \times & \Gamma(g_j) & = & \Gamma(g_i \cdot g_j) \end{array}$
gilmore-lie-groups_ EQ0587 (11.8)	189	■ 0.860	$\begin{gathered} H_1 \quad \frac{1}{\sqrt{2}}(E_{+1} + E_{-1}) \quad \frac{1}{\sqrt{2}}(E_{+1} - E_{-1}) \\ \left(E_{+1} + E_{-1}\right) \quad \frac{1}{\sqrt{2}}(E_{+1} + E_{-1}) \\ \left(E_{+1} - E_{-1}\right) \quad \frac{1}{\sqrt{2}}(E_{+1} - E_{-1}) \\ \left(h_r + i h_i\right) \left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right] + (a_r + i a_i) \left[\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right] + (b_r + i b_i) \left[\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right] \\ (h_r, a_r, b_r) \longrightarrow \left[\begin{array}{cc} h_r & a_r + b_r \\ a_r - b_r & -h_r \end{array}\right] \quad \mathfrak{sl}(2; \mathbb{R}) \\ (i h_i, i a_i, b_r) \longrightarrow i \left[\begin{array}{cc} h_i & a_i - i b_r \\ a_i + i b_r & -h_i \end{array}\right] \quad \mathfrak{su}(2) \end{gathered}$	$\begin{aligned} H_1 \quad \frac{1}{\sqrt{2}}(E_{+1} + E_{-1}) \quad \frac{1}{\sqrt{2}}(E_{+1} - E_{-1}) \\ (h_r + i h_i) \left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right] + (a_r + i a_i) \left[\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right] + (b_r + i b_i) \left[\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right] \\ (h_r, a_r, b_r) \longrightarrow \left[\begin{array}{cc} h_r & a_r + b_r \\ a_r - b_r & -h_r \end{array}\right] \quad \mathfrak{sl}(2; \mathbb{R}) \\ (i h_i, i a_i, b_r) \longrightarrow i \left[\begin{array}{cc} h_i & a_i - i b_r \\ a_i + i b_r & -h_i \end{array}\right] \quad \mathfrak{su}(2) \end{aligned}$	$\begin{aligned} H_1 \quad \frac{1}{\sqrt{2}}(E_{+1} + E_{-1}) \quad \frac{1}{\sqrt{2}}(E_{+1} - E_{-1}) \\ (h_r + i h_i) \left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right] + (a_r + i a_i) \left[\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right] + (b_r + i b_i) \left[\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right] \\ (h_r, a_r, b_r) \longrightarrow \left[\begin{array}{cc} h_r & a_r + b_r \\ a_r - b_r & -h_r \end{array}\right] \quad \mathfrak{sl}(2; \mathbb{R}) \\ (i h_i, i a_i, b_r) \longrightarrow i \left[\begin{array}{cc} h_i & a_i - i b_r \\ a_i + i b_r & -h_i \end{array}\right] \quad \mathfrak{su}(2) \end{aligned}$
gilmore-lie-groups_ EQ0178	66	■ 0.861	$\begin{aligned} & \text{\texttt{\textbackslash left[\textbackslash begin{array}{c} A_L \text{\texttt{\textbackslash\ \&}} B_L \text{\texttt{\textbackslash end{array}}} \\ & \text{\texttt{\textbackslash right}}] = T \text{\texttt{\textbackslash left[\textbackslash begin{array}{c} A_R \text{\texttt{\textbackslash\ \&}} B_R \text{\texttt{\textbackslash end{array}}} \text{\texttt{\textbackslash right}}} \quad \text{\texttt{\textbackslash quad}} \text{\texttt{\textbackslash left[\textbackslash begin{array}{c} A_R \text{\texttt{\textbackslash\ \&}} B_L \text{\texttt{\textbackslash end{array}}} \text{\texttt{\textbackslash right}}} = S \text{\texttt{\textbackslash left[\textbackslash begin{array}{c} A_L \text{\texttt{\textbackslash\ \&}} B_R \text{\texttt{\textbackslash end{array}}} \text{\texttt{\textbackslash right}}}\end{aligned}$	$\left[\begin{array}{c} A_L \\ B_L \end{array}\right] = T \left[\begin{array}{c} A_R \\ B_R \end{array}\right] \quad \left[\begin{array}{c} A_R \\ B_L \end{array}\right] = S \left[\begin{array}{c} A_L \\ B_R \end{array}\right]$	$\left[\begin{array}{c} A_L \\ B_L \end{array}\right] = T \left[\begin{array}{c} A_R \\ B_R \end{array}\right] \quad \left[\begin{array}{c} A_R \\ B_L \end{array}\right] = S \left[\begin{array}{c} A_L \\ B_R \end{array}\right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00645 (12.14)	210	0.861	$\begin{aligned} & \Gamma_{\mu\nu}^{\sigma} \rightarrow \frac{1}{2} \left(\frac{\partial g_{\mu\sigma}}{\partial x^{\nu}} + \frac{\partial g_{\nu\sigma}}{\partial x^{\mu}} - \frac{\partial g_{\mu\nu}}{\partial x^{\sigma}} \right) \\ & = \frac{1}{2} \left\{ \delta_{\nu\mu} x^{\sigma} + \delta_{\mu\nu} x^{\sigma} - \delta_{\sigma\mu} x^{\nu} \right\} \\ & \rightarrow \delta_{\mu\nu} x^{\sigma} \quad (\rightarrow 0 \text{ at origin}) \end{aligned}$	$\Gamma_{\mu\nu}^{\sigma} \rightarrow \frac{1}{2} \left(\frac{\partial g_{\mu\sigma}}{\partial x^{\nu}} + \frac{\partial g_{\nu\sigma}}{\partial x^{\mu}} - \frac{\partial g_{\mu\nu}}{\partial x^{\sigma}} \right)$ $= \frac{1}{2} \left\{ \delta_{\nu\mu} x^{\sigma} + \delta_{\mu\nu} x^{\sigma} - \delta_{\sigma\mu} x^{\nu} \right\}$ $\rightarrow \delta_{\mu\nu} x^{\sigma} \quad (\rightarrow 0 \text{ at origin})$	$\Gamma_{\mu\nu}^{\sigma} \rightarrow \frac{1}{2} \left(\frac{\partial g_{\mu\sigma}}{\partial x^{\nu}} + \frac{\partial g_{\nu\sigma}}{\partial x^{\mu}} - \frac{\partial g_{\mu\nu}}{\partial x^{\sigma}} \right)$ $= \frac{1}{2} \left\{ \delta_{\nu\mu} x^{\sigma} + \delta_{\mu\nu} x^{\sigma} - \delta_{\sigma\mu} x^{\nu} \right\}$ $\rightarrow \delta_{\mu\nu} x^{\sigma} \quad (\rightarrow 0 \text{ at origin})$
gilmore-lie-groups_ E00125	53	0.867	$\begin{array}{ll} A_{i,j} = \delta_{i,j} & 1 \leq i, j \leq p \\ A_{\alpha,\beta} = \delta_{\alpha,\beta} & p+1 \leq \alpha, \beta \leq p+q \\ A_{\alpha,j} = 0 \\ A_{i,\beta} = \text{arbitrary} \end{array}$	$\begin{array}{ll} A_{i,j} = \delta_{i,j} & 1 \leq i, j \leq p \\ A_{\alpha,\beta} = \delta_{\alpha,\beta} & p+1 \leq \alpha, \beta \leq p+q \\ A_{\alpha,j} = 0 \\ A_{i,\beta} = \text{arbitrary} \end{array}$	$\begin{array}{ll} A_{i,j} = \delta_{i,j} & 1 \leq i, j \leq p \\ A_{\alpha,\beta} = \delta_{\alpha,\beta} & p+1 \leq \alpha, \beta \leq p+q \\ A_{\alpha,j} = 0 \\ A_{i,\beta} = \text{arbitrary} \end{array}$
gilmore-lie-groups_ E00290 (5.29)	94	0.869	$\mathfrak{su}(2) = \mathfrak{u}(2) \cap \mathfrak{sl}(2; \mathbb{C}) = \frac{i}{2} \left[\begin{array}{cc} x_3 & x_1 - ix_2 \\ x_1 + ix_2 & -x_3 \end{array} \right]$	$\mathfrak{su}(2) = \mathfrak{u}(2) \cap \mathfrak{sl}(2; \mathbb{C}) = \frac{i}{2} \left[\begin{array}{cc} x_3 & x_1 - ix_2 \\ x_1 + ix_2 & -x_3 \end{array} \right]$	$\mathfrak{su}(2) = \mathfrak{u}(2) \cap \mathfrak{sl}(2; \mathbb{C}) = \frac{i}{2} \left[\begin{array}{cc} x_3 & x_1 - ix_2 \\ x_1 + ix_2 & -x_3 \end{array} \right]$
gilmore-lie-groups_ E00197 (4.10)	73	0.872	$\text{Lie group} \stackrel{\ln}{\rightleftharpoons} \text{Lie algebra}$	$\text{Lie group} \stackrel{\ln}{\rightleftharpoons} \text{Lie algebra}$	$\text{Lie group} \stackrel{\ln}{\rightleftharpoons} \text{Lie algebra}$
gilmore-lie-groups_ E00790 (14.32)	247	0.874	$\begin{array}{l} \mathbf{L} = \mathbf{r} \times \mathbf{p} \quad \mathbf{L} \cdot \mathbf{L} = L^2 \\ \mathbf{M} = \frac{\mathbf{p} \times \mathbf{L}}{m} - K \frac{\mathbf{r}}{r} \quad \mathbf{M} \cdot \mathbf{M} = M^2 = \frac{2E}{m} L^2 + K^2 \\ \mathbf{W} = \frac{\mathbf{p}}{m} L^2 - K \frac{\mathbf{L} \times \mathbf{r}}{r} \quad \mathbf{W} \cdot \mathbf{W} = L^2 M^2 \end{array}$	$\begin{array}{l} \mathbf{L} = \mathbf{r} \times \mathbf{p} \quad \mathbf{L} \cdot \mathbf{L} = L^2 \\ \mathbf{M} = \frac{\mathbf{p} \times \mathbf{L}}{m} - K \frac{\mathbf{r}}{r} \quad \mathbf{M} \cdot \mathbf{M} = M^2 = \frac{2E}{m} L^2 + K^2 \\ \mathbf{W} = \frac{\mathbf{p}}{m} L^2 - K \frac{\mathbf{L} \times \mathbf{r}}{r} \quad \mathbf{W} \cdot \mathbf{W} = L^2 M^2 \end{array}$	$\begin{array}{l} \mathbf{L} = \mathbf{r} \times \mathbf{p} \quad \mathbf{L} \cdot \mathbf{L} = L^2 \\ \mathbf{M} = \frac{\mathbf{p} \times \mathbf{L}}{m} - K \frac{\mathbf{r}}{r} \quad \mathbf{M} \cdot \mathbf{M} = M^2 = \frac{2E}{m} L^2 + K^2 \\ \mathbf{W} = \frac{\mathbf{p}}{m} L^2 - K \frac{\mathbf{L} \times \mathbf{r}}{r} \quad \mathbf{W} \cdot \mathbf{W} = L^2 M^2 \end{array}$
gilmore-lie-groups_ E00120 (3.8)	51	0.874	$m_{ij} - \delta_{ij} = 0 \quad p+1 \leq i \leq p+q \quad p+1 \leq j \leq p+q$	$m_{ij} - \delta_{ij} = 0 \quad p+1 \leq i \leq p+q \quad p+1 \leq j \leq p+q$	$m_{ij} - \delta_{ij} = 0 \quad p+1 \leq i \leq p+q \quad p+1 \leq j \leq p+q$
gilmore-lie-groups_ E00237 (4.47)	81	0.874	$g(x) = M^{-1}(x)^t g(\text{Id}) M^{-1}(x)$	$g(x) = M^{-1}(x)^t g(\text{Id}) M^{-1}(x)$	$g(x) = M^{-1}(x)^t g(\text{Id}) M^{-1}(x)$
gilmore-lie-groups_ E00389 (7.25)	123	0.874	$(w, z) = \text{EXP} \left[\begin{array}{cc} w & z \\ 0 & 1 \end{array} \right] = \left[\begin{array}{cc} e^w & (e^w - 1)z/w \\ 0 & 1 \end{array} \right]$	$(w, z) = \text{EXP} \left[\begin{array}{cc} w & z \\ 0 & 1 \end{array} \right] = \left[\begin{array}{cc} e^w & (e^w - 1)z/w \\ 0 & 1 \end{array} \right]$	$(w, z) = \text{EXP} \left[\begin{array}{cc} w & z \\ 0 & 1 \end{array} \right] = \left[\begin{array}{cc} e^w & (e^w - 1)z/w \\ 0 & 1 \end{array} \right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E09973 (16.16)	303	■ 0.874	$\begin{array}{r} \left(\xi(x,y)\frac{\partial}{\partial x}+\eta(x,y)\frac{\partial}{\partial y}\right)R(x,y)=0 \\ \left(\xi(x,y)\frac{\partial}{\partial x}+\eta(x,y)\frac{\partial}{\partial y}+\zeta(x,y)\frac{\partial}{\partial p}\right)T(x,y,p)=0 \end{array}$	$\begin{array}{r} \left(\xi(x,y)\frac{\partial}{\partial x}+\eta(x,y)\frac{\partial}{\partial y}\right)R(x,y)=0 \\ \left(\xi(x,y)\frac{\partial}{\partial x}+\eta(x,y)\frac{\partial}{\partial y}+\zeta(x,y)\frac{\partial}{\partial p}\right)T(x,y,p)=0 \end{array}$	$\begin{array}{r} \left(\xi(x,y)\frac{\partial}{\partial x}+\eta(x,y)\frac{\partial}{\partial y}\right)R(x,y)=0 \\ \left(\xi(x,y)\frac{\partial}{\partial x}+\eta(x,y)\frac{\partial}{\partial y}+\zeta(x,y)\frac{\partial}{\partial p}\right)T(x,y,p)=0 \end{array}$
gilmore-lie-groups_ E09471 (8.2)	143	■ 0.876	$S(Z)_r^s=\left(A^{-1}\right)_r^iR(Z)_i^jA_j^s$	$S(Z)_r^s=\left(A^{-1}\right)_r^iR(Z)_i^jA_j^s$	$S(Z)_r^s=\left(A^{-1}\right)_r^iR(Z)_i^jA_j^s$
gilmore-lie-groups_ E08869 (15.2)	275	■ 0.878	$\left[\begin{array}{c} x \\ y \\ z \\ ict \end{array}\right]=\left[\begin{array}{c} \Lambda \end{array}\right]\left[\begin{array}{c} x \\ y \\ z \\ ict \end{array}\right]'$	$\left[\begin{array}{c} x \\ y \\ z \\ ict \end{array}\right]=\left[\begin{array}{c} \Lambda \end{array}\right]\left[\begin{array}{c} x \\ y \\ z \\ ict \end{array}\right]'$	$\left[\begin{array}{c} x \\ y \\ z \\ ict \end{array}\right]=\left[\begin{array}{c} \Lambda \end{array}\right]\left[\begin{array}{c} x \\ y \\ z \\ ict \end{array}\right]'$
gilmore-lie-groups_ E09385 (7.22)	121	■ 0.887	$\begin{gathered} \mathfrak{su}(2)=\frac{i}{2}\left[\begin{array}{cc} 0 & b_1-i b_2 \\ b_1+i b_2 & 0 \end{array}\right]+\frac{i}{2}\left[\begin{array}{cc} b_3 & 0 \\ 0 & -b_3 \end{array}\right] \\ \text{EXP}\downarrow\text{EXP} \\ \left[\begin{array}{cc} z' & i\left(x'-i y'\right) \\ i\left(x'+i y'\right) & z' \end{array}\right] \times\left[\begin{array}{cc} e^{i b_3 / 2} & 0 \\ 0 & e^{-i b_3 / 2} \end{array}\right] \end{gathered}$	$\begin{array}{c} \mathfrak{su}(2)=\frac{i}{2}\left[\begin{array}{cc} 0 & b_1-i b_2 \\ b_1+i b_2 & 0 \end{array}\right]+\frac{i}{2}\left[\begin{array}{cc} b_3 & 0 \\ 0 & -b_3 \end{array}\right] \\ \text{EXP}\downarrow\text{EXP} \\ \left[\begin{array}{cc} z' & i\left(x'-i y'\right) \\ i\left(x'+i y'\right) & z' \end{array}\right] \times\left[\begin{array}{cc} e^{i b_3 / 2} & 0 \\ 0 & e^{-i b_3 / 2} \end{array}\right] \end{array}$	$\begin{array}{c} \mathfrak{su}(2)=\frac{i}{2}\left[\begin{array}{cc} 0 & b_1-i b_2 \\ b_1+i b_2 & 0 \end{array}\right]+\frac{i}{2}\left[\begin{array}{cc} b_3 & 0 \\ 0 & -b_3 \end{array}\right] \\ \text{EXP}\downarrow\text{EXP} \\ \left[\begin{array}{cc} z' & i\left(x'-i y'\right) \\ i\left(x'+i y'\right) & z' \end{array}\right] \times\left[\begin{array}{cc} e^{i b_3 / 2} & 0 \\ 0 & e^{-i b_3 / 2} \end{array}\right] \end{array}$
gilmore-lie-groups_ E09185	68	■ 0.888	$\left[\begin{array}{c} o_i \\ o_e \end{array}\right]=\left[\begin{array}{cc} A & B \\ C & D \end{array}\right]\left[\begin{array}{c} i_i \\ i_e \end{array}\right] \quad\left[i_i\right]=\Gamma\left[o_i\right]$	$\left[\begin{array}{c} o_i \\ o_e \end{array}\right]=\left[\begin{array}{cc} A & B \\ C & D \end{array}\right]\left[\begin{array}{c} i_i \\ i_e \end{array}\right] \quad\left[i_i\right]=\Gamma\left[o_i\right]$	$\left[\begin{array}{c} o_i \\ o_e \end{array}\right]=\left[\begin{array}{cc} A & B \\ C & D \end{array}\right]\left[\begin{array}{c} i_i \\ i_e \end{array}\right] \quad\left[i_i\right]=\Gamma\left[o_i\right]$
gilmore-lie-groups_ E09527 (9.36)	163	■ 0.890	$\sum_{\alpha\neq 0}\alpha_i\alpha_j=\delta_{ij} \quad\text{or}\quad \sum_{\alpha\neq 0}\alpha\cdot\alpha=\text{rank}=l$	$\sum_{\alpha\neq 0}\alpha_i\alpha_j=\delta_{ij} \quad\text{or}\quad \sum_{\alpha\neq 0}\alpha\cdot\alpha=\text{rank}=l$	$\sum_{\alpha\neq 0}\alpha_i\alpha_j=\delta_{ij} \quad\text{or}\quad \sum_{\alpha\neq 0}\alpha\cdot\alpha=\text{rank}=l$
gilmore-lie-groups_ E09953	295	■ 0.890	$\begin{aligned} x_1&=r\cos\theta_2 \\ x_2&=r\sin\theta_2\cos\theta_3 \\ &\vdots \\ x_{n-1}&=r\sin\theta_2\sin\theta_3\cdots\sin\theta_{n-1}\cos\theta_n \\ x_n&=r\sin\theta_2\sin\theta_3\cdots\sin\theta_{n-1}\sin\theta_n \end{aligned}$	$\begin{array}{c} x_1=r\cos\theta_2 \\ x_2=r\sin\theta_2\cos\theta_3 \\ \vdots \\ x_{n-1}=r\sin\theta_2\sin\theta_3\cdots\sin\theta_{n-1}\cos\theta_n \\ x_n=r\sin\theta_2\sin\theta_3\cdots\sin\theta_{n-1}\sin\theta_n \end{array}$	$\begin{array}{c} x_1=r\cos\theta_2 \\ x_2=r\sin\theta_2\cos\theta_3 \\ \vdots \\ x_{n-1}=r\sin\theta_2\sin\theta_3\cdots\sin\theta_{n-1}\cos\theta_n \\ x_n=r\sin\theta_2\sin\theta_3\cdots\sin\theta_{n-1}\sin\theta_n \end{array}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0731 (13.50)	229	0.890	$\begin{aligned} & \lim_{c \rightarrow 0} l^{-1/4} P_m^l(u = x/\sqrt{l}) \\ &= \lim_{c \rightarrow 0} (-)^n \sqrt{\frac{(2l)! l^{1/2}}{2^{(2l)} l! l!}} \sqrt{\frac{1}{2^n n! (2lc^2)^n}} \\ &\quad \times [1 - 2c^2 x^2]^{(-1/2c^2)/2} \frac{d^n}{dx^n} [1 - 2c^2 x^2]^{1/2c^2} \end{aligned}$	$\lim_{c \rightarrow 0} l^{-1/4} P_m^l(u = x/\sqrt{l})$ $= \lim_{c \rightarrow 0} (-)^n \sqrt{\frac{(2l)! l^{1/2}}{2^{(2l)} l! l!}} \sqrt{\frac{1}{2^n n! (2lc^2)^n}}$ $\times [1 - 2c^2 x^2]^{(-1/2c^2)/2} \frac{d^n}{dx^n} [1 - 2c^2 x^2]^{1/2c^2}$	$\lim_{c \rightarrow 0} l^{-1/4} P_m^l(u = x/\sqrt{l})$ $= \lim_{c \rightarrow 0} (-)^n \sqrt{\frac{(2l)! l^{1/2}}{2^{(2l)} l! l!}} \sqrt{\frac{1}{2^n n! (2lc^2)^n}}$ $\times [1 - 2c^2 x^2]^{(-1/2c^2)/2} \frac{d^n}{dx^n} [1 - 2c^2 x^2]^{1/2c^2}$
gilmore-lie-groups_ EQ0399 (7.35)	124	0.891	$e^{\{r \ a^{\dagger}\}} e^{\{+\frac{1}{2}\} \operatorname{I}r} I e^{la} = e^{ra^{\dagger}+la} = e^{la} e^{-\frac{1}{2} \operatorname{I}r} e^{ra^{\dagger}}$	$e^{ra^{\dagger}} e^{+\frac{1}{2} \operatorname{I}r} I e^{la} = e^{ra^{\dagger}+la} = e^{la} e^{-\frac{1}{2} \operatorname{I}r} e^{ra^{\dagger}}$	$e^{ra^{\dagger}} e^{+\frac{1}{2} \operatorname{I}r} I e^{la} = e^{ra^{\dagger}+la} = e^{la} e^{-\frac{1}{2} \operatorname{I}r} e^{ra^{\dagger}}$
gilmore-lie-groups_ EQ0664	214	0.894	$\Delta = \frac{1}{\sqrt{\ g\ }} \partial_{\mu} \sqrt{\ g\ } g^{\mu\nu} \partial_{\nu} \quad \text{where} \quad \partial_{\nu} = \partial/\partial\phi_{\mu}$	$\Delta = \frac{1}{\sqrt{\ g\ }} \partial_{\mu} \sqrt{\ g\ } g^{\mu\nu} \partial_{\nu} \quad \text{where} \quad \partial_{\nu} = \partial/\partial\phi_{\mu}$	$\Delta = \frac{1}{\sqrt{\ g\ }} \partial_{\mu} \sqrt{\ g\ } g^{\mu\nu} \partial_{\nu} \quad \text{where} \quad \partial_{\nu} = \partial/\partial\phi_{\mu}$
gilmore-lie-groups_ EQ0417 (7.53)	127	0.896	$\left\langle \begin{array}{c} j \\ j \end{array} \left J_+^r J_-^s \right \begin{array}{c} j \\ j \end{array} \right\rangle = \frac{(2j)! r!}{(2j-r)!} \delta_{rs}$	$\left\langle \begin{array}{c} j \\ j \end{array} \left J_+^r J_-^s \right \begin{array}{c} j \\ j \end{array} \right\rangle = \frac{(2j)! r!}{(2j-r)!} \delta_{rs}$	$\left\langle \begin{array}{c} j \\ j \end{array} \left J_+^r J_-^s \right \begin{array}{c} j \\ j \end{array} \right\rangle = \frac{(2j)! r!}{(2j-r)!} \delta_{rs}$
gilmore-lie-groups_ EQ0265 (5.3)	89	0.897	$m_{i\alpha} = 0 \quad \begin{array}{c} p+1 \leq i \leq p+q \\ 1 \leq \alpha \leq p \end{array}$	$m_{i\alpha} = 0 \quad \begin{array}{c} p+1 \leq i \leq p+q \\ 1 \leq \alpha \leq p \end{array}$	$m_{i\alpha} = 0 \quad \begin{array}{c} p+1 \leq i \leq p+q \\ 1 \leq \alpha \leq p \end{array}$
gilmore-lie-groups_ EQ0424 (7.60)	130	0.900	$\begin{aligned} & \frac{d}{dt} \begin{bmatrix} 1 & r^* \frac{(e^{in}-1)}{(in)} & r^* r \left(\frac{(e^{in}-1-in)}{(in)^2} \right) + id \\ 0 & e^{in} & r \frac{(e^{in}-1)}{(in)} \\ 0 & 0 & 1 \end{bmatrix} \\ &= -\frac{i}{\hbar} \begin{bmatrix} 0 & \alpha^*(t) & \delta(t) \\ 0 & \omega(t) & \alpha(t) \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & r^* \frac{(e^{in}-1)}{(in)} & r^* r \left(\frac{(e^{in}-1-in)}{(in)^2} \right) + id \\ 0 & e^{in} & r \frac{(e^{in}-1)}{(in)} \\ 0 & 0 & 1 \end{bmatrix} \end{aligned}$	$\frac{d}{dt} \begin{bmatrix} 1 & r^* \frac{(e^{in}-1)}{(in)} & r^* r \left(\frac{(e^{in}-1-in)}{(in)^2} \right) + id \\ 0 & e^{in} & r \frac{(e^{in}-1)}{(in)} \\ 0 & 0 & 1 \end{bmatrix}$ $= -\frac{i}{\hbar} \begin{bmatrix} 0 & \alpha^*(t) & \delta(t) \\ 0 & \omega(t) & \alpha(t) \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & r^* \frac{(e^{in}-1)}{(in)} & r^* r \left(\frac{(e^{in}-1-in)}{(in)^2} \right) + id \\ 0 & e^{in} & r \frac{(e^{in}-1)}{(in)} \\ 0 & 0 & 1 \end{bmatrix}$	$\frac{d}{dt} \begin{bmatrix} 1 & r^* \frac{(e^{in}-1)}{(in)} & r^* r \left(\frac{(e^{in}-1-in)}{(in)^2} \right) + id \\ 0 & e^{in} & r \frac{(e^{in}-1)}{(in)} \\ 0 & 0 & 1 \end{bmatrix}$ $= -\frac{i}{\hbar} \begin{bmatrix} 0 & \alpha^*(t) & \delta(t) \\ 0 & \omega(t) & \alpha(t) \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & r^* \frac{(e^{in}-1)}{(in)} & r^* r \left(\frac{(e^{in}-1-in)}{(in)^2} \right) + id \\ 0 & e^{in} & r \frac{(e^{in}-1)}{(in)} \\ 0 & 0 & 1 \end{bmatrix}$
gilmore-lie-groups_ EQ0391 (7.27)	123	0.900	$\begin{aligned} & x = e^w \\ & y = (e^w - 1)z/w = z \left(1 + \frac{w}{2!} + \frac{w^2}{3!} + \dots \right) \end{aligned}$	$x = e^w$ $y = (e^w - 1)z/w = z \left(1 + \frac{w}{2!} + \frac{w^2}{3!} + \dots \right)$	$x = e^w$ $y = (e^w - 1)z/w = z \left(1 + \frac{w}{2!} + \frac{w^2}{3!} + \dots \right)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0653 (12.22)	211	■ 0.900	$\begin{aligned} & T_1 \mathfrak{g}_{\pm,*} = \pm \mathfrak{g}_{\pm,*} \\ & T_2 \mathfrak{g}_{*,\pm} = \pm \mathfrak{g}_{*,\pm} \end{aligned}$	$T_1 \mathfrak{g}_{\pm,*} = \pm \mathfrak{g}_{\pm,*}$ $T_2 \mathfrak{g}_{*,\pm} = \pm \mathfrak{g}_{*,\pm}$	$T_1 \mathfrak{g}_{\pm,*} = \pm \mathfrak{g}_{\pm,*}$ $T_2 \mathfrak{g}_{*,\pm} = \pm \mathfrak{g}_{*,\pm}$
gilmore-lie-groups_ EQ0601 (11.22)	193	■ 0.901	$\begin{array}{r} \mathfrak{su}(2) = \frac{1}{2} \left[\begin{array}{cc} i a_3 & i a_1 + a_2 \\ i a_1 - a_2 & -i a_3 \end{array} \right] \rightarrow \frac{1}{2} \left[\begin{array}{cc} 0 & +a_2 \\ -a_2 & 0 \end{array} \right] + \frac{i}{2} \left[\begin{array}{cc} a_3 & a_1 \\ a_1 & -a_3 \end{array} \right] \\ \mathfrak{sl}(2; \mathbb{R}) = \frac{1}{2} \left[\begin{array}{cc} a_3 & a_1 + a_2 \\ a_1 - a_2 & -a_3 \end{array} \right] \leftarrow \frac{1}{2} \left[\begin{array}{cc} 0 & +a_2 \\ -a_2 & 0 \end{array} \right] + \frac{1}{2} \left[\begin{array}{cc} a_3 & a_1 \\ a_1 & -a_3 \end{array} \right] \end{array}$	$\mathfrak{su}(2) = \frac{1}{2} \left[\begin{array}{cc} i a_3 & i a_1 + a_2 \\ i a_1 - a_2 & -i a_3 \end{array} \right] \rightarrow \frac{1}{2} \left[\begin{array}{cc} 0 & +a_2 \\ -a_2 & 0 \end{array} \right] + \frac{i}{2} \left[\begin{array}{cc} a_3 & a_1 \\ a_1 & -a_3 \end{array} \right]$ $\mathfrak{sl}(2; \mathbb{R}) = \frac{1}{2} \left[\begin{array}{cc} a_3 & a_1 + a_2 \\ a_1 - a_2 & -a_3 \end{array} \right] \leftarrow \frac{1}{2} \left[\begin{array}{cc} 0 & +a_2 \\ -a_2 & 0 \end{array} \right] + \frac{1}{2} \left[\begin{array}{cc} a_3 & a_1 \\ a_1 & -a_3 \end{array} \right]$	$\mathfrak{su}(2) = \frac{1}{2} \left[\begin{array}{cc} i a_3 & i a_1 + a_2 \\ i a_1 - a_2 & -i a_3 \end{array} \right] \rightarrow \frac{1}{2} \left[\begin{array}{cc} 0 & +a_2 \\ -a_2 & 0 \end{array} \right] + \frac{i}{2} \left[\begin{array}{cc} a_3 & a_1 \\ a_1 & -a_3 \end{array} \right]$ $\mathfrak{sl}(2; \mathbb{R}) = \frac{1}{2} \left[\begin{array}{cc} a_3 & a_1 + a_2 \\ a_1 - a_2 & -a_3 \end{array} \right] \leftarrow \frac{1}{2} \left[\begin{array}{cc} 0 & +a_2 \\ -a_2 & 0 \end{array} \right] + \frac{1}{2} \left[\begin{array}{cc} a_3 & a_1 \\ a_1 & -a_3 \end{array} \right]$
gilmore-lie-groups_ EQ0983 (16.26)	306	■ 0.902	$\begin{gathered} \text{monomial} \\ y^2/x \\ y/x \\ 1 \\ x \\ y \\ xy \\ x^2y \\ xy^2 \\ xy^3 \\ xy^4 \end{gathered}$	$\begin{array}{c} \text{monomial} \\ y^2/x \\ y/x \\ 1 \\ x \\ y \\ xy \\ x^2y \\ xy^2 \\ xy^3 \\ xy^4 \end{array} \left[\begin{array}{cccccc} \xi_{00} & \xi_{10} & \xi_{01} & \eta_{00} & \eta_{10} & \eta_{01} \\ 0 & 0 & -2 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & +1 & 0 & 0 \\ 0 & 0 & 0 & 0 & +2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2 & 0 & 0 \\ 0 & 0 & 0 & 0 & -2 & 0 \\ 0 & -1 & 0 & 0 & 0 & -1 \\ 0 & 0 & -2 & 0 & 0 & 0 \\ 0 & 0 & +1 & 0 & 0 & 0 \end{array} \right] \left[\begin{array}{c} \xi_{00} \\ \xi_{10} \\ \xi_{01} \\ \eta_{00} \\ \eta_{10} \\ \eta_{01} \end{array} \right] = \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{array} \right]$	$\begin{array}{c} \text{monomial} \\ y^2/x \\ y/x \\ 1 \\ x \\ y \\ xy \\ x^2y \\ xy^2 \\ xy^3 \\ xy^4 \end{array} \left[\begin{array}{cccccc} \xi_{00} & \xi_{10} & \xi_{01} & \eta_{00} & \eta_{10} & \eta_{01} \\ 0 & 0 & -2 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & +1 & 0 & 0 \\ 0 & 0 & 0 & 0 & +2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2 & 0 & 0 \\ 0 & 0 & 0 & 0 & -2 & 0 \\ 0 & -1 & 0 & 0 & 0 & -1 \\ 0 & 0 & -2 & 0 & 0 & 0 \\ 0 & 0 & +1 & 0 & 0 & 0 \end{array} \right] \left[\begin{array}{c} \xi_{00} \\ \xi_{10} \\ \xi_{01} \\ \eta_{00} \\ \eta_{10} \\ \eta_{01} \end{array} \right] = \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{array} \right]$
gilmore-lie-groups_ EQ0990 (16.33)	308	■ 0.904	$\begin{array}{l} \text{surface equation} \\ \text{constraint equation} \end{array}$	$\begin{array}{l} \text{surface equation} \\ \text{constraint equation} \end{array}$	$\begin{array}{l} \text{surface equation} \\ \text{constraint equation} \end{array}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0436	132	■ 0.905	$\left[\begin{array}{lll} 1 & 0 & B \\ 0 & A & 0 \\ 0 & 0 & 1 \end{array}\right] \rightarrow e^{Aa^\dagger a + BI}$	$\begin{bmatrix} 1 & 0 & B \\ 0 & A & 0 \\ 0 & 0 & 1 \end{bmatrix} \rightarrow e^{Aa^\dagger a + BI}$	$\begin{bmatrix} 1 & 0 & B \\ 0 & A & 0 \\ 0 & 0 & 1 \end{bmatrix} \rightarrow e^{Aa^\dagger a + BI}$
gilmore-lie-groups_ EQ0924 (15.58)	287	■ 0.911	$\begin{aligned} M^{jj'}(k^0) \left\langle k^0; \begin{array}{c} j \\ \mu \end{array} \begin{array}{c} j' \\ \mu' \end{array} \middle \psi \right\rangle &= 0 \\ C_k M^{jj'}(k^0) C_k^{-1} \left\langle k; \begin{array}{c} j \\ \mu \end{array} \begin{array}{c} j' \\ \mu' \end{array} \middle \psi \right\rangle &= 0 \end{aligned}$	$M^{jj'}(k^0) \left\langle k^0; \begin{array}{c} j \\ \mu \end{array} \begin{array}{c} j' \\ \mu' \end{array} \middle \psi \right\rangle = 0$ $C_k M^{jj'}(k^0) C_k^{-1} \left\langle k; \begin{array}{c} j \\ \mu \end{array} \begin{array}{c} j' \\ \mu' \end{array} \middle \psi \right\rangle = 0$	$M^{jj'}(k^0) \left\langle k^0; \begin{array}{c} j \\ \mu \end{array} \begin{array}{c} j' \\ \mu' \end{array} \middle \psi \right\rangle = 0$ $C_k M^{jj'}(k^0) C_k^{-1} \left\langle k; \begin{array}{c} j \\ \mu \end{array} \begin{array}{c} j' \\ \mu' \end{array} \middle \psi \right\rangle = 0$
gilmore-lie-groups_ EQ0181	66	■ 0.911	$\left[\begin{array}{l} i_1 \\ o_2 \end{array}\right] = T_1 \left[\begin{array}{l} o_1 \\ i_2 \end{array}\right] \quad \left[\begin{array}{l} i_3 \\ o_4 \end{array}\right] = T_2 \left[\begin{array}{l} o_3 \\ i_4 \end{array}\right]$	$\begin{bmatrix} i_1 \\ o_2 \end{bmatrix} = T_1 \begin{bmatrix} o_1 \\ i_2 \end{bmatrix} \quad \begin{bmatrix} i_3 \\ o_4 \end{bmatrix} = T_2 \begin{bmatrix} o_3 \\ i_4 \end{bmatrix}$	$\begin{bmatrix} i_1 \\ o_2 \end{bmatrix} = T_1 \begin{bmatrix} o_1 \\ i_2 \end{bmatrix} \quad \begin{bmatrix} i_3 \\ o_4 \end{bmatrix} = T_2 \begin{bmatrix} o_3 \\ i_4 \end{bmatrix}$
gilmore-lie-groups_ EQ0279 (5.18)	93	■ 0.919	$\left[\begin{array}{ccc} 0 & \theta & v_1 \\ -\theta & 0 & v_2 \\ v_1 & v_2 & 0 \end{array}\right]$	$\begin{bmatrix} 0 & \theta & v_1 \\ -\theta & 0 & v_2 \\ v_1 & v_2 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \theta & v_1 \\ -\theta & 0 & v_2 \\ v_1 & v_2 & 0 \end{bmatrix}$
gilmore-lie-groups_ EQ0803	252	■ 0.923	$\begin{aligned} \mathbf{B} &= \mathbf{M}' u_4 + \mathbf{L} \times \mathbf{u} - \frac{3}{2} i \mathbf{u} = \frac{i}{2} [\mathbf{u}, \mathbf{L}^2 + \mathbf{M}'^2] \\ B_4 &= \mathbf{M}' \cdot \mathbf{u} + \frac{3}{2} i \mathbf{u} \\ &= \frac{i}{2} [u_4, \mathbf{L}^2 + \mathbf{M}'^2] \end{aligned}$	$\mathbf{B} = \mathbf{M}' u_4 + \mathbf{L} \times \mathbf{u} - \frac{3}{2} i \mathbf{u} = \frac{i}{2} [\mathbf{u}, \mathbf{L}^2 + \mathbf{M}'^2]$ $B_4 = \mathbf{M}' \cdot \mathbf{u} + \frac{3}{2} i \mathbf{u} = \frac{i}{2} [u_4, \mathbf{L}^2 + \mathbf{M}'^2]$	$\mathbf{B} = \mathbf{M}' u_4 + \mathbf{L} \times \mathbf{u} - \frac{3}{2} i \mathbf{u} = \frac{i}{2} [\mathbf{u}, \mathbf{L}^2 + \mathbf{M}'^2]$ $B_4 = \mathbf{M}' \cdot \mathbf{u} + \frac{3}{2} i \mathbf{u} = \frac{i}{2} [u_4, \mathbf{L}^2 + \mathbf{M}'^2]$
gilmore-lie-groups_ EQ0962 (16.5)	300	■ 0.930	$\left[\begin{array}{c} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial p} \end{array}\right] [p - g(x)] = \begin{bmatrix} * \\ 0 \\ * \end{bmatrix} \quad \frac{\partial}{\partial y} \begin{bmatrix} x \\ y \\ p \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$	$\begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial p} \end{bmatrix} [p - g(x)] = \begin{bmatrix} * \\ 0 \\ * \end{bmatrix} \quad \frac{\partial}{\partial y} \begin{bmatrix} x \\ y \\ p \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$	$\begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial p} \end{bmatrix} [p - g(x)] = \begin{bmatrix} * \\ 0 \\ * \end{bmatrix} \quad \frac{\partial}{\partial y} \begin{bmatrix} x \\ y \\ p \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08776 (14.19)	243	■ 0.932	$\begin{aligned} L_z &= \frac{\hbar}{i} \frac{\partial}{\partial \phi} \\ L_{\pm} &= \hbar \left(\pm \frac{\partial}{\partial \theta} + i \frac{\cos \theta}{\sin \theta} \frac{\partial}{\partial \phi} \right) \end{aligned}$		
gilmore-lie-groups_ E08096	44	■ 0.934	$\begin{aligned} \left[\begin{array}{l} x' \\ y' \\ \theta' \end{array} \right] &= - \begin{bmatrix} \cos(2\theta) & -\sin(2\theta) & 0 \\ \sin(2\theta) & \cos(2\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ \theta \end{bmatrix} \end{aligned}$		
gilmore-lie-groups_ E08545	170	■ 0.936	$\left(A_1, A_2, \dots, A_r \right)_{\Re e g} = \operatorname{tr} \Re e g \left(A_1 \right) \Re e g \left(A_2 \right) \cdots \Re e g \left(A_r \right)$		
gilmore-lie-groups_ E08084 (1.1)	18	■ 0.938	$\begin{aligned} \left(z - z_1 \right) \left(z - z_2 \right) \cdots \left(z - z_n \right) &= 0 \\ z^n - I_1 z^{n-1} + I_2 z^{n-2} + \cdots + (-1)^n I_n &= 0 \end{aligned}$		
gilmore-lie-groups_ E08784 (13.22)	224	■ 0.938	$\begin{aligned} X_{ij} &= x_i \partial_j - x_j \partial_i = \epsilon_{ijk} L_k \quad 1 \leq i, j, k \leq 3 \\ B_{i4} &= x_i \partial_4 + x_4 \partial_i \quad 1 \leq i \leq 3 \\ T_{i5} &= x_i \partial_5 \pm x_5 \partial_i \quad i = 1, 2, 3 \\ & \quad i = 4 \end{aligned}$		
gilmore-lie-groups_ E08148 (3.31)	57	■ 0.939	$U(n) \xrightarrow{\mathbb{C} \rightarrow 2 \times 2} \mathbb{R} O U(2n) \subset S O(2n)$		
gilmore-lie-groups_ E08689 (13.7)	221	■ 0.940	$\left[\begin{array}{l} \epsilon L_1 \\ \epsilon L_2 \end{array} \right] \xrightarrow{\epsilon=1/R} \left[\begin{array}{l} (1/R) L_1 \\ (1/R) L_2 \end{array} \right] \rightarrow \left[\begin{array}{l} -P_2 \\ +P_1 \end{array} \right]$		
gilmore-lie-groups_ E08932 (15.67)	288	■ 0.941	$\begin{aligned} \operatorname{Re} &+ \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} - \nabla \times \mathbf{B} = 0 \\ \operatorname{Im} &- \frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} - \nabla \times \mathbf{E} = 0 \end{aligned}$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0006 (1.3)	18	■ 0.944	$\begin{array}{l} \begin{array}{r} I \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \\ \quad (123) \rightarrow \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \\ \quad (321) \rightarrow \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \\ \quad (12) \rightarrow \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \\ \quad (23) \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \\ \quad (13) \rightarrow \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \end{array} \end{array}$	$I \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (123) \rightarrow \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \quad (321) \rightarrow \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$ $(12) \rightarrow \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (23) \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad (13) \rightarrow \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$	
gilmore-lie-groups_ EQ0416 (7.52)	127	■ 0.946	$\left\langle \begin{array}{c} j \\ j \end{array} \left e^{\beta' J_-} e^{n' J_z} e^{\alpha' J_+} \right \begin{array}{c} j \\ j \end{array} \right\rangle = e^{jn'} = (1 + \alpha\beta)^{2j}$	$\left\langle \begin{array}{c} j \\ j \end{array} \left e^{\beta' J_-} e^{n' J_z} e^{\alpha' J_+} \right \begin{array}{c} j \\ j \end{array} \right\rangle = e^{jn'} = (1 + \alpha\beta)^{2j}$	
gilmore-lie-groups_ EQ0490 (8.24)	150	■ 0.946	$\mathfrak{g} = \underbrace{\left(\hat{n} + \frac{1}{2}, a^{\dagger 2}, a^2 \right)}_{\mathfrak{su}(1,1)} + \underbrace{(a^{\dagger}, a, I)}_{\text{nilpotent}}$	$\mathfrak{g} = \underbrace{\left(\hat{n} + \frac{1}{2}, a^{\dagger 2}, a^2 \right)}_{\mathfrak{su}(1,1)} + \underbrace{(a^{\dagger}, a, I)}_{\text{nilpotent invariant subalgebra}}$	
gilmore-lie-groups_ EQ0620 (12.1)	204	■ 0.947	$\begin{aligned} \mathfrak{g} &= \mathfrak{h} + \mathfrak{p} \quad (\mathfrak{p}, \mathfrak{p}) < 0 \\ \mathfrak{g}' &= \mathfrak{h} + i\mathfrak{p} \quad (i\mathfrak{p}, i\mathfrak{p}) > 0 \end{aligned}$	$\mathfrak{g} = \mathfrak{h} + \mathfrak{p} \quad (\mathfrak{p}, \mathfrak{p}) < 0$ $\mathfrak{g}' = \mathfrak{h} + i\mathfrak{p} \quad (i\mathfrak{p}, i\mathfrak{p}) > 0$	
gilmore-lie-groups_ EQ0225 (4.35)	78	■ 0.948	$(A, A) = \sum_i \sum_j \left A_i^j \right ^2 \geq 0 \quad = 0 \Rightarrow A = 0$	$(A, A) = \sum_i \sum_j \left A_i^j \right ^2 \geq 0 \quad = 0 \Rightarrow A = 0$	
gilmore-lie-groups_ EQ0903 (15.37)	282	■ 0.950	$\begin{aligned} b_3 &= 0 \\ b_2 &= +\theta_1 \\ b_1 &= -\theta_2 \end{aligned}$	$b_3 = 0$ $b_2 = +\theta_1$ $b_1 = -\theta_2$	
gilmore-lie-groups_ EQ0016 (1.12)	21	■ 0.951	$\begin{array}{l} g_i \cdot g_j = g_{ij} \\ \downarrow \quad \downarrow \quad \downarrow \\ f(g_i) \times f(g_j) = f(g_{ij}) \end{array}$	$\begin{array}{l} g_i \cdot g_j = g_{ij} \\ \downarrow \quad \downarrow \quad \downarrow \\ f(g_i) \times f(g_j) = f(g_{ij}) \end{array}$	
gilmore-lie-groups_ EQ0167	64	■ 0.951	$\begin{aligned} UT(p, q) &= UT(q, p) & SO(p, q) &= SO(q, p) \\ A(p, q) &= A(q, p) & U(p, q) &= U(q, p) \\ Sp(p, q) &= Sp(q, p) \end{aligned}$	$UT(p, q) = UT(q, p) \quad SO(p, q) = SO(q, p)$ $A(p, q) = A(q, p) \quad U(p, q) = U(q, p)$ $Sp(p, q) = Sp(q, p)$	

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00211 (4.21)	76	■ 0.951	$X_{\{i\}}=A_{\{i\}}^{\{r\}} Y_{\{r\}}$	$X_i = A_i^r Y_r$	$X_i = A_i^r Y_r$
gilmore-lie-groups_ E00214 (4.24)	76	■ 0.952	$\left[Z, X_{\{i\}} \right] = R(Z)_{\{i\}}^{\{j\}} X_{\{j\}}$	$[Z, X_i] = R(Z)_i^j X_j$	$[Z, X_i] = R(Z)_i^j X_j$
gilmore-lie-groups_ E00433 (7.69)	132	■ 0.952	$\begin{aligned} e^{\{-\beta M(H)\}} = & \mathrm{EXP} \{-\beta \left[\begin{array}{ccc} 0 & \alpha^* & \delta \\ 0 & \hbar \omega & \alpha \\ 0 & 0 & 0 \end{array} \right] \\ & \left[\begin{array}{ccc} 1 & \alpha^* & \frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \\ 0 & e^{-\beta \hbar \omega} & \frac{e^{-\beta \hbar \omega} - 1 + \beta \hbar \omega}{(\hbar \omega)^2} \alpha^* \alpha - \beta \delta \\ 0 & 0 & \alpha \frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \end{array} \right] \right] \\ & \frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \alpha^* \alpha - \beta \delta \end{aligned}$	$e^{-\beta M(H)} = \mathrm{EXP} - \beta \left[\begin{array}{ccc} 0 & \alpha^* & \delta \\ 0 & \hbar \omega & \alpha \\ 0 & 0 & 0 \end{array} \right] = \left[\begin{array}{ccc} 1 & \alpha^* & \frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \\ 0 & e^{-\beta \hbar \omega} & \frac{e^{-\beta \hbar \omega} - 1 + \beta \hbar \omega}{(\hbar \omega)^2} \alpha^* \alpha - \beta \delta \\ 0 & 0 & \alpha \frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \end{array} \right]$	$e^{-\beta M(H)} = \mathrm{EXP} - \beta \left[\begin{array}{ccc} 0 & \alpha^* & \delta \\ 0 & \hbar \omega & \alpha \\ 0 & 0 & 0 \end{array} \right] = \left[\begin{array}{ccc} 1 & \alpha^* & \frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \\ 0 & e^{-\beta \hbar \omega} & \frac{e^{-\beta \hbar \omega} - 1 + \beta \hbar \omega}{(\hbar \omega)^2} \alpha^* \alpha - \beta \delta \\ 0 & 0 & \alpha \frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \end{array} \right]$
gilmore-lie-groups_ E00354	110	■ 0.953	$\left[\begin{array}{c} a \\ a^\dagger \end{array} \right] = \frac{1}{\sqrt{2}} \left[\begin{array}{cc} 1 & 1 \\ -1 & 1 \end{array} \right] \left[\begin{array}{c} \partial \\ x \end{array} \right]$	$\left[\begin{array}{c} a \\ a^\dagger \end{array} \right] = \frac{1}{\sqrt{2}} \left[\begin{array}{cc} 1 & 1 \\ -1 & 1 \end{array} \right] \left[\begin{array}{c} \partial \\ x \end{array} \right]$	$\left[\begin{array}{c} a \\ a^\dagger \end{array} \right] = \frac{1}{\sqrt{2}} \left[\begin{array}{cc} 1 & 1 \\ -1 & 1 \end{array} \right] \left[\begin{array}{c} \partial \\ x \end{array} \right]$
gilmore-lie-groups_ E00414 (7.50)	127	■ 0.954	$\left\langle \begin{array}{c} j \\ j \end{array} \middle J_+^k J_-^k \middle \begin{array}{c} j \\ j \end{array} \right\rangle$	$\left\langle \begin{array}{c} j \\ j \end{array} \middle J_+^k J_-^k \middle \begin{array}{c} j \\ j \end{array} \right\rangle$	$\left\langle \begin{array}{c} j \\ j \end{array} \middle J_+^k J_-^k \middle \begin{array}{c} j \\ j \end{array} \right\rangle$
gilmore-lie-groups_ E00918 (15.53)	285	■ 0.954	$\begin{aligned} & D^{\{0\}}(H_{k^0}) = \mathrm{EXP} \left(\theta_3 J_3^{(j')} + \theta_1 (J_1^{(j')} - i J_2^{(j')}) + \theta_2 (J_2^{(j')} + i J_1^{(j')}) \right) \\ & = \mathrm{EXP} \left(\theta_3 J_3^{(j')} + (\theta_1 + i \theta_2) (J_1^{(j')} - i J_2^{(j')}) \right) \\ & = \left[\begin{array}{cccc} e^{ij'\theta_3} & & & \\ * & e^{i(j'-1)\theta_3} & & \\ * & * & \ddots & \\ * & * & * & \ddots \\ * & * & * & * & e^{-ij'\theta_3} \end{array} \right] \end{aligned}$	$D^{0j'}(H_{k^0}) = \mathrm{EXP} \left(\theta_3 J_3^{(j')} + \theta_1 (J_1^{(j')} - i J_2^{(j')}) + \theta_2 (J_2^{(j')} + i J_1^{(j')}) \right) \\ = \mathrm{EXP} \left(\theta_3 J_3^{(j')} + (\theta_1 + i \theta_2) (J_1^{(j')} - i J_2^{(j')}) \right) \\ = \left[\begin{array}{cccc} e^{ij'\theta_3} & & & \\ * & e^{i(j'-1)\theta_3} & & \\ * & * & \ddots & \\ * & * & * & \ddots \\ * & * & * & * & e^{-ij'\theta_3} \end{array} \right]$	$D^{0j'}(H_{k^0}) = \mathrm{EXP} \left(\theta_3 J_3^{(j')} + \theta_1 (J_1^{(j')} - i J_2^{(j')}) + \theta_2 (J_2^{(j')} + i J_1^{(j')}) \right) \\ = \mathrm{EXP} \left(\theta_3 J_3^{(j')} + (\theta_1 + i \theta_2) (J_1^{(j')} - i J_2^{(j')}) \right) \\ = \left[\begin{array}{cccc} e^{ij'\theta_3} & & & \\ * & e^{i(j'-1)\theta_3} & & \\ * & * & \ddots & \\ * & * & * & \ddots \\ * & * & * & * & e^{-ij'\theta_3} \end{array} \right]$
gilmore-lie-groups_ E00215 (4.25)	76	■ 0.955	$\left[Z, Y_{\{r\}} \right] = S(Z)_{\{r\}}^{\{s\}} Y_{\{s\}}$	$[Z, Y_r] = S(Z)_r^s Y_s$	$[Z, Y_r] = S(Z)_r^s Y_s$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08145 (3.28)	57	■ 0.955	$\operatorname{det} M=+1 \mathbb{F}=\begin{cases} \mathbb{R} & SL(n,\mathbb{R}) \\ \mathbb{C} & SL(n,\mathbb{C}) \\ \mathbb{Q} & SL(n,\mathbb{Q}) \end{cases}$	$\det M = +1\mathbb{F} = \begin{cases} \mathbb{R} & SL(n,\mathbb{R}) \\ \mathbb{C} & SL(n,\mathbb{C}) \\ \mathbb{Q} & SL(n,\mathbb{Q}) \end{cases}$	$\det M = +1 \quad \mathbb{F} = \begin{cases} \mathbb{R} & SL(n,\mathbb{R}) \\ \mathbb{C} & SL(n,\mathbb{C}) \\ \mathbb{Q} & SL(n,\mathbb{Q}) \end{cases}$
gilmore-lie-groups_ E08138 (3.22)	55	■ 0.957	$\begin{array}{l} \mathbb{R} & O(n;G) \\ \mathbb{C} & U(n;G) \\ \mathbb{Q} & Sp(n;G) \end{array}$	$\begin{array}{l} \mathbb{R} & O(n;G) \\ \mathbb{C} & U(n;G) \\ \mathbb{Q} & Sp(n;G) \end{array}$	$\begin{array}{l} \mathbb{R} & O(n;G) \\ \mathbb{C} & U(n;G) \\ \mathbb{Q} & Sp(n;G) \end{array}$
gilmore-lie-groups_ E06692 (13.10)	221	■ 0.957	$\begin{aligned} & \frac{1}{R} L_1 = \frac{1}{R} (x^2 \partial_3 - x^3 \partial_2) = \frac{1}{R} (y \partial / \partial z - R \partial / \partial y) \rightarrow -\partial / \partial y = -\partial_2 = -P_2 \\ & \frac{1}{R} L_2 = \frac{1}{R} (x^3 \partial_1 - x^1 \partial_3) = \frac{1}{R} (R \partial / \partial x - x \partial / \partial z) \rightarrow +\partial / \partial x = +\partial_1 = +P_1 \end{aligned}$	$\begin{aligned} \frac{1}{R} L_1 &= \frac{1}{R} (x^2 \partial_3 - x^3 \partial_2) = \frac{1}{R} (y \partial / \partial z - R \partial / \partial y) \rightarrow -\partial / \partial y = -\partial_2 = -P_2 \\ \frac{1}{R} L_2 &= \frac{1}{R} (x^3 \partial_1 - x^1 \partial_3) = \frac{1}{R} (R \partial / \partial x - x \partial / \partial z) \rightarrow +\partial / \partial x = +\partial_1 = +P_1 \end{aligned}$	$\frac{1}{R} L_1 = \frac{1}{R} (x^2 \partial_3 - x^3 \partial_2) = \frac{1}{R} (y \partial / \partial z - R \partial / \partial y) \rightarrow -\partial / \partial y = -\partial_2 = -P_2$ $\frac{1}{R} L_2 = \frac{1}{R} (x^3 \partial_1 - x^1 \partial_3) = \frac{1}{R} (R \partial / \partial x - x \partial / \partial z) \rightarrow +\partial / \partial x = +\partial_1 = +P_1$
gilmore-lie-groups_ E06631 (12.4)	206	■ 0.957	$\begin{aligned} & \operatorname{SU}(p,q) / \operatorname{S}[U(p) \times U(q)], \operatorname{Sp}(p,q) / \operatorname{Sp}(p) \times \operatorname{Sp}(q) \text{ are} \\ & \left\ \begin{bmatrix} 0 & B \\ B^\dagger & 0 \end{bmatrix} - \lambda I_{p+q} \right\ = \sum_{j=0}^{n=p+q} (-\lambda)^{n-j} \phi_j(B, B^\dagger) \end{aligned}$	$\operatorname{SU}(p,q) / \operatorname{S}[U(p) \times U(q)], \operatorname{Sp}(p,q) / \operatorname{Sp}(p) \times \operatorname{Sp}(q) \text{ are}$ $\left\ \begin{bmatrix} 0 & B \\ B^\dagger & 0 \end{bmatrix} - \lambda I_{p+q} \right\ = \sum_{j=0}^{n=p+q} (-\lambda)^{n-j} \phi_j(B, B^\dagger)$	$SU(p,q)/S[U(p) \times U(q)], Sp(p,q)/Sp(p) \times Sp(q) \text{ are}$ $\left\ \begin{bmatrix} 0 & B \\ B^\dagger & 0 \end{bmatrix} - \lambda I_{p+q} \right\ = \sum_{j=0}^{n=p+q} (-\lambda)^{n-j} \phi_j(B, B^\dagger)$
gilmore-lie-groups_ E0531	163	■ 0.958	$\beta - m\alpha \quad \beta - (m-1)\alpha \quad \cdots \quad \beta \quad \beta + \alpha \quad \cdots \quad \beta + n\alpha$	$\beta - m\alpha \quad \beta - (m-1)\alpha \quad \cdots \quad \beta \quad \beta + \alpha \quad \cdots \quad \beta + n\alpha$	$\beta - m\alpha \quad \beta - (m-1)\alpha \quad \cdots \quad \beta \quad \beta + \alpha \quad \cdots \quad \beta + n\alpha$
gilmore-lie-groups_ E08010	19	■ 0.959	$\begin{aligned} & S_2(12) = \{I, (12)\} \\ & S_2(23) = \{I, (23)\} \\ & S_2(13) = \{I, (13)\} \end{aligned}$	$\begin{aligned} S_2(12) &= \{I, (12)\} \\ S_2(23) &= \{I, (23)\} \\ S_2(13) &= \{I, (13)\} \end{aligned}$	$\begin{aligned} S_2(12) &= \{I, (12)\} \\ S_2(23) &= \{I, (23)\} \\ S_2(13) &= \{I, (13)\} \end{aligned}$
gilmore-lie-groups_ E08140 (3.24)	55	■ 0.960	$\begin{aligned} & \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A^t & C^t \\ B^t & D \end{bmatrix} = \begin{bmatrix} AA^t & AC^t \\ CA^t & CC^t \end{bmatrix} = \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix} \\ & \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A^t & C^t \\ B^t & D \end{bmatrix} = \begin{bmatrix} AA^t & AC^t \\ CA^t & CC^t \end{bmatrix} = \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix} \end{aligned}$	$\begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A^t & C^t \\ B^t & D \end{bmatrix} = \begin{bmatrix} AA^t & AC^t \\ CA^t & CC^t \end{bmatrix} = \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix}$	$\begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A^t & C^t \\ B^t & D \end{bmatrix} = \begin{bmatrix} AA^t & AC^t \\ CA^t & CC^t \end{bmatrix} = \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix}$
gilmore-lie-groups_ E08163	63	■ 0.960	$SO(2) \longrightarrow \begin{bmatrix} \pm\sqrt{1-x^2} & x \\ -x & \pm\sqrt{1-x^2} \end{bmatrix}$	$SO(2) \longrightarrow \begin{bmatrix} \pm\sqrt{1-x^2} & x \\ -x & \pm\sqrt{1-x^2} \end{bmatrix}$	$SO(2) \longrightarrow \begin{bmatrix} \pm\sqrt{1-x^2} & x \\ -x & \pm\sqrt{1-x^2} \end{bmatrix}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0133 (3.19)	54	■ 0.961	$\begin{aligned} & M^t I_{3,1} M = I_{3,1} \quad \& \& \left[\begin{array}{cc} A^t & C^t \\ B^t & D^t \end{array} \right] \left[\begin{array}{cc} I_3 & 0 \\ 0 & -1 \end{array} \right] \left[\begin{array}{cc} A & B \\ C & D \end{array} \right] \\ & = \left[\begin{array}{cc} A^t A - C^t C & A^t B - C^t D \\ B^t A - D^t C & B^t B - D^t D \end{array} \right] = \left[\begin{array}{cc} I_3 & 0 \\ 0 & -1 \end{array} \right] \end{aligned}$	$M^t I_{3,1} M = I_{3,1}$ $\begin{bmatrix} A^t & C^t \\ B^t & D^t \end{bmatrix} \begin{bmatrix} I_3 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} A^t A - C^t C & A^t B - C^t D \\ B^t A - D^t C & B^t B - D^t D \end{bmatrix} = \begin{bmatrix} I_3 & 0 \\ 0 & -1 \end{bmatrix}$	$M^t I_{3,1} M = I_{3,1}$ $\begin{bmatrix} A^t & C^t \\ B^t & D^t \end{bmatrix} \begin{bmatrix} I_3 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} A^t A - C^t C & A^t B - C^t D \\ B^t A - D^t C & B^t B - D^t D \end{bmatrix} = \begin{bmatrix} I_3 & 0 \\ 0 & -1 \end{bmatrix}$
gilmore-lie-groups_ EQ0122 (3.10)	52	■ 0.961	$\left[\begin{array}{c cc c} 1 & * & * & * \\ \hline 0 & SU(1,1) & * & \\ 0 & & * & \\ \hline 0 & 0 & 0 & 1 \end{array} \right]$	$\left[\begin{array}{c cc c} 1 & * & * & * \\ \hline 0 & SU(1,1) & * & \\ 0 & & * & \\ \hline 0 & 0 & 0 & 1 \end{array} \right]$	$\left[\begin{array}{c cc c} 1 & * & * & * \\ \hline 0 & SU(1,1) & * & \\ 0 & & * & \\ \hline 0 & 0 & 0 & 1 \end{array} \right]$
gilmore-lie-groups_ EQ0706 (13.23)	224	■ 0.964	$\begin{aligned} & \mathcal{C}^4[SO(5)] = W^\alpha W_\alpha \\ & W^\alpha = \epsilon^{\alpha\beta\gamma\mu\nu} X_{\beta\gamma} X_{\mu\nu} \end{aligned}$	$\mathcal{C}^4[SO(5)] = W^\alpha W_\alpha$ $W^\alpha = \epsilon^{\alpha\beta\gamma\mu\nu} X_{\beta\gamma} X_{\mu\nu}$	$\mathcal{C}^4[SO(5)] = W^\alpha W_\alpha$ $W^\alpha = \epsilon^{\alpha\beta\gamma\mu\nu} X_{\beta\gamma} X_{\mu\nu}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00944	292	0.965	$\begin{aligned} & \left[\begin{array}{ccccc} 0 & \partial_y & -\partial_x & 2\partial_z & 0 \\ -\partial_y & 0 & \partial_z & -\partial_x & -\sqrt{3}\partial_x \\ \partial_x & -\partial_z & 0 & -\partial_y & \sqrt{3}\partial_y \\ -2\partial_z & \partial_x & \partial_y & 0 & 0 \\ 0 & \sqrt{3}\partial_x & -\sqrt{3}\partial_y & 0 & 0 \end{array} \right] \begin{pmatrix} F_1 \\ F_2 \\ F_3 \\ F_4 \\ F_5 \end{pmatrix} + \frac{2}{c} \frac{\partial}{\partial t} \begin{pmatrix} G_1 \\ G_2 \\ G_3 \\ G_4 \\ G_5 \end{pmatrix} = 0 \\ & \left[\begin{array}{ccccc} 0 & \partial_y & -\partial_x & 2\partial_z & 0 \\ -\partial_y & 0 & \partial_z & -\partial_x & -\sqrt{3}\partial_x \\ \partial_x & -\partial_z & 0 & -\partial_y & \sqrt{3}\partial_y \\ -2\partial_z & \partial_x & \partial_y & 0 & 0 \\ 0 & \sqrt{3}\partial_x & -\sqrt{3}\partial_y & 0 & 0 \end{array} \right] \begin{pmatrix} G_1 \\ G_2 \\ G_3 \\ G_4 \\ G_5 \end{pmatrix} - \frac{2}{c} \frac{\partial}{\partial t} \begin{pmatrix} F_1 \\ F_2 \\ F_3 \\ F_4 \\ F_5 \end{pmatrix} = 0 \end{aligned}$	$\begin{bmatrix} 0 & \partial_y & -\partial_x & 2\partial_z & 0 \\ -\partial_y & 0 & \partial_z & -\partial_x & -\sqrt{3}\partial_x \\ \partial_x & -\partial_z & 0 & -\partial_y & \sqrt{3}\partial_y \\ -2\partial_z & \partial_x & \partial_y & 0 & 0 \\ 0 & \sqrt{3}\partial_x & -\sqrt{3}\partial_y & 0 & 0 \end{bmatrix} \begin{pmatrix} F_1 \\ F_2 \\ F_3 \\ F_4 \\ F_5 \end{pmatrix} + \frac{2}{c} \frac{\partial}{\partial t} \begin{pmatrix} G_1 \\ G_2 \\ G_3 \\ G_4 \\ G_5 \end{pmatrix} = 0$ $\begin{bmatrix} 0 & \partial_y & -\partial_x & 2\partial_z & 0 \\ -\partial_y & 0 & \partial_z & -\partial_x & -\sqrt{3}\partial_x \\ \partial_x & -\partial_z & 0 & -\partial_y & \sqrt{3}\partial_y \\ -2\partial_z & \partial_x & \partial_y & 0 & 0 \\ 0 & \sqrt{3}\partial_x & -\sqrt{3}\partial_y & 0 & 0 \end{bmatrix} \begin{pmatrix} G_1 \\ G_2 \\ G_3 \\ G_4 \\ G_5 \end{pmatrix} - \frac{2}{c} \frac{\partial}{\partial t} \begin{pmatrix} F_1 \\ F_2 \\ F_3 \\ F_4 \\ F_5 \end{pmatrix} = 0$	
gilmore-lie-groups_ E00105	47	0.965	$\begin{aligned} & \left[\begin{array}{c} X \\ Y \end{array} \right] \rightarrow \left[\begin{array}{c} X' \\ Y' \end{array} \right] \\ & = \left(\left[\begin{array}{cc} m_{11} & m_{12} \\ m_{21} & m_{22} \end{array} \right] \left[\begin{array}{c} X \\ Y \end{array} \right] + \left[\begin{array}{c} m_{13} \\ m_{23} \end{array} \right] \right) / \left(\left[\begin{array}{cc} m_{31} & m_{32} \end{array} \right] \left[\begin{array}{c} X \\ Y \end{array} \right] + m_{33} \right) \end{aligned}$	$\left[\begin{array}{c} X \\ Y \end{array} \right] \rightarrow \left[\begin{array}{c} X' \\ Y' \end{array} \right]$ $= \left(\left[\begin{array}{cc} m_{11} & m_{12} \\ m_{21} & m_{22} \end{array} \right] \left[\begin{array}{c} X \\ Y \end{array} \right] + \left[\begin{array}{c} m_{13} \\ m_{23} \end{array} \right] \right) / \left(\left[\begin{array}{cc} m_{31} & m_{32} \end{array} \right] \left[\begin{array}{c} X \\ Y \end{array} \right] + m_{33} \right)$	$\left[\begin{array}{c} X \\ Y \end{array} \right] \rightarrow \left[\begin{array}{c} X' \\ Y' \end{array} \right]$ $= \left(\left[\begin{array}{cc} m_{11} & m_{12} \\ m_{21} & m_{22} \end{array} \right] \left[\begin{array}{c} X \\ Y \end{array} \right] + \left[\begin{array}{c} m_{13} \\ m_{23} \end{array} \right] \right) / \left(\left[\begin{array}{cc} m_{31} & m_{32} \end{array} \right] \left[\begin{array}{c} X \\ Y \end{array} \right] + m_{33} \right)$
gilmore-lie-groups_ E00682	218	0.965	$\begin{aligned} & \left[\begin{array}{c} a \\ b \\ c \\ d \end{array} \right] S^{-1} = \left[\begin{array}{cc} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{array} \right] \quad S = \frac{1}{\sqrt{2}} \left[\begin{array}{cc} 1 & -i \\ -i & 1 \end{array} \right] \\ & S^{-1} = \left[\begin{array}{cc} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{array} \right] \quad S = \frac{1}{\sqrt{2}} \left[\begin{array}{cc} 1 & -i \\ -i & 1 \end{array} \right] \end{aligned}$	$S \left[\begin{array}{cc} a & b \\ c & d \end{array} \right] S^{-1} = \left[\begin{array}{cc} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{array} \right] \quad S = \frac{1}{\sqrt{2}} \left[\begin{array}{cc} 1 & -i \\ -i & 1 \end{array} \right]$	$S \left[\begin{array}{cc} a & b \\ c & d \end{array} \right] S^{-1} = \left[\begin{array}{cc} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{array} \right] \quad S = \frac{1}{\sqrt{2}} \left[\begin{array}{cc} 1 & -i \\ -i & 1 \end{array} \right]$
gilmore-lie-groups_ E00392 (7.28)	123	0.966	$\begin{aligned} & w = \ln x \\ & z = y \ln(x)/(x-1) \quad z = 0 \text{ for } x = 1 \end{aligned}$	$w = \ln x$ $z = y \ln(x)/(x-1) \quad z = 0 \text{ for } x = 1$	$w = \ln x$ $z = y \ln(x)/(x-1) \quad z = 0 \text{ for } x = 1$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08393 (7.29)	123	■ 0.966	$\left[\begin{array}{lll} 0 & l & \delta \\ 0 & 0 & r \\ 0 & 0 & 0 \end{array}\right]=l X_l+r X_r+\delta X_\delta$	$\begin{bmatrix} 0 & l & \delta \\ 0 & 0 & r \\ 0 & 0 & 0 \end{bmatrix} = lX_l + rX_r + \delta X_\delta$	$\begin{bmatrix} 0 & l & \delta \\ 0 & 0 & r \\ 0 & 0 & 0 \end{bmatrix} = lX_l + rX_r + \delta X_\delta$
gilmore-lie-groups_ E08657	213	■ 0.966	$\operatorname{EXP}\left[\begin{array}{c cc} 0 & t_1 & t_2 \\ t_1 & 0 & 0 \\ t_2 & 0 & 0 \end{array}\right]=\left[\begin{array}{c cc} x_0 & x_1 & x_2 \\ x_1 & * & * \\ x_2 & * & * \end{array}\right] \quad x_0^2-x_1^2-x_2^2=1$	$\operatorname{EXP}\left[\begin{array}{c cc} 0 & t_1 & t_2 \\ t_1 & 0 & 0 \\ t_2 & 0 & 0 \end{array}\right]=\left[\begin{array}{c cc} x_0 & x_1 & x_2 \\ x_1 & * & * \\ x_2 & * & * \end{array}\right] \quad x_0^2-x_1^2-x_2^2=1$	$\operatorname{EXP}\left[\begin{array}{c cc} 0 & t_1 & t_2 \\ t_1 & 0 & 0 \\ t_2 & 0 & 0 \end{array}\right]=\left[\begin{array}{c cc} x_0 & x_1 & x_2 \\ x_1 & * & * \\ x_2 & * & * \end{array}\right] \quad x_0^2-x_1^2-x_2^2=1$
gilmore-lie-groups_ E08157 (3.39)	61	■ 0.967	$l=\left[\begin{array}{ll} 1 & 0 \\ 0 & 1 \end{array}\right] \quad \mathcal{I}=i\left[\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array}\right] \quad \mathcal{J}=i\left[\begin{array}{cc} 0 & -i \\ i & 0 \end{array}\right] \quad \mathcal{K}=i\left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right]$	$1=\left[\begin{array}{ll} 1 & 0 \\ 0 & 1 \end{array}\right] \quad \mathcal{I}=i\left[\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array}\right] \quad \mathcal{J}=i\left[\begin{array}{cc} 0 & -i \\ i & 0 \end{array}\right] \quad \mathcal{K}=i\left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right]$	$1=\left[\begin{array}{ll} 1 & 0 \\ 0 & 1 \end{array}\right] \quad \mathcal{I}=i\left[\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array}\right] \quad \mathcal{J}=i\left[\begin{array}{cc} 0 & -i \\ i & 0 \end{array}\right] \quad \mathcal{K}=i\left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right]$
gilmore-lie-groups_ E08216 (4.26)	76	■ 0.969	$S_{\{r\}^{\{S\}}(Z)=\left(A^{-1}\right)_r^i R(Z)_i^j A_j^s$	$S_r^S(Z)=\left(A^{-1}\right)_r^i R(Z)_i^j A_j^s$	$S_r^S(Z)=\left(A^{-1}\right)_r^i R(Z)_i^j A_j^s$
gilmore-lie-groups_ E08468 (7.86)	138	■ 0.970	$\left \begin{array}{c} j \\ \theta \end{array}\right\rangle=e^{i\left(\theta_x J_x+\theta_y J_y\right)}\left \begin{array}{c} j \\ -j \end{array}\right\rangle \quad i\left(\theta_x J_x+\theta_y J_y\right)=\frac{i}{2}\left[\begin{array}{cc} 0 & \theta_x-i \theta_y \\ \theta_x+i \theta_y & 0 \end{array}\right]$	$\left \begin{array}{c} j \\ \theta \end{array}\right\rangle=e^{i\left(\theta_x J_x+\theta_y J_y\right)}\left \begin{array}{c} j \\ -j \end{array}\right\rangle \quad i\left(\theta_x J_x+\theta_y J_y\right)=\frac{i}{2}\left[\begin{array}{cc} 0 & \theta_x-i \theta_y \\ \theta_x+i \theta_y & 0 \end{array}\right]$	$\left \begin{array}{c} j \\ \theta \end{array}\right\rangle=e^{i\left(\theta_x J_x+\theta_y J_y\right)}\left \begin{array}{c} j \\ -j \end{array}\right\rangle \quad i\left(\theta_x J_x+\theta_y J_y\right)=\frac{i}{2}\left[\begin{array}{cc} 0 & \theta_x-i \theta_y \\ \theta_x+i \theta_y & 0 \end{array}\right]$
gilmore-lie-groups_ E08662	214	■ 0.970	$\Gamma_{rs}^{t}=\frac{1}{2}g^{tu}\left(\partial_sg_{ru}+\partial_rg_{su}-\partial_ug_{rs}\right)$	$\Gamma_{rs}^{t}=\frac{1}{2}g^{tu}\left(\partial_sg_{ru}+\partial_rg_{su}-\partial_ug_{rs}\right)$	$\Gamma_{rs}^{t}=\frac{1}{2}g^{tu}\left(\partial_sg_{ru}+\partial_rg_{su}-\partial_ug_{rs}\right)$
gilmore-lie-groups_ E08818 (1.14)	22	■ 0.970	$\begin{array}{c} G/H \\ \{I,(12)\} \\ \{I,(12),(321)\} \end{array} \cdot \begin{array}{c} H \\ \{I,(123),(321)\} \\ \{I,(23)\} \end{array} = \begin{array}{c} G \\ \{I,(12),(321)\} \\ \{I,(12),(321)\} \cdot \{I,(23)\} \end{array} = \begin{array}{c} H \\ \{I,(123),(321),(12),(13),(23)\} \\ \{I,(23)\} \end{array} = \begin{array}{c} G \\ \{I,(123),(321),(12),(13),(23)\} \\ \{I,(23),(12),(123),(321),(13)\} \end{array}$	$\begin{array}{c} G/H \\ \{I,(12)\} \\ \{I,(12),(321)\} \end{array} \cdot \begin{array}{c} H \\ \{I,(123),(321)\} \\ \{I,(23)\} \end{array} = \begin{array}{c} H \\ \{I,(123),(321),(12),(13),(23)\} \\ \{I,(23)\} \end{array} = \begin{array}{c} G \\ \{I,(123),(321),(12),(13),(23)\} \\ \{I,(23),(12),(123),(321),(13)\} \end{array}$	$\begin{array}{c} G/H \\ \{I,(12)\} \\ \{I,(12),(321)\} \end{array} \cdot \begin{array}{c} H \\ \{I,(123),(321)\} \\ \{I,(23)\} \end{array} = \begin{array}{c} G \\ \{I,(123),(321),(12),(13),(23)\} \\ \{I,(23),(12),(123),(321),(13)\} \end{array}$
gilmore-lie-groups_ E08415 (7.51)	127	■ 0.971	$\left\langle\begin{array}{c} j \\ j \end{array}\right e^{\alpha J_+}e^{\beta J_-}\left \begin{array}{c} j \\ j \end{array}\right\rangle=\sum_{rs}\frac{\alpha^r\beta^s}{r!s!}\left\langle\begin{array}{c} j \\ j \end{array}\right J_+^rJ_-^s\left \begin{array}{c} j \\ j \end{array}\right\rangle$	$\left\langle\begin{array}{c} j \\ j \end{array}\right e^{\alpha J_+}e^{\beta J_-}\left \begin{array}{c} j \\ j \end{array}\right\rangle=\sum_{rs}\frac{\alpha^r\beta^s}{r!s!}\left\langle\begin{array}{c} j \\ j \end{array}\right J_+^rJ_-^s\left \begin{array}{c} j \\ j \end{array}\right\rangle$	$\left\langle\begin{array}{c} j \\ j \end{array}\right e^{\alpha J_+}e^{\beta J_-}\left \begin{array}{c} j \\ j \end{array}\right\rangle=\sum_{rs}\frac{\alpha^r\beta^s}{r!s!}\left\langle\begin{array}{c} j \\ j \end{array}\right J_+^rJ_-^s\left \begin{array}{c} j \\ j \end{array}\right\rangle$
gilmore-lie-groups_ E08121 (3.9)	51	■ 0.972	$\left[\begin{array}{c} x' \\ 1 \end{array}\right]=\left[\begin{array}{cc} a & b \\ 0 & 1 \end{array}\right]\left[\begin{array}{c} x \\ 1 \end{array}\right]=\left[\begin{array}{c} ax+b \\ 1 \end{array}\right]$	$\left[\begin{array}{c} x' \\ 1 \end{array}\right]=\left[\begin{array}{cc} a & b \\ 0 & 1 \end{array}\right]\left[\begin{array}{c} x \\ 1 \end{array}\right]=\left[\begin{array}{c} ax+b \\ 1 \end{array}\right]$	$\left[\begin{array}{c} x' \\ 1 \end{array}\right]=\left[\begin{array}{cc} a & b \\ 0 & 1 \end{array}\right]\left[\begin{array}{c} x \\ 1 \end{array}\right]=\left[\begin{array}{c} ax+b \\ 1 \end{array}\right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0661	214	■0.972	$\Delta=g^{\mathrm{r}\,s}\,\partial_r\partial_s-g^{\mathrm{r}\,s}\Gamma_{rs}^{\,\,t}\partial_t$ $\Gamma_{rs}^{\,\,t}\partial_t$	$\Delta = g^{rs}\partial_r\partial_s - g^{rs}\Gamma_{rs}^{\,\,t}\partial_t$	$\Delta = g^{rs}\partial_r\partial_s - g^{rs}\Gamma_{rs}^{\,\,t}\partial_t$
gilmore-lie-groups_ EQ9987 (16.30)	307	■0.973	$\left(\begin{array}{c} R \\ S \\ T \end{array}\right)=\left(\begin{array}{c} xy \\ \ln(y) \\ x^2p \end{array}\right)\quad\left(\begin{array}{c} x \\ y \\ p \end{array}\right)=\left(\begin{array}{c} Re^{-S} \\ e^S \\ Te^{2S}/R^2 \end{array}\right)$	$\left(\begin{array}{c} R \\ S \\ T \end{array}\right)=\left(\begin{array}{c} xy \\ \ln(y) \\ x^2p \end{array}\right)\quad\left(\begin{array}{c} x \\ y \\ p \end{array}\right)=\left(\begin{array}{c} Re^{-S} \\ e^S \\ Te^{2S}/R^2 \end{array}\right)$	$\left(\begin{array}{c} R \\ S \\ T \end{array}\right)=\left(\begin{array}{c} xy \\ \ln(y) \\ x^2p \end{array}\right)\quad\left(\begin{array}{c} x \\ y \\ p \end{array}\right)=\left(\begin{array}{c} Re^{-S} \\ e^S \\ Te^{2S}/R^2 \end{array}\right)$
gilmore-lie-groups_ EQ0674	217	■0.973	$d\,s^{\mathrm{2}}=\left[\begin{array}{ll} d\,x&d\,y \end{array}\right]\left[\begin{array}{cc} \frac{1}{y}&0 \\ 0&\frac{1}{y} \end{array}\right]\left[\begin{array}{c} dx \\ dy \end{array}\right]=\frac{d\bar{z}dz}{y^2}$	$ds^2 = \left[\begin{array}{cc} dx & dy \end{array}\right] \left[\begin{array}{cc} \frac{1}{y} & 0 \\ 0 & \frac{1}{y} \end{array}\right] \left[\begin{array}{c} dx \\ dy \end{array}\right] = \frac{d\bar{z}dz}{y^2}$	$ds^2 = \left[\begin{array}{cc} dx & dy \end{array}\right] \left[\begin{array}{cc} \frac{1}{y} & 0 \\ 0 & \frac{1}{y} \end{array}\right] \left[\begin{array}{c} dx \\ dy \end{array}\right] = \frac{d\bar{z}dz}{y^2}$
gilmore-lie-groups_ EQ0218 (4.28)	77	■0.976	$\left[X_{\mathrm{i}},\,X_{\mathrm{j}}\right]=R\left(X_{\mathrm{i}}\right)_{\mathrm{j}}^{\mathrm{k}}X_{\mathrm{k}}$ $X_{\mathrm{k}}=C_{\mathrm{i}\,\mathrm{j}}^{\mathrm{k}}X_{\mathrm{k}}$	$[X_i,X_j]=R\left(X_i\right)_j^kX_k=C_{ij}^kX_k$	$[X_i,X_j]=R\left(X_i\right)_j^kX_k=C_{ij}^kX_k$
gilmore-lie-groups_ EQ0469	141	■0.976	$\mathrm{mathcal{H}}_{\mathrm{atoms}}=\epsilon J_z+\frac{\lambda}{\sqrt{N}}\left(J_+\langle a(t)\rangle+J_-\langle a(t)\rangle^*\right)$	$\mathcal{H}_{\mathrm{atoms}}=\epsilon J_z+\frac{\lambda}{\sqrt{N}}\left(J_+\langle a(t)\rangle+J_-\langle a(t)\rangle^*\right)$	$\mathcal{H}_{\mathrm{atoms}}=\epsilon J_z+\frac{\lambda}{\sqrt{N}}\left(J_+\langle a(t)\rangle+J_-\langle a(t)\rangle^*\right)$
gilmore-lie-groups_ EQ0509 (9.18)	157	■0.976	$M: \sum a^{\mathrm{i}}_{\mathrm{j}}\,M_{\mathrm{i}}^{\mathrm{j}}$	$M: \sum a^i_j M_i^j$	$M: \sum a^i_j M_i^j$
gilmore-lie-groups_ EQ9906 (15.40)	282	■0.976	$\left(\begin{array}{cccc} 0&+\theta_3&-\theta_2&ib_1 \\ -\theta_3&0&+\theta_1&ib_2 \\ +\theta_2&-\theta_1&0&ib_3 \\ -ib_1&-ib_2&-ib_3&0 \end{array}\right)\left[\begin{array}{c} 0 \\ 0 \\ 0 \\ i \end{array}\right]=\left[\begin{array}{c} -b_1 \\ -b_2 \\ -b_3 \\ 0 \end{array}\right]=\left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array}\right]$	$\left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & ib_1 \\ -\theta_3 & 0 & +\theta_1 & ib_2 \\ +\theta_2 & -\theta_1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{array}\right] \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ i \end{array}\right] = \left[\begin{array}{c} -b_1 \\ -b_2 \\ -b_3 \\ 0 \end{array}\right] = \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array}\right]$	$\left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & ib_1 \\ -\theta_3 & 0 & +\theta_1 & ib_2 \\ +\theta_2 & -\theta_1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{array}\right] \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ i \end{array}\right] = \left[\begin{array}{c} -b_1 \\ -b_2 \\ -b_3 \\ 0 \end{array}\right] = \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array}\right]$
gilmore-lie-groups_ EQ0347	108	■0.978	$\begin{aligned} &J_z=\frac{1}{2}\left(a_1^{\dagger}a_1-a_2^{\dagger}a_2\right) \\ &a_1-a_2^{\dagger}a_2\right)\,\,\&J_x=\frac{1}{2}\left(J_++J_-\right)=\frac{1}{2}\left(a_1^{\dagger}a_2+a_2^{\dagger}a_1\right) \\ &\left(J_++J_-\right)=\frac{1}{2}\left(a_1^{\dagger}a_2+a_2^{\dagger}a_1\right)\,\,\&J_y=\frac{1}{2i}\left(J_+-J_-\right)=\frac{1}{2i}\left(a_1^{\dagger}a_2-a_2^{\dagger}a_1\right) \end{aligned}$	$J_z = \frac{1}{2}\left(a_1^{\dagger}a_1 - a_2^{\dagger}a_2\right)$ $J_x = \frac{1}{2}\left(J_+ + J_-\right) = \frac{1}{2}\left(a_1^{\dagger}a_2 + a_2^{\dagger}a_1\right)$ $J_y = \frac{1}{2i}\left(J_+ - J_-\right) = \frac{1}{2i}\left(a_1^{\dagger}a_2 - a_2^{\dagger}a_1\right)$	$J_z = \frac{1}{2}(a_1^{\dagger}a_1 - a_2^{\dagger}a_2)$ $J_x = \frac{1}{2}(J_+ + J_-) = \frac{1}{2}(a_1^{\dagger}a_2 + a_2^{\dagger}a_1)$ $J_y = \frac{1}{2i}(J_+ - J_-) = \frac{1}{2i}(a_1^{\dagger}a_2 - a_2^{\dagger}a_1)$
gilmore-lie-groups_ EQ0270 (5.9)	91	■0.979	$\left(\begin{array}{lll} 0&l&\delta \\ 0&\eta&r \\ 0&0&0 \end{array}\right)=\eta X_{\eta}+lX_l+rX_r+\delta X_{\delta}$	$\left[\begin{array}{ccc} 0 & l & \delta \\ 0 & \eta & r \\ 0 & 0 & 0 \end{array}\right] = \eta X_{\eta} + lX_l + rX_r + \delta X_{\delta}$	$\left[\begin{array}{ccc} 0 & l & \delta \\ 0 & \eta & r \\ 0 & 0 & 0 \end{array}\right] = \eta X_{\eta} + lX_l + rX_r + \delta X_{\delta}$
gilmore-lie-groups_ EQ0435	132	■0.979	$M(S)\left(\begin{array}{ccc} 1&Z_l&* \\ 0&* & Z_r \\ 0&0&1 \end{array}\right)M\left(S^{-1}\right)=\left(\begin{array}{ccc} 1&0&B \\ 0&A&0 \\ 0&0&1 \end{array}\right)$	$M(S)\left[\begin{array}{ccc} 1 & Z_l & * \\ 0 & * & Z_r \\ 0 & 0 & 1 \end{array}\right]M\left(S^{-1}\right)=\left[\begin{array}{ccc} 1 & 0 & B \\ 0 & A & 0 \\ 0 & 0 & 1 \end{array}\right]$	$M(S)\left[\begin{array}{ccc} 1 & Z_l & * \\ 0 & * & Z_r \\ 0 & 0 & 1 \end{array}\right]M(S^{-1})=\left[\begin{array}{ccc} 1 & 0 & B \\ 0 & A & 0 \\ 0 & 0 & 1 \end{array}\right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0976 (16.19)	304	■ 0.979	$\begin{array}{lr} \text{surface equation} & F(R,-,T)=0 \\ \text{constraint equation} & dS/dR=f(R,-,T) \end{array}$	surface equation constraint equation $F(R,-,T)=0$ $dS/dR=f(R,-,T)$	surface equation constraint equation $F(R,-,T)=0$ $dS/dR=f(R,-,T)$
gilmore-lie-groups_ EQ0649 (12.18)	210	■ 0.981	$\begin{array}{cc} S_{0(2+1)} / S_{0(2)} & \& S_{0(2,1)} / S_{0(2)} \\ g_{ij} & \rightarrow \delta_{ij} + x^i x^j \end{array}$	$SO(2+1)/SO(2)$ $SO(2,1)/SO(2)$ $g_{ij} \rightarrow \delta_{ij} + x^i x^j \rightarrow g_{ij} \rightarrow \delta_{ij} - x^i x^j$	$SO(2+1)/SO(2)$ $SO(2,1)/SO(2)$ $g_{ij} \rightarrow \delta_{ij} + x^i x^j \rightarrow g_{ij} \rightarrow \delta_{ij} - x^i x^j$
gilmore-lie-groups_ EQ0462 (7.88)	139	■ 0.981	$\left\langle \begin{array}{c} j \\ \theta'_x \theta'_y \end{array} \middle \begin{array}{c} j \\ \theta_x \theta_y \end{array} \right\rangle = \left[\cos\left(\frac{\theta'}{2}\right) \cos\left(\frac{\theta}{2}\right) + e^{i(\phi' - \phi)} \sin\left(\frac{\theta'}{2}\right) \sin\left(\frac{\theta}{2}\right) \right]^{2j}$	$\left\langle \begin{array}{c} j \\ \theta'_x \theta'_y \end{array} \middle \begin{array}{c} j \\ \theta_x \theta_y \end{array} \right\rangle = \left[\cos\left(\frac{\theta'}{2}\right) \cos\left(\frac{\theta}{2}\right) + e^{i(\phi' - \phi)} \sin\left(\frac{\theta'}{2}\right) \sin\left(\frac{\theta}{2}\right) \right]^{2j}$	$\left\langle \begin{array}{c} j \\ \theta'_x \theta'_y \end{array} \middle \begin{array}{c} j \\ \theta_x \theta_y \end{array} \right\rangle = \left[\cos\left(\frac{\theta'}{2}\right) \cos\left(\frac{\theta}{2}\right) + e^{i(\phi' - \phi)} \sin\left(\frac{\theta'}{2}\right) \sin\left(\frac{\theta}{2}\right) \right]^{2j}$
gilmore-lie-groups_ EQ0183	67	■ 0.981	$\left[\begin{array}{l} o_1 \\ o_2 \\ o_3 \\ o_4 \end{array} \right] \left[\begin{array}{cc cc} s_{11} & s_{12} & 0 & 0 \\ s_{21} & s_{22} & 0 & 0 \\ \hline 0 & 0 & s_{33} & s_{34} \\ 0 & 0 & s_{43} & s_{44} \end{array} \right] \left[\begin{array}{l} i_1 \\ i_2 \\ i_3 \\ i_4 \end{array} \right]$	$\left[\begin{array}{l} o_1 \\ o_2 \\ o_3 \\ o_4 \end{array} \right] \left[\begin{array}{cc cc} s_{11} & s_{12} & 0 & 0 \\ s_{21} & s_{22} & 0 & 0 \\ \hline 0 & 0 & s_{33} & s_{34} \\ 0 & 0 & s_{43} & s_{44} \end{array} \right] \left[\begin{array}{l} i_1 \\ i_2 \\ i_3 \\ i_4 \end{array} \right]$	$\left[\begin{array}{l} o_1 \\ o_2 \\ o_3 \\ o_4 \end{array} \right] \left[\begin{array}{cc cc} s_{11} & s_{12} & 0 & 0 \\ s_{21} & s_{22} & 0 & 0 \\ \hline 0 & 0 & s_{33} & s_{34} \\ 0 & 0 & s_{43} & s_{44} \end{array} \right] \left[\begin{array}{l} i_1 \\ i_2 \\ i_3 \\ i_4 \end{array} \right]$
gilmore-lie-groups_ EQ0498 (9.7)	155	■ 0.981	$\left \Re e \, g \left(r a^{\dagger} + l a + \delta I \right) - \lambda I_3 \right = (-\lambda)^3 = 0$	$\ \Re e \, g \left(r a^{\dagger} + l a + \delta I \right) - \lambda I_3 \ = (-\lambda)^3 = 0$	$\ \Re e \, g \left(r a^{\dagger} + l a + \delta I \right) - \lambda I_3 \ = (-\lambda)^3 = 0$
gilmore-lie-groups_ EQ0626	201	■ 0.982	$\begin{array}{l} m=+ j +1,+ j +2,\ldots \\ m=- j -1,- j -2,\ldots \end{array}$	$m=+ j +1,+ j +2,\ldots$ $m=- j -1,- j -2,\ldots$	$m=+ j +1,+ j +2,\ldots$ $m=- j -1,- j -2,\ldots$
gilmore-lie-groups_ EQ0934 (15.69)	288	■ 0.982	$\left[\begin{array}{ccc} +\frac{i}{c} \frac{\partial}{\partial t} & +\partial_3 & -\partial_2 \\ -\partial_3 & +\frac{i}{c} \frac{\partial}{\partial t} & +\partial_1 \\ +\partial_2 & -\partial_1 & +\frac{i}{c} \frac{\partial}{\partial t} \end{array} \right] \left[\begin{array}{l} B_1 - i E_1 \\ B_2 - i E_2 \\ B_3 - i E_3 \end{array} \right] = 0$	$\left[\begin{array}{ccc} +\frac{i}{c} \frac{\partial}{\partial t} & +\partial_3 & -\partial_2 \\ -\partial_3 & +\frac{i}{c} \frac{\partial}{\partial t} & +\partial_1 \\ +\partial_2 & -\partial_1 & +\frac{i}{c} \frac{\partial}{\partial t} \end{array} \right] \left[\begin{array}{l} B_1 - i E_1 \\ B_2 - i E_2 \\ B_3 - i E_3 \end{array} \right] = 0$	$\left[\begin{array}{ccc} +\frac{i}{c} \frac{\partial}{\partial t} & +\partial_3 & -\partial_2 \\ -\partial_3 & +\frac{i}{c} \frac{\partial}{\partial t} & +\partial_1 \\ +\partial_2 & -\partial_1 & +\frac{i}{c} \frac{\partial}{\partial t} \end{array} \right] \left[\begin{array}{l} B_1 - i E_1 \\ B_2 - i E_2 \\ B_3 - i E_3 \end{array} \right] = 0$
gilmore-lie-groups_ EQ0126 (3.13)	53	■ 0.982	$\left[\begin{array}{ll} 1 & a \\ 0 & 1 \end{array} \right] \left[\begin{array}{ll} 1 & b \\ 0 & 1 \end{array} \right] = \left[\begin{array}{ll} 1 & a+b \\ 0 & 1 \end{array} \right] = \left[\begin{array}{ll} 1 & b \\ 0 & 1 \end{array} \right] \left[\begin{array}{ll} 1 & a \\ 0 & 1 \end{array} \right]$	$\left[\begin{array}{ll} 1 & a \\ 0 & 1 \end{array} \right] \left[\begin{array}{ll} 1 & b \\ 0 & 1 \end{array} \right] = \left[\begin{array}{ll} 1 & a+b \\ 0 & 1 \end{array} \right] = \left[\begin{array}{ll} 1 & b \\ 0 & 1 \end{array} \right] \left[\begin{array}{ll} 1 & a \\ 0 & 1 \end{array} \right]$	$\left[\begin{array}{ll} 1 & a \\ 0 & 1 \end{array} \right] \left[\begin{array}{ll} 1 & b \\ 0 & 1 \end{array} \right] = \left[\begin{array}{ll} 1 & a+b \\ 0 & 1 \end{array} \right] = \left[\begin{array}{ll} 1 & b \\ 0 & 1 \end{array} \right] \left[\begin{array}{ll} 1 & a \\ 0 & 1 \end{array} \right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E01036	321	■ 0.982	$\begin{aligned} & \int d\rho(g)\left\langle \begin{array}{c} \lambda' \\ \mu'\nu' \end{array} \middle g \right\rangle \left\langle g \middle \begin{array}{c} \lambda \\ \mu\nu \end{array} \right\rangle = \left\langle \begin{array}{c} \lambda' \\ \mu'\nu' \end{array} \middle \begin{array}{c} \lambda \\ \mu\nu \end{array} \right\rangle \\ & \sum_{\lambda} \sum_{\mu} \sum_{\nu} \left\langle g' \middle \begin{array}{c} \lambda \\ \mu\nu \end{array} \right\rangle \left\langle \begin{array}{c} \lambda \\ \mu\nu \end{array} \middle g \right\rangle = \langle g' g \rangle \end{aligned}$	$\int d\rho(g)\left\langle \begin{array}{c} \lambda' \\ \mu'\nu' \end{array} \middle g \right\rangle \left\langle g \middle \begin{array}{c} \lambda \\ \mu\nu \end{array} \right\rangle = \left\langle \begin{array}{c} \lambda' \\ \mu'\nu' \end{array} \middle \begin{array}{c} \lambda \\ \mu\nu \end{array} \right\rangle$ $\sum_{\lambda} \sum_{\mu} \sum_{\nu} \left\langle g' \middle \begin{array}{c} \lambda \\ \mu\nu \end{array} \right\rangle \left\langle \begin{array}{c} \lambda \\ \mu\nu \end{array} \middle g \right\rangle = \langle g' g \rangle$	$\int d\rho(g)\left\langle \begin{array}{c} \lambda' \\ \mu'\nu' \end{array} \middle g \right\rangle \left\langle g \middle \begin{array}{c} \lambda \\ \mu\nu \end{array} \right\rangle = \left\langle \begin{array}{c} \lambda' \\ \mu'\nu' \end{array} \middle \begin{array}{c} \lambda \\ \mu\nu \end{array} \right\rangle$ $\sum_{\lambda} \sum_{\mu} \sum_{\nu} \left\langle g' \middle \begin{array}{c} \lambda \\ \mu\nu \end{array} \right\rangle \left\langle \begin{array}{c} \lambda \\ \mu\nu \end{array} \middle g \right\rangle = \langle g' g \rangle$
gilmore-lie-groups_ E00400 (7.36)	124	■ 0.983	$\left[\begin{array}{lll} 0 & l & \delta \\ 0 & \eta & r \\ 0 & 0 & 0 \end{array}\right] = \eta X_{\eta} + lX_l + rX_r + \delta X_{\delta}$	$\left[\begin{array}{ccc} 0 & l & \delta \\ 0 & \eta & r \\ 0 & 0 & 0 \end{array}\right] = \eta X_{\eta} + lX_l + rX_r + \delta X_{\delta}$	$\left[\begin{array}{ccc} 0 & l & \delta \\ 0 & \eta & r \\ 0 & 0 & 0 \end{array}\right] = \eta X_{\eta} + lX_l + rX_r + \delta X_{\delta}$
gilmore-lie-groups_ E00634	207	■ 0.983	$\begin{aligned} & dV = \rho(\mathrm{Id}) dx^1(\mathrm{Id}) \wedge dx^2(\mathrm{Id}) \wedge \cdots \wedge dx^n(\mathrm{Id}) \\ & = \rho(p) dx^1(p) \wedge dx^2(p) \wedge \cdots \wedge dx^n(p) \\ & \Rightarrow \rho(p) = \ M(p)\ \rho(\mathrm{Id}) \sim \sqrt{\det g(p)} \end{aligned}$	$dV = \rho(\mathrm{Id}) dx^1(\mathrm{Id}) \wedge dx^2(\mathrm{Id}) \wedge \cdots \wedge dx^n(\mathrm{Id})$ $= \rho(p) dx^1(p) \wedge dx^2(p) \wedge \cdots \wedge dx^n(p)$ $\Rightarrow \rho(p) = \ M(p)\ \rho(\mathrm{Id}) \sim \sqrt{\det g(p)}$	$dV = \rho(\mathrm{Id}) dx^1(\mathrm{Id}) \wedge dx^2(\mathrm{Id}) \wedge \cdots \wedge dx^n(\mathrm{Id})$ $= \rho(p) dx^1(p) \wedge dx^2(p) \wedge \cdots \wedge dx^n(p)$ $\Rightarrow \rho(p) = \ M(p)\ \rho(\mathrm{Id}) \sim \sqrt{\det g(p)}$
gilmore-lie-groups_ E00656	213	■ 0.984	$\operatorname{det} \left \operatorname{Re} e g \left(\frac{p}{\lambda} \right) - \lambda I_3 \right = -\lambda \left(\lambda^2 + b_1^2 + b_2^2 \right) = 0$	$\det \Re g(\mathfrak{p}) - \lambda I_3 = -\lambda \left[\lambda^2 + (b_1^2 + b_2^2) \right] = 0$	$\det \Re g(\mathfrak{p}) - \lambda I_3 = -\lambda \left[\lambda^2 + (b_1^2 + b_2^2) \right] = 0$
gilmore-lie-groups_ E00623	201	■ 0.986	$j+\frac{1}{2}=i\beta \quad \beta \text{ real} \quad m=0,\pm 1,\pm 2,\dots$	$j+\frac{1}{2}=i\beta \quad \beta \text{ real} \quad m=0,\pm 1,\pm 2,\dots$	$j+\frac{1}{2}=i\beta \quad \beta \text{ real} \quad m=0,\pm 1,\pm 2,\dots$
gilmore-lie-groups_ E00810 (14.48)	254	■ 0.987	$\left[\begin{array}{l} Q_1 \\ Q_2 \\ Q_3 \\ Q_4 \end{array}\right] = M_{KS} \left[\begin{array}{l} q_1 \\ q_2 \\ q_3 \\ q_4 \end{array}\right] = \left[\begin{array}{cccc} q_1 & -q_2 & -q_3 & q_4 \\ q_2 & q_1 & -q_4 & -q_3 \\ q_3 & q_4 & q_1 & q_2 \\ q_4 & -q_3 & q_2 & -q_1 \end{array}\right] \left[\begin{array}{l} q_1 \\ q_2 \\ q_3 \\ q_4 \end{array}\right]$	$\left[\begin{array}{l} Q_1 \\ Q_2 \\ Q_3 \\ Q_4 \end{array}\right] = M_{KS} \left[\begin{array}{l} q_1 \\ q_2 \\ q_3 \\ q_4 \end{array}\right] = \left[\begin{array}{cccc} q_1 & -q_2 & -q_3 & q_4 \\ q_2 & q_1 & -q_4 & -q_3 \\ q_3 & q_4 & q_1 & q_2 \\ q_4 & -q_3 & q_2 & -q_1 \end{array}\right] \left[\begin{array}{l} q_1 \\ q_2 \\ q_3 \\ q_4 \end{array}\right]$	$\left[\begin{array}{l} Q_1 \\ Q_2 \\ Q_3 \\ Q_4 \end{array}\right] = M_{KS} \left[\begin{array}{l} q_1 \\ q_2 \\ q_3 \\ q_4 \end{array}\right] = \left[\begin{array}{cccc} q_1 & -q_2 & -q_3 & q_4 \\ q_2 & q_1 & -q_4 & -q_3 \\ q_3 & q_4 & q_1 & q_2 \\ q_4 & -q_3 & q_2 & -q_1 \end{array}\right] \left[\begin{array}{l} q_1 \\ q_2 \\ q_3 \\ q_4 \end{array}\right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ1005 (16.48)	313	■ 0.987	\begin{aligned} \eta^{\{0\}}(x,y) &= \eta(x,y) \\ \eta^{\{1\}}(x,y,y^{(1)}) &= D^{(0)}\eta^{\{0\}} - y^{(1)}D^{(0)}\xi \\ \eta^{\{2\}}(x,y,y^{(1)},y^{(2)}) &= D^{(1)}\eta^{\{1\}} - y^{(2)}D^{(0)}\xi \\ \eta^{\{3\}}(x,y,y^{(1)},y^{(2)},y^{(3)}) &= D^{(2)}\eta^{\{2\}} - y^{(3)}D^{(0)}\xi \\ &\vdots \\ &\vdots \end{aligned}	$\eta^{(0)}(x,y) = \eta(x,y)$ $\eta^{(1)}\left(x,y,y^{(1)}\right) = D^{(0)}\eta^{(0)} - y^{(1)}D^{(0)}\xi$ $\eta^{(2)}\left(x,y,y^{(1)},y^{(2)}\right) = D^{(1)}\eta^{(1)} - y^{(2)}D^{(0)}\xi$ $\eta^{(3)}\left(x,y,y^{(1)},y^{(2)},y^{(3)}\right) = D^{(2)}\eta^{(2)} - y^{(3)}D^{(0)}\xi$ $\vdots$ $\vdots$	$\eta^{(0)}(x,y) = \eta(x,y)$ $\eta^{(1)}(x,y,y^{(1)}) = D^{(0)}\eta^{(0)} - y^{(1)}D^{(0)}\xi$ $\eta^{(2)}(x,y,y^{(1)},y^{(2)}) = D^{(1)}\eta^{(1)} - y^{(2)}D^{(0)}\xi$ $\eta^{(3)}(x,y,y^{(1)},y^{(2)},y^{(3)}) = D^{(2)}\eta^{(2)} - y^{(3)}D^{(0)}\xi$ $\vdots$ $\vdots$
gilmore-lie-groups_ EQ0143 (3.26)	56	■ 0.988	\text { Poincaré group }\left[\begin{array}{c c} 0(3,1) & t \\ \hline 0 & 1 \end{array}\right]	Poincaré group $\left[\begin{array}{c c} O(3,1) & t \\ \hline 0 & 1 \end{array}\right]$	Poincaré group $\left[\begin{array}{c c} O(3,1) & t \\ \hline 0 & 1 \end{array}\right]$
gilmore-lie-groups_ EQ0239 (4.49)	81	■ 0.989	d \mu(x)=\rho(x) \, d V(x)=\rho(x)\, M(x) \, d V(\mathrm{Id}) \\ \rightarrow \rho(x)= M(x) ^{-1}	$d\mu(x) = \rho(x)dV(x) = \rho(x)\ M(x)\ dV(\text{Id}) \Rightarrow \rho(x) = \ M(x)\ ^{-1}$	$d\mu(x) = \rho(x)dV(x) = \rho(x) \ M(x) \ dV(\text{Id}) \Rightarrow \rho(x) = \ M(x) \ ^{-1}$
gilmore-lie-groups_ EQ0046	28	■ 0.990	\begin{aligned} & v_2 \quad \& v_3 \end{aligned}=\left\{-\frac{27}{2} I_3^{\prime} \pm \frac{1}{2}\left[\left(27 I_3^{\prime}\right)^2+4 \times 27 I_2^{\prime 3}\right]^{1 / 2}\right\}^{1 / 3}	$v_2 = \left\{-\frac{27}{2}I_3' \pm \frac{1}{2}\left[(27I_3')^2 + 4 \times 27I_2'^3\right]^{1/2}\right\}^{1/3}$	$\frac{v_2}{v_3} = \left\{-\frac{27}{2}I_3' \pm \frac{1}{2}\left[(27I_3')^2 + 4 \times 27I_2'^3\right]^{1/2}\right\}^{1/3}$
gilmore-lie-groups_ EQ0071 (1.66)	34	■ 0.990	\begin{gathered} x^2-1399680 \, x+552983334912=0 \\ K_1=1399680 \quad K_2=552983334912 \end{gathered}	$x^2-1399680x+552983334912=0$ $K_1=1399680$ $K_2=552983334912$	$x^2-1399680x+552983334912=0$ $K_1=1399680$ $K_2=552983334912$
gilmore-lie-groups_ EQ0236 (4.46)	81	■ 0.991	g_{rs}(x) \, dx^r \, dx^s=g_{ij}(\mathrm{Id}) \, d\alpha^i \, d\alpha^j	$g_{rs}(x)dx^r dx^s = g_{ij}(\text{Id})d\alpha^i d\alpha^j$	$g_{rs}(x)dx^r dx^s = g_{ij}(\text{Id})d\alpha^i d\alpha^j$
gilmore-lie-groups_ EQ0608	197	■ 0.991	\operatorname{tr} \operatorname{Re} e \, \mathfrak{g}^2\left(w \, a_1^{\dagger} a_1+x \, a_1^{\dagger} a_2+y \, a_2^{\dagger} a_1+z \, a_2^{\dagger} a_2\right)=2(w-z)^2+8 \, x \, y	$\operatorname{tr} \Re \mathfrak{g}^2\left(w a_1^\dagger a_1+x a_1^\dagger a_2+y a_2^\dagger a_1+z a_2^\dagger a_2\right)=2(w-z)^2+8xy$	$\operatorname{tr} \Re \mathfrak{g}^2(w a_1^\dagger a_1+x a_1^\dagger a_2+y a_2^\dagger a_1+z a_2^\dagger a_2)=2(w-z)^2+8xy$
gilmore-lie-groups_ EQ0113 (3.3)	49	■ 0.991	\begin{array}{cc} c=(a,b)=a+ib & a\in\mathbb{R},\,b\in\mathbb{R},\,i^2=-1 \\ (a,b)\longrightarrow\left[\begin{array}{cc} a & b \\ -b & a \end{array}\right] & i=(0,1)\longrightarrow\left[\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right] \end{array}	$c=(a,b)=a+ib \qquad a\in\mathbb{R},b\in\mathbb{R},i^2=-1$ $(a,b)\longrightarrow\left[\begin{array}{cc} a & b \\ -b & a \end{array}\right] \qquad i=(0,1)\longrightarrow\left[\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right]$	$c=(a,b)=a+ib \qquad a\in\mathbb{R},\,b\in\mathbb{R},\,i^2=-1$ $(a,b)\longrightarrow\left[\begin{array}{cc} a & b \\ -b & a \end{array}\right] \qquad i=(0,1)\longrightarrow\left[\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right]$
gilmore-lie-groups_ EQ0115	49	■ 0.991	q_{\{0\}}+q_{\{1\}}\,\mathcal{I}+q_{\{2\}}\,\mathcal{J}+q_{\{3\}}\,\mathcal{K}\rightarrow\left[\begin{array}{cc} q_0+i\,q_3 & i\,q_1+q_2 \\ i\,q_1-q_2 & q_0-i\,q_3 \end{array}\right]	$q_0+q_1\mathcal{I}+q_2\mathcal{J}+q_3\mathcal{K}\rightarrow\left[\begin{array}{cc} q_0+iq_3 & iq_1+q_2 \\ iq_1-q_2 & q_0-iq_3 \end{array}\right]$	$q_0+q_1\mathcal{I}+q_2\mathcal{J}+q_3\mathcal{K}\rightarrow\left[\begin{array}{cc} q_0+iq_3 & iq_1+q_2 \\ iq_1-q_2 & q_0-iq_3 \end{array}\right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0497 (9.6)	155	■ 0.991	$r\ a^{\dagger}+l\ a+\delta I\ \xrightarrow[\text{representation}]{\text{regular}}\left[\begin{array}{ccc}0&l&0\\0&0&0\\0&-r&0\end{array}\right]\begin{array}{l}a^{\dagger}\\I\\a\end{array}$		
gilmore-lie-groups_ EQ0512 (9.21)	159	■ 0.991	$\mathcal{C}_2[\mathfrak{so}(n)]=\sum X_{ij}^2$		
gilmore-lie-groups_ EQ0261 (4.62b)	87	■ 0.992	$\begin{array}{l}2\ \alpha_1\ &=+m_{11}+\frac{\kappa_R}{\kappa_L}m_{22}\quad 2\ \alpha_2\ &=-m_{12}\kappa_R-\kappa_L^{-1}m_{21}\\2\ \beta_1\ &=+m_{11}-\frac{\kappa_R}{\kappa_L}m_{22}\quad 2\ \beta_2\ &=+m_{12}\kappa_R-\kappa_L^{-1}m_{21}\end{array}$		
gilmore-lie-groups_ EQ0035 (1.32)	26	■ 0.992	$\left[\begin{array}{l}v_1\\v_2\\v_3\end{array}\right]=\left[\begin{array}{lll}1&1&1\\1&\omega&\omega^2\\1&\omega^2&\omega\end{array}\right]\left[\begin{array}{l}s_1\\s_2\\s_3\end{array}\right]=\left[\begin{array}{l}s_1+s_2+s_3\\s_1+\omega s_2+\omega^2 s_3\\s_1+\omega^2 s_2+\omega s_3\end{array}\right]$		
gilmore-lie-groups_ EQ0878 (15.11)	277	■ 0.992	$\begin{aligned}&\&\left[\mathbf{J}_{i}^{(1)},\mathbf{J}_{j}^{(1)}\right]=-\epsilon_{ijk}\mathbf{J}_k^{(1)},\\&\left[\mathbf{J}_{i}^{(2)},\mathbf{J}_{j}^{(2)}\right]=-\epsilon_{ijk}\mathbf{J}_k^{(2)},\\&\left[\mathbf{J}_i^{(1)},\mathbf{J}_j^{(2)}\right]=0\end{aligned}$		
gilmore-lie-groups_ EQ0668	215	■ 0.992	$M=\left[\begin{array}{c cc}m_{ij}&&\\ \hline &0&-\frac{1}{2}\\ &-\frac{1}{2}&0\end{array}\right]$		
gilmore-lie-groups_ EQ0743	232	■ 0.993	$l^{1/4}P_{n-l,0}^l\left(\frac{\pi}{2}-\sqrt{2}\epsilon x\right)\xrightarrow{\lim}\psi_n(x)=N_nH_n(x)e^{-x^2/2}$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0188	68	■ 0.993	$\begin{array}{l} H_2 = \frac{H_1 + I_n}{H_1 - I_n} \\ \& \left[H_1, H_2 \right] = 0 \\ \& H_1 = \frac{H_2 + I_n}{H_2 - I_n} \\ U = \frac{I_n + iH}{I_n - iH} \\ \& [H, U] = 0 \\ \& H = i \frac{I_n - U}{I_n + U} \\ A = \frac{I_n + iU}{I_n - iU} \\ \& [U, A] = 0 \\ \& U = i \frac{I_n - A}{I_n + A} \\ H = \frac{I_n - iA}{I_n + iA} \\ \& [A, H] = 0 \\ \& A = i \frac{H - I_n}{H + I_n} \end{array}$	$\begin{array}{l} H_2 = \frac{H_1 + I_n}{H_1 - I_n} \\ U = \frac{I_n + iH}{I_n - iH} \\ A = \frac{I_n + iU}{I_n - iU} \\ H = \frac{I_n - iA}{I_n + iA} \end{array} \quad \begin{array}{l} [H_1, H_2] = 0 \\ [H, U] = 0 \\ [U, A] = 0 \\ [A, H] = 0 \end{array} \quad \begin{array}{l} H_1 = \frac{H_2 + I_n}{H_2 - I_n} \\ H = i \frac{I_n - U}{I_n + U} \\ U = i \frac{I_n - A}{I_n + A} \\ A = i \frac{H - I_n}{H + I_n} \end{array}$	$\begin{array}{l} H_2 = \frac{H_1 + I_n}{H_1 - I_n} \\ U = \frac{I_n + iH}{I_n - iH} \\ A = \frac{I_n + iU}{I_n - iU} \\ H = \frac{I_n - iA}{I_n + iA} \end{array} \quad \begin{array}{l} [H_1, H_2] = 0 \\ [H, U] = 0 \\ [U, A] = 0 \\ [A, H] = 0 \end{array} \quad \begin{array}{l} H_1 = \frac{H_2 + I_n}{H_2 - I_n} \\ H = i \frac{I_n - U}{I_n + U} \\ U = i \frac{I_n - A}{I_n + A} \\ A = i \frac{H - I_n}{H + I_n} \end{array}$
gilmore-lie-groups_ EQ0386 (7.23)	121	■ 0.993	$g \in G, \quad d_i \in D, \quad G \text{ simple, } \quad gd_i = d_i g \Rightarrow d_i = \lambda I_n$	$g \in G, \quad d_i \in D, \quad G \text{ simple, } \quad gd_i = d_i g \Rightarrow d_i = \lambda I_n$	$g \in G, \quad d_i \in D, \quad G \text{ simple, } \quad gd_i = d_i g \Rightarrow d_i = \lambda I_n$
gilmore-lie-groups_ EQ0826 (14.61)	258	■ 0.993	$\begin{aligned} M_4 &= \mathbf{r} \cdot \mathbf{p} - i \\ \Gamma_4 &= \frac{1}{2} \left\{ \left(r \mathbf{p} \cdot \mathbf{p} - r - \frac{\alpha^2}{r} - \frac{i \alpha \gamma \cdot \mathbf{r}}{r^2} \right) \right\} \\ \Gamma_5 &= \frac{1}{2} \left\{ \left(r \mathbf{p} \cdot \mathbf{p} + r - \frac{\alpha^2}{r} - \frac{i \alpha \gamma \cdot \mathbf{r}}{r^2} \right) \right\} \end{aligned}$	$\begin{aligned} M_4 &= \mathbf{r} \cdot \mathbf{p} - i \\ \Gamma_4 &= \frac{1}{2} \left\{ \left(r \mathbf{p} \cdot \mathbf{p} - r - \frac{\alpha^2}{r} - \frac{i \alpha \gamma \cdot \mathbf{r}}{r^2} \right) \right\} \\ \Gamma_5 &= \frac{1}{2} \left\{ \left(r \mathbf{p} \cdot \mathbf{p} + r - \frac{\alpha^2}{r} - \frac{i \alpha \gamma \cdot \mathbf{r}}{r^2} \right) \right\} \end{aligned}$	$\begin{aligned} M_4 &= \mathbf{r} \cdot \mathbf{p} - i \\ \Gamma_4 &= \frac{1}{2} \left\{ \left(r \mathbf{p} \cdot \mathbf{p} - r - \frac{\alpha^2}{r} - \frac{i \alpha \gamma \cdot \mathbf{r}}{r^2} \right) \right\} \\ \Gamma_5 &= \frac{1}{2} \left\{ \left(r \mathbf{p} \cdot \mathbf{p} + r - \frac{\alpha^2}{r} - \frac{i \alpha \gamma \cdot \mathbf{r}}{r^2} \right) \right\} \end{aligned}$
gilmore-lie-groups_ EQ0328 (6.13)	104	■ 0.994	$\mathfrak{so}(3) = \left[\begin{array}{ccc} 0 & \theta_3 & -\theta_2 \\ -\theta_3 & 0 & \theta_1 \\ \theta_2 & -\theta_1 & 0 \end{array} \right] = \theta \cdot \mathbf{L}$	$\mathfrak{so}(3) = \left[\begin{array}{ccc} 0 & \theta_3 & -\theta_2 \\ -\theta_3 & 0 & \theta_1 \\ \theta_2 & -\theta_1 & 0 \end{array} \right] = \theta \cdot \mathbf{L}$	$\mathfrak{so}(3) = \left[\begin{array}{ccc} 0 & \theta_3 & -\theta_2 \\ -\theta_3 & 0 & \theta_1 \\ \theta_2 & -\theta_1 & 0 \end{array} \right] = \theta \cdot \mathbf{L}$
gilmore-lie-groups_ EQ0456 (7.82)	138	■ 0.994	$\left[\begin{array}{ccc} 1 & \beta' & \delta \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{array} \right]$	$\left[\begin{array}{ccc} 1 & \beta' & \delta \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{array} \right]$	$\left[\begin{array}{ccc} 1 & \beta' & \delta \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{array} \right]$
gilmore-lie-groups_ EQ0882 (15.15)	278	■ 0.994	$\begin{aligned} D^{j0} [\theta \cdot \mathbf{J} + \mathbf{b} \cdot \mathbf{K}] &= \text{EXP} \left[\theta \cdot \mathbf{J}^{(j)} + \mathbf{b} \cdot (+i\mathbf{J})^{(j)} \right] = \text{EXP} \left[(\theta + i\mathbf{b}) \cdot \mathbf{J}^{(j)} \right] \\ D^{0j} [\theta \cdot \mathbf{J} + \mathbf{b} \cdot \mathbf{K}] &= \text{EXP} \left[\theta \cdot \mathbf{J}^{(j)} + \mathbf{b} \cdot (-i\mathbf{J})^{(j)} \right] = \text{EXP} \left[(\theta - i\mathbf{b}) \cdot \mathbf{J}^{(j)} \right] \end{aligned}$	$\begin{aligned} D^{j0} [\theta \cdot \mathbf{J} + \mathbf{b} \cdot \mathbf{K}] &= \text{EXP} \left[\theta \cdot \mathbf{J}^{(j)} + \mathbf{b} \cdot (+i\mathbf{J})^{(j)} \right] = \text{EXP} \left[(\theta + i\mathbf{b}) \cdot \mathbf{J}^{(j)} \right] \\ D^{0j} [\theta \cdot \mathbf{J} + \mathbf{b} \cdot \mathbf{K}] &= \text{EXP} \left[\theta \cdot \mathbf{J}^{(j)} + \mathbf{b} \cdot (-i\mathbf{J})^{(j)} \right] = \text{EXP} \left[(\theta - i\mathbf{b}) \cdot \mathbf{J}^{(j)} \right] \end{aligned}$	$\begin{aligned} D^{j0} [\theta \cdot \mathbf{J} + \mathbf{b} \cdot \mathbf{K}] &= \text{EXP} \left[\theta \cdot \mathbf{J}^{(j)} + \mathbf{b} \cdot (+i\mathbf{J})^{(j)} \right] = \text{EXP} \left[(\theta + i\mathbf{b}) \cdot \mathbf{J}^{(j)} \right] \\ D^{0j} [\theta \cdot \mathbf{J} + \mathbf{b} \cdot \mathbf{K}] &= \text{EXP} \left[\theta \cdot \mathbf{J}^{(j)} + \mathbf{b} \cdot (-i\mathbf{J})^{(j)} \right] = \text{EXP} \left[(\theta - i\mathbf{b}) \cdot \mathbf{J}^{(j)} \right] \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0823	257	■0.994	$\left[\begin{array}{cccc cc} 0 & J_3 & -J_2 & M_1 & A_1 & \Gamma_1 \\ -J_3 & 0 & J_1 & M_2 & A_2 & \Gamma_2 \\ J_2 & -J_1 & 0 & M_3 & A_3 & \Gamma_3 \\ -M_1 & -M_2 & -M_3 & 0 & A_4 & \Gamma_4 \\ \hline A_1 & A_2 & A_3 & A_4 & 0 & \Gamma_5 \\ \Gamma_1 & \Gamma_2 & \Gamma_3 & \Gamma_4 & -\Gamma_5 & 0 \end{array}\right] +$	$\left[\begin{array}{cccc cc} 0 & J_3 & -J_2 & M_1 & A_1 & \Gamma_1 \\ -J_3 & 0 & J_1 & M_2 & A_2 & \Gamma_2 \\ J_2 & -J_1 & 0 & M_3 & A_3 & \Gamma_3 \\ -M_1 & -M_2 & -M_3 & 0 & A_4 & \Gamma_4 \\ \hline A_1 & A_2 & A_3 & A_4 & 0 & \Gamma_5 \\ \Gamma_1 & \Gamma_2 & \Gamma_3 & \Gamma_4 & -\Gamma_5 & 0 \end{array}\right] +$	
gilmore-lie-groups_ EQ0888 (15.21)	279	■0.994	$\begin{aligned} \{I, a\} k \rangle &= k \rangle e^{ik \cdot a} \\ \{I, a\} \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle &= \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle \end{aligned}$	$\{I, a\} k \rangle = k \rangle e^{ik \cdot a}$ $\{I, a\} \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle = \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle$	$\{I, a\} k \rangle = k \rangle e^{ik \cdot a}$ $\{I, a\} \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle = \left \begin{array}{cc} j & j' \\ \mu & \mu' \end{array} \right\rangle$
gilmore-lie-groups_ EQ1010 (16.53)	314	■0.995	$\int_V k \nabla^2 x ^{2-n} dV = \int_{S=\partial V} k(2-n) \frac{\hat{\mathbf{n}} \cdot d\mathbf{S}}{ x ^{n-1}} = (2-n)kV(S^n) = 1$	$\int_V k \nabla^2 x ^{2-n} dV = \int_{S=\partial V} k(2-n) \frac{\hat{\mathbf{n}} \cdot d\mathbf{S}}{ x ^{n-1}} = (2-n)kV(S^n) = 1$	$\int_V k \nabla^2 x ^{2-n} dV = \int_{S=\partial V} k(2-n) \frac{\hat{\mathbf{n}} \cdot d\mathbf{S}}{ x ^{n-1}} = (2-n)kV(S^n) = 1$
gilmore-lie-groups_ EQ0194 (4.7)	72	■0.995	$\begin{aligned} \operatorname{EXP}(X) &= \sum_{n=0}^{\infty} \frac{1}{n!} \left[\begin{array}{cc} a & b \\ c & -a \end{array} \right]^n = I_2 \cosh \theta + X \frac{\sinh \theta}{\theta} \\ \operatorname{EXP}(aX_a + bX_b + cX_c) &= \sum_{n=0}^{\infty} \frac{1}{n!} \left[\begin{array}{cc} a & b \\ c & -a \end{array} \right]^n = I_2 \cosh \theta + X \frac{\sinh \theta}{\theta} \\ \cosh \theta + a \sinh(\theta)/\theta &= \cosh \theta - a \sinh(\theta)/\theta \\ c \sinh(\theta)/\theta &= c \sinh(\theta)/\theta \end{aligned}$	$\operatorname{EXP}(X) = \operatorname{EXP}(aX_a + bX_b + cX_c) = \sum_{n=0}^{\infty} \frac{1}{n!} \left[\begin{array}{cc} a & b \\ c & -a \end{array} \right]^n = I_2 \cosh \theta + X \frac{\sinh \theta}{\theta}$ $= \left[\begin{array}{cc} \cosh \theta + a \sinh(\theta)/\theta & b \sinh(\theta)/\theta \\ c \sinh(\theta)/\theta & \cosh \theta - a \sinh(\theta)/\theta \end{array} \right]$ $\theta^2 = a^2 + bc$	$\operatorname{EXP}(X) = \operatorname{EXP}(aX_a + bX_b + cX_c) = \sum_{n=0}^{\infty} \frac{1}{n!} \left[\begin{array}{cc} a & b \\ c & -a \end{array} \right]^n = I_2 \cosh \theta + X \frac{\sinh \theta}{\theta}$ $= \left[\begin{array}{cc} \cosh \theta + a \sinh(\theta)/\theta & b \sinh(\theta)/\theta \\ c \sinh(\theta)/\theta & \cosh \theta - a \sinh(\theta)/\theta \end{array} \right]$ $\theta^2 = a^2 + bc$ (4.7)
gilmore-lie-groups_ EQ0549	171	■0.995	$M = \left[\begin{array}{ccc} 0 & \theta_3 & -\theta_2 \\ -\theta_3 & 0 & \theta_1 \\ \theta_2 & -\theta_1 & 0 \end{array}\right] \rightarrow \mathcal{M} = \left[\begin{array}{ccc} 0 & J_3 & -J_2 \\ -J_3 & 0 & J_1 \\ J_2 & -J_1 & 0 \end{array}\right]$	$M = \left[\begin{array}{ccc} 0 & \theta_3 & -\theta_2 \\ -\theta_3 & 0 & \theta_1 \\ \theta_2 & -\theta_1 & 0 \end{array}\right] \rightarrow \mathcal{M} = \left[\begin{array}{ccc} 0 & J_3 & -J_2 \\ -J_3 & 0 & J_1 \\ J_2 & -J_1 & 0 \end{array}\right]$	$M = \left[\begin{array}{ccc} 0 & \theta_3 & -\theta_2 \\ -\theta_3 & 0 & \theta_1 \\ \theta_2 & -\theta_1 & 0 \end{array}\right] \rightarrow \mathcal{M} = \left[\begin{array}{ccc} 0 & J_3 & -J_2 \\ -J_3 & 0 & J_1 \\ J_2 & -J_1 & 0 \end{array}\right]$
gilmore-lie-groups_ EQ0688 (13.6)	220	■0.995	$\begin{aligned} [L_1, L_2] &= -L_3 \\ [L_2, L_3] &= -L_1 \\ [L_3, L_1] &= -L_2 \end{aligned}$	$\begin{aligned} [L_1, L_2] &= -L_3 \\ [L_2, L_3] &= -L_1 \\ [L_3, L_1] &= -L_2 \end{aligned}$	$\begin{aligned} [L_1, L_2] &= -L_3 \\ [L_2, L_3] &= -L_1 \\ [L_3, L_1] &= -L_2 \end{aligned}$
gilmore-lie-groups_ EQ0210 (4.20)	76	■0.996	$[X_i, X_j] + [X_j, X_i] = 0 \quad C_{ij}^k + C_{ji}^k = 0$	$[X_i, X_j] + [X_j, X_i] = 0 \quad C_{ij}^k + C_{ji}^k = 0$	$[X_i, X_j] + [X_j, X_i] = 0 \quad C_{ij}^k + C_{ji}^k = 0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E09366 (7.4)	114	■ 0.996	$\begin{array}{cc} \left\{ \begin{array}{c} -\lambda \\ 0 \end{array} \right\} & \begin{array}{c} -\lambda > 0 \\ -1/\lambda > 0 \end{array} \end{array}$	$\begin{bmatrix} -\lambda & 0 \\ 0 & -1/\lambda \end{bmatrix} \quad \lambda > 1$	$\begin{bmatrix} -\lambda & 0 \\ 0 & -1/\lambda \end{bmatrix} \quad \lambda > 1$
gilmore-lie-groups_ E09589 (11.10)	190	■ 0.996	$\begin{array}{l} \mathfrak{g} = \mathfrak{h} + \mathfrak{p} \\ \mathfrak{h} \cap \mathfrak{p} = \{0\} \\ \mathfrak{h} \cap \mathfrak{p} = \{0\} \\ \mathfrak{h} \cap \mathfrak{p} = \{0\} \end{array}$	$\begin{array}{l} \mathfrak{g} = \mathfrak{h} + \mathfrak{p} \\ [\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h} \quad (\mathfrak{h}, \mathfrak{h}) < 0 \\ [\mathfrak{h}, \mathfrak{p}] \subseteq \mathfrak{p} \quad (\mathfrak{h}, \mathfrak{p}) = 0 \\ [\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h} \quad (\mathfrak{p}, \mathfrak{p}) < 0 \end{array}$	$\begin{array}{l} \mathfrak{g} = \mathfrak{h} + \mathfrak{p} \\ [\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h} \quad (\mathfrak{h}, \mathfrak{h}) < 0 \\ [\mathfrak{h}, \mathfrak{p}] \subseteq \mathfrak{p} \quad (\mathfrak{h}, \mathfrak{p}) = 0 \\ [\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h} \quad (\mathfrak{p}, \mathfrak{p}) < 0 \end{array}$
gilmore-lie-groups_ E09754	234	■ 0.996	$g_{\alpha, \beta} = C_{\alpha, \gamma}^i C_{\beta, i}^{\gamma} + C_{\alpha, i}^{\gamma} C_{\beta, \gamma}^i$	$g_{\alpha, \beta} = C_{\alpha, \gamma}^i C_{\beta, i}^{\gamma} + C_{\alpha, i}^{\gamma} C_{\beta, \gamma}^i$	$g_{\alpha, \beta} = C_{\alpha, \gamma}^i C_{\beta, i}^{\gamma} + C_{\alpha, i}^{\gamma} C_{\beta, \gamma}^i$
gilmore-lie-groups_ E09787 (14.29)	246	■ 0.996	$\left \begin{array}{c} l \\ m \end{array} \right\rangle = \left \begin{array}{cc} j/2 & j/2 \\ m_a & m_b \end{array} \right\rangle \left\langle \begin{array}{cc} j/2 & j/2 \\ m_a & m_b \end{array} \right \begin{array}{c} l \\ m \end{array} \right\rangle$	$\left \begin{array}{c} l \\ m \end{array} \right\rangle = \left \begin{array}{cc} j/2 & j/2 \\ m_a & m_b \end{array} \right\rangle \left\langle \begin{array}{cc} j/2 & j/2 \\ m_a & m_b \end{array} \right \begin{array}{c} l \\ m \end{array} \right\rangle$	$\left \begin{array}{c} l \\ m \end{array} \right\rangle = \left \begin{array}{cc} j/2 & j/2 \\ m_a & m_b \end{array} \right\rangle \left\langle \begin{array}{cc} j/2 & j/2 \\ m_a & m_b \end{array} \right \begin{array}{c} l \\ m \end{array} \right\rangle$
gilmore-lie-groups_ E09856 (1.52)	31	■ 0.996	$\begin{array}{l} w_2 = \pm \sqrt{y_2} \\ w_3 = \pm \sqrt{y_3} \\ w_4 = \pm \sqrt{y_4} \end{array}$	$\begin{array}{l} w_2 = \pm \sqrt{y_2} \\ w_3 = \pm \sqrt{y_3} \\ w_4 = \pm \sqrt{y_4} \end{array}$	$\begin{array}{l} w_2 = \pm \sqrt{y_2} \\ w_3 = \pm \sqrt{y_3} \\ w_4 = \pm \sqrt{y_4} \end{array}$
gilmore-lie-groups_ E09630 (12.3)	205	■ 0.996	$\left \Re \mathfrak{g}(\mathfrak{h} + \mathfrak{p}) - \lambda I_n \right = \sum_{j=0}^n (-\lambda)^{n-j} \phi_j(\mathfrak{h}, \mathfrak{p})$	$\left \Re \mathfrak{g}(\mathfrak{h} + \mathfrak{p}) - \lambda I_n \right = \sum_{j=0}^n (-\lambda)^{n-j} \phi_j(\mathfrak{h}, \mathfrak{p})$	$\left \Re \mathfrak{g}(\mathfrak{h} + \mathfrak{p}) - \lambda I_n \right = \sum_{j=0}^n (-\lambda)^{n-j} \phi_j(\mathfrak{h}, \mathfrak{p})$
gilmore-lie-groups_ E09486 (8.20) [re- fined: census]	149	■ 0.996	$\mathfrak{g} = \underbrace{\underbrace{\underbrace{V_{-}^{\prime}}_{\text{compact subalgebra}} + \underbrace{V_{+}^{\prime}}_{\text{noncompact generators}}}_{\text{nonsemisimple Lie algebra}} + \underbrace{V_0^{\prime} + V_0}_{\text{maximum solvable invariant subalgebra}}}_{\text{nonsemisimple Lie algebra}}$	$\mathfrak{g} = \underbrace{\underbrace{\underbrace{V_{-}^{\prime}}_{\text{compact subalgebra}} + \underbrace{V_{+}^{\prime}}_{\text{noncompact generators}}}_{\text{nonsemisimple Lie algebra}} + \underbrace{V_0^{\prime} + V_0}_{\text{maximum solvable invariant subalgebra}}}_{\text{nonsemisimple Lie algebra}}$	$\mathfrak{g} = \underbrace{\underbrace{\underbrace{V_{-}^{\prime}}_{\text{compact subalgebra}} + \underbrace{V_{+}^{\prime}}_{\text{noncompact generators}}}_{\text{semisimple}} + \underbrace{\underbrace{V_0^{\prime} + V_0}_{\text{abelian nilpotent}}}_{\text{maximum solvable invariant subalgebra}}}_{\text{nonsemisimple Lie algebra}}$
gilmore-lie-groups_ E09879	36	■ 0.997	$\begin{array}{l} I_1 = 0 \quad J_1 = 162 \\ I_2 = -7 \quad J_2 = 9261 \\ I_3 = -6 \end{array}$	$\begin{array}{l} I_1 = 0 \quad J_1 = 162 \\ I_2 = -7 \quad J_2 = 9261 \\ I_3 = -6 \end{array}$	$\begin{array}{l} I_1 = 0 \quad J_1 = 162 \\ I_2 = -7 \quad J_2 = 9261 \\ I_3 = -6 \end{array}$
gilmore-lie-groups_ E09872 (1.67)	34	■ 0.997	$x^{\prime 2} + 63207309312 = 0 \quad K_1^{\prime} = 0 \quad K_2 = 63207309312$	$x^{\prime 2} + 63207309312 = 0 \quad K_1^{\prime} = 0 \quad K_2 = 63207309312$	$x^{\prime 2} + 63207309312 = 0 \quad K_1^{\prime} = 0 \quad K_2 = 63207309312$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00094 (2.6)	43	0.998	$\left[\begin{array}{cc} x_1 & x_2 \\ x_3 & (1+x_2x_3)/x_1 \end{array}\right]^{-1} = \left[\begin{array}{cc} (1+x_2x_3)/x_1 & -x_2 \\ -x_3 & x_1 \end{array}\right]$	$\begin{bmatrix} x_1 & x_2 \\ x_3 & (1+x_2x_3)/x_1 \end{bmatrix}^{-1} = \begin{bmatrix} (1+x_2x_3)/x_1 & -x_2 \\ -x_3 & x_1 \end{bmatrix}$	$\begin{bmatrix} x_1 & x_2 \\ x_3 & (1+x_2x_3)/x_1 \end{bmatrix}^{-1} = \begin{bmatrix} (1+x_2x_3)/x_1 & -x_2 \\ -x_3 & x_1 \end{bmatrix}$
gilmore-lie-groups_ E00301	98	0.998	$\begin{aligned} & \{\frac{h}{p}, \frac{h}{p}\} \subseteq \frac{h}{p} \\ & \{\frac{h}{p}, \frac{p}{p}\} \subseteq \frac{p}{p} \\ & \{\frac{p}{p}, \frac{p}{p}\} \subseteq \frac{h}{p} \end{aligned}$	$\begin{aligned} [h, h] &\subseteq h \\ [h, p] &\subseteq p \\ [p, p] &\subseteq h \end{aligned}$	$\begin{aligned} [h, h] &\subseteq h \\ [h, p] &\subseteq p \\ [p, p] &\subseteq h \end{aligned}$
gilmore-lie-groups_ E00937	290	0.998	$\begin{aligned} & \nabla \times \mathbf{B} - \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} = + \frac{4\pi}{c} \mathbf{J}_e \\ & \nabla \times \mathbf{E} + \frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} = - \frac{4\pi}{c} \mathbf{J}_m \end{aligned}$	$\begin{aligned} \nabla \times \mathbf{B} - \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} &= + \frac{4\pi}{c} \mathbf{J}_e \\ \nabla \times \mathbf{E} + \frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} &= - \frac{4\pi}{c} \mathbf{J}_m \end{aligned}$	$\begin{aligned} \nabla \times \mathbf{B} - \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} &= + \frac{4\pi}{c} \mathbf{J}_e \\ \nabla \times \mathbf{E} + \frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} &= - \frac{4\pi}{c} \mathbf{J}_m \end{aligned}$
gilmore-lie-groups_ E00676	217	0.998	$w \rightarrow w' = \frac{\alpha w + \beta}{\bar{\beta} w + \bar{\alpha}} \quad \left[\begin{array}{cc} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{array} \right] \in SU(1,1), \quad \bar{\alpha}\alpha - \bar{\beta}\beta = 1$	$w \rightarrow w' = \frac{\alpha w + \beta}{\bar{\beta} w + \bar{\alpha}} \quad \left[\begin{array}{cc} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{array} \right] \in SU(1,1), \quad \bar{\alpha}\alpha - \bar{\beta}\beta = 1$	$w \rightarrow w' = \frac{\alpha w + \beta}{\bar{\beta} w + \bar{\alpha}} \quad \left[\begin{array}{cc} \alpha & \beta \\ \bar{\beta} & \bar{\alpha} \end{array} \right] \in SU(1,1), \quad \bar{\alpha}\alpha - \bar{\beta}\beta = 1$
gilmore-lie-groups_ E00063 (1.58)	33	0.998	$\begin{aligned} & w_2^2 = y_2 \rightarrow (-10)^2 = 100 \\ & w_3^2 = y_3 \rightarrow (-4)^2 = 16 \\ & w_4^2 = y_4 \rightarrow (+2)^2 = 4 \end{aligned}$	$\begin{aligned} w_2^2 &= y_2 \rightarrow (-10)^2 = 100 \\ w_3^2 &= y_3 \rightarrow (-4)^2 = 16 \\ w_4^2 &= y_4 \rightarrow (+2)^2 = 4 \end{aligned}$	$\begin{aligned} w_2^2 &= y_2 \rightarrow (-10)^2 = 100 \\ w_3^2 &= y_3 \rightarrow (-4)^2 = 16 \\ w_4^2 &= y_4 \rightarrow (+2)^2 = 4 \end{aligned}$
gilmore-lie-groups_ E00144 (3.27)	56	0.998	$\text{Galilei group} \left[\begin{array}{c c c} O(3) & \mathbf{v} & \mathbf{t} \\ \hline 0 & 1 & t_4 \\ \hline 0 & 0 & 1 \end{array} \right]$	$\text{Galilei group} \left[\begin{array}{c c c} O(3) & \mathbf{v} & \mathbf{t} \\ \hline 0 & 1 & t_4 \\ \hline 0 & 0 & 1 \end{array} \right]$	$\text{Galilei group} \left[\begin{array}{c c c} O(3) & \mathbf{v} & \mathbf{t} \\ \hline 0 & 1 & t_4 \\ \hline 0 & 0 & 1 \end{array} \right]$
gilmore-lie-groups_ E00445	135	0.998	$\begin{aligned} & \langle n' x^4 n \rangle = \frac{d^4}{d\lambda^4} \sum_{p,q,r} \frac{\lambda^{p+2q+r}}{p!q!r!} 2^{-(p/2+2q+r/2)} \langle n' (a^\dagger)^p a^r n \rangle_{\lambda=0} \\ & \rightarrow \sum_{p+2q+r=4} \frac{4!}{p!q!r!} \frac{\langle n' (a^\dagger)^p (a)^r n \rangle}{2^{(p/2+q+r/2)}} \end{aligned}$	$\begin{aligned} \langle n' x^4 n \rangle &= \frac{d^4}{d\lambda^4} \sum_{p,q,r} \frac{\lambda^{p+2q+r}}{p!q!r!} 2^{-(p/2+2q+r/2)} \langle n' (a^\dagger)^p a^r n \rangle_{\lambda=0} \\ &\rightarrow \sum_{p+2q+r=4} \frac{4!}{p!q!r!} \frac{\langle n' (a^\dagger)^p (a)^r n \rangle}{2^{(p/2+q+r/2)}} \end{aligned}$	$\begin{aligned} \langle n' x^4 n \rangle &= \frac{d^4}{d\lambda^4} \sum_{p,q,r} \frac{\lambda^{p+2q+r}}{p!q!r!} 2^{-(p/2+2q+r/2)} \langle n' (a^\dagger)^p a^r n \rangle_{\lambda=0} \\ &\rightarrow \sum_{p+2q+r=4} \frac{4!}{p!q!r!} \frac{\langle n' (a^\dagger)^p (a)^r n \rangle}{2^{(p/2+q+r/2)}} \end{aligned}$
gilmore-lie-groups_ E00452 (7.78)	137	0.998	$S e^{x^j X_j} e^{-\beta a^i X_i} S^{-1} = e^{-\beta d^i(x,a) H_i}$	$S e^{x^j X_j} e^{-\beta a^i X_i} S^{-1} = e^{-\beta d^i(x,a) H_i}$	$S e^{x^j X_j} e^{-\beta a^i X_i} S^{-1} = e^{-\beta d^i(x,a) H_i}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0566	176	■ 0.998	$\begin{aligned} & E_{\{6\}} \pm \mathbf{e}_{\{i\}} \pm \mathbf{e}_{\{j\}} \\ & \frac{1}{2} \underbrace{(\pm \mathbf{e}_1 \pm \mathbf{e}_2 \pm \mathbf{e}_3 \pm \mathbf{e}_4 \pm \mathbf{e}_5)}_{\text{even number of + signs}} \pm \frac{\sqrt{3}}{4} \mathbf{e}_6 \quad 1 \leq i \neq j \leq 5 \end{aligned}$	$E_6 \pm \mathbf{e}_i \pm \mathbf{e}_j$ $\frac{1}{2} \underbrace{(\pm \mathbf{e}_1 \pm \mathbf{e}_2 \pm \mathbf{e}_3 \pm \mathbf{e}_4 \pm \mathbf{e}_5)}_{\text{even number of + signs}} \pm \frac{\sqrt{3}}{4} \mathbf{e}_6 \quad 1 \leq i \neq j \leq 5$	$E_6 \pm \mathbf{e}_i \pm \mathbf{e}_j$ $\frac{1}{2} \underbrace{(\pm \mathbf{e}_1 \pm \mathbf{e}_2 \pm \mathbf{e}_3 \pm \mathbf{e}_4 \pm \mathbf{e}_5)}_{\text{even number of + signs}} \pm \frac{\sqrt{3}}{4} \mathbf{e}_6 \quad 1 \leq i \neq j \leq 5$
gilmore-lie-groups_ EQ0632 (12.5)	207	■ 0.998	$d \, x^{\{i\}}(\mathrm{Id}) = M^i_{\{\mu\}} \, d \, x^{\{\mu\}}(p)$	$dx^i(\mathrm{Id}) = M^i_{\mu} dx^{\mu}(p)$	$dx^i(\mathrm{Id}) = M^i_{\mu} dx^{\mu}(p)$
gilmore-lie-groups_ EQ0926 (15.60)	287	■ 0.998	$R(k)(0,0,k,i\,k)=\left(k_{\{1\}},k_{\{2\}},k_{\{3\}},i\,k_{\{4\}}\right)\,\,\,\,\,\text{quad}\,\,\,\,\,k_{\{1\}}^2+k_{\{2\}}^2+k_{\{3\}}^2=k_{\{4\}}^2=k^2$	$R(k)(0,0,k,ik) = (k_1,k_2,k_3,ik_4) \quad k_1^2 + k_2^2 + k_3^2 = k_4^2 = k^2$	$R(k)(0,0,k,ik) = (k_1,k_2,k_3,ik_4) \quad k_1^2 + k_2^2 + k_3^2 = k_4^2 = k^2$
gilmore-lie-groups_ EQ0966 (16.9)	301	■ 0.998	$\begin{aligned} & F(x,y,p)=0 \rightarrow F(\bar{x}(\epsilon),\bar{y}(\epsilon),\bar{p}(\epsilon)) \stackrel{\epsilon \text{ small}}{\longrightarrow} F(x+\epsilon \xi,y+\epsilon \eta,p+\epsilon \zeta) \\ & = F(x,y,p)+\epsilon \left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} + \zeta \frac{\partial}{\partial p} \right) F(x,y,p) + \text{h.o.t.} \end{aligned}$	$F(x,y,p)=0 \rightarrow F(\bar{x}(\epsilon),\bar{y}(\epsilon),\bar{p}(\epsilon)) \stackrel{\epsilon \text{ small}}{\longrightarrow} F(x+\epsilon \xi,y+\epsilon \eta,p+\epsilon \zeta)$ $= F(x,y,p)+\epsilon \left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} + \zeta \frac{\partial}{\partial p} \right) F(x,y,p) + \text{h.o.t.}$	$F(x,y,p)=0 \rightarrow F(\bar{x}(\epsilon),\bar{y}(\epsilon),\bar{p}(\epsilon)) \stackrel{\epsilon \text{ small}}{\longrightarrow} F(x+\epsilon \xi,y+\epsilon \eta,p+\epsilon \zeta)$ $= F(x,y,p)+\epsilon \left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} + \zeta \frac{\partial}{\partial p} \right) F(x,y,p) + \text{h.o.t.} \quad (1)$
gilmore-lie-groups_ EQ1024	319	■ 0.998	$\begin{aligned} & \xi^1=a_1+a_2x+a_3t+a_4xt \\ & \xi^2=2a_2t+a_4t^2+a_5 \\ & \eta=-\frac{1}{2}a_3xu-a_4\left(\frac{1}{2}t+\frac{1}{4}x^2\right)u+a_6u+h(x,t) \end{aligned}$	$\xi^1=a_1+a_2x+a_3t+a_4xt$ $\xi^2=2a_2t+a_4t^2+a_5$ $\eta=-\frac{1}{2}a_3xu-a_4\left(\frac{1}{2}t+\frac{1}{4}x^2\right)u+a_6u+h(x,t)$	$\xi^1=a_1+a_2x+a_3t+a_4xt$ $\xi^2=2a_2t+a_4t^2+a_5$ $\eta=-\frac{1}{2}a_3xu-a_4(\frac{1}{2}t+\frac{1}{4}x^2)u+a_6u+h(x,t)$
gilmore-lie-groups_ EQ0017 (1.13)	21	■ 0.998	$\begin{array}{lcl} S_3 & \xrightarrow{f} & S_2 \\ I,(123),(321) & \longrightarrow & I \\ (12),(23),(31) & \longrightarrow & (12) \end{array}$	$\begin{array}{lcl} S_3 & \xrightarrow{f} & S_2 \\ I,(123),(321) & \longrightarrow & I \\ (12),(23),(31) & \longrightarrow & (12) \end{array}$	$\begin{array}{lcl} S_3 & \xrightarrow{f} & S_2 \\ I,(123),(321) & \longrightarrow & I \\ (12),(23),(31) & \longrightarrow & (12) \end{array}$
gilmore-lie-groups_ EQ0182	67	■ 0.998	$\left[\begin{array}{c} i_1 \\ o_2 \end{array}\right]=T_{\text{Tot}}\left[\begin{array}{c} o_3 \\ i_4 \end{array}\right] \quad T_{\text{Tot}}=T_1T_2$	$\left[\begin{array}{c} i_1 \\ o_2 \end{array}\right]=T_{\text{Tot}}\left[\begin{array}{c} o_3 \\ i_4 \end{array}\right] \quad T_{\text{Tot}}=T_1T_2$	$\left[\begin{array}{c} i_1 \\ o_2 \end{array}\right]=T_{\text{Tot}}\left[\begin{array}{c} o_3 \\ i_4 \end{array}\right] \quad T_{\text{Tot}}=T_1T_2$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00777 (14.20)	243	■ 0.998	$\begin{gathered} \langle \theta \phi L_- \theta' \phi' \rangle \\ \begin{array}{c} l \\ m \end{array} \end{gathered}$	$\langle \theta \phi L_- \theta' \phi' \rangle \left\langle \theta' \phi' \left \begin{array}{c} l \\ m \end{array} \right\rangle = \left\langle \theta \phi \left \begin{array}{c} l' \\ m' \end{array} \right\rangle \left\langle \begin{array}{c} l' \\ m' \end{array} \right L_- \left \begin{array}{c} l \\ m \end{array} \right\rangle \right.$	$\langle \theta \phi L_- \theta' \phi' \rangle \left\langle \theta' \phi' \left \begin{array}{c} l \\ m \end{array} \right\rangle = \langle \theta \phi \left \begin{array}{c} l' \\ m' \end{array} \right\rangle \left\langle \begin{array}{c} l' \\ m' \end{array} \right L_- \left \begin{array}{c} l \\ m \end{array} \right\rangle$
gilmore-lie-groups_ E00047 (1.43)	28	■ 0.999	$\left[\begin{array}{ccc} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{array} \right] \left[\begin{array}{c} s_1 \\ s_2 \\ s_3 \end{array} \right] = \left[\begin{array}{c} v_1 \\ v_2 \\ v_3 \end{array} \right]$	$\left[\begin{array}{ccc} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{array} \right] \left[\begin{array}{c} s_1 \\ s_2 \\ s_3 \end{array} \right] = \left[\begin{array}{c} v_1 \\ v_2 \\ v_3 \end{array} \right]$	$\left[\begin{array}{ccc} 1 & 1 & 1 \\ 1 & \omega & \omega^2 \\ 1 & \omega^2 & \omega \end{array} \right] \left[\begin{array}{c} s_1 \\ s_2 \\ s_3 \end{array} \right] = \left[\begin{array}{c} v_1 \\ v_2 \\ v_3 \end{array} \right]$
gilmore-lie-groups_ E00132 (3.18)	54	■ 0.999	$\begin{array}{l} \text{orthogonal group} \\ \text{unitary group} \\ \text{symplectic group} \end{array}$	$G = I_{p,q} \begin{array}{l} \mathbb{R} \\ \mathbb{C} \\ \mathbb{Q} \end{array} \begin{array}{l} O(p, q) \\ U(p, q) \\ Sp(p, q) \end{array} \begin{array}{l} \text{orthogonal group} \\ \text{unitary group} \\ \text{symplectic group} \end{array}$	$G = I_{p,q} \begin{array}{l} \mathbb{R} \\ \mathbb{C} \\ \mathbb{Q} \end{array} \begin{array}{l} O(p, q) \\ U(p, q) \\ Sp(p, q) \end{array} \begin{array}{l} \text{orthogonal group} \\ \text{unitary group} \\ \text{symplectic group} \end{array}$
gilmore-lie-groups_ E00177	66	■ 0.999	$\left[\begin{array}{c} A_L \\ B_L \end{array} \right] = \left[\begin{array}{cc} t_{11} & t_{12} \\ t_{21} & t_{22} \end{array} \right] \left[\begin{array}{c} A_R \\ B_R \end{array} \right]$	$\left[\begin{array}{c} A_L \\ B_L \end{array} \right] = \left[\begin{array}{cc} t_{11} & t_{12} \\ t_{21} & t_{22} \end{array} \right] \left[\begin{array}{c} A_R \\ B_R \end{array} \right]$	$\left[\begin{array}{c} A_L \\ B_L \end{array} \right] = \left[\begin{array}{cc} t_{11} & t_{12} \\ t_{21} & t_{22} \end{array} \right] \left[\begin{array}{c} A_R \\ B_R \end{array} \right]$
gilmore-lie-groups_ E00234 (4.44)	81	■ 0.999	$dx^r = M(x)^r_i d\alpha^i$	$dx^r = M(x)^r_i d\alpha^i$	$dx^r = M(x)^r_i d\alpha^i$
gilmore-lie-groups_ E00318 (6.3)	102	■ 0.999	$\begin{aligned} & \left[\begin{array}{c} A \\ B \end{array} \right] = \left[\begin{array}{c} A \\ B \end{array} \right] \\ & \left[\begin{array}{c} A \\ B \end{array} \right] = \left[\begin{array}{c} A \\ B \end{array} \right] \end{aligned}$	$\begin{aligned} [\mathcal{A}, \mathcal{B}] &= [b_i^\dagger A_{ij} b_j, b_r^\dagger B_{rs} b_s] \\ &= A_{ij} B_{rs} [b_i^\dagger b_j, b_r^\dagger b_s] \\ &= A_{ij} B_{rs} (b_i^\dagger \delta_{jr} b_s - b_r^\dagger \delta_{si} b_j) \\ &= b_i^\dagger A_{ij} B_{js} b_s - b_r^\dagger B_{rs} A_{sj} b_j \\ &= b_i^\dagger [A, B]_{ij} b_j \\ &= \mathcal{C} \end{aligned}$	$\begin{aligned} [\mathcal{A}, \mathcal{B}] &= [b_i^\dagger A_{ij} b_j, b_r^\dagger B_{rs} b_s] \\ &= A_{ij} B_{rs} [b_i^\dagger b_j, b_r^\dagger b_s] \\ &= A_{ij} B_{rs} (b_i^\dagger \delta_{jr} b_s - b_r^\dagger \delta_{si} b_j) \\ &= b_i^\dagger A_{ij} B_{js} b_s - b_r^\dagger B_{rs} A_{sj} b_j \\ &= b_i^\dagger [A, B]_{ij} b_j \\ &= \mathcal{C} \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0353	109	■ 0.999	$\left[\begin{array}{c} d \\ d^\dagger \end{array}\right] = \left[\begin{array}{cc} A & B \\ C & D \end{array}\right] \left[\begin{array}{c} a \\ a^\dagger \end{array}\right]$		$\left[\begin{array}{c} d \\ d^\dagger \end{array}\right] = \left[\begin{array}{cc} A & B \\ C & D \end{array}\right] \left[\begin{array}{c} a \\ a^\dagger \end{array}\right]$
gilmore-lie-groups_ EQ0461 (7.87)	139	■ 0.999	$U(\alpha) 0\rangle = e^{i\alpha_+ J_+} \left \begin{array}{c} j \\ -j \end{array} \right\rangle e^{-ij\alpha_z} = \sum_{m=-j}^{m=+j} \frac{(i\alpha_+ J_+)^{j+m}}{(j+m)!} \left \begin{array}{c} j \\ -j \end{array} \right\rangle e^{-ij\alpha_z}$		$U(\alpha) 0\rangle = e^{i\alpha_+ J_+} \left \begin{array}{c} j \\ -j \end{array} \right\rangle e^{-ij\alpha_z} = \sum_{m=-j}^{m=+j} \frac{(i\alpha_+ J_+)^{j+m}}{(j+m)!} \left \begin{array}{c} j \\ -j \end{array} \right\rangle e^{-ij\alpha_z}$
gilmore-lie-groups_ EQ0752	234	■ 0.999	$\begin{aligned} & \mathcal{C}^2 \rightarrow \mathbf{A} \cdot \mathbf{A} + \mathbf{B} \cdot \mathbf{B} \\ & \mathcal{C}^3 \rightarrow \epsilon^{ijkl} X_{ij} A_k B_l \\ & \mathcal{C}^4 \rightarrow \sum_{ij} (\epsilon^{ijkl} A_k B_l)^2 \end{aligned}$		$\begin{aligned} \mathcal{C}^2 & \rightarrow \mathbf{A} \cdot \mathbf{A} + \mathbf{B} \cdot \mathbf{B} \\ \mathcal{C}^3 & \rightarrow \epsilon^{ijkl} X_{ij} A_k B_l \\ \mathcal{C}^4 & \rightarrow \sum_{ij} (\epsilon^{ijkl} A_k B_l)^2 \end{aligned}$
gilmore-lie-groups_ EQ0929 (15.63)	287	■ 0.999	$\langle k x \rangle \left\{ \mathbf{J} \cdot \frac{1}{i} \nabla + 1 \frac{1}{i} \frac{\partial}{\partial(ict)} I_3 \right\} \langle x k \rangle \left\langle k; \begin{array}{cc} 1 & 0 \\ m & 0 \end{array} \middle \psi \right\rangle = 0$		$\langle k x \rangle \left\{ \mathbf{J} \cdot \frac{1}{i} \nabla + 1 \frac{1}{i} \frac{\partial}{\partial(ict)} I_3 \right\} \langle x k \rangle \left\langle k; \begin{array}{cc} 1 & 0 \\ m & 0 \end{array} \middle \psi \right\rangle = 0$
gilmore-lie-groups_ EQ0403 (7.39)	125	■ 0.999	$\begin{aligned} \eta' &= \eta & l' &= (e^\eta - 1)l/\eta \\ \delta' &= (e^\eta - 1 - \eta)lr/\eta^2 & r' &= (e^\eta - 1)r/\eta \end{aligned}$		$\begin{aligned} \eta' &= \eta & l' &= (e^\eta - 1)l/\eta \\ \delta' &= (e^\eta - 1 - \eta)lr/\eta^2 & r' &= (e^\eta - 1)r/\eta \end{aligned}$
gilmore-lie-groups_ EQ0067 (1.62)	33	■ 0.999	$\begin{aligned} & y'^3 - J_1' y'^2 + J_2' y' - J_3' = 0 \\ & J_1' = 0 \\ & J_2' = -2736 \\ & J_3' = 51840 \end{aligned}$		$\begin{aligned} y'^3 - J_1' y'^2 + J_2' y' - J_3' &= 0 \\ J_1' &= 0 \\ J_2' &= -2736 \\ J_3' &= 51840 \end{aligned}$
gilmore-lie-groups_ EQ0098	44	■ 0.999	$\begin{aligned} & M_1^2 = I_2 + \begin{bmatrix} a \\ b \end{bmatrix} \begin{bmatrix} a & b \end{bmatrix} = \begin{bmatrix} 1+a^2 & ab \\ ba & 1+b^2 \end{bmatrix} \text{ and} \\ & M_2^2 = I_1 + \begin{bmatrix} a & b \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 1+a^2+b^2 \end{bmatrix} \end{aligned}$		$\begin{aligned} M_1^2 &= I_2 + \begin{bmatrix} a \\ b \end{bmatrix} \begin{bmatrix} a & b \end{bmatrix} = \begin{bmatrix} 1+a^2 & ab \\ ba & 1+b^2 \end{bmatrix} \text{ and} \\ M_2^2 &= I_1 + \begin{bmatrix} a & b \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} 1+a^2+b^2 \end{bmatrix} \end{aligned}$
gilmore-lie-groups_ EQ0161	62	■ 0.999	$SU(2) = \left[\begin{array}{cc} c_1 & c_2 \\ -c_2^* & c_1^* \end{array} \right]$		$SU(2) = \left[\begin{array}{cc} c_1 & c_2 \\ -c_2^* & c_1^* \end{array} \right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0241 (4.51)	82	■ 0.999	$\left[\begin{array}{l} dx \\ dy \\ dz \end{array}\right] = \left[\begin{array}{ccc} 1+x & 0 & y \\ -y & 1+x & 0 \\ z & 0 & \frac{1+yz}{1+x} \end{array}\right] \left[\begin{array}{l} d\alpha^1 \\ d\alpha^2 \\ d\alpha^3 \end{array}\right]$		
gilmore-lie-groups_ EQ0253 (4.59)	85	■ 0.999	$X = \sum_{ij} a_{ij} X_{ij} \rightarrow \left\{ \begin{array}{l} \mathfrak{def}(X) = \sum a_{ij} \mathfrak{def}(X_{ij}) \\ \mathfrak{reg}(X) = \sum a_{ij} \mathfrak{reg}(X_{ij}) \end{array} \right.$		
gilmore-lie-groups_ EQ0304	99	■ 0.999	$\begin{aligned} & (\mathfrak{h}, \mathfrak{h}) \leq 0 \\ & (\mathfrak{h}, \mathfrak{p}) = 0 \\ & (\mathfrak{p}, \mathfrak{p}) \geq 0 \end{aligned}$		
gilmore-lie-groups_ EQ0302 (7.20)	120	■ 0.999	$\mathfrak{s o}(3) = \left[\begin{array}{ccc} 0 & a_3 & -a_2 \\ -a_3 & 0 & a_1 \\ a_2 & -a_1 & 0 \end{array}\right] \quad \mathfrak{s u}(2) = \frac{i}{2} \left[\begin{array}{cc} b_3 & b_1 - ib_2 \\ b_1 + ib_2 & -b_3 \end{array}\right]$		
gilmore-lie-groups_ EQ0473 (8.5)	145	■ 0.999	$\mathfrak{Re g}(\eta \hat{n} + l a + r a^\dagger + \delta I) = \left[\begin{array}{cccc} 0 & -r & l & 0 \\ 0 & \eta & 0 & l \\ 0 & 0 & -\eta & -r \\ 0 & 0 & 0 & 0 \end{array}\right] \begin{array}{l} \hat{n} \\ a^\dagger \\ a \\ I \end{array}$		
gilmore-lie-groups_ EQ0524 (9.33)	162	■ 0.999	$\begin{aligned} & [H_i, E_\alpha] = \alpha_i E_\alpha \\ & [\mathbf{H}, E_\alpha] = \alpha E_\alpha \end{aligned}$		
gilmore-lie-groups_ EQ0540 (9.47)	166	■ 0.999	$\begin{array}{l} \text{normalization} \\ \text{Jacobi} \\ \text{symmetry} \end{array}$		
gilmore-lie-groups_ EQ0898 (15.32)	281	■ 0.999	$\Lambda = C_k H_{k^0}$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E09933 (15.68)	288	■ 0.999	$\begin{aligned} & \nabla \times \mathbf{B} - \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} = 0 \\ & \frac{\partial}{\partial t} (\nabla \times \mathbf{E} + \frac{1}{c} \frac{\partial \mathbf{B}}{\partial t}) = 0 \end{aligned}$	$\nabla \times \mathbf{B} - \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} = 0$ $\nabla \times \mathbf{E} + \frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} = 0$	$\nabla \times \mathbf{B} - \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} = 0$ $\nabla \times \mathbf{E} + \frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} = 0$
gilmore-lie-groups_ E09064 (1.59)	33	■ 0.999	$\begin{gathered} y^3 - J_1 y^2 + J_2 y - J_3 = 0 \\ J_1 = (-4)^1 [2I_2'] = (-4)(-30) = 120 \\ J_2 = (-4)^2 [I_2'^2 - 4I_4'] = 16(225 - 4 \times 24) = 2064 \\ J_3 = (-4)^3 [-I_3'^2] = (-64)(-100) = 6400 \end{gathered}$	$y^3 - J_1 y^2 + J_2 y - J_3 = 0$ $J_1 = (-4)^1 [2I_2'] = (-4)(-30) = 120$ $J_2 = (-4)^2 [I_2'^2 - 4I_4'] = 16(225 - 4 \times 24) = 2064$ $J_3 = (-4)^3 [-I_3'^2] = (-64)(-100) = 6400$	$y^3 - J_1 y^2 + J_2 y - J_3 = 0$ $J_1 = (-4)^1 [2I_2'] = (-4)(-30) = 120$ $J_2 = (-4)^2 [I_2'^2 - 4I_4'] = 16(225 - 4 \times 24) = 2064$ $J_3 = (-4)^3 [-I_3'^2] = (-64)(-100) = 6400$
gilmore-lie-groups_ E09297 (5.36)	96	■ 0.999	$\begin{aligned} & X = \begin{bmatrix} 0 & +1 \\ -1 & 0 \end{bmatrix} \quad (X, X) = -2 \quad e^{\theta X} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \\ & Y = \begin{bmatrix} 0 & +1 \\ +1 & 0 \end{bmatrix} \quad (Y, Y) = +2 \quad e^{\theta Y} = \begin{bmatrix} \cosh \theta & \sinh \theta \\ \sinh \theta & \cosh \theta \end{bmatrix} \end{aligned}$	$X = \begin{bmatrix} 0 & +1 \\ -1 & 0 \end{bmatrix} \quad (X, X) = -2 \quad e^{\theta X} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ $Y = \begin{bmatrix} 0 & +1 \\ +1 & 0 \end{bmatrix} \quad (Y, Y) = +2 \quad e^{\theta Y} = \begin{bmatrix} \cosh \theta & \sinh \theta \\ \sinh \theta & \cosh \theta \end{bmatrix}$	$X = \begin{bmatrix} 0 & +1 \\ -1 & 0 \end{bmatrix} \quad (X, X) = -2 \quad e^{\theta X} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ $Y = \begin{bmatrix} 0 & +1 \\ +1 & 0 \end{bmatrix} \quad (Y, Y) = +2 \quad e^{\theta Y} = \begin{bmatrix} \cosh \theta & \sinh \theta \\ \sinh \theta & \cosh \theta \end{bmatrix}$
gilmore-lie-groups_ E09670	216	■ 0.999	$\begin{aligned} & L_{\mu\nu} = x_\mu \partial_\nu - x_\nu \partial_\mu \\ & P_\mu = \partial_\mu \\ & K_\mu = 2x_\mu (x^\nu \partial_\nu) - (x^\nu x_\nu) \partial_\mu = 2x_\mu (x, \partial) - (x, x) \partial_\mu \\ & S = x^\nu \partial_\nu \end{aligned}$	$L_{\mu\nu} = x_\mu \partial_\nu - x_\nu \partial_\mu$ $P_\mu = \partial_\mu$ $K_\mu = 2x_\mu (x^\nu \partial_\nu) - (x^\nu x_\nu) \partial_\mu = 2x_\mu (x, \partial) - (x, x) \partial_\mu$ $S = x^\nu \partial_\nu$	$L_{\mu\nu} = x_\mu \partial_\nu - x_\nu \partial_\mu$ $P_\mu = \partial_\mu$ $K_\mu = 2x_\mu (x^\nu \partial_\nu) - (x^\nu x_\nu) \partial_\mu = 2x_\mu (x, \partial) - (x, x) \partial_\mu$ $S = x^\nu \partial_\nu$
gilmore-lie-groups_ E09045 (1.42)	28	■ 1.000	$v_2^3, v_3^3 = -\frac{27}{2} I_3' \pm \frac{1}{2} [(27I_3')^2 + 4 \times 27I_2'^3]^{1/2}$	$v_2^3, v_3^3 = -\frac{27}{2} I_3' \pm \frac{1}{2} [(27I_3')^2 + 4 \times 27I_2'^3]^{1/2}$	$v_2^3, v_3^3 = -\frac{27}{2} I_3' \pm \frac{1}{2} [(27I_3')^2 + 4 \times 27I_2'^3]^{1/2}$
gilmore-lie-groups_ E09048 (1.44)	28	■ 1.000	$\begin{aligned} & \left[\begin{array}{l} s_1 \\ s_2 \\ s_3 \end{array} \right] = \frac{1}{3} \left[\begin{array}{ccc} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{array} \right] \left[\begin{array}{l} I_1 \\ v_2 \\ v_3 \end{array} \right] = \frac{1}{3} \left[\begin{array}{l} v_1 + v_2 + v_3 \\ v_1 + \omega^2 v_2 + \omega v_3 \\ v_1 + \omega v_2 + \omega^2 v_3 \end{array} \right] \end{aligned}$	$\left[\begin{array}{l} s_1 \\ s_2 \\ s_3 \end{array} \right] = \frac{1}{3} \left[\begin{array}{ccc} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{array} \right] \left[\begin{array}{l} I_1 \\ v_2 \\ v_3 \end{array} \right] = \frac{1}{3} \left[\begin{array}{l} v_1 + v_2 + v_3 \\ v_1 + \omega^2 v_2 + \omega v_3 \\ v_1 + \omega v_2 + \omega^2 v_3 \end{array} \right]$	$\left[\begin{array}{l} s_1 \\ s_2 \\ s_3 \end{array} \right] = \frac{1}{3} \left[\begin{array}{ccc} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{array} \right] \left[\begin{array}{l} I_1 \\ v_2 \\ v_3 \end{array} \right] = \frac{1}{3} \left[\begin{array}{l} v_1 + v_2 + v_3 \\ v_1 + \omega^2 v_2 + \omega v_3 \\ v_1 + \omega v_2 + \omega^2 v_3 \end{array} \right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0057 (1.53)	31	■ 1.000	$\left[\begin{array}{l} t_1 \\ t_2 \\ t_3 \\ t_4 \end{array}\right] = \frac{1}{4} \left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{array}\right] \left[\begin{array}{c} I_1 \\ w_2 \\ w_3 \\ w_4 \end{array}\right]$		
gilmore-lie-groups_ EQ0066 (1.61)	33	■ 1.000	$\begin{aligned} & \& y_{1'} = y_1 - 40 \rightarrow 4 - 40 = -36 \\ & \& y_{2'} = y_2 - 40 \rightarrow 16 - 40 = -24 \\ & \& y_{3'} = y_3 - 40 \rightarrow 100 - 40 = 60 \end{aligned}$	$\begin{aligned} y_1' &= y_1 - 40 \rightarrow 4 - 40 = -36 \\ y_2' &= y_2 - 40 \rightarrow 16 - 40 = -24 \\ y_3' &= y_3 - 40 \rightarrow 100 - 40 = 60 \end{aligned}$	$\begin{aligned} y_1' &= y_1 - 40 \rightarrow 4 - 40 = -36 \\ y_2' &= y_2 - 40 \rightarrow 16 - 40 = -24 \\ y_3' &= y_3 - 40 \rightarrow 100 - 40 = 60 \end{aligned}$
gilmore-lie-groups_ EQ0075 (1.70)	34	■ 1.000	$\begin{aligned} & \& v_2^3 = 699840 + i 145152 \sqrt{3} \\ & \& v_3^3 = 699840 - i 145152 \sqrt{3} \end{aligned}$	$\begin{aligned} v_2^3 &= 699840 + i 145152 \sqrt{3} \\ v_3^3 &= 699840 - i 145152 \sqrt{3} \end{aligned}$	$\begin{aligned} v_2^3 &= 699840 + i 145152 \sqrt{3} \\ v_3^3 &= 699840 - i 145152 \sqrt{3} \end{aligned}$
gilmore-lie-groups_ EQ0077 (1.73)	35	■ 1.000	$\begin{aligned} & \& w_2^2 = y_1 \quad w_2 = \pm 4 \\ & \& w_3^2 = y_2 \quad w_3 = \pm 10 \\ & \& w_4^2 = y_3 \quad w_4 = \pm 2 \end{aligned}$	$\begin{aligned} \left[\begin{array}{c} y_1 \\ y_2 \\ y_3 \end{array}\right] &= \frac{1}{3} \left[\begin{array}{ccc} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{array}\right] \left[\begin{array}{c} J_1 = 120 \\ v_2 = -36 + i 48 \sqrt{3} \\ v_3 = -36 - i 48 \sqrt{3} \end{array}\right] = \left[\begin{array}{c} 16 \\ 100 \\ 4 \end{array}\right] \end{aligned}$	$\begin{aligned} \left[\begin{array}{c} y_1 \\ y_2 \\ y_3 \end{array}\right] &= \frac{1}{3} \left[\begin{array}{ccc} 1 & 1 & 1 \\ 1 & \omega^2 & \omega \\ 1 & \omega & \omega^2 \end{array}\right] \left[\begin{array}{c} J_1 = 120 \\ v_2 = -36 + i 48 \sqrt{3} \\ v_3 = -36 - i 48 \sqrt{3} \end{array}\right] = \left[\begin{array}{c} 16 \\ 100 \\ 4 \end{array}\right] \end{aligned}$
gilmore-lie-groups_ EQ0086 (2.1)	39	■ 1.000	$A = \left[\begin{array}{ll} \alpha & \beta \\ \gamma & \delta \end{array}\right] \quad \det(A) = \alpha\delta - \beta\gamma = +1$	$A = \left[\begin{array}{cc} \alpha & \beta \\ \gamma & \delta \end{array}\right] \quad \det(A) = \alpha\delta - \beta\gamma = +1$	$A = \left[\begin{array}{cc} \alpha & \beta \\ \gamma & \delta \end{array}\right] \quad \det(A) = \alpha\delta - \beta\gamma = +1$
gilmore-lie-groups_ EQ0108	47	■ 1.000	$\begin{aligned} & \& -ds^2 = dx^2 + dy^2 - \left(d\sqrt{1+x^2+y^2}\right)^2 \\ & \& = \frac{1}{1+x^2+y^2} \left(\begin{array}{cc} dx & dy \end{array}\right) \left[\begin{array}{cc} 1+y^2 & -xy \\ -yx & 1+x^2 \end{array}\right] \left(\begin{array}{c} dx \\ dy \end{array}\right) \end{aligned}$	$\begin{aligned} -ds^2 &= dx^2 + dy^2 - \left(d\sqrt{1+x^2+y^2}\right)^2 \\ &= \frac{1}{1+x^2+y^2} \left(\begin{array}{cc} dx & dy \end{array}\right) \left[\begin{array}{cc} 1+y^2 & -xy \\ -yx & 1+x^2 \end{array}\right] \left(\begin{array}{c} dx \\ dy \end{array}\right) \end{aligned}$	$\begin{aligned} -ds^2 &= dx^2 + dy^2 - \left(d\sqrt{1+x^2+y^2}\right)^2 \\ &= \frac{1}{1+x^2+y^2} \left(\begin{array}{cc} dx & dy \end{array}\right) \left[\begin{array}{cc} 1+y^2 & -xy \\ -yx & 1+x^2 \end{array}\right] \left(\begin{array}{c} dx \\ dy \end{array}\right) \end{aligned}$
gilmore-lie-groups_ EQ0111 (3.1)	48	■ 1.000	$\left[\begin{array}{cc} a & b \\ 0 & \frac{1}{a} \end{array}\right]$	$\left[\begin{array}{cc} a & b \\ 0 & \frac{1}{a} \end{array}\right]$	$\left[\begin{array}{cc} a & b \\ 0 & \frac{1}{a} \end{array}\right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0175	65	■ 1.000	$\psi_L(x)=A_L e^{+i k x}+B_L e^{-i k x}=\left[\begin{array}{l} e^{+i k x} \\ e^{-i k x} \end{array}\right]\left[\begin{array}{l} A_L \\ B_L \end{array}\right]$	$\psi_L(x)=A_L e^{+i k x}+B_L e^{-i k x}=\left[\begin{array}{cc} e^{+i k x} & e^{-i k x} \end{array}\right]\left[\begin{array}{c} A_L \\ B_L \end{array}\right]$	$\psi_L(x)=A_L e^{+i k x}+B_L e^{-i k x}=\left[\begin{array}{cc} e^{+i k x} & e^{-i k x} \end{array}\right]\left[\begin{array}{c} A_L \\ B_L \end{array}\right]$
gilmore-lie-groups_ EQ0187	68	■ 1.000	$S_{\text{Network}}=D+C\Gamma B+C\Gamma A\Gamma B+C\Gamma A\Gamma A\Gamma B+C\Gamma A\Gamma A\Gamma A\Gamma B+\cdots$	$S_{\text{Network}}=D+C\Gamma B+C\Gamma A\Gamma B+C\Gamma A\Gamma A\Gamma B+C\Gamma A\Gamma A\Gamma A\Gamma B+\cdots$	$S_{\text{Network}}=D+C\Gamma B+C\Gamma A\Gamma B+C\Gamma A\Gamma A\Gamma B+C\Gamma A\Gamma A\Gamma A\Gamma B+\cdots$
gilmore-lie-groups_ EQ0191 (4.4)	71	■ 1.000	$\begin{gathered} (\delta a, \delta b, \delta c) \rightarrow I_2 + \delta a X_a + \delta b X_b + \delta c X_c = \left[\begin{array}{cc} 1 + \delta a & \delta b \\ \delta c & 1 - \delta a \end{array}\right] \\ X_a = \left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right] = \frac{\partial M(a, b, c)}{\partial a} \Big _{(a, b, c)=(0, 0, 0)} \\ X_b = \left[\begin{array}{cc} 0 & 1 \\ 0 & 0 \end{array}\right] = \frac{\partial M(a, b, c)}{\partial b} \Big _{(a, b, c)=(0, 0, 0)} \\ X_c = \left[\begin{array}{cc} 0 & 0 \\ 1 & 0 \end{array}\right] = \frac{\partial M(a, b, c)}{\partial c} \Big _{(a, b, c)=(0, 0, 0)} \end{gathered}$	$(\delta a, \delta b, \delta c) \rightarrow I_2 + \delta a X_a + \delta b X_b + \delta c X_c = \left[\begin{array}{cc} 1 + \delta a & \delta b \\ \delta c & 1 - \delta a \end{array}\right]$ $X_a = \left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right] = \frac{\partial M(a, b, c)}{\partial a} \Big _{(a, b, c)=(0, 0, 0)}$ $X_b = \left[\begin{array}{cc} 0 & 1 \\ 0 & 0 \end{array}\right] = \frac{\partial M(a, b, c)}{\partial b} \Big _{(a, b, c)=(0, 0, 0)}$ $X_c = \left[\begin{array}{cc} 0 & 0 \\ 1 & 0 \end{array}\right] = \frac{\partial M(a, b, c)}{\partial c} \Big _{(a, b, c)=(0, 0, 0)}$	$(\delta a, \delta b, \delta c) \rightarrow I_2 + \delta a X_a + \delta b X_b + \delta c X_c = \left[\begin{array}{cc} 1 + \delta a & \delta b \\ \delta c & 1 - \delta a \end{array}\right]$ $X_a = \left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right] = \frac{\partial M(a, b, c)}{\partial a} \Big _{(a, b, c)=(0, 0, 0)}$ $X_b = \left[\begin{array}{cc} 0 & 1 \\ 0 & 0 \end{array}\right] = \frac{\partial M(a, b, c)}{\partial b} \Big _{(a, b, c)=(0, 0, 0)}$ $X_c = \left[\begin{array}{cc} 0 & 0 \\ 1 & 0 \end{array}\right] = \frac{\partial M(a, b, c)}{\partial c} \Big _{(a, b, c)=(0, 0, 0)}$
gilmore-lie-groups_ EQ0213 (4.23)	76	■ 1.000	$C_{ab}^b = -C_{ba}^b = +2 \quad C_{ac}^c = -C_{ca}^c = -2 \quad C_{bc}^a = -C_{cb}^a = +1$	$C_{ab}^b = -C_{ba}^b = +2 \quad C_{ac}^c = -C_{ca}^c = -2 \quad C_{bc}^a = -C_{cb}^a = +1$	$C_{ab}^b = -C_{ba}^b = +2 \quad C_{ac}^c = -C_{ca}^c = -2 \quad C_{bc}^a = -C_{cb}^a = +1$
gilmore-lie-groups_ EQ0222 (4.32)	78	■ 1.000	$L_z = \left[\begin{array}{ccc} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{array}\right] \quad P_x = \left[\begin{array}{ccc} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array}\right] \quad P_y = \left[\begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{array}\right]$	$L_z = \left[\begin{array}{ccc} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{array}\right] \quad P_x = \left[\begin{array}{ccc} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array}\right] \quad P_y = \left[\begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{array}\right]$	$L_z = \left[\begin{array}{ccc} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{array}\right] \quad P_x = \left[\begin{array}{ccc} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array}\right] \quad P_y = \left[\begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{array}\right]$
gilmore-lie-groups_ EQ0235 (4.45)	81	■ 1.000	$ds^2 = g_{ij}(\text{Id}) d\alpha^i d\alpha^j$	$ds^2 = g_{ij}(\text{Id}) d\alpha^i d\alpha^j$	$ds^2 = g_{ij}(\text{Id}) d\alpha^i d\alpha^j$
gilmore-lie-groups_ EQ0243 (4.53)	82	■ 1.000	$\left[\begin{array}{ccc} 2 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{array}\right] \text{left translation by } (x, y, z) \left[\begin{array}{ccc} \frac{2(1+yz)}{(1+x)^2} & -\frac{z}{(1+x)} & -\frac{y}{(1+x)} \\ -\frac{z}{(1+x)} & 0 & 1 \\ -\frac{y}{(1+x)} & 1 & 0 \end{array}\right]$	$\left[\begin{array}{ccc} 2 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{array}\right] \text{left translation by } (x, y, z) \left[\begin{array}{ccc} \frac{2(1+yz)}{(1+x)^2} & -\frac{z}{(1+x)} & -\frac{y}{(1+x)} \\ -\frac{z}{(1+x)} & 0 & 1 \\ -\frac{y}{(1+x)} & 1 & 0 \end{array}\right]$	$\left[\begin{array}{ccc} 2 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{array}\right] \xrightarrow[\text{by } (x, y, z)]{\text{left translation}} \left[\begin{array}{ccc} \frac{2(1+yz)}{(1+x)^2} & -\frac{z}{(1+x)} & -\frac{y}{(1+x)} \\ -\frac{z}{(1+x)} & 0 & 1 \\ -\frac{y}{(1+x)} & 1 & 0 \end{array}\right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00254 (4.60)	85	■1.000	$\operatorname{tr} \frac{\mathrm{d} e f}{\mathrm{d} e f}(X) \frac{\mathrm{d} e f}{\mathrm{d} e f}(X) \leftarrow (X, X) \rightarrow \operatorname{tr} \frac{\mathrm{d} e f}{\mathrm{d} e f}(X) \frac{\mathrm{d} e f}{\mathrm{d} e f}(X)$	$\operatorname{tr} \mathfrak{d} \operatorname{ef}(X) \mathfrak{d} \operatorname{ef}(X) \leftarrow (X, X) \rightarrow \operatorname{tr} \operatorname{reg}(X) \operatorname{reg}(X)$	$\operatorname{tr} \mathfrak{d} \operatorname{ef}(X) \mathfrak{d} \operatorname{ef}(X) \leftarrow (X, X) \rightarrow \operatorname{tr} \operatorname{reg}(X) \operatorname{reg}(X)$
gilmore-lie-groups_ E00281 (5.20)	93	■1.000	$M^{\dagger} G + G M = 0 \quad G^t = -G \quad \mathbb{F} = \begin{cases} \mathbb{R} & \mathfrak{sp}(G; \mathbb{R}) \\ \mathbb{C} & \mathfrak{sp}(G; \mathbb{C}) \end{cases}$	$M^{\dagger} G + G M = 0 \quad G^t = -G \quad \mathbb{F} = \begin{cases} \mathbb{R} & \mathfrak{sp}(G; \mathbb{R}) \\ \mathbb{C} & \mathfrak{sp}(G; \mathbb{C}) \end{cases}$	$M^{\dagger} G + G M = 0 \quad G^t = -G \quad \mathbb{F} = \begin{cases} \mathbb{R} & \mathfrak{sp}(G; \mathbb{R}) \\ \mathbb{C} & \mathfrak{sp}(G; \mathbb{C}) \end{cases}$
gilmore-lie-groups_ E00329 (6.14)	104	■1.000	$\left(\begin{array}{lll} x_1 & x_2 & x_3 \end{array} \right) \left(\begin{array}{ccc} 0 & \theta_3 & -\theta_2 \\ -\theta_3 & 0 & \theta_1 \\ \theta_2 & -\theta_1 & 0 \end{array} \right) \left(\begin{array}{c} \partial_1 \\ \partial_2 \\ \partial_3 \end{array} \right) = \theta \cdot \mathcal{L}$	$\left(\begin{array}{ccc} x_1 & x_2 & x_3 \end{array} \right) \left(\begin{array}{ccc} 0 & \theta_3 & -\theta_2 \\ -\theta_3 & 0 & \theta_1 \\ \theta_2 & -\theta_1 & 0 \end{array} \right) \left(\begin{array}{c} \partial_1 \\ \partial_2 \\ \partial_3 \end{array} \right) = \theta \cdot \mathcal{L}$	$(x_1 \quad x_2 \quad x_3) \begin{bmatrix} 0 & \theta_3 & -\theta_2 \\ -\theta_3 & 0 & \theta_1 \\ \theta_2 & -\theta_1 & 0 \end{bmatrix} \begin{bmatrix} \partial_1 \\ \partial_2 \\ \partial_3 \end{bmatrix} = \theta \cdot \mathcal{L}$
gilmore-lie-groups_ E00332 (6.17)	105	■1.000	$\left(\begin{array}{lll} x_1 & x_2 & 1 \end{array} \right) \left(\begin{array}{ccc} 0 & \theta & 0 \\ -\theta & 0 & 0 \\ t_1 & t_2 & 0 \end{array} \right) \left(\begin{array}{c} \partial_1 \\ \partial_2 \\ 1 \end{array} \right) = \theta \mathcal{L}_z + t_i \mathcal{T}_i$	$\left(\begin{array}{ccc} x_1 & x_2 & 1 \end{array} \right) \left(\begin{array}{ccc} 0 & \theta & 0 \\ -\theta & 0 & 0 \\ t_1 & t_2 & 0 \end{array} \right) \left(\begin{array}{c} \partial_1 \\ \partial_2 \\ 1 \end{array} \right) = \theta \mathcal{L}_z + t_i \mathcal{T}_i$	$(x_1 \ x_2 \ 1) \begin{bmatrix} 0 & \theta & 0 \\ -\theta & 0 & 0 \\ t_1 & t_2 & 0 \end{bmatrix} \begin{bmatrix} \partial_1 \\ \partial_2 \\ 1 \end{bmatrix} = \theta \mathcal{L}_z + t_i \mathcal{T}_i$
gilmore-lie-groups_ E00404 (7.40)	125	■1.000	$\langle 0 e^{\eta a^{\dagger} a + r a^{\dagger} + l a} 0 \rangle = \langle 0 e^{r' a^{\dagger}} e^{\eta' a^{\dagger} a + \delta' I} e^{l' a} 0 \rangle$	$\langle 0 e^{\eta a^{\dagger} a + r a^{\dagger} + l a} 0 \rangle = \langle 0 e^{r' a^{\dagger}} e^{\eta' a^{\dagger} a + \delta' I} e^{l' a} 0 \rangle$	$\langle 0 e^{\eta a^{\dagger} a + r a^{\dagger} + l a} 0 \rangle = \langle 0 e^{r' a^{\dagger}} e^{\eta' a^{\dagger} a + \delta' I} e^{l' a} 0 \rangle$
gilmore-lie-groups_ E00443	135	■1.000	$e^{r a^{\dagger}} e^{\delta I} e^{l a} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & r \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & \delta \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & l & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & l & \delta \\ 0 & 1 & r \\ 0 & 0 & 1 \end{bmatrix}$	$e^{r a^{\dagger}} e^{\delta I} e^{l a} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & r \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & \delta \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & l & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & l & \delta \\ 0 & 1 & r \\ 0 & 0 & 1 \end{bmatrix}$	$e^{r a^{\dagger}} e^{\delta I} e^{l a} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & r \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & \delta \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & l & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & l & \delta \\ 0 & 1 & r \\ 0 & 0 & 1 \end{bmatrix}$
gilmore-lie-groups_ E00476 (8.8)	146	■1.000	$\begin{array}{l} Y_i = \frac{1}{2} (X_{i4} + \frac{1}{2} \epsilon_{irs} X_{rs}) \\ Z_i = \frac{1}{2} (X_{i4} - \frac{1}{2} \epsilon_{irs} X_{rs}) \end{array}$	$\begin{array}{l} Y_i = \frac{1}{2} (X_{i4} + \frac{1}{2} \epsilon_{irs} X_{rs}) \\ Z_i = \frac{1}{2} (X_{i4} - \frac{1}{2} \epsilon_{irs} X_{rs}) \end{array}$	$\begin{array}{l} Y_i = \frac{1}{2} (X_{i4} + \frac{1}{2} \epsilon_{irs} X_{rs}) \\ Z_i = \frac{1}{2} (X_{i4} - \frac{1}{2} \epsilon_{irs} X_{rs}) \end{array}$
gilmore-lie-groups_ E00493 (9.2)	154	■1.000	$\left[Z, \sum a^i X_i \right] = \lambda \sum a^j X_j$	$\sum \sum a^i (R(Z)_i^j - \lambda \delta_i^j) X_j = 0$	$\left[Z, \sum a^i X_i \right] = \lambda \sum a^j X_j$ $\sum \sum a^i (R(Z)_i^j - \lambda \delta_i^j) X_j = 0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08518 (9.27)	161	■ 1.000	$\begin{aligned} & \phi_2(z_1, z_2) = -5z_1^2 z_1^2 \quad \& \phi_4(z_1, z_2) = 4z_1^4 = 4(-\phi_2(z_1, z_2)/5)^2 \\ & z_1^4 = 4\left(-\phi_2(z_1, z_2) / 5\right)^2 \end{aligned}$	$\begin{aligned} \phi_2(z_1, z_2) &= -5z_1^2 \\ \phi_4(z_1, z_2) &= 4z_1^4 = 4(-\phi_2(z_1, z_2)/5)^2 \end{aligned}$	$\begin{aligned} \phi_2(z_1, z_2) &= -5z_1^2 \\ \phi_4(z_1, z_2) &= 4z_1^4 = 4(-\phi_2(z_1, z_2)/5)^2 \end{aligned}$
gilmore-lie-groups_ E08534 (9.41)	164	■ 1.000	$\begin{aligned} & N_{-\alpha, \beta - m\alpha} = 0 \\ & N_{+\alpha, \beta + n\alpha} = 0 \end{aligned}$	$\begin{aligned} N_{-\alpha, \beta - m\alpha} &= 0 \\ N_{+\alpha, \beta + n\alpha} &= 0 \end{aligned}$	$\begin{aligned} N_{-\alpha, \beta - m\alpha}^2 &= 0 \\ N_{+\alpha, \beta + n\alpha}^2 &= 0 \end{aligned}$
gilmore-lie-groups_ E08546	170	■ 1.000	$\frac{\left(\left(A_1, A_2, \ldots, A_r\right)_{\Re e g}\right)_{\left\{f_r\left(\Re e g\right)\right\}}}{f_r\left(\Re e g\right)}=\left(A_1, A_2, \ldots, A_r\right)=\frac{\left(A_1, A_2, \ldots, A_r\right)_{\Gamma}}{f_r\left(\Gamma\right)}$	$\frac{\left(A_1, A_2, \ldots, A_r\right)_{\Re e g}}{f_r\left(\Re e g\right)}=\left(A_1, A_2, \ldots, A_r\right)=\frac{\left(A_1, A_2, \ldots, A_r\right)_{\Gamma}}{f_r\left(\Gamma\right)}$	$\frac{\left(A_1, A_2, \ldots, A_r\right)_{\Re e g}}{f_r\left(\Re e g\right)}=\left(A_1, A_2, \ldots, A_r\right)=\frac{\left(A_1, A_2, \ldots, A_r\right)_{\Gamma}}{f_r\left(\Gamma\right)}$
gilmore-lie-groups_ E08687	197	■ 1.000	$\begin{aligned} & \Re e g\left(w a_1^{\dagger} a_1+x a_1^{\dagger} a_2+y a_2^{\dagger} a_1+z a_2^{\dagger} a_2\right)=\left[\begin{array}{cccc} 0 & -x & y & 0 \\ -y & w-z & 0 & y \\ x & 0 & -w+z & -x \\ 0 & x & -y & 0 \end{array}\right] \end{aligned}$	$\Re e g\left(w a_1^{\dagger} a_1+x a_1^{\dagger} a_2+y a_2^{\dagger} a_1+z a_2^{\dagger} a_2\right)=\left[\begin{array}{cccc} 0 & -x & y & 0 \\ -y & w-z & 0 & y \\ x & 0 & -w+z & -x \\ 0 & x & -y & 0 \end{array}\right]$	$\Re e g\left(w a_1^{\dagger} a_1+x a_1^{\dagger} a_2+y a_2^{\dagger} a_1+z a_2^{\dagger} a_2\right)=\left[\begin{array}{cccc} 0 & -x & y & 0 \\ -y & w-z & 0 & y \\ x & 0 & -w+z & -x \\ 0 & x & -y & 0 \end{array}\right]$
gilmore-lie-groups_ E08667	215	■ 1.000	$\begin{aligned} & y^i=s x^i \quad\left(1 \leq i \leq n\right) \\ & y^{n+1}=s \\ & y^{n+2}=s(x, x)_m \end{aligned}$	$\begin{aligned} y^i &=s x^i \quad\left(1 \leq i \leq n\right) \\ y^{n+1} &=s \\ y^{n+2} &=s(x, x)_m \end{aligned}$	$\begin{aligned} y^i &=s x^i \quad\left(1 \leq i \leq n\right) \\ y^{n+1} &=s \\ y^{n+2} &=s(x, x)_m \end{aligned}$
gilmore-lie-groups_ E08675	217	■ 1.000	$\begin{aligned} & s\left(z_1, z_2\right)=2 \tanh ^{-1} \frac{\left \left z_1-z_2\right \right }{\left \left z_1-\bar{z}_2\right \right }=\log \left\{\frac{\left \left z_1-\bar{z}_2\right \right +\left \left z_1-z_2\right \right }{\left \left z_1-\bar{z}_2\right \right -\left \left z_1-z_2\right \right }\right\} \end{aligned}$	$s\left(z_1, z_2\right)=2 \tanh ^{-1} \frac{\left \left z_1-z_2\right \right }{\left \left z_1-\bar{z}_2\right \right }=\log \left\{\frac{\left \left z_1-\bar{z}_2\right \right +\left \left z_1-z_2\right \right }{\left \left z_1-\bar{z}_2\right \right -\left \left z_1-z_2\right \right }\right\}$	$s\left(z_1, z_2\right)=2 \tanh ^{-1} \frac{\left \left z_1-z_2\right \right }{\left \left z_1-\bar{z}_2\right \right }=\log \left\{\frac{\left \left z_1-\bar{z}_2\right \right +\left \left z_1-z_2\right \right }{\left \left z_1-\bar{z}_2\right \right -\left \left z_1-z_2\right \right }\right\}$
gilmore-lie-groups_ E08714 (13.31)	226	■ 1.000	$\begin{aligned} & \left[\begin{array}{c} h_3 \\ h_+ \\ h_- \\ h_0 \end{array}\right] \xrightarrow{c \rightarrow 0}\left[\begin{array}{c} \hat{n}+\frac{1}{2} I=\frac{1}{2}\left\{a, a^{\dagger}\right\} \\ a^{\dagger} \\ a \\ I \end{array}\right] \end{aligned}$	$\left[\begin{array}{c} h_3 \\ h_+ \\ h_- \\ h_0 \end{array}\right] \xrightarrow{c \rightarrow 0}\left[\begin{array}{c} \hat{n}+\frac{1}{2} I=\frac{1}{2}\left\{a, a^{\dagger}\right\} \\ a^{\dagger} \\ a \\ I \end{array}\right]$	$\left[\begin{array}{c} h_3 \\ h_+ \\ h_- \\ h_0 \end{array}\right] \xrightarrow{c \rightarrow 0}\left[\begin{array}{c} \hat{n}+\frac{1}{2} I=\frac{1}{2}\left\{a, a^{\dagger}\right\} \\ a^{\dagger} \\ a \\ I \end{array}\right]$
gilmore-lie-groups_ E08746	233	■ 1.000	$\begin{aligned} & e^{\alpha a^{\dagger}-\alpha^* a}=e^{\alpha a^{\dagger}} e^{-\frac{1}{2} \alpha^* \alpha I} e^{-\alpha^* a} \end{aligned}$	$e^{\alpha a^{\dagger}-\alpha^* a}=e^{\alpha a^{\dagger}} e^{-\frac{1}{2} \alpha^* \alpha I} e^{-\alpha^* a}$	$e^{\alpha a^{\dagger}-\alpha^* a}=e^{\alpha a^{\dagger}} e^{-\frac{1}{2} \alpha^* \alpha I} e^{-\alpha^* a}$
gilmore-lie-groups_ E08769	240	■ 1.000	$\begin{aligned} & E=\frac{m c^2}{\sqrt{1+\left(\alpha / N'\right)^2}}, W=-\frac{1}{2} m c^2 \alpha^2 \frac{1}{N^2} \\ & N'=n+\frac{1}{2}+\sqrt{\left(l+\frac{1}{2}\right)^2-\alpha^2}, \quad N=n+l+1 \end{aligned}$	$\begin{aligned} E &= \frac{m c^2}{\sqrt{1+\left(\alpha / N'\right)^2}}, W=-\frac{1}{2} m c^2 \alpha^2 \frac{1}{N^2} \\ N' &=n+\frac{1}{2}+\sqrt{\left(l+\frac{1}{2}\right)^2-\alpha^2}, \quad N=n+l+1 \end{aligned}$	$\begin{aligned} E &= \frac{m c^2}{\sqrt{1+\left(\alpha / N'\right)^2}}, W=-\frac{1}{2} m c^2 \alpha^2 \frac{1}{N^2} \\ N' &=n+\frac{1}{2}+\sqrt{\left(l+\frac{1}{2}\right)^2-\alpha^2}, \quad N=n+l+1 \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08882 (14.45)	251	■ 1.000	$\begin{aligned} & \mathbf{u}=2\mathbf{p}_4\left(\mathbf{p}\cdot\mathbf{p}+\mathbf{p}_4^2\right)^{-1}\mathbf{p} \\ & \mathbf{u}_4=\left(\mathbf{p}\cdot\mathbf{p}-\mathbf{p}_4^2\right)\left(\mathbf{p}\cdot\mathbf{p}+\mathbf{p}_4^2\right)^{-1} \end{aligned}$	$\mathbf{u}=2p_4\left(\mathbf{p}\cdot\mathbf{p}+p_4^2\right)^{-1}\mathbf{p}$ $u_4=\left(\mathbf{p}\cdot\mathbf{p}-p_4^2\right)\left(\mathbf{p}\cdot\mathbf{p}+p_4^2\right)^{-1}$	$\mathbf{u}=2p_4(\mathbf{p}\cdot\mathbf{p}+p_4^2)^{-1}\mathbf{p}$ $u_4=(\mathbf{p}\cdot\mathbf{p}-p_4^2)(\mathbf{p}\cdot\mathbf{p}+p_4^2)^{-1}$
gilmore-lie-groups_ E08818 (14.54)	256	■ 1.000	$S\left(8;\mathbb{R}\right)\cap S\left(0(4,4)=S\left(U(2,2)\right)\simeq S\left(0(4,2)\right)\right)$	$Sp(8;\mathbb{R})\cap SO(4,4)=SU(2,2)\simeq SO(4,2)$	$Sp(8;\mathbb{R})\cap SO(4,4)=SU(2,2)\simeq SO(4,2)$
gilmore-lie-groups_ E08833 (14.67)	260	■ 1.000	$e^{\theta M_4}\binom{\Gamma_5}{\Gamma_4}e^{-\theta M_4}=\left[\begin{array}{cc}\cosh\theta & -\sinh\theta \\ -\sinh\theta & \cosh\theta\end{array}\right]\binom{\Gamma_5}{\Gamma_4}$	$e^{\theta M_4}\binom{\Gamma_5}{\Gamma_4}e^{-\theta M_4}=\left[\begin{array}{cc}\cosh\theta & -\sinh\theta \\ -\sinh\theta & \cosh\theta\end{array}\right]\binom{\Gamma_5}{\Gamma_4}$	$e^{\theta M_4}\binom{\Gamma_5}{\Gamma_4}e^{-\theta M_4}=\left[\begin{array}{cc}\cosh\theta & -\sinh\theta \\ -\sinh\theta & \cosh\theta\end{array}\right]\binom{\Gamma_5}{\Gamma_4}$
gilmore-lie-groups_ E0886 (15.19)	278	■ 1.000	$\Lambda T_{\mu\nu}(x)=T_{\mu^{\prime}\nu^{\prime}}(x\Lambda^{-1})\Lambda_{\mu^{\prime}\mu}\Lambda_{\nu^{\prime}\nu}$	$\Lambda T_{\mu\nu}(x)=T_{\mu^{\prime}\nu^{\prime}}(x\Lambda^{-1})\Lambda_{\mu^{\prime}\mu}\Lambda_{\nu^{\prime}\nu}$	$\Lambda T_{\mu\nu}(x)=T_{\mu^{\prime}\nu^{\prime}}(x\Lambda^{-1})\Lambda_{\mu^{\prime}\mu}\Lambda_{\nu^{\prime}\nu}$
gilmore-lie-groups_ E0921 (15.55)	286	■ 1.000	$\left\{J_3^{(j)}k_3^0+jk_4^0I_{2j+1}\right\}\left\langle k^0;\begin{array}{c}0\\0\end{array}\begin{array}{c}j'\\m'\end{array}\middle \psi\right\rangle=0$	$\left\{J_3^{(j)}k_3^0+jk_4^0I_{2j+1}\right\}\left\langle k^0;\begin{array}{c}0\\0\end{array}\begin{array}{c}j'\\m'\end{array}\middle \psi\right\rangle=0$	$\left\{J_3^{(j)}k_3^0+jk_4^0I_{2j+1}\right\}\left\langle k^0;\begin{array}{c}0\\0\end{array}\begin{array}{c}j'\\m'\end{array}\middle \psi\right\rangle=0$
gilmore-lie-groups_ E0946	291	■ 1.000	$\begin{aligned} & \nabla\cdot\mathbf{B}(\mathbf{x},t)=4\pi\rho_m(\mathbf{x},t)+C_m(\mathbf{x}) \\ & \nabla\cdot\mathbf{E}(\mathbf{x},t)=4\pi\rho_e(\mathbf{x},t)+C_e(\mathbf{x}) \end{aligned}$	$\nabla\cdot\mathbf{B}(\mathbf{x},t)=4\pi\rho_m(\mathbf{x},t)+C_m(\mathbf{x})$ $\nabla\cdot\mathbf{E}(\mathbf{x},t)=4\pi\rho_e(\mathbf{x},t)+C_e(\mathbf{x})$	$\nabla\cdot\mathbf{B}(\mathbf{x},t)=4\pi\rho_m(\mathbf{x},t)+C_m(\mathbf{x})$ $\nabla\cdot\mathbf{E}(\mathbf{x},t)=4\pi\rho_e(\mathbf{x},t)+C_e(\mathbf{x})$
gilmore-lie-groups_ E0970 (16.13)	302	■ 1.000	$\xi(x,y)=\sum_{i,j}\xi_{ij}x^iy^j\quad 0\leq i,j,\quad i+j\leq d_\xi$	$\xi(x,y)=\sum_{i,j}\xi_{ij}x^iy^j\quad 0\leq i,j,\quad i+j\leq d_\xi$	$\xi(x,y)=\sum_{i,j}\xi_{ij}x^iy^j\quad 0\leq i,j,\quad i+j\leq d_\xi$
gilmore-lie-groups_ E01009 (16.52)	314	■ 1.000	$\begin{aligned} & X_{ij}=x^i\partial_j-x^j\partial_i \\ & Z=x^i\partial_i+(2-n)u\frac{\partial}{\partial u} \end{aligned}$	$X_{ij}=x^i\partial_j-x^j\partial_i$ $Z=x^i\partial_i+(2-n)u\frac{\partial}{\partial u}$	$X_{ij}=x^i\partial_j-x^j\partial_i$ $Z=x^i\partial_i+(2-n)u\frac{\partial}{\partial u}$
gilmore-lie-groups_ E0001	18	■ 1.000	$\left(g_i\cdot g_j\right)\cdot g_k=g_i\cdot\left(g_j\cdot g_k\right)$	$(g_i\cdot g_j)\cdot g_k=g_i\cdot(g_j\cdot g_k)$	$(g_i\cdot g_j)\cdot g_k=g_i\cdot(g_j\cdot g_k)$
gilmore-lie-groups_ E0002	18	■ 1.000	$g_i\cdot I=g_i=I\cdot g_i$	$g_i\cdot I=g_i=I\cdot g_i$	$g_i\cdot I=g_i=I\cdot g_i$
gilmore-lie-groups_ E0003	18	■ 1.000	$g_i\cdot g_i^{-1}=I=g_i^{-1}\cdot g_i$	$g_i\cdot g_i^{-1}=I=g_i^{-1}\cdot g_i$	$g_i\cdot g_i^{-1}=I=g_i^{-1}\cdot g_i$
gilmore-lie-groups_ E0005 (1.2)	18	■ 1.000	$\left[\begin{array}{c}z_1\\z_2\\\vdots\\z_n\end{array}\right]\longrightarrow\left[\begin{array}{c}z_{i_1}\\z_{i_2}\\\vdots\\z_{i_n}\end{array}\right]$	$\left[\begin{array}{c}z_1\\z_2\\\vdots\\z_n\end{array}\right]\longrightarrow\left[\begin{array}{c}z_{i_1}\\z_{i_2}\\\vdots\\z_{i_n}\end{array}\right]$	$\left[\begin{array}{c}z_1\\z_2\\\vdots\\z_n\end{array}\right]\longrightarrow\left[\begin{array}{c}z_{i_1}\\z_{i_2}\\\vdots\\z_{i_n}\end{array}\right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0007 (1.4)	19	■ 1.000	$\left[\begin{array}{l} z_1 \\ z_2 \\ z_3 \end{array}\right] \xrightarrow{(123)} \left[\begin{array}{l} z_2 \\ z_3 \\ z_1 \end{array}\right]$	$\begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix} \xrightarrow{(123)} \begin{bmatrix} z_2 \\ z_3 \\ z_1 \end{bmatrix}$	$\begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix} \xrightarrow{(123)} \begin{bmatrix} z_2 \\ z_3 \\ z_1 \end{bmatrix}$
gilmore-lie-groups_ EQ0008 (1.5)	19	■ 1.000	$\left[\begin{array}{l} z_2 \\ z_3 \\ z_1 \end{array}\right] = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix}$	$\begin{bmatrix} z_2 \\ z_3 \\ z_1 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix}$	$\begin{bmatrix} z_2 \\ z_3 \\ z_1 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix}$
gilmore-lie-groups_ EQ0011 (1.7)	20	■ 1.000	$g_i \cdot g_j = g_j \cdot g_i$	$g_i \cdot g_j = g_j \cdot g_i$	$g_i \cdot g_j = g_j \cdot g_i$
gilmore-lie-groups_ EQ0013 (1.9)	20	■ 1.000	$g H_1 g^{-1} = H_2$	$g H_1 g^{-1} = H_2$	$g H_1 g^{-1} = H_2$
gilmore-lie-groups_ EQ0014 (1.10)	20	■ 1.000	$(23) S_2(12)(23)^{-1} = (23)\{I, (12)\}(23)^{-1} = \{I, (13)\} = S_2(13)$	$(23)S_2(12)(23)^{-1} = (23)\{I, (12)\}(23)^{-1} = \{I, (13)\} = S_2(13)$	$(23)S_2(12)(23)^{-1} = (23)\{I, (12)\}(23)^{-1} = \{I, (13)\} = S_2(13)$
gilmore-lie-groups_ EQ0015 (1.11)	20	■ 1.000	$(23) A_3(23)^{-1} = (23)\{I, (123), (321)\}(23)^{-1} = \{I, (321), (123)\} = A_3$	$(23)A_3(23)^{-1} = (23)\{I, (123), (321)\}(23)^{-1} = \{I, (321), (123)\} = A_3$	$(23)A_3(23)^{-1} = (23)\{I, (123), (321)\}(23)^{-1} = \{I, (321), (123)\} = A_3$
gilmore-lie-groups_ EQ0020 (1.16)	23	■ 1.000	$G = G_0 \supset G_1 \supset \dots \supset G_\omega = I$	$G = G_0 \supset G_1 \supset \dots \supset G_\omega = I$	$G = G_0 \supset G_1 \supset \dots \supset G_\omega = I$
gilmore-lie-groups_ EQ0021 (1.17)	24	■ 1.000	$\begin{aligned} &\left(z - r_1\right)\left(z - r_2\right) = z^2 - I_1 z + I_2 = 0 \\ &I_1 = r_1 + r_2 \\ &I_2 = r_1 r_2 \end{aligned}$	$\begin{aligned} (z - r_1)(z - r_2) &= z^2 - I_1 z + I_2 = 0 \\ I_1 &= r_1 + r_2 \\ I_2 &= r_1 r_2 \end{aligned}$	$\begin{aligned} (z - r_1)(z - r_2) &= z^2 - I_1 z + I_2 = 0 \\ I_1 &= r_1 + r_2 \\ I_2 &= r_1 r_2 \end{aligned}$
gilmore-lie-groups_ EQ0023 (1.19)	24	■ 1.000	$\left[\begin{array}{l} u_1 \\ u_2 \end{array}\right] = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} r_1 \\ r_2 \end{bmatrix} = \begin{bmatrix} r_1 + r_2 \\ r_1 - r_2 \end{bmatrix}$	$\begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} r_1 \\ r_2 \end{bmatrix} = \begin{bmatrix} r_1 + r_2 \\ r_1 - r_2 \end{bmatrix}$	$\begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} r_1 \\ r_2 \end{bmatrix} = \begin{bmatrix} r_1 + r_2 \\ r_1 - r_2 \end{bmatrix}$
gilmore-lie-groups_ EQ0025 (1.21)	24	■ 1.000	$\begin{aligned} &(r_1 - r_2)^2 = r_1^2 - 2r_1 r_2 + r_2^2 \\ &= r_1^2 + 2r_1 r_2 + r_2^2 - 4r_1 r_2 = I_1^2 - 4I_2 = D \end{aligned}$	$\begin{aligned} (r_1 - r_2)^2 &= r_1^2 - 2r_1 r_2 + r_2^2 \\ &= r_1^2 + 2r_1 r_2 + r_2^2 - 4r_1 r_2 = I_1^2 - 4I_2 = D \end{aligned}$	$\begin{aligned} (r_1 - r_2)^2 &= r_1^2 - 2r_1 r_2 + r_2^2 \\ &= r_1^2 + 2r_1 r_2 + r_2^2 - 4r_1 r_2 = I_1^2 - 4I_2 = D \end{aligned}$
gilmore-lie-groups_ EQ0027 (1.23)	25	■ 1.000	$\left[\begin{array}{l} r_1 \\ r_2 \end{array}\right] = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} I_1 \\ \pm \sqrt{D} \end{bmatrix}$	$\begin{bmatrix} r_1 \\ r_2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} I_1 \\ \pm \sqrt{D} \end{bmatrix}$	$\begin{bmatrix} r_1 \\ r_2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} I_1 \\ \pm \sqrt{D} \end{bmatrix}$
gilmore-lie-groups_ EQ0028 (1.24)	25	■ 1.000	$r_1, r_2 = \frac{1}{2} (I_1 \pm \sqrt{D})$	$r_1, r_2 = \frac{1}{2} (I_1 \pm \sqrt{D})$	$r_1, r_2 = \frac{1}{2} (I_1 \pm \sqrt{D})$
gilmore-lie-groups_ EQ0029 (1.25)	25	■ 1.000	$z = x + \frac{1}{2} I_1$	$z = x + \frac{1}{2} I_1$	$z = x + \frac{1}{2} I_1$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0030 (1.26)	25	■ 1.000	$\begin{aligned} x^2 - I_1 x + I_2 &= x^2 + I_2 = 0 \\ x^2 + I_2 &= 0 \quad \text{if } I_1 = 0 \\ I_2 &= I_2 - \left(\frac{1}{2}I_1\right)^2 \end{aligned}$	$\begin{aligned} x^2 - I_1 x + I_2 &= x^2 + I_2 = 0 \\ I_1 &= 0 \\ I_2 &= I_2 - \left(\frac{1}{2}I_1\right)^2 \end{aligned}$	$\begin{aligned} x^2 - I_1 x + I_2 &= x^2 + I_2 = 0 \\ I_1 &= 0 \\ I_2 &= I_2 - \left(\frac{1}{2}I_1\right)^2 \end{aligned}$
gilmore-lie-groups_ EQ0031 (1.27)	25	■ 1.000	$x = \pm \sqrt{-I_2}$	$x = \pm \sqrt{-I_2}$	$x = \pm \sqrt{-I_2}$
gilmore-lie-groups_ EQ0032 (1.28)	25	■ 1.000	$r_{1,2} = \frac{1}{2} \left(I_1 \pm \sqrt{I_1^2 - 4I_2} \right)$	$r_{1,2} = \frac{1}{2} \left(I_1 \pm \sqrt{I_1^2 - 4I_2} \right)$	$r_{1,2} = \frac{1}{2} \left(I_1 \pm \sqrt{I_1^2 - 4I_2} \right)$
gilmore-lie-groups_ EQ0033 (1.29)	25	■ 1.000	$\begin{aligned} &\left(z - s_1 \right) \left(z - s_2 \right) \left(z - s_3 \right) = z^3 - I_1 z^2 + I_2 z - I_3 = 0 \\ &I_1 = s_1 + s_2 + s_3 \\ &I_2 = s_1 s_2 + s_1 s_3 + s_2 s_3 \\ &I_3 = s_1 s_2 s_3 \end{aligned}$	$\begin{aligned} &\left(z - s_1 \right) \left(z - s_2 \right) \left(z - s_3 \right) = z^3 - I_1 z^2 + I_2 z - I_3 = 0 \\ &I_1 = s_1 + s_2 + s_3 \\ &I_2 = s_1 s_2 + s_1 s_3 + s_2 s_3 \\ &I_3 = s_1 s_2 s_3 \end{aligned}$	$\begin{aligned} &\left(z - s_1 \right) \left(z - s_2 \right) \left(z - s_3 \right) = z^3 - I_1 z^2 + I_2 z - I_3 = 0 \\ &I_1 = s_1 + s_2 + s_3 \\ &I_2 = s_1 s_2 + s_1 s_3 + s_2 s_3 \\ &I_3 = s_1 s_2 s_3 \end{aligned}$
gilmore-lie-groups_ EQ0034 (1.31)	26	■ 1.000	$\omega^3 = +1 \quad \omega = e^{2\pi i/3} = \frac{-1 + i\sqrt{3}}{2}$	$\omega^3 = +1 \quad \omega = e^{2\pi i/3} = \frac{-1 + i\sqrt{3}}{2}$	$\omega^3 = +1 \quad \omega = e^{2\pi i/3} = \frac{-1 + i\sqrt{3}}{2}$
gilmore-lie-groups_ EQ0036 (1.33)	26	■ 1.000	$\begin{aligned} &(123)^{-1} v_2 = (321) v_2 = (321) (s_1 + \omega s_2 + \omega^2 s_3) \\ &= s_3 + \omega s_1 + \omega^2 s_2 = \omega (s_1 + \omega s_2 + \omega^2 s_3) = \omega v_2 \end{aligned}$	$\begin{aligned} &(123)^{-1} v_2 = (321) v_2 = (321) (s_1 + \omega s_2 + \omega^2 s_3) \\ &= s_3 + \omega s_1 + \omega^2 s_2 = \omega (s_1 + \omega s_2 + \omega^2 s_3) = \omega v_2 \end{aligned}$	$\begin{aligned} &(123)^{-1} v_2 = (321) v_2 = (321) (s_1 + \omega s_2 + \omega^2 s_3) \\ &= s_3 + \omega s_1 + \omega^2 s_2 = \omega (s_1 + \omega s_2 + \omega^2 s_3) = \omega v_2 \end{aligned}$
gilmore-lie-groups_ EQ0037 (1.34)	27	■ 1.000	$v_1 = I_1$	$v_1 = I_1$	$v_1 = I_1$
gilmore-lie-groups_ EQ0038 (1.35)	27	■ 1.000	$\begin{aligned} &(12) \left(v_2 \right)^3 = (12) (s_1 + \omega s_2 + \omega^2 s_3)^3 = (s_2 + \omega s_1 + \omega^2 s_3)^3 \\ &= \omega^3 (s_1 + \omega^2 s_2 + \omega s_3)^3 = (v_3)^3 \\ &(12) \left(v_3 \right)^3 = (12) (s_1 + \omega^2 s_2 + \omega s_3)^3 = (s_2 + \omega^2 s_1 + \omega s_3)^3 \\ &= \omega^6 (s_1 + \omega s_2 + \omega^2 s_3)^3 = (v_2)^3 \end{aligned}$	$\begin{aligned} &(12) \left(v_2 \right)^3 = (12) (s_1 + \omega s_2 + \omega^2 s_3)^3 = (s_2 + \omega s_1 + \omega^2 s_3)^3 \\ &= \omega^3 (s_1 + \omega^2 s_2 + \omega s_3)^3 = (v_3)^3 \\ &(12) \left(v_3 \right)^3 = (12) (s_1 + \omega^2 s_2 + \omega s_3)^3 = (s_2 + \omega^2 s_1 + \omega s_3)^3 \\ &= \omega^6 (s_1 + \omega s_2 + \omega^2 s_3)^3 = (v_2)^3 \end{aligned}$	$\begin{aligned} &(12) \left(v_2 \right)^3 = (12) (s_1 + \omega s_2 + \omega^2 s_3)^3 = (s_2 + \omega s_1 + \omega^2 s_3)^3 \\ &= \omega^3 (s_1 + \omega^2 s_2 + \omega s_3)^3 = (v_3)^3 \\ &(12) \left(v_3 \right)^3 = (12) (s_1 + \omega^2 s_2 + \omega s_3)^3 = (s_2 + \omega^2 s_1 + \omega s_3)^3 \\ &= \omega^6 (s_1 + \omega s_2 + \omega^2 s_3)^3 = (v_2)^3 \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0039 (1.36)	27	■ 1.000	\begin{aligned} \left(x-v_2^3\right)\left(x-v_3^3\right) &=x^2-J_1 x+J_2=0 \\ \& \ J_1 &=v_2^3+v_3^3 \\ \& \ J_2 &=v_2^3 v_3^3 \end{aligned}	$\begin{aligned} (x-v_2^3)(x-v_3^3) &=x^2-J_1 x+J_2=0 \\ J_1 &=v_2^3+v_3^3 \\ J_2 &=v_2^3 v_3^3 \end{aligned}$	$\begin{aligned} (x-v_2^3)(x-v_3^3) &=x^2-J_1 x+J_2=0 \\ J_1 &=v_2^3+v_3^3 \\ J_2 &=v_2^3 v_3^3 \end{aligned}$
gilmore-lie-groups_ EQ0041 (1.38)	27	■ 1.000	z=y+\frac{1}{3} I_1	$z=y+\frac{1}{3} I_1$	$z=y+\frac{1}{3} I_1$
gilmore-lie-groups_ EQ0043 (1.40)	28	■ 1.000	\begin{aligned} \& \ J_1 &=v_2^3+v_3^3=-27 I_3^{\prime} \\ \& \ J_2 &=v_2^3 v_3^3=-27 I_2^{\prime 3} \end{aligned}	$\begin{aligned} J_1 &=v_2^3+v_3^3=-27 I_3' \\ J_2 &=v_2^3 v_3^3=-27 I_2'^3 \end{aligned}$	$\begin{aligned} J_1 &=v_2^3+v_3^3=-27 I_3' \\ J_2 &=v_2^3 v_3^3=-27 I_2'^3 \end{aligned}$
gilmore-lie-groups_ EQ0044 (1.41)	28	■ 1.000	x^2-\left(-27 I_3^{\prime}\right) x+\left(-27 I_2^{\prime 3}\right)=0	$x^2-(-27 I_3') x+(-27 I_2'^3)=0$	$x^2-(-27 I_3') x+(-27 I_2'^3)=0$
gilmore-lie-groups_ EQ0049 (1.45)	29	■ 1.000	\begin{aligned} \left(z-t_1\right)\left(z-t_2\right)\left(z-t_3\right)\left(z-t_4\right) &=z^4-I_1 z^3+I_2 z^2-I_3 z+I_4=0 \\ \& \ I_1 &=t_1+t_2+t_3+t_4 \\ \& \ I_2 &=t_1 t_2+t_1 t_3+t_1 t_4+t_2 t_3+t_2 t_4+t_3 t_4 \\ \& \ I_3 &=t_1 t_2 t_3+t_1 t_2 t_4+t_1 t_3 t_4+t_2 t_3 t_4 \\ \& \ I_4 &=t_1 t_2 t_3 t_4 \end{aligned}	$\begin{aligned} (z-t_1)(z-t_2)(z-t_3)(z-t_4) &=z^4-I_1 z^3+I_2 z^2-I_3 z+I_4=0 \\ I_1 &=t_1+t_2+t_3+t_4 \\ I_2 &=t_1 t_2+t_1 t_3+t_1 t_4+t_2 t_3+t_2 t_4+t_3 t_4 \\ I_3 &=t_1 t_2 t_3+t_1 t_2 t_4+t_1 t_3 t_4+t_2 t_3 t_4 \\ I_4 &=t_1 t_2 t_3 t_4 \end{aligned}$	$\begin{aligned} (z-t_1)(z-t_2)(z-t_3)(z-t_4) &=z^4-I_1 z^3+I_2 z^2-I_3 z+I_4=0 \\ I_1 &=t_1+t_2+t_3+t_4 \\ I_2 &=t_1 t_2+t_1 t_3+t_1 t_4+t_2 t_3+t_2 t_4+t_3 t_4 \\ I_3 &=t_1 t_2 t_3+t_1 t_2 t_4+t_1 t_3 t_4+t_2 t_3 t_4 \\ I_4 &=t_1 t_2 t_3 t_4 \end{aligned}$
gilmore-lie-groups_ EQ0051	29	■ 1.000	S_4 \supset A_4 \supset V_4 \supset I	$S_4 \supset A_4 \supset V_4 \supset I$	$S_4 \supset A_4 \supset V_4 \supset I$
gilmore-lie-groups_ EQ0052 (1.48)	30	■ 1.000	\left[\begin{array}{l} w_1 \\ w_2 \\ w_3 \\ w_4 \end{array}\right]=\left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{array}\right]\left[\begin{array}{l} t_1 \\ t_2 \\ t_3 \\ t_4 \end{array}\right]=\left[\begin{array}{l} t_1+t_2+t_3+t_4 \\ t_1+t_2-t_3-t_4 \\ t_1-t_2+t_3-t_4 \\ t_1-t_2-t_3+t_4 \end{array}\right]	$\left[\begin{array}{l} w_1 \\ w_2 \\ w_3 \\ w_4 \end{array}\right]=\left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{array}\right]\left[\begin{array}{l} t_1 \\ t_2 \\ t_3 \\ t_4 \end{array}\right]=\left[\begin{array}{l} t_1+t_2+t_3+t_4 \\ t_1+t_2-t_3-t_4 \\ t_1-t_2+t_3-t_4 \\ t_1-t_2-t_3+t_4 \end{array}\right]$	$\left[\begin{array}{l} w_1 \\ w_2 \\ w_3 \\ w_4 \end{array}\right]=\left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{array}\right]\left[\begin{array}{l} t_1 \\ t_2 \\ t_3 \\ t_4 \end{array}\right]=\left[\begin{array}{l} t_1+t_2+t_3+t_4 \\ t_1+t_2-t_3-t_4 \\ t_1-t_2+t_3-t_4 \\ t_1-t_2-t_3+t_4 \end{array}\right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0062 (1.57)	33	■ 1.000	$\left[\begin{array}{l} w_1 \\ w_2 \\ w_3 \\ w_4 \end{array}\right] = \left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{array}\right] \left[\begin{array}{l} t_1' \\ t_2' \\ t_3' \\ t_4' \end{array}\right] \rightarrow \left[\begin{array}{l} 0 \\ -10 \\ -4 \\ 2 \end{array}\right]$		$\begin{bmatrix} w_1 \\ w_2 \\ w_3 \\ w_4 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} t_1' \\ t_2' \\ t_3' \\ t_4' \end{bmatrix} \rightarrow \begin{bmatrix} 0 \\ -10 \\ -4 \\ 2 \end{bmatrix}$
gilmore-lie-groups_ EQ0065 (1.60)	33	■ 1.000	$\left(y-2^2\right)\left(y-4^2\right)\left(y-10^2\right)=y^3-120y^2+2064y-6400=0$	$(y-2^2)(y-4^2)(y-10^2)=y^3-120y^2+2064y-6400=0$	$(y-2^2)(y-4^2)(y-10^2)=y^3-120y^2+2064y-6400=0$
gilmore-lie-groups_ EQ0068 (1.63)	33	■ 1.000	$\left(y^{\prime}+36\right)\left(y^{\prime}+24\right)\left(y^{\prime}-60\right)=0$	$(y'+36)(y'+24)(y'-60)=0$	$(y'+36)(y'+24)(y'-60)=0$
gilmore-lie-groups_ EQ0074 (1.69)	34	■ 1.000	$\begin{aligned} &\left[\begin{array}{l} x_1 \\ x_2 \end{array}\right] = \frac{1}{2} \left[\begin{array}{cc} 1 & 1 \\ 1 & -1 \end{array}\right] \left[\begin{array}{l} K_1=1399680 \\ 2\sqrt{-K_2}=i \times 290304\sqrt{3} \end{array}\right] \\ &= 699840 \pm i \times 145152\sqrt{3} \end{aligned}$	$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} K_1 = 1399680 \\ 2\sqrt{-K_2} = i \times 290304\sqrt{3} \end{bmatrix} \\ = 699840 \pm i \times 145152\sqrt{3}$	$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} K_1 = 1399680 \\ 2\sqrt{-K_2} = i \times 290304\sqrt{3} \end{bmatrix} \\ = 699840 \pm i \times 145152\sqrt{3}$
gilmore-lie-groups_ EQ0076 (1.71)	35	■ 1.000	$\begin{aligned} &v_2 = -36 + i48\sqrt{3} \\ &v_3 = -36 - i48\sqrt{3} \end{aligned}$	$v_2 = -36 + i48\sqrt{3}$ $v_3 = -36 - i48\sqrt{3}$	$v_2 = -36 + i48\sqrt{3}$ $v_3 = -36 - i48\sqrt{3}$
gilmore-lie-groups_ EQ0078 (1.74)	35	■ 1.000	$\left[\begin{array}{l} x_1 \\ x_2 \\ x_3 \\ x_4 \end{array}\right] = \frac{1}{4} \left[\begin{array}{cccc} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{array}\right] \left[\begin{array}{l} I_1=4 \\ w_2=4 \\ w_3=10 \\ w_4=2 \end{array}\right] = \left[\begin{array}{l} 20/4=+5 \\ -4/4=-1 \\ 8/4=+2 \\ -8/4=-2 \end{array}\right]$	$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} I_1 = 4 \\ w_2 = 4 \\ w_3 = 10 \\ w_4 = 2 \end{bmatrix} = \begin{bmatrix} 20/4 = +5 \\ -4/4 = -1 \\ 8/4 = +2 \\ -8/4 = -2 \end{bmatrix}$	$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} I_1 = 4 \\ w_2 = 4 \\ w_3 = 10 \\ w_4 = 2 \end{bmatrix} = \begin{bmatrix} 20/4 = +5 \\ -4/4 = -1 \\ 8/4 = +2 \\ -8/4 = -2 \end{bmatrix}$
gilmore-lie-groups_ EQ0080	36	■ 1.000	$\begin{aligned} &e^{i3\theta} = (e^{i\theta})^3 \\ &\cos(3\theta) + i\sin(3\theta) = (\cos^3(\theta) - 3\cos(\theta)\sin^2(\theta)) \\ &\quad + i(3\cos^2(\theta)\sin(\theta) - \sin^3(\theta)) \end{aligned}$	$e^{i3\theta} = (e^{i\theta})^3$ $\cos(3\theta) + i\sin(3\theta) = (\cos^3(\theta) - 3\cos(\theta)\sin^2(\theta)) \\ + i(3\cos^2(\theta)\sin(\theta) - \sin^3(\theta))$	$e^{i3\theta} = (e^{i\theta})^3$ $\cos(3\theta) + i\sin(3\theta) = (\cos^3(\theta) - 3\cos(\theta)\sin^2(\theta)) \\ + i(3\cos^2(\theta)\sin(\theta) - \sin^3(\theta))$
gilmore-lie-groups_ EQ0081	37	■ 1.000	$\cos(3\theta) = 4\cos^3(\theta) - 3\cos(\theta)$	$\cos(3\theta) = 4\cos^3(\theta) - 3\cos(\theta)$	$\cos(3\theta) = 4\cos^3(\theta) - 3\cos(\theta)$
gilmore-lie-groups_ EQ0082	37	■ 1.000	$4\cos^3(\theta) - 3\cos(\theta) - \cos(3\theta) = 0$	$4\cos^3(\theta) - 3\cos(\theta) - \cos(3\theta) = 0$	$4\cos^3(\theta) - 3\cos(\theta) - \cos(3\theta) = 0$
gilmore-lie-groups_ EQ0083	38	■ 1.000	$(g_i \circ g_j) \circ g_k = g_i \circ (g_j \circ g_k)$	$(g_i \circ g_j) \circ g_k = g_i \circ (g_j \circ g_k)$	$(g_i \circ g_j) \circ g_k = g_i \circ (g_j \circ g_k)$
gilmore-lie-groups_ EQ0084	38	■ 1.000	$g_i \circ e = g_i = e \circ g_i$	$g_i \circ e = g_i = e \circ g_i$	$g_i \circ e = g_i = e \circ g_i$
gilmore-lie-groups_ EQ0085	38	■ 1.000	$g_i \circ g_i^{-1} = e = g_i^{-1} \circ g_i$	$g_i \circ g_i^{-1} = e = g_i^{-1} \circ g_i$	$g_i \circ g_i^{-1} = e = g_i^{-1} \circ g_i$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0088	39	■ 1.000	$e \longrightarrow I_2 = \left[\begin{array}{ll} 1 & 0 \\ 0 & 1 \end{array} \right]$	$e \longrightarrow I_2 = \left[\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array} \right]$	$e \longrightarrow I_2 = \left[\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array} \right]$
gilmore-lie-groups_ EQ0089	39	■ 1.000	$\left[\begin{array}{ll} A_{11} & A_{12} \\ A_{21} & A_{22} \end{array} \right] \longrightarrow \left[\begin{array}{ll} A_{11} & A_{12} \\ A_{21} & A_{22} \end{array} \right]^{-1} = \frac{1}{A_{11}A_{22} - A_{12}A_{21}} \left[\begin{array}{cc} A_{22} & -A_{12} \\ -A_{21} & A_{11} \end{array} \right]$	$\left[\begin{array}{cc} A_{11} & A_{12} \\ A_{21} & A_{22} \end{array} \right] \rightarrow \left[\begin{array}{cc} A_{11} & A_{12} \\ A_{21} & A_{22} \end{array} \right]^{-1} = \frac{1}{A_{11}A_{22} - A_{12}A_{21}} \left[\begin{array}{cc} A_{22} & -A_{12} \\ -A_{21} & A_{11} \end{array} \right]$	$\left[\begin{array}{cc} A_{11} & A_{12} \\ A_{21} & A_{22} \end{array} \right] \rightarrow \left[\begin{array}{cc} A_{11} & A_{12} \\ A_{21} & A_{22} \end{array} \right]^{-1} = \frac{1}{A_{11}A_{22} - A_{12}A_{21}} \left[\begin{array}{cc} A_{22} & -A_{12} \\ -A_{21} & A_{11} \end{array} \right]$
gilmore-lie-groups_ EQ0090 (2.3)	40	■ 1.000	$\left(x_1, x_2, x_3 \right) \longrightarrow \left[\begin{array}{cc} x_1 & x_2 \\ x_3 & \frac{1+x_2x_3}{x_1} \end{array} \right] \quad x_1 \neq 0$	$(x_1, x_2, x_3) \longrightarrow \left[\begin{array}{cc} x_1 & x_2 \\ x_3 & \frac{1+x_2x_3}{x_1} \end{array} \right] \quad x_1 \neq 0$	$(x_1, x_2, x_3) \longrightarrow \left[\begin{array}{cc} x_1 & x_2 \\ x_3 & \frac{1+x_2x_3}{x_1} \end{array} \right] \quad x_1 \neq 0$
gilmore-lie-groups_ EQ0093 (2.5)	42	■ 1.000	$\begin{aligned} & z_1 = \phi_1(x_1, x_2, x_3; y_1, y_2, y_3) = x_1y_1 + x_2y_3 \\ & z_2 = \phi_2(x_1, x_2, x_3; y_1, y_2, y_3) = x_1y_2 + x_2 \frac{1+y_2y_3}{y_1} \\ & z_3 = \phi_3(x_1, x_2, x_3; y_1, y_2, y_3) = x_3y_1 + \frac{1+x_2x_3}{x_1} y_3 \end{aligned}$	$\begin{aligned} z_1 &= \phi_1(x_1, x_2, x_3; y_1, y_2, y_3) = x_1y_1 + x_2y_3 \\ z_2 &= \phi_2(x_1, x_2, x_3; y_1, y_2, y_3) = x_1y_2 + x_2 \frac{1+y_2y_3}{y_1} \\ z_3 &= \phi_3(x_1, x_2, x_3; y_1, y_2, y_3) = x_3y_1 + \frac{1+x_2x_3}{x_1} y_3 \end{aligned}$	$\begin{aligned} z_1 &= \phi_1(x_1, x_2, x_3; y_1, y_2, y_3) = x_1y_1 + x_2y_3 \\ z_2 &= \phi_2(x_1, x_2, x_3; y_1, y_2, y_3) = x_1y_2 + x_2 \frac{1+y_2y_3}{y_1} \\ z_3 &= \phi_3(x_1, x_2, x_3; y_1, y_2, y_3) = x_3y_1 + \frac{1+x_2x_3}{x_1} y_3 \end{aligned}$
gilmore-lie-groups_ EQ0095 (2.7)	43	■ 1.000	$\begin{aligned} & \psi_1(x_1, x_2, x_3) = y_1 = (1+x_2x_3)/x_1 \\ & \psi_2(x_1, x_2, x_3) = y_2 = -x_2 \\ & \psi_3(x_1, x_2, x_3) = y_3 = -x_3 \end{aligned}$	$\begin{aligned} \psi_1(x_1, x_2, x_3) &= y_1 = (1+x_2x_3)/x_1 \\ \psi_2(x_1, x_2, x_3) &= y_2 = -x_2 \\ \psi_3(x_1, x_2, x_3) &= y_3 = -x_3 \end{aligned}$	$\begin{aligned} \psi_1(x_1, x_2, x_3) &= y_1 = (1+x_2x_3)/x_1 \\ \psi_2(x_1, x_2, x_3) &= y_2 = -x_2 \\ \psi_3(x_1, x_2, x_3) &= y_3 = -x_3 \end{aligned}$
gilmore-lie-groups_ EQ0099	45	■ 1.000	$\left[\begin{array}{cccc} \pm 1 & 0 & 0 & 0 \\ 0 & \pm 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right]$	$\left[\begin{array}{cccc} \pm 1 & 0 & 0 & 0 \\ 0 & \pm 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right]$	$\left[\begin{array}{cccc} \pm 1 & 0 & 0 & 0 \\ 0 & \pm 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array} \right]$
gilmore-lie-groups_ EQ0101	46	■ 1.000	$x \longrightarrow x' = (a, b, c, d) x = \frac{ax+b}{cx+d}$	$x \rightarrow x' = (a, b, c, d)x = \frac{ax+b}{cx+d}$	$x \rightarrow x' = (a, b, c, d)x = \frac{ax+b}{cx+d}$
gilmore-lie-groups_ EQ0102	46	■ 1.000	$(A, B, C, D) = (a', b', c', d') \circ (a, b, c, d)$	$(A, B, C, D) = (a', b', c', d') \circ (a, b, c, d)$	$(A, B, C, D) = (a', b', c', d') \circ (a, b, c, d)$
gilmore-lie-groups_ EQ0103	46	■ 1.000	$\left[\begin{array}{l} y' \\ z' \end{array} \right] = \left[\begin{array}{cc} a & b \\ c & d \end{array} \right] \left[\begin{array}{l} y \\ z \end{array} \right] \Rightarrow x' = \frac{y'}{z'} = \frac{a(y/z) + b}{c(y/z) + d} = \frac{ax+b}{cx+d}$	$\left[\begin{array}{l} y' \\ z' \end{array} \right] = \left[\begin{array}{cc} a & b \\ c & d \end{array} \right] \left[\begin{array}{l} y \\ z \end{array} \right] \Rightarrow x' = \frac{y'}{z'} = \frac{a(y/z) + b}{c(y/z) + d} = \frac{ax+b}{cx+d}$	$\left[\begin{array}{l} y' \\ z' \end{array} \right] = \left[\begin{array}{cc} a & b \\ c & d \end{array} \right] \left[\begin{array}{l} y \\ z \end{array} \right] \Rightarrow x' = \frac{y'}{z'} = \frac{a(y/z) + b}{c(y/z) + d} = \frac{ax+b}{cx+d}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0130 (3.16)	54	■ 1.000	$a_{1}^{2}+b_{1}^{2}+a_{2}^{2}+b_{2}^{2}=1$	$a_1^2 + b_1^2 + a_2^2 + b_2^2 = 1$	$a_1^2 + b_1^2 + a_2^2 + b_2^2 = 1$
gilmore-lie-groups_ EQ0134 (3.20)	55	■ 1.000	$M^{\mathrm{t}}GM=G\quad \mathbb{F}=\begin{cases}\mathbb{R} & \mathrm{Sp}(N,\mathbb{R}) \\ \mathbb{C} & \mathrm{Sp}(N,\mathbb{C})\end{cases}$	$M^tGM = G \quad \mathbb{F} = \begin{cases} \mathbb{R} & \mathrm{Sp}(N, \mathbb{R}) \\ \mathbb{C} & \mathrm{Sp}(N, \mathbb{C}) \end{cases}$	$M^tGM = G \quad \mathbb{F} = \begin{cases} \mathbb{R} & \mathrm{Sp}(N, \mathbb{R}) \\ \mathbb{C} & \mathrm{Sp}(N, \mathbb{C}) \end{cases}$
gilmore-lie-groups_ EQ0135	55	■ 1.000	$G=\left[\begin{array}{cc} 0 & I_n \\ -I_n & 0 \end{array}\right]$	$G = \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}$	$G = \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}$
gilmore-lie-groups_ EQ0136	55	■ 1.000	$G=\sum_{\alpha=1}^n\oplus\left[i\,\sigma_y\right]_{\alpha}$	$G = \sum_{\alpha=1}^n \oplus [i\sigma_y]_{\alpha}$	$G = \sum_{\alpha=1}^n \oplus [i\sigma_y]_{\alpha}$
gilmore-lie-groups_ EQ0137 (3.21)	55	■ 1.000	$\left[\begin{array}{ll} a & c \\ b & d \end{array}\right]\left[\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right]\left[\begin{array}{ll} a & b \\ c & d \end{array}\right]=\left[\begin{array}{cc} 0 & ad-bc \\ bc-ad & 0 \end{array}\right]=\left[\begin{array}{ll} 0 & 1 \\ -1 & 0 \end{array}\right]$	$\begin{bmatrix} a & c \\ b & d \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 0 & ad-bc \\ bc-ad & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$	$\begin{bmatrix} a & c \\ b & d \end{bmatrix} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 0 & ad-bc \\ bc-ad & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$
gilmore-lie-groups_ EQ0139 (3.23)	55	■ 1.000	$G=\left[\begin{array}{cc} I_3 & 0 \\ 0 & 0 \end{array}\right]$	$G = \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix}$	$G = \begin{bmatrix} I_3 & 0 \\ 0 & 0 \end{bmatrix}$
gilmore-lie-groups_ EQ0147 (3.30)	57	■ 1.000	$SU(1,1)=U(1,1)\cap SL(2;\mathbb{C})\rightarrow\left[\begin{array}{cc} a & b \\ b^* & a^* \end{array}\right]$	$SU(1,1) = U(1,1) \cap SL(2; \mathbb{C}) \rightarrow \begin{bmatrix} a & b \\ b^* & a^* \end{bmatrix}$	$SU(1,1) = U(1,1) \cap SL(2; \mathbb{C}) \rightarrow \begin{bmatrix} a & b \\ b^* & a^* \end{bmatrix}$
gilmore-lie-groups_ EQ0150 (3.33)	58	■ 1.000	$\left[\begin{array}{ll} 1 & 1 \\ 1 & 0 \end{array}\right]\in GL(n;\mathbb{Z})$	$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \in GL(n; \mathbb{Z})$	$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \in GL(n; \mathbb{Z})$
gilmore-lie-groups_ EQ0151 (3.34)	59	■ 1.000	$\left[\begin{array}{ll} 1 & 1 \\ 1 & 0 \end{array}\right]^n=\left[\begin{array}{cc} F(n+1) & F(n) \\ F(n) & F(n-1) \end{array}\right]$	$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^n = \begin{bmatrix} F(n+1) & F(n) \\ F(n) & F(n-1) \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^n = \begin{bmatrix} F(n+1) & F(n) \\ F(n) & F(n-1) \end{bmatrix}$
gilmore-lie-groups_ EQ0153 (3.35)	59	■ 1.000	$\begin{aligned} & \left[\begin{array}{ll} 1 & 1 \\ 1 & 0 \end{array}\right]^{n+1} = \left[\begin{array}{ll} 1 & 1 \\ 1 & 0 \end{array}\right] \left[\begin{array}{cc} F(n+1) & F(n) \\ F(n) & F(n-1) \end{array}\right] \\ & = \left[\begin{array}{cc} F(n+1)+F(n) & F(n)+F(n-1) \\ F(n+1) & F(n) \end{array}\right] \\ & = \left[\begin{array}{cc} F(n+2) & F(n+1) \\ F(n+1) & F(n) \end{array}\right] \end{aligned}$	$\begin{aligned} & \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^{n+1} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} F(n+1) & F(n) \\ F(n) & F(n-1) \end{bmatrix} \\ & = \begin{bmatrix} F(n+1)+F(n) & F(n)+F(n-1) \\ F(n+1) & F(n) \end{bmatrix} \\ & = \begin{bmatrix} F(n+2) & F(n+1) \\ F(n+1) & F(n) \end{bmatrix} \end{aligned}$	$\begin{aligned} & \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^{n+1} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} F(n+1) & F(n) \\ F(n) & F(n-1) \end{bmatrix} \\ & = \begin{bmatrix} F(n+1)+F(n) & F(n)+F(n-1) \\ F(n+1) & F(n) \end{bmatrix} \\ & = \begin{bmatrix} F(n+2) & F(n+1) \\ F(n+1) & F(n) \end{bmatrix} \end{aligned}$
gilmore-lie-groups_ EQ0154 (3.36)	60	■ 1.000	$A_n=P_n\cap SL(n;\mathbb{Z})$	$A_n = P_n \cap SL(n; \mathbb{Z})$	$A_n = P_n \cap SL(n; \mathbb{Z})$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0155 (3.37)	60	■ 1.000	$\begin{array}{l} \pm\left[\begin{array}{ll} 1 & 0 \\ 0 & 1 \end{array}\right] \quad \pm\left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right] \quad \pm\left[\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array}\right] \quad \pm\left[\begin{array}{ll} 0 & 1 \\ -1 & 0 \end{array}\right] \\ \end{array}$		
gilmore-lie-groups_ EQ0156 (3.38)	60	■ 1.000	$\begin{array}{l} \begin{array}{l} \left[\begin{array}{lll} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array}\right] \left[\begin{array}{lll} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{array}\right] \left[\begin{array}{lll} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{array}\right] \\ \left[\begin{array}{lll} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{array}\right] \left[\begin{array}{lll} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{array}\right] \left[\begin{array}{lll} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{array}\right] \end{array} \\ \end{array}$		
gilmore-lie-groups_ EQ0158	61	■ 1.000	$\begin{array}{l} \begin{array}{l} \text{GL}(n; \mathbb{R}) = n^2 \quad \text{SL}(n; \mathbb{R}) = n^2 - 1 \\ \text{GL}(n; \mathbb{C}) = 2n^2 \quad \text{SL}(n; \mathbb{C}) = 2n^2 - 2 \\ \text{GL}(n; \mathbb{Q}) = 4n^2 \end{array} \end{array}$		
gilmore-lie-groups_ EQ0159	61	■ 1.000	$\begin{array}{l} \begin{array}{l} \text{SL}_1(n; \mathbb{C}) \det(M) = e^{i\phi} \quad 2n^2 - 1 \\ \text{SL}_2(n; \mathbb{C}) \det(M) = e^\lambda \quad 2n^2 - 1 \\ \text{SL}_3(n; \mathbb{C}) \det(M) = r \quad 2n^2 - 1 \\ \text{SL}(n; \mathbb{C}) \det(M) = +1 \quad 2n^2 - 2 \end{array} \end{array}$		
gilmore-lie-groups_ EQ0160	62	■ 1.000	$\left(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2}\right) A_\mu(x, t) = -\frac{4\pi}{c} j_\mu$		
gilmore-lie-groups_ EQ0164	63	■ 1.000	$SO(3) \longrightarrow \begin{bmatrix} m_{11} & x & y \\ m_{21} & m_{22} & z \\ m_{31} & m_{32} & m_{33} \end{bmatrix}$		
gilmore-lie-groups_ EQ0166	64	■ 1.000	$E = -E_0 Z^2 \{1 + \delta * (n - l - 1)\} / n^2$		
gilmore-lie-groups_ EQ0168	64	■ 1.000	$G_1 = \begin{bmatrix} I_n & 0 \\ 0 & I_n \end{bmatrix} \quad G_2 = \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00169	64	■ 1.000	$\begin{array}{ccc} G_{\{1\}} & \& G_{\{2\}} & \& G_{\{1\}} \end{array} \text{ and } G_{\{2\}} \setminus 0(2\,n\,;\,\mathbb{R}) \& \operatorname{Sp}(2\,n\,;\,\mathbb{R}) \& 0\,U(2\,n\,;\,\mathbb{R}) \end{array}$	$\begin{array}{ccc} G_1 & G_2 & G_1 \text{ and } G_2 \\ O(2n;\mathbb{R}) & \operatorname{Sp}(2n;\mathbb{R}) & OU(2n;\mathbb{R}) \end{array}$	$\begin{array}{ccc} G_1 & G_2 & G_1 \text{ and } G_2 \\ O(2n;\mathbb{R}) & \operatorname{Sp}(2n;\mathbb{R}) & OU(2n;\mathbb{R}) \end{array}$
gilmore-lie-groups_ E00171	64	■ 1.000	$\{f,g\}=\sum_k\frac{\partial f}{\partial q_k}\frac{\partial g}{\partial p_k}-\frac{\partial g}{\partial q_k}\frac{\partial f}{\partial p_k}$	$\{f,g\}=\sum_k\frac{\partial f}{\partial q_k}\frac{\partial g}{\partial p_k}-\frac{\partial g}{\partial q_k}\frac{\partial f}{\partial p_k}$	$\{f,g\}=\sum_k\frac{\partial f}{\partial q_k}\frac{\partial g}{\partial p_k}-\frac{\partial g}{\partial q_k}\frac{\partial f}{\partial p_k}$
gilmore-lie-groups_ E00173	65	■ 1.000	$\left[\begin{array}{l} \partial g/\partial Q \\ \partial g/\partial P \end{array}\right]=\left[\begin{array}{cc} A & B \\ C & D \end{array}\right]\left[\begin{array}{l} \partial g/\partial q \\ \partial g/\partial p \end{array}\right]$	$\left[\begin{array}{l} \partial g/\partial Q \\ \partial g/\partial P \end{array}\right]=\left[\begin{array}{cc} A & B \\ C & D \end{array}\right]\left[\begin{array}{l} \partial g/\partial q \\ \partial g/\partial p \end{array}\right]$	$\left[\begin{array}{l} \partial g/\partial Q \\ \partial g/\partial P \end{array}\right]=\left[\begin{array}{cc} A & B \\ C & D \end{array}\right]\left[\begin{array}{l} \partial g/\partial q \\ \partial g/\partial p \end{array}\right]$
gilmore-lie-groups_ E00174	65	■ 1.000	$\{u(q,p),v(q,p)\}\stackrel{\text{Leftrightarrow}}{\frac{[u(\hat{q},\hat{p}),v(\hat{q},\hat{p})]}{i\hbar}}$	$\{u(q,p),v(q,p)\}\leftrightarrow\frac{[u(\hat{q},\hat{p}),v(\hat{q},\hat{p})]}{i\hbar}$	$\{u(q,p),v(q,p)\}\leftrightarrow\frac{[u(\hat{q},\hat{p}),v(\hat{q},\hat{p})]}{i\hbar}$
gilmore-lie-groups_ E00176	65	■ 1.000	$ A_L ^2- B_L ^2= A_R ^2- B_R ^2$	$ A_L ^2- B_L ^2= A_R ^2- B_R ^2$	$ A_L ^2- B_L ^2= A_R ^2- B_R ^2$
gilmore-lie-groups_ E00179	66	■ 1.000	$\left[\begin{array}{ll} s_{\{11\}} & s_{\{12\}} \\ s_{\{21\}} & s_{\{22\}} \end{array}\right]=\left[\begin{array}{cc} \frac{1}{t_{11}} & -\frac{t_{12}}{t_{11}} \\ \frac{t_{21}}{t_{11}} & \frac{t_{11}t_{22}-t_{12}t_{21}}{t_{11}} \end{array}\right]$	$\left[\begin{array}{cc} s_{11} & s_{12} \\ s_{21} & s_{22} \end{array}\right]=\left[\begin{array}{cc} \frac{1}{t_{11}} & -\frac{t_{12}}{t_{11}} \\ \frac{t_{21}}{t_{11}} & \frac{t_{11}t_{22}-t_{12}t_{21}}{t_{11}} \end{array}\right]$	$\left[\begin{array}{cc} s_{11} & s_{12} \\ s_{21} & s_{22} \end{array}\right]=\left[\begin{array}{cc} \frac{1}{t_{11}} & -\frac{t_{12}}{t_{11}} \\ \frac{t_{21}}{t_{11}} & \frac{t_{11}t_{22}-t_{12}t_{21}}{t_{11}} \end{array}\right]$
gilmore-lie-groups_ E00180	66	■ 1.000	$\left[\begin{array}{l} i_2 \\ i_3 \end{array}\right]=\left[\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right]\left[\begin{array}{l} o_1 \\ o_4 \end{array}\right]$	$\left[\begin{array}{l} i_2 \\ i_3 \end{array}\right]=\left[\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right]\left[\begin{array}{l} o_1 \\ o_4 \end{array}\right]$	$\left[\begin{array}{l} i_2 \\ i_3 \end{array}\right]=\left[\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right]\left[\begin{array}{l} o_1 \\ o_4 \end{array}\right]$
gilmore-lie-groups_ E00189 (4.1)	71	■ 1.000	$(a,b,c)\longrightarrow M(a,b,c)=\left[\begin{array}{cc} 1+a & b \\ c & (1+bc)/(1+a) \end{array}\right]$	$(a,b,c)\longrightarrow M(a,b,c)=\left[\begin{array}{cc} 1+a & b \\ c & (1+bc)/(1+a) \end{array}\right]$	$(a,b,c)\longrightarrow M(a,b,c)=\left[\begin{array}{cc} 1+a & b \\ c & (1+bc)/(1+a) \end{array}\right]$
gilmore-lie-groups_ E00190 (4.2)	71	■ 1.000	$\begin{gathered} (a,b,c)\rightarrow(\delta a,\delta b,\delta c)\rightarrow M(\delta a,\delta b,\delta c) \\ =\left[\begin{array}{cc} 1+\delta a & \delta b \\ \delta c & (1+\delta b\delta c)/(1+\delta a) \end{array}\right] \end{gathered}$	$(a,b,c)\rightarrow(\delta a,\delta b,\delta c)\rightarrow M(\delta a,\delta b,\delta c) \\ =\left[\begin{array}{cc} 1+\delta a & \delta b \\ \delta c & (1+\delta b\delta c)/(1+\delta a) \end{array}\right]$	$(a,b,c)\rightarrow(\delta a,\delta b,\delta c)\rightarrow M(\delta a,\delta b,\delta c) \\ =\left[\begin{array}{cc} 1+\delta a & \delta b \\ \delta c & (1+\delta b\delta c)/(1+\delta a) \end{array}\right]$
gilmore-lie-groups_ E00192 (4.5)	72	■ 1.000	$\lim_{k\rightarrow\infty}\left(I+\frac{1}{k}X\right)^k=\sum_{n=0}^{\infty}\frac{X^n}{n!}=\operatorname{EXP}(X)$	$\lim_{k\rightarrow\infty}\left(I+\frac{1}{k}X\right)^k=\sum_{n=0}^{\infty}\frac{X^n}{n!}=\operatorname{EXP}(X)$	$\lim_{k\rightarrow\infty}\left(I+\frac{1}{k}X\right)^k=\sum_{n=0}^{\infty}\frac{X^n}{n!}=\operatorname{EXP}(X)$
gilmore-lie-groups_ E00193 (4.6)	72	■ 1.000	$X=aX_{\{a\}}+bX_{\{b\}}+cX_{\{c\}}=\left[\begin{array}{cc} a & b \\ c & -a \end{array}\right]$	$X=aX_a+bX_b+cX_c=\left[\begin{array}{cc} a & b \\ c & -a \end{array}\right]$	$X=aX_a+bX_b+cX_c=\left[\begin{array}{cc} a & b \\ c & -a \end{array}\right]$
gilmore-lie-groups_ E00195 (4.8)	72	■ 1.000	$F(X)=f_{\{0\}}(a,b,c)X^{\{0\}}+f_{\{1\}}(a,b,c)X^{\{1\}}$	$F(X)=f_0(a,b,c)X^0+f_1(a,b,c)X^1$	$F(X)=f_0(a,b,c)X^0+f_1(a,b,c)X^1$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0196 (4.9)	73	■ 1.000	$\operatorname{EXP}(X)=f_{\{0\}}\left(\theta^2\right) I_{\{2\}}+f_{\{1\}}\left(\theta^2\right) X$	$\operatorname{EXP}(X)=f_0\left(\theta^2\right) I_2+f_1\left(\theta^2\right) X$	$\operatorname{EXP}(X)=f_0(\theta^2)I_2+f_1(\theta^2)X$
gilmore-lie-groups_ EQ0198 (4.11)	74	■ 1.000	$(I+\epsilon\alpha X)(I+\epsilon\beta X)=I+\epsilon(\alpha X+\beta Y)+\text{ higher order terms }$	$(I+\epsilon\alpha X)(I+\epsilon\beta X)=I+\epsilon(\alpha X+\beta Y)+\text{ higher order terms }$	$(I+\epsilon\alpha X)(I+\epsilon\beta X)=I+\epsilon(\alpha X+\beta Y)+\text{ higher order terms }$
gilmore-lie-groups_ EQ0199 (4.12)	74	■ 1.000	$\text{ commutator of } g_{\{1\}} \text{ and } g_{\{2\}} \text{ is } g_{\{1\}} g_{\{2\}} g_{\{1\}}^{-1} g_{\{2\}}^{-1}$	commutator of g_1 and g_2 is $g_1g_2g_1^{-1}g_2^{-1}$	commutator of g_1 and g_2 is $g_1g_2g_1^{-1}g_2^{-1}$
gilmore-lie-groups_ EQ0201 (4.14)	74	■ 1.000	$g_{\{1\}} g_{\{2\}} g_{\{1\}}^{-1} g_{\{2\}}^{-1} \in G$ $\Leftrightarrow [X, Y] \in \mathfrak{g}$	$g_1g_2g_1^{-1}g_2^{-1} \in G \Leftrightarrow [X, Y] \in \mathfrak{g}$	$g_1g_2g_1^{-1}g_2^{-1} \in G \Leftrightarrow [X, Y] \in \mathfrak{g}$
gilmore-lie-groups_ EQ0202 (4.15)	74	■ 1.000	$g_{\{1\}} g_{\{2\}} g_{\{1\}}^{-1} g_{\{2\}}^{-1}=I \Rightarrow [X, Y]=0$	$g_1g_2g_1^{-1}g_2^{-1}=I \Rightarrow [X, Y]=0$	$g_1g_2g_1^{-1}g_2^{-1}=I \Rightarrow [X, Y]=0$
gilmore-lie-groups_ EQ0203 (4.16)	74	■ 1.000	$g_{\{1\}} H g_{\{1\}}^{-1} \in H \Rightarrow [X, Y] \in \text{ Lie algebra of } H$	$g_1Hg_1^{-1} \in H \Rightarrow [X, Y] \in \text{ Lie algebra of } H$	$g_1Hg_1^{-1} \in H \Rightarrow [X, Y] \in \text{ Lie algebra of } H$
gilmore-lie-groups_ EQ0204 (4.17)	74	■ 1.000	$[X,[Y,Z]]+[Y,[Z,X]]+[Z,[X,Y]]=0$	$[X,[Y,Z]]+[Y,[Z,X]]+[Z,[X,Y]]=0$	$[X,[Y,Z]]+[Y,[Z,X]]+[Z,[X,Y]]=0$
gilmore-lie-groups_ EQ0205	75	■ 1.000	$X \in \mathfrak{g}, \quad Y \in \mathfrak{g}, \quad \alpha X+\beta Y \in \mathfrak{g}$	$X \in \mathfrak{g}, \quad Y \in \mathfrak{g}, \quad \alpha X+\beta Y \in \mathfrak{g}$	$X \in \mathfrak{g}, \quad Y \in \mathfrak{g}, \quad \alpha X+\beta Y \in \mathfrak{g}$
gilmore-lie-groups_ EQ0206	75	■ 1.000	$X \in \mathfrak{g}, \quad Y \in \mathfrak{g}, \quad [X, Y] \in \mathfrak{g}$	$X \in \mathfrak{g}, \quad Y \in \mathfrak{g}, \quad [X, Y] \in \mathfrak{g}$	$X \in \mathfrak{g}, \quad Y \in \mathfrak{g}, \quad [X, Y] \in \mathfrak{g}$
gilmore-lie-groups_ EQ0207	75	■ 1.000	$[X,[Y,Z]]+[Y,[Z,X]]+[Z,[X,Y]]=0$	$[X,[Y,Z]]+[Y,[Z,X]]+[Z,[X,Y]]=0$	$[X,[Y,Z]]+[Y,[Z,X]]+[Z,[X,Y]]=0$
gilmore-lie-groups_ EQ0209 (4.19)	75	■ 1.000	$\left[X_{\{i\}}, X_{\{j\}}\right]=C_{\{i j\}}^{\{k\}} X_{\{k\}}$	$[X_i, X_j]=C_{ij}^k X_k$	$[X_i, X_j]=C_{ij}^k X_k$
gilmore-lie-groups_ EQ0219 (4.29)	77	■ 1.000	$R\left(X_{\{i\}}\right)_{\{j\}}^{\{k\}}=C_{\{i j\}}^{\{k\}}$	$R\left(X_i\right)_j^k=C_{ij}^k$	$R\left(X_i\right)_j^k=C_{ij}^k$
gilmore-lie-groups_ EQ0220 (4.30)	77	■ 1.000	$\begin{aligned} R(X) &=R\left(a X_{\{a\}}+b X_{\{b\}}+c X_{\{c\}}\right) \\ &=a R\left(X_{\{a\}}\right)+b R\left(X_{\{b\}}\right)+c R\left(X_{\{c\}}\right) \\ &\quad \&= \begin{bmatrix} 0 & -2 b & 2 c \\ -c & 2 a & 0 \\ b & 0 & -2 a \end{bmatrix} \end{aligned}$	$R(X)=R\left(a X_a+b X_b+c X_c\right)=a R\left(X_a\right)+b R\left(X_b\right)+c R\left(X_c\right) \\ =\left[\begin{array}{ccc} 0 & -2 b & 2 c \\ -c & 2 a & 0 \\ b & 0 & -2 a \end{array}\right]$	$R(X)=R\left(a X_a+b X_b+c X_c\right)=a R\left(X_a\right)+b R\left(X_b\right)+c R\left(X_c\right) \\ =\left[\begin{array}{ccc} 0 & -2 b & 2 c \\ -c & 2 a & 0 \\ b & 0 & -2 a \end{array}\right]$
gilmore-lie-groups_ EQ0221 (4.31)	78	■ 1.000	$E(2)=\left[\begin{array}{ccc} \cos \theta & \sin \theta & t_1 \\ -\sin \theta & \cos \theta & t_2 \\ 0 & 0 & 1 \end{array}\right]$	$E(2)=\left[\begin{array}{ccc} \cos \theta & \sin \theta & t_1 \\ -\sin \theta & \cos \theta & t_2 \\ 0 & 0 & 1 \end{array}\right]$	$E(2)=\left[\begin{array}{ccc} \cos \theta & \sin \theta & t_1 \\ -\sin \theta & \cos \theta & t_2 \\ 0 & 0 & 1 \end{array}\right]$
gilmore-lie-groups_ EQ0223 (4.33)	78	■ 1.000	$R\left(\theta L_z+t_1 P_x+t_2 P_y\right)=\left[\begin{array}{ccc} 0 & -\theta & 0 \\ \theta & 0 & 0 \\ -t_2 & t_1 & 0 \end{array}\right]$	$R\left(\theta L_z+t_1 P_x+t_2 P_y\right)=\left[\begin{array}{ccc} 0 & -\theta & 0 \\ \theta & 0 & 0 \\ -t_2 & t_1 & 0 \end{array}\right]$	$R\left(\theta L_z+t_1 P_x+t_2 P_y\right)=\left[\begin{array}{ccc} 0 & -\theta & 0 \\ \theta & 0 & 0 \\ -t_2 & t_1 & 0 \end{array}\right]$
gilmore-lie-groups_ EQ0224 (4.34)	78	■ 1.000	$(A, B)=\operatorname{tr} A^{\dagger} B$	$(A, B)=\operatorname{tr} A^{\dagger} B$	$(A, B)=\operatorname{tr} A^{\dagger} B$
gilmore-lie-groups_ EQ0228 (4.38)	79	■ 1.000	$\mathfrak{g}=V_{\{+\}}+V_{\{-\}}+V_{\{0\}}$	$\mathfrak{g}=V_++V_-+V_0$	$\mathfrak{g}=V_++V_-+V_0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00229 (4.39)	80	■ 1.000	$\frac{\mathfrak{g}}{\underset{\text{inner product}}{\text{Cartan-Killing}}}\begin{cases} V_0 & \text{nilpotent invariant subalgebra} \\ V_- & \text{compact subalgebra} \\ V_+ & \text{noncompact operators} \end{cases}$	$\mathfrak{g} \xrightarrow[\text{inner product}]{\text{Cartan-Killing}} \begin{cases} V_0 & \text{nilpotent invariant subalgebra} \\ V_- & \text{compact subalgebra} \\ V_+ & \text{noncompact operators} \end{cases}$	$\mathfrak{g} \xrightarrow[\text{inner product}]{\text{Cartan-Killing}} \begin{cases} V_0 & \text{nilpotent invariant subalgebra} \\ V_- & \text{compact subalgebra} \\ V_+ & \text{noncompact operators} \end{cases}$
gilmore-lie-groups_ E00230 (4.40)	80	■ 1.000	$(X, X) = \operatorname{tr} R(X) R(X) = \operatorname{tr} \begin{bmatrix} 0 & -2b & 2c \\ -c & 2a & 0 \\ b & 0 & -2a \end{bmatrix} = 8(a^2 + bc)$	$(X, X) = \operatorname{tr} R(X) R(X) = \operatorname{tr} \begin{bmatrix} 0 & -2b & 2c \\ -c & 2a & 0 \\ b & 0 & -2a \end{bmatrix} = 8(a^2 + bc)$	$(X, X) = \operatorname{tr} R(X) R(X) = \operatorname{tr} \begin{bmatrix} 0 & -2b & 2c \\ -c & 2a & 0 \\ b & 0 & -2a \end{bmatrix} = 8(a^2 + bc)$
gilmore-lie-groups_ E00232 (4.41)	80	■ 1.000	$8(a^2 + bc) = \begin{bmatrix} a & b & c \end{bmatrix} \begin{bmatrix} 8 & 0 & 0 \\ 0 & 0 & 4 \\ 0 & 4 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix}$	$8(a^2 + bc) = \begin{bmatrix} a & b & c \end{bmatrix} \begin{bmatrix} 8 & 0 & 0 \\ 0 & 0 & 4 \\ 0 & 4 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix}$	$8(a^2 + bc) = \begin{bmatrix} a & b & c \end{bmatrix} \begin{bmatrix} 8 & 0 & 0 \\ 0 & 0 & 4 \\ 0 & 4 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix}$
gilmore-lie-groups_ E00232 (4.42)	80	■ 1.000	$\begin{bmatrix} 8 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & -8 \end{bmatrix} \begin{bmatrix} X_a \\ X_+ \\ X_- \end{bmatrix}$	$\begin{bmatrix} 8 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & -8 \end{bmatrix} \begin{bmatrix} X_a \\ X_+ \\ X_- \end{bmatrix}$	$\begin{bmatrix} 8 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & -8 \end{bmatrix} \begin{bmatrix} X_a \\ X_+ \\ X_- \end{bmatrix}$
gilmore-lie-groups_ E00233 (4.43)	80	■ 1.000	$(X, X) = \operatorname{tr} \begin{bmatrix} a & b \\ c & -a \end{bmatrix} = 2(a^2 + bc)$	$(X, X) = \operatorname{tr} \begin{bmatrix} a & b \\ c & -a \end{bmatrix} = 2(a^2 + bc)$	$(X, X) = \operatorname{tr} \begin{bmatrix} a & b \\ c & -a \end{bmatrix} = 2(a^2 + bc)$
gilmore-lie-groups_ E00238 (4.48)	81	■ 1.000	$\begin{aligned} dV(\operatorname{Id}) &= d\alpha^1 \wedge d\alpha^2 \wedge \cdots \wedge d\alpha^n \\ dV(x) &= dx^1 \wedge dx^2 \wedge \cdots \wedge dx^n = \ M\ d\alpha^1 \wedge d\alpha^2 \wedge \cdots \wedge d\alpha^n \end{aligned}$	$\begin{aligned} dV(\operatorname{Id}) &= d\alpha^1 \wedge d\alpha^2 \wedge \cdots \wedge d\alpha^n \\ dV(x) &= dx^1 \wedge dx^2 \wedge \cdots \wedge dx^n = \ M\ d\alpha^1 \wedge d\alpha^2 \wedge \cdots \wedge d\alpha^n \end{aligned}$	$\begin{aligned} dV(\operatorname{Id}) &= d\alpha^1 \wedge d\alpha^2 \wedge \cdots \wedge d\alpha^n \\ dV(x) &= dx^1 \wedge dx^2 \wedge \cdots \wedge dx^n = \ M\ d\alpha^1 \wedge d\alpha^2 \wedge \cdots \wedge d\alpha^n \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00240 (4.50)	82	■ 1.000	$\begin{aligned} & \left[\begin{array}{cc} 1+x & y \\ z & \frac{1+yz}{1+x} \end{array} \right] \times \left[\begin{array}{cc} 1+d\alpha^1 & d\alpha^2 \\ d\alpha^3 & 1-d\alpha^1 \end{array} \right] \\ & = \left[\begin{array}{cc} 1+(x+dx) & y+dy \\ z+dz & \frac{1+(y+dy)(z+dz)}{1+(x+dx)} \end{array} \right] \\ & = \left[\begin{array}{cc} (1+x)(1+d\alpha^1)+yd\alpha^3 & (1+x)d\alpha^2+y(1-d\alpha^1) \\ z(1+d\alpha^1)+\frac{(1+yz)d\alpha^3}{(1+x)} & zd\alpha^2+\frac{(1+yz)(1-d\alpha^1)}{(1+x)} \end{array} \right] \end{aligned}$	$\left[\begin{array}{cc} 1+x & y \\ z & \frac{1+yz}{1+x} \end{array} \right] \times \left[\begin{array}{cc} 1+d\alpha^1 & d\alpha^2 \\ d\alpha^3 & 1-d\alpha^1 \end{array} \right] \\ = \left[\begin{array}{cc} 1+(x+dx) & y+dy \\ z+dz & \frac{1+(y+dy)(z+dz)}{1+(x+dx)} \end{array} \right] \\ = \left[\begin{array}{cc} (1+x)(1+d\alpha^1)+yd\alpha^3 & (1+x)d\alpha^2+y(1-d\alpha^1) \\ z(1+d\alpha^1)+\frac{(1+yz)d\alpha^3}{(1+x)} & zd\alpha^2+\frac{(1+yz)(1-d\alpha^1)}{(1+x)} \end{array} \right]$	$\left[\begin{array}{cc} 1+x & y \\ z & \frac{1+yz}{1+x} \end{array} \right] \times \left[\begin{array}{cc} 1+d\alpha^1 & d\alpha^2 \\ d\alpha^3 & 1-d\alpha^1 \end{array} \right] \\ = \left[\begin{array}{cc} 1+(x+dx) & y+dy \\ z+dz & \frac{1+(y+dy)(z+dz)}{1+(x+dx)} \end{array} \right] \\ = \left[\begin{array}{cc} (1+x)(1+d\alpha^1)+yd\alpha^3 & (1+x)d\alpha^2+y(1-d\alpha^1) \\ z(1+d\alpha^1)+\frac{(1+yz)d\alpha^3}{(1+x)} & zd\alpha^2+\frac{(1+yz)(1-d\alpha^1)}{(1+x)} \end{array} \right]$
gilmore-lie-groups_ E00242 (4.52)	82	■ 1.000	$d\mu(x)=\rho(x,y,z)dx\wedge dy\wedge dz=\frac{dx\wedge dy\wedge dz}{1+x}$	$d\mu(x)=\rho(x,y,z)dx\wedge dy\wedge dz=\frac{dx\wedge dy\wedge dz}{1+x}$	$d\mu(x)=\rho(x,y,z)dx\wedge dy\wedge dz=\frac{dx\wedge dy\wedge dz}{1+x}$
gilmore-lie-groups_ E00244	83	■ 1.000	$\left(X^{\prime},X\right)=\frac{1}{2}\left[\left(X^{\prime}+X,X^{\prime}+X\right)-\left(X^{\prime},X^{\prime}\right)-\left(X,X\right)\right]$	$(X',X)=\frac{1}{2}[(X'+X,X'+X)-(X',X')-(X,X)]$	$(X',X)=\frac{1}{2}[(X'+X,X'+X)-(X',X')-(X,X)]$
gilmore-lie-groups_ E00245	83	■ 1.000	$\left(X^{\prime},X\right)=2a^{\prime}a+b^{\prime}c+c^{\prime}b$	$(X',X)=2a'a+b'c+c'b$	$(X',X)=2a'a+b'c+c'b$
gilmore-lie-groups_ E00246 (4.54)	83	■ 1.000	$X=-\sum_{n=1}^{\infty}\frac{(I-Y)^n}{n}$	$X=-\sum_{n=1}^{\infty}\frac{(I-Y)^n}{n}$	$X=-\sum_{n=1}^{\infty}\frac{(I-Y)^n}{n}$
gilmore-lie-groups_ E00247 (4.55)	84	■ 1.000	$\theta\cdot\mathbf{L}=\left[\begin{array}{ccc}0&\theta_3&-\theta_2\\-\theta_3&0&\theta_1\\\theta_2&-\theta_1&0\end{array}\right]=\left[\begin{array}{ccc}0&\theta_{12}&\theta_{13}\\\theta_{21}&0&\theta_{23}\\\theta_{31}&\theta_{32}&0\end{array}\right]=\mathbf{X}$	$\theta\cdot\mathbf{L}=\left[\begin{array}{ccc}0&\theta_3&-\theta_2\\-\theta_3&0&\theta_1\\\theta_2&-\theta_1&0\end{array}\right]=\left[\begin{array}{ccc}0&\theta_{12}&\theta_{13}\\\theta_{21}&0&\theta_{23}\\\theta_{31}&\theta_{32}&0\end{array}\right]=\mathbf{X}$	$\theta\cdot\mathbf{L}=\left[\begin{array}{ccc}0&\theta_3&-\theta_2\\-\theta_3&0&\theta_1\\\theta_2&-\theta_1&0\end{array}\right]=\left[\begin{array}{ccc}0&\theta_{12}&\theta_{13}\\\theta_{21}&0&\theta_{23}\\\theta_{31}&\theta_{32}&0\end{array}\right]=\mathbf{X}$
gilmore-lie-groups_ E00248 (4.56)	84	■ 1.000	$e^{\theta\cdot\mathbf{L}}=I_3f_0(\theta)+\mathbf{X}f_1(\theta)+\mathbf{X}^2f_2(\theta)$	$e^{\theta\cdot\mathbf{L}}=I_3f_0(\theta)+\mathbf{X}f_1(\theta)+\mathbf{X}^2f_2(\theta)$	$e^{\theta\cdot\mathbf{L}}=I_3f_0(\theta)+\mathbf{X}f_1(\theta)+\mathbf{X}^2f_2(\theta)$
gilmore-lie-groups_ E00250 (4.57)	84	■ 1.000	$[X_{ij},X_{rs}]=X_{is}\delta_{jr}+X_{jr}\delta_{is}-X_{ir}\delta_{js}-X_{js}\delta_{ir}$	$[X_{ij},X_{rs}]=X_{is}\delta_{jr}+X_{jr}\delta_{is}-X_{ir}\delta_{js}-X_{js}\delta_{ir}$	$[X_{ij},X_{rs}]=X_{is}\delta_{jr}+X_{jr}\delta_{is}-X_{ir}\delta_{js}-X_{js}\delta_{ir}$
gilmore-lie-groups_ E00251 (4.58)	84	■ 1.000	$[D,[Y,Z]]=[[D,Y],Z]+[Y,[D,Z]]$	$[D,[Y,Z]]=[[D,Y],Z]+[Y,[D,Z]]$	$[D,[Y,Z]]=[[D,Y],Z]+[Y,[D,Z]]$
gilmore-lie-groups_ E00252	84	■ 1.000	$d(f\wedge g)=(df)\wedge g+f\wedge(dg)$	$d(f\wedge g)=(df)\wedge g+f\wedge(dg)$	$d(f\wedge g)=(df)\wedge g+f\wedge(dg)$
gilmore-lie-groups_ E00255	85	■ 1.000	$\mathfrak{g}\rightarrow V_0+V_-+V_+$	$\mathfrak{g}\rightarrow V_0+V_-+V_+$	$\mathfrak{g}\rightarrow V_0+V_-+V_+$
gilmore-lie-groups_ E00256	86	■ 1.000	$[X_{ij},X_{rs}]=+X_{is}g_{jr}+X_{jr}g_{is}-X_{ir}g_{js}+X_{js}g_{ir}$	$[X_{ij},X_{rs}]=+X_{is}g_{jr}+X_{jr}g_{is}-X_{ir}g_{js}+X_{js}g_{ir}$	$[X_{ij},X_{rs}]=+X_{is}g_{jr}+X_{jr}g_{is}-X_{ir}g_{js}+X_{js}g_{ir}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0259 (4.61c)	86	■ 1.000	$\left[\begin{array}{cc} i & 0 \\ 0 & -i \end{array}\right] \quad \left[\begin{array}{cc} 0 & i \\ -i & 0 \end{array}\right] \quad \left[\begin{array}{cc} 0 & -1 \\ -1 & 0 \end{array}\right]$		
gilmore-lie-groups_ EQ0260 (4.62a)	86	■ 1.000	$\left[\begin{array}{ll} \alpha_1 + \alpha_2 & \beta_1 + \beta_2 \\ \beta_1 - \beta_2 & \alpha_1 - \alpha_2 \end{array}\right]$		
gilmore-lie-groups_ EQ0262 (4.62c)	87	■ 1.000	$\left[\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right] \quad \left[\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right] \quad \left[\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right]$		
gilmore-lie-groups_ EQ0263 (5.1)	88	■ 1.000	$\begin{aligned} &(I + \epsilon A)^\dagger G(I + \epsilon A) = G \\ &G + \epsilon(A^\dagger G + GA) + \mathcal{O}(\epsilon^2) = G \\ &A^\dagger G + GA = 0 \end{aligned}$		
gilmore-lie-groups_ EQ0264 (5.2)	88	■ 1.000	$\begin{aligned} &\operatorname{det}(I + \epsilon A) = 1 + \epsilon \operatorname{tr}(A) + \mathcal{O}(\epsilon^2) = 1 \\ &\operatorname{tr}(A) = 0 \end{aligned}$		
gilmore-lie-groups_ EQ0267 (5.5)	90	■ 1.000	$M(a, b) = \begin{bmatrix} a & b \\ 0 & 1 \end{bmatrix}$		
gilmore-lie-groups_ EQ0268 (5.6)	90	■ 1.000	$\left.\frac{\partial M}{\partial a}\right _{(1,0)} = X_a = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \quad \left.\frac{\partial M}{\partial b}\right _{(1,0)} = X_b = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$		
gilmore-lie-groups_ EQ0269 (5.7)	90	■ 1.000	$\begin{bmatrix} 0 & l & r & -2\delta \\ 0 & \eta & 2R & -r \\ 0 & -2L & -\eta & l \\ 0 & 0 & 0 & 0 \end{bmatrix} = \eta X_\eta + RX_R + LX_L + rX_r + lX_l + \delta X_\delta$		
gilmore-lie-groups_ EQ0271 (5.10)	91	■ 1.000	$\begin{aligned} &\left[\begin{array}{l} \left[X_\eta, X_r\right] = +X_r \\ \left[X_\eta, X_l\right] = -X_l \\ \left[X_l, X_r\right] = X_\delta \\ \left[X_\eta, X_\delta\right] = 0 \end{array}\right] \quad \left[\begin{array}{l} \left[a^\dagger a, a^\dagger\right] = +a^\dagger \\ \left[a^\dagger a, a\right] = -a \\ \left[a, a^\dagger\right] = I \\ \left[a^\dagger a, I\right] = 0 \end{array}\right] \\ &\left[\begin{array}{l} \left[X_\eta, X_r\right] = +X_r \\ \left[X_\eta, X_l\right] = -X_l \\ \left[X_l, X_r\right] = X_\delta \\ \left[X_\eta, X_\delta\right] = 0 \end{array}\right] \quad \left[\begin{array}{l} \left[a^\dagger a, a^\dagger\right] = +a^\dagger \\ \left[a^\dagger a, a\right] = -a \\ \left[a, a^\dagger\right] = I \\ \left[a^\dagger a, I\right] = 0 \end{array}\right] \end{aligned}$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0273 (5.12)	91	■ 1.000	$\begin{aligned} & \left\{ \left[X_l, X_r \right] \right\} = X_{\delta} \\ & \left\{ \left[a, a^{\dagger} \right] \right\} = I \\ & \left\{ \left[X_l, X_{\delta} \right] \right\} = 0 \\ & \left\{ \left[X_r, X_{\delta} \right] \right\} = 0 \end{aligned}$	$\begin{aligned} & [X_l, X_r] = X_{\delta} \quad [a, a^{\dagger}] = I \\ & [X_l, X_{\delta}] = 0 \quad [a, I] = 0 \\ & [X_r, X_{\delta}] = 0 \quad [a^{\dagger}, I] = 0 \end{aligned}$	$\begin{aligned} & [X_l, X_r] = X_{\delta} \quad [a, a^{\dagger}] = I \\ & [X_l, X_{\delta}] = 0 \quad [a, I] = 0 \\ & [X_r, X_{\delta}] = 0 \quad [a^{\dagger}, I] = 0 \end{aligned}$
gilmore-lie-groups_ EQ0275 (5.14)	92	■ 1.000	$\begin{aligned} & \frac{o(3)}{2} = \frac{1}{2} \sum_i \theta_i L_i \\ & \frac{u(2)}{2} = \frac{1}{2} \sum_{\mu} x_{\mu} \sigma_{\mu} \end{aligned}$	$\begin{aligned} & o(3) = \begin{bmatrix} 0 & \theta_3 & -\theta_2 \\ -\theta_3 & 0 & \theta_1 \\ \theta_2 & -\theta_1 & 0 \end{bmatrix} = \sum_i \theta_i L_i \\ & u(2) = \frac{1}{2} \begin{bmatrix} ix_0 + ix_3 & ix_1 + x_2 \\ ix_1 - x_2 & ix_0 - ix_3 \end{bmatrix} = \frac{i}{2} \sum_{\mu} x_{\mu} \sigma_{\mu} \end{aligned}$	$\begin{aligned} & o(3) = \begin{bmatrix} 0 & \theta_3 & -\theta_2 \\ -\theta_3 & 0 & \theta_1 \\ \theta_2 & -\theta_1 & 0 \end{bmatrix} = \sum_i \theta_i L_i \\ & u(2) = \frac{1}{2} \begin{bmatrix} ix_0 + ix_3 & ix_1 + x_2 \\ ix_1 - x_2 & ix_0 - ix_3 \end{bmatrix} = \frac{i}{2} \sum_{\mu} x_{\mu} \sigma_{\mu} \end{aligned}$
gilmore-lie-groups_ EQ0278 (5.17)	92	■ 1.000	$\begin{aligned} & A = -A^t \\ & D = -D = 0 \\ & B^t = C \end{aligned}$	$\begin{aligned} & A = -A^t \\ & D = -D = 0 \\ & B^t = C \end{aligned}$	$\begin{aligned} & A = -A^t \\ & D = -D = 0 \\ & B^t = C \end{aligned}$
gilmore-lie-groups_ EQ0280 (5.19)	93	■ 1.000	$\begin{aligned} & \left[\begin{array}{cccc} 0 & \theta_3 & -\theta_2 & v_1 \\ -\theta_3 & 0 & \theta_1 & v_2 \\ \theta_2 & -\theta_1 & 0 & v_3 \\ v_1 & v_2 & v_3 & 0 \end{array} \right] \end{aligned}$	$\begin{bmatrix} 0 & \theta_3 & -\theta_2 & v_1 \\ -\theta_3 & 0 & \theta_1 & v_2 \\ \theta_2 & -\theta_1 & 0 & v_3 \\ v_1 & v_2 & v_3 & 0 \end{bmatrix}$	$\begin{bmatrix} 0 & \theta_3 & -\theta_2 & v_1 \\ -\theta_3 & 0 & \theta_1 & v_2 \\ \theta_2 & -\theta_1 & 0 & v_3 \\ v_1 & v_2 & v_3 & 0 \end{bmatrix}$
gilmore-lie-groups_ EQ0282 (5.21)	93	■ 1.000	$\begin{aligned} & \text{tr } M = -\text{tr } G^{-1} M^{\dagger} G = -\text{tr } M^{\dagger} = -\text{tr } M^* = 0 \end{aligned}$	$\text{tr } M = -\text{tr } G^{-1} M^{\dagger} G = -\text{tr } M^{\dagger} = -\text{tr } M^* = 0$	$\text{tr } M = -\text{tr } G^{-1} M^{\dagger} G = -\text{tr } M^{\dagger} = -\text{tr } M^* = 0$
gilmore-lie-groups_ EQ0284 (5.23)	93	■ 1.000	$\begin{aligned} & \left[\begin{array}{cc} A & B \\ C & D \end{array} \right] \left[\begin{array}{cc} g & 0 \\ 0 & 0 \end{array} \right] + \left[\begin{array}{cc} g & 0 \\ 0 & 0 \end{array} \right] \left[\begin{array}{cc} A^{\dagger} & C^{\dagger} \\ B^{\dagger} & D^{\dagger} \end{array} \right] = \left[\begin{array}{cc} 0 & 0 \\ 0 & 0 \end{array} \right] \\ & \Rightarrow Ag + gA^{\dagger} = 0 \quad C = C^{\dagger} = 0 \quad B, D \text{ arbitrary} \end{aligned}$	$\begin{aligned} & \left[\begin{array}{cc} A & B \\ C & D \end{array} \right] \left[\begin{array}{cc} g & 0 \\ 0 & 0 \end{array} \right] + \left[\begin{array}{cc} g & 0 \\ 0 & 0 \end{array} \right] \left[\begin{array}{cc} A^{\dagger} & C^{\dagger} \\ B^{\dagger} & D^{\dagger} \end{array} \right] = \left[\begin{array}{cc} 0 & 0 \\ 0 & 0 \end{array} \right] \\ & \Rightarrow Ag + gA^{\dagger} = 0 \quad C = C^{\dagger} = 0 \quad B, D \text{ arbitrary} \end{aligned}$	$\begin{aligned} & \left[\begin{array}{cc} A & B \\ C & D \end{array} \right] \left[\begin{array}{cc} g & 0 \\ 0 & 0 \end{array} \right] + \left[\begin{array}{cc} g & 0 \\ 0 & 0 \end{array} \right] \left[\begin{array}{cc} A^{\dagger} & C^{\dagger} \\ B^{\dagger} & D^{\dagger} \end{array} \right] = \left[\begin{array}{cc} 0 & 0 \\ 0 & 0 \end{array} \right] \\ & \Rightarrow Ag + gA^{\dagger} = 0 \quad C = C^{\dagger} = 0 \quad B, D \text{ arbitrary} \end{aligned}$
gilmore-lie-groups_ EQ0285 (5.24)	93	■ 1.000	$\begin{aligned} & \left[\begin{array}{cccc} 0 & \theta_3 & -\theta_2 & t_1 \\ -\theta_3 & 0 & \theta_1 & t_2 \\ \theta_2 & -\theta_1 & 0 & t_3 \\ 0 & 0 & 0 & s_4 \end{array} \right] \end{aligned}$	$\begin{bmatrix} 0 & \theta_3 & -\theta_2 & t_1 \\ -\theta_3 & 0 & \theta_1 & t_2 \\ \theta_2 & -\theta_1 & 0 & t_3 \\ 0 & 0 & 0 & s_4 \end{bmatrix}$	$\begin{bmatrix} 0 & \theta_3 & -\theta_2 & t_1 \\ -\theta_3 & 0 & \theta_1 & t_2 \\ \theta_2 & -\theta_1 & 0 & t_3 \\ 0 & 0 & 0 & s_4 \end{bmatrix}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00286 (5.25)	94	1.000	$\left[\begin{array}{ccccc} 0 & \theta_3 & -\theta_2 & v_1 & t_1 \\ -\theta_3 & 0 & \theta_1 & v_2 & t_2 \\ \theta_2 & -\theta_1 & 0 & v_3 & t_3 \\ v_1 & v_2 & v_3 & 0 & t_4 \\ 0 & 0 & 0 & 0 & 0 \end{array}\right]$	$\left[\begin{array}{ccccc} 0 & \theta_3 & -\theta_2 & v_1 & t_1 \\ -\theta_3 & 0 & \theta_1 & v_2 & t_2 \\ \theta_2 & -\theta_1 & 0 & v_3 & t_3 \\ v_1 & v_2 & v_3 & 0 & t_4 \\ 0 & 0 & 0 & 0 & 0 \end{array}\right]$	
gilmore-lie-groups_ E00287 (5.26)	94	1.000	$\left[\begin{array}{ccccc} 0 & \theta_3 & -\theta_2 & v_1 & t_1 \\ -\theta_3 & 0 & \theta_1 & v_2 & t_2 \\ \theta_2 & -\theta_1 & 0 & v_3 & t_3 \\ 0 & 0 & 0 & 0 & t_4 \\ 0 & 0 & 0 & 0 & 0 \end{array}\right]$	$\left[\begin{array}{ccccc} 0 & \theta_3 & -\theta_2 & v_1 & t_1 \\ -\theta_3 & 0 & \theta_1 & v_2 & t_2 \\ \theta_2 & -\theta_1 & 0 & v_3 & t_3 \\ 0 & 0 & 0 & 0 & t_4 \\ 0 & 0 & 0 & 0 & 0 \end{array}\right]$	
gilmore-lie-groups_ E00288 (5.27)	94	1.000	$\operatorname{tr} M=0 \quad \mathbb{F}=\begin{cases} \mathbb{R} & \mathfrak{sl}(n, \mathbb{R}) \\ \mathbb{C} & \mathfrak{sl}(n, \mathbb{C}) \\ \mathbb{Q} & \mathfrak{sl}(n, \mathbb{Q}) \end{cases}$	$\operatorname{tr} M=0 \quad \mathbb{F}=\begin{cases} \mathbb{R} & \mathfrak{sl}(n, \mathbb{R}) \\ \mathbb{C} & \mathfrak{sl}(n, \mathbb{C}) \\ \mathbb{Q} & \mathfrak{sl}(n, \mathbb{Q}) \end{cases}$	$\operatorname{tr} M=0 \quad \mathbb{F}=\begin{cases} \mathbb{R} & \mathfrak{sl}(n, \mathbb{R}) \\ \mathbb{C} & \mathfrak{sl}(n, \mathbb{C}) \\ \mathbb{Q} & \mathfrak{sl}(n, \mathbb{Q}) \end{cases}$
gilmore-lie-groups_ E00289 (5.28)	94	1.000	$\operatorname{det}(e^M)=e^{\operatorname{tr} M}$	$\det(e^M)=e^{\operatorname{tr} M}$	$\det(e^M)=e^{\operatorname{tr} M}$
gilmore-lie-groups_ E00291 (5.30)	95	1.000	$M \in \mathfrak{u}(n) \Rightarrow (M^\dagger)_{ij} = -M_{ji}^*$	$M \in \mathfrak{u}(n) \Rightarrow (M^\dagger)_{ij} = -M_{ji}^*$	$M \in \mathfrak{u}(n) \Rightarrow (M^\dagger)_{ij} = -M_{ji}^*$
gilmore-lie-groups_ E00292 (5.31)	95	1.000	$M \in \mathfrak{sp}(n) \Rightarrow (M^\dagger)_{ij} = -M_{ji}^*$	$M \in \mathfrak{sp}(n) \Rightarrow (M^\dagger)_{ij} = -M_{ji}^*$	$M \in \mathfrak{sp}(n) \Rightarrow (M^\dagger)_{ij} = -M_{ji}^*$
gilmore-lie-groups_ E00293 (5.32)	95	1.000	$(X, Y)=\operatorname{tr} X^\dagger Y$	$(X, Y)=\operatorname{tr} X^\dagger Y$	$(X, Y)=\operatorname{tr} X^\dagger Y$
gilmore-lie-groups_ E00294 (5.33)	96	1.000	$\left(X^{\prime}, X\right)=\operatorname{tr} X^{\prime \dagger} X=2\left(+\theta^{\prime} \theta+v_1^{\prime} v_1+v_2^{\prime} v_2\right)$	$\left(X^{\prime}, X\right)=\operatorname{tr} X^{\prime \dagger} X=2\left(+\theta^{\prime} \theta+v_1^{\prime} v_1+v_2^{\prime} v_2\right)$	$\left(X^{\prime}, X\right)=\operatorname{tr} X^{\prime \dagger} X=2\left(+\theta^{\prime} \theta+v_1^{\prime} v_1+v_2^{\prime} v_2\right)$
gilmore-lie-groups_ E00295 (5.34)	96	1.000	$(X, Y)=\operatorname{tr} X Y$	$(X, Y)=\operatorname{tr} X Y$	$(X, Y)=\operatorname{tr} X Y$
gilmore-lie-groups_ E00296 (5.35)	96	1.000	$\left(X^{\prime}, X\right)=\operatorname{tr} X^{\prime} X=2\left(-\theta^{\prime} \theta+v_1^{\prime} v_1+v_2^{\prime} v_2\right)$	$\left(X^{\prime}, X\right)=\operatorname{tr} X^{\prime} X=2\left(-\theta^{\prime} \theta+v_1^{\prime} v_1+v_2^{\prime} v_2\right)$	$\left(X^{\prime}, X\right)=\operatorname{tr} X^{\prime} X=2\left(-\theta^{\prime} \theta+v_1^{\prime} v_1+v_2^{\prime} v_2\right)$
gilmore-lie-groups_ E00298	97	1.000	$A=\left[\begin{array}{ll} a & b \\ 0 & c \end{array}\right]=a X_a+b X_b+c X_c$	$A=\left[\begin{array}{ll} a & b \\ 0 & c \end{array}\right]=a X_a+b X_b+c X_c$	$A=\left[\begin{array}{ll} a & b \\ 0 & c \end{array}\right]=a X_a+b X_b+c X_c$
gilmore-lie-groups_ E00302	98	1.000	$(Z, Z)=\operatorname{tr} Z^2=\operatorname{tr} A^2+\operatorname{tr} B C+\operatorname{tr} C B+\operatorname{tr} D^2$	$(Z, Z)=\operatorname{tr} Z^2=\operatorname{tr} A^2+\operatorname{tr} B C+\operatorname{tr} C B+\operatorname{tr} D^2$	$(Z, Z)=\operatorname{tr} Z^2=\operatorname{tr} A^2+\operatorname{tr} B C+\operatorname{tr} C B+\operatorname{tr} D^2$
gilmore-lie-groups_ E00303	98	1.000	$(X, X)=\operatorname{tr}\left(A^2+D^2\right) \quad(Y, Y)=\operatorname{tr}(B C+C B)$	$(X, X)=\operatorname{tr}\left(A^2+D^2\right) \quad(Y, Y)=\operatorname{tr}(B C+C B)$	$(X, X)=\operatorname{tr}\left(A^2+D^2\right) \quad(Y, Y)=\operatorname{tr}(B C+C B)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0305	99	■ 1.000	$\begin{aligned} & \left(\frac{\mathfrak{h}}{\mathfrak{h}}, \frac{\mathfrak{h}}{\mathfrak{h}} \right) \leq 0 \\ & \left(\frac{\mathfrak{h}}{\mathfrak{h}}, \frac{\mathfrak{p}}{\mathfrak{p}} \right) = 0 \\ & \left(\frac{\mathfrak{p}}{\mathfrak{p}}, \frac{\mathfrak{p}}{\mathfrak{p}} \right) \geq 0 \end{aligned}$	$\begin{aligned} (\mathfrak{h}, \mathfrak{h}) &\leq 0 \\ (\mathfrak{h}, \mathfrak{p}) &= 0 \\ (\mathfrak{p}, \mathfrak{p}) &\geq 0 \end{aligned}$	$\begin{aligned} (\mathfrak{h}, \mathfrak{h}) &\leq 0 \\ (\mathfrak{h}, \mathfrak{p}) &= 0 \\ (\mathfrak{p}, \mathfrak{p}) &\geq 0 \end{aligned}$
gilmore-lie-groups_ EQ0306	99	■ 1.000	$\begin{aligned} & \frac{\mathfrak{s} \, \mathfrak{l}}{\mathfrak{g}} (n ; \mathbb{R}) \, \& \, = A + B \\ & \mathfrak{g} = \mathfrak{h} + \mathfrak{p} \end{aligned}$	$\mathfrak{s}l(n; \mathbb{R}) = A + B$ $\mathfrak{g} = \mathfrak{h} + \mathfrak{p}$	$\mathfrak{s}l(n; \mathbb{R}) = A + B$ $\mathfrak{g} = \mathfrak{h} + \mathfrak{p}$
gilmore-lie-groups_ EQ0312	100	■ 1.000	$\begin{aligned} A = & \left[\begin{array}{ccc c c} 0 & \theta_3 & -\theta_2 & b_1 & t_1 \\ -\theta_3 & 0 & \theta_1 & b_2 & t_2 \\ \theta_2 & -\theta_1 & 0 & b_3 & t_3 \\ \hline \mu b_1 & \mu b_2 & \mu b_3 & 0 & t_4 \\ \sigma t_1 & \sigma t_2 & \sigma t_3 & -\mu \sigma t_4 & 0 \end{array} \right] \\ & \frac{1}{2} (A, A) = -(\theta \cdot \theta) + \mu (\mathbf{b} \cdot \mathbf{b}) + \sigma (\mathbf{t} \cdot \mathbf{t}) - \mu \sigma t_4^2 \end{aligned}$	$A = \left[\begin{array}{ccc c c} 0 & \theta_3 & -\theta_2 & b_1 & t_1 \\ -\theta_3 & 0 & \theta_1 & b_2 & t_2 \\ \theta_2 & -\theta_1 & 0 & b_3 & t_3 \\ \hline \mu b_1 & \mu b_2 & \mu b_3 & 0 & t_4 \\ \sigma t_1 & \sigma t_2 & \sigma t_3 & -\mu \sigma t_4 & 0 \end{array} \right]$ $\frac{1}{2} (A, A) = -(\theta \cdot \theta) + \mu (\mathbf{b} \cdot \mathbf{b}) + \sigma (\mathbf{t} \cdot \mathbf{t}) - \mu \sigma t_4^2$	$A = \left[\begin{array}{ccc c c} 0 & \theta_3 & -\theta_2 & b_1 & t_1 \\ -\theta_3 & 0 & \theta_1 & b_2 & t_2 \\ \theta_2 & -\theta_1 & 0 & b_3 & t_3 \\ \hline \mu b_1 & \mu b_2 & \mu b_3 & 0 & t_4 \\ \sigma t_1 & \sigma t_2 & \sigma t_3 & -\mu \sigma t_4 & 0 \end{array} \right]$ $\frac{1}{2} (A, A) = -(\theta \cdot \theta) + \mu (\mathbf{b} \cdot \mathbf{b}) + \sigma (\mathbf{t} \cdot \mathbf{t}) - \mu \sigma t_4^2$
gilmore-lie-groups_ EQ0313	100	■ 1.000	$\left[\begin{array}{l} d \, x \\ d \, y \\ d \, z \end{array} \right] = \left[\begin{array}{ccc} m_{11} & 0 & -y \\ 0 & m_{11} & x \\ 0 & m_{21} & m_{22} \end{array} \right] \left[\begin{array}{l} \delta x \\ \delta y \\ \delta z \end{array} \right]$	$\left[\begin{array}{l} dx \\ dy \\ dz \end{array} \right] = \left[\begin{array}{ccc} m_{11} & 0 & -y \\ 0 & m_{11} & x \\ 0 & m_{21} & m_{22} \end{array} \right] \left[\begin{array}{l} \delta x \\ \delta y \\ \delta z \end{array} \right]$	$\left[\begin{array}{l} dx \\ dy \\ dz \end{array} \right] = \left[\begin{array}{ccc} m_{11} & 0 & -y \\ 0 & m_{11} & x \\ 0 & m_{21} & m_{22} \end{array} \right] \left[\begin{array}{l} \delta x \\ \delta y \\ \delta z \end{array} \right]$
gilmore-lie-groups_ EQ0314	101	■ 1.000	$\begin{array}{cc} \frac{\mathfrak{h}}{\mathfrak{g} + \mathfrak{g}^*} & \frac{\mathfrak{p}}{\mathfrak{g} - \mathfrak{g}^*} \\ \hline \frac{\mathfrak{h}}{\mathfrak{g} - \mathfrak{g}^t} & \frac{\mathfrak{p}}{\mathfrak{g} + \mathfrak{g}^t} \\ \frac{\mathfrak{h}}{\mathfrak{g} - \mathfrak{g}^\dagger} & \frac{\mathfrak{p}}{\mathfrak{g} + \mathfrak{g}^\dagger} \end{array}$	$\begin{array}{cc} \mathfrak{h} & \mathfrak{p} \\ \hline \mathfrak{g} + \mathfrak{g}^* & \mathfrak{g} - \mathfrak{g}^* \\ \mathfrak{g} - \mathfrak{g}^t & \mathfrak{g} + \mathfrak{g}^t \\ \mathfrak{g} - \mathfrak{g}^\dagger & \mathfrak{g} + \mathfrak{g}^\dagger \end{array}$	$\begin{array}{cc} \mathfrak{h} & \mathfrak{p} \\ \hline \mathfrak{g} + \mathfrak{g}^* & \mathfrak{g} - \mathfrak{g}^* \\ \mathfrak{g} - \mathfrak{g}^t & \mathfrak{g} + \mathfrak{g}^t \\ \mathfrak{g} - \mathfrak{g}^\dagger & \mathfrak{g} + \mathfrak{g}^\dagger \end{array}$
gilmore-lie-groups_ EQ0315	101	■ 1.000	$\begin{aligned} & \left[\frac{\mathfrak{h}}{\mathfrak{h}}, \frac{\mathfrak{h}}{\mathfrak{h}} \right] \subseteq \frac{\mathfrak{h}}{\mathfrak{h}} \quad \left[\frac{\mathfrak{h}}{\mathfrak{h}}, \frac{\mathfrak{p}}{\mathfrak{p}} \right] \subseteq \frac{\mathfrak{h}}{\mathfrak{h}} \quad \left[\frac{\mathfrak{p}}{\mathfrak{p}}, \frac{\mathfrak{p}}{\mathfrak{p}} \right] \subseteq \frac{\mathfrak{h}}{\mathfrak{h}} \\ & \left[\frac{\mathfrak{h}}{\mathfrak{h}}, \frac{\mathfrak{p}}{\mathfrak{p}} \right] \subseteq \frac{\mathfrak{h}}{\mathfrak{h}} \quad \left[\frac{\mathfrak{h}}{\mathfrak{h}}, \frac{\mathfrak{p}}{\mathfrak{p}} \right] \subseteq \frac{\mathfrak{p}}{\mathfrak{p}} \quad \left[\frac{\mathfrak{p}}{\mathfrak{p}}, \frac{\mathfrak{p}}{\mathfrak{p}} \right] \subseteq \frac{\mathfrak{h}}{\mathfrak{h}} \end{aligned}$	$[\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h} \quad [\mathfrak{h}, \mathfrak{p}] \subseteq \mathfrak{p} \quad [\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h}$	$[\mathfrak{h}, \mathfrak{h}] \subseteq \mathfrak{h} \quad [\mathfrak{h}, \mathfrak{p}] \subseteq \mathfrak{p} \quad [\mathfrak{p}, \mathfrak{p}] \subseteq \mathfrak{h}$
gilmore-lie-groups_ EQ0316 (6.1)	102	■ 1.000	$\left[b_i, b_j^\dagger \right] = I \, \delta_{ij}$	$\left[b_i, b_j^\dagger \right] = I \delta_{ij}$	$[b_i, b_j^\dagger] = I \delta_{ij}$
gilmore-lie-groups_ EQ0317 (6.2)	102	■ 1.000	$A \rightarrow \mathcal{A} = b^\dagger A b = \sum_i \sum_j b_i^\dagger A_{ij} b_j$	$A \rightarrow \mathcal{A} = b^\dagger A b = \sum_i \sum_j b_i^\dagger A_{ij} b_j$	$A \rightarrow \mathcal{A} = b^\dagger A b = \sum_i \sum_j b_i^\dagger A_{ij} b_j$
gilmore-lie-groups_ EQ0319 (6.4)	103	■ 1.000	$\begin{aligned} & [A, B] = C \quad \Leftrightarrow \quad [\mathcal{A}, \mathcal{B}] = C \quad \mathcal{A} = \sum_{ij} b_i^\dagger A_{ij} b_j \\ & \left[\sum_i \sum_j b_i^\dagger A_{ij} b_j, \sum_i \sum_j b_i^\dagger A_{ij} b_j \right] = \sum_i \sum_j b_i^\dagger A_{ij} b_j \end{aligned}$	$[A, B] = C \Leftrightarrow [\mathcal{A}, \mathcal{B}] = C \quad \mathcal{A} = \sum_{ij} b_i^\dagger A_{ij} b_j$	$[A, B] = C \Leftrightarrow [\mathcal{A}, \mathcal{B}] = C \quad \mathcal{A} = \sum_{ij} b_i^\dagger A_{ij} b_j$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_E00320 (6.5)	103	1.000	$\left[b_i^\dagger b_j, b_r^\dagger b_s\right] = b_i^\dagger b_s \delta_{jr} - b_r^\dagger b_j \delta_{si}$	$\left[b_i^\dagger b_j, b_r^\dagger b_s\right] = b_i^\dagger b_s \delta_{jr} - b_r^\dagger b_j \delta_{si}$	$[b_i^\dagger b_j, b_r^\dagger b_s] = b_i^\dagger b_s \delta_{jr} - b_r^\dagger b_j \delta_{si}$
gilmore-lie-groups_E00321 (6.6)	103	1.000	$\left[X_{ij}, X_{rs}\right] = X_{is} \delta_{jr} - X_{rj} \delta_{si}$	$[X_{ij}, X_{rs}] = X_{is} \delta_{jr} - X_{rj} \delta_{si}$	$[X_{ij}, X_{rs}] = X_{is} \delta_{jr} - X_{rj} \delta_{si}$
gilmore-lie-groups_E00322 (6.7)	103	1.000	$A \rightarrow \mathcal{A} = \sum_{ij} A_{ij} X_{ij}$	$A \rightarrow \mathcal{A} = \sum_{ij} A_{ij} X_{ij}$	$A \rightarrow \mathcal{A} = \sum_{ij} A_{ij} X_{ij}$
gilmore-lie-groups_E00323 (6.8)	103	1.000	$\left\{f_i, f_j^\dagger\right\} = f_i f_j^\dagger + f_j^\dagger f_i = I \delta_{ij}$	$\left\{f_i, f_j^\dagger\right\} = f_i f_j^\dagger + f_j^\dagger f_i = I \delta_{ij}$	$\left\{f_i, f_j^\dagger\right\} = f_i f_j^\dagger + f_j^\dagger f_i = I \delta_{ij}$
gilmore-lie-groups_E00324 (6.9)	103	1.000	$\begin{aligned} &\left[f_i^\dagger f_j, f_r^\dagger f_s\right] = f_i^\dagger f_j f_r^\dagger f_s - f_r^\dagger f_s f_i^\dagger f_j \\ &= f_i^\dagger (\delta_{jr} - f_r^\dagger f_j) f_s - f_r^\dagger (\delta_{is} - f_i^\dagger f_s) f_j \\ &= f_i^\dagger f_s \delta_{jr} - f_r^\dagger f_j \delta_{si} \end{aligned}$	$\begin{aligned} &\left[f_i^\dagger f_j, f_r^\dagger f_s\right] = f_i^\dagger f_j f_r^\dagger f_s - f_r^\dagger f_s f_i^\dagger f_j \\ &= f_i^\dagger (\delta_{jr} - f_r^\dagger f_j) f_s - f_r^\dagger (\delta_{is} - f_i^\dagger f_s) f_j \\ &= f_i^\dagger f_s \delta_{jr} - f_r^\dagger f_j \delta_{si} \end{aligned}$	$\begin{aligned} &\left[f_i^\dagger f_j, f_r^\dagger f_s\right] = f_i^\dagger f_j f_r^\dagger f_s - f_r^\dagger f_s f_i^\dagger f_j \\ &= f_i^\dagger (\delta_{jr} - f_r^\dagger f_j) f_s - f_r^\dagger (\delta_{is} - f_i^\dagger f_s) f_j \\ &= f_i^\dagger f_s \delta_{jr} - f_r^\dagger f_j \delta_{si} \end{aligned}$
gilmore-lie-groups_E00325 (6.10)	104	1.000	$[A, B] = C \rightarrow \mathcal{A} = \sum_{ij} f_i^\dagger A_{ij} f_j$	$[A, B] = C \Leftrightarrow [\mathcal{A}, \mathcal{B}] = C \quad \mathcal{A} = \sum_{ij} f_i^\dagger A_{ij} f_j$	$[A, B] = C \Leftrightarrow [\mathcal{A}, \mathcal{B}] = C \quad \mathcal{A} = \sum_{ij} f_i^\dagger A_{ij} f_j$
gilmore-lie-groups_E00326 (6.11)	104	1.000	$X_{ij} \rightarrow x_i \partial_j = x_i \frac{\partial}{\partial x_j}$	$X_{ij} \rightarrow x_i \partial_j = x_i \frac{\partial}{\partial x_j}$	$X_{ij} \rightarrow x_i \partial_j = x_i \frac{\partial}{\partial x_j}$
gilmore-lie-groups_E00327 (6.12)	104	1.000	$[A, B] = C \rightarrow \mathcal{A} = \sum_{ij} x_i A_{ij} \partial_j = \sum_{ij} A_{ij} X_{ij}$	$[A, B] = C \Leftrightarrow [\mathcal{A}, \mathcal{B}] = C \quad \mathcal{A} = \sum_{ij} x_i A_{ij} \partial_j = \sum_{ij} A_{ij} X_{ij}$	$[A, B] = C \Leftrightarrow [\mathcal{A}, \mathcal{B}] = C \quad \mathcal{A} = \sum_{ij} x_i A_{ij} \partial_j = \sum_{ij} A_{ij} X_{ij}$
gilmore-lie-groups_E00330 (6.15)	104	1.000	$\left[L_i, L_j\right] = -\epsilon_{ijk} L_k$	$[L_i, L_j] = -\epsilon_{ijk} L_k$	$[L_i, L_j] = -\epsilon_{ijk} L_k$
gilmore-lie-groups_E00331 (6.16)	105	1.000	$\left[\begin{array}{ccc} 0 & \theta & 0 \\ -\theta & 0 & 0 \\ t_1 & t_2 & 0 \end{array}\right] = \theta L_z + t_i T_i$	$\left[\begin{array}{ccc} 0 & \theta & 0 \\ -\theta & 0 & 0 \\ t_1 & t_2 & 0 \end{array}\right] = \theta L_z + t_i T_i$	$\left[\begin{array}{ccc} 0 & \theta & 0 \\ -\theta & 0 & 0 \\ t_1 & t_2 & 0 \end{array}\right] = \theta L_z + t_i T_i$
gilmore-lie-groups_E00333 (6.18)	105	1.000	$(a, b) \rightarrow \left[\begin{array}{cc} e^a & b \\ 0 & 1 \end{array}\right]$	$(a, b) \rightarrow \left[\begin{array}{cc} e^a & b \\ 0 & 1 \end{array}\right]$	$(a, b) \rightarrow \left[\begin{array}{cc} e^a & b \\ 0 & 1 \end{array}\right]$
gilmore-lie-groups_E00336 (6.21)	106	1.000	$f_{S'}[x'(p)] = f_S[x(x'(p))]$	$f_{S'}[x'(p)] = f_S[x(x'(p))]$	$f_{S'}[x'(p)] = f_S[x(x'(p))]$
gilmore-lie-groups_E00337 (6.22)	106	1.000	$\begin{aligned} &f_{S'}[x'(p)] = f_S[x'(1 - da) - db] \\ &= f_S[x'] - da x' \frac{\partial f_S}{\partial x'} - db \frac{\partial f_S}{\partial x'} \end{aligned}$	$\begin{aligned} &f_{S'}[x'(p)] = f_S[x'(1 - da) - db] \\ &= f_S[x'] - da x' \frac{\partial f_S}{\partial x'} - db \frac{\partial f_S}{\partial x'} \end{aligned}$	$\begin{aligned} &f_{S'}[x'(p)] = f_S[x'(1 - da) - db] \\ &= f_S[x'] - da x' \frac{\partial f_S}{\partial x'} - db \frac{\partial f_S}{\partial x'} \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0338 (6.23)	106	■ 1.000	$\mathcal{X}_a=-x^{\prime}\frac{\partial}{\partial x^{\prime}}\frac{\partial}{\partial x^{\prime}}\mathcal{X}_b=-\frac{\partial}{\partial x^{\prime}}\frac{\partial}{\partial x^{\prime}}\mathcal{X}_a$	$\mathcal{X}_a=-x^{\prime}\frac{\partial}{\partial x^{\prime}}\quad\mathcal{X}_b=-\frac{\partial}{\partial x^{\prime}}$	$\mathcal{X}_a=-x^{\prime}\frac{\partial}{\partial x^{\prime}}\quad\mathcal{X}_b=-\frac{\partial}{\partial x^{\prime}}$
gilmore-lie-groups_ EQ0339 (6.24)	106	■ 1.000	$\left[X_a,X_b\right]=X_b\Leftrightarrow\left[\mathcal{X}_a,\mathcal{X}_b\right]=\mathcal{X}_b$	$[X_a,X_b]=X_b\Leftrightarrow[\mathcal{X}_a,\mathcal{X}_b]=\mathcal{X}_b$	$[X_a,X_b]=X_b\Leftrightarrow[\mathcal{X}_a,\mathcal{X}_b]=\mathcal{X}_b$
gilmore-lie-groups_ EQ0340 (6.25)	106	■ 1.000	$G_{S^{\prime}}(x^{\prime},y^{\prime})=G_S(x,y)$	$G_{S^{\prime}}(x^{\prime},y^{\prime})=G_S(x,y)$	$G_{S^{\prime}}(x^{\prime},y^{\prime})=G_S(x,y)$
gilmore-lie-groups_ EQ0342 (6.27)	106	■ 1.000	$\begin{aligned} G_{S^{\prime}}(x^{\prime},y^{\prime})&=G_S\left[x=(1-da)x^{\prime},y=(1-d)y^{\prime}-db\right] \\ &=G_S(x^{\prime},y^{\prime})+\left\{da\left(-x^{\prime}\frac{\partial}{\partial x^{\prime}}-y^{\prime}\frac{\partial}{\partial y^{\prime}}\right)+db\left(-\frac{\partial}{\partial y^{\prime}}\right)\right\}G_S(x^{\prime},y^{\prime}) \end{aligned}$	$G_{S^{\prime}}(x^{\prime},y^{\prime})=G_S\left[x=(1-da)x^{\prime},y=(1-d)y^{\prime}-db\right] \\ =G_S(x^{\prime},y^{\prime})+\left\{da\left(-x^{\prime}\frac{\partial}{\partial x^{\prime}}-y^{\prime}\frac{\partial}{\partial y^{\prime}}\right)+db\left(-\frac{\partial}{\partial y^{\prime}}\right)\right\}G_S(x^{\prime},y^{\prime})$	$G_{S^{\prime}}(x^{\prime},y^{\prime})=G_S[x=(1-da)x^{\prime},y=(1-d)y^{\prime}-db] \\ =G_S(x^{\prime},y^{\prime})+\left\{da\left(-x^{\prime}\frac{\partial}{\partial x^{\prime}}-y^{\prime}\frac{\partial}{\partial y^{\prime}}\right)+db\left(-\frac{\partial}{\partial y^{\prime}}\right)\right\}G_S(x^{\prime},y^{\prime})$
gilmore-lie-groups_ EQ0344 (6.29)	106	■ 1.000	$\left[X_a,X_b\right]=X_b\Leftrightarrow\left[\mathcal{X}_x,\mathcal{X}_b\right]=\mathcal{X}_b$	$[X_a,X_b]=X_b\Leftrightarrow[\mathcal{X}_x,\mathcal{X}_b]=\mathcal{X}_b$	$[X_a,X_b]=X_b\Leftrightarrow[\mathcal{X}_x,\mathcal{X}_b]=\mathcal{X}_b$
gilmore-lie-groups_ EQ0345 (6.30)	107	■ 1.000	$f_{S^{\prime}}[x(p)]=f_S\left[x^{\prime}(p)\right]$	$f_{S^{\prime}}[x(p)]=f_S\left[x^{\prime}(p)\right]$	$f_{S^{\prime}}[x(p)]=f_S[x^{\prime}(p)]$
gilmore-lie-groups_ EQ0348	108	■ 1.000	$J^2=\left(\frac{1}{2}\hat{n}\right)\left(\frac{1}{2}\hat{n}+1\right)$	$J^2=\left(\frac{1}{2}\hat{n}\right)\left(\frac{1}{2}\hat{n}+1\right)$	$J^2=\left(\frac{1}{2}\hat{n}\right)\left(\frac{1}{2}\hat{n}+1\right)$
gilmore-lie-groups_ EQ0349	108	■ 1.000	$a_i^{\dagger} n_i\rangle=\sqrt{n_i+1} n_i+1\rangle\quad a_i n_i\rangle=\sqrt{n_i} n_i-1\rangle$	$a_i^{\dagger} n_i\rangle=\sqrt{n_i+1} n_i+1\rangle\quad a_i n_i\rangle=\sqrt{n_i} n_i-1\rangle$	$a_i^{\dagger} n_i\rangle=\sqrt{n_i+1} n_i+1\rangle\quad a_i n_i\rangle=\sqrt{n_i} n_i-1\rangle$
gilmore-lie-groups_ EQ0351 (6.31)	109	■ 1.000	$\langle n_1^{\prime},n_2^{\prime},n_3^{\prime} \hat{b}_i^{\dagger}\hat{b}_j n_1,n_2,n_3\rangle$	$\langle n_1^{\prime},n_2^{\prime},n_3^{\prime} \hat{b}_i^{\dagger}\hat{b}_j n_1,n_2,n_3\rangle$	$\langle n_1^{\prime},n_2^{\prime},n_3^{\prime} \hat{b}_i^{\dagger}\hat{b}_j n_1,n_2,n_3\rangle$
gilmore-lie-groups_ EQ0352 (6.32)	109	■ 1.000	$\langle n_1^{\prime},n_2^{\prime},n_3^{\prime} \mathcal{O} n_1,n_2,n_3\rangle=I_D\delta_{n_1^{\prime},n_1}\delta_{n_2^{\prime},n_2}\delta_{n_3^{\prime},n_3}$	$\langle n_1^{\prime},n_2^{\prime},n_3^{\prime} \mathcal{O} n_1,n_2,n_3\rangle=I_D\delta_{n_1^{\prime},n_1}\delta_{n_2^{\prime},n_2}\delta_{n_3^{\prime},n_3}$	$\langle n_1^{\prime},n_2^{\prime},n_3^{\prime} \mathcal{O} n_1,n_2,n_3\rangle=I_D\delta_{n_1^{\prime},n_1}\delta_{n_2^{\prime},n_2}\delta_{n_3^{\prime},n_3}$
gilmore-lie-groups_ EQ0355	110	■ 1.000	$A^tD^t-B^tC^t=I_n\quad A^tB^t=B^tA^t\quad C^tD^t=D^tC^t$	$AD^t-BC^t=I_n\quad AB^t=BA^t\quad CD^t=DC^t$	$AD^t-BC^t=I_n\quad AB^t=BA^t\quad CD^t=DC^t$
gilmore-lie-groups_ EQ0356	110	■ 1.000	$\mathcal{H}=\hbar\omega\sum_{i=1}^N\left(a_i^{\dagger}a_i+\frac{1}{2}\right)$	$\mathcal{H}=\hbar\omega\sum_{i=1}^N\left(a_i^{\dagger}a_i+\frac{1}{2}\right)$	$\mathcal{H}=\hbar\omega\sum_{i=1}^N\left(a_i^{\dagger}a_i+\frac{1}{2}\right)$
gilmore-lie-groups_ EQ0357	111	■ 1.000	$H_n(x)=e^{x^2}\left(-\frac{d}{dx}\right)^ne^{-x^2}$	$H_n(x)=e^{x^2}\left(-\frac{d}{dx}\right)^ne^{-x^2}$	$H_n(x)=e^{x^2}\left(-\frac{d}{dx}\right)^ne^{-x^2}$
gilmore-lie-groups_ EQ0359	111	■ 1.000	$H_n(x)e^{-x^2/2}=e^{+x^2/2}\left(-\frac{d}{dx}\right)^ne^{-x^2}=\left(x-\frac{d}{dx}\right)^ne^{-x^2/2}$	$H_n(x)e^{-x^2/2}=e^{+x^2/2}\left(-\frac{d}{dx}\right)^ne^{-x^2}=\left(x-\frac{d}{dx}\right)^ne^{-x^2/2}$	$H_n(x)e^{-x^2/2}=e^{+x^2/2}\left(-\frac{d}{dx}\right)^ne^{-x^2}=\left(x-\frac{d}{dx}\right)^ne^{-x^2/2}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08360 (6.33)	111	■ 1.000	$\langle x \mid n \rangle = \frac{\left(\sqrt{2} a^\dagger \right)^n}{\sqrt{2^n n!}} \langle x \mid 0 \rangle = \frac{H_n(x) e^{-x^2/2}}{\sqrt{2^n n!} \sqrt{\pi}} = \psi_n(x)$	$\langle x \mid n \rangle = \frac{(\sqrt{2} a^\dagger)^n}{\sqrt{2^n n!}} \langle x \mid 0 \rangle = \frac{H_n(x) e^{-x^2/2}}{\sqrt{2^n n!} \sqrt{\pi}} = \psi_n(x)$	$\langle x \mid n \rangle = \frac{(\sqrt{2} a^\dagger)^n}{\sqrt{2^n n!}} \langle x \mid 0 \rangle = \frac{H_n(x) e^{-x^2/2}}{\sqrt{2^n n!} \sqrt{\pi}} = \psi_n(x)$
gilmore-lie-groups_ E08361	111	■ 1.000	$H = \sum_{i=1}^n \hbar \omega_i \left(a_i^\dagger a_i + \frac{1}{2} \right) + i \sum_{i < j} \theta_{ij} \left(a_i^\dagger a_j - a_j^\dagger a_i \right) + \sum_{i \leq j} q_{ij} \left(a_i^\dagger a_j + a_j^\dagger a_i \right)$	$H = \sum_{i=1}^n \hbar \omega_i \left(a_i^\dagger a_i + \frac{1}{2} \right) + i \sum_{i < j} \theta_{ij} \left(a_i^\dagger a_j - a_j^\dagger a_i \right) + \sum_{i \leq j} q_{ij} \left(a_i^\dagger a_j + a_j^\dagger a_i \right)$	$H = \sum_{i=1}^n \hbar \omega_i \left(a_i^\dagger a_i + \frac{1}{2} \right) + i \sum_{i < j} \theta_{ij} \left(a_i^\dagger a_j - a_j^\dagger a_i \right) + \sum_{i \leq j} q_{ij} \left(a_i^\dagger a_j + a_j^\dagger a_i \right)$
gilmore-lie-groups_ E08362	111	■ 1.000	$\Gamma_{ij} = \hbar \omega_i \delta_{ij} + (q + i \theta)_{ij}$	$\Gamma_{ij} = \hbar \omega_i \delta_{ij} + (q + i \theta)_{ij}$	$\Gamma_{ij} = \hbar \omega_i \delta_{ij} + (q + i \theta)_{ij}$
gilmore-lie-groups_ E08364 (7.2)	114	■ 1.000	$X = \begin{bmatrix} a & b+c \\ b-c & -a \end{bmatrix} \in \mathfrak{sl}(2; \mathbb{R})$	$X = \begin{bmatrix} a & b+c \\ b-c & -a \end{bmatrix} \in \mathfrak{sl}(2; \mathbb{R})$	$X = \begin{bmatrix} a & b+c \\ b-c & -a \end{bmatrix} \in \mathfrak{sl}(2; \mathbb{R})$
gilmore-lie-groups_ E08365 (7.3)	114	■ 1.000	$\operatorname{tr} \operatorname{EXP}(X) \geq -2$	$\operatorname{tr} \operatorname{EXP}(X) \geq -2$	$\operatorname{tr} \operatorname{EXP}(X) \geq -2$
gilmore-lie-groups_ E08367 (7.5)	114	■ 1.000	$\operatorname{tr} e^X = \operatorname{tr} S e^X S^{-1} = \operatorname{tr} e^{S X S^{-1}}$	$\operatorname{tr} e^X = \operatorname{tr} S e^X S^{-1} = \operatorname{tr} e^{S X S^{-1}}$	$\operatorname{tr} e^X = \operatorname{tr} S e^X S^{-1} = \operatorname{tr} e^{S X S^{-1}}$
gilmore-lie-groups_ E08370 (7.8)	116	■ 1.000	$(0, 0, c) \longrightarrow \begin{bmatrix} \cos c & \sin c \\ -\sin c & \cos c \end{bmatrix}$	$(0, 0, c) \longrightarrow \begin{bmatrix} \cos c & \sin c \\ -\sin c & \cos c \end{bmatrix}$	$(0, 0, c) \longrightarrow \begin{bmatrix} \cos c & \sin c \\ -\sin c & \cos c \end{bmatrix}$
gilmore-lie-groups_ E08371 (7.9)	116	■ 1.000	$(a, b, 0) \longrightarrow \begin{bmatrix} \cosh r + (a/r) \sinh r & (b/r) \sinh r \\ (b/r) \sinh r & \cosh r - (a/r) \sinh r \end{bmatrix} \quad r^2 = a^2 + b^2$	$(a, b, 0) \longrightarrow \begin{bmatrix} \cosh r + (a/r) \sinh r & (b/r) \sinh r \\ (b/r) \sinh r & \cosh r - (a/r) \sinh r \end{bmatrix} \quad r^2 = a^2 + b^2$	$(a, b, 0) \longrightarrow \begin{bmatrix} \cosh r + (a/r) \sinh r & (b/r) \sinh r \\ (b/r) \sinh r & \cosh r - (a/r) \sinh r \end{bmatrix} \quad r^2 = a^2 + b^2$
gilmore-lie-groups_ E08372 (7.10)	116	■ 1.000	$(X, X) = \operatorname{tr} X^2 = 2(a^2 + b^2 - c^2)$	$(X, X) = \operatorname{tr} X^2 = 2(a^2 + b^2 - c^2)$	$(X, X) = \operatorname{tr} X^2 = 2(a^2 + b^2 - c^2)$
gilmore-lie-groups_ E08373 (7.11)	116	■ 1.000	$\begin{array}{c} \begin{bmatrix} a & b \\ b & -a \end{bmatrix} \\ \text{EXP} \downarrow \end{array} + \begin{array}{c} \begin{bmatrix} 0 & c \\ -c & 0 \end{bmatrix} \\ \downarrow \text{EXP} \end{array}$	$\begin{array}{c} \begin{bmatrix} a & b \\ b & -a \end{bmatrix} \\ \text{EXP} \downarrow \end{array} + \begin{array}{c} \begin{bmatrix} 0 & c \\ -c & 0 \end{bmatrix} \\ \downarrow \text{EXP} \end{array}$	$\begin{array}{c} \begin{bmatrix} a & b \\ b & -a \end{bmatrix} \\ \text{EXP} \downarrow \end{array} + \begin{array}{c} \begin{bmatrix} 0 & c \\ -c & 0 \end{bmatrix} \\ \downarrow \text{EXP} \end{array}$
gilmore-lie-groups_ E08374 (7.12)	116	■ 1.000	$z^2 - x^2 - y^2 = 1$	$z^2 - x^2 - y^2 = 1$	$z^2 - x^2 - y^2 = 1$
gilmore-lie-groups_ E08376 (7.14)	117	■ 1.000	$z^2 + x^2 - y^2 = 1$	$z^2 + x^2 - y^2 = 1$	$z^2 + x^2 - y^2 = 1$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E09379 (7.17)	119	■ 1.000	$\frac{s}{o}(2,1)=\left[\begin{array}{ccc} 0 & a_3 & a_2 \\ & -a_3 & 0 \\ & a_2 & a_1 \end{array}\right] \quad \frac{s}{u}(1,1)=\frac{i}{2}\left[\begin{array}{cc} b_3 & ib_1+b_2 \\ ib_1-b_2 & -b_3 \end{array}\right]$	$\mathfrak{so}(2,1)=\left[\begin{array}{ccc} 0 & a_3 & a_2 \\ -a_3 & 0 & a_1 \\ a_2 & a_1 & 0 \end{array}\right] \quad \mathfrak{su}(1,1)=\frac{i}{2}\left[\begin{array}{cc} b_3 & ib_1+b_2 \\ ib_1-b_2 & -b_3 \end{array}\right]$	$\mathfrak{so}(2,1)=\left[\begin{array}{ccc} 0 & a_3 & a_2 \\ -a_3 & 0 & a_1 \\ a_2 & a_1 & 0 \end{array}\right] \quad \mathfrak{su}(1,1)=\frac{i}{2}\left[\begin{array}{cc} b_3 & ib_1+b_2 \\ ib_1-b_2 & -b_3 \end{array}\right]$
gilmore-lie-groups_ E09383	120	■ 1.000	$\pi(\sin\theta\cos\phi,\sin\theta\sin\phi,\cos\theta)\sim-\pi(\sin\theta\cos\phi,\sin\theta\sin\phi,\cos\theta)$	$\pi(\sin\theta\cos\phi,\sin\theta\sin\phi,\cos\theta)\sim-\pi(\sin\theta\cos\phi,\sin\theta\sin\phi,\cos\theta)$	$\pi(\sin\theta\cos\phi,\sin\theta\sin\phi,\cos\theta)\sim-\pi(\sin\theta\cos\phi,\sin\theta\sin\phi,\cos\theta)$
gilmore-lie-groups_ E09387	121	■ 1.000	$G_1=\bar{G}/D_1\rightarrow\bar{G}/D_{\mathrm{MAX}}\leftarrow\bar{G}/D_2=G_2$	$G_1=\bar{G}/D_1\rightarrow\bar{G}/D_{\mathrm{MAX}}\leftarrow\bar{G}/D_2=G_2$	$G_1=\bar{G}/D_1\rightarrow\bar{G}/D_{\mathrm{MAX}}\leftarrow\bar{G}/D_2=G_2$
gilmore-lie-groups_ E09394 (7.30)	124	■ 1.000	$\left[X_l,X_r\right]=X_\delta \quad \left[X_l,X_\delta\right]=\left[X_r,X_\delta\right]=0$	$\left[X_l,X_r\right]=X_\delta \quad \left[X_l,X_\delta\right]=\left[X_r,X_\delta\right]=0$	$\left[X_l,X_r\right]=X_\delta \quad \left[X_l,X_\delta\right]=\left[X_r,X_\delta\right]=0$
gilmore-lie-groups_ E09395 (7.31)	124	■ 1.000	$\left[a,a^\dagger\right]=I \quad \left[a,I\right]=\left[a^\dagger,I\right]=0$	$\left[a,a^\dagger\right]=I \quad \left[a,I\right]=\left[a^\dagger,I\right]=0$	$\left[a,a^\dagger\right]=I \quad \left[a,I\right]=\left[a^\dagger,I\right]=0$
gilmore-lie-groups_ E09396 (7.32)	124	■ 1.000	$\begin{aligned} &X_l\rightarrow a \\ &X_r\rightarrow a^\dagger \\ &X_\delta\rightarrow I \end{aligned}$	$\begin{aligned} &X_l\rightarrow a \\ &X_r\rightarrow a^\dagger \\ &X_\delta\rightarrow I \end{aligned}$	$\begin{aligned} &X_l\rightarrow a \\ &X_r\rightarrow a^\dagger \\ &X_\delta\rightarrow I \end{aligned}$
gilmore-lie-groups_ E09401 (7.37)	125	■ 1.000	$\begin{aligned} &X_\eta\rightarrow\hat{n} \\ &X_l\rightarrow a \\ &X_r\rightarrow a^\dagger \\ &X_\delta\rightarrow I \end{aligned}$	$\begin{aligned} &X_\eta\rightarrow\hat{n} \\ &X_l\rightarrow a \\ &X_r\rightarrow a^\dagger \\ &X_\delta\rightarrow I \end{aligned}$	$\begin{aligned} &X_\eta\rightarrow\hat{n} \\ &X_l\rightarrow a \\ &X_r\rightarrow a^\dagger \\ &X_\delta\rightarrow I \end{aligned}$
gilmore-lie-groups_ E09402 (7.38)	125	■ 1.000	$\begin{aligned} &\operatorname{EXP}\left(\eta a^\dagger a+r a^\dagger+l a\right)=\operatorname{EXP}\left(r^\prime a^\dagger\right) \operatorname{EXP}\left(\eta^\prime a^\dagger a+\delta^\prime I\right) \operatorname{EXP}\left(l^\prime a\right) \\ &\parallel \\ &\left[\begin{array}{ccc} 1 & \left(e^\eta-1\right) l / \eta & \left(e^\eta-1-\eta\right) l r / \eta^2 \\ 0 & e^\eta & \left(e^\eta-1\right) r / \eta \\ 0 & 0 & 1 \end{array}\right]=\left[\begin{array}{ccc} 1 & l^\prime & \delta^\prime \\ 0 & e^{\eta^\prime} & r^\prime \\ 0 & 0 & 1 \end{array}\right] \end{aligned}$	$\begin{aligned} &\operatorname{EXP}\left(\eta a^\dagger a+r a^\dagger+l a\right)=\operatorname{EXP}\left(r^\prime a^\dagger\right) \operatorname{EXP}\left(\eta^\prime a^\dagger a+\delta^\prime I\right) \operatorname{EXP}\left(l^\prime a\right) \\ &\parallel \\ &\left[\begin{array}{ccc} 1 & \left(e^\eta-1\right) l / \eta & \left(e^\eta-1-\eta\right) l r / \eta^2 \\ 0 & e^\eta & \left(e^\eta-1\right) r / \eta \\ 0 & 0 & 1 \end{array}\right]=\left[\begin{array}{ccc} 1 & l^\prime & \delta^\prime \\ 0 & e^{\eta^\prime} & r^\prime \\ 0 & 0 & 1 \end{array}\right] \end{aligned}$	$\begin{aligned} &\operatorname{EXP}\left(\eta a^\dagger a+r a^\dagger+l a\right)=\operatorname{EXP}\left(r^\prime a^\dagger\right) \operatorname{EXP}\left(\eta^\prime a^\dagger a+\delta^\prime I\right) \operatorname{EXP}\left(l^\prime a\right) \\ &\parallel \\ &\left[\begin{array}{ccc} 1 & \left(e^\eta-1\right) l / \eta & \left(e^\eta-1-\eta\right) l r / \eta^2 \\ 0 & e^\eta & \left(e^\eta-1\right) r / \eta \\ 0 & 0 & 1 \end{array}\right]=\left[\begin{array}{ccc} 1 & l^\prime & \delta^\prime \\ 0 & e^{\eta^\prime} & r^\prime \\ 0 & 0 & 1 \end{array}\right] \end{aligned}$
gilmore-lie-groups_ E09405 (7.41)	125	■ 1.000	$\langle 0 e^{\eta a^\dagger a+r a^\dagger+l a} 0\rangle=e^{\delta^\prime}=\operatorname{EXP}\left(\frac{\left(e^\eta-1-\eta\right) l r}{\eta^2}\right)$	$\langle 0 e^{\eta a^\dagger a+r a^\dagger+l a} 0\rangle=e^{\delta^\prime}=\operatorname{EXP}\left(\frac{\left(e^\eta-1-\eta\right) l r}{\eta^2}\right)$	$\langle 0 e^{\eta a^\dagger a+r a^\dagger+l a} 0\rangle=e^{\delta^\prime}=\operatorname{EXP}\left(\frac{\left(e^\eta-1-\eta\right) l r}{\eta^2}\right)$
gilmore-lie-groups_ E09406 (7.42)	125	■ 1.000	$\left\langle\begin{array}{c} j \\ -j \end{array}\right e^{i \theta \cdot \mathbf{J}}\left \begin{array}{c} j \\ -j \end{array}\right\rangle$	$\left\langle\begin{array}{c} j \\ -j \end{array}\right e^{i \theta \cdot \mathbf{J}}\left \begin{array}{c} j \\ -j \end{array}\right\rangle$	$\left\langle\begin{array}{c} j \\ -j \end{array}\right e^{i \theta \cdot \mathbf{J}}\left \begin{array}{c} j \\ -j \end{array}\right\rangle$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E04497 (7.43)	126	■ 1.000	$\left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta \cdot \mathbf{J}} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = \left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta'_+ J_+} e^{i\theta'_z J_z} e^{i\theta'_- J_-} \left \begin{array}{c} j \\ -j \end{array} \right\rangle$	$\left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta \cdot \mathbf{J}} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = \left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta'_+ J_+} e^{i\theta'_z J_z} e^{i\theta'_- J_-} \left \begin{array}{c} j \\ -j \end{array} \right\rangle$	$\left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta \cdot \mathbf{J}} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = \left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta'_+ J_+} e^{i\theta'_z J_z} e^{i\theta'_- J_-} \left \begin{array}{c} j \\ -j \end{array} \right\rangle$
gilmore-lie-groups_ E04498 (7.44)	126	■ 1.000	$e^{i\theta \cdot \mathbf{J}} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = (I + i\theta'_- J_- + \dots) \left \begin{array}{c} j \\ -j \end{array} \right\rangle = \left \begin{array}{c} j \\ -j \end{array} \right\rangle$	$e^{i\theta'_- J_-} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = (I + i\theta'_- J_- + \dots) \left \begin{array}{c} j \\ -j \end{array} \right\rangle = \left \begin{array}{c} j \\ -j \end{array} \right\rangle$	$e^{i\theta'_- J_-} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = (I + i\theta'_- J_- + \dots) \left \begin{array}{c} j \\ -j \end{array} \right\rangle = \left \begin{array}{c} j \\ -j \end{array} \right\rangle$
gilmore-lie-groups_ E04499 (7.45)	126	■ 1.000	$\left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta \cdot \mathbf{J}} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = \left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta'_z J_z} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = e^{-ij\theta'_z}$	$\left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta \cdot \mathbf{J}} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = \left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta'_z J_z} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = e^{-ij\theta'_z}$	$\left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta \cdot \mathbf{J}} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = \left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta'_z J_z} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = e^{-ij\theta'_z}$
gilmore-lie-groups_ E04410 (7.46)	126	■ 1.000	$\begin{aligned} & e^{i\theta \cdot \mathbf{J}} \rightarrow \text{EXP} \frac{i}{2} \left[\begin{array}{cc} \theta_z & \theta_x - i\theta_y \\ \theta_x + i\theta_y & -\theta_z \end{array} \right] \\ & = \left[\begin{array}{cc} \cos(\theta/2) + i(\theta_z/\theta) \sin(\theta/2) & i[(\theta_x - i\theta_y)/\theta] \sin(\theta/2) \\ i[(\theta_x + i\theta_y)/\theta] \sin(\theta/2) & \cos(\theta/2) - i(\theta_z/\theta) \sin(\theta/2) \end{array} \right] \end{aligned}$	$\begin{aligned} & e^{i\theta \cdot \mathbf{J}} \rightarrow \text{EXP} \frac{i}{2} \left[\begin{array}{cc} \theta_z & \theta_x - i\theta_y \\ \theta_x + i\theta_y & -\theta_z \end{array} \right] \\ & = \left[\begin{array}{cc} \cos(\theta/2) + i(\theta_z/\theta) \sin(\theta/2) & i[(\theta_x - i\theta_y)/\theta] \sin(\theta/2) \\ i[(\theta_x + i\theta_y)/\theta] \sin(\theta/2) & \cos(\theta/2) - i(\theta_z/\theta) \sin(\theta/2) \end{array} \right] \end{aligned}$	$\begin{aligned} & e^{i\theta \cdot \mathbf{J}} \rightarrow \text{EXP} \frac{i}{2} \left[\begin{array}{cc} \theta_z & \theta_x - i\theta_y \\ \theta_x + i\theta_y & -\theta_z \end{array} \right] \\ & = \left[\begin{array}{cc} \cos(\theta/2) + i(\theta_z/\theta) \sin(\theta/2) & i[(\theta_x - i\theta_y)/\theta] \sin(\theta/2) \\ i[(\theta_x + i\theta_y)/\theta] \sin(\theta/2) & \cos(\theta/2) - i(\theta_z/\theta) \sin(\theta/2) \end{array} \right] \end{aligned}$
gilmore-lie-groups_ E04412 (7.48) [re-fined: census]	126	■ 1.000	$e^{-i\theta'_z J_z} = \cos(\theta/2) - i(\theta_z/\theta) \sin(\theta/2)$	$e^{-i\theta'_z J_z} = \cos(\theta/2) - i(\theta_z/\theta) \sin(\theta/2)$	$e^{-i\theta'_z J_z} = \cos(\theta/2) - i(\theta_z/\theta) \sin(\theta/2)$
gilmore-lie-groups_ E04413 (7.49) [re-fined: census]	126	■ 1.000	$\left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta \cdot \mathbf{J}} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = e^{-ij\theta'_z} = \left(e^{-i\theta'_z/2} \right)^{2j} = [\cos(\theta/2) - i(\theta_z/\theta) \sin(\theta/2)]^{2j}$	$\left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta \cdot \mathbf{J}} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = e^{-ij\theta'_z} = \left(e^{-i\theta'_z/2} \right)^{2j} = [\cos(\theta/2) - i(\theta_z/\theta) \sin(\theta/2)]^{2j}$	$\left\langle \begin{array}{c} j \\ -j \end{array} \right e^{i\theta \cdot \mathbf{J}} \left \begin{array}{c} j \\ -j \end{array} \right\rangle = e^{-ij\theta'_z} = (e^{-i\theta'_z/2})^{2j} = [\cos(\theta/2) - i(\theta_z/\theta) \sin(\theta/2)]^{2j}$
gilmore-lie-groups_ E04418 (7.54)	128	■ 1.000	$H \psi\rangle = i\hbar \frac{\partial}{\partial t} \psi\rangle$	$H \psi\rangle = i\hbar \frac{\partial}{\partial t} \psi\rangle$	$H \psi\rangle = i\hbar \frac{\partial}{\partial t} \psi\rangle$
gilmore-lie-groups_ E04419 (7.55)	128	■ 1.000	$ \psi(t+\delta t)\rangle = \left(I - \frac{i}{\hbar} H \delta t \right) \psi(t)\rangle = e^{-\frac{i}{\hbar} H \delta t} \psi(t)\rangle$	$ \psi(t+\delta t)\rangle = \left(I - \frac{i}{\hbar} H \delta t \right) \psi(t)\rangle = e^{-\frac{i}{\hbar} H \delta t} \psi(t)\rangle$	$ \psi(t+\delta t)\rangle = \left(I - \frac{i}{\hbar} H \delta t \right) \psi(t)\rangle = e^{-\frac{i}{\hbar} H \delta t} \psi(t)\rangle$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00420 (7.56)	128	■ 1.000	$\begin{aligned} U(t_f, t_i) &= U(t_f, t_f - \delta t) \cdots U(t_i + 2\delta t, t_i + \delta t) U(t_i + \delta t, t_i) \\ &= \prod U(t_i + (n+1)\delta t, t_i + n\delta t) = \int_{t_i}^{t_f} U(\tau) d\tau \\ &= T \int_{t_i}^{t_f} e^{-\frac{i}{\hbar} H(t)} dt \end{aligned}$	$U(t_f, t_i) = U(t_f, t_f - \delta t) \cdots U(t_i + 2\delta t, t_i + \delta t) U(t_i + \delta t, t_i)$ $= \prod U(t_i + (n+1)\delta t, t_i + n\delta t) = \int_{t_i}^{t_f} U(\tau) d\tau$ $= T \int_{t_i}^{t_f} e^{-\frac{i}{\hbar} H(t)} dt$	$U(t_f, t_i) = U(t_f, t_f - \delta t) \cdots U(t_i + 2\delta t, t_i + \delta t) U(t_i + \delta t, t_i)$ $= \prod U(t_i + (n+1)\delta t, t_i + n\delta t) = \int_{t_i}^{t_f} U(\tau) d\tau$ $= T \int_{t_i}^{t_f} e^{-\frac{i}{\hbar} H(t)} dt$
gilmore-lie-groups_ E00421 (7.57)	129	■ 1.000	$H = \epsilon(t) J_z + \alpha(t) J_+ + \alpha^*(t) J_-$ $\xrightarrow{j \rightarrow \frac{1}{2}} \begin{bmatrix} \frac{1}{2}\epsilon(t) & \alpha(t) \\ \alpha^*(t) & -\frac{1}{2}\epsilon(t) \end{bmatrix}$	$H = \epsilon(t) J_z + \alpha(t) J_+ + \alpha^*(t) J_- \xrightarrow{j \rightarrow \frac{1}{2}} \begin{bmatrix} \frac{1}{2}\epsilon(t) & \alpha(t) \\ \alpha^*(t) & -\frac{1}{2}\epsilon(t) \end{bmatrix}$	$H = \epsilon(t) J_z + \alpha(t) J_+ + \alpha^*(t) J_- \xrightarrow{j \rightarrow \frac{1}{2}} \begin{bmatrix} \frac{1}{2}\epsilon(t) & \alpha(t) \\ \alpha^*(t) & -\frac{1}{2}\epsilon(t) \end{bmatrix}$
gilmore-lie-groups_ E00422 (7.58)	129	■ 1.000	$\frac{d}{dt} \begin{bmatrix} a(t) & b(t) \\ -b^*(t) & a^*(t) \end{bmatrix} = -\frac{i}{\hbar} \begin{bmatrix} \frac{1}{2}\epsilon(t) & \alpha(t) \\ \alpha^*(t) & -\frac{1}{2}\epsilon(t) \end{bmatrix} \begin{bmatrix} a(t) & b(t) \\ -b^*(t) & a^*(t) \end{bmatrix}$	$\frac{d}{dt} \begin{bmatrix} a(t) & b(t) \\ -b^*(t) & a^*(t) \end{bmatrix} = -\frac{i}{\hbar} \begin{bmatrix} \frac{1}{2}\epsilon(t) & \alpha(t) \\ \alpha^*(t) & -\frac{1}{2}\epsilon(t) \end{bmatrix} \begin{bmatrix} a(t) & b(t) \\ -b^*(t) & a^*(t) \end{bmatrix}$	$\frac{d}{dt} \begin{bmatrix} a(t) & b(t) \\ -b^*(t) & a^*(t) \end{bmatrix} = -\frac{i}{\hbar} \begin{bmatrix} \frac{1}{2}\epsilon(t) & \alpha(t) \\ \alpha^*(t) & -\frac{1}{2}\epsilon(t) \end{bmatrix} \begin{bmatrix} a(t) & b(t) \\ -b^*(t) & a^*(t) \end{bmatrix}$
gilmore-lie-groups_ E00423 (7.59)	129	■ 1.000	$H = \omega(t) a^\dagger a + \alpha(t) a^\dagger + \alpha^*(t) a + \delta(t) I$ $\rightarrow \begin{bmatrix} 0 & \alpha^*(t) & \delta(t) \\ 0 & \omega(t) & \alpha(t) \\ 0 & 0 & 0 \end{bmatrix}$	$H = \omega(t) a^\dagger a + \alpha(t) a^\dagger + \alpha^*(t) a + \delta(t) I \rightarrow \begin{bmatrix} 0 & \alpha^*(t) & \delta(t) \\ 0 & \omega(t) & \alpha(t) \\ 0 & 0 & 0 \end{bmatrix}$	$H = \omega(t) a^\dagger a + \alpha(t) a^\dagger + \alpha^*(t) a + \delta(t) I \rightarrow \begin{bmatrix} 0 & \alpha^*(t) & \delta(t) \\ 0 & \omega(t) & \alpha(t) \\ 0 & 0 & 0 \end{bmatrix}$
gilmore-lie-groups_ E00425 (7.61)	130	■ 1.000	$\langle \mathcal{O} \rangle = \text{tr } \rho \mathcal{O}$	$\langle \mathcal{O} \rangle = \text{tr } \rho \mathcal{O}$	$\langle \mathcal{O} \rangle = \text{tr } \rho \mathcal{O}$
gilmore-lie-groups_ E00426 (7.62)	130	■ 1.000	$H = \epsilon J_z + \alpha J_+ + \alpha^* J_-$ $\xrightarrow{j \rightarrow \frac{1}{2}} \begin{bmatrix} \frac{1}{2}\epsilon(t) & \alpha(t) \\ \alpha^*(t) & -\frac{1}{2}\epsilon(t) \end{bmatrix}$	$H = \epsilon J_z + \alpha J_+ + \alpha^* J_- \xrightarrow{j \rightarrow \frac{1}{2}} \begin{bmatrix} \frac{1}{2}\epsilon(t) & \alpha(t) \\ \alpha^*(t) & -\frac{1}{2}\epsilon(t) \end{bmatrix}$	$H = \epsilon J_z + \alpha J_+ + \alpha^* J_- \xrightarrow{j \rightarrow \frac{1}{2}} \begin{bmatrix} \frac{1}{2}\epsilon(t) & \alpha(t) \\ \alpha^*(t) & -\frac{1}{2}\epsilon(t) \end{bmatrix}$
gilmore-lie-groups_ E00427 (7.63)	131	■ 1.000	$e^{-\beta H} e^\Lambda \rightarrow \left(I_2 \cosh(\beta H) - \beta \begin{bmatrix} \epsilon/2 & \alpha \\ \alpha^* & -\epsilon/2 \end{bmatrix} \frac{\sinh(\beta H)}{\beta H } \right) \times \left(I_2 \cosh(\Lambda) + \begin{bmatrix} \lambda_3/2 & \lambda \\ \lambda^* & -\lambda_3/2 \end{bmatrix} \frac{\sinh(\Lambda)}{ \Lambda } \right)$	$e^{-\beta H} e^\Lambda \rightarrow \left(I_2 \cosh(\beta H) - \beta \begin{bmatrix} \epsilon/2 & \alpha \\ \alpha^* & -\epsilon/2 \end{bmatrix} \frac{\sinh(\beta H)}{\beta H } \right) \times \left(I_2 \cosh(\Lambda) + \begin{bmatrix} \lambda_3/2 & \lambda \\ \lambda^* & -\lambda_3/2 \end{bmatrix} \frac{\sinh(\Lambda)}{ \Lambda } \right)$	$e^{-\beta H} e^\Lambda \rightarrow \left(I_2 \cosh(\beta H) - \beta \begin{bmatrix} \epsilon/2 & \alpha \\ \alpha^* & -\epsilon/2 \end{bmatrix} \frac{\sinh(\beta H)}{\beta H } \right) \times \left(I_2 \cosh(\Lambda) + \begin{bmatrix} \lambda_3/2 & \lambda \\ \lambda^* & -\lambda_3/2 \end{bmatrix} \frac{\sinh(\Lambda)}{ \Lambda } \right)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0439 (7.70)	133	■ 1.000	$\begin{aligned} -A = & \beta \hbar \omega - \lambda_n \\ B = & \frac{e^{-\beta \hbar \omega} - 1 + \beta \hbar \omega}{(\hbar \omega)^2} \alpha^* \alpha - \beta \delta + d + \frac{e^{\lambda_n} - 1 - \lambda_n}{\lambda_n^2} \lambda^* \lambda \\ & + \frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \frac{e^{\lambda_n} - 1}{\lambda_n} (\alpha^* \lambda + \alpha \lambda^*) / (1 - e^{-(\beta \hbar \omega - \lambda_n)}) \\ & + \left[e^{-\beta \hbar \omega} \left(\frac{e^{\lambda_n} - 1}{\lambda_n} \right)^2 \lambda^* \lambda + e^{\lambda_n} \left(\frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \right)^2 \alpha^* \alpha \right] / [1 - e^{-(\beta \hbar \omega - \lambda_n)}] \end{aligned}$	$-A = \beta \hbar \omega - \lambda_n$ $B = \frac{e^{-\beta \hbar \omega} - 1 + \beta \hbar \omega}{(\hbar \omega)^2} \alpha^* \alpha - \beta \delta + d + \frac{e^{\lambda_n} - 1 - \lambda_n}{\lambda_n^2} \lambda^* \lambda$ $+ \frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \frac{e^{\lambda_n} - 1}{\lambda_n} (\alpha^* \lambda + \alpha \lambda^*) / (1 - e^{-(\beta \hbar \omega - \lambda_n)})$ $+ \left[e^{-\beta \hbar \omega} \left(\frac{e^{\lambda_n} - 1}{\lambda_n} \right)^2 \lambda^* \lambda + e^{\lambda_n} \left(\frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \right)^2 \alpha^* \alpha \right] / [1 - e^{-(\beta \hbar \omega - \lambda_n)}]$	$-A = \beta \hbar \omega - \lambda_n$ $B = \frac{e^{-\beta \hbar \omega} - 1 + \beta \hbar \omega}{(\hbar \omega)^2} \alpha^* \alpha - \beta \delta + d + \frac{e^{\lambda_n} - 1 - \lambda_n}{\lambda_n^2} \lambda^* \lambda$ $+ \frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \frac{e^{\lambda_n} - 1}{\lambda_n} (\alpha^* \lambda + \alpha \lambda^*) / (1 - e^{-(\beta \hbar \omega - \lambda_n)})$ $+ \left[e^{-\beta \hbar \omega} \left(\frac{e^{\lambda_n} - 1}{\lambda_n} \right)^2 \lambda^* \lambda + e^{\lambda_n} \left(\frac{e^{-\beta \hbar \omega} - 1}{\hbar \omega} \right)^2 \alpha^* \alpha \right] / [1 - e^{-(\beta \hbar \omega - \lambda_n)}]$
gilmore-lie-groups_ EQ0440	135	■ 1.000	$a^{\dagger} = \frac{1}{\sqrt{2}} \left(x - \frac{d}{dx} \right) \quad a = \frac{1}{\sqrt{2}} \left(x + \frac{d}{dx} \right)$	$a^{\dagger} = \frac{1}{\sqrt{2}} \left(x - \frac{d}{dx} \right) \quad a = \frac{1}{\sqrt{2}} \left(x + \frac{d}{dx} \right)$	$a^{\dagger} = \frac{1}{\sqrt{2}} \left(x - \frac{d}{dx} \right) \quad a = \frac{1}{\sqrt{2}} \left(x + \frac{d}{dx} \right)$
gilmore-lie-groups_ EQ0441	135	■ 1.000	$\langle n' x^k n \rangle = \frac{d^k}{d\lambda^k} \langle n' e^{\lambda x} n \rangle_{\lambda=0}$	$\langle n' x^k n \rangle = \frac{d^k}{d\lambda^k} \langle n' e^{\lambda x} n \rangle_{\lambda=0}$	$\langle n' x^k n \rangle = \frac{d^k}{d\lambda^k} \langle n' e^{\lambda x} n \rangle_{\lambda=0}$
gilmore-lie-groups_ EQ0442	135	■ 1.000	$e^{\lambda x} = e^{\lambda(a^{\dagger} + a)/\sqrt{2}} = \text{EXP} \left(\frac{\lambda}{\sqrt{2}} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \right) = \begin{bmatrix} 1 & \lambda/\sqrt{2} & \lambda^2/4 \\ 0 & 1 & \lambda/\sqrt{2} \\ 0 & 0 & 1 \end{bmatrix}$	$e^{\lambda x} = e^{\lambda(a^{\dagger} + a)/\sqrt{2}} = \text{EXP} \left(\frac{\lambda}{\sqrt{2}} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \right) = \begin{bmatrix} 1 & \lambda/\sqrt{2} & \lambda^2/4 \\ 0 & 1 & \lambda/\sqrt{2} \\ 0 & 0 & 1 \end{bmatrix}$	$e^{\lambda x} = e^{\lambda(a^{\dagger} + a)/\sqrt{2}} = \text{EXP} \left(\frac{\lambda}{\sqrt{2}} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \right) = \begin{bmatrix} 1 & \lambda/\sqrt{2} & \lambda^2/4 \\ 0 & 1 & \lambda/\sqrt{2} \\ 0 & 0 & 1 \end{bmatrix}$
gilmore-lie-groups_ EQ0444	135	■ 1.000	$e^{\lambda(a^{\dagger} + a)/\sqrt{2}} = e^{\lambda a^{\dagger}/\sqrt{2}} e^{\lambda^2/4} e^{\lambda a/\sqrt{2}}$	$e^{\lambda(a^{\dagger} + a)/\sqrt{2}} = e^{\lambda a^{\dagger}/\sqrt{2}} e^{\lambda^2/4} e^{\lambda a/\sqrt{2}}$	$e^{\lambda(a^{\dagger} + a)/\sqrt{2}} = e^{\lambda a^{\dagger}/\sqrt{2}} e^{\lambda^2/4} e^{\lambda a/\sqrt{2}}$
gilmore-lie-groups_ EQ0446 (7.71)	136	■ 1.000	$\langle e^{ikx} \rangle = \frac{\text{tr} e^{ikx} e^{-\beta \mathcal{H}}}{\text{tr} e^{-\beta \mathcal{H}}}$	$\langle e^{ikx} \rangle = \frac{\text{tr} e^{ikx} e^{-\beta \mathcal{H}}}{\text{tr} e^{-\beta \mathcal{H}}}$	$\langle e^{ikx} \rangle = \frac{\text{tr} e^{ikx} e^{-\beta \mathcal{H}}}{\text{tr} e^{-\beta \mathcal{H}}}$
gilmore-lie-groups_ EQ0447 (7.72)	136	■ 1.000	$\text{tr} e^{ikx} e^{-\beta \mathcal{H}} = \sum_{n=0}^{\infty} \langle n e^{ikx} n \rangle e^{-n\beta \hbar \omega} = \sum_{n=0}^{\infty} \langle n e^{ikx} e^{-n\beta \hbar \omega} n \rangle$	$\text{tr} e^{ikx} e^{-\beta \mathcal{H}} = \sum_{n=0}^{\infty} \langle n e^{ikx} n \rangle e^{-n\beta \hbar \omega} = \sum_{n=0}^{\infty} \langle n e^{ikx} e^{-n\beta \hbar \omega} n \rangle$	$\text{tr} e^{ikx} e^{-\beta \mathcal{H}} = \sum_{n=0}^{\infty} \langle n e^{ikx} n \rangle e^{-n\beta \hbar \omega} = \sum_{n=0}^{\infty} \langle n e^{ikx} e^{-n\beta \hbar \omega} n \rangle$
gilmore-lie-groups_ EQ0448 (7.73)	136	■ 1.000	$\text{tr} e^{ikx} e^{-\beta \hbar \omega a^{\dagger} a} = \text{tr} e^{-\beta \hbar \omega a^{\dagger} a} e^{\delta} = e^{\delta} \text{tr} e^{-\beta \hbar \omega a^{\dagger} a}$	$\text{tr} e^{ikx} e^{-\beta \hbar \omega a^{\dagger} a} = \text{tr} e^{-\beta \hbar \omega a^{\dagger} a} e^{\delta} = e^{\delta} \text{tr} e^{-\beta \hbar \omega a^{\dagger} a}$	$\text{tr} e^{ikx} e^{-\beta \hbar \omega a^{\dagger} a} = \text{tr} e^{-\beta \hbar \omega a^{\dagger} a} e^{\delta} = e^{\delta} \text{tr} e^{-\beta \hbar \omega a^{\dagger} a}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00449 (7.74)	136	■ 1.000	$M(S) M\left(e^{\mathrm{i} \, k \, x}\right) M\left(e^{-\beta \hbar \omega \, \hbar \omega \, \hat{n}}\right) M\left(S^{-1}\right)=M\left(e^{-\beta \hbar \omega \, \hbar \omega \, \hat{n}}\right) M\left(e^{\delta}\right)$	$M(S)M\left(e^{ikx}\right)M\left(e^{-\beta\hbar\omega\hat{n}}\right)M\left(S^{-1}\right)=M\left(e^{-\beta\hbar\omega'\hat{n}}\right)M\left(e^{\delta}\right)$	$M(S) M(e^{ikx}) M(e^{-\beta\hbar\omega\hat{n}}) M(S^{-1}) = M(e^{-\beta\hbar\omega'\hat{n}}) M(e^{\delta})$
gilmore-lie-groups_ E00450 (7.76)	136	■ 1.000	$\left\langle e^{\mathrm{i} \, k \, x}\right\rangle=e^{\delta} \quad \delta=-\frac{1}{2} k^2 \coth \left(\frac{1}{2} \beta \hbar \omega\right)$	$\left\langle e^{ikx}\right\rangle=e^{\delta} \quad \delta=-\frac{1}{2}k^2\coth\left(\frac{1}{2}\beta\hbar\omega\right)$	$\langle e^{ikx}\rangle=e^{\delta} \quad \delta=-\frac{1}{2}k^2\coth\left(\frac{1}{2}\beta\hbar\omega\right)$
gilmore-lie-groups_ E00451 (7.77)	137	■ 1.000	$\left\langle e^{x^i X_i}\right\rangle=\frac{\mathrm{tr} e^{R\left(x^i X_i\right)} e^{-\beta \mathcal{H}}}{\mathrm{tr} e^{-\beta \mathcal{H}}} \rightarrow \frac{\mathrm{tr} R\left(e^{x^i X_i}\right) R\left(e^{-\beta a^i X_i}\right)}{\mathrm{tr} R\left(e^{-\beta a^i X_i}\right)}$	$\left\langle e^{x^i X_i}\right\rangle=\frac{\mathrm{tr} e^{R\left(x^i X_i\right)} e^{-\beta \mathcal{H}}}{\mathrm{tr} e^{-\beta \mathcal{H}}} \rightarrow \frac{\mathrm{tr} R\left(e^{x^i X_i}\right) R\left(e^{-\beta a^i X_i}\right)}{\mathrm{tr} R\left(e^{-\beta a^i X_i}\right)}$	$\langle e^{x^i X_i}\rangle=\frac{\mathrm{tr} e^{R(x^i X_i)} e^{-\beta \mathcal{H}}}{\mathrm{tr} e^{-\beta \mathcal{H}}} \rightarrow \frac{\mathrm{tr} R(e^{x^i X_i}) R(e^{-\beta a^i X_i})}{\mathrm{tr} R(e^{-\beta a^i X_i})}$
gilmore-lie-groups_ E00453 (7.79)	137	■ 1.000	$\left\langle e^{x^i X_i}\right\rangle=\frac{\mathrm{tr} R\left(e^{-\beta d^i(x, a) H_i}\right)}{\mathrm{tr} R\left(e^{-\beta d^i(0, a) H_i}\right)}$	$\left\langle e^{x^i X_i}\right\rangle=\frac{\mathrm{tr} R\left(e^{-\beta d^i(x, a) H_i}\right)}{\mathrm{tr} R\left(e^{-\beta d^i(0, a) H_i}\right)}$	$\langle e^{x^i X_i}\rangle=\frac{\mathrm{tr} R(e^{-\beta d^i(x, a) H_i})}{\mathrm{tr} R(e^{-\beta d^i(0, a) H_i})}$
gilmore-lie-groups_ E00454 (7.80)	137	■ 1.000	$U(\alpha) 0\rangle=e^{\left(\alpha a^{\dagger}-\alpha^* a\right)} 0\rangle=e^{\beta a^{\dagger}} e^{\delta I} e^{\beta' a} 0\rangle$	$U(\alpha) 0\rangle=e^{\left(\alpha a^{\dagger}-\alpha^* a\right)} 0\rangle=e^{\beta a^{\dagger}} e^{\delta I} e^{\beta' a} 0\rangle$	$U(\alpha) 0\rangle=e^{\left(\alpha a^{\dagger}-\alpha^* a\right)} 0\rangle=e^{\beta a^{\dagger}} e^{\delta I} e^{\beta' a} 0\rangle$
gilmore-lie-groups_ E00455 (7.81)	138	■ 1.000	$\operatorname{EXP}\left[\begin{array}{ccc} 1 & -\alpha^* & 0 \\ 0 & 1 & \alpha \\ 0 & 0 & 1 \end{array}\right]=\left[\begin{array}{ccc} 1 & -\alpha^* & -\alpha^* \alpha / 2 \\ 0 & 1 & \alpha \\ 0 & 0 & 1 \end{array}\right]$	$\operatorname{EXP}\left[\begin{array}{ccc} 1 & -\alpha^* & 0 \\ 0 & 1 & \alpha \\ 0 & 0 & 1 \end{array}\right]=\left[\begin{array}{ccc} 1 & -\alpha^* & -\alpha^* \alpha / 2 \\ 0 & 1 & \alpha \\ 0 & 0 & 1 \end{array}\right]$	$\operatorname{EXP}\left[\begin{array}{ccc} 1 & -\alpha^* & 0 \\ 0 & 1 & \alpha \\ 0 & 0 & 1 \end{array}\right]=\left[\begin{array}{ccc} 1 & -\alpha^* & -\alpha^* \alpha / 2 \\ 0 & 1 & \alpha \\ 0 & 0 & 1 \end{array}\right]$
gilmore-lie-groups_ E00457 (7.83)	138	■ 1.000	$e^{\left(\alpha a^{\dagger}-\alpha^* a\right)} 0\rangle=e^{\alpha a^{\dagger}} e^{-\alpha^* \alpha I / 2} e^{-\alpha^* a} 0\rangle=\sum \frac{\left(\alpha a^{\dagger}\right)^n}{n!} 0\rangle e^{-\alpha^* \alpha / 2}$	$e^{\left(\alpha a^{\dagger}-\alpha^* a\right)} 0\rangle=e^{\alpha a^{\dagger}} e^{-\alpha^* \alpha I / 2} e^{-\alpha^* a} 0\rangle=\sum \frac{\left(\alpha a^{\dagger}\right)^n}{n!} 0\rangle e^{-\alpha^* \alpha / 2}$	$e^{\left(\alpha a^{\dagger}-\alpha^* a\right)} 0\rangle=e^{\alpha a^{\dagger}} e^{-\alpha^* \alpha I / 2} e^{-\alpha^* a} 0\rangle=\sum \frac{\left(\alpha a^{\dagger}\right)^n}{n!} 0\rangle e^{-\alpha^* \alpha / 2}$
gilmore-lie-groups_ E00458 (7.84)	138	■ 1.000	$ \alpha\rangle=U(\alpha) 0\rangle=e^{-\alpha^* \alpha / 2} \sum \frac{\alpha^n}{\sqrt{n!}} n\rangle$	$ \alpha\rangle=U(\alpha) 0\rangle=e^{-\alpha^* \alpha / 2} \sum \frac{\alpha^n}{\sqrt{n!}} n\rangle$	$ \alpha\rangle=U(\alpha) 0\rangle=e^{-\alpha^* \alpha / 2} \sum \frac{\alpha^n}{\sqrt{n!}} n\rangle$
gilmore-lie-groups_ E00463	139	■ 1.000	$\left(E J_3+p J_1-Z\right) u\rangle=0$	$(E J_3+p J_1-Z) u\rangle=0$	$(E J_3+p J_1-Z) u\rangle=0$
gilmore-lie-groups_ E00464	139	■ 1.000	$\begin{aligned} \operatorname{EXP}\left(\mathrm{i} \beta J_y\right)_{m n}= & D_{m n}^j(\beta)=P_{m n}^j(z)=\frac{(-)^{j-m}}{2^j(j-n)!}\left[\frac{(j-n)!(j+m)!}{(j+n)!(j-m)!}\right]^{1 / 2} \\ & \times(1+z)^{-(m+n) / 2}(1-z)^{-(m-n) / 2}\left(\frac{d}{d z}\right)^{j-m}\left[(1-z)^{j-n}(1+z)^{j+n}\right] \end{aligned}$	$\operatorname{EXP}\left(\mathrm{i} \beta J_y\right)_{m n}=D_{m n}^j(\beta)=P_{m n}^j(z)=\frac{(-)^{j-m}}{2^j(j-n)!}\left[\frac{(j-n)!(j+m)!}{(j+n)!(j-m)!}\right]^{1 / 2} \\ \times(1+z)^{-(m+n) / 2}(1-z)^{-(m-n) / 2}\left(\frac{d}{d z}\right)^{j-m}\left[(1-z)^{j-n}(1+z)^{j+n}\right]$	$\operatorname{EXP}(i\beta J_y)_{mn}=D_{mn}^j(\beta)=P_{mn}^j(z)=\frac{(-)^{j-m}}{2^j(j-n)!}\left[\frac{(j-n)!(j+m)!}{(j+n)!(j-m)!}\right]^{1/2} \\ \times(1+z)^{-(m+n)/2}(1-z)^{-(m-n)/2}\left(\frac{d}{dz}\right)^{j-m}\left[(1-z)^{j-n}(1+z)^{j+n}\right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E04465	140	■ 1.000	$\left[\begin{array}{cc} e^{i\theta/2} & 0 \\ 0 & e^{-i\theta/2} \end{array}\right] \xrightarrow{2\rightarrow 1} \left[\begin{array}{cc} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{array}\right]$		
gilmore-lie-groups_ E04466	140	■ 1.000	$\mathcal{H} = \sum_{i=1}^N \frac{\epsilon}{2} \sigma_z^{(i)} + \hbar\omega a^\dagger a + \frac{\lambda}{\sqrt{N}} \sum_{i=1}^N \sigma_+^{(i)} a + \sigma_-^{(i)} a^\dagger$		
gilmore-lie-groups_ E04467	140	■ 1.000	$\left[\sigma_z^{(i)}, \sigma_\pm^{(j)}\right] = \pm \sigma_\pm^{(i)} \delta_{ij} \quad \left[\sigma_+^{(i)}, \sigma_-^{(j)}\right] = \sigma_z^{(i)} \delta_{ij}$		
gilmore-lie-groups_ E04468	141	■ 1.000	$\mathcal{H}_{\text{field}} = \epsilon \langle J_z(t) \rangle + \hbar\omega a^\dagger a + \frac{\lambda}{\sqrt{N}} (\langle J_+(t) \rangle a + \langle J_-(t) \rangle a^\dagger)$		
gilmore-lie-groups_ E04472 (8.4)	144	■ 1.000	$\Re e g \left(l a + r a^\dagger + \delta I \right) = \left[\begin{array}{ccc} 0 & 0 & l \\ 0 & 0 & -r \\ 0 & 0 & 0 \end{array} \right] \begin{array}{c} a^\dagger \\ a \\ I \end{array}$		
gilmore-lie-groups_ E04475 (8.7)	145	■ 1.000	$[X_{ij}, X_{rs}] = X_{is} \delta_{jr} + X_{jr} \delta_{is} - X_{ir} \delta_{js} - X_{js} \delta_{ir}$		
gilmore-lie-groups_ E04477 (8.9)	146	■ 1.000	$\begin{aligned} & \left\{ \left[Y_i, Y_j \right] \right\} = -\epsilon_{ijk} Y_k \\ & \left\{ \left[Z_i, Z_j \right] \right\} = +\epsilon_{ijk} Z_k \\ & \left\{ \left[Y_i, Z_j \right] \right\} = 0 \end{aligned}$		
gilmore-lie-groups_ E04480 (8.12)	147	■ 1.000	$[X_i, X_j] = 0$		
gilmore-lie-groups_ E04481 (8.13)	147	■ 1.000	$\begin{array}{l} [Z, X_i] = R(Z)_i^j X_j \\ R(Z)_i^j = 0 \text{ unless } j \geq i \text{ solvable} \end{array} \quad \begin{array}{l} j > i \text{ nilpotent} \\ j \geq i \text{ solvable} \end{array}$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0482 (8.14)	147	■ 1.000	$\begin{array}{c} V_1 \text{ \texttt{\text{spanned by}} } X_n, \\ X_{n-1}, X_{n-2}, \ldots, X_2, X_1 \\ \cup \\ V_2 \text{ \texttt{\text{spanned by}} } X_n, X_{n-1}, X_{n-2}, \ldots, X_2 \\ \cup \\ \vdots \qquad \qquad \qquad \vdots \\ \cup \\ V_{n-2} \text{ \texttt{\text{spanned by}} } X_n, X_{n-1}, X_{n-2} \\ \cup \\ V_{n-1} \text{ \texttt{\text{spanned by}} } X_n, X_{n-1} \\ \cup \\ V_n \text{ \texttt{\text{spanned by}} } X_n \end{array}$	$\begin{array}{c} V_1 \text{ \texttt{\text{spanned by}} } X_n, X_{n-1}, X_{n-2}, \ldots, X_2, X_1 \\ \cup \\ V_2 \text{ \texttt{\text{spanned by}} } X_n, X_{n-1}, X_{n-2}, \ldots, X_2 \\ \cup \vdots \\ \vdots \cup \\ V_{n-2} \text{ \texttt{\text{spanned by}} } X_n, X_{n-1}, X_{n-2} \\ \cup \\ V_{n-1} \text{ \texttt{\text{spanned by}} } X_n, X_{n-1} \\ \cup \\ V_n \text{ \texttt{\text{spanned by}} } X_n \end{array}$	$\begin{array}{c} V_1 \text{ \texttt{\text{spanned by}} } X_n, X_{n-1}, X_{n-2}, \ldots, X_2, X_1 \\ \cup \\ V_2 \text{ \texttt{\text{spanned by}} } X_n, X_{n-1}, X_{n-2}, \ldots, X_2 \\ \cup \\ \vdots \qquad \qquad \qquad \vdots \\ \cup \\ V_{n-2} \text{ \texttt{\text{spanned by}} } X_n, X_{n-1}, X_{n-2} \\ \cup \\ V_{n-1} \text{ \texttt{\text{spanned by}} } X_n, X_{n-1} \\ \cup \\ V_n \text{ \texttt{\text{spanned by}} } X_n \end{array}$
gilmore-lie-groups_ EQ0483 (8.17)	148	■ 1.000	$X=[Y, \ Z]$	$X = [Y, Z]$	$X = [Y, Z]$
gilmore-lie-groups_ EQ0484 (8.18)	149	■ 1.000	$\mathfrak{g}=\left(V_{-}+V_{+}\right)+V_{0}$	$\mathfrak{g} = (V_- + V_+) + V_0$	$\mathfrak{g} = (V_- + V_+) + V_0$
gilmore-lie-groups_ EQ0485 (8.19)	149	■ 1.000	$\mathfrak{g}^{\prime}=\left(V_{-}^{\prime}+V_{+}^{\prime}\right)+V_{0}^{\prime}$	$\mathfrak{g}' = (V'_- + V'_+) + V'_0$	$\mathfrak{g}' = (V'_- + V'_+) + V'_0$
gilmore-lie-groups_ EQ0487 (8.21)	150	■ 1.000	$(X, \ X)=-40 \ R \ L+10 \ \eta^2$	$(X, X) = -40RL + 10\eta^2$	$(X, X) = -40RL + 10\eta^2$
gilmore-lie-groups_ EQ0488 (8.22)	150	■ 1.000	$\eta\left(a^{\dagger}a+\frac{1}{2}\right)+Ra^{\dagger 2}+La^2\longrightarrow\left[\begin{array}{ccc} 0 & -2R & 2L \\ 4L & 2\eta & 0 \\ -4R & 0 & -2\eta \end{array}\right]\begin{array}{c} \hat{n}+\frac{1}{2} \\ a^{\dagger 2} \\ a^2 \end{array}$	$\eta\left(a^{\dagger}a+\frac{1}{2}\right)+Ra^{\dagger 2}+La^2\longrightarrow\left[\begin{array}{ccc} 0 & -2R & 2L \\ 4L & 2\eta & 0 \\ -4R & 0 & -2\eta \end{array}\right]\begin{array}{c} \hat{n}+\frac{1}{2} \\ a^{\dagger 2} \\ a^2 \end{array}$	$\eta\left(a^{\dagger}a+\frac{1}{2}\right)+Ra^{\dagger 2}+La^2\longrightarrow\left[\begin{array}{ccc} 0 & -2R & 2L \\ 4L & 2\eta & 0 \\ -4R & 0 & -2\eta \end{array}\right]\begin{array}{c} \hat{n}+\frac{1}{2} \\ a^{\dagger 2} \\ a^2 \end{array}$
gilmore-lie-groups_ EQ0489 (8.23)	150	■ 1.000	$(X, \ X)^{\prime}=-32 \ R \ L+8 \ \eta^2$	$(X, X)' = -32RL + 8\eta^2$	$(X, X)' = -32RL + 8\eta^2$
gilmore-lie-groups_ EQ0491 (8.26)	152	■ 1.000	$\begin{array}{l} X_{ij} = x^i \partial_j - x^j \partial_i \\ Y_i = 2t \frac{\partial}{\partial x^i} - x^i u \frac{\partial}{\partial u} \\ Z = 2t \frac{\partial}{\partial t} + x^i \frac{\partial}{\partial x^i} - nu \frac{\partial}{\partial u} \end{array}$	$\begin{array}{l} X_{ij} = x^i \partial_j - x^j \partial_i \\ Y_i = 2t \frac{\partial}{\partial x^i} - x^i u \frac{\partial}{\partial u} \\ Z = 2t \frac{\partial}{\partial t} + x^i \frac{\partial}{\partial x^i} - nu \frac{\partial}{\partial u} \end{array}$	$\begin{array}{l} X_{ij} = x^i \partial_j - x^j \partial_i \\ Y_i = 2t \frac{\partial}{\partial x^i} - x^i u \frac{\partial}{\partial u} \\ Z = 2t \frac{\partial}{\partial t} + x^i \frac{\partial}{\partial x^i} - nu \frac{\partial}{\partial u} \end{array}$
gilmore-lie-groups_ EQ0492 (9.1)	154	■ 1.000	$[Z, \ X]=\lambda X$	$[Z, X] = \lambda X$	$[Z, X] = \lambda X$
gilmore-lie-groups_ EQ0494 (9.3)	154	■ 1.000	$\left\ R(Z)-\lambda I_N\right\ =0$	$\ R(Z) - \lambda I_N\ = 0$	$\ R(Z) - \lambda I_N\ = 0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08495 (9.4)	154	■ 1.000	$\sum_{j=0}^N (-\lambda)^{N-j} \phi_j(Z)=0$	$\sum_{j=0}^N (-\lambda)^{N-j} \phi_j(Z) = 0$	$\sum_{j=0}^N (-\lambda)^{N-j} \phi_j(Z) = 0$
gilmore-lie-groups_ E08496 (9.5)	154	■ 1.000	$\phi_j(Z) \rightarrow \phi_j(z^i)$	$\phi_j(Z) \rightarrow \phi_j(z^i)$	$\phi_j(Z) \rightarrow \phi_j(z^i)$
gilmore-lie-groups_ E08503 (9.12)	156	■ 1.000	$\left \mathfrak{d e f}(X)-\lambda I_2\right =\lambda^2+\left(\frac{1}{2}\right)^2\left(a_1^2+a_2^2+a_3^2\right)=0$	$\ \mathfrak{d e f}(X)-\lambda I_2\ =\lambda^2+\left(\frac{1}{2}\right)^2\left(a_1^2+a_2^2+a_3^2\right)=0$	$\ \mathfrak{d e f}(X)-\lambda I_2\ =\lambda^2+\left(\frac{1}{2}\right)^2\left(a_1^2+a_2^2+a_3^2\right)=0$
gilmore-lie-groups_ E08504 (9.13)	156	■ 1.000	$\left \mathfrak{d e f}(Y)-\lambda I_2\right =\lambda^2+\left(\frac{1}{2}\right)^2\left(-b_1^2-b_2^2+b_3^2\right)=0$	$\ \mathfrak{d e f}(Y)-\lambda I_2\ =\lambda^2+\left(\frac{1}{2}\right)^2\left(-b_1^2-b_2^2+b_3^2\right)=0$	$\ \mathfrak{d e f}(Y)-\lambda I_2\ =\lambda^2+\left(\frac{1}{2}\right)^2\left(-b_1^2-b_2^2+b_3^2\right)=0$
gilmore-lie-groups_ E08505 (9.14)	157	■ 1.000	$z^i \rightarrow X_i \quad \phi_j\left(z^i\right) \longrightarrow \phi_j\left(X_i\right)$	$z^i \rightarrow X_i \quad \phi_j\left(z^i\right) \longrightarrow \phi_j\left(X_i\right)$	$z^i \rightarrow X_i \quad \phi_j\left(z^i\right) \longrightarrow \phi_j\left(X_i\right)$
gilmore-lie-groups_ E08506 (9.15)	157	■ 1.000	$\left[\phi_j\left(X_i\right), X_k\right]=0$	$\left[\phi_j\left(X_i\right), X_k\right]=0$	$\left[\phi_j\left(X_i\right), X_k\right]=0$
gilmore-lie-groups_ E08507 (9.16)	157	■ 1.000	$\phi_2(\mathbf{a})=+a_1^2+a_2^2+a_3^2 \longrightarrow \phi_2(X)=+X_1^2+X_2^2+X_3^2$	$\phi_2(\mathbf{a})=+a_1^2+a_2^2+a_3^2 \longrightarrow \phi_2(X)=+X_1^2+X_2^2+X_3^2$	$\phi_2(\mathbf{a})=+a_1^2+a_2^2+a_3^2 \longrightarrow \phi_2(X)=+X_1^2+X_2^2+X_3^2$
gilmore-lie-groups_ E08508 (9.17)	157	■ 1.000	$\phi_2(\mathbf{b})=-b_1^2-b_2^2+b_3^2 \longrightarrow \phi_2(Y)=-Y_1^2-Y_2^2+Y_3^2$	$\phi_2(\mathbf{b})=-b_1^2-b_2^2+b_3^2 \longrightarrow \phi_2(Y)=-Y_1^2-Y_2^2+Y_3^2$	$\phi_2(\mathbf{b})=-b_1^2-b_2^2+b_3^2 \longrightarrow \phi_2(Y)=-Y_1^2-Y_2^2+Y_3^2$
gilmore-lie-groups_ E08510 (9.19)	159	■ 1.000	$\left \mathfrak{d e f}(X)-\lambda I_n\right =\sum(-\lambda)^{n-j} \phi_j(X)=0$	$\ \mathfrak{d e f}(X)-\lambda I_n\ =\sum(-\lambda)^{n-j} \phi_j(X)=0$	$\ \mathfrak{d e f}(X)-\lambda I_n\ =\sum(-\lambda)^{n-j} \phi_j(X)=0$
gilmore-lie-groups_ E08511 (9.20)	159	■ 1.000	$\phi_j(X)=\phi_j(-X)=(-)^j \phi_j(X)$	$\phi_j(X)=\phi_j(-X)=(-)^j \phi_j(X)$	$\phi_j(X)=\phi_j(-X)=(-)^j \phi_j(X)$
gilmore-lie-groups_ E08513 (9.22)	160	■ 1.000	$\mathcal{C}_4[\mathfrak{so}(5)]=\sum_{i=1}^5 v_i^2$	$\mathcal{C}_4[\mathfrak{so}(5)]=\sum_{i=1}^5 v_i^2$	$\mathcal{C}_4[\mathfrak{so}(5)]=\sum_{i=1}^5 v_i^2$
gilmore-lie-groups_ E08514 (9.23)	160	■ 1.000	$v^5=\epsilon^{ijkl5} X_{ij} X_{kl} \sim X_{12} X_{34}-X_{13} X_{24}+X_{14} X_{23}$	$v^5=\epsilon^{ijkl5} X_{ij} X_{kl} \sim X_{12} X_{34}-X_{13} X_{24}+X_{14} X_{23}$	$v^5=\epsilon^{ijkl5} X_{ij} X_{kl} \sim X_{12} X_{34}-X_{13} X_{24}+X_{14} X_{23}$
gilmore-lie-groups_ E08515 (9.24)	160	■ 1.000	$\mathcal{C}_4[\mathfrak{so}(4)]=\left(\epsilon^{ijkl} X_{ij} X_{kl}\right)^2 \sim\left(X_{12} X_{34}-X_{13} X_{24}+X_{14} X_{23}\right)^2$	$\mathcal{C}_4[\mathfrak{so}(4)]=\left(\epsilon^{ijkl} X_{ij} X_{kl}\right)^2 \sim\left(X_{12} X_{34}-X_{13} X_{24}+X_{14} X_{23}\right)^2$	$\mathcal{C}_4[\mathfrak{so}(4)]=\left(\epsilon^{ijkl} X_{ij} X_{kl}\right)^2 \sim\left(X_{12} X_{34}-X_{13} X_{24}+X_{14} X_{23}\right)^2$
gilmore-lie-groups_ E08516 (9.25)	160	■ 1.000	$Z=z_1\left(\hat{n}+\frac{1}{2}\right)+z_2 I$	$Z=z_1\left(\hat{n}+\frac{1}{2}\right)+z_2 I$	$Z=z_1\left(\hat{n}+\frac{1}{2}\right)+z_2 I$
gilmore-lie-groups_ E08519 (9.28)	161	■ 1.000	$\phi_2\left(z_1, z_2\right)=-z_1^2$	$\phi_2\left(z_1, z_2\right)=-z_1^2$	$\phi_2\left(z_1, z_2\right)=-z_1^2$
gilmore-lie-groups_ E08520 (9.29)	161	■ 1.000	$\left[\hat{n}+\frac{1}{2}, X_{(k, 0)}\right]=k X_{(k, 0)} \quad\left[I, X_{(k, 0)}\right]=0 X_{(k, 0)}$	$\left[\hat{n}+\frac{1}{2}, X_{(k, 0)}\right]=k X_{(k, 0)} \quad\left[I, X_{(k, 0)}\right]=0 X_{(k, 0)}$	$\left[\hat{n}+\frac{1}{2}, X_{(k, 0)}\right]=k X_{(k, 0)} \quad\left[I, X_{(k, 0)}\right]=0 X_{(k, 0)}$
gilmore-lie-groups_ E08521 (9.30)	161	■ 1.000	$\left[X_{(k, 0)}, X_{(l, 0)}\right] \sim X_{(k+l, 0)}$	$\left[X_{(k, 0)}, X_{(l, 0)}\right] \sim X_{(k+l, 0)}$	$\left[X_{(k, 0)}, X_{(l, 0)}\right] \sim X_{(k+l, 0)}$
gilmore-lie-groups_ E08522 (9.31)	162	■ 1.000	$\left[V_{\alpha}, V_{\beta}\right] \subset V_{\alpha+\beta}$	$\left[V_{\alpha}, V_{\beta}\right] \subset V_{\alpha+\beta}$	$\left[V_{\alpha}, V_{\beta}\right] \subset V_{\alpha+\beta}$
gilmore-lie-groups_ E08523 (9.32)	162	■ 1.000	$\left[H_i, H_j\right]=0 \quad 1 \leq i, j \leq l$	$\left[H_i, H_j\right]=0 \quad 1 \leq i, j \leq l$	$\left[H_i, H_j\right]=0 \quad 1 \leq i, j \leq l$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08525 (9.34)	162	■ 1.000	$\left[E_{\{\alpha\}}, E_{\{-\alpha\}}\right]=\alpha^i H_{\{i\}}$	$[E_{\alpha}, E_{-\alpha}] = \alpha^i H_i$	$[E_{\alpha}, E_{-\alpha}] = \alpha^i H_i$
gilmore-lie-groups_ E08526 (9.35)	162	■ 1.000	$(\alpha, \beta)=\alpha_{\{i\}} \beta_{\{i\}}=\alpha^j \beta_j$ $\beta_{\{j\}}=\alpha_{\{i\}} h^{\{i j\}} \beta_j$	$(\alpha, \beta) = \alpha_i \beta^i = \alpha^j \beta_j = \alpha_i h^{ij} \beta_j$	$(\alpha, \beta) = \alpha_i \beta^i = \alpha^j \beta_j = \alpha_i h^{ij} \beta_j$
gilmore-lie-groups_ E08528	163	■ 1.000	$\left[E_{\{\alpha\}}, E_{\{\beta\}}\right] \rightarrow$ $\begin{cases} 0 & \alpha+\beta \text { not a root } \\ N_{\alpha, \beta} E_{\alpha+\beta} & \alpha+\beta \text { a root } \\ \alpha \cdot \mathbf{H} & \alpha+\beta=0 \end{cases}$	$[E_{\alpha}, E_{\beta}] \rightarrow \begin{cases} 0 & \alpha+\beta \text { not a root } \\ N_{\alpha, \beta} E_{\alpha+\beta} & \alpha+\beta \text { a root } \\ \alpha \cdot \mathbf{H} & \alpha+\beta=0 \end{cases}$	$[E_{\alpha}, E_{\beta}] \rightarrow \begin{cases} 0 & \alpha+\beta \text { not a root } \\ N_{\alpha, \beta} E_{\alpha+\beta} & \alpha+\beta \text { a root } \\ \alpha \cdot \mathbf{H} & \alpha+\beta=0 \end{cases}$
gilmore-lie-groups_ E08529 (9.37)	163	■ 1.000	$\left[\left[E_{\{\alpha\}}, E_{\{\beta\}}\right], E_{\{\gamma\}}\right. \\ \left.\left.\right]+\left[\left[E_{\{\beta\}}, E_{\{\gamma\}}\right], E_{\{\alpha\}}\right]+\left[\left[E_{\{\gamma\}}, E_{\{\alpha\}}\right], E_{\{\beta\}}\right]=0$	$[[E_{\alpha}, E_{\beta}], E_{\gamma}] + [[E_{\beta}, E_{\gamma}], E_{\alpha}] + [[E_{\gamma}, E_{\alpha}], E_{\beta}] = 0$	$[[E_{\alpha}, E_{\beta}], E_{\gamma}] + [[E_{\beta}, E_{\gamma}], E_{\alpha}] + [[E_{\gamma}, E_{\alpha}], E_{\beta}] = 0$
gilmore-lie-groups_ E08532 (9.39)	163	■ 1.000	$\begin{aligned} & \beta-(m+1) \alpha \quad \& \quad \beta+(n+1) \alpha \\ & \alpha \end{aligned}$	$\beta-(m+1) \alpha$ $\beta+(n+1) \alpha$	$\beta-(m+1) \alpha$ $\beta+(n+1) \alpha$
gilmore-lie-groups_ E08533 (9.40)	163	■ 1.000	$N_{\{\alpha, \beta+(k-1) \alpha\}}^2=N_{\{\alpha, \beta+k \alpha\}}^2+\alpha \cdot(\beta+k \alpha)$	$N_{\alpha, \beta+(k-1) \alpha}^2=N_{\alpha, \beta+k \alpha}^2+\alpha \cdot(\beta+k \alpha)$	$N_{\alpha, \beta+(k-1) \alpha}^2=N_{\alpha, \beta+k \alpha}^2+\alpha \cdot(\beta+k \alpha)$
gilmore-lie-groups_ E08535 (9.42)	164	■ 1.000	$N_{\{\alpha, \beta+(k-1) \alpha\}}^2=(n-k+1)\left(\alpha \cdot \beta+\frac{1}{2}(n+k) \alpha \cdot \alpha\right)$	$N_{\alpha, \beta+(k-1) \alpha}^2=(n-k+1)\left(\alpha \cdot \beta+\frac{1}{2}(n+k) \alpha \cdot \alpha\right)$	$N_{\alpha, \beta+(k-1) \alpha}^2=(n-k+1)\left(\alpha \cdot \beta+\frac{1}{2}(n+k) \alpha \cdot \alpha\right)$
gilmore-lie-groups_ E08536 (9.43)	164	■ 1.000	$N_{\{\alpha, \beta-(m+1) \alpha\}}^2=(n+m+1)\left(\alpha \cdot \beta+\frac{1}{2}(n-m) \alpha \cdot \alpha\right)=0$	$N_{\alpha, \beta-(m+1) \alpha}^2=(n+m+1)\left(\alpha \cdot \beta+\frac{1}{2}(n-m) \alpha \cdot \alpha\right)=0$	$N_{\alpha, \beta-(m+1) \alpha}^2=(n+m+1)\left(\alpha \cdot \beta+\frac{1}{2}(n-m) \alpha \cdot \alpha\right)=0$
gilmore-lie-groups_ E08537 (9.44)	164	■ 1.000	$-n \leq \frac{2 \alpha \cdot \beta}{\alpha \cdot \alpha}=-n+m \leq m$	$-n \leq \frac{2 \alpha \cdot \beta}{\alpha \cdot \alpha}=-n+m \leq m$	$-n \leq \frac{2 \alpha \cdot \beta}{\alpha \cdot \alpha}=-n+m \leq m$
gilmore-lie-groups_ E08538 (9.45)	164	■ 1.000	$\beta^{\prime}=\beta-2 \frac{\beta \cdot \alpha}{\alpha \cdot \alpha} \alpha$	$\beta' = \beta - 2 \frac{\beta \cdot \alpha}{\alpha \cdot \alpha} \alpha$	$\beta' = \beta - 2 \frac{\beta \cdot \alpha}{\alpha \cdot \alpha} \alpha$
gilmore-lie-groups_ E08539 (9.46)	165	■ 1.000	$\begin{aligned} & \left\{\left[H_{\{i\}}, H_{\{j\}}\right]\right\} \&=0 \& \& \backslash \\ & \backslash\left\{\left[\mathbf{H}, E_{\{\alpha\}}\right]\right\} \&=\alpha E_{\{\alpha\}} \& \& \backslash \\ & \backslash\left\{\left[E_{\{\alpha\}}, E_{\{\beta\}}\right]\right\} \& \\ & =\alpha \cdot \mathbf{H} \& \& \alpha+\beta=0 \& \& =N_{\{\alpha, \beta\}} E_{\{\alpha+\beta\}} \& \& \alpha+\beta \neq 0, \text { a root } \\ & \backslash\&=0 \& \& \alpha+\beta \text { not a root } \end{aligned}$	$\begin{aligned} & \left[H_i, H_j\right]=0 \\ & \left[\mathbf{H}, E_{\alpha}\right]=\alpha E_{\alpha} \\ & \left[E_{\alpha}, E_{\beta}\right]=\alpha \cdot \mathbf{H} \quad \alpha+\beta=0 \\ & \quad =N_{\alpha, \beta} E_{\alpha+\beta} \quad \alpha+\beta \neq 0, \text { a root } \\ & \quad =0 \quad \alpha+\beta \text { not a root } \end{aligned}$	$\begin{aligned} & \left[H_i, H_j\right]=0 \\ & \left[\mathbf{H}, E_{\alpha}\right]=\alpha E_{\alpha} \\ & \left[E_{\alpha}, E_{\beta}\right]=\alpha \cdot \mathbf{H} \quad \alpha+\beta=0 \\ & \quad =N_{\alpha, \beta} E_{\alpha+\beta} \quad \alpha+\beta \neq 0, \text { a root } \\ & \quad =0 \quad \alpha+\beta \text { not a root } \end{aligned}$
gilmore-lie-groups_ E08541 (9.48)	166	■ 1.000	$\left(\mathbf{e}_i, \mathbf{e}_j\right)=\frac{1}{12} \delta_{i j}$	$(\mathbf{e}_i, \mathbf{e}_j) = \frac{1}{12} \delta_{ij}$	$(\mathbf{e}_i, \mathbf{e}_j) = \frac{1}{12} \delta_{ij}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08542	169	■ 1.000	$\begin{array}{cccc} b_{\{i\}}^{\dagger} b_{\{j\}} & b_{\{i\}}^{\dagger} b_{\{j\}} + \frac{1}{2} \delta_{ij} & b_{\{i\}}^{\dagger} b_{\{j\}}^{\dagger} & b_{\{i\}}^{\dagger} b_{\{j\}} \\ b_{\{j\}}^{\dagger} b_{\{i\}} + \frac{1}{2} \delta_{ij} & b_{\{j\}}^{\dagger} b_{\{i\}}^{\dagger} & b_{\{j\}}^{\dagger} b_{\{i\}} & b_{\{j\}}^{\dagger} b_{\{i\}} \\ b_{\{i\}}^{\dagger} b_{\{j\}}^{\dagger} & b_{\{i\}}^{\dagger} b_{\{j\}} & b_{\{i\}}^{\dagger} b_{\{j\}} & b_{\{i\}}^{\dagger} b_{\{j\}} \\ b_{\{j\}}^{\dagger} b_{\{i\}} & b_{\{j\}}^{\dagger} b_{\{i\}} & b_{\{j\}}^{\dagger} b_{\{i\}} & b_{\{j\}}^{\dagger} b_{\{i\}} \end{array}$	$\begin{array}{cccc} b_i^{\dagger} b_j & b_i^{\dagger} b_j + \frac{1}{2} \delta_{ij} & b_i^{\dagger} b_j^{\dagger} & b_i^{\dagger} b_j \\ f_i^{\dagger} f_j & f_i^{\dagger} f_j + \frac{1}{2} \delta_{ij} & f_i^{\dagger} f_j^{\dagger} & f_i^{\dagger} f_j \\ x^i \partial_j & x^i \partial_j + \frac{1}{2} \delta_{ij} & x^i x^j & \partial_i \partial_j \end{array}$	$\begin{array}{cccc} b_i^{\dagger} b_j & b_i^{\dagger} b_j + \frac{1}{2} \delta_{ij} & b_i^{\dagger} b_j^{\dagger} & b_i^{\dagger} b_j \\ f_i^{\dagger} f_j & f_i^{\dagger} f_j + \frac{1}{2} \delta_{ij} & f_i^{\dagger} f_j^{\dagger} & f_i^{\dagger} f_j \\ x^i \partial_j & x^i \partial_j + \frac{1}{2} \delta_{ij} & x^i x^j & \partial_i \partial_j \end{array}$
gilmore-lie-groups_ E08547	170	■ 1.000	$\frac{f_r(\Gamma)}{f_r(\text{def})} = \frac{\text{tr}(\Gamma(A))^r}{\text{tr}(\text{def}(A))^r} = \frac{\dim(\Gamma) \mathcal{C}^r(\Gamma)}{\dim(\text{def}) \mathcal{C}^r(\text{def})}$	$\frac{f_r(\Gamma)}{f_r(\text{def})} = \frac{\text{tr}(\Gamma(A))^r}{\text{tr}(\text{def}(A))^r} = \frac{\dim(\Gamma) \mathcal{C}^r(\Gamma)}{\dim(\text{def}) \mathcal{C}^r(\text{def})}$	$\frac{f_r(\Gamma)}{f_r(\text{def})} = \frac{\text{tr}(\Gamma(A))^r}{\text{tr}(\text{def}(A))^r} = \frac{\dim(\Gamma) \mathcal{C}^r(\Gamma)}{\dim(\text{def}) \mathcal{C}^r(\text{def})}$
gilmore-lie-groups_ E08548	170	■ 1.000	$f_{\{2\}}(j) = \frac{1}{6} \{(2j)(2j+1)(2j+2)\} f_{\{2\}} \left(j = \frac{1}{2} \right)$	$f_2(j) = \frac{1}{6} \{(2j)(2j+1)(2j+2)\} f_2 \left(j = \frac{1}{2} \right)$	$f_2(j) = \frac{1}{6} \{(2j)(2j+1)(2j+2)\} f_2(j = \frac{1}{2})$
gilmore-lie-groups_ E08550	171	■ 1.000	$f(X) = f_{\{0\}} I_{\{n\}} + f_{\{1\}} X^{\{1\}} + f_{\{2\}} X^{\{2\}} + \dots + f_{\{n-1\}} X^{\{n-1\}} = \sum_{j=0}^{n-1} f_j X^j$	$f(X) = f_0 I_n + f_1 X^1 + f_2 X^2 + \dots + f_{n-1} X^{n-1} = \sum_{j=0}^{j=n-1} f_j X^j$	$f(X) = f_0 I_n + f_1 X^1 + f_2 X^2 + \dots + f_{n-1} X^{n-1} = \sum_{j=0}^{j=n-1} f_j X^j$
gilmore-lie-groups_ E08551	171	■ 1.000	$\left[\begin{array}{cccccc} 1 & \lambda_1 & \lambda_1^2 & \dots & \lambda_1^{n-1} \\ 1 & \lambda_2 & \lambda_2^2 & \dots & \lambda_2^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \lambda_n & \lambda_n^2 & \dots & \lambda_n^{n-1} \end{array} \right] \left[\begin{array}{c} f_0 \\ f_1 \\ \vdots \\ f_{n-1} \end{array} \right] = \left[\begin{array}{c} f(\lambda_1) \\ f(\lambda_2) \\ \vdots \\ f(\lambda_n) \end{array} \right]$	$\left[\begin{array}{ccccc} 1 & \lambda_1 & \lambda_1^2 & \dots & \lambda_1^{n-1} \\ 1 & \lambda_2 & \lambda_2^2 & \dots & \lambda_2^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \lambda_n & \lambda_n^2 & \dots & \lambda_n^{n-1} \end{array} \right] \left[\begin{array}{c} f_0 \\ f_1 \\ \vdots \\ f_{n-1} \end{array} \right] = \left[\begin{array}{c} f(\lambda_1) \\ f(\lambda_2) \\ \vdots \\ f(\lambda_n) \end{array} \right]$	$\left[\begin{array}{ccccc} 1 & \lambda_1 & \lambda_1^2 & \dots & \lambda_1^{n-1} \\ 1 & \lambda_2 & \lambda_2^2 & \dots & \lambda_2^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \lambda_n & \lambda_n^2 & \dots & \lambda_n^{n-1} \end{array} \right] \left[\begin{array}{c} f_0 \\ f_1 \\ \vdots \\ f_{n-1} \end{array} \right] = \left[\begin{array}{c} f(\lambda_1) \\ f(\lambda_2) \\ \vdots \\ f(\lambda_n) \end{array} \right]$
gilmore-lie-groups_ E08552	172	■ 1.000	$\left[\begin{array}{cc} 1 & \frac{1}{2} \\ 1 & -\frac{1}{2} \end{array} \right] \left[\begin{array}{ccc} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 1 & -1 & 1 \end{array} \right] \left[\begin{array}{ccc} 1 & \frac{3}{2} & \left(\frac{3}{2}\right)^2 \\ 1 & \frac{1}{2} & \left(\frac{1}{2}\right)^2 \\ 1 & -\frac{1}{2} & \left(-\frac{1}{2}\right)^2 \\ 1 & -\frac{3}{2} & \left(-\frac{3}{2}\right)^2 \end{array} \right] \left[\begin{array}{ccc} \left(\frac{3}{2}\right)^3 & \left(\frac{1}{2}\right)^3 & \left(-\frac{1}{2}\right)^3 \\ \left(\frac{1}{2}\right)^3 & \left(-\frac{1}{2}\right)^3 & \left(-\frac{3}{2}\right)^3 \end{array} \right]$	$\left[\begin{array}{ccc} 1 & \frac{3}{2} & \left(\frac{3}{2}\right)^2 \\ 1 & \frac{1}{2} & \left(\frac{1}{2}\right)^2 \\ 1 & -\frac{1}{2} & \left(-\frac{1}{2}\right)^2 \\ 1 & -\frac{3}{2} & \left(-\frac{3}{2}\right)^2 \end{array} \right] \left[\begin{array}{ccc} \left(\frac{3}{2}\right)^3 & \left(\frac{1}{2}\right)^3 & \left(-\frac{1}{2}\right)^3 \\ \left(\frac{1}{2}\right)^3 & \left(-\frac{1}{2}\right)^3 & \left(-\frac{3}{2}\right)^3 \end{array} \right]$	$\left[\begin{array}{ccc} 1 & \frac{3}{2} & \left(\frac{3}{2}\right)^2 \\ 1 & \frac{1}{2} & \left(\frac{1}{2}\right)^2 \\ 1 & -\frac{1}{2} & \left(-\frac{1}{2}\right)^2 \\ 1 & -\frac{3}{2} & \left(-\frac{3}{2}\right)^2 \end{array} \right] \left[\begin{array}{ccc} \left(\frac{3}{2}\right)^3 & \left(\frac{1}{2}\right)^3 & \left(-\frac{1}{2}\right)^3 \\ \left(\frac{1}{2}\right)^3 & \left(-\frac{1}{2}\right)^3 & \left(-\frac{3}{2}\right)^3 \end{array} \right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0553	172	■ 1.000	$\left[\begin{array}{ccccc} 1 & 2 & 4 & 8 & 16 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 1 & -1 & 1 & -1 & 1 \\ 1 & -2 & 4 & -8 & 16 \end{array}\right] \left[\begin{array}{c} f_0 \\ (i\phi)^1 f_1 \\ (i\phi)^2 f_2 \\ (i\phi)^3 f_3 \\ (i\phi)^4 f_4 \end{array}\right] = \left[\begin{array}{c} e^{2i\phi} \\ e^{i\phi} \\ 1 \\ e^{-i\phi} \\ e^{-2i\phi} \end{array}\right]$	$\left[\begin{array}{ccccc} 1 & 2 & 4 & 8 & 16 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 1 & -1 & 1 & -1 & 1 \\ 1 & -2 & 4 & -8 & 16 \end{array}\right] \left[\begin{array}{c} f_0 \\ (i\phi)^1 f_1 \\ (i\phi)^2 f_2 \\ (i\phi)^3 f_3 \\ (i\phi)^4 f_4 \end{array}\right] = \left[\begin{array}{c} e^{2i\phi} \\ e^{i\phi} \\ 1 \\ e^{-i\phi} \\ e^{-2i\phi} \end{array}\right]$	$\left[\begin{array}{ccccc} 1 & 2 & 4 & 8 & 16 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 1 & -1 & 1 & -1 & 1 \\ 1 & -2 & 4 & -8 & 16 \end{array}\right] \left[\begin{array}{c} f_0 \\ (i\phi)^1 f_1 \\ (i\phi)^2 f_2 \\ (i\phi)^3 f_3 \\ (i\phi)^4 f_4 \end{array}\right] = \left[\begin{array}{c} e^{2i\phi} \\ e^{i\phi} \\ 1 \\ e^{-i\phi} \\ e^{-2i\phi} \end{array}\right]$
gilmore-lie-groups_ EQ0554	172	■ 1.000	$\begin{array}{ll} j=\frac{1}{2} & e^X = \cos\left(\frac{\phi}{2}\right) I_2 + \frac{\sin(\phi/2)}{\phi/2} X \\ l=1 & e^X = I_3 + \frac{\sin(\phi)}{\phi} X + \frac{1-\cos(\phi)}{\phi^2} X^2 \end{array}$	$j = \frac{1}{2} \quad e^X = \cos\left(\frac{\phi}{2}\right) I_2 + \frac{\sin(\phi/2)}{\phi/2} X$ $l = 1 \quad e^X = I_3 + \frac{\sin(\phi)}{\phi} X + \frac{1-\cos(\phi)}{\phi^2} X^2$	$j = \frac{1}{2} \quad e^X = \cos\left(\frac{\phi}{2}\right) I_2 + \frac{\sin(\phi/2)}{\phi/2} X$ $l = 1 \quad e^X = I_3 + \frac{\sin(\phi)}{\phi} X + \frac{1-\cos(\phi)}{\phi^2} X^2$
gilmore-lie-groups_ EQ0555	172	■ 1.000	$\operatorname{tr} X^2 = -\frac{j(j+1)(2j+1)}{3} \phi^2$	$\operatorname{tr} X^2 = -\frac{j(j+1)(2j+1)}{3} \phi^2$	$\operatorname{tr} X^2 = -\frac{j(j+1)(2j+1)}{3} \phi^2$
gilmore-lie-groups_ EQ0557 (10.2)	174	■ 1.000	$N_{\alpha, \beta+k\alpha} = (n-k)(m+k+1)(\alpha \cdot \alpha)/2$	$N_{\alpha, \beta+k\alpha} = (n-k)(m+k+1)(\alpha \cdot \alpha)/2$	$N_{\alpha, \beta+k\alpha} = (n-k)(m+k+1)(\alpha \cdot \alpha)/2$
gilmore-lie-groups_ EQ0559 (10.4)	174	■ 1.000	$\frac{2(\alpha \cdot \beta)}{\alpha \cdot \alpha}$ is an integer $\beta' = \beta - \frac{2(\alpha \cdot \beta)}{\alpha \cdot \alpha} \alpha$ is a root	$\frac{2(\alpha \cdot \beta)}{\alpha \cdot \alpha}$ is an integer $\beta' = \beta - \frac{2(\alpha \cdot \beta)}{\alpha \cdot \alpha} \alpha$ is a root	$\frac{2(\alpha \cdot \beta)}{\alpha \cdot \alpha}$ is an integer $\beta' = \beta - \frac{2(\alpha \cdot \beta)}{\alpha \cdot \alpha} \alpha$ is a root
gilmore-lie-groups_ EQ0560 (10.5)	174	■ 1.000	$0 \leq \cos^2(\alpha, \beta) = \left(\frac{\alpha \cdot \beta}{\alpha \cdot \alpha}\right) \left(\frac{\alpha \cdot \beta}{\beta \cdot \beta}\right) = \frac{n}{2} \frac{n'}{2} \leq 1$	$0 \leq \cos^2(\alpha, \beta) = \left(\frac{\alpha \cdot \beta}{\alpha \cdot \alpha}\right) \left(\frac{\alpha \cdot \beta}{\beta \cdot \beta}\right) = \frac{n}{2} \frac{n'}{2} \leq 1$	$0 \leq \cos^2(\alpha, \beta) = \left(\frac{\alpha \cdot \beta}{\alpha \cdot \alpha}\right) \left(\frac{\alpha \cdot \beta}{\beta \cdot \beta}\right) = \frac{n}{2} \frac{n'}{2} \leq 1$
gilmore-lie-groups_ EQ0561 (10.6)	175	■ 1.000	$\begin{array}{llll} A_{l-1} & +\mathbf{e}_i - \mathbf{e}_j & 1 \leq i \neq j \leq l & l-1 \geq 1 \\ D_l & \pm \mathbf{e}_i \pm \mathbf{e}_j & 1 \leq i \neq j \leq l & l > 3 \\ B_l & \pm \mathbf{e}_i \pm \mathbf{e}_j, \pm \mathbf{e}_i & 1 \leq i \neq j \leq l & l > 2 \\ C_l & \pm \mathbf{e}_i \pm \mathbf{e}_j, \pm 2\mathbf{e}_i & 1 \leq i \neq j \leq l & l > 1 \end{array}$	$\begin{array}{llll} A_{l-1} & +\mathbf{e}_i - \mathbf{e}_j & 1 \leq i \neq j \leq l & l-1 \geq 1 \\ D_l & \pm \mathbf{e}_i \pm \mathbf{e}_j & 1 \leq i \neq j \leq l & l > 3 \\ B_l & \pm \mathbf{e}_i \pm \mathbf{e}_j, \pm \mathbf{e}_i & 1 \leq i \neq j \leq l & l > 2 \\ C_l & \pm \mathbf{e}_i \pm \mathbf{e}_j, \pm 2\mathbf{e}_i & 1 \leq i \neq j \leq l & l > 1 \end{array}$	$\begin{array}{llll} A_{l-1} & +\mathbf{e}_i - \mathbf{e}_j & 1 \leq i \neq j \leq l & l-1 \geq 1 \\ D_l & \pm \mathbf{e}_i \pm \mathbf{e}_j & 1 \leq i \neq j \leq l & l > 3 \\ B_l & \pm \mathbf{e}_i \pm \mathbf{e}_j, \pm \mathbf{e}_i & 1 \leq i \neq j \leq l & l > 2 \\ C_l & \pm \mathbf{e}_i \pm \mathbf{e}_j, \pm 2\mathbf{e}_i & 1 \leq i \neq j \leq l & l > 1 \end{array}$
gilmore-lie-groups_ EQ0562 (10.7)	175	■ 1.000	$A_1 = B_1 = C_1$ $B_2 = C_2$ $A_3 = D_3$	$A_1 = B_1 = C_1$ $B_2 = C_2$ $A_3 = D_3$	$A_1 = B_1 = C_1$ $B_2 = C_2$ $A_3 = D_3$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08563 (10.8)	175	■ 1.000	$D_{\{2\}}=A_{\{1\}}+A_{\{1\}}$	$D_2 = A_1 + A_1$	$D_2 = A_1 + A_1$
gilmore-lie-groups_ E08569 (10.10)	177	■ 1.000	$\begin{array}{l} A_{l-1} \quad SU(l), SL(l;\mathbb{R}), SU(p,q) \\ D_l \quad SO(2l), SO(p,q) \\ B_l \quad SO(2l+1), SO(p,q) \\ C_l \quad Sp(l), Sp(p,q) \end{array} \quad \begin{array}{l} p+q=l \\ p+q=2l \\ p+q=2l+1 \\ p+q=l \end{array}$	$\begin{array}{l} A_{l-1} \quad SU(l), SL(l;\mathbb{R}), SU(p,q) \\ D_l \quad SO(2l), SO(p,q) \\ B_l \quad SO(2l+1), SO(p,q) \\ C_l \quad Sp(l), Sp(p,q) \end{array} \quad \begin{array}{l} p+q=l \\ p+q=2l \\ p+q=2l+1 \\ p+q=l \end{array}$	$\begin{array}{l} A_{l-1} \quad SU(l), SL(l;\mathbb{R}), SU(p,q) \\ D_l \quad SO(2l), SO(p,q) \\ B_l \quad SO(2l+1), SO(p,q) \\ C_l \quad Sp(l), Sp(p,q) \end{array} \quad \begin{array}{l} p+q=l \\ p+q=2l \\ p+q=2l+1 \\ p+q=l \end{array}$
gilmore-lie-groups_ E08570 (10.12)	180	■ 1.000	$\left(\alpha_i, \alpha_j\right)=-\sqrt{n_{ij} / 4}$	$\left(\alpha_i, \alpha_j\right)=-\sqrt{n_{ij} / 4}$	$\left(\alpha_i, \alpha_j\right)=-\sqrt{n_{ij} / 4}$
gilmore-lie-groups_ E08571 (10.13)	180	■ 1.000	$\sum\left(\mathbf{u} \cdot \mathbf{v}\right)^2 \leq 1$	$\sum(\mathbf{u} \cdot \mathbf{v})^2 \leq 1$	$\sum(\mathbf{u} \cdot \mathbf{v})^2 \leq 1$
gilmore-lie-groups_ E08572 (10.14)	181	■ 1.000	$\left(\sum \mathbf{u}_i, \sum \mathbf{u}_j\right)=n+2 \sum \sum \mathbf{u}_i \cdot \mathbf{u}_j \geq 0$	$\left(\sum \mathbf{u}_i, \sum \mathbf{u}_j\right)=n+2 \sum \sum \mathbf{u}_i \cdot \mathbf{u}_j \geq 0$	$\left(\sum \mathbf{u}_i, \sum \mathbf{u}_j\right)=n+2 \sum \sum \mathbf{u}_i \cdot \mathbf{u}_j \geq 0$
gilmore-lie-groups_ E08573 (10.15)	181	■ 1.000	$\sum\left(\mathbf{u} \cdot \mathbf{v}\right)^2=\sum n_{i} / 4<1$	$\sum(\mathbf{u} \cdot \mathbf{v})^2=\sum n_i / 4 < 1$	$\sum(\mathbf{u} \cdot \mathbf{v})^2=\sum n_i / 4 < 1$
gilmore-lie-groups_ E08574 (10.16)	181	■ 1.000	$\mathbf{u}=\sum_{i=1}^p i \mathbf{u}_i \quad \mathbf{v}=\sum_{j=1}^q j \mathbf{v}_j$	$\mathbf{u}=\sum_{i=1}^p i \mathbf{u}_i \quad \mathbf{v}=\sum_{j=1}^q j \mathbf{v}_j$	$\mathbf{u}=\sum_{i=1}^p i \mathbf{u}_i \quad \mathbf{v}=\sum_{j=1}^q j \mathbf{v}_j$
gilmore-lie-groups_ E08575 (10.17)	182	■ 1.000	$\left(1+\frac{1}{p}\right)\left(1+\frac{1}{q}\right)>2$	$\left(1+\frac{1}{p}\right)\left(1+\frac{1}{q}\right)>2$	$\left(1+\frac{1}{p}\right)\left(1+\frac{1}{q}\right)>2$
gilmore-lie-groups_ E08577 (10.19)	182	■ 1.000	$\frac{1}{p}+\frac{1}{q}+\frac{1}{r}>1$	$\frac{1}{p}+\frac{1}{q}+\frac{1}{r}>1$	$\frac{1}{p}+\frac{1}{q}+\frac{1}{r}>1$
gilmore-lie-groups_ E08579	185	■ 1.000	$\begin{aligned} & \left[\mathbf{L}, \mathbf{L}\right]=\mathbf{L}\left[L_{ij}, L_{rs}\right]=+\delta_{jr} L_{is}+\delta_{is} L_{jr}-\delta_{ir} L_{js}-\delta_{js} L_{ir} \\ & \left[\mathbf{L}, \mathbf{Q}\right]=\mathbf{Q}\left[L_{ij}, Q_{rs}\right]=+\delta_{jr} Q_{is}-\delta_{is} Q_{jr}-\delta_{ir} Q_{js}+\delta_{js} Q_{ir} \\ & \left[\mathbf{Q}, \mathbf{Q}\right]=\mathbf{L}\left[Q_{ij}, Q_{rs}\right]=+\delta_{jr} L_{is}+\delta_{is} L_{jr}+\delta_{ir} L_{js}+\delta_{js} L_{ir} \end{aligned}$	$\begin{aligned} & \left[\mathbf{L}, \mathbf{L}\right]=\mathbf{L}\left[L_{ij}, L_{rs}\right]=+\delta_{jr} L_{is}+\delta_{is} L_{jr}-\delta_{ir} L_{js}-\delta_{js} L_{ir} \\ & \left[\mathbf{L}, \mathbf{Q}\right]=\mathbf{Q}\left[L_{ij}, Q_{rs}\right]=+\delta_{jr} Q_{is}-\delta_{is} Q_{jr}-\delta_{ir} Q_{js}+\delta_{js} Q_{ir} \\ & \left[\mathbf{Q}, \mathbf{Q}\right]=\mathbf{L}\left[Q_{ij}, Q_{rs}\right]=+\delta_{jr} L_{is}+\delta_{is} L_{jr}+\delta_{ir} L_{js}+\delta_{js} L_{ir} \end{aligned}$	$\begin{aligned} & \left[\mathbf{L}, \mathbf{L}\right]=\mathbf{L}\left[L_{ij}, L_{rs}\right]=+\delta_{jr} L_{is}+\delta_{is} L_{jr}-\delta_{ir} L_{js}-\delta_{js} L_{ir} \\ & \left[\mathbf{L}, \mathbf{Q}\right]=\mathbf{Q}\left[L_{ij}, Q_{rs}\right]=+\delta_{jr} Q_{is}-\delta_{is} Q_{jr}-\delta_{ir} Q_{js}+\delta_{js} Q_{ir} \\ & \left[\mathbf{Q}, \mathbf{Q}\right]=\mathbf{L}\left[Q_{ij}, Q_{rs}\right]=+\delta_{jr} L_{is}+\delta_{is} L_{jr}+\delta_{ir} L_{js}+\delta_{js} L_{ir} \end{aligned}$
gilmore-lie-groups_ E08580	185	■ 1.000	$\hat{N}=\sum_{i=1}^3 a_i^{\dagger} a_i \quad \hat{Q}_{ij}=Q_{ij}-\frac{2}{3} \hat{N} \delta_{ij}$	$\hat{N}=\sum_{i=1}^3 a_i^{\dagger} a_i \quad \hat{Q}_{ij}=Q_{ij}-\frac{2}{3} \hat{N} \delta_{ij}$	$\hat{N}=\sum_{i=1}^3 a_i^{\dagger} a_i \quad \hat{Q}_{ij}=Q_{ij}-\frac{2}{3} \hat{N} \delta_{ij}$
gilmore-lie-groups_ E08585 (11.5)	189	■ 1.000	$i h^i H_i+i e^{\alpha} \frac{1}{\sqrt{2}}\left(E_{\alpha}+E_{-\alpha}\right)+e^{-\alpha} \frac{1}{\sqrt{2}}\left(E_{\alpha}-E_{-\alpha}\right)$	$i h^i H_i+i e^{\alpha} \frac{1}{\sqrt{2}}\left(E_{\alpha}+E_{-\alpha}\right)+e^{-\alpha} \frac{1}{\sqrt{2}}\left(E_{\alpha}-E_{-\alpha}\right)$	$i h^i H_i+i e^{\alpha} \frac{1}{\sqrt{2}}\left(E_{\alpha}+E_{-\alpha}\right)+e^{-\alpha} \frac{1}{\sqrt{2}}\left(E_{\alpha}-E_{-\alpha}\right)$
gilmore-lie-groups_ E08588 (11.9)	190	■ 1.000	$\text { - dimension } \leq \chi=\text { character } \leq+\text { rank }$	$\text { - dimension } \leq \chi=\text { character } \leq+\text { rank }$	$\text { - dimension } \leq \chi=\text { character } \leq+\text { rank }$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E09592 (11.13)	191	■ 1.000	$\chi\!\left(\!\!\mathfrak{g}^{\!\!\prime}\!\!\right)\!=\!\operatorname{dim}\!\left(\!\!\mathfrak{p}^{\!\!\prime}\!\!\right)\!-\!\operatorname{dim}\!\left(\!\!\mathfrak{h}\!\!\right)\!=\!\operatorname{dim}\!\left(\!\!\mathfrak{g}\!\!\right)\!-\!2\operatorname{dim}\!\left(\!\!\mathfrak{h}\!\!\right)\!=\!2\operatorname{dim}\!\left(\!\!\mathfrak{p}^{\!\!\prime}\!\!\right)\!-\!\operatorname{dim}\!\left(\!\!\mathfrak{g}\!\!\right)$	$\chi(\mathfrak{g}') = \dim(\mathfrak{p}') - \dim(\mathfrak{h}) = \dim(\mathfrak{g}) - 2 \dim(\mathfrak{h}) = 2 \dim(\mathfrak{p}') - \dim(\mathfrak{g})$	$\chi(\mathfrak{g}') = \dim(\mathfrak{p}') - \dim(\mathfrak{h}) = \dim(\mathfrak{g}) - 2 \dim(\mathfrak{h}) = 2 \dim(\mathfrak{p}') - \dim(\mathfrak{g})$
gilmore-lie-groups_ E09593 (11.14)	191	■ 1.000	$\mathfrak{su}(2) \rightarrow \mathfrak{su}(1,1) : \quad \frac{i}{2} \begin{bmatrix} a_3 & 0 \\ 0 & -a_3 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 0 & a_1 - ia_2 \\ a_1 + ia_2 & 0 \end{bmatrix}$	$\mathfrak{su}(2) \rightarrow \mathfrak{su}(1,1) : \quad \frac{i}{2} \begin{bmatrix} a_3 & 0 \\ 0 & -a_3 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 0 & a_1 - ia_2 \\ a_1 + ia_2 & 0 \end{bmatrix}$	$\mathfrak{su}(2) \rightarrow \mathfrak{su}(1,1) : \quad \frac{i}{2} \begin{bmatrix} a_3 & 0 \\ 0 & -a_3 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 0 & a_1 - ia_2 \\ a_1 + ia_2 & 0 \end{bmatrix}$
gilmore-lie-groups_ E09594 (11.15)	191	■ 1.000	$T^2=I$	$T^2 = I$	$T^2 = I$
gilmore-lie-groups_ E09596 (11.17)	191	■ 1.000	$(\mathfrak{h},\mathfrak{p})=(T^2\mathfrak{h},\mathfrak{p})=(T\mathfrak{h},T\mathfrak{p})=(\mathfrak{h},-\mathfrak{p})=- (\mathfrak{h},\mathfrak{p})=0$	$(\mathfrak{h},\mathfrak{p}) = (T^2\mathfrak{h},\mathfrak{p}) = (T\mathfrak{h},T\mathfrak{p}) = (\mathfrak{h},-\mathfrak{p}) = -(\mathfrak{h},\mathfrak{p}) = 0$	$(\mathfrak{h},\mathfrak{p}) = (T^2\mathfrak{h},\mathfrak{p}) = (T\mathfrak{h},T\mathfrak{p}) = (\mathfrak{h},-\mathfrak{p}) = -(\mathfrak{h},\mathfrak{p}) = 0$
gilmore-lie-groups_ E09597 (11.18)	192	■ 1.000	$\mathfrak{u}(n;\mathbb{F}) \quad \begin{bmatrix} A_p & 0 \\ 0 & A_q \end{bmatrix} + \begin{bmatrix} 0 & B \\ -B^\dagger & 0 \end{bmatrix}$	$\mathfrak{u}(n;\mathbb{F}) \quad \begin{bmatrix} A_p & 0 \\ 0 & A_q \end{bmatrix} + \begin{bmatrix} 0 & B \\ -B^\dagger & 0 \end{bmatrix}$	$\mathfrak{u}(n;\mathbb{F}) \quad \begin{bmatrix} A_p & 0 \\ 0 & A_q \end{bmatrix} + \begin{bmatrix} 0 & B \\ -B^\dagger & 0 \end{bmatrix}$
gilmore-lie-groups_ E09598 (11.19)	192	■ 1.000	$\mathfrak{u}(p,q;\mathbb{F}) \quad \begin{bmatrix} A_p & 0 \\ 0 & A_q \end{bmatrix} + \begin{bmatrix} 0 & B \\ +B^\dagger & 0 \end{bmatrix}$	$\mathfrak{u}(p,q;\mathbb{F}) \quad \begin{bmatrix} A_p & 0 \\ 0 & A_q \end{bmatrix} + \begin{bmatrix} 0 & B \\ +B^\dagger & 0 \end{bmatrix}$	$\mathfrak{u}(p,q;\mathbb{F}) \quad \begin{bmatrix} A_p & 0 \\ 0 & A_q \end{bmatrix} + \begin{bmatrix} 0 & B \\ +B^\dagger & 0 \end{bmatrix}$
gilmore-lie-groups_ E09602 (11.23)	193	■ 1.000	$\begin{aligned} &\alpha+i\beta \longrightarrow \begin{bmatrix} \alpha & \beta \\ -\beta & \alpha \end{bmatrix} \\ &\alpha+\mathcal{I}\beta+\mathcal{J}\gamma+\mathcal{K}\delta \longrightarrow \begin{bmatrix} \alpha+i\delta & i\beta+\gamma \\ i\beta-\gamma & \alpha-i\delta \end{bmatrix} \end{aligned}$	$\begin{aligned} &\alpha+i\beta \longrightarrow \begin{bmatrix} \alpha & \beta \\ -\beta & \alpha \end{bmatrix} \\ &\alpha+\mathcal{I}\beta+\mathcal{J}\gamma+\mathcal{K}\delta \longrightarrow \begin{bmatrix} \alpha+i\delta & i\beta+\gamma \\ i\beta-\gamma & \alpha-i\delta \end{bmatrix} \end{aligned}$	$\begin{aligned} &\alpha+i\beta \longrightarrow \begin{bmatrix} \alpha & \beta \\ -\beta & \alpha \end{bmatrix} \\ &\alpha+\mathcal{I}\beta+\mathcal{J}\gamma+\mathcal{K}\delta \longrightarrow \begin{bmatrix} \alpha+i\delta & i\beta+\gamma \\ i\beta-\gamma & \alpha-i\delta \end{bmatrix} \end{aligned}$
gilmore-lie-groups_ E09610	197	■ 1.000	$w\,a_1^{\dagger}a_1+x\,a_1^{\dagger}a_2+y\,a_2^{\dagger}a_1+z\,a_2^{\dagger}a_2=\alpha\left(a_1^{\dagger}a_1+a_2^{\dagger}a_2\right)$	$wa_1^\dagger a_1 + xa_1^\dagger a_2 + ya_2^\dagger a_1 + za_2^\dagger a_2 = \alpha \left(a_1^\dagger a_1 + a_2^\dagger a_2 \right)$	$wa_1^\dagger a_1 + xa_1^\dagger a_2 + ya_2^\dagger a_1 + za_2^\dagger a_2 = \alpha (a_1^\dagger a_1 + a_2^\dagger a_2)$
gilmore-lie-groups_ E09611	197	■ 1.000	$+\beta\left(a_1^{\dagger}a_1-a_2^{\dagger}a_2\right)+\gamma\left(a_1^{\dagger}a_2+a_2^{\dagger}a_1\right)+\delta\left(a_1^{\dagger}a_2-a_2^{\dagger}a_1\right)$	$+\beta \left(a_1^\dagger a_1 - a_2^\dagger a_2 \right) + \gamma \left(a_1^\dagger a_2 + a_2^\dagger a_1 \right) + \delta \left(a_1^\dagger a_2 - a_2^\dagger a_1 \right)$	$+\beta (a_1^\dagger a_1 - a_2^\dagger a_2) + \gamma (a_1^\dagger a_2 + a_2^\dagger a_1) + \delta (a_1^\dagger a_2 - a_2^\dagger a_1)$
gilmore-lie-groups_ E09613	198	■ 1.000	$\mathcal{C}^2=\sum H_i^2+\sum E_{\alpha}E_{-\alpha}+E_{-\alpha}E_{\alpha}$	$\mathcal{C}^2 = \sum H_i^2 + \sum E_\alpha E_{-\alpha} + E_{-\alpha} E_\alpha$	$\mathcal{C}^2 = \sum H_i^2 + \sum E_\alpha E_{-\alpha} + E_{-\alpha} E_\alpha$
gilmore-lie-groups_ E09614	198	■ 1.000	$H_i\left n_1,n_2,\ldots,n_r\right\rangle=n_i\left n_1,n_2,\ldots,n_r\right\rangle$	$H_i n_1,n_2,\ldots,n_r\rangle = n_i n_1,n_2,\ldots,n_r\rangle$	$H_i n_1,n_2,\ldots,n_r\rangle = n_i n_1,n_2,\ldots,n_r\rangle$
gilmore-lie-groups_ E09615	198	■ 1.000	$\mathcal{C}^2\mid\text{state}\rangle=-l(l+n-2)\mid\text{state}\rangle$	$\mathcal{C}^2 \mid \text{state} \rangle = -l(l+n-2) \mid \text{state} \rangle$	$\mathcal{C}^2 \text{state}\rangle = -l(l+n-2) \text{state}\rangle$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0659	214	■ 1.000	$d\,\mu=\frac{d\,x_{\{1\}}\,d\,x_{\{2\}}}{\sqrt{1+x_{\{1\}}^2+x_{\{2\}}^2}}$	$d\mu=\frac{dx_1dx_2}{\sqrt{1+x_1^2+x_2^2}}$	$d\mu=\frac{dx_1dx_2}{\sqrt{1+x_1^2+x_2^2}}$
gilmore-lie-groups_ EQ0665	215	■ 1.000	$\mathrm{cal}{C}_{\{2\}}[S\,0(n+1)\,/\,S\,0(n)]=\sum_{1\leq r<s}^nX_{r,s}^2(\phi)$	$\mathcal{C}_2[SO(n+1)/SO(n)]=\sum_{1\leq r<s}^{n+1}X_{r,s}^2(\phi)$	$\mathcal{C}_2[SO(n+1)/SO(n)]=\sum_{1\leq r<s}^{n+1}X_{r,s}^2(\phi)$
gilmore-lie-groups_ EQ0666	215	■ 1.000	$\Delta\left(S^{\{n\}}\right)=\partial_{\{n\}}\left(f_{\{1\}}(\phi)\right)\partial_{\{n\}}\left(f_{\{2\}}(\phi)\right)+\Delta\left(S^{\{n-1\}}\right)$	$\Delta\left(S^n\right)=\partial_n\left(f_1(\phi)\partial_n\right)+f_2(\phi)\Delta\left(S^{n-1}\right)$	$\Delta(S^n)=\partial_n(f_1(\phi)\partial_n)+f_2(\phi)\Delta(S^{n-1})$
gilmore-lie-groups_ EQ0669	215	■ 1.000	$E^{\{2\}}-(p\,c)^{\{2\}}=\left(m\,c^{\{2\}}\right)^{\{2\}}$	$E^2-(pc)^2=\left(mc^2\right)^2$	$E^2-(pc)^2=(mc^2)^2$
gilmore-lie-groups_ EQ0673	216	■ 1.000	$z^{\{\prime\}}=\frac{a\,c\left(x^{\{2\}}+y^{\{2\}}\right)+(a\,d+b\,c)\,x+b\,d+i\,y}{ c\,z+d ^{\{2\}}}$	$z'=\frac{ac\left(x^2+y^2\right)+(ad+bc)x+bd+iy}{ cz+d ^2}$	$z'=\frac{ac(x^2+y^2)+(ad+bc)x+bd+iy}{ cz+d ^2}$
gilmore-lie-groups_ EQ0677	217	■ 1.000	$d\,s^{\{2\}}=\left(\begin{array}{ll}d\,x&d\,y\end{array}\right)\left[\begin{array}{cc}\frac{1}{(1-\overline{w}w)^2}&0\\0&\frac{1}{(1-\overline{w}w)^2}\end{array}\right]\left(\begin{array}{l}dx\\dy\end{array}\right)=\frac{d\overline{w}dw}{(1-\overline{w}w)^2}$	$ds^2=\left(\begin{array}{ll}dx&dy\end{array}\right)\left[\begin{array}{cc}\frac{1}{(1-\overline{w}w)^2}&0\\0&\frac{1}{(1-\overline{w}w)^2}\end{array}\right]\left(\begin{array}{l}dx\\dy\end{array}\right)=\frac{d\overline{w}dw}{(1-\overline{w}w)^2}$	$ds^2=\left(\begin{array}{ll}dx&dy\end{array}\right)\left[\begin{array}{cc}\frac{1}{(1-\overline{w}w)^2}&0\\0&\frac{1}{(1-\overline{w}w)^2}\end{array}\right]\left(\begin{array}{l}dx\\dy\end{array}\right)=\frac{d\overline{w}dw}{(1-\overline{w}w)^2}$
gilmore-lie-groups_ EQ0678	217	■ 1.000	$d\,\mu=\frac{d\,x\,d\,y}{(1-\overline{w}w)^{\{2\}}}=\frac{d\overline{w}dw}{(1-\overline{w}w)^{\{2\}}}$	$d\mu=\frac{xdy}{(1-\overline{w}w)^2}=\frac{d\overline{w}dw}{(1-\overline{w}w)^2}$	$d\mu=\frac{dx\,dy}{(1-\overline{w}w)^2}=\frac{d\overline{w}dw}{(1-\overline{w}w)^2}$
gilmore-lie-groups_ EQ0679	217	■ 1.000	$s\left(w_{\{1\}},w_{\{2\}}\right)=\tanh^{-1}\left\{\frac{\left w_{\{1\}}-w_{\{2\}}\right }{\left 1-w_{\{1\}}\overline{w}_{\{2\}}\right }\right\}$	$s\left(w_1,w_2\right)=\tanh^{-1}\left\{\frac{\left w_1-w_2\right }{\left 1-w_1\overline{w}_2\right }\right\}$	$s(w_1,w_2)=\tanh^{-1}\left\{\frac{\left w_1-w_2\right }{\left 1-w_1\overline{w}_2\right }\right\}$
gilmore-lie-groups_ EQ0680	217	■ 1.000	$w=e^{i\,\phi}\frac{z-z_{\{0\}}}{z-\overline{z}_{\{0\}}}$	$w=e^{i\phi}\frac{z-z_0}{z-\overline{z}_0}$	$w=e^{i\phi}\frac{z-z_0}{z-\overline{z}_0}$
gilmore-lie-groups_ EQ0681	218	■ 1.000	$w=\frac{i\,z+1}{z+i}$	$w=\frac{iz+1}{z+i}$	$w=\frac{iz+1}{z+i}$
gilmore-lie-groups_ EQ0691 (13.9)	221	■ 1.000	$\theta_{\{1\}}L_{\{1\}}+\theta_{\{2\}}L_{\{2\}}+\theta_{\{3\}}L_{\{3\}}\longrightarrow(-d_{\{2\}})\left(\frac{L_{\{1\}}}{R}\right)+(+d_{\{1\}})\left(\frac{L_{\{2\}}}{R}\right)+\theta_{\{3\}}L_{\{3\}}$	$\theta_1L_1+\theta_2L_2+\theta_3L_3\longrightarrow(-d_2)\left(\frac{L_1}{R}\right)+(+d_1)\left(\frac{L_2}{R}\right)+\theta_3L_3$	$\theta_1L_1+\theta_2L_2+\theta_3L_3\longrightarrow(-d_2)\left(\frac{L_1}{R}\right)+(+d_1)\left(\frac{L_2}{R}\right)+\theta_3L_3$
gilmore-lie-groups_ EQ0694 (13.12)	222	■ 1.000	$\begin{aligned}&\mathrm{cal}{C}^{\{2\}}[S\,0(3)]\,\&\\&=L_{\{1\}}^{\{2\}}+L_{\{2\}}^{\{2\}}+L_{\{3\}}^{\{3\}}\,\&\mathrm{cal}{C}^{\{2\}}[I\,S\,0(2)]\,\&=\lim\left(1/R^{\{2\}}\right)\mathrm{cal}{C}^{\{2\}}[S\,0(3)]\,\&\&=\lim\left[\left(L_{\{1\}}/R\right)^{\{2\}}+\left(L_{\{2\}}/R\right)^{\{2\}}+\left(L_{\{3\}}/R\right)^{\{2\}}\right]\,\&\\&=\left(-P_{\{2\}}\right)^{\{2\}}+\left(+P_{\{1\}}\right)^{\{2\}}+\theta=\frac{\partial^{\{2\}}}{\partial y^{\{2\}}}+\frac{\partial^{\{2\}}}{\partial x^{\{2\}}}\end{aligned}$	$\begin{aligned}\mathcal{C}^2[SO(3)]&=L_1^2+L_2^2+L_3^3\\\mathcal{C}^2[ISO(2)]&=\lim\left(1/R^2\right)\mathcal{C}^2[SO(3)]\\&=\lim\left[\left(L_1/R\right)^2+\left(L_2/R\right)^2+\left(L_3/R\right)^2\right]\\&=\left(-P_2\right)^2+\left(+P_1\right)^2+0=\frac{\partial^2}{\partial y^2}+\frac{\partial^2}{\partial x^2}\end{aligned}$	$\begin{aligned}\mathcal{C}^2[SO(3)]&=L_1^2+L_2^2+L_3^3\\\mathcal{C}^2[ISO(2)]&=\lim(1/R^2)\mathcal{C}^2[SO(3)]\\&=\lim[(L_1/R)^2+(L_2/R)^2+(L_3/R)^2]\\&=\left(-P_2\right)^2+\left(+P_1\right)^2+0=\frac{\partial^2}{\partial y^2}+\frac{\partial^2}{\partial x^2}\end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0696 (13.14)	223	■ 1.000	$\begin{aligned} \left\{ \left[L_i, L_j \right] \right\} &= -\epsilon_{ijk} L_k \\ \left\{ \left[L_i, V_j \right] \right\} &= -\epsilon_{ijk} V_k \\ \left\{ \left[V_i, V_j \right] \right\} &= -\epsilon_{ijk} L_k \end{aligned}$	$\begin{aligned} [L_i, L_j] &= -\epsilon_{ijk} L_k \\ [L_i, V_j] &= -\epsilon_{ijk} V_k \\ [V_i, V_j] &= -\epsilon_{ijk} L_k \end{aligned}$	$\begin{aligned} [L_i, L_j] &= -\epsilon_{ijk} L_k \\ [L_i, V_j] &= -\epsilon_{ijk} V_k \\ [V_i, V_j] &= -\epsilon_{ijk} L_k \end{aligned}$
gilmore-lie-groups_ EQ0697 (13.15)	223	■ 1.000	$-P_i = \lim_{R \rightarrow \infty} \frac{1}{R} V_i = \lim_{R \rightarrow \infty} \frac{1}{R} (x_i \partial_4 - x_4 \partial_i) = -\partial_i$	$-P_i = \lim_{R \rightarrow \infty} \frac{1}{R} V_i = \lim_{R \rightarrow \infty} \frac{1}{R} (x_i \partial_4 - x_4 \partial_i) = -\partial_i$	$-P_i = \lim_{R \rightarrow \infty} \frac{1}{R} V_i = \lim_{R \rightarrow \infty} \frac{1}{R} (x_i \partial_4 - x_4 \partial_i) = -\partial_i$
gilmore-lie-groups_ EQ0698 (13.16)	223	■ 1.000	$\begin{aligned} \left\{ \left[L_i, L_j \right] \right\} &= -\epsilon_{ijk} L_k \\ \left\{ \left[L_i, P_j \right] \right\} &= -\epsilon_{ijk} P_k \\ \left\{ \left[P_i, P_j \right] \right\} &= 0 \end{aligned}$	$\begin{aligned} [L_i, L_j] &= -\epsilon_{ijk} L_k \\ [L_i, P_j] &= -\epsilon_{ijk} P_k \\ [P_i, P_j] &= 0 \end{aligned}$	$\begin{aligned} [L_i, L_j] &= -\epsilon_{ijk} L_k \\ [L_i, P_j] &= -\epsilon_{ijk} P_k \\ [P_i, P_j] &= 0 \end{aligned}$
gilmore-lie-groups_ EQ0699 (13.17)	223	■ 1.000	$\begin{aligned} \mathcal{C}_1^2[ISO(3)] &= \lim_{R \rightarrow \infty} (1/R^2) (\mathbf{L} \cdot \mathbf{L} + \mathbf{V} \cdot \mathbf{V}) \\ &= \lim_{R \rightarrow \infty} [(\mathbf{L}/R) \cdot (\mathbf{L}/R) + (\mathbf{V}/R) \cdot (\mathbf{V}/R)] \\ &= 0 + \mathbf{P} \cdot \mathbf{P} = \nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \end{aligned}$	$\begin{aligned} \mathcal{C}_1^2[ISO(3)] &= \lim_{R \rightarrow \infty} (1/R^2) (\mathbf{L} \cdot \mathbf{L} + \mathbf{V} \cdot \mathbf{V}) \\ &= \lim_{R \rightarrow \infty} [(\mathbf{L}/R) \cdot (\mathbf{L}/R) + (\mathbf{V}/R) \cdot (\mathbf{V}/R)] \\ &= 0 + \mathbf{P} \cdot \mathbf{P} = \nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \end{aligned}$	$\begin{aligned} \mathcal{C}_1^2[ISO(3)] &= \lim_{R \rightarrow \infty} (1/R^2) (\mathbf{L} \cdot \mathbf{L} + \mathbf{V} \cdot \mathbf{V}) \\ &= \lim_{R \rightarrow \infty} [(\mathbf{L}/R) \cdot (\mathbf{L}/R) + (\mathbf{V}/R) \cdot (\mathbf{V}/R)] \\ &= 0 + \mathbf{P} \cdot \mathbf{P} = \nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \end{aligned}$
gilmore-lie-groups_ EQ0700 (13.18)	223	■ 1.000	$\mathcal{C}_2^2[SO(4)] = \epsilon^{ijkl} X_{ij} X_{kl} \rightarrow 8 \mathbf{L} \cdot \mathbf{V}$	$\mathcal{C}_2^2[SO(4)] = \epsilon^{ijkl} X_{ij} X_{kl} \rightarrow 8 \mathbf{L} \cdot \mathbf{V}$	$\mathcal{C}_2^2[SO(4)] = \epsilon^{ijkl} X_{ij} X_{kl} \rightarrow 8 \mathbf{L} \cdot \mathbf{V}$
gilmore-lie-groups_ EQ0701 (13.19)	223	■ 1.000	$\begin{aligned} \mathcal{C}_2^2[ISO(3)]/8 &= \lim_{R \rightarrow \infty} (1/R) (\mathbf{L} \cdot \mathbf{V}) \\ &= \lim_{R \rightarrow \infty} [\mathbf{L} \cdot (\mathbf{V}/R)] = -\mathbf{L} \cdot \mathbf{P} \end{aligned}$	$\begin{aligned} \mathcal{C}_2^2[ISO(3)]/8 &= \lim_{R \rightarrow \infty} (1/R) (\mathbf{L} \cdot \mathbf{V}) \\ &= \lim_{R \rightarrow \infty} [\mathbf{L} \cdot (\mathbf{V}/R)] = -\mathbf{L} \cdot \mathbf{P} \end{aligned}$	$\begin{aligned} \mathcal{C}_2^2[ISO(3)]/8 &= \lim_{R \rightarrow \infty} (1/R) (\mathbf{L} \cdot \mathbf{V}) \\ &= \lim_{R \rightarrow \infty} [\mathbf{L} \cdot (\mathbf{V}/R)] = -\mathbf{L} \cdot \mathbf{P} \end{aligned}$
gilmore-lie-groups_ EQ0702 (13.20)	224	■ 1.000	$x^2 + y^2 + z^2 - (ct)^2$	$x^2 + y^2 + z^2 - (ct)^2$	$x^2 + y^2 + z^2 - (ct)^2$
gilmore-lie-groups_ EQ0703 (13.21)	224	■ 1.000	$(x - x')^2 + (y - y')^2 + (z - z')^2 - (ct - ct')^2$	$(x - x')^2 + (y - y')^2 + (z - z')^2 - (ct - ct')^2$	$(x - x')^2 + (y - y')^2 + (z - z')^2 - (ct - ct')^2$
gilmore-lie-groups_ EQ0707 (13.24)	224	■ 1.000	$\begin{aligned} \lim_{R \rightarrow \infty} (1/R) W^\alpha &\rightarrow \lim_{R \rightarrow \infty} \epsilon^{\alpha\beta\gamma\mu 5} X_{\beta\gamma} [(1/R) X_{\mu 5}] \\ &= \epsilon^{\alpha\beta\gamma\mu 5} X_{\beta\gamma} (\partial/\partial x^\mu) \end{aligned}$	$\begin{aligned} \lim_{R \rightarrow \infty} (1/R) W^\alpha &\rightarrow \lim_{R \rightarrow \infty} \epsilon^{\alpha\beta\gamma\mu 5} X_{\beta\gamma} [(1/R) X_{\mu 5}] \\ &= \epsilon^{\alpha\beta\gamma\mu 5} X_{\beta\gamma} (\partial/\partial x^\mu) \end{aligned}$	$\begin{aligned} \lim_{R \rightarrow \infty} (1/R) W^\alpha &\rightarrow \lim_{R \rightarrow \infty} \epsilon^{\alpha\beta\gamma\mu 5} X_{\beta\gamma} [(1/R) X_{\mu 5}] \\ &= \epsilon^{\alpha\beta\gamma\mu 5} X_{\beta\gamma} (\partial/\partial x^\mu) \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_E08708 (13.25)	225	1.000	$P_{\mu}=(0,0,0,\,m\,c)$	$P_{\mu}=(0,0,0,mc)$	$P_{\mu}=(0,0,0,mc)$
gilmore-lie-groups_E08709 (13.26)	225	1.000	$W^{\alpha}=\epsilon^{\alpha\beta\gamma\mu}\,X_{\beta\gamma}\,X_{\mu}\,L_{\alpha}\,m\,c$	$W^{\alpha}=\epsilon^{\alpha\beta\gamma\mu}X_{\beta\gamma}mc=L_{\alpha}mc$	$W^{\alpha}=\epsilon^{\alpha\beta\gamma\mu}X_{\beta\gamma}mc=L_{\alpha}mc$
gilmore-lie-groups_E08710 (13.27)	225	1.000	$W^{\alpha}\,W_{\alpha}=(\mathbf{L}\cdot\mathbf{L})(\mathbf{P}\cdot\mathbf{P})$	$W^{\alpha}W_{\alpha}=(\mathbf{L}\cdot\mathbf{L})(\mathbf{P}\cdot\mathbf{P})$	$W^{\alpha}W_{\alpha}=(\mathbf{L}\cdot\mathbf{L})(\mathbf{P}\cdot\mathbf{P})$
gilmore-lie-groups_E08711 (13.28)	225	1.000	$\begin{aligned} \left[J_3,J_{\pm}\right]&=\pm J_{\pm} \\ \left[J_+,J_{-}\right]&=2J_3 \\ \left[J_0,\mathbf{J}\right]&=0 \end{aligned}$	$\begin{aligned} \left[J_3,J_{\pm}\right]&=\pm J_{\pm} \\ \left[J_+,J_{-}\right]&=2J_3 \\ \left[J_0,\mathbf{J}\right]&=0 \end{aligned}$	$\begin{aligned} \left[J_3,J_{\pm}\right]&=\pm J_{\pm} \\ \left[J_+,J_{-}\right]&=2J_3 \\ \left[J_0,\mathbf{J}\right]&=0 \end{aligned}$
gilmore-lie-groups_E08713 (13.30)	226	1.000	$\begin{aligned} \left[h_3,h_{\pm}\right]&=\pm h_{\pm} \\ \left[h_+,h_{-}\right]&=2c^2h_3-h_0 \\ \left[h_0,\mathbf{h}\right]&=0 \end{aligned}$	$\begin{aligned} \left[h_3,h_{\pm}\right]&=\pm h_{\pm} \\ \left[h_+,h_{-}\right]&=2c^2h_3-h_0 \\ \left[h_0,\mathbf{h}\right]&=0 \end{aligned}$	$\begin{aligned} \left[h_3,h_{\pm}\right]&=\pm h_{\pm} \\ \left[h_+,h_{-}\right]&=2c^2h_3-h_0 \\ \left[h_0,\mathbf{h}\right]&=0 \end{aligned}$
gilmore-lie-groups_E08715 (13.32)	226	1.000	$\begin{aligned} \mathcal{C}^1&=J_0 \\ \mathcal{C}^2&=J_3^2+\frac{1}{2}(J_+J_{-}+J_{-}J_+) \end{aligned}$	$\begin{aligned} \mathcal{C}^1&=J_0 \\ \mathcal{C}^2&=J_3^2+\frac{1}{2}(J_+J_{-}+J_{-}J_+) \end{aligned}$	$\begin{aligned} \mathcal{C}^1&=J_0 \\ \mathcal{C}^2&=J_3^2+\frac{1}{2}(J_+J_{-}+J_{-}J_+) \end{aligned}$
gilmore-lie-groups_E08716 (13.33)	226	1.000	$\begin{aligned} \lim_{c\rightarrow 0}\mathcal{C}^2&=\lim_{c\rightarrow 0}\left(h_3-\frac{1}{2c^2}h_0\right)^2+\frac{1}{2}[(cJ_+)(cJ_{-})+(cJ_{-})(cJ_+)] \\ &=\lim_{c\rightarrow 0}c^2h_3^2-\frac{1}{2}(h_3h_0+h_0h_3)+c^2\left(-\frac{1}{2c^2}h_0\right)^2 \\ &\quad +\frac{1}{2}[(h_+)(h_{-})+(h_{-})(h_+)] \end{aligned}$	$\begin{aligned} \lim_{c\rightarrow 0}c^2\mathcal{C}^2&=\lim_{c\rightarrow 0}c^2\left(h_3-\frac{1}{2c^2}h_0\right)^2+\frac{1}{2}[(cJ_+)(cJ_{-})+(cJ_{-})(cJ_+)] \\ &=\lim_{c\rightarrow 0}c^2h_3^2-\frac{1}{2}(h_3h_0+h_0h_3)+c^2\left(-\frac{1}{2c^2}h_0\right)^2 \\ &\quad +\frac{1}{2}[(h_+)(h_{-})+(h_{-})(h_+)] \end{aligned}$	$\begin{aligned} \lim_{c\rightarrow 0}c^2\mathcal{C}^2&=\lim_{c\rightarrow 0}c^2\left(h_3-\frac{1}{2c^2}h_0\right)^2+\frac{1}{2}[(cJ_+)(cJ_{-})+(cJ_{-})(cJ_+)] \\ &=\lim_{c\rightarrow 0}c^2h_3^2-\frac{1}{2}(h_3h_0+h_0h_3)+c^2\left(-\frac{1}{2c^2}h_0\right)^2 \\ &\quad +\frac{1}{2}[(h_+)(h_{-})+(h_{-})(h_+)] \end{aligned}$
gilmore-lie-groups_E08717 (13.34)	226	1.000	$\begin{aligned} \lim_{c\rightarrow 0}\mathcal{C}^2[U(2)]&=\lim_{c\rightarrow 0}\left(h_0/2c\right)^2\rightarrow\mathcal{C}^2[H_4]=-\frac{1}{2}\left[\left(\hat{n}+\frac{1}{2}I\right)I+I\left(\hat{n}+\frac{1}{2}I\right)\right] \\ &\quad +\frac{1}{2}(aa^{\dagger}+a^{\dagger}a) \end{aligned}$	$\begin{aligned} \lim_{c\rightarrow 0}c^2\mathcal{C}^2[U(2)]&=\lim_{c\rightarrow 0}\left(h_0/2c\right)^2\rightarrow\mathcal{C}^2[H_4]=-\frac{1}{2}\left[\left(\hat{n}+\frac{1}{2}I\right)I+I\left(\hat{n}+\frac{1}{2}I\right)\right] \\ &\quad +\frac{1}{2}(aa^{\dagger}+a^{\dagger}a) \end{aligned}$	$\begin{aligned} \lim_{c\rightarrow 0}c^2\mathcal{C}^2[U(2)]&=\lim_{c\rightarrow 0}\left(h_0/2c\right)^2\rightarrow\mathcal{C}^2[H_4]=-\frac{1}{2}\left[\left(\hat{n}+\frac{1}{2}I\right)I+I\left(\hat{n}+\frac{1}{2}I\right)\right] \\ &\quad +\frac{1}{2}(aa^{\dagger}+a^{\dagger}a) \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0718 (13.35)	227	■ 1.000	$\begin{aligned} i \, \theta_{\mu} J_{\mu} &= \frac{1}{2} \theta e^{-i \phi} J_{+} - \frac{1}{2} \theta e^{+i \phi} J_{-} + i \theta_3 J_3 + i \theta_0 J_0 \\ \theta e^{-i \phi} J_{+} &= \frac{\theta}{2c} e^{-i \phi} h_{+} - \frac{\theta}{2c} e^{+i \phi} h_{-} + i \theta_3 h_3 + i \left(\theta_0 - \frac{\theta_3}{2c^2} \right) h_0 \end{aligned}$	$i \theta_{\mu} J_{\mu} = \frac{1}{2} \theta e^{-i \phi} J_{+} - \frac{1}{2} \theta e^{+i \phi} J_{-} + i \theta_3 J_3 + i \theta_0 J_0$ $= \frac{\theta}{2c} e^{-i \phi} h_{+} - \frac{\theta}{2c} e^{+i \phi} h_{-} + i \theta_3 h_3 + i \left(\theta_0 - \frac{\theta_3}{2c^2} \right) h_0$	$i \theta_{\mu} J_{\mu} = \frac{1}{2} \theta e^{-i \phi} J_{+} - \frac{1}{2} \theta e^{+i \phi} J_{-} + i \theta_3 J_3 + i \theta_0 J_0$ $= \frac{\theta}{2c} e^{-i \phi} h_{+} - \frac{\theta}{2c} e^{+i \phi} h_{-} + i \theta_3 h_3 + i \left(\theta_0 - \frac{\theta_3}{2c^2} \right) h_0$
gilmore-lie-groups_ EQ0719 (13.36)	227	■ 1.000	$\begin{aligned} &\lim_{c \rightarrow 0} \left(\theta e^{-i \phi} J_{+} - \frac{1}{2} \theta e^{+i \phi} J_{-} + i \theta_3 J_3 + i \theta_0 J_0 \right) \\ &= \lim_{c \rightarrow 0} \left(\frac{\theta}{2c} e^{-i \phi} h_{+} - \frac{\theta}{2c} e^{+i \phi} h_{-} + i \theta_3 h_3 + i \left(\theta_0 - \frac{\theta_3}{2c^2} \right) h_0 \right) \end{aligned}$	$\lim_{c \rightarrow 0} \left(\theta e^{-i \phi} J_{+} - \frac{1}{2} \theta e^{+i \phi} J_{-} + i \theta_3 J_3 + i \theta_0 J_0 \right)$ $= \lim_{c \rightarrow 0} \left(\frac{\theta}{2c} e^{-i \phi} h_{+} - \frac{\theta}{2c} e^{+i \phi} h_{-} + i \theta_3 h_3 + i \left(\theta_0 - \frac{\theta_3}{2c^2} \right) h_0 \right)$	$\lim_{c \rightarrow 0} \left(\theta e^{-i \phi} J_{+} - \frac{1}{2} \theta e^{+i \phi} J_{-} + i \theta_3 J_3 + i \theta_0 J_0 \right)$ $= \lim_{c \rightarrow 0} \left(\frac{\theta}{2c} e^{-i \phi} h_{+} - \frac{\theta}{2c} e^{+i \phi} h_{-} + i \theta_3 h_3 + i \left(\theta_0 - \frac{\theta_3}{2c^2} \right) h_0 \right)$
gilmore-lie-groups_ EQ0721 (13.38)	227	■ 1.000	$n = J + M$	$n = J + M$	$n = J + M$
gilmore-lie-groups_ EQ0722 (13.39)	227	■ 1.000	$\lim_{c \rightarrow 0} \left(M + \frac{1}{2c^2} \right) = \lim_{c \rightarrow 0} \left(n - J + \frac{1}{2c^2} \right)$	$\lim_{c \rightarrow 0} \left(M + \frac{1}{2c^2} \right) = \lim_{c \rightarrow 0} \left(n - J + \frac{1}{2c^2} \right)$	$\lim_{c \rightarrow 0} \left(M + \frac{1}{2c^2} \right) = \lim_{c \rightarrow 0} \left(n - J + \frac{1}{2c^2} \right)$
gilmore-lie-groups_ EQ0727 (13.46)	229	■ 1.000	$e^{\left(\zeta J_{+} - \zeta^{*} J_{-} \right)} = e^{\tau J_{+}} e^{\ln(1 + \tau^{*} \tau) J_3} e^{-\tau^{*} J_{-}}$ $\zeta \tan \zeta = \tau$	$e^{\left(\zeta J_{+} - \zeta^{*} J_{-} \right)} = e^{\tau J_{+}} e^{\ln(1 + \tau^{*} \tau) J_3} e^{-\tau^{*} J_{-}}$ $\zeta \tan \zeta = \tau$	$e^{\left(\zeta J_{+} - \zeta^{*} J_{-} \right)} = e^{\tau J_{+}} e^{\ln(1 + \tau^{*} \tau) J_3} e^{-\tau^{*} J_{-}}$ $\zeta \tan \zeta = \tau$
gilmore-lie-groups_ EQ0728 (13.47)	229	■ 1.000	$e^{\left(\alpha a^{\dagger} - \alpha^{*} a \right)} = e^{\alpha a^{\dagger}} e^{-\frac{1}{2} \alpha^{*} \alpha I} e^{-\alpha^{*} a}$ $\alpha = \lim_{c \rightarrow 0} \zeta / c$	$e^{\left(\alpha a^{\dagger} - \alpha^{*} a \right)} = e^{\alpha a^{\dagger}} e^{-\frac{1}{2} \alpha^{*} \alpha I} e^{-\alpha^{*} a}$ $\alpha = \lim_{c \rightarrow 0} \zeta / c$	$e^{\left(\alpha a^{\dagger} - \alpha^{*} a \right)} = e^{\alpha a^{\dagger}} e^{-\frac{1}{2} \alpha^{*} \alpha I} e^{-\alpha^{*} a}$ $\alpha = \lim_{c \rightarrow 0} \zeta / c$
gilmore-lie-groups_ EQ0729 (13.48)	229	■ 1.000	$Y_m^l(\theta, \phi) = \frac{e^{im\phi}}{\sqrt{2\pi}} P_m^l(\cos \theta)$ $Y_{-m}^l(\theta, \phi) = (-)^m Y_{+m}^l(\theta, \phi)^{*}$	$Y_m^l(\theta, \phi) = \frac{e^{im\phi}}{\sqrt{2\pi}} P_m^l(\cos \theta)$ $Y_{-m}^l(\theta, \phi) = (-)^m Y_{+m}^l(\theta, \phi)^{*}$	$Y_m^l(\theta, \phi) = \frac{e^{im\phi}}{\sqrt{2\pi}} P_m^l(\cos \theta)$ $Y_{-m}^l(\theta, \phi) = (-)^m Y_{+m}^l(\theta, \phi)^{*}$
gilmore-lie-groups_ EQ0730 (13.49)	229	■ 1.000	$P_m^l(u) = (-)^{l+m} \frac{1}{2^l l!} \sqrt{\frac{2l+1}{2}} \sqrt{\frac{(l-m)!}{(l+m)!}} (1-u^2)^{+m/2} \frac{d^{l+m}}{du^{l+m}} (1-u^2)^l$	$P_m^l(u) = (-)^{l+m} \frac{1}{2^l l!} \sqrt{\frac{2l+1}{2}} \sqrt{\frac{(l-m)!}{(l+m)!}} (1-u^2)^{+m/2} \frac{d^{l+m}}{du^{l+m}} (1-u^2)^l$	$P_m^l(u) = (-)^{l+m} \frac{1}{2^l l!} \sqrt{\frac{2l+1}{2}} \sqrt{\frac{(l-m)!}{(l+m)!}} (1-u^2)^{+m/2} \frac{d^{l+m}}{du^{l+m}} (1-u^2)^l$
gilmore-lie-groups_ EQ0732 (13.51)	230	■ 1.000	$\lim_{c \rightarrow 0} l^{-1/4} P_m^l(u = x/\sqrt{l}) = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} e^{x^2/2} \left(-\frac{d}{dx} \right)^n e^{-x^2} = \psi_n(x)$	$\lim_{c \rightarrow 0} l^{-1/4} P_m^l(u = x/\sqrt{l}) = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} e^{x^2/2} \left(-\frac{d}{dx} \right)^n e^{-x^2} = \psi_n(x)$	$\lim_{c \rightarrow 0} l^{-1/4} P_m^l(u = x/\sqrt{l}) = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} e^{x^2/2} \left(-\frac{d}{dx} \right)^n e^{-x^2} = \psi_n(x)$
gilmore-lie-groups_ EQ0733 (13.52)	230	■ 1.000	$\psi_n(x) = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} H_n(x) e^{-x^2/2}$	$\psi_n(x) = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} H_n(x) e^{-x^2/2}$	$\psi_n(x) = \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} H_n(x) e^{-x^2/2}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0734 (13.53)	230	■ 1.000	$\begin{aligned} \delta_{mm'} &= \int_{-1}^{+1} P_m^l(u) P_{m'}^l(u) du \\ &\rightarrow \lim_{l \rightarrow \infty} \int_{-\sqrt{l}}^{+\sqrt{l}} \left(\frac{1}{l^{1/4}} P_m^l(x/\sqrt{l}) \right) \left(\frac{1}{l^{1/4}} P_{m'}^l(x/\sqrt{l}) \right) d(u\sqrt{l}) \\ &\rightarrow \int_{-\infty}^{+\infty} \psi_n(x) \psi_{n'}(x) dx = \delta_{nn'} \end{aligned}$	$\delta_{mm'} = \int_{-1}^{+1} P_m^l(u) P_{m'}^l(u) du$ $\rightarrow \lim_{l \rightarrow \infty} \int_{-\sqrt{l}}^{+\sqrt{l}} \left(\frac{1}{l^{1/4}} P_m^l(x/\sqrt{l}) \right) \left(\frac{1}{l^{1/4}} P_{m'}^l(x/\sqrt{l}) \right) d(u\sqrt{l})$ $\rightarrow \int_{-\infty}^{+\infty} \psi_n(x) \psi_{n'}(x) dx = \delta_{nn'}$	$\delta_{mm'} = \int_{-1}^{+1} P_m^l(u) P_{m'}^l(u) du$ $\rightarrow \lim_{l \rightarrow \infty} \int_{-\sqrt{l}}^{+\sqrt{l}} \left(\frac{1}{l^{1/4}} P_m^l(x/\sqrt{l}) \right) \left(\frac{1}{l^{1/4}} P_{m'}^l(x/\sqrt{l}) \right) d(u\sqrt{l})$ $\rightarrow \int_{-\infty}^{+\infty} \psi_n(x) \psi_{n'}(x) dx = \delta_{nn'}$
gilmore-lie-groups_ EQ0737	231	■ 1.000	$\lim_{l \rightarrow \infty} P_{mn}^l(\cos(x/l)) = (-1)^{m-n} J_{m-n}(x)$	$\lim_{l \rightarrow \infty} P_{mn}^l(\cos(x/l)) = (-1)^{m-n} J_{m-n}(x)$	$\lim_{l \rightarrow \infty} P_{mn}^l(\cos(x/l)) = (-1)^{m-n} J_{m-n}(x)$
gilmore-lie-groups_ EQ0738	231	■ 1.000	$\lim_{l \rightarrow \infty} \sqrt{\frac{2\pi}{l}} Y_m^l(\beta = x/l) \rightarrow J_m(x)$	$\lim_{l \rightarrow \infty} \sqrt{\frac{2\pi}{l}} Y_m^l(\beta = x/l) \rightarrow J_m(x)$	$\lim_{l \rightarrow \infty} \sqrt{\frac{2\pi}{l}} Y_m^l(\beta = x/l) \rightarrow J_m(x)$
gilmore-lie-groups_ EQ0739	231	■ 1.000	$\lim_{l \rightarrow \infty} P^l(\cos(\beta = x/l)) \rightarrow J_0(x)$	$\lim_{l \rightarrow \infty} P^l(\cos(\beta = x/l)) \rightarrow J_0(x)$	$\lim_{l \rightarrow \infty} P^l(\cos(\beta = x/l)) \rightarrow J_0(x)$
gilmore-lie-groups_ EQ0740	231	■ 1.000	$e^{\alpha J_+} Y_m^l(\theta, \phi) = \sum_{k \geq 0} A_k^l Y_{m+k}^l(\theta, \phi) = Y_m^l(\theta', \phi')$	$e^{\alpha J_+} Y_m^l(\theta, \phi) = \sum_{k \geq 0} A_k^l Y_{m+k}^l(\theta, \phi) = Y_m^l(\theta', \phi')$	$e^{\alpha J_+} Y_m^l(\theta, \phi) = \sum_{k \geq 0} A_k^l Y_{m+k}^l(\theta, \phi) = Y_m^l(\theta', \phi')$
gilmore-lie-groups_ EQ0744	232	■ 1.000	$\begin{aligned} \lim_{\epsilon \rightarrow 0} \epsilon (2J_3) &\rightarrow P \quad P' = \partial_x \\ \lim_{\epsilon \rightarrow 0} \epsilon (J_+) &\rightarrow T \quad T' = \partial_t \\ \lim_{\epsilon \rightarrow 0} (J_-) &\rightarrow V \quad V' = t \partial_x \end{aligned}$	$\lim_{\epsilon \rightarrow 0} \epsilon (2J_3) \rightarrow P \quad P' = \partial_x$ $\lim_{\epsilon \rightarrow 0} \epsilon (J_+) \rightarrow T \quad T' = \partial_t$ $\lim_{\epsilon \rightarrow 0} (J_-) \rightarrow V \quad V' = t \partial_x$	$\lim_{\epsilon \rightarrow 0} \epsilon (2J_3) \rightarrow P \quad P' = \partial_x$ $\lim_{\epsilon \rightarrow 0} \epsilon (J_+) \rightarrow T \quad T' = \partial_t$ $\lim_{\epsilon \rightarrow 0} (J_-) \rightarrow V \quad V' = t \partial_x$
gilmore-lie-groups_ EQ0745	233	■ 1.000	$e^{\zeta J_+ - \zeta^* J_-} = e^{\tau J_+} e^{\log(1 + \tau^* \tau) J_3} e^{-\tau^* J_-}$	$e^{\zeta J_+ - \zeta^* J_-} = e^{\tau J_+} e^{\log(1 + \tau^* \tau) J_3} e^{-\tau^* J_-}$	$e^{\zeta J_+ - \zeta^* J_-} = e^{\tau J_+} e^{\log(1 + \tau^* \tau) J_3} e^{-\tau^* J_-}$
gilmore-lie-groups_ EQ0747	233	■ 1.000	$\begin{aligned} e^{\theta \cdot J} &\rightarrow \begin{bmatrix} \cosh(\theta/2) + (\theta_z/\theta) \sinh(\theta/2) & (\theta_x - i\theta_y)/\theta \sinh(\theta/2) \\ (\theta_x + i\theta_y)/\theta \sinh(\theta/2) & \cosh(\theta/2) - (\theta_z/\theta) \sinh(\theta/2) \end{bmatrix} \\ e^{-\beta \mathcal{H}} &\rightarrow \begin{bmatrix} e^{-\beta \epsilon/2} & 0 \\ 0 & e^{+\beta \epsilon/2} \end{bmatrix} \end{aligned}$	$e^{\theta \cdot J} \rightarrow \begin{bmatrix} \cosh(\theta/2) + (\theta_z/\theta) \sinh(\theta/2) & (\theta_x - i\theta_y)/\theta \sinh(\theta/2) \\ (\theta_x + i\theta_y)/\theta \sinh(\theta/2) & \cosh(\theta/2) - (\theta_z/\theta) \sinh(\theta/2) \end{bmatrix}$ $e^{-\beta \mathcal{H}} \rightarrow \begin{bmatrix} e^{-\beta \epsilon/2} & 0 \\ 0 & e^{+\beta \epsilon/2} \end{bmatrix}$	$e^{\theta \cdot J} \rightarrow \begin{bmatrix} \cosh(\theta/2) + (\theta_z/\theta) \sinh(\theta/2) & (\theta_x - i\theta_y)/\theta \sinh(\theta/2) \\ (\theta_x + i\theta_y)/\theta \sinh(\theta/2) & \cosh(\theta/2) - (\theta_z/\theta) \sinh(\theta/2) \end{bmatrix}$ $e^{-\beta \mathcal{H}} \rightarrow \begin{bmatrix} e^{-\beta \epsilon/2} & 0 \\ 0 & e^{+\beta \epsilon/2} \end{bmatrix}$
gilmore-lie-groups_ EQ0748	233	■ 1.000	$2 \cosh(\theta/2) \cosh(\beta \epsilon/2) - 2(\theta_z/\theta) \sinh(\theta/2) \sinh(\beta \epsilon/2) \quad (= 2 \cosh(\psi/2))$	$2 \cosh(\theta/2) \cosh(\beta \epsilon/2) - 2(\theta_z/\theta) \sinh(\theta/2) \sinh(\beta \epsilon/2) \quad (= 2 \cosh(\psi/2))$	$2 \cosh(\theta/2) \cosh(\beta \epsilon/2) - 2(\theta_z/\theta) \sinh(\theta/2) \sinh(\beta \epsilon/2) \quad (= 2 \cosh(\psi/2))$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0749	233	■ 1.000	$\left\langle e^{\mathrm{i} \theta J} \right\rangle = \frac{\sinh \psi / \sinh (\psi / 2)}{\sinh \beta \epsilon / \sinh (\beta \epsilon / 2)}$	$\langle e^{\theta \cdot J} \rangle = (\sinh \psi / \sinh (\psi / 2)) / (\sinh \beta \epsilon / \sinh (\beta \epsilon / 2))$	$\langle e^{\theta \cdot J} \rangle = (\sinh \psi / \sinh (\psi / 2)) / (\sinh \beta \epsilon / \sinh (\beta \epsilon / 2))$
gilmore-lie-groups_ EQ0750	233	■ 1.000	$\left\langle e^{\mathrm{i} \theta J} \right\rangle = \frac{\sinh ((2 j+1) \psi / 2) / \sinh (\psi / 2)}{\sinh ((2 j+1) \beta \epsilon / 2) / \sinh (\beta \epsilon / 2)}$	$\langle e^{\theta \cdot J} \rangle = \frac{\sinh ((2 j+1) \psi / 2) / \sinh (\psi / 2)}{\sinh ((2 j+1) \beta \epsilon / 2) / \sinh (\beta \epsilon / 2)}$	$\langle e^{\theta \cdot J} \rangle = \frac{\sinh ((2 j+1) \psi / 2) / \sinh (\psi / 2)}{\sinh ((2 j+1) \beta \epsilon / 2) / \sinh (\beta \epsilon / 2)}$
gilmore-lie-groups_ EQ0751	233	■ 1.000	$\left\langle e^{\mathrm{i} \theta J} \right\rangle \xrightarrow{j \rightarrow \infty} \sinh (j \psi) / \sinh (j \beta \epsilon)$	$\langle e^{\theta \cdot J} \rangle \xrightarrow{j \rightarrow \infty} \sinh (j \psi) / \sinh (j \beta \epsilon)$	$\langle e^{\theta \cdot J} \rangle \xrightarrow{j \rightarrow \infty} \sinh (j \psi) / \sinh (j \beta \epsilon)$
gilmore-lie-groups_ EQ0755	234	■ 1.000	$g(\mathfrak{p}')_{\alpha, \beta} = \lim_{\epsilon \rightarrow 0} (Y_{\alpha}, Y_{\beta}) / \epsilon^2 = (X_{\alpha}, X_{\beta})$	$g(\mathfrak{p}')_{\alpha, \beta} = \lim_{\epsilon \rightarrow 0} (Y_{\alpha}, Y_{\beta}) / \epsilon^2 = (X_{\alpha}, X_{\beta})$	$g(\mathfrak{p}')_{\alpha, \beta} = \lim_{\epsilon \rightarrow 0} (Y_{\alpha}, Y_{\beta}) / \epsilon^2 = (X_{\alpha}, X_{\beta})$
gilmore-lie-groups_ EQ0756 (14.1)	237	■ 1.000	$\langle S \mid \psi \rangle = \langle S \mid S' \rangle \langle S' \mid \psi \rangle$	$\langle S \mid \psi \rangle = \langle S \mid S' \rangle \langle S' \mid \psi \rangle$	$\langle S \mid \psi \rangle = \langle S \mid S' \rangle \langle S' \mid \psi \rangle$
gilmore-lie-groups_ EQ0757	237	■ 1.000	the rest of the universe looks the same	<i>therestoftheuniverselooksthesame</i>	the rest of the universe looks the same
gilmore-lie-groups_ EQ0758 (14.2)	237	■ 1.000	$\langle S \mid \psi' \rangle = \langle S' \mid \psi \rangle$	$\langle S \mid \psi' \rangle = \langle S' \mid \psi \rangle$	$\langle S \mid \psi' \rangle = \langle S' \mid \psi \rangle$
gilmore-lie-groups_ EQ0759 (14.3)	237	■ 1.000	$E^2 - (\mathbf{p} c)^2 = (m c^2)^2$	$E^2 - (\mathbf{p} c)^2 = (m c^2)^2$	$E^2 - (\mathbf{p} c)^2 = (m c^2)^2$
gilmore-lie-groups_ EQ0760 (14.4)	238	■ 1.000	$\left\{ E^2 - (m c^2)^2 + 2 E \left(\frac{e^2}{r} \right) + \left(\frac{e^2}{r} \right)^2 - (-i \hbar c \nabla)^2 \right\} \psi(x) = 0$	$\left\{ E^2 - (m c^2)^2 + 2 E \left(\frac{e^2}{r} \right) + \left(\frac{e^2}{r} \right)^2 - (-i \hbar c \nabla)^2 \right\} \psi(x) = 0$	$\left\{ E^2 - (m c^2)^2 + 2 E \left(\frac{e^2}{r} \right) + \left(\frac{e^2}{r} \right)^2 - (-i \hbar c \nabla)^2 \right\} \psi(x) = 0$
gilmore-lie-groups_ EQ0761 (14.5)	238	■ 1.000	$\left\{ \frac{\mathbf{p} \cdot \mathbf{p}}{2 m} - \frac{e^2}{r} - W \right\} \psi(\mathbf{x}) = \left\{ -\frac{\hbar^2}{2 m} \nabla^2 - \frac{e^2}{r} - W \right\} \psi(\mathbf{x}) = 0$	$\left\{ \frac{\mathbf{p} \cdot \mathbf{p}}{2 m} - \frac{e^2}{r} - W \right\} \psi(\mathbf{x}) = \left\{ -\frac{\hbar^2}{2 m} \nabla^2 - \frac{e^2}{r} - W \right\} \psi(\mathbf{x}) = 0$	$\left\{ \frac{\mathbf{p} \cdot \mathbf{p}}{2 m} - \frac{e^2}{r} - W \right\} \psi(\mathbf{x}) = \left\{ -\frac{\hbar^2}{2 m} \nabla^2 - \frac{e^2}{r} - W \right\} \psi(\mathbf{x}) = 0$
gilmore-lie-groups_ EQ0762 (14.6)	239	■ 1.000	$\nabla^2 = \left(\frac{1}{r} \frac{\partial}{\partial r} r \right)^2 + \frac{\mathcal{L}^2(S^2)}{r^2}$	$\nabla^2 = \left(\frac{1}{r} \frac{\partial}{\partial r} r \right)^2 + \frac{\mathcal{L}^2(S^2)}{r^2}$	$\nabla^2 = \left(\frac{1}{r} \frac{\partial}{\partial r} r \right)^2 + \frac{\mathcal{L}^2(S^2)}{r^2}$
gilmore-lie-groups_ EQ0763 (14.8)	239	■ 1.000	$\psi(r, \theta, \phi) \rightarrow \frac{1}{r} R(r) Y_m^l(\theta, \phi)$	$\psi(r, \theta, \phi) \rightarrow \frac{1}{r} R(r) Y_m^l(\theta, \phi)$	$\psi(r, \theta, \phi) \rightarrow \frac{1}{r} R(r) Y_m^l(\theta, \phi)$
gilmore-lie-groups_ EQ0764 (14.9)	239	■ 1.000	$\left(\frac{d^2}{dr^2} + \frac{A}{r^2} + \frac{B}{r} + C \right) R(r) = 0$	$\left(\frac{d^2}{dr^2} + \frac{A}{r^2} + \frac{B}{r} + C \right) R(r) = 0$	$\left(\frac{d^2}{dr^2} + \frac{A}{r^2} + \frac{B}{r} + C \right) R(r) = 0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0767	240	■ 1.000	$\begin{aligned} & \& f \simeq e^{-2 \lambda r} \text{ if series does not terminate } \\ & \& \simeq e^{+1 \lambda r} \text{ if series does terminate } (\lambda < 0) \end{aligned}$	$f \simeq e^{-2 \lambda r} \text{ if series does not terminate}$ $\simeq e^{+1 \lambda r} \text{ if series does terminate } (\lambda < 0)$	$f \simeq e^{-2 \lambda r} \text{ if series does not terminate}$ $\simeq e^{+1 \lambda r} \text{ if series does terminate } (\lambda < 0)$
gilmore-lie-groups_ EQ0768	240	■ 1.000	$\begin{aligned} & 2 \lambda(n+\gamma)+B=0 \text{ or } \\ & n+\frac{1}{2}+\sqrt{\left(\frac{1}{2}\right)^2-A}=\frac{B}{2 \sqrt{-C}} \end{aligned}$	$2 \lambda(n+\gamma)+B=0 \text{ or}$ $n+\frac{1}{2}+\sqrt{\left(\frac{1}{2}\right)^2-A}=\frac{B}{2 \sqrt{-C}}$	$2 \lambda(n+\gamma)+B=0 \text{ or}$ $n+\frac{1}{2}+\sqrt{\left(\frac{1}{2}\right)^2-A}=\frac{B}{2 \sqrt{-C}}$
gilmore-lie-groups_ EQ0771 (14.13)	241	■ 1.000	$g_{\{i\}} H g_{\{i\}}^{-1}=H \quad \text{or} \quad g_{\{i\}} H=H g_{\{i\}}$	$g_i H g_i^{-1}=H \quad \text{or} \quad g_i H=H g_i$	$g_i H g_i^{-1}=H \quad \text{or} \quad g_i H=H g_i$
gilmore-lie-groups_ EQ0772 (14.14)	242	■ 1.000	$H \left(g_{\{i\}} \mid \psi \right)=\left(H g_{\{i\}} \right) \mid \psi \rangle \mid \psi \rangle \mid \psi \rangle =g_{\{i\}} \left(H \mid \psi \rangle \right) \mid \psi \rangle \mid \psi \rangle =g_{\{i\}} \left(E \mid \psi \rangle \right) \mid \psi \rangle \mid \psi \rangle =E \left(g_{\{i\}} \mid \psi \rangle \right) \mid \psi \rangle \mid \psi \rangle$	$H \left(g_i \mid \psi \right)=\left(H g_i \right) \mid \psi \rangle =\left(g_i H \right) \mid \psi \rangle =g_i \left(H \mid \psi \right)=g_i \left(E \mid \psi \right)=E \left(g_i \mid \psi \right)$	$H \left(g_i \mid \psi \right)=\left(H g_i \right) \mid \psi \rangle =\left(g_i H \right) \mid \psi \rangle =g_i \left(H \mid \psi \right)=g_i \left(E \mid \psi \right)=E \left(g_i \mid \psi \right)$
gilmore-lie-groups_ EQ0773 (14.15)	242	■ 1.000	$R(\theta)=e^{\epsilon_{i j k} \theta_i x_j \partial_k}=e^{i \theta \cdot \mathbf{L} / \hbar}$	$R(\theta)=e^{\epsilon_{i j k} \theta_i x_j \partial_k}=e^{i \theta \cdot \mathbf{L} / \hbar}$	$R(\theta)=e^{\epsilon_{i j k} \theta_i x_j \partial_k}=e^{i \theta \cdot \mathbf{L} / \hbar}$
gilmore-lie-groups_ EQ0774 (14.16)	242	■ 1.000	$\left[L_{\{i\}}, L_{\{j\}}\right]=i \hbar \epsilon_{i j k} L_{\{k\}}$	$\left[L_i, L_j\right]=i \hbar \epsilon_{i j k} L_k$	$\left[L_i, L_j\right]=i \hbar \epsilon_{i j k} L_k$
gilmore-lie-groups_ EQ0775 (14.18)	242	■ 1.000	$\begin{aligned} & \left\{\left[L_{\{z\}}, L_{\{\pm\}}\right]\right\}=\pm \hbar L_{\pm} \\ & \left\{\left[L_{\{+\}}, L_{\{-\}}\right]\right\}=2 \hbar L_z \end{aligned}$	$\left[L_z, L_{\pm}\right]=\pm \hbar L_{\pm}$ $\left[L_{+}, L_{-}\right]=2 \hbar L_z$	$\left[L_z, L_{\pm}\right]=\pm \hbar L_{\pm}$ $\left[L_{+}, L_{-}\right]=2 \hbar L_z$
gilmore-lie-groups_ EQ0779 (14.21)	244	■ 1.000	$L_{\{+\}} Y_{\{m\}}^l(\theta, \phi)=Y_{\{m+1\}}^l(\theta, \phi) \sqrt{(l-m)(l+m+1)}$	$L_{+} Y_m^l(\theta, \phi)=Y_{m+1}^l(\theta, \phi) \sqrt{(l-m)(l+m+1)}$	$L_{+} Y_m^l(\theta, \phi)=Y_{m+1}^l(\theta, \phi) \sqrt{(l-m)(l+m+1)}$
gilmore-lie-groups_ EQ0781 (14.23)	245	■ 1.000	$\mathbf{M}=\frac{\mathbf{p} \times \mathbf{L}}{m}-K \frac{\mathbf{r}}{r}$	$\mathbf{M}=\frac{\mathbf{p} \times \mathbf{L}}{m}-K \frac{\mathbf{r}}{r}$	$\mathbf{M}=\frac{\mathbf{p} \times \mathbf{L}}{m}-K \frac{\mathbf{r}}{r}$
gilmore-lie-groups_ EQ0782 (14.24)	245	■ 1.000	$\hat{\mathbf{M}}=\frac{\hat{\mathbf{p}} \times \hat{\mathbf{L}}-\hat{\mathbf{L}} \times \hat{\mathbf{p}}}{2 m}-K \frac{\hat{\mathbf{r}}}{r}$	$\hat{\mathbf{M}}=\frac{\hat{\mathbf{p}} \times \hat{\mathbf{L}}-\hat{\mathbf{L}} \times \hat{\mathbf{p}}}{2 m}-K \frac{\hat{\mathbf{r}}}{r}$	$\hat{\mathbf{M}}=\frac{\hat{\mathbf{p}} \times \hat{\mathbf{L}}-\hat{\mathbf{L}} \times \hat{\mathbf{p}}}{2 m}-K \frac{\hat{\mathbf{r}}}{r}$
gilmore-lie-groups_ EQ0783 (14.25)	245	■ 1.000	$\begin{aligned} & \left\{\left[L_{\{i\}}, L_{\{j\}}\right]\right\}=i \hbar \epsilon_{i j k} L_{\{k\}} \\ & \left\{\left[L_{\{i\}}, M_{\{j\}}\right]\right\}=i \hbar \epsilon_{i j k} M_{\{k\}} \\ & \left\{\left[M_{\{i\}}, M_{\{j\}}\right]\right\}=\left(-\frac{2 H}{m}\right) i \hbar \epsilon_{i j k} L_{\{k\}} \end{aligned}$	$\left[L_i, L_j\right]=i \hbar \epsilon_{i j k} L_k$ $\left[L_i, M_j\right]=i \hbar \epsilon_{i j k} M_k$ $\left[M_i, M_j\right]=\left(-\frac{2 H}{m}\right) i \hbar \epsilon_{i j k} L_k$	$\left[L_i, L_j\right]=i \hbar \epsilon_{i j k} L_k$ $\left[L_i, M_j\right]=i \hbar \epsilon_{i j k} M_k$ $\left[M_i, M_j\right]=\left(-\frac{2 H}{m}\right) i \hbar \epsilon_{i j k} L_k$
gilmore-lie-groups_ EQ0784 (14.26)	246	■ 1.000	$\begin{aligned} & \mathbf{L} \cdot \mathbf{M}=\mathbf{M} \cdot \mathbf{L}=0 \\ & \mathbf{M} \cdot \mathbf{M}=\frac{2 H}{m}\left(\mathbf{L} \cdot \mathbf{L}+\hbar^2\right)+K^2 \end{aligned}$	$\mathbf{L} \cdot \mathbf{M}=\mathbf{M} \cdot \mathbf{L}=0$ $\mathbf{M} \cdot \mathbf{M}=\frac{2 H}{m}\left(\mathbf{L} \cdot \mathbf{L}+\hbar^2\right)+K^2$	$\mathbf{L} \cdot \mathbf{M}=\mathbf{M} \cdot \mathbf{L}=0$ $\mathbf{M} \cdot \mathbf{M}=\frac{2 H}{m}\left(\mathbf{L} \cdot \mathbf{L}+\hbar^2\right)+K^2$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00785 (14.27)	246	■ 1.000	$\begin{aligned} \{\left[L_i, L_j\right] \} &= i \hbar \epsilon_{ijk} L_k \\ \{\left[L_i, M_j'\right] \} &= i \hbar \epsilon_{ijk} M_k' \\ \{\left[M_i', M_j'\right] \} &= i \hbar \epsilon_{ijk} L_k \end{aligned}$	$\begin{aligned} [L_i, L_j] &= i \hbar \epsilon_{ijk} L_k \\ [L_i, M_j'] &= i \hbar \epsilon_{ijk} M_k' \\ [M_i', M_j'] &= i \hbar \epsilon_{ijk} L_k \end{aligned}$	$\begin{aligned} [L_i, L_j] &= i \hbar \epsilon_{ijk} L_k \\ [L_i, M_j'] &= i \hbar \epsilon_{ijk} M_k' \\ [M_i', M_j'] &= i \hbar \epsilon_{ijk} L_k \end{aligned}$
gilmore-lie-groups_ E00786 (14.28)	246	■ 1.000	$\begin{aligned} &\mathbf{A} = \frac{1}{2} (\mathbf{L} + \mathbf{M}') \\ &\mathbf{B} = \frac{1}{2} (\mathbf{L} - \mathbf{M}') \end{aligned}$	$\begin{aligned} \mathbf{A} &= \frac{1}{2} (\mathbf{L} + \mathbf{M}') \\ \mathbf{B} &= \frac{1}{2} (\mathbf{L} - \mathbf{M}') \end{aligned}$	$\begin{aligned} \mathbf{A} &= \frac{1}{2} (\mathbf{L} + \mathbf{M}') \\ \mathbf{B} &= \frac{1}{2} (\mathbf{L} - \mathbf{M}') \end{aligned}$
gilmore-lie-groups_ E00791 (14.33)	248	■ 1.000	$p_x^2 + \left(p_y - \frac{mM}{L}\right)^2 = \left(\frac{mK}{L}\right)^2$	$p_x^2 + \left(p_y - \frac{mM}{L}\right)^2 = \left(\frac{mK}{L}\right)^2$	$p_x^2 + \left(p_y - \frac{mM}{L}\right)^2 = \left(\frac{mK}{L}\right)^2$
gilmore-lie-groups_ E00793 (14.35)	248	■ 1.000	$\hat{\mathbf{u}} = \frac{\mathbf{p} \cdot \mathbf{p} - p_0^2}{\mathbf{p} \cdot \mathbf{p} + p_0^2} \hat{\mathbf{w}} + \frac{2p_0}{\mathbf{p} \cdot \mathbf{p} + p_0^2} \mathbf{p}$	$\hat{\mathbf{u}} = \frac{\mathbf{p} \cdot \mathbf{p} - p_0^2}{\mathbf{p} \cdot \mathbf{p} + p_0^2} \hat{\mathbf{w}} + \frac{2p_0}{\mathbf{p} \cdot \mathbf{p} + p_0^2} \mathbf{p}$	$\hat{\mathbf{u}} = \frac{\mathbf{p} \cdot \mathbf{p} - p_0^2}{\mathbf{p} \cdot \mathbf{p} + p_0^2} \hat{\mathbf{w}} + \frac{2p_0}{\mathbf{p} \cdot \mathbf{p} + p_0^2} \mathbf{p}$
gilmore-lie-groups_ E00794 (14.36)	248	■ 1.000	$\text{circle in } R^3 \xrightarrow{T} \text{circle in } S^3 \xrightarrow{SO(4)} \text{circle in } S^3 \xrightarrow{T^{-1}} \text{circle in } R^3$	$\text{circle in } R^3 \xrightarrow{T} \text{circle in } S^3 \xrightarrow{SO(4)} \text{circle in } S^3 \xrightarrow{T^{-1}} \text{circle in } R^3$	$\text{circle in } R^3 \xrightarrow{T} \text{circle in } S^3 \xrightarrow{SO(4)} \text{circle in } S^3 \xrightarrow{T^{-1}} \text{circle in } R^3$
gilmore-lie-groups_ E00795 (14.37)	249	■ 1.000	$(\mathbf{y}, \mathbf{y})_N - z_1 z_2 = (\lambda \mathbf{x}, \lambda \mathbf{x})_N - \lambda [\lambda(\mathbf{x}, \mathbf{x})_N] = 0$	$(\mathbf{y}, \mathbf{y})_N - z_1 z_2 = (\lambda \mathbf{x}, \lambda \mathbf{x})_N - \lambda [\lambda(\mathbf{x}, \mathbf{x})_N] = 0$	$(\mathbf{y}, \mathbf{y})_N - z_1 z_2 = (\lambda \mathbf{x}, \lambda \mathbf{x})_N - \lambda [\lambda(\mathbf{x}, \mathbf{x})_N] = 0$
gilmore-lie-groups_ E00796 (14.39)	250	■ 1.000	$\begin{aligned} &\hat{\mathbf{u}} = \frac{\frac{1}{2}(p_0^2 - \mathbf{p} \cdot \mathbf{p})}{\frac{1}{2}(p_0^2 + \mathbf{p} \cdot \mathbf{p})} \mathbf{w} + \frac{p_0 \mathbf{p}}{\frac{1}{2}(p_0^2 + \mathbf{p} \cdot \mathbf{p})} \quad E < 0 \\ &\hat{\mathbf{u}} = \frac{\frac{1}{2}(p_0^2 + \mathbf{p} \cdot \mathbf{p})}{\frac{1}{2}(p_0^2 - \mathbf{p} \cdot \mathbf{p})} \mathbf{w} + \frac{p_0 \mathbf{p}}{\frac{1}{2}(p_0^2 - \mathbf{p} \cdot \mathbf{p})} \quad E > 0 \end{aligned}$	$\begin{aligned} \hat{\mathbf{u}} &= \frac{\frac{1}{2}(p_0^2 - \mathbf{p} \cdot \mathbf{p})}{\frac{1}{2}(p_0^2 + \mathbf{p} \cdot \mathbf{p})} \mathbf{w} + \frac{p_0 \mathbf{p}}{\frac{1}{2}(p_0^2 + \mathbf{p} \cdot \mathbf{p})} E < 0 \\ \hat{\mathbf{u}} &= \frac{\frac{1}{2}(p_0^2 + \mathbf{p} \cdot \mathbf{p})}{\frac{1}{2}(p_0^2 - \mathbf{p} \cdot \mathbf{p})} \mathbf{w} + \frac{p_0 \mathbf{p}}{\frac{1}{2}(p_0^2 - \mathbf{p} \cdot \mathbf{p})} E > 0 \end{aligned}$	$\begin{aligned} \hat{\mathbf{u}} &= \frac{\frac{1}{2}(p_0^2 - \mathbf{p} \cdot \mathbf{p})}{\frac{1}{2}(p_0^2 + \mathbf{p} \cdot \mathbf{p})} \mathbf{w} + \frac{p_0 \mathbf{p}}{\frac{1}{2}(p_0^2 + \mathbf{p} \cdot \mathbf{p})} \quad E < 0 \\ \hat{\mathbf{u}} &= \frac{\frac{1}{2}(p_0^2 + \mathbf{p} \cdot \mathbf{p})}{\frac{1}{2}(p_0^2 - \mathbf{p} \cdot \mathbf{p})} \mathbf{w} + \frac{p_0 \mathbf{p}}{\frac{1}{2}(p_0^2 - \mathbf{p} \cdot \mathbf{p})} \quad E > 0 \end{aligned}$
gilmore-lie-groups_ E00797 (14.40)	250	■ 1.000	$\mathbf{u}^t G \mathbf{u} = u_0^2 \pm \sum_{i=1}^3 u_i^2 \quad \begin{aligned} &+ \text{ for } E < 0 \\ &- \text{ for } E > 0 \end{aligned}$	$\mathbf{u}^t G \mathbf{u} = u_0^2 \pm \sum_{i=1}^3 u_i^2 \quad \begin{aligned} &+ \text{ for } E < 0 \\ &- \text{ for } E > 0 \end{aligned}$	$\mathbf{u}^t G \mathbf{u} = u_0^2 \pm \sum_{i=1}^3 u_i^2 \quad \begin{aligned} &+ \text{ for } E < 0 \\ &- \text{ for } E > 0 \end{aligned}$
gilmore-lie-groups_ E00799 (14.42)	251	■ 1.000	$\mathbf{u}' = \frac{A(\lambda \mathbf{u}) + B\lambda}{C(\lambda \mathbf{u}) + D\lambda}$	$\mathbf{u}' = \frac{A(\lambda \mathbf{u}) + B\lambda}{C(\lambda \mathbf{u}) + D\lambda}$	$\mathbf{u}' = \frac{A(\lambda \mathbf{u}) + B\lambda}{C(\lambda \mathbf{u}) + D\lambda}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0800 (14.43)	251	■ 1.000	$\left(\mathbf{u}^{\prime}\right)^t G \mathbf{u}^{\prime}-1=\frac{(A \lambda \mathbf{u}+B \lambda)^t G(A \lambda \mathbf{u}+B \lambda)-(C \lambda \mathbf{u}+D \lambda)^t(C \lambda \mathbf{u}+D \lambda)}{(C \lambda \mathbf{u}+D \lambda)^t(C \lambda \mathbf{u}+D \lambda)}$	$\left(\mathbf{u}^{\prime}\right)^t G \mathbf{u}^{\prime}-1=\frac{(A \lambda \mathbf{u}+B \lambda)^t G(A \lambda \mathbf{u}+B \lambda)-(C \lambda \mathbf{u}+D \lambda)^t(C \lambda \mathbf{u}+D \lambda)}{(C \lambda \mathbf{u}+D \lambda)^t(C \lambda \mathbf{u}+D \lambda)}$	$\left(\mathbf{u}^{\prime}\right)^t G \mathbf{u}^{\prime}-1=\frac{(A \lambda \mathbf{u}+B \lambda)^t G(A \lambda \mathbf{u}+B \lambda)-(C \lambda \mathbf{u}+D \lambda)^t(C \lambda \mathbf{u}+D \lambda)}{(C \lambda \mathbf{u}+D \lambda)^t(C \lambda \mathbf{u}+D \lambda)}$
gilmore-lie-groups_ EQ0801 (14.44)	251	■ 1.000	$\left(\mathbf{u}^{\prime}\right)^t G \mathbf{u}^{\prime}-1=\frac{(A \lambda \mathbf{u}+B \lambda)^t G(A \lambda \mathbf{u}+B \lambda)-(C \lambda \mathbf{u}+D \lambda)^t(C \lambda \mathbf{u}+D \lambda)}{(C \lambda \mathbf{u}+D \lambda)^t(C \lambda \mathbf{u}+D \lambda)}$	$\left(\mathbf{u}^{\prime}, \mathbf{u}^{\prime}\right)_N-1=\frac{(\mathbf{u}, \mathbf{u})_N-1}{(C \mathbf{u}+D)^t(C \mathbf{u}+D)}$	$\left(\mathbf{u}^{\prime}, \mathbf{u}^{\prime}\right)_N-1=\frac{(\mathbf{u}, \mathbf{u})_N-1}{(C \mathbf{u}+D)^t(C \mathbf{u}+D)}$
gilmore-lie-groups_ EQ0805	252	■ 1.000	$\begin{aligned} & a_1^{\dagger} a_2^{\dagger} a_1 a_2 \left m_1, m_2 \right\rangle \\ & ; n_1, n_2 \right\rangle \left m_1+1, m_2-1 \right\rangle \\ & n_1, n_2 \right\rangle \left m_1, m_2 \right\rangle \sqrt{m_1+1} \sqrt{m_2} \\ & \backslash \& b_1^{\dagger} b_1^{\dagger} b_1 b_1 \left m_1, m_2 \right\rangle ; n_1, \\ & n_2 \right\rangle \left m_1, m_2 \right\rangle ; n_1, n_2 \right\rangle \left m_1, m_2 \right\rangle \sqrt{n_1} \sqrt{n_2} \end{aligned}$	$a_1^{\dagger} a_2^{\dagger} \left m_1, m_2 ; n_1, n_2 \right\rangle = \left m_1+1, m_2-1 ; n_1 n_2 \right\rangle \sqrt{m_1+1} \sqrt{m_2}$ $b_1^{\dagger} b_1^{\dagger} \left m_1, m_2 ; n_1, n_2 \right\rangle = \left m_1, m_2 ; n_1 n_2 \right\rangle (\sqrt{n_1})^2$	$a_1^{\dagger} a_2^{\dagger} \left m_1, m_2 ; n_1, n_2 \right\rangle = \left m_1+1, m_2-1 ; n_1 n_2 \right\rangle \sqrt{m_1+1} \sqrt{m_2}$ $b_1^{\dagger} b_1^{\dagger} \left m_1, m_2 ; n_1, n_2 \right\rangle = \left m_1, m_2 ; n_1 n_2 \right\rangle (\sqrt{n_1})^2$
gilmore-lie-groups_ EQ0806 (14.46)	253	■ 1.000	$\begin{array}{l} \text{operators} \\ \text{subalgebra} \\ \text{number} \end{array} \begin{array}{l} a_i^{\dagger} a_j \\ u(2) \\ 4 \end{array} \begin{array}{l} b_i^{\dagger} b_j \\ u(2) \\ 4 \end{array} \begin{array}{l} a_i^{\dagger} b_j^{\dagger} \\ 4 \end{array} \begin{array}{l} a_i b_j \\ 4 \end{array}$	$\begin{array}{l} \text{operators} \\ \text{subalgebra} \\ \text{number} \end{array} \begin{array}{l} a_i^{\dagger} a_j \\ u(2) \\ 4 \end{array} \begin{array}{l} b_i^{\dagger} b_j \\ u(2) \\ 4 \end{array} \begin{array}{l} a_i^{\dagger} b_j^{\dagger} \\ 4 \end{array} \begin{array}{l} a_i b_j \\ 4 \end{array}$	$\begin{array}{l} \text{operators} \\ \text{subalgebra} \\ \text{number} \end{array} \begin{array}{l} a_i^{\dagger} a_j \\ u(2) \\ 4 \end{array} \begin{array}{l} b_i^{\dagger} b_j \\ u(2) \\ 4 \end{array} \begin{array}{l} a_i^{\dagger} b_j^{\dagger} \\ 4 \end{array} \begin{array}{l} a_i b_j \\ 4 \end{array}$
gilmore-lie-groups_ EQ0808	253	■ 1.000	$\left(a_1^{\dagger} a_1+a_2^{\dagger} a_2\right)-\left(b_1^{\dagger} b_1+b_2^{\dagger} b_2\right)$	$\left(a_1^{\dagger} a_1+a_2^{\dagger} a_2\right)-\left(b_1^{\dagger} b_1+b_2^{\dagger} b_2\right)$	$\left(a_1^{\dagger} a_1+a_2^{\dagger} a_2\right)-\left(b_1^{\dagger} b_1+b_2^{\dagger} b_2\right)$
gilmore-lie-groups_ EQ0811	254	■ 1.000	$\left(P_1, P_2, P_3, P_4\right)^t=\frac{1}{2 R} M_{K S}\left(p_1, p_2, p_3, p_4\right)^t$	$\left(P_1, P_2, P_3, P_4\right)^t=\frac{1}{2 R} M_{K S}\left(p_1, p_2, p_3, p_4\right)^t$	$\left(P_1, P_2, P_3, P_4\right)^t=\frac{1}{2 R} M_{K S}\left(p_1, p_2, p_3, p_4\right)^t$
gilmore-lie-groups_ EQ0812 (14.49)	254	■ 1.000	$\zeta=-2 R P_4=\left(q_1 p_4-q_4 p_1\right)+\left(q_3 p_2-q_2 p_3\right)=0$	$\zeta=-2 R P_4=\left(q_1 p_4-q_4 p_1\right)+\left(q_3 p_2-q_2 p_3\right)=0$	$\zeta=-2 R P_4=\left(q_1 p_4-q_4 p_1\right)+\left(q_3 p_2-q_2 p_3\right)=0$
gilmore-lie-groups_ EQ0813 (14.50)	255	■ 1.000	$\frac{P^2}{2 m}-\frac{e^2}{R}=E \xrightarrow{\times R} \frac{R P^2}{2 m}-e^2=E R \xrightarrow{K S} \frac{p^2}{8 m}-e^2=E q^2$	$\frac{P^2}{2 m}-\frac{e^2}{R}=E \xrightarrow{\times R} \frac{R P^2}{2 m}-e^2=E R \xrightarrow{K S} \frac{p^2}{8 m}-e^2=E q^2$	$\frac{P^2}{2 m}-\frac{e^2}{R}=E \xrightarrow{\times R} \frac{R P^2}{2 m}-e^2=E R \xrightarrow{K S} \frac{p^2}{8 m}-e^2=E q^2$
gilmore-lie-groups_ EQ0814 (14.51)	255	■ 1.000	$\frac{p^2}{2 m}-4 E q^2=4 e^2$	$\frac{p^2}{2 m}-4 E q^2=4 e^2$	$\frac{p^2}{2 m}-4 E q^2=4 e^2$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08815 (14.52)	255	■1.000	$\begin{aligned} & \left(Q_1, Q_2, Q_3, Q_4 \right) \\ & \left[\begin{array}{cccc} 0 & \theta_3 & -\theta_2 & * \\ * & -\theta_3 & 0 & \theta_1 \\ * & \theta_2 & -\theta_1 & 0 \\ * & * & * & 0 \end{array} \right] \left[\begin{array}{c} P_1 \\ P_2 \\ P_3 \\ P_4 \end{array} \right] \\ & \frac{1}{2} (q_1, q_2, q_3, q_4) \left[\begin{array}{cccc} 0 & \theta_3 & -\theta_2 & \theta_1 \\ -\theta_3 & 0 & \theta_1 & \theta_2 \\ \theta_2 & -\theta_1 & 0 & \theta_3 \\ -\theta_1 & -\theta_2 & -\theta_3 & 0 \end{array} \right] \left[\begin{array}{c} p_1 \\ p_2 \\ p_3 \\ p_4 \end{array} \right] \end{aligned}$	$(Q_1, Q_2, Q_3, Q_4) \left[\begin{array}{cccc} 0 & \theta_3 & -\theta_2 & * \\ -\theta_3 & 0 & \theta_1 & * \\ \theta_2 & -\theta_1 & 0 & * \\ -* & -* & -* & 0 \end{array} \right] \left[\begin{array}{c} P_1 \\ P_2 \\ P_3 \\ P_4 \end{array} \right]$ $\frac{1}{2} (q_1, q_2, q_3, q_4) \left[\begin{array}{cccc} 0 & \theta_3 & -\theta_2 & \theta_1 \\ -\theta_3 & 0 & \theta_1 & \theta_2 \\ \theta_2 & -\theta_1 & 0 & \theta_3 \\ -\theta_1 & -\theta_2 & -\theta_3 & 0 \end{array} \right] \left[\begin{array}{c} p_1 \\ p_2 \\ p_3 \\ p_4 \end{array} \right]$	
gilmore-lie-groups_ E08816	255	■1.000	$G_i=\left[\begin{array}{cc} 0 & M_i \\ -M_i & 0 \end{array}\right]$	$G_i = \left[\begin{array}{cc} 0 & M_i \\ -M_i & 0 \end{array} \right]$	$G_i = \left[\begin{array}{cc} 0 & M_i \\ -M_i & 0 \end{array} \right]$
gilmore-lie-groups_ E08820 (14.56)	256	■1.000	$\begin{aligned} & z_1=\frac{1}{\sqrt{2}}\left(s_1+i s_2\right) \quad z_2=\frac{1}{\sqrt{2}}\left(r_1+i r_2\right) \\ & z_3=\frac{1}{\sqrt{2}}\left(s_3+i s_4\right) \quad z_4=\frac{1}{\sqrt{2}}\left(r_3+i r_4\right) \end{aligned}$ $z_1^* z_1-z_2^* z_2+z_3^* z_3-z_4^* z_4=\zeta$	$z_1=\frac{1}{\sqrt{2}}\left(s_1+i s_2\right) \quad z_2=\frac{1}{\sqrt{2}}\left(r_1+i r_2\right)$ $z_3=\frac{1}{\sqrt{2}}\left(s_3+i s_4\right) \quad z_4=\frac{1}{\sqrt{2}}\left(r_3+i r_4\right)$ $z_1^* z_1-z_2^* z_2+z_3^* z_3-z_4^* z_4=\zeta$	$z_1=\frac{1}{\sqrt{2}}\left(s_1+i s_2\right) \quad z_2=\frac{1}{\sqrt{2}}\left(r_1+i r_2\right)$ $z_3=\frac{1}{\sqrt{2}}\left(s_3+i s_4\right) \quad z_4=\frac{1}{\sqrt{2}}\left(r_3+i r_4\right)$ $z_1^* z_1-z_2^* z_2+z_3^* z_3-z_4^* z_4=\zeta$
gilmore-lie-groups_ E08821 (14.57)	257	■1.000	$\begin{aligned} & \mathbf{J}=\mathbf{r} \times \mathbf{p} \\ & \mathbf{M}=\frac{1}{2 m}(\mathbf{p} \times \mathbf{L}-\mathbf{L} \times \mathbf{p})-K \frac{\mathbf{r}}{r} \\ & \mathbf{A}=\frac{1}{2 m}(\mathbf{p} \times \mathbf{L}-\mathbf{L} \times \mathbf{p})+K \frac{\mathbf{r}}{r} \\ & A_4=\mathbf{r} \cdot \mathbf{p}+\frac{3}{2} \frac{\hbar}{i} \end{aligned}$	$\mathbf{J}=\mathbf{r} \times \mathbf{p}$ angular momentum $\mathbf{M}=\frac{1}{2 m}(\mathbf{p} \times \mathbf{L}-\mathbf{L} \times \mathbf{p})-K \frac{\mathbf{r}}{r}$ Laplace-Runge-Lenz vector $\mathbf{A}=\frac{1}{2 m}(\mathbf{p} \times \mathbf{L}-\mathbf{L} \times \mathbf{p})+K \frac{\mathbf{r}}{r}$ dual vector $A_4=\mathbf{r} \cdot \mathbf{p}+\frac{3}{2} \frac{\hbar}{i}$ dual scalar	$\mathbf{J}=\mathbf{r} \times \mathbf{p}$ angular momentum $\mathbf{M}=\frac{1}{2 m}(\mathbf{p} \times \mathbf{L}-\mathbf{L} \times \mathbf{p})-K \frac{\mathbf{r}}{r}$ Laplace–Runge–Lenz vector $\mathbf{A}=\frac{1}{2 m}(\mathbf{p} \times \mathbf{L}-\mathbf{L} \times \mathbf{p})+K \frac{\mathbf{r}}{r}$ dual vector $A_4=\mathbf{r} \cdot \mathbf{p}+\frac{3}{2} \frac{\hbar}{i}$ dual scalar
gilmore-lie-groups_ E08824 (14.59)	258	■1.000	$H_D=-e \phi(r)+\beta m c^2+\gamma \cdot(c \mathbf{p}+e \mathbf{A})$	$H_D=-e \phi(r)+\beta m c^2+\gamma \cdot(c \mathbf{p}+e \mathbf{A})$	$H_D=-e \phi(r)+\beta m c^2+\gamma \cdot(c \mathbf{p}+e \mathbf{A})$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0825 (14.60)	258	■ 1.000	$\backslash beta=\backslash left[\backslash begin{array}{cc} I_{\{2\}} \ \& \ 0 \ \backslash \ 0 \ \& \ -I_{\{2\}} \ \backslash end{array}\backslash right] \quad \backslash gamma_{\{i\}}=\backslash left[\backslash begin{array}{cc} 0 \ \& \ \backslash sigma_{\{i\}} \ \backslash \ \backslash sigma_{\{i\}} \ \& \ 0 \ \backslash end{array}\backslash right]$	$\beta = \left[\begin{array}{cc} I_2 & 0 \\ 0 & -I_2 \end{array} \right] \quad \gamma_i = \left[\begin{array}{cc} 0 & \sigma_i \\ \sigma_i & 0 \end{array} \right]$	$\beta = \left[\begin{array}{cc} I_2 & 0 \\ 0 & -I_2 \end{array} \right] \quad \gamma_i = \left[\begin{array}{cc} 0 & \sigma_i \\ \sigma_i & 0 \end{array} \right]$
gilmore-lie-groups_ EQ0827 (14.62)	259	■ 1.000	$\backslash begin{aligned} \backslash Theta \ \& \ =r\backslash left\backslash \backslash left(H_{\{D\}}+\backslash frac{\backslash alpha}{\{r\}}\backslash right)^{\{2\}}-\backslash left(E+\backslash frac{\backslash alpha}{\{r\}}\backslash right)^{\{2\}} \backslash right\backslash \ \backslash \ \& \ =A\backslash left(\backslash Gamma_{\{5\}}+\backslash Gamma_{\{4\}}\backslash right)+B \backslash left(\backslash Gamma_{\{5\}}-\backslash Gamma_{\{4\}}\backslash right)+C \ \backslash end{aligned}$	$\Theta = r \left\{ \left(H_D + \frac{\alpha}{r} \right)^2 - \left(E + \frac{\alpha}{r} \right)^2 \right\} \\ = A (\Gamma_5 + \Gamma_4) + B (\Gamma_5 - \Gamma_4) + C$	$\Theta = r \left\{ \left(H_D + \frac{\alpha}{r} \right)^2 - \left(E + \frac{\alpha}{r} \right)^2 \right\} \\ = A (\Gamma_5 + \Gamma_4) + B (\Gamma_5 - \Gamma_4) + C$
gilmore-lie-groups_ EQ0828 (14.63)	259	■ 1.000	$\backslash left(r \ D^{\{2\}}+\backslash frac{A}{\{r\}}+B+C \ r\backslash right) \ R(r)=0$	$\left(r D^2 + \frac{A}{r} + B + C r \right) R(r) = 0$	$\left(r D^2 + \frac{A}{r} + B + C r \right) R(r) = 0$
gilmore-lie-groups_ EQ0830	260	■ 1.000	$\backslash begin{aligned} \ \& \ \backslash left[r \ D, \ \backslash frac{1}{\{r\}}\backslash right]=-\backslash frac{1}{\{r\}} \ \backslash \ \& \ \backslash left[r \ D^{\{2\}}, \ \backslash frac{1}{\{r\}}\backslash right]=\backslash frac{2}{\{r^{\{2\}}\}}-\backslash frac{1}{\{r\}} \ D \ \backslash end{aligned}$	$\left[r D, \frac{1}{r} \right] = -\frac{1}{r} \\ \left[r D^2, \frac{1}{r} \right] = \frac{2}{r^2} - \frac{1}{r} D$	$\left[r D, \frac{1}{r} \right] = -\frac{1}{r} \\ \left[r D^2, \frac{1}{r} \right] = \frac{2}{r^2} - \frac{1}{r} D$
gilmore-lie-groups_ EQ0831 (14.65)	260	■ 1.000	$\backslash begin{aligned} \backslash Gamma_{\{5\}} \ \& \ =\backslash frac{1}{\{2\}}\backslash left(r \ D^{\{2\}}+\backslash frac{a}{\{r\}}-r\backslash right) \ \backslash \ \backslash Gamma_{\{4\}} \ \& \ =\backslash frac{1}{\{2\}}\backslash left(r \ D^{\{2\}}+\backslash frac{a}{\{r\}}+r\backslash right) \ \backslash \ M_{\{4\}} \ \& \ =r \ D \ \backslash end{aligned}$	$\Gamma_5 = \frac{1}{2} \left(r D^2 + \frac{a}{r} - r \right) \\ \Gamma_4 = \frac{1}{2} \left(r D^2 + \frac{a}{r} + r \right) \\ M_4 = r D$	$\Gamma_5 = \frac{1}{2} \left(r D^2 + \frac{a}{r} - r \right) \\ \Gamma_4 = \frac{1}{2} \left(r D^2 + \frac{a}{r} + r \right) \\ M_4 = r D$
gilmore-lie-groups_ EQ0832 (14.66)	260	■ 1.000	$\backslash left(\backslash left(\backslash Gamma_{\{5\}}+\backslash Gamma_{\{4\}}\backslash right)+B+C\backslash left(\backslash Gamma_{\{4\}}-\backslash Gamma_{\{5\}}\backslash right)\backslash right) \ R(r)=0$	$((\Gamma_5 + \Gamma_4) + B + C (\Gamma_4 - \Gamma_5)) R(r) = 0$	$((\Gamma_5 + \Gamma_4) + B + C (\Gamma_4 - \Gamma_5)) R(r) = 0$
gilmore-lie-groups_ EQ0834 (14.68)	260	■ 1.000	$\backslash left[\backslash left(e^{\{-\backslash theta\}}-C \ e^{\{\backslash theta\}}\backslash right) \ \backslash Gamma_{\{5\}}+\backslash left(e^{\{-\backslash theta\}}+C \ e^{\{\backslash theta\}}\backslash right) \ \backslash Gamma_{\{4\}}+B\backslash right] \ e^{\{\backslash theta \ M_{\{4\}}\}} \ R(r)=0$	$[(e^{-\theta} - C e^{\theta}) \Gamma_5 + (e^{-\theta} + C e^{\theta}) \Gamma_4 + B] e^{\theta M_4} R(r) = 0$	$[(e^{-\theta} - C e^{\theta}) \Gamma_5 + (e^{-\theta} + C e^{\theta}) \Gamma_4 + B] e^{\theta M_4} R(r) = 0$
gilmore-lie-groups_ EQ0835 (14.69)	260	■ 1.000	$\backslash left(2 \ \backslash sqrt{-C} \ \backslash Gamma_{\{5\}}+B\backslash right) \ u(r)=0$	$\left(2\sqrt{-C} \Gamma_5 + B \right) u(r) = 0$	$\left(2\sqrt{-C} \Gamma_5 + B \right) u(r) = 0$
gilmore-lie-groups_ EQ0836 (14.70)	261	■ 1.000	$\backslash left(2 \ \backslash sqrt{C} \ \backslash Gamma_{\{4\}}+B\backslash right) \ u(r)=0$	$\left(2\sqrt{C} \Gamma_4 + B \right) u(r) = 0$	$\left(2\sqrt{C} \Gamma_4 + B \right) u(r) = 0$
gilmore-lie-groups_ EQ0837 (14.71)	261	■ 1.000	$R_{\{k, \ l\}}(r) \ \backslash sim \ \backslash sqrt{\backslash frac{2}{\{ \pi \}}} \ \backslash sin \ \backslash left(k \ r-\backslash frac{\{ \pi \}}{\{2\}} \ j+\backslash frac{\backslash alpha}{\{k\}}(\log(2 \ k \ r)+\backslash delta(j)) \ \backslash right)$	$R_{k,l}(r) \sim \sqrt{\frac{2}{\pi}} \sin \left(k r - \frac{\pi}{2} j + \frac{\alpha}{k} (\log(2 k r) + \delta(j)) \right)$	$R_{k,l}(r) \sim \sqrt{\frac{2}{\pi}} \sin \left(k r - \frac{\pi}{2} j + \frac{\alpha}{k} (\log(2 k r) + \delta(j)) \right)$
gilmore-lie-groups_ EQ0838 (14.72)	262	■ 1.000	$A \ \backslash rightharpoonup A^{\{\backslash prime\}}=A+\backslash mu_{\{l\}}$	$A \rightarrow A' = A + \mu_l$	$A \rightarrow A' = A + \mu_l$
gilmore-lie-groups_ EQ0839 (14.73)	262	■ 1.000	$E_{\{N=n+l+1\}}=-\backslash frac{m \ c^{\{2\}} \ \backslash alpha^{\{2\}}}{\{2 \ N^{\{2\}}\}} \ \backslash rightharpoonup -\backslash frac{m \ c^{\{2\}} \ \backslash alpha^{\{2\}}}{\{2(N+\backslash Delta \ j)\}^{\{2\}}}$	$E_{N=n+l+1} = -\frac{m c^2 \alpha^2}{2 N^2} \rightarrow -\frac{m c^2 \alpha^2}{2 (N + \Delta j)^2}$	$E_{N=n+l+1} = -\frac{m c^2 \alpha^2}{2 N^2} \rightarrow -\frac{m c^2 \alpha^2}{2 (N + \Delta j)^2}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08840 (14.74)	262	■ 1.000	$\Delta \phi = -\frac{\pi}{2} \Delta j + \frac{\alpha}{\pi} \arg(\Gamma[j+1+\Delta j-i(\alpha/k)]-\Gamma[j+1-i(\alpha/k)])$	$\Delta \phi = -\frac{\pi}{2} \Delta j + \frac{\alpha}{\pi} \arg(\Gamma[j+1+\Delta j-i(\alpha/k)]-\Gamma[j+1-i(\alpha/k)])$	$\Delta \phi = -\frac{\pi}{2} \Delta j + \frac{\alpha}{\pi} \arg(\Gamma[j+1+\Delta j-i(\alpha/k)]-\Gamma[j+1-i(\alpha/k)])$
gilmore-lie-groups_ E08841	264	■ 1.000	$\begin{aligned} & z = x_3 = \theta_3 \cos \theta_2 \\ & y = x_2 = \theta_3 \sin \theta_2 \cos \theta_1 \\ & x = x_1 = \theta_3 \sin \theta_2 \sin \theta_1 \end{aligned}$	$\begin{aligned} z &= x_3 = \theta_3 \cos \theta_2 \\ y &= x_2 = \theta_3 \sin \theta_2 \cos \theta_1 \\ x &= x_1 = \theta_3 \sin \theta_2 \sin \theta_1 \end{aligned}$	$\begin{aligned} z &= x_3 = \theta_3 \cos \theta_2 \\ y &= x_2 = \theta_3 \sin \theta_2 \cos \theta_1 \\ x &= x_1 = \theta_3 \sin \theta_2 \sin \theta_1 \end{aligned}$
gilmore-lie-groups_ E08842	264	■ 1.000	$\sin^2 \theta_2 \mathcal{L}^2(S^2) = \left(\sin \theta_2 \frac{\partial}{\partial \theta_2} \right)^2 + \mathcal{L}^2(S^1)$	$\sin^2 \theta_2 \mathcal{L}^2(S^2) = \left(\sin \theta_2 \frac{\partial}{\partial \theta_2} \right)^2 + \mathcal{L}^2(S^1)$	$\sin^2 \theta_2 \mathcal{L}^2(S^2) = \left(\sin \theta_2 \frac{\partial}{\partial \theta_2} \right)^2 + \mathcal{L}^2(S^1)$
gilmore-lie-groups_ E08843	265	■ 1.000	$\begin{aligned} r &\rightarrow 0 \quad \left(\frac{d^2}{dr^2} + \frac{A}{r^2} + \frac{B}{r} + C \right) R(r) \rightarrow \left(\frac{d^2}{dr^2} + \frac{A}{r^2} \right) R(r) = 0 \\ R(r) &\simeq r^\gamma \quad \gamma(\gamma-1) + A = 0 \\ r &\rightarrow \infty \quad \left(\frac{d^2}{dr^2} + \frac{A}{r^2} + \frac{B}{r} + C \right) R(r) \rightarrow \left(\frac{d^2}{dr^2} + C \right) R(r) = 0 \\ R(r) &\simeq e^{\lambda r} \quad \lambda^2 + C = 0 \end{aligned}$	$\begin{aligned} r &\rightarrow 0 \quad \left(\frac{d^2}{dr^2} + \frac{A}{r^2} + \frac{B}{r} + C \right) R(r) \rightarrow \left(\frac{d^2}{dr^2} + \frac{A}{r^2} \right) R(r) = 0 \\ R(r) &\simeq r^\gamma \quad \gamma(\gamma-1) + A = 0 \\ r &\rightarrow \infty \quad \left(\frac{d^2}{dr^2} + \frac{A}{r^2} + \frac{B}{r} + C \right) R(r) \rightarrow \left(\frac{d^2}{dr^2} + C \right) R(r) = 0 \\ R(r) &\simeq e^{\lambda r} \quad \lambda^2 + C = 0 \end{aligned}$	$\begin{aligned} r &\rightarrow 0 \quad \left(\frac{d^2}{dr^2} + \frac{A}{r^2} + \frac{B}{r} + C \right) R(r) \rightarrow \left(\frac{d^2}{dr^2} + \frac{A}{r^2} \right) R(r) = 0 \\ R(r) &\simeq r^\gamma \quad \gamma(\gamma-1) + A = 0 \\ r &\rightarrow \infty \quad \left(\frac{d^2}{dr^2} + \frac{A}{r^2} + \frac{B}{r} + C \right) R(r) \rightarrow \left(\frac{d^2}{dr^2} + C \right) R(r) = 0 \\ R(r) &\simeq e^{\lambda r} \quad \lambda^2 + C = 0 \end{aligned}$
gilmore-lie-groups_ E08844	265	■ 1.000	$[(j+1)j+2\gamma(j+1)]f_{j+1} + (2\lambda\gamma+2\lambda j+B)f_j = 0$	$[(j+1)j+2\gamma(j+1)]f_{j+1} + (2\lambda\gamma+2\lambda j+B)f_j = 0$	$[(j+1)j+2\gamma(j+1)]f_{j+1} + (2\lambda\gamma+2\lambda j+B)f_j = 0$
gilmore-lie-groups_ E08845	265	■ 1.000	$f(r) = \sum_{j=0}^{\infty} \frac{\Gamma(j+\gamma+(B/2\lambda))}{\Gamma(\gamma+(B/2\lambda))} \frac{\Gamma(2\gamma)}{\Gamma(j+2\gamma)} \frac{(-2\lambda r)^j}{j!}$	$f(r) = \sum_{j=0}^{\infty} \frac{\Gamma(j+\gamma+(B/2\lambda))}{\Gamma(\gamma+(B/2\lambda))} \frac{\Gamma(2\gamma)}{\Gamma(j+2\gamma)} \frac{(-2\lambda r)^j}{j!}$	$f(r) = \sum_{j=0}^{\infty} \frac{\Gamma(j+\gamma+(B/2\lambda))}{\Gamma(\gamma+(B/2\lambda))} \frac{\Gamma(2\gamma)}{\Gamma(j+2\gamma)} \frac{(-2\lambda r)^j}{j!}$
gilmore-lie-groups_ E08846	265	■ 1.000	$n + \frac{1}{2} + \sqrt{\left(\frac{1}{2}\right)^2 - A} = \frac{B}{2\sqrt{-C}}$	$n + \frac{1}{2} + \sqrt{\left(\frac{1}{2}\right)^2 - A} = \frac{B}{2\sqrt{-C}}$	$n + \frac{1}{2} + \sqrt{\left(\frac{1}{2}\right)^2 - A} = \frac{B}{2\sqrt{-C}}$
gilmore-lie-groups_ E08847	266	■ 1.000	$\begin{aligned} E(n, l) &= \frac{mc^2}{\sqrt{1 + \frac{\alpha^2}{\left(n + \frac{1}{2} + \sqrt{(l + \frac{1}{2})^2 - \alpha^2}\right)^2}}} \\ W(n, l) &= -\frac{1}{2} mc^2 \alpha^2 \frac{1}{(n+l+1)^2} \end{aligned}$	$\begin{aligned} E(n, l) &= \frac{mc^2}{\sqrt{1 + \frac{\alpha^2}{\left(n + \frac{1}{2} + \sqrt{(l + \frac{1}{2})^2 - \alpha^2}\right)^2}}} \\ W(n, l) &= -\frac{1}{2} mc^2 \alpha^2 \frac{1}{(n+l+1)^2} \end{aligned}$	$\begin{aligned} E(n, l) &= \frac{mc^2}{\sqrt{1 + \frac{\alpha^2}{\left(n + \frac{1}{2} + \sqrt{(l + \frac{1}{2})^2 - \alpha^2}\right)^2}}} \\ W(n, l) &= -\frac{1}{2} mc^2 \alpha^2 \frac{1}{(n+l+1)^2} \end{aligned}$
gilmore-lie-groups_ E08848	266	■ 1.000	$f(r) = \sum_{j=0}^n \frac{\Gamma(2\gamma)}{\Gamma(j+2\gamma)} \frac{n!}{(n-j)!j!} (2\lambda r)^j$	$f(r) = \sum_{j=0}^n \frac{\Gamma(2\gamma)}{\Gamma(j+2\gamma)} \frac{n!}{(n-j)!j!} (2\lambda r)^j$	$f(r) = \sum_{j=0}^n \frac{\Gamma(2\gamma)}{\Gamma(j+2\gamma)} \frac{n!}{(n-j)!j!} (2\lambda r)^j$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00849	266	■ 1.000	$\begin{aligned} E(n, l) = & \frac{m c^2}{\sqrt{1 + \left(\frac{\alpha}{N'} \right)^2}} \rightarrow m c^2 - m c^2 \frac{1}{2 N^2} \alpha^2 + m c^2 \left(\frac{3}{8 N^4} - \frac{1}{N^3 (2 l + 1)} \right) \alpha^4 \\ & + m c^2 \left(-\frac{5}{16 N^6} + \frac{3}{2 N^5 (2 l + 1)} - \frac{2 N + 3 (2 l + 1)}{2 N^4 (2 l + 1)^3} \right) \alpha^6 + m c^2 \\ & \times \left(\frac{35}{128 N^8} - \frac{15}{8 N^7 (2 l + 1)} + \frac{6 N + 9 (2 l + 1)}{4 N^6 (2 l + 1)^3} \right. \\ & \left. - \frac{2 N^2 + 3 N (2 l + 1) + 2 (2 l + 1)^2}{N^5 (2 l + 1)^5} \right) \alpha^8 + \mathcal{O}(\alpha^{10}) \end{aligned}$	$E(n, l) = \frac{m c^2}{\sqrt{1 + \left(\frac{\alpha}{N'} \right)^2}} \rightarrow m c^2 - m c^2 \frac{1}{2 N^2} \alpha^2 + m c^2 \left(\frac{3}{8 N^4} - \frac{1}{N^3 (2 l + 1)} \right) \alpha^4$ $+ m c^2 \left(-\frac{5}{16 N^6} + \frac{3}{2 N^5 (2 l + 1)} - \frac{2 N + 3 (2 l + 1)}{2 N^4 (2 l + 1)^3} \right) \alpha^6 + m c^2$ $\times \left(\frac{35}{128 N^8} - \frac{15}{8 N^7 (2 l + 1)} + \frac{6 N + 9 (2 l + 1)}{4 N^6 (2 l + 1)^3} \right.$ $\left. - \frac{2 N^2 + 3 N (2 l + 1) + 2 (2 l + 1)^2}{N^5 (2 l + 1)^5} \right) \alpha^8 + \mathcal{O}(\alpha^{10})$	$E(n, l) = \frac{m c^2}{\sqrt{1 + \left(\frac{\alpha}{N'} \right)^2}} \rightarrow m c^2 - m c^2 \frac{1}{2 N^2} \alpha^2 + m c^2 \left(\frac{3}{8 N^4} - \frac{1}{N^3 (2 l + 1)} \right) \alpha^4$ $+ m c^2 \left(-\frac{5}{16 N^6} + \frac{3}{2 N^5 (2 l + 1)} - \frac{2 N + 3 (2 l + 1)}{2 N^4 (2 l + 1)^3} \right) \alpha^6 + m c^2$ $\times \left(\frac{35}{128 N^8} - \frac{15}{8 N^7 (2 l + 1)} + \frac{6 N + 9 (2 l + 1)}{4 N^6 (2 l + 1)^3} \right.$ $\left. - \frac{2 N^2 + 3 N (2 l + 1) + 2 (2 l + 1)^2}{N^5 (2 l + 1)^5} \right) \alpha^8 + \mathcal{O}(\alpha^{10})$
gilmore-lie-groups_ E00850	266	■ 1.000	$\begin{array}{l} \text{\texttt{\textbackslash begin{array}{lll} \text{ { relativistic } } \& a(n, l)=} \\ \text{\texttt{\textbackslash sqrt{\left(N^{\text{\texttt{\textbackslash prime}}}\right)^2+\text{\texttt{\textbackslash alpha}^2}} } a_{\text{\texttt{B}}} } \\ \text{\texttt{\& N^{\text{\texttt{\textbackslash prime}}}=n+\frac{1}{2}+\sqrt{\left(l+\frac{1}{2}\right)^2-\text{\texttt{\textbackslash alpha}^2}} } \\ \text{\texttt{\textbackslash right)^2-\text{\texttt{\textbackslash alpha}^2}} } \text{\texttt{\textbackslash \text{ { nonrelativistic } } \& } } \\ \text{\texttt{a(n, l)=N a_{\text{\texttt{B}}} \& N=n+l+1 \text{\texttt{\textbackslash end{array}}} } \end{array}$	$\begin{array}{ll} \text{relativistic} & a(n, l) = \sqrt{(N')^2 + \alpha^2} a_B \\ \text{nonrelativistic} & a(n, l) = N a_B \end{array} \quad \begin{array}{ll} N' = n + \frac{1}{2} + \sqrt{(l + \frac{1}{2})^2 - \alpha^2} \\ N = n + l + 1 \end{array}$	$\begin{array}{ll} \text{relativistic} & a(n, l) = \sqrt{(N')^2 + \alpha^2} a_B \\ \text{nonrelativistic} & a(n, l) = N a_B \end{array} \quad \begin{array}{ll} N' = n + \frac{1}{2} + \sqrt{(l + \frac{1}{2})^2 - \alpha^2} \\ N = n + l + 1 \end{array}$
gilmore-lie-groups_ E00851	267	■ 1.000	$f = -\frac{K}{r^2} + \frac{C}{r^3}$	$f = -\frac{K}{r^2} + \frac{C}{r^3}$	$f = -\frac{K}{r^2} + \frac{C}{r^3}$
gilmore-lie-groups_ E00852	267	■ 1.000	$r = \frac{a \left(1 - \epsilon^2 \right)}{1 + \epsilon \cos(\alpha \theta)}$	$r = \frac{a \left(1 - \epsilon^2 \right)}{1 + \epsilon \cos(\alpha \theta)}$	$r = \frac{a \left(1 - \epsilon^2 \right)}{1 + \epsilon \cos(\alpha \theta)}$
gilmore-lie-groups_ E00853	268	■ 1.000	$H = m c^2 + \frac{p^2}{2 m} - \frac{K'}{r} + \frac{C'}{r^2}$	$H = m c^2 + \frac{p^2}{2 m} - \frac{K'}{r} + \frac{C'}{r^2}$	$H = m c^2 + \frac{p^2}{2 m} - \frac{K'}{r} + \frac{C'}{r^2}$
gilmore-lie-groups_ E00854	268	■ 1.000	$\omega = \frac{\partial}{\partial L} \left(\frac{1}{T} \int_0^T p(r) dt \right) = \frac{\partial}{\partial L} \left(\frac{m}{L T} \oint r^2 p(r) d\theta \right)$	$\omega = \frac{\partial}{\partial L} \left(\frac{1}{T} \int_0^T p(r) dt \right) = \frac{\partial}{\partial L} \left(\frac{m}{L T} \oint r^2 p(r) d\theta \right)$	$\omega = \frac{\partial}{\partial L} \left(\frac{1}{T} \int_0^T p(r) dt \right) = \frac{\partial}{\partial L} \left(\frac{m}{L T} \oint r^2 p(r) d\theta \right)$
gilmore-lie-groups_ E00855	268	■ 1.000	$\begin{array}{l} \text{\texttt{\textbackslash begin{array}{ll} \text{ { special relativity } } C=\frac{K}{L^2} \frac{1}{2 m^2} c^2 \text{\texttt{\& } } \omega=\frac{\pi K^2}{T L^2 c^2} \\ \text{\texttt{\textbackslash c^2}} \text{\texttt{\textbackslash \text{ { general relativity } } } C=6 \times \frac{K}{L^2} \frac{1}{2 m^2} c^2 \text{\texttt{\& } } \omega=\frac{6 \pi K^2}{T L^2 c^2} \\ \text{\texttt{\textbackslash end{array}}} \end{array}$	$\begin{array}{ll} \text{special relativity} & C = \frac{K L^2}{2 m^2 c^2} \quad \omega = \frac{\pi K^2}{T L^2 c^2} \\ \text{general relativity} & C = 6 \times \frac{K L^2}{2 m^2 c^2} \quad \omega = \frac{6 \pi K^2}{T L^2 c^2} \end{array}$	$\begin{array}{ll} \text{special relativity} & C = \frac{K L^2}{2 m^2 c^2} \quad \omega = \frac{\pi K^2}{T L^2 c^2} \\ \text{general relativity} & C = 6 \times \frac{K L^2}{2 m^2 c^2} \quad \omega = \frac{6 \pi K^2}{T L^2 c^2} \end{array}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0856	269	■ 1.000	$\left(-\frac{\partial}{\partial\theta}+\frac{\cos\theta}{\sin\theta}\frac{\partial}{\partial\phi}\right)P_{-l}^l(\theta)e^{-il\phi}=e^{-il\phi}\left(-\frac{\partial}{\partial\theta}+l\frac{\cos\theta}{\sin\theta}\frac{\partial}{\partial\phi}\right)P_{-l}^l(\theta)$	$\left(-\frac{\partial}{\partial\theta}+i\frac{\cos\theta}{\sin\theta}\frac{\partial}{\partial\phi}\right)P_{-l}^l(\theta)e^{-il\phi}=e^{-il\phi}\left(-\frac{\partial}{\partial\theta}+l\frac{\cos\theta}{\sin\theta}\frac{\partial}{\partial\phi}\right)P_{-l}^l(\theta)$	$\left(-\frac{\partial}{\partial\theta}+i\frac{\cos\theta}{\sin\theta}\frac{\partial}{\partial\phi}\right)P_{-l}^l(\theta)e^{-il\phi}=e^{-il\phi}\left(-\frac{\partial}{\partial\theta}+l\frac{\cos\theta}{\sin\theta}\frac{\partial}{\partial\phi}\right)P_{-l}^l(\theta)$
gilmore-lie-groups_ EQ0857	269	■ 1.000	$\int_0^\pi d\theta \sin\theta \int_0^{2\pi} d\phi N_l \sin^l\theta e^{-il\phi} ^2 = 1$	$\int_0^\pi d\theta \sin\theta \int_0^{2\pi} d\phi N_l \sin^l\theta e^{-il\phi} ^2 = 1$	$\int_0^\pi d\theta \sin\theta \int_0^{2\pi} d\phi N_l \sin^l\theta e^{-il\phi} ^2 = 1$
gilmore-lie-groups_ EQ0858	269	■ 1.000	$N_l=\sqrt{\frac{1}{4\pi}}\sqrt{\frac{(2l+1)!!}{(2l)!!}}$	$N_l=\sqrt{\frac{1}{4\pi}}\sqrt{\frac{(2l+1)!!}{(2l)!!}}$	$N_l=\sqrt{\frac{1}{4\pi}}\sqrt{\frac{(2l+1)!!}{(2l)!!}}$
gilmore-lie-groups_ EQ0859	269	■ 1.000	$N_l=\sqrt{\frac{2l+1}{2l}}N_{l-1}$	$N_l=\sqrt{\frac{2l+1}{2l}}N_{l-1}$	$N_l=\sqrt{\frac{2l+1}{2l}}N_{l-1}$
gilmore-lie-groups_ EQ0860	269	■ 1.000	$\frac{d}{dt}\left(\frac{\mathbf{r}}{r}\right)=\frac{\dot{\mathbf{r}}(\mathbf{r}\cdot\mathbf{r})-\mathbf{r}(\mathbf{r}\cdot\dot{\mathbf{r}})}{r^3}=-\frac{\mathbf{r}\times(\mathbf{r}\times\dot{\mathbf{r}})}{r^3}$	$\frac{d}{dt}\left(\frac{\mathbf{r}}{r}\right)=\frac{\dot{\mathbf{r}}(\mathbf{r}\cdot\mathbf{r})-\mathbf{r}(\mathbf{r}\cdot\dot{\mathbf{r}})}{r^3}=-\frac{\mathbf{r}\times(\mathbf{r}\times\dot{\mathbf{r}})}{r^3}$	$\frac{d}{dt}\left(\frac{\mathbf{r}}{r}\right)=\frac{\dot{\mathbf{r}}(\mathbf{r}\cdot\mathbf{r})-\mathbf{r}(\mathbf{r}\cdot\dot{\mathbf{r}})}{r^3}=-\frac{\mathbf{r}\times(\mathbf{r}\times\dot{\mathbf{r}})}{r^3}$
gilmore-lie-groups_ EQ0861	270	■ 1.000	$\frac{\mathbf{p}}{p_0}=\frac{\mathbf{u}}{1-\mathbf{u}\cdot\mathbf{w}}$	$\frac{\mathbf{p}}{p_0}=\frac{\mathbf{u}}{1-\mathbf{u}\cdot\mathbf{w}}$	$\frac{\mathbf{p}}{p_0}=\frac{\mathbf{u}}{1-\mathbf{u}\cdot\mathbf{w}}$
gilmore-lie-groups_ EQ0862	271	■ 1.000	$\frac{(a+b+1)!}{(a+\frac{1}{2})!(b+\frac{1}{2})!}=\frac{(a+b)!}{(a-\frac{1}{2})!(b+\frac{1}{2})!}+\frac{(a+b)!}{(a+\frac{1}{2})!(b-\frac{1}{2})!}$	$\frac{(a+b+1)!}{(a+\frac{1}{2})!(b+\frac{1}{2})!}=\frac{(a+b)!}{(a-\frac{1}{2})!(b+\frac{1}{2})!}+\frac{(a+b)!}{(a+\frac{1}{2})!(b-\frac{1}{2})!}$	$\frac{(a+b+1)!}{(a+\frac{1}{2})!(b+\frac{1}{2})!}=\frac{(a+b)!}{(a-\frac{1}{2})!(b+\frac{1}{2})!}+\frac{(a+b)!}{(a+\frac{1}{2})!(b-\frac{1}{2})!}$
gilmore-lie-groups_ EQ0863	271	■ 1.000	$\dim \mathcal{Y}^l(S^n)=\dim \mathcal{Y}^{l-1}(S^n)+\dim \mathcal{Y}^l(S^{n-1})$	$\dim \mathcal{Y}^l(S^n)=\dim \mathcal{Y}^{l-1}(S^n)+\dim \mathcal{Y}^l(S^{n-1})$	$\dim \mathcal{Y}^l(S^n)=\dim \mathcal{Y}^{l-1}(S^n)+\dim \mathcal{Y}^l(S^{n-1})$
gilmore-lie-groups_ EQ0864	271	■ 1.000	$\nabla^2=\left(\frac{1}{r^{D-1/2}}\frac{\partial}{\partial r}r^{D-1/2}\right)^2+\frac{\mathcal{L}^2}{r^2}$	$\nabla^2=\left(\frac{1}{r^{D-1/2}}\frac{\partial}{\partial r}r^{D-1/2}\right)^2+\frac{\mathcal{L}^2}{r^2}$	$\nabla^2=\left(\frac{1}{r^{D-1/2}}\frac{\partial}{\partial r}r^{D-1/2}\right)^2+\frac{\mathcal{L}^2}{r^2}$
gilmore-lie-groups_ EQ0865	271	■ 1.000	$\left(l+\frac{1}{2}\right)^2-\left(\frac{1}{2}\right)^2\rightarrow\left(l+\frac{D-2}{2}\right)^2-\left(\frac{D-2}{2}\right)^2$	$\left(l+\frac{1}{2}\right)^2-\left(\frac{1}{2}\right)^2\rightarrow\left(l+\frac{D-2}{2}\right)^2-\left(\frac{D-2}{2}\right)^2$	$\left(l+\frac{1}{2}\right)^2-\left(\frac{1}{2}\right)^2\rightarrow\left(l+\frac{D-2}{2}\right)^2-\left(\frac{D-2}{2}\right)^2$
gilmore-lie-groups_ EQ0866	271	■ 1.000	$\begin{array}{ll} \text{relativistic} & N'\rightarrow n+\frac{1}{2}+\sqrt{\left(l+\frac{1}{2}\right)^2+l(D-3)-\alpha^2} \\ \text{nonrelativistic} & N\rightarrow n+\frac{1}{2}+\sqrt{\left(l+\frac{1}{2}\right)^2+l(D-3)} \end{array}$	$\begin{array}{ll} \text{relativistic} & N'\rightarrow n+\frac{1}{2}+\sqrt{\left(l+\frac{1}{2}\right)^2+l(D-3)-\alpha^2} \\ \text{nonrelativistic} & N\rightarrow n+\frac{1}{2}+\sqrt{\left(l+\frac{1}{2}\right)^2+l(D-3)} \end{array}$	$\begin{array}{ll} \text{relativistic} & N'\rightarrow n+\frac{1}{2}+\sqrt{\left(l+\frac{1}{2}\right)^2+l(D-3)-\alpha^2} \\ \text{nonrelativistic} & N\rightarrow n+\frac{1}{2}+\sqrt{\left(l+\frac{1}{2}\right)^2+l(D-3)} \end{array}$
gilmore-lie-groups_ EQ0867	272	■ 1.000	$H=\sum_{i=1}^n\hbar\omega\left(a_i^\dagger a_i+\frac{1}{2}\right)$	$H=\sum_{i=1}^n\hbar\omega\left(a_i^\dagger a_i+\frac{1}{2}\right)$	$H=\sum_{i=1}^n\hbar\omega\left(a_i^\dagger a_i+\frac{1}{2}\right)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E00868 (15.1)	275	■ 1.000	$x^2+y^2+z^2-(c\,t)^2=x^{\prime 2}+y^{\prime 2}+z^{\prime 2}-\left(c\,t^{\prime }\right)^2=0$	$x^2+y^2+z^2-(ct)^2=x'^2+y'^2+z'^2-(ct')^2=0$	$x^2+y^2+z^2-(ct)^2=x'^2+y'^2+z'^2-(ct')^2=0$
gilmore-lie-groups_ E00870 (15.3)	275	■ 1.000	$\Lambda \rightarrowtail I_4+\epsilon \left[\begin{array}{cccc} 0 & +\theta _3 & -\theta _2 & ib_1 \\ -\theta _3 & 0 & +\theta _1 & ib_2 \\ +\theta _2 & -\theta _1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{array}\right]=I_4+\epsilon (\theta \cdot \mathbf{J}+\mathbf{b}\cdot \mathbf{K})$	$\Lambda \rightarrow I_4 + \epsilon \left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & ib_1 \\ -\theta_3 & 0 & +\theta_1 & ib_2 \\ +\theta_2 & -\theta_1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{array} \right] = I_4 + \epsilon (\theta \cdot \mathbf{J} + \mathbf{b} \cdot \mathbf{K})$	$\Lambda \rightarrow I_4 + \epsilon \left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & ib_1 \\ -\theta_3 & 0 & +\theta_1 & ib_2 \\ +\theta_2 & -\theta_1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{array} \right] = I_4 + \epsilon (\theta \cdot \mathbf{J} + \mathbf{b} \cdot \mathbf{K})$
gilmore-lie-groups_ E00871 (15.4)	275	■ 1.000	$\begin{aligned} \left\{ \left[J_i, J_j \right] \right\} &= -\epsilon_{ijk} J_k \\ \left\{ \left[J_i, K_j \right] \right\} &= -\epsilon_{ijk} K_k \\ \left\{ \left[K_i, K_j \right] \right\} &= +\epsilon_{ijk} J_k \end{aligned}$	$\begin{aligned} [J_i, J_j] &= -\epsilon_{ijk} J_k \\ [J_i, K_j] &= -\epsilon_{ijk} K_k \\ [K_i, K_j] &= +\epsilon_{ijk} J_k \end{aligned}$	$\begin{aligned} [J_i, J_j] &= -\epsilon_{ijk} J_k \\ [J_i, K_j] &= -\epsilon_{ijk} K_k \\ [K_i, K_j] &= +\epsilon_{ijk} J_k \end{aligned}$
gilmore-lie-groups_ E00872 (15.5)	275	■ 1.000	$\left(x_2-x_1\right)^2+\left(y_2-y_1\right)^2+\left(z_2-z_1\right)^2-\left(c\,t_2-c\,t_1\right)^2=\text {invariant }$	$(x_2-x_1)^2+(y_2-y_1)^2+(z_2-z_1)^2-(ct_2-ct_1)^2=\text{invariant}$	$(x_2-x_1)^2+(y_2-y_1)^2+(z_2-z_1)^2-(ct_2-ct_1)^2=\text{invariant}$
gilmore-lie-groups_ E00874 (15.7)	276	■ 1.000	$\begin{gathered} \left\{ \Lambda _2, a_2 \right\} \left\{ \Lambda _1, a_1 \right\} = \left\{ \Lambda _2 \Lambda _1, a_2 + \Lambda _2 a_1 \right\} \\ \left\{ \Lambda _2, a_2 \right\} \left\{ \Lambda _1, a_1 \right\} = \left\{ \Lambda _2 \Lambda _1, a_2 + \Lambda _2 a_1 \right\} \\ \left\{ \Lambda _2, a_2 \right\} \left\{ \Lambda _1, a_1 \right\} = \left\{ \Lambda _2 \Lambda _1, a_2 + \Lambda _2 a_1 \right\} \end{gathered}$	$\begin{aligned} \{ \Lambda_2, a_2 \} \{ \Lambda_1, a_1 \} &= \{ \Lambda_2 \Lambda_1, a_2 + \Lambda_2 a_1 \} \\ \{ I, a \} \{ \Lambda, 0 \} &= \{ \Lambda, a \} = \{ \Lambda, 0 \} \{ I, \Lambda^{-1} a \} \end{aligned}$	$\begin{aligned} \{ \Lambda_2, a_2 \} \{ \Lambda_1, a_1 \} &= \{ \Lambda_2 \Lambda_1, a_2 + \Lambda_2 a_1 \} \\ \{ I, a \} \{ \Lambda, 0 \} &= \{ \Lambda, a \} = \{ \Lambda, 0 \} \{ I, \Lambda^{-1} a \} \end{aligned}$
gilmore-lie-groups_ E00875 (15.8)	276	■ 1.000	$\Gamma^k(\{I, a\})=e^{ik\cdot a}$	$\Gamma^k(\{I, a\})=e^{ik\cdot a}$	$\Gamma^k(\{I, a\})=e^{ik\cdot a}$
gilmore-lie-groups_ E00876 (15.9)	276	■ 1.000	$\{I, a\} k\rangle = k'\rangle \langle k' \{I, a\} k\rangle = k'\rangle \delta(k' - k) \Gamma^k(\{I, a\}) = k\rangle e^{ik\cdot a}$	$\{I, a\} k\rangle = k'\rangle \langle k' \{I, a\} k\rangle = k'\rangle \delta(k' - k) \Gamma^k(\{I, a\}) = k\rangle e^{ik\cdot a}$	$\{I, a\} k\rangle = k'\rangle \langle k' \{I, a\} k\rangle = k'\rangle \delta(k' - k) \Gamma^k(\{I, a\}) = k\rangle e^{ik\cdot a}$
gilmore-lie-groups_ E00877 (15.10)	277	■ 1.000	$\begin{aligned} \mathbf{J}^{\{1\}} &= \frac{1}{2}(\mathbf{J} - i\mathbf{K}) \\ \mathbf{J}^{\{2\}} &= \frac{1}{2}(\mathbf{J} + i\mathbf{K}) \end{aligned}$	$\begin{aligned} \mathbf{J}^{(1)} &= \frac{1}{2}(\mathbf{J} - i\mathbf{K}) \\ \mathbf{J}^{(2)} &= \frac{1}{2}(\mathbf{J} + i\mathbf{K}) \end{aligned}$	$\begin{aligned} \mathbf{J}^{(1)} &= \frac{1}{2}(\mathbf{J} - i\mathbf{K}) \\ \mathbf{J}^{(2)} &= \frac{1}{2}(\mathbf{J} + i\mathbf{K}) \end{aligned}$
gilmore-lie-groups_ E00879 (15.12)	277	■ 1.000	$\begin{aligned} \operatorname{EXP}(\theta \cdot \mathbf{J} + \mathbf{b} \cdot \mathbf{K}) &= \operatorname{EXP} \left[(\theta + i\mathbf{b}) \cdot \mathbf{J}^{(1)} + (\theta - i\mathbf{b}) \cdot \mathbf{J}^{(2)} \right] \\ &= D^j \left[(\theta + i\mathbf{b}) \cdot \mathbf{J}^{(1)} \right] D^{j'} \left[(\theta - i\mathbf{b}) \cdot \mathbf{J}^{(2)} \right] \end{aligned}$	$\begin{aligned} \operatorname{EXP}(\theta \cdot \mathbf{J} + \mathbf{b} \cdot \mathbf{K}) &= \operatorname{EXP} \left[(\theta + i\mathbf{b}) \cdot \mathbf{J}^{(1)} + (\theta - i\mathbf{b}) \cdot \mathbf{J}^{(2)} \right] \\ &= D^j \left[(\theta + i\mathbf{b}) \cdot \mathbf{J}^{(1)} \right] D^{j'} \left[(\theta - i\mathbf{b}) \cdot \mathbf{J}^{(2)} \right] \end{aligned}$	$\begin{aligned} \operatorname{EXP}(\theta \cdot \mathbf{J} + \mathbf{b} \cdot \mathbf{K}) &= \operatorname{EXP} \left[(\theta + i\mathbf{b}) \cdot \mathbf{J}^{(1)} + (\theta - i\mathbf{b}) \cdot \mathbf{J}^{(2)} \right] \\ &= D^j \left[(\theta + i\mathbf{b}) \cdot \mathbf{J}^{(1)} \right] D^{j'} \left[(\theta - i\mathbf{b}) \cdot \mathbf{J}^{(2)} \right] \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0880 (15.13)	277	■ 1.000	$\begin{array}{ll} \mathbf{J}=\frac{i}{2}\sigma & \mathbf{J}=\frac{i}{2}\sigma \\ \mathbf{K}=-\frac{1}{2}\sigma & \mathbf{K}=-\frac{1}{2}\sigma \end{array}$	$\mathbf{J}=\frac{i}{2}\sigma \quad \mathbf{J}=\frac{i}{2}\sigma$ $\mathbf{K}=-\frac{1}{2}\sigma \quad \mathbf{K}=-\frac{1}{2}\sigma$	$\mathbf{J}=\frac{i}{2}\sigma \quad \mathbf{J}=\frac{i}{2}\sigma$ $\mathbf{K}=-\frac{1}{2}\sigma \quad \mathbf{K}=-\frac{1}{2}\sigma$
gilmore-lie-groups_ EQ0881 (15.14)	277	■ 1.000	$\begin{array}{cc} D^{\{j\} \emptyset} & D^{\{0\} j} \\ \mathbf{K}^{\{j\}}=i\mathbf{J}^{\{j\}} & \mathbf{K}^{\{j\}}=-i\mathbf{J}^{\{j\}} \end{array}$	$D^{j0} \quad D^{0j}$ $\mathbf{K}^{(j)}=i\mathbf{J}^{(j)} \quad \mathbf{K}^{(j)}=-i\mathbf{J}^{(j)}$	$D^{j0} \quad D^{0j}$ $\mathbf{K}^{(j)}=i\mathbf{J}^{(j)} \quad \mathbf{K}^{(j)}=-i\mathbf{J}^{(j)}$
gilmore-lie-groups_ EQ0883 (15.16)	278	■ 1.000	$D^{\{j\} j^{\prime}}(\theta \cdot \mathbf{J}+\mathbf{b} \cdot \mathbf{K})=\operatorname{EXP}\left[(\theta+i \mathbf{b}) \cdot \mathbf{J}^{(j)}\right] \operatorname{EXP}\left[(\theta-i \mathbf{b}) \cdot \mathbf{J}^{(j^{\prime})}\right]=D^{jj^{\prime}}(\Lambda)$	$D^{jj^{\prime}}(\theta \cdot \mathbf{J}+\mathbf{b} \cdot \mathbf{K})=\operatorname{EXP}\left[(\theta+i \mathbf{b}) \cdot \mathbf{J}^{(j)}\right] \operatorname{EXP}\left[(\theta-i \mathbf{b}) \cdot \mathbf{J}^{(j^{\prime})}\right]=D^{jj^{\prime}}(\Lambda)$	$D^{jj^{\prime}}(\theta \cdot \mathbf{J}+\mathbf{b} \cdot \mathbf{K})=\operatorname{EXP}\left[(\theta+i \mathbf{b}) \cdot \mathbf{J}^{(j)}\right] \operatorname{EXP}\left[(\theta-i \mathbf{b}) \cdot \mathbf{J}^{(j^{\prime})}\right]=D^{jj^{\prime}}(\Lambda)$
gilmore-lie-groups_ EQ0885 (15.18)	278	■ 1.000	$\begin{gathered} D^{\{j\} j^{\prime}}(\Lambda) \xrightarrow{\Lambda \downarrow SO(3)} D^j[SO(3)] \times D^{j^{\prime}}[SO(3)]=\sum_{j^{\prime \prime}} D^{jj^{\prime \prime}}[SO(3)] \\ j-j^{\prime} \leq j^{\prime \prime} \leq j+j^{\prime} \end{gathered}$	$D^{jj^{\prime}}(\Lambda) \xrightarrow{\Lambda \downarrow SO(3)} D^j[SO(3)] \times D^{j^{\prime}}[SO(3)]=\sum_{j^{\prime \prime}} D^{jj^{\prime \prime}}[SO(3)]$ $ j-j^{\prime} \leq j^{\prime \prime} \leq j+j^{\prime}$	$D^{jj^{\prime}}(\Lambda) \xrightarrow{\Lambda \downarrow SO(3)} D^j[SO(3)] \times D^{j^{\prime}}[SO(3)]=\sum_{j^{\prime \prime}} D^{jj^{\prime \prime}}[SO(3)]$ $ j-j^{\prime} \leq j^{\prime \prime} \leq j+j^{\prime}$
gilmore-lie-groups_ EQ0889 (15.22)	279	■ 1.000	$\begin{aligned} \{I, a\}[\{\Lambda, 0\} k\rangle] &= \{\Lambda, 0\} \{I, \Lambda^{-1} a\} k\rangle \\ &= [\{\Lambda, 0\} k\rangle] e^{i k \cdot \Lambda^{-1} a} \\ &= [\{\Lambda, 0\} k\rangle] e^{i \Lambda k \cdot a}= \Lambda k\rangle e^{i \Lambda k \cdot a} \end{aligned}$	$\{I, a\}[\{\Lambda, 0\} k\rangle]=\{\Lambda, 0\} \{I, \Lambda^{-1} a\} k\rangle$ $=[\{\Lambda, 0\} k\rangle] e^{i k \cdot \Lambda^{-1} a}$ $=[\{\Lambda, 0\} k\rangle] e^{i \Lambda k \cdot a}= \Lambda k\rangle e^{i \Lambda k \cdot a}$	$\{I, a\}[\{\Lambda, 0\} k\rangle]=\{\Lambda, 0\} \{I, \Lambda^{-1} a\} k\rangle$ $=[\{\Lambda, 0\} k\rangle] e^{i k \cdot \Lambda^{-1} a}$ $=[\{\Lambda, 0\} k\rangle] e^{i \Lambda k \cdot a}= \Lambda k\rangle e^{i \Lambda k \cdot a}$
gilmore-lie-groups_ EQ0893 (15.27)	280	■ 1.000	$\{I, a\} k ; \xi\rangle= k ; \xi\rangle e^{i k \cdot a}$	$\{I, a\} k ; \xi\rangle= k ; \xi\rangle e^{i k \cdot a}$	$\{I, a\} k ; \xi\rangle= k ; \xi\rangle e^{i k \cdot a}$
gilmore-lie-groups_ EQ0894 (15.28)	280	■ 1.000	$\begin{aligned} \{I, a\}[\{\Lambda, 0\} k ; \xi\rangle] &= \{\Lambda, 0\} \{I, \Lambda^{-1} a\} k ; \xi\rangle \\ &= [\{\Lambda, 0\} k ; \xi\rangle] e^{i k \cdot \Lambda^{-1} a} \\ &= [\{\Lambda, 0\} k ; \xi\rangle] e^{i \Lambda k \cdot a} \end{aligned}$	$\{I, a\}[\{\Lambda, 0\} k ; \xi\rangle]=\{\Lambda, 0\} \{I, \Lambda^{-1} a\} k ; \xi\rangle$ $=[\{\Lambda, 0\} k ; \xi\rangle] e^{i k \cdot \Lambda^{-1} a}$ $=[\{\Lambda, 0\} k ; \xi\rangle] e^{i \Lambda k \cdot a}$	$\{I, a\}[\{\Lambda, 0\} k ; \xi\rangle]=\{\Lambda, 0\} \{I, \Lambda^{-1} a\} k ; \xi\rangle$ $=[\{\Lambda, 0\} k ; \xi\rangle] e^{i k \cdot \Lambda^{-1} a}$ $=[\{\Lambda, 0\} k ; \xi\rangle] e^{i \Lambda k \cdot a}$
gilmore-lie-groups_ EQ0895 (15.29)	280	■ 1.000	$\{\Lambda, 0\} k ; \xi\rangle=\left \Lambda k ; \xi^{\prime}\right\rangle M_{\xi^{\prime} \xi}(\Lambda)$	$\{\Lambda, 0\} k ; \xi\rangle= \Lambda k ; \xi^{\prime}\rangle M_{\xi^{\prime} \xi}(\Lambda)$	$\{\Lambda, 0\} k ; \xi\rangle= \Lambda k ; \xi^{\prime}\rangle M_{\xi^{\prime} \xi}(\Lambda)$
gilmore-lie-groups_ EQ0896 (15.30)	280	■ 1.000	$k^{\prime}=\Lambda k$	$k^{\prime}=\Lambda k$	$k^{\prime}=\Lambda k$
gilmore-lie-groups_ EQ0899 (15.33)	281	■ 1.000	$\begin{aligned} H_k k^0 &=k^0 \\ C_k k^0 &=k \end{aligned}$	$H_k k^0=k^0$ $C_k k^0=k$	$H_k k^0=k^0$ $C_k k^0=k$
gilmore-lie-groups_ EQ0900 (15.34)	281	■ 1.000	$C_k k^0=k=\Lambda k^0$	$C_k k^0=k=\Lambda k^0$	$C_k k^0=k=\Lambda k^0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ9901 (15.35)	281	■ 1.000	$\left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & ib_1 \\ & -\theta_3 & 0 & +\theta_1 \\ +\theta_2 & -\theta_1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{array}\right] \left[\begin{array}{c} 0 \\ 0 \\ 1 \\ 0 \end{array}\right] = \left[\begin{array}{c} -\theta_2 \\ +\theta_1 \\ 0 \\ -ib_3 \end{array}\right] = \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array}\right]$	$\left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & ib_1 \\ & -\theta_3 & 0 & +\theta_1 \\ +\theta_2 & -\theta_1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{array}\right] \left[\begin{array}{c} 0 \\ 0 \\ 1 \\ 0 \end{array}\right] = \left[\begin{array}{c} -\theta_2 \\ +\theta_1 \\ 0 \\ -ib_3 \end{array}\right] = \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array}\right]$	$\left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & ib_1 \\ -\theta_3 & 0 & +\theta_1 & ib_2 \\ +\theta_2 & -\theta_1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{array}\right] \left[\begin{array}{c} 0 \\ 0 \\ 1 \\ 0 \end{array}\right] = \left[\begin{array}{c} -\theta_2 \\ +\theta_1 \\ 0 \\ -ib_3 \end{array}\right] = \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array}\right]$
gilmore-lie-groups_ EQ9904 (15.38)	282	■ 1.000	$\left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & -\theta_2 \\ & -i\theta_2 & -\theta_3 & 0 \\ +\theta_2 & -\theta_1 & 0 & +\theta_1 \\ i\theta_2 & -i\theta_1 & 0 & 0 \end{array}\right] = \sum_i \theta_i Y_i \quad \begin{array}{l} Y_1 = J_1 + K_2 \\ Y_2 = J_2 - K_1 \\ Y_3 = J_3 \end{array}$	$\left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & -i\theta_2 \\ -\theta_3 & 0 & +\theta_1 & i\theta_1 \\ +\theta_2 & -\theta_1 & 0 & 0 \\ i\theta_2 & -i\theta_1 & 0 & 0 \end{array}\right] = \sum_i \theta_i Y_i \quad \begin{array}{l} Y_1 = J_1 + K_2 \\ Y_2 = J_2 - K_1 \\ Y_3 = J_3 \end{array}$	$\left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & -i\theta_2 \\ -\theta_3 & 0 & +\theta_1 & i\theta_1 \\ +\theta_2 & -\theta_1 & 0 & 0 \\ i\theta_2 & -i\theta_1 & 0 & 0 \end{array}\right] = \sum_i \theta_i Y_i \quad \begin{array}{l} Y_1 = J_1 + K_2 \\ Y_2 = J_2 - K_1 \\ Y_3 = J_3 \end{array}$
gilmore-lie-groups_ EQ9907 (15.41)	283	■ 1.000	$\left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & ib_1 \\ & -\theta_3 & 0 & +\theta_1 \\ +\theta_2 & -\theta_1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{array}\right] \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array}\right] = \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array}\right]$	$\left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & ib_1 \\ & -\theta_3 & 0 & +\theta_1 \\ +\theta_2 & -\theta_1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{array}\right] \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array}\right] = \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array}\right]$	$\left[\begin{array}{cccc} 0 & +\theta_3 & -\theta_2 & ib_1 \\ -\theta_3 & 0 & +\theta_1 & ib_2 \\ +\theta_2 & -\theta_1 & 0 & ib_3 \\ -ib_1 & -ib_2 & -ib_3 & 0 \end{array}\right] \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array}\right] = \left[\begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array}\right]$
gilmore-lie-groups_ EQ9908 (15.42)	283	■ 1.000	$\begin{array}{l} \operatorname{H}_{k^0} \left k^0 \right\rangle ; \xi \rangle = \left H_{k^0} k^0 \right\rangle ; \xi' \rangle D_{\xi' \xi} \left(H_{k^0} \right) \\ = \left k^0 \right\rangle ; \xi' \rangle D_{\xi' \xi} \left(H_{k^0} \right) \end{array}$	$H_{k^0} \left k^0 \right\rangle ; \xi \rangle = \left H_{k^0} k^0 \right\rangle ; \xi' \rangle D_{\xi' \xi} \left(H_{k^0} \right) = \left k^0 \right\rangle ; \xi' \rangle D_{\xi' \xi} \left(H_{k^0} \right)$	$H_{k^0} \left k^0 \right\rangle ; \xi \rangle = \left H_{k^0} k^0 \right\rangle ; \xi' \rangle D_{\xi' \xi} \left(H_{k^0} \right) = \left k^0 \right\rangle ; \xi' \rangle D_{\xi' \xi} \left(H_{k^0} \right)$
gilmore-lie-groups_ EQ9909 (15.44)	284	■ 1.000	$\operatorname{EXP} \left(\theta_3 Y_3 + \theta_1 Y_1 + \theta_2 Y_2 \right) = e^{i \xi \theta_3}$	$\operatorname{EXP} \left(\theta_3 Y_3 + \theta_1 Y_1 + \theta_2 Y_2 \right) = e^{i \xi \theta_3}$	$\operatorname{EXP} \left(\theta_3 Y_3 + \theta_1 Y_1 + \theta_2 Y_2 \right) = e^{i \xi \theta_3}$
gilmore-lie-groups_ EQ9910 (15.45)	284	■ 1.000	$C_k \left k^0 \right\rangle ; \xi \rangle = \left k \right\rangle ; \xi \rangle$	$C_k \left k^0 \right\rangle ; \xi \rangle = \left k \right\rangle ; \xi \rangle$	$C_k \left k^0 \right\rangle ; \xi \rangle = \left k \right\rangle ; \xi \rangle$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0911 (15.46)	284	■1.000	$\begin{aligned} & \backslash\{\Lambda, a\} k; \xi\rangle = \{\Lambda, 0\} \{I, \Lambda^{-1} a\} k; \xi\rangle \\ & = \{\Lambda, 0\} k; \xi\rangle e^{ik \cdot \Lambda^{-1} a} \\ & = \{\Lambda, 0\} C_k k^0; \xi\rangle e^{i \Lambda k \cdot a} \\ & = \{\Lambda C_k, 0\} k^0; \xi\rangle e^{i \Lambda k \cdot a} \\ & = \{C_{k'} H_{k^0}, 0\} k^0; \xi\rangle e^{i \Lambda k \cdot a} \\ & = k'; \xi\rangle e^{i \xi \Theta} e^{i \Lambda k \cdot a} \end{aligned}$	$\begin{aligned} \{\Lambda, a\} k; \xi\rangle &= \{\Lambda, 0\} \{I, \Lambda^{-1} a\} k; \xi\rangle \\ &= \{\Lambda, 0\} k; \xi\rangle e^{ik \cdot \Lambda^{-1} a} \\ &= \{\Lambda, 0\} C_k k^0; \xi\rangle e^{i \Lambda k \cdot a} \\ &= \{\Lambda C_k, 0\} k^0; \xi\rangle e^{i \Lambda k \cdot a} \\ &= \{C_{k'} H_{k^0}, 0\} k^0; \xi\rangle e^{i \Lambda k \cdot a} \\ &= k'; \xi\rangle e^{i \xi \Theta} e^{i \Lambda k \cdot a} \end{aligned}$	$\begin{aligned} \{\Lambda, a\} k; \xi\rangle &= \{\Lambda, 0\} \{I, \Lambda^{-1} a\} k; \xi\rangle \\ &= \{\Lambda, 0\} k; \xi\rangle e^{ik \cdot \Lambda^{-1} a} \\ &= \{\Lambda, 0\} C_k k^0; \xi\rangle e^{i \Lambda k \cdot a} \\ &= \{\Lambda C_k, 0\} k^0; \xi\rangle e^{i \Lambda k \cdot a} \\ &= \{C_{k'} H_{k^0}, 0\} k^0; \xi\rangle e^{i \Lambda k \cdot a} \\ &= k'; \xi\rangle e^{i \xi \Theta} e^{i \Lambda k \cdot a} \end{aligned}$
gilmore-lie-groups_ EQ0912 (15.47)	284	■1.000	$C_{k'}^{-1} \Lambda C_k = H_{k^0} = \text{EXP}(\Theta J_3 + \theta_1 Y_1 + \theta_2 Y_2) \longrightarrow e^{i \xi \Theta}$	$C_{k'}^{-1} \Lambda C_k = H_{k^0} = \text{EXP}(\Theta J_3 + \theta_1 Y_1 + \theta_2 Y_2) \longrightarrow e^{i \xi \Theta}$	$C_{k'}^{-1} \Lambda C_k = H_{k^0} = \text{EXP}(\Theta J_3 + \theta_1 Y_1 + \theta_2 Y_2) \longrightarrow e^{i \xi \Theta}$
gilmore-lie-groups_ EQ0913 (15.48)	284	■1.000	$\begin{aligned} & k; \xi\rangle \quad k = \Lambda k^0 \\ & \quad k^0 = (0, 0, 1, \pm i) \end{aligned}$	$\begin{aligned} k; \xi\rangle \quad k &= \Lambda k^0 \\ k^0 &= (0, 0, 1, \pm i) \end{aligned}$	$\begin{aligned} k; \xi\rangle \quad k &= \Lambda k^0 \\ k^0 &= (0, 0, 1, \pm i) \end{aligned}$
gilmore-lie-groups_ EQ0915 (15.50)	285	■1.000	$\left\{ H_{k^0}, 0 \right\} k^0; \xi\rangle = k^0; \xi\rangle e^{i \xi \Theta}$	$\{H_{k^0}, 0\} k^0; \xi\rangle = k^0; \xi\rangle e^{i \xi \Theta}$	$\{H_{k^0}, 0\} k^0; \xi\rangle = k^0; \xi\rangle e^{i \xi \Theta}$
gilmore-lie-groups_ EQ0922 (15.56)	286	■1.000	$C_k k^0; \xi\rangle = k; \xi\rangle$	$C_k k^0; \xi\rangle = k; \xi\rangle$	$C_k k^0; \xi\rangle = k; \xi\rangle$
gilmore-lie-groups_ EQ0925 (15.59)	287	■1.000	$B_z(k)(0, 0, 1, i) = (0, 0, k, i k)$	$B_z(k)(0, 0, 1, i) = (0, 0, k, i k)$	$B_z(k)(0, 0, 1, i) = (0, 0, k, i k)$
gilmore-lie-groups_ EQ0928 (15.62)	287	■1.000	$\left\{ \mathbf{J} \cdot \mathbf{k} + 1 k_4 I_3 \right\} \left\langle k; \begin{array}{cc} 0 & 1 \\ 0 & \mu' \end{array} \middle \psi \right\rangle = 0$	$\left\{ \mathbf{J} \cdot \mathbf{k} + 1 k_4 I_3 \right\} \left\langle k; \begin{array}{cc} 0 & 1 \\ 0 & \mu' \end{array} \middle \psi \right\rangle = 0$	$\left\{ \mathbf{J} \cdot \mathbf{k} + 1 k_4 I_3 \right\} \left\langle k; \begin{array}{cc} 0 & 1 \\ 0 & \mu' \end{array} \middle \psi \right\rangle = 0$
gilmore-lie-groups_ EQ0931 (15.66)	288	■1.000	$-\frac{i}{c} \frac{\partial}{\partial t} (\mathbf{B} + i \mathbf{E}) - \nabla \times (\mathbf{B} + i \mathbf{E}) = 0$	$-\frac{i}{c} \frac{\partial}{\partial t} (\mathbf{B} + i \mathbf{E}) - \nabla \times (\mathbf{B} + i \mathbf{E}) = 0$	$-\frac{i}{c} \frac{\partial}{\partial t} (\mathbf{B} + i \mathbf{E}) - \nabla \times (\mathbf{B} + i \mathbf{E}) = 0$
gilmore-lie-groups_ EQ0935	290	■1.000	$\mathbf{k} \cdot \mathbf{v}(\mathbf{k}) = 0 \xrightarrow{\text{FT}^{-1}} \nabla \cdot (\mathbf{B} + i \mathbf{E}) = 0$	$\mathbf{k} \cdot \mathbf{v}(\mathbf{k}) = 0 \xrightarrow{\text{FT}^{-1}} \nabla \cdot (\mathbf{B} + i \mathbf{E}) = 0$	$\mathbf{k} \cdot \mathbf{v}(\mathbf{k}) = 0 \xrightarrow{\text{FT}^{-1}} \nabla \cdot (\mathbf{B} + i \mathbf{E}) = 0$
gilmore-lie-groups_ EQ0936 (15.70)	290	■1.000	$\left(\nabla \times + \frac{i}{c} \frac{\partial}{\partial t} \right) (\mathbf{B} + i \mathbf{E}) = \frac{1}{i} \frac{4\pi}{c} (\mathbf{J}_m + i \mathbf{J}_e)$	$\left(\nabla \times + \frac{i}{c} \frac{\partial}{\partial t} \right) (\mathbf{B} + i \mathbf{E}) = \frac{1}{i} \frac{4\pi}{c} (\mathbf{J}_m + i \mathbf{J}_e)$	$\left(\nabla \times + \frac{i}{c} \frac{\partial}{\partial t} \right) (\mathbf{B} + i \mathbf{E}) = \frac{1}{i} \frac{4\pi}{c} (\mathbf{J}_m + i \mathbf{J}_e)$
gilmore-lie-groups_ EQ0939	291	■1.000	$\frac{i}{c} \frac{\partial}{\partial t} \{ \nabla \cdot (\mathbf{B} + i \mathbf{E}) - 4\pi (\rho_m + i \rho_e) \} = 0$	$\frac{i}{c} \frac{\partial}{\partial t} \{ \nabla \cdot (\mathbf{B} + i \mathbf{E}) - 4\pi (\rho_m + i \rho_e) \} = 0$	$\frac{i}{c} \frac{\partial}{\partial t} \{ \nabla \cdot (\mathbf{B} + i \mathbf{E}) - 4\pi (\rho_m + i \rho_e) \} = 0$
gilmore-lie-groups_ EQ0941 (15.71)	291	■1.000	$-\frac{2i}{c} \frac{\partial}{\partial t} (\mathbf{G}_m + i \mathbf{G}_e) - \nabla \times (\mathbf{G}_m + i \mathbf{G}_e) = 0$	$-\frac{2i}{c} \frac{\partial}{\partial t} (\mathbf{G}_m + i \mathbf{G}_e) - \nabla \times (\mathbf{G}_m + i \mathbf{G}_e) = 0$	$-\frac{2i}{c} \frac{\partial}{\partial t} (\mathbf{G}_m + i \mathbf{G}_e) - \nabla \times (\mathbf{G}_m + i \mathbf{G}_e) = 0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E08943	291	■ 1.000	$\mathbf{J} \cdot \nabla = \left[\begin{array}{ccccc} +2\partial_0 & \sqrt{4}\partial_+ & 0 & 0 & 0 \\ \sqrt{4}\partial_- & +1\partial_0 & \sqrt{6}\partial_+ & 0 & 0 \\ 0 & \sqrt{6}\partial_- & 0\partial_0 & \sqrt{6}\partial_+ & 0 \\ 0 & 0 & \sqrt{6}\partial_- & -1\partial_0 & \sqrt{4}\partial_+ \\ 0 & 0 & 0 & \sqrt{6}\partial_- & -2\partial_0 \end{array} \right]$	$\mathbf{J} \cdot \nabla = \left[\begin{array}{ccccc} +2\partial_0 & \sqrt{4}\partial_+ & 0 & 0 & 0 \\ \sqrt{4}\partial_- & +1\partial_0 & \sqrt{6}\partial_+ & 0 & 0 \\ 0 & \sqrt{6}\partial_- & 0\partial_0 & \sqrt{6}\partial_+ & 0 \\ 0 & 0 & \sqrt{6}\partial_- & -1\partial_0 & \sqrt{4}\partial_+ \\ 0 & 0 & 0 & \sqrt{6}\partial_- & -2\partial_0 \end{array} \right]$	
gilmore-lie-groups_ E08945	292	■ 1.000	$F_{ij} = \left(\begin{array}{ccc} F_{11} & F_{12} & F_{13} \\ F_{21} & F_{22} & F_{23} \\ F_{31} & F_{32} & F_{33} \end{array} \right) = \left(\begin{array}{ccc} F_4 - \frac{1}{\sqrt{3}}F_5 & F_1 & F_3 \\ F_1 & -F_4 - \frac{1}{\sqrt{3}}F_5 & F_2 \\ F_3 & F_2 & +\frac{2}{\sqrt{3}}F_5 \end{array} \right)$	$F_{ij} = \left(\begin{array}{ccc} F_{11} & F_{12} & F_{13} \\ F_{21} & F_{22} & F_{23} \\ F_{31} & F_{32} & F_{33} \end{array} \right) = \left(\begin{array}{ccc} F_4 - \frac{1}{\sqrt{3}}F_5 & F_1 & F_3 \\ F_1 & -F_4 - \frac{1}{\sqrt{3}}F_5 & F_2 \\ F_3 & F_2 & +\frac{2}{\sqrt{3}}F_5 \end{array} \right)$	$F_{ij} = \left(\begin{array}{ccc} F_{11} & F_{12} & F_{13} \\ F_{21} & F_{22} & F_{23} \\ F_{31} & F_{32} & F_{33} \end{array} \right) = \left(\begin{array}{ccc} F_4 - \frac{1}{\sqrt{3}}F_5 & F_1 & F_3 \\ F_1 & -F_4 - \frac{1}{\sqrt{3}}F_5 & F_2 \\ F_3 & F_2 & +\frac{2}{\sqrt{3}}F_5 \end{array} \right)$
gilmore-lie-groups_ E08947	294	■ 1.000	$\oint \mathbf{E} \cdot d\mathbf{S} = \int 4\pi \rho dV$	$\oint \mathbf{E} \cdot d\mathbf{S} = \int 4\pi \rho dV$	$\oint \mathbf{E} \cdot d\mathbf{S} = \int 4\pi \rho dV$
gilmore-lie-groups_ E08948	294	■ 1.000	$4\pi a^2 \mathbf{E}(a) = 4\pi q$	$4\pi a^2 \mathbf{E}(a) = 4\pi q$	$4\pi a^2 \mathbf{E}(a) = 4\pi q$
gilmore-lie-groups_ E08949	294	■ 1.000	$\mathbf{E}(a) = \frac{q}{a^2} \frac{\mathbf{a}}{ \mathbf{a} }$	$\mathbf{E}(a) = \frac{q}{a^2} \frac{\mathbf{a}}{ \mathbf{a} }$	$\mathbf{E}(a) = \frac{q}{a^2} \frac{\mathbf{a}}{ \mathbf{a} }$
gilmore-lie-groups_ E08950	295	■ 1.000	$ \mathbf{E} = \frac{q}{R \sin \theta} = \frac{q}{R \sin (a / R)}$	$ \mathbf{E} = \frac{q}{R \sin \theta} = \frac{q}{R \sin (a / R)}$	$ \mathbf{E} = \frac{q}{R \sin \theta} = \frac{q}{R \sin (a / R)}$
gilmore-lie-groups_ E08951	295	■ 1.000	$ \mathbf{E} = \frac{q}{R \sin (a / R)} = \frac{q(a)}{a} \quad q(a) = q \left(\frac{a / R}{\sin (a / R)} \right)$	$ \mathbf{E} = \frac{q}{R \sin (a / R)} = \frac{q(a)}{a} \quad q(a) = q \left(\frac{a / R}{\sin (a / R)} \right)$	$ \mathbf{E} = \frac{q}{R \sin (a / R)} = \frac{q(a)}{a} \quad q(a) = q \left(\frac{a / R}{\sin (a / R)} \right)$
gilmore-lie-groups_ E08952	295	■ 1.000	$ds^2 = \frac{dr^2}{1 - kr^2} + r^2 \sum_{j=2}^n (\sin \theta_2 \sin \theta_3 \cdots \sin \theta_{j-1} d\theta_j)^2$	$ds^2 = \frac{dr^2}{1 - kr^2} + r^2 \sum_{j=2}^n (\sin \theta_2 \sin \theta_3 \cdots \sin \theta_{j-1} d\theta_j)^2$	$ds^2 = \frac{dr^2}{1 - kr^2} + r^2 \sum_{j=2}^n (\sin \theta_2 \sin \theta_3 \cdots \sin \theta_{j-1} d\theta_j)^2$
gilmore-lie-groups_ E08954	295	■ 1.000	$\oint \mathbf{E} \cdot d\mathbf{S} = \int \Omega \rho(x) dV$	$\oint \mathbf{E} \cdot d\mathbf{S} = \int \Omega \rho(x) dV$	$\oint \mathbf{E} \cdot d\mathbf{S} = \int \Omega \rho(x) dV$
gilmore-lie-groups_ E08955	295	■ 1.000	$ \mathbf{E} a^{n-1} = q$	$ \mathbf{E} a^{n-1} = q$	$ \mathbf{E} a^{n-1} = q$
gilmore-lie-groups_ E08956	296	■ 1.000	$d(a) = \int_0^a \frac{dr}{\sqrt{1 - kr^2}} \longrightarrow \begin{cases} \sinh^{-1} a & k = -1 \\ a & k = 0 \\ \sin^{-1} a & k = +1 \end{cases}$	$d(a) = \int_0^a \frac{dr}{\sqrt{1 - kr^2}} \longrightarrow \begin{cases} \sinh^{-1} a & k = -1 \\ a & k = 0 \\ \sin^{-1} a & k = +1 \end{cases}$	$d(a) = \int_0^a \frac{dr}{\sqrt{1 - kr^2}} \longrightarrow \begin{cases} \sinh^{-1} a & k = -1 \\ a & k = 0 \\ \sin^{-1} a & k = +1 \end{cases}$
gilmore-lie-groups_ E08958 (16.1)	299	■ 1.000	$\frac{dy}{dx} = g(x)$	$\frac{dy}{dx} = g(x)$	$\frac{dy}{dx} = g(x)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0959 (16.2)	299	■ 1.000	$y=G(x)=\int g(x) \, d x \quad(+ \, \text{additive constant})=G(x)+c$	$y = G(x) = \int g(x) dx \quad (+ \, \text{additive constant}) = G(x) + c$	$y = G(x) = \int g(x) \, dx \quad (+ \, \text{additive constant}) = G(x) + c$
gilmore-lie-groups_ EQ0960 (16.3)	299	■ 1.000	$e^{c \, \partial / \, \partial y} [y - G(x) = 0] = y + c - G(x) = 0$	$e^{c \partial / \partial y} [y - G(x) = 0] = y + c - G(x) = 0$	$e^{c \partial / \partial y} [y - G(x) = 0] = y + c - G(x) = 0$
gilmore-lie-groups_ EQ0963 (16.6)	301	■ 1.000	$\begin{array}{ll} x \rightarrow \bar{x}(\epsilon) = x + \epsilon \xi(x, y) + \mathcal{O}(\epsilon^2) & \bar{x}(\epsilon = 0) = x \\ y \rightarrow \bar{y}(\epsilon) = y + \epsilon \eta(x, y) + \mathcal{O}(\epsilon^2) & \bar{y}(\epsilon = 0) = y \\ p \rightarrow \bar{p}(\epsilon) = p + \epsilon \zeta(x, y, p) + \mathcal{O}(\epsilon^2) & \bar{p}(\epsilon = 0) = p \end{array}$	$\begin{array}{ll} x \rightarrow \bar{x}(\epsilon) = x + \epsilon \xi(x, y) + \mathcal{O}(\epsilon^2) & \bar{x}(\epsilon = 0) = x \\ y \rightarrow \bar{y}(\epsilon) = y + \epsilon \eta(x, y) + \mathcal{O}(\epsilon^2) & \bar{y}(\epsilon = 0) = y \\ p \rightarrow \bar{p}(\epsilon) = p + \epsilon \zeta(x, y, p) + \mathcal{O}(\epsilon^2) & \bar{p}(\epsilon = 0) = p \end{array}$	$\begin{array}{ll} x \rightarrow \bar{x}(\epsilon) = x + \epsilon \xi(x, y) + \mathcal{O}(\epsilon^2) & \bar{x}(\epsilon = 0) = x \\ y \rightarrow \bar{y}(\epsilon) = y + \epsilon \eta(x, y) + \mathcal{O}(\epsilon^2) & \bar{y}(\epsilon = 0) = y \\ p \rightarrow \bar{p}(\epsilon) = p + \epsilon \zeta(x, y, p) + \mathcal{O}(\epsilon^2) & \bar{p}(\epsilon = 0) = p \end{array}$
gilmore-lie-groups_ EQ0964 (16.7)	301	■ 1.000	$\bar{p} = \frac{d \bar{y}}{d \bar{x}} = \frac{d \bar{y} / d x}{d \bar{x} / d x} = \frac{p + \epsilon (\eta_x + \eta_y p)}{1 + \epsilon (\xi_x + \xi_y p)} \longrightarrow p + \epsilon [\eta_x + (\eta_y - \xi_x) p - \xi_y p^2]$	$\bar{p} = \frac{d \bar{y}}{d \bar{x}} = \frac{d \bar{y} / d x}{d \bar{x} / d x} = \frac{p + \epsilon (\eta_x + \eta_y p)}{1 + \epsilon (\xi_x + \xi_y p)} \longrightarrow p + \epsilon [\eta_x + (\eta_y - \xi_x) p - \xi_y p^2]$	$\bar{p} = \frac{d \bar{y}}{d \bar{x}} = \frac{d \bar{y} / d x}{d \bar{x} / d x} = \frac{p + \epsilon (\eta_x + \eta_y p)}{1 + \epsilon (\xi_x + \xi_y p)} \longrightarrow p + \epsilon [\eta_x + (\eta_y - \xi_x) p - \xi_y p^2]$
gilmore-lie-groups_ EQ0965 (16.8)	301	■ 1.000	$\zeta(x, y, p) = \eta^{(1)}(x, y, y^{(1)}) = \eta_x + (\eta_y - \xi_x) p - \xi_y p^2$	$\zeta(x, y, p) = \eta^{(1)}(x, y, y^{(1)}) = \eta_x + (\eta_y - \xi_x) p - \xi_y p^2$	$\zeta(x, y, p) = \eta^{(1)}(x, y, y^{(1)}) = \eta_x + (\eta_y - \xi_x) p - \xi_y p^2$
gilmore-lie-groups_ EQ0967 (16.10)	301	■ 1.000	$F(\bar{x}(\epsilon), \bar{y}(\epsilon), \bar{p}(\epsilon)) = e^{\epsilon X} F(x, y, p) = 0$	$F(\bar{x}(\epsilon), \bar{y}(\epsilon), \bar{p}(\epsilon)) = e^{\epsilon X} F(x, y, p) = 0$	$F(\bar{x}(\epsilon), \bar{y}(\epsilon), \bar{p}(\epsilon)) = e^{\epsilon X} F(x, y, p) = 0$
gilmore-lie-groups_ EQ0968 (16.11)	302	■ 1.000	$X = \xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} + \zeta \frac{\partial}{\partial p}$	$X = \xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} + \zeta \frac{\partial}{\partial p}$	$X = \xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} + \zeta \frac{\partial}{\partial p}$
gilmore-lie-groups_ EQ0969 (16.12)	302	■ 1.000	$F(x, y, p) = 0 \quad \text{and} \quad X F(x, y, p) = 0$	$F(x, y, p) = 0 \quad \text{and} \quad X F(x, y, p) = 0$	$F(x, y, p) = 0 \quad \text{and} \quad X F(x, y, p) = 0$
gilmore-lie-groups_ EQ0971 (16.14)	303	■ 1.000	$X F = 0 \quad X \left[\begin{array}{c} R(x, y) \\ S(x, y) \\ T(x, y, p) \end{array} \right] = \left[\begin{array}{c} 0 \\ 1 \\ 0 \end{array} \right]$	$X F = 0 \quad X \left[\begin{array}{c} R(x, y) \\ S(x, y) \\ T(x, y, p) \end{array} \right] = \left[\begin{array}{c} 0 \\ 1 \\ 0 \end{array} \right]$	$X F = 0 \quad X \left[\begin{array}{c} R(x, y) \\ S(x, y) \\ T(x, y, p) \end{array} \right] = \left[\begin{array}{c} 0 \\ 1 \\ 0 \end{array} \right]$
gilmore-lie-groups_ EQ0972 (16.15)	303	■ 1.000	$\left(\xi(x, y) \frac{\partial}{\partial x} + \eta(x, y) \frac{\partial}{\partial y} \right) S(x, y) = 1$	$\left(\xi(x, y) \frac{\partial}{\partial x} + \eta(x, y) \frac{\partial}{\partial y} \right) S(x, y) = 1$	$\left(\xi(x, y) \frac{\partial}{\partial x} + \eta(x, y) \frac{\partial}{\partial y} \right) S(x, y) = 1$
gilmore-lie-groups_ EQ0974 (16.17)	303	■ 1.000	$\frac{dx}{\xi(x, y)} = \frac{dy}{\eta(x, y)} = \frac{dp}{\zeta(x, y, p)}$	$\frac{dx}{\xi(x, y)} = \frac{dy}{\eta(x, y)} = \frac{dp}{\zeta(x, y, p)}$	$\frac{dx}{\xi(x, y)} = \frac{dy}{\eta(x, y)} = \frac{dp}{\zeta(x, y, p)}$
gilmore-lie-groups_ EQ0975 (16.18)	304	■ 1.000	$\frac{dS}{dR} = \frac{dS(x, y)}{dR(x, y)} = \frac{dS/dx}{dR/dx} = \frac{S_x + S_y p}{R_x + R_y p}$	$\frac{dS}{dR} = \frac{dS(x, y)}{dR(x, y)} = \frac{dS/dx}{dR/dx} = \frac{S_x + S_y p}{R_x + R_y p}$	$\frac{dS}{dR} = \frac{dS(x, y)}{dR(x, y)} = \frac{dS/dx}{dR/dx} = \frac{S_x + S_y p}{R_x + R_y p}$
gilmore-lie-groups_ EQ0977 (16.20)	304	■ 1.000	$S = \int f(R, -, T(R)) \, dR + c$	$S = \int f(R, -, T(R)) dR + c$	$S = \int f(R, -, T(R)) \, dR + c$
gilmore-lie-groups_ EQ0978 (16.21)	305	■ 1.000	$F(x, y, p) = xp + y - xy^2 = 0$	$F(x, y, p) = xp + y - xy^2 = 0$	$F(x, y, p) = xp + y - xy^2 = 0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ0979 (16.22)	305	■ 1.000	$\xi\left(p-y^2\right)+\eta(1-2 x y)+\left[\eta_x+\left(\eta_y-\xi_x\right) p-\xi_y p^2\right] x=0$	$\xi\left(p-y^2\right)+\eta(1-2 x y)+\left[\eta_x+\left(\eta_y-\xi_x\right) p-\xi_y p^2\right] x=0$	$\xi(p-y^2)+\eta(1-2xy)+[\eta_x+(\eta_y-\xi_x)p-\xi_y p^2]x=0$
gilmore-lie-groups_ EQ0980 (16.23)	306	■ 1.000	$\xi\left(-\frac{y}{x}\right)+\eta(1-2 x y)+\eta_x x+\left(\eta_y-\xi_x\right)\left(x y^2-y\right)-\xi_y\left(x y^4-2 y^3+\frac{y^2}{x}\right)=0$	$\xi\left(-\frac{y}{x}\right)+\eta(1-2 x y)+\eta_x x+\left(\eta_y-\xi_x\right)\left(x y^2-y\right)-\xi_y\left(x y^4-2 y^3+\frac{y^2}{x}\right)=0$	$\xi\left(-\frac{y}{x}\right)+\eta(1-2 x y)+\eta_x x+\left(\eta_y-\xi_x\right)\left(x y^2-y\right)-\xi_y\left(x y^4-2 y^3+\frac{y^2}{x}\right)=0$
gilmore-lie-groups_ EQ0982 (16.25)	306	■ 1.000	$\begin{aligned} & \xi\left(-\frac{y}{x}\right)+\eta(1-2 x y)+\eta_x x+\left(\eta_y-\xi_x\right)\left(x y^2-y\right)-\xi_y\left(x y^4-2 y^3+\frac{y^2}{x}\right) \\ & +\left(\xi_{00}+\xi_{10} x+\xi_{01} y\right)\left(-\frac{y}{x}\right)+\left(\eta_{00}+\eta_{10} x+\eta_{01} y\right)(1-2 x y)+\eta_{10} x \\ & +\left(\eta_{01}-\xi_{10}\right)\left(x y^2-y\right)-\xi_{01}\left(x y^4-2 y^3+\frac{y^2}{x}\right)=0 \end{aligned}$	$\begin{aligned} & \left(\xi_{00}+\xi_{10} x+\xi_{01} y\right)\left(-\frac{y}{x}\right)+\left(\eta_{00}+\eta_{10} x+\eta_{01} y\right)(1-2 x y)+\eta_{10} x \\ & +\left(\eta_{01}-\xi_{10}\right)\left(x y^2-y\right)-\xi_{01}\left(x y^4-2 y^3+\frac{y^2}{x}\right)=0 \end{aligned}$	$\begin{aligned} & \left(\xi_{00}+\xi_{10} x+\xi_{01} y\right)\left(-\frac{y}{x}\right)+\left(\eta_{00}+\eta_{10} x+\eta_{01} y\right)(1-2 x y)+\eta_{10} x \\ & +\left(\eta_{01}-\xi_{10}\right)\left(x y^2-y\right)-\xi_{01}\left(x y^4-2 y^3+\frac{y^2}{x}\right)=0 \end{aligned}$
gilmore-lie-groups_ EQ0984 (16.27)	307	■ 1.000	$X=x \frac{\partial}{\partial x}-y \frac{\partial}{\partial y}-2 p \frac{\partial}{\partial p}$	$X=x \frac{\partial}{\partial x}-y \frac{\partial}{\partial y}-2 p \frac{\partial}{\partial p}$	$X=x \frac{\partial}{\partial x}-y \frac{\partial}{\partial y}-2 p \frac{\partial}{\partial p}$
gilmore-lie-groups_ EQ0985 (16.28)	307	■ 1.000	$\left(x \frac{\partial}{\partial x}-y \frac{\partial}{\partial y}\right) S(x, y)=1$	$\left(x \frac{\partial}{\partial x}-y \frac{\partial}{\partial y}\right) S(x, y)=1$	$\left(x \frac{\partial}{\partial x}-y \frac{\partial}{\partial y}\right) S(x, y)=1$
gilmore-lie-groups_ EQ0986 (16.29)	307	■ 1.000	$\frac{d x}{x}=\frac{d y}{-y}=\frac{d p}{-2 p}$	$\frac{d x}{x}=\frac{d y}{-y}=\frac{d p}{-2 p}$	$\frac{d x}{x}=\frac{d y}{-y}=\frac{d p}{-2 p}$
gilmore-lie-groups_ EQ0988 (16.31)	307	■ 1.000	$F(x, y, p)=x p+y-x y^2=0 \longrightarrow e^S\left[\frac{T}{R}+1-R\right]=0$	$F(x, y, p)=x p+y-x y^2=0 \longrightarrow e^S\left[\frac{T}{R}+1-R\right]=0$	$F(x, y, p)=x p+y-x y^2=0 \longrightarrow e^S\left[\frac{T}{R}+1-R\right]=0$
gilmore-lie-groups_ EQ0989 (16.32)	308	■ 1.000	$\frac{d S}{d R}=\frac{d(\ln y)}{d(x y)}=\frac{p / y}{y+x p}=\frac{T e^S / R^2}{e^S+\left(R e^{-S}\right)\left(T e^{2 S} / R^2\right)}$	$\frac{d S}{d R}=\frac{d(\ln y)}{d(x y)}=\frac{p / y}{y+x p}=\frac{T e^S / R^2}{e^S+\left(R e^{-S}\right)\left(T e^{2 S} / R^2\right)}$	$\frac{d S}{d R}=\frac{d(\ln y)}{d(x y)}=\frac{p / y}{y+x p}=\frac{T e^S / R^2}{e^S+\left(R e^{-S}\right)\left(T e^{2 S} / R^2\right)}$
gilmore-lie-groups_ EQ0991 (16.34)	308	■ 1.000	$\frac{d S}{d R}=\frac{1}{R}-\frac{1}{R^2} \Longrightarrow S=\ln (R)+\frac{1}{R}+c$	$\frac{d S}{d R}=\frac{1}{R}-\frac{1}{R^2} \Longrightarrow S=\ln (R)+\frac{1}{R}+c$	$\frac{d S}{d R}=\frac{1}{R}-\frac{1}{R^2} \Longrightarrow S=\ln (R)+\frac{1}{R}+c$
gilmore-lie-groups_ EQ0992 (16.35)	309	■ 1.000	$y=\frac{-1}{x(c+\ln x)}$	$y=\frac{-1}{x(c+\ln x)}$	$y=\frac{-1}{x(c+\ln x)}$
gilmore-lie-groups_ EQ0994 (16.37)	310	■ 1.000	$X=\xi \frac{\partial}{\partial x}+\eta \frac{\partial}{\partial y}+\eta^{(1)} \frac{\partial}{\partial y^{(1)}}+\eta^{(2)} \frac{\partial}{\partial y^{(2)}}$	$X=\xi \frac{\partial}{\partial x}+\eta \frac{\partial}{\partial y}+\eta^{(1)} \frac{\partial}{\partial y^{(1)}}+\eta^{(2)} \frac{\partial}{\partial y^{(2)}}$	$X=\xi \frac{\partial}{\partial x}+\eta \frac{\partial}{\partial y}+\eta^{(1)} \frac{\partial}{\partial y^{(1)}}+\eta^{(2)} \frac{\partial}{\partial y^{(2)}}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ E09995 (16.38)	310	■ 1.000	$\begin{aligned} \frac{d^2\bar{y}}{d\bar{x}^2} &= \frac{d}{d\bar{x}} \left(\frac{d\bar{y}}{d\bar{x}} \right) = \frac{d}{d\bar{x}} \left(p + \epsilon \eta^{(1)} \right) = \frac{D^{(1)}(p + \epsilon \eta^{(1)})}{D^{(0)}(x + \epsilon \xi)} = \frac{y^{(2)} + \epsilon D^{(1)} \eta^{(1)}}{1 + \epsilon D^{(0)} \xi} \\ &= y^{(2)} + \epsilon \left(D^{(1)} \eta^{(1)} - y^{(2)} D^{(0)} \xi \right) \end{aligned} \quad (16.)$	$\frac{d^2\bar{y}}{d\bar{x}^2} = \frac{d}{d\bar{x}} \left(\frac{d\bar{y}}{d\bar{x}} \right) = \frac{d}{d\bar{x}} \left(p + \epsilon \eta^{(1)} \right) = \frac{D^{(1)}(p + \epsilon \eta^{(1)})}{D^{(0)}(x + \epsilon \xi)} = \frac{y^{(2)} + \epsilon D^{(1)} \eta^{(1)}}{1 + \epsilon D^{(0)} \xi}$	$\frac{d^2\bar{y}}{d\bar{x}^2} = \frac{d}{d\bar{x}} \left(\frac{d\bar{y}}{d\bar{x}} \right) = \frac{d}{d\bar{x}} (p + \epsilon \eta^{(1)}) = \frac{D^{(1)}(p + \epsilon \eta^{(1)})}{D^{(0)}(x + \epsilon \xi)} = \frac{y^{(2)} + \epsilon D^{(1)} \eta^{(1)}}{1 + \epsilon D^{(0)} \xi}$
gilmore-lie-groups_ E09996 (16.39)	310	■ 1.000	$\eta^{(2)}(x, y, y^{(1)}, y^{(2)}) = D^{(1)} \eta^{(1)} - y^{(2)} D^{(0)} \xi$	$\eta^{(2)}(x, y, y^{(1)}, y^{(2)}) = D^{(1)} \eta^{(1)} - y^{(2)} D^{(0)} \xi$	$\eta^{(2)}(x, y, y^{(1)}, y^{(2)}) = D^{(1)} \eta^{(1)} - y^{(2)} D^{(0)} \xi$
gilmore-lie-groups_ E09997 (16.40)	311	■ 1.000	$D^{(n)} = \frac{\partial}{\partial x} + \frac{dy}{dx} \frac{\partial}{\partial y} + \frac{dy^{(1)}}{dx} \frac{\partial}{\partial y^{(1)}} + \cdots + y^{(n+1)} \frac{\partial}{\partial y^{(n)}}$	$D^{(n)} = \frac{\partial}{\partial x} + \frac{dy}{dx} \frac{\partial}{\partial y} + \frac{dy^{(1)}}{dx} \frac{\partial}{\partial y^{(1)}} + \cdots + y^{(n+1)} \frac{\partial}{\partial y^{(n)}}$	$D^{(n)} = \frac{\partial}{\partial x} + \frac{dy}{dx} \frac{\partial}{\partial y} + \frac{dy^{(1)}}{dx} \frac{\partial}{\partial y^{(1)}} + \cdots + y^{(n+1)} \frac{\partial}{\partial y^{(n)}}$
gilmore-lie-groups_ E09998 (16.41)	311	■ 1.000	$\eta^{(2)} = \eta_{xx} + (2\eta_{xy} - \xi_{xx}) y' + (\eta_{yy} - 2\xi_{xy}) y'^2 - \xi_{yy} y'^3 + (\eta_y - 2\xi_x - 3\xi_{yy'}) y''$	$\eta^{(2)} = \eta_{xx} + (2\eta_{xy} - \xi_{xx}) y' + (\eta_{yy} - 2\xi_{xy}) y'^2 - \xi_{yy} y'^3 + (\eta_y - 2\xi_x - 3\xi_{yy'}) y''$	$\eta^{(2)} = \eta_{xx} + (2\eta_{xy} - \xi_{xx}) y' + (\eta_{yy} - 2\xi_{xy}) y'^2 - \xi_{yy} y'^3 + (\eta_y - 2\xi_x - 3\xi_{yy'}) y''$
gilmore-lie-groups_ E09999 (16.42)	311	■ 1.000	$F \left(x, y, y^{(1)}, y^{(2)} \right) = 0 \quad X \left(x, y, y^{(1)}, y^{(2)} \right) F \left(x, y, y^{(1)}, y^{(2)} \right) = 0$	$F \left(x, y, y^{(1)}, y^{(2)} \right) = 0 \quad X \left(x, y, y^{(1)}, y^{(2)} \right) F \left(x, y, y^{(1)}, y^{(2)} \right) = 0$	$F(x, y, y^{(1)}, y^{(2)}) = 0 \quad X(x, y, y^{(1)}, y^{(2)}) F(x, y, y^{(1)}, y^{(2)}) = 0$
gilmore-lie-groups_ E01000 (16.43)	312	■ 1.000	$\frac{dT}{dR} = \frac{dT/dx}{dR/dx} = \frac{T_x + T_y y^{(1)} + T_{y^{(1)}} y^{(2)}}{R_x + R_y y^{(1)}}$	$\frac{dT}{dR} = \frac{dT/dx}{dR/dx} = \frac{T_x + T_y y^{(1)} + T_{y^{(1)}} y^{(2)}}{R_x + R_y y^{(1)}}$	$\frac{dT}{dR} = \frac{dT/dx}{dR/dx} = \frac{T_x + T_y y^{(1)} + T_{y^{(1)}} y^{(2)}}{R_x + R_y y^{(1)}}$
gilmore-lie-groups_ E01001 (16.44)	312	■ 1.000	$\frac{dT}{dR} = \frac{2xy' + x^2 y''}{y + xy'}$	$\frac{dT}{dR} = \frac{2xy' + x^2 y''}{y + xy'}$	$\frac{dT}{dR} = \frac{2xy' + x^2 y''}{y + xy'}$
gilmore-lie-groups_ E01002 (16.45)	312	■ 1.000	$G \left(R, -, T, \frac{dT}{dR} \right) = 0$	$G \left(R, -, T, \frac{dT}{dR} \right) = 0$	$G \left(R, -, T, \frac{dT}{dR} \right) = 0$
gilmore-lie-groups_ E01003 (16.46)	312	■ 1.000	$\frac{dT}{dR} = \frac{x^2 y'' + 2xy'}{y + xy'} = g(xy, x^2 y') = g(R, T)$	$\frac{dT}{dR} = \frac{x^2 y'' + 2xy'}{y + xy'} = g(xy, x^2 y') = g(R, T)$	$\frac{dT}{dR} = \frac{x^2 y'' + 2xy'}{y + xy'} = g(xy, x^2 y') = g(R, T)$
gilmore-lie-groups_ E01004 (16.47)	313	■ 1.000	$X = \xi \frac{\partial}{\partial x} + \eta^{(0)} \frac{\partial}{\partial y^{(0)}} + \eta^{(1)} \frac{\partial}{\partial y^{(1)}} + \cdots + \eta^{(n)} \frac{\partial}{\partial y^{(n)}} = \xi \frac{\partial}{\partial x} + \sum_{j=0}^n \eta^{(j)} \frac{\partial}{\partial y^{(j)}}$	$X = \xi \frac{\partial}{\partial x} + \eta^{(0)} \frac{\partial}{\partial y^{(0)}} + \eta^{(1)} \frac{\partial}{\partial y^{(1)}} + \cdots + \eta^{(n)} \frac{\partial}{\partial y^{(n)}} = \xi \frac{\partial}{\partial x} + \sum_{j=0}^n \eta^{(j)} \frac{\partial}{\partial y^{(j)}}$	$X = \xi \frac{\partial}{\partial x} + \eta^{(0)} \frac{\partial}{\partial y^{(0)}} + \eta^{(1)} \frac{\partial}{\partial y^{(1)}} + \cdots + \eta^{(n)} \frac{\partial}{\partial y^{(n)}} = \xi \frac{\partial}{\partial x} + \sum_{j=0}^n \eta^{(j)} \frac{\partial}{\partial y^{(j)}}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ1023	319	■1.000	$\left[\begin{array}{c} i_x \\ i_y \\ i_t \end{array}\right]=\left[\begin{array}{cccc} x^2-y^2+t^2 & 2xy & 2xt & -xu \\ 2yx & -x^2+y^2+t^2 & 2yt & -yu \\ 2tx & 2ty & x^2+y^2+t^2 & -tu \end{array}\right]\left[\begin{array}{c} \partial_x \\ \partial_y \\ \partial_t \\ \partial_u \end{array}\right]$		
gilmore-lie-groups_ EQ1025	320	■1.000	$S=t^2\frac{\partial}{\partial t}+t\mathbf{x}\cdot\nabla-\frac{1}{4}(\mathbf{x}\cdot\mathbf{x}+2nt)u\frac{\partial}{\partial u}$		
gilmore-lie-groups_ EQ1026	320	■1.000	$I=\int_{t_1}^{t_2}L(t,x,\dot{x})dt\quad\delta I=0$		
gilmore-lie-groups_ EQ1027	320	■1.000	$\frac{d}{dt}\left(\frac{\partial L}{\partial\dot{x}_i}\right)-\frac{\partial L}{\partial x_i}=0$		
gilmore-lie-groups_ EQ1028	320	■1.000	$I=\int_{t_1'}^{t_2'}L(t',x',\dot{x}')dt'=\int_{t_1}^{t_2}L(t',x',\dot{x}')\frac{dt'}{dt}dt$		
gilmore-lie-groups_ EQ1029	320	■1.000	$\int_{t_1}^{t_2}\left(\xi\frac{\partial L}{\partial t}+\eta_i\frac{\partial L}{\partial x_i}+\eta_i^{(1)}\frac{\partial L}{\partial\dot{x}_i}+\frac{d\xi}{dt}L\right)dt=0$		
gilmore-lie-groups_ EQ1030	320	■1.000	$\frac{d}{dt}\left[\xi L+(\eta_i-\xi\dot{x}_i)L_{\dot{x}_i}\right]=0$		
gilmore-lie-groups_ EQ1031	321	■1.000	$\mathbf{v}=\sum_{i=1}^p\xi^i(x,u)\frac{\partial}{\partial x^i}+\sum_{\alpha=1}^q\phi^\alpha(x,u)\frac{\partial}{\partial u^\alpha}$		
gilmore-lie-groups_ EQ1032	321	■1.000	$P^i=\xi^i\mathcal{L}+\sum_{\alpha=1}^q\phi^\alpha(x,u)\frac{\partial\mathcal{L}}{\partial u_i^\alpha}-\sum_{\alpha=1}^q\sum_{j=1}^p\xi^ju_j^\alpha\frac{\partial\mathcal{L}}{\partial u_i^\alpha}$		
gilmore-lie-groups_ EQ1033	321	■1.000	$\nabla P=\operatorname{div}P=\frac{\partial P^i}{\partial x^i}=0$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
gilmore-lie-groups_ EQ1035	321	■ 1.000	$\left\langle g \left \begin{array}{c} \lambda \\ \mu \nu \end{array} \right. \right\rangle = \sqrt{\frac{\dim \lambda}{\text{Vol}(G)}} \Gamma_{\mu \nu}^{\lambda}(g) \quad \left\langle \begin{array}{c} \lambda \\ \mu \nu \end{array} \left g \right. \right\rangle = \sqrt{\frac{\dim \lambda}{\text{Vol}(G)}} \Gamma_{\mu \nu}^{\lambda*}(g)$	$\left\langle g \left \begin{array}{c} \lambda \\ \mu \nu \end{array} \right. \right\rangle = \sqrt{\frac{\dim \lambda}{\text{Vol}(G)}} \Gamma_{\mu \nu}^{\lambda}(g) \quad \left\langle \begin{array}{c} \lambda \\ \mu \nu \end{array} \left g \right. \right\rangle = \sqrt{\frac{\dim \lambda}{\text{Vol}(G)}} \Gamma_{\mu \nu}^{\lambda*}(g)$	$\left\langle g \left \begin{array}{c} \lambda \\ \mu \nu \end{array} \right. \right\rangle = \sqrt{\frac{\dim \lambda}{\text{Vol}(G)}} \Gamma_{\mu \nu}^{\lambda}(g) \quad \left\langle \begin{array}{c} \lambda \\ \mu \nu \end{array} \left g \right. \right\rangle = \sqrt{\frac{\dim \lambda}{\text{Vol}(G)}} \Gamma_{\mu \nu}^{\lambda*}(g)$