

0707.4470: residuals

equations measured 2026-09-05T10:57:49Z against the model of 2026-09-02

Corrected (1)

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0707.4470_F00175 (was)	29	—	g	g	Given a discrete 1-form A and dual source 3-form $\mathcal{J}$, define the discrete
0707.4470_F00175 (now)	29	—	J	$\mathcal{J}$	Given a discrete 1-form A and dual source 3-form $\mathcal{J}$, define the discrete
basis: source / source+ink					
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value					
Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Flagged, not acted on

Identifier	Page	Conf.	LaTeX source	Rendered	Image
0707.4470_E00035	20	■ 0.705 ● C +22	$\begin{aligned} & \frac{\left(D_x\right)_{k+\frac{1}{2}, l, m}^{n+\frac{1}{2}} - \left(D_x\right)_{k+\frac{1}{2}, l, m}^{n-\frac{1}{2}}}{\Delta t} \\ & \frac{\left(H_z\right)_{k+\frac{1}{2}, l+\frac{1}{2}, m}^n - \left(H_z\right)_{k+\frac{1}{2}, l-\frac{1}{2}, m}^n}{\Delta y} - \frac{\left(H_y\right)_{k+\frac{1}{2}, l, m+\frac{1}{2}}^n - \left(H_y\right)_{k+\frac{1}{2}, l, m-\frac{1}{2}}^n}{\Delta z} - J_x\left _{k+\frac{1}{2}, l, m}^n\right. \\ & \frac{\left(D_y\right)_{k, l+\frac{1}{2}, m}^{n+\frac{1}{2}} - \left(D_y\right)_{k, l+\frac{1}{2}, m}^{n-\frac{1}{2}}}{\Delta t} \\ & \frac{\left(H_x\right)_{k, l+\frac{1}{2}, m+\frac{1}{2}}^n - \left(H_x\right)_{k, l+\frac{1}{2}, m-\frac{1}{2}}^n}{\Delta z} - \frac{\left(H_z\right)_{k+\frac{1}{2}, l+\frac{1}{2}, m}^n - \left(H_z\right)_{k-\frac{1}{2}, l+\frac{1}{2}, m}^n}{\Delta x} - J_y\left _{k, l+\frac{1}{2}, m}^n\right. \\ & \frac{\left(D_z\right)_{k, l, m+\frac{1}{2}}^{n+\frac{1}{2}} - \left(D_z\right)_{k, l, m+\frac{1}{2}}^{n-\frac{1}{2}}}{\Delta t} \\ & \frac{\left(H_y\right)_{k+\frac{1}{2}, l, m+\frac{1}{2}}^n - \left(H_y\right)_{k-\frac{1}{2}, l, m+\frac{1}{2}}^n}{\Delta x} - \frac{\left(H_x\right)_{k, l+\frac{1}{2}, m+\frac{1}{2}}^n - \left(H_x\right)_{k, l-\frac{1}{2}, m+\frac{1}{2}}^n}{\Delta y} - J_z\left _{k, l, m+\frac{1}{2}}^n\right. \end{aligned}$	$\begin{aligned} & \frac{\left(D_x\right)_{k+\frac{1}{2}, l, m}^{n+\frac{1}{2}} - \left(D_x\right)_{k+\frac{1}{2}, l, m}^{n-\frac{1}{2}}}{\Delta t} \\ & \frac{\left(H_z\right)_{k+\frac{1}{2}, l+\frac{1}{2}, m}^n - \left(H_z\right)_{k+\frac{1}{2}, l-\frac{1}{2}, m}^n}{\Delta y} - \frac{\left(H_y\right)_{k+\frac{1}{2}, l, m+\frac{1}{2}}^n - \left(H_y\right)_{k+\frac{1}{2}, l, m-\frac{1}{2}}^n}{\Delta z} - J_x\left _{k+\frac{1}{2}, l, m}^n\right. \\ & \frac{\left(D_y\right)_{k, l+\frac{1}{2}, m}^{n+\frac{1}{2}} - \left(D_y\right)_{k, l+\frac{1}{2}, m}^{n-\frac{1}{2}}}{\Delta t} \\ & \frac{\left(H_x\right)_{k, l+\frac{1}{2}, m+\frac{1}{2}}^n - \left(H_x\right)_{k, l+\frac{1}{2}, m-\frac{1}{2}}^n}{\Delta z} - \frac{\left(H_z\right)_{k+\frac{1}{2}, l+\frac{1}{2}, m}^n - \left(H_z\right)_{k-\frac{1}{2}, l+\frac{1}{2}, m}^n}{\Delta x} = J_y\left _{k, l+\frac{1}{2}, m}^n\right. \\ & \frac{\left(H_y\right)_{k+\frac{1}{2}, l, m+\frac{1}{2}}^n - \left(H_y\right)_{k-\frac{1}{2}, l, m+\frac{1}{2}}^n}{\Delta x} - \frac{\left(H_x\right)_{k, l+\frac{1}{2}, m+\frac{1}{2}}^n - \left(H_x\right)_{k, l-\frac{1}{2}, m+\frac{1}{2}}^n}{\Delta y} - J_z\left _{k, l, m+\frac{1}{2}}^n\right. \end{aligned}$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

9 more flagged rows below the band, stated as a count: 0 component, 9 weak; confidence high 7, mid 1, low 1, none 0