

cardona-qft-methods: residuals

equations measured 2026-09-05T11:34:14Z against the model of 2026-09-03

Unresolved

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_E00654 (4)	261	■0.167	$\begin{aligned} &\backslash\mathrm{phi_}\{\backslash\mathrm{theta}\}(x) \backslash\mathrm{phi_}\{\backslash\mathrm{theta}\}(y)=e^{\backslash\mathrm{frac}\{1\}{2} \backslash\mathrm{frac}\{ \\ &\backslash\mathrm{partial}\}{\backslash\mathrm{partial} \ x^{\backslash\mathrm{mu}}}\} \backslash\mathrm{theta}^{\backslash\mathrm{mu} \ \backslash\mathrm{nu}} \backslash\mathrm{frac}\{ \\ &\backslash\mathrm{partial}\}{\backslash\mathrm{partial} \ y^{\backslash\mathrm{nu}}}\} \backslash\mathrm{phi_}\{0\}(x) \backslash\mathrm{phi_}\{0\}(y) \ e^{\backslash\mathrm{frac}\{1\}{2} \backslash\mathrm{left}\{\backslash\mathrm{frac}\{\backslash\mathrm{overleftarrow}\{\backslash\mathrm{delta}\}\}{\backslash\mathrm{partial} \\ &x^{\backslash\mathrm{mu}}\}+\backslash\mathrm{frac}\{\backslash\mathrm{overleftarrow}\{\square\}\}{\backslash\mathrm{partial} \ y^{\backslash\mathrm{mu}}}\} \\ &\backslash\mathrm{right}\} \backslash\mathrm{theta}^{\backslash\mathrm{mu} \ \backslash\mathrm{nu}} \ P_{\backslash\mathrm{nu}}\} \ \backslash\mathrm{neq} \ \backslash\mathrm{phi_}\{0\}(x) \\ &\backslash\mathrm{phi_}\{0\}(y) \ . \end{aligned}$	(not rendered)	$\phi_{\theta}(x)\phi_{\theta}(y)=e^{-\frac{1}{2}\frac{\partial}{\partial x^{\mu}}\theta^{\mu\nu}\frac{\partial}{\partial y^{\nu}}}\phi_0(x)\phi_0(y)e^{\frac{1}{2}\left(\frac{\overleftarrow{\partial}}{\partial x^{\mu}}+\frac{\overleftarrow{\partial}}{\partial y^{\mu}}\right)\theta^{\mu\nu}P_{\nu}}\neq\phi_0(x)\phi_0(y).$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Flagged, not acted on

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0677	46	■ 0.150 ● C +44	\left.\begin{array}{cccccc} 1 & \rightarrow & \operatorname{Inn}(\mathcal{A}) & \rightarrow & \operatorname{Aut}(\mathcal{A}) & \rightarrow & \operatorname{Aut}(\mathcal{A})/\operatorname{Inn}(\mathcal{A}) & \rightarrow & 1, \\ 1 & \rightarrow & \mathcal{G} & \rightarrow & \mathcal{G} \rtimes \operatorname{Diff}(M) & \rightarrow & \operatorname{Diff}(M) & \rightarrow & 1. \end{array}\right\}	(not rendered)	$\begin{array}{lcl} AdS_5 : & -R^2 = -Z_0^2 + Z_1^2 + Z_2^2 + Z_3^2 + Z_4^2 - Z_5^2, \\ S^5 : & R^2 = Y_1^2 + Y_2^2 + Y_3^2 + Y_4^2 + Y_5^2 + Y_6^2. \end{array}$
cardona-qft-methods_ EQ0748	284	■ 0.547 ● C -27	\begin{aligned} A d S_{\{5\}} & : & R^2 & = -Z_{\{0\}}^2 + Z_{\{1\}}^2 + Z_{\{2\}}^2 + Z_{\{3\}}^2 + Z_{\{4\}}^2 - Z_{\{5\}}^2, \\ S^5 & : & R^2 & = Y_{\{1\}}^2 + Y_{\{2\}}^2 + Y_{\{3\}}^2 + Y_{\{4\}}^2 + Y_{\{5\}}^2 + Y_{\{6\}}^2. \end{aligned}		
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

88 more flagged rows below the band, stated as a count: 18 component, 70 weak; confidence high 60, mid 14, low 14, none 0

Low confidence

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ9908 [conf 0.080]	350	0.080 U +5	$\begin{array}{rlc} P_{\mathrm{g}} M\colon {}_{\mathrm{1}}\times {}_{\mathrm{0}}P_{\mathrm{h}} M & \longrightarrow & P_{\mathrm{g}} M\times P_{\mathrm{h}} M \\ & \downarrow & \downarrow \\ & M & \downarrow \\ & \downarrow & \downarrow \times M \end{array}$	$\begin{array}{ccc} P_g M \colon {}_1 \times {}_0 P_h M & \longrightarrow & P_g M \times P_h M \\ \downarrow e_0 \circ \pi_1 & & \downarrow e_0 \times e_0 \\ M & \xrightarrow{\Delta_g} & M \times M \end{array}$	
Conf. MathPix confidence: Residual render vs scan (inkdrill):					