

Display equations

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_ E08282 [conf 0.001]	291	■ 0.001	G_{1}=\begin{array}{ c } \hline \text{ 李 } \\ \hline f=g \\ \end{array} \quad G_{2}=\begin{array}{ c } \hline f \\ \stackrel{\text{Q}}{\bullet} \\ a \\ \hline f=a \\ \end{array} \quad G_{3}=\begin{array}{ ccc } \bullet & \xrightarrow{f} & \bullet \\ g \downarrow & & \downarrow h \\ \bullet & \xrightarrow{i} & \bullet \\ f \circ h = g \circ i \\ \hline \end{array} \quad G_{4}=\begin{array}{ ccc } \bullet & \xrightarrow{f} & \bullet \\ g \downarrow & & \downarrow h \\ \bullet & & \bullet \\ \hline \text{no equations} \\ \hline \end{array}	(not rendered)	
fong-spivak-invitation_ E08962 [conf 0.001]	113	■ 0.001	I=\left\lvert\lvert\begin{array}{ccc} 1 & a & 2 \\ \downarrow & \downarrow & \\ \triangleright & 3 & \\ & 4 & b \end{array}\right\rvert.	$I = \left \begin{array}{ccc} 1 & a & 2 \\ & \downarrow & \downarrow \\ & \triangleright & 3 \\ & & 4 \\ & & b \end{array} \right $	
fong-spivak-invitation_ E08188 [conf 0.013]	273	■ 0.013	G=\left \begin{array}{ccc} 1 & & 2 \\ & a^{\downarrow}e^{\downarrow}_c & \\ & & v_3 \end{array}\right	$G = \left \begin{array}{ccc} 1 & & 2 \\ & a^{\downarrow}e^{\downarrow}_c & \\ & & v_3 \end{array} \right $	
fong-spivak-invitation_ E08135 [conf 0.019]	215	■ 0.019	\begin{aligned} \mu_X &:= \left(X + X \xrightarrow{\left(\mathrm{id}_X, \mathrm{id}_X \right)} X \xrightarrow{\mathrm{id}_X} X \right) \\ \eta_X &:= \left(\emptyset \xrightarrow{!_X} X \xleftarrow{\mathrm{id}_X} X \right) \\ \delta_X &:= \left(X \xrightarrow{\mathrm{id}_X} X \xrightarrow{\left(\mathrm{id}_X, \mathrm{id}_X \right)} X + X \right) \\ \epsilon_X &:= \left(X \xrightarrow{\mathrm{id}_X} X \xleftarrow{!_X} \emptyset \right) \end{aligned}	$\begin{aligned} \mu_X &:= \left(X + X \xrightarrow{[\mathrm{id}_X, \mathrm{id}_X]} X \xleftarrow{\mathrm{id}_X} X \right) \\ \eta_X &:= \left(\emptyset \xrightarrow{!_X} X \xleftarrow{\mathrm{id}_X} X \right) \\ \delta_X &:= \left(X \xrightarrow{\mathrm{id}_X} X \xrightarrow{[\mathrm{id}_X, \mathrm{id}_X]} X + X \right) \\ \epsilon_X &:= \left(X \xrightarrow{\mathrm{id}_X} X \xleftarrow{!_X} \emptyset \right) \end{aligned}$	$\begin{aligned} \mu_X &:= \left(X + X \xrightarrow{[\mathrm{id}_X, \mathrm{id}_X]} X \xleftarrow{\mathrm{id}_X} X \right) \\ \eta_X &:= \left(\emptyset \xrightarrow{!_X} X \xleftarrow{\mathrm{id}_X} X \right) \\ \delta_X &:= \left(X \xrightarrow{\mathrm{id}_X} X \xleftarrow{[\mathrm{id}_X, \mathrm{id}_X]} X + X \right) \\ \epsilon_X &:= \left(X \xrightarrow{\mathrm{id}_X} X \xleftarrow{!_X} \emptyset \right) \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_EQ0118	184	■ 0.327	$G_{\mathcal{R}} := \{\supset-, \circ-, \subset, \cdots\} \cup \{-\mathbf{a}- \mid \mathbf{a} \in R\},$	$G_R := \{\supset-, \circ-, \subset, \cdots\} \cup \{-\mathbf{a}- \mid \mathbf{a} \in R\},$	$G_R := \left\{ \supset-, \circ-, \subset, \cdots \right\} \cup \left\{ -\mathbf{a}- \mid a \in R \right\},$
fong-spivak-invitation_EQ0037	74	■ 0.341	$X := \operatorname{Ob}(\mathcal{X}), \quad \text{d}(x, y) := \mathcal{X}(x, y).$	$X := \operatorname{Ob}(\mathcal{X}), \quad d(x, y) := \mathcal{X}(x, y).$	$X := \operatorname{Ob}(\mathcal{X}), \quad d(x, y) := \mathcal{X}(x, y).$
fong-spivak-invitation_EQ0260	332	■ 0.372	$\bigcup_{i \in I} A_i = \bigcup_{i \in I} \left(B_i \cap Y \right) \cap Y \in \mathbf{Op}_{? \cap Y}.$	$\bigcup_{i \in I} A_i = \bigcup_{i \in I} (B_i \cap Y) = \left(\bigcup_{i \in I} B_i \right) \cap Y \in \mathbf{Op}_{? \cap Y}.$	$\bigcup_{i \in I} A_i = \bigcup_{i \in I} (B_i \cap Y) = \left(\bigcup_{i \in I} B_i \right) \cap Y \in \mathbf{Op}_{? \cap Y}.$
fong-spivak-invitation_EQ0075 (3.15)	121	■ 0.380	$\mathcal{G} := \text{mid} \stackrel{\text{Email}}{\underset{\text{received_by}}{\longrightarrow}} \text{Address}$	(not rendered)	$\mathcal{G} := \begin{array}{c} \text{Email} \xrightarrow{\text{sent_by}} \text{Address} \\ \text{received_by} \\ \text{no equations} \end{array}$
fong-spivak-invitation_EQ0141	221	■ 0.382	$\begin{aligned} \operatorname{Circ}(f) : \operatorname{Circ}(V) &\longrightarrow \operatorname{Circ}(V') \\ \operatorname{Circ}(V) &\longrightarrow \operatorname{Circ}(V') \\ \operatorname{Circ}(V) &\longrightarrow \operatorname{Circ}(V') \end{aligned}$	$\operatorname{Circ}(f) : \operatorname{Circ}(V) \longrightarrow \operatorname{Circ}(V')$ $(V, A, s, t, \ell) \mapsto (V', A, (s \div f), (t; f), \ell).$	$\operatorname{Circ}(f) : \operatorname{Circ}(V) \longrightarrow \operatorname{Circ}(V');$ $(V, A, s, t, \ell) \mapsto (V', A, (s \circ f), (t \circ f), \ell).$
fong-spivak-invitation_EQ0214	299	■ 0.401	$\begin{aligned} \Lambda(\text{monoid}, \text{nothing}) &= \Lambda(\text{monoid}, \text{this book}) \\ \Lambda(\text{monoid}, \text{nothing}) &= \Lambda(\text{preorder}, \text{this book}) \\ \Lambda(\text{preorder}, \text{nothing}) &= \Lambda(\text{category}, \text{nothing}) = \text{false}. \end{aligned}$	$\Lambda(\text{monoid}, \text{nothing}) = \Lambda(\text{monoid}, \text{this book})$ $\Lambda(\text{monoid}, \text{nothing}) = \Lambda(\text{preorder}, \text{this book})$ $\Lambda(\text{preorder}, \text{nothing}) = \Lambda(\text{category}, \text{nothing}) = \text{false}.$	$\Lambda(\text{monoid}, \text{nothing}) = \Lambda(\text{monoid}, \text{this book})$ $\Lambda(\text{monoid}, \text{nothing}) = \Lambda(\text{preorder}, \text{this book})$ $\Lambda(\text{preorder}, \text{nothing}) = \Lambda(\text{category}, \text{nothing}) = \text{false}.$
fong-spivak-invitation_EQ0024	59	■ 0.473	$\left. \left. \frac{x}{y} \right \right _z = \frac{x \otimes y}{z}$	$\left. \left. \frac{x}{y} \right \right _z = \frac{x \otimes y}{z}$	$\left. \left. \frac{x}{y} \right \right _z = \frac{x \otimes y}{z}$
fong-spivak-invitation_EQ0255	320	■ 0.480	$\operatorname{id}_A + \operatorname{id}_B = [\operatorname{id}_A \circ \iota_A, \operatorname{id}_B \circ \iota_B] = [\iota_A, \iota_B] = \operatorname{id}_{A+B}.$	$\operatorname{id}_A + \operatorname{id}_B = [\operatorname{id}_A \circ \iota_A, \operatorname{id}_B \circ \iota_B] = [\iota_A, \iota_B] = \operatorname{id}_{A+B}.$	$\operatorname{id}_A + \operatorname{id}_B = [\operatorname{id}_A \circ \iota_A, \operatorname{id}_B \circ \iota_B] = [\iota_A, \iota_B] = \operatorname{id}_{A+B}.$
fong-spivak-invitation_EQ0210	296	■ 0.483	$H := \stackrel{4}{\leftarrow} \stackrel{d}{\rightarrow} \stackrel{5}{\rightarrow} \circ e$	$H := \stackrel{4}{\leftarrow} \stackrel{d}{\rightarrow} \stackrel{5}{\rightarrow} \circ e$	$H := \stackrel{4}{\leftarrow} \stackrel{d}{\rightarrow} \stackrel{5}{\rightarrow} \circ e$
fong-spivak-invitation_EQ0186	284	■ 0.486	$\begin{aligned} g(\text{false}) + g(\text{false}) &= \infty + \infty \geq \infty = g(\text{false} \wedge \text{false}), \\ g(\text{false}) + g(\text{true}) &= \infty + 0 \geq \infty = g(\text{false} \wedge \text{true}), \\ g(\text{true}) + g(\text{false}) &= 0 + \infty \geq \infty = g(\text{true} \wedge \text{false}), \\ g(\text{true}) + g(\text{true}) &= 0 + 0 \geq 0 = g(\text{true} \wedge \text{true}). \end{aligned}$	$g(\text{false}) + g(\text{false}) = \infty + \infty \geq \infty = g(\text{false} \wedge \text{false}),$ $g(\text{false}) + g(\text{true}) = \infty + 0 \geq \infty = g(\text{false} \wedge \text{true}),$ $g(\text{true}) + g(\text{false}) = 0 + \infty \geq \infty = g(\text{true} \wedge \text{false}),$ $g(\text{true}) + g(\text{true}) = 0 + 0 \geq 0 = g(\text{true} \wedge \text{true})$	$g(\text{false}) + g(\text{false}) = \infty + \infty \geq \infty = g(\text{false} \wedge \text{false}),$ $g(\text{false}) + g(\text{true}) = \infty + 0 \geq \infty = g(\text{false} \wedge \text{true}),$ $g(\text{true}) + g(\text{false}) = 0 + \infty \geq \infty = g(\text{true} \wedge \text{false}),$ $g(\text{true}) + g(\text{true}) = 0 + 0 \geq 0 = g(\text{true} \wedge \text{true}).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_E00092	143	■ 0.504	$(\backslash\Phi\ ;\ \backslash\Psi)\ ;\ \backslash\Upsilon=\backslash\Phi\ ;(\backslash\Psi\ ;\ \backslash\Upsilon)\ .$	$(\Phi;\Psi);\Upsilon = \Phi;(\Psi;\Upsilon).$	$(\Phi \circ \Psi) \circ \Upsilon = \Phi \circ (\Psi \circ \Upsilon).$
fong-spivak-invitation_E00077	122	■ 0.507	$\left.\frac{\mathbf{1}}{\text{Em}_6}\right\lvert$	<i>(not rendered)</i>	$\frac{\mathbf{1}}{\text{Em}_6}$
fong-spivak-invitation_E00071	118	■ 0.519	$\text{Set} \underset{(-)^B}{\overset{-\times B}{\rightleftarrows}} \text{Set} \quad \text{Set}(A \times B, C) \cong \text{Set}(A, C^B).$	$\text{Set} \overset{-\times B}{\underset{(-)^B}{\rightleftarrows}} \text{Set} \quad \text{Set}(A \times B, C) \cong \text{Set}(A, C^B).$	$\text{Set} \overset{-\times B}{\underset{(-)^B}{\rightleftarrows}} \text{Set} \quad \text{Set}(A \times B, C) \cong \text{Set}(A, C^B).$
fong-spivak-invitation_E00206	293	■ 0.529	$F(a):=a^{\prime}, \quad \backslash\text{quad } G(a):=a^{\prime}, \quad \backslash\text{quad } F(b):=b^{\prime}, \quad \backslash\text{quad } G(b):=b^{\prime}, \quad \backslash\text{quad } F(f):=f_1, \quad \backslash\text{quad } G(f):=f_2.$	$F(a) := a', \quad G(a) := a', \quad F(b) := b', \quad G(b) := b', \quad F(f) := f_1, \quad \text{and } G(f) := f_2.$	$F(a) := a', \quad G(a) := a', \quad F(b) := b', \quad G(b) := b', \quad F(f) := f_1, \text{ and } G(f) := f_2.$
fong-spivak-invitation_E00074	121	■ 0.559	$\Sigma_! : \mathcal{C} - \text{Inst} \rightarrow \text{Set} \quad \text{and} \quad \Pi_! : \mathcal{C} - \text{Inst} \rightarrow \text{Set}.$	$\Sigma_! : \mathcal{C} - \text{Inst} \rightarrow \text{Set} \quad \text{and} \quad \Pi_! : \mathcal{C} - \text{Inst} \rightarrow \text{Set}.$	$\Sigma_! : \mathcal{C}\text{-Inst} \rightarrow \text{Set} \quad \text{and} \quad \Pi_! : \mathcal{C}\text{-Inst} \rightarrow \text{Set}.$
fong-spivak-invitation_E00239	313	■ 0.561	$c_n := -c : (1 + -c) : (1 + 1 + -c) : \ldots : (1 + \cdots + 1 + < c) : 1 \rightarrow n.$	$c_n := -c : (1 + -c) : (1 + 1 + -c) : \ldots : (1 + \cdots + 1 + < c) : 1 \rightarrow n.$	$c_n := -< \circ (1 + -<) \circ (1 + 1 + -<) \circ \ldots \circ (1 + \cdots + 1 + -<) : 1 \rightarrow n.$
fong-spivak-invitation_E00073 (3.14)	121	■ 0.576	$! : \mathcal{C} \rightarrow \mathbf{1}.$	$! : \mathcal{C} \rightarrow \mathbf{1}.$	$! : \mathcal{C} \rightarrow \mathbf{1}.$
fong-spivak-invitation_E00025	59	■ 0.583	$\frac{\frac{x}{y}}{z} = \frac{\frac{x}{y}}{\frac{z}{z}}$	$\frac{\frac{x}{y}}{z} = \frac{\frac{x}{y}}{\frac{z}{z}}$	$\frac{\frac{x}{y}}{z} = \frac{\frac{x}{y}}{\frac{z}{z}}$
fong-spivak-invitation_E00132	206	■ 0.584	$\left\{ (v, d) \mid d \in D(v), \text{ where } v = v_1 \text{ or } v = v_2 \right\}.$	$\{(v, d) \mid d \in D(v), \text{ where } v = v_1 \text{ or } v = v_2\}.$	$\{(v, d) \mid d \in D(v), \text{ where } v = v_1 \text{ or } v = v_2\}.$
fong-spivak-invitation_E00057	98	■ 0.592	$\{A, B, C, D, f, g, h, i, f; h\}.$	$\{A, B, C, D, f, g, h, i, f; h\}.$	$\{A, B, C, D, f, g, h, i, f \circ h\}.$
fong-spivak-invitation_E00123 (5.14)	189	■ 0.602	$\mathbf{SFG}_R^+ := \text{Free} \left(G_R \sqcup G_R^{\text{op}} \right).$	$\mathbf{SFG}_R^+ := \text{Free} \left(G_R \sqcup G_R^{\text{op}} \right).$	$\mathbf{SFG}_R^+ := \text{Free} \left(G_R \sqcup G_R^{\text{op}} \right).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation-EQ0221	301	■ 0.622	$\begin{aligned} \Phi: \Psi &= (M_X^3 * M_\Phi * M_Y^2) * (M_Y^2 * M_\Psi * M_Z^3) \\ &= M_X^3 * M_\Phi * M_Y^4 * M_\Psi * M_Z^3 \\ &= (M_X^3 * M_\Phi * M_Y^2) * M_\Psi * M_Z^3 \\ &= \Phi * M_\Psi * M_Z^3 \\ &= \begin{pmatrix} 17 & 20 & 20 \\ 11 & 14 & 14 \\ 14 & 17 & 17 \\ 12 & 9 & 15 \end{pmatrix} \begin{pmatrix} \infty & \infty & \infty & \infty \\ \infty & \infty & 0 & \infty \\ 4 & \infty & \infty & 4 \end{pmatrix} \begin{pmatrix} 0 & 2 & 4 & 5 \\ 4 & 0 & 2 & 3 \\ 2 & 4 & 0 & 1 \\ 1 & 3 & 5 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 17 & 20 & 20 \\ 11 & 14 & 14 \\ 14 & 17 & 17 \\ 12 & 9 & 15 \end{pmatrix} \begin{pmatrix} \infty & \infty & \infty & \infty \\ 2 & 4 & 0 & 1 \\ 4 & 6 & 8 & 4 \end{pmatrix} \\ &= \begin{pmatrix} 22 & 24 & 20 & 21 \\ 16 & 18 & 14 & 15 \\ 19 & 21 & 17 & 18 \\ 11 & 13 & 9 & 10 \end{pmatrix}. \end{aligned}$	$\begin{aligned} \Phi: \Psi &= (M_X^3 * M_\Phi * M_Y^2) * (M_Y^2 * M_\Psi * M_Z^3) \\ &= M_X^3 * M_\Phi * M_Y^4 * M_\Psi * M_Z^3 \\ &= (M_X^3 * M_\Phi * M_Y^2) * M_\Psi * M_Z^3 \\ &= \Phi * M_\Psi * M_Z^3 \\ &= \begin{pmatrix} 17 & 20 & 20 \\ 11 & 14 & 14 \\ 14 & 17 & 17 \\ 12 & 9 & 15 \end{pmatrix} \begin{pmatrix} \infty & \infty & \infty & \infty \\ \infty & \infty & 0 & \infty \\ 4 & \infty & \infty & 4 \end{pmatrix} \begin{pmatrix} 0 & 2 & 4 & 5 \\ 4 & 0 & 2 & 3 \\ 2 & 4 & 0 & 1 \\ 1 & 3 & 5 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 17 & 20 & 20 \\ 11 & 14 & 14 \\ 14 & 17 & 17 \\ 12 & 9 & 15 \end{pmatrix} \begin{pmatrix} \infty & \infty & \infty & \infty \\ 2 & 4 & 0 & 1 \\ 4 & 6 & 8 & 4 \end{pmatrix} \\ &= \begin{pmatrix} 22 & 24 & 20 & 21 \\ 16 & 18 & 14 & 15 \\ 19 & 21 & 17 & 18 \\ 11 & 13 & 9 & 10 \end{pmatrix}. \end{aligned}$	$\begin{aligned} \Phi \circ \Psi &= (M_X^3 * M_\Phi * M_Y^2) * (M_Y^2 * M_\Psi * M_Z^3) \\ &= M_X^3 * M_\Phi * M_Y^4 * M_\Psi * M_Z^3 \\ &= (M_X^3 * M_\Phi * M_Y^2) * M_\Psi * M_Z^3 \\ &= \Phi * M_\Psi * M_Z^3 \\ &= \begin{pmatrix} 17 & 20 & 20 \\ 11 & 14 & 14 \\ 14 & 17 & 17 \\ 12 & 9 & 15 \end{pmatrix} \begin{pmatrix} \infty & \infty & \infty & \infty \\ \infty & \infty & 0 & \infty \\ 4 & \infty & \infty & 4 \end{pmatrix} \begin{pmatrix} 0 & 2 & 4 & 5 \\ 4 & 0 & 2 & 3 \\ 2 & 4 & 0 & 1 \\ 1 & 3 & 5 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 17 & 20 & 20 \\ 11 & 14 & 14 \\ 14 & 17 & 17 \\ 12 & 9 & 15 \end{pmatrix} \begin{pmatrix} \infty & \infty & \infty & \infty \\ 2 & 4 & 0 & 1 \\ 4 & 6 & 8 & 4 \end{pmatrix} \\ &= \begin{pmatrix} 22 & 24 & 20 & 21 \\ 16 & 18 & 14 & 15 \\ 19 & 21 & 17 & 18 \\ 11 & 13 & 9 & 10 \end{pmatrix}. \end{aligned}$
fong-spivak-invitation-EQ0010 (1.7)	44	■ 0.623	$p \leq g(f(p)) \quad \text{and} \quad f(g(q)) \leq q.$	$p \leq g(f(p)) \quad \text{and} \quad f(g(q)) \leq q.$	$p \leq g(f(p)) \quad \text{and} \quad f(g(q)) \leq q.$
fong-spivak-invitation-EQ0155 (7.7)	242	■ 0.623	$X = \{(\text{true}, \text{true}), (\text{true}, \text{false}), (\text{false}, \text{true})\} \subseteq \mathbb{B} \times \mathbb{B}.$	$X = \{(\text{true}, \text{true}), (\text{true}, \text{false}), (\text{false}, \text{true})\} \subseteq \mathbb{B} \times \mathbb{B}.$	$X = \{(\text{true}, \text{true}), (\text{true}, \text{false}), (\text{false}, \text{true})\} \subseteq \mathbb{B} \times \mathbb{B}.$
fong-spivak-invitation-EQ0156	247	■ 0.628	$B(p; \epsilon) := \{p' \in \mathbb{R}^2 \mid d(p, p') < \epsilon\}.$	$B(p; \epsilon) := \{p' \in \mathbb{R}^2 \mid d(p, p') < \epsilon\}.$	$B(p; \epsilon) := \{p' \in \mathbb{R}^2 \mid d(p, p') < \epsilon\}.$
fong-spivak-invitation-EQ0124	190	■ 0.641	$\mathrm{B}(g \circ (h^{\mathrm{op}})) = \{(x, y) \mid S(g)(x) = S(h)(y)\}.$	$\mathrm{B}(g \circ (h^{\mathrm{op}})) = \{(x, y) \mid S(g)(x) = S(h)(y)\}.$	$\mathrm{B}(g \circ (h^{\mathrm{op}})) = \{(x, y) \mid S(g)(x) = S(h)(y)\}.$
fong-spivak-invitation-EQ0080 (4.2)	134	■ 0.641	$\Phi: X^{\mathrm{op}} \times y \rightarrow \mathbf{Bool}.$	$\Phi: X^{\mathrm{op}} \times y \rightarrow \mathbf{Bool}.$	$\Phi: \mathcal{X}^{\mathrm{op}} \times \mathcal{Y} \rightarrow \mathbf{Bool}.$
fong-spivak-invitation-EQ0097	150	■ 0.644	$S \times (T \times U) := \{(s, (t, u)) \mid s \in S, t \in T, u \in U\} \\ =? (S \times T) \times U := \{((s, t), u) \mid s \in S, t \in T, u \in U\}.$	$S \times (T \times U) := \{(s, (t, u)) \mid s \in S, t \in T, u \in U\} \\ =? (S \times T) \times U := \{((s, t), u) \mid s \in S, t \in T, u \in U\}.$	$S \times (T \times U) := \{(s, (t, u)) \mid s \in S, t \in T, u \in U\} \\ =? (S \times T) \times U := \{((s, t), u) \mid s \in S, t \in T, u \in U\}.$
fong-spivak-invitation-EQ0224	305	■ 0.649	$\bigvee_{y \in \mathcal{X}} \mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \leq \bigvee_{y \in \mathcal{X}} \mathcal{X}(x, z) = \mathcal{X}(x, z).$	$\bigvee_{y \in \mathcal{X}} \mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \leq \bigvee_{y \in \mathcal{X}} \mathcal{X}(x, z) = \mathcal{X}(x, z).$	$\bigvee_{y \in \mathcal{X}} \mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \leq \bigvee_{y \in \mathcal{X}} \mathcal{X}(x, z) = \mathcal{X}(x, z).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_EQ0055 (3.5)	96	■ 0.649	$\mathbf{2} := \operatorname{Free} \left(\begin{array}{ ccc } \hline v_1 & f_1 & v_2 \\ \bullet & & \bullet \\ \hline \end{array} \right)$	$\mathbf{2} := \operatorname{Free} \left(\begin{array}{ ccc } \hline v_1 & f_1 & v_2 \\ \bullet & & \bullet \\ \hline \end{array} \right)$	$\mathbf{2} := \operatorname{Free} \left(\begin{array}{ ccc } \hline v_1 & f_1 & v_2 \\ \bullet & \longrightarrow & \bullet \\ \hline \end{array} \right)$
fong-spivak-invitation_EQ0107 (5.4)	167	■ 0.670	$G + G^{\prime} := \left(\left(V \sqcup V^{\prime} \right), [\text{in}, \text{in}^{\prime}], [\text{out}, \text{out}^{\prime}], (\iota \sqcup \iota^{\prime}) \right)$	$G + G' := ((V \sqcup V'), [\text{in}, \text{in}'], [\text{out}, \text{out}'], (\iota \sqcup \iota')).$	$G + G' := ((V \sqcup V'), [\text{in}, \text{in}'], [\text{out}, \text{out}'], (\iota \sqcup \iota')).$
fong-spivak-invitation_EQ0151 (7.3)	239	■ 0.683	$\mathcal{C}(A \times C, D) \cong \mathcal{C}(A, D^C)$	$\mathcal{C}(A \times C, D) \cong \mathcal{C}(A, D^C).$	$\mathcal{C}(A \times C, D) \cong \mathcal{C}(A, D^C).$
fong-spivak-invitation_EQ0191	288	■ 0.683	$\emptyset \otimes v \cong v \otimes \emptyset \cong \bigvee_{a \in \emptyset} (v \otimes a) = 0$	$0 \otimes v \cong v \otimes 0 \cong \left(v \otimes \bigvee_{a \in \emptyset} a \right) = \bigvee_{a \in \emptyset} (v \otimes a) = 0.$	$0 \otimes v \cong v \otimes 0 \cong \left(v \otimes \bigvee_{a \in \emptyset} a \right) = \bigvee_{a \in \emptyset} (v \otimes a) = 0.$
fong-spivak-invitation_EQ0200	291	■ 0.687	$A, \quad A: f, \quad A: g, \quad A: j, \quad B, \quad B: h, \quad C, \quad C: i, \quad D$	$A, \quad A: f, \quad A: g, \quad A: j, \quad B, \quad B: h, \quad C, \quad C: i, \quad D$	$A, \quad A \circ f, \quad A \circ g, \quad A \circ j, \quad B, \quad B \circ h, \quad C, \quad C \circ i, \quad D$
fong-spivak-invitation_EQ0089	142	■ 0.688	$\mathrm{U}_P; \Phi = \Phi = \Phi; \mathrm{U}_Q.$	$\mathrm{U}_P; \Phi = \Phi = \Phi; \mathrm{U}_Q.$	$\mathrm{U}_P \circ \Phi = \Phi = \Phi \circ \mathrm{U}_Q.$
fong-spivak-invitation_EQ0170 (7.18)	263	■ 0.706	$\forall (t : T) . \exists (s : S) . f(s) = t$	$\forall (t : T) . \exists (s : S) . f(s) = t$	$\forall (t : T) . \exists (s : S) . f(s) = t$
fong-spivak-invitation_EQ0061 (3.10)	113	■ 0.708	$\mathbf{Gr} := \mid \stackrel{\text{Arrow}}{\text{source}} \stackrel{\text{Vertex}}{\text{target}} \bullet$	$\mathbf{Gr} := \mid \stackrel{\text{Arrow}}{\text{source}} \stackrel{\text{Vertex}}{\text{target}} \bullet$	$\mathbf{Gr} := \begin{array}{ c } \hline \begin{array}{ccc} \text{Arrow} & \xrightarrow[\text{target}]{\text{source}} & \text{Vertex} \\ \bullet & & \bullet \end{array} \\ \hline \text{no equations} \end{array}$
fong-spivak-invitation_EQ0018	49	■ 0.726	$\text{fix}_j := \{p \in P \mid j(p) \cong p\}$	$\text{fix}_j := \{p \in P \mid j(p) \cong p\}.$	$\text{fix}_j := \{p \in P \mid j(p) \cong p\}.$
fong-spivak-invitation_EQ0149 (7.2)	234	■ 0.727	$((A = \text{go}) \Rightarrow \ddot{S} > 1) \wedge ((A = \text{stop}) \Rightarrow \ddot{S} = 0)$	$((A = \text{go}) \Rightarrow \ddot{S} > 1) \wedge ((A = \text{stop}) \Rightarrow \ddot{S} = 0).$	$((A = \text{go}) \Rightarrow \ddot{S} > 1) \wedge ((A = \text{stop}) \Rightarrow \ddot{S} = 0).$
fong-spivak-invitation_EQ0082	135	■ 0.730	$\Phi: x^{\operatorname{op}} \times y \rightarrow \mathcal{V}$	$\Phi: x^{\operatorname{op}} \times y \rightarrow \mathcal{V}.$	$\Phi: \mathcal{X}^{\operatorname{op}} \times \mathcal{Y} \rightarrow \mathcal{V}.$
fong-spivak-invitation_EQ0248	316	■ 0.734	$\mathbf{B}(g_9^{\circ}(h^{\operatorname{op}})) = \{(x, y) \mid S(g)(x) = S(h)(y)\}$	$\mathbf{B}(g_9^{\circ}(h^{\operatorname{op}})) = \{(x, y) \mid S(g)(x) = S(h)(y)\}.$	$\mathbf{B}(g \circ (h^{\operatorname{op}})) = \{(x, y) \mid S(g)(x) = S(h)(y)\}.$
fong-spivak-invitation_EQ0114	178	■ 0.740	$\left(\begin{array}{cc} A & 0 \\ 0 & B \end{array} \right)$	$\left(\begin{array}{cc} A & 0 \\ 0 & B \end{array} \right)$	$\left(\begin{array}{cc} A & 0 \\ 0 & B \end{array} \right)$
fong-spivak-invitation_EQ0138	220	■ 0.748	$C := \{ \text{light, switch, battery} \} \sqcup \{ x \Omega \mid x \in \mathbb{R}^+ \}$	$C := \{ \text{light, switch, battery} \} \sqcup \{ x \Omega \mid x \in \mathbb{R}^+ \}.$	$C := \{ \text{light, switch, battery} \} \sqcup \{ x \Omega \mid x \in \mathbb{R}^+ \}.$
fong-spivak-invitation_EQ0143 (6.21)	226	■ 0.754	$\begin{aligned} \circ_i &: \mathcal{O}(s_1, \dots, s_m; t_i) \times \mathcal{O}(t_1, \dots, t_n; t) \\ &\rightarrow \mathcal{O}(t_1, \dots, t_{i-1}, s_1, \dots, s_m, t_{i+1}, \dots, t_n; t); \end{aligned}$	$\circ_i: \mathcal{O}(s_1, \dots, s_m; t_i) \times \mathcal{O}(t_1, \dots, t_n; t) \rightarrow \mathcal{O}(t_1, \dots, t_{i-1}, s_1, \dots, s_m, t_{i+1}, \dots, t_n; t);$	$\circ_i: \mathcal{O}(s_1, \dots, s_m; t_i) \times \mathcal{O}(t_1, \dots, t_n; t) \rightarrow \mathcal{O}(t_1, \dots, t_{i-1}, s_1, \dots, s_m, t_{i+1}, \dots, t_n; t);$
fong-spivak-invitation_EQ0198	290	■ 0.762	$\mathbf{v}_1 \xrightarrow{f_1} \bullet \xrightarrow{f_2} \dots \xrightarrow{f_{n-1}} \mathbf{v}_n$	$\mathbf{v}_1 \xrightarrow{f_1} \bullet \xrightarrow{f_2} \dots \xrightarrow{f_{n-1}} \mathbf{v}_n$	$\mathbf{v}_1 \xrightarrow{f_1} \bullet \xrightarrow{f_2} \dots \xrightarrow{f_{n-1}} \mathbf{v}_n$
fong-spivak-invitation_EQ0209	296	■ 0.772	$G := \mathbf{1} \xrightarrow{a} \mathbf{2} \xrightarrow{b}$	$G := \mathbf{1} \xrightarrow{a} \mathbf{2} \xrightarrow{b}$	$G := \mathbf{1} \xrightarrow{a} \mathbf{2} \xrightarrow{b} \mathbf{3}$
fong-spivak-invitation_EQ0160	253	■ 0.775	$\operatorname{Sec}_f(U) := \{g: U \rightarrow X \mid g \text{ is continuous and } (g \circ f)(u) = u \text{ for all } u \in U\}$	$\operatorname{Sec}_f(U) := \{g: U \rightarrow X \mid g \text{ is continuous and } (g \circ f)(u) = u \text{ for all } u \in U\}.$	$\operatorname{Sec}_f(U) := \{g: U \rightarrow X \mid g \text{ is continuous and } (g \circ f)(u) = u \text{ for all } u \in U\}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_E00193	289	■ 0.936	$\begin{aligned} ((M * N) * P)(w, z) &= \bigvee_{y \in Y} \left(\bigvee_{x \in X} M(w, x) \otimes N(x, y) \right) \otimes P(y, z) \\ &\cong \bigvee_{y \in Y, x \in X} M(w, x) \otimes N(x, y) \otimes P(y, z) \\ &\cong \bigvee_{x \in X} M(w, x) \otimes \left(\bigvee_{y \in Y} N(x, y) \otimes P(y, z) \right) \\ &= (M * (N * P))(w, z) \end{aligned}$	$\begin{aligned} ((M * N) * P)(w, z) &= \bigvee_{y \in Y} \left(\bigvee_{x \in X} M(w, x) \otimes N(x, y) \right) \otimes P(y, z) \\ &\cong \bigvee_{y \in Y, x \in X} M(w, x) \otimes N(x, y) \otimes P(y, z) \\ &\cong \bigvee_{x \in X} M(w, x) \otimes \left(\bigvee_{y \in Y} N(x, y) \otimes P(y, z) \right) \\ &= (M * (N * P))(w, z) \end{aligned}$	$\begin{aligned} ((M * N) * P)(w, z) &= \bigvee_{y \in Y} \left(\bigvee_{x \in X} M(w, x) \otimes N(x, y) \right) \otimes P(y, z) \\ &\cong \bigvee_{y \in Y, x \in X} M(w, x) \otimes N(x, y) \otimes P(y, z) \\ &\cong \bigvee_{x \in X} M(w, x) \otimes \left(\bigvee_{y \in Y} N(x, y) \otimes P(y, z) \right) \\ &= (M * (N * P))(w, z). \end{aligned}$
fong-spivak-invitation_E00093	144	■ 0.937	$\widehat{F}(p, q) := \mathcal{Q}(F(p), q) \quad \text{and} \quad \check{F}(q, p) := \mathcal{Q}(q, F(p)).$	$\widehat{F}(p, q) := \mathcal{Q}(F(p), q) \quad \text{and} \quad \check{F}(q, p) := \mathcal{Q}(q, F(p)).$	$\widehat{F}(p, q) := \mathcal{Q}(F(p), q) \quad \text{and} \quad \check{F}(q, p) := \mathcal{Q}(q, F(p)).$
fong-spivak-invitation_E00174 (7.19)	268	■ 0.937	$\begin{aligned} &\forall (t : \mathbb{R}). @_t(\text{bad_pos}(D)) \Rightarrow \\ &\exists (r : \mathbb{R}). (0 < r < 1) \wedge \forall (r' : \mathbb{R}). 0 \leq r' \leq 5 \Rightarrow @_{t+r+r'}(\text{engaged}(T)). \end{aligned}$	$\forall (t : \mathbb{R}). @_t(\text{bad_pos}(D)) \Rightarrow \\ \exists (r : \mathbb{R}). (0 < r < 1) \wedge \forall (r' : \mathbb{R}). 0 \leq r' \leq 5 \Rightarrow @_{t+r+r'}(\text{engaged}(T)).$	$\forall (t : \mathbb{R}). @_t(\text{bad_pos}(D)) \Rightarrow \\ \exists (r : \mathbb{R}). (0 < r < 1) \wedge \forall (r' : \mathbb{R}). 0 \leq r' \leq 5 \Rightarrow @_{t+r+r'}(\text{engaged}(T)).$
fong-spivak-invitation_E00079	127	■ 0.943	$\lim_{\mathcal{J}} D := \{(d_1, \dots, d_n) \mid d_i \in D(v_i) \text{ for all } 1 \leq i \leq n \text{ and for all } a : v_i \rightarrow v_j \in A, \text{ we have } D(a)(d_i) = d_j\}$	$\lim_{\mathcal{J}} D := \{(d_1, \dots, d_n) \mid d_i \in D(v_i) \text{ for all } 1 \leq i \leq n \text{ and for all } a : v_i \rightarrow v_j \in A, \text{ we have } D(a)(d_i) = d_j\}$	$\lim_{\mathcal{J}} D := \{(d_1, \dots, d_n) \mid d_i \in D(v_i) \text{ for all } 1 \leq i \leq n \text{ and for all } a : v_i \rightarrow v_j \in A, \text{ we have } D(a)(d_i) = d_j\}$
fong-spivak-invitation_E00226	305	■ 0.948	$x \xrightarrow{\alpha^{-1}} x \times \mathbf{1} \xrightarrow{U_x \times \eta_x} x \times x^{\text{op}} \times x \xrightarrow{\epsilon_x \times U_x} \mathbf{1} \times x \xrightarrow{\alpha} x$	$x \xrightarrow{\alpha^{-1}} x \times \mathbf{1} \xrightarrow{U_x \times \eta_x} x \times x^{\text{op}} \times x \xrightarrow{\epsilon_x \times U_x} \mathbf{1} \times x \xrightarrow{\alpha} x$	$\mathcal{X} \xrightarrow{\alpha^{-1}} \mathcal{X} \times \mathbf{1} \xrightarrow{U_{\mathcal{X}} \times \eta_{\mathcal{X}}} \mathcal{X} \times \mathcal{X}^{\text{op}} \times \mathcal{X} \xrightarrow{\epsilon_{\mathcal{X}} \times U_{\mathcal{X}}} \mathbf{1} \times \mathcal{X} \xrightarrow{\alpha} \mathcal{X}$
fong-spivak-invitation_E00145 (6.22)	228	■ 0.950	$(g \circ i f) : X_1 \times \dots \times X_{i-1} \times W_1 \times \dots \times W_m \times X_{i+1} \times \dots \times X_n \longrightarrow Y,$	$(g \circ i f) : X_1 \times \dots \times X_{i-1} \times W_1 \times \dots \times W_m \times X_{i+1} \times \dots \times X_n \longrightarrow Y,$	$(g \circ_i f) : X_1 \times \dots \times X_{i-1} \times W_1 \times \dots \times W_m \times X_{i+1} \times \dots \times X_n \longrightarrow Y,$
fong-spivak-invitation_E00207	293	■ 0.954	$(F ; G)(f ; g) = G(F(f) ; F(g)) = G(F(f)) ; G(F(g))$	$(F;G)(f;g) = G(F(f);F(g)) = G(F(f));G(F(g))$	$(F \circ G)(f \circ g) = G(F(f) \circ F(g)) = G(F(f)) \circ G(F(g))$
fong-spivak-invitation_E00048 (2.24)	84	■ 0.954	$\left(v \otimes \bigvee_{a \in A} a \right) \cong \bigvee_{a \in A} (v \otimes a).$	$\left(v \otimes \bigvee_{a \in A} a \right) \cong \bigvee_{a \in A} (v \otimes a).$	$\left(v \otimes \bigvee_{a \in A} a \right) \cong \bigvee_{a \in A} (v \otimes a).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_EQ0219	300	■ 0.957	$\begin{aligned} \Phi &= M_X^3 * M_\Phi * M_Y^2 = \begin{pmatrix} 0 & 6 & 3 & 11 \\ 2 & 0 & 5 & 5 \\ 5 & 3 & 0 & 8 \\ 11 & 9 & 6 & 0 \end{pmatrix} \begin{pmatrix} \infty & \infty & \infty \\ 11 & \infty & \infty \\ \infty & \infty & \infty \\ \infty & 9 & \infty \end{pmatrix} \begin{pmatrix} 0 & 4 & 3 \\ 3 & 0 & 6 \\ 7 & 4 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 17 & 20 & \infty \\ 11 & 14 & \infty \\ 14 & 17 & \infty \\ 20 & 9 & \infty \end{pmatrix} \begin{pmatrix} 0 & 4 & 3 \\ 3 & 0 & 6 \\ 7 & 4 & 0 \end{pmatrix} \end{aligned}$	$\Phi = M_X^3 * M_\Phi * M_Y^2 = \begin{pmatrix} 0 & 6 & 3 & 11 \\ 2 & 0 & 5 & 5 \\ 5 & 3 & 0 & 8 \\ 11 & 9 & 6 & 0 \end{pmatrix} \begin{pmatrix} \infty & \infty & \infty \\ 11 & \infty & \infty \\ \infty & \infty & \infty \\ \infty & 9 & \infty \end{pmatrix} \begin{pmatrix} 0 & 4 & 3 \\ 3 & 0 & 6 \\ 7 & 4 & 0 \end{pmatrix} \\ = \begin{pmatrix} 17 & 20 & \infty \\ 11 & 14 & \infty \\ 14 & 17 & \infty \\ 20 & 9 & \infty \end{pmatrix} \begin{pmatrix} 0 & 4 & 3 \\ 3 & 0 & 6 \\ 7 & 4 & 0 \end{pmatrix}$	$\Phi = M_X^3 * M_\Phi * M_Y^2 = \begin{pmatrix} 0 & 6 & 3 & 11 \\ 2 & 0 & 5 & 5 \\ 5 & 3 & 0 & 8 \\ 11 & 9 & 6 & 0 \end{pmatrix} \begin{pmatrix} \infty & \infty & \infty \\ 11 & \infty & \infty \\ \infty & \infty & \infty \\ \infty & 9 & \infty \end{pmatrix} \begin{pmatrix} 0 & 4 & 3 \\ 3 & 0 & 6 \\ 7 & 4 & 0 \end{pmatrix} \\ = \begin{pmatrix} 17 & 20 & \infty \\ 11 & 14 & \infty \\ 14 & 17 & \infty \\ 20 & 9 & \infty \end{pmatrix} \begin{pmatrix} 0 & 4 & 3 \\ 3 & 0 & 6 \\ 7 & 4 & 0 \end{pmatrix}$
fong-spivak-invitation_EQ0072	120	■ 0.959	$\Sigma_{\{I\}}(\text{Airline Seat}) = I(\text{Economy}) \sqcup I(\text{First Class}).$	$\Sigma_F(I)(\text{Airline Seat}) = I(\text{Economy}) \sqcup I(\text{First Class}).$	$\Sigma_F(I)(\text{Airline Seat}) = I(\text{Economy}) \sqcup I(\text{First Class}).$
fong-spivak-invitation_EQ0103	159	■ 0.962	$\eta_X: \mathbf{1} \times \mathcal{X}^{\text{op}} \times \mathcal{X} \rightarrow \mathcal{V};$	$\eta_X: \mathbf{1} \times \mathcal{X}^{\text{op}} \times \mathcal{X} \rightarrow \mathcal{V};$	$\eta_X: \mathbf{1} \times \mathcal{X}^{\text{op}} \times \mathcal{X} \rightarrow \mathcal{V};$
fong-spivak-invitation_EQ0046	83	■ 0.964	$a + c \rightarrow d \quad \text{iff} \quad a \rightarrow (c \multimap d).$	$a + c \rightarrow d \quad \text{iff} \quad a \rightarrow (c \multimap d).$	$a + c \rightarrow d \quad \text{iff} \quad a \rightarrow (c \multimap d).$
fong-spivak-invitation_EQ0110	176	■ 0.965	$\dot{x} + 3\ddot{y} - 2z = 0, \quad \ddot{y} + 5\dot{z} = 0.$	$\dot{x} + 3\ddot{y} - 2z = 0, \quad \ddot{y} + 5\dot{z} = 0.$	$\dot{x} + 3\ddot{y} - 2z = 0, \quad \ddot{y} + 5\dot{z} = 0.$
fong-spivak-invitation_EQ0172	265	■ 0.965	$\mathbb{R} \cong \{[d, u] \in \mathbb{R} \mid d = u\}.$	$\mathbb{R} \cong \{[d, u] \in \mathbb{R} \mid d = u\}.$	$\mathbb{R} \cong \{[d, u] \in \mathbb{R} \mid d = u\}.$
fong-spivak-invitation_EQ0216	300	■ 0.965	$X(x', x) \otimes y(y, y') \leq \Phi(x, y) \multimap \Phi(x', y').$	$X(x', x) \otimes y(y, y') \leq \Phi(x, y) \multimap \Phi(x', y').$	$\mathcal{X}(x', x) \otimes \mathcal{Y}(y, y') \leq \Phi(x, y) \multimap \Phi(x', y').$
fong-spivak-invitation_EQ0250	316	■ 0.966	$\{(y, z) \mid \text{there exists } x \in R^m \text{ such that } y = S(g)(x) \text{ and } S(h)(x) = z\}.$	$\{(y, z) \mid \text{there exists } x \in R^m \text{ such that } y = S(g)(x) \text{ and } S(h)(x) = z\}.$	$\{(y, z) \mid \text{there exists } x \in R^m \text{ such that } y = S(g)(x) \text{ and } S(h)(x) = z\}.$
fong-spivak-invitation_EQ0168	261	■ 0.967	$\forall (t : T). p(s, t).$	$\forall (t : T). p(s, t).$	$\forall (t : T). p(s, t).$
fong-spivak-invitation_EQ0007	40	■ 0.967	$f(a) \vee f(b) \supseteq f(a \vee b).$	$f(a) \vee f(b) \not\supseteq f(a \vee b).$	$f(a) \vee f(b) \not\supseteq f(a \vee b).$
fong-spivak-invitation_EQ0101	158	■ 0.969	$(\mathcal{X} \times \mathcal{Y})((x, y), (x', y')) := \mathcal{X}(x, x') \otimes \mathcal{Y}(y, y').$	$(\mathcal{X} \times \mathcal{Y})((x, y), (x', y')) := \mathcal{X}(x, x') \otimes \mathcal{Y}(y, y').$	$(\mathcal{X} \times \mathcal{Y})((x, y), (x', y')) := \mathcal{X}(x, x') \otimes \mathcal{Y}(y, y').$
fong-spivak-invitation_EQ0094 (4.11)	145	■ 0.970	$\mathcal{P}(p, G(q)) \cong Q(F(p), q).$	$\mathcal{P}(p, G(q)) \cong Q(F(p), q).$	$\mathcal{P}(p, G(q)) \cong Q(F(p), q).$
fong-spivak-invitation_EQ0213	299	■ 0.971	$\lim_{\mathcal{J}} D := \{(d_1, d_2) \mid d_1 \in X \text{ and } d_2 \in Y\},$	$\lim_{\mathcal{J}} D := \{(d_1, d_2) \mid d_1 \in X \text{ and } d_2 \in Y\},$	$\lim_{\mathcal{J}} D := \{(d_1, d_2) \mid d_1 \in X \text{ and } d_2 \in Y\},$
fong-spivak-invitation_EQ0259	328	■ 0.972	$p; h_3 = r; g; h_3 = r; h_2; g' = q; f'; g',$	$p; h_3 = r; g; h_3 = r; h_2; g' = q; f'; g',$	$p \circ h_3 = r \circ g \circ h_3 = r \circ h_2 \circ g' = q \circ f' \circ g',$
fong-spivak-invitation_EQ0152 (7.5)	241	■ 0.972	$\Omega_{\text{Set}} := \mathbb{B} = \{\text{true}, \text{false}\}.$	$\Omega_{\text{Set}} := \mathbb{B} = \{\text{true}, \text{false}\}.$	$\Omega_{\text{Set}} := \mathbb{B} = \{\text{true}, \text{false}\}.$
fong-spivak-invitation_EQ0068 (3.13)	117	■ 0.979	$f(p) \leq q \text{ iff } p \leq g(q).$	$f(p) \leq q \text{ iff } p \leq g(q).$	$f(p) \leq q \text{ iff } p \leq g(q).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_EQ0111	176	■ 0.979	$\begin{array}{r} D \ x+3 \ D^{\{2\}} \ y-2 \ z=0 \ \backslash \ D^{\{2\}} \ y+5 \ D \ z=0 \\ \end{array}$	$\begin{array}{l} Dx+3D^2y-2z=0 \\ D^2y+5Dz=0 \end{array}$	$\begin{array}{l} Dx+3D^2y-2z=0, \\ D^2y+5Dz=0, \end{array}$
fong-spivak-invitation_EQ0104 (5.2)	163	■ 0.979	$i \longmapsto \begin{cases} f(i) & \text{if } 1 \leq i \leq m, \\ m'+g(i) & \text{if } m+1 \leq i \leq m+n. \end{cases}$	$i \mapsto \begin{cases} f(i) & \text{if } 1 \leq i \leq m, \\ m'+g(i) & \text{if } m+1 \leq i \leq m+n. \end{cases}$	$i \mapsto \begin{cases} f(i) & \text{if } 1 \leq i \leq m, \\ m'+g(i) & \text{if } m+1 \leq i \leq m+n. \end{cases}$
fong-spivak-invitation_EQ0016	47	■ 0.980	$f_{\{!\}}\!\left(A^{\{\prime\}}\!\right)\!:=\!\left\{b\in B\mid \text{there exists } a\in A^{\{\prime\}}\text{ such that } f(a)=b\right\}.$	$f_!(A') := \{b \in B \mid \text{there exists } a \in A' \text{ such that } f(a) = b\}.$	$f_!(A') := \{b \in B \mid \text{there exists } a \in A' \text{ such that } f(a) = b\}.$
fong-spivak-invitation_EQ0076	122	■ 0.982	$\begin{array}{c c} 1 & \\ \hline \text{Bob-Grace-Pat-Emmy} & \\ \text{Sue-Doug} & \end{array}$	$\begin{array}{c c} 1 & \\ \hline \text{Bob-Grace-Pat-Emmy} & \\ \text{Sue-Doug} & \end{array}$	$\begin{array}{c c} \mathbf{1} & \\ \hline \text{Bob-Grace-Pat-Emmy} & \\ \text{Sue-Doug} & \end{array}$
fong-spivak-invitation_EQ0034	67	■ 0.983	$\text{Cost} := ([0, \infty], \geq, 0, +),$	$\text{Cost} := ([0, \infty], \geq, 0, +),$	$\text{Cost} := ([0, \infty], \geq, 0, +),$
fong-spivak-invitation_EQ0131	206	■ 0.985	$\operatorname{colim}_{\mathcal{J}} D := \{(v, d) \mid v \in V \text{ and } d \in D(v)\} / \sim$	$\operatorname{colim}_{\mathcal{J}} D := \{(v, d) \mid v \in V \text{ and } d \in D(v)\} / \sim$	$\operatorname{colim}_{\mathcal{J}} D := \{(v, d) \mid v \in V \text{ and } d \in D(v)\} / \sim$
fong-spivak-invitation_EQ0247	316	■ 0.986	$\left\{ (x, y) \mid \text{there exists } z \in R^n \text{ such that } S(g)(x) = z \text{ and } z = S(h)(y) \right\}.$	$\{(x, y) \mid \text{there exists } z \in R^n \text{ such that } S(g)(x) = z \text{ and } z = S(h)(y)\}.$	$\{(x, y) \mid \text{there exists } z \in R^n \text{ such that } S(g)(x) = z \text{ and } z = S(h)(y)\}.$
fong-spivak-invitation_EQ0017	47	■ 0.988	$f_{\{*\}}\!\left(A^{\{\prime\}}\!\right)\!:=\!\left\{b\in B\mid \text{for all } a\text{ such that } f(a)=b, \text{ we have } a\in A^{\{\prime\}}\right\}.$	$f_*(A') := \{b \in B \mid \text{for all } a \text{ such that } f(a) = b, \text{ we have } a \in A'\}.$	$f_*(A') := \{b \in B \mid \text{for all } a \text{ such that } f(a) = b, \text{ we have } a \in A'\}.$
fong-spivak-invitation_EQ0069	118	■ 0.988	$\alpha_{c,d} : \mathcal{C}(c, R(d)) \xrightarrow{\cong} \mathcal{D}(L(c), d)$	$\alpha_{c,d} : \mathcal{C}(c, R(d)) \xrightarrow{\cong} \mathcal{D}(L(c), d)$	$\alpha_{c,d} : \mathcal{C}(c, R(d)) \xrightarrow{\cong} \mathcal{D}(L(c), d)$
fong-spivak-invitation_EQ0228	307	■ 0.992	$\sigma_{m,n}(i) := \begin{cases} i+n & \text{if } i \leq m, \\ i-m & \text{if } m+1 \leq i. \end{cases}$	$\sigma_{m,n}(i) := \begin{cases} i+n & \text{if } i \leq m, \\ i-m & \text{if } m+1 \leq i. \end{cases}$	$\sigma_{m,n}(i) := \begin{cases} i+n & \text{if } i \leq m, \\ i-m & \text{if } m+1 \leq i. \end{cases}$
fong-spivak-invitation_EQ0126 (and)	191	■ 0.992	$\{(0, (x, x)) \mid x \in R\} \subseteq R^0 \times R^2 \quad \text{and} \quad \{((x, x), 0) \mid x \in R\} \subseteq R^2 \times R^0.$	$\{(0, (x, x)) \mid x \in R\} \subseteq R^0 \times R^2 \quad \text{and} \quad \{((x, x), 0) \mid x \in R\} \subseteq R^2 \times R^0.$	$\{(0, (x, x)) \mid x \in R\} \subseteq R^0 \times R^2 \quad \text{and} \quad \{((x, x), 0) \mid x \in R\} \subseteq R^2 \times R^0.$
fong-spivak-invitation_EQ0161 (7.13)	255	■ 0.992	$\Omega(U) := \{U' \in \mathbf{Op} \mid U' \subseteq U\}.$	$\Omega(U) := \{U' \in \mathbf{Op} \mid U' \subseteq U\}.$	$\Omega(U) := \{U' \in \mathbf{Op} \mid U' \subseteq U\}.$
fong-spivak-invitation_EQ0042	79	■ 0.993	$\mathcal{X}(x_1, x_2) \leq y(F(x_1), F(x_2)),$	$\mathcal{X}(x_1, x_2) \leq y(F(x_1), F(x_2)),$	$\mathcal{X}(x_1, x_2) \leq \mathcal{Y}(F(x_1), F(x_2)),$
fong-spivak-invitation_EQ0164 (7.15)	257	■ 0.994	$(U \wedge V) := U \cap V \quad \text{and} \quad (U \vee V) := U \cup V.$	$(U \wedge V) := U \cap V \quad \text{and} \quad (U \vee V) := U \cup V.$	$(U \wedge V) := U \cap V \quad \text{and} \quad (U \vee V) := U \cup V.$
fong-spivak-invitation_EQ0171	265	■ 0.994	$\mathbb{I} \mathbb{R} := \{[d, u] \subseteq \mathbb{R} \mid d \leq u\}.$	$\mathbb{I} \mathbb{R} := \{[d, u] \subseteq \mathbb{R} \mid d \leq u\}.$	$\mathbb{I} \mathbb{R} := \{[d, u] \subseteq \mathbb{R} \mid d \leq u\}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_EQ0192	289	■ 0.994	$\begin{aligned} I_X * M(x, y) &= \bigvee_{x' \in X} I_X(x, x') \otimes M(x', y) \\ &= (I_X(x, x) \otimes M(x, y)) \vee \left(\bigvee_{x' \in X, x' \neq x} I_X(x, x') \otimes M(x', y) \right) \\ &= (I \otimes M(x, y)) \vee \left(\bigvee_{x' \in X, x' \neq x} 0 \otimes M(x', y) \right) \\ &= M(x, y) \vee 0 = M(x, y). \end{aligned}$	$\begin{aligned} I_X * M(x, y) &= \bigvee_{x' \in X} I_X(x, x') \otimes M(x', y) \\ &= (I_X(x, x) \otimes M(x, y)) \vee \left(\bigvee_{x' \in X, x' \neq x} I_X(x, x') \otimes M(x', y) \right) \\ &= (I \otimes M(x, y)) \vee \left(\bigvee_{x' \in X, x' \neq x} 0 \otimes M(x', y) \right) \\ &= M(x, y) \vee 0 = M(x, y) \end{aligned}$	$\begin{aligned} I_X * M(x, y) &= \bigvee_{x' \in X} I_X(x, x') \otimes M(x', y) \\ &= (I_X(x, x) \otimes M(x, y)) \vee \left(\bigvee_{x' \in X, x' \neq x} I_X(x, x') \otimes M(x', y) \right) \\ &= (I \otimes M(x, y)) \vee \left(\bigvee_{x' \in X, x' \neq x} 0 \otimes M(x', y) \right) \\ &= M(x, y) \vee 0 = M(x, y). \end{aligned}$
fong-spivak-invitation_EQ0113 (5.7)	178	■ 0.995	$M ; N(a, c) := \sum_{b \in \underline{n}} M(a, b) \times N(b, c).$	$M; N(a, c) := \sum_{b \in \underline{n}} M(a, b) \times N(b, c).$	$M \circ N(a, c) := \sum_{b \in \underline{n}} M(a, b) \times N(b, c).$
fong-spivak-invitation_EQ0056 (3.6)	96	■ 0.995	$\mathbf{3} := \text{Free} \mid \bullet \xrightarrow{f_1} \bullet \xrightarrow{f_2} \bullet$	$\mathbf{3} := \text{Free} \mid \bullet \xrightarrow{v_1} \bullet \xrightarrow{v_2} \bullet \xrightarrow{v_3}$	$\mathbf{3} := \text{Free} \mid \begin{array}{c} v_1 \xrightarrow{f_1} v_2 \xrightarrow{f_2} v_3 \end{array}$
fong-spivak-invitation_EQ0195	290	■ 0.996	$t(p) := \begin{cases} v & \text{if } n = 0, \\ t(a_n) & \text{if } n \geq 1. \end{cases}$	$t(p) := \begin{cases} v & \text{if } n = 0, \\ t(a_n) & \text{if } n \geq 1. \end{cases}$	$t(p) := \begin{cases} v & \text{if } n = 0, \\ t(a_n) & \text{if } n \geq 1. \end{cases}$
fong-spivak-invitation_EQ0002 (1.5)	23	■ 0.996	$A = \bigcup_{p \in P} A_p \quad \text{and} \quad \text{if } p \neq q \text{ then } A_p \cap A_q = \emptyset.$	$A = \bigcup_{p \in P} A_p \quad \text{and} \quad \text{if } p \neq q \text{ then } A_p \cap A_q = \emptyset.$	$A = \bigcup_{p \in P} A_p \quad \text{and} \quad \text{if } p \neq q \text{ then } A_p \cap A_q = \emptyset.$
fong-spivak-invitation_EQ0189	288	■ 0.996	$A \cap B \subseteq (\bar{B} \cup C) \cap B = (\bar{B} \cap B) \cup (C \cap B) = C \cap B \subseteq C.$	$A \cap B \subseteq (\bar{B} \cup C) \cap B = (\bar{B} \cap B) \cup (C \cap B) = C \cap B \subseteq C.$	$A \cap B \subseteq (\bar{B} \cup C) \cap B = (\bar{B} \cap B) \cup (C \cap B) = C \cap B \subseteq C.$
fong-spivak-invitation_EQ0136	217	■ 0.996	$\varphi_{c_1, c_2}: F(c_1) \otimes_{\mathcal{D}} F(c_2) \rightarrow F(c_1 \otimes_{\mathcal{C}} c_2),$	$\varphi_{c_1, c_2}: F(c_1) \otimes_{\mathcal{D}} F(c_2) \rightarrow F(c_1 \otimes_{\mathcal{C}} c_2),$	$\varphi_{c_1, c_2}: F(c_1) \otimes_{\mathcal{D}} F(c_2) \rightarrow F(c_1 \otimes_{\mathcal{C}} c_2),$
fong-spivak-invitation_EQ0009	42	■ 0.997	$\lceil x/3 \rceil \leq y \text{ if and only if } x \leq 3y.$	$\lceil x/3 \rceil \leq y \text{ if and only if } x \leq 3y.$	$\lceil x/3 \rceil \leq y \text{ if and only if } x \leq 3y.$
fong-spivak-invitation_EQ0053	87	■ 0.997	$\begin{pmatrix} \text{false} & \text{false} \\ \text{false} & \text{true} \\ \text{true} & \text{true} \end{pmatrix} * \begin{pmatrix} \text{true} & \text{true} & \text{false} \\ \text{true} & \text{false} & \text{true} \end{pmatrix} = \begin{pmatrix} \text{false} & \text{false} & \text{false} \\ \text{true} & \text{false} & \text{true} \\ \text{true} & \text{true} & \text{true} \end{pmatrix}$	$\begin{pmatrix} \text{false} & \text{false} \\ \text{false} & \text{true} \\ \text{true} & \text{true} \end{pmatrix} * \begin{pmatrix} \text{true} & \text{true} & \text{false} \\ \text{true} & \text{false} & \text{true} \end{pmatrix} = \begin{pmatrix} \text{false} & \text{false} & \text{false} \\ \text{true} & \text{false} & \text{true} \\ \text{true} & \text{true} & \text{true} \end{pmatrix}$	$\begin{pmatrix} \text{false} & \text{false} \\ \text{false} & \text{true} \\ \text{true} & \text{true} \end{pmatrix} * \begin{pmatrix} \text{true} & \text{true} & \text{false} \\ \text{true} & \text{false} & \text{true} \end{pmatrix} = \begin{pmatrix} \text{false} & \text{false} & \text{false} \\ \text{true} & \text{false} & \text{true} \\ \text{true} & \text{true} & \text{true} \end{pmatrix}$
fong-spivak-invitation_EQ0231	310	■ 0.997	$[], \quad [a], \quad [b], \quad [a, a], \quad [a, b], \quad \dots, \quad [b, a, b, b, a, b, a, a, a, a], \quad \dots$	$[], \quad [a], \quad [b], \quad [a, a], \quad [a, b], \quad \dots, \quad [b, a, b, b, a, b, a, a, a, a], \quad \dots$	$[], \quad [a], \quad [b], \quad [a, a], \quad [a, b], \quad \dots, \quad [b, a, b, b, a, b, a, a, a, a], \quad \dots$
fong-spivak-invitation_EQ0049	86	■ 0.997	$d(U, V) := \sup_{u \in U} \inf_{v \in V} d(u, v), \quad X(U, V) := \bigwedge_{u \in U} \bigvee_{v \in V} X(u, v)$	$d(U, V) := \sup_{u \in U} \inf_{v \in V} d(u, v), \quad X(U, V) := \bigwedge_{u \in U} \bigvee_{v \in V} X(u, v)$	$d(U, V) := \sup_{u \in U} \inf_{v \in V} d(u, v), \quad X(U, V) := \bigwedge_{u \in U} \bigvee_{v \in V} X(u, v).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_EQ0215	300	■ 0.997	$\left(\mathrm{X}^{\mathrm{op}} \times \mathrm{Y}\right) \left((x, y), \left(x', y'\right)\right) \leq \mathcal{V}\left(\Phi(x, y), \Phi\left(x', y'\right)\right).$	$(\mathcal{X}^{\mathrm{op}} \times \mathcal{Y})((x, y), (x', y')) \leq \mathcal{V}(\Phi(x, y), \Phi(x', y')).$	$(\mathcal{X}^{\mathrm{op}} \times \mathcal{Y})((x, y), (x', y')) \leq \mathcal{V}(\Phi(x, y), \Phi(x', y')).$
fong-spivak-invitation_EQ0108	169	■ 0.998	$a \subset \bullet \supset b$	$a \subset \bullet \supset b$	$a \curvearrowright \bullet \curvearrowleft b$
fong-spivak-invitation_EQ0166 (7.17)	260	■ 0.998	$\left(\left \Omega^S\right ,\leq^S\right).$	$(\Omega^S , \leq^S).$	$(\Omega^S , \leq^S).$
fong-spivak-invitation_EQ0011	45	■ 0.998	$g(\bigwedge A) \cong \bigwedge g(A).$	$g(\bigwedge A) \cong \bigwedge g(A).$	$g\left(\bigwedge A\right) \cong \bigwedge g(A).$
fong-spivak-invitation_EQ0106	166	■ 0.998	$\iota^{\mathrm{\prime}}(x)=\begin{cases}\iota(x) & \text{if } \iota(x) \in I, \\ \iota'(\iota(x)) & \text{if } \iota(x) \in \underline{n}, \\ \iota'(x) & \text{if } x \in O'.\end{cases}$	$\iota''(x) = \begin{cases} \iota(x) & \text{if } \iota(x) \in I, \\ \iota'(\iota(x)) & \text{if } \iota(x) \in \underline{n}, \\ \iota'(x) & \text{if } x \in O'. \end{cases}$	$\iota''(x) = \begin{cases} \iota(x) & \text{if } \iota(x) \in I, \\ \iota'(\iota(x)) & \text{if } \iota(x) \in \underline{n}, \\ \iota'(x) & \text{if } x \in O'. \end{cases}$
fong-spivak-invitation_EQ0008 (1.6)	41	■ 0.999	$f(p) \leq q \quad \text{if and only if} \quad p \leq g(q).$	$f(p) \leq q \quad \text{if and only if} \quad p \leq g(q).$	$f(p) \leq q \quad \text{if and only if} \quad p \leq g(q).$
fong-spivak-invitation_EQ0212	298	■ 0.999	$\left(c_1, d_1\right) \xrightarrow{\left(f_1, g_1\right)}\left(c_2, d_2\right) \xrightarrow{\left(f_2, g_2\right)}\left(c_3, d_3\right) \xrightarrow{\left(f_3, g_3\right)}\left(c_4, d_4\right).$	$(c_1, d_1) \xrightarrow{(f_1, g_1)} (c_2, d_2) \xrightarrow{(f_2, g_2)} (c_3, d_3) \xrightarrow{(f_3, g_3)} (c_4, d_4).$	$(c_1, d_1) \xrightarrow{(f_1, g_1)} (c_2, d_2) \xrightarrow{(f_2, g_2)} (c_3, d_3) \xrightarrow{(f_3, g_3)} (c_4, d_4).$
fong-spivak-invitation_EQ0232	310	■ 0.999	$G:=\left\{\rho _{m,n}:m\rightarrow n\mid m,n\in \mathbb{N}\right\},\quad s\left(\rho _{m,n}\right):=m,\quad t\left(\rho _{m,n}\right):=n;$	$G:=\{\rho_{m,n}:m\rightarrow n\mid m,n\in\mathbb{N}\},\quad s(\rho_{m,n}):=m,\quad t(\rho_{m,n}):=n;$	$G:=\{\rho_{m,n}:m\rightarrow n\mid m,n\in\mathbb{N}\},\quad s(\rho_{m,n}):=m,\quad t(\rho_{m,n}):=n;$
fong-spivak-invitation_EQ0253	318	■ 0.999	$\begin{aligned} f(m) *_R f(n) &= (\underbrace{1_R + \cdots + 1_R}_{m \text{ summands}} + \underbrace{\cdots + 1_R + \cdots + 1_R}_{n \text{ summands}}) *_R (\underbrace{1_R + \cdots + 1_R}_{m \text{ summands}} + \underbrace{\cdots + 1_R + \cdots + 1_R}_{n \text{ summands}}) \\ &= \underbrace{1_R * (\underbrace{1_R + \cdots + 1_R}_{n \text{ summands}}) + \cdots + 1_R * (\underbrace{1_R + \cdots + 1_R}_{n \text{ summands}})}_{m \text{ summands}} \\ &= \underbrace{1_R + \cdots + 1_R}_{mn \text{ summands}} = f(m * n). \end{aligned}$	$\begin{aligned} f(m) *_R f(n) &= (\underbrace{1_R + \cdots + 1_R}_{m \text{ summands}} *_R \underbrace{(1_R + \cdots + 1_R)_{n \text{ summands}}}_{n \text{ summands}}) \\ &= \underbrace{1_R * (\underbrace{1_R + \cdots + 1_R}_{n \text{ summands}}) + \cdots + 1_R * (\underbrace{1_R + \cdots + 1_R}_{n \text{ summands}})}_{m \text{ summands}} \\ &= \underbrace{1_R + \cdots + 1_R}_{mn \text{ summands}} = f(m * n). \end{aligned}$	$\begin{aligned} f(m) *_R f(n) &= (\underbrace{1_R + \cdots + 1_R}_{m \text{ summands}} *_R \underbrace{(1_R + \cdots + 1_R)_{n \text{ summands}}}_{n \text{ summands}}) \\ &= \underbrace{1_R * (\underbrace{1_R + \cdots + 1_R}_{n \text{ summands}}) + \cdots + 1_R * (\underbrace{1_R + \cdots + 1_R}_{n \text{ summands}})}_{m \text{ summands}} \\ &= \underbrace{1_R + \cdots + 1_R}_{mn \text{ summands}} = f(m * n). \end{aligned}$
fong-spivak-invitation_EQ0146	230	■ 0.999	$(f \circ i g) := (\mathrm{id} \otimes, \dots, \otimes \mathrm{id} \otimes g \otimes \mathrm{id} \otimes, \dots, \otimes \mathrm{id}); f,$	$(f \circ i g) := (\mathrm{id} \otimes, \dots, \otimes \mathrm{id} \otimes g \otimes \mathrm{id} \otimes, \dots, \otimes \mathrm{id}); f,$	$(f \circ_i g) := (\mathrm{id} \otimes, \dots, \otimes \mathrm{id} \otimes g \otimes \mathrm{id} \otimes, \dots, \otimes \mathrm{id}) \circ f,$
fong-spivak-invitation_EQ0015	47	■ 0.999	$f\left(g\left(q_{\{0\}}\right)\right)=\bigwedge\left\{q \in Q \mid g\left(q_{\{0\}}\right) \leq g(q)\right\} \leq \bigwedge\left\{q_{\{0\}}=q_{\{0\}},\right.$	$f(g(q_0)) = \bigwedge \{q \in Q \mid g(q_0) \leq g(q)\} \leq \bigwedge \{q_0\} = q_0,$	$f(g(q_0)) = \bigwedge \{q \in Q \mid g(q_0) \leq g(q)\} \leq \bigwedge \{q_0\} = q_0,$
fong-spivak-invitation_EQ0081 (4.3)	134	■ 0.999	$b \wedge c \leq d \quad \text{iff} \quad b \leq (c \Rightarrow d).$	$b \wedge c \leq d \quad \text{iff} \quad b \leq (c \Rightarrow d).$	$b \wedge c \leq d \quad \text{iff} \quad b \leq (c \Rightarrow d).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_EQ0091 (4.10)	142	■ 0.999	$\begin{aligned} \mathcal{P}\left(p, p_{\{1\}}\right) \otimes \Phi\left(p_{\{1\}}, q\right) &= \mathcal{P}\left(p, p_{\{1\}}\right) \otimes \Phi\left(p_{\{1\}}, q\right) \otimes I \\ &\leq \mathcal{P}\left(p, p_{\{1\}}\right) \otimes \Phi\left(p_{\{1\}}, q\right) \otimes \mathcal{Q}(q, q) \leq \Phi(p, q) . \end{aligned}$	$\mathcal{P}\left(p, p_{\{1\}}\right) \otimes \Phi\left(p_{\{1\}}, q\right) = \mathcal{P}\left(p, p_{\{1\}}\right) \otimes \Phi\left(p_{\{1\}}, q\right) \otimes I \\ \leq \mathcal{P}\left(p, p_{\{1\}}\right) \otimes \Phi\left(p_{\{1\}}, q\right) \otimes \mathcal{Q}(q, q) \leq \Phi(p, q) .$	$\mathcal{P}(p, p_1) \otimes \Phi(p_1, q) = \mathcal{P}(p, p_1) \otimes \Phi(p_1, q) \otimes I \\ \leq \mathcal{P}(p, p_1) \otimes \Phi(p_1, q) \otimes \mathcal{Q}(q, q) \leq \Phi(p, q).$
fong-spivak-invitation_EQ0157	248	■ 0.999	$\mathbf{p}_{\{1\}}:=\{\text{varnothing},\{1\},X\} \quad \text{and} \quad \mathbf{p}_{\{2\}}:=\{\text{varnothing},\{2\},X\} .$	$\mathbf{Op}_1 := \{\emptyset, \{1\}, X\} \quad \text{and} \quad \mathbf{Op}_2 := \{\emptyset, \{2\}, X\}.$	$\mathbf{Op}_1 := \{\emptyset, \{1\}, X\} \quad \text{and} \quad \mathbf{Op}_2 := \{\emptyset, \{2\}, X\} .$
fong-spivak-invitation_EQ0183	276	■ 0.999	$f^{\ast}(P)=\{\{x \in \mathbb{Z} \mid x \leq 0\},\{x \in \mathbb{Z} \mid x \geq 1\}\} \quad \text{and} \quad f^{\ast}(Q)=\{\{0\},\{x \in \mathbb{Z} \mid x \neq 0\}\} .$	$f^{\ast}(P)=\{\{x \in \mathbb{Z} \mid x \leq 0\},\{x \in \mathbb{Z} \mid x \geq 1\}\} \quad \text{and} \quad f^{\ast}(Q)=\{\{0\},\{x \in \mathbb{Z} \mid x \neq 0\}\}.$	$f^{\ast}(P)=\{\{x \in \mathbb{Z} \mid x \leq 0\},\{x \in \mathbb{Z} \mid x \geq 1\}\} \quad \text{and} \quad f^{\ast}(Q)=\{\{0\},\{x \in \mathbb{Z} \mid x \neq 0\}\}.$
fong-spivak-invitation_EQ0245	316	■ 0.999	$B+C:=\left\{(w, y, x, z) \in R^{m+p} \times R^{n+q} \mid (w, x) \in B \text { and } (y, z) \in C\right\} .$	$B+C:=\left\{(w, y, x, z) \in R^{m+p} \times R^{n+q} \mid (w, x) \in B \text { and } (y, z) \in C\right\} .$	$B+C:=\{(w, y, x, z) \in R^{m+p} \times R^{n+q} \mid (w, x) \in B \text { and } (y, z) \in C\}.$
fong-spivak-invitation_EQ0039	78	■ 0.999	$\begin{array}{l} I_W \leq f(I_V) \quad (f \text { is monoidal monotone }) \\ \leq f(\mathcal{C}(c, c)) \quad (\mathcal{C} \text { is } \mathcal{V}\text{-category }) \\ = \mathcal{C}_f(c, c) \quad (\text { definition of } \mathcal{C}_f); \end{array}$	$\begin{array}{l} I_W \leq f(I_V) \quad (f \text { is monoidal monotone }) \\ \leq f(\mathcal{C}(c, c)) \quad (\mathcal{C} \text { is } \mathcal{V}\text{-category }) \\ = \mathcal{C}_f(c, c) \quad (\text { definition of } \mathcal{C}_f); \end{array}$	$\begin{array}{l} I_W \leq f(I_V) \quad (f \text { is monoidal monotone }) \\ \leq f(\mathcal{C}(c, c)) \quad (\mathcal{C} \text { is } \mathcal{V}\text{-category }) \\ = \mathcal{C}_f(c, c) \quad (\text { definition of } \mathcal{C}_f); \end{array}$
fong-spivak-invitation_EQ0142 (6.16)	221	■ 0.999	$\begin{aligned} \psi_{V, V'}: \text{Circ}(V) \times \text{Circ}(V') &\longrightarrow \text{Circ}(V+V'); \\ ((V, A, s, t, \ell), (V', A', s', t', \ell')) &\longmapsto (V+V', A+A', s+s', t+t', [\ell, \ell']). \end{aligned}$	$\begin{array}{l} \psi_{V, V'}: \text{Circ}(V) \times \text{Circ}(V') \longrightarrow \text{Circ}(V+V'); \\ ((V, A, s, t, \ell), (V', A', s', t', \ell')) \longmapsto (V+V', A+A', s+s', t+t', [\ell, \ell']). \end{array}$	$\begin{array}{l} \psi_{V, V'}: \text{Circ}(V) \times \text{Circ}(V') \longrightarrow \text{Circ}(V+V'); \\ ((V, A, s, t, \ell), (V', A', s', t', \ell')) \longmapsto (V+V', A+A', s+s', t+t', [\ell, \ell']). \end{array}$
fong-spivak-invitation_EQ0052 (2.26)	87	■ 1.000	$(M * N)(x, z):=\bigvee_{y \in Y} M(x, y) \otimes N(y, z) .$	$(M * N)(x, z):=\bigvee_{y \in Y} M(x, y) \otimes N(y, z) .$	$(M * N)(x, z):=\bigvee_{y \in Y} M(x, y) \otimes N(y, z) .$
fong-spivak-invitation_EQ0054	87	■ 1.000	$I_X(x, y):=\begin{cases} I & \text { if } x=y \\ 0 & \text { if } x \neq y \end{cases}$	$I_X(x, y):=\begin{cases} I & \text { if } x=y \\ 0 & \text { if } x \neq y \end{cases}$	$I_X(x, y):=\begin{cases} I & \text { if } x=y, \\ 0 & \text { if } x \neq y. \end{cases}$
fong-spivak-invitation_EQ0063	113	■ 1.000	$\alpha_{\text {Vertex }}: G(\text {Vertex }) \rightarrow H(\text {Vertex }) \quad \text {and} \quad \alpha_{\text {Arrow }}: G(\text {Arrow }) \rightarrow H(\text {Arrow }),$	$\alpha_{\text {Vertex }}: G(\text {Vertex }) \rightarrow H(\text {Vertex }) \quad \text {and} \quad \alpha_{\text {Arrow }}: G(\text {Arrow }) \rightarrow H(\text {Arrow }),$	$\alpha_{\text {Vertex }}: G(\text {Vertex }) \rightarrow H(\text {Vertex }) \quad \text {and} \quad \alpha_{\text {Arrow }}: G(\text {Arrow }) \rightarrow H(\text {Arrow }),$
fong-spivak-invitation_EQ0067 (3.12)	116	■ 1.000	$\text{Gr} \xrightarrow{F} \text{DDS} \xrightarrow{I} \text{Set}.$	$\text{Gr} \xrightarrow{F} \text{DDS} \xrightarrow{I} \text{Set}.$	$\text{Gr} \xrightarrow{F} \text{DDS} \xrightarrow{I} \text{Set}.$
fong-spivak-invitation_EQ0181	273	■ 1.000	$1 \leq 1, \quad 1 \leq 2, \quad 1 \leq 3, \quad 2 \leq 2, \quad 2 \leq 3, \quad 3 \leq 3, \quad 4 \leq 4 .$	$1 \leq 1, \quad 1 \leq 2, \quad 1 \leq 3, \quad 2 \leq 2, \quad 2 \leq 3, \quad 3 \leq 3, \quad 4 \leq 4 .$	$1 \leq 1, \quad 1 \leq 2, \quad 1 \leq 3, \quad 2 \leq 2, \quad 2 \leq 3, \quad 3 \leq 3, \quad 4 \leq 4 .$
fong-spivak-invitation_EQ0220	301	■ 1.000	$\begin{aligned} &= \left(\begin{array}{ccc} 17 & 20 & 20 \\ 11 & 14 & 14 \\ 14 & 17 & 17 \\ 12 & 9 & 15 \end{array} \right) . \end{aligned}$	$= \left(\begin{array}{ccc} 17 & 20 & 20 \\ 11 & 14 & 14 \\ 14 & 17 & 17 \\ 12 & 9 & 15 \end{array} \right) .$	$= \left(\begin{array}{ccc} 17 & 20 & 20 \\ 11 & 14 & 14 \\ 14 & 17 & 17 \\ 12 & 9 & 15 \end{array} \right) .$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_EQ0258	324	1.000	$\begin{aligned} \varphi_{S',T'}(\varphi_{S,T}(A,B)) &= \varphi_{S',T'}(\{f(a) \mid a \in A\} \times \{g(b) \mid b \in B\}) \\ &= \{(f(a),g(b)) \mid a \in A, b \in B\} \\ &= \operatorname{im}_{f \times g}(A \times B) \\ &= \operatorname{im}_{f \times g}(\varphi_{S,T}(A,B)). \end{aligned}$	$\varphi_{S',T'}((\operatorname{im}_f \times \operatorname{im}_g)(A \times B)) = \varphi_{S',T'}(\{f(a) \mid a \in A\} \times \{g(b) \mid b \in B\})$ $= \{(f(a),g(b)) \mid a \in A, b \in B\}$ $= \operatorname{im}_{f \times g}(A \times B)$ $= \operatorname{im}_{f \times g}(\varphi_{S,T}(A,B)).$	$\varphi_{S',T'}((\operatorname{im}_f \times \operatorname{im}_g)(A \times B)) = \varphi_{S',T'}(\{f(a) \mid a \in A\} \times \{g(b) \mid b \in B\})$ $= \{(f(a),g(b)) \mid a \in A, b \in B\}$ $= \operatorname{im}_{f \times g}(A \times B)$ $= \operatorname{im}_{f \times g}(\varphi_{S,T}(A,B)).$
fong-spivak-invitation_EQ0001 (1.4)	20	1.000	$\Phi(A) \vee \Phi(B) \leq \Phi(A \vee B) .$	$\Phi(A) \vee \Phi(B) \leq \Phi(A \vee B).$	$\Phi(A) \vee \Phi(B) \leq \Phi(A \vee B).$
fong-spivak-invitation_EQ0004	26	1.000	$\begin{aligned} f(11) &= a, \quad f(12) = a, \quad f(13) = b, \quad f(21) = c, \\ f(22) &= d, \quad f(23) = d . \end{aligned}$	$f(11) = a, \quad f(12) = a, \quad f(13) = b, \quad f(21) = c, \quad f(22) = d, \quad f(23) = d.$	$f(11) = a, \quad f(12) = a, \quad f(13) = b, \quad f(21) = c, \quad f(22) = d, \quad f(23) = d.$
fong-spivak-invitation_EQ0006	37	1.000	$\left\{ \left. \frac{1}{n+1} \right n \in \mathbb{N} \right\} = \left\{ 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right\} \subseteq \mathbb{R}$	$\left\{ \left. \frac{1}{n+1} \right n \in \mathbb{N} \right\} = \left\{ 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right\} \subseteq \mathbb{R}$	$\left\{ \frac{1}{n+1} \mid n \in \mathbb{N} \right\} = \left\{ 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots \right\} \subseteq \mathbb{R}$
fong-spivak-invitation_EQ0012	45	1.000	$f(\bigvee A) \cong \bigvee f(A)$	$f(\bigvee A) \cong \bigvee f(A)$	$f\left(\bigvee A\right) \cong \bigvee f(A).$
fong-spivak-invitation_EQ0013 (1.8)	46	1.000	$f(p) := \bigwedge \{q \in Q \mid p \leq g(q)\} ;$	$f(p) := \bigwedge \{q \in Q \mid p \leq g(q)\};$	$f(p) := \bigwedge \{q \in Q \mid p \leq g(q)\};$
fong-spivak-invitation_EQ0014	47	1.000	$p_{\{0\}} \leq \bigwedge \{g(q) \in P \mid p_{\{0\}} \leq g(q)\} \cong g\left(\bigwedge \{q \in Q \mid p_{\{0\}} \leq g(q)\}\right) = g(f(p_0)),$	$p_0 \leq \bigwedge \{g(q) \in P \mid p_0 \leq g(q)\} \cong g\left(\bigwedge \{q \in Q \mid p_0 \leq g(q)\}\right) = g(f(p_0)),$	$p_0 \leq \bigwedge \{g(q) \in P \mid p_0 \leq g(q)\} \cong g\left(\bigwedge \{q \in Q \mid p_0 \leq g(q)\}\right) = g(f(p_0)),$
fong-spivak-invitation_EQ0019 (2.2)	55	1.000	$m * e = m, \quad e * m = m, \quad (m * n) * p = m * (n * p)$	$m * e = m, \quad e * m = m, \quad (m * n) * p = m * (n * p)$	$m * e = m, \quad e * m = m, \quad (m * n) * p = m * (n * p)$
fong-spivak-invitation_EQ0026 (2.8)	60	1.000	$t \leq v + w, \quad w + u \leq x + z, \quad v + x \leq y,$	$t \leq v + w, \quad w + u \leq x + z, \quad v + x \leq y,$	$t \leq v + w, \quad w + u \leq x + z, \quad v + x \leq y,$
fong-spivak-invitation_EQ0027 (2.9)	60	1.000	$t + u \leq y + z .$	$t + u \leq y + z .$	$t + u \leq y + z .$
fong-spivak-invitation_EQ0028 (2.10)	60	1.000	$t + u \leq v + w + u \leq v + x + z \leq y + z .$	$t + u \leq v + w + u \leq v + x + z \leq y + z .$	$t + u \leq v + w + u \leq v + x + z \leq y + z .$
fong-spivak-invitation_EQ0029	61	1.000	$\mathrm{H}_2 \mathrm{O}, \quad \mathrm{NaCl}, \quad 2 \mathrm{NaOH}, \quad \mathrm{CH}_4 + 3 \mathrm{O}_2$	$\mathrm{H}_2 \mathrm{O}, \quad \mathrm{NaCl}, \quad 2 \mathrm{NaOH}, \quad \mathrm{CH}_4 + 3 \mathrm{O}_2$	$\mathrm{H}_2 \mathrm{O}, \quad \mathrm{NaCl}, \quad 2 \mathrm{NaOH}, \quad \mathrm{CH}_4 + 3 \mathrm{O}_2$
fong-spivak-invitation_EQ0030	61	1.000	$2 \mathrm{H}_2 \mathrm{O} + 2 \mathrm{Na} \rightarrow 2 \mathrm{NaOH} + \mathrm{H}_2.$	$2 \mathrm{H}_2 \mathrm{O} + 2 \mathrm{Na} \rightarrow 2 \mathrm{NaOH} + \mathrm{H}_2.$	$2 \mathrm{H}_2 \mathrm{O} + 2 \mathrm{Na} \rightarrow 2 \mathrm{NaOH} + \mathrm{H}_2.$
fong-spivak-invitation_EQ0031 (2.11)	62	1.000	$y + k \rightarrow y' + k', \quad x + y' \rightarrow z', \quad z' + k' \rightarrow z + k.$	$y + k \rightarrow y' + k', \quad x + y' \rightarrow z', \quad z' + k' \rightarrow z + k.$	$y + k \rightarrow y' + k', \quad x + y' \rightarrow z', \quad z' + k' \rightarrow z + k.$
fong-spivak-invitation_EQ0032 (2.12)	62	1.000	$x + y + k \rightarrow x + y' + k' \rightarrow z' + k' \rightarrow z + k.$	$x + y + k \rightarrow x + y' + k' \rightarrow z' + k' \rightarrow z + k.$	$x + y + k \rightarrow x + y' + k' \rightarrow z' + k' \rightarrow z + k.$
fong-spivak-invitation_EQ0033 (2.14)	63	1.000	$\mathrm{H}_2 \mathrm{O} + \mathrm{NaCl} \rightarrow ? \mathrm{H}_2 \mathrm{O}$	$\mathrm{H}_2 \mathrm{O} + \mathrm{NaCl} \rightarrow ? \mathrm{H}_2 \mathrm{O}$	$\mathrm{H}_2 \mathrm{O} + \mathrm{NaCl} \rightarrow ? \mathrm{H}_2 \mathrm{O}$
fong-spivak-invitation_EQ0035	69	1.000	$d(x) := \begin{cases} \text{false} & \text{if } x > 0, \\ \text{true} & \text{if } x = 0, \end{cases} \quad u(x) := \begin{cases} \text{false} & \text{if } x = \infty, \\ \text{true} & \text{if } x < \infty. \end{cases}$	$d(x) := \begin{cases} \text{false} & \text{if } x > 0, \\ \text{true} & \text{if } x = 0, \end{cases} \quad u(x) := \begin{cases} \text{false} & \text{if } x = \infty, \\ \text{true} & \text{if } x < \infty. \end{cases}$	$d(x) := \begin{cases} \text{false} & \text{if } x > 0, \\ \text{true} & \text{if } x = 0, \end{cases} \quad u(x) := \begin{cases} \text{false} & \text{if } x = \infty, \\ \text{true} & \text{if } x < \infty. \end{cases}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_EQ0036	73	■ 1.000	$d_{\{L\}}(U, V) := \sup_{u \in U} \inf_{v \in V} d(u, v),$	$d_L(U, V) := \sup_{u \in U} \inf_{v \in V} d(u, v),$	$d_L(U, V) := \sup_{u \in U} \inf_{v \in V} d(u, v),$
fong-spivak-invitation_EQ0038 (2.20)	76	■ 1.000	$M_{\{Y\}} := \begin{array}{c ccc} & x & y & z \\ \hline x & 0 & 4 & 3 \\ y & 3 & 0 & \infty \\ z & \infty & 4 & 0 \end{array}$	$M_Y := \begin{array}{c ccc} \nearrow & x & y & z \\ \hline x & 0 & 4 & 3 \\ y & 3 & 0 & \infty \\ z & \infty & 4 & 0 \end{array}$	$M_Y := \begin{array}{c ccc} \nearrow & x & y & z \\ \hline x & 0 & 4 & 3 \\ y & 3 & 0 & \infty \\ z & \infty & 4 & 0 \end{array}$
fong-spivak-invitation_EQ0041 (2.21)	78	■ 1.000	$f(x) := \begin{cases} \text{true} & \text{if } x = 0, \\ \text{false} & \text{if } x > 0. \end{cases}$	$f(x) := \begin{cases} \text{true} & \text{if } x = 0, \\ \text{false} & \text{if } x > 0. \end{cases}$	$f(x) := \begin{cases} \text{true} & \text{if } x = 0, \\ \text{false} & \text{if } x > 0. \end{cases}$
fong-spivak-invitation_EQ0044 (2.23)	82	■ 1.000	$(a \otimes v) \leq w \quad \text{iff} \quad a \leq (v \multimap w)$	$(a \otimes v) \leq w \quad \text{iff} \quad a \leq (v \multimap w)$	$(a \otimes v) \leq w \quad \text{iff} \quad a \leq (v \multimap w)$
fong-spivak-invitation_EQ0045	83	■ 1.000	$a + x \geq y \quad \text{iff} \quad a \geq y - x \quad \text{iff} \quad \max(0, a) \geq \max(0, y - x) \quad \text{iff} \quad a \geq (x \multimap y)$	$a + x \geq y \quad \text{iff} \quad a \geq y - x \quad \text{iff} \quad \max(0, a) \geq \max(0, y - x) \quad \text{iff} \quad a \geq (x \multimap y)$	$a + x \geq y \quad \text{iff} \quad a \geq y - x \quad \text{iff} \quad \max(0, a) \geq \max(0, y - x) \quad \text{iff} \quad a \geq (x \multimap y)$
fong-spivak-invitation_EQ0050	86	■ 1.000	$v \multimap w := \bigvee_{\{a \in V \mid a \otimes v \leq w\}} a.$	$v \multimap w := \bigvee_{\{a \in V \mid a \otimes v \leq w\}} a.$	$v \multimap w := \bigvee_{\{a \in V \mid a \otimes v \leq w\}} a.$
fong-spivak-invitation_EQ0051 (2.25)	87	■ 1.000	$(M * N)(i, k) = \sum_j M(i, j) * N(j, k).$	$(M * N)(i, k) = \sum_j M(i, j) * N(j, k).$	$(M * N)(i, k) = \sum_j M(i, j) * N(j, k).$
fong-spivak-invitation_EQ0058	107	■ 1.000	$f^{\prime}; h^{\prime} = F(f); F(h) = F(f; h) = F(g; i) = F(g); F(i) = g^{\prime}; i^{\prime}.$	$f^{\prime}; h^{\prime} = F(f); F(h) = F(f; h) = F(g; i) = F(g); F(i) = g^{\prime}; i^{\prime}.$	$f^{\prime}; h^{\prime} = F(f); F(h) = F(f; h) = F(g; i) = F(g); F(i) = g^{\prime}; i^{\prime}.$
fong-spivak-invitation_EQ0060 (3.9)	110	■ 1.000	$F(f); \alpha_d = \alpha_c; G(f).$	$F(f); \alpha_d = \alpha_c; G(f).$	$F(f); \alpha_d = \alpha_c; G(f).$
fong-spivak-invitation_EQ0078	124	■ 1.000	$X \times Y := \{(x, y) \mid x \in X, y \in Y\},$	$X \times Y := \{(x, y) \mid x \in X, y \in Y\},$	$X \times Y := \{(x, y) \mid x \in X, y \in Y\},$
fong-spivak-invitation_EQ0084 (4.5)	137	■ 1.000	$\Phi(B, x) = 11, \quad \Phi(A, z) = 20, \quad \Phi(C, y) = 17.$	$\Phi(B, x) = 11, \quad \Phi(A, z) = 20, \quad \Phi(C, y) = 17.$	$\Phi(B, x) = 11, \quad \Phi(A, z) = 20, \quad \Phi(C, y) = 17.$
fong-spivak-invitation_EQ0086 (4.7)	140	■ 1.000	$(\Phi; \Psi)(p, r) := \bigvee_{q \in Q} \Phi(p, q) \wedge \Psi(q, r).$	$(\Phi; \Psi)(p, r) := \bigvee_{q \in Q} \Phi(p, q) \wedge \Psi(q, r).$	$(\Phi; \Psi)(p, r) := \bigvee_{q \in Q} \Phi(p, q) \wedge \Psi(q, r).$
fong-spivak-invitation_EQ0087	141	■ 1.000	$(\Phi; \Psi)(p, r) = \bigvee_{q \in Q} (\Phi(p, q) \otimes \Psi(q, r)).$	$(\Phi; \Psi)(p, r) = \bigvee_{q \in Q} (\Phi(p, q) \otimes \Psi(q, r)).$	$(\Phi; \Psi)(p, r) = \bigvee_{q \in Q} (\Phi(p, q) \otimes \Psi(q, r)).$
fong-spivak-invitation_EQ0098	150	■ 1.000	$\alpha_{s,t,u} : \{(s, (t, u)) \mid s \in S, t \in T, u \in U\} \xrightarrow{\cong} \{((s, t), u) \mid s \in S, t \in T, u \in U\}.$	$\alpha_{s,t,u} : \{(s, (t, u)) \mid s \in S, t \in T, u \in U\} \xrightarrow{\cong} \{((s, t), u) \mid s \in S, t \in T, u \in U\}.$	$\alpha_{s,t,u} : \{(s, (t, u)) \mid s \in S, t \in T, u \in U\} \xrightarrow{\cong} \{((s, t), u) \mid s \in S, t \in T, u \in U\}.$
fong-spivak-invitation_EQ0102	158	■ 1.000	$(\Phi \times \Psi)(\left(\left(x_1, y_1\right), \left(x_2, y_2\right)\right) := \Phi\left(x_1, x_2\right) \otimes \Psi\left(y_1, y_2\right).$	$(\Phi \times \Psi)((x_1, y_1), (x_2, y_2)) := \Phi(x_1, x_2) \otimes \Psi(y_1, y_2).$	$(\Phi \times \Psi)((x_1, y_1), (x_2, y_2)) := \Phi(x_1, x_2) \otimes \Psi(y_1, y_2).$
fong-spivak-invitation_EQ0105	164	■ 1.000	$R; S := \{(i, k) \in \underline{m} \times \underline{p} \mid \exists(j \in \underline{n}).(i, j) \in R \text{ and } (j, k) \in S\}.$	$R; S := \{(i, k) \in \underline{m} \times \underline{p} \mid \exists(j \in \underline{n}).(i, j) \in R \text{ and } (j, k) \in S\}.$	$R; S := \{(i, k) \in \underline{m} \times \underline{p} \mid \exists(j \in \underline{n}).(i, j) \in R \text{ and } (j, k) \in S\}.$
fong-spivak-invitation_EQ0109 (5.5)	170	■ 1.000	$G := \left\{ \rho_{m,n} \mid m, n \in \mathbb{N} \right\}, \quad s(\rho_{m,n}) := m, \quad t(\rho_{m,n}) := n,$	$G := \{ \rho_{m,n} \mid m, n \in \mathbb{N} \}, \quad s(\rho_{m,n}) := m, \quad t(\rho_{m,n}) := n,$	$G := \{ \rho_{m,n} \mid m, n \in \mathbb{N} \}, \quad s(\rho_{m,n}) := m, \quad t(\rho_{m,n}) := n,$
fong-spivak-invitation_EQ0115	178	■ 1.000	$A = \left(\begin{array}{ccc} 3 & 3 & 1 \\ 2 & 0 & 4 \end{array} \right), \quad B = \left(\begin{array}{cccc} 2 & 5 & 6 & 1 \end{array} \right).$	$A = \left(\begin{array}{ccc} 3 & 3 & 1 \\ 2 & 0 & 4 \end{array} \right), \quad B = \left(\begin{array}{cccc} 2 & 5 & 6 & 1 \end{array} \right).$	$A = \left(\begin{array}{ccc} 3 & 3 & 1 \\ 2 & 0 & 4 \end{array} \right), \quad B = \left(\begin{array}{cccc} 2 & 5 & 6 & 1 \end{array} \right).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation_EQ0117	181	■1.000	$\sum_{j \in \underline{n}} \alpha(i, j) * \beta(j, k) \text{ .}$	$\sum_{j \in \underline{n}} \alpha(i, j) * \beta(j, k).$	$\sum_{j \in \underline{n}} \alpha(i, j) * \beta(j, k).$
fong-spivak-invitation_EQ0120 (5.11)	188	■1.000	$\mathrm{B}(g)=\left\{(x, S(g)(x)) \mid x \in R^m\right\} \subseteq R^m \times R^n \text{ .}$	$B(g)=\{(x, S(g)(x)) \mid x \in R^m\} \subseteq R^m \times R^n.$	$B(g)=\{(x, S(g)(x)) \mid x \in R^m\} \subseteq R^m \times R^n.$
fong-spivak-invitation_EQ0121 (5.12)	188	■1.000	$\mathrm{B}\backslash\left(g^{\mathrm{op}}\backslash\right):=\left\{(S(g)(x), x) \mid x \in R^m\right\} \subseteq R^n \times R^m \text{ .}$	$B\left(g^{\mathrm{op}}\right):=\{(S(g)(x), x) \mid x \in R^m\} \subseteq R^n \times R^m.$	$B(g^{\mathrm{op}}):=\{(S(g)(x), x) \mid x \in R^m\} \subseteq R^n \times R^m.$
fong-spivak-invitation_EQ0125	190	■1.000	$\mathrm{B}\backslash\left(\left(\left(g^{\mathrm{op}}\backslash\right) ; h\right)=\left\{(S(g)(x), S(h)(x)) \mid x \in R^m\right\} \text{ .}$	$B\left(\left(g^{\mathrm{op}}\right) ; h\right)=\{(S(g)(x), S(h)(x)) \mid x \in R^m\}.$	$B((g^{\mathrm{op}}) \circledast h)=\{(S(g)(x), S(h)(x)) \mid x \in R^m\}.$
fong-spivak-invitation_EQ0127	192	■1.000	$v=\int \frac{1}{m} F(t) d t+u(t)+p \int a u(t)+b v(t) d t \text{ .}$	$v=\int \frac{1}{m} F(t) d t+u(t)+p \int a u(t)+b v(t) d t.$	$v=\int \frac{1}{m} F(t) d t+u(t)+p \int a u(t)+b v(t) d t.$
fong-spivak-invitation_EQ0130	204	■1.000	$y(\theta)=y(g(\theta))=x(f(\theta))=x(\theta) \text{ .}$	$y(0)=y(g(0))=x(f(0))=x(0).$	$y(0)=y(g(0))=x(f(0))=x(0).$
fong-spivak-invitation_EQ0133	207	■1.000	$\{(v, d) \mid d \in D(v)\} \text{ .}$	$\{(v, d) \mid d \in D(v)\}.$	$\{(v, d) \mid d \in D(v)\}.$
fong-spivak-invitation_EQ0137 (6.15)	219	■1.000	$F\left(\left(\left[\iota_N, \iota_P\right]\right) \backslash\right)\left(\varphi_{N, P}(s, t)\right) \text{ .}$	$F\left([\iota_N, \iota_P]\right)\left(\varphi_{N, P}(s, t)\right) \text{ .}$	$F([\iota_N, \iota_P])\left(\varphi_{N, P}(s, t)\right).$
fong-spivak-invitation_EQ0140	221	■1.000	$\operatorname{Circ}(V):=\{(V, A, s, t, \ell) \mid \text { where } s, t: A \rightarrow V, \ell: E \rightarrow C\} \text{ .}$	$\operatorname{Circ}(V):=\{(V, A, s, t, \ell) \mid \textbf{ where } s, t: A \rightarrow V, \ell: E \rightarrow C\}.$	$\operatorname{Circ}(V):=\{(V, A, s, t, \ell) \mid \text { where } s, t: A \rightarrow V, \ell: E \rightarrow C\}.$
fong-spivak-invitation_EQ0147	230	■1.000	$F: \mathcal{O}\left(t_1, \ldots, t_n ; t\right) \rightarrow \mathcal{P}\left(f\left(t_1\right), \ldots, f\left(t_n\right) ; f(t)\right) \rightarrow \mathcal{P}\left(f\left(t_1\right), \ldots, f\left(t_n\right) ; f(t)\right) \text{ ; } f(t) \text{)}$	$F: \mathcal{O}\left(t_1, \ldots, t_n ; t\right) \rightarrow \mathcal{P}\left(f\left(t_1\right), \ldots, f\left(t_n\right) ; f(t)\right)$	$F: \mathcal{O}\left(t_1, \ldots, t_n ; t\right) \rightarrow \mathcal{P}\left(f\left(t_1\right), \ldots, f\left(t_n\right) ; f(t)\right)$
fong-spivak-invitation_EQ0154 (7.6)	241	■1.000	$X=\{y \in Y \mid p(y)=\text { true }\} \text{ .}$	$X=\{y \in Y \mid p(y)=\textbf{ true }\}.$	$X=\{y \in Y \mid p(y)=\textbf{true}\}.$
fong-spivak-invitation_EQ0158	249	■1.000	$\left.s_i\right _{U_i \cap U_j}=\left.s_j\right _{U_i \cap U_j} \text{ .}$	$s_i _{U_i \cap U_j}=s_j _{U_i \cap U_j} \text{ .}$	$s_i _{U_i \cap U_j}=s_j _{U_i \cap U_j}.$
fong-spivak-invitation_EQ0162 (7.14)	255	■1.000	$\begin{aligned} & \Omega(U) \rightarrow \Omega(V), \\ & U^{\prime} \mapsto U^{\prime} \cap V. \end{aligned}$	$\Omega(U) \rightarrow \Omega(V),$ $U' \mapsto U' \cap V.$	$\Omega(U) \rightarrow \Omega(V),$ $U' \mapsto U' \cap V.$
fong-spivak-invitation_EQ0163	256	■1.000	$V \cap U_j=\left(\bigcup_{i \in I} V_i\right) \cap U_j=\bigcup_{i \in I}\left(V_i \cap U_j\right)=\bigcup_{i \in I}\left(V_j \cap U_i\right)=\left(\bigcup_{i \in I} U_i\right) \cap V_j=V_j \text{ .}$	$V \cap U_j=\left(\bigcup_{i \in I} V_i\right) \cap U_j=\bigcup_{i \in I}\left(V_i \cap U_j\right)=\bigcup_{i \in I}\left(V_j \cap U_i\right)=\left(\bigcup_{i \in I} U_i\right) \cap V_j=V_j.$	$V \cap U_j=\left(\bigcup_{i \in I} V_i\right) \cap U_j=\bigcup_{i \in I}\left(V_i \cap U_j\right)=\bigcup_{i \in I}\left(V_j \cap U_i\right)=\left(\bigcup_{i \in I} U_i\right) \cap V_j=V_j.$
fong-spivak-invitation_EQ0169	261	■1.000	$p(n, z)=\begin{cases} \text { true } & \text { if } n \leq z , \\ \text { false } & \text { if } n> z \end{cases} \text{ .}$	$p(n, z)=\left\{\begin{array}{ll} \textbf{true} & \text { if } n \leq z , \\ \textbf{false} & \text { if } n> z . \end{array}\right.$	$p(n, z)=\left\{\begin{array}{ll} \textbf{true} & \text { if } n \leq z , \\ \textbf{false} & \text { if } n> z . \end{array}\right.$
fong-spivak-invitation_EQ0173	266	■1.000	$S\left(o_{[a, b]}\right) \cong \lim _{\epsilon>0} S\left(o_{[a-\epsilon, b+\epsilon]}\right)$	$S\left(o_{[a, b]}\right) \cong \lim _{\epsilon>0} S\left(o_{[a-\epsilon, b+\epsilon]}\right)$	$S(o_{[a, b]}) \cong \lim _{\epsilon>0} S(o_{[a-\epsilon, b+\epsilon]}).$
fong-spivak-invitation_EQ0175	271	■1.000	$\varnothing, \quad\{1\}, \quad\{2\}, \quad\{3\}, \quad\{1,2\}, \quad\{1,3\}, \quad\{2,3\}, \quad\{1,2,3\} \text{ .}$	$\varnothing, \quad\{1\}, \quad\{2\}, \quad\{3\}, \quad\{1,2\}, \quad\{1,3\}, \quad\{2,3\}, \quad\{1,2,3\}.$	$\varnothing, \quad\{1\}, \quad\{2\}, \quad\{3\}, \quad\{1,2\}, \quad\{1,3\}, \quad\{2,3\}, \quad\{1,2,3\}.$
fong-spivak-invitation_EQ0176	271	■1.000	$(h, 1), \quad\quad(h, 2), \quad\quad(h, 3), \quad\quad(1,1), \quad\quad(1,2), \quad\quad(1,3) \text{ .}$	$(h,1), \quad (h,2), \quad (h,3), \quad (1,1), \quad (1,2), \quad (1,3).$	$(h,1), \quad (h,2), \quad (h,3), \quad (1,1), \quad (1,2), \quad (1,3).$
fong-spivak-invitation_EQ0177	271	■1.000	$(h, 1), \quad\quad\quad(1,1), \quad\quad\quad(1,2), \quad\quad\quad(2,2), \quad\quad\quad(3,2) \text{ .}$	$(h,1), \quad (1,1), \quad (1,2), \quad (2,2), \quad (3,2).$	$(h,1), \quad (1,1), \quad (1,2), \quad (2,2), \quad (3,2).$
fong-spivak-invitation_EQ0178	271	■1.000	$h, \quad\quad 1, \quad\quad 2, \quad\quad 3 \text{ .}$	$h, \quad 1, \quad 2, \quad 3.$	$h, \quad 1, \quad 2, \quad 3.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation-EQ0179	272	■ 1.000	$\begin{array}{lllll} (11,11), & \& (11,12), & \& (12,11), & \& \\ (12,12), & \& (13,13), & \& (21,21), & \& (22,22), & \& (22,23), & \& \\ (23,22), & \& (23,23) . \end{array}$	$(11,11), (11,12), (12,11), (12,12), (13,13), (21,21), (22,22), (22,23), (23,22), (23,23).$	$(11,11), (11,12), (12,11), (12,12), (13,13), (21,21), (22,22), (22,23), (23,22), (23,23).$
fong-spivak-invitation-EQ0182	273	■ 1.000	$(\bullet)(\circ)(*), \quad \quad (\bullet)(\circ)(*), \quad \quad (\bullet)(\circ)(*), \quad \quad (\bullet)(\circ)(*), \quad \quad (\bullet)(\circ)(*) .$	$(\bullet)(\circ)(*), (\bullet)(\circ)(*), (\bullet)(\circ)(*), (\bullet)(\circ)(*), (\bullet)(\circ)(*) .$	$(\bullet)(\circ)(*), (\bullet)(\circ)(*), (\bullet)(\circ)(*), (\bullet)(\circ)(*), (\bullet)(\circ)(*) .$
fong-spivak-invitation-EQ0184	280	■ 1.000	$g(q) \leq g^{\prime}(f(g(q))) \leq g^{\prime}(q) .$	$g(q) \leq g'(f(g(q))) \leq g'(q).$	$g(q) \leq g'(f(g(q))) \leq g'(q).$
fong-spivak-invitation-EQ0188	288	■ 1.000	$A=(A \cap \bar{B}) \cup (A \cap B) \subseteq \bar{B} \cup C .$	$A = (A \cap \bar{B}) \cup (A \cap B) \subseteq \bar{B} \cup C.$	$A = (A \cap \bar{B}) \cup (A \cap B) \subseteq \bar{B} \cup C.$
fong-spivak-invitation-EQ0190	288	■ 1.000	$\left(\begin{array}{ll} 1 & 0 \\ 0 & 1 \end{array} \right), \quad \left(\begin{array}{ll} \text{true} & \text{false} \\ \text{false} & \text{true} \end{array} \right), \quad \text{and} \quad \left(\begin{array}{ll} 0 & \infty \\ \infty & 0 \end{array} \right) .$	$\left(\begin{array}{ll} 1 & 0 \\ 0 & 1 \end{array} \right), \quad \left(\begin{array}{ll} \text{true} & \text{false} \\ \text{false} & \text{true} \end{array} \right), \quad \text{and} \quad \left(\begin{array}{ll} 0 & \infty \\ \infty & 0 \end{array} \right) .$	$\left(\begin{array}{ll} 1 & 0 \\ 0 & 1 \end{array} \right), \quad \left(\begin{array}{ll} \text{true} & \text{false} \\ \text{false} & \text{true} \end{array} \right), \quad \text{and} \quad \left(\begin{array}{ll} 0 & \infty \\ \infty & 0 \end{array} \right) .$
fong-spivak-invitation-EQ0194	289	■ 1.000	$\begin{gathered} M_X = \left(\begin{array}{cccc} 0 & \infty & 3 & \infty \\ \infty & 3 & 0 & \infty \\ \infty & \infty & 6 & 0 \end{array} \right), \quad M_X^2 = \left(\begin{array}{cccc} 0 & 6 & 3 & \infty \\ 2 & 0 & 5 & 5 \\ 5 & 3 & 0 & 8 \\ \infty & 9 & 6 & 0 \end{array} \right), \\ M_X^3 = M_X^4 = \left(\begin{array}{cccc} 0 & 6 & 3 & 11 \\ 2 & 0 & 5 & 5 \\ 5 & 3 & 0 & 8 \\ 11 & 9 & 6 & 0 \end{array} \right) . \end{gathered}$	$M_X = \left(\begin{array}{cccc} 0 & \infty & 3 & \infty \\ 2 & 0 & \infty & 5 \\ \infty & 3 & 0 & \infty \\ \infty & \infty & 6 & 0 \end{array} \right), \quad M_X^2 = \left(\begin{array}{cccc} 0 & 6 & 3 & \infty \\ 2 & 0 & 5 & 5 \\ 5 & 3 & 0 & 8 \\ \infty & 9 & 6 & 0 \end{array} \right),$ $M_X^3 = M_X^4 = \left(\begin{array}{cccc} 0 & 6 & 3 & 11 \\ 2 & 0 & 5 & 5 \\ 5 & 3 & 0 & 8 \\ 11 & 9 & 6 & 0 \end{array} \right) .$	$M_X = \left(\begin{array}{cccc} 0 & \infty & 3 & \infty \\ 2 & 0 & \infty & 5 \\ \infty & 3 & 0 & \infty \\ \infty & \infty & 6 & 0 \end{array} \right), \quad M_X^2 = \left(\begin{array}{cccc} 0 & 6 & 3 & \infty \\ 2 & 0 & 5 & 5 \\ 5 & 3 & 0 & 8 \\ \infty & 9 & 6 & 0 \end{array} \right),$ $M_X^3 = M_X^4 = \left(\begin{array}{cccc} 0 & 6 & 3 & 11 \\ 2 & 0 & 5 & 5 \\ 5 & 3 & 0 & 8 \\ 11 & 9 & 6 & 0 \end{array} \right) .$
fong-spivak-invitation-EQ0196	290	■ 1.000	$(p ; q) := \left(v, a_1, \dots, a_m, b_1, \dots, b_n \right) .$	$(p;q) := (v,a_1,\dots,a_m,b_1,\dots,b_n) .$	$(p \circ q) := (v,a_1,\dots,a_m,b_1,\dots,b_n).$
fong-spivak-invitation-EQ0197	290	■ 1.000	$\left(v, a_1, \dots, a_m, b_1, \dots, b_n, c_1, \dots, c_o \right) .$	$(v,a_1,\dots,a_m,b_1,\dots,b_n,c_1,\dots,c_o) .$	$(v,a_1,\dots,a_m,b_1,\dots,b_n,c_1,\dots,c_o).$
fong-spivak-invitation-EQ0203	292	■ 1.000	$\begin{array}{lll} (1,1) & \& (1,2) & \& (1,3) \\ (2,1) & \& (2,2) & \& (2,3) \\ (3,1) & \& (3,2) & \& (3,3) \end{array}$	$\begin{array}{lll} (1,1) & (1,2) & (1,3) \\ (2,1) & (2,2) & (2,3) \\ (3,1) & (3,2) & (3,3) \end{array}$	$\begin{array}{lll} (1,1) & (1,2) & (1,3) \\ (2,1) & (2,2) & (2,3) \\ (3,1) & (3,2) & (3,3) \end{array}$
fong-spivak-invitation-EQ0204	292	■ 1.000	$f^{-1}(1)=b, \quad \quad f^{-1}(2)=a, \quad \quad f^{-1}(3)=c .$	$f^{-1}(1) = b, \quad f^{-1}(2) = a, \quad f^{-1}(3) = c.$	$f^{-1}(1) = b, \quad f^{-1}(2) = a, \quad f^{-1}(3) = c.$
fong-spivak-invitation-EQ0205	293	■ 1.000	$\begin{gathered} A^{\prime} \mapsto A, \quad \quad f^{\prime} \mapsto f, \quad \quad g^{\prime} \mapsto g, \quad \quad f^{\prime} ; h^{\prime} \mapsto f ; h, \quad \quad g^{\prime} ; i^{\prime} \mapsto f ; h, \\ B^{\prime} \mapsto B, \quad \quad h^{\prime} \mapsto h, \quad \quad C^{\prime} \mapsto C, \quad \quad i^{\prime} \mapsto i, \quad \quad D^{\prime} \mapsto D. \end{gathered}$	$A' \mapsto A, \quad f' \mapsto f, \quad g' \mapsto g, \quad f';h' \mapsto f;h, \quad g';i' \mapsto f;h, \\ B' \mapsto B, \quad h' \mapsto h, \quad C' \mapsto C, \quad i' \mapsto i, \quad D' \mapsto D.$	$A' \mapsto A, \quad f' \mapsto f, \quad g' \mapsto g, \quad f' \circ h' \mapsto f \circ h, \quad g' \circ i' \mapsto f \circ h, \\ B' \mapsto B, \quad h' \mapsto h, \quad C' \mapsto C, \quad i' \mapsto i, \quad D' \mapsto D.$
fong-spivak-invitation-EQ0208	293	■ 1.000	$((F ; G) ; H)(c)=H((F ; G)(c))=H(G(F(c)))=(G ; H)(F(c))=(F ; (G ; H))(c) .$	$((F;G);H)(c) = H((F;G)(c)) = H(G(F(c))) = (G;H)(F(c)) = (F;(G;H))(c).$	$((F \circ G) \circ H)(c) = H((F \circ G)(c)) = H(G(F(c))) = (G \circ H)(F(c)) = (F \circ (G \circ H))(c).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation-EQ0211	297	■ 1.000	$\left(f_{\{1\}} ; f_{\{2\}}\right)^{\{B\}}(g)=g ; \left(f_{\{1\}} ; f_{\{2\}}\right)=\left(g ; f_{\{1\}}\right) ; f_{\{2\}}=\left(f_{\{1\}}^{\{B\}} ; f_{\{2\}}^{\{B\}}\right)(g)$	$(f_1;f_2)^B(g)=g;(f_1;f_2)=(g;f_1);f_2=(f_1^B;f_2^B)(g)$	$(f_1\circ f_2)^B(g)=g\circ(f_1\circ f_2)=(g\circ f_1)\circ f_2=(f_1^B\circ f_2^B)(g)$
fong-spivak-invitation-EQ0218	300	■ 1.000	$\begin{array}{c ccc} \Phi & x & y & z \\ \hline A & 17 & 20 & 20 \\ B & 11 & 14 & 14 \\ C & 14 & 17 & 17 \\ D & 12 & 9 & 15 \end{array}$	$\begin{array}{c ccc} \Phi & x & y & z \\ \hline A & 17 & 20 & 20 \\ B & 11 & 14 & 14 \\ C & 14 & 17 & 17 \\ D & 12 & 9 & 15 \end{array}$	$\begin{array}{c ccc} \Phi & x & y & z \\ \hline A & 17 & 20 & 20 \\ B & 11 & 14 & 14 \\ C & 14 & 17 & 17 \\ D & 12 & 9 & 15 \end{array}$
fong-spivak-invitation-EQ0229	307	■ 1.000	$(f+g)(i):=\begin{cases} f(i) & \text{if } i\leq m, \\ g(i-m)+m^{\prime} & \text{if } m+1\leq i. \end{cases}$	$(f+g)(i):=\begin{cases} f(i) & \text{if } i\leq m, \\ g(i-m)+m' & \text{if } m+1\leq i. \end{cases}$	$(f+g)(i):=\begin{cases} f(i) & \text{if } i\leq m, \\ g(i-m)+m' & \text{if } m+1\leq i. \end{cases}$
fong-spivak-invitation-EQ0230	310	■ 1.000	$a^{\{0\}}, a^{\{1\}}, a^{\{2\}}, a^{\{3\}}, \ldots, a^{\{2019\}}, \ldots$	$a^0,a^1,a^2,a^3,\ldots,a^{2019},\ldots$	$a^0,a^1,a^2,a^3,\ldots,a^{2019},\ldots$
fong-spivak-invitation-EQ0233	311	■ 1.000	$\left(\begin{array}{llll} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array}\right)$	$\left(\begin{array}{llll} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array}\right)$	$\left(\begin{array}{llll} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{array}\right)$
fong-spivak-invitation-EQ0234	311	■ 1.000	$A*B=\left(\begin{array}{ll} 0 & 1 \\ 0 & 0 \end{array}\right)*\left(\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array}\right)\neq\left(\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array}\right)*\left(\begin{array}{ll} 0 & 1 \\ 0 & 0 \end{array}\right)=B*A$	$A*B=\left(\begin{array}{ll} 0 & 1 \\ 0 & 0 \end{array}\right)*\left(\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array}\right)\neq\left(\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array}\right)*\left(\begin{array}{ll} 0 & 1 \\ 0 & 0 \end{array}\right)=B*A$	$A*B=\left(\begin{array}{ll} 0 & 1 \\ 0 & 0 \end{array}\right)*\left(\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array}\right)\neq\left(\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array}\right)*\left(\begin{array}{ll} 0 & 1 \\ 0 & 0 \end{array}\right)=B*A$
fong-spivak-invitation-EQ0236	312	■ 1.000	$\left(\begin{array}{lll} 1 & 1 & 1 \end{array}\right)$	$\left(\begin{array}{lll} 1 & 1 & 1 \end{array}\right)$	$\left(\begin{array}{lll} 1 & 1 & 1 \end{array}\right)$
fong-spivak-invitation-EQ0237	312	■ 1.000	$\left(\begin{array}{lll} 1 & 1 & 1 \end{array}\right)$	$\left(\begin{array}{lll} 1 & 1 & 1 \end{array}\right)$	$\left(\begin{array}{lll} 1 & 1 & 1 \end{array}\right)$
fong-spivak-invitation-EQ0238	313	■ 1.000	$g_{\{1\}}:=c_{\{n\}}+\cdots+c_{\{n\}}:m\rightarrow(m\times n),$	$g_1:=c_n+\cdots+c_n:m\rightarrow(m\times n),$	$g_1:=c_n+\cdots+c_n:m\rightarrow(m\times n),$
fong-spivak-invitation-EQ0240	313	■ 1.000	$\begin{aligned} g_{\{2\}}:=& s_{\{M(1,1)\}}+\cdots+s_{\{M(1,n)\}} \\ & +s_{\{M(2,1)\}}+\cdots+s_{\{M(2,n)\}} \\ & +\cdots \\ & +s_{\{M(m,1)\}}+\cdots+s_{\{M(m,n)\}}:(m\times n) \\ & \rightarrow(m\times n), \end{aligned}$	$\begin{aligned} g_2:=& s_{M(1,1)}+\cdots+s_{M(1,n)} \\ & +s_{M(2,1)}+\cdots+s_{M(2,n)} \\ & +\cdots \\ & +s_{M(m,1)}+\cdots+s_{M(m,n)}:(m\times n)\rightarrow(m\times n), \end{aligned}$	$\begin{aligned} g_2:=& s_{M(1,1)}+\cdots+s_{M(1,n)} \\ & +s_{M(2,1)}+\cdots+s_{M(2,n)} \\ & +\cdots \\ & +s_{M(m,1)}+\cdots+s_{M(m,n)}:(m\times n)\rightarrow(m\times n), \end{aligned}$
fong-spivak-invitation-EQ0241	313	■ 1.000	$g_{\{4\}}:=a_{\{m\}}+\cdots+a_{\{m\}}:(m\times n)\rightarrow n,$	$g_4:=a_m+\cdots+a_m:(m\times n)\rightarrow n,$	$g_4:=a_m+\cdots+a_m:(m\times n)\rightarrow n,$
fong-spivak-invitation-EQ0243	315	■ 1.000	$\mu:=\left(\begin{array}{ll} 1 & 0 \\ 0 & 1 \\ 1 & 0 \\ 0 & 1 \end{array}\right)$	$\mu:=\left(\begin{array}{ll} 1 & 0 \\ 0 & 1 \\ 1 & 0 \\ 0 & 1 \end{array}\right)$	$\mu:=\left(\begin{array}{ll} 1 & 0 \\ 0 & 1 \\ 1 & 0 \\ 0 & 1 \end{array}\right)$
fong-spivak-invitation-EQ0244	315	■ 1.000	$U(\mu)\left(\left(a_1,\ldots,a_m\right),\left(b_1,\ldots,b_m\right)\right):=\left(a_1+b_1,\ldots,a_m+b_m\right).$	$U(\mu)\left(\left(a_1,\ldots,a_m\right),\left(b_1,\ldots,b_m\right)\right):=\left(a_1+b_1,\ldots,a_m+b_m\right).$	$U(\mu)\left(\left(a_1,\ldots,a_m\right),\left(b_1,\ldots,b_m\right)\right):=\left(a_1+b_1,\ldots,a_m+b_m\right).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-invitation-EQ0246	316	1.000	$\begin{aligned} \mathrm{B}(g) &= \{(x, z) \in R^m \times R^n \mid S(g)(x) = z\}, \\ \mathrm{B}(h^{\mathrm{op}}) &= \{(z, y) \in R^n \times R^\ell \mid z = S(h)(y)\}, \end{aligned}$	$\begin{aligned} \mathrm{B}(g) &= \{(x, z) \in R^m \times R^n \mid S(g)(x) = z\}, \\ \mathrm{B}(h^{\mathrm{op}}) &= \{(z, y) \in R^n \times R^\ell \mid z = S(h)(y)\}, \end{aligned}$	$\begin{aligned} \mathrm{B}(g) &= \{(x, z) \in R^m \times R^n \mid S(g)(x) = z\}, \\ \mathrm{B}(h^{\mathrm{op}}) &= \{(z, y) \in R^n \times R^\ell \mid z = S(h)(y)\}, \end{aligned}$
fong-spivak-invitation-EQ0249	316	1.000	$\begin{aligned} \mathrm{B}\left(g^{\mathrm{op}}\right) &= \{(y, x) \in R^n \times R^m \mid y = S(g)(x)\}, \\ \mathrm{B}(h) &= \{(x, z) \in R^m \times R^p \mid S(h)(x) = z\}, \end{aligned}$	$\begin{aligned} \mathrm{B}(g^{\mathrm{op}}) &= \{(y, x) \in R^n \times R^m \mid y = S(g)(x)\}, \\ \mathrm{B}(h) &= \{(x, z) \in R^m \times R^p \mid S(h)(x) = z\}, \end{aligned}$	$\begin{aligned} \mathrm{B}(g^{\mathrm{op}}) &= \{(y, x) \in R^n \times R^m \mid y = S(g)(x)\}, \\ \mathrm{B}(h) &= \{(x, z) \in R^m \times R^p \mid S(h)(x) = z\}, \end{aligned}$
fong-spivak-invitation-EQ0251	316	1.000	$\mathrm{B}\left(\left(\left(g^{\mathrm{op}}\right) ; h\right) ; \right)=\left(\left(S(g)(x), S(h)(x)\right) \mid x \in R^m\right) .$	$\mathrm{B}\left(\left(g^{\mathrm{op}}\right) ; h\right)=\{(S(g)(x), S(h)(x)) \mid x \in R^m\} .$	$\mathrm{B}\left(\left(g^{\mathrm{op}}\right) ; h\right)=\{(S(g)(x), S(h)(x)) \mid x \in R^m\} .$
fong-spivak-invitation-EQ0252	318	1.000	$f(m)=f(\underbrace{1+1+\cdots+1}_m\text{ summands})=\underbrace{f(1)+f(1)+\cdots+f(1)}_m\text{ summands}=\underbrace{1_R+1_R+\cdots+1_R}_m\text{ summands} .$	$f(m)=f(\underbrace{1+1+\cdots+1}_m\text{ summands})=\underbrace{f(1)+f(1)+\cdots+f(1)}_m\text{ summands}=\underbrace{1_R+1_R+\cdots+1_R}_m\text{ summands} .$	$f(m)=f(\underbrace{1+1+\cdots+1}_m\text{ summands})=\underbrace{f(1)+f(1)+\cdots+f(1)}_m\text{ summands}=\underbrace{1_R+1_R+\cdots+1_R}_m\text{ summands} .$
fong-spivak-invitation-EQ0254	318	1.000	$\begin{aligned} [f, g]: A \sqcup B &\longrightarrow T \\ \text{apple } 1 &\longmapsto a \end{aligned}$	$\begin{aligned} [f, g]: A \sqcup B &\longrightarrow T \\ \text{apple } 1 &\longmapsto a \end{aligned}$	$\begin{aligned} [f, g]: A \sqcup B &\longrightarrow T \\ \text{apple } 1 &\longmapsto a \end{aligned}$
fong-spivak-invitation-EQ0261	338	1.000	$U=\bigcup_{i \in I}\left\{x \in \mathbb{R} \mid a_i<x<b_i\right\}$	$U=\bigcup_{i \in I}\left\{x \in \mathbb{R} \mid a_i<x<b_i\right\}$	$U=\bigcup_{i \in I}\left\{x \in \mathbb{R} \mid a_i<x<b_i\right\}$
fong-spivak-invitation-EQ0262	338	1.000	$U=\bigcup_{j \in J}\left\{x \in \mathbb{R} \mid a_{\{j\}}<x<b_{\{j\}}\right\}=\bigcup_{j \in J}\left(o_{\left\{\left[a_{\{j\}}, b_{\{j\}}\right]\right\}} \cap \mathbb{R}\right)=\left(\bigcup_{j \in J} o_{\left\{\left[a_{\{j\}}, b_{\{j\}}\right]\right\}}\right) \cap \mathbb{R}$	$U=\bigcup_{j \in J}\left\{x \in \mathbb{R} \mid a_j<x<b_j\right\}=\bigcup_{j \in J}\left(o_{\left[a_j, b_j\right]} \cap \mathbb{R}\right)=\left(\bigcup_{j \in J} o_{\left[a_j, b_j\right]}\right) \cap \mathbb{R}$	$U=\bigcup_{j \in J}\left\{x \in \mathbb{R} \mid a_j<x<b_j\right\}=\bigcup_{j \in J}\left(o_{\left[a_j, b_j\right]} \cap \mathbb{R}\right)=\left(\bigcup_{j \in J} o_{\left[a_j, b_j\right]}\right) \cap \mathbb{R}$
fong-spivak-invitation-EQ0064	114	—	—	—	$G :=$ $\begin{array}{c} 1 \\ \bullet \end{array} \xrightarrow{a} \begin{array}{c} 2 \\ \bullet \end{array} \xrightarrow{b} \begin{array}{c} 3 \\ \bullet \end{array}$ $H :=$ $\begin{array}{c} 4 \\ \bullet \end{array} \xrightarrow[c]{d} \begin{array}{c} 5 \\ \bullet \end{array} \curvearrowright e$
fong-spivak-invitation-EQ0065	114	—	—	—	$G :=$ Arrow source target a 1 b 2 2 3 3 4 $H :=$ Arrow source target c 4 d 4 e 5 5 5 4 5