

penev_B: formula evidence

The page, the confidence and the picture of an inline formula are its HOST LINE’s — a formula has none of its own. A line’s confidence is not a formula’s.

Inline formulas (first occurrence)

Identifier	Page	Conf.	LaTeX source	Rendered	Image
penev_B_F00001	1	■ 1.000	$\left\langle \mathbf{s}_{\parallel}, \mathbf{s}_{\perp} \right\rangle = 0$	$\langle \mathbf{s}_{\parallel}, \mathbf{s}_{\perp} \rangle = 0$	from (4.57), $\langle \mathbf{s}_{\parallel}, \mathbf{s}_{\perp} \rangle = 0$, hence $\ \mathbf{s}\ ^2 = \ \mathbf{s}_{\parallel}\ ^2 + \ \mathbf{s}_{\perp}\ ^2$ (4.44). Analogously, the
penev_B_F00002	1	■ 1.000	$\ \mathbf{s}\ ^2 = \ \mathbf{s}_{\parallel}\ ^2 + \ \mathbf{s}_{\perp}\ ^2$	$\ \mathbf{s}\ ^2 = \ \mathbf{s}_{\parallel}\ ^2 + \ \mathbf{s}_{\perp}\ ^2$	from (4.57), $\langle \mathbf{s}_{\parallel}, \mathbf{s}_{\perp} \rangle = 0$, hence $\ \mathbf{s}\ ^2 = \ \mathbf{s}_{\parallel}\ ^2 + \ \mathbf{s}_{\perp}\ ^2$ (4.44). Analogously, the
penev_B_F00003	1	■ 1.000	O_m	O_m	Notably, the change of a single O_m does not change any of the other projections
penev_B_F00004	1	■ 0.430	$\mathbf{s}^{\text{rec}}$	$\mathbf{s}^{\text{rec}}$	of $\mathbf{s}^{\text{rec}}$ to $\{\mathbf{q}_m\}_{\mathbf{x}_m \in \mathcal{M}}$ (4.56); this comes directly from (4.54).
penev_B_F00005	1	■ 0.430	$\{\mathbf{q}_m\}_{\mathbf{x}_m \in \mathcal{M}}$	$\{\mathbf{q}_m\}_{\mathbf{x}_m \in \mathcal{M}}$	of $\mathbf{s}^{\text{rec}}$ to $\{\mathbf{q}_m\}_{\mathbf{x}_m \in \mathcal{M}}$ (4.56); this comes directly from (4.54).
penev_B_F00006	1	■ 0.577	$O^{\text{rec}}(\mathbf{x})$ (4.45)	$O^{\text{rec}}(\mathbf{x})$ (4.45)	Now we are prepared to calculate $O^{\text{rec}}(\mathbf{x})$ (4.45):
penev_B_F00007	1	■ 0.606	$\{\mathbf{p}_m\}_{\mathbf{x}_m \in \mathcal{M}}$	$\{\mathbf{p}_m\}_{\mathbf{x}_m \in \mathcal{M}}$	with the action of $\mathbf{O}^*$ on the dual-basis vectors $\{\mathbf{p}_m\}_{\mathbf{x}_m \in \mathcal{M}}$, only this time, the
penev_B_F00008	1	■ 0.486	$\mathbf{Q}_{nm} = \langle \mathbf{q}_n, \mathbf{q}_m \rangle$	$\mathbf{Q}_{nm} = \langle \mathbf{q}_n, \mathbf{q}_m \rangle$	with $\mathbf{Q}_{nm} = \langle \mathbf{q}_n, \mathbf{q}_m \rangle$. Therefore, $\mathbf{B} = \mathbf{Q}^{-1}$ and
penev_B_F00009	1	■ 0.486	$\mathbf{B} = \mathbf{Q}^{-1}$	$\mathbf{B} = \mathbf{Q}^{-1}$	with $\mathbf{Q}_{nm} = \langle \mathbf{q}_n, \mathbf{q}_m \rangle$. Therefore, $\mathbf{B} = \mathbf{Q}^{-1}$ and
penev_B_F00010	1	■ 0.686	(4.62)	(4.62)	(4.62)
penev_B_F00011	1	■ 0.592	$\mathbf{Q}$	$\mathbf{Q}$	Notably, the matrix $\mathbf{Q}$ is the <i>restriction</i> of the matrix $\mathbf{P}$ on the set of reference
penev_B_F00012	1	■ 0.592	$\mathbf{P}$	$\mathbf{P}$	Notably, the matrix $\mathbf{Q}$ is the <i>restriction</i> of the matrix $\mathbf{P}$ on the set of reference
penev_B_F00013	1	■ 1.000	$\mathcal{M}$	$\mathcal{M}$	points, $\mathcal{M}$:
penev_B_F00014	1	■ 0.826	$\mathbf{x}$	$\mathbf{x}$	Analogously to (4.48–4.49), the $\mathbf{x}$ dependence of the reconstructors $a_m(\mathbf{x})$ de-
penev_B_F00015	1	■ 0.826	$a_m(\mathbf{x})$	$a_m(\mathbf{x})$	Analogously to (4.48–4.49), the $\mathbf{x}$ dependence of the reconstructors $a_m(\mathbf{x})$ de-
penev_B_F00016	1	■ 1.000	ϕ	ϕ	ends only on the locations of the reference points $\mathcal{M}$, but not on ϕ itself.
penev_B_F00017	1	■ 0.999	ϕ is ³⁵	ϕ is ³⁵	The reconstruction of the example ϕ is ³⁵
penev_B_F00018	1	■ 0.654	$K^{(-1)}_n(\mathbf{x}) = \left(\tilde{\mathbf{x}}, \mathbf{K}^{(-1)} \hat{\mathbf{x}}_n \right)$	$K^{(-1)}_n(\mathbf{x}) = \left(\tilde{\mathbf{x}}, \mathbf{K}^{(-1)} \hat{\mathbf{x}}_n \right)$	with $K^{(-1)}_n(\mathbf{x}) = \left(\tilde{\mathbf{x}}, \mathbf{K}^{(-1)} \hat{\mathbf{x}}_n \right)$.
penev_B_F00019	1	■ 0.269	$ \mathcal{M} $	$ \mathcal{M} $	this might not seem natural—when $ \mathcal{M} $ is large, $\mathbf{Q}$ is also large, and inverting
penev_B_F00020	1	■ 0.999	Q	Q	it is difficult. Nevertheless, $\mathbf{Q}$ is a special matrix—most of its values are van-
penev_B_F00021	1	■ 0.623	$\{P_m(\mathbf{x})\}_{\mathbf{x}_m \in \mathcal{M}}$	$\{P_m(\mathbf{x})\}_{\mathbf{x}_m \in \mathcal{M}}$	ishingly small; because the supports of $\{P_m(\mathbf{x})\}_{\mathbf{x}_m \in \mathcal{M}}$ are local, $\mathbf{Q}$ is <i>sparse</i> .
penev_B_F00022	1	■ 0.990	$\mathbf{x}_m$	$\mathbf{x}_m$	For each point $\mathbf{x}_m$, the m -th column of $\mathbf{Q}$ has substantially nonvanishing val-
penev_B_F00023	1	■ 0.990	m	m	For each point $\mathbf{x}_m$, the m -th column of $\mathbf{Q}$ has substantially nonvanishing val-
penev_B_F00024	1	■ 0.446	$P_m(\mathbf{x}_m)$	$P_m(\mathbf{x}_m)$	$P_m(\mathbf{x}_m)$. Moreover, this is true even when the volume of the receptor grid V
penev_B_F00025	1	■ 0.446	V	V	$P_m(\mathbf{x}_m)$. Moreover, this is true even when the volume of the receptor grid V
penev_B_F00026	1	■ 0.961	$\mathbf{x}_m$	$\mathbf{x}_m$	increases—adding more and more receptors distant to $\mathbf{x}_m$ does not add non-
penev_B_F00027	1	■ 0.905	N	N	small, regardless of N and V . ³⁶
penev_B_F00028	1	■ 0.905	V	V	small, regardless of N and V . ³⁶
penev_B_F00029	1	■ 0.709	$\phi(\mathbf{x})$	$\phi(\mathbf{x})$	In this Section, we have shown how to represent a sensory stimulus $\phi(\mathbf{x})$
penev_B_F00030	1	■ 1.000	$\mathcal{P}[\phi(\mathbf{x})]$	$\mathcal{P}[\phi(\mathbf{x})]$	and its probability $\mathcal{P}[\phi(\mathbf{x})]$, in the context of an ensemble with low dimension-
penev_B_F00031	1	■ 0.968	$O^{\text{rec}}(\mathbf{x})$	$O^{\text{rec}}(\mathbf{x})$	will result in a large error in $O^{\text{rec}}(\mathbf{x})$.
penev_B_F00032	1	■ 0.743	$\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$	$\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$	So far, we have addressed the question, given a sparsification $\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$,
penev_B_F00033	1	■ 1.000	$O(\mathbf{x})$	$O(\mathbf{x})$	how to reconstruct $O(\mathbf{x})$ and, from there, the example ϕ and its probabil-
penev_B_F00034	1	■ 1.000	$\mathcal{P}[\phi]$	$\mathcal{P}[\phi]$	ity $\mathcal{P}[\phi]$.
penev_B_F00035	1	■ 0.999	$\mathcal{M}^{(0)} = \emptyset$	$\mathcal{M}^{(0)} = \emptyset$	In the sparsification stage, we start with the empty set $\mathcal{M}^{(0)} = \emptyset$ and
penev_B_F00036	1	■ 0.866	n	n	perform a series of iterative steps; at the n -th step, we add to $\mathcal{M}^{(n)}$ one reference
penev_B_F00037	1	■ 0.866	$\mathcal{M}^{(n)}$	$\mathcal{M}^{(n)}$	perform a series of iterative steps; at the n -th step, we add to $\mathcal{M}^{(n)}$ one reference
penev_B_F00038	1	■ 0.806	$\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$	$\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$	The resulting set $\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$ is an LFA representation of ϕ .

Identifier	Page	Conf.	LaTeX source	Rendered	Image
penev_B_F00039	1	■ 1.000	$\backslash\mathrm{mathcal}\{M\}=\backslash\mathrm{mathcal}\{M\}^{\wedge\{n\}}$	$\mathcal{M} = \mathcal{M}^{(n)}$	First, given the current set $\mathcal{M} = \mathcal{M}^{(n)}$, we calculate the current maximum-
penev_B_F00040	1	■ 0.479	$0^{\wedge\{r\}}\mathrm{e}\mathrm{c}\}\backslash\mathrm{mathrm}\{x\}$	$O^{rec}(\mathbf{x})$	likelihood estimate $O^{rec}(\mathbf{x})$ by (4.67–4.68) and calculate the current error
penev_B_F00041	1	■ 0.718	$\backslash\mathrm{mathrm}\{x\}_{m+1}$	$\mathbf{x}_{m+1}$	Second, we look for the grid point $\mathbf{x}_{m+1}$, such that the value of $O^{err}(\mathbf{x}_{m+1})$
penev_B_F00042	1	■ 0.718	$0^{\wedge\{\text{\texttt{err}}\}}\backslash\mathrm{left}\backslash\mathrm{mathrm}\{x\}_{m+1}\backslash\mathrm{right}$	$O^{err}(\mathbf{x}_{m+1})$	Second, we look for the grid point $\mathbf{x}_{m+1}$, such that the value of $O^{err}(\mathbf{x}_{m+1})$
penev_B_F00043	1	■ 0.980	$0^{\wedge\{\text{\texttt{err}}\}}\backslash\mathrm{left}\backslash\mathrm{mathbf}\{x\}_{m+1}\backslash\mathrm{right}\wedge\{2\}$	$O^{err}(\mathbf{x}_{m+1})^2$	is maximum. If its square, $O^{err}(\mathbf{x}_{m+1})^2$, is below a predetermined threshold,
penev_B_F00044	1	■ 1.000	$-\backslash\log\backslash\mathrm{mathcal}\{P\}[\phi]$	$-\log\mathcal{P}[\phi]$	terion. We will remember that the information content of the example, $-\log\mathcal{P}[\phi]$, is de-
penev_B_F00045	1	■ 0.810	$-\backslash\log\backslash\mathrm{mathcal}\{P\}[\phi]-\backslash\mathrm{left}(-\backslash\log\backslash\mathrm{mathcal}\{P\}\backslash\mathrm{left}[\phi^{\text{\texttt{rec}}}\backslash\mathrm{right}]\backslash\mathrm{right})$	$-\log\mathcal{P}[\phi] - (-\log\mathcal{P}[\phi^{rec}])$	$-\log\mathcal{P}[\phi] - (-\log\mathcal{P}[\phi^{rec}])$, is equal to $\ O^{err}\ _F^2$. Therefore, by bounding of the maximum
penev_B_F00046	1	■ 0.810	$\backslash\mathrm{left}\backslash 0^{\wedge\{\text{\texttt{err}}\}}\backslash\mathrm{right}\backslash_U\wedge\{2\}$	$\ O^{err}\ _U^2$	$-\log\mathcal{P}[\phi] - (-\log\mathcal{P}[\phi^{rec}])$, is equal to $\ O^{err}\ _F^2$. Therefore, by bounding of the maximum
penev_B_F00047	1	■ 0.771	$\backslash\mathrm{left}\backslash 0^{\wedge\{\text{\texttt{err}}\}}\backslash\mathrm{right}\backslash_E\wedge\{2\}=\backslash\mathrm{frac}\{1\}\{V\}\backslash\mathrm{sum}_x0^{\wedge\{\text{\texttt{err}}\}}\backslash\mathrm{mathbf}\{x\}\wedge\{2\}(2.8)$	$\ O^{err}\ _E^2 = \frac{1}{V} \sum_x O^{err}(\mathbf{x})^2 (2.8)$	term in the sum $\ O^{err}\ _E^2 = \frac{1}{V} \sum_x O^{err}(\mathbf{x})^2$ (2.8), we bound the whole sum and, therefore,
penev_B_F00048	1	■ 0.779	$\backslash\mathrm{mathbf}\{x\}_{m+1}$	$\mathbf{x}_{m+1}$	Third, if the algorithm has not terminated yet, we add $\mathbf{x}_{m+1}$ to $\mathcal{M}^{(n)}$, which
penev_B_F00049	1	■ 0.779	$\backslash\mathrm{mathcal}\{M\}^{\wedge\{(\backslash\mathrm{mathbf}\{n\})\}}$	$\mathcal{M}^{(\mathbf{n})}$	Third, if the algorithm has not terminated yet, we add $\mathbf{x}_{m+1}$ to $\mathcal{M}^{(n)}$, which
penev_B_F00050	1	■ 0.974	$\backslash\mathrm{mathcal}\{M\}^{\wedge\{(n+1)\}}$	$\mathcal{M}^{(n+1)}$	results in $\mathcal{M}^{(n+1)}$, and go to step $n + 1$. ³⁹
penev_B_F00051	1	■ 0.974	$n+1\text{\texttt{.}}^{\wedge\{39\}}$	$n + 1$. ³⁹	results in $\mathcal{M}^{(n+1)}$, and go to step $n + 1$. ³⁹
penev_B_F00052	1	■ 0.978	$ \backslash\mathrm{mathcal}\{M\} =N$	$ \mathcal{M} = N$	dicted well, therefore they are linearly independent, and by the time $ \mathcal{M} = N$,
penev_B_F00053	1	■ 0.972	$0^{\wedge\{e\}}\mathrm{r}\mathrm{r}\}\{x\}=0$	$O^{err}(x) = 0$	$O^{err}(\mathbf{x}) = 0$ (4.36).
penev_B_F00054	1	■ 1.000	$\backslash\mathrm{mathcal}\{M\}^{\wedge\{(25)\}}$	$\mathcal{M}^{(25)}$	The location and the order of first 25 points, those in $\mathcal{M}^{(25)}$ (Fig. 4.4a),
penev_B_F00055	1	■ 0.990	$\backslash\mathrm{mathcal}\{M\}^{\wedge\{(64)\}}$	$\mathcal{M}^{(64)}$	which is a desirable property. The points in $\mathcal{M}^{(64)}$ enough for acceptable
penev_B_F00056	1	■ 0.822	$a_{\{r\}}$	a_r	number of values 64. We calculated the PCA coefficients a_r (2.47–2.52) and
penev_B_F00057	1	■ 0.996	$x_{\{m+1\}}$	x_{m+1}	footnote 39 we argue that this is a costly calculation. Moreover, if we add the point $\mathbf{x}_{m+1}$,
penev_B_F00058	1	■ 0.958	$0^{\wedge\{\text{\texttt{err}}\}}\backslash\mathrm{left}\backslash\mathrm{mathbf}\{x\}_{m+1}\backslash\mathrm{right}$	$O^{err}(\mathbf{x}_{m+1})$	we are making a correction, $O^{err}(\mathbf{x}_{m+1})$, to the estimated value $O^{rec}(\mathbf{x}_{m+1})$; this has to be
penev_B_F00059	1	■ 0.958	$0^{\wedge\{\text{\texttt{rec}}\}}\backslash\mathrm{left}\backslash\mathrm{mathbf}\{x\}_{m+1}\backslash\mathrm{right}$	$O^{rec}(\mathbf{x}_{m+1})$	we are making a correction, $O^{err}(\mathbf{x}_{m+1})$, to the estimated value $O^{rec}(\mathbf{x}_{m+1})$; this has to be
penev_B_F00060	1	■ 0.637	$0\backslash\mathrm{left}\backslash\mathrm{mathbf}\{x\}_{m+1}\backslash\mathrm{right}$	$O(\mathbf{x}_{m+1})$	compared to the error in the measurement of $O(\mathbf{x}_{m+1})$ itself, so a criterion defined at the
penev_B_F00061	1	■ 0.938	$0^{\wedge\{e\}}\mathrm{r}\mathrm{r}\}\backslash\mathrm{left}\backslash x_{m+1}\backslash\mathrm{right}$	$O^{err}(x_{m+1})$	level of the individual error, $O^{err}(\mathbf{x}_{m+1})$, might not be such a bad idea after all. Also, we
penev_B_F00062	1	■ 0.515	$\backslash\mathrm{left}\backslash 0^{\wedge\{e\}}\mathrm{r}\mathrm{r}\}\backslash\mathrm{right}\backslash_U\wedge\{2\}$	$\ O^{err}\ _U^2$	provisionally taken; $\mathbf{Q}$, inverted; the resulting $\ O^{err}\ _U^2$, stored; and, after all $\sim V$ provisional
penev_B_F00063	1	■ 0.515	$\backslash\mathrm{sim}\ V$	$\sim V$	provisionally taken; $\mathbf{Q}$, inverted; the resulting $\ O^{err}\ _U^2$, stored; and, after all $\sim V$ provisional
penev_B_F00064	2	■ 0.869	$\backslash\mathrm{mathbb}\{I\}$	$\mathbb{I}$	The building of the mask $\mathcal{M}$ for Example 1, reconstructed in the context of
penev_B_F00065	2	■ 1.000	$N=220, n=\backslash\mathrm{sigma}_{400}$	$N = 220, n = \sigma_{400}$	Ensemble 1. $N = 220, n = \sigma_{400}$ (see Fig. 4.1). a : the first 25 points— $\mathcal{M}^{(25)}$,
penev_B_F00066	2	■ 1.000	a	a	Ensemble 1. $N = 220, n = \sigma_{400}$ (see Fig. 4.1). a : the first 25 points— $\mathcal{M}^{(25)}$,
penev_B_F00067	2	■ 0.999	$\phi(\backslash\mathrm{mathbf}\{x\})$	$\phi(\mathbf{x})$	overlayed on $\phi(\mathbf{x})$ and numbered sequentially. (The label for point 11, in the
penev_B_F00068	2	■ 0.999	b	b	is the bottommost point on the right edge, shown on b .) b : the points in $\mathcal{M}^{(64)}$.
penev_B_F00069	2	■ 0.995	$N=220$	$N = 220$	the LFA ones $O(\mathbf{x})$ (4.31) for $N = 220$, then sparsified to get $\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$.
penev_B_F00070	2	■ 0.995	$\phi^{\text{\texttt{rec}}}\backslash\mathrm{r}\mathrm{e}\mathrm{c}\}\backslash\mathrm{mathbf}\{x\}$	$\phi^{rec}(\mathbf{x})$	We calculated the reconstruction $\phi^{rec}(\mathbf{x})$ either (a), from (2.50) with $N = 64$, or
penev_B_F00071	2	■ 0.995	$N=64$	$N = 64$	We calculated the reconstruction $\phi^{rec}(\mathbf{x})$ either (a), from (2.50) with $N = 64$, or
penev_B_F00072	2	■ 0.965	$8\ \backslash\mathrm{times}\ 8$	8×8	For comparison, we subsampled $\phi(\mathbf{x})$ on a uniform 8×8 grid and reconstructed
penev_B_F00073	2	■ 1.000	$\backslash\mathrm{Psi}_{\{1\}}$	Ψ_1	it with those 64 points (c). In all cases we reconstructed about Ψ_1 the average
penev_B_F00074	2	■ 1.000	$a_{\{1\}}$	a_1	of the ensemble; we kept a_1 fixed. Evidentially, the perceptual quality of the
penev_B_F00075	2	■ 0.999	$\phi^{\text{\texttt{rec}}}\backslash\mathrm{phi}^{\wedge\{\text{\texttt{rec}}\}}\backslash\mathrm{mathbf}\{x\}$	$\phi^{rec}(\mathbf{x})$	reconstruction $\phi^{rec}(\mathbf{x})$ (top row) is the best, and the error $\phi(\mathbf{x}) - \phi^{rec}(\mathbf{x})$ (bot-
penev_B_F00076	2	■ 0.999	$\phi(\backslash\mathrm{mathbf}\{x\})-\phi^{\text{\texttt{rec}}}\backslash\mathrm{phi}^{\wedge\{\text{\texttt{rec}}\}}\backslash\mathrm{mathbf}\{x\}$	$\phi(\mathbf{x}) - \phi^{rec}(\mathbf{x})$	reconstruction $\phi^{rec}(\mathbf{x})$ (top row) is the best, and the error $\phi(\mathbf{x}) - \phi^{rec}(\mathbf{x})$ (bot-
penev_B_F00077	2	■ 0.499	c	c	and 508 for (a), (b), and (c), respectively, out of 560 total power in $(\phi(\mathbf{x}) - a_1\Psi_1)$. Notably,
penev_B_F00078	2	■ 0.499	$\backslash\mathrm{left}\backslash\phi(\backslash\mathrm{mathrm}\{x\})-a_{\{1\}}\backslash\mathrm{Psi}_{\{1\}}\backslash\mathrm{right}.$	$(\phi(\mathbf{x}) - a_1\Psi_1)$	and 508 for (a), (b), and (c), respectively, out of 560 total power in $(\phi(\mathbf{x}) - a_1\Psi_1)$. Notably,
penev_B_F00079	2	■ 0.990	$\backslash\mathrm{Psi}_{\{2\}}$	Ψ_2	as Ψ_2 on Fig. 3.9. Indeed, the m.s.e. due to the error in a_2 in (b) is 75. If one fixes not
penev_B_F00080	2	■ 0.990	$a_{\{2\}}$	a_2	as Ψ_2 on Fig. 3.9. Indeed, the m.s.e. due to the error in a_2 in (b) is 75. If one fixes not
penev_B_F00081	2	■ 0.474	$\backslash\mathrm{left}\backslash\{0\backslash\mathrm{left}\backslash\mathrm{mathbf}\{x\}_m\backslash\mathrm{right}\}\backslash\mathrm{right}\backslash_m\backslash\mathrm{mathbf}\{x\}_m\in\backslash\mathrm{mathcal}\{M\}\}(4.674\text{--}4.68)$	$\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}} (4.674.68)$	(2.48). b : reconstruction from the sparsification $\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$ (4.67–4.68) with
penev_B_F00082	2	■ 1.000	$\backslash\mathrm{mathcal}\{M\}=\backslash\mathrm{mathcal}\{M\}^{\wedge\{(64)\}}$	$\mathcal{M} = \mathcal{M}^{(64)}$	$\mathcal{M} = \mathcal{M}^{(64)}$ shown on Fig. 4.4b, which was produced by the serial algorithm
penev_B_F00083	2	■ 0.788	$\phi_{\{64\}}^{\text{\texttt{err}}}\backslash\mathrm{phi}_{\{64\}}^{\text{\texttt{err}}}\backslash\mathrm{phi}_{\{64\}}^{\text{\texttt{err}}}$	ϕ_{64}^{err}	The error on (a) is exactly equal to ϕ_{64}^{err} , and the error on (b) is perceptually
penev_B_F00084	2	■ 0.830	$\phi_{\{100\}}^{\text{\texttt{err}}}\backslash\mathrm{phi}_{\{100\}}^{\text{\texttt{err}}}\backslash\mathrm{phi}_{\{100\}}^{\text{\texttt{err}}}$	ϕ_{100}^{err}	somewhere between ϕ_{100}^{err} and ϕ_{200}^{err} (cf. also the reconstruction with ϕ_{200}^{err}), both
penev_B_F00085	2	■ 0.830	$\phi_{\{200\}}^{\text{\texttt{err}}}\backslash\mathrm{phi}_{\{200\}}^{\text{\texttt{err}}}\backslash\mathrm{phi}_{\{200\}}^{\text{\texttt{err}}}$	ϕ_{200}^{err}	somewhere between ϕ_{100}^{err} and ϕ_{200}^{err} (cf. also the reconstruction with ϕ_{200}^{err}), both
penev_B_F00086	2	■ 0.830	$\phi_{\{200\}}^{\text{\texttt{rec}}}\backslash\mathrm{phi}_{\{200\}}^{\text{\texttt{rec}}}\backslash\mathrm{phi}_{\{200\}}^{\text{\texttt{rec}}}$	ϕ_{200}^{rec}	somewhere between ϕ_{100}^{err} and ϕ_{200}^{err} (cf. also the reconstruction with ϕ_{200}^{err}), both
penev_B_F00087	2	■ 0.992	$5\ \backslash\mathrm{times}$	$5 \times$	are magnified $5 \times$ (cf. Fig. 3.12, Fig. 4.3). The error on (c) is not magnified.
penev_B_F00088	2	■ 1.000	$(m-1)$	$(m - 1)$	the $(m - 1)$ -st step through a simple algebraic formula. The algorithm for inversion through
penev_B_F00089	2	■ 0.585	$ \backslash\mathrm{mathcal}\{M\} 20\text{--}25\ \backslash\%$	$ \mathcal{M} 20\% - 25\%$	ence points $ \mathcal{M} $ 20%–25% of their respective PCA dimensionalities N ; 99%,
penev_B_F00090	2	■ 0.585	$N;99\ \backslash\%$	$N;99\%$	ence points $ \mathcal{M} $ 20%–25% of their respective PCA dimensionalities N ; 99%,
penev_B_F00091	2	■ 0.970	$45\text{--}50\ \backslash\%$	$45\% - 50\%$	with 45%–50%. Therefore, the necessary LFA dimensionality for a given “in-
penev_B_F00092	2	■ 0.560	$99\ \backslash\%$	99%	to get 99% of the fidelity?—the incremental benefit of adding a reference point in the 50%

Identifier	Page	Conf.	LaTeX source	Rendered	Image
penev_B_F00093	2	0.560	50 \%	50%	to get 99% of the fidelity?—the incremental benefit of adding a reference point in the 50%
penev_B_F00094	2	1.000	r	r	LFA representations by taking the sums in (4.29) in several separate bands for r , instead of
penev_B_F00095	2	1.000	\approx 1	≈ 1	contributes on the average ≈ 1 .
penev_B_F00096	2	0.999	N \in	$N \in$	the number of referce points $ \mathcal{M} $ is shown with dashed lines for $N \in$
penev_B_F00097	3	0.977	S-N	$S - N$	Figure 4.7: Scaling of LFA S - N Diagrams
penev_B_F00098	3	0.682	\ O\ _{\tilde{V}}^2=\left\ \Phi^{*+}\phi\right\ _S^2	$\ O\ _{\tilde{V}}^2 = \ \Phi^{*+}\phi\ _S^2$	$\ O\ _{\tilde{V}}^2 = \ \Phi^{*+}\phi\ _S^2$ (4.22), as a function of $ \mathcal{M} /N$, the relative dimensionalities
penev_B_F00099	3	0.682	\mathcal{M} / N	$ \mathcal{M} /N$	$\ O\ _{\tilde{V}}^2 = \ \Phi^{*+}\phi\ _S^2$ (4.22), as a function of $ \mathcal{M} /N$, the relative dimensionalities
penev_B_F00100	3	0.977	N \in\{600,400,300,220,110,64,32\}	$N \in \{600, 400, 300, 220, 110, 64, 32\}$	of the LFA and PCA representations for $N \in \{600, 400, 300, 220, 110, 64, 32\}$
penev_B_F00101	3	1.000	\mathcal{M}^{\{(50)\}}	$\mathcal{M}^{(50)}$	on Fig. 4.8. Almost all of the points in the respective sets $\mathcal{M}^{(50)}$ are on the
penev_B_F00102	3	0.496	\left\{O\left(x_m\right)\right\}_{x_m \in \mathcal{M}}	$\{O(x_m)\}_{x_m \in \mathcal{M}}$	given a sparsification $\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$ of the dense LFA outputs $O(\mathbf{x})$ for an
penev_B_F00103	3	0.496	O(x)	$O(x)$	given a sparsification $\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$ of the dense LFA outputs $O(\mathbf{x})$ for an
penev_B_F00104	3	0.951	\left\{a_m(\mathbf{x})\right\}_{x_m \in \mathcal{M}}	$\{a_m(\mathbf{x})\}_{x_m \in \mathcal{M}}$	given by the set of interpolating reconstructors $\{a_m(\mathbf{x})\}_{\mathbf{x}_m \in \mathcal{M}}$ (4.68).
penev_B_F00105	3	0.342	\sum_{n=1}^{\mathcal{M}} Q_{n n n}^{-1} K^{(-1)}_n(\mathbf{x})	$\sum_{n=1}^{\mathcal{M}} Q_{n n n}^{-1} K^{(-1)}_n(\mathbf{x})$	combination, only with a different set of reconstructors, $\sum_{n=1}^{ \mathcal{M} } \mathbf{Q}_{n n n}^{-1} K^{(-1)}_n(\mathbf{x})$.
penev_B_F00106	3	1.000	V=8640	$V = 8640$	in the context of Ensemble 3 (see Appendix A.1 for definition) with $V = 8640$,
penev_B_F00107	3	1.000	T=1038, N=400, n=\sigma_{400}	$T = 1038, N = 400, n = \sigma_{400}$	$T = 1038, N = 400, n = \sigma_{400}$ (see Fig. 4.1).
penev_B_F00108	3	0.938	\mathbf{Q}^{-1}	$\mathbf{Q}^{-1}$	operation. By the same argument, $\mathbf{Q}^{-1}$ will be sparse, and acting with it on O_m
penev_B_F00109	3	0.907	\mathcal{O}_I	$\mathcal{O}_I$	will be a local operation too. Moreover, one may try to solve for $\mathcal{O}_I$ directly
penev_B_F00110	3	1.000	\mathbf{Q}^{-1}	$\mathbf{Q}^{-1}$	and never explicitly store $\mathbf{Q}^{-1}$, which will eliminate one computational step
penev_B_F00111	3	0.644	\mathcal{O}_1	$\mathcal{O}_1$	Some intuition into what is involved in the calculation of $\mathcal{O}_1$ may be de-
penev_B_F00112	3	1.000	\mathcal{O}_l	$\mathcal{O}_l$	may be solved for $\mathcal{O}_l$ (cf. 4.72) by a rapid iteration.
penev_B_F00113	4	0.601	G(\mathcal{O})(\mathbf{x})=\mathbf{g}(\mathcal{O}(\mathbf{x}))	$G(\mathcal{O})(\mathbf{x}) = \mathbf{g}(\mathcal{O}(\mathbf{x}))$	where $G(\mathcal{O})(\mathbf{x}) = \mathbf{g}(\mathcal{O}(\mathbf{x}))$ with $\mathbf{g}$, a suitably chosen <i>sigmoid</i> function, ⁴⁶ and
penev_B_F00114	4	1.000	\tau	τ	τ is the characteristic time scale.
penev_B_F00115	4	1.000	\mathcal{O}(\mathbf{x})	$\mathcal{O}(\mathbf{x})$	to pass the large ones intact. Then, if all $\mathcal{O}(\mathbf{x})$ either vanish, or are sufficiently
penev_B_F00116	4	0.555	\frac{d}{dt} \mathcal{O}(\mathbf{x})=0	$\frac{d}{dt} \mathcal{O}(\mathbf{x}) = 0$	large, $\mathbf{g}$ will be the identity. In this regime, for the steady states, $\frac{d}{dt} \mathcal{O}(\mathbf{x}) = 0$,
penev_B_F00117	4	0.465	O^{\mathrm{rec}}=\mathbf{P}\mathcal{O}=\mathbf{K}\phi=O	$O^{\mathrm{rec}} = \mathbf{P}\mathcal{O} = \mathbf{K}\phi = O$	The steady-state equation (4.76) is equivalent to $O^{\mathrm{rec}} = \mathbf{P}\mathcal{O} = \mathbf{K}\phi = O$
penev_B_F00118	4	0.997	\left\{\mathcal{O}_m\right\}_{x_m \in \mathcal{M}}	$\{\mathcal{O}_m\}_{x_m \in \mathcal{M}}$	hence, the sparsified outputs $\{\mathcal{O}_m\}_{\mathbf{x}_m \in \mathcal{M}}$ are solutions of (4.76) and, therefore,
penev_B_F00119	4	0.840	\mathbf{g}: \mathcal{R} \rightarrow[a, b]	$\mathbf{g}: \mathcal{R} \rightarrow[a, b]$	⁴⁶ A sigmoid function is a monotonic nondecreasing function $\mathbf{g}: \mathcal{R} \rightarrow[a, b]$, often $a=-b$,
penev_B_F00120	4	0.840	a=-b	$a=-b$	⁴⁶ A sigmoid function is a monotonic nondecreasing function $\mathbf{g}: \mathcal{R} \rightarrow[a, b]$, often $a=-b$,
penev_B_F00121	4	0.809	\mathbf{g}'	$\mathbf{g}'$	and usually $\mathbf{g}'$ is non-zero only near the origin, or a suitably chosen <i>threshold</i> θ .
penev_B_F00122	4	0.809	\theta	θ	and usually $\mathbf{g}'$ is non-zero only near the origin, or a suitably chosen <i>threshold</i> θ .
penev_B_F00123	4	1.000	\left\{\mathcal{O}(\mathbf{u})\right\}_{\mathbf{u} \in V}	$\{\mathcal{O}(\mathbf{u})\}_{\mathbf{u} \in V}$	The input layer stores the example $\phi(\mathbf{x})$; the output layer, $\{\mathcal{O}(\mathbf{u})\}_{\mathbf{u} \in V}$.
penev_B_F00124	4	0.803	\mathbf{u}	$\mathbf{u}$	The output layer unit centered at $\mathbf{u}$ has a feed-forward receptive field $\mathbf{K}_{\mathbf{u}}(\mathbf{x})$
penev_B_F00125	4	0.803	\mathbf{K}_{\mathbf{u}}(\mathbf{x})	$\mathbf{K}_{\mathbf{u}}(\mathbf{x})$	The output layer unit centered at $\mathbf{u}$ has a feed-forward receptive field $\mathbf{K}_{\mathbf{u}}(\mathbf{x})$
penev_B_F00126	4	0.998	O^{\mathrm{err}}(\mathbf{u})	$O^{\mathrm{err}}(\mathbf{u})$	The current error signal $O^{\mathrm{err}}(\mathbf{u})$ (4.71) is calculated from the current output
penev_B_F00127	4	0.962	\mathcal{O}(\mathbf{u})	$\mathcal{O}(\mathbf{u})$	variables $\mathcal{O}(\mathbf{u})$ by (4.72); the residual correlations $\mathbf{P}_{\mathbf{u}}(\mathbf{v})$ (cf. Fig. 4.1–4.2) serve
penev_B_F00128	4	0.962	\mathbf{P}_{\mathbf{u}}(\mathbf{v})	$\mathbf{P}_{\mathbf{u}}(\mathbf{v})$	variables $\mathcal{O}(\mathbf{u})$ by (4.72); the residual correlations $\mathbf{P}_{\mathbf{u}}(\mathbf{v})$ (cf. Fig. 4.1–4.2) serve
penev_B_F00129	4	0.772	\mathcal{O}(\mathbf{u})=0	$\mathcal{O}(\mathbf{u}) = 0$	to $\mathcal{O}(\mathbf{u}) = 0$ everywhere on the grid, with the exception of $\{\mathcal{O}_{\mathbf{m}}\}_{\mathbf{m}=1}^{ \mathcal{M} }$ (4.74).
penev_B_F00130	4	0.772	\left\{\mathcal{O}_{\mathbf{m}}\right\}_{\mathbf{m}=\mathbf{I}}^{ \mathcal{M} }	$\{\mathcal{O}_{\mathbf{m}}\}_{\mathbf{m}=\mathbf{I}}^{ \mathcal{M} }$	to $\mathcal{O}(\mathbf{u}) = 0$ everywhere on the grid, with the exception of $\{\mathcal{O}_{\mathbf{m}}\}_{\mathbf{m}=1}^{ \mathcal{M} }$ (4.74).
penev_B_F00131	4	0.800	g	g	If output dynamic range equalization is required, $\mathbf{g}$ may be chosen to preclude
penev_B_F00132	4	0.607	\mathcal{O}_i	$\mathcal{O}_i$	large $\mathcal{O}_i$.
penev_B_F00133	4	0.863	\mathcal{P}_{\mathrm{objects}}	$\mathcal{P}_{\mathrm{objects}}$	ability distribution, $\mathcal{P}_{\mathrm{objects}}$, is entirely contained in the probability distribution
penev_B_F00134	4	0.992	\mathcal{P}_{\mathrm{images}}	$\mathcal{P}_{\mathrm{images}}$	of natural images, $\mathcal{P}_{\mathrm{images}}$.
penev_B_F00135	4	0.445	\mathcal{P}_{\mathrm{object}}^{\mathrm{can}}	$\mathcal{P}_{\mathrm{object}}^{\mathrm{can}}$	approximation and parameterization. In that sense, $\mathcal{P}_{\mathrm{objects}}$ can be considered
penev_B_F00136	5	0.945	\left\{I^t\right\}_{t \in T}	$\{I^t\}_{t \in T}$	of images $\{I^t\}_{t \in T}$ was prepared, wherein each image contained a human face
penev_B_F00137	5	1.000	I^t	I^t	in a relatively frontal pose. For each image in the ensemble, I^t , the locations
penev_B_F00138	5	1.000	\left(X_l^t, Y_l^t\right)	(X_l^t, Y_l^t)	(X_l^t, Y_l^t) and (X_r^t, Y_r^t) of the left and right eye, respectively (see discussion in
penev_B_F00139	5	1.000	\left(X_r^t, Y_r^t\right)	(X_r^t, Y_r^t)	(X_l^t, Y_l^t) and (X_r^t, Y_r^t) of the left and right eye, respectively (see discussion in
penev_B_F00140	5	0.698	\theta^i	θ^i	Subsequently, the image I^t was <i>rotated</i> , by the angle θ^t , to line the eyes
penev_B_F00141	5	1.000	S^i	S^i	horizontally; <i>scaled</i> , by the scale factor S^t , to bring the inter-eye distance to a
penev_B_F00142	5	0.998	X^t	X^t	pre-defined length; and <i>translated</i> , horizontally by X^t and vertically by Y^t , to
penev_B_F00143	5	0.998	Y^t	Y^t	pre-defined length; and <i>translated</i> , horizontally by X^t and vertically by Y^t , to
penev_B_F00144	5	0.998	\phi^t	ϕ^t	for a given cropping, the example ϕ^t depends on the image I^t , as well as on the

Identifier	Page	Conf.	LaTeX source	Rendered	Image
penev_B_F00200	6	■ 0.994	S	S	determined in the context of PCA by the projector on the PCA S -subspace $\mathbf{P}'$
penev_B_F00201	6	■ 0.994	$\mathbf{P}^{\prime}$	$\mathbf{P}'$	determined in the context of PCA by the projector on the PCA S -subspace $\mathbf{P}'$
penev_B_F00202	6	■ 0.966	I	I	given image I and an initial guess α_0 , calculate the entropic costs (2.49)
penev_B_F00203	6	■ 0.966	α_0	α_0	given image I and an initial guess α_0 , calculate the entropic costs (2.49)
penev_B_F00204	6	■ 0.860	$\{\phi(\alpha)\}_{\alpha \in \Delta_{\alpha_0}}$	$\{\phi(\alpha)\}_{\alpha \in \Delta_{\alpha_0}}$	for all images $\{\phi(\alpha)\}_{\alpha \in \Delta_{\alpha_0}}$. The maximum likelihood estimate is α that mini-
penev_B_F00205	6	■ 0.769	$\langle S \rangle(5.4)$	$\langle S \rangle(5.4)$	mizes $\langle S \rangle$ (5.4) and, therefore, maximizes the probability $\mathcal{P}[\phi(\alpha)]$ (2.20).
penev_B_F00206	6	■ 0.769	$\mathcal{P}[\phi(\alpha)]$	$\mathcal{P}[\phi(\alpha)]$	mizes $\langle S \rangle$ (5.4) and, therefore, maximizes the probability $\mathcal{P}[\phi(\alpha)]$ (2.20).
penev_B_F00207	6	■ 0.998	α^2	α^2	the parameters of the symmetry breaking α^t . For the ensembles used in Chapter 3, the
penev_B_F00208	6	■ 0.723	$\left\{ \left(X_l^t, Y_l^t; X_r^t, Y_r^t \right) \right\}_{t \in T}$	$\left\{ \left(X_l^t, Y_l^t; X_r^t, Y_r^t \right) \right\}_{t \in T}$	$\{(X_l^t, Y_l^t; X_r^t, Y_r^t)\}_{t \in T}$. From them, the wonderful set of high-quality coefficients $\{\alpha_{G,M}^t\}_{t \in T}$
penev_B_F00209	6	■ 0.723	$\left\{ \alpha_{G,M}^t \right\}_{t \in T}$	$\left\{ \alpha_{G,M}^t \right\}_{t \in T}$	$\{(X_l^t, Y_l^t; X_r^t, Y_r^t)\}_{t \in T}$. From them, the wonderful set of high-quality coefficients $\{\alpha_{G,M}^t\}_{t \in T}$
penev_B_F00210	7	■ 0.861	$\langle S \rangle(N, \alpha_h^t)$	$\langle S \rangle(N, \alpha_h^t)$	The average entropic cost $\langle S \rangle(N, \alpha_h^t)$ (5.4) is plotted as a function of N for 7
penev_B_F00211	7	■ 0.998	α_h^t	α_h^t	different choices of α_h^t (5.5) for one of the images I^t in Ensemble 6 (see Ap-
penev_B_F00212	7	■ 0.999	$\phi^t(0)$	$\phi^t(0)$	The vertical axis is additionally scaled so that the in-sample example $\phi^t(0)$ (=)
penev_B_F00213	7	■ 0.985	$\langle S \rangle(N) \rightarrow \frac{\sigma_0^2}{\sum_{r=1}^N \sigma_r^2}$	$\langle S \rangle(N) \rightarrow \frac{\sigma_0^2}{\sum_{r=1}^N \sigma_r^2}$	has $\langle S \rangle(N) \rightarrow \frac{\sigma_0^2}{\sum_{r=1}^N \sigma_r^2}$ when $N \rightarrow T$, instead of the more usual, 1.
penev_B_F00214	7	■ 0.985	$N \rightarrow T$	$N \rightarrow T$	has $\langle S \rangle(N) \rightarrow \frac{\sigma_0^2}{\sum_{r=1}^N \sigma_r^2}$ when $N \rightarrow T$, instead of the more usual, 1.
penev_B_F00215	7	■ 0.999	$\phi^t(h_n)$	$\phi^t(h_n)$	The six out-of-sample examples $\phi^t(h_n)$ were produced with $h_n = 1 - s_n$, $s_n =$
penev_B_F00216	7	■ 0.999	$h_n = 1 - s_n$	$h_n = 1 - s_n$	The six out-of-sample examples $\phi^t(h_n)$ were produced with $h_n = 1 - s_n$, $s_n =$
penev_B_F00217	7	■ 1.000	$1.05^{n/2}$	$1.05^{n/2}$	$1.05^{n/2}$, for $n \in \{-3, -2, -1, 1, 2, 3\}$, as outlined in Appendix A.1.
penev_B_F00218	7	■ 1.000	$n \in \{-3, -2, -1, 1, 2, 3\}$	$n \in \{-3, -2, -1, 1, 2, 3\}$	$1.05^{n/2}$, for $n \in \{-3, -2, -1, 1, 2, 3\}$, as outlined in Appendix A.1.
penev_B_F00219	7	■ 0.998	I^t, α^t	I^t, α^t	for definition) are shown on Fig. 5.5. For all 1216 images I^t , α^t was determined.
penev_B_F00220	7	■ 1.000	$\phi^t(h) = H(I^t; \alpha_h^t)$	$\phi^t(h) = H(I^t; \alpha_h^t)$	From each α^t , several examples $\phi^t(h) = H(I^t; \alpha_h^t)$ were produced with
penev_B_F00221	7	■ 1.000	$\phi^t(0)$	$\phi^t(0)$	as outlined in Appendix A.1. Only the examples $\{\phi^t(0)\}$ were used to derive
penev_B_F00222	7	■ 0.997	$T = 1216$	$T = 1216$	the modes, hence $T = 1216$.
penev_B_F00223	7	■ 1.000	$\langle S \rangle(N) \rightarrow 1$	$\langle S \rangle(N) \rightarrow 1$	Two effects can be observed. First, the in-sample effect causes $\langle S \rangle(N) \rightarrow 1$
penev_B_F00224	7	■ 1.000	$h = 0$	$h = 0$	with $N \rightarrow T$ for $h = 0$. Also, for smaller deviations from in-sample-ness $ h $,
penev_B_F00225	7	■ 1.000	$ h $	$ h $	with $N \rightarrow T$ for $h = 0$. Also, for smaller deviations from in-sample-ness $ h $,
penev_B_F00226	7	■ 0.509	$\langle S \rangle(T, \alpha_h)$	$\langle S \rangle(T, \alpha_h)$	the entropic cost of the un-truncated representation, $\langle S \rangle(T, \alpha_h)$, is generally
penev_B_F00227	7	■ 1.000	$N \in [350, 750]$	$N \in [350, 750]$	Second, for the “reasonable regime” $N \in [350, 750]$, the out-of-sample ex-
penev_B_F00228	7	■ 0.999	$\phi^t(h_{+1})$	$\phi^t(h_{+1})$	ample $\phi^t(h_{+1})$ is more probable than the in-sample example $\phi^t(0)$. Therefore,
penev_B_F00229	7	■ 0.999	$\phi^t(0)$	$\phi^t(0)$	ample $\phi^t(h_{+1})$ is more probable than the in-sample example $\phi^t(0)$. Therefore,
penev_B_F00230	7	■ 0.922	$S - \log N$	$S - \log N$	Figure 5.6: $S - \log N$ Diagrams and Scale Symmetry Breaking (+)
penev_B_F00231	7	■ 1.000	$n \in \{-4, -3, 3, 4\}$	$n \in \{-4, -3, 3, 4\}$	the examples $\phi^t(h_n)$, with $n \in \{-4, -3, 3, 4\}$, have been substituted with the
penev_B_F00232	7	■ 0.996	$\phi^t(h_m)$	$\phi^t(h_m)$	examples $\phi^t(h_m)$ from Ensemble 7 (see Appendix A.1 for definition) with $h_m =$
penev_B_F00233	7	■ 0.996	$h_m =$	$h_m =$	examples $\phi^t(h_m)$ from Ensemble 7 (see Appendix A.1 for definition) with $h_m =$
penev_B_F00234	7	■ 0.990	$1 - s_m, s_m = 1.025^{m/3}$	$1 - s_m, s_m = 1.025^{m/3}$	$1 - s_m$, $s_m = 1.025^{m/3}$. In this notation, $n = 1$ produces almost the same
penev_B_F00235	7	■ 0.990	$n = 1$	$n = 1$	$1 - s_m$, $s_m = 1.025^{m/3}$. In this notation, $n = 1$ produces almost the same
penev_B_F00236	7	■ 1.000	$m = 3$	$m = 3$	example as $m = 3$.
penev_B_F00237	7	■ 0.911	$(- -); -1(- =); +1(+ =)$	$(- -); -1(- =); +1(+ =)$	The values of m are: -2 , labeled with $(- -)$; -1 $(- =)$; $+1$ $(+ =)$; and $+2$ $(++)$.
penev_B_F00238	7	■ 0.911	$+2(++)$	$+2(++)$	The values of m are: -2 , labeled with $(- -)$; -1 $(- =)$; $+1$ $(+ =)$; and $+2$ $(++)$.
penev_B_F00239	7	■ 0.813	α_{+1}^t	α_{+1}^t	for a reasonable truncation, α_{+1}^t is a better parameter of symmetry breaking
penev_B_F00240	7	■ 1.000	α^t	α^t	than α^t .
penev_B_F00241	7	■ 1.000	h	h	optimal value for h .
penev_B_F00242	7	■ 0.990	$N, h_{\text{best}}(N) > 0$	$N, h_{\text{best}}(N) > 0$	mensionalities N , $h_{\text{best}}(N) > 0$. Also, very big values of $ h $ result in very
penev_B_F00243	7	■ 0.963	$N, \langle S \rangle(N, \alpha_h^t)$	$N, \langle S \rangle(N, \alpha_h^t)$	Evidently, for all N , $\langle S \rangle(N, \alpha_h^t)$ has a local minimum for $h > 0$. As N
penev_B_F00244	7	■ 0.963	$h > 0$	$h > 0$	Evidently, for all N , $\langle S \rangle(N, \alpha_h^t)$ has a local minimum for $h > 0$. As N
penev_B_F00245	7	■ 1.000	$h \approx +1.5\%$	$h \approx +1.5\%$	increases, this minimum is achieved for $h \approx +1.5\%$, and is forced to $h = 0$ as
penev_B_F00246	7	■ 0.998	$h = 1.5\%$	$h = 1.5\%$	the in-sample effect takes over. Therefore, one can conclude that $h = 1.5\%$
penev_B_F00247	7	■ 1.000	$T > N$	$T > N$	which will result in $T > N$), there will be one optimal value of h for all N . A
penev_B_F00248	7	■ 0.777	$\phi^t(h)$	$\phi^t(h)$	studied $\phi^t(h)$. In this manner, for each image I^t , we produced the <i>trajectory</i>
penev_B_F00249	7	■ 1.000	$\{\Phi^{*+} \phi^t(h)\} \subset S$	$\{\Phi^{*+} \phi^t(h)\} \subset S$	$\{\Phi^{*+} \phi^t(h)\} \subset S$.
penev_B_F00250	7	■ 0.998	$\ \Phi^{*+} \phi^t(h)\ _S^2$	$\ \Phi^{*+} \phi^t(h)\ _S^2$	jectory and the origin, $\ \Phi^{*+} \phi^t(h)\ _S^2$ as the entropy (2.20) of the example $\phi^t(h)$
penev_B_F00251	7	■ 0.998	$\phi^t(h)$	$\phi^t(h)$	jectory and the origin, $\ \Phi^{*+} \phi^t(h)\ _S^2$ as the entropy (2.20) of the example $\phi^t(h)$
penev_B_F00252	7	■ 1.000	k	k	In general, α has k , more than one, independent components for the
penev_B_F00253	7	■ 0.937	$\phi^t(\alpha)$	$\phi^t(\alpha)$	special case discussed in Section 5.1 they are four (5.1)—and $\phi^t(\alpha)$ is a k -
penev_B_F00254	7	■ 0.941	$\Phi^{*+} \phi(\alpha)$	$\Phi^{*+} \phi(\alpha)$	points on the manifold $\Phi^{*+} \phi(\alpha)$ is a function of α — $S(\alpha)$. On one hand, for

Identifier	Page	Conf.	LaTeX source	Rendered	Image
penev_B_F00255	7	■ 0.941	α - $S(\alpha)$	$\alpha - S(\alpha)$	points on the manifold $\Phi^{*+}\phi(\alpha)$ is a function of $\alpha \rightarrow S(\alpha)$. On one hand, for
penev_B_F00256	7	■ 1.000	$\phi(\alpha)$	$\phi(\alpha)$	values of α that are very wrong, the probability that $\phi(\alpha)$ is in the ensemble
penev_B_F00257	7	■ 0.996	$S(\alpha)$	$S(\alpha)$	will be very low and $S(\alpha)$ will be very large a tree is not a face, as well as
penev_B_F00258	7	■ 0.995	$S_{\{\text{max}\}}$	$S_{\max}$	one can find a reasonable threshold $S_{\max}$ and a finite compact region on the
penev_B_F00259	7	■ 0.904	Δ	Δ	manifold Δ , such that $\{\alpha S(\alpha) < S_{\max}\} \subset \Delta$.
penev_B_F00260	7	■ 0.904	$\left\{ \alpha \mid S(\alpha) < S_{\max} \right\} \subset \Delta$	$\{\alpha \mid S(\alpha) < S_{\max}\} \subset \Delta$	manifold Δ , such that $\{\alpha S(\alpha) < S_{\max}\} \subset \Delta$.
penev_B_F00261	8	■ 0.787	α_{GM}^t	α_{GM}^t	and for a reasonable choice for N , and for a reasonable initial estimate $\alpha_{G,M}^t$.
penev_B_F00262	8	■ 1.000	$N \approx 10$	$N \approx 10$	Suppose that the knowledge extends only up to $N \approx 10$. In that context, only
penev_B_F00263	8	■ 0.791	X, Y	X, Y	contact, the example is centered at the fovea and, therefore, the (X, Y) part
penev_B_F00264	8	■ 1.000	Ψ_3	Ψ_3	due to the strong “Lambertian” mode Ψ_3 (Fig. 3.9) and its symmetrizations.
penev_B_F00265	8	■ 0.893	Ψ_2, Ψ_4	Ψ_2, Ψ_4	that other strong irrelevant modes—the background modes Ψ_2, Ψ_4 , and Ψ_5 ,
penev_B_F00266	8	■ 0.893	Ψ_5	Ψ_5	that other strong irrelevant modes—the background modes Ψ_2, Ψ_4 , and Ψ_5 ,
penev_B_F00267	8	■ 0.993	Ψ_1^{12}	Ψ_1^{12}	will suffice—a very frugal definition of Ψ_1 . ¹²
penev_B_F00268	8	■ 1.000	$\alpha, \mathcal{P}_\alpha$	$\alpha, \mathcal{P}_\alpha$	$\mathcal{P}_{objects}$, and the symmetric density of the parameter of the breaking $\alpha, \mathcal{P}_\alpha$.
penev_B_F00269	8	■ 0.988	$\beta_{feature}$	$\beta_{feature}$	$\beta_{feature}$, as a parameter for the breaking of the local symmetry. Analogously,
penev_B_F00270	8	■ 0.994	$\mathcal{P}_{feature}$	$\mathcal{P}_{feature}$	ensemble, this time of features, and also, its probability density $\mathcal{P}_{feature}$, along
penev_B_F00271	8	■ 1.000	$\mathcal{P}_\beta$	$\mathcal{P}_\beta$	with the probability density $\mathcal{P}_\beta$, as a parameterization of $\mathcal{P}_{objects}$.
penev_B_F00272	8	■ 1.000	Ψ_r	Ψ_r	at the fovea; this obviates the need for any genetic storage of Ψ_r .
penev_B_F00273	8	■ 0.429	$\phi^t(\mathbf{x})$	$\phi^t(\mathbf{x})$	receptors”— $\phi^t(\mathbf{x})$ was interpreted as the luminosity value at the grid point $\mathbf{x}$. This worked
penev_B_F00274	8	■ 1.000	A	A	weak assumptions, if area A projects forward to area B , then almost always
penev_B_F00275	8	■ 1.000	B	B	weak assumptions, if area A projects forward to area B , then almost always
penev_B_F00276	8	■ 0.932	20 %	20%	Ensemble 6, where about 20% of the faces have eyeglasses, the “cheek-bone”
penev_B_F00277	8	■ 0.996	$P_m(\mathbf{x})$	$P_m(\mathbf{x})$	Unfortunately, its reconstructor $P_m(\mathbf{x})$ (Fig. 5.4e) will tend to create an illusion
penev_B_F00278	8	■ 0.662	$\phi_{100}^{err} - \phi_{500}^{err}$	$\phi_{100}^{err} - \phi_{500}^{err}$	of faint eyeglasses (cf. $\phi_{100}^{err} - \phi_{500}^{err}$ on Fig. 5.2), ¹⁵ and it will take many more
penev_B_F00279	8	■ 0.919	$\{\alpha_{G,M}^t\}_{t \in T}$	$\{\alpha_{G,M}^t\}_{t \in T}$	well for two reasons: we had a good initial guess $\{\alpha_{G,M}^t\}_{t \in T}$ about the symmetry breaking
penev_B_F00280	8	■ 0.769	$\mathcal{P}_{objects}$	$\mathcal{P}_{objects}$	$\mathcal{P}_{objects}$, probability distributions in one representational module (Fig. 5.1). In general, when
penev_B_F00281	8	■ 0.916	A_3	A_3	the forward, or <i>ascending</i> , pathway originate in the superficial pyramidal cells of area A_1 ,
penev_B_F00282	9	■ 0.985	$O^{err}(\mathbf{x})$	$O^{err}(\mathbf{x})$	ule will then calculate $O^{err}(\mathbf{x})$ and will feed it to the cheekbone module for
penev_B_F00283	9	■ 1.000	U	U	achieved by topography and the simplistic fixing of the U -symmetry (4.16); as long as we
penev_B_F00284	9	■ 0.996	$\mathcal{M} \subset V$	$\mathcal{M} \subset V$	values in a sparse subset $\mathcal{M} \subset V$ of the receptor grid $\{\mathbf{x}\}_{\mathbf{x} \in V}$ (Fig. 4.6). An
penev_B_F00285	9	■ 0.996	$\{\mathbf{x}\}_{\mathbf{x} \in V}$	$\{\mathbf{x}\}_{\mathbf{x} \in V}$	values in a sparse subset $\mathcal{M} \subset V$ of the receptor grid $\{\mathbf{x}\}_{\mathbf{x} \in V}$ (Fig. 4.6). An
penev_B_F00286	9	■ 0.868	$\mathbf{P} = \mathbf{O}^* \mathbf{O} (4.28)$	$\mathbf{P} = \mathbf{O}^* \mathbf{O} (4.28)$	residual correlations $\mathbf{P} = \mathbf{O}^* \mathbf{O}$ (4.28), and, in the symmetric case, their support
penev_B_F00287	9	■ 0.949	$\mathbf{O} * \mathbf{O} = \mathbf{1} (4.12)$	$\mathbf{O} * \mathbf{O} = \mathbf{1} (4.12)$	collapses to the point in the center, $\mathbf{O}^* \mathbf{O} = \mathbf{1}$ (4.12). Therefore, the knowledge
penev_B_F00288	9	■ 0.536	$\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$	$\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$	of the values $\{O(\mathbf{x}_m)\}_{\mathbf{x}_m \in \mathcal{M}}$ does not provide any information about the rest
penev_B_F00289	9	■ 0.961	$\mathbf{f} \in \Delta_j$	$\mathbf{f} \in \Delta_j$	$\mathbf{f} \in \Delta_j$. Therefore, the correlation function of the sensory ensemble Φ_j , as seen
penev_B_F00290	9	■ 0.961	Φ_j	Φ_j	$\mathbf{f} \in \Delta_j$. Therefore, the correlation function of the sensory ensemble Φ_j , as seen
penev_B_F00291	9	■ 0.994	j	j	by a neuron in the j -th frequency band, is $\mathbf{R}_j \equiv \mathbf{K}_j^{(-2)}$ with (cf. 4.27)
penev_B_F00292	9	■ 0.994	$\mathbf{R}_j \equiv \mathbf{K}_j^{(-2)}$	$\mathbf{R}_j \equiv \mathbf{K}_j^{(-2)}$	by a neuron in the j -th frequency band, is $\mathbf{R}_j \equiv \mathbf{K}_j^{(-2)}$ with (cf. 4.27)
penev_B_F00293	9	■ 0.692	$\phi_j = \mathbf{P}_j \phi$	$\phi_j = \mathbf{P}_j \phi$	The sensory signal that a multi-scale neuron sees is $\phi_j = \mathbf{P}_j \phi$ with $\mathbf{P}_j = \mathbf{K}_j^{(0)}$,
penev_B_F00294	9	■ 0.692	$\mathbf{P}_j = \mathbf{K}_j^{(0)}$	$\mathbf{P}_j = \mathbf{K}_j^{(0)}$	The sensory signal that a multi-scale neuron sees is $\phi_j = \mathbf{P}_j \phi$ with $\mathbf{P}_j = \mathbf{K}_j^{(0)}$,
penev_B_F00295	9	■ 0.910	$\mathbf{K}_j = \mathbf{K}_j^{(1)}$	$\mathbf{K}_j = \mathbf{K}_j^{(1)}$	Now, the outputs of the action of $\mathbf{K}_j = \mathbf{K}_j^{(1)}$ on ϕ_j is ²⁰
penev_B_F00296	9	■ 0.910	ϕ_j	ϕ_j	Now, the outputs of the action of $\mathbf{K}_j = \mathbf{K}_j^{(1)}$ on ϕ_j is ²⁰
penev_B_F00297	9	■ 0.499	$O(\mathbf{x})$	$O(\mathbf{x})$	²⁰ Notably, j is not a substitute for the receptor grid index $\mathbf{x}$ in $O(\mathbf{x})$; there are several band-
penev_B_F00298	9	■ 1.000	O_j	O_j	passed dense LFA outputs O_j , which collectively carry the information that was previously
penev_B_F00299	9	■ 1.000	O	O	carried entirely by O . When we want to emphasize their dependence on $\{\mathbf{x}\}$, we will write
penev_B_F00300	9	■ 1.000	$\{\mathbf{x}\}$	$\{\mathbf{x}\}$	carried entirely by O . When we want to emphasize their dependence on $\{\mathbf{x}\}$, we will write
penev_B_F00301	9	■ 0.439	$(\hat{\mathbf{x}}, O_j) = O_j(\mathbf{x})$	$(\hat{\mathbf{x}}, O_j) = O_j(\mathbf{x})$	$(\hat{\mathbf{x}}, O_j) = O_j(\mathbf{x})$. For the ensemble of monaural sound, $\mathbf{x}$ is the time index t ; for the static
penev_B_F00302	9	■ 0.439	t	t	$(\hat{\mathbf{x}}, O_j) = O_j(\mathbf{x})$. For the ensemble of monaural sound, $\mathbf{x}$ is the time index t ; for the static
penev_B_F00303	9	■ 0.999	(x, y)	(x, y)	natural images, the pair (x, y) ; for the dynamic natural “movies,” the triplet (x, y, t) .
penev_B_F00304	9	■ 0.999	(x, y, t)	(x, y, t)	natural images, the pair (x, y) ; for the dynamic natural “movies,” the triplet (x, y, t) .
penev_B_F00305	9	■ 1.000	$\{\Delta_j\}$	$\{\Delta_j\}$	²¹ Here, we are not addressing the possibility that the bands $\{\Delta_j\}$ may happen to be non-
penev_B_F00306	9	■ 0.999	$\mathbf{O}_j^* \mathbf{O}_k$	$\mathbf{O}_j^* \mathbf{O}_k$	question what $\mathbf{O}_j^* \mathbf{O}_k$ is for $j \neq k$.
penev_B_F00307	9	■ 0.999	$j \neq k$	$j \neq k$	question what $\mathbf{O}_j^* \mathbf{O}_k$ is for $j \neq k$.
penev_B_F00308	9	■ 1.000	F_r	F_r	One way to deal uniformly with all cases is through the “noise suppression” factor F_r .
penev_B_F00309	9	■ 0.995	$(\mathbf{P} = \mathbf{1})$	$(\mathbf{P} = \mathbf{1})$	page 61); it could be decorrelated fully ($\mathbf{P} = \mathbf{1}$) by the retinal filters (4.26), but

Identifier	Page	Conf.	LaTeX source	Rendered	Image
penev_B_F00310	9	■ 1.000	$\left\{\phi_j\right\}$	$\{\phi_j\}$	among the bands $\{\Delta_j\}$, resulting in $\{\phi_j\}$, so is the dense LFA output, $\{O_j\}$.
penev_B_F00311	9	■ 1.000	$\left\{0_j\right\}$	$\{O_j\}$	among the bands $\{\Delta_j\}$, resulting in $\{\phi_j\}$, so is the dense LFA output, $\{O_j\}$.
penev_B_F00312	9	■ 1.000	$\mathrm{0}_j$	O_j	the respective output ensembles O_j , are symmetric. The multi-scale process-
penev_B_F00313	9	■ 1.000	$0_{\{3\}}$	O_3	after a specific example ϕ has been presented and the degenerate outputs O_j
penev_B_F00314	9	■ 0.858	$\mathbf{K}_{\{j\}}^{\{(\mathbf{n})\}}=\sum_{r\in\Delta_j}\psi_r\left(\frac{1}{\sigma_r}\right)^n(\psi_r,)$	$\mathbf{K}_j^{(\mathbf{n})}=\sum_{r\in\Delta_j}\psi_r\left(\frac{1}{\sigma_r}\right)^n(\psi_r,)$	representation of the type $\mathbf{K}_j^{(\mathbf{n})}=\sum_{r\in\Delta_j}\psi_r\left(\frac{1}{\sigma_r}\right)^n(\psi_r,)$ (cf. 4.30) can result in additional
penev_B_F00315	9	■ 0.695	$\{ \}^{\{23\}}\mathrm{As}$	$^{23}\mathrm{As}$	$^{23}\mathrm{As}$ with any symmetry breaking, it can be understood as a definition of a restricted
penev_B_F00316	10	■ 0.723	$\left\{a_m\right\}(\mathbf{x})\right\}_{x_m\in\mathcal{M}}\left\{x_{\{m\}}\in\mathcal{M}\right\}(4.68)$	$\{a_m(\mathbf{x})\}_{x_m\in\mathcal{M}}(4.68)$	struction as either the set of interpolation reconstructors $\{a_m(\mathbf{x})\}_{\mathbf{x}_m\in\mathcal{M}}(4.68)$
penev_B_F00317	10	■ 0.869	$P_{\{m\}}(\mathbf{x})(4.73)$	$P_m(\mathbf{x})(4.73)$	$P_m(\mathbf{x})(4.73)$.
penev_B_F00318	10	■ 1.000	$ \mathcal{M} =M-1$	$ \mathcal{M} =M-1$	We will consider the case when $ \mathcal{M} =M-1$ units are currently active and the
penev_B_F00319	10	■ 1.000	M	M	M -th one is being activated. ²⁷ Then, two things happen in the reconstruction
penev_B_F00320	10	■ 1.000	$M-1$	$M-1$	generated; and corrections to the old $M-1$ reconstructors are made.
penev_B_F00321	10	■ 0.403	$\mathbf{Q}(4.63)$	$\mathbf{Q}(4.63)$	inversion of the matrix $\mathbf{Q}(4.63)$ and the correlators $P_m(\mathbf{x})(4.68)$; the latter
penev_B_F00322	10	■ 0.403	$P_{\{m\}}(\mathbf{x})(4.68)$	$P_m(\mathbf{x})(4.68)$	inversion of the matrix $\mathbf{Q}(4.63)$ and the correlators $P_m(\mathbf{x})(4.68)$; the latter
penev_B_F00323	10	■ 0.999	$n=n^{\prime}+n^{\prime\prime}$	$n=n^{\prime}+n^{\prime\prime}$	In general, for $n=n^{\prime}+n^{\prime\prime}$, any $n\times n$ matrix W can be partitioned as
penev_B_F00324	10	■ 0.999	$n\times n$	$n\times n$	In general, for $n=n^{\prime}+n^{\prime\prime}$, any $n\times n$ matrix W can be partitioned as
penev_B_F00325	10	■ 0.999	W	W	In general, for $n=n^{\prime}+n^{\prime\prime}$, any $n\times n$ matrix W can be partitioned as
penev_B_F00326	10	■ 0.745	$\mathbf{1}_{n^{\prime}}^{\{n^{\prime}\}}$	$\mathbf{1}_{n^{\prime}}$	where $\mathbf{1}_{n^{\prime}}$ is the $n^{\prime}\times n^{\prime}$ unit matrix; analogously for $\mathbf{1}_{n^{\prime\prime}}$; and the partition-
penev_B_F00327	10	■ 0.745	$n^{\{n^{\prime}\}}\times n^{\{n^{\prime}\}}$	$n^{\prime}\times n^{\prime}$	where $\mathbf{1}_{n^{\prime}}$ is the $n^{\prime}\times n^{\prime}$ unit matrix; analogously for $\mathbf{1}_{n^{\prime\prime}}$; and the partition-
penev_B_F00328	10	■ 0.745	$\mathbf{1}_{n^{\prime\prime}}^{\{n^{\prime}\prime}\}$	$\mathbf{1}_{n^{\prime\prime}}$	where $\mathbf{1}_{n^{\prime}}$ is the $n^{\prime}\times n^{\prime}$ unit matrix; analogously for $\mathbf{1}_{n^{\prime\prime}}$; and the partition-
penev_B_F00329	10	■ 1.000	W^{-1}	W^{-1}	derstanding that they are as in (5.9). Then, for the inverse W^{-1} , we have
penev_B_F00330	10	■ 0.999	w^{-1}	w^{-1}	Thus, W^{-1} is calculated by a correction to w^{-1} , and only the quantities x , y ,
penev_B_F00331	10	■ 0.999	x , y	x, y	Thus, W^{-1} is calculated by a correction to w^{-1} , and only the quantities x , y ,
penev_B_F00332	10	■ 1.000	z	z	and z are needed additionally; W and w are never needed explicitly.
penev_B_F00333	10	■ 1.000	w	w	and z are needed additionally; W and w are never needed explicitly.
penev_B_F00334	10	■ 1.000	$M\times M$	$M\times M$	After the M -th unit becomes active, we identify W with the $M\times M$ ma-
penev_B_F00335	10	■ 0.513	$(n=M)$	$(n=M)$	trix $\mathbf{Q}(n=M)$, whose matrix elements are (4.63)
penev_B_F00336	10	■ 0.933	$(M-1)\times(M-1)$	$(M-1)\times(M-1)$	Also, we identify w with the $(M-1)\times(M-1)$ matrix from the previous step,
penev_B_F00337	10	■ 0.861	$w\equiv q$	$w\equiv q$	$w\equiv q$, comprised by the first $M-1$ rows and columns of $\mathbf{Q}$, $q_{ml}=Q_{ml} _{l,m=1}^{M-1}$.
penev_B_F00338	10	■ 0.861	$\mathbf{Q}, q_{ml}=\left.Q_{\{m\}}\right _{\{l,n\}=1}^{M-1}$	$\mathbf{Q}, q_{ml}=Q_{ml} _{l,ns=1}^{M-1}$	$w\equiv q$, comprised by the first $M-1$ rows and columns of $\mathbf{Q}$, $q_{ml}=Q_{ml} _{l,m=1}^{M-1}$.
penev_B_F00339	10	■ 0.994	$w, n^{\{n^{\prime}\}}=M-1$	$w, n^{\prime}=M-1$	With these W and w , $n^{\prime}=M-1$, hence $n^{\prime\prime}=1$; then x is a $(M-1)$ -
penev_B_F00340	10	■ 0.994	$n^{\{n^{\prime}\prime}\}=1$	$n^{\prime\prime}=1$	With these W and w , $n^{\prime}=M-1$, hence $n^{\prime\prime}=1$; then x is a $(M-1)$ -
penev_B_F00341	10	■ 0.994	x	x	With these W and w , $n^{\prime}=M-1$, hence $n^{\prime\prime}=1$; then x is a $(M-1)$ -
penev_B_F00342	10	■ 0.994	$(M-1)$	$(M-1)$	With these W and w , $n^{\prime}=M-1$, hence $n^{\prime\prime}=1$; then x is a $(M-1)$ -
penev_B_F00343	10	■ 0.908	y	y	dimensional column vector; y , a row vector; z and $\bar{z}$, scalars. Because $\mathbf{P}(4.29)$
penev_B_F00344	10	■ 0.908	$\bar{z}$	$\bar{z}$	dimensional column vector; y , a row vector; z and $\bar{z}$, scalars. Because $\mathbf{P}(4.29)$
penev_B_F00345	10	■ 0.806	p	p	and, therefore, $\mathbf{Q}$ are symmetric, we can define p as the row vector ²⁸
penev_B_F00346	10	■ 0.917	$\tilde{z}$	$\tilde{z}$	Because $\mathbf{Q}$ is symmetric and $\tilde{z}$ is a scalar, the second term in (5.11) factorizes
penev_B_F00347	10	■ 0.870	$O^{\{r\}}(\mathbf{x})$	$O^{ree}(\mathbf{x})$	means that $O^{rec}(\mathbf{x})$ is a V -dimensional column vector, which is a linear com-
penev_B_F00348	10	■ 1.000	$A^{\{m\}}(\mathbf{x})$	$A^m(\mathbf{x})$	bination of the M (uppercase) column vectors $A^m(\mathbf{x})$ with the M coefficients
penev_B_F00349	10	■ 1.000	$O_{\{m\}}=O\left(\mathbf{x}_m\right)$	$O_m=O(\mathbf{x}_m)$	$O_m=O(\mathbf{x}_m)$. Analogously, the column vector $A^m(\mathbf{x})$ for the m -th reconstruc-
penev_B_F00350	10	■ 1.000	$\{ \}^{\{28\}}p^{\{T\}}$	$^{28}p^T$	$^{28}p^T$ is a column vector, which is the <i>transpose</i> of the row vector p .
penev_B_F00351	10	■ 1.000	$P^{\{l\}}(\mathbf{x})$	$P^l(\mathbf{x})$	is a linear combination of the M correlators $P^l(\mathbf{x})$, centered at the active grid
penev_B_F00352	10	■ 0.967	$\mathbf{x}_l$	$\mathbf{x}_l$	points, $\mathbf{x}_i$, with the coefficients $(\mathbf{Q}^{-1})_i^m$, from the m -th column of $\mathbf{Q}^{-1}$.
penev_B_F00353	10	■ 0.967	$\left(\mathbf{Q}^{-1}\right)_l^m$	$(\mathbf{Q}^{-1})_l^m$	points, $\mathbf{x}_i$, with the coefficients $(\mathbf{Q}^{-1})_i^m$, from the m -th column of $\mathbf{Q}^{-1}$.
penev_B_F00354	10	■ 0.994	$\left[a^{\{m\}}(\mathbf{x})\quad 0\right]$	$[a^m(\mathbf{x})\ 0]$	Thus, the new reconstructors $A^m(\mathbf{x})$ are the old ones, $\left[\begin{array}{c} a^m(\mathbf{x}) \\ 0 \end{array}\right]$, with a
penev_B_F00355	10	■ 0.880	$A(\mathbf{x})$	$A(\mathbf{x})$	whose entire $\mathbf{x}$ -dependence is given by $A(\mathbf{x})$. Indeed, the new reconstructor
penev_B_F00356	10	■ 1.000	$A^{\{M\}}(\mathbf{x})$	$A^M(\mathbf{x})$	$A^M(\mathbf{x})$ is proportional to $A(\mathbf{x})$ with a coefficient $\tilde{z}$,
penev_B_F00357	10	■ 1.000	$\tilde{z}_1$	$\tilde{z}_1$	$A^M(\mathbf{x})$ is proportional to $A(\mathbf{x})$ with a coefficient $\tilde{z}$,
penev_B_F00358	10	■ 0.886	$a^{\{m\}}(\mathbf{x})$	$a^m(\mathbf{x})$	and the correction to any old reconstructor $a^m(\mathbf{x})$, is proportional to $A^M(\mathbf{x})$
penev_B_F00359	10	■ 0.886	$A^{\{M\}}(\mathbf{x})$	$A^M(\mathbf{x})$	and the correction to any old reconstructor $a^m(\mathbf{x})$, is proportional to $A^M(\mathbf{x})$
penev_B_F00360	10	■ 1.000	$-u^m$	$-u^m$	with a coefficient $-u^m$:
penev_B_F00361	10	■ 0.981	$u=pq^1$	$u=pq^1$	Now, we are well motivated to inquire into the meaning of $u=pq^{-1}$ (5.15).
penev_B_F00362	10	■ 0.993	u^m	u^m	From (5.13) (cf. 5.18), u^m is the old reconstructor $a^m(\mathbf{x})$, evaluated at the

Identifier	Page	Conf.	LaTeX source	Rendered	Image
penev_B_F00363	10	0.679	$\mathrm{x}_{[M]}$	$\mathbf{x}_M$	newly active point, $\mathbf{x}_M$,
penev_B_F00364	10	0.815	$p^{\{l\}}(\mathbf{u}^{\{T\}}) = p^{\{l\}}(\mathbf{x}) \left(q^{-1}\right)_l^m P^M(\mathbf{x}_m) = a^m(\mathbf{x}) P^M(\mathbf{x}_m)$	$p^l(\mathbf{x}) u^{T_l} = p^l(\mathbf{x}) (q^{-1})_l^m P^M(\mathbf{x}_m) = a^m(\mathbf{x}) P^M(\mathbf{x}_m)$	Also, (cf. 5.20), $p^l(\mathbf{x}) u^{T_l} = p^l(\mathbf{x}) (q^{-1})_l^m P^M(\mathbf{x}_m) = a^m(\mathbf{x}) P^M(\mathbf{x}_m)$; hence
penev_B_F00365	10	0.943	$a^{\{m\}}(\mathbf{x}) P^{\{M\}}(\mathbf{x}_{[m]})$	$a^m(\mathbf{x}) P^M(\mathbf{x}_m)$	The last term in (5.25), $a^m(\mathbf{x}) P^M(\mathbf{x}_m)$, is analogous to (5.17) with the
penev_B_F00366	10	0.629	$\left\{\mathbf{x}_{[m]}\right\}$	$\{\mathbf{x}_m\}$	evaluated at $\{\mathbf{x}_m\}$ is $P^M(\mathbf{x})$, rather than $O(\mathbf{x})$. Therefore, it is useful, given
penev_B_F00367	10	0.629	$P^{\{M\}}(\mathbf{x})$	$P^M(\mathbf{x})$	evaluated at $\{\mathbf{x}_m\}$ is $P^M(\mathbf{x})$, rather than $O(\mathbf{x})$. Therefore, it is useful, given
penev_B_F00368	10	1.000	$V(\mathbf{x})$	$V(\mathbf{x})$	the current restriction $\mathcal{M}$, to define, for any vector $V(\mathbf{x})$, the <i>reconstruction</i>
penev_B_F00369	11	0.926	$P^{\{M\}}(\mathbf{x})$	$P^M(\mathbf{x})$	Remarkably, $A(\mathbf{x})$ is the difference between the true value of $P^M(\mathbf{x})$ and its
penev_B_F00370	11	1.000	$\bar{z}^{-1} = \bar{z} - u p^T$	$\bar{z}^{-1} = \bar{z} - u p^T$	$\bar{z}^{-1} = \bar{z} - u p^T$, and from (5.14), (5.24), (5.13), and (5.29),
penev_B_F00371	11	0.868	$a^{\{m\}}(\mathbf{x})$	$a^m(\mathbf{x})$	predictors, $a^m(\mathbf{x})$, are corrected by a negative feedback of this error, in propor-
penev_B_F00372	11	0.870	$a^{\{m\}}(\mathbf{x}_{[M]})$	$a^m(\mathbf{x}_M)$	tionality with their original prediction strengths, $a^m(\mathbf{x}_M)$.
penev_B_F00373	11	0.582	$O^{\{\text{rec}\}}(\mathbf{x}) = \mathcal{R}_{\mathcal{O}}^{\{M-1\}}(\mathbf{x})$	$O^{\text{rec}}(\mathbf{x}) = \mathcal{R}_O^{M-1}(\mathbf{x})$	the current estimate $O^{\text{rec}}(\mathbf{x}) = \mathcal{R}_O^{M-1}(\mathbf{x})$, the current error $O^{\text{err}}(\mathbf{x}) = \mathcal{D}_O(\mathbf{x})$
penev_B_F00374	11	0.582	$O^{\{\text{err}\}}(\mathbf{x}) = \mathcal{D}_{\mathcal{O}}(\mathbf{x})$	$O^{\text{err}}(\mathbf{x}) = \mathcal{D}_O(\mathbf{x})$	the current estimate $O^{\text{rec}}(\mathbf{x}) = \mathcal{R}_O^{M-1}(\mathbf{x})$, the current error $O^{\text{err}}(\mathbf{x}) = \mathcal{D}_O(\mathbf{x})$
penev_B_F00375	11	0.381	$\Delta O^{\{\text{rec}\}}(\mathbf{x})$	$\Delta O^{\text{rec}}(\mathbf{x})$	introduces a correction $\Delta O^{\text{rec}}(\mathbf{x})$, which is exactly equal to the error at that
penev_B_F00376	11	0.716	$\mathcal{D}_{\mathcal{O}}(\mathbf{x}_{[M]})$	$\mathcal{D}_O(\mathbf{x}_M)$	on the current reconstruction error, $\mathcal{D}_O(\mathbf{x}_M)$.
penev_B_F00377	11	0.490	$O^{\{\text{operatorname{err}}\}}(\mathbf{x}_{[M]}) = \mathcal{D}_{\mathcal{O}}(\mathbf{x}_{[M]})$	$O^{\text{err}}(\mathbf{x}_M) = \mathcal{D}_O(\mathbf{x}_M)$	$O^{\text{err}}(\mathbf{x}_M) = \mathcal{D}_O(\mathbf{x}_M)$.
penev_B_F00378	11	1.000	$f(t)$	$f(t)$	like to understand it as an infinite source $f(t)$ (cf. Appendix A.6), from which
penev_B_F00379	11	0.551	$\phi^{\{t_n\}}(\mathbf{x}_{[k]}) = f(t_{\mathbf{n}-k})$	$\phi^{t_n}(\mathbf{x}_k) = f(t_{\mathbf{n}-k})$	we will construct finite-dimensional ensembles of examples $\phi^{t_n}(\mathbf{x}_k) = f(t_{n-k})$
penev_B_F00380	11	1.000	$t_{[n]}$	t_n	by imposing a finite discrete window of size V around the time t_n (A.13).
penev_B_F00381	11	0.975	$\left\{f_{\chi}\right\}_{\chi \in X}$	$\{f_{\chi}\}_{\chi \in X}$	The <i>Encoder</i> (a) has access to the signal $\{f_{\chi}\}_{\chi \in X}$. Based on the part of the
penev_B_F00382	11	0.467	$\mathbf{x}_{[n]}$	$\mathbf{x}_n$	location $\mathbf{x}_n$, an estimate $f^{\text{rec}}(\mathbf{x}_n) \equiv f_n^{\text{rec}}$, and the <i>Quantizer</i> quantizes it to $\hat{f}_n$.
penev_B_F00383	11	0.467	$f^{\{\text{rec}\}}(\mathbf{x}_{[n]}) \equiv f_n^{\{\text{rec}\}}$	$f^{\text{rec}}(\mathbf{x}_n) \equiv f_n^{\text{rec}}$	location $\mathbf{x}_n$, an estimate $f^{\text{rec}}(\mathbf{x}_n) \equiv f_n^{\text{rec}}$, and the <i>Quantizer</i> quantizes it to $\hat{f}_n$.
penev_B_F00384	11	0.467	$f_{[n]}$	f_n	location $\mathbf{x}_n$, an estimate $f^{\text{rec}}(\mathbf{x}_n) \equiv f_n^{\text{rec}}$, and the <i>Quantizer</i> quantizes it to $\hat{f}_n$.
penev_B_F00385	11	0.996	$e_{[n]} = f_{[n]} - \hat{f}_{[n]}$	$e_n = f_n - \hat{f}_n$	The error $e_n = f_n - \hat{f}_n$ is calculated and is fed to the <i>Symbol Encoder</i> , which
penev_B_F00386	11	1.000	$e_{[n]}$	e_n	error signal e_n . This error is added to the current reconstruction $\hat{f}_n$, which
penev_B_F00387	11	1.000	$f_{[n]} = \hat{f}_{[n]} + e_{[n]}$	$f_n = \hat{f}_n + e_n$	The result, $f_n = \hat{f}_n + e_n$, is an identical copy of f_n .
penev_B_F00388	11	0.934	$f^{\{\text{rec}\}}(t_{[n]}) = f_{[n]}$	$f^{\text{rec}}(t_n) = f_n$	mates of the of the current value, $f^{\text{rec}}(t_n) = \hat{f}_n$. The encoder, which has access
penev_B_F00389	11	0.824	$e_{[n]} = f_{[n]} - \hat{f}_{[n]}$	$e_n = f_n - \hat{f}_n$	to the true value f_n , calculates the error $e_n = f_n - \hat{f}_n$ and feeds it, via the com-
penev_B_F00390	11	0.999	$t_{[n]}, \left\{f_{[n]}(m) = f(t_{[n-m]}) = f_{[n-m]}\right\}_{m=1}^M$	$t_n, \{f_n(m) = f(t_{n-m}) = f_{n-m}\}_{m=1}^M$	only a small time window, <i>preceding</i> t_n , $\{f_n(m) = f(t_{n-m}) = f_{n-m}\}_{m=1}^M$. The
penev_B_F00391	11	0.995	$\hat{f}_{[n]} = \alpha^{\{m\}} f_{[n-m]}$	$\hat{f}_n = \alpha^m f_{n-m}$	similar to (5.17)— $\hat{f}_n = \alpha^m f_{n-m}$.
penev_B_F00392	11	1.000	$\alpha_{[m]}$	α_m	There are many possible choices for the coefficients α_m ; when they are
penev_B_F00393	11	0.998	$E\left\{e_{[n]}^2\right\}$	$E\left\{e_n^2\right\}$	chosen to minimize the expectation of the square of the error signal, $E\{e_n^2\}$, the
penev_B_F00394	11	1.000	$\left\{t_{[n-m]}\right\}_{m=1}^{M-1}$	$\{t_{n-m}\}_{m=1}^{M-1}$	is known, $\{t_{n-m}\}_{m=1}^{M-1}$, with $\{\mathbf{x}_m\}_{m=1}^{M-1}$, and the currently reconstructed grid
penev_B_F00395	11	1.000	$\left\{\mathbf{x}_{[m]}\right\}_{m=1}^{M-1}$	$\{\mathbf{x}_m\}_{m=1}^{M-1}$	is known, $\{t_{n-m}\}_{m=1}^{M-1}$, with $\{\mathbf{x}_m\}_{m=1}^{M-1}$, and the currently reconstructed grid
penev_B_F00396	11	0.964	$\mathbf{x}_{[M]}$	$\mathbf{x}_M$	point t_n , with $\mathbf{x}_M$. Then $\mathbf{Q}$ (5.40) is the correlation matrix of the signal,
penev_B_F00397	12	1.000	Φ	Φ	of experimentally well-controlled sensory stimuli Φ , interpreted as the horizon-
penev_B_F00398	12	1.000	$v(t)$	$v(t)$	tal velocity $v(t)$; simultaneously, the spike train $\{t_m\}$ that results from this
penev_B_F00399	12	1.000	$\left\{t_{[m]}\right\}$	$\{t_m\}$	tal velocity $v(t)$; simultaneously, the spike train $\{t_m\}$ that results from this
penev_B_F00400	12	0.702	$A^{\{M\}}(t) = F_1(t - t_{[M]})$	$A^M(t) = F_1(t - t_M)$	enforced, $A^M(t) = F_1(t - t_M)$, and the m.s.e., $E\{(v(t) - v^{\text{rec}}(t))^2\}$, is minimized
penev_B_F00401	12	0.702	$E\left\{\left(v(t) - v^{\{\text{rec}\}}(t)\right)^2\right\}$	$E\left\{(v(t) - v^{\text{rec}}(t))^2\right\}$	enforced, $A^M(t) = F_1(t - t_M)$, and the m.s.e., $E\{(v(t) - v^{\text{rec}}(t))^2\}$, is minimized
penev_B_F00402	12	1.000	$F(\tau)$	$F(\tau)$	over the experimental data set, the best $F(\tau)$ is determined. It is used to
penev_B_F00403	12	0.946	$v^{\{\text{rec}\}}(t)$	$v^{\text{rec}}(t)$	calculate the mutual information between $v(t)$ and $v^{\text{rec}}(t)$, which is a lower
penev_B_F00404	12	1.000	$A^{\{m\}}(t)$	$A^m(t)$	lar to (5.36), there are two major differences. The reconstructors $A^m(t)$ are
penev_B_F00405	12	0.424	$t_{[M]}$. ³⁶	t_M . ³⁶	identical—they are based on a single grid point, t_M . ³⁶
penev_B_F00406	12	0.974	100% . ³⁷	100% . ³⁷	about the actual signal will be much higher, close to 100% . ³⁷
penev_B_F00407	12	0.974	$\left\{^{\{36\}}\mathrm{An}\right\}$	^{36}An	^{36}An attempt to base them on two grid points, t_M and t_{M-1} , albeit also in a time-
penev_B_F00408	12	0.974	$t_{[M]}$	t_M	^{36}An attempt to base them on two grid points, t_M and t_{M-1} , albeit also in a time-
penev_B_F00409	12	0.974	$t_{[M-1]}$	t_{M-1}	^{36}An attempt to base them on two grid points, t_M and t_{M-1} , albeit also in a time-
penev_B_F00410	12	0.998	$5-15\%$	$5-15\%$	(5–15%, depending on the system under study).
penev_B_F00411	12	0.996	$\mathcal{P}_{\{\text{objects}\}}$	$\mathcal{P}_{\text{objects}}$	of the probability densities of images, $\mathcal{P}_{\text{images}}$, and of objects, $\mathcal{P}_{\text{objects}}$. We

Identifier	Page	Conf.	LaTeX source	Rendered	Image
penev_B_F00412	12	■ 0.964	$\mathcal{P}_{\alpha}$	$\mathcal{P}_{\alpha}$	understood the probability distributions $\mathcal{P}_{\alpha}$ and $\mathcal{P}_{object}$ as part of the explicit
penev_B_F00413	12	■ 0.515	$\mathcal{P}_{\text{imnges}}$	$\mathcal{P}_{\text{imnges}}$	parameterization of $\mathcal{P}_{images}$.
penev_B_F00414	12	■ 0.800	$\mathcal{P}_{\text{o b j e c t s}}$	$\mathcal{P}_{objects}$	the re-parameterization of $\mathcal{P}_{objects}$ with $\mathcal{P}_{\beta}$ and $\mathcal{P}_{future}$. This hierarchical pa-
penev_B_F00415	13	■ 0.989	$0^{\{r \text{ e c}\}}(\mathbf{x}) = \mathcal{R}_{\{0\}^{M-1}}(\mathbf{x})$	$O^{rec}(\mathbf{x}) = \mathcal{R}_O^{M-1}(\mathbf{x})$	identical reconstructions $O^{rec}(\mathbf{x}) = \mathcal{R}_O^{M-1}(\mathbf{x})$; the encoder calculates the recon-
penev_B_F00416	13	■ 0.991	$\mathcal{D}_{\{\mathcal{O}\}}(\mathbf{x})$	$\mathcal{D}_O(\mathbf{x})$	$\mathcal{D}_O(\mathbf{x})$ was largest. Here, we pose the question, Is it possible to find another
penev_B_F00417	13	■ 0.945	$\mathcal{D}_{\{0\}}(\mathbf{x}_{\text{spike}}) = \theta$	$\mathcal{D}_O(\mathbf{x}_{\text{spike}}) = \theta$	time the current reconstruction error is equal to the threshold, $\mathcal{D}_O(\mathbf{x}_{\text{spike}}) = \theta$.
penev_B_F00418	13	■ 0.999	$\mathcal{D}_{\{0\}}(\mathbf{x})^2$	$\mathcal{D}_O(\mathbf{x})^2$	the individual grid points, $\mathcal{D}_O(\mathbf{x})^2$. On one hand, when θ is relatively large,
penev_B_F00419	13	■ 0.938	$\mathcal{R}_{\{0\}^M}(\mathbf{x})$	$\mathcal{R}_O^M(\mathbf{x})$	tion, $\mathcal{R}_O^M(\mathbf{x})$, but also, a current expectation of the variance of the error,
penev_B_F00420	13	■ 0.867	$D(\mathbf{x})^2 = E\{\mathcal{D}_{\{0\}}(\mathbf{x})^2\}$	$D(\mathbf{x})^2 = E\{\mathcal{D}_O(\mathbf{x})^2\}$	$D(\mathbf{x})^2 = E\{\mathcal{D}_O(\mathbf{x})^2\}$ (see discussion in footnote 32 on page 77). Then, θ may
penev_B_F00421	13	■ 0.495	$\theta(\mathbf{x} - \mathbf{x}_{\text{vpike}})$	$\theta(\mathbf{x} - \mathbf{x}_{\text{vpike}})$	decays with the time since the last spike, $\theta(\mathbf{x} - \mathbf{x}_{\text{vpike}})$.
penev_B_F00422	13	■ 0.508	$\mathcal{D}_{\{0\}}(\mathbf{x})$	$\mathcal{D}_O(\mathbf{x})$	the timing, which generates an error in the decoder's estimate of $\mathcal{D}_O(\mathbf{x})$ and,
penev_B_F00423	13	■ 0.842	$0^{\{r \text{ e c}\}}(\mathbf{x})$	$O^{rec}(\mathbf{x})$	therefore, of $O^{rec}(\mathbf{x})$, which depends on the <i>derivative</i> of $O(\mathbf{x})$. ³¹ One way to
penev_B_F00424	13	■ 0.842	$0(\mathbf{x}) .^{31}$	$O(\mathbf{x}) .^{31}$	therefore, of $O^{rec}(\mathbf{x})$, which depends on the <i>derivative</i> of $O(\mathbf{x})$. ³¹ One way to
penev_B_F00425	13	■ 1.000	Δt	Δt	is determined with precision Δt , we may consider that a spike can be triggered
penev_B_F00426	13	■ 1.000	$D(\mathbf{x})$	$D(\mathbf{x})$	compares the target function, for example $D(\mathbf{x})$, with the threshold $\theta(\mathbf{x})$. Ob-
penev_B_F00427	13	■ 1.000	$\theta(\mathbf{x})$	$\theta(\mathbf{x})$	compares the target function, for example $D(\mathbf{x})$, with the threshold $\theta(\mathbf{x})$. Ob-
penev_B_F00428	13	■ 0.985	$\mathbf{x}_{\text{m}}$	$\mathbf{x}_m$	the difference is positive at the moment of the spike, $\mathbf{x}_m$, but was negative the
penev_B_F00429	13	■ 0.999	$\mathbf{x}'$	$\mathbf{x}'$	preceding moment, $\mathbf{x}'$.
penev_B_F00430	13	■ 0.991	$\mathbf{q}_{\mathbf{x}_m} \in S$	$\mathbf{q}_{\mathbf{x}_m} \in S$	The grid location $\mathbf{x}_m$ determines a vector $\mathbf{q}_{\mathbf{x}_m} \in S$ (4.38), which in turn de-
penev_B_F00431	13	■ 1.000	$U_m \subset S$	$U_m \subset S$	termines a one-dimensional subspace $U_m \subset S$ (vertical line), with a projector
penev_B_F00432	13	■ 0.997	$\mathbf{Q}_{\mathbf{x}_m}$	$\mathbf{Q}_{\mathbf{x}_m}$	$\mathbf{Q}_{\mathbf{x}_m}$ (4.39).
penev_B_F00433	13	■ 0.893	$\phi = \Phi^* s$	$\phi = \Phi^* s$	For the example $\phi = \Phi^* s$, in Section 4.4.2 we interpreted the choice of the
penev_B_F00434	13	■ 0.737	$O(\mathbf{x}_m) = O_m = \langle \mathbf{q}_{\mathbf{x}_m}, \mathbf{s} \rangle$	$O(\mathbf{x}_m) = O_m = \langle \mathbf{q}_{\mathbf{x}_m}, \mathbf{s} \rangle$	grid point $\mathbf{x}_m$ as the measurement $O(\mathbf{x}_m) = O_m = \langle \mathbf{q}_{\mathbf{x}_m}, \mathbf{s} \rangle$ (4.37) and we
penev_B_F00435	13	■ 0.863	$\mathbf{s}_{\parallel} = O_m \mathbf{p}_{\mathbf{x}_m} \in U_m$	$\mathbf{s}_{\parallel} = O_m \mathbf{p}_{\mathbf{x}_m} \in U_m$	introduced $\mathbf{s}_{\parallel} = O_m \mathbf{p}_{\mathbf{x}_m} \in U_m$ (4.42) (light vertical arrow, ending at A).
penev_B_F00436	13	■ 0.961	$\mathbf{q}' \in S$	$\mathbf{q}' \in S$	Analogously, any non-selected grid location $\mathbf{x}'$ determines the vector $\mathbf{q}' \in S$
penev_B_F00437	13	■ 0.659	$\mathbf{q}_{\mathbf{x}}$	$\mathbf{q}_{\mathbf{x}}$	(along the light slanted arrow). Together with $\mathbf{q}_{\mathbf{x}_m}$, they determine a 2-
penev_B_F00438	13	■ 0.659	2-	2-	(along the light slanted arrow). Together with $\mathbf{q}_{\mathbf{x}_m}$, they determine a 2-
penev_B_F00439	13	■ 1.000	U_m	U_m	U_m and U'_m .
penev_B_F00440	13	■ 1.000	U'_m	U'_m	U_m and U'_m .
penev_B_F00441	13	■ 0.843	$N - 2$	$N - 2$	All other $N - 2$ dimensions of the PCA S -subspace are collapsed to $U^{\perp}$.
penev_B_F00442	13	■ 0.843	$U^{\perp}$	$U^{\perp}$	All other $N - 2$ dimensions of the PCA S -subspace are collapsed to $U^{\perp}$.
penev_B_F00443	13	■ 0.958	$O_m \mathbf{p}'$	$O_m \mathbf{p}'$	For a given measurement O_m , we can define $O_m \mathbf{p}'$ (light slanted arrow, ending
penev_B_F00444	13	■ 1.000	$N - 1$	$N - 1$	at B). It can be treated as a normal to an $N - 1$ dimensional hyper-plane, whose
penev_B_F00445	13	■ 0.990	$(B - C)$	$(B - C)$	intersection with the plane of the circle is the slanted line $(B - C)$, tangential
penev_B_F00446	13	■ 1.000	C	C	to the circle, and which intersects with U'_m at point C . This hyper-plane's
penev_B_F00447	13	■ 0.914	$\phi = \Phi^* s$	$\phi = \Phi^* s$	the context of sparsification. Given the example $\phi = \Phi^* s$, and the choice
penev_B_F00448	13	■ 0.380	$\mathbf{s}_{\parallel} = O_m \mathbf{p}_{\mathbf{x}_m} \in U_m$ (4.42)	$\mathbf{s}_{\parallel} = O_m \mathbf{p}_{\mathbf{x}_m} \in U_m$ (4.42)	$\mathbf{s}_{\parallel} = O_m \mathbf{p}_{\mathbf{x}_m} \in U_m$ (4.42), point A on Fig. 5.10, and a vector $\mathbf{s}_{\perp}$, orthogonal
penev_B_F00449	13	■ 0.380	$\mathbf{s}_{\perp}$	$\mathbf{s}_{\perp}$	$\mathbf{s}_{\parallel} = O_m \mathbf{p}_{\mathbf{x}_m} \in U_m$ (4.42), point A on Fig. 5.10, and a vector $\mathbf{s}_{\perp}$, orthogonal
penev_B_F00450	13	■ 0.829	$\mathbf{s}_{\parallel}$	$\mathbf{s}_{\parallel}$	to it. We interpreted $\mathbf{s}_{\parallel}$ as the maximum likelihood estimate.
penev_B_F00451	13	■ 1.000	$O(\mathbf{x}_m) = O_m$	$O(\mathbf{x}_m) = O_m$	decoder cannot assume that $O(\mathbf{x}_m) = O_m$ exactly, but only that $O(\mathbf{x}_m) \geq O_m$.
penev_B_F00452	13	■ 1.000	$O(\mathbf{x}_m) \geq O_m$	$O(\mathbf{x}_m) \geq O_m$	decoder cannot assume that $O(\mathbf{x}_m) = O_m$ exactly, but only that $O(\mathbf{x}_m) \geq O_m$.
penev_B_F00453	13	■ 0.989	$O(\mathbf{x}')$	$O(\mathbf{x}')$	positive transition—that the value $O(\mathbf{x}')$, immediately prior to the spike, was
penev_B_F00454	13	■ 0.822	$O' .^{33}$	$O' .^{33}$	below a certain threshold, $O' .^{33}$ We can decompose $\mathbf{s}$ into a sum of three mutu-
penev_B_F00455	13	■ 1.000	$\mathbf{s}_{\parallel} \in U_m, \mathbf{s}' \in U'_m$	$\mathbf{s}_{\parallel} \in U_m, \mathbf{s}' \in U'_m$	ally orthogonal vectors, $\mathbf{s}_{\parallel} \in U_m, \mathbf{s}' \in U'_m$, and $\mathbf{s}_{\perp} \in U^{\perp}$. Since both $\mathbf{q}'$ and $\mathbf{q}_{\mathbf{x}_m}$
penev_B_F00456	13	■ 1.000	$\mathbf{s}_{\perp} \in U^{\perp}$	$\mathbf{s}_{\perp} \in U^{\perp}$	ally orthogonal vectors, $\mathbf{s}_{\parallel} \in U_m, \mathbf{s}' \in U'_m$, and $\mathbf{s}_{\perp} \in U^{\perp}$. Since both $\mathbf{q}'$ and $\mathbf{q}_{\mathbf{x}_m}$
penev_B_F00457	13	■ 1.000	$\mathbf{q}'$	$\mathbf{q}'$	ally orthogonal vectors, $\mathbf{s}_{\parallel} \in U_m, \mathbf{s}' \in U'_m$, and $\mathbf{s}_{\perp} \in U^{\perp}$. Since both $\mathbf{q}'$ and $\mathbf{q}_{\mathbf{x}_m}$
penev_B_F00458	13	■ 1.000	$\mathbf{q}_{\mathbf{x}_m}$	$\mathbf{q}_{\mathbf{x}_m}$	ally orthogonal vectors, $\mathbf{s}_{\parallel} \in U_m, \mathbf{s}' \in U'_m$, and $\mathbf{s}_{\perp} \in U^{\perp}$. Since both $\mathbf{q}'$ and $\mathbf{q}_{\mathbf{x}_m}$
penev_B_F00459	13	■ 0.999	$U^{\perp}, \mathbf{s}_{\perp}$	$U^{\perp}, \mathbf{s}_{\perp}$	are orthogonal to $U^{\perp}$, $\mathbf{s}_{\perp}$ can be anywhere in $U^{\perp}$, and a maximum likelihood
penev_B_F00460	13	■ 0.972	$\mathbf{s}_{\perp} = \mathbf{0}$	$\mathbf{s}_{\perp} = \mathbf{0}$	estimate of it is $\mathbf{s}_{\perp} = \mathbf{0}$.
penev_B_F00461	13	■ 0.989	$\mathbf{s}'$	$\mathbf{s}'$	On the other hand, $\mathbf{s}'$ can be anywhere on the ray in U'_m , starting at C ,
penev_B_F00462	13	■ 0.998	$\mathbf{s}' = \mathbf{0}$	$\mathbf{s}' = \mathbf{0}$	which is the mean of the respective distribution, is not $\mathbf{s}' = \mathbf{0}$, but is to the
penev_B_F00463	13	■ 0.994	D	D	left of A , at D . Also, the line $B - C$ intersects with U_m at some place, and that

Identifier	Page	Conf.	LaTeX source	Rendered	Image
penev_B_F00464	13	■ 0.994	B C	BC	left of A , at D . Also, the line $B C$ intersects with U_m at some place, and that
penev_B_F00465	13	■ 0.943	$\mathbf{o}_m = \left\langle \mathbf{q}_m, \mathbf{s}_m \right\rangle$	$o_m = \langle \mathbf{q}_m, \mathbf{s}_m \rangle$	is an upper bound for $o_m = \langle \mathbf{q}_m, \mathbf{s}_m \rangle$; hence the maximum likelihood estimate
penev_B_F00466	13	■ 0.952	$\mathbf{o}_m = \mathbf{p}_m(A)$	$O_m \mathbf{p}_m(A)$	for $\mathbf{s}_m$ is not $O_m \mathbf{p}_m(A)$, but is a bit higher. ³⁴
penev_B_F00467	13	■ 0.350	$\mathbf{a}_m = \left\langle \mathbf{q}_m, \mathbf{s}_m \right\rangle$	$a_m = \langle \mathbf{q}_m, \mathbf{s}_m \rangle$	³² The values of the projection $a_m = \langle \mathbf{q}_m, \mathbf{s}_m \rangle$ will be distributed according to the orig-
penev_B_F00468	13	■ 0.966	$\mathbf{o}_m \geq \mathbf{o}_m$	$o_m \geq O_m$	inal probability distribution, normal by assumption (2.49), <i>conditional</i> on $o_m \geq O_m$. The
penev_B_F00469	13	■ 0.553	$\mathbf{o}_m \mathbf{p}_m$	$O_m \mathbf{p}_m$	³³ For simplicity, $O_m \mathbf{p}_m$ and $O' \mathbf{p}'$ are shown to have the same lengths on Fig. 5.10. In
penev_B_F00470	13	■ 0.553	$\mathbf{o}^{\prime \prime} \mathbf{p}^{\prime \prime}$	$O' \mathbf{p}'$	³³ For simplicity, $O_m \mathbf{p}_m$ and $O' \mathbf{p}'$ are shown to have the same lengths on Fig. 5.10. In
penev_B_F00471	13	■ 0.810	$\mathbf{q}^{\prime \prime}$	$\mathbf{q}'$	probability distribution along U_m ; the smaller the angle α between $\mathbf{q}'$ and $\mathbf{q}_m$, the more
penev_B_F00472	13	■ 0.810	$\mathbf{q}_m$	$\mathbf{q}_m$	probability distribution along U_m ; the smaller the angle α between $\mathbf{q}'$ and $\mathbf{q}_m$, the more
penev_B_F00473	13	■ 1.000	$C \rightarrow A$	$C \rightarrow A$	informative the lack of a spike. Ultimately, when $C \rightarrow A$, half of the axis U_m is excluded,
penev_B_F00474	13	■ 0.848	$\mathbf{o}_m$	o_m	o_m approaches O_m , and the information due to the localization of $\mathbf{s}_m$ in U_m increases.
penev_B_F00475	13	■ 0.848	$\mathbf{s}_m$	s_m	o_m approaches O_m , and the information due to the localization of $\mathbf{s}_m$ in U_m increases.
penev_B_F00476	13	■ 0.810	$\cos \alpha = P_m(\mathbf{x}')$	$\cos \alpha = P_m(x')$	This is intuitively correct; $\cos \alpha = P_m(\mathbf{x}')$, the output correlation between two adjacent
penev_B_F00477	14	■ 0.867	$\mathcal{P}_{\text{world}}$	$\mathcal{P}_{\text{world}}$	parameterization of the probability density $\mathcal{P}_{\text{world}}$. Using this parameterization,
penev_B_F00478	14	■ 1.000	$T=1038$	$T = 1038$	We selected from the FERET database $T = 1038$ examples without glasses
penev_B_F00479	14	■ 1.000	64×60	64×60	that line. Finally, the examples in the data set were cropped through a 64×60
penev_B_F00480	14	■ 0.973	$r(\theta, \ell)$	$r(\theta, \ell)$	ordinates $r(\theta, \ell)$ with 512 samples spanning a full revolution in the θ direction
penev_B_F00481	14	■ 1.000	ℓ	ℓ	and 256 samples in the ℓ direction. Anthropological landmarks on the surfaces
penev_B_F00482	14	■ 0.823	$\theta_{\text{fixed}}, \ell_{\text{fixed}}$	$\theta_{\text{fixed}}, \ell_{\text{fixed}}$	The examples were aligned by us, so that the fixed point $(\theta_{\text{fixed}}, \ell_{\text{fixed}})$
penev_B_F00483	14	■ 0.998	3×3	3×3	smoothed with a 3×3 Gaussian filter and under-sampled twice to produce a
penev_B_F00484	14	■ 0.981	256×128	256×128	256×128 representation. All examples in the <i>Mini Survey study</i> database were
penev_B_F00485	14	■ 1.000	$T=348$	$T = 348$	chosen to produce a $T = 348$ data set.
penev_B_F00486	14	■ 0.999	128×64	128×64	Finally, the examples in the data set were cropped through a 128×64
penev_B_F00487	14	■ 1.000	96×90	96×90	cropped through a 96×90 window centered horizontally about the fixed point
penev_B_F00488	14	■ 1.000	96×120	96×120	cropped through a 96×120 window centered horizontally about the fixed point
penev_B_F00489	14	■ 1.000	$\left\{s_n = 1.05^{n/2}\right\}_{n=-4}^{+4}$	$\left\{s_n = 1.05^{n/2}\right\}_{n=-4}^{+4}$	ing parameter $\{s_n = 1.05^{n/2}\}_{n=-4}^{+4}$, i.e., the up and down deviations from the
penev_B_F00490	14	■ 1.000	$\approx \{10\%, 7.5\%, 5\%, 2.5\%, 0\%\}$	$\approx \{10\%, 7.5\%, 5\%, 2.5\%, 0\%\}$	hand-determined scale were $\approx \{10\%, 7.5\%, 5\%, 2.5\%, 0\%\}$, resulting in dimen-
penev_B_F00491	14	■ 0.982	$T=10944, V=3840$	$T = 10944, V = 3840$	sionalities ($T = 10944, V = 3840$).
penev_B_F00492	14	■ 0.983	$\left\{s_m = 1.025^{m/3}\right\}_{m \in \{-2, -1, 1, 2\}}$	$\left\{s_m = 1.025^{m/3}\right\}_{m \in \{-2, -1, 1, 2\}}$	additional scalings $\{s_m = 1.025^{m/3}\}_{m \in \{-2, -1, 1, 2\}}$, resulting in deviations of $\approx$
penev_B_F00493	14	■ 0.983	$\approx$	$\approx$	additional scalings $\{s_m = 1.025^{m/3}\}_{m \in \{-2, -1, 1, 2\}}$, resulting in deviations of $\approx$
penev_B_F00494	14	■ 0.651	$V=96 \times 120=11,520$	$V = 96 \times 120 = 11,520$	Three croppings were prepared—full with $V = 96 \times 120 = 11,520$, b3q
penev_B_F00495	14	■ 0.992	$V=96 \times 90=8,640$	$V = 96 \times 90 = 8,640$	with $V = 96 \times 90 = 8,640$ which is the same as that for Ensemble 3, and with
penev_B_F00496	14	■ 0.881	$V=64 \times 60=3,840$	$V = 64 \times 60 = 3,840$	third $V = 64 \times 60 = 3,840$ is the same as Ensemble 1.
penev_B_F00497	14	■ 0.992	$10 \times$	$10 \times$	The image was converted to grayscale, oversampled $10 \times$, and then under-