

voloshin-hypergraph: equation evidence

Display equations

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0124	159	■ 0.310	$a_{ij} = \begin{cases} 1 & \text{if } \mathcal{D}(x_i) \cap \mathcal{D}(x_j) \neq \emptyset \\ 0 & \text{otherwise.} \end{cases}$	$a_{ij} = \begin{cases} 1 & \text{if } \mathcal{D}(x_i) \cap \mathcal{D}(x_j) \neq \emptyset \\ 0 & \text{otherwise.} \end{cases}$	$a_{ij} = \begin{cases} 1 & \text{if } \mathcal{D}(x_i) \cap \mathcal{D}(x_j) \neq \emptyset, \\ 0 & \text{otherwise.} \end{cases}$
voloshin-hypergraph_ EQ0133	170	■ 0.340	$(\mathcal{H})_2 = (X, \mathcal{E}) \text{ where } \{x_i, x_j\} \in \mathcal{E} \Leftrightarrow \mathcal{D}(x_i) \cap \mathcal{D}(x_j) \neq \emptyset.$	$(\mathcal{H})_2 = (X, \mathcal{E}) \text{ where } \{x_i, x_j\} \in \mathcal{E} \Leftrightarrow \mathcal{D}(x_i) \cap \mathcal{D}(x_j) \neq \emptyset.$	$(\mathcal{H})_2 = (X, \mathcal{E}) \text{ where } \{x_i, x_j\} \in \mathcal{E} \Leftrightarrow \mathcal{D}(x_i) \cap \mathcal{D}(x_j) \neq \emptyset.$
voloshin-hypergraph_ EQ0141 (8.1)	185	■ 0.473	$\tau(\mathcal{H}) = \nu(\mathcal{H}).$	$\tau(\mathcal{H}) = \nu(\mathcal{H}).$	$\tau(\mathcal{H}) = \nu(\mathcal{H}).$
voloshin-hypergraph_ EQ0197	220	■ 0.500	$\chi(\mathcal{H}) = \bar{\chi}(\mathcal{H}) = 0, \quad R(\mathcal{H}) = (0, 0, \dots, 0), \quad P(\mathcal{H}, \lambda) = 0.$	$\chi(\mathcal{H}) = \bar{\chi}(\mathcal{H}) = 0, \quad R(\mathcal{H}) = (0, 0, \dots, 0), \quad P(\mathcal{H}, \lambda) = 0.$	$\chi(\mathcal{H}) = \bar{\chi}(\mathcal{H}) = 0, \quad R(\mathcal{H}) = (0, 0, \dots, 0), \quad P(\mathcal{H}, \lambda) = 0.$
voloshin-hypergraph_ EQ0244	258	■ 0.530	$\bar{\chi}(\mathcal{H}) = \alpha_C(\mathcal{H}) = X - \tau(\mathcal{H}_C) = X - \nu(\mathcal{H}_C).$	$\bar{\chi}(\mathcal{H}) = \alpha_C(\mathcal{H}) = X - \tau(\mathcal{H}_C) = X - \nu(\mathcal{H}_C).$	$\bar{\chi}(\mathcal{H}) = \alpha_C(\mathcal{H}) = X - \tau(\mathcal{H}_C) = X - \nu(\mathcal{H}_C).$
voloshin-hypergraph_ EQ0207	230	■ 0.531	$P(\mathcal{H}, \lambda) = P(\mathcal{H}_1, \lambda), \quad R(\mathcal{H}) = R(\mathcal{H}_1), \text{ where}$	$P(\mathcal{H}, \lambda) = P(\mathcal{H}_1, \lambda), \quad R(\mathcal{H}) = R(\mathcal{H}_1), \text{ where}$	$P(\mathcal{H}, \lambda) = P(\mathcal{H}_1, \lambda), \quad R(\mathcal{H}) = R(\mathcal{H}_1), \text{ where}$
voloshin-hypergraph_ EQ0233	253	■ 0.543	$\alpha_c(\mathcal{H}) - \bar{\chi}(\mathcal{H}) > k.$	$\alpha_c(\mathcal{H}) - \bar{\chi}(\mathcal{H}) > k.$	$\alpha_c(\mathcal{H}) - \bar{\chi}(\mathcal{H}) > k.$
voloshin-hypergraph_ EQ0130 (7.3)	168	■ 0.556	$\tau(\mathcal{H}) = \nu(\mathcal{H}).$	$\tau(\mathcal{H}) = \nu(\mathcal{H}).$	$\tau(\mathcal{H}) = \nu(\mathcal{H}).$
voloshin-hypergraph_ EQ0190	216	■ 0.566	$2, 3, 1, 4, 5,$	$2, 3, 1, 4, 5,$	$2, 3, 1, 4, 5,$
voloshin-hypergraph_ EQ0029	85	■ 0.576	$n - m + f = 2.$	$n - m + f = 2.$	$n - m + f = 2.$
voloshin-hypergraph_ EQ0051	107	■ 0.608	$P(G \cdot e, \lambda) = \sum_{i=0}^{n-1} (-1)^i b_i \lambda^{n-1-i}.$	$P(G \cdot e, \lambda) = \sum_{i=0}^{n-1} (-1)^i b_i \lambda^{n-1-i}.$	$P(G \cdot e, \lambda) = \sum_{i=0}^{n-1} (-1)^i b_i \lambda^{n-1-i}.$
voloshin-hypergraph_ EQ0120	156	■ 0.613	$\begin{gathered} \mathcal{H} = (X, \mathcal{D}), \quad X = \{1, 2, 3, 4, 5, 6\}, \quad \mathcal{D} = \{D_1, D_2, D_3, D_4, D_5\}, \\ D_1 = \{1\}, \quad D_2 = \{1, 2\}, \quad D_3 = \{1, 2, 4\}, \\ D_4 = \{2, 3, 5\}, \quad D_5 = \{3, 4, 5\}. \end{gathered}$	$\mathcal{H} = (X, \mathcal{D}), \quad X = \{1, 2, 3, 4, 5, 6\}, \quad \mathcal{D} = \{D_1, D_2, D_3, D_4, D_5\}, \\ D_1 = \{1\}, \quad D_2 = \{1, 2\}, \quad D_3 = \{1, 2, 4\}, \\ D_4 = \{2, 3, 5\}, \quad D_5 = \{3, 4, 5\}.$	$\mathcal{H} = (X, \mathcal{D}), \quad X = \{1, 2, 3, 4, 5, 6\}, \quad \mathcal{D} = \{D_1, D_2, D_3, D_4, D_5\}, \\ D_1 = \{1\}, \quad D_2 = \{1, 2\}, \quad D_3 = \{1, 2, 4\}, \\ D_4 = \{2, 3, 5\}, \quad D_5 = \{3, 4, 5\}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0056	109	■ 0.653	$\begin{gathered} P(G, \lambda) = \frac{P(\text{left}(G_1), \lambda) P(\text{left}(G_2), \lambda)}{\lambda(2)} = \\ \frac{[\lambda(\lambda-1)(\lambda^2-3\lambda+3)] [\lambda(\lambda-1)(\lambda-2)]}{\lambda(\lambda-1)} = \\ \lambda(\lambda-1)(\lambda-2)(\lambda^2-3\lambda+3). \end{gathered}$	$P(G, \lambda) = \frac{P(G_1, \lambda) P(G_2, \lambda)}{\lambda(2)} = \\ \frac{[\lambda(\lambda-1)(\lambda^2-3\lambda+3)] [\lambda(\lambda-1)(\lambda-2)]}{\lambda(\lambda-1)} = \\ \lambda(\lambda-1)(\lambda-2)(\lambda^2-3\lambda+3)$	$P(G, \lambda) = \frac{P(G_1, \lambda) P(G_2, \lambda)}{\lambda(2)} = \\ \frac{[\lambda(\lambda-1)(\lambda^2-3\lambda+3)] [\lambda(\lambda-1)(\lambda-2)]}{\lambda(\lambda-1)} = \\ \lambda(\lambda-1)(\lambda-2)(\lambda^2-3\lambda+3).$
voloshin-hypergraph_ EQ0125	160	■ 0.655	$\left \mathcal{D}(\text{left}(K_n^r)) \right = \frac{\binom{n}{r}}{r!} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)(n-2) \cdots (n-r+1)}{r!}.$	$ \mathcal{D}(K_n^r) = \binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)(n-2) \cdots (n-r+1)}{r!}.$	$ \mathcal{D}(K_n^r) = \binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)(n-2) \cdots (n-r+1)}{r!}.$
voloshin-hypergraph_ EQ0189	216	■ 0.679	$5, 4, 1, 3, 2.$	$5, 4, 1, 3, 2.$	$5, 4, 1, 3, 2.$
voloshin-hypergraph_ EQ0198 (10.3)	221	■ 0.699	$1 \leq \chi(\text{left}(\mathcal{H}_{\text{left}(\mathcal{H})}) \leq \chi(\mathcal{H}) \leq \bar{\chi}(\mathcal{H}) \leq \bar{\chi}(\mathcal{H}_C) \leq n.$	$1 \leq \chi(\mathcal{H}_{\mathcal{D}}) \leq \chi(\mathcal{H}) \leq \bar{\chi}(\mathcal{H}) \leq \bar{\chi}(\mathcal{H}_C) \leq n.$	$1 \leq \chi(\mathcal{H}_{\mathcal{D}}) \leq \chi(\mathcal{H}) \leq \bar{\chi}(\mathcal{H}) \leq \bar{\chi}(\mathcal{H}_C) \leq n.$
voloshin-hypergraph_ EQ0160	195	■ 0.709	$\wedge(\mathcal{H}) = \min_{T} \wedge(\mathcal{H}, T) = \sum_{D \in \mathcal{D}} (D - 1) - w(\mathcal{H}).$	$\wedge(\mathcal{H}) = \min_T \wedge(\mathcal{H}, T) = \sum_{D \in \mathcal{D}} (D - 1) - w(\mathcal{H}).$	$\wedge(\mathcal{H}) = \min_T \wedge(\mathcal{H}, T) = \sum_{D \in \mathcal{D}} (D - 1) - w(\mathcal{H}).$
voloshin-hypergraph_ EQ0040	102	■ 0.712	$\begin{gathered} P(G, \lambda) = P(\text{left}(K_3), \lambda) + P(\text{left}(K_2), \lambda) = \lambda^3 + \lambda^2 = \\ \lambda(\lambda-1)(\lambda-2) + \lambda(\lambda-1) = \lambda(\lambda-1)^2. \end{gathered}$	$P(G, \lambda) = P(K_3, \lambda) + P(K_2, \lambda) = \lambda^3 + \lambda^2 = \\ \lambda(\lambda-1)(\lambda-2) + \lambda(\lambda-1) = \lambda(\lambda-1)^2$	$P(G, \lambda) = P(K_3, \lambda) + P(K_2, \lambda) = \lambda^3 + \lambda^2 = \\ \lambda(\lambda-1)(\lambda-2) + \lambda(\lambda-1) = \lambda(\lambda-1)^2.$
voloshin-hypergraph_ EQ0216	239	■ 0.715	$\mathcal{K}(n, l, m) = (X, \mathcal{C}, \mathcal{D}) = \left(X, \binom{X}{l}, \binom{X}{m} \right).$	$\mathcal{K}(n, l, m) = (X, \mathcal{C}, \mathcal{D}) = \left(X, \binom{X}{l}, \binom{X}{m} \right).$	$\mathcal{K}(n, l, m) = (X, \mathcal{C}, \mathcal{D}) = \left(X, \binom{X}{l}, \binom{X}{m} \right).$
voloshin-hypergraph_ EQ0266	287	■ 0.722	$I(\text{left}(G_1)) = \left(\begin{array}{llll} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{array} \right)$	$I(G_1) = \left(\begin{array}{llll} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{array} \right)$	$I(G_1) = \left(\begin{array}{llll} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{array} \right)$
voloshin-hypergraph_ EQ0009	33	■ 0.738	$A(G) = \left(\begin{array}{llll} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 2 & 0 & 1 & 0 \end{array} \right);$	$A(G) = \left(\begin{array}{llll} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 2 & 0 & 1 & 0 \end{array} \right);$	$A(G) = \left(\begin{array}{llll} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 2 & 0 & 1 & 0 \end{array} \right);$
voloshin-hypergraph_ EQ0137 (7.4)	173	■ 0.743	$\left(D_1 \cap D_2 \right) \cup \left(D_1 \cap D_3 \right) \cup \left(D_2 \cap D_3 \right) \subseteq D.$	$(D_1 \cap D_2) \cup (D_1 \cap D_3) \cup (D_2 \cap D_3) \subseteq D.$	$(D_1 \cap D_2) \cup (D_1 \cap D_3) \cup (D_2 \cap D_3) \subseteq D.$
voloshin-hypergraph_ EQ0119	156	■ 0.755	$\begin{gathered} \mathcal{Z} = \{X_1, X_2, \dots, X_n\}, \\ X_i = \{d_j : x_i \in D_j \text{ in } \mathcal{H}\}. \end{gathered}$	$\mathcal{Z} = \{X_1, X_2, \dots, X_n\}, \\ X_i = \{d_j : x_i \in D_j \text{ in } \mathcal{H}\}.$	$\mathcal{Z} = \{X_1, X_2, \dots, X_n\}, \\ X_i = \{d_j : x_i \in D_j \text{ in } \mathcal{H}\}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0052	107	■ 0.759	$\begin{gathered} P(G, \lambda) = P(G - e, \lambda) - P(G \cdot e, \lambda) \\ P(G - e, \lambda) = \lambda^n - (m-1)\lambda^{n-1} + a_2\lambda^{n-2} - \dots + (-1)^i a_i \lambda^{n-i} \dots \\ P(G \cdot e, \lambda) = \lambda^{n-1} - b_1\lambda^{n-2} + b_2\lambda^{n-3} - \dots + (-1)^i b_{i-1} \lambda^{n-i} \dots \\ P(G, \lambda) = \lambda^n - m(G)\lambda^{n-1} + (a_2 + b_1)\lambda^{n-2} \dots (-1)^i (a_i + b_{i-1}) \lambda^{n-i} \dots \end{gathered}$	$P(G, \lambda) = P(G - e, \lambda) - P(G \cdot e, \lambda) \\ = \lambda^n - (m-1)\lambda^{n-1} + a_2\lambda^{n-2} - \dots + (-1)^i a_i \lambda^{n-i} \dots \\ - (\lambda^{n-1} - b_1\lambda^{n-2} + b_2\lambda^{n-3} - \dots + (-1)^i b_{i-1} \lambda^{n-i} \dots) \\ = \lambda^n - m(G)\lambda^{n-1} + (a_2 + b_1)\lambda^{n-2} \dots (-1)^i (a_i + b_{i-1}) \lambda^{n-i} \dots$	$P(G, \lambda) = P(G - e, \lambda) - P(G \cdot e, \lambda) \\ = \lambda^n - (m-1)\lambda^{n-1} + a_2\lambda^{n-2} - \dots + (-1)^i a_i \lambda^{n-i} \dots \\ - (\lambda^{n-1} - b_1\lambda^{n-2} + b_2\lambda^{n-3} - \dots + (-1)^i b_{i-1} \lambda^{n-i} \dots) \\ = \lambda^n - m(G)\lambda^{n-1} + (a_2 + b_1)\lambda^{n-2} \dots (-1)^i (a_i + b_{i-1}) \lambda^{n-i} \dots$
voloshin-hypergraph_ EQ0064	111	■ 0.763	$\begin{gathered} P(G_1, \lambda) = (\lambda - k_1) P(G_2, \lambda) = (\lambda - k_1)(\lambda - k_2) P(G_3, \lambda) \dots = \\ (\lambda - k_1)(\lambda - k_2)(\lambda - k_3) \dots (\lambda - k_n). \end{gathered}$	$P(G_1, \lambda) = (\lambda - k_1) P(G_2, \lambda) = (\lambda - k_1)(\lambda - k_2) P(G_3, \lambda) \dots = \\ (\lambda - k_1)(\lambda - k_2)(\lambda - k_3) \dots (\lambda - k_n)$	$P(G_1, \lambda) = (\lambda - k_1) P(G_2, \lambda) = (\lambda - k_1)(\lambda - k_2) P(G_3, \lambda) \dots = \\ (\lambda - k_1)(\lambda - k_2)(\lambda - k_3) \dots (\lambda - k_n).$
voloshin-hypergraph_ EQ0206	230	■ 0.771	$\begin{gathered} \mathcal{H}_1 = (X, \mathcal{C}, \mathcal{D}_1), \mathcal{D}_1 = \mathcal{D} \cup \{x_k, x_l\}, \\ \mathcal{H}_2 = (X, \mathcal{C}_1, \mathcal{D}), \mathcal{C}_1 = \mathcal{C} \cup \{x_k, x_l\} \text{ (splitting)}. \end{gathered}$	$\mathcal{H}_1 = (X, \mathcal{C}, \mathcal{D}_1), \mathcal{D}_1 = \mathcal{D} \cup \{x_k, x_l\}, \\ \mathcal{H}_2 = (X, \mathcal{C}_1, \mathcal{D}), \mathcal{C}_1 = \mathcal{C} \cup \{x_k, x_l\} \text{ (splitting)}.$	$\mathcal{H}_1 = (X, \mathcal{C}, \mathcal{D}_1), \mathcal{D}_1 = \mathcal{D} \cup \{x_k, x_l\}, \\ \mathcal{H}_2 = (X, \mathcal{C}_1, \mathcal{D}), \mathcal{C}_1 = \mathcal{C} \cup \{x_k, x_l\} \text{ (splitting)}.$
voloshin-hypergraph_ EQ0007	30	■ 0.775	$\begin{aligned} J(G) = & \left\{ \left\{ x_1, x_2 \right\}, \left\{ x_2, x_3 \right\}, \left\{ x_3, x_4 \right\}, \right. \\ & \left. \left\{ x_4, x_5 \right\}, \left\{ x_1, x_5 \right\}, \left\{ x_2, x_5 \right\}, \left\{ x_2, x_4 \right\} \right\}. \end{aligned}$	$J(G) = \left\{ \left\{ x_1, x_2 \right\}, \left\{ x_2, x_3 \right\}, \left\{ x_3, x_4 \right\}, \right. \\ \left. \left\{ x_4, x_5 \right\}, \left\{ x_1, x_5 \right\}, \left\{ x_2, x_5 \right\}, \left\{ x_2, x_4 \right\} \right\}.$	$J(G) = \left\{ \left\{ x_1, x_2 \right\}, \left\{ x_2, x_3 \right\}, \left\{ x_3, x_4 \right\}, \right. \\ \left. \left\{ x_4, x_5 \right\}, \left\{ x_1, x_5 \right\}, \left\{ x_2, x_5 \right\}, \left\{ x_2, x_4 \right\} \right\}.$
voloshin-hypergraph_ EQ0143 (8.3)	190	■ 0.779	$\bigwedge (\mathcal{H}, T) = \sum_{D \in \mathcal{D}} (D - 1) - w(T) + l(\mathcal{H}, T).$	$\bigwedge (\mathcal{H}, T) = \sum_{D \in \mathcal{D}} (D - 1) - w(T) + l(\mathcal{H}, T).$	$\bigwedge (\mathcal{H}, T) = \sum_{D \in \mathcal{D}} (D - 1) - w(T) + l(\mathcal{H}, T).$
voloshin-hypergraph_ EQ0245	258	■ 0.790	$\nu(\mathcal{H}_C) = \tau(\mathcal{H}_C).$	$\nu(\mathcal{H}_C) = \tau(\mathcal{H}_C).$	$\nu(\mathcal{H}_C) = \tau(\mathcal{H}_C).$
voloshin-hypergraph_ EQ0161	195	■ 0.792	$\bigwedge (\mathcal{H}) = \bigwedge \left(\mathcal{H}_1 \right).$	$\bigwedge (\mathcal{H}) = \bigwedge (\mathcal{H}_1).$	$\bigwedge (\mathcal{H}) = \bigwedge (\mathcal{H}_1).$
voloshin-hypergraph_ EQ0118	155	■ 0.808	$(i, j) \text{-entry} = \begin{cases} 1 & \text{if } x_i \in D_j, \\ 0 & \text{if } x_i \notin D_j. \end{cases}$	$(i, j)\text{-entry} = \begin{cases} 1 & \text{if } x_i \in D_j, \\ 0 & \text{if } x_i \notin D_j. \end{cases}$	$(i, j)\text{-entry} = \begin{cases} 1 & \text{if } x_i \in D_j, \\ 0 & \text{if } x_i \notin D_j. \end{cases}$
voloshin-hypergraph_ EQ0138	173	■ 0.808	$(D_1 \cap D_2) \cup (D_1 \cap D_3) \cup (D_2 \cap D_3) \subseteq D.$	$(D_1 \cap D_2) \cup (D_1 \cap D_3) \cup (D_2 \cap D_3) \subseteq D.$	$(D_1 \cap D_2) \cup (D_1 \cap D_3) \cup (D_2 \cap D_3) \subseteq D.$
voloshin-hypergraph_ EQ0162	195	■ 0.828	$\bigwedge (\mathcal{H}) = \bigwedge (\mathcal{H}, T) = \bigwedge \left(\mathcal{H}_1, T_1 \right).$	$\bigwedge (\mathcal{H}) = \bigwedge (\mathcal{H}, T) = \bigwedge (\mathcal{H}_1, T_1).$	$\bigwedge (\mathcal{H}) = \bigwedge (\mathcal{H}, T) = \bigwedge (\mathcal{H}_1, T_1).$
voloshin-hypergraph_ EQ0030	85	■ 0.840	$m(G) \leq 3n(G) - 6.$	$m(G) \leq 3n(G) - 6.$	$m(G) \leq 3n(G) - 6.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0199	221	■ 0.852	$1 \leq \chi(\mathcal{H}) = 2 \leq \bar{\chi}(\mathcal{H}) = 4 \leq \bar{\chi}(\mathcal{H}_C) = 4 \leq n = 5$.	$1 \leq \chi(\mathcal{H}_D) = 2 \leq \chi(\mathcal{H}) = 2 \leq \bar{\chi}(\mathcal{H}) = 4 \leq \bar{\chi}(\mathcal{H}_C) = 4 \leq n = 5$.	$1 \leq \chi(\mathcal{H}_D) = 2 \leq \chi(\mathcal{H}) = 2 \leq \bar{\chi}(\mathcal{H}) = 4 \leq \bar{\chi}(\mathcal{H}_C) = 4 \leq n = 5$.
voloshin-hypergraph_ EQ0087	116	■ 0.863	$\begin{gathered} P(G, \lambda) = \lambda P(G-x, \lambda-1) = \\ \frac{P(G-x, \lambda)}{P(G-x, \lambda-1)} \lambda P(G-x, \lambda-1) = \\ P(G-x, \lambda) W(G_{N(x)}, \lambda) \end{gathered}$	$P(G, \lambda) = \lambda P(G-x, \lambda-1) = \frac{P(G-x, \lambda)}{P(G-x, \lambda-1)} \lambda P(G-x, \lambda-1) = \\ P(G-x, \lambda) W(G_{N(x)}, \lambda)$	$P(G, \lambda) = \lambda P(G-x, \lambda-1) = \frac{P(G-x, \lambda)}{P(G-x, \lambda-1)} \lambda P(G-x, \lambda-1) = \\ P(G-x, \lambda) W(G_{N(x)}, \lambda)$
voloshin-hypergraph_ EQ0153	193	■ 0.873	$\wedge(\mathcal{H}, T) = \wedge(\mathcal{H}_1, T_1)$ if and only if $ \mathcal{D}(x) = \mathcal{D}_T(x) + m_2$.	$\wedge(\mathcal{H}, T) = \wedge(\mathcal{H}_1, T_1)$ if and only if $ \mathcal{D}(x) = \mathcal{D}_T(x) + m_2$.	$\wedge(\mathcal{H}, T) = \wedge(\mathcal{H}_1, T_1)$ if and only if $ \mathcal{D}(x) = \mathcal{D}_T(x) + m_2$.
voloshin-hypergraph_ EQ0214	233	■ 0.876	$\begin{gathered} \chi(\mathcal{H}) = \max \left\{ \chi(\mathcal{H}_1), \chi(\mathcal{H}_2), \dots, \chi(\mathcal{H}_k) \right\}, \\ \bar{\chi}(\mathcal{H}) = \bar{\chi}(\mathcal{H}_1) + \bar{\chi}(\mathcal{H}_2) + \dots + \bar{\chi}(\mathcal{H}_k), \\ P(\mathcal{H}, \lambda) = P(\mathcal{H}_1, \lambda) P(\mathcal{H}_2, \lambda) \dots P(\mathcal{H}_k, \lambda). \end{gathered}$	$\chi(\mathcal{H}) = \max \left\{ \chi(\mathcal{H}_1), \chi(\mathcal{H}_2), \dots, \chi(\mathcal{H}_k) \right\}, \\ \bar{\chi}(\mathcal{H}) = \bar{\chi}(\mathcal{H}_1) + \bar{\chi}(\mathcal{H}_2) + \dots + \bar{\chi}(\mathcal{H}_k), \\ P(\mathcal{H}, \lambda) = P(\mathcal{H}_1, \lambda) P(\mathcal{H}_2, \lambda) \dots P(\mathcal{H}_k, \lambda).$	$\chi(\mathcal{H}) = \max \left\{ \chi(\mathcal{H}_1), \chi(\mathcal{H}_2), \dots, \chi(\mathcal{H}_k) \right\}, \\ \bar{\chi}(\mathcal{H}) = \bar{\chi}(\mathcal{H}_1) + \bar{\chi}(\mathcal{H}_2) + \dots + \bar{\chi}(\mathcal{H}_k), \\ P(\mathcal{H}, \lambda) = P(\mathcal{H}_1, \lambda) P(\mathcal{H}_2, \lambda) \dots P(\mathcal{H}_k, \lambda).$
voloshin-hypergraph_ EQ0099	129	■ 0.890	$\chi'(G) = \Delta(G)$, or $\chi'(G) = \Delta(G) + 1$.	$\chi'(G) = \Delta(G)$, or $\chi'(G) = \Delta(G) + 1$.	$\chi'(G) = \Delta(G)$, or $\chi'(G) = \Delta(G) + 1$.
voloshin-hypergraph_ EQ0235 (10.5)	253	■ 0.901	$\bar{\chi}(\mathcal{H}') = \alpha_c(\mathcal{H}')$.	$\bar{\chi}(\mathcal{H}') = \alpha_c(\mathcal{H}')$.	$\bar{\chi}(\mathcal{H}') = \alpha_c(\mathcal{H}')$.
voloshin-hypergraph_ EQ0201	224	■ 0.913	$o(x, \mathcal{H}) = \mathcal{C}(x) - b(x, \mathcal{H}) \geq 0$	$o(x, \mathcal{H}) = \mathcal{C}(x) - b(x, \mathcal{H}) \geq 0$	$o(x, \mathcal{H}) = \mathcal{C}(x) - b(x, \mathcal{H}) \geq 0$
voloshin-hypergraph_ EQ0232 (10.4)	252	■ 0.932	$\bar{\chi}(\mathcal{H}) \leq \alpha_c(\mathcal{H})$.	$\bar{\chi}(\mathcal{H}) \leq \alpha_c(\mathcal{H})$.	$\bar{\chi}(\mathcal{H}) \leq \alpha_c(\mathcal{H})$.
voloshin-hypergraph_ EQ0259	273	■ 0.935	$P(\mathcal{H}, \lambda) = \sum_{i \geq 1} f_i(G^*) \lambda(\lambda-1)^i$.	$P(\mathcal{H}, \lambda) = \sum_{i \geq 1} f_i(G^*) \lambda(\lambda-1)^i$.	$P(\mathcal{H}, \lambda) = \sum_{i \geq 1} f_i(G^*) \lambda(\lambda-1)^i$.
voloshin-hypergraph_ EQ0226	248	■ 0.938	$ C \geq n - \chi$.	$ C \geq n - \chi$.	$ C \geq n - \chi$.
voloshin-hypergraph_ EQ0231	250	■ 0.941	$\sigma = (x_5, x_3, x_4, x_2, x_1)$.	$\sigma = (x_5, x_3, x_4, x_2, x_1)$.	$\sigma = (x_5, x_3, x_4, x_2, x_1)$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0265	286	■ 0.955	$A\left(G_{1}\right)=\left(\begin{array}{cccc} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{array}\right)$		
voloshin-hypergraph_ EQ0136	171	■ 0.957	$\nu(\mathcal{H})=\alpha(L(\mathcal{H})) \text{ and } \nu(\mathcal{H}^*)=\alpha((\mathcal{H})_2).$		
voloshin-hypergraph_ EQ0152 (8.6)	193	■ 0.970	$\wedge(\mathcal{H},T)=\wedge(\mathcal{H}_1,T_1)+ \mathcal{D}(x) - \mathcal{D}_T(x) -m_2.$		
voloshin-hypergraph_ EQ0218	240	■ 0.971	$(l-1)(m-1)\leq n.$		
voloshin-hypergraph_ EQ0243	257	■ 0.973	$\bigcap_{C\in\mathcal{C}_1}C\neq\emptyset.$		
voloshin-hypergraph_ EQ0242	256	■ 0.975	$\bar{\chi}(\mathcal{H}^r)=2r-4<2r-3=\alpha_{\mathcal{C}}(\mathcal{H}^r).$		
voloshin-hypergraph_ EQ0072	114	■ 0.983	$P(G',\lambda)=\lambda^{s'_0}(\lambda-1)^{s'_1}\dots(\lambda-\chi'+1)^{s'_{\chi'-1}},$		
voloshin-hypergraph_ EQ0015	48	■ 0.988	$\tau(G)\geq v(G).$		
voloshin-hypergraph_ EQ0096	118	■ 0.988	$P(G_1,\lambda)W(G_{N(4)},\lambda)=\lambda(\lambda-1)^2(\lambda-2)^3(\lambda-3)\frac{(\lambda-2)^2}{\lambda-1}=\\ \lambda(\lambda-1)(\lambda-2)^5(\lambda-3)=P(G,\lambda)$		
voloshin-hypergraph_ EQ0228	248	■ 0.988	$ X_1 -1+ X_2 -1+\dots+ X_{\chi} -1= X -\chi.$		
voloshin-hypergraph_ EQ0107	136	■ 0.989	$\bar{\chi}'(G)=c+m-n+p=1+5-4+1=3,$		
voloshin-hypergraph_ EQ0186	215	■ 0.990	$M(\mathcal{H})=m(y,\mathcal{H}_Y)\leq m(x_k,\mathcal{H}_k)\leq t.$		
voloshin-hypergraph_ EQ0254	266	■ 0.990	$R(\mathcal{H}')=(0,r_2(\mathcal{H}'),0,\dots,0,r_{\chi},\dots,r_{\bar{\chi}},0,\dots,0).$		
voloshin-hypergraph_ EQ0054	108	■ 0.991	$P(G,\lambda)=\frac{P(G_{X_1\cup S},\lambda)}{\lambda^{(k)}}\frac{P(G_{X_2\cup S},\lambda)}{\lambda^{(k)}}\lambda^{(k)}=\\ \frac{P(G_{X_1\cup S},\lambda)P(G_{X_2\cup S},\lambda)}{\lambda^{(k)}}.$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0132	170	■ 0.993	$L(\mathcal{H})=(\mathcal{D},\mathcal{E}),\text{ where } \left(D_i,D_j\right)\in \mathcal{E}\Leftrightarrow D_i\cap D_j\neq \emptyset$.	$L(\mathcal{H})=(\mathcal{D},\mathcal{E}),\text{ where } (D_i,D_j)\in \mathcal{E}\Leftrightarrow D_i\cap D_j\neq \emptyset$.	$L(\mathcal{H})=(\mathcal{D},\mathcal{E}),\text{ where } (D_i,D_j)\in \mathcal{E}\Leftrightarrow D_i\cap D_j\neq \emptyset$.
voloshin-hypergraph_ EQ0150	192	■ 0.994	$\wedge(\mathcal{H},T)=(2-1)+(1-1)+(1-1)+1=2$.	$\wedge(\mathcal{H},T)=(2-1)+(1-1)+(1-1)+1=2$.	$\wedge(\mathcal{H},T)=(2-1)+(1-1)+(1-1)+1=2$.
voloshin-hypergraph_ EQ0065 (5.11)	112	■ 0.994	$P(G,\lambda)=\lambda^{\lambda^{\{0\}}}(\lambda-1)^{\lambda^{\{1\}}}(\lambda-2)^{\lambda^{\{2\}}}\cdots(\lambda-\chi(G)+1)^{\lambda^{\{\chi-1\}}}$	$P(G,\lambda)=\lambda^{S_0}(\lambda-1)^{S_1}(\lambda-2)^{S_2}\cdots(\lambda-\chi(G)+1)^{S_{\chi-1}}$	$P(G,\lambda)=\lambda^{s_0}(\lambda-1)^{s_1}(\lambda-2)^{s_2}\cdots(\lambda-\chi(G)+1)^{s_{\chi-1}}$
voloshin-hypergraph_ EQ0223	243	■ 0.994	$P(\mathcal{H},\lambda)=\lambda(\lambda-1)(\lambda-2)\cdots(\lambda-\chi+1)=\lambda^{(\chi)}$.	$P(\mathcal{H},\lambda)=\lambda(\lambda-1)(\lambda-2)\cdots(\lambda-\chi+1)=\lambda^{(\chi)}$.	$P(\mathcal{H},\lambda)=\lambda(\lambda-1)(\lambda-2)\cdots(\lambda-\chi+1)=\lambda^{(\chi)}$.
voloshin-hypergraph_ EQ0182	212	■ 0.995	$\chi(\mathcal{H})\leq \wedge(\mathcal{H})+2$.	$\chi(\mathcal{H})\leq \wedge(\mathcal{H})+2$.	$\chi(\mathcal{H})\leq \wedge(\mathcal{H})+2$.
voloshin-hypergraph_ EQ0257	268	■ 0.996	$\begin{array}{lll} 123 & \& 341 & \& 561 \\ 124 & \& 342 & \& 562 \\ 125 & \& 345 & \& 563 \\ 126 & \& 346 & \& 564 \end{array}$	$\begin{array}{lll} 123 & 341 & 561 \\ 124 & 342 & 562 \\ 125 & 345 & 563 \\ 126 & 346 & 564 \end{array}$	$\begin{array}{lll} 123 & 341 & 561 \\ 124 & 342 & 562 \\ 125 & 345 & 563 \\ 126 & 346 & 564 \end{array}$
voloshin-hypergraph_ EQ0176	211	■ 0.996	$\left\lfloor\frac{ D }{\lambda}\right\rfloor\leq D\cap S_i \leq\left\lceil\frac{ D }{\lambda}\right\rceil$,	$\left\lfloor\frac{ D }{\lambda}\right\rfloor\leq D\cap S_i \leq\left\lceil\frac{ D }{\lambda}\right\rceil$,	$\left\lfloor\frac{ D }{\lambda}\right\rfloor\leq D\cap S_i \leq\left\lceil\frac{ D }{\lambda}\right\rceil$,
voloshin-hypergraph_ EQ0085	116	■ 0.996	$W(G,\lambda)=\frac{\lambda P(G,\lambda-1)}{P(G,\lambda)}$.	$W(G,\lambda)=\frac{\lambda P(G,\lambda-1)}{P(G,\lambda)}$.	$W(G,\lambda)=\frac{\lambda P(G,\lambda-1)}{P(G,\lambda)}$.
voloshin-hypergraph_ EQ0156	193	■ 0.996	$\wedge(\mathcal{H}-x,T-x)+ \mathcal{D}(x) - \mathcal{D}_T(x) -m_2=2+3-1-1=3=\wedge(\mathcal{H},T)$.	$\wedge(\mathcal{H}-x,T-x)+ \mathcal{D}(x) - \mathcal{D}_T(x) -m_2=2+3-1-1=3=\wedge(\mathcal{H},T)$.	$\wedge(\mathcal{H}-x,T-x)+ \mathcal{D}(x) - \mathcal{D}_T(x) -m_2=2+3-1-1=3=\wedge(\mathcal{H},T)$.
voloshin-hypergraph_ EQ0258	273	■ 0.996	$r_k(G)=\sum_{i\geq 1}S(i,k-1)f_i(G^*),1\leq k\leq n(G)$,	$r_k(G)=\sum_{i\geq 1}S(i,k-1)f_i(G^*),1\leq k\leq n(G)$,	$r_k(G)=\sum_{i\geq 1}S(i,k-1)f_i(G^*),1\leq k\leq n(G)$,
voloshin-hypergraph_ EQ0208	230	■ 0.996	$\mathcal{C}_1=\mathcal{C}-C_t$ (contraction).	$\mathcal{C}_1=\mathcal{C}-C_t$ (contraction).	$\mathcal{C}_1=\mathcal{C}-C_t$ (contraction).
voloshin-hypergraph_ EQ0246	259	■ 0.997	$\begin{aligned} &V_1=(\{1,2,3,4\},\{\{1,2,3\},\{1,3,4\}\})\text{ (monostar); }\\ &V_2=(\{1,2,3,4,5\},\{\{1,2,3\},\{1,4,5\}\})\text{ (monostar); }\\ &V_3=(\{1,2,3,4,5\},\{\{1,2,3\},\{1,3,4\},\{1,4,5\}\})\text{ (monostar); }\\ &V_4=(\{1,2,3,4,5\},\{\{1,2,3\},\{1,3,4\},\{1,4,5\},\{1,2,5\}\})\text{ (monostar); }\\ &V_5=(\{1,2,3,4,5\},\{\{1,2,3\},\{2,3,4\},\{3,4,5\},\{4,5,1\},\{5,1,2\}\})\text{ (cycloid }C_5^3);\\ &K_1=(\{1,2,3,4,5,6\},\{\{1,2,4\},\{2,3,5\},\{3,4,6\},\{4,5,1\},\{5,6,2\},\{6,1,3\},\{3,5\},\{2,4,6\}\})\text{ (Král's example). } \end{aligned}$	$\begin{aligned} &V_1=(\{1,2,3,4\},\{\{1,2,3\},\{1,3,4\},\{1,2,4\}\})\text{ (monostar); }\\ &V_2=(\{1,2,3,4,5\},\{\{1,2,3\},\{1,4,5\}\})\text{ (monostar); }\\ &V_3=(\{1,2,3,4,5\},\{\{1,2,3\},\{1,3,4\},\{1,4,5\}\})\text{ (monostar); }\\ &V_4=(\{1,2,3,4,5\},\{\{1,2,3\},\{1,3,4\},\{1,4,5\},\{1,2,5\}\})\text{ (monostar); }\\ &V_5=(\{1,2,3,4,5\},\{\{1,2,3\},\{2,3,4\},\{3,4,5\},\{4,5,1\},\{5,1,2\}\})\text{ (cycloid }C_5^3);\\ &K_1=(\{1,2,3,4,5,6\},\{\{1,2,4\},\{2,3,5\},\{3,4,6\},\{4,5,1\},\{5,6,2\},\{6,1,3\},\{3,5\},\{2,4,6\}\})\text{ (Král's example). } \end{aligned}$	$\begin{aligned} &V_1=(\{1,2,3,4\},\{\{1,2,3\},\{1,3,4\},\{1,2,4\}\})\text{ (monostar); }\\ &V_2=(\{1,2,3,4,5\},\{\{1,2,3\},\{1,4,5\}\})\text{ (monostar); }\\ &V_3=(\{1,2,3,4,5\},\{\{1,2,3\},\{1,3,4\},\{1,4,5\}\})\text{ (monostar); }\\ &V_4=(\{1,2,3,4,5\},\{\{1,2,3\},\{1,3,4\},\{1,4,5\},\{1,2,5\}\})\text{ (monostar); }\\ &V_5=(\{1,2,3,4,5\},\{\{1,2,3\},\{2,3,4\},\{3,4,5\},\{4,5,1\},\{5,1,2\}\})\text{ (cycloid }C_5^3);\\ &K_1=(\{1,2,3,4,5,6\},\{\{1,2,4\},\{2,3,5\},\{3,4,6\},\{4,5,1\},\{5,6,2\},\{6,1,3\},\{3,5\},\{2,4,6\}\})\text{ (Král's example). } \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0195 (10.2)	220	■ 0.997	$P(\mathcal{H}, \lambda) = \sum_{i=\chi(\mathcal{H})}^{\bar{\chi}(\mathcal{H})} r_i(\mathcal{H}) \lambda^{(i)}.$		
voloshin-hypergraph_ EQ0261	274	■ 0.997	$X=\{1,2,3,4\}, \mathcal{C}=\mathcal{D}=\{\{1,2,3\}, \{2,3,4\}, \{3,4,1\}, \{4,1,2\}\};$		
voloshin-hypergraph_ EQ0026	73	■ 0.997	$M(G)=\max_{X'\subseteq X} \min_{x\in G'} d(x).$		
voloshin-hypergraph_ EQ0018	63	■ 0.998	$ N_G(S) \geq S .$		
voloshin-hypergraph_ EQ0071	113	■ 0.998	$\lambda(\lambda-1)^{n-1} - [(\lambda-1)^{n-1} + (-1)^{n-1}(\lambda-1)] = (\lambda-1)^n + (-1)^n(\lambda-1).$		
voloshin-hypergraph_ EQ0154	193	■ 0.999	$\wedge(\mathcal{H}, T) = 2 + 2 + 1 + 0 + 0 - 3 + 1 = 3.$		
voloshin-hypergraph_ EQ0155	193	■ 0.999	$\wedge(\mathcal{H} - x, T - x) = 2 + 1 + 0 + 0 - 2 + 1 = 2.$		
voloshin-hypergraph_ EQ0185	215	■ 0.999	$m(y, \mathcal{H}_Y) = \min_z m(z, \mathcal{H}_Y) = M(\mathcal{H}).$		
voloshin-hypergraph_ EQ0203	227	■ 0.999	$o(y, \mathcal{H}') = \min_z o(z, \mathcal{H}') = O(\mathcal{H}) \geq t + 1.$		
voloshin-hypergraph_ EQ0025	64	■ 0.999	$\tau(G) = \mathcal{V}(G).$		
voloshin-hypergraph_ EQ0013	40	■ 0.999	$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ a & b & c & d & e & f \end{pmatrix}.$		
voloshin-hypergraph_ EQ0224	244	■ 0.999	$x_j \neq x_{j+1} \neq x_{j+2} \neq x_j \Rightarrow \{x_j, x_{j+1}, x_{j+2}\} \in \mathcal{C}, j = 0, 1, \dots, k - 2.$		
voloshin-hypergraph_ EQ0248	260	■ 0.999	1122, 1112, 1121, 1222, 2122, 1212, 1221.	1122, 1112, 1121, 1222, 2122, 1212, 1221.	1122, 1112, 1121, 1222, 2122, 1212, 1221.
voloshin-hypergraph_ EQ0179	211	■ 0.999	$\left\lfloor \frac{n}{\lambda} \right\rfloor \leq S_i \leq \left\lceil \frac{n}{\lambda} \right\rceil.$		
voloshin-hypergraph_ EQ0004	29	■ 1.000	$L(G) = \{\{x_2, x_5\}, \{x_1, x_3, x_4, x_5\}, \{x_2, x_4\}, \{x_2, x_3, x_5\}, \{x_1, x_2, x_4\}\}.$		
voloshin-hypergraph_ EQ0006	30	■ 1.000	$I(G) = \begin{pmatrix} 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 \end{pmatrix}.$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0019 (2.1)	63	■ 1.000	$ S < \mathcal{N}_G(S) $.	$ S < N_G(S) $.	$ S < N_G(S) $.
voloshin-hypergraph_ EQ0039 (5.3)	102	■ 1.000	$P(G, \lambda) = \sum_{i=\chi}^n r_i(G) \lambda^{\{i\}}$	$P(G, \lambda) = \sum_{i=\chi}^n r_i(G) \lambda^{(i)}$	$P(G, \lambda) = \sum_{i=\chi}^n r_i(G) \lambda^{(i)}$.
voloshin-hypergraph_ EQ0050	106	■ 1.000	$P(G-e, \lambda) = \sum_{i=0}^n (-1)^i a_i \lambda^{n-i}$	$P(G-e, \lambda) = \sum_{i=0}^n (-1)^i a_i \lambda^{n-i}$	$P(G-e, \lambda) = \sum_{i=0}^n (-1)^i a_i \lambda^{n-i}$
voloshin-hypergraph_ EQ0066 (5.12)	112	■ 1.000	$\begin{gathered} P(\text{left}(G_1), \lambda) = (\\ \lambda - 2)(\lambda - 2)(\lambda - 3)(\lambda - 2)(\lambda - 2)(\\ \lambda - 2)(\lambda - 1)(\lambda - 0) = \lambda(\lambda - 1)(\lambda - 2)^5(\lambda - 3) \\ \end{gathered}$	$P(G_1, \lambda) = (\lambda - 2)(\lambda - 2)(\lambda - 3)(\lambda - 2)(\lambda - 2)(\lambda - 2)(\lambda - 1)(\lambda - 0) = \lambda(\lambda - 1)(\lambda - 2)^5(\lambda - 3).$	$P(G_1, \lambda) = (\lambda - 2)(\lambda - 2)(\lambda - 3)(\lambda - 2)(\lambda - 2)(\lambda - 2)(\lambda - 1)(\lambda - 0) = \lambda(\lambda - 1)(\lambda - 2)^5(\lambda - 3).$
voloshin-hypergraph_ EQ0067	113	■ 1.000	$P(\text{left}(G_1), 4) = 4(4-1)(4-2)^5(4-3) = 384$.	$P(G_1, 4) = 4(4-1)(4-2)^5(4-3) = 384.$	$P(G_1, 4) = 4(4-1)(4-2)^5(4-3) = 384.$
voloshin-hypergraph_ EQ0079	115	■ 1.000	$\text{left}\{x_0\}\cup N(x_0) \subseteq X_1 \cup A, N(x_0) \cap A \subseteq X_i \cup A, 1 \leq i \leq k.$	$\{x_0\} \cup N(x_0) \subseteq X_1 \cup A, N(x_0) \cap A \subseteq X_i \cup A, 1 \leq i \leq k.$	$\{x_0\} \cup N(x_0) \subseteq X_1 \cup A, N(x_0) \cap A \subseteq X_i \cup A, 1 \leq i \leq k.$
voloshin-hypergraph_ EQ0084	116	■ 1.000	$P(\text{left}(C_k), \lambda) = \lambda(\lambda-1)^2 \lambda(\lambda-1)^{k-2} \lambda^{-2} = (\lambda-1)^k \neq P(C_k, \lambda),$	$P(C_k, \lambda) = \lambda(\lambda-1)^2 \lambda(\lambda-1)^{k-2} \lambda^{-2} = (\lambda-1)^k \neq P(C_k, \lambda),$	$P(C_k, \lambda) = \lambda(\lambda-1)^2 \lambda(\lambda-1)^{k-2} \lambda^{-2} = (\lambda-1)^k \neq P(C_k, \lambda),$
voloshin-hypergraph_ EQ0090	117	■ 1.000	$P(C, \lambda) = P(C-x, \lambda) W(\text{left}(C_{N(x)}), \lambda) = (\lambda-1)^k \neq P(C, \lambda),$	$P(C, \lambda) = P(C-x, \lambda) W(C_{N(x)}, \lambda) = (\lambda-1)^k \neq P(C, \lambda),$	$P(C, \lambda) = P(C-x, \lambda) W(C_{N(x)}, \lambda) = (\lambda-1)^k \neq P(C, \lambda),$
voloshin-hypergraph_ EQ0106	135	■ 1.000	$\bar{\chi}^{\prime}(\mathcal{A}_1) \geq \bar{\chi}^{\prime}(\mathcal{A}_0) + 1 \geq c + m(\mathcal{A}_1) - n(\mathcal{A}_1) + p(\mathcal{A}_1),$	$\bar{\chi}^{\prime}(\mathcal{A}_1) \geq \bar{\chi}^{\prime}(\mathcal{A}_0) + 1 \geq c + m(\mathcal{A}_1) - n(\mathcal{A}_1) + p(\mathcal{A}_1),$	$\bar{\mathcal{X}}^{\prime}(\mathcal{A}_1) \geq \bar{\mathcal{X}}^{\prime}(\mathcal{A}_0) + 1 \geq c + m(\mathcal{A}_1) - n(\mathcal{A}_1) + p(\mathcal{A}_1),$
voloshin-hypergraph_ EQ0146 (8.4)	192	■ 1.000	$\wedge(\mathcal{H}, T) = \sum_{D \in \mathcal{D}} (c(T, D) - 1) + l(\mathcal{H}, T).$	$\wedge(\mathcal{H}, T) = \sum_{D \in \mathcal{D}} (c(T, D) - 1) + l(\mathcal{H}, T).$	$\wedge(\mathcal{H}, T) = \sum_{D \in \mathcal{D}} (c(T, D) - 1) + l(\mathcal{H}, T).$
voloshin-hypergraph_ EQ0180	211	■ 1.000	$\emptyset \leq a_{\{j\}} \leq b_{\{j\}} \leq \mathcal{D}_{\{j\}} $.	$0 \leq a_j \leq b_j \leq D_j $.	$0 \leq a_j \leq b_j \leq D_j $.
voloshin-hypergraph_ EQ0194	220	■ 1.000	$R(\mathcal{H}) = (\emptyset, \ldots, \emptyset, r_{\chi}, \ldots, r_{\bar{\chi}}, \emptyset, \ldots, \emptyset).$	$R(\mathcal{H}) = (0, \ldots, 0, r_{\chi}, \ldots, r_{\bar{\chi}}, 0, \ldots, 0).$	$R(\mathcal{H}) = (0, \ldots, 0, r_{\chi}, \ldots, r_{\bar{\chi}}, 0, \ldots, 0).$
voloshin-hypergraph_ EQ0200	223	■ 1.000	$b(x, \mathcal{H}) = \max_y \{ \mathcal{C}(x) \cap \mathcal{C}(y) : y \in X, y \neq x \}$	$b(x, \mathcal{H}) = \max_y \{ \mathcal{C}(x) \cap \mathcal{C}(y) : y \in X, y \neq x \}$	$b(x, \mathcal{H}) = \max_y \{ \mathcal{C}(x) \cap \mathcal{C}(y) : y \in X, y \neq x \}$
voloshin-hypergraph_ EQ0204	227	■ 1.000	$o(\mathcal{x}_i, \mathcal{H}_i) \leq O(\mathcal{H}) = \max_{Y \subseteq X} \min_{x \in Y} o(x, \mathcal{H}/Y)$	$o(x_i, \mathcal{H}_i) \leq O(\mathcal{H}) = \max_{Y \subseteq X} \min_{x \in Y} o(x, \mathcal{H}/Y)$	$o(x_i, \mathcal{H}_i) \leq O(\mathcal{H}) = \max_{Y \subseteq X} \min_{x \in Y} o(x, \mathcal{H}/Y)$
voloshin-hypergraph_ EQ0229	248	■ 1.000	$M = \sum_{\substack{x \in T \\ d(x) \geq 2}} \binom{d(x)}{2},$	$M = \sum_{\substack{x \in T \\ d(x) \geq 2}} \binom{d(x)}{2},$	$M = \sum_{\substack{x \in T \\ d(x) \geq 2}} \binom{d(x)}{2},$
voloshin-hypergraph_ EQ0237	255	■ 1.000	$\begin{aligned} & n_{12} + 1 + n_{23} < r, \quad \& \\ & n_{23} + 1 + n_{31} < r, \quad \end{aligned}$	$\begin{aligned} & n_{12} + 1 + n_{23} < r, \\ & n_{23} + 1 + n_{31} < r, \end{aligned}$	$\begin{aligned} & n_{12} + 1 + n_{23} < r, \\ & n_{23} + 1 + n_{31} < r, \end{aligned}$
voloshin-hypergraph_ EQ0001	22	■ 1.000	$G=(X, E)$.	$G=(X, E).$	$G=(X, E).$
voloshin-hypergraph_ EQ0002 (1.1)	23	■ 1.000	$\sum_{i=1}^n d(\mathcal{x}_i) = 2m$.	$\sum_{i=1}^n d(x_i) = 2m.$	$\sum_{i=1}^n d(x_i) = 2m.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0003	23	■ 1.000	$2+4+2+3+3=14=2 \times 7$.	$2 + 4 + 2 + 3 + 3 = 14 = 2 \times 7.$	$2 + 4 + 2 + 3 + 3 = 14 = 2 \times 7.$
voloshin-hypergraph_ EQ0005	29	■ 1.000	$A(G)=\left(\begin{array}{lllll} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 \end{array}\right)$.	$A(G)=\left(\begin{array}{ccccc} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 \end{array}\right).$	$A(G)=\left(\begin{array}{ccccc} 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 \end{array}\right).$
voloshin-hypergraph_ EQ0008	32	■ 1.000	$L(G)=\{\emptyset,\{2\},\{2,4\},\{1,1,3\}\}$;	$L(G)=\{\emptyset,\{2\},\{2,4\},\{1,1,3\}\};$	$L(G)=\{\emptyset,\{2\},\{2,4\},\{1,1,3\}\};$
voloshin-hypergraph_ EQ0010	33	■ 1.000	$I(G)=\left(\begin{array}{rrrrrr} 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & l & 0 & 0 \\ 0 & 0 & -1 & 0 & 1 & -1 \\ -1 & -1 & 0 & 0 & -1 & 1 \end{array}\right)$;	$I(G)=\left(\begin{array}{ccccc} 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & l & 0 \\ 0 & 0 & -1 & 0 & 1 \\ -1 & -1 & 0 & 0 & -1 \end{array}\right);$	$I(G)=\left(\begin{array}{ccccc} 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & l & 0 \\ 0 & 0 & -1 & 0 & 1 \\ -1 & -1 & 0 & 0 & -1 \end{array}\right);$
voloshin-hypergraph_ EQ0011	33	■ 1.000	$J(G)=\{(4,1),(4,1),(3,2),(2,2),(4,3),(3,4)\}$.	$J(G)=\{(4,1),(4,1),(3,2),(2,2),(4,3),(3,4)\}.$	$J(G)=\{(4,1),(4,1),(3,2),(2,2),(4,3),(3,4)\}.$
voloshin-hypergraph_ EQ0012	36	■ 1.000	$m=\binom{n}{2}=\frac{n(n-1)}{2}$.	$m=\binom{n}{2}=\frac{n(n-1)}{2}.$	$m=\binom{n}{2}=\frac{n(n-1)}{2}.$
voloshin-hypergraph_ EQ0014	47	■ 1.000	$\alpha(G)+\tau(G)= X $.	$\alpha(G)+\tau(G)= X .$	$\alpha(G)+\tau(G)= X .$
voloshin-hypergraph_ EQ0016	61	■ 1.000	$X=N_{\{0\}} \cup N_{\{1\}} \cup N_{\{2\}} \cup N_{\{3\}} \cup \ldots \cup N_{\{k\}}$.	$X=N_0 \cup N_1 \cup N_2 \cup N_3 \cup \ldots \cup N_k.$	$X=N_0 \cup N_1 \cup N_2 \cup N_3 \cup \ldots \cup N_k.$
voloshin-hypergraph_ EQ0017	62	■ 1.000	$N(S)=\cup_{x \in S} N(x) \backslash S$.	$N(S)=\cup_{x \in S} N(x) \backslash S.$	$N(S)=\cup_{x \in S} N(x) \backslash S.$
voloshin-hypergraph_ EQ0020	63	■ 1.000	$\left S^{\prime}\right \leq\left N_{\{G\}}\left(S^{\prime}\right)\right $.	$ S' \leq N_G(S') .$	$ S' \leq N_G(S') .$
voloshin-hypergraph_ EQ0021 (2.2)	63	■ 1.000	$\left S_{\{0\}}\right =\left N_{\{G\}}\left(S_{\{0\}}\right)\right $.	$ S_0 = N_G(S_0) .$	$ S_0 = N_G(S_0) .$
voloshin-hypergraph_ EQ0022	63	■ 1.000	$A=S_{\{0\}} \cup N_{\{G\}}\left(S_{\{0\}}\right) \text { and } B=\left(X \cup Y\right) \backslash A$	$A=S_0 \cup N_G\left(S_0\right) \text { and } B=\left(X \cup Y\right) \backslash A$	$A=S_0 \cup N_G\left(S_0\right) \text { and } B=\left(X \cup Y\right) \backslash A$
voloshin-hypergraph_ EQ0023	63	■ 1.000	$N_{\{G\}\{B\}}(S)=N_{\{G\}}(S) \backslash N_{\{G\}}\left(S_{\{0\}}\right),$	$N_{G_B}(S)=N_G(S) \backslash N_G\left(S_0\right),$	$N_{G_B}(S)=N_G(S) \backslash N_G\left(S_0\right),$
voloshin-hypergraph_ EQ0024	63	■ 1.000	$\left S_{\{0\}}\right + S =\left S_{\{0\}} \cup S\right \leq\left N_{\{G\}}\left(S_{\{0\}} \cup S\right)\right =\left N_{\{G\}}\left(S_{\{0\}}\right)\right +\left N_{\{G\}\{B\}}(S)\right $.	$ S_0 + S = S_0 \cup S \leq N_G\left(S_0 \cup S\right) = N_G\left(S_0\right) + N_{G_B}(S) .$	$ S_0 + S = S_0 \cup S \leq N_G\left(S_0 \cup S\right) = N_G\left(S_0\right) + N_{G_B}(S) .$
voloshin-hypergraph_ EQ0027	74	■ 1.000	$M(G)-(\omega(G)-1)$.	$M(G)-(\omega(G)-1).$	$M(G)-(\omega(G)-1).$
voloshin-hypergraph_ EQ0028	75	■ 1.000	$X=N_{\{0\}} \cup N_{\{1\}} \cup N_{\{2\}} \cup \cdots \cup N_{\{k-1\}} \cup N_{\{k\}}$	$X=N_0 \cup N_1 \cup N_2 \cup \cdots \cup N_{k-1} \cup N_k$	$X=N_0 \cup N_1 \cup N_2 \cup \cdots \cup N_{k-1} \cup N_k$
voloshin-hypergraph_ EQ0031	97	■ 1.000	$R(G)=\left(r_{\{1\}}, r_{\{2\}}, \ldots, r_{\{n\}}\right)$	$R(G)=\left(r_1, r_2, \ldots, r_n\right)$	$R(G)=\left(r_1, r_2, \ldots, r_n\right)$
voloshin-hypergraph_ EQ0032	97	■ 1.000	$R(G)=\left(0,0, \ldots, 0, r_{\chi}, r_{\chi+1}, \ldots, r_{\{n\}}\right)$.	$R(G)=\left(0,0, \ldots, 0, r_{\chi}, r_{\chi+1}, \ldots, r_n\right).$	$R(G)=\left(0,0, \ldots, 0, r_{\chi}, r_{\chi+1}, \ldots, r_n\right).$
voloshin-hypergraph_ EQ0033	98	■ 1.000	$R(G)=(0,1,1)$.	$R(G)=(0,1,1).$	$R(G)=(0,1,1).$
voloshin-hypergraph_ EQ0034	99	■ 1.000	$P\left(K_{\{n\}}, \lambda\right)=\lambda(\lambda-1)(\lambda-2) \cdots(\lambda-n+1)$.	$P\left(K_n, \lambda\right)=\lambda(\lambda-1)(\lambda-2) \cdots(\lambda-n+1).$	$P\left(K_n, \lambda\right)=\lambda(\lambda-1)(\lambda-2) \cdots(\lambda-n+1).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0035 (5.1)	100	■ 1.000	$P(G, \lambda) = P(G_1, \lambda) + P(G_2, \lambda) + \dots + P(K_s, \lambda)$	$P(G, \lambda) = P(G_1, \lambda) + P(G_2, \lambda)$	$P(G, \lambda) = P(G_1, \lambda) + P(G_2, \lambda)$
voloshin-hypergraph_ EQ0036	101	■ 1.000	$P(G, \lambda) = P(K_{i_1}, \lambda) + P(K_{i_2}, \lambda) + \dots + P(K_{i_s}, \lambda)$	$P(G, \lambda) = P(K_{i_1}, \lambda) + P(K_{i_2}, \lambda) + \dots + P(K_{i_s}, \lambda)$	$P(G, \lambda) = P(K_{i_1}, \lambda) + P(K_{i_2}, \lambda) + \dots + P(K_{i_s}, \lambda)$
voloshin-hypergraph_ EQ0037	101	■ 1.000	$P(G, \lambda) = P(K_{i_1}, \lambda) + P(K_{i_2}, \lambda) + \dots + P(K_{i_s}, \lambda) = r_1(G)P(K_1, \lambda) + r_2(G)P(K_2, \lambda) + \dots + r_n(G)P(K_n, \lambda)$	$P(G, \lambda) = P(K_{i_1}, \lambda) + P(K_{i_2}, \lambda) + \dots + P(K_{i_s}, \lambda) = r_1(G)P(K_1, \lambda) + r_2(G)P(K_2, \lambda) + \dots + r_n(G)P(K_n, \lambda)$	$P(G, \lambda) = P(K_{i_1}, \lambda) + P(K_{i_2}, \lambda) + \dots + P(K_{i_s}, \lambda) = r_1(G)P(K_1, \lambda) + r_2(G)P(K_2, \lambda) + \dots + r_n(G)P(K_n, \lambda)$
voloshin-hypergraph_ EQ0038 (5.2)	102	■ 1.000	$\sum_{i=1}^n r_i(G) P(K_i, \lambda)$	$\sum_{i=1}^n r_i(G) P(K_i, \lambda)$	$\sum_{i=1}^n r_i(G) P(K_i, \lambda)$
voloshin-hypergraph_ EQ0042	103	■ 1.000	$R(C_4) = (0, 1, 2, 1)$	$R(C_4) = (0, 1, 2, 1)$	$R(C_4) = (0, 1, 2, 1)$
voloshin-hypergraph_ EQ0043 (5.5)	105	■ 1.000	$P(G, \lambda) = P(G_1, \lambda) P(G_2, \lambda) \dots P(G_k, \lambda)$	$P(G, \lambda) = P(G_1, \lambda) P(G_2, \lambda) \dots P(G_k, \lambda)$	$P(G, \lambda) = P(G_1, \lambda) P(G_2, \lambda) \dots P(G_k, \lambda)$
voloshin-hypergraph_ EQ0044	105	■ 1.000	$P(E_n, \lambda) = \lambda \cdot \lambda \cdot \dots \cdot \lambda = \lambda^n$	$P(E_n, \lambda) = \lambda \cdot \lambda \cdot \dots \cdot \lambda = \lambda^n$	$P(E_n, \lambda) = \lambda \cdot \lambda \cdot \dots \cdot \lambda = \lambda^n$
voloshin-hypergraph_ EQ0045 (5.6)	105	■ 1.000	$\lambda^n = S(n, 1)\lambda^{(1)} + S(n, 2)\lambda^{(2)} + \dots + S(n, n)\lambda^{(n)}$	$\lambda^n = S(n, 1)\lambda^{(1)} + S(n, 2)\lambda^{(2)} + \dots + S(n, n)\lambda^{(n)}$	$\lambda^n = S(n, 1)\lambda^{(1)} + S(n, 2)\lambda^{(2)} + \dots + S(n, n)\lambda^{(n)}$
voloshin-hypergraph_ EQ0046	106	■ 1.000	$R(E_n) = (S(n, 1), S(n, 2), \dots, S(n, n))$	$R(E_n) = (S(n, 1), S(n, 2), \dots, S(n, n))$	$R(E_n) = (S(n, 1), S(n, 2), \dots, S(n, n))$
voloshin-hypergraph_ EQ0047	106	■ 1.000	$\lambda^{(n)} = s(n, 1)\lambda + s(n, 2)\lambda^2 + \dots + s(n, n)\lambda^n$	$\lambda^{(n)} = s(n, 1)\lambda + s(n, 2)\lambda^2 + \dots + s(n, n)\lambda^n$	$\lambda^{(n)} = s(n, 1)\lambda + s(n, 2)\lambda^2 + \dots + s(n, n)\lambda^n$
voloshin-hypergraph_ EQ0048	106	■ 1.000	$P(G, \lambda) = P(G_1, \lambda) + P(G_2, \lambda)$	$P(G, \lambda) = P(G_1, \lambda) + P(G_2, \lambda)$	$P(G, \lambda) = P(G_1, \lambda) + P(G_2, \lambda)$
voloshin-hypergraph_ EQ0049 (5.7)	106	■ 1.000	$P(G_1, \lambda) = P(G, \lambda) - P(G_2, \lambda)$	$P(G_1, \lambda) = P(G, \lambda) - P(G_2, \lambda)$	$P(G_1, \lambda) = P(G, \lambda) - P(G_2, \lambda)$
voloshin-hypergraph_ EQ0053 (5.8)	108	■ 1.000	$P(G, \lambda) = \frac{P(G_1, \lambda) P(G_2, \lambda)}{\lambda^{(k)}}$	$P(G, \lambda) = \frac{P(G_1, \lambda) P(G_2, \lambda)}{\lambda^{(k)}}$	$P(G, \lambda) = \frac{P(G_1, \lambda) P(G_2, \lambda)}{\lambda^{(k)}}$
voloshin-hypergraph_ EQ0055 (5.9)	109	■ 1.000	$P(G, \lambda) = \frac{P(G_1, \lambda) P(G_2, \lambda) \dots P(G_l, \lambda)}{[\lambda^{(k)}]^{l-1}}$	$P(G, \lambda) = \frac{P(G_1, \lambda) P(G_2, \lambda) \dots P(G_l, \lambda)}{[\lambda^{(k)}]^{l-1}}$	$P(G, \lambda) = \frac{P(G_1, \lambda) P(G_2, \lambda) \dots P(G_l, \lambda)}{[\lambda^{(k)}]^{l-1}}$
voloshin-hypergraph_ EQ0057 (5.10)	109	■ 1.000	$P(G, \lambda) = (\lambda - k) P(G - x, \lambda)$	$P(G, \lambda) = (\lambda - k) P(G - x, \lambda)$	$P(G, \lambda) = (\lambda - k) P(G - x, \lambda)$
voloshin-hypergraph_ EQ0058	109	■ 1.000	$P(G, \lambda) = \frac{P(G_1, \lambda) P(K_{k+1}, \lambda)}{\lambda^{(k)}} =$	$P(G, \lambda) = \frac{P(G_1, \lambda) P(K_{k+1}, \lambda)}{\lambda^{(k)}} =$	$P(G, \lambda) = \frac{P(G_1, \lambda) P(K_{k+1}, \lambda)}{\lambda^{(k)}} =$
voloshin-hypergraph_ EQ0059	110	■ 1.000	$\frac{P(G - x, \lambda) \lambda^{(k+1)}}{\lambda^{(k)}} = (\lambda - k) P(G - x, \lambda)$	$\frac{P(G - x, \lambda) \lambda^{(k+1)}}{\lambda^{(k)}} = (\lambda - k) P(G - x, \lambda)$	$\frac{P(G - x, \lambda) \lambda^{(k+1)}}{\lambda^{(k)}} = (\lambda - k) P(G - x, \lambda)$
voloshin-hypergraph_ EQ0060	110	■ 1.000	$P(G, \lambda) = (\lambda - 2) P(G - x, \lambda) = \lambda(\lambda - 1)(\lambda - 2)(\lambda^2 - 3\lambda + 3)$	$P(G, \lambda) = (\lambda - 2) P(G - x, \lambda) = \lambda(\lambda - 1)(\lambda - 2)(\lambda^2 - 3\lambda + 3)$	$P(G, \lambda) = (\lambda - 2) P(G - x, \lambda) = \lambda(\lambda - 1)(\lambda - 2)(\lambda^2 - 3\lambda + 3)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0061	110	■ 1.000	$r_{\{i\}}(G)=(i-k) \ r_{\{i\}}(G-x)+r_{\{i-1\}}(G-x) \ .$	$r_i(G) = (i - k)r_i(G - x) + r_{i-1}(G - x).$	$r_i(G) = (i - k)r_i(G - x) + r_{i-1}(G - x).$
voloshin-hypergraph_ EQ0062	110	■ 1.000	$S(n, i)=i \ S(n-1, i)+S(n-1, i-1) \ .$	$S(n, i) = iS(n - 1, i) + S(n - 1, i - 1).$	$S(n, i) = iS(n - 1, i) + S(n - 1, i - 1).$
voloshin-hypergraph_ EQ0063	110	■ 1.000	$S(5,3)=25=3 \ \cdot \ S(4,3)+S(4,2)=3 \ \cdot \ 6+7=25 \ .$	$S(5, 3) = 25 = 3 \cdot S(4, 3) + S(4, 2) = 3 \cdot 6 + 7 = 25.$	$S(5, 3) = 25 = 3 \cdot S(4, 3) + S(4, 2) = 3 \cdot 6 + 7 = 25.$
voloshin-hypergraph_ EQ0068 (5.13)	113	■ 1.000	$P\left(T_{\{n\}}, \ \lambda\right)=\lambda(\lambda-1)^{\{n-1\}}$	$P(T_n, \lambda) = \lambda(\lambda - 1)^{n-1}$	$P(T_n, \lambda) = \lambda(\lambda - 1)^{n-1}.$
voloshin-hypergraph_ EQ0069 (5.14)	113	■ 1.000	$P\left(C_{\{n\}}, \ \lambda\right)=(\lambda-1)^{\{n\}}+(-1)^{\{n\}}(\lambda-1) \ .$	$P(C_n, \lambda) = (\lambda - 1)^n + (-1)^n(\lambda - 1).$	$P(C_n, \lambda) = (\lambda - 1)^n + (-1)^n(\lambda - 1).$
voloshin-hypergraph_ EQ0070	113	■ 1.000	$P\left(C_{\{n\}}, \ \lambda\right)=P\left(T_{\{n\}}, \ \lambda\right)-P\left(C_{\{n-1\}}, \ \lambda\right)=$	$P(C_n, \lambda) = P(T_n, \lambda) - P(C_{n-1}, \lambda) =$	$P(C_n, \lambda) = P(T_n, \lambda) - P(C_{n-1}, \lambda) =$
voloshin-hypergraph_ EQ0073	114	■ 1.000	$G_{\{X_{\{1\}} \ \cup \ A\}}=G_{\{1\}}, \ G_{\{X_{\{2\}} \ \cup \ A\}}=G_{\{2\}}, \ \ldots, \ G_{\{X_{\{k\}} \ \cup \ A\}}=G_{\{k\}}, \ G_{\{A\}}=G_{\{0\}} \ .$	$G_{X_1 \cup A} = G_1, G_{X_2 \cup A} = G_2, \dots, G_{X_k \cup A} = G_k, G_A = G_0.$	$G_{X_1 \cup A} = G_1, \ G_{X_2 \cup A} = G_2, \ \dots, \ G_{X_k \cup A} = G_k, \ G_A = G_0.$
voloshin-hypergraph_ EQ0074 (5.15)	114	■ 1.000	$P\left(G^{\{\prime\}}, \ \lambda\right)=\frac{P\left(G_{\{1\}}^{\{\prime\}}, \ \lambda\right) \ P\left(G_{\{2\}}^{\{\prime\}}, \ \lambda\right) \ \ldots \ P\left(G_{\{k\}}^{\{\prime\}}, \ \lambda\right)}{P\left(G_{\{0\}}^{\{\prime\}}, \ \lambda\right)^{\{k-1\}}}.$	$P(G', \lambda) = \frac{P(G'_1, \lambda) P(G'_2, \lambda) \dots P(G'_k, \lambda)}{P(G'_0, \lambda)^{k-1}}.$	$P(G', \lambda) = \frac{P(G'_1, \lambda)P(G'_2, \lambda) \dots P(G'_k, \lambda)}{P(G'_0, \lambda)^{k-1}}.$
voloshin-hypergraph_ EQ0075	114	■ 1.000	$P\left(G_{\{1\}}, \ \lambda\right)=(\lambda-p) \ P\left(G_{\{1\}}-x_{\{0\}}, \ \lambda\right) \ .$	$P(G_1, \lambda) = (\lambda - p)P(G_1 - x_0, \lambda) \ .$	$P(G_1, \lambda) = (\lambda - p)P(G_1 - x_0, \lambda).$
voloshin-hypergraph_ EQ0076	114	■ 1.000	$P(G, \ \lambda)=(\lambda-p) \ P\left(G-x_{\{0\}}, \ \lambda\right) \ .$	$P(G, \lambda) = (\lambda - p)P(G - x_0, \lambda) \ .$	$P(G, \lambda) = (\lambda - p)P(G - x_0, \lambda).$
voloshin-hypergraph_ EQ0077 (5.16)	114	■ 1.000	$P\left(G-x_{\{0\}}, \ \lambda\right)=\frac{P\left(G_{\{1\}}-x_{\{0\}}, \ \lambda\right) \ P\left(G_{\{2\}}, \ \lambda\right) \ \ldots \ P\left(G_{\{k\}}, \ \lambda\right)}{P\left(G_{\{0\}}, \ \lambda\right)^{\{k-1\}}}.$	$P(G - x_0, \lambda) = \frac{P(G_1 - x_0, \lambda) P(G_2, \lambda) \dots P(G_k, \lambda)}{P(G_0, \lambda)^{k-1}}.$	$P(G - x_0, \lambda) = \frac{P(G_1 - x_0, \lambda)P(G_2, \lambda) \dots P(G_k, \lambda)}{P(G_0, \lambda)^{k-1}}.$
voloshin-hypergraph_ EQ0078	115	■ 1.000	$P(G, \ \lambda)=\frac{P\left(G_{\{1\}}, \ \lambda\right) \ P\left(G_{\{2\}}, \ \lambda\right) \ \ldots \ P\left(G_{\{k\}}, \ \lambda\right)}{P\left(G_{\{0\}}, \ \lambda\right)^{\{k-1\}}}.$	$P(G, \lambda) = \frac{P(G_1, \lambda) P(G_2, \lambda) \dots P(G_k, \lambda)}{P(G_0, \lambda)^{k-1}}.$	$P(G, \lambda) = \frac{P(G_1, \lambda)P(G_2, \lambda) \dots P(G_k, \lambda)}{P(G_0, \lambda)^{k-1}}.$
voloshin-hypergraph_ EQ0080	115	■ 1.000	$P(G, \ \lambda)=(\lambda-p) \ P\left(G-x_{\{0\}}, \ \lambda\right) \ .$	$P(G, \lambda) = (\lambda - p)P(G - x_0, \lambda) \ .$	$P(G, \lambda) = (\lambda - p)P(G - x_0, \lambda).$
voloshin-hypergraph_ EQ0081 (5.17)	115	■ 1.000	$P\left(G-x_{\{0\}}, \ \lambda\right)=\frac{P\left(G_{\{1\}}^{\{\prime\}}, \ \lambda\right) \ P\left(G_{\{2\}}^{\{\prime\}}, \ \lambda\right) \ \ldots \ P\left(G_{\{k\}}^{\{\prime\}}, \ \lambda\right)}{P\left(G_{\{0\}}^{\{\prime\}}, \ \lambda\right)^{\{k-1\}}},$	$P(G - x_0, \lambda) = \frac{P(G'_1, \lambda) P(G'_2, \lambda) \dots P(G'_k, \lambda)}{P(G'_0, \lambda)^{k-1}},$	$P(G - x_0, \lambda) = \frac{P(G'_1, \lambda)P(G'_2, \lambda) \dots P(G'_k, \lambda)}{P(G'_0, \lambda)^{k-1}},$
voloshin-hypergraph_ EQ0082	115	■ 1.000	$P\left(G_{\{1\}}, \ \lambda\right)=(\lambda-p) \ P\left(G_{\{1\}}^{\{\prime\}}, \ \lambda\right), \ \text{and} \ P\left(G_{\{i\}}, \ \lambda\right)=(\lambda-p_1) \ P\left(G_{\{i\}}^{\{\prime\}}, \ \lambda\right)$	$P(G_1, \lambda) = (\lambda - p)P(G'_1, \lambda), \ \text{and} \ P(G_i, \lambda) = (\lambda - p_1) P(G'_i, \lambda)$	$P(G_1, \lambda) = (\lambda - p)P(G'_1, \lambda), \ \text{and} \ P(G_i, \lambda) = (\lambda - p_1)P(G'_i, \lambda)$
voloshin-hypergraph_ EQ0083	116	■ 1.000	$P(G, \ \lambda)=\frac{P\left(G_{\{1\}}, \ \lambda\right) \ P\left(G_{\{2\}}, \ \lambda\right) \ \ldots \ P\left(G_{\{k\}}, \ \lambda\right)}{P\left(G_{\{0\}}, \ \lambda\right)^{\{k-1\}}}.$	$P(G, \lambda) = \frac{P(G_1, \lambda) P(G_2, \lambda) \dots P(G_k, \lambda)}{P(G_0, \lambda)^{k-1}}.$	$P(G, \lambda) = \frac{P(G_1, \lambda)P(G_2, \lambda) \dots P(G_k, \lambda)}{P(G_0, \lambda)^{k-1}}.$
voloshin-hypergraph_ EQ0086 (5.18)	116	■ 1.000	$P\left(G^{\{\prime\}}, \ \lambda\right)=P\left(G^{\{\prime\}}-x, \ \lambda\right) \ W\left(G_{N(x)}^{\{\prime\}}, \ \lambda\right) \ .$	$P(G', \lambda) = P(G' - x, \lambda) W\left(G'_{N(x)}, \lambda\right) \ .$	$P(G', \lambda) = P(G' - x, \lambda)W(G'_{N(x)}, \lambda).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0088	116	■ 1.000	$P(G, \lambda) = \frac{P(G-x, \lambda)}{P(G_{N(x)}, \lambda)}$	$P(G, \lambda) = \frac{P(G-x, \lambda)P(G_{\{x\} \cup N(x)}, \lambda)}{P(G_{N(x)}, \lambda)} =$	$P(G, \lambda) = \frac{P(G-x, \lambda)P(G_{\{x\} \cup N(x)}, \lambda)}{P(G_{N(x)}, \lambda)} =$
voloshin-hypergraph_ EQ0089	117	■ 1.000	$P(G-x, \lambda) \frac{\lambda P(G_{N(x)}, \lambda-1)}{P(G_{N(x)}, \lambda)} = P(G-x, \lambda)W(G_{N(x)}, \lambda).$	$P(G-x, \lambda) \frac{\lambda P(G_{N(x)}, \lambda-1)}{P(G_{N(x)}, \lambda)} = P(G-x, \lambda)W(G_{N(x)}, \lambda).$	$P(G-x, \lambda) \frac{\lambda P(G_{N(x)}, \lambda-1)}{P(G_{N(x)}, \lambda)} = P(G-x, \lambda)W(G_{N(x)}, \lambda).$
voloshin-hypergraph_ EQ0091	117	■ 1.000	$P(G, \lambda) = (\lambda - k) P(G-x, \lambda).$	$P(G, \lambda) = (\lambda - k)P(G-x, \lambda).$	$P(G, \lambda) = (\lambda - k)P(G-x, \lambda).$
voloshin-hypergraph_ EQ0092	117	■ 1.000	$W(G_{N(x)}, \lambda) = W(K_k, \lambda) = \frac{\lambda(\lambda-1)^{(k)}}{\lambda^{(k)}} = (\lambda - k).$	$W(G_{N(x)}, \lambda) = W(K_k, \lambda) = \frac{\lambda(\lambda-1)^{(k)}}{\lambda^{(k)}} = (\lambda - k).$	$W(G_{N(x)}, \lambda) = W(K_k, \lambda) = \frac{\lambda(\lambda-1)^{(k)}}{\lambda^{(k)}} = (\lambda - k).$
voloshin-hypergraph_ EQ0093	117	■ 1.000	$P(G, \lambda) = P(G-x, \lambda) W(G_{N(x)}, \lambda) = (\lambda - k)P(G-x, \lambda).$	$P(G, \lambda) = P(G-x, \lambda)W(G_{N(x)}, \lambda) = (\lambda - k)P(G-x, \lambda).$	$P(G, \lambda) = P(G-x, \lambda)W(G_{N(x)}, \lambda) = (\lambda - k)P(G-x, \lambda).$
voloshin-hypergraph_ EQ0094	117	■ 1.000	$W(G_{N(4)}, \lambda) = \frac{\lambda P(T_3, \lambda-1)}{P(T_3, \lambda)} = \frac{\lambda(\lambda-1)(\lambda-2)^2}{\lambda(\lambda-1)^2} = \frac{(\lambda-2)^2}{\lambda-1}.$	$W(G_{N(4)}, \lambda) = \frac{\lambda P(T_3, \lambda-1)}{P(T_3, \lambda)} = \frac{\lambda(\lambda-1)(\lambda-2)^2}{\lambda(\lambda-1)^2} = \frac{(\lambda-2)^2}{\lambda-1}.$	$W(G_{N(4)}, \lambda) = \frac{\lambda P(T_3, \lambda-1)}{P(T_3, \lambda)} = \frac{\lambda(\lambda-1)(\lambda-2)^2}{\lambda(\lambda-1)^2} = \frac{(\lambda-2)^2}{\lambda-1}.$
voloshin-hypergraph_ EQ0095	117	■ 1.000	$P(G_1, \lambda) = \lambda(\lambda-1)^2(\lambda-2)^3(\lambda-3).$	$P(G_1, \lambda) = \lambda(\lambda-1)^2(\lambda-2)^3(\lambda-3).$	$P(G_1, \lambda) = \lambda(\lambda-1)^2(\lambda-2)^3(\lambda-3).$
voloshin-hypergraph_ EQ0097	118	■ 1.000	$M(G) = \max_{X' \subseteq X} \min_{x \in G'} d(x)$	$M(G) = \max_{X' \subseteq X} \min_{x \in G'} d(x)$	$M(G) = \max_{X' \subseteq X} \min_{x \in G'} d(x)$
voloshin-hypergraph_ EQ0098 (5.19)	124	■ 1.000	$\chi(G) \geq \omega(G).$	$\chi(G) \geq \omega(G).$	$\chi(G) \geq \omega(G).$
voloshin-hypergraph_ EQ0100	133	■ 1.000	$\bar{\chi}^i(G_i) = 1, i=1, \dots,$	$\bar{\chi}^i(G_i) = 1, i=1, \dots, \bar{\chi}^i(G).$	$\bar{\chi}^i(G_i) = 1, i=1, \dots, \bar{\chi}^i(G).$
voloshin-hypergraph_ EQ0101	134	■ 1.000	$\bar{\chi}^i(G) = c+m-n+p.$	$\bar{\chi}^i(G) = c+m-n+p.$	$\bar{\chi}^i(G) = c+m-n+p.$
voloshin-hypergraph_ EQ0102	135	■ 1.000	$n-v'-p+r.$	$n-v'-p+r.$	$n-v'-p+r.$
voloshin-hypergraph_ EQ0103	135	■ 1.000	$m'=v'+(n-v'-p+r)=n-p+r.$	$m'=v'+(n-v'-p+r)=n-p+r.$	$m'=v'+(n-v'-p+r)=n-p+r.$
voloshin-hypergraph_ EQ0104	135	■ 1.000	$\bar{\chi}^i(G) = c'+r+(m-m')=c'+m-n+p,$	$\bar{\chi}^i(G) = c'+r+(m-m')=c'+m-n+p,$	$\bar{\chi}^i(G) = c'+r+(m-m')=c'+m-n+p,$
voloshin-hypergraph_ EQ0105	135	■ 1.000	$\bar{\chi}^i(G) = c'+m-n+p \leq c+m-n+p.$	$\bar{\chi}^i(G) = c'+m-n+p \leq c+m-n+p.$	$\bar{\chi}^i(G) = c'+m-n+p \leq c+m-n+p.$
voloshin-hypergraph_ EQ0108	136	■ 1.000	$\bar{\chi}^i(G) = p-1.$	$\bar{\chi}^i(G) = p-1.$	$\bar{\chi}^i(G) = p-1.$
voloshin-hypergraph_ EQ0109	141	■ 1.000	$d(x)+d(y) \geq n,$	$d(x)+d(y) \geq n,$	$d(x)+d(y) \geq n,$
voloshin-hypergraph_ EQ0110	141	■ 1.000	$d(x_1)+d(x_n) \leq (n-2)+(n-2)-(n-3)=n-1$	$d(x_1)+d(x_n) \leq (n-2)+(n-2)-(n-3)=n-1$	$d(x_1)+d(x_n) \leq (n-2)+(n-2)-(n-3)=n-1$
voloshin-hypergraph_ EQ0111	142	■ 1.000	$x_1, x_{i+1}, x_{i+2}, \dots, x_n, x_i, x_{i-1}, \dots, x_1$	$x_1, x_{i+1}, x_{i+2}, \dots, x_n, x_i, x_{i-1}, \dots, x_1$	$x_1, x_{i+1}, x_{i+2}, \dots, x_n, x_i, x_{i-1}, \dots, x_1$
voloshin-hypergraph_ EQ0112	144	■ 1.000	$u \rightarrow x_4 \rightarrow x_3 \rightarrow x_1 \rightarrow x_2 \rightarrow v$	$u \rightarrow x_4 \rightarrow x_3 \rightarrow x_1 \rightarrow x_2 \rightarrow v$	$u \rightarrow x_4 \rightarrow x_3 \rightarrow x_1 \rightarrow x_2 \rightarrow v$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0113	144	■ 1.000	$u \rightarrow x_{\{1\}} \rightarrow x_{\{2\}} \rightarrow v$.	$u \rightarrow x_1 \rightarrow x_2 \rightarrow v$.	$u \rightarrow x_1 \rightarrow x_2 \rightarrow v$.
voloshin-hypergraph_ EQ0114	144	■ 1.000	$u \rightarrow x_{\{4\}} \rightarrow x_{\{3\}} \rightarrow v$.	$u \rightarrow x_4 \rightarrow x_3 \rightarrow v$.	$u \rightarrow x_4 \rightarrow x_3 \rightarrow v$.
voloshin-hypergraph_ EQ0115	144	■ 1.000	$u \rightarrow x_{\{4\}} \rightarrow x_{\{3\}} \rightarrow x_{\{1\}} \rightarrow x_{\{2\}} \rightarrow v$.	$u \rightarrow x_4 \rightarrow x_3 \rightarrow x_1 \rightarrow x_2 \rightarrow v$.	$u \rightarrow x_4 \rightarrow x_3 \rightarrow x_1 \rightarrow x_2 \rightarrow v$.
voloshin-hypergraph_ EQ0116	152	■ 1.000	$\Delta(\mathcal{H}) = \max_{x \in X} \mathcal{D}(x) $.	$\Delta(\mathcal{H}) = \max_{x \in X} \mathcal{D}(x) $.	$\Delta(\mathcal{H}) = \max_{x \in X} \mathcal{D}(x) $.
voloshin-hypergraph_ EQ0117	152	■ 1.000	$r(\mathcal{H}) = \max_{D \in \mathcal{D}} D $.	$r(\mathcal{H}) = \max_{D \in \mathcal{D}} D $.	$r(\mathcal{H}) = \max_{D \in \mathcal{D}} D $.
voloshin-hypergraph_ EQ0121 (7.1)	158	■ 1.000	$\sum_{i=1}^n \mathcal{D}(x_i) = \sum_{j=1}^m D_j $.	$\sum_{i=1}^n \mathcal{D}(x_i) = \sum_{j=1}^m D_j $.	$\sum_{i=1}^n \mathcal{D}(x_i) = \sum_{j=1}^m D_j $.
voloshin-hypergraph_ EQ0122	158	■ 1.000	$3+3+2+2+0=12=1+2+3+3$.	$3 + 3 + 2 + 2 + 2 + 0 = 12 = 1 + 2 + 3 + 3 + 3$.	$3 + 3 + 2 + 2 + 2 + 0 = 12 = 1 + 2 + 3 + 3 + 3$.
voloshin-hypergraph_ EQ0123	159	■ 1.000	$L = \{\{1\}, \{1, 2\}, \{1, 2, 4\}, \{2, 3, 5\}, \{3, 4, 5\}\}$.	$L = \{\{1\}, \{1, 2\}, \{1, 2, 4\}, \{2, 3, 5\}, \{3, 4, 5\}\}$.	$L = \{\{1\}, \{1, 2\}, \{1, 2, 4\}, \{2, 3, 5\}, \{3, 4, 5\}\}$.
voloshin-hypergraph_ EQ0126	160	■ 1.000	$ \mathcal{D}(K_4^3) = \frac{\binom{4}{3}}{1 \cdot 2 \cdot 3} = 4$.	$ \mathcal{D}(K_4^3) = \binom{4}{3} = \frac{4 \cdot 3 \cdot 2}{1 \cdot 2 \cdot 3} = 4$.	$ \mathcal{D}(K_4^3) = \binom{4}{3} = \frac{4 \cdot 3 \cdot 2}{1 \cdot 2 \cdot 3} = 4$.
voloshin-hypergraph_ EQ0127	160	■ 1.000	$\mu = x_{\{0\}} D_{\{0\}} x_{\{1\}} D_{\{1\}} x_{\{2\}} \dots x_{\{t-1\}} D_{\{t-1\}} x_{\{t\}}$	$\mu = x_0 D_0 x_1 D_1 x_2 \dots x_{t-1} D_{t-1} x_t$	$\mu = x_0 D_0 x_1 D_1 x_2 \dots x_{t-1} D_{t-1} x_t$
voloshin-hypergraph_ EQ0128	168	■ 1.000	$\alpha(\mathcal{H}) + \tau(\mathcal{H}) = X $.	$\alpha(\mathcal{H}) + \tau(\mathcal{H}) = X $.	$\alpha(\mathcal{H}) + \tau(\mathcal{H}) = X $.
voloshin-hypergraph_ EQ0129 (7.2)	168	■ 1.000	$\tau(\mathcal{H}) \geq v(\mathcal{H})$.	$\tau(\mathcal{H}) \geq v(\mathcal{H})$.	$\tau(\mathcal{H}) \geq v(\mathcal{H})$.
voloshin-hypergraph_ EQ0131	169	■ 1.000	$\rho(\mathcal{H}) = \tau(\mathcal{H}^*)$.	$\rho(\mathcal{H}) = \tau(\mathcal{H}^*)$.	$\rho(\mathcal{H}) = \tau(\mathcal{H}^*)$.
voloshin-hypergraph_ EQ0134	170	■ 1.000	$(\mathcal{H})_2 = L(\mathcal{H}^*)$.	$(\mathcal{H})_2 = L(\mathcal{H}^*)$.	$(\mathcal{H})_2 = L(\mathcal{H}^*)$.
voloshin-hypergraph_ EQ0135	170	■ 1.000	$L(\mathcal{H}) = (\mathcal{H}^*)_2$.	$L(\mathcal{H}) = (\mathcal{H}^*)_2$.	$L(\mathcal{H}) = (\mathcal{H}^*)_2$.
voloshin-hypergraph_ EQ0139	173	■ 1.000	$Y = (Y_{\{1\}} \cap Y_{\{2\}}) \cup (Y_{\{1\}} \cap Y_{\{3\}}) \cup (Y_{\{2\}} \cap Y_{\{3\}})$.	$Y = (Y_1 \cap Y_2) \cup (Y_1 \cap Y_3) \cup (Y_2 \cap Y_3)$.	$Y = (Y_1 \cap Y_2) \cup (Y_1 \cap Y_3) \cup (Y_2 \cap Y_3)$.
voloshin-hypergraph_ EQ0140	173	■ 1.000	$Y \subseteq (D_{\{1\}} \cap D_{\{2\}}) \cup (D_{\{1\}} \cap D_{\{3\}}) \cup (D_{\{2\}} \cap D_{\{3\}})$.	$Y \subseteq (D_1 \cap D_2) \cup (D_1 \cap D_3) \cup (D_2 \cap D_3)$.	$Y \subseteq (D_1 \cap D_2) \cup (D_1 \cap D_3) \cup (D_2 \cap D_3)$.
voloshin-hypergraph_ EQ0142 (8.2)	190	■ 1.000	$w(T) \leq \sum_{D \in \mathcal{D}} (D - 1)$.	$w(T) \leq \sum_{D \in \mathcal{D}} (D - 1)$.	$w(T) \leq \sum_{D \in \mathcal{D}} (D - 1)$.
voloshin-hypergraph_ EQ0147 (8.5)	192	■ 1.000	$c(T, D) = D - E_D $.	$c(T, D) = D - E_D $.	$c(T, D) = D - E_D $.
voloshin-hypergraph_ EQ0148	192	■ 1.000	$w(T) = \sum_{ D \geq 2} E_D $.	$w(T) = \sum_{ D \geq 2} E_D $.	$w(T) = \sum_{ D \geq 2} E_D $.
voloshin-hypergraph_ EQ0151	193	■ 1.000	$w(T) = \sum_{D \in \mathcal{D}} (D - 1)$.	$w(T) = \sum_{D \in \mathcal{D}} (D - 1)$.	$w(T) = \sum_{D \in \mathcal{D}} (D - 1)$.
voloshin-hypergraph_ EQ0157	194	■ 1.000	$w(T_1) = w(T) - \mathcal{D}_T(x) $.	$w(T_1) = w(T) - \mathcal{D}_T(x) $.	$w(T_1) = w(T) - \mathcal{D}_T(x) $.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0158	194	■ 1.000	$w(\mathcal{T}^{\prime})=w(\mathcal{T}_1)+ \mathcal{D}_T(x) \geq w(\mathcal{T}_1)+1+ \mathcal{D}_T(x) =w(T)+1$	$w(T') = w(T_1) + \mathcal{D}_T(x) \geq w(T_1) + 1 + \mathcal{D}_T(x) = w(T) + 1$	$w(T') = w(T_1) + \mathcal{D}_T(x) \geq w(T_1) + 1 + \mathcal{D}_T(x) = w(T) + 1$
voloshin-hypergraph_ EQ0159	194	■ 1.000	$w(\mathcal{H})=\max_{\mathcal{T}} w(T) \text{ .}$	$w(\mathcal{H}) = \max_T w(T).$	$w(\mathcal{H}) = \max_T w(T).$
voloshin-hypergraph_ EQ0163	197	■ 1.000	$\mu=x_{\{0\}} D_{\{0\}} x_{\{1\}} D_{\{1\}} x_{\{2\}} \ldots x_{\{t-1\}} D_{\{t-1\}} x_{\{t\}}$	$\mu = x_0 D_0 x_1 D_1 x_2 \dots x_{t-1} D_{t-1} x_t$	$\mu = x_0 D_0 x_1 D_1 x_2 \dots x_{t-1} D_{t-1} x_t$
voloshin-hypergraph_ EQ0164	201	■ 1.000	$\chi^{\prime}(\mathcal{H}) (\mathcal{H}) \geq \omega(L(\mathcal{H})) \geq \Delta(\mathcal{H}) \text{ .}$	$\chi'(\mathcal{H}) \geq \omega(L(\mathcal{H})) \geq \Delta(\mathcal{H}).$	$\chi'(\mathcal{H}) \geq \omega(L(\mathcal{H})) \geq \Delta(\mathcal{H}).$
voloshin-hypergraph_ EQ0165	201	■ 1.000	$\chi^{\prime}(\mathcal{H}^{\prime}) (\mathcal{H}^{\prime}) = \Delta(\mathcal{H}^{\prime}) \text{ .}$	$\chi'(\mathcal{H}') = \Delta(\mathcal{H}').$	$\chi'(\mathcal{H}') = \Delta(\mathcal{H}').$
voloshin-hypergraph_ EQ0166	201	■ 1.000	$\chi^{\prime}(\mathcal{H}) (\mathcal{H}) = \chi(L(\mathcal{H})) = \omega(L(\mathcal{H})) = \Delta(\mathcal{H}) \text{ .}$	$\chi'(\mathcal{H}) = \chi(L(\mathcal{H})) = \omega(L(\mathcal{H})) = \Delta(\mathcal{H}).$	$\chi'(\mathcal{H}) = \chi(L(\mathcal{H})) = \omega(L(\mathcal{H})) = \Delta(\mathcal{H}).$
voloshin-hypergraph_ EQ0167	204	■ 1.000	$n-\sum_{i=1}^m \left(\left \mathcal{E}_i\right -1\right)+f=m-\sum_{j=1}^n \left(\left \mathcal{D}\left(x_j\right)\right -1\right)+f=2 \text{ .}$	$n-\sum_{i=1}^m (E_i -1)+f=m-\sum_{j=1}^n (\mathcal{D}(x_j) -1)+f=2.$	$n-\sum_{i=1}^m (E_i -1)+f=m-\sum_{j=1}^n (\mathcal{D}(x_j) -1)+f=2.$
voloshin-hypergraph_ EQ0168 (9.1)	204	■ 1.000	$m^{\prime}(\mathcal{H})=\sum_{j=1}^m \left \mathcal{D}_j\right =\sum_{i=1}^n \left \mathcal{D}\left(x_i\right)\right $	$m' = \sum_{j=1}^m D_j = \sum_{i=1}^n \mathcal{D}(x_i) $	$m' = \sum_{j=1}^m D_j = \sum_{i=1}^n \mathcal{D}(x_i) $
voloshin-hypergraph_ EQ0169	205	■ 1.000	$n^{\prime}(\mathcal{H})-m^{\prime}(\mathcal{H})+f^{\prime}(\mathcal{H})=2$	$n' - m' + f' = 2$	$n' - m' + f' = 2.$
voloshin-hypergraph_ EQ0170 (9.2)	205	■ 1.000	$n-\sum_{i=1}^m \left(\left \mathcal{D}_i\right -1\right)+f=2$	$n-\sum_{i=1}^m (D_i -1)+f=2$	$n-\sum_{i=1}^m (D_i -1)+f=2,$
voloshin-hypergraph_ EQ0171 (9.3)	205	■ 1.000	$m-\sum_{j=1}^n \left(\left \mathcal{D}\left(x_j\right)\right -1\right)+f=2$	$m-\sum_{j=1}^n (\mathcal{D}(x_j) -1)+f=2$	$m-\sum_{j=1}^n (\mathcal{D}(x_j) -1)+f=2.$
voloshin-hypergraph_ EQ0172	205	■ 1.000	$n-\sum_{i=1}^m \left(\left \mathcal{E}_i\right -1\right)+f=4-(2+2)+2=2$	$n-\sum_{i=1}^m (E_i -1)+f=4-(2+2)+2=2$	$n-\sum_{i=1}^m (E_i -1)+f=4-(2+2)+2=2$
voloshin-hypergraph_ EQ0173	205	■ 1.000	$\left(m-\sum_{j=1}^n \left(\left \mathcal{D}\left(x_j\right)\right -1\right)+f=2-(0+1+0+1)+2\right)=2 \text{ .}$	$m-\sum_{j=1}^n (\mathcal{D}(x_j) -1)+f=2-(0+1+0+1)+2 \Bigg) = 2.$	$m-\sum_{j=1}^n (\mathcal{D}(x_j) -1)+f=2-(0+1+0+1)+2)=2.$
voloshin-hypergraph_ EQ0174 (10.1)	209	■ 1.000	$X=\cup_{i=1}^k S_i, \quad S_i \neq \emptyset, \quad S_i \cap S_j = \emptyset, i \neq j. \quad S_i \cap S_j = \emptyset, i \neq j.$	$X = \cup_{i=1}^k S_i, \quad S_i \neq \emptyset, \quad S_i \cap S_j = \emptyset, i \neq j.$	$X = \cup_{i=1}^k S_i, \quad S_i \neq \emptyset, \quad S_i \cap S_j = \emptyset, i \neq j.$
voloshin-hypergraph_ EQ0175	210	■ 1.000	$\begin{gathered} \alpha(\mathcal{H}) \chi(\mathcal{H}) \geq n; \\ \chi(\mathcal{H}) \leq \tau(\mathcal{H}) + 1 = n - \alpha(\mathcal{H}) + 1. \end{gathered}$	$\alpha(\mathcal{H}) \chi(\mathcal{H}) \geq n; \\ \chi(\mathcal{H}) \leq \tau(\mathcal{H}) + 1 = n - \alpha(\mathcal{H}) + 1.$	$\alpha(\mathcal{H}) \chi(\mathcal{H}) \geq n; \\ \chi(\mathcal{H}) \leq \tau(\mathcal{H}) + 1 = n - \alpha(\mathcal{H}) + 1.$
voloshin-hypergraph_ EQ0177	211	■ 1.000	$\min \{ D , \lambda \}$	$\min\{ D , \lambda\}$	$\min\{ D , \lambda\}$
voloshin-hypergraph_ EQ0178	211	■ 1.000	$\lambda \leq \min_{D \in \mathcal{D}} D ,$	$\lambda \leq \min_{D \in \mathcal{D}} D ,$	$\lambda \leq \min_{D \in \mathcal{D}} D ,$
voloshin-hypergraph_ EQ0181	211	■ 1.000	$a_j \leq \left \mathcal{D}_j \cap S_i \right \leq b_j \text{ .}$	$a_j \leq D_j \cap S_i \leq b_j.$	$a_j \leq D_j \cap S_i \leq b_j.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
voloshin-hypergraph_ EQ0225	244	■ 1.000	$R\left(\mathrm{mathcal{H}}_{\{1\}}\right)=R\left(\mathrm{mathcal{H}}_{\{2\}}\right)=(0,1,0,0),\ P\left(\mathrm{mathcal{H}}_{\{1\}},\ \lambda\right)=P\left(\mathrm{mathcal{H}}_{\{2\}},\ \lambda\right)=\lambda(\lambda-1)$.	$R(\mathcal{H}_1) = R(\mathcal{H}_2) = (0, 1, 0, 0), P(\mathcal{H}_1, \lambda) = P(\mathcal{H}_2, \lambda) = \lambda(\lambda - 1).$	$R(\mathcal{H}_1) = R(\mathcal{H}_2) = (0, 1, 0, 0), P(\mathcal{H}_1, \lambda) = P(\mathcal{H}_2, \lambda) = \lambda(\lambda - 1).$
voloshin-hypergraph_ EQ0227	248	■ 1.000	$L_{\{1\}}=\left(X_{\{1\}},\ E_{\{1\}}\right),\ L_{\{2\}}=\left(X_{\{2\}},\ E_{\{2\}}\right),\ \ldots,\ L_{\{\chi\}}=\left(X_{\{\chi\}},\ E_{\{\chi\}}\right)$.	$L_1 = (X_1, E_1), L_2 = (X_2, E_2), \dots, L_\chi = (X_\chi, E_\chi).$	$L_1 = (X_1, E_1), L_2 = (X_2, E_2), \dots, L_\chi = (X_\chi, E_\chi).$
voloshin-hypergraph_ EQ0230	249	■ 1.000	$\left\{x_{\{1\}},\ x_{\{2\}},\ \ldots,\ x_{\{i\}}\right\},\ i=1,2,\ \ldots,\ n$	$\{x_1, x_2, \dots, x_i\}, i = 1, 2, \dots, n$	$\{x_1, x_2, \dots, x_i\}, i = 1, 2, \dots, n$
voloshin-hypergraph_ EQ0234	253	■ 1.000	$C_{\{i\}}\ \cap\ C_{\{j\}}=\{1\},\ \text{quad } i,\ j\ \in\ I=\{1,2,\ \ldots\ k+2\},\ \text{quad } i\ \not\equiv\ j$.	$C_i \cap C_j = \{1\},\ \ i, j \in I = \{1, 2, \dots k + 2\},\ \ i \neq j.$	$C_i \cap C_j = \{1\},\ \ i, j \in I = \{1, 2, \dots k + 2\},\ \ i \neq j.$
voloshin-hypergraph_ EQ0236	255	■ 1.000	$\begin{aligned} &\& n_{\{12\}}+n_{\{23\}}+n_{\{34\}}+2<r, \\ &\& n_{\{12\}}+n_{\{14\}}+n_{\{34\}}+2<r \end{aligned}$.	$\begin{aligned} &n_{12} + n_{23} + n_{34} + 2 < r, \\ &n_{12} + n_{14} + n_{34} + 2 < r. \end{aligned}$	$\begin{aligned} &n_{12} + n_{23} + n_{34} + 2 < r, \\ &n_{12} + n_{14} + n_{34} + 2 < r. \end{aligned}$
voloshin-hypergraph_ EQ0238	256	■ 1.000	$n_{\{31\}}+1+n_{\{12\}}<r$.	$n_{31} + 1 + n_{12} < r.$	$n_{31} + 1 + n_{12} < r.$
voloshin-hypergraph_ EQ0239	256	■ 1.000	$X=\{1,2,3,\ \ldots,\ 2\ r\},$	$X = \{1, 2, 3, \dots, 2r\},$	$X = \{1, 2, 3, \dots, 2r\},$
voloshin-hypergraph_ EQ0240	256	■ 1.000	$\mathrm{mathcal{C}}=\left\{C_{\{1\}},\ C_{\{2\}},\ \ldots,\ C_{\{2\ r\}},\ C_{\{o\}},\ C_{\{e\}}\right\}$	$\mathcal{C} = \{C_1, C_2, \dots, C_{2r}, C_o, C_e\}$	$\mathcal{C} = \{C_1, C_2, \dots, C_{2r}, C_o, C_e\}$
voloshin-hypergraph_ EQ0241	256	■ 1.000	$C_{\{1\}}=\{1,2,4,6,8,\ \ldots,\ 2\ r-2\}$.	$C_1 = \{1, 2, 4, 6, 8, \dots, 2r - 2\}.$	$C_1 = \{1, 2, 4, 6, 8, \dots, 2r - 2\}.$
voloshin-hypergraph_ EQ0247	260	■ 1.000	$R(\mathrm{mathcal{H}})=(0,7,0,0)$	$R(\mathcal{H}) = (0, 7, 0, 0)$	$R(\mathcal{H}) = (0, 7, 0, 0)$
voloshin-hypergraph_ EQ0249	261	■ 1.000	$R(\mathrm{mathcal{H}})=\left(0,\ r_{\{2\}},\ 0,1,0,0,0,0\right)$.	$R(\mathcal{H}) = (0, r_2, 0, 1, 0, 0, 0).$	$R(\mathcal{H}) = (0, r_2, 0, 1, 0, 0, 0).$
voloshin-hypergraph_ EQ0250	263	■ 1.000	$\left\{x_{\{1\}},\ x_{\{2\}},\ a_{\{1\}},\ \ldots,\ a_{\{t-2\}},\ b_{\{1\}},\ \ldots,\ b_{\{t-2\}}\right\}$.	$\{x_1, x_2, a_1, \dots, a_{t-2}, b_1, \dots, b_{t-2}\}.$	$\{x_1, x_2, a_1, \dots, a_{t-2}, b_1, \dots, b_{t-2}\}.$
voloshin-hypergraph_ EQ0251	264	■ 1.000	$R\left(\mathrm{mathcal{H}}_{\{2,4\}}\right)=(0,4,0,1,0,0),$	$R(\mathcal{H}_{2,4}) = (0, 4, 0, 1, 0, 0),$	$R(\mathcal{H}_{2,4}) = (0, 4, 0, 1, 0, 0),$
voloshin-hypergraph_ EQ0252	264	■ 1.000	$P(\mathrm{mathcal{H}},\ \lambda)=\lambda(\lambda-1)\left(\lambda^2-5\lambda+10\right)$.	$P(\mathcal{H}, \lambda) = \lambda(\lambda - 1) (\lambda^2 - 5\lambda + 10).$	$P(\mathcal{H}, \lambda) = \lambda(\lambda - 1) (\lambda^2 - 5\lambda + 10).$
voloshin-hypergraph_ EQ0253	266	■ 1.000	$R(\mathrm{mathcal{H}})=\left(0,\ \ldots,\ 0,\ r_{\{\chi\}},\ \ldots,\ r_{\{\bar{\chi}\}},\ 0,\ \ldots,\ 0\right)$.	$R(\mathcal{H}) = (0, \dots, 0, r_\chi, \dots, r_{\bar{\chi}}, 0, \dots, 0).$	$R(\mathcal{H}) = (0, \dots, 0, r_\chi, \dots, r_{\bar{\chi}}, 0, \dots, 0).$
voloshin-hypergraph_ EQ0255	268	■ 1.000	$(A,\ A,\ B,\ B,\ C,\ C,\ D)$.	$(A, A, B, B, C, C, D).$	$(A, A, B, B, C, C, D).$
voloshin-hypergraph_ EQ0256	268	■ 1.000	$\{1,2,7\},\{3,4,7\},\{5,6,7\}\ \in\ \mathrm{mathcal{E}}$.	$\{1, 2, 7\}, \{3, 4, 7\}, \{5, 6, 7\} \in \mathcal{E}.$	$\{1, 2, 7\}, \{3, 4, 7\}, \{5, 6, 7\} \in \mathcal{E}.$
voloshin-hypergraph_ EQ0260	274	■ 1.000	$\mathrm{mathcal{H}}=\left(X,\ \binom{X}{3},\ \binom{X}{3}\right)$	$\mathcal{H} = \left(X, \binom{X}{3}, \binom{X}{3}\right)$	$\mathcal{H} = (X, \binom{X}{3}, \binom{X}{3})$
voloshin-hypergraph_ EQ0262	276	■ 1.000	$R\left(\mathrm{mathcal{H}}_{\{2,4\}}^{\prime}\right)=(0,1,0,1,0,0),$	$R(\mathcal{H}_{2,4}') = (0, 1, 0, 1, 0, 0),$	$R(\mathcal{H}_{2,4}') = (0, 1, 0, 1, 0, 0),$
voloshin-hypergraph_ EQ0263	276	■ 1.000	$P\left(\mathrm{mathcal{H}}_{\{2,4\}}^{\prime},\ \lambda\right)=\lambda(\lambda-1)\left(\lambda^2-5\lambda+7\right)$	$P(\mathcal{H}_{2,4}', \lambda) = \lambda(\lambda - 1) (\lambda^2 - 5\lambda + 7)$	$P(\mathcal{H}_{2,4}', \lambda) = \lambda(\lambda - 1) (\lambda^2 - 5\lambda + 7)$
voloshin-hypergraph_ EQ0264	285	■ 1.000	$f(n)\ \leq\ C\ g(n)$	$f(n) \leq Cg(n)$	$f(n) \leq Cg(n)$