

0707.4470 — formula report

213 inline formulas (section omitted; all rendered), 62 display equations, 0 tables, 12 unrecovered image regions.

Corrected (1)

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0707.4470_F00175 (was)	29	—	\mathscr{g}	<i>g</i>	—
0707.4470_F00175 (now)	29	—	\mathscr{J}	<i>J</i>	—
basis: source / source+ink					
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value					
Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Flagged, not acted on — 1 of 10 shown

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0707.4470_ E00035 ● C +22	020	■ 0.705	$\begin{aligned} & \frac{\left(D_x\right)_{k+\frac{1}{2}, l, m}^n - \left(D_x\right)_{k+\frac{1}{2}, l, m}^{n-\frac{1}{2}}}{\Delta t} = \\ & \frac{H_z\left _{k+\frac{1}{2}, l+\frac{1}{2}, m}^n - H_z\left _{k+\frac{1}{2}, l-\frac{1}{2}, m}^n}{\Delta y} - \frac{H_y\left _{k+\frac{1}{2}, l, m+\frac{1}{2}}^n - H_y\left _{k+\frac{1}{2}, l, m-\frac{1}{2}}^n}{\Delta z} - J_x\right _{k+\frac{1}{2}, l, m}^n \\ & \frac{D_y\left _{k, l+\frac{1}{2}, m}^{n+\frac{1}{2}} - D_y\left _{k, l+\frac{1}{2}, m}^{n-\frac{1}{2}}}{\Delta t} = \\ & \frac{H_x\left _{k, l+\frac{1}{2}, m+\frac{1}{2}}^n - H_x\left _{k, l+\frac{1}{2}, m-\frac{1}{2}}^n}{\Delta z} - \frac{H_z\left _{k+\frac{1}{2}, l+\frac{1}{2}, m}^n - H_z\left _{k-\frac{1}{2}, l+\frac{1}{2}, m}^n}{\Delta x} - J_y\right _{k, l+\frac{1}{2}, m}^n \\ & \frac{D_z\left _{k, l, m+\frac{1}{2}}^{n+\frac{1}{2}} - D_z\left _{k, l, m+\frac{1}{2}}^{n-\frac{1}{2}}}{\Delta t} = \\ & \frac{H_y\left _{k+\frac{1}{2}, l, m+\frac{1}{2}}^n - H_y\left _{k-\frac{1}{2}, l, m+\frac{1}{2}}^n}{\Delta x} - \frac{H_x\left _{k, l+\frac{1}{2}, m+\frac{1}{2}}^n - H_x\left _{k, l-\frac{1}{2}, m+\frac{1}{2}}^n}{\Delta y} - J_z\right _{k, l, m+\frac{1}{2}}^n \end{aligned}$	$\frac{D_x _{k+\frac{1}{2}, l, m}^{n+\frac{1}{2}} - D_x _{k+\frac{1}{2}, l, m}^{n-\frac{1}{2}}}{\Delta t} =$ $\frac{H_z _{k+\frac{1}{2}, l+\frac{1}{2}, m}^n - H_z _{k+\frac{1}{2}, l-\frac{1}{2}, m}^n}{\Delta y} - \frac{H_y _{k+\frac{1}{2}, l, m+\frac{1}{2}}^n - H_y _{k+\frac{1}{2}, l, m-\frac{1}{2}}^n}{\Delta z} - J_x _{k+\frac{1}{2}, l, m}^n$ $\frac{D_y _{k, l+\frac{1}{2}, m}^{n+\frac{1}{2}} - D_y _{k, l+\frac{1}{2}, m}^{n-\frac{1}{2}}}{\Delta t} =$ $\frac{H_x _{k, l+\frac{1}{2}, m+\frac{1}{2}}^n - H_x _{k, l+\frac{1}{2}, m-\frac{1}{2}}^n}{\Delta z} - \frac{H_z _{k+\frac{1}{2}, l+\frac{1}{2}, m}^n - H_z _{k-\frac{1}{2}, l+\frac{1}{2}, m}^n}{\Delta x} - J_y _{k, l+\frac{1}{2}, m}^n$ $\frac{D_z _{k, l, m+\frac{1}{2}}^{n+\frac{1}{2}} - D_z _{k, l, m+\frac{1}{2}}^{n-\frac{1}{2}}}{\Delta t} =$ $\frac{H_y _{k+\frac{1}{2}, l, m+\frac{1}{2}}^n - H_y _{k-\frac{1}{2}, l, m+\frac{1}{2}}^n}{\Delta x} - \frac{H_x _{k, l+\frac{1}{2}, m+\frac{1}{2}}^n - H_x _{k, l-\frac{1}{2}, m+\frac{1}{2}}^n}{\Delta y} - J_z _{k, l, m+\frac{1}{2}}^n$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

The remaining 9 flagged rows are stated as a count.

Each differs from its scan by fewer than 20 components, which is the tail of the distribution rather than its bulk: 0 are component (C), 9 weak (W). By MathPix confidence: 7 at ≥0.9, 1 in 0.6-0.9, 1 below 0.6, 0 with none. Out/465 sampled twenty rows drawn from the ≥0.9 band corpus-wide and found eighteen purely typographic differences, one typographic plus a crop overrun, one borderline and *no* unambiguous content errors — so a high-confidence flag here is evidence of a rendering difference, not of a defect. Every row is in report.ink.json.