

mielke-geometrodynamics: residuals

equations measured 2026-09-14T14:45:38Z against the model of 2026-09-14

Corrected (36)

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_E00120 (was)	60	■ 0.651	$d\,\Sigma_{\alpha}\simeq\left(e_{\alpha}\downharpoonleft F\right)\wedge j$	$d\Sigma_{\alpha}\simeq(e_{\alpha}\downharpoonleft F)\wedge j$	$d\Sigma_{\alpha}\simeq(e_{\alpha}\downharpoonleft F)\wedge j$
mielke-geometrodynamics_E00120 (now)	60	—	$d\,\Sigma_{\alpha}\simeq\left(e_{\alpha}\rfloor F\right)\wedge j$	$d\Sigma_{\alpha}\simeq(e_{\alpha}\rfloor F)\wedge j$	$d\Sigma_{\alpha}\simeq(e_{\alpha}\rfloor F)\wedge j$
basis: census / census					
mielke-geometrodynamics_E00292 (was)	100	■ 0.165	$\begin{aligned} \mathfrak{f}_{\xi} L &= \left(\mathfrak{f}_{\xi} g_{\alpha\beta} \right) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \left(\mathfrak{f}_{\xi} \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} \\ &+ \left(\mathfrak{f}_{\xi} \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\mathfrak{f}_{\xi} D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \\ &= D \left[\left(\xi \downharpoonleft \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \downharpoonleft \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \downharpoonleft D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \right] \\ &- \left(\xi \downharpoonleft \vartheta^{\alpha} \right) D \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \downharpoonleft T^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \downharpoonleft R^{\gamma}_{\beta} \right) \wedge I^{\beta}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} \\ &+ \left(\xi \downharpoonleft D\Psi \right) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p \left(\xi \downharpoonleft \Psi \right) \wedge D \frac{\delta L}{\delta \Psi} . \end{aligned}$	$\begin{aligned} \mathfrak{f}_{\xi} L &= \left(\mathfrak{f}_{\xi} g_{\alpha\beta} \right) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \left(\mathfrak{f}_{\xi} \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} \\ &+ \left(\mathfrak{f}_{\xi} \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\mathfrak{f}_{\xi} D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \\ &= D \left[\left(\xi \downharpoonleft \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \downharpoonleft \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \downharpoonleft D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \right] \\ &- \left(\xi \downharpoonleft \vartheta^{\alpha} \right) D \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \downharpoonleft T^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \downharpoonleft R^{\gamma}_{\beta} \right) \wedge I^{\beta}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} \\ &+ \left(\xi \downharpoonleft D\Psi \right) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p \left(\xi \downharpoonleft \Psi \right) \wedge D \frac{\delta L}{\delta \Psi} . \end{aligned}$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: basis] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_E00292 (now)	100	—	$\begin{aligned} \mathfrak{f}_{\xi} L &= \left(\mathfrak{f}_{\xi} g_{\alpha\beta} \right) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \left(\mathfrak{f}_{\xi} \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} \\ &+ \left(\mathfrak{f}_{\xi} \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\mathfrak{f}_{\xi} D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \\ &= D \left[\left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \right] \\ &- \left(\xi \rfloor \vartheta^{\alpha} \right) D \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor T^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor R_{\beta}^{\gamma} \right) \wedge I^{\beta}{}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} \\ &+ \left(\xi \rfloor D\Psi \right) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p \left(\xi \rfloor \Psi \right) \wedge D \frac{\delta L}{\delta \Psi} . \end{aligned}$	$\begin{aligned} \mathfrak{f}_{\xi} L &= \left(\mathfrak{f}_{\xi} g_{\alpha\beta} \right) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \left(\mathfrak{f}_{\xi} \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} \\ &+ \left(\mathfrak{f}_{\xi} \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\mathfrak{f}_{\xi} D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \\ &= D \left[\left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \right] \\ &- \left(\xi \rfloor \vartheta^{\alpha} \right) D \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor T^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor R_{\beta}^{\gamma} \right) \wedge I^{\beta}{}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} \\ &+ \left(\xi \rfloor D\Psi \right) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p \left(\xi \rfloor \Psi \right) \wedge D \frac{\delta L}{\delta \Psi} . \end{aligned}$	
basis: census / census					
mielke-geometrodynamics_E00293 (was)	101	■0.052	$\xi \downharpoonleft L \equiv \left(\xi \downharpoonleft \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \downharpoonleft \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \downharpoonleft D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} .$	$\xi \downharpoonleft L \equiv \left(\xi \downharpoonleft \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \downharpoonleft \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \downharpoonleft D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} .$	
mielke-geometrodynamics_E00293 (now)	101	—	$\xi \rfloor L \equiv \left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} .$	$\xi \rfloor L \equiv \left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} .$	
basis: census / census					
mielke-geometrodynamics_E00294 (was)	101	■0.510	$\Sigma_{\alpha} = e_{\alpha} \downharpoonleft L - \left(e_{\alpha} \downharpoonleft D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} - \left(e_{\alpha} \downharpoonleft \Psi \right) \wedge \frac{\partial L}{\partial \Psi} .$	$\Sigma_{\alpha} = e_{\alpha} \downharpoonleft L - \left(e_{\alpha} \downharpoonleft D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} - \left(e_{\alpha} \downharpoonleft \Psi \right) \wedge \frac{\partial L}{\partial \Psi} .$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_101 EQ0294 (now)	101	—	$\left.\Sigma_{\alpha}=\mathrm{e}_{\alpha}\right\rfloor L-\left(\mathrm{e}_{\alpha}\right\rfloor D\,\Psi\right)\wedge\frac{\partial L}{\partial D\,\Psi}-\left(\mathrm{e}_{\alpha}\right\rfloor\Psi\right)\wedge\frac{\partial L}{\partial\Psi}.$	$\Sigma_{\alpha} = e_{\alpha}\rfloor L - (e_{\alpha}\rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha}\rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$	$\Sigma_{\alpha} = e_{\alpha}\rfloor L - (e_{\alpha}\rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha}\rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$
basis: census / census					
mielke-geometrodynamics_101 EQ0296 (was)	101	■ 0.847	$F_{\alpha}=\left(\mathrm{e}_{\alpha}\right\downarrow D\,\Psi\right)\frac{\delta L}{\delta\Psi}+(-1)^p\left(\mathrm{e}_{\alpha}\right\downarrow\Psi\right)\wedge D\frac{\delta L}{\delta\Psi}$	$F_{\alpha} = (e_{\alpha}\downarrow D\Psi) \frac{\delta L}{\delta\Psi} + (-1)^p (e_{\alpha}\downarrow \Psi) \wedge D \frac{\delta L}{\delta\Psi}$	$F_{\alpha} = (e_{\alpha}\rfloor D\Psi) \frac{\delta L}{\delta\Psi} + (-1)^p (e_{\alpha}\rfloor \Psi) \wedge D \frac{\delta L}{\delta\Psi}$
mielke-geometrodynamics_101 EQ0296 (now)	101	—	$F_{\alpha}=\left(\mathrm{e}_{\alpha}\right\rfloor D\,\Psi\right)\frac{\delta L}{\delta\Psi}+(-1)^p\left(\mathrm{e}_{\alpha}\right\rfloor\Psi\right)\wedge D\frac{\delta L}{\delta\Psi}$	$F_{\alpha} = (e_{\alpha}\rfloor D\Psi) \frac{\delta L}{\delta\Psi} + (-1)^p (e_{\alpha}\rfloor \Psi) \wedge D \frac{\delta L}{\delta\Psi}$	$F_{\alpha} = (e_{\alpha}\rfloor D\Psi) \frac{\delta L}{\delta\Psi} + (-1)^p (e_{\alpha}\rfloor \Psi) \wedge D \frac{\delta L}{\delta\Psi}$
basis: census / census					
mielke-geometrodynamics_103 EQ0305 (was)	103	■ 0.793	$\varepsilon_{\mathrm{RC}}:=\xi^{\alpha}\Sigma_{\alpha}+\left(\mathrm{e}_{\beta}\right\downarrow\widehat{D}\xi^{\gamma}\right)\tau^{\beta}_{\gamma},$	$\varepsilon_{\mathrm{RC}} := \xi^{\alpha}\Sigma_{\alpha} + \left(e_{\beta}\downarrow \widehat{D}\xi^{\gamma}\right) \tau^{\beta}_{\gamma},$	$\varepsilon_{\mathrm{RC}} := \xi^{\alpha}\Sigma_{\alpha} + (e_{\beta}\rfloor \widehat{D}\xi^{\gamma}) \tau^{\beta}_{\gamma},$
mielke-geometrodynamics_103 EQ0305 (now)	103	—	$\varepsilon_{\mathrm{RC}}:=\xi^{\alpha}\Sigma_{\alpha}+\left(\mathrm{e}_{\beta}\right\rfloor\widehat{D}\xi^{\gamma}\right)\tau^{\beta}_{\gamma},$	$\varepsilon_{\mathrm{RC}} := \xi^{\alpha}\Sigma_{\alpha} + \left(e_{\beta}\rfloor \widehat{D}\xi^{\gamma}\right) \tau^{\beta}_{\gamma},$	$\varepsilon_{\mathrm{RC}} := \xi^{\alpha}\Sigma_{\alpha} + (e_{\beta}\rfloor \widehat{D}\xi^{\gamma}) \tau^{\beta}_{\gamma},$
basis: census / census					
mielke-geometrodynamics_103 EQ0306 (was)	103	■ 0.406	$\left.\widetilde{\varepsilon}_{\mathrm{MA}}:=\varepsilon_{\mathrm{RC}}+\xi\right\right\lrcorner L\cong-\left(\mathscr{L}_{\xi}\Gamma_{\beta}^{\gamma}\right)\wedge H^{\beta}_{\gamma}+dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}} := \varepsilon_{\mathrm{RC}} + \xi\lrcorner L \cong - \left(\mathcal{L}_{\xi}\Gamma_{\beta}^{\gamma}\right) \wedge H^{\beta}_{\gamma} + dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}} := \varepsilon_{\mathrm{RC}} + \xi\rfloor L \cong - \left(\mathcal{L}_{\xi}\Gamma_{\beta}^{\gamma}\right) \wedge H^{\beta}_{\gamma} + dH_{\infty}$
mielke-geometrodynamics_103 EQ0306 (now)	103	—	$\left.\widetilde{\varepsilon}_{\mathrm{MA}}:=\varepsilon_{\mathrm{RC}}+\xi\right\rfloor L\cong-\left(\mathscr{L}_{\xi}\Gamma_{\beta}^{\gamma}\right)\wedge H^{\beta}_{\gamma}+dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}} := \varepsilon_{\mathrm{RC}} + \xi\rfloor L \cong - \left(\mathcal{L}_{\xi}\Gamma_{\beta}^{\gamma}\right) \wedge H^{\beta}_{\gamma} + dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}} := \varepsilon_{\mathrm{RC}} + \xi\rfloor L \cong - \left(\mathcal{L}_{\xi}\Gamma_{\beta}^{\gamma}\right) \wedge H^{\beta}_{\gamma} + dH_{\infty}$
basis: census / census					
mielke-geometrodynamics_104 EQ0311 (was)	104	■ 0.111	$\begin{aligned} \widetilde{\leftarrow}_{\widetilde{y}}\widetilde{D}\widetilde{\xi}^{\alpha} &= \widetilde{y}\rfloor\left(\widetilde{D}\widetilde{D}\widetilde{\xi}^{\alpha}\right)+\widetilde{D}\left(\widetilde{y}\rfloor\widetilde{D}\widetilde{\xi}^{\alpha}\right) \\ &= yJ\left(DD\xi^{\alpha}+R^{(T)\alpha}\right)=0, \end{aligned}$	$\widetilde{\leftarrow}_{\widetilde{y}}\widetilde{D}\widetilde{\xi}^{\alpha} = \widetilde{y}\rfloor\left(\widetilde{D}\widetilde{D}\widetilde{\xi}^{\alpha}\right)+\widetilde{D}\left(\widetilde{y}\rfloor\widetilde{D}\widetilde{\xi}^{\alpha}\right) \\ = y\rfloor(DD\xi^{\alpha}+R^{(T)\alpha}) = 0,$	$\widetilde{\leftarrow}_{\widetilde{y}}\widetilde{D}\widetilde{\xi}^{\alpha} = \widetilde{y}\rfloor\left(\widetilde{D}\widetilde{D}\widetilde{\xi}^{\alpha}\right)+\widetilde{D}\left(\widetilde{y}\rfloor\widetilde{D}\widetilde{\xi}^{\alpha}\right) \\ = y\rfloor(DD\xi^{\alpha}+R^{(T)\alpha}) = 0,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_104E08311 (now)	104	—	$\begin{aligned} & \widetilde{\leftarrow}_{\widetilde{y}} \widetilde{\widetilde{D}} \widetilde{\xi}^\alpha = \widetilde{y} \rfloor \left(\widetilde{\widetilde{D}} \widetilde{\widetilde{D}} \widetilde{\xi}^\alpha \right) + \widetilde{\widetilde{D}} (\widetilde{y} \rfloor \widetilde{\widetilde{D}} \widetilde{\xi}^\alpha) \\ & = y \rfloor \left(DD \xi^\alpha + R^{(T)\alpha} \right) = 0, \end{aligned}$	$\begin{aligned} \widetilde{\widetilde{\mathbf{L}}}_{\widetilde{y}} \widetilde{\widetilde{D}} \widetilde{\xi}^\alpha &= \widetilde{y} \rfloor \left(\widetilde{\widetilde{D}} \widetilde{\widetilde{D}} \widetilde{\xi}^\alpha \right) + \widetilde{\widetilde{D}} (\widetilde{y} \rfloor \widetilde{\widetilde{D}} \widetilde{\xi}^\alpha) \\ &= y \rfloor (DD \xi^\alpha + R^{(T)\alpha}) = 0, \end{aligned}$	
basis: census / census					
mielke-geometrodynamics_128E08378 (was)	128	■ 0.136	$\begin{aligned} D D \vartheta^\alpha &= R_\beta^\alpha \wedge \vartheta^\beta = 0, \quad \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha = - \stackrel{(\pm)}{R}_\alpha{}^\beta \wedge \stackrel{(\pm)}{\Pi}_\beta = \pm \frac{i}{2} \ell^2 \left(e_\alpha \rfloor \stackrel{(\pm)}{\Pi}^\beta \right) \wedge \Sigma_\beta \cong 0. \\ \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha &= - \stackrel{(\pm)}{R}_\alpha{}^\beta \wedge \stackrel{(\pm)}{\Pi}_\beta = \pm \frac{i}{2} \ell^2 \left(e_\alpha \rfloor \stackrel{(\pm)}{\Pi}^\beta \right) \wedge \Sigma_\beta \cong 0. \end{aligned}$	$DD\vartheta^\alpha = R_\beta^\alpha \wedge \vartheta^\beta = 0, \quad \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha = - \stackrel{(\pm)}{R}_\alpha{}^\beta \wedge \stackrel{(\pm)}{\Pi}_\beta = \pm \frac{i}{2} \ell^2 \left(e_\alpha \rfloor \stackrel{(\pm)}{\Pi}^\beta \right) \wedge \Sigma_\beta \cong 0.$	$DD\vartheta^\alpha = R_\beta^\alpha \wedge \vartheta^\beta = 0, \quad \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha = - \stackrel{(\pm)}{R}_\alpha{}^\beta \wedge \stackrel{(\pm)}{\Pi}_\beta = \pm \frac{i}{2} \ell^2 (e_\alpha \rfloor \stackrel{(\pm)}{\Pi}^\beta) \wedge \Sigma_\beta \cong 0.$
mielke-geometrodynamics_128E08378 (now)	128	—	$\begin{aligned} D D \vartheta^\alpha &= R_\beta^\alpha \wedge \vartheta^\beta = 0, \quad \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha = - \stackrel{(\pm)}{R}_\alpha{}^\beta \wedge \stackrel{(\pm)}{\Pi}_\beta = \pm \frac{i}{2} \ell^2 \left(e_\alpha \rfloor \stackrel{(\pm)}{\Pi}^\beta \right) \wedge \Sigma_\beta \cong 0. \\ \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha &= - \stackrel{(\pm)}{R}_\alpha{}^\beta \wedge \stackrel{(\pm)}{\Pi}_\beta = \pm \frac{i}{2} \ell^2 \left(e_\alpha \rfloor \stackrel{(\pm)}{\Pi}^\beta \right) \wedge \Sigma_\beta \cong 0. \end{aligned}$	$DD\vartheta^\alpha = R_\beta^\alpha \wedge \vartheta^\beta = 0, \quad \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha = - \stackrel{(\pm)}{R}_\alpha{}^\beta \wedge \stackrel{(\pm)}{\Pi}_\beta = \pm \frac{i}{2} \ell^2 \left(e_\alpha \rfloor \stackrel{(\pm)}{\Pi}^\beta \right) \wedge \Sigma_\beta \cong 0.$	$DD\vartheta^\alpha = R_\beta^\alpha \wedge \vartheta^\beta = 0, \quad \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha = - \stackrel{(\pm)}{R}_\alpha{}^\beta \wedge \stackrel{(\pm)}{\Pi}_\beta = \pm \frac{i}{2} \ell^2 (e_\alpha \rfloor \stackrel{(\pm)}{\Pi}^\beta) \wedge \Sigma_\beta \cong 0.$
basis: census / census					
mielke-geometrodynamics_129E08381 (was)	129	■ 0.588	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha := D \stackrel{(\pm)}{\Pi}_\alpha \pm \frac{i}{2} \ell^2 \left(e_\alpha \rfloor \stackrel{(\mp)}{\Pi}^\beta \right) \wedge \stackrel{(\pm)}{\Pi}_\beta,$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha := D \stackrel{(\pm)}{\Pi}_\alpha \pm \frac{i}{2} \ell^2 \left(e_\alpha \rfloor \stackrel{(\mp)}{\Pi}^\beta \right) \wedge \stackrel{(\pm)}{\Pi}_\beta,$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha := D \stackrel{(\pm)}{\Pi}_\alpha \pm \frac{i}{2} \ell^2 (e_\alpha \rfloor \stackrel{(\mp)}{\Pi}^\beta) \wedge \stackrel{(\pm)}{\Pi}_\beta,$
mielke-geometrodynamics_129E08381 (now)	129	—	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha := D \stackrel{(\pm)}{\Pi}_\alpha \pm \frac{i}{2} \ell^2 \left(e_\alpha \rfloor \stackrel{(\mp)}{\Pi}^\beta \right) \wedge \stackrel{(\pm)}{\Pi}_\beta,$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha := D \stackrel{(\pm)}{\Pi}_\alpha \pm \frac{i}{2} \ell^2 \left(e_\alpha \rfloor \stackrel{(\mp)}{\Pi}^\beta \right) \wedge \stackrel{(\pm)}{\Pi}_\beta,$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_\alpha := D \stackrel{(\pm)}{\Pi}_\alpha \pm \frac{i}{2} \ell^2 (e_\alpha \rfloor \stackrel{(\mp)}{\Pi}^\beta) \wedge \stackrel{(\pm)}{\Pi}_\beta,$
basis: census / census					
mielke-geometrodynamics_129E08382 (was)	129	■ 0.087	$\begin{aligned} \stackrel{(\pm)}{\Gamma}_\alpha{}^\beta &:= \Gamma_\alpha{}^\beta + i A_\alpha{}^\beta, \quad A_\alpha{}^\beta := \pm \frac{1}{2} \ell^2 e_\alpha \rfloor \Pi^\beta, \\ \stackrel{(\pm)}{A}_\alpha{}^\beta &:= \pm \frac{1}{2} \ell^2 e_\alpha \rfloor \Pi^\beta, \end{aligned}$	$\stackrel{(\pm)}{\Gamma}_\alpha{}^\beta := \Gamma_\alpha{}^\beta + i A_\alpha{}^\beta, \quad A_\alpha{}^\beta := \pm \frac{1}{2} \ell^2 e_\alpha \rfloor \Pi^\beta,$	$\stackrel{(\pm)}{\Gamma}_\alpha{}^\beta := \Gamma_\alpha{}^\beta + i A_\alpha{}^\beta, \quad A_\alpha{}^\beta := \pm \frac{1}{2} \ell^2 e_\alpha \rfloor \Pi^\beta,$
mielke-geometrodynamics_129E08382 (now)	129	—	$\begin{aligned} \stackrel{(\pm)}{\Gamma}_\alpha{}^\beta &:= \Gamma_\alpha{}^\beta + i A_\alpha{}^\beta, \quad A_\alpha{}^\beta := \pm \frac{1}{2} \ell^2 e_\alpha \rfloor \Pi^\beta, \\ \stackrel{(\pm)}{A}_\alpha{}^\beta &:= \pm \frac{1}{2} \ell^2 e_\alpha \rfloor \Pi^\beta, \end{aligned}$	$\stackrel{(\pm)}{\Gamma}_\alpha{}^\beta := \Gamma_\alpha{}^\beta + i A_\alpha{}^\beta, \quad A_\alpha{}^\beta := \pm \frac{1}{2} \ell^2 e_\alpha \rfloor \Pi^\beta,$	$\stackrel{(\pm)}{\Gamma}_\alpha{}^\beta := \Gamma_\alpha{}^\beta + i A_\alpha{}^\beta, \quad A_\alpha{}^\beta := \pm \frac{1}{2} \ell^2 e_\alpha \rfloor \Pi^\beta,$
basis: census / census					
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_131 EQ0391 (was)	131	■ 0.066	$\begin{aligned} \Sigma_{\alpha} &:= \frac{\delta L}{\delta \vartheta^{\alpha}} = e_{\alpha} \rfloor L - (e_{\alpha} \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} \\ &\quad - (e_{\alpha} \rfloor T^{\beta}) \wedge \frac{\partial L}{\partial T^{\beta}} + D \frac{\partial L}{\partial T^{\alpha}} - (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \frac{\partial L}{\partial R_{\beta}^{\gamma}}. \end{aligned}$	$\Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = e_{\alpha} \rfloor L - (e_{\alpha} \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} \\ - (e_{\alpha} \rfloor T^{\beta}) \wedge \frac{\partial L}{\partial T^{\beta}} + D \frac{\partial L}{\partial T^{\alpha}} - (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \frac{\partial L}{\partial R_{\beta}^{\gamma}}.$	$\Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = e_{\alpha} \rfloor L - (e_{\alpha} \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} \\ - (e_{\alpha} \rfloor T^{\beta}) \wedge \frac{\partial L}{\partial T^{\beta}} + D \frac{\partial L}{\partial T^{\alpha}} - (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \frac{\partial L}{\partial R_{\beta}^{\gamma}}.$
mielke-geometrodynamics_131 EQ0391 (now)	131	—	$\begin{aligned} \Sigma_{\alpha} &:= \frac{\delta L}{\delta \vartheta^{\alpha}} = e_{\alpha} \rfloor L - (e_{\alpha} \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} \\ &\quad - (e_{\alpha} \rfloor T^{\beta}) \wedge \frac{\partial L}{\partial T^{\beta}} + D \frac{\partial L}{\partial T^{\alpha}} - (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \frac{\partial L}{\partial R_{\beta}^{\gamma}}. \end{aligned}$	$\Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = e_{\alpha} \rfloor L - (e_{\alpha} \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} \\ - (e_{\alpha} \rfloor T^{\beta}) \wedge \frac{\partial L}{\partial T^{\beta}} + D \frac{\partial L}{\partial T^{\alpha}} - (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \frac{\partial L}{\partial R_{\beta}^{\gamma}}.$	$\Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = e_{\alpha} \rfloor L - (e_{\alpha} \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} \\ - (e_{\alpha} \rfloor T^{\beta}) \wedge \frac{\partial L}{\partial T^{\beta}} + D \frac{\partial L}{\partial T^{\alpha}} - (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \frac{\partial L}{\partial R_{\beta}^{\gamma}}.$
basis: census / census					
mielke-geometrodynamics_132 EQ0392 (was)	132	■ 0.552	$\left. \left. n^{\alpha} \right. \Sigma_{\alpha} = n \rfloor L - (n \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (n \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} \right.$	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - (n \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (n \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - (n \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (n \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$
mielke-geometrodynamics_132 EQ0392 (now)	132	—	$\left. \left. n^{\alpha} \right. \Sigma_{\alpha} = n \rfloor L - (n \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (n \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} \right.$	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - (n \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (n \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - (n \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (n \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$
basis: census / census					
mielke-geometrodynamics_132 EQ0393 (was)	132	■ 0.382	$\begin{aligned} n^{\alpha} \Sigma_{\alpha} &= n \rfloor L - (t_n \Psi) \wedge \frac{\partial L}{\partial D\Psi} - (n \rfloor \Psi) \wedge \frac{\delta L}{\delta \Psi} + D[(n \rfloor \Psi) \wedge \frac{\partial L}{\partial D\Psi}] \\ &\cong n \rfloor L - \left(\ell_n \Psi + \Gamma_{\perp}^{\alpha\beta} \rho(I_{\alpha\beta}) \Psi \right) \wedge \frac{\partial L}{\partial D\Psi} + D[(n \rfloor \Psi) \wedge \frac{\partial L}{\partial D\Psi}]. \end{aligned}$	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - (t_n \Psi) \wedge \frac{\partial L}{\partial D\Psi} - (n \rfloor \Psi) \wedge \frac{\delta L}{\delta \Psi} + D[(n \rfloor \Psi) \wedge \frac{\partial L}{\partial D\Psi}] \\ \cong n \rfloor L - \left(\ell_n \Psi + \Gamma_{\perp}^{\alpha\beta} \rho(I_{\alpha\beta}) \Psi \right) \wedge \frac{\partial L}{\partial D\Psi} + D[(n \rfloor \Psi) \wedge \frac{\partial L}{\partial D\Psi}].$	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - (\mathbb{E}_n \Psi) \wedge \frac{\partial L}{\partial D\Psi} - (n \rfloor \Psi) \wedge \frac{\delta L}{\delta \Psi} + D[(n \rfloor \Psi) \wedge \frac{\partial L}{\partial D\Psi}] \\ \cong n \rfloor L - \left(\ell_n \Psi + \Gamma_{\perp}^{\alpha\beta} \rho(I_{\alpha\beta}) \Psi \right) \wedge \frac{\partial L}{\partial D\Psi} + D[(n \rfloor \Psi) \wedge \frac{\partial L}{\partial D\Psi}].$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: basis] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — basis names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamicsEQ0393 (now)	132	—	$\begin{aligned} n^{\alpha} \Sigma_{\alpha} &= n \rfloor L - (\mathfrak{L}_n \Psi) \wedge \frac{\partial L}{\partial D \Psi} - (n \rfloor \Psi) \wedge \frac{\delta L}{\delta \Psi} + D \Big[(n \rfloor \Psi) \wedge \frac{\partial L}{\partial D \Psi} \Big] \\ &\cong n \rfloor L - \Big(\ell_n \Psi + \Gamma_{\perp}^{\alpha \beta} \rho(I_{\alpha \beta}) \Psi \Big) \wedge \frac{\partial L}{\partial D \Psi} + D \Big[(n \rfloor \Psi) \wedge \frac{\partial L}{\partial D \Psi} \Big]. \end{aligned}$	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - (\mathfrak{L}_n \Psi) \wedge \frac{\partial L}{\partial D \Psi} - (n \rfloor \Psi) \wedge \frac{\delta L}{\delta \Psi} + D[(n \rfloor \Psi) \wedge \frac{\partial L}{\partial D \Psi}]$ $\cong n \rfloor L - \Big(\ell_n \Psi + \Gamma_{\perp}^{\alpha \beta} \rho(I_{\alpha \beta}) \Psi \Big) \wedge \frac{\partial L}{\partial D \Psi} + D[(n \rfloor \Psi) \wedge \frac{\partial L}{\partial D \Psi}].$	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - (\mathfrak{L}_n \Psi) \wedge \frac{\partial L}{\partial D \Psi} - (n \rfloor \Psi) \wedge \frac{\delta L}{\delta \Psi} + D \Big[(n \rfloor \Psi) \wedge \frac{\partial L}{\partial D \Psi} \Big]$ $\cong n \rfloor L - \Big(\ell_n \Psi + \Gamma_{\perp}^{\alpha \beta} \rho(I_{\alpha \beta}) \Psi \Big) \wedge \frac{\partial L}{\partial D \Psi} + D \Big[(n \rfloor \Psi) \wedge \frac{\partial L}{\partial D \Psi} \Big].$
basis: census / census					
mielke-geometrodynamicsEQ0402 (was)	134	■0.042	$\begin{array}{ll} \text{q: } & \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = \underline{*}'^{\vartheta} \Psi_{\parallel}(\vartheta), \\ & \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = -i \ell^2 \frac{\delta}{\delta \underline{*}^{\vartheta}} \Psi_{\parallel}(\vartheta), \\ \text{p: } & \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = -i \ell^2 \frac{\delta}{\delta \underline{*}^{\vartheta}} \Psi_{\parallel}(\vartheta), \end{array}$	$q: \quad \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = \underline{*}'^{\vartheta} \Psi_{\parallel}(\vartheta),$ $\underline{\pm} p: \quad \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = -i \ell^2 \frac{\delta}{\delta \underline{*}^{\vartheta}} \Psi_{\parallel}(\vartheta),$	$q: \quad \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = \underline{*}'^{\vartheta} \Psi_{\parallel}(\vartheta),$ $\underline{\pm} p: \quad \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = -i \ell^2 \frac{\delta}{\delta \underline{*}^{\vartheta}} \Psi_{\parallel}(\vartheta),$
mielke-geometrodynamicsEQ0402 (now)	134	—	$\begin{array}{ll} \text{q: } & \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = \underline{*}'^{\vartheta} \Psi_{\parallel}(\vartheta), \\ & \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = -i \ell^2 \frac{\delta}{\delta \underline{*}^{\vartheta}} \Psi_{\parallel}(\vartheta), \\ \text{p: } & \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = -i \ell^2 \frac{\delta}{\delta \underline{*}^{\vartheta}} \Psi_{\parallel}(\vartheta), \end{array}$	$q: \quad \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = \underline{*}'^{\vartheta} \Psi_{\parallel}(\vartheta),$ $\underline{\pm} p: \quad \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = -i \ell^2 \frac{\delta}{\delta \underline{*}^{\vartheta}} \Psi_{\parallel}(\vartheta),$	$q: \quad \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = \underline{*}'^{\vartheta} \Psi_{\parallel}(\vartheta),$ $\underline{\pm} p: \quad \underline{*}^{\vartheta} \Psi_{\parallel}(\vartheta) = -i \ell^2 \frac{\delta}{\delta \underline{*}^{\vartheta}} \Psi_{\parallel}(\vartheta),$
basis: inferred / ink					
mielke-geometrodynamicsEQ0437 (was)	142	■0.532	$E_{\alpha} := \frac{\partial L_g}{\partial \vartheta^{\alpha}} = e_{\alpha} \rfloor L_g + (e_{\alpha} \rfloor T^{\beta}) \wedge H_{\beta} + (e_{\alpha} \rfloor R^{\beta \gamma}) \wedge H_{\beta \gamma},$	$E_{\alpha} := \frac{\partial L_g}{\partial \vartheta^{\alpha}} = e_{\alpha} \rfloor L_g + (e_{\alpha} \rfloor T^{\beta}) \wedge H_{\beta} + (e_{\alpha} \rfloor R^{\beta \gamma}) \wedge H_{\beta \gamma},$	$E_{\alpha} := \frac{\partial L_g}{\partial \vartheta^{\alpha}} = e_{\alpha} \rfloor L_g + (e_{\alpha} \rfloor T^{\beta}) \wedge H_{\beta} + (e_{\alpha} \rfloor R^{\beta \gamma}) \wedge H_{\beta \gamma},$
mielke-geometrodynamicsEQ0437 (now)	142	—	$E_{\alpha} := \frac{\partial L_g}{\partial \vartheta^{\alpha}} = e_{\alpha} \rfloor L_g + (e_{\alpha} \rfloor T^{\beta}) \wedge H_{\beta} + (e_{\alpha} \rfloor R^{\beta \gamma}) \wedge H_{\beta \gamma},$	$E_{\alpha} := \frac{\partial L_g}{\partial \vartheta^{\alpha}} = e_{\alpha} \rfloor L_g + (e_{\alpha} \rfloor T^{\beta}) \wedge H_{\beta} + (e_{\alpha} \rfloor R^{\beta \gamma}) \wedge H_{\beta \gamma},$	$E_{\alpha} := \frac{\partial L_g}{\partial \vartheta^{\alpha}} = e_{\alpha} \rfloor L_g + (e_{\alpha} \rfloor T^{\beta}) \wedge H_{\beta} + (e_{\alpha} \rfloor R^{\beta \gamma}) \wedge H_{\beta \gamma},$
basis: census / census					
mielke-geometrodynamicsEQ0440 (was)	143	■0.288	$D \Sigma_{\alpha} \cong (e_{\alpha} \rfloor T^{\gamma}) \wedge \Sigma_{\gamma} + (e_{\alpha} \rfloor R^{\gamma \delta}) \wedge \tau_{\gamma \delta}$	$D \Sigma_{\alpha} \cong (e_{\alpha} \rfloor T^{\gamma}) \wedge \Sigma_{\gamma} + (e_{\alpha} \rfloor R^{\gamma \delta}) \wedge \tau_{\gamma \delta}$	$D \Sigma_{\alpha} \cong (e_{\alpha} \rfloor T^{\gamma}) \wedge \Sigma_{\gamma} + (e_{\alpha} \rfloor R^{\gamma \delta}) \wedge \tau_{\gamma \delta}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

