

1605.05775 — formula report

138 inline formulas (section omitted; all rendered), 26 display equations, 0 tables, 15 unrecovered image regions.

Display equations

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
1605.05775_ E00014 (17)	007	■0.927 ●W +1	$\begin{aligned} & W_{s_1 s_2 \cdots s_N}^{\ell} = \\ & \sum_{\{\alpha\}} U_{s_1}^{\alpha_1} \cdots U_{\alpha_{i-1} s_i}^{\alpha_i} C_{\alpha_i \alpha_{i+1}}^{\ell} V_{s_{i+1} \alpha_{i+2}}^{\alpha_{i+1}} \cdots V_{s_N}^{\alpha_{N-1}} \end{aligned}$	$W_{s_1 s_2 \cdots s_N}^{\ell} = \sum_{\{\alpha\}} U_{s_1}^{\alpha_1} \cdots U_{\alpha_{i-1} s_i}^{\alpha_i} C_{\alpha_i \alpha_{i+1}}^{\ell} V_{s_{i+1} \alpha_{i+2}}^{\alpha_{i+1}} \cdots V_{s_N}^{\alpha_{N-1}}$	$W_{s_1 s_2 \cdots s_N}^{\ell} = \sum_{\{\alpha\}} U_{s_1}^{\alpha_1} \cdots U_{\alpha_{i-1} s_i}^{\alpha_i} C_{\alpha_i \alpha_{i+1}}^{\ell} V_{s_{i+1} \alpha_{i+2}}^{\alpha_{i+1}} \cdots V_{s_N}^{\alpha_{N-1}}$
1605.05775_ E00026 (B4)	010	■0.988 ●N +0	$\phi^{\wedge s_j}(x_j) = \sqrt{\binom{d-1}{s_j-1}} \left(\cos\left(\frac{\pi}{2} x_j\right) \right)^{d-s_j} \left(\sin\left(\frac{\pi}{2} x_j\right) \right)^{s_j-1}$	$\phi^{s_j}(x_j) = \sqrt{\binom{d-1}{s_j-1}} \left(\cos\left(\frac{\pi}{2} x_j\right) \right)^{d-s_j} \left(\sin\left(\frac{\pi}{2} x_j\right) \right)^{s_j-1}$	$\phi^{s_j}(x_j) = \sqrt{\binom{d-1}{s_j-1}} \left(\cos\left(\frac{\pi}{2} x_j\right) \right)^{d-s_j} \left(\sin\left(\frac{\pi}{2} x_j\right) \right)^{s_j-1}$
1605.05775_ E00013 (15)	007	■0.989 ●N +0	$\begin{aligned} & \phi(0) = [1, 0] \\ & \phi(1) = [0, 1] . \end{aligned}$	$\begin{aligned} \phi(0) &= [1, 0] \\ \phi(1) &= [0, 1]. \end{aligned}$	$\begin{aligned} \phi(0) &= [1, 0] \\ \phi(1) &= [0, 1] . \end{aligned}$
1605.05775_ E00025 (B3)	010	■0.990 ●N +0	$\begin{aligned} & \left(\cos^2(\theta_j) + \sin^2(\theta_j) \right)^{d-1} = 1 \\ & = \sum_{p=0}^{d-1} \binom{d-1}{p} (\cos \theta_j)^{2(d-1-p)} (\sin \theta_j)^{2p} . \end{aligned}$	$\begin{aligned} & \left(\cos^2(\theta_j) + \sin^2(\theta_j) \right)^{d-1} = 1 \\ & = \sum_{p=0}^{d-1} \binom{d-1}{p} (\cos \theta_j)^{2(d-1-p)} (\sin \theta_j)^{2p} . \end{aligned}$	$\begin{aligned} & (\cos^2(\theta_j) + \sin^2(\theta_j))^{d-1} = 1 \\ & = \sum_{p=0}^{d-1} \binom{d-1}{p} (\cos \theta_j)^{2(d-1-p)} (\sin \theta_j)^{2p} . \end{aligned}$
1605.05775_ E00005 (5)	003	■0.995 ●N +0	$W_{s_1 s_2 \cdots s_N}^{\ell} = \sum_{\{\alpha\}} A_{s_1}^{\alpha_1} A_{s_2}^{\alpha_1 \alpha_2} \cdots A_{s_j}^{\ell; \alpha_j \alpha_{j+1}} \cdots A_{s_N}^{\alpha_{N-1}}$	$W_{s_1 s_2 \cdots s_N}^{\ell} = \sum_{\{\alpha\}} A_{s_1}^{\alpha_1} A_{s_2}^{\alpha_1 \alpha_2} \cdots A_{s_j}^{\ell; \alpha_j \alpha_{j+1}} \cdots A_{s_N}^{\alpha_{N-1}}$	$W_{s_1 s_2 \cdots s_N}^{\ell} = \sum_{\{\alpha\}} A_{s_1}^{\alpha_1} A_{s_2}^{\alpha_1 \alpha_2} \cdots A_{s_j}^{\ell; \alpha_j \alpha_{j+1}} \cdots A_{s_N}^{\alpha_{N-1}}$
1605.05775_ E00008 (9)	004	■0.999 ●W +2	$\begin{aligned} & \Delta B^{\ell} \stackrel{\text{def}}{=} -\frac{\partial C}{\partial B^{\ell}} \\ & = \sum_{n=1}^{N_T} \sum_{\ell'} \left(\delta_{L_n}^{\ell'} - f^{\ell'}(\mathbf{x}_n) \right) \frac{\partial f^{\ell'}(\mathbf{x}_n)}{\partial B^{\ell}} \\ & = \sum_{n=1}^{N_T} \left(\delta_{L_n}^{\ell} - f^{\ell}(\mathbf{x}_n) \right) \tilde{\Phi}_n \end{aligned}$	$\begin{aligned} \Delta B^{\ell} &\stackrel{\text{def}}{=} -\frac{\partial C}{\partial B^{\ell}} \\ &= \sum_{n=1}^{N_T} \sum_{\ell'} \left(\delta_{L_n}^{\ell'} - f^{\ell'}(\mathbf{x}_n) \right) \frac{\partial f^{\ell'}(\mathbf{x}_n)}{\partial B^{\ell}} \\ &= \sum_{n=1}^{N_T} \left(\delta_{L_n}^{\ell} - f^{\ell}(\mathbf{x}_n) \right) \tilde{\Phi}_n \end{aligned}$	$\begin{aligned} \Delta B^{\ell} &\stackrel{\text{def}}{=} -\frac{\partial C}{\partial B^{\ell}} \\ &= \sum_{n=1}^{N_T} \sum_{\ell'} (\delta_{L_n}^{\ell'} - f^{\ell'}(\mathbf{x}_n)) \frac{\partial f^{\ell'}(\mathbf{x}_n)}{\partial B^{\ell}} \\ &= \sum_{n=1}^{N_T} (\delta_{L_n}^{\ell} - f^{\ell}(\mathbf{x}_n)) \tilde{\Phi}_n . \end{aligned}$
Conf. MathPix confidence: ■≥0.9 ■0.5-0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					










Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
1605.05775_ EQ0009 (11)	004	■ 0.999 ● N +1	$B_{s_j s_{j+1}}^{\alpha_j s_{j+1}} = \sum_{\alpha_j} U_{s_j \alpha_j'}^{\alpha_j-1} S_{\alpha_j}^{\alpha_j'} V_{s_{j+1}}^{\alpha_j \ell \alpha_{j+1}}$	$B_{s_j s_{j+1}}^{\alpha_j-1 \ell \alpha_{j+1}} = \sum_{\alpha_j'} U_{s_j \alpha_j'}^{\alpha_j-1} S_{\alpha_j}^{\alpha_j'} V_{s_{j+1}}^{\alpha_j \ell \alpha_{j+1}}$	$B_{s_j s_{j+1}}^{\alpha_j-1 \ell \alpha_{j+1}} = \sum_{\alpha_j'} U_{s_j \alpha_j'}^{\alpha_j-1} S_{\alpha_j}^{\alpha_j'} V_{s_{j+1}}^{\alpha_j \ell \alpha_{j+1}},$
1605.05775_ EQ0001 (1)	001	■ 1.000 ● K +0	$f(\mathbf{x})=W \cdot \Phi(\mathbf{x})$	$f(\mathbf{x}) = W \cdot \Phi(\mathbf{x}).$	$f(\mathbf{x}) = W \cdot \Phi(\mathbf{x})$
1605.05775_ EQ0002 (2)	002	■ 1.000 ● N +1	$\Phi^{s_1 s_2 \cdots s_N}(\mathbf{x}) = \phi^{s_1}(x_1) \otimes \phi^{s_2}(x_2) \otimes \cdots \phi^{s_N}(x_N)$	$\Phi^{s_1 s_2 \cdots s_N}(\mathbf{x}) = \phi^{s_1}(x_1) \otimes \phi^{s_2}(x_2) \otimes \cdots \phi^{s_N}(x_N).$	$\Phi^{s_1 s_2 \cdots s_N}(\mathbf{x}) = \phi^{s_1}(x_1) \otimes \phi^{s_2}(x_2) \otimes \cdots \phi^{s_N}(x_N)$
1605.05775_ EQ0003 (3)	002	■ 1.000 ● N +0	$\phi^{s_j}(x_j) = \left[\cos\left(\frac{\pi}{2}x_j\right), \sin\left(\frac{\pi}{2}x_j\right) \right]$	$\phi^{s_j}(x_j) = \left[\cos\left(\frac{\pi}{2}x_j\right), \sin\left(\frac{\pi}{2}x_j\right) \right]$	$\phi^{s_j}(x_j) = \left[\cos\left(\frac{\pi}{2}x_j\right), \sin\left(\frac{\pi}{2}x_j\right) \right]$
1605.05775_ EQ0004 (4)	002	■ 1.000 ● N +0	$f^{\ell}(\mathbf{x})=W^{\ell} \cdot \Phi(\mathbf{x})$	$f^{\ell}(\mathbf{x}) = W^{\ell} \cdot \Phi(\mathbf{x})$	$f^{\ell}(\mathbf{x}) = W^{\ell} \cdot \Phi(\mathbf{x})$
1605.05775_ EQ0006 (6)	003	■ 1.000 ● N +0	$C=\frac{1}{2} \sum_{n=1}^{N_T} \sum_{\ell} (f^{\ell}(\mathbf{x}_n) - \delta_{L_n}^{\ell})^2$	$C = \frac{1}{2} \sum_{n=1}^{N_T} \sum_{\ell} (f^{\ell}(\mathbf{x}_n) - \delta_{L_n}^{\ell})^2$	$C = \frac{1}{2} \sum_{n=1}^{N_T} \sum_{\ell} (f^{\ell}(\mathbf{x}_n) - \delta_{L_n}^{\ell})^2$
1605.05775_ EQ0007 (7)	004	■ 1.000 ● N +0	$f^{\ell}(\mathbf{x}_n) = \sum_{\alpha_j} U_{s_j \alpha_j'}^{\alpha_j-1} S_{\alpha_j}^{\alpha_j'} V_{s_{j+1}}^{\alpha_j \ell \alpha_{j+1}}$	$f^{\ell}(\mathbf{x}_n) = \sum_{\alpha_j-1 \alpha_{j+1}} \sum_{s_j s_{j+1}} B_{s_j s_{j+1}}^{\alpha_j-1 \ell \alpha_{j+1}} \left(\tilde{\Phi}_n \right)_{\alpha_j-1 \ell \alpha_{j+1}}^{s_j s_{j+1}}$	$f^{\ell}(\mathbf{x}_n) = \sum_{\alpha_j-1 \alpha_{j+1}} \sum_{s_j s_{j+1}} B_{s_j s_{j+1}}^{\alpha_j-1 \ell \alpha_{j+1}} \left(\tilde{\Phi}_n \right)_{\alpha_j-1 \ell \alpha_{j+1}}^{s_j s_{j+1}}$
1605.05775_ EQ0010 (12)	005	■ 1.000 ● N +0	$\Phi(x_1,x_2)=\phi^{s_1}(x_1)\otimes\phi^{s_2}(x_2)$	$\Phi(x_1,x_2) = \phi^{s_1}(x_1) \otimes \phi^{s_2}(x_2)$	$\Phi(x_1,x_2) = \phi^{s_1}(x_1) \otimes \phi^{s_2}(x_2)$
1605.05775_ EQ0011 (13)	007	■ 1.000 ● N +0	$f(\mathbf{x})=\sum_{\{s\}} W_{s_1 s_2 \cdots s_N} \phi^{s_1}(x_1) \otimes \phi^{s_2}(x_2) \otimes \cdots \phi^{s_N}(x_N)$	$f(\mathbf{x}) = \sum_{\{s\}} W_{s_1 s_2 \cdots s_N} \phi^{s_1}(x_1) \otimes \phi^{s_2}(x_2) \otimes \cdots \phi^{s_N}(x_N).$	$f(\mathbf{x}) = \sum_{\{s\}} W_{s_1 s_2 \cdots s_N} \phi^{s_1}(x_1) \otimes \phi^{s_2}(x_2) \otimes \cdots \phi^{s_N}(x_N)$
1605.05775_ EQ0012 (14)	007	■ 1.000 ● N +0	$\phi^{s_1}(x_1) \otimes \phi^{s_2}(x_2) \otimes \cdots \phi^{s_N}(x_N)$	$\phi^{s_1}(x_1) \otimes \phi^{s_2}(x_2) \otimes \cdots \phi^{s_N}(x_N)$	$\phi^{s_1}(x_1) \otimes \phi^{s_2}(x_2) \otimes \cdots \phi^{s_N}(x_N)$
1605.05775_ EQ0015 (18)	008	■ 1.000 ● N +0	$\sum_{\ell} \int_{\mathbf{x}} f^{\ell}(\mathbf{x}) ^2 d\mu(x) = 1$	$\sum_{\ell} \int_{\mathbf{x}} f^{\ell}(\mathbf{x}) ^2 d\mu(x) = 1$	$\sum_{\ell} \int_{\mathbf{x}} f^{\ell}(\mathbf{x}) ^2 d\mu(x) = 1$
1605.05775_ EQ0016 (19)	008	■ 1.000 ● N +0	$\sum_{\ell} \sum_{s_1, s_2, \dots, s_N} \bar{W}_{s_1 s_2 \cdots s_N}^{\ell} W_{s_1 s_2 \cdots s_N}^{\ell} = 1.$	$\sum_{\ell} \sum_{s_1, s_2, \dots, s_N} \bar{W}_{s_1 s_2 \cdots s_N}^{\ell} W_{s_1 s_2 \cdots s_N}^{\ell} = 1.$	$\sum_{\ell} \sum_{s_1, s_2, \dots, s_N} \bar{W}_{s_1 s_2 \cdots s_N}^{\ell} W_{s_1 s_2 \cdots s_N}^{\ell} = 1$
1605.05775_ EQ0017 (20)	008	■ 1.000 ● N +1	$\int_x \bar{\phi}^s(x) \phi^{s'}(x) d\mu(x) = \delta_{ss'}$	$\int_x \bar{\phi}^s(x) \phi^{s'}(x) d\mu(x) = \delta_{ss'}$	$\int_x \bar{\phi}^s(x) \phi^{s'}(x) d\mu(x) = \delta_{ss'},$
1605.05775_ EQ0018 (21)	008	■ 1.000 ● K +0	$\sum_s \phi^s(x) ^2 = 1$	$\sum_s \phi^s(x) ^2 = 1$	$\sum_s \phi^s(x) ^2 = 1$
1605.05775_ EQ0019 (22)	008	■ 1.000 ● N +0	$\phi(x)=[\cos(\pi x), \sin(\pi x)]$	$\phi(x) = [\cos(\pi x), \sin(\pi x)].$	$\phi(x) = [\cos(\pi x), \sin(\pi x)]$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					
















Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
1605.05775_ EQ0020 (23)	008	■ 1.000 ● N +0	$\phi(x)=\left[e^{i(3\pi/2)x}\cos\left(\frac{\pi}{2}x\right),e^{-i(3\pi/2)x}\sin\left(\frac{\pi}{2}x\right)\right]$.	$\phi(x)=\left[e^{i(3\pi/2)x}\cos\left(\frac{\pi}{2}x\right),e^{-i(3\pi/2)x}\sin\left(\frac{\pi}{2}x\right)\right]$.	$\phi(x)=\left[e^{i(3\pi/2)x}\cos\left(\frac{\pi}{2}x\right),e^{-i(3\pi/2)x}\sin\left(\frac{\pi}{2}x\right)\right]$.
1605.05775_ EQ0021 (24)	009	■ 1.000 ● N +1	$C=-\sum_{n=1}^{N_s}\log\left f^{L_n}(\mathbf{x}_n)\right ^2$	$C=-\sum_{n=1}^{N_s}\log\left f^{L_n}(\mathbf{x}_n)\right ^2$	$C=-\sum_{n=1}^{N_s}\log\left f^{L_n}(\mathbf{x}_n)\right ^2$
1605.05775_ EQ0022 (25)	009	■ 1.000 ● N +1	$D_{\mathrm{KL}}=\sum_{\ell}\int_{\mathbf{x}}p(\ell,\mathbf{x})\log\left(\frac{p(\ell,\mathbf{x})}{\tilde{p}(\ell,\mathbf{x})}\right)$	$D_{\mathrm{KL}}=\sum_{\ell}\int_{\mathbf{x}}p(\ell,\mathbf{x})\log\left(\frac{p(\ell,\mathbf{x})}{\tilde{p}(\ell,\mathbf{x})}\right)$	$D_{\mathrm{KL}}=\sum_{\ell}\int_{\mathbf{x}}p(\ell,\mathbf{x})\log\left(\frac{p(\ell,\mathbf{x})}{\tilde{p}(\ell,\mathbf{x})}\right)$
1605.05775_ EQ0023 (B1)	010	■ 1.000 ● N +0	$\phi^{s_j}(x_j)=\left[\cos\left(\frac{\pi}{2}x_j\right),\sin\left(\frac{\pi}{2}x_j\right)\right]$.	$\phi^{s_j}(x_j)=\left[\cos\left(\frac{\pi}{2}x_j\right),\sin\left(\frac{\pi}{2}x_j\right)\right]$.	$\phi^{s_j}(x_j)=\left[\cos\left(\frac{\pi}{2}x_j\right),\sin\left(\frac{\pi}{2}x_j\right)\right]$.
1605.05775_ EQ0024 (B2)	010	■ 1.000 ● K +0	$\left(\cos^2(\theta_j)+\sin^2(\theta_j)\right)^{d-1}=1$.	$\left(\cos^2(\theta_j)+\sin^2(\theta_j)\right)^{d-1}=1$.	$(\cos^2(\theta_j)+\sin^2(\theta_j))^{d-1}=1$.
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					





Image regions — rendered against scan

Two comparable cells per row. **Rendered**: the region’s own LaTeX compiled as its OWN document, or — where no LaTeX exists — the scan again, so every row has two cells and the ink difference reads as a floor rather than as missing-versus-present. A duplicated row is marked *(dup)* and its distance is a SELF-comparison, not agreement between two sources. **Scan**: the tex.zip image whose filename region 5-tuple matches this row, else the CDN crop. The Class column carries the residual class beside MathPix’s confidence, as the equation rows do. Setting a region’s LaTeX does NOT say it renders to what is on the page — that is what the two columns are for, and 281 is the open question they exist to let a reader answer. A region with no LaTeX can be reconstructed with pdfdrill vision; verify any LLM result against the real ink with inkdrill.

Identifier	Page	Class	Source	Author source	Rendered	Scan
1605.05775_PIC_0001	009	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-09_554_726_176_249.jpg			
1605.05775_DIA_0001	001	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-01_79_857_2241_185.jpg			
1605.05775_DIA_0002	002	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-02_170_559_178_1246.jpg	$\Phi = \begin{matrix} s_1 & s_2 & s_3 & s_4 & s_5 & s_6 \\ \bigcirc & \bigcirc & \bigcirc & \bigcirc & \bigcirc & \bigcirc \\ \phi^{s_1} & \phi^{s_2} & \phi^{s_3} & \phi^{s_4} & \phi^{s_5} & \phi^{s_6} \end{matrix}$	$\Phi = \begin{matrix} s_1 & s_2 & s_3 & s_4 & s_5 & s_6 \\ \bigcirc & \bigcirc & \bigcirc & \bigcirc & \bigcirc & \bigcirc \\ \phi^{s_1} & \phi^{s_2} & \phi^{s_3} & \phi^{s_4} & \phi^{s_5} & \phi^{s_6} \end{matrix}$	$\Phi = \begin{matrix} s_1 & s_2 & s_3 & s_4 & s_5 & s_6 \\ \bigcirc & \bigcirc & \bigcirc & \bigcirc & \bigcirc & \bigcirc \\ \phi^{s_1} & \phi^{s_2} & \phi^{s_3} & \phi^{s_4} & \phi^{s_5} & \phi^{s_6} \end{matrix}$
1605.05775_DIA_0003	002	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-02_374_539_510_1257.jpg			
1605.05775_DIA_0004	003	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-03_185_726_183_251.jpg	$\begin{matrix} \ell \\ \text{---} \end{matrix} W^\ell = \bigcirc_{\Phi(\mathbf{x})}^\ell f^\ell(\mathbf{x})$	$\begin{matrix} \ell \\ \text{---} \end{matrix} W^\ell = \bigcirc_{\Phi(\mathbf{x})}^\ell f^\ell(\mathbf{x})$	$\begin{matrix} \ell \\ \text{---} \end{matrix} W^\ell = \bigcirc_{\Phi(\mathbf{x})}^\ell f^\ell(\mathbf{x})$
1605.05775_DIA_0005	003	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-03_155_855_180_1100.jpg			

1605.05775_DIA_0006	004	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-04_622_848_176_192.jpg			
1605.05775_DIA_0007	004	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-04_615_855_181_1100.jpg			
1605.05775_DIA_0008	005	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-05_213_679_188_1186.jpg			
1605.05775_DIA_0009	006	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-06_561_561_1756_333.jpg			

1605.05775_DIA_0010	006	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-06_1514_584_204_1231.jpg	(a)  (b)  (c) 	(a)  (b)  (c) 	(a)  (b)  (c) 
1605.05775_DIA_0011	007	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-07_360_844_188_190.jpg	(a)  (b) 	(a)  (b) 	(a)  (b) 
1605.05775_DIA_0012	008	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-08_525_857_176_181.jpg	(a) $W_{s_1 \dots s_6}^\ell = \text{Diagram with nodes } U_{s_1}, U_{s_2}, U_{s_3}, \text{ and } V_{s_4}, V_{s_5}, V_{s_6} \text{ connected by lines.}$ (b) $\text{Diagram with nodes } U_{s_j} \text{ and } U_{s_j}^\dagger = \text{Diagram with nodes } V_{s_j} \text{ and } V_{s_j}^\dagger = \text{Diagram}$ (c) $\text{Diagram with nodes } U_{s_1}, U_{s_2}, U_{s_3}, \text{ and } V_{s_4}, V_{s_5}, V_{s_6} \text{ connected by lines.} = \text{Diagram with nodes } U_{s_1}, U_{s_2}, U_{s_3}, \text{ and } V_{s_4}, V_{s_5}, V_{s_6} \text{ connected by lines.}$	(a) $W_{s_1 \dots s_6}^\ell = \text{Diagram with nodes } U_{s_1}, U_{s_2}, U_{s_3}, \text{ and } V_{s_4}, V_{s_5}, V_{s_6} \text{ connected by lines.}$ (b) $\text{Diagram with nodes } U_{s_j} \text{ and } U_{s_j}^\dagger = \text{Diagram with nodes } V_{s_j} \text{ and } V_{s_j}^\dagger = \text{Diagram}$ (c) $\text{Diagram with nodes } U_{s_1}, U_{s_2}, U_{s_3}, \text{ and } V_{s_4}, V_{s_5}, V_{s_6} \text{ connected by lines.} = \text{Diagram with nodes } U_{s_1}, U_{s_2}, U_{s_3}, \text{ and } V_{s_4}, V_{s_5}, V_{s_6} \text{ connected by lines.}$	(a) $W_{s_1 \dots s_6}^\ell = \text{Diagram with nodes } U_{s_1}, U_{s_2}, U_{s_3}, \text{ and } V_{s_4}, V_{s_5}, V_{s_6} \text{ connected by lines.}$ (b) $\text{Diagram with nodes } U_{s_j} \text{ and } U_{s_j}^\dagger = \text{Diagram with nodes } V_{s_j} \text{ and } V_{s_j}^\dagger = \text{Diagram}$ (c) $\text{Diagram with nodes } U_{s_1}, U_{s_2}, U_{s_3}, \text{ and } V_{s_4}, V_{s_5}, V_{s_6} \text{ connected by lines.} = \text{Diagram with nodes } U_{s_1}, U_{s_2}, U_{s_3}, \text{ and } V_{s_4}, V_{s_5}, V_{s_6} \text{ connected by lines.}$

1605.05775_DIA_0013	010	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-10_217_597_178_318.jpg			
1605.05775_DIA_0014	010	— (dup)	90117dc3-f9f1-431f-89f9-03fd8cfa01c-10_423_505_563_363.jpg	(a)  =  $\sum_j M_{ij} v_j = w_i$ (b)  =  $\sum_{kl} A_{ijkl} B_{klm} = C_{ijm}$	(a)  =  $\sum_j M_{ij} v_j = w_i$ (b)  =  $\sum_{kl} A_{ijkl} B_{klm} = C_{ijm}$	(a)  =  $\sum_j M_{ij} v_j = w_i$ (b)  =  $\sum_{kl} A_{ijkl} B_{klm} = C_{ijm}$