

voloshin-hypergraph: residuals

equations measured 2026-09-13T13:16:46Z against the model of 2026-09-13

Corrected (2)

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
voloshin-hypergraph_ EQ0133 (was)	170	■ 0.340	$(\mathcal{H})_2=(X, \mathcal{E}) \text{ where } \left\{x_i, x_j\right\} \in \mathcal{E} \Leftrightarrow \mathcal{D}\left(x_i\right) \cap \mathcal{D}\left(x_j\right) \neq \emptyset$	$(\mathcal{H})_2=(X, \mathcal{E}) \text{ where } \left\{x_i, x_j\right\} \in \mathcal{E} \Leftrightarrow \mathcal{D}\left(x_i\right) \cap \mathcal{D}\left(x_j\right) \neq \emptyset$	$(\mathcal{H})_2=(X, \mathcal{E}) \text{ where } \left\{x_i, x_j\right\} \in \mathcal{E} \Leftrightarrow \mathcal{D}\left(x_i\right) \cap \mathcal{D}\left(x_j\right) \neq \emptyset$
voloshin-hypergraph_ EQ0133 (now)	170	—	$(\mathcal{H})_2=(X, \mathcal{E}) \text{ where } \left\{x_i, x_j\right\} \in \mathcal{E} \Leftrightarrow \mathcal{D}\left(x_i\right) \cap \mathcal{D}\left(x_j\right) \neq \emptyset$	$(\mathcal{H})_2=(X, \mathcal{E}) \text{ where } \left\{x_i, x_j\right\} \in \mathcal{E} \Leftrightarrow \mathcal{D}\left(x_i\right) \cap \mathcal{D}\left(x_j\right) \neq \emptyset$	$(\mathcal{H})_2=(X, \mathcal{E}) \text{ where } \left\{x_i, x_j\right\} \in \mathcal{E} \Leftrightarrow \mathcal{D}\left(x_i\right) \cap \mathcal{D}\left(x_j\right) \neq \emptyset$
basis: inferred / ink					
voloshin-hypergraph_ EQ0141 (was)	185	■ 0.473	$\tau(\mathcal{H})=\nu(\mathcal{H})$	$\tau(\mathcal{H})=\nu(\mathcal{H})$	$\tau(\mathcal{H})=\nu(\mathcal{H})$
voloshin-hypergraph_ EQ0141 (now)	185	—	$\tau(\mathcal{H})=\nu(\mathcal{H})$. a hypergraph, draw	$\tau(\mathcal{H})=\nu(\mathcal{H}).ahypergraph, draw$	$\tau(\mathcal{H})=\nu(\mathcal{H})$
basis: inferred / ink					
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5-0.9 ■ < 0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					