

Geometrodynamics of Gauge Fields On the Geometry of Yang-Mills and Gravitational Gauge Theories (Eckehard W. Mielke) (Z-Library) — formula report

1794 inline formulas, 1149 display equations, 5 tables, 18 unrecovered image regions.

3 rows show a REFINEMENT below, not the OCR reading: 3 verified against the inferred. Each is marked [refined] beside its identifier. The model is unchanged — the original reading is still what ‘latex’ holds; this projection chose the refinement for them. Rows: Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories_Eckehard_W_Mielke_Z-Library_EQ0378, Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories_Eckehard_W_Mielke_Z-Library_EQ0402, Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories_Eckehard_W_Mielke_Z-Library_EQ0972

Display equations

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0966 (14.2.7) [conf 0.024]	313	■ 0.024 ● W +1	$\begin{aligned} d\hat{s}^2 &= \frac{1}{4+K} \operatorname{Tr} \left(\hat{\vartheta} \otimes_s \hat{\vartheta}^T \right) = \hat{g}_{AB} \hat{\vartheta}^A \otimes_s \hat{\vartheta}^B; \quad A, B, = 0, \dots, 3+K \\ &= \frac{1}{4+K} \operatorname{Tr} \left[\begin{array}{c c} \vartheta \otimes \vartheta + \hat{\ell}^2 \omega \otimes \omega & \hat{\ell} \omega \otimes \overset{\circ}{\vartheta} \\ \hline \hat{\ell} \vartheta \otimes \omega & \begin{array}{c} \ddots \\ \vartheta \otimes \vartheta \end{array} \end{array} \right]. \end{aligned}$	$d\hat{s}^2 = \frac{1}{4+K} \operatorname{Tr} \left(\hat{\vartheta} \otimes_s \hat{\vartheta}^T \right) = \hat{g}_{AB} \hat{\vartheta}^A \otimes_s \hat{\vartheta}^B; \quad A, B, = 0, \dots, 3+K$ $= \frac{1}{4+K} \operatorname{Tr} \left[\begin{array}{c c} \vartheta \otimes \vartheta + \hat{\ell}^2 \omega \otimes \omega & \hat{\ell} \omega \otimes \overset{\circ}{\vartheta} \\ \hline \hat{\ell} \vartheta \otimes \omega & \begin{array}{c} \ddots \\ \vartheta \otimes \vartheta \end{array} \end{array} \right]. \quad (1)$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0403 (6.4.23) [conf 0.028]	134	■ 0.028 ● C +4	$\Psi_{\parallel}(\vartheta) = \exp \left(- \int_{M_3} \overset{(\pm)}{C}_{\mathrm{TT}} \right) = \exp \left(\frac{-1}{2\ell^2} \int_{M_3} \left[\overset{*}{+} \underline{\vartheta}_B \wedge \overset{*}{T}^B \right] \right)$	$\Psi_{\parallel}(\vartheta) = \exp \left(- \int_{M_3} \overset{(\pm)}{C}_{\mathrm{TT}} \right) = \exp \left(\frac{-1}{2\ell^2} \int_{M_3} \left[\overset{*}{+} \underline{\vartheta}_B \wedge \overset{*}{T}^B \right] \right)$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0850 (12.3.3) [conf 0.032]	275	■0.032 C +3	$\begin{aligned} \not{D}^2 &= -\frac{1}{2}\gamma^\alpha\gamma^\beta\left(\{D_\alpha^{\{\}},D_\beta^{\{\}}\}+\left[D_\alpha^{\{\}},D_\beta^{\{\}}\right]\right)-2im\not{D} \\ &= -\frac{i}{4}\gamma_5\left(\not{D}\not{\not{A}}\right)+\frac{1}{2}\gamma_5\sigma^{\alpha\beta}\not{A}_\alpha D_\beta^{\{\}}+m^2-\frac{1}{2}m\gamma_5\not{\not{A}}-\frac{1}{16}\not{\not{A}}\not{\not{A}} \\ &\cong-\square-\frac{1}{8}\sigma^{\alpha\beta}R_{\alpha\beta\mu\nu}^{\{\}}\sigma^{\mu\nu} \\ &= -\frac{i}{4}\gamma_5\left(\not{D}\not{B}\right)+\frac{1}{2}\gamma_5\sigma^{\alpha\beta}\not{A}_\alpha D_\beta^{\{\}}-\frac{1}{16}\not{A}_\alpha\not{A}^\alpha-m^2 \end{aligned}$	$\begin{aligned} \not{D}^2 &= -\frac{1}{2}\gamma^\alpha\gamma^\beta\left(\{D_\alpha^{\{\}},D_\beta^{\{\}}\}+\left[D_\alpha^{\{\}},D_\beta^{\{\}}\right]\right)-2im\not{D} \\ &= -\frac{i}{4}\gamma_5\left(\not{D}\not{\not{A}}\right)+\frac{1}{2}\gamma_5\sigma^{\alpha\beta}\not{A}_\alpha D_\beta^{\{\}}+m^2-\frac{1}{2}m\gamma_5\not{\not{A}}-\frac{1}{16}\not{\not{A}}\not{\not{A}} \\ &\cong-\square-\frac{1}{8}\sigma^{\alpha\beta}R_{\alpha\beta\mu\nu}^{\{\}}\sigma^{\mu\nu} \\ &= -\frac{i}{4}\gamma_5\left(\not{D}\not{B}\right)+\frac{1}{2}\gamma_5\sigma^{\alpha\beta}\not{A}_\alpha D_\beta^{\{\}}-\frac{1}{16}\not{A}_\alpha\not{A}^\alpha-m^2 \end{aligned}$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0402 (6.4.22) [conf 0.042] [refined: inferred]	134	■0.042 N +2	$\begin{array}{ll} \text{q: } & \backslash\stackrel{\ast}{\operatorname{rel}}\{\ast\}\{\operatorname{underline}\{\vartheta\}^{\wedge\beta}\}\backslash\Psi_{\{\backslash\}\backslash(\vartheta)}=\backslash\stackrel{\ast}{\operatorname{rel}}\{\ast'\}\{\operatorname{underline}\{\vartheta\}^{\wedge\beta}\}\backslash\Psi_{\{\backslash\}\backslash(\vartheta)},\backslash \\ & \backslash\backslash\stackrel{\ast}{\operatorname{rel}}\{\backslash\text{pm}\}\{\text{p}\}:& \backslash\stackrel{\ast}{\operatorname{rel}}\{\ast^{\wedge(\backslash\text{pm})}\}\}\{\operatorname{underline}\{\Pi_{\{\alpha\}}\}\}\backslash\Psi_{\{\backslash\}\backslash(\vartheta)}=-i\ell^2\frac{\delta}{\delta\backslash\vartheta_{\alpha}}\Psi_{\{\backslash\}\backslash(\vartheta)},\backslash \\ & \backslash\backslash\stackrel{\ast}{\operatorname{rel}}\{\ast\}\{\operatorname{underline}\{\vartheta\}^{\wedge\beta}\}\backslash\Psi_{\{\backslash\}\backslash(\vartheta)},\backslash\end{array}$	$\begin{aligned} q:& \quad \vartheta^{\ast\beta}\Psi_{\parallel}(\vartheta)=\vartheta^{\ast'\beta}\Psi_{\parallel}(\vartheta), \\ \pm p:& \quad \vartheta^{\ast(\pm)}_{\alpha}\Psi_{\parallel}(\vartheta)=-i\ell^2\frac{\delta}{\delta\vartheta^{\ast}_{\alpha}}\Psi_{\parallel}(\vartheta), \end{aligned}$	$\begin{aligned} q:& \quad \vartheta^{\ast\beta}\Psi_{\parallel}(\vartheta)=\vartheta^{\ast'\beta}\Psi_{\parallel}(\vartheta), \\ \pm p:& \quad \vartheta^{\ast(\pm)}_{\alpha}\Psi_{\parallel}(\vartheta)=-i\ell^2\frac{\delta}{\delta\vartheta^{\ast}_{\alpha}}\Psi_{\parallel}(\vartheta), \end{aligned}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0293 (4.5.4) [conf 0.052]	101	■0.052 N +0	$\xi\downharpoonleft L\equiv(\xi\downharpoonleft\vartheta^\alpha)\wedge\frac{\partial L}{\partial\vartheta^\alpha}+(\xi\downharpoonleft\Psi)\wedge\frac{\partial L}{\partial\Psi}+(\xi\downharpoonleft D\Psi)\wedge\frac{\partial L}{\partial D\Psi}.$	$\xi\downharpoonleft L\equiv(\xi\downharpoonleft\vartheta^\alpha)\wedge\frac{\partial L}{\partial\vartheta^\alpha}+(\xi\downharpoonleft\Psi)\wedge\frac{\partial L}{\partial\Psi}+(\xi\downharpoonleft D\Psi)\wedge\frac{\partial L}{\partial D\Psi}.$	

Conf. MathPix confidence:

■≥0.9

■0.5–0.9

■<0.5

— no value

Residual render vs scan (inkdrill):

●C component

●W weak

●S stable

●N noise

●K clean

●not measured

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0664 (2.8.6) [conf 0.054]	047	■ 0.054 ● N -2	$\begin{aligned} & \stackrel{*}{\pi}(\gamma_1) = -\frac{1}{2\pi i} \text{Tr}(\Omega) \quad (=0) \\ & \stackrel{*}{\pi}(\gamma_2) = -\frac{1}{8\pi^2} \{ \text{Tr}(\Omega \wedge \Omega) - \text{Tr}(\Omega) \wedge \text{Tr}(\Omega) \} \\ & \quad \left(= -\frac{1}{32\pi^2} \epsilon^{\alpha\beta\mu\nu} F_{\alpha\beta}^j F_{\mu\nu j} \sqrt{ g } d^4x \right) \\ & \stackrel{*}{\pi}(\gamma_k) = 0 \quad \text{for } k \geq 3. \end{aligned}$	$\begin{aligned} & \stackrel{*}{\pi}(\gamma_1) = -\frac{1}{2\pi i} \text{Tr} \Omega \quad (=0) \\ & \stackrel{*}{\pi}(\gamma_2) = -\frac{1}{8\pi^2} \{ \text{Tr}(\Omega \wedge \Omega) - \text{Tr}(\Omega) \wedge \text{Tr}(\Omega) \} \\ & \quad \left(= -\frac{1}{32\pi^2} \epsilon^{\alpha\beta\mu\nu} F_{\alpha\beta}^j F_{\mu\nu j} \sqrt{ g } d^4x \right) \\ & \stackrel{*}{\pi}(\gamma_k) = 0 \text{ for } k \geq 3. \end{aligned}$	$\begin{aligned} & \stackrel{*}{\pi}(\gamma_1) = -\frac{1}{2\pi i} \text{Tr} \Omega \quad (=0) \\ & \stackrel{*}{\pi}(\gamma_2) = -\frac{1}{8\pi^2} \{ \text{Tr}(\Omega \wedge \Omega) - \text{Tr}(\Omega) \wedge \text{Tr}(\Omega) \} \\ & \quad \left(= -\frac{1}{32\pi^2} \epsilon^{\alpha\beta\mu\nu} F_{\alpha\beta}^j F_{\mu\nu j} \sqrt{ g } d^4x \right) \\ & \stackrel{*}{\pi}(\gamma_k) = 0 \quad \text{for } k \geq 3. \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0400 (6.4.19) [conf 0.060]	133	■ 0.060 ● W +1	$\stackrel{(\pm)}{\mathcal{G}}_{\alpha} := \underline{D}^{(\pm)} \stackrel{(\pm)}{\Pi}_{\alpha} = \underline{D}^{(\pm)} \stackrel{(\pm)}{A}_{\alpha} = 0,$	$\stackrel{(\pm)}{\mathcal{G}}_{\alpha} := \underline{D}^{(\pm)} \stackrel{(\pm)}{\Pi}_{\alpha} = \underline{D}^{(\pm)} \stackrel{(\pm)}{A}_{\alpha} = 0,$	$\stackrel{(\pm)}{\mathcal{G}}_{\alpha} := \underline{D}^{(\pm)} \stackrel{(\pm)}{\Pi}_{\alpha} = \underline{D}^{(\pm)} \stackrel{(\pm)}{A}_{\alpha} = 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0718 (10.9.4) [conf 0.066]	229	■ 0.066 ● N +0	$\begin{aligned} & \Sigma_{\alpha} := e_{\alpha} \lrcorner L_{\Psi} - (e_{\alpha} \lrcorner \Psi) \wedge \frac{\partial L_{\Psi}}{\partial \Psi} - (e_{\alpha} \lrcorner \bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial \bar{\Psi}} \\ & \quad - (e_{\alpha} \lrcorner D\Psi) \wedge \frac{\partial L_{\Psi}}{\partial D\Psi} - (e_{\alpha} \lrcorner D\bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial D\bar{\Psi}}; \end{aligned}$	$\begin{aligned} & \Sigma_{\alpha} := e_{\alpha} \lrcorner L_{\Psi} - (e_{\alpha} \lrcorner \Psi) \wedge \frac{\partial L_{\Psi}}{\partial \Psi} - (e_{\alpha} \lrcorner \bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial \bar{\Psi}} \\ & \quad - (e_{\alpha} \lrcorner D\Psi) \wedge \frac{\partial L_{\Psi}}{\partial D\Psi} - (e_{\alpha} \lrcorner D\bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial D\bar{\Psi}}; \end{aligned}$	$\begin{aligned} & \Sigma_{\alpha} := e_{\alpha} \lrcorner L_{\Psi} - (e_{\alpha} \lrcorner \Psi) \wedge \frac{\partial L_{\Psi}}{\partial \Psi} - (e_{\alpha} \lrcorner \bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial \bar{\Psi}} \\ & \quad - (e_{\alpha} \lrcorner D\Psi) \wedge \frac{\partial L_{\Psi}}{\partial D\Psi} - (e_{\alpha} \lrcorner D\bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial D\bar{\Psi}}; \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0391 (6.4.6) [conf 0.066]	131	■ 0.066 ● W +0	$\begin{aligned} & \Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = e_{\alpha} \lrcorner L - (e_{\alpha} \lrcorner D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha} \lrcorner \Psi) \wedge \frac{\partial L}{\partial \Psi} \\ & \quad - (e_{\alpha} \lrcorner T^{\beta}) \wedge \frac{\partial L}{\partial T^{\beta}} + D \frac{\partial L}{\partial T^{\alpha}} - (e_{\alpha} \lrcorner R_{\beta}^{\gamma}) \wedge \frac{\partial L}{\partial R_{\beta}^{\gamma}}. \end{aligned}$	$\begin{aligned} & \Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = e_{\alpha} \lrcorner L - (e_{\alpha} \lrcorner D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha} \lrcorner \Psi) \wedge \frac{\partial L}{\partial \Psi} \\ & \quad - (e_{\alpha} \lrcorner T^{\beta}) \wedge \frac{\partial L}{\partial T^{\beta}} + D \frac{\partial L}{\partial T^{\alpha}} - (e_{\alpha} \lrcorner R_{\beta}^{\gamma}) \wedge \frac{\partial L}{\partial R_{\beta}^{\gamma}}. \end{aligned}$	$\begin{aligned} & \Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = e_{\alpha} \lrcorner L - (e_{\alpha} \lrcorner D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha} \lrcorner \Psi) \wedge \frac{\partial L}{\partial \Psi} \\ & \quad - (e_{\alpha} \lrcorner T^{\beta}) \wedge \frac{\partial L}{\partial T^{\beta}} + D \frac{\partial L}{\partial T^{\alpha}} - (e_{\alpha} \lrcorner R_{\beta}^{\gamma}) \wedge \frac{\partial L}{\partial R_{\beta}^{\gamma}}. \end{aligned}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ E08528 (7.7.48) [conf 0.078]	168	■ 0.078 ● C +3	$\begin{aligned} 2 \ell^2 d C_{\mathrm{T}} &:= 2 d \left(\vartheta^{\alpha} \wedge \eta_{\alpha\beta} - R^{\alpha\beta} \wedge \eta_{\alpha\beta} - T^{\alpha} \wedge \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta} \rfloor T^{\beta}) - \frac{1}{2} e_{\alpha} \rfloor (T^{\beta} \wedge \vartheta_{\beta}) \right] \right. \\ &\quad \left. - T^{\alpha} \wedge \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta} \rfloor T^{\beta}) - \frac{1}{2} e_{\alpha} \rfloor (T^{\beta} \wedge \vartheta_{\beta}) \right] \right) \\ &= R^{\alpha\beta} \wedge \eta_{\alpha\beta} - R^{\alpha\beta} \wedge \eta_{\alpha\beta} - T^{\alpha} \wedge \left(- {}^{(1)}T_{\alpha} + 2 {}^{(2)}T_{\alpha} + \frac{1}{2} {}^{(3)}T_{\alpha} \right), \end{aligned}$	$2\ell^2 dC_{\mathrm{T}} := 2d(\vartheta^{\alpha} \wedge T_{\alpha})$ $= R^{\alpha\beta} \wedge \eta_{\alpha\beta} - R^{\alpha\beta} \wedge \eta_{\alpha\beta} - T^{\alpha} \wedge \left(- {}^{(1)}T_{\alpha} + 2 {}^{(2)}T_{\alpha} + \frac{1}{2} {}^{(3)}T_{\alpha} \right),$	$2\ell^2 dC_{\mathrm{T}} := 2d(\vartheta^{\alpha} \wedge T_{\alpha})$ $= R^{\alpha\beta} \wedge \eta_{\alpha\beta} - R^{\alpha\beta} \wedge \eta_{\alpha\beta} - T^{\alpha} \wedge \left(- {}^{(1)}T_{\alpha} + 2 {}^{(2)}T_{\alpha} + \frac{1}{2} {}^{(3)}T_{\alpha} \right),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ E08382 (6.2.33) [conf 0.087]	129	■ 0.087 ● W +1	$\stackrel{\text{pm}}{\Gamma}_{\alpha}^{\beta} := \Gamma_{\alpha}^{\beta} + i A^{\beta}, \quad \stackrel{\text{pm}}{A}_{\alpha}^{\beta} := \pm \frac{1}{2} \ell^2 e_{\alpha} \rfloor \Pi^{\beta},$	$\stackrel{\text{pm}}{\Gamma}_{\alpha}^{\beta} := \Gamma_{\alpha}^{\beta} + i A^{\beta}, \quad \stackrel{\text{pm}}{A}_{\alpha}^{\beta} := \pm \frac{1}{2} \ell^2 e_{\alpha} \rfloor \Pi^{\beta},$	$\stackrel{\text{pm}}{\Gamma}_{\alpha}^{\beta} := \Gamma_{\alpha}^{\beta} + i A^{\beta}, \quad \stackrel{\text{pm}}{A}_{\alpha}^{\beta} := \pm \frac{1}{2} \ell^2 e_{\alpha} \rfloor \Pi^{\beta},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ E08485 (6.4.25) [conf 0.098]	135	■ 0.098 ● C +8	$\underline{D}^{\pm} \left(\frac{\delta}{\delta^* \vartheta_B} \right) = \underline{D}^{\pm} \underline{T}^B \equiv \underline{R}_C^{\pm B} \wedge \underline{\vartheta}^C = 0,$	$-\ell^2 \underline{D}^{\pm} \left(\frac{\delta}{\delta^* \vartheta_B} \right) = \underline{D}^{\pm} \underline{T}^B \equiv \underline{R}_C^{\pm B} \wedge \underline{\vartheta}^C = 0,$	$-\ell^2 \underline{D}^{\pm} \left(\frac{\delta}{\delta^* \vartheta_B} \right) = \underline{D}^{\pm} \underline{T}^B \equiv \underline{R}_C^{\pm B} \wedge \underline{\vartheta}^C = 0,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0529 (7.7.49) [conf 0.099]	168	■ 0.099 ● C +6	$\begin{aligned} L_{\text{Euler}} &:= (-1)^{\text{sig}+1} \text{Tr} \left\{ \Omega \wedge \Omega^{(*)} \right\} = \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= dC_{\text{RR}^{(*)}} = \frac{(-1)^{\text{sig}}}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta(*)} - \frac{1}{3} \Gamma_{\alpha}{}^{\beta(*)} \wedge \Gamma_{\beta}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\alpha} \right) \\ &\equiv \frac{1}{2} R_{\alpha\beta} \wedge {}^* R^{\alpha\beta} - 2 \text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} + \frac{1}{2} \text{Ric}_{\alpha}{}^{\alpha} \wedge {}^* \text{Ric}_{\beta}{}^{\beta} \\ &\equiv -L_{\text{SKY}} \\ &\quad - 2 \left(\text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} - \frac{1}{4} \text{Ric}_{\alpha}{}^{\alpha} \wedge {}^* \text{Ric}_{\beta}{}^{\beta} \right) \end{aligned}$	$\begin{aligned} L_{\text{Euler}} &:= (-1)^{\text{sig}+1} \text{Tr} \left\{ \Omega \wedge \Omega^{(*)} \right\} = \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= dC_{\text{RR}^{(*)}} = \frac{(-1)^{\text{sig}}}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta(*)} - \frac{1}{3} \Gamma_{\alpha}{}^{\beta(*)} \wedge \Gamma_{\beta}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\alpha} \right) \\ &\equiv \frac{1}{2} R_{\alpha\beta} \wedge {}^* R^{\alpha\beta} - 2 \text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} + \frac{1}{2} \text{Ric}_{\alpha}{}^{\alpha} \wedge {}^* \text{Ric}_{\beta}{}^{\beta} \\ &\equiv -L_{\text{SKY}} \\ &\quad - 2 \left(\text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} - \frac{1}{4} \text{Ric}_{\alpha}{}^{\alpha} \wedge {}^* \text{Ric}_{\beta}{}^{\beta} \right) \end{aligned}$	$\begin{aligned} L_{\text{Euler}} &:= (-1)^{\text{sig}+1} \text{Tr} \{ \varOmega \wedge \varOmega^{(*)} \} = \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= dC_{\text{RR}^{(*)}} = \frac{(-1)^{\text{sig}}}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta(*)} - \frac{1}{3} \Gamma_{\alpha}{}^{\beta(*)} \wedge \Gamma_{\beta}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\alpha} \right) \\ &\equiv \frac{1}{2} R_{\alpha\beta} \wedge {}^* R^{\alpha\beta} - 2 \text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} + \frac{1}{2} \text{Ric}_{\alpha}{}^{\alpha} \wedge {}^* \text{Ric}_{\beta}{}^{\beta} \\ &\equiv -L_{\text{SKY}} \\ &\quad - 2 \left(\text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} - \frac{1}{4} \text{Ric}_{\alpha}{}^{\alpha} \wedge {}^* \text{Ric}_{\beta}{}^{\beta} \right) \end{aligned}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0974 (14.2.15) [conf 0.099]	315	■ 0.099 ● N +0	$\Theta^{\perp} := \pi(\hat{\Theta} - [\hat{\alpha}, \hat{\vartheta}]) = \left[\frac{\Theta}{0} \middle \frac{\hat{\ell}\Omega'}{\circ\Theta} = 0 \right],$	$\Theta^{\perp} := \pi(\hat{\Theta} - [\hat{\alpha}, \hat{\vartheta}]) = \left[\frac{\Theta}{0} \middle \frac{\hat{\ell}\Omega'}{\circ\Theta} = 0 \right],$	$\Theta^{\perp} := \pi(\hat{\Theta} - [\hat{\alpha}, \hat{\vartheta}]) = \left[\frac{\Theta}{0} \middle \frac{\hat{\ell}\varOmega'}{\pi\overset{\circ}{\Theta} = 0} \right],$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0553 (8.4.7)	180	■ 0.100 ● W +2	$\begin{aligned} L_{\text{SKY}}^{(*)} &= \frac{1}{2} \text{Tr} \left(\overset{\pm}{\Omega}^g \wedge {}^* \overset{\pm}{\Omega}^g \right) \\ &= -\frac{1}{2} R^{\alpha\beta} \wedge {}^* R_{\alpha\beta} \mp \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= -\frac{1}{4} \left(R_{\alpha\beta} \pm {}^* R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} \pm {}^* R^{\alpha\beta(*)} \right). \end{aligned}$	$\begin{aligned} L_{\text{SKY}}^{(*)} &= \frac{1}{2} \text{Tr} \left(\overset{\pm}{\Omega}^g \wedge {}^* \overset{\pm}{\Omega}^g \right) \\ &= -\frac{1}{2} R^{\alpha\beta} \wedge {}^* R_{\alpha\beta} \mp \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= -\frac{1}{4} \left(R_{\alpha\beta} \pm {}^* R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} \pm {}^* R^{\alpha\beta(*)} \right). \end{aligned}$	$\begin{aligned} L_{\text{SKY}}^{(*)} &= \frac{1}{2} \text{Tr} \left(\overset{\pm}{\varOmega}^g \wedge {}^* \overset{\pm}{\varOmega}^g \right) \\ &= -\frac{1}{2} R^{\alpha\beta} \wedge {}^* R_{\alpha\beta} \mp \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= -\frac{1}{4} \left(R_{\alpha\beta} \pm {}^* R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} \pm {}^* R^{\alpha\beta(*)} \right). \end{aligned}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0311 (4.6.3)	104	■ 0.111 ● C +6	$\begin{aligned} & \widetilde{\leftarrow}_{\widetilde{y}} \widetilde{\widetilde{D}} \widetilde{\xi}^\alpha = \widetilde{y} \rfloor \left(\widetilde{\widetilde{D}} \widetilde{\widetilde{D}} \widetilde{\xi}^\alpha \right) + \widetilde{\widetilde{D}} (\widetilde{y} \rfloor \widetilde{\widetilde{D}} \widetilde{\xi}^\alpha) \\ & = y J \left(D D \xi^\alpha + R^{(T)\alpha} \right) = 0, \end{aligned}$	$\begin{aligned} \widetilde{\widetilde{\mathbf{L}}}_{\widetilde{y}} \widetilde{\widetilde{D}} \widetilde{\xi}^\alpha &= \widetilde{y} \rfloor (\widetilde{\widetilde{D}} \widetilde{\widetilde{D}} \widetilde{\xi}^\alpha) + \widetilde{\widetilde{D}} (\widetilde{y} \rfloor \widetilde{\widetilde{D}} \widetilde{\xi}^\alpha) \\ &= y \rfloor (D D \xi^\alpha + R^{(T)\alpha}) = 0, \end{aligned}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0378 (6.2.29) [re- fined: inferred]	128	■ 0.136 ● C -66	preserve the term Pontryagin index for the classification of real associated vector bundles (\textsc{Mayer} \& \textsc{Drechsler} 1977). Nevertheless, the denotation of \textsc{Belavin}	<i>preservethetermPontryaginindexfortheclassificationofrealassociatedvectorbundles(Mayer&Drechsler1977).Nevertheless,thedenotationofBelavin</i>	$D D \vartheta^\alpha = R_\beta{}^\alpha \wedge \vartheta^\beta = 0, \quad D \overset{(\pm)(\pm)(\pm)}{D} \overset{(\pm)}{D} \overset{(\pm)}{\Pi}_\alpha = - R_\alpha{}^\beta \wedge \overset{(\pm)}{\Pi}_\beta = \pm \frac{i}{2} \ell^2 (e_\alpha \rfloor \overset{(\pm)}{\Pi}^\beta) \wedge \Sigma_\beta \cong 0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ1059 (15.2.16)	346	■ 0.147 ● C +12	$\begin{aligned} \psi = & \left\{ \begin{array}{llll} \mathcal{P}_a & \mathcal{P}_b & \mathcal{P}_c & \nu_e \\ \eta_a & \eta_b & \eta_c & e^- \\ \lambda_a & \lambda_b & \lambda_c & \mu^- \\ c_a & c_b & c_c & \nu_\mu \end{array} \right\} \begin{array}{l} \text{up} \\ \text{down} \\ \text{strange} \\ \text{charm} \end{array} \begin{array}{l} \uparrow \\ \text{flavors} \\ \downarrow \end{array} \\ & \text{red} \quad \text{yellow} \quad \text{blue} \quad \text{lilac} \end{aligned}$	$\psi = \left[\begin{array}{ccc c} \mathcal{P}_a & \mathcal{P}_b & \mathcal{P}_c & \nu_e \\ \eta_a & \eta_b & \eta_c & e^- \\ \lambda_a & \lambda_b & \lambda_c & \mu^- \\ c_a & c_b & c_c & \nu_\mu \end{array} \right] \begin{array}{l} \text{up} \\ \text{down} \\ \text{strange} \\ \text{charm} \end{array} \begin{array}{l} \uparrow \\ \text{flavors} \\ \downarrow \end{array}$	$\psi = \left[\begin{array}{ccc c} \mathcal{P}_a & \mathcal{P}_b & \mathcal{P}_c & \nu_e \\ \eta_a & \eta_b & \eta_c & e^- \\ \lambda_a & \lambda_b & \lambda_c & \mu^- \\ c_a & c_b & c_c & \nu_\mu \end{array} \right] \begin{array}{l} \text{up} \\ \text{down} \\ \text{strange} \\ \text{charm} \end{array} \begin{array}{l} \uparrow \\ \text{flavors} \\ \downarrow \end{array}$ red yellow blue lilac ← (body) – colors →
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0972 (14.2.13) [re- fined: inferred]	314	■ 0.153 ● N +0	$\overset{(1)}{D} \alpha := \overset{(0)}{d} \alpha - \omega^g \wedge \alpha - \alpha \wedge \omega^g$	$\overset{(1)}{D} \alpha := \overset{(0)}{d} \alpha - \omega^g \wedge \alpha - \alpha \wedge \omega^g$	$\overset{(\circ)}{D} \alpha := \overset{(\circ)}{d} \alpha - \omega^g \wedge \alpha - \alpha \wedge \overset{\circ}{\omega}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00264 (4.3.29)	095	■ 0.154 ● W +2	$\Gamma_{ij}^{\prime} = \left\{ \begin{array}{c} \kappa \\ ij \end{array} \right\} - K_{ij}^{\cdot\kappa},$	$\Gamma_{ij}^{\cdot,K} = \left\{ \begin{array}{c} \kappa \\ ij \end{array} \right\} - K_{ij}^{\cdot,K},$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00292 (4.5.3)	100	■ 0.165 ● W +2	$\begin{aligned} \mathfrak{f}_{\xi} L &= \left(\mathfrak{f}_{\xi} g_{\alpha\beta} \right) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \left(\mathfrak{f}_{\xi} \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} \\ &+ \left(\mathfrak{f}_{\xi} \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\mathfrak{f}_{\xi} D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \\ &= D \left[\left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \right] \\ &- \left(\xi \rfloor \vartheta^{\alpha} \right) D \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor T^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor R_{\beta}^{\gamma} \right) \wedge I^{\beta}{}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} \\ &+ \left(\xi \rfloor D\Psi \right) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p \left(\xi \rfloor \Psi \right) \wedge D \frac{\delta L}{\delta \Psi} \end{aligned}$	$\begin{aligned} \mathfrak{f}_{\xi} L &= \left(\mathfrak{f}_{\xi} g_{\alpha\beta} \right) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \left(\mathfrak{f}_{\xi} \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} \\ &+ \left(\mathfrak{f}_{\xi} \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\mathfrak{f}_{\xi} D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \\ &= D \left[\left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \right] \\ &- \left(\xi \rfloor \vartheta^{\alpha} \right) D \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor T^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor R_{\beta}^{\gamma} \right) \wedge I^{\beta}{}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} \\ &+ \left(\xi \rfloor D\Psi \right) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p \left(\xi \rfloor \Psi \right) \wedge D \frac{\delta L}{\delta \Psi} \end{aligned}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00734 (11.1.1)	237	■ 0.220 ● W +1	$\pi_{\{1\}} \left(S_{0_{\circ}}(s, n-s), e \right) = \begin{cases} \mathbb{Z}_2 & n=2k \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2 & n=4k \\ \mathbb{Z}_4 & n=2(2k+1) \end{cases}$	$\pi_1(SO_{\circ}(s, n-s), e) = \begin{cases} \mathbb{Z}_2 & n=2k \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2 & n=4k \\ \mathbb{Z}_4 & n=2(2k+1) \end{cases}$	$\pi_1(SO_{\circ}(s, n-s), e) = \begin{cases} \mathbb{Z}_2 & n=2k \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2 & n=4k \\ \mathbb{Z}_4 & n=2(2k+1). \end{cases}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0961 (14.2.1)	312	■ 0.220 ● C +3	$\hat{\vartheta}(m, g) = \underbrace{\left[\begin{array}{c c} \vartheta & \hat{\ell}\omega \\ \hline 0 & \grave{\vartheta} \end{array} \right]}_4 \}^4 K$	$\hat{\vartheta} = \hat{\vartheta}(m, g) = \left[\begin{array}{c c} \vartheta & \hat{\ell}\omega \\ \hline 0 & \grave{\vartheta} \end{array} \right] \}^4 K$	$\hat{\vartheta} = \hat{\vartheta}(m, g) = \left[\begin{array}{c c} \vartheta & \hat{\ell}\omega \\ \hline 0 & \grave{\vartheta} \end{array} \right] \}^4 K$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0211 (4.2.3)	085	■ 0.223 ● C +4	$\begin{aligned} & * \widetilde{\Omega} = \chi \left(d\widetilde{\omega} - \frac{1}{2} [\widetilde{\omega}, \widetilde{\omega}] \right) \\ & = d\omega^L - \omega^L \wedge \omega^L + d\omega^T - \omega^T \wedge \omega^L - \omega^L \wedge \omega^T + \frac{1}{2} [\omega^T, \omega^T] \\ & = \Omega^L + \Omega^T \end{aligned}$	$\begin{aligned} * \widetilde{\Omega} &= \chi \left(d\widetilde{\omega} - \frac{1}{2} [\widetilde{\omega}, \widetilde{\omega}] \right) \\ &= d\omega^L - \omega^L \wedge \omega^L + d\omega^T - \omega^T \wedge \omega^L - \omega^L \wedge \omega^T + \frac{1}{2} [\omega^T, \omega^T] \\ &= \Omega^L + \Omega^T \end{aligned}$	$\begin{aligned} * \widetilde{\Omega} &= \chi \left(d\widetilde{\omega} - \frac{1}{2} [\widetilde{\omega}, \widetilde{\omega}] \right) \\ &= d\omega^L - \omega^L \wedge \omega^L + d\omega^T - \omega^T \wedge \omega^L - \omega^L \wedge \omega^T + \frac{1}{2} [\omega^T, \omega^T] \\ &= \Omega^L + \Omega^T \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0266 (4.3.31)	095	■ 0.227 ● W -2	$R_{ijc}^{::d} = R_{ijc}^{\{!...^d} - 2D_{[i}K_{j]c}^{::d} + 2K_{[i c}^{::h}K_{j]h}^{::d}$	$R_{ijc}^{::d} = R_{ijc}^{\{!...^d} - 2D_{[i}K_{j]c}^{::d} + 2K_{[i c}^{::h}K_{j]h}^{::d}$	$R_{ijc}^{::d} = R_{ijc}^{\{!...^d} - 2D_{[i}K_{j]c}^{::d} + 2K_{[i c}^{::h}K_{j]h}^{::d}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0286 (4.4.15)	099	■ 0.229 ● K +0	$D \Pi_{[]}^T - E_{[]} = \Sigma_{[]}$	$D \Pi_{[]}^T - E_{[]} = \Sigma_{[]}$	$D \Pi_{[]}^T - E_{[]} = \Sigma_{[]}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5-0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0337 (5.3.14)	114	■ 0.239 ● C +4	$\begin{aligned} R^{\{\alpha\beta\}} \wedge \eta_{\alpha\beta} &\equiv R^{\alpha\beta} \wedge \eta_{\alpha\beta} - K^{\alpha\mu} \wedge K_{\mu}{}^{\beta} \wedge \eta_{\alpha\beta} + K^{\alpha\beta} \wedge T^{\gamma} \wedge \eta_{\alpha\beta\gamma} \\ &\quad + d(K^{\alpha\beta} \wedge \eta_{\alpha\beta}) \\ &= R^{\alpha\beta} \wedge \eta_{\alpha\beta} + T^{\alpha} \wedge * \left(-^{(1)}T_{\alpha} + 2^{(2)}T_{\alpha} + \frac{1}{2}^{(3)}T_{\alpha} \right) + 2dC_{\text{TT}}* \end{aligned}$	$\begin{aligned} R^{\{\alpha\beta\}} \wedge \eta_{\alpha\beta} &\equiv R^{\alpha\beta} \wedge \eta_{\alpha\beta} - K^{\alpha\mu} \wedge K_{\mu}{}^{\beta} \wedge \eta_{\alpha\beta} + K^{\alpha\beta} \wedge T^{\gamma} \wedge \eta_{\alpha\beta\gamma} \\ &\quad + d(K^{\alpha\beta} \wedge \eta_{\alpha\beta}) \\ &= R^{\alpha\beta} \wedge \eta_{\alpha\beta} + T^{\alpha} \wedge * \left(-^{(1)}T_{\alpha} + 2^{(2)}T_{\alpha} + \frac{1}{2}^{(3)}T_{\alpha} \right) + 2dC_{\text{TT}}* \end{aligned} \quad (5.3)$	$\begin{aligned} R^{ \alpha\beta} \wedge \eta_{\alpha\beta} &\equiv R^{\alpha\beta} \wedge \eta_{\alpha\beta} - K^{\alpha\mu} \wedge K_{\mu}{}^{\beta} \wedge \eta_{\alpha\beta} + K^{\alpha\beta} \wedge T^{\gamma} \wedge \eta_{\alpha\beta\gamma} \\ &\quad + d(K^{\alpha\beta} \wedge \eta_{\alpha\beta}) \\ &= R^{\alpha\beta} \wedge \eta_{\alpha\beta} + T^{\alpha} \wedge * \left(-^{(1)}T_{\alpha} + 2^{(2)}T_{\alpha} + \frac{1}{2}^{(3)}T_{\alpha} \right) + 2dC_{\text{TT}}* \end{aligned} \quad (5.3)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0835 (12.2.3)	271	■ 0.247 ● C -3	$\begin{aligned} L_{\text{D}} &= L(\gamma, \psi, D^{(f)}\psi) - \frac{i}{2} \bar{\psi} (*\gamma \wedge K - K \wedge *\gamma) \psi + A \wedge j \\ &= L(\gamma, \psi, D^{(\dagger)}\psi) + \frac{1}{4} \mathcal{A} \wedge j_5 + A \wedge j \\ &= L(\gamma, \psi, D^{(i)}\psi) - T^{\alpha} \wedge \mu_{\alpha} + A \wedge j. \end{aligned}$	$\begin{aligned} L_{\text{D}} &= L(\gamma, \psi, D^{(f)}\psi) - \frac{i}{2} \bar{\psi} (*\gamma \wedge K - K \wedge *\gamma) \psi + A \wedge j \\ &= L(\gamma, \psi, D^{(\dagger)}\psi) + \frac{1}{4} \mathcal{A} \wedge j_5 + A \wedge j \\ &= L(\gamma, \psi, D^{(i)}\psi) - T^{\alpha} \wedge \mu_{\alpha} + A \wedge j. \end{aligned}$	$\begin{aligned} L_{\text{D}} &= L(\gamma, \psi, D^{ \psi}) - \frac{i}{2} \bar{\psi} (*\gamma \wedge K - K \wedge *\gamma) \psi + A \wedge j \\ &= L(\gamma, \psi, D^{ \psi}) + \frac{1}{4} \mathcal{A} \wedge j_5 + A \wedge j \\ &= L(\gamma, \psi, D^{ \psi}) - T^{\alpha} \wedge \mu_{\alpha} + A \wedge j. \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0787 (11.3.6)	248	■ 0.258 ● C +3	$i \gamma \wedge * D^{[1} \psi - 6 \ell^{*2} \gamma^{n+1} \gamma \wedge * (\bar{\psi} \gamma^{n+1} \gamma \psi) \psi - m \psi \eta = 0$	$i \gamma \wedge * D^{[1} \psi - 6 \ell^{*2} \gamma^{n+1} \gamma \wedge * (\bar{\psi} \gamma^{n+1} \gamma \psi) \psi - m \psi \eta = 0$	$i \gamma \wedge * D^{ \psi} - 6 \ell^{*2} \gamma^{n+1} \gamma \wedge * (\bar{\psi} \gamma^{n+1} \gamma \psi) \psi - m \psi \eta = 0 \Big ,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1078 (A.3)	365	■ 0.258 ● W +1	$\begin{aligned} &\left(\sigma^a \right)^{\text{T}} := \left(\begin{array}{l} \sigma^0 \\ \sigma^1 \\ \sigma^2 \end{array} \right) \\ &\quad \sigma^0 := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \equiv \mathbb{1}, \quad \sigma^1 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 := \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \end{aligned}$	$\begin{aligned} &\left(\sigma^a \right)^{\text{T}} := \begin{pmatrix} \sigma^0, \sigma^k \\ \sigma^0 \\ \sigma^2 \end{pmatrix} \quad k = 1, 2, 3 \\ &\quad \sigma^0 := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \equiv \mathbb{1}, \quad \sigma^1 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 := \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \end{aligned}$	$\begin{aligned} &\left(\sigma^a \right)^{\text{T}} := (\sigma^0, \sigma^k) \quad k = 1, 2, 3. \\ &\quad \sigma^0 := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \equiv \mathbb{1}, \quad \sigma^1 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\ &\quad \sigma^2 := \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \end{aligned}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0018 (2.3.4)	037	■ 0.261 ● N +0	$\delta:\left\{\begin{array}{lll} G\times P&\rightarrow&P\\ w&\psi&\psi\\ g_p\cdot&p_\circ=p& \end{array}\right e\cdot p_\circ=p_\circ\in P\right\}.$	$\delta:\left\{\begin{array}{lll} G\times P&\rightarrow&P\\ w&\psi&\psi\\ g_p\cdot&p_\circ=p& \end{array}\right e\cdot p_\circ=p_\circ\in P\right\}.$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0544 (8.3.5)	177	■ 0.261 ● W +1	$\left.\left.\left.\exp\left(\left\lfloor z\right\rfloor\right):=\mathbf{1}+z+\frac{1}{2!}z^2+\frac{1}{3!}z^3+\dots\right.\right.\right.$	$\exp(\zeta\rfloor):=\mathbf{1}+\zeta\rfloor+\frac{1}{2!}\zeta\rfloor\zeta\rfloor+\frac{1}{3!}\zeta\rfloor\zeta\rfloor\zeta\rfloor+\cdots.$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0605 (9.2.4)	199	■ 0.263 ● N +0	$\chi(M):=\sum_{p=0}^n(-1)^pb_p=\int_M\pi^*(\gamma_2^g)$	$\chi(M):=\sum_{p=0}^n(-1)^pb_p=\int_M\pi^*(\gamma_2^g)$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0627 (10.1.15)	211	■ 0.271 ● N +0	$E_{\alpha}:=\frac{\partial L}{\partial \vartheta^{\alpha}}=e_{\alpha}\lrcorner L+(e_{\alpha}\lrcorner T^{\beta})\wedge H_{\beta}+2(-1)^s(e_{\alpha}\lrcorner R^{\star\beta})\wedge H_{\beta}^{\star},$	$E_{\alpha}:=\frac{\partial L}{\partial \vartheta^{\alpha}}=e_{\alpha}\lrcorner L+(e_{\alpha}\lrcorner T^{\beta})\wedge H_{\beta}+2(-1)^s(e_{\alpha}\lrcorner R^{\star\beta})\wedge H_{\beta}^{\star},$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0979 (14.2.20)	316	■ 0.272 ● N +0	$\hat{\ell}=\left(\begin{array}{c}+\\-\end{array}\right)\ell^*/\sqrt{\alpha_g}=\left(\begin{array}{c}+\\-\end{array}\right)\ell^*\frac{\sqrt{\hbar c}}{g}$	$\hat{\ell}=\left(\begin{array}{c}+\\-\end{array}\right)\ell^*/\sqrt{\alpha_g}=\left(\begin{array}{c}+\\-\end{array}\right)\ell^*\frac{\sqrt{\hbar c}}{g}$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00654 (10.3.11)	216	■ 0.274 ● N +0	$\left.\left.\left.E_{\alpha}=e_{\alpha}\right\downarrow\right\downarrow L+\left[\left(e_{\alpha}\right\downarrow T^{\beta}\right)+\left(-1\right)^s2\delta\ell\left(e_{\alpha}\right\downarrow R^{*\beta}\right)\right]\wedge H_{\beta}+\left(-1\right)^s\frac{2\gamma}{\ell}\left(e_{\alpha}\right\downarrow R^{*\beta}\right)\wedge\vartheta_{\beta},$	$E_{\alpha}=e_{\alpha}\downarrow L+\left[\left(e_{\alpha}\downarrow T^{\beta}\right)+\left(-1\right)^s2\delta\ell\left(e_{\alpha}\downarrow R^{*\beta}\right)\right]\wedge H_{\beta}+\left(-1\right)^s\frac{2\gamma}{\ell}\left(e_{\alpha}\downarrow R^{*\beta}\right)\wedge\vartheta_{\beta},$	$E_{\alpha}=e_{\alpha}\downarrow L+\left[\left(e_{\alpha}\downarrow T^{\beta}\right)+\left(-1\right)^s2\delta\ell\left(e_{\alpha}\downarrow R^{*\beta}\right)\right]\wedge H_{\beta}+\left(-1\right)^s\frac{2\gamma}{\ell}\left(e_{\alpha}\downarrow R^{*\beta}\right)\wedge\vartheta_{\beta},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00260 (4.3.33)	095	■ 0.276 ● W -2	$R_{\{i[j\kappa l]\}^{\{l\}}}\equiv 0$	$R_{i[j\kappa l]}^{(l)}\equiv 0$	$R_{i[j\kappa l]}^{(l)}\equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00639 (10.2.10)	213	■ 0.278 ● N -1	$R_{\{\alpha\}^{\{i\star\}}=-\Lambda_{\mathrm{eff}}\}\eta_{\alpha},$	$R_{\alpha}^{i\star}=-\Lambda_{\mathrm{eff}}\eta_{\alpha},$	$R_{\alpha}^{(i)\star}=-\Lambda_{\mathrm{eff}}\eta_{\alpha},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00527 (7.7.47)	168	■ 0.282 ● N -1	$\begin{aligned} L_{\mathrm{Pontr}}:=& \mathrm{d} C_{\mathrm{RR}}=\frac{1}{2} R_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha}=\frac{1}{2} R_{\alpha \beta}^{\{\}} \wedge R^{\{j \alpha \beta} \\ & -\frac{1}{12} d\left[\ast \mathscr{A} \wedge R^{\{\}}-\frac{1}{3} \mathscr{A} \wedge d \mathscr{A}+\frac{1}{9} \ast \mathscr{A} \wedge^{\ast}\left(\mathscr{A} \wedge^{\ast} \mathscr{A}\right)\right] . \end{aligned}$	$L_{\mathrm{Pontr}}:=dC_{\mathrm{RR}}=\frac{1}{2}R_{\alpha}^{\beta}\wedge R_{\beta}^{\alpha}=-\frac{1}{2}R_{\alpha\beta}^{\{\}}\wedge R^{\{j\alpha\beta}$ $-\frac{1}{12}d\left[\ast\mathscr{A}\wedge R^{\{\}}-\frac{1}{3}\mathscr{A}\wedge d\mathscr{A}+\frac{1}{9}\ast\mathscr{A}\wedge^{\ast}(\mathscr{A}\wedge^{\ast}\mathscr{A})\right].$	$L_{\mathrm{Pontr}}:=dC_{\mathrm{RR}}=\frac{1}{2}R_{\alpha}^{\beta}\wedge R_{\beta}^{\alpha}=-\frac{1}{2}R_{\alpha\beta}^{\{\}}\wedge R^{\{\} \alpha\beta}$ $-\frac{1}{12}d\left[\ast\mathscr{A}\wedge R^{\{\}}-\frac{1}{3}\mathscr{A}\wedge d\mathscr{A}+\frac{1}{9}\ast\mathscr{A}\wedge^{\ast}(\mathscr{A}\wedge^{\ast}\mathscr{A})\right].$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00942 (14.1.23)	306	■ 0.287 ● W +0	$\xi\wedge^*\xi=\pm\frac{\Omega\wedge^*\Omega\mathrm{Tr}^*(\Omega\wedge^*\Omega)-(-1)^{2z}\Omega\wedge\Omega\mathrm{Tr}^*(\Omega\wedge\Omega)}{\sqrt{(\mathrm{Tr}^*(\Omega\wedge^*\Omega))^2+(-1)^{2z}(\mathrm{Tr}^*(\Omega\wedge\Omega))^2}}.$	$\xi\wedge^*\xi=\pm\frac{\Omega\wedge^*\Omega\mathrm{Tr}^*(\Omega\wedge^*\Omega)-(-1)^{2z}\Omega\wedge\Omega\mathrm{Tr}^*(\Omega\wedge\Omega)}{\sqrt{(\mathrm{Tr}^*(\Omega\wedge^*\Omega))^2+(-1)^{2z}(\mathrm{Tr}^*(\Omega\wedge\Omega))^2}}.$	$\xi\wedge^*\xi=\pm\frac{\Omega\wedge^*\Omega\mathrm{Tr}^*(\Omega\wedge^*\Omega)-(-1)^{2z}\Omega\wedge\Omega\mathrm{Tr}^*(\Omega\wedge\Omega)}{\sqrt{(\mathrm{Tr}^*(\Omega\wedge^*\Omega))^2+(-1)^{2z}(\mathrm{Tr}^*(\Omega\wedge\Omega))^2}}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0440 (6.8.11)	143	■ 0.288 ● N +0	$\left.D\,\Sigma_{\alpha}\right.\cong\left(e_{\alpha}\downharpoonleft T^{\gamma}\right)\wedge\Sigma_{\gamma}+\left(e_{\alpha}\downharpoonleft R^{\gamma\delta}\right)\wedge\tau_{\gamma\delta}$	$D\Sigma_{\alpha}\cong(e_{\alpha}\rfloor T^{\gamma})\wedge\Sigma_{\gamma}+(e_{\alpha}\rfloor R^{\gamma\delta})\wedge\tau_{\gamma\delta}$	$D\Sigma_{\alpha}\cong(e_{\alpha}\rfloor T^{\gamma})\wedge\Sigma_{\gamma}+(e_{\alpha}\rfloor R^{\gamma\delta})\wedge\tau_{\gamma\delta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0315 (4.6.7)	105	■ 0.292 ● N +1	$\Delta\Phi(\widetilde{\Gamma},\gamma)\simeq\frac{i}{\hbar}\int_S\Big(R^{(T)\alpha}P_{\alpha}+R^{(L)\beta}_{\alpha}L^{\alpha}_{\beta}\Big)$	$\Delta\Phi(\widetilde{\Gamma},\gamma)\simeq\frac{i}{\hbar}\int_S\Big(R^{(T)\alpha}P_{\alpha}+R^{(L)\beta}_{\alpha}L^{\alpha}_{\beta}\Big)$	$\Delta\Phi(\widetilde{\Gamma},\gamma)\simeq\frac{i}{\hbar}\int_S(R^{(T)\alpha}P_{\alpha}+R^{(L)\beta}_{\alpha}L^{\alpha}_{\beta}).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0965 (14.2.5)	313	■ 0.299 ● N +0	$\begin{aligned} &\stackrel{\circ}{\stackrel{\circ}{\Theta}}:=\dot{\stackrel{\circ}{\vartheta}}-[\stackrel{\circ}{\omega},\stackrel{\circ}{\vartheta}]=\left(1-2b\right)\stackrel{\circ}{\vartheta}\wedge\stackrel{\circ}{\vartheta},\\ &\stackrel{\circ}{\stackrel{\circ}{\Omega}}:=\dot{\stackrel{\circ}{\omega}}-\stackrel{\circ}{\omega}\wedge\stackrel{\circ}{\omega}=b\left(1-b\right)\stackrel{\circ}{\vartheta}\wedge\stackrel{\circ}{\vartheta}. \end{aligned}$	$\begin{aligned} \stackrel{\circ}{\Theta} &:= \dot{\stackrel{\circ}{\vartheta}} - [\stackrel{\circ}{\omega}, \stackrel{\circ}{\vartheta}] = (1 - 2b) \stackrel{\circ}{\vartheta} \wedge \stackrel{\circ}{\vartheta}, \\ \stackrel{\circ}{\Omega} &:= \dot{\stackrel{\circ}{\omega}} - \stackrel{\circ}{\omega} \wedge \stackrel{\circ}{\omega} = b(1 - b) \stackrel{\circ}{\vartheta} \wedge \stackrel{\circ}{\vartheta}. \end{aligned}$	$\begin{aligned} \stackrel{\circ}{\Theta} &:= \dot{\stackrel{\circ}{\vartheta}} - [\stackrel{\circ}{\omega}, \stackrel{\circ}{\vartheta}] = (1 - 2b) \stackrel{\circ}{\vartheta} \wedge \stackrel{\circ}{\vartheta}, \\ \stackrel{\circ}{\Omega} &:= \dot{\stackrel{\circ}{\omega}} - \stackrel{\circ}{\omega} \wedge \stackrel{\circ}{\omega} = b(1 - b) \stackrel{\circ}{\vartheta} \wedge \stackrel{\circ}{\vartheta}. \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0226 (4.2.19)	087	■ 0.299 ● C +4	$\begin{aligned} A^{-1}\omega^{\mathrm{T}}&=A^{-1}\widetilde{\vartheta}-G^{-1}DG^{A^{-1}}\widetilde{\varphi}\\ &=G^{-1}\widetilde{\vartheta}-G^{-1}D\widetilde{\varphi}, \end{aligned}$	$\begin{aligned} A^{-1}\omega^{\mathrm{T}} &= A^{-1}\widetilde{\vartheta} - G^{-1}DG^{A^{-1}}\widetilde{\varphi} \\ &= G^{-1}\widetilde{\vartheta} - G^{-1}D\widetilde{\varphi}, \end{aligned}$	$\begin{aligned} A^{-1}\omega^{\mathrm{T}} &= A^{-1}\widetilde{\vartheta} - G^{-1}DG^{A^{-1}}\widetilde{\varphi} \\ &= G^{-1}\widetilde{\vartheta} - G^{-1}D\widetilde{\varphi}, \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0407 (6.4.27)	135	■ 0.315 ● N +0	$\Psi_{\mathbf{i}_{\Lambda}}(A)=\exp\left(\frac{3}{\Lambda}\int_{M_3}\stackrel{(\pm)}{C}_{\mathrm{RR}}\left(\right)\right),$	$\Psi_{\Lambda}(A)=\exp\left(\frac{3}{\Lambda}\int_{M_3}\stackrel{(\pm)}{C}_{\mathrm{RR}}\left(\right)\right),$	$\Psi_A(A)=\exp\left(\frac{3}{\Lambda}\int_{M_3}\stackrel{(\pm)}{C}_{\mathrm{RR}}{}^{(\mathfrak{l})}\right),$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0262 (4.3.27)	094	■ 0.399 ● C +3	$\begin{aligned} \Omega^g &= \Omega^{\{\}} - dK + K \wedge \omega^{\{\}} + \omega^{\{\}} \wedge K - K \wedge K \\ &= \Omega^{\{\}} - D^{\{\}} K - K \wedge K = \Omega^{\{\}} - DK + K \wedge K \end{aligned}$	$\Omega^g = \Omega^{\{\}} - dK + K \wedge \omega^{\{\}} + \omega^{\{\}} \wedge K - K \wedge K$ $= \Omega^{\{\}} - D^{\{\}} K - K \wedge K = \Omega^{\{\}} - DK + K \wedge K$	$\Omega^g = \Omega^{\{\}} - dK + K \wedge \omega^{\{\}} + \omega^{\{\}} \wedge K - K \wedge K$ $= \Omega^{\{\}} - D^{\{\}} K - K \wedge K = \Omega^{\{\}} - DK + K \wedge K$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0570 (8.6.8)	184	■ 0.403 ● K +0	$\left. E_{\alpha} := \partial L / \partial \vartheta^{\alpha} \right L + \left(e_{\alpha} \rfloor T^{\beta} \right) \wedge H_{\beta} + \left(e_{\alpha} \rfloor R^{\beta \gamma} \right) \wedge H_{\beta \gamma}$	$E_{\alpha} := \partial L / \partial \vartheta^{\alpha} = e_{\alpha} \rfloor L + \left(e_{\alpha} \rfloor T^{\beta} \right) \wedge H_{\beta} + \left(e_{\alpha} \rfloor R^{\beta \gamma} \right) \wedge H_{\beta \gamma}$	$E_{\alpha} := \partial L / \partial \vartheta^{\alpha} = e_{\alpha} \rfloor L + \left(e_{\alpha} \rfloor T^{\beta} \right) \wedge H_{\beta} + \left(e_{\alpha} \rfloor R^{\beta \gamma} \right) \wedge H_{\beta \gamma}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0306 (4.5.16)	103	■ 0.406 ● N +0	$\left. \widetilde{\varepsilon}_{\mathrm{MA}} := \varepsilon_{\mathrm{RC}} + \xi \rfloor L \right L \cong - \left(\mathcal{L}_{\xi} \Gamma_{\beta}^{\gamma} \right) \wedge H^{\beta}{}_{\gamma} + dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}} := \varepsilon_{\mathrm{RC}} + \xi \rfloor L \cong - \left(\mathcal{L}_{\xi} \Gamma_{\beta}^{\gamma} \right) \wedge H^{\beta}{}_{\gamma} + dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}} := \varepsilon_{\mathrm{RC}} + \xi \rfloor L \cong - \left(\mathcal{L}_{\xi} \Gamma_{\beta}^{\gamma} \right) \wedge H^{\beta}{}_{\gamma} + dH_{\infty}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0336 (5.3.13)	114	■ 0.407 ● N -1	$R^{\alpha \beta} = R^{\{j \alpha \beta\}} - D^{\{\}} K^{\alpha}{}_{\mu} \wedge K^{\mu \beta}$	$R^{\alpha \beta} = R^{\{j \alpha \beta} - D^{\{\}} K^{\alpha \beta} - K^{\alpha}{}_{\mu} \wedge K^{\mu \beta}$	$R^{\alpha \beta} = R^{\{\} \alpha \beta} - D^{\{\}} K^{\alpha \beta} - K^{\alpha}{}_{\mu} \wedge K^{\mu \beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1136 (C.4)	375	■ 0.409 ● N +0	$c_{ij}^k = -c_{ji}^k$	$c_{ij}^k = -c_{ji}^k$	$c_{ij}^k = -c_{ji}^k$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._Mielke_ __Z-Library__ EQ0861 (12.4.8)	279	■0.420 ● N -1	$\begin{aligned} d\,C_{\mathrm{RR}} &= -\operatorname{Tr}(\Omega \wedge \Omega) \\ &= \frac{1}{2}R_{\alpha\beta}^{\{\}} \wedge R^{\{\beta\alpha\beta\}} \\ &+ \frac{1}{12}d\left[{}^*\mathcal{A} \wedge R^{\{\}} - \frac{1}{3}\mathcal{A} \wedge d\mathcal{A} + \frac{1}{9}{}^*\mathcal{A} \wedge {}^*(\mathcal{A} \wedge {}^*\mathcal{A})\right]. \end{aligned}$	$dC_{\mathrm{RR}} = -\mathrm{Tr}(\Omega \wedge \Omega)$ $= \frac{1}{2}R_{\alpha\beta}^{\{\}} \wedge R^{\{\beta\alpha\beta\}}$ $+ \frac{1}{12}d\left[{}^*\mathcal{A} \wedge R^{\{\}} - \frac{1}{3}\mathcal{A} \wedge d\mathcal{A} + \frac{1}{9}{}^*\mathcal{A} \wedge {}^*(\mathcal{A} \wedge {}^*\mathcal{A})\right].$	$dC_{\mathrm{RR}} = -\mathrm{Tr}(\varOmega \wedge \varOmega)$ $= \frac{1}{2}R_{\alpha\beta}^{\{\}} \wedge R^{\{\}\alpha\beta}$ $+ \frac{1}{12}d\left[{}^*\mathcal{A} \wedge R^{\{\}} - \frac{1}{3}\mathcal{A} \wedge d\mathcal{A} + \frac{1}{9}{}^*\mathcal{A} \wedge {}^*(\mathcal{A} \wedge {}^*\mathcal{A})\right].$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._Mielke_ __Z-Library__ EQ0451 (7.2.4)	151	■0.424 ● N +0	$\stackrel{+}{\Omega} := \frac{1}{2}\left(\Omega^g \pm {}^*\Omega^{g(*)}\right),$	$\stackrel{+g}{\Omega} := \frac{1}{2}\left(\Omega^g \pm {}^*\Omega^{g(*)}\right),$	$\stackrel{\pm}{\varOmega}^g := \frac{1}{2}\left(\varOmega^g \pm {}^*\varOmega^{g(*)}\right),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._Mielke_ __Z-Library__ EQ0372 (6.2.23)	127	■0.437 ● C -3	$\left.\stackrel{(\pm)}{\Pi}_\alpha \mp \frac{i}{4}\ell^2 e_\alpha\right]\left(\stackrel{(+)}{\Pi}^\beta \wedge \stackrel{(-)}{\Pi}_\beta\right) = \Sigma_\alpha,$	$\stackrel{(\pm)}{D}\stackrel{(\pm)}{\Pi}_\alpha \mp \frac{i}{4}\ell^2 e_\alpha\left]\left(\stackrel{(+)}{\Pi}^\beta \wedge \stackrel{(-)}{\Pi}_\beta\right) = \Sigma_\alpha,$	$\stackrel{(\pm)}{D}\stackrel{(\pm)}{\Pi}_\alpha \mp \frac{i}{4}\ell^2 e_\alpha\left]\left(\stackrel{(+)}{\Pi}^\beta \wedge \stackrel{(-)}{\Pi}_\beta\right) = \Sigma_\alpha,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._Mielke_ __Z-Library__ EQ0997 (14.3.10)	322	■0.442 ● N -1	$D^{\circ}(\text{int})=\stackrel{\circ}{D}=\stackrel{\circ}{\gamma}\wedge{}^* \stackrel{\circ}{D}=\stackrel{\circ}{\gamma}\wedge{}^*(\stackrel{\circ}{d}+i\stackrel{\circ}{\tilde{\omega}})$	$D^{(\mathrm{int})}=\stackrel{\circ}{\gamma}\wedge{}^* \stackrel{\circ}{D}=\stackrel{\circ}{\gamma}\wedge{}^*(\stackrel{\circ}{d}+i\stackrel{\circ}{\tilde{\omega}})$	$\mathcal{D}^{(int)}=\stackrel{\circ}{\gamma}\wedge{}^* \stackrel{\circ}{D}=\stackrel{\circ}{\gamma}\wedge{}^*(\stackrel{\circ}{d}+i\stackrel{\circ}{\tilde{\omega}})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._Mielke_ __Z-Library__ EQ0474 (7.4.2)	156	■0.444 ● N -1	$R_{\alpha\beta} \equiv C_{\alpha\beta} + \frac{1}{2}Ri^{\tau}_{\alpha\beta} - \frac{R}{12}\vartheta_{\alpha} \wedge \vartheta_{\beta},$	$R_{\alpha\beta} \equiv C_{\alpha\beta} + \frac{1}{2}Ri^{\tau}_{\alpha\beta} - \frac{R}{12}\vartheta_{\alpha} \wedge \vartheta_{\beta},$	$R_{\alpha\beta} \equiv C_{\alpha\beta} + \frac{1}{2}Rj^{\tau}_{\alpha\beta} - \frac{R}{12}\vartheta_{\alpha} \wedge \vartheta_{\beta},$

Conf.

MathPix confidence:

■

≥0.9

■

0.5–0.9

■

<0.5

— no value

Residual

render vs scan (inkdrill):

●

C component

●

W weak

●

S stable

●

N noise

●

K clean

●

not measured

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0681 (10.3.38)	221	■ 0.452 ● C +4	$\begin{gathered} \chi=\theta_{\mathrm{L}}\,\varepsilon-\tilde{\theta}_{\mathrm{L}}\,\tilde{\varepsilon},\quad \Lambda=\frac{(-1)^s}{4\ell^2}\left(\theta_{\mathrm{L}}\varepsilon^3-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^3\right),\quad \theta_{\mathrm{T}}=\frac{(-1)^s}{2}\left(\theta_{\mathrm{L}}\varepsilon^2-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^2\right),\\ \Delta\theta_{\mathrm{L}}=\theta_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}},\quad \Delta\theta_{\mathrm{TL}}=\frac{1}{2}\left(\theta_{\mathrm{L}}\varepsilon-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}\right). \end{gathered}$	$\chi=\theta_{\mathrm{L}}\varepsilon-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon},\quad \Lambda=\frac{(-1)^s}{4\ell^2}\left(\theta_{\mathrm{L}}\varepsilon^3-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^3\right),\quad \theta_{\mathrm{T}}=\frac{(-1)^s}{2}\left(\theta_{\mathrm{L}}\varepsilon^2-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^2\right),$ $\Delta\theta_{\mathrm{L}}=\theta_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}},\quad \Delta\theta_{\mathrm{TL}}=\frac{1}{2}\left(\theta_{\mathrm{L}}\varepsilon-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}\right).$	$\chi=\theta_{\mathrm{L}}\varepsilon-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon},\quad \Lambda=\frac{(-1)^s}{4\ell^2}(\theta_{\mathrm{L}}\varepsilon^3-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^3),\quad \theta_{\mathrm{T}}=\frac{(-1)^s}{2}(\theta_{\mathrm{L}}\varepsilon^2-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^2),$ $\Delta\theta_{\mathrm{L}}=\theta_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}},\quad \Delta\theta_{\mathrm{TL}}=\frac{1}{2}(\theta_{\mathrm{L}}\varepsilon-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0845 (12.2.13)	273	■ 0.457 ● N +0	$C_{\mathrm{TT}}\cong\frac{1}{4}j_5,\quad \mathscr{A}=2\ell^2\ast C_{\mathrm{TT}}=(\ell^2/2)\,\bar{\psi}\gamma_5\gamma\psi.$	$C_{\mathrm{TT}}\cong\frac{1}{4}j_5,\quad \mathscr{A}=2\ell^2\ast C_{\mathrm{TT}}=(\ell^2/2)\,\bar{\psi}\gamma_5\gamma\psi.$	$C_{\mathrm{TT}}\cong\frac{1}{4}j_5,\quad \mathscr{A}=2\ell^2\ast C_{\mathrm{TT}}=(\ell^2/2)\overline{\psi}\gamma_5\gamma\psi.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0375 (6.2.26)	128	■ 0.466 ● W +2	$\stackrel{\scriptscriptstyle\pm}{L}_{\parallel}=\mp\frac{i}{4}\ell^2\stackrel{\scriptscriptstyle\pm}{\Pi}^{\alpha}\wedge\stackrel{\scriptscriptstyle\pm}{\Pi}_{\alpha},\quad \overline{\stackrel{\scriptscriptstyle\pm}{L}_{\parallel}}=\stackrel{\scriptscriptstyle-}{L}_{\parallel},$	$\stackrel{\scriptscriptstyle\pm}{L}_{\parallel}=\mp\frac{i}{4}\ell^2\stackrel{\scriptscriptstyle\pm}{\Pi}^{\alpha}\wedge\stackrel{\scriptscriptstyle\pm}{\Pi}_{\alpha},\quad \overline{\stackrel{\scriptscriptstyle\pm}{L}_{\parallel}}=\stackrel{\scriptscriptstyle-}{L}_{\parallel}$	$\stackrel{\scriptscriptstyle\pm}{L}_{\parallel}=\mp\frac{i}{4}\ell^2\stackrel{\scriptscriptstyle\pm}{\Pi}^{\alpha}\wedge\stackrel{\scriptscriptstyle\pm}{\Pi}_{\alpha},\quad \overline{\stackrel{\scriptscriptstyle\pm}{L}_{\parallel}}=\stackrel{\scriptscriptstyle-}{L}_{\parallel},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0222 (4.2.15)	087	■ 0.466 ● N +0	$\widetilde{\varphi}:A(M)\rightarrow C^{\infty}(\overset{\circ}{V})$	$\widetilde{\varphi}:A(M)\rightarrow C^{\infty}(\overset{\circ}{V})$	$\widetilde{\varphi}:A(M)\rightarrow C^{\infty}(\overset{\circ}{V})$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0604 (9.2.3)	199	■ 0.474 ● N +1	\begin{aligned} L_{\text{Euler}}, &= \frac{1}{(4\pi)^2} \text{Tr} \left(\Omega^g \wedge \Omega^{g(*)} \right) \\ &= \frac{(-1)^{s/4}}{32\pi^2} R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta}^* \sqrt{ g } d^4x. \end{aligned}	$L_{\text{Euler}}, = \pi^*(\gamma_2^g) = \frac{1}{(4\pi)^2} \text{Tr} \left(\Omega^g \wedge \Omega^{g(*)} \right)$ $= \frac{(-1)^{s/4}}{32\pi^2} R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta}^* \sqrt{ g } d^4x.$	$L_{\text{Euler}}, = \pi^*(\gamma_2^g) = \frac{1}{(4\pi)^2} \text{Tr}(\Omega^g \wedge \Omega^{g(*)})$ $= \frac{(-1)^{s/4}}{32\pi^2} R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta}^* \sqrt{ g } d^4x.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0401 (6.4.20)	133	■ 0.482 ● W +1	\mathscr{H}_{\parallel} = \stackrel{\scriptscriptstyle (\pm)}{\underline{D}} - \stackrel{\scriptscriptstyle (\pm)}{\mathscr{A}}_{\hat{0}} = 0,	$\mathscr{H}_{\parallel} = \stackrel{\scriptscriptstyle (\pm)}{\underline{D}} - \stackrel{\scriptscriptstyle (\pm)}{\mathscr{A}}_{\hat{0}} = 0,$	$\mathscr{H}_{\parallel} = \stackrel{\scriptscriptstyle (\pm)}{\underline{D}} - \stackrel{\scriptscriptstyle (\pm)}{\mathscr{A}}_{\hat{0}} = 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0750 (11.1.17)	241	■ 0.484 ● C +23	D_{\left.\left.\left.\right.\right.\overparen{S_0(1,3)}}^{\mathrm{F}} \mathrm{F} = \underbrace{D^{\frac{1}{2}} \times \cdots \times D^{\frac{1}{2}}}_{(n-4)/4} \times \cdots	(not rendered)	$D_{[\overline{SO}_0(1,3)}^{\mathrm{F}} = \underbrace{D^{\frac{1}{2}} \times \cdots \times D^{\frac{1}{2}}}_{(n-4)/4}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0686 (10.3.44)	222	■ 0.497 ● N +0	\left.\left.\left.C_l = E_l^\alpha * \left\{ D^{\{ \right\}} \left[2 \left(e^\beta \lrcorner D^{\{ \right\}} H_{[\alpha\beta]} \right) + \frac{1}{2} \vartheta_\alpha \wedge \left(e^\beta \lrcorner e^\gamma \lrcorner D^{\{ \right\}} H_{[\beta\gamma]} \right) \right] \right\} \right.	$C_l = E_l^\alpha * \left\{ D^{\{ \right\}} \left[2 \left(e^\beta \lrcorner D^{\{ \right\}} H_{[\alpha\beta]} \right) + \frac{1}{2} \vartheta_\alpha \wedge \left(e^\beta \lrcorner e^\gamma \lrcorner D^{\{ \right\}} H_{[\beta\gamma]} \right) \right] \right\}$	$C_l = E_l^\alpha * \left\{ D^{\{ \right\}} \left[2 \left(e^\beta \lrcorner D^{\{ \right\}} H_{[\alpha\beta]} \right) + \frac{1}{2} \vartheta_\alpha \wedge \left(e^\beta \lrcorner e^\gamma \lrcorner D^{\{ \right\}} H_{[\beta\gamma]} \right) \right] \right\}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0515 (7.7.33)	166	■ 0.497 ● N +0	\left.\left.T := \frac{1}{4} \text{Tr}(\check{\gamma} \rfloor \Theta) = e_\alpha \rfloor T^\alpha, \quad \mathscr{A} := \frac{1}{4} {}^* \text{Tr}(\gamma \wedge \Theta) = {}^*(\vartheta_\alpha \wedge T^\alpha).\right.	$T := \frac{1}{4} \text{Tr}(\check{\gamma} \rfloor \Theta) = e_\alpha \rfloor T^\alpha, \quad \mathscr{A} := \frac{1}{4} {}^* \text{Tr}(\gamma \wedge \Theta) = {}^*(\vartheta_\alpha \wedge T^\alpha).$	$T := \frac{1}{4} \text{Tr}(\check{\gamma} \rfloor \Theta) = e_\alpha \rfloor T^\alpha, \quad \mathscr{A} := \frac{1}{4} {}^* \text{Tr}(\gamma \wedge \Theta) = {}^*(\vartheta_\alpha \wedge T^\alpha).$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value
Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0799 (11.4.11)	254	0.505 K +0	$ds^2=\frac{1}{\lambda^2}\left(R_{\circ}^2d\lambda^2-\mathrm{o}_{ij}dy^i\otimes_sdy^j\right).$	$ds^2=\frac{1}{\lambda^2}\left(R_{\circ}^2d\lambda^2-\mathrm{o}_{ij}dy^i\otimes_sdy^j\right).$	$ds^2=\frac{1}{\lambda^2}\left(R_{\circ}^2d\lambda^2-\mathrm{o}_{ij}dy^i\otimes_sdy^j\right).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0960 (14.1.42)	309	0.508 W -1	$\xi_{\mu\nu}\xi_{\rho\sigma}=-\frac{1}{2}\frac{\bar{R}_{\mu\nu\rho\sigma}}{\bar{R}_{\alpha\beta}R^{\{\j\}\alpha\beta}}-\frac{1}{2}\frac{\bar{R}_{\mu\nu\gamma\delta}\bar{R}_{\rho\sigma}^{\{\gamma\delta\}}}{\bar{R}_{\alpha\beta}^{\{\}\gamma\delta}R^{\{\j\}\alpha\beta}}.$	$\xi_{\mu\nu}\xi_{\rho\sigma}=-\frac{1}{2}\frac{\bar{R}_{\mu\nu\rho\sigma}}{\bar{R}_{\alpha\beta}R^{\{\j\}\alpha\beta}}-\frac{1}{2}\frac{\bar{R}_{\mu\nu\gamma\delta}\bar{R}_{\rho\sigma}^{\{\gamma\delta\}}}{\bar{R}_{\alpha\beta}^{\{\}\gamma\delta}R^{\{\j\}\alpha\beta}}.$	$\xi_{\mu\nu}\xi_{\rho\sigma}=-\frac{1}{2}\frac{\bar{R}_{\mu\nu\rho\sigma}}{\bar{R}_{\alpha\beta}R^{\{\j\}\alpha\beta}}-\frac{1}{2}\frac{\bar{R}_{\mu\nu\gamma\delta}\bar{R}_{\rho\sigma}^{\{\gamma\delta\}}}{\bar{R}_{\alpha\beta}^{\{\}\gamma\delta}R^{\{\j\}\alpha\beta}}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1131 (B.43)	373	0.509 W +2	$\langle\,,\rangle_M:\bigwedge^pT^*(M)\times C_p(M,\mathbb{R})\rightarrow\mathbb{R}$ $\langle\alpha^{(p)},c_p\rangle_M\rightarrow\int_{c_p}\alpha^{(p)}$	$\langle\,,\rangle_M:\begin{cases}\bigwedge^pT^*(M)\times C_p(M,\mathbb{R})\rightarrow\mathbb{R}\\\langle\alpha^{(p)},c_p\rangle_M\rightarrow\int_{c_p}\alpha^{(p)}\end{cases}$	$\langle\,,\rangle_M:\begin{cases}\bigwedge^pT^*(M)\times C_p(M,\mathbb{R})\rightarrow\mathbb{R}\\\langle\alpha^{(p)},c_p\rangle_M\rightarrow\int_{c_p}\alpha^{(p)}\end{cases}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0294 (4.5.5)	101	0.510 N +0	$\left(e_{\alpha}\lrcorner D\Psi\right)\wedge\frac{\partial L}{\partial D\Psi}-\left(e_{\alpha}\lrcorner\Psi\right)\wedge\frac{\partial L}{\partial\Psi}.$	$\Sigma_{\alpha}=e_{\alpha}\lrcorner L-\left(e_{\alpha}\lrcorner D\Psi\right)\wedge\frac{\partial L}{\partial D\Psi}-\left(e_{\alpha}\lrcorner\Psi\right)\wedge\frac{\partial L}{\partial\Psi}.$	$\Sigma_{\alpha}=e_{\alpha}\lrcorner L-\left(e_{\alpha}\lrcorner D\Psi\right)\wedge\frac{\partial L}{\partial D\Psi}-\left(e_{\alpha}\lrcorner\Psi\right)\wedge\frac{\partial L}{\partial\Psi}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0118 (3.3.19)	059	0.513 N -1	$L_{\mathrm{Pontr}}:=-\frac{1}{2}F\wedge F=\mathbf{E}\cdot\mathbf{B}\eta.$	$L_{\mathrm{Pontr}}:=-\frac{1}{2}F\wedge F=\mathbf{E}\cdot\mathbf{B}\eta.$	$L_{\mathrm{Pontr}}:=-\frac{1}{2}F\wedge F=\mathbf{E}\cdot\mathbf{B}\eta.$
Conf. MathPix confidence: 0.5-0.9 0.5-0.9 <0.5 no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E08533 (7.7.53)	169	■0.513 ●C +5	$\begin{aligned} G &:= \frac{1}{2} R^{\beta \gamma} \wedge \eta_{\alpha \beta \gamma} \gamma^{\alpha} \\ &= G^{(\beta)} + \frac{(-1)^{\text{sig}}}{12} \left(e_{\alpha} \rfloor \mathcal{A} \wedge {}^* \mathcal{A} - \frac{1}{3} \mathcal{A} \wedge e_{\alpha} \rfloor {}^* \mathcal{A} \right) \gamma^{\alpha} + \frac{(-1)^{\text{sig}}}{6} \gamma \wedge d \mathcal{A} \end{aligned}$	$G := \frac{1}{2} R^{\beta \gamma} \wedge \eta_{\alpha \beta \gamma} \gamma^{\alpha}$ $= G^{(\beta)} + \frac{(-1)^{\text{sig}}}{12} \left(e_{\alpha} \rfloor \mathcal{A} \wedge {}^* \mathcal{A} - \frac{1}{3} \mathcal{A} \wedge e_{\alpha} \rfloor {}^* \mathcal{A} \right) \gamma^{\alpha} + \frac{(-1)^{\text{sig}}}{6} \gamma \wedge d \mathcal{A}$	$G := \frac{1}{2} R^{\beta \gamma} \wedge \eta_{\alpha \beta \gamma} \gamma^{\alpha} \quad (7.7.5)$ $= G^{(\beta)} + \frac{(-1)^{\text{sig}}}{12} \left(e_{\alpha} \rfloor \mathcal{A} \wedge {}^* \mathcal{A} - \frac{1}{3} \mathcal{A} \wedge e_{\alpha} \rfloor {}^* \mathcal{A} \right) \gamma^{\alpha} + \frac{(-1)^{\text{sig}}}{6} \gamma \wedge d \mathcal{A}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E0898 (13.5.3)	288	■0.515 ●N +0	$\hat{B} := B_A{}^B A^A{}_B = \left(b_{\alpha} P_{*}^{\alpha} + \hat{b}^{\alpha} P_{\alpha} \right) \ell^2 + B_{\alpha}{}^{\beta} L^{\alpha}{}_{\beta}$	$\hat{B} := B_A{}^B A^A{}_B = \left(b_{\alpha} P_{*}^{\alpha} + \hat{b}^{\alpha} P_{\alpha} \right) \ell^2 + B_{\alpha}{}^{\beta} L^{\alpha}{}_{\beta}$	$\hat{B} := B_A{}^B A^A{}_B = (b_{\alpha} P_{*}^{\alpha} + \hat{b}^{\alpha} P_{\alpha}) \ell^2 + B_{\alpha}{}^{\beta} L^{\alpha}{}_{\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E0845 (2.6.8)	043	■0.516 ●N +0	$(d \, b)_{\text{hor}} = D b = \omega \otimes b$	$(db)_{\text{hor}} = Db = \omega \otimes b$	$(db)_{\text{hor}} = Db = \omega \otimes b$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E0876 (13.3.5)	285	■0.518 ●W -1	$\begin{aligned} &\left\{ \left[P_{\alpha}, P_{\beta} \right] = 0, \left[P_{*}^{\alpha}, P_{*}^{\beta} \right] = 0, \right. \\ &\left. \left[L_{\beta}^{\alpha}, P_{\gamma} \right] = \delta_{\gamma}^{\alpha} P_{\beta}, \left[L_{\beta}^{\alpha}, P_{*}^{\gamma} \right] = -\delta_{\beta}^{\gamma} P_{*}^{\alpha} \right\} \end{aligned}$	$\left[P_{\alpha}, P_{\beta} \right] = 0, \left[P_{*}^{\alpha}, P_{*}^{\beta} \right] = 0,$ $\left[L_{\beta}^{\alpha}, P_{\gamma} \right] = \delta_{\gamma}^{\alpha} P_{\beta}, \left[L_{\beta}^{\alpha}, P_{*}^{\gamma} \right] = -\delta_{\beta}^{\gamma} P_{*}^{\alpha}.$	$\left[P_{\alpha}, P_{\beta} \right] = 0, \quad \left[P_{*}^{\alpha}, P_{*}^{\beta} \right] = 0,$ $\left[L^{\alpha}{}_{\beta}, P_{\gamma} \right] = \delta_{\gamma}^{\alpha} P_{\beta}, \quad \left[L^{\alpha}{}_{\beta}, P_{*}^{\gamma} \right] = -\delta_{\beta}^{\gamma} P_{*}^{\alpha}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E0890 (14.3.3)	320	■0.526 ●W +0	$\begin{aligned} &\hat{L}_{\text{D}} := \frac{i}{2} \left(\overline{\hat{\psi}} \hat{\gamma} \wedge {}^* \hat{D}^{g+0} \hat{\psi} + \overline{\left(\hat{D}^{g+0} \hat{\psi} \right)} \wedge {}^* \hat{\gamma} \hat{\psi} \right) - \hat{m} \overline{\hat{\psi}} \hat{\psi} \eta \\ &+ \frac{1}{2} \overline{\hat{\psi}} \left[\hat{\alpha}, {}^* \hat{\gamma} \right] \hat{\psi}. \end{aligned}$	$\hat{L}_{\text{D}} := \frac{i}{2} \left(\overline{\hat{\psi}} \hat{\gamma} \wedge {}^* \hat{D}^{g+0} \hat{\psi} + \overline{\left(\hat{D}^{g+0} \hat{\psi} \right)} \wedge {}^* \hat{\gamma} \hat{\psi} \right) - \hat{m} \overline{\hat{\psi}} \hat{\psi} \eta$ $+ \frac{1}{2} \overline{\hat{\psi}} \left[\hat{\alpha}, {}^* \hat{\gamma} \right] \hat{\psi}.$	$\hat{L}_{\text{D}} := \frac{i}{2} \left(\overline{\hat{\psi}} \hat{\gamma} \wedge {}^* \hat{D}^{g+0} \hat{\psi} + \overline{\left(\hat{D}^{g+0} \hat{\psi} \right)} \wedge {}^* \hat{\gamma} \hat{\psi} \right) - \hat{m} \overline{\hat{\psi}} \hat{\psi} \eta$ $+ \frac{1}{2} \overline{\hat{\psi}} \left[\hat{\alpha}, {}^* \hat{\gamma} \right] \hat{\psi}.$

Conf. MathPix confidence:

■ ≥0.9

■ 0.5-0.9

■ <0.5

— no value

Residual render vs scan (inkdrill):

● C component

● W weak

● S stable

● N noise

● K clean

● not measured

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0456 (7.2.8)	152	0.527 W +0	$\bar{R}_{\{. \ . \ c \ d\}^{\{a \ b\}}=2 \ R_{\{.[c]^{\{[a] \ \delta_{[d]}^{\{b]\}} \}} \ ;$ $\quad R_{\{. \ b\}^{\{a\}}:=R_{\{. \ b\}^{\{a\}}-\frac{1}{4} \ \delta_{[b]^{\{a\}}}$ $R,$	$\bar{R}_{\cdot cd}{}^{ab}=2\mathcal{R}_{[c}{}^{[a}{}_{\delta d]}{}^{b]}, \quad \mathcal{R}^a{}_{\cdot b}:=R^a{}_{\cdot b}-\frac{1}{4}\delta_b^a R,$	$\bar{R}_{\cdot cd}{}^{ab}=2\mathcal{R}_{[c}{}^{[a}{}_{\delta d]}{}^{b]}, \quad \mathcal{R}^a{}_{\cdot b}:=R^a{}_{\cdot b}-\frac{1}{4}\delta_b^a R,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0360 (6.2.11)	125	0.531 N +0	$L_{\{ \}} \equiv L_{\{\mathrm{HE}\}}+\frac{1}{2} \ \kappa \ R_{\{\alpha \ \beta \} \ \wedge^{\{*\}} \left(\vartheta^{\alpha} \wedge \vartheta^{\beta} \right. \\ \left. \vartheta^{\alpha} \wedge \vartheta^{\beta} \right) + \frac{2}{\kappa} \ d C_{\mathrm{TT}}^{\{*\}},$	$L_{\parallel} \equiv L_{\mathrm{HE}} + \frac{1}{2\kappa} R_{\alpha\beta} \wedge^* (\vartheta^\alpha \wedge \vartheta^\beta) + \frac{2\ell^2}{\kappa} dC_{\mathrm{TT}}^*,$	$L_{\parallel} \equiv L_{\mathrm{HE}} + \frac{1}{2\kappa} R_{\alpha\beta} \wedge^* (\vartheta^\alpha \wedge \vartheta^\beta) + \frac{2\ell^2}{\kappa} dC_{\mathrm{TT}}^*,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0437 (6.8.8)	142	0.532 N +0	$E_{\{\alpha\}}:=\frac{\partial L_g}{\partial \vartheta^\alpha}=e_\alpha \lrcorner L_g + \left(e_\alpha \lrcorner T^\beta\right) \wedge H_\beta + \left(e_\alpha \lrcorner R^{\beta\gamma}\right) \wedge H_{\beta\gamma},$	$E_\alpha := \frac{\partial L_g}{\partial \vartheta^\alpha} = e_\alpha \lrcorner L_g + (e_\alpha \lrcorner T^\beta) \wedge H_\beta + (e_\alpha \lrcorner R^{\beta\gamma}) \wedge H_{\beta\gamma},$	$E_\alpha := \frac{\partial L_g}{\partial \vartheta^\alpha} = e_\alpha \lrcorner L_g + (e_\alpha \lrcorner T^\beta) \wedge H_\beta + (e_\alpha \lrcorner R^{\beta\gamma}) \wedge H_{\beta\gamma},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0198 (4.1.11)	082	0.533 W +0	$\begin{aligned} e_{\{\alpha\}}(m) &\rightarrowtail e_{\{\alpha\}}^{\{\prime\}}(m)=g_{\{\alpha\}}^{\{\}}\wedge^{\{\beta\}}e_{\{\beta\}}(m), \quad \& \ g= \\ &\left[g_{\{\alpha\}}^{\{\}}\wedge^{\{\beta\}}\right] \in G \ L(n, \ \mathbb{R}) \\ &\end{aligned}$	$e_\alpha(m) \rightarrowtail e_\alpha{}'(m) = g_\alpha{}^\beta e_\beta(m),$ $g = [g_\alpha{}^\beta] \in GL(n, \mathbb{R}).$	$e_\alpha(m) \rightarrowtail e_\alpha{}'(m) = g_\alpha{}^\beta e_\beta(m),$ $g = [g_\alpha{}^\beta] \in GL(n, \mathbb{R}).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0599 (9.1.17)	196	0.534 W +2	$\begin{aligned} \& f=-\frac{1}{r}\left(1+\frac{\kappa}{2}\ell^2\right) e^{\{-\nu / 2\}}, \ g=\frac{1}{r} \\ e^{\{\nu / 2\}} \quad \& h=-\frac{1}{(+)\ell^*} \frac{\sqrt{-\kappa}}{3}, \quad k=+\frac{1}{(-)2\ell^*} \sqrt{-3\kappa} \\ \& \end{aligned}$	$f=-\frac{1}{r}\left(1+\frac{\kappa}{2\ell^{*2}}r^2\right) e^{-\nu/2}, g=\frac{1}{r}e^{\nu/2}$ $h=-\frac{1}{(+)\ell^*}\frac{\sqrt{-\kappa}}{3}, \quad k=+\frac{1}{(-)2\ell^*}\sqrt{-3\kappa}$	$f=-\frac{1}{r}\left(1+\frac{\kappa}{2\ell^{*2}}r^2\right) e^{-\nu/2}, g=\frac{1}{r}e^{\nu/2}$ $h=-\frac{1}{(+)\ell^*}\frac{\sqrt{-\kappa}}{3}, \quad k=+\frac{1}{(-)2\ell^*}\sqrt{-3\kappa}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0752 (11.1.19)	241	0.535 K +0	$\Psi=\Psi \otimes \tilde{b} \in C^{\infty}\left(V^{\frac{1}{2}}\right) \text{ .}$	$\Psi = \Psi \otimes \tilde{b} \in C^{\infty}\left(V^{\frac{1}{2}}\right) \text{ .}$	$\Psi = \psi \otimes \tilde{b} \in C^{\infty}(V^{\frac{1}{2}}) \text{ .}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ1047 (15.2.4)	344	0.536 W +0	$\begin{aligned} \bar{E}_{\{i \ \alpha\}^{\{f \ \} j} \ \gamma^{\alpha}+i \ \}^{\{* \} E_{\{i \ \alpha\}^{\{f \ \} j} \ \gamma^{\alpha} \ \gamma^5\} \right\} \lambda_j^{(f)} \\ \oplus \frac{1}{2} \left\{ E_{i \alpha}^{(c) j} \gamma^{\alpha}+i \ ^* E_{i \alpha}^{(c) j} \gamma^{\alpha} \gamma^5\right\} \lambda_j^{(c)} \end{aligned}$	$\bar{E}_i=\frac{1}{2}\left\{E_{i \alpha}^{(f) j} \gamma^{\alpha}+i^* E_{i \alpha}^{(f) j} \gamma^{\alpha} \gamma^5\right\} \lambda_j^{(f)} \\ \oplus \frac{1}{2}\left\{E_{i \alpha}^{(c) j} \gamma^{\alpha}+i^* E_{i \alpha}^{(c) j} \gamma^{\alpha} \gamma^5\right\} \lambda_j^{(c)}$	$\bar{E}_i=\frac{1}{2}\left\{E_{i \alpha}^{(f) j} \gamma^{\alpha}+i^* E_{i \alpha}^{(f) j} \gamma^{\alpha} \gamma^5\right\} \lambda_j^{(f)} \\ \oplus \frac{1}{2}\left\{E_{i \alpha}^{(c) j} \gamma^{\alpha}+i^* E_{i \alpha}^{(c) j} \gamma^{\alpha} \gamma^5\right\} \lambda_j^{(c)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0982 (14.2.23)	317	0.537 N +0	$\int_G \pi \hat{L}_{\mathrm{E.}}=\frac{1}{\ell^{* 2}} L_{\mathrm{W.}}+L_{\mathrm{YM}}^{\prime}+\frac{1}{2 \ell^{* 2}} \vartheta \wedge \vartheta \wedge \vartheta \wedge \vartheta \int_G \stackrel{\circ}{R} d \mu(G) \text{ .}$	$\int_G \pi \hat{L}_{\mathrm{E.}}=\frac{1}{\ell^{* 2}} L_{\mathrm{W.}}+L_{\mathrm{YM}}^{\prime}+\frac{1}{2 \ell^{* 2}} \vartheta \wedge \vartheta \wedge \vartheta \wedge \vartheta \int_G \stackrel{\circ}{R} d \mu(G) \text{ .}$	$\int_G \pi \hat{L}_{\mathrm{E.}}=\frac{1}{\ell^{* 2}} L_{\mathrm{W.}}+L_{\mathrm{YM}}^{\prime}+\frac{1}{2 \ell^{* 2}} \vartheta \wedge \vartheta \wedge \vartheta \wedge \vartheta \int_G \stackrel{\circ}{R} d \mu(G) \text{ .}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ1134 (C.2)	375	0.539 K +0	$\exp : \mathfrak{g} \rightarrow G \ni g=\exp \left(\iota \alpha \xi^j I_j\right)$	$\exp : \mathfrak{g} \rightarrow G \ni g=\exp \left(\iota \xi^j I_j\right)$	$\exp : \mathfrak{g} \rightarrow G \ni g=\exp (\iota \xi^j I_j)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ1002 (14.3.15)	323	0.541 N +0	$(i \not{D})^2=(-1)^{3 n-1+s} \operatorname{Tr}\left\{\gamma \wedge \gamma \wedge^*(D D)-\gamma \wedge(D \gamma) \wedge^* D\right\}$	$(i \not{D})^2=(-1)^{3 n-1+s} \operatorname{Tr}\left\{\gamma \wedge \gamma \wedge^*(D D)-\gamma \wedge(D \gamma) \wedge^* D\right\}$	$(i \not{D})^2=(-1)^{3 n-1+s} \operatorname{Tr}\left\{\gamma \wedge \gamma \wedge^*(D D)-\gamma \wedge(D \gamma) \wedge^* D\right\}$
Conf. MathPix confidence: ≥0.9 0.5–0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ1085 (A.11)	365	0.541 K +0	$\begin{aligned} & \mathrm{C}(0,1)=\mathbb{C} \quad (\text{field of complex numbers}), \\ & \mathrm{C}(1,0)=\mathbb{R} \oplus \mathbb{R}, \end{aligned}$	$\begin{aligned} C(0,1) &= \mathbb{C} \quad (\text{field of complex numbers}), \\ C(1,0) &= \mathbb{R} \oplus \mathbb{R}, \end{aligned}$	$\begin{aligned} C(0,1) &= \mathbb{C} \quad (\text{field of complex numbers}), \\ C(1,0) &= \mathbb{R} \oplus \mathbb{R}, \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0004 (2.1.3)	033	0.543 N +0	$e_{\{\alpha\}}(m)=e_{\{,\alpha\}^{\{i\}}(m)\,\partial_{\{i\}}\,.$	$e_{\alpha}(m)=e^i_{,\alpha}(m)\partial_i.$	$e_{\alpha}(m)=e^i_{.\alpha}(m)\,\partial_i.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0558 (8.5.1)	181	0.544 N +0	$\Delta L_{\mathrm{FP}}=s\operatorname{Tr}\!\left\{\!\frac{\theta_{\mathrm{T}}^{*i}}{4\ell^2}\bar{\chi}\wedge^*\sigma-\bar{c}d^*\gamma\right\}$	$\Delta L_{\mathrm{FP}}=s\operatorname{Tr}\!\left\{\!\frac{\theta_{\mathrm{T}}^{*i}}{4\ell^2}\bar{\chi}\wedge^*\sigma-\bar{c}d^*\gamma\right\}$	$\Delta L_{\mathrm{FP}}=s\operatorname{Tr}\!\left\{\!\frac{\theta_{\mathrm{T}}^{*i}}{4\ell^2}\bar{\chi}\wedge^*\sigma-\bar{c}d^*\gamma\right\}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0304 (4.5.14)	102	0.545 N +1	$\begin{gathered} D\,\Sigma_{\{\alpha\}}\simeq 0\,\,\,\, D\!\left(\tau_{\{\alpha\beta\}}+x_{[\alpha}\wedge\Sigma_{\beta]}\right)\simeq 0\,\,\,\,.\,\,\,\,\end{gathered}$	$\begin{aligned} D\Sigma_{\alpha} &\simeq 0 \\ D\left(\tau_{\alpha\beta}+x_{[\alpha}\wedge\Sigma_{\beta]}\right) &\simeq 0. \end{aligned}$	$\begin{aligned} D\Sigma_{\alpha} &\simeq 0, \\ D(\tau_{\alpha\beta}+x_{[\alpha}\wedge\Sigma_{\beta]}) &\simeq 0. \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0270 (4.3.35)	095	0.546 N +0	$R:=R_{\mu}^{\mu}=\mathrm{R}_{\alpha\mu}^{\mu\alpha}=\ast\left(\Omega^g\wedge\vartheta\wedge\vartheta\right).$	$R:=R_{\mu}^{\mu}=R_{\alpha\mu}^{\cdot\mu\alpha}=\ast(\Omega^g\wedge\vartheta\wedge\vartheta).$	$R:=R_{\mu}^{\mu}=R_{\alpha\mu}^{\cdot\mu\alpha}=\ast(\Omega^g\wedge\vartheta\wedge\vartheta).$
Conf. MathPix confidence: ≥0.9 0.5–0.9 <0.5 — no value					
Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ1011 (14.4.6)	325	0.554 N +0	$r_{(-)}^{+}=\frac{\hbar}{M^{*2}c}\left(M_{(-)}^{+}\sqrt{M^2-J(J+1)\frac{M^{*4}}{M^2}-\alpha_eM^{*2}\mathrm{Tr}\bar{Q}^2}\right)$	$r_{(-)}^{+}=\frac{\hbar}{M^{*2}c}\left(M_{(-)}^{+}\sqrt{M^2-J(J+1)\frac{M^{*4}}{M^2}-\alpha_eM^{*2}\mathrm{Tr}\bar{Q}^2}\right)$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0385 (6.3.3)	130	0.561 W +2	$\stackrel{\scriptscriptstyle{(\pm)}}{\Gamma}_{\alpha}^{\scriptscriptstyle{(\pm)}}\beta\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\beta}+\Sigma_{\alpha}\cong d\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},\quad\stackrel{\scriptscriptstyle{(\mathrm{A})}}{S}_{\alpha}=\stackrel{\scriptscriptstyle{(\mp)}}{S}_{\alpha}$	$\stackrel{\scriptscriptstyle{(\pm)}}{S}_{\alpha}:=\stackrel{\scriptscriptstyle{(\pm)}}{\Gamma}_{\alpha}^{\scriptscriptstyle{(\pm)}}\beta\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\beta}+\Sigma_{\alpha}\cong d\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},\quad\stackrel{\scriptscriptstyle{(\mathrm{A})}}{S}_{\alpha}=\stackrel{\scriptscriptstyle{(\mp)}}{S}_{\alpha}$	$\stackrel{\scriptscriptstyle{(\pm)}}{S}_{\alpha}:=\stackrel{\scriptscriptstyle{(\pm)}}{\Gamma}_{\alpha}^{\scriptscriptstyle{(\pm)}}\beta\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\beta}+\Sigma_{\alpha}\cong d\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},\quad\stackrel{\scriptscriptstyle{(\pm)}}{S}_{\alpha}=\stackrel{\scriptscriptstyle{(\mp)}}{S}_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0596 (9.1.14)	195	0.564 N +0	$(\stackrel{\scriptscriptstyle{+}}{D}K)-K\wedge K=\stackrel{\scriptscriptstyle{+}}{\Omega}_{\mathrm{C}}\{ \}.$	$(\stackrel{\scriptscriptstyle{+}}{D}K)-K\wedge K=\stackrel{\scriptscriptstyle{+}}{\Omega}_{\mathrm{C}}\{ \}.$	$(\stackrel{\scriptscriptstyle{+}}{D}K)-K\wedge K=\stackrel{\scriptscriptstyle{+}}{\Omega}_{\mathrm{C}}\{ \}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0169 (3.6.4)	070	0.570 W +1	$D_{\pm}\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)=\mp\frac{i}{2}\bar{\psi}^*D_{\pm}\psi=\mp\frac{1}{2}\bar{\psi}^*\gamma\wedge^*(i^*\gamma\wedge D_{\pm}\psi).$	$D_{\pm}\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)=\mp\frac{i}{2}\bar{\psi}^*D_{\pm}\psi=\mp\frac{1}{2}\bar{\psi}^*\gamma\wedge^*(i^*\gamma\wedge D_{\pm}\psi).$	$D_{\pm}\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)=\mp\frac{i}{2}\bar{\psi}^*D_{\pm}\psi=\mp\frac{1}{2}\bar{\psi}^*\gamma\wedge^*(i^*\gamma\wedge D_{\pm}\psi).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0889 (13.4.8)	286	0.576 N +0	$\begin{aligned} & \Gamma^{\scriptscriptstyle{(T)}}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}}=\Lambda^{-1}(x)\left[\Gamma^{\scriptscriptstyle{(T)}}+D\tau(x)\right], \\ & \Gamma^{\scriptscriptstyle{(T)}}_{*}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}}_{*}=\left[\Gamma^{\scriptscriptstyle{(T)}}_{*}+D\tau_{*}(x)\right]\Lambda(x); \end{aligned}$	$\Gamma^{\scriptscriptstyle{(T)}}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}}=\Lambda^{-1}(x)\left[\Gamma^{\scriptscriptstyle{(T)}}+D\tau(x)\right],$ $\Gamma^{\scriptscriptstyle{(T)}}_{*}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}}_{*}=\left[\Gamma^{\scriptscriptstyle{(T)}}_{*}+D\tau_{*}(x)\right]\Lambda(x);$	$\Gamma^{\scriptscriptstyle{(T)}}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}}=\Lambda^{-1}(x)\left[\Gamma^{\scriptscriptstyle{(T)}}+D\tau(x)\right],$ $\Gamma^{\scriptscriptstyle{(T)}}_{*}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}}_{*}=\left[\Gamma^{\scriptscriptstyle{(T)}}_{*}+D\tau_{*}(x)\right]\Lambda(x);$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library_ E00116 (3.3.17)	059	0.585 N +0	$\mathbf{H}=\mathbf{j}\left[\begin{array}{l} \boldsymbol{\nabla}\times\mathbf{B}-\frac{\partial\mathbf{E}}{c^2\partial t}=\mu\mathbf{J} \\ \boldsymbol{\nabla}\cdot\mathbf{E}=\rho/\varepsilon. \end{array}\right].$	$dH=j\left\{\begin{array}{l} \boldsymbol{\nabla}\times\mathbf{B}-\frac{\partial\mathbf{E}}{c^2\partial t}=\mu\mathbf{J} \\ \boldsymbol{\nabla}\cdot\mathbf{E}=\rho/\varepsilon. \end{array}\right.$	$dH=j\left\{\begin{array}{l} \boldsymbol{\nabla}\times\mathbf{B}-\frac{\partial\mathbf{E}}{c^2\partial t}=\mu\mathbf{J} \\ \boldsymbol{\nabla}\cdot\mathbf{E}=\rho/\varepsilon. \end{array}\right.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library_ E00381 (6.2.32)	129	0.588 W +1	$\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\alpha:=D\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\alpha\pm\frac{i}{2}\ell^2\left(e_\alpha\downarrow\stackrel{\scriptscriptstyle{(\mp)}}{\Pi}\right)\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\beta,$	$\stackrel{\scriptscriptstyle{(\pm)}}{D}\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\alpha:=D\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\alpha\pm\frac{i}{2}\ell^2\left(e_\alpha\downarrow\stackrel{\scriptscriptstyle{(\mp)}}{\Pi}\right)\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\beta,$	$\stackrel{\scriptscriptstyle{(\pm)}}{D}\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\alpha:=D\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\alpha\pm\frac{i}{2}\ell^2\left(e_\alpha\downarrow\stackrel{\scriptscriptstyle{(\mp)}}{\Pi}^\beta\right)\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\beta,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library_ E00056 (2.7.6)	045	0.592 N +0	$\varphi=\varphi\otimes b=G^{-1}\varphi\otimes^Gb\in C^\infty(V^\rho).$	$\phi=\varphi\otimes b=G^{-1}\varphi\otimes^Gb\in C^\infty(V^\rho).$	$\phi=\varphi\otimes b=G^{-1}\varphi\otimes^Gb\in C^\infty(V^\rho).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library_ E00187 (3.7.17)	074	0.593 N +0	$\begin{aligned} \left(D^*\Omega\right)&=\vec{D}^*\vec{\Omega}+\left[\omega_0,\Omega_0\right] \\ &=\vec{D}^*\vec{\Omega}+\left[\omega_0,\Theta^\perp\right]=0. \end{aligned}$	$\begin{aligned} (D^*\Omega)&=\vec{D}^*\vec{\Omega}+\left[\omega_0,\Omega_0\right] \\ &=\vec{D}^*\vec{\Omega}+\left[\omega_0,\Theta^\perp\right]=0. \end{aligned}$	$\begin{aligned} (D^*\Omega)&=\vec{D}^*\vec{\Omega}+\left[\omega_0,\Omega_0\right] \\ &=\vec{D}^*\vec{\Omega}+\left[\omega_0,\Theta^\perp\right]=0. \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library_ E00581 (8.6.19)	186	0.593 K +0	$\widehat{\Sigma}_\alpha:=\Sigma_\alpha-D^{\{\}}\mu_\alpha,$	$\widehat{\Sigma}_\alpha:=\Sigma_\alpha-D^{\{\}}\mu_\alpha,$	$\widehat{\Sigma}_\alpha:=\Sigma_\alpha-D^{\{\}}\mu_\alpha,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0948 (14.1.29)	307	0.593 ● N +0	$\begin{aligned} D^{\{ * \}} \Omega &= \left(D^{\{ * \}} \xi - (-1)^{z} \xi \wedge D \delta \right) \cos \delta \\ &\quad - (-1)^z (D \xi + (-1)^{z * } \xi \wedge D \delta) \sin \delta = \tau_e \end{aligned}$	$D^* \Omega = (D^* \xi - (-1)^z \xi \wedge D \delta) \cos \delta - (-1)^z (D \xi + (-1)^{z * } \xi \wedge D \delta) \sin \delta = \tau_e$	$D^* \Omega = (D^* \xi - (-1)^z \xi \wedge D \delta) \cos \delta - (-1)^z (D \xi + (-1)^{z * } \xi \wedge D \delta) \sin \delta = \tau_e$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0357 (6.2.8)	125	0.600 ● N +0	$L_{\mathrm{HE}} = -\frac{1}{2\kappa} R_{\alpha\beta}^{\{ \}} \wedge^* (\vartheta^\alpha \wedge \vartheta^\beta)$	$L_{\mathrm{HE}} = -\frac{1}{2\kappa} R_{\alpha\beta}^{\{ \}} \wedge^* (\vartheta^\alpha \wedge \vartheta^\beta)$	$L_{\mathrm{HE}} = -\frac{1}{2\kappa} R_{\alpha\beta}^{\{ \}} \wedge^* (\vartheta^\alpha \wedge \vartheta^\beta)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0704 (10.6.3)	225	0.601 ● N +0	$\square F - m_{\mathrm{photon}}^2 F = 0,$	$\square F - m_{\mathrm{photon}}^2 F = 0,$	$\square F - m_{\mathrm{photon}}^2 F = 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0927 (14.1.8)	303	0.604 ● W -1	$\begin{aligned} \Sigma_{\mathrm{YM}} &= \frac{(-1)^s}{2\alpha_g} \mathrm{Tr} \{ \Omega \wedge^* (\vartheta \wedge^* \Omega) +^* \vartheta \wedge^* (\Omega \wedge^* \Omega) \} \\ &= \frac{(-1)^s}{4\alpha_g} \mathrm{Tr} \{ \Omega \wedge^* (\vartheta \wedge^* \Omega) -^* \Omega \wedge^* (\vartheta \wedge \Omega) \} . \end{aligned}$	$\Sigma_{\mathrm{YM}} = \frac{(-1)^s}{2\alpha_g} \mathrm{Tr} \{ \Omega \wedge^* (\vartheta \wedge^* \Omega) +^* \vartheta \wedge^* (\Omega \wedge^* \Omega) \} = \frac{(-1)^s}{4\alpha_g} \mathrm{Tr} \{ \Omega \wedge^* (\vartheta \wedge^* \Omega) -^* \Omega \wedge^* (\vartheta \wedge \Omega) \} .$	$\Sigma_{\mathrm{YM}} = \frac{(-1)^s}{2\alpha_g} \mathrm{Tr} \{ \Omega \wedge^* (\vartheta \wedge^* \Omega) +^* \vartheta \wedge^* (\Omega \wedge^* \Omega) \} = \frac{(-1)^s}{4\alpha_g} \mathrm{Tr} \{ \Omega \wedge^* (\vartheta \wedge^* \Omega) -^* \Omega \wedge^* (\vartheta \wedge \Omega) \} .$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0326 (5.3.3)	112	0.613 ● N +0	$\left. \widehat{D} G_{\alpha} \right _{\mathrm{R}^{\beta \gamma}} \equiv \frac{1}{2} (e_{\alpha} \rfloor R^{\beta \gamma}) \wedge \eta_{\beta \gamma \mu} \wedge T^{\mu}$	$\widehat{D} G_{\alpha} \equiv \frac{1}{2} (e_{\alpha} \rfloor R^{\beta \gamma}) \wedge \eta_{\beta \gamma \mu} \wedge T^{\mu}$	$\widehat{D} G_{\alpha} \equiv \frac{1}{2} (e_{\alpha} \rfloor R^{\beta \gamma}) \wedge \eta_{\beta \gamma \mu} \wedge T^{\mu}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0050 (2.6.13)	044	0.626 W +2	$\begin{aligned} D \, D \, D \, b = & \, D(\backslash\Omega \, \otimes \, b) = (d \, \backslash\Omega + [\backslash\Omega, \, \omega]) \, \otimes \, b \\ & = (d \, \backslash\Omega + \backslash\Omega \wedge \omega - \omega \wedge \backslash\Omega) \, \otimes \, b \\ & = (dd\omega - d\omega \wedge \omega + \omega \wedge d\omega \\ & \quad + d\omega \wedge \omega - \omega \wedge \omega \wedge \omega \\ & \quad + \omega \wedge \omega \wedge \omega - \omega \wedge d\omega) \, \otimes \, b \equiv 0. \end{aligned}$	$DDDb = D(\Omega \otimes b) = (d\Omega + [\Omega, \omega]) \otimes b$ $= (d\Omega + \Omega \wedge \omega - \omega \wedge \Omega) \otimes b$ $= (dd\omega - d\omega \wedge \omega + \omega \wedge d\omega$ $+ d\omega \wedge \omega - \omega \wedge \omega \wedge \omega$ $+ \omega \wedge \omega \wedge \omega - \omega \wedge d\omega) \otimes b \equiv 0.$	$DDDb = D(\Omega \otimes b) = (d\Omega + [\Omega, \omega]) \otimes b$ $= (d\Omega + \Omega \wedge \omega - \omega \wedge \Omega) \otimes b$ $= (dd\omega - d\omega \wedge \omega + \omega \wedge d\omega$ $+ d\omega \wedge \omega - \omega \wedge \omega \wedge \omega$ $+ \omega \wedge \omega \wedge \omega - \omega \wedge d\omega) \otimes b \equiv 0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0115 (3.3.16)	059	0.630 N +1	$d \, F \equiv 0 \quad \left\{ \begin{array}{l} \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} \equiv 0 \\ \nabla \cdot \mathbf{B} \equiv 0 \end{array} \right.$	$dF \equiv 0 \quad \left\{ \begin{array}{l} \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} \equiv 0 \\ \nabla \cdot \mathbf{B} \equiv 0 \end{array} \right.$	$dF \equiv 0 \quad \left\{ \begin{array}{l} \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} \equiv 0 \\ \nabla \cdot \mathbf{B} \equiv 0. \end{array} \right.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0171 (3.6.6)	070	0.630 N +0	$i^* \gamma \wedge D_{\pm} \psi = 0 \quad \text{and} \quad F^{\pm} = \frac{1}{2} \bar{\psi} \sigma_{\pm} \psi = m \mathcal{P}_{\pm},$	$i^* \gamma \wedge D_{\pm} \psi = 0 \quad \text{and} \quad F^{\pm} = \frac{1}{2} \bar{\psi} \sigma_{\pm} \psi = m \mathcal{P}_{\pm},$	$i^* \gamma \wedge D_{\pm} \psi = 0 \quad \text{and} \quad F^{\pm} = \frac{1}{2} \bar{\psi} \sigma_{\pm} \psi = m \mathcal{P}_{\pm},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0189 (3.7.19)	074	0.631 N +0	$\{ \}^* \backslash \Omega = (\stackrel{+}{-}) \backslash \Omega \text{ \textit{ \{ . \} } }$	$^* \Omega = (\overset{+}{-}) \Omega.$	$^* \Omega = (\overset{+}{-}) \Omega.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0788 (11.3.7)	249	0.635 W +0	$\left\{ i \gamma^{\kappa} \nabla_{\kappa}^{()} - \frac{6}{n^2} \ell^{*2} \gamma^{n+1} \gamma^{\mu} \bar{\psi} \gamma^{n+1} \gamma_{\mu} \psi - m \right\} \psi = 0.$	$\left\{ i \gamma^{\kappa} \nabla_{\kappa}^{()} - \frac{6}{n^2} \ell^{*2} \gamma^{n+1} \gamma^{\mu} \bar{\psi} \gamma^{n+1} \gamma_{\mu} \psi - m \right\} \psi = 0.$	$\left\{ i \gamma^{\kappa} \nabla_{\kappa}^{()} - \frac{6}{n^2} \ell^{*2} \gamma^{n+1} \gamma^{\mu} \bar{\psi} \gamma^{n+1} \gamma_{\mu} \psi - m \right\} \psi = 0.$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0164 (3.5.12)	068	0.636 N +0	$L_{\mathrm{YM}}+L_{\mathrm{inf}}=(\stackrel{+}{-})\frac{1}{2}\alpha_g\operatorname{Tr}\left(\Omega\left(\stackrel{+}{-}\right)\Omega\right)^2\geq 0$	$L_{\mathrm{YM}}+L_{\infty}=\left(\stackrel{+}{-}\right)\frac{1}{2\alpha_g}\operatorname{Tr}\left({}^*\Omega\left(\stackrel{+}{-}\right)\Omega\right)^2\geq 0$	$L_{\mathrm{YM}}+L_{\infty}=\left(\stackrel{+}{-}\right)\frac{1}{2\alpha_g}\operatorname{Tr}({}^*\Omega(\stackrel{-}{+})\Omega)^2\geq 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0549 (8.4.3)	179	0.637 W +2	$\begin{aligned} L_{\mathrm{FP}}\& \;:=-s\operatorname{Tr}\left\{\bar{\chi}\wedge{}^*\Omega^g+\bar{\Phi}D^*\Psi+\frac{1}{2}\rho\bar{\chi}\wedge{}^*\beta+\bar{C}d^*\Gamma+\frac{1}{2}\bar{C}\wedge{}^*\mathbb{B}\right\} \\ &=\frac{1}{2}s\left[\bar{\chi}^{\alpha\beta}\wedge{}^*R_{\alpha\beta}^{(\pm)}+\bar{\Phi}_{\alpha\beta}D^*\Psi^{\alpha\beta}+\frac{1}{2}\rho\bar{\chi}^{\alpha\beta}\wedge{}^*b_{\alpha\beta}+\tilde{\Psi}_{\mathrm{F}}\right] \end{aligned}$	$L_{\mathrm{FP}}:=-s\operatorname{Tr}\left\{\bar{\chi}\wedge{}^*\Omega^g+\bar{\Phi}D^*\Psi+\frac{1}{2}\rho\bar{\chi}\wedge{}^*\beta+\bar{C}d^*\Gamma+\frac{1}{2}\bar{C}\wedge{}^*\mathbb{B}\right\} \\ =\frac{1}{2}s\left[\bar{\chi}^{\alpha\beta}\wedge{}^*R_{\alpha\beta}^{(\pm)}+\bar{\Phi}_{\alpha\beta}D^*\Psi^{\alpha\beta}+\frac{1}{2}\rho\bar{\chi}^{\alpha\beta}\wedge{}^*b_{\alpha\beta}+\tilde{\Psi}_{\mathrm{F}}\right]$	$L_{\mathrm{FP}}:=-s\operatorname{Tr}\left\{\bar{\chi}\wedge{}^*\bar{\Omega}^g+\bar{\Phi}D^*\Psi+\frac{1}{2}\rho\bar{\chi}\wedge{}^*\beta+\bar{C}d^*\Gamma+\frac{1}{2}\bar{C}\wedge{}^*\mathbb{B}\right\} \\ =\frac{1}{2}s\left[\bar{\chi}^{\alpha\beta}\wedge{}^*R_{\alpha\beta}^{(\pm)}+\bar{\Phi}_{\alpha\beta}D^*\Psi^{\alpha\beta}+\frac{1}{2}\rho\bar{\chi}^{\alpha\beta}\wedge{}^*b_{\alpha\beta}+\tilde{\Psi}_{\mathrm{F}}\right]. \quad (8)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1058 (15.2.15)	346	0.638 K +0	$\bar{D}^{\{\}}:=d+i\bar{\omega}^{\{\}}$	$\bar{D}^{\{\}}:=d+i\bar{\omega}^{\{\}}$	$\bar{D}^{\{\}}:=d+i\bar{\omega}^{\{\}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1008 (14.4.3)	325	0.640 N +1	$\begin{aligned} ds^2& \;=-\frac{\Delta}{\rho^2}\left[c\,dt-J_3\frac{\hbar}{M}\sin^2\vartheta\,d\phi\right]^2+\frac{\rho^2}{\Delta}dr+\rho^2d\vartheta^2 \\ & \;+\frac{\sin^2\vartheta}{\rho^2}\left\{\left[r^2+J(J+1)\frac{\hbar^2}{M^2c^2}\right]d\phi-J_3\frac{\hbar}{M}dt\right\}^2 \end{aligned}$	$ds^2=-\frac{\Delta}{\rho^2}\left[c\,dt-J_3\frac{\hbar}{M}\sin^2\vartheta\,d\phi\right]^2+\frac{\rho^2}{\Delta}dr+\rho^2d\vartheta^2 \\ +\frac{\sin^2\vartheta}{\rho^2}\left\{\left[r^2+J(J+1)\frac{\hbar^2}{M^2c^2}\right]d\phi-J_3\frac{\hbar}{M}dt\right\}^2$	$ds^2=-\frac{\Delta}{\rho^2}\left[c\,dt-J_3\frac{\hbar}{M}\sin^2\vartheta\,d\phi\right]^2+\frac{\rho^2}{\Delta}dr+\rho^2d\vartheta^2 \\ +\frac{\sin^2\vartheta}{\rho^2}\left\{\left[r^2+J(J+1)\frac{\hbar^2}{M^2c^2}\right]d\phi-J_3\frac{\hbar}{M}dt\right\}^2.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0255 (4.3.20)	093	0.642 N +0	$\stackrel{*}{\sigma}\Omega^g=\frac{1}{4}R_{\alpha\beta}^{..cd}L_{cd}\otimes\vartheta^\alpha\wedge\vartheta^\beta.$	$\stackrel{*}{\sigma}\Omega^g=\frac{1}{4}R_{\alpha\beta}^{..cd}L_{cd}\otimes\vartheta^\alpha\wedge\vartheta^\beta.$	$\stackrel{*}{\sigma}\Omega^g=\frac{1}{4}R_{\alpha\beta}^{..cd}L_{cd}\otimes\vartheta^\alpha\wedge\vartheta^\beta.$
Conf. MathPix confidence: ≥0.9 0.5–0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0316 (4.6.8)	105	■ 0.664 ● N +0	$\Phi(\widetilde{\Gamma},\gamma)=\left(2\pi\hbar\mathscr{M}_nG/\hbar\ell c^2\right)=2\pi n$	$\Phi(\widetilde{\Gamma},\gamma)=\left(2\pi\hbar\mathscr{M}_nG/\hbar\ell c^2\right)=2\pi n$	$\Phi(\widetilde{\Gamma},\gamma)=\left(2\pi\hbar\mathscr{M}_nG/\hbar\ell c^2\right)=2\pi n,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0335 (5.3.12)	113	■ 0.665 ● W +0	$\begin{aligned} \widetilde{T}_{\mu\nu} &= T_{\mu\nu} + \ell^{*2} \left\{ -4\tau_{\mu\cdot[\lambda]}^{\cdot\kappa}\tau_{\nu\cdot \kappa]}^{\cdot\lambda} - 2\tau_{\mu}^{\cdot\kappa\lambda}\tau_{\nu\kappa\lambda} + \tau_{\cdot,\mu}^{\kappa\lambda}\tau_{\kappa\lambda\nu} \right. \\ &\quad \left. + \frac{1}{2}g_{\mu\nu}\left(4\tau_{\delta\cdot[\lambda]}^{\cdot\kappa}\tau_{\cdot,\kappa]}^{\delta\lambda} + \tau^{\delta\kappa\lambda}\tau_{\delta\kappa\lambda}\right) \right\}. \end{aligned}$	$\widetilde{T}_{\mu\nu} = T_{\mu\nu} + \ell^{*2} \left\{ -4\tau_{\mu\cdot[\lambda]}^{\cdot\kappa}\tau_{\nu\cdot \kappa]}^{\cdot\lambda} - 2\tau_{\mu}^{\cdot\kappa\lambda}\tau_{\nu\kappa\lambda} + \tau_{\cdot,\mu}^{\kappa\lambda}\tau_{\kappa\lambda\nu} \right. \\ \left. + \frac{1}{2}g_{\mu\nu}\left(4\tau_{\delta\cdot[\lambda]}^{\cdot\kappa}\tau_{\cdot,\kappa]}^{\delta\lambda} + \tau^{\delta\kappa\lambda}\tau_{\delta\kappa\lambda}\right) \right\}.$	$\widetilde{T}_{\mu\nu} = T_{\mu\nu} + \ell^{*2} \left\{ -4\tau_{\mu\cdot[\lambda]}^{\cdot\kappa}\tau_{\nu\cdot \kappa]}^{\cdot\lambda} - 2\tau_{\mu}^{\cdot\kappa\lambda}\tau_{\nu\kappa\lambda} + \tau_{\cdot,\mu}^{\kappa\lambda}\tau_{\kappa\lambda\nu} \right. \\ \left. + \frac{1}{2}g_{\mu\nu}(4\tau_{\delta\cdot[\lambda]}^{\cdot\kappa}\tau_{\cdot,\kappa]}^{\delta\lambda} + \tau^{\delta\kappa\lambda}\tau_{\delta\kappa\lambda}) \right\}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1055 (15.2.12)	345	■ 0.665 ● N +0	$\bar{\mathrm{L}}_{\mathrm{w}}(f)=\frac{i}{8}\operatorname{Tr}\left(\bar{\gamma}\wedge\bar{\gamma}\wedge^{*}\bar{\Omega}\right).$	$\bar{L}_{\mathrm{w}}(f)=\frac{i}{8}\operatorname{Tr}\left(\bar{\gamma}\wedge\bar{\gamma}\wedge^{*}\bar{\Omega}\right).$	$\bar{L}_{\mathrm{w}}(f)=\frac{i}{8}\operatorname{Tr}\left(\bar{\gamma}\wedge\bar{\gamma}\wedge^{*}\bar{\Omega}\right).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1145 (C.13)	377	■ 0.667 ● N +0	$\stackrel{\circ}{g}_{lm}:=c_{lk}{}^ic_{mi}{}^k$	$\stackrel{\circ}{g}_{lm}:=c_{lk}{}^ic_{mi}{}^k$	$\stackrel{\circ}{g}_{lm}:=c_{lk}{}^ic_{mi}{}^k$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0450 (7.2.3)	151	■ 0.676 ● N +0	$\Omega^g=\stackrel{+}{\Omega}^g+\bar{\Omega}^g$	$\Omega^g=\stackrel{+}{\Omega}^g+\bar{\Omega}^g$	$\Omega^g=\stackrel{+}{\Omega}^g+\bar{\Omega}^g$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0949 (14.1.30)	307	0.688 N +0	$\begin{aligned} D \, \Omega &\equiv \left(D \, \xi + (-1)^z * \xi \wedge D \delta \right) \cos \delta \\ &+ (-1)^z \left(D^* \xi - (-1)^z \xi \wedge D \delta \right) \sin \delta = 0. \end{aligned}$	$D\Omega = (D\xi + (-1)^z * \xi \wedge D\delta) \cos \delta + (-1)^z (D^* \xi - (-1)^z \xi \wedge D\delta) \sin \delta = 0.$	$D\Omega = (D\xi + (-1)^z * \xi \wedge D\delta) \cos \delta + (-1)^z (D^* \xi - (-1)^z \xi \wedge D\delta) \sin \delta = 0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0696 (10.5.7)	224	0.694 N +0	$\left. D^* \gamma = * \gamma \wedge T, \quad T := e_\alpha \rfloor T^\alpha, \right. \\ \left. \gamma = \left\lfloor T^\alpha e_\alpha, \right. \right.$	$D^* \gamma = * \gamma \wedge T, \quad T := e_\alpha \rfloor T^\alpha,$	$D^* \gamma = * \gamma \wedge T, \quad T := e_\alpha \rfloor T^\alpha,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0534 (8.2.1)	175	0.699 C +5	$\begin{aligned} s \, \Gamma &\equiv \Psi - DC \\ s \, C &\equiv \Phi - \frac{1}{2} [C, C] \\ s \, \Omega^g &\equiv -D\Psi - [C, \Omega^g] \\ s \, \Psi &\equiv -D\Phi - [C, \Psi] \\ s \, \Phi &\equiv -[C, \Phi] \end{aligned}$	$\begin{aligned} s \Gamma &= \Psi - DC \\ s C &= \Phi - \frac{1}{2} [C, C] \\ s \Omega^g &= -D\Psi - [C, \Omega^g] \\ s \Psi &= -D\Phi - [C, \Psi] \\ s \Phi &= -[C, \Phi] \end{aligned}$	$\begin{aligned} s \Gamma &= \Psi - DC, \\ s C &= \Phi - \frac{1}{2} [C, C], \\ s \Omega^g &= -D\Psi - [C, \Omega^g], \\ s \Psi &= -D\Phi - [C, \Psi], \\ s \Phi &= -[C, \Phi]. \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0329 (5.3.6)	112	0.699 N +0	$D \, \Theta \equiv \left[\Omega^g, \gamma \right], \quad D \, \Omega^g \equiv 0,$	$D\Theta \equiv [\Omega^g, \gamma], \quad D\Omega^g \equiv 0,$	$D\Theta \equiv [\Omega^g, \gamma], \quad D\Omega^g \equiv 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0471 (7.3.10)	155	0.706 N +0	$\bar{\Omega}^{\{\}} = 0 \Longleftrightarrow \Omega_C^{\{\}} = 0 \text{ and } R^{\{\}} = 0.$	$\bar{\Omega}^{\{\}} = 0 \Longleftrightarrow \Omega_C^{\{\}} = 0 \text{ and } R^{\{\}} = 0.$	$\bar{\Omega}^{\{\}} = 0 \Longleftrightarrow \Omega_C^{\{\}} = 0 \text{ and } R^{\{\}} = 0.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0747 (11.1.14)	240	0.706 N +0	$M=\mathbb{R}\times S^3\#^vW^3$.	$M=\mathbb{R}\times S^3\#^vW^3.$	$M=\mathbb{R}\times S^3\#^vW^3.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0314 (4.6.6)	105	0.710 N +0	$\Phi(\stackrel{\sim}{\Gamma},\gamma)=P\exp\left[(i/\hbar)\oint\left(\Gamma^{(T)\alpha}P_\alpha+\Gamma_\alpha^{(L)\beta}L^\alpha_\beta\right)\right]$	$\Phi(\stackrel{\sim}{\Gamma},\gamma)=P\exp\left[(i/\hbar)\oint\left(\Gamma^{(T)\alpha}P_\alpha+\Gamma_\alpha^{(L)\beta}L^\alpha_\beta\right)\right]$	$\Phi(\stackrel{\sim}{\Gamma},\gamma)=P\exp\left[(i/\hbar)\oint\left(\Gamma^{(T)\alpha}P_\alpha+\Gamma_\alpha^{(L)\beta}L^\alpha_\beta\right)\right]$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0108 (3.3.9)	058	0.711 N -1	$L_{\mathrm{Pontr}}=-\frac{1}{2}F\wedge F=-\frac{1}{2}dC,$	$L_{\mathrm{Pontr}}=-\frac{1}{2}F\wedge F=-\frac{1}{2}dC,$	$L_{\mathrm{Pontr}}=-\frac{1}{2}F\wedge F=-\frac{1}{2}dC,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0545 (8.3.6)	177	0.711 N +0	$d\rightarrowtail\exp(\zeta\rfloor)d\exp(-\zeta\rfloor)$.	$d\rightarrow\exp(\zeta\rfloor)d\exp(-\zeta\rfloor).$	$d\rightarrowtail\exp(\zeta\rfloor)d\exp(-\zeta\rfloor).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0583 (9.1.1)	192	0.712 N +0	$d\,s^2=-e^vd\,t^2+e^{\lambda}d\,r^2+r^2d\,\Omega^{n-2}$	$ds^2=-e^vdt^2+e^\lambda dr^2+r^2d\Omega^{n-2}$	$ds^2=-e^vdt^2+e^\lambda dr^2+r^2d\Omega^{n-2}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0260 (4.3.25)	094	0.718 N +0	$\stackrel{*}{\mathrm{K}}=\frac{1}{2}\mathrm{K}_{\alpha}^{\beta}\vartheta^{\alpha}\vartheta^{\beta}$	$\sigma^*K=\frac{1}{2}K_{\alpha}^{\beta}L_{\beta}\otimes\vartheta^{\alpha}.$	$\sigma^*K=\frac{1}{2}K_{\alpha}^{\beta}L_{\beta}\otimes\vartheta^{\alpha}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0878 (13.3.7)	285	0.722 N +0	$\mathrm{SL}(n+1,\mathbb{R})\approx\mathrm{A}_*(n,\mathbb{R})\otimes\mathrm{GL}(n,\mathbb{R})\otimes\mathbb{R}_*^n$	$\mathrm{SL}(n+1,\mathbb{R})\approx\mathrm{A}_*(n,\mathbb{R}):=\mathbb{R}^n\otimes\mathrm{GL}(n,\mathbb{R})\otimes\mathbb{R}_*^n.$	$\mathrm{SL}(n+1,\mathbb{R})\approx\mathrm{A}_*(n,\mathbb{R}):=\mathbb{R}^n\otimes\mathrm{GL}(n,\mathbb{R})\otimes\mathbb{R}_*^n.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0041 (2.6.4)	042	0.725 N +0	$\Omega=\frac{1}{2}F_{\alpha\beta}L_j\otimes\vartheta^{\alpha}\wedge\vartheta^{\beta}$	$\sigma^*\Omega=\frac{1}{2}F_{\alpha\beta}L_j\otimes\vartheta^{\alpha}\wedge\vartheta^{\beta}$	$\sigma^*\Omega=\frac{1}{2}F_{\alpha\beta}L_j\otimes\vartheta^{\alpha}\wedge\vartheta^{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0574 (8.6.12)	185	0.725 W -2	$L_{\mathrm{eff}}:=L_{\mathrm{qPG}}-dC_{\theta}=-\frac{\theta_{\mathrm{T}}^*}{2\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}+\frac{\theta_{\mathrm{T}}^*\Lambda_{\mathrm{eff}}}{\ell^2}\eta.$	$L_{\mathrm{eff}}:=L_{\mathrm{qPG}}-dC_{\theta}=-\frac{\theta_{\mathrm{T}}^*}{2\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}+\frac{\theta_{\mathrm{T}}^*\Lambda_{\mathrm{eff}}}{\ell^2}\eta.$	$L_{\mathrm{eff}}:=L_{\mathrm{qPG}}-dC_{\theta}=-\frac{\theta_{\mathrm{T}}^*}{2\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}+\frac{\theta_{\mathrm{T}}^*\Lambda_{\mathrm{eff}}}{\ell^2}\eta.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0366 (6.2.13)	126	0.725 N -1	$L_{\mathrm{EC}}:=L_{\mathrm{EC}}\pm idC_{\mathrm{TT}}=-\frac{1}{2\ell^2}R^{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}+\frac{\Lambda}{\ell^2}\eta.$	$L_{\mathrm{EC}}:=L_{\mathrm{EC}}\pm idC_{\mathrm{TT}}=-\frac{1}{2\ell^2}R^{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}+\frac{\Lambda}{\ell^2}\eta.$	$L_{\mathrm{EC}}:=L_{\mathrm{EC}}\pm idC_{\mathrm{TT}}=-\frac{1}{2\ell^2}R^{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}+\frac{\Lambda}{\ell^2}\eta.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Ekehard_W._Mielke_ __Z-Library_ EQ0585 (9.1.3)	193	0.725 W +0	$d\,s^2=\frac{n-2}{2\,\Lambda_{\text{eff}}}\,t^2\left(-d\,t^2+d\,r^2\right)+\frac{(n-2)(n-3)}{2\Lambda_{\text{eff.}}}\,d\,\Omega^{n-2}$	$ds^2=\frac{n-2}{2\Lambda_{\text{eff}}\,t^2}\left(-dt^2+dr^2\right)+\frac{(n-2)(n-3)}{2\Lambda_{\text{eff.}}}d\Omega^{n-2}$	$ds^2=\frac{n-2}{2\Lambda_{\text{eff}}t^2}(-dt^2+dr^2)+\frac{(n-2)(n-3)}{2\Lambda_{\text{eff.}}}d\Omega^{n-2}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Ekehard_W._Mielke_ __Z-Library_ EQ0021 (2.4.3)	038	0.726 N -2	$\varphi^{\{p\}}:=\varphi^p\otimes b\in C^{\infty}(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^{\rho}),\quad b\in C^{\infty}(V^{\rho}).$	$\phi^{(p)}:=\varphi^p\otimes b\in C^{\infty}(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^{\rho}),\quad b\in C^{\infty}(V^{\rho}).$	$\phi^{(p)}:=\varphi^p\otimes b\in C^{\infty}(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^{\rho}),\quad b\in C^{\infty}(V^{\rho}).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Ekehard_W._Mielke_ __Z-Library_ EQ0209 (4.2.1)	085	0.728 N +0	$\stackrel{*}{\chi}\widetilde{\omega}=\omega^{\mathrm{L}}\otimes\omega^{\mathrm{T}},$	$\stackrel{*}{\chi}\widetilde{\omega}=\omega^{\mathrm{L}}\otimes\omega^{\mathrm{T}},$	$\stackrel{*}{\chi}\widetilde{\omega}=\omega^{\mathrm{L}}\otimes\omega^{\mathrm{T}},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Ekehard_W._Mielke_ __Z-Library_ EQ0309 (4.6.1)	104	0.728 N +0	$\widetilde{\Gamma}:=\Gamma^{(T)\alpha}P_{\alpha}+\Gamma_{\alpha}^{(L)\beta}L^{\alpha}{}_{\beta}$	$\widetilde{\Gamma}:=\Gamma^{(T)\alpha}P_{\alpha}+\Gamma_{\alpha}^{(L)\beta}L^{\alpha}{}_{\beta}$	$\widetilde{\Gamma}:=\Gamma^{(T)\alpha}P_{\alpha}+\Gamma_{\alpha}^{(L)\beta}L^{\alpha}{}_{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Ekehard_W._Mielke_ __Z-Library_ EQ0009 (2.2.4)	034	0.729 N +1	$E_{\dot{i}}^{\alpha}e^{\dot{i}}{}_{\dot{\beta}}=\delta^{\alpha}{}_{\dot{\beta}}.$	$E_i^{\alpha}e^i{}_{\beta}=\delta^{\alpha}{}_{\beta}.$	$E_i^{\alpha}e^i{}_{\beta}=\delta^{\alpha}{}_{\beta}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0384 (6.3.2)	130	■ 0.730 ● C +3	$H_{\alpha}^{\wedge\{\}}=\frac{1}{\ell^2}\left[\vartheta^{\wedge\{\}}\wedge\left(d\vartheta_{\beta}\wedge\vartheta_{\alpha}\right)-\frac{1}{2}\vartheta_{\alpha}\wedge\left(d\vartheta^{\beta}\wedge\vartheta_{\beta}\right)\right]=\frac{1}{2\ell^2}\Gamma^{\{\}\beta\gamma}\wedge\eta_{\alpha\beta\gamma}$	$H_{\alpha}^{\parallel}=\frac{1}{\ell^2}\left[\vartheta^{\beta}\wedge\left(d\vartheta_{\beta}\wedge\vartheta_{\alpha}\right)-\frac{1}{2}\vartheta_{\alpha}\wedge\left(d\vartheta^{\beta}\wedge\vartheta_{\beta}\right)\right]=\frac{1}{2\ell^2}\Gamma^{\{\}\beta\gamma}\wedge\eta_{\alpha\beta\gamma}$	$H_{\alpha}^{\parallel}=\frac{1}{\ell^2}\left[\vartheta^{\beta}\wedge\left(d\vartheta_{\beta}\wedge\vartheta_{\alpha}\right)-\frac{1}{2}\vartheta_{\alpha}\wedge\left(d\vartheta^{\beta}\wedge\vartheta_{\beta}\right)\right]=\frac{1}{2\ell^2}\Gamma^{\{\}\beta\gamma}\wedge\eta_{\alpha\beta\gamma}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0689 (10.3.47)	222	■ 0.731 ● N +0	$\tilde{\Lambda}=(-1)^s\{L(\star)+\frac{1}{3\ell}\left[4\gamma+(-1)^s\ell\right]\star\left(R^{\{\}\star\alpha}\wedge\vartheta_{\alpha}\right),$	$\tilde{\Lambda}=(-1)^sL(\star)+\frac{1}{3\ell}\left[4\gamma+(-1)^s\ell\right]\star\left(R^{\{\}\star\alpha}\wedge\vartheta_{\alpha}\right),$	$\tilde{\Lambda}=(-1)^sL(\star)+\frac{1}{3\ell}\left[4\gamma+(-1)^s\ell\right]\star\left(R^{\{\}\star\alpha}\wedge\vartheta_{\alpha}\right),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0662 (10.3.19)	217	■ 0.733 ● N -1	$\square:=(-1)^{pn+s}\left[\left\{\right\}^{\wedge\{\}}D\left\{\right\}^{\wedge\{\}}D+(-1)^nD^{\wedge\{\}}D^{\wedge\{\}}\right]$	$\square:=(-1)^{pn+s}\left[\left\{\right\}^{\wedge\{\}}D\left\{\right\}^{\wedge\{\}}D+(-1)^nD^{\wedge\{\}}D^{\wedge\{\}}\right]$	$\square:=(-1)^{pn+s}\left[\left\{\right\}^{\wedge\{\}}D\left\{\right\}^{\wedge\{\}}D+(-1)^nD^{\wedge\{\}}D^{\wedge\{\}}\right]$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ1128 (B.40)	373	■ 0.735 ● N +0	$\text{d}\vartheta\equiv 0,$	$\text{d}\vartheta\equiv 0,$	$\vartheta\vartheta\equiv 0,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0374 (6.2.25)	127	■ 0.742 ● W +0	$\begin{aligned}\stackrel{\wedge\{\}}{\Pi}L&= & -\frac{i}{4}\ell^2\left[\stackrel{\wedge\{\}}{\Pi}^{\alpha}\vartheta_{\alpha}-\frac{1}{2}\ell^2\left[\stackrel{\wedge\{\}}{\Pi}^{\alpha\beta}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\frac{a_0}{2\ell^2}\eta_{\alpha\beta}\right]\right]\wedge\left[\stackrel{\wedge\{\}}{\Pi}^{\alpha}\vartheta_{\alpha}\right] \\ & -\frac{i}{4}\left[\stackrel{\wedge\{\}}{\Pi}^{\alpha\beta}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\frac{a_0}{2\ell^2}\eta_{\alpha\beta}\right]\wedge\left[\stackrel{\wedge\{\}}{\Pi}^{\alpha}\vartheta_{\alpha}\right]\end{aligned}$	$\stackrel{\wedge\{\}}{L}=-\frac{i}{4}\ell^2\left[\stackrel{\wedge\{\}}{\Pi}^{\alpha}\vartheta_{\alpha}-\frac{1}{2}\ell^2\left[\stackrel{\wedge\{\}}{\Pi}^{\alpha\beta}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\frac{a_0}{2\ell^2}\eta_{\alpha\beta}\right]\right]\wedge\left[\stackrel{\wedge\{\}}{\Pi}^{\alpha}\vartheta_{\alpha}\right]$	$\stackrel{\wedge\{\}}{L}=-\frac{i}{4}\ell^2\left[\stackrel{\wedge\{\}}{\Pi}^{\alpha}\vartheta_{\alpha}-\frac{1}{2}\ell^2\left[\stackrel{\wedge\{\}}{\Pi}^{\alpha\beta}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\frac{a_0}{2\ell^2}\eta_{\alpha\beta}\right]\right]\wedge\left[\stackrel{\wedge\{\}}{\Pi}^{\alpha}\vartheta_{\alpha}\right]$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0524 (7.7.43)	167	0.743 C +6	$\Theta^{\pm} := \frac{1}{2} \left(\Theta \pm \Theta^* \right), \quad \Omega^{\pm} := \frac{1}{2} \left(\Omega \pm \Omega^* \right), \quad \Omega^{(\pm)} := \frac{1}{2} \left(\Omega \pm \Omega^{(*)} \right),$	$\Theta^{\pm} := \frac{1}{2} (\Theta \pm {}^*\Theta), \quad \Omega^{g\pm} := \frac{1}{2} (\Omega^g \pm {}^*\Omega^g), \quad \Omega^{(\pm)} := \frac{1}{2} (\Omega^g \pm \Omega^{g(*)}),$	$\Theta^{\pm} := \frac{1}{2} (\Theta \pm {}^*\Theta), \quad \Omega^{g\pm} := \frac{1}{2} (\Omega^g \pm {}^*\Omega^g), \quad \Omega^{(\pm)} := \frac{1}{2} (\Omega^g \pm \Omega^{g(*)}),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0157 (3.5.5)	067	0.746 K +0	$\{ \}^{\{ * \}} F_{\{ \alpha \beta \mu \nu \}} = (\stackrel{+}{-}) F_{\{ \alpha \beta \mu \nu \}}$	${}^*F_{\alpha\beta\mu\nu}^* = (-^+){F_{\alpha\beta\mu\nu}}$	${}^*F_{\alpha\beta\mu\nu}^* = (-^+){F_{\alpha\beta\mu\nu}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1099 (B.10)	369	0.746 C +3	$\begin{aligned} & \sigma_* \left(\alpha^{(p)} + \beta^{(p)} \right) = \sigma_* \left(\alpha^{(p)} \right) + \sigma_* \left(\beta^{(p)} \right) \\ & \sigma_* \left(\alpha^{(p)} \wedge \beta^{(q)} \right) = \sigma_* \left(\alpha^{(p)} \right) \wedge \sigma_* \left(\beta^{(q)} \right) \\ & d \left(\sigma_* \alpha^{(p)} \right) = \sigma_* \left(d\alpha^{(p)} \right) \end{aligned}$	$\begin{aligned} \sigma_* \left(\alpha^{(p)} + \beta^{(p)} \right) &= \sigma_* \left(\alpha^{(p)} \right) + \sigma_* \left(\beta^{(p)} \right) \\ \sigma_* \left(\alpha^{(p)} \wedge \beta^{(q)} \right) &= \sigma_* \left(\alpha^{(p)} \right) \wedge \sigma_* \left(\beta^{(q)} \right) \\ d \left(\sigma_* \alpha^{(p)} \right) &= \sigma_* \left(d\alpha^{(p)} \right) \end{aligned}$	$\begin{aligned} \sigma_*(\alpha^{(p)} + \beta^{(p)}) &= \sigma_*(\alpha^{(p)}) + \sigma_*(\beta^{(p)}), \\ \sigma_*(\alpha^{(p)} \wedge \beta^{(q)}) &= \sigma_*(\alpha^{(p)}) \wedge \sigma_*(\beta^{(q)}), \\ d(\sigma_*\alpha^{(p)}) &= \sigma_*(d\alpha^{(p)}). \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0655 (10.3.12)	216	0.746 N +0	$\left. \left. \left. E_{\alpha} = e_{\alpha} \right\rfloor L(\star) - \frac{2\gamma\varepsilon^2}{\ell^3(1+\delta\varepsilon)}\eta_{\alpha} + (-1)^s \frac{2\gamma}{\ell(1+\delta\varepsilon)} \left[(e_{\alpha} \right\rfloor R^{\star\beta}) \wedge \vartheta_{\beta} \right], \right. \left. \right.$	$E_{\alpha} = e_{\alpha} \rfloor L(\star) - \frac{2\gamma\varepsilon^2}{\ell^3(1+\delta\varepsilon)}\eta_{\alpha} + (-1)^s \frac{2\gamma}{\ell(1+\delta\varepsilon)} \left[(e_{\alpha} \rfloor R^{\star\beta}) \wedge \vartheta_{\beta} \right],$	$E_{\alpha} = e_{\alpha} \rfloor L(\star) - \frac{2\gamma\varepsilon^2}{\ell^3(1+\delta\varepsilon)}\eta_{\alpha} + (-1)^s \frac{2\gamma}{\ell(1+\delta\varepsilon)} \left[(e_{\alpha} \rfloor R^{\star\beta}) \wedge \vartheta_{\beta} \right],$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0477 (7.4.5)	156	0.748 N +0	$g_{ij} = \Omega^2 o_{ij}^{\mathrm{Mink}}$	$g_{ij} = \Omega^2 o_{ij}^{\mathrm{Mink}}$	$g_{ij} = \Omega^2 o_{ij}^{\mathrm{Mink}},$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0963 (14.2.3)	312	■0.750 ● N +1	$\stackrel{\circ}{\circ}\stackrel{\circ}{\circ} \stackrel{\circ}{\circ}\stackrel{\circ}{\circ} {\mathcal{U}}=\stackrel{\circ}{\circ}\stackrel{\circ}{\circ} {\vartheta} \wedge \stackrel{\circ}{\circ}\stackrel{\circ}{\circ} {\vartheta}$		
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0687 (10.3.45)	222	■0.753 ● N +0	$\left.\left.-E_{\alpha}=e_{\alpha}\right\rceil L(\star)-(-1)^s\frac{2}{\ell}\left(e_{\alpha}\right\rceil R^{\{\}\star\beta}\right)\wedge\vartheta_{\beta}=\ell\widetilde{\Sigma}_{\alpha}$		
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0606 (9.2.5)	199	■0.753 ● N +1	$\chi(M)=\int_M\{{\gamma}_2^g\}-\frac{1}{(4\pi)^2}\int_{\partial M}\Bigl\{\alpha\wedge\Omega^{g(\star)}+\frac{2}{3}\alpha\wedge(\alpha\wedge\alpha)^{(\star)}\Bigr\}.$		
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0992 (14.3.5)	320	■0.754 ● N +0	$\stackrel{\circ}{\circ}\stackrel{\circ}{\circ} {\gamma}^{K+1}\chi_{\rho F}^\pm(\xi)=\pm\chi_{\rho F}^\pm(\xi)$		
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0249 (4.3.14)	092	■0.755 ● N +0	$D_k g_{ij}=E_{\cdot k}^\alpha\nabla_\alpha g_{ij}=0$		

Conf. MathPix confidence:

</

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ8399 (6.4.17)	133	0.755 W +0	$\begin{aligned} & \left[D_{\alpha}, D_{\beta} \right] = -T_{\alpha\beta}{}^{\gamma}(x) D_{\gamma} + R_{\alpha\beta\gamma}{}^{\delta}(x) L^{\gamma}{}_{\delta}, \\ & \left[L^{\alpha}{}_{\beta}, D_{\gamma} \right] = \delta_{\gamma}^{\alpha} D_{\beta}, \\ & \left[L^{\alpha}{}_{\beta}, L^{\gamma}{}_{\delta} \right] = \delta_{\delta}^{\alpha} L^{\gamma}{}_{\beta} - \delta_{\beta}^{\gamma} L^{\alpha}{}_{\delta}, \end{aligned}$	$\begin{aligned} [D_{\alpha}, D_{\beta}] &= -T_{\alpha\beta}{}^{\gamma}(x) D_{\gamma} + R_{\alpha\beta\gamma}{}^{\delta}(x) L^{\gamma}{}_{\delta}, \\ [L^{\alpha}{}_{\beta}, D_{\gamma}] &= \delta_{\gamma}^{\alpha} D_{\beta}, \\ [L^{\alpha}{}_{\beta}, L^{\gamma}{}_{\delta}] &= \delta_{\delta}^{\alpha} L^{\gamma}{}_{\beta} - \delta_{\beta}^{\gamma} L^{\alpha}{}_{\delta}, \end{aligned}$	$\begin{aligned} [D_{\alpha}, D_{\beta}] &= -T_{\alpha\beta}{}^{\gamma}(x) D_{\gamma} + R_{\alpha\beta\gamma}{}^{\delta}(x) L^{\gamma}{}_{\delta}, \\ [L^{\alpha}{}_{\beta}, D_{\gamma}] &= \delta_{\gamma}^{\alpha} D_{\beta}, \\ [L^{\alpha}{}_{\beta}, L^{\gamma}{}_{\delta}] &= \delta_{\delta}^{\alpha} L^{\gamma}{}_{\beta} - \delta_{\beta}^{\gamma} L^{\alpha}{}_{\delta}, \end{aligned}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ8957 (14.1.39)	309	0.756 N +0	$\partial_{\kappa} \delta = \sqrt{ g } \varepsilon_{\kappa\lambda\rho\sigma} R^{\{\sigma}{}_{\nu} \nabla^{\rho} R^{\}\nu\lambda} / \left(R^{\{\sigma}{}_{\alpha\beta} R^{\}\alpha\beta} \right).$	$\partial_{\kappa} \delta = \sqrt{ g } \varepsilon_{\kappa\lambda\rho\sigma} R^{\{\sigma}{}_{\nu} \nabla^{\rho} R^{\}\nu\lambda} / \left(R^{\{\sigma}{}_{\alpha\beta} R^{\}\alpha\beta} \right).$	$\partial_{\kappa} \delta = \sqrt{ g } \varepsilon_{\kappa\lambda\rho\sigma} R^{\{\sigma}{}_{\nu} \nabla^{\rho} R^{\}\nu\lambda} / \left(R^{\{\sigma}{}_{\alpha\beta} R^{\}\alpha\beta} \right).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ8970 (14.2.11)	314	0.758 C +6	$\hat{\Omega} = \hat{d}\hat{\omega} - \hat{\omega} - \hat{\omega} \wedge \hat{\omega} = \left[\begin{array}{c c} \Omega^g + \alpha \wedge \alpha^T & -D\alpha \\ \hline 0 & \overset{\circ}{\Omega} + \alpha^T \wedge \alpha \end{array} \right].$	$\hat{\Omega} = \hat{d}\hat{\omega} - \hat{\omega} - \hat{\omega} \wedge \hat{\omega} = \left[\begin{array}{c c} \Omega^g + \alpha \wedge \alpha^T & -D\alpha \\ \hline 0 & \overset{\circ}{\Omega} + \alpha^T \wedge \alpha \end{array} \right].$	$\hat{\Omega} = \hat{d}\hat{\omega} - \hat{\omega} - \hat{\omega} \wedge \hat{\omega} = \left[\begin{array}{c c} \Omega^g + \alpha \wedge \alpha^T & -D\alpha \\ \hline 0 & \overset{\circ}{\Omega} + \alpha^T \wedge \alpha \end{array} \right].$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ1016 (14.4.11)	327	0.758 N +0	$d s^2 = - \left(1 - \left(\frac{r_{\circ}}{r} \right)^4 \right) \frac{\Delta}{r^2} dt^2 + \frac{r^2}{\Delta} dr^2 + r^2 d\Omega^2.$	$d s^2 = - \left(1 - \left(\frac{r_{\circ}}{r} \right)^4 \right) \frac{\Delta}{r^2} dt^2 + \frac{r^2}{\Delta} dr^2 + r^2 d\Omega^2.$	$d s^2 = - \left(1 - \left(\frac{r_{\circ}}{r} \right)^4 \right) \frac{\Delta}{r^2} dt^2 + \frac{r^2}{\Delta} dr^2 + r^2 d\Omega^2.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ8310 (4.6.2)	104	0.760 W +2	$\begin{aligned} & \xi^{\alpha} \wedge \xi^{\beta} = D^{\alpha} \xi^{\beta} - D^{\beta} \xi^{\alpha} \\ & \xi^{\alpha} \wedge \xi^{\beta} = D^{\alpha} \xi^{\beta} - D^{\beta} \xi^{\alpha} \\ & \xi^{\alpha} \wedge \xi^{\beta} = D^{\alpha} \xi^{\beta} - D^{\beta} \xi^{\alpha} \end{aligned}$	$\begin{aligned} \widetilde{D} \widetilde{\xi}^{\alpha} &= d\xi^{\alpha} + \Gamma_{\beta}^{(L)\alpha} \wedge \xi^{\beta} + \Gamma^{(T)\alpha} \\ &= D\xi^{\alpha} + \Gamma^{(T)\alpha} = 0 \end{aligned}$	$\begin{aligned} \widetilde{D} \widetilde{\xi}^{\alpha} &= d\xi^{\alpha} + \Gamma_{\beta}^{(L)\alpha} \wedge \xi^{\beta} + \Gamma^{(T)\alpha} \\ &= D\xi^{\alpha} + \Gamma^{(T)\alpha} = 0. \end{aligned}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__E08832 (12.1.4)	270	■ 0.763 ● N +0	$\left\langle d\widetilde{j}_5\right\rangle =0,$	$\left\langle d\widetilde{j}_5\right\rangle =0,$	$\langle d\widetilde{j}_5\rangle =0,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__E08231 (4.2.24)	088	■ 0.766 ● N +2	$T_{\{\alpha\ \beta\}}^{\{.\ .\ c\}}=2\ D_{\{[\alpha\} E_{\{\cdot\dot{\mid}\ \beta\}}\mid\ \beta\}}^{\{c\}}=2\left(\partial_{[\alpha}\partial_{\cdot\mid\beta]} E_{\{\cdot\dot{\mid}\ \beta\}}\mid\ \beta\}+\Gamma_{[\alpha\mid d}\partial_{\cdot\mid\beta]} E_{\{\cdot\dot{\mid}\ \beta\}}^d\right)\ .$	$T_{\alpha\beta}^{\cdot\cdot c}=2D_{[\alpha}E_{\cdot \beta]}^c=2\left(\partial_{[\alpha}E_{\cdot \beta]}^c+\Gamma_{[\alpha d}^{\cdot c}E_{\cdot \beta]}^d\right).$	$T_{\alpha\beta}^{\cdot\cdot c}=2D_{[\alpha}E_{\cdot \beta]}^c=2\left(\partial_{[\alpha}E_{\cdot \beta]}^c+\Gamma_{[\alpha d}^{\cdot c}E_{\cdot \beta]}^d\right).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__E01879 (A.4)	365	■ 0.766 ● N +0	$\left[\sigma^i,\sigma^j\right]=\varepsilon^ij\sigma^k$	$\left[\sigma^i,\sigma^j\right]=\varepsilon^{ij}_k\sigma^k$	$[\sigma^i,\sigma^j]=\varepsilon^{ij}_k\sigma^k$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__E08276 (4.4.4)	097	■ 0.767 ● W +0	$\begin{aligned} &\&\ \delta L_g=\delta\vartheta\wedge E-\delta\vartheta\wedge D\Pi^T+d\left(\delta\vartheta\wedge\Pi^T\right)\\ &E-\delta\vartheta\wedge D\Pi^T+d\left(\delta\vartheta\wedge\Pi^T\right)\\ &\&\ -\left[\delta\omega^g,\vartheta\right]\wedge\left[\Pi_\varnothing^T,\vartheta\right]-\delta\omega^g\wedge D\Pi^L+d\left(\delta\omega^g\wedge\Pi^L\right), \end{aligned}$	$\delta L_g=\delta\vartheta\wedge E-\delta\vartheta\wedge D\Pi^T+d\left(\delta\vartheta\wedge\Pi^T\right)\\-\left[\delta\omega^g,\vartheta\right]\wedge\left[\Pi_\varnothing^T,\vartheta\right]-\delta\omega^g\wedge D\Pi^L+d\left(\delta\omega^g\wedge\Pi^L\right),$	$\delta L_g=\delta\vartheta\wedge E-\delta\vartheta\wedge D\Pi^T+d(\delta\vartheta\wedge\Pi^T)\\-\left[\delta\omega^g,\vartheta\right]\wedge\left[\Pi_{\left[\right]}^T,\vartheta\right]-\delta\omega^g\wedge D\Pi^L+d(\delta\omega^g\wedge\Pi^L),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__E08613 (10.1.1)	209	■ 0.769 ● N +0	$\vartheta^{\alpha}=E_i^{\alpha}dx^i\quad\text{and}\quad\Gamma_{\alpha}^{\star}:=\frac{1}{2}\eta_{\alpha\beta\gamma}\Gamma^{\beta\gamma}.$	$\vartheta^{\alpha}=E_i^{\alpha}dx^i\quad\text{and}\quad\Gamma_{\alpha}^{\star}:=\frac{1}{2}\eta_{\alpha\beta\gamma}\Gamma^{\beta\gamma}.$	$\vartheta^{\alpha}=E_i^{\alpha}dx^i\quad\text{and}\quad\Gamma_{\alpha}^{\star}:=\frac{1}{2}\eta_{\alpha\beta\gamma}\Gamma^{\beta\gamma}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1066 (16.2.5)	355	0.770 C -3	$\rho^*=\int_0^{\rho}\frac{dx}{1\pm x^2}=\left\{\begin{array}{l}\operatorname{artan}\rho,\kappa>0\\\operatorname{Arth}\rho,\kappa<0\end{array}\right\}.$	$\rho^*=\int_o^{\rho}\frac{dx}{1\pm x^2}=\left\{\begin{array}{l}\operatorname{artan}\rho,\kappa>0\\\operatorname{Arth}\rho,\kappa<0\end{array}\right\}.$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1030 (15.1.6)	337	0.771 W -1	$\begin{aligned}&\frac{1}{2}\vartheta^g\wedge^*\Omega^g+\Lambda^*\vartheta_g=\ell^{*2}\left\{\Sigma_{\mathrm{m}}(\mathrm{leptons})+\Sigma^{\mathrm{g}}\right\},\\&\frac{1}{2}\vartheta^f\wedge^*\Omega^f=\ell^2\left\{\Sigma_{\mathrm{m}}(\mathrm{hadrons})+\Sigma^{\mathrm{f}}\right\}.\end{aligned}$	$\frac{1}{2}\vartheta^g\wedge^*\Omega^g+\Lambda^*\vartheta_g=\ell^{*2}\left\{\Sigma_{\mathrm{m}}(\mathrm{leptons})+\Sigma^{\mathrm{g}}\right\},$ $\frac{1}{2}\vartheta^f\wedge^*\Omega^f=\ell^2\left\{\Sigma_{\mathrm{m}}(\mathrm{hadrons})+\Sigma^{\mathrm{f}}\right\}.$	$\frac{1}{2}\vartheta^g\wedge^*\Omega^g+\Lambda^*\vartheta_g=\ell^{*2}\left\{\Sigma_{\mathrm{m}}(\mathrm{leptons})+\Sigma^{\mathrm{g}}\right\},$ $\frac{1}{2}\vartheta^f\wedge^*\Omega^f=\ell^2\left\{\Sigma_{\mathrm{m}}(\mathrm{hadrons})+\Sigma^{\mathrm{f}}\right\}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0112 (3.3.13)	059	0.772 W -2	$\begin{aligned}j:=&\&\mathscr{J}\wedge dt+\hat{\rho}\\=&\&\left(J_xdy\wedge dz+J_ydz\wedge dx+J_zdx\wedge dy\right)\wedge dt\\&+\rho dx\wedge dy\wedge dz\end{aligned}$	$j:=-\mathscr{J}\wedge dt+\hat{\rho}$ $=-(J_xdy\wedge dz+J_ydz\wedge dx+J_zdx\wedge dy)\wedge dt$ $+\rho dx\wedge dy\wedge dz.$	$j:=-\mathscr{J}\wedge dt+\hat{\rho}$ $=-(J_xdy\wedge dz+J_ydz\wedge dx+J_zdx\wedge dy)\wedge dt$ $+\rho dx\wedge dy\wedge dz.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0130 (3.4.1)	062	0.780 N -1	$\mathscr{F}:=\frac{1}{\alpha_g}\operatorname{Tr}\left\{\left\{\right\}^*\left(\Omega\wedge\right)\right\}=\frac{1}{4\alpha_g}F_{\alpha\beta}{}^jF^{\alpha\beta}{}_j$	$\mathscr{F}:=\frac{1}{\alpha_g}\operatorname{Tr}\left\{\left\{\right\}^*(\Omega\wedge^*\Omega)\right\}=\frac{1}{4\alpha_g}F_{\alpha\beta}{}^jF^{\alpha\beta}{}_j$	$\mathscr{F}:=\frac{1}{\alpha_g}\operatorname{Tr}\left\{\left\{\right\}^*(\Omega\wedge^*\Omega)\right\}=\frac{1}{4\alpha_g}F_{\alpha\beta}{}^jF^{\alpha\beta}{}_j$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0470 (7.3.9)	155	0.782 N +0	$\left\{\right\}^*\Omega^{\mathrm{g}(\ast)}=\zeta\Omega^{\mathrm{g}},$	$^*\Omega^{g(\ast)}=\zeta\Omega^g,$	$^*\Omega^{g(\ast)}=\zeta\Omega^g,$

Conf. MathPix confidence:

≥0.9

0.5–0.9

<0.5

— no value

Residual render vs scan (inkdrill):

C component

W weak

S stable

N noise

K clean

not measured

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0046 (2.6.9)	043	0.783 N +0	$\begin{aligned} \Delta b &= \int_{\partial U} \omega \otimes b = \int_U d(\omega \otimes b)_{\mathrm{hor}} = \int_U D(\omega \otimes b) \\ &= \int_U \Omega \otimes b \simeq \Omega \otimes b \end{aligned}$	$\Delta b = \int_{\partial U} \omega \otimes b = \int_U d(\omega \otimes b)_{\mathrm{hor}} = \int_U D(\omega \otimes b) \\ = \int_U \Omega \otimes b \simeq \Omega \otimes b$	$\Delta b = \int_{\partial U} \omega \otimes b = \int_U d(\omega \otimes b)_{\mathrm{hor}} = \int_U D(\omega \otimes b) \\ = \int_U \Omega \otimes b \simeq \Omega \otimes b$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1049 (15.2.6)	344	0.785 N +1	$\{ \}^{*} \bar{\omega} = \bar{\omega}_{\{i} \} \bar{L}_{\{j} \} \otimes dx^i = \bar{\omega}_i dx^i$	${}^* \bar{\omega} = \bar{\omega}_i^j \bar{L}_j \otimes dx^i = \bar{\omega}_i dx^i$	${}^* \bar{\omega} = \bar{\omega}_i^j \bar{L}_j \otimes dx^i = \bar{\omega}_i dx^i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0133 (3.4.4)	063	0.785 N -2	$L_{\{\omega\}} = L(\mathcal{F}, \{ \}^{*} \mathcal{F} \backslash \eta)$	$L_{\omega} = L(\mathcal{F}, {}^* \mathcal{F}) \eta.$	$L_{\omega} = L(\mathcal{F}, {}^* \mathcal{F}) \eta.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0312 (4.6.4)	104	0.786 W +0	$\begin{aligned} \Delta \xi^{\alpha} &= - \oint_C y \rfloor (DD \xi^{\alpha}) = \oint_C y \rfloor R^{(T) \alpha} = \int_S R^{(T) \alpha} \\ &\simeq \frac{1}{2} (T_{ij}^{\alpha} - R_{ij \beta}^{\alpha} \xi^{\beta}) \int_S dy^i \wedge dy^j \end{aligned}$	$\Delta \xi^{\alpha} = - \oint_C y \rfloor (DD \xi^{\alpha}) = \oint_C y \rfloor R^{(T) \alpha} = \int_S R^{(T) \alpha} \\ \simeq \frac{1}{2} (T_{ij}^{\alpha} - R_{ij \beta}^{\alpha} \xi^{\beta}) \int_S dy^i \wedge dy^j$	$\Delta \xi^{\alpha} = - \oint_C y \rfloor (DD \xi^{\alpha}) = \oint_C y \rfloor R^{(T) \alpha} = \int_S R^{(T) \alpha} \\ \simeq \frac{1}{2} (T_{ij}^{\alpha} - R_{ij \beta}^{\alpha} \xi^{\beta}) \int_S dy^i \wedge dy^j .$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0224 (4.2.17)	087	0.787 N +0	$\widetilde{\varphi} \rightarrow A^{-1} \widetilde{\varphi} = G^{-1} (\widetilde{\varphi} - T).$	$\widetilde{\varphi} \rightarrow A^{-1} \widetilde{\varphi} = G^{-1} (\widetilde{\varphi} - T).$	$\widetilde{\varphi} \rightarrow A^{-1} \widetilde{\varphi} = G^{-1} (\widetilde{\varphi} - T).$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0941 (14.1.22)	306	0.788 N +0	$\tan 2\delta=(-1)^z\frac{\operatorname{Tr}^*(\Omega\wedge\Omega\wedge\operatorname{Tr}^*(\Omega\wedge\Omega))}{\operatorname{Tr}^*(\Omega\wedge\Omega)}.$	$\tan 2\delta=(-1)^z\frac{\operatorname{Tr}^*(\Omega\wedge\Omega)}{\operatorname{Tr}^*(\Omega\wedge\Omega)}.$	$\tan 2\delta=(-1)^z\frac{\operatorname{Tr}^*(\Omega\wedge\Omega)}{\operatorname{Tr}^*(\Omega\wedge\Omega)}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0760 (11.1.27)	242	0.791 N +0	$\left(\gamma^{n+1}\right)^2=\mp \mathbb{1}.$	$\left(\gamma^{n+1}\right)^2=\mp \mathbb{1}.$	$(\gamma^{n+1})^2=\mp \mathbb{1}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0725 (10.10.5)	230	0.793 N +0	$\delta_{\text{susy}}L\cong\bar{\sigma}\left(i\gamma^{\alpha}\Psi\wedge\frac{\delta L}{\delta\vartheta^{\alpha}}-2D\frac{\delta L}{\delta\Psi}+c\gamma\wedge\frac{\delta L}{\delta\bar{\Psi}}\right)+2d\left(\bar{\sigma}\wedge\frac{\delta L}{\delta\Psi}\right).$	$\delta_{\text{susy}}L\cong\bar{\sigma}\left(i\gamma^{\alpha}\Psi\wedge\frac{\delta L}{\delta\vartheta^{\alpha}}-2D\frac{\delta L}{\delta\Psi}+c\gamma\wedge\frac{\delta L}{\delta\bar{\Psi}}\right)+2d\left(\bar{\sigma}\wedge\frac{\delta L}{\delta\Psi}\right).$	$\delta_{\text{susy}}L\cong\bar{\sigma}\left(i\gamma^{\alpha}\Psi\wedge\frac{\delta L}{\delta\vartheta^{\alpha}}-2D\frac{\delta L}{\delta\Psi}+c\gamma\wedge\frac{\delta L}{\delta\bar{\Psi}}\right)+2d\left(\bar{\sigma}\wedge\frac{\delta L}{\delta\Psi}\right).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0810 (11.4.22)	256	0.793 N +0	$\frac{3}{8}\ell^2\bar{\psi}\psi=4\pi me^{-\mu-\nu/2}\left(H^2-F^2\right)\left Y_{ \kappa -1}^{ \kappa -1}(\vartheta,\phi)\right ^2$	$\frac{3}{8}\ell^2\bar{\psi}\psi=4\pi me^{-\mu-\nu/2}\left(H^2-F^2\right)\left Y_{ \kappa -1}^{ \kappa -1}(\vartheta,\phi)\right ^2$	$\frac{3}{8}\ell^2\bar{\psi}\psi=4\pi me^{-\mu-\nu/2}(H^2-F^2)\left Y_{ \kappa -1}^{ \kappa -1}(\vartheta,\phi)\right ^2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ1032 (15.1.9)	338	0.793 N +0	$\frac{1}{2}*(\vartheta^g\wedge^*\Omega^g)+\Lambda\vartheta^g=0,$	$\frac{1}{2}*(\vartheta^g\wedge^*\Omega^g)+\Lambda\vartheta^g=0,$	$\frac{1}{2}*(\vartheta^g\wedge^*\Omega^g)+\Lambda\vartheta^g=0,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__EQ0305 (4.5.15)	103	■0.793 ●C +12	\varepsilon_{\mathrm{RC}}:=\xi^{\alpha}\,\Sigma_\alpha\,\Sigma_{\alpha}+\left(e_\beta\downharpoonleft\overparen{D}\,\xi^\gamma\right)\tau^\beta{}_\gamma,	(not rendered)	$\varepsilon_{\text{RC}} := \xi^\alpha \Sigma_\alpha + (e_\beta \downharpoonleft \widehat{D} \xi^\gamma) \tau^\beta{}_\gamma,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__EQ0848 (12.3.1)	274	■0.794 ●C +5	\left\langle d j_5\right\rangle=2 i m\left\langle\bar{\psi} \gamma_5 \psi\right\rangle \eta-\frac{1}{4 \pi^2} \operatorname{Tr}(G \wedge G)-\frac{1}{96 \pi^2}\left[2 R_{\alpha \beta}^{(1)} \wedge R^{\{\beta \alpha \beta\}}+d \mathscr{A} \wedge d \mathscr{A}\right]	$\langle dj_5 \rangle = 2im \langle \bar{\psi} \gamma_5 \psi \rangle \eta - \frac{1}{4\pi^2} \operatorname{Tr}(G \wedge G) - \frac{1}{96\pi^2} \left[2R_{\alpha\beta}^{(1)} \wedge R^{\{\beta\alpha\beta\}} + d\mathcal{A} \wedge d\mathcal{A} \right]$	$\langle dj_5 \rangle = 2im \langle \bar{\psi} \gamma_5 \psi \rangle \eta - \frac{1}{4\pi^2} Tr(G \wedge G) - \frac{1}{96\pi^2} \left[2R_{\alpha\beta}^{(1)} \wedge R^{\{\alpha\beta\}} + d\mathcal{A} \wedge d\mathcal{A} \right]$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__EQ0319 (5.2.3)	110	■0.795 ●N +1	\begin{aligned} L_{\mathrm{D}} &=L\left(\gamma,\psi,D^{(1)}\psi\right)-\frac{i}{2} \bar{\psi}\left(*\gamma \wedge K-K \wedge *\gamma\right) \psi \\ &=L\left(\gamma,\psi,D^{(t)}\psi\right)+\frac{1}{4} \mathscr{A} \wedge j_5 . \end{aligned}	$L_D=L\left(\gamma,\psi,D^{(1)}\psi\right)-\frac{i}{2}\bar{\psi}\left(*\gamma\wedge K-K\wedge*\gamma\right)\psi\\=L\left(\gamma,\psi,D^{(t)}\psi\right)+\frac{1}{4}\mathcal{A}\wedge j_5.$	$L_D=L(\gamma,\psi,D^{(1)}\psi)-\frac{i}{2}\bar{\psi}\left(*\gamma\wedge K-K\wedge*\gamma\right)\psi\\=L(\gamma,\psi,D^{(1)}\psi)+\frac{1}{4}\mathcal{A}\wedge j_5.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__EQ0552 (8.4.6)	179	■0.797 ●N +0	\stackrel{+}{\Omega}:=\frac{1}{2}\left(\Omega^g\pm*\Omega^{g(*)}\right)=0	$\overset{+}{\Omega}:=\frac{1}{2}\left(\Omega^g\pm*\Omega^{g(*)}\right)=0$	$\overset{\pm}{\Omega}^g:=\frac{1}{2}\left(\Omega^g\pm*\Omega^{g(*)}\right)=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__EQ1001 (14.3.14)	323	■0.798 ●N -1	(i\not D)^2:-\operatorname{Tr}\left(\gamma\wedge\gamma\wedge D^*(\gamma\wedge D)\right).	$(i\,D)^2:=-\operatorname{Tr}\left\{\gamma\wedge*D^*(\gamma\wedge*D)\right\}.$	$(i\,\mathbb{D})^2:=-\operatorname{Tr}\left\{\gamma\wedge*D^*(\gamma\wedge*D)\right\}.$
Conf. MathPix confidence: ■≥0.9 ■0.5–0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0065 (2.8.8)	047	■ 0.803 ● N +0	$c_{\{2\}}(M)=-\int_M^{\pi}\pi\left(\gamma_2\right)=\frac{1}{8\pi^2}\int_M\operatorname{Tr}(\Omega\wedge\Omega)$	$c_2(M)=-\int_M^{\pi}\pi(\gamma_2)=\frac{1}{8\pi^2}\int_M\operatorname{Tr}(\Omega\wedge\Omega).$	$c_2(M)=-\int_M^{\pi}\pi(\gamma_2)=\frac{1}{8\pi^2}\int_M\operatorname{Tr}(\Omega\wedge\Omega).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0652 (10.3.9)	216	■ 0.803 ● K +0	$H_{\{\alpha\}}=-\frac{\gamma\varepsilon}{\ell^2(1+\delta\varepsilon)}\vartheta_{\alpha}\Leftrightarrow T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},$	$H_{\alpha}=-\frac{\gamma\varepsilon}{\ell^2(1+\delta\varepsilon)}\vartheta_{\alpha}\Leftrightarrow T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},$	$H_{\alpha}=-\frac{\gamma\varepsilon}{\ell^2(1+\delta\varepsilon)}\vartheta_{\alpha}\Leftrightarrow T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1065 (16.2.4)	355	■ 0.803 ● N +0	$\rho:=\sqrt{\frac{1}{3}\Lambda_{\text{eff}}}\,r,\quad r= \mathbf{x} $	$\rho:=\sqrt{\frac{1}{3}\Lambda_{\text{eff}}}\,r,\quad r= \mathbf{x} $	$\rho:=\sqrt{\frac{1}{3}\Lambda_{\text{eff}}}\,r,\quad r= \mathbf{x} $
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0166 (3.6.1)	069	■ 0.806 ● W +0	$\begin{aligned} L_{\mathrm{SW}}=&\frac{1}{2}\overline{D_{\pm}\psi}\wedge D_{\pm}\psi\mp i\left(F^{\pm}-\frac{1}{2}\overline{\psi}\sigma_{\pm}\psi\right)^2\\&=\mp i\operatorname{Tr}\left(F^{\pm}\wedge F^{\pm}\right)+\frac{1}{2}\overline{D_{\pm}\psi}\wedge D_{\pm}\psi\\&\quad\pm i\overline{\psi}\sigma_{\pm}\psi\wedge F^{\pm}\mp\frac{i}{4}\overline{\psi}\sigma_{\pm}\psi\wedge\overline{\psi}\sigma_{\pm}\psi; \end{aligned}$	$L_{\text{SW}}=\frac{1}{2}\overline{D_{\pm}\psi}\wedge D_{\pm}\psi\mp i\left(F^{\pm}-\frac{1}{2}\overline{\psi}\sigma_{\pm}\psi\right)^2$ $=\mp i\operatorname{Tr}\left(F^{\pm}\wedge F^{\pm}\right)+\frac{1}{2}\overline{D_{\pm}\psi}\wedge D_{\pm}\psi$ $\pm i\overline{\psi}\sigma_{\pm}\psi\wedge F^{\pm}\mp\frac{i}{4}\overline{\psi}\sigma_{\pm}\psi\wedge\overline{\psi}\sigma_{\pm}\psi;$	$L_{\text{SW}}=\frac{1}{2}\overline{D_{\pm}\psi}\wedge D_{\pm}\psi\mp i\left(F^{\pm}-\frac{1}{2}\overline{\psi}\sigma_{\pm}\psi\right)^2$ $=\mp i\operatorname{Tr}\left(F^{\pm}\wedge F^{\pm}\right)+\frac{1}{2}\overline{D_{\pm}\psi}\wedge D_{\pm}\psi$ $\pm i\overline{\psi}\sigma_{\pm}\psi\wedge F^{\pm}\mp\frac{i}{4}\overline{\psi}\sigma_{\pm}\psi\wedge\overline{\psi}\sigma_{\pm}\psi;$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1070 (16.2.9)	356	■ 0.806 ● N -1	$V_{\mathrm{eff}}^o(\rho^*)=\left(\frac{1}{\beta^2}-2\right)\left[1+\tan^2\rho^*\right]$	$V_{\text{eff}}^o(\rho^*)=\left(\frac{1}{\beta^2}-2\right)\left[1+\tan^2\rho^*\right]$	$V_{\text{eff}}^o(\rho^*)=\left(\frac{1}{\beta^2}-2\right)\left[1+\tan^2\rho^*\right]$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0446 (6.9.2)	144	■ 0.806 ● W +0	$\begin{aligned} \stackrel{\theta}{L} &= \frac{1}{2} \left[T^\alpha \wedge {}^* T_\alpha + \theta_{\mathrm{T}} \left(T^\alpha \wedge T_\alpha + R_{\alpha\beta} \wedge \vartheta^\alpha \wedge \vartheta^\beta \right) \right] \\ &+ \frac{1}{2} \left[R^{\alpha\beta} \wedge {}^* R_{\alpha\beta} + \theta_{\mathrm{L}} R^{\alpha\beta} \wedge R_{\alpha\beta} \right]. \end{aligned}$	$\stackrel{\theta}{L} = \frac{1}{2\ell} \left[T^\alpha \wedge {}^* T_\alpha + \theta_{\mathrm{T}} \left(T^\alpha \wedge T_\alpha + R_{\alpha\beta} \wedge \vartheta^\alpha \wedge \vartheta^\beta \right) \right] + \frac{1}{2} \left[R^{\alpha\beta} \wedge {}^* R_{\alpha\beta} + \theta_{\mathrm{L}} R^{\alpha\beta} \wedge R_{\alpha\beta} \right].$	$\stackrel{\theta}{L} = \frac{1}{2\ell} \left[T^\alpha \wedge {}^* T_\alpha + \theta_{\mathrm{T}} \left(T^\alpha \wedge T_\alpha + R_{\alpha\beta} \wedge \vartheta^\alpha \wedge \vartheta^\beta \right) \right] + \frac{1}{2} \left[R^{\alpha\beta} \wedge {}^* R_{\alpha\beta} + \theta_{\mathrm{L}} R^{\alpha\beta} \wedge R_{\alpha\beta} \right].$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0610 (9.3.1)	202	■ 0.806 ● N +0	$\begin{aligned} d\,C_{\mathrm{RR}^{(*)}} &:= \frac{1}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta(*)} - \frac{1}{3} \Gamma_\alpha{}^{\beta(*)} \wedge \Gamma_\beta{}^\gamma \wedge \Gamma_\gamma{}^\alpha \right) \\ &\equiv -L_{\mathrm{SKY}} - 2\,\mathrm{Ric}_{\alpha\beta} \wedge {}^*\mathrm{Ric}^{\alpha\beta} + \frac{1}{2} \mathrm{Ric}_\alpha{}^\alpha \wedge {}^*\mathrm{Ric}_\beta{}^\beta \end{aligned}$	$dC_{\mathrm{RR}^{(*)}} := \frac{1}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta(*)} - \frac{1}{3} \Gamma_\alpha{}^{\beta(*)} \wedge \Gamma_\beta{}^\gamma \wedge \Gamma_\gamma{}^\alpha \right) \equiv -L_{\mathrm{SKY}} - 2\,\mathrm{Ric}_{\alpha\beta} \wedge {}^*\mathrm{Ric}^{\alpha\beta} + \frac{1}{2} \mathrm{Ric}_\alpha{}^\alpha \wedge {}^*\mathrm{Ric}_\beta{}^\beta$	$dC_{\mathrm{RR}^{(*)}} := \frac{1}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta(*)} - \frac{1}{3} \Gamma_\alpha{}^{\beta(*)} \wedge \Gamma_\beta{}^\gamma \wedge \Gamma_\gamma{}^\alpha \right) \equiv -L_{\mathrm{SKY}} - 2\,\mathrm{Ric}_{\alpha\beta} \wedge {}^*\mathrm{Ric}^{\alpha\beta} + \frac{1}{2} \mathrm{Ric}_\alpha{}^\alpha \wedge {}^*\mathrm{Ric}_\beta{}^\beta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0767 (11.1.34)	244	■ 0.807 ● N +0	$\psi \longmapsto S^{-1} \psi = S^{-1} \psi$	$\psi \longrightarrow S^{-1} \psi = S^{-1} \psi$	$\psi \longrightarrow S^{-1} \psi = S^{-1} \psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0267 (4.3.32)	095	■ 0.808 ● W +2	$\begin{aligned} &R_{(ij)\kappa l} = R_{ij(\kappa l)} = 0 \\ &R_{ij\kappa l} = R_{\kappa lij}. \end{aligned}$	$R_{(ij)\kappa l} = R_{ij(\kappa l)} = 0$ $R_{ij\kappa l} = R_{\kappa lij}.$	$R_{(ij)\kappa l} = R_{ij(\kappa l)} = 0$ $R_{ij\kappa l} = R_{\kappa lij}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0298 (4.5.8)	101	■ 0.811 ● N +1	$\begin{aligned} \delta L &= \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha} \\ &+ \delta \Gamma_\alpha{}^\beta \wedge I^\alpha{}_\beta \Psi \wedge \frac{\partial L}{\partial D\Psi} + \delta \Psi \wedge \frac{\delta L}{\delta \Psi} + D \left(\delta \Psi \wedge \frac{\partial L}{\partial D\Psi} \right) \end{aligned}$	$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha} + \delta \Gamma_\alpha{}^\beta \wedge I^\alpha{}_\beta \Psi \wedge \frac{\partial L}{\partial D\Psi} + \delta \Psi \wedge \frac{\delta L}{\delta \Psi} + D \left(\delta \Psi \wedge \frac{\partial L}{\partial D\Psi} \right)$	$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha} + \delta \Gamma_\alpha{}^\beta \wedge I^\alpha{}_\beta \Psi \wedge \frac{\partial L}{\partial D\Psi} + \delta \Psi \wedge \frac{\delta L}{\delta \Psi} + D \left(\delta \Psi \wedge \frac{\partial L}{\partial D\Psi} \right).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0919 (13.8.5)	295	■ 0.811 ● N +0	$D \, T^{\{\alpha\}} \equiv R_{\{\beta\}}{}^{\{\alpha\}} \wedge \vartheta^\beta$	$DT^\alpha \equiv R_\beta{}^\alpha \wedge \vartheta^\beta$	$DT^\alpha \equiv R_\beta{}^\alpha \wedge \vartheta^\beta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0802 (11.4.14)	254	■ 0.812 ● W +0	$\begin{aligned} &\left[i \gamma^0 \partial_0 - e^{(\nu-\mu)/2} \left[i \vec{\gamma} \cdot \vec{\partial} + i \gamma^1 \partial_r \left(\frac{\mu}{2} + \frac{\nu}{4} \right) \right] \right. \\ &\left. + \frac{3\varepsilon}{8} \ell^2 e^{\nu/2} (\bar{\psi} \psi - \mathbf{a} \bar{\psi} \gamma^5 \psi \gamma^5) - e^{\nu/2} m \right] \psi = 0 \end{aligned}$	$\left\{ i \gamma^0 \partial_0 - e^{(\nu-\mu)/2} \left[i \vec{\gamma} \cdot \vec{\partial} + i \gamma^1 \partial_r \left(\frac{\mu}{2} + \frac{\nu}{4} \right) \right] + \frac{3\varepsilon}{8} \ell^2 e^{\nu/2} (\bar{\psi} \psi - \mathbf{a} \bar{\psi} \gamma^5 \psi \gamma^5) - e^{\nu/2} m \right\} \psi = 0$	$\left\{ i \gamma^0 \partial_0 - e^{(v-\mu)/2} \left[i \vec{\gamma} \cdot \vec{\partial} + i \gamma^1 \partial_r \left(\frac{\mu}{2} + \frac{\nu}{4} \right) \right] + \frac{3\varepsilon}{8} \ell^2 e^{\nu/2} (\bar{\psi} \psi - \mathbf{a} \bar{\psi} \gamma^5 \psi \gamma^5) - e^{\nu/2} m \right\} \psi = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0807 (11.4.19)	255	■ 0.812 ● N +0	$\vec{\sigma} \cdot \vec{L} \chi_\kappa^m = -(\kappa + 1) \chi_\kappa^m.$	$\vec{\sigma} \cdot \vec{L} \chi_\kappa^m = -(\kappa + 1) \chi_\kappa^m.$	$\vec{\sigma} \cdot \vec{L} \chi_\kappa^m = -(\kappa + 1) \chi_\kappa^m.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0397 (6.4.12)	132	■ 0.813 ● N +0	$\mathscr{H} \cong n^\alpha \mathscr{G}_\alpha + \Gamma_\perp{}^{\alpha\beta} \mathscr{G}_{\alpha\beta},$	$\mathscr{H} \cong n^\alpha \mathscr{G}_\alpha + \Gamma_\perp{}^{\alpha\beta} \mathscr{G}_{\alpha\beta},$	$\mathscr{H} \cong n^\alpha \mathscr{G}_\alpha + \Gamma_\perp{}^{\alpha\beta} \mathscr{G}_{\alpha\beta},$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0709 (10.7.5)	227	0.813 N +0	$R_{\{\beta\}^{\{\alpha\}}\wedge\vartheta^{\beta}}=\frac{\ell^2}{2}\varepsilon(x^{\alpha}Y\wedge X\wedge Y-y^{\alpha}X\wedge X\wedge Y)=0\;.$	$R_{\beta}^{\alpha}\wedge\vartheta^{\beta}=\frac{\ell^2}{2}\varepsilon(x^{\alpha}Y\wedge X\wedge Y-y^{\alpha}X\wedge X\wedge Y)=0\;.$	$R_{\beta}^{\alpha}\wedge\vartheta^{\beta}=\frac{\ell^2}{2}\varepsilon(x^{\alpha}Y\wedge X\wedge Y-y^{\alpha}X\wedge X\wedge Y)=0\;.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0210 (4.2.2)	085	0.814 N +1	$\stackrel{*}{\chi}\approx\left[\begin{array}{c c}\omega^{\mathrm{L}}&\omega^{\mathrm{T}}\\ \hline 0&0\end{array}\right]\;.$	$^*\tilde{\chi}=\left[\begin{array}{c c}\omega^{\mathrm{L}}&\omega^{\mathrm{T}}\\ \hline 0&0\end{array}\right]\;.$	$^*\tilde{\chi}\tilde{\omega}=\left[\begin{array}{c c}\omega^{\mathrm{L}}&\omega^{\mathrm{T}}\\ \hline 0&0\end{array}\right]\;.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0769 (11.1.36)	244	0.817 N +0	$\widetilde{\omega}\longrightarrow\widetilde{\omega}=S\widetilde{\omega}S^{-1}+(dS)S^{-1}$	$\widetilde{\omega}\longrightarrow{}^S\widetilde{\omega}=S\widetilde{\omega}S^{-1}+(dS)S^{-1}$	$\widetilde{\omega}\longrightarrow{}^S\widetilde{\omega}=S\widetilde{\omega}S^{-1}+(dS)S^{-1}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0398 (6.4.14)	133	0.820 W +0	$\begin{aligned} &\left(\mathscr{G}_{\alpha}(t,\mathbf{x}),\mathscr{G}_{\beta}(t,\mathbf{y})\right)=\left(-T_{\alpha\beta}{}^{\gamma}\mathscr{G}_{\gamma}+R_{\alpha\beta\gamma}{}^{\delta}\mathscr{G}_{\delta}\right)\cdot\delta(\mathbf{x}-\mathbf{y}),\\ &\left(\mathscr{G}^{\alpha}_{\beta}(t,\mathbf{x}),\mathscr{G}_{\gamma}(t,\mathbf{y})\right)=\delta^{\gamma}_{\beta}\mathscr{G}_{\beta}\delta(\mathbf{x}-\mathbf{y}),\\ &\left(\mathscr{G}^{\alpha}_{\beta}(t,\mathbf{x}),\mathscr{G}^{\gamma}_{\delta}(t,\mathbf{y})\right)=\left(\delta^{\alpha}_{\delta}\mathscr{G}^{\gamma}_{\beta}-\delta^{\gamma}_{\beta}\mathscr{G}^{\alpha}_{\delta}\right)\cdot\delta(\mathbf{x}-\mathbf{y}). \end{aligned}$	$\begin{aligned} &\left(\mathscr{G}_{\alpha}(t,\mathbf{x}),\mathscr{G}_{\beta}(t,\mathbf{y})\right)=\left(-T_{\alpha\beta}{}^{\gamma}\mathscr{G}_{\gamma}+R_{\alpha\beta\gamma}{}^{\delta}\mathscr{G}_{\delta}\right)\cdot\delta(\mathbf{x}-\mathbf{y}),\\ &\left(\mathscr{G}^{\alpha}_{\beta}(t,\mathbf{x}),\mathscr{G}_{\gamma}(t,\mathbf{y})\right)=\delta^{\gamma}_{\beta}\mathscr{G}_{\beta}\delta(\mathbf{x}-\mathbf{y}),\\ &\left(\mathscr{G}^{\alpha}_{\beta}(t,\mathbf{x}),\mathscr{G}^{\gamma}_{\delta}(t,\mathbf{y})\right)=\left(\delta^{\alpha}_{\delta}\mathscr{G}^{\gamma}_{\beta}-\delta^{\gamma}_{\beta}\mathscr{G}^{\alpha}_{\delta}\right)\cdot\delta(\mathbf{x}-\mathbf{y}). \end{aligned}$	$\begin{aligned} &\left(\mathscr{G}_{\alpha}(t,\mathbf{x}),\mathscr{G}_{\beta}(t,\mathbf{y})\right)=\left(-T_{\alpha\beta}{}^{\gamma}\mathscr{G}_{\gamma}+R_{\alpha\beta\gamma}{}^{\delta}\mathscr{G}_{\delta}\right)\cdot\delta(\mathbf{x}-\mathbf{y}),\\ &\left(\mathscr{G}^{\alpha}_{\beta}(t,\mathbf{x}),\mathscr{G}_{\gamma}(t,\mathbf{y})\right)=\delta^{\gamma}_{\beta}\mathscr{G}_{\beta}\delta(\mathbf{x}-\mathbf{y}),\\ &\left(\mathscr{G}^{\alpha}_{\beta}(t,\mathbf{x}),\mathscr{G}^{\gamma}_{\delta}(t,\mathbf{y})\right)=\left(\delta^{\alpha}_{\delta}\mathscr{G}^{\gamma}_{\beta}-\delta^{\gamma}_{\beta}\mathscr{G}^{\alpha}_{\delta}\right)\cdot\delta(\mathbf{x}-\mathbf{y}). \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0059 (2.7.9)	045	0.820 K +0	$\bar{\varphi}\rightarrow{}^{G^{-1}}\bar{\varphi}:=\bar{\varphi}G\quad\in C^{\infty}\left(V^{\bar{s}}\right).$	$\bar{\varphi}\rightarrow{}^{G^{-1}}\bar{\varphi}:=\bar{\varphi}G\quad\in C^{\infty}\left(V^{\bar{s}}\right).$	$\bar{\varphi}\rightarrow{}^{G^{-1}}\bar{\varphi}:=\bar{\varphi}G\quad\in C^{\infty}\left(V^{\bar{s}}\right).$
Conf. MathPix confidence: 0.9 0.5–0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0993 (14.3.6)	320	0.822 W +0	$\begin{aligned} \int_G \pi \hat{L}_D. &= \frac{i}{2} \left(\bar{\psi} \gamma \wedge^* D^{g+\omega} \psi + \overline{(D^{g+\omega} \psi)} \wedge^* \gamma \psi \right) \\ &+ \frac{\hat{\ell}}{2} \bar{\psi} \gamma \wedge \gamma \wedge^* \Omega \psi - \widehat{m} \bar{\psi} \psi \eta, \end{aligned}$	$\int_G \pi \hat{L}_D. = \frac{i}{2} \left(\bar{\psi} \gamma \wedge^* D^{g+\omega} \psi + \overline{(D^{g+\omega} \psi)} \wedge^* \gamma \psi \right) + \frac{\hat{\ell}}{2} \bar{\psi} \gamma \wedge \gamma \wedge^* \Omega \psi - \widehat{m} \bar{\psi} \psi \eta,$	$\int_G \pi \hat{L}_D. = \frac{i}{2} \left(\bar{\psi} \gamma \wedge^* D^{g+\omega} \psi + \overline{(D^{g+\omega} \psi)} \wedge^* \gamma \psi \right) + \frac{\hat{\ell}}{2} \bar{\psi} \gamma \wedge \gamma \wedge^* \Omega \psi - \widehat{m} \bar{\psi} \psi \eta,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0844 (12.2.12)	273	0.831 N +0	$\mathrm{L} = \frac{\mathrm{i}}{2\ell^2} \mathrm{Tr}(\Omega \wedge^* \sigma) + \mathrm{L}_D = \frac{1}{2\ell^2} \mathbf{R}^{\alpha\beta} \wedge \eta_{\alpha\beta} + \mathrm{L}_D,$	$\mathrm{L} = \frac{\mathrm{i}}{2\ell^2} \mathrm{Tr}(\Omega \wedge^* \sigma) + \mathrm{L}_D = \frac{1}{2\ell^2} \mathbf{R}^{\alpha\beta} \wedge \eta_{\alpha\beta} + \mathrm{L}_D,$	$\mathrm{L} = \frac{\mathrm{i}}{2\ell^2} \mathrm{Tr}(\Omega \wedge^* \sigma) + \mathrm{L}_D = \frac{1}{2\ell^2} \mathbf{R}^{\alpha\beta} \wedge \eta_{\alpha\beta} + \mathrm{L}_D,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0218 (4.2.10)	086	0.836 N +0	$\begin{aligned} &\omega^{\mathrm{L}} \rightarrow A^{-1} \omega^{\mathrm{L}} = G^{-1} \omega^{\mathrm{L}} G - G^{-1} dG, \\ &\omega^{\mathrm{T}} \rightarrow A^{-1} \omega^{\mathrm{T}} = G^{-1} \omega^{\mathrm{T}} - G^{-1} DT. \end{aligned}$	$\omega^{\mathrm{L}} \rightarrow A^{-1} \omega^{\mathrm{L}} = G^{-1} \omega^{\mathrm{L}} G - G^{-1} dG,$ $\omega^{\mathrm{T}} \rightarrow A^{-1} \omega^{\mathrm{T}} = G^{-1} \omega^{\mathrm{T}} - G^{-1} DT.$	$\omega^{\mathrm{L}} \rightarrow A^{-1} \omega^{\mathrm{L}} = G^{-1} \omega^{\mathrm{L}} G - G^{-1} dG,$ $\omega^{\mathrm{T}} \rightarrow A^{-1} \omega^{\mathrm{T}} = G^{-1} \omega^{\mathrm{T}} - G^{-1} DT.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0240 (4.3.5)	090	0.836 W -1	$\begin{aligned} &F(m, f\vartheta) = f F(m, \vartheta), \\ &F(m, \vartheta) > 0 \quad \text{if } \vartheta \neq 0. \end{aligned}$	$F(m, f\vartheta) = f F(m, \vartheta),$ $F(m, \vartheta) > 0 \quad \text{if } \vartheta \neq 0.$	$F(m, f\vartheta) = f F(m, \vartheta),$ $F(m, \vartheta) > 0 \quad \text{if } \vartheta \neq 0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0445 (6.9.1)	144	0.837 N +0	$ds^2 = - \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right)^{-1} dr^2 + r^2 d\Omega^2,$	$ds^2 = - \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right)^{-1} dr^2 + r^2 d\Omega^2,$	$ds^2 = - \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right)^{-1} dr^2 + r^2 d\Omega^2,$
Conf. MathPix confidence: 0.9 0.5–0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0227 (4.2.20)	088	■ 0.839 ● N +1	$\widetilde{\vartheta} \rightarrow^{A^{-1}} \widetilde{\vartheta} = G^{-1} \widetilde{\vartheta}.$		$\widetilde{\vartheta} \rightarrow^{A^{-1}} \widetilde{\vartheta} = G^{-1} \widetilde{\vartheta}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0773 (11.2.2)	245	■ 0.840 ● N +0	$i \gamma \wedge^* D \psi - \frac{i}{2} (D \wedge^* \gamma) \psi - m \psi \eta = 0$	$i \gamma \wedge^* D \psi - \frac{i}{2} (D \wedge^* \gamma) \psi - m \psi \eta = 0$	$i \gamma \wedge^* D \psi - \frac{i}{2} (D \wedge^* \gamma) \psi - m \psi \eta = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0811 (11.4.23)	256	■ 0.840 ● N +0	$\frac{3}{8} \ell^2 \bar{\psi} \gamma^5 \psi = 4 \pi m e^{-\mu - v/2} \frac{2}{3} H F \left Y_{ \kappa -1}^{ \kappa -1}(\vartheta, \phi) \right ^2$	$\frac{3}{8} \ell^2 \bar{\psi} \gamma^5 \psi = 4 \pi m e^{-\mu - v/2} \frac{2}{3} H F \left Y_{ \kappa -1}^{ \kappa -1}(\vartheta, \phi) \right ^2$	$\frac{3}{8} \ell^2 \bar{\psi} \gamma^5 \psi = 4 \pi m e^{-\mu - v/2} \frac{2}{3} H F \left Y_{ \kappa -1}^{ \kappa -1}(\vartheta, \phi) \right ^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0504 (9.1.2)	192	■ 0.843 ● N +0	$\begin{aligned} e^{\{\nu\}} &= e^{\{-\lambda\}} \wedge \frac{2}{n-1} \left(\frac{2}{r} \right)^{n-3} - \frac{2 \Lambda_{\text{eff.}}}{(n-1)(n-2)} r^2 \\ \mu &= \frac{M}{m^{*2}}; \quad d\Omega^2 := d\vartheta^2 + \sin^2 \vartheta d\phi^2 \end{aligned}$	$e^\nu = e^{-\lambda} = 1 - \left(\frac{2\mu}{r} \right)^{n-3} - \frac{2\Lambda_{\text{eff.}}}{(n-1)(n-2)} r^2$ $\mu = \frac{M}{m^{*2}}; \quad d\Omega^2 := d\vartheta^2 + \sin^2 \vartheta d\phi^2$	$e^\nu = e^{-\lambda} = 1 - \left(\frac{2\mu}{r} \right)^{n-3} - \frac{2\Lambda_{\text{eff.}}}{(n-1)(n-2)} r^2$ $\mu = \frac{M}{m^{*2}}; \quad d\Omega^2 := d\vartheta^2 + \sin^2 \vartheta d\phi^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0349 (5.4.7)	117	■ 0.845 ● N +0	$\left\langle \text{d j}_5 \right\rangle \text{rangle} = 2 \text{ i m} \left\langle \text{d j}_5 \right\rangle \text{rangle} \eta - \frac{1}{96 \pi^2} \left[2 R_{\alpha \beta}^{\{ \} } \wedge R^{\{ \beta \alpha \beta \} } + d \mathscr{A} \wedge d \mathscr{A} \right]$	$\langle dj_5 \rangle = 2 i m \langle \bar{\psi} \gamma_5 \psi \rangle \eta - \frac{1}{96 \pi^2} \left[2 R_{\alpha \beta}^{\{ \} } \wedge R^{\{ \beta \alpha \beta \} } + d \mathscr{A} \wedge d \mathscr{A} \right]$	$\langle dj_5 \rangle = 2 i m \langle \bar{\psi} \gamma_5 \psi \rangle \eta - \frac{1}{96 \pi^2} \left[2 R_{\alpha \beta}^{\{ \} } \wedge R^{\{ \beta \alpha \beta \} } + d \mathscr{A} \wedge d \mathscr{A} \right]$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0792 (11.4.4)	253	■0.845 ●N +0	$\begin{aligned} d\,s^{\wedge 2} &\,=\,g_{kl}\,d\,x^{\wedge k}\,\otimes_{\scriptscriptstyle S}\,d\,x^{\wedge l} \\ &\,=\,e^{\vartheta}\,dt^2\,-\,e^{\mu}\,\Bigl(dr^2\,+\,r^2d\vartheta^2\,+\,r^2\sin^2\vartheta d\phi^2\Bigr)\,.\end{aligned}$	$ds^2 = g_{kl}dx^k \otimes_s dx^l = e^{\vartheta}dt^2 - e^{\mu}\left(dr^2 + r^2d\vartheta^2 + r^2\sin^2\vartheta d\phi^2\right).$	$ds^2 = g_{kl}dx^k \otimes_s dx^l = e^{\vartheta}dt^2 - e^{\mu}\left(dr^2 + r^2d\vartheta^2 + r^2\sin^2\vartheta d\phi^2\right).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0742 (11.1.9)	238	■0.846 ●W +1	$\Lambda\!:\!\left\{\begin{array}{ll}\widetilde{SO}_{\circ}(s,n-s) &\longrightarrow SO_{\circ}(s,n-s),\\ \Psi &\longrightarrow g^{\alpha}{}_{\beta} = -\frac{i}{N}\,\mathrm{Tr}\left(\gamma^{\alpha}\,\mathrm{Si}\,\gamma_{\beta}S^{-1}\right),\end{array}\right.$	$\Lambda : \left\{ \begin{array}{ll} \widetilde{SO}_{\circ}(s,n-s) & \longrightarrow SO_{\circ}(s,n-s), \\ \Psi & \longrightarrow g^{\alpha}{}_{\beta} = -\frac{i}{N}\,\mathrm{Tr}\left(\gamma^{\alpha}\,\mathrm{Si}\,\gamma_{\beta}S^{-1}\right), \end{array} \right.$	$\Lambda : \left\{ \begin{array}{ll} \widetilde{SO}_{\circ}(s,n-s) & \longrightarrow SO_{\circ}(s,n-s), \\ \Psi & \longrightarrow g^{\alpha}{}_{\beta} = -\frac{i}{N}\mathrm{Tr}(\gamma^{\alpha}Si\gamma_{\beta}S^{-1}), \end{array} \right.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0897 (13.5.2)	288	■0.846 ●W +1	$\begin{aligned} \hat{R} &\, &:= R_{\scriptscriptstyle A}{}^{\scriptscriptstyle B}E^{\scriptscriptstyle A}{}_{\scriptscriptstyle B} &:= \left[d\Gamma_{\scriptscriptstyle A}{}^{\scriptscriptstyle B} - \Gamma_{\scriptscriptstyle A}{}^{\scriptscriptstyle C}\wedge\Gamma_{\scriptscriptstyle C}{}^{\scriptscriptstyle B}\right]E^{\scriptscriptstyle A}{}_{\scriptscriptstyle B} \\ &= \left[R_{\alpha}{}^{\beta} - \frac{\Lambda}{3}\theta_{\alpha}\wedge\vartheta^{\beta}\right]L^{\alpha}{}_{\beta} + T^{\beta}P_{\beta} + D\theta_{\alpha}P^{\alpha}_{\ast} \end{aligned}$	$\hat{R} := R_A{}^B E^A{}_B := \left[d\Gamma_A{}^B - \Gamma_A{}^C \wedge \Gamma_C{}^B\right] E^A{}_B = \left[R_{\alpha}{}^{\beta} - \frac{\Lambda}{3}\theta_{\alpha} \wedge \vartheta^{\beta}\right] L^{\alpha}{}_{\beta} + T^{\beta}P_{\beta} + D\theta_{\alpha}P^{\alpha}_{*}$	$\hat{R} := R_A{}^B \not{E}^A{}_B := \left[d\Gamma_A{}^B - \Gamma_A{}^C \wedge \Gamma_C{}^B\right] \not{E}^A{}_B = \left[R_{\alpha}{}^{\beta} - \frac{\Lambda}{3}\theta_{\alpha} \wedge \vartheta^{\beta}\right] L^{\alpha}{}_{\beta} + T^{\beta}P_{\beta} + D\theta_{\alpha}P^{\alpha}_{*},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0296 (4.5.7)	101	■0.847 ●N +0	$F_{\alpha}=\left(e_{\alpha}\lrcorner D\Psi\right)\frac{\delta L}{\delta\Psi}+(-1)^p\left(e_{\alpha}\lrcorner\Psi\right)\wedge D\frac{\delta L}{\delta\Psi}$	$F_{\alpha} = (e_{\alpha} \lrcorner D\Psi) \frac{\delta L}{\delta\Psi} + (-1)^p (e_{\alpha} \lrcorner \Psi) \wedge D\frac{\delta L}{\delta\Psi}$	$F_{\alpha} = (e_{\alpha} \lrcorner D\Psi) \frac{\delta L}{\delta\Psi} + (-1)^p (e_{\alpha} \lrcorner \Psi) \wedge D\frac{\delta L}{\delta\Psi}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0241 (4.3.6)	091	■0.848 ●N +0	$g_{\alpha\beta}(m,\vartheta):=\frac{1}{2}\frac{\partial^2F^2(m,\vartheta)}{\partial\vartheta^{\alpha}\otimes_s\partial\vartheta^{\beta}}.$	$g_{\alpha\beta}(m,\vartheta):=\frac{1}{2}\frac{\partial^2F^2(m,\vartheta)}{\partial\vartheta^{\alpha}\otimes_s\partial\vartheta^{\beta}}.$	$g_{\alpha\beta}(m,\vartheta):=\frac{1}{2}\frac{\partial^2F^2(m,\vartheta)}{\partial\vartheta^{\alpha}\otimes_s\partial\vartheta^{\beta}}.$
Conf. MathPix confidence: ■ ■ ■ — ≥0.90.5–0.9<0.5no value Residual render vs scan (inkdrill): ● ● ● ● ● ● C componentW weakS stableN noiseK cleannot measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0478 (7.4.6)	157	0.849 N -1	$\operatorname{Ric}_{\alpha\beta}:=\operatorname{Ric}_{\alpha\beta}-\frac{1}{4}\operatorname{Rg}_{\alpha\beta}=0.$	$\operatorname{Ric}_{\alpha\beta}:=\operatorname{Ric}_{\alpha\beta}-\frac{1}{4}\operatorname{Rg}_{\alpha\beta}=0.$	$R\check{\mathcal{L}}_{\alpha\beta}:=\operatorname{Ric}_{\alpha\beta}-\frac{1}{4}Rg_{\alpha\beta}=0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0295 (4.5.6)	101	0.850 K +0	$\begin{aligned} D\,\Sigma_{\alpha} & \& \left.\left.=\right.\right. \\ & \left.\left(e_{\alpha}\right\rfloor T^{\beta}\right) \wedge \Sigma_{\beta}+\left(e_{\alpha}\right\rfloor R^{\beta \gamma}\right) \wedge \tau_{\beta \gamma}+F_{\alpha} \\ & \& \left.\left(e_{\alpha}\right\rfloor T^{\beta}\right) \wedge \Sigma_{\beta}+\left(e_{\alpha}\right\rfloor R^{\beta \gamma}\right) \wedge \tau_{\beta \gamma} . \end{aligned}$	$D \Sigma_{\alpha}=\left(e_{\alpha}\right\rfloor T^{\beta}) \wedge \Sigma_{\beta}+\left(e_{\alpha}\right\rfloor R^{\beta \gamma}) \wedge \tau_{\beta \gamma}+F_{\alpha} \\ \simeq\left(e_{\alpha}\right\rfloor T^{\beta}) \wedge \Sigma_{\beta}+\left(e_{\alpha}\right\rfloor R^{\beta \gamma}) \wedge \tau_{\beta \gamma} .$	$D \Sigma_{\alpha}=\left(e_{\alpha}\right\rfloor T^{\beta}) \wedge \Sigma_{\beta}+\left(e_{\alpha}\right\rfloor R^{\beta \gamma}) \wedge \tau_{\beta \gamma}+F_{\alpha} \\ \simeq\left(e_{\alpha}\right\rfloor T^{\beta}) \wedge \Sigma_{\beta}+\left(e_{\alpha}\right\rfloor R^{\beta \gamma}) \wedge \tau_{\beta \gamma} .$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0842 (12.2.10)	272	0.852 N +0	$\mu:=e_{\alpha}\downharpoonleft\mu^{\alpha}=\frac{3}{4}j_5$	$\mu:=e_{\alpha}\downharpoonleft\mu^{\alpha}=\frac{3}{4}j_5$	$\mu:=e_{\alpha}\downharpoonleft\mu^{\alpha}=\frac{3}{4}j_5$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0506 (7.7.24)	164	0.852 K +0	$\gamma_5:=(i/4!)(\gamma\wedge\gamma\wedge\gamma\wedge\gamma)$	$\gamma_5:=(i/4!)(\gamma\wedge\gamma\wedge\gamma\wedge\gamma)$	$\gamma_5:=(i/4!)(\gamma\wedge\gamma\wedge\gamma\wedge\gamma)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ1014 (14.4.9)	326	0.856 N +0	$M^2=\left(M_{\mathrm{ir}}+\frac{\alpha_eM^{*2}}{4M_{\mathrm{ir}}}\mathrm{Tr}\,\bar{Q}^2\right)^2+\frac{1}{4}J(J+1)\frac{M^{*4}}{M_{\mathrm{ir}}^2},$	$M^2=\left(M_{\mathrm{ir}}+\frac{\alpha_eM^{*2}}{4M_{\mathrm{ir}}}\mathrm{Tr}\,\bar{Q}^2\right)^2+\frac{1}{4}J(J+1)\frac{M^{*4}}{M_{\mathrm{ir}}^2},$	$M^2=\left(M_{\mathrm{ir}}+\frac{\alpha_eM^{*2}}{4M_{\mathrm{ir}}}\mathrm{Tr}\,\bar{Q}^2\right)^2+\frac{1}{4}J(J+1)\frac{M^{*4}}{M_{\mathrm{ir}}^2},$
Conf. MathPix confidence: ■≥0.9 ■0.5–0.9 ■<0.5 —no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0753 (11.1.20)	241	0.859 C +17	$\hat{\Psi}_{\mid \overparen{S\,0}\circ(1,3)}=\left[\begin{array}{l}\Psi_1\\\Psi_2\\\vdots\\\Psi_{[\frac{n}{4}]}\end{array}\right].$	(not rendered)	$\hat{\Psi}_{\mid \overparen{S\,0}\circ(1,3)}=\left[\begin{array}{l}\Psi_1\\\Psi_2\\\vdots\\\Psi_{[\frac{n}{4}]}\end{array}\right].$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0783 (11.3.2)	247	0.862 N +0	$\begin{aligned} L_{\mathrm{D}} &= \frac{i}{2} \left(\bar{\psi} \gamma \wedge^* D^{\{3\}} \psi + \overline{(D^{\{3\}} \psi)} \wedge^* \gamma \psi \right) - m \bar{\psi} \psi \eta \\ &+ \frac{1}{2} \bar{\psi} \left[\tilde{K}, * \gamma \right] \psi =: L_{\mathrm{D}}^{\{3\}} + L_{\mathrm{NL}}. \end{aligned}$	$L_{\mathrm{D}} = \frac{i}{2} \left(\bar{\psi} \gamma \wedge^* D^{\{3\}} \psi + \overline{(D^{\{3\}} \psi)} \wedge^* \gamma \psi \right) - m \bar{\psi} \psi \eta \\ + \frac{1}{2} \bar{\psi} \left[\tilde{K}, * \gamma \right] \psi =: L_{\mathrm{D}}^{\{3\}} + L_{\mathrm{NL}}.$	$L_{\mathrm{D}} = \frac{i}{2} \left(\bar{\psi} \gamma \wedge^* D^{\{3\}} \psi + \overline{(D^{\{3\}} \psi)} \wedge^* \gamma \psi \right) - m \bar{\psi} \psi \eta \\ + \frac{1}{2} \bar{\psi} [\tilde{K}, * \gamma] \psi =: L_{\mathrm{D}}^{\{3\}} + L_{\mathrm{NL}}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1100 (B.12)	369	0.864 N +0	$\begin{aligned} \{ \}^{*} \alpha^{\{p\}} &:= \frac{1}{(n-p)! \cdot (p!)} \sqrt{ \det g_{ij} } \varepsilon_{\alpha_1 \cdots \alpha_p \beta_1 \cdots \beta_{n-p}} \\ &\times A^{\alpha_1 \cdots \alpha_p} \vartheta^{\beta_1} \wedge \cdots \wedge \vartheta^{\beta_{n-p}}. \end{aligned}$	${}^* \alpha^{(p)} := \frac{1}{(n-p)! \cdot (p!)} \sqrt{ \det g_{ij} } \varepsilon_{\alpha_1 \cdots \alpha_p \beta_1 \cdots \beta_{n-p}} \\ \times A^{\alpha_1 \cdots \alpha_p} \vartheta^{\beta_1} \wedge \cdots \wedge \vartheta^{\beta_{n-p}}.$	${}^* \alpha^{(p)} := \frac{1}{(n-p)! \cdot (p!)} \sqrt{ \det g_{ij} } \varepsilon_{\alpha_1 \cdots \alpha_p \beta_1 \cdots \beta_{n-p}} \\ \times A^{\alpha_1 \cdots \alpha_p} \vartheta^{\beta_1} \wedge \cdots \wedge \vartheta^{\beta_{n-p}}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0184 (3.7.14)	074	0.864 C +16	$\vec{\Omega} = d\vec{\omega} - (\overrightarrow{\omega} \wedge)$	$\vec{\Omega} = d\vec{\omega} - (\vec{\omega} \wedge)$	$\vec{\Omega} = d\vec{\omega} - (\overrightarrow{\omega \wedge \vec{\omega}}) = d\vec{\omega} - \vec{\omega} \wedge \vec{\omega}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0165 (3.5.13)	069	0.865 N +0	$\{ \}^{*} \Omega = (\stackrel{+}{-}) \Omega ;$	${}^* \Omega = (\overset{+}{-}) \Omega ;$	${}^* \Omega = (\overset{+}{-}) \Omega ;$
Conf. MathPix confidence: ≥0.9 0.5–0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0737 (11.1.4)	238	■0.867 ●N +0	x^{\alpha} \rightarrow x^{\prime\,\alpha}=g_{\beta}^{\alpha}\,x^\beta, \left[g_\beta{}^\alpha\right] \in S\,O(s,\,-n-s), \quad x^\alpha \in {\rm E}^n \; .	$x^\alpha \rightarrow x'^\alpha = g_\beta^\alpha \;\; x^\beta , \,[g_\beta^\alpha] \in SO(s,n-s) , \;\; x^\alpha \in {\bf E}^n .$	$x^\alpha \rightarrow x'^\alpha = g_\beta^\alpha \; x^\beta , [g_\beta^\alpha] \in SO(s,n-s) , \; x^\alpha \in {\bf E}^n .$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0853 (12.3.6)	276	■0.868 ●C +3	\left\langle\vartheta^\alpha\wedge\sigma_\alpha\right\rangle=-\frac{1}{3\pi^2}\left[\mathrm{Tr}(G\wedge {}^* G)+\frac{1}{24}\left(2R^{\alpha\beta\{\}}\wedge R^{j(\star)}_{\alpha\beta}+d\mathscr A\wedge {}^* d\mathscr A\right)\right],	$\langle\vartheta^\alpha\wedge\sigma_\alpha\rangle= -\frac{1}{3\pi^2}\left[{\rm Tr}\,(G\wedge {}^*G)+\frac{1}{24}\left(2R^{\alpha\beta\{\}}\wedge R^{j(\star)}_{\alpha\beta}+d\mathcal{A}\wedge {}^*d\mathcal{A}\right)\right],$	$\langle\vartheta^\alpha\wedge\sigma_\alpha\rangle= -\frac{1}{3\pi^2}\left[Tr(G\wedge {}^*G)+\frac{1}{24}\left(2R^{\alpha\beta\{\}}\wedge R^{j(\star)}_{\alpha\beta}+d\mathcal{A}\wedge {}^*d\mathcal{A}\right)\right],$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0774 (11.2.3)	245	■0.868 ●N +0	i(\overline{D}\psi)\wedge^*\gamma+\frac{i}{2}\bar\psi D\wedge^*\gamma+m\bar\psi\eta=0,	$i(\overline{D}\psi)\wedge^*\gamma +\frac{i}{2}\bar\psi D\wedge^*\gamma+m\bar\psi\eta=0,$	$i(\overline{D}\psi)\wedge^*\gamma +\frac{i}{2}\bar\psi D\wedge^*\gamma+m\bar\psi\eta=0,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0548 (8.4.2)	178	■0.871 ●N -1	s_L_{\text {Pontr }}=-2\,d\,{\operatorname{Tr}}\!\left({\Psi _1}\wedge {{\Omega ^g}_1}\right)= -d\left({{\Psi _1}_{\alpha }\wedge {{R_\beta }^\alpha }} \right).	$sL_{\text{Pontr}} \;=\;-2d\,{\rm Tr}\,\{\Psi\wedge\Omega^g\} \;=\;-d\left(\Psi_\alpha{}^\beta\wedge R_\beta{}^\alpha\right).$	$sL_{\text{Pontr}} \;=\;-2d{\rm Tr}\{\Psi\wedge\Omega^g\} \;=\;-d(\Psi_\alpha{}^\beta\wedge R_\beta{}^\alpha).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0244 (4.3.9)	091	■0.873 ●N +1	g_{ij}\, e_{\alpha}^i\cdots e_{\beta}^j=g_{\alpha\beta}	$g_{ij}e_{\alpha}^ie_{\beta}^j=g_{\alpha\beta}.$	$g_{ij}e_a^ie_b^j=g_{ab}.$
Conf. MathPix confidence: ■≥0.9 ■0.5–0.9 ■<0.5 — no value					
Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0890 (13.4.9)	287	0.876 N +0	$\hat{\mathrm{x}}_i=\mathrm{binom}\{\mathrm{x}_i\}{1}=\mathrm{binom}\{\mathrm{x}_i^{\alpha}\}{\mathrm{P}_{\alpha}}_{\{1\}},\quad \hat{\mathrm{x}}_i^{\ast}=\mathrm{left}(\mathrm{x}_i^{\ast},\,1\mathrm{right})=\mathrm{left}(\mathrm{x}_i^{\beta}\mathrm{P}_{\ast}^{\ast})_{\{\beta\}},\,1\mathrm{right})$	$\hat{\xi}=\begin{pmatrix}\xi\\1\end{pmatrix}=\begin{pmatrix}\xi^{\alpha}P_{\alpha}\\1\end{pmatrix},\quad \hat{\xi}_{\ast}=(\xi_{\ast},1)=\begin{pmatrix}\xi_{\beta}^{\ast}P_{\ast}^{\beta}\\1\end{pmatrix}$	$\hat{\xi}=\begin{pmatrix}\xi\\1\end{pmatrix}=\begin{pmatrix}\xi^{\alpha}P_{\alpha}\\1\end{pmatrix},\quad \hat{\xi}_{\ast}=(\xi_{\ast},1)=\begin{pmatrix}\xi_{\beta}^{\ast}P_{\ast}^{\beta}\\1\end{pmatrix}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1007 (14.4.2)	325	0.877 N +1	$\begin{aligned} A^{\{(\circ)\}}&=\stackrel{\ast}{\omega}^{\{(\circ)\}}=-\frac{r^2}{\rho^2}\left[dt-J_3\frac{\hbar}{Mc}\sin^2\vartheta d\phi\right]Q^{\{(\circ)\}}\cos\delta\\ &-\frac{\cos\vartheta}{\rho^2}\left[\left(r^2+J(J+1)\frac{\hbar^2}{M^2c^2}\right)d\phi-J_3\frac{\hbar}{M}dt\right]P^{\{(\circ)\}}\sin\delta \end{aligned}$	$A^{\{(\circ)\}}=\stackrel{\ast}{\omega}^{\{(\circ)\}}=-\frac{r^2}{\rho^2}\left[dt-J_3\frac{\hbar}{Mc}\sin^2\vartheta d\phi\right]Q^{\{(\circ)\}}\cos\delta\\ -\frac{\cos\vartheta}{\rho^2}\left[\left(r^2+J(J+1)\frac{\hbar^2}{M^2c^2}\right)d\phi-J_3\frac{\hbar}{M}dt\right]P^{\{(\circ)\}}\sin\delta$	$A^{\{(\circ)\}}=\stackrel{\ast}{\omega}^{\{(\circ)\}}=-\frac{r^2}{\rho^2}\left[dt-J_3\frac{\hbar}{Mc}\sin^2\vartheta d\phi\right]Q^{\{(\circ)\}}\cos\delta\\ -\frac{\cos\vartheta}{\rho^2}\left[\left(r^2+J(J+1)\frac{\hbar^2}{M^2c^2}\right)d\phi-J_3\frac{\hbar}{M}dt\right]P^{\{(\circ)\}}\sin\delta.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1062 (16.2.1)	355	0.877 N +0	$\mathrm{left}[\mathrm{square}-(\mathrm{mc}/\hbar)^2\mathrm{right}]\,\mathrm{varphi}=0$	$[\square-(\mathrm{mc}/\hbar)^2]\varphi=0$	$[\square-(mc/\hbar)^2]\varphi=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0051 (2.7.1)	044	0.881 N -2	$G(p)\colon P\rightarrowtail P,\quad G(p)\in\mathrm{mathscr{G}}_{\{p\}}$	$G(p):P\rightarrow P,\quad G(p)\in\mathscr{G}_p$	$G(p):P\rightarrow P,\quad G(p)\in\mathscr{G}_p$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0178 (3.7.8)	072	0.881 K +0	$\mathrm{left}\langle\mathrm{rangle}\mathrm{left.}\,0\mathrm{right} ^{\{\lambda\}}\,\mathrm{varphi}\,\mathrm{mid}\,0\,\mathrm{right}\rangle\mathrm{rangle}=\mu/\sqrt{\lambda}$	$\left\langle O\right ^{\lambda}\varphi\left O\right\rangle =\mu/\sqrt{\lambda}$	$\langle O\left ^{\lambda}\varphi\left O\right\rangle =\mu/\sqrt{\lambda}$
Conf. MathPix confidence: ≥0.9 0.5–0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0028 (2.5.1)	040	■0.885 ● N +0	$V_{\{p\}^{\wedge\{*\}}=T_{\{p\}^{\wedge\{*\}}\left(F_{\{m\}}(M)\right),$	$V_p^*=T_p^*(F_m(M)),$	$V_p^*=T_p^*(F_m(M)),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0920 (13.8.6)	295	■0.888 ● N -2	$\begin{aligned} & d\,C_{\{\mathrm{RR}\}(\star)}\,\&:=\frac{1}{2}\,d\left(\Gamma_{\alpha\beta}\wedge R^{\alpha\beta(\star)}-\frac{1}{3}\Gamma_{\alpha}{}^{\beta(\star)}\wedge\Gamma_{\beta}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\alpha}\right)\\ & \equiv -L_{\mathrm{SKY}}-2\,\mathrm{Ric}_{\alpha\beta}\wedge{}^*\mathrm{Ric}{}^{\alpha\beta}+\frac{1}{2}\,\mathrm{Ric}_{\alpha}{}^{\alpha}\wedge{}^*\mathrm{Ric}_{\beta}{}^{\beta} \end{aligned}$	$dC_{\mathrm{RR}(\star)}:=\frac{1}{2}d\left(\Gamma_{\alpha\beta}\wedge R^{\alpha\beta(\star)}-\frac{1}{3}\Gamma_{\alpha}{}^{\beta(\star)}\wedge\Gamma_{\beta}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\alpha}\right)\\ \equiv -L_{\mathrm{SKY}}-2\,\mathrm{Ric}_{\alpha\beta}\wedge{}^*\mathrm{Ric}{}^{\alpha\beta}+\frac{1}{2}\,\mathrm{Ric}_{\alpha}{}^{\alpha}\wedge{}^*\mathrm{Ric}_{\beta}{}^{\beta}$	$dC_{\mathrm{RR}(\star)}:=\frac{1}{2}d\left(\Gamma_{\alpha\beta}\wedge R^{\alpha\beta(\star)}-\frac{1}{3}\Gamma_{\alpha}{}^{\beta(\star)}\wedge\Gamma_{\beta}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\alpha}\right)\\ \equiv -L_{\mathrm{SKY}}-2\mathrm{Ric}_{\alpha\beta}\wedge{}^*\mathrm{Ric}^{\alpha\beta}+\frac{1}{2}\mathrm{Ric}_{\alpha}{}^{\alpha}\wedge{}^*\mathrm{Ric}_{\beta}{}^{\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0674 (10.3.31)	220	■0.890 ● C -5	$\check{\mathrm{E}}_{\{\alpha\}}=-\frac{\theta_{\mathrm{T}}}{\theta_{\mathrm{TL}}}\frac{\theta_{\mathrm{TL}}}{2\ell^2}\left(\frac{1}{\theta_{\mathrm{TL}}}d\vartheta_{\alpha}-\frac{4\ell}{\theta_{\mathrm{T}}\theta_{\mathrm{L}}}\check{\mathcal{J}}_{\alpha}^{\star}\right),$	$\check{\mathcal{E}}_{\alpha}=-\frac{\theta_{\mathrm{T}}\theta_{\mathrm{TL}}}{2\ell^2}\left(\frac{1}{\theta_{\mathrm{TL}}}d\vartheta_{\alpha}-\frac{4\ell}{\theta_{\mathrm{T}}\theta_{\mathrm{L}}}\check{\mathcal{J}}_{\alpha}^{\star}\right),$	$\check{\mathcal{E}}_{\alpha}=-\frac{\theta_{\mathrm{T}}\theta_{\mathrm{TL}}}{2\ell^2}\left(\frac{1}{\theta_{\mathrm{TL}}}d\vartheta_{\alpha}-\frac{4\ell}{\theta_{\mathrm{T}}\theta_{\mathrm{L}}}\check{\mathcal{J}}_{\alpha}^{\star}\right),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0531 (7.7.51)	169	■0.890 ● N +0	$\left.\left.\left.R:=e_{\{\beta\}}\right\rfloor\right\rfloor e_{\{\alpha\}}\right\rfloor R^{\alpha\beta},\quad{}^*R^{\alpha\beta}\wedge\vartheta_{\alpha}\wedge\vartheta_{\beta}\equiv R^{\alpha\beta}\wedge\eta_{\alpha\beta}\equiv-R\eta,$	$R:=e_{\beta}\rfloor e_{\alpha}\rfloor R^{\alpha\beta},\quad{}^*R^{\alpha\beta}\wedge\vartheta_{\alpha}\wedge\vartheta_{\beta}\equiv R^{\alpha\beta}\wedge\eta_{\alpha\beta}\equiv-R\eta,$	$R:=e_{\beta}\rfloor e_{\alpha}\rfloor R^{\alpha\beta},\quad{}^*R^{\alpha\beta}\wedge\vartheta_{\alpha}\wedge\vartheta_{\beta}\equiv R^{\alpha\beta}\wedge\eta_{\alpha\beta}\equiv-R\eta,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0540 (8.3.1)	176	■0.891 ● N +2	$\begin{aligned} & (d\oplus s)\,\gamma+[\widetilde{\Gamma},\gamma]=\widetilde{\Theta} \\ & (d\oplus s)\widetilde{\Theta}+[\widetilde{\Gamma},\widetilde{\Theta}]\equiv\left[\widetilde{\Omega}^g,\gamma\right] \end{aligned}$	$(d\oplus s)\gamma+[\widetilde{\Gamma},\gamma]=\widetilde{\Theta} \\ (d\oplus s)\widetilde{\Theta}+[\widetilde{\Gamma},\widetilde{\Theta}]\equiv\left[\widetilde{\Omega}^g,\gamma\right]$	$(d\oplus s)\gamma+[\widetilde{F},\gamma]=\widetilde{\Theta}, \\ (d\oplus s)\widetilde{\Theta}+[\widetilde{F},\widetilde{\Theta}]\equiv[\widetilde{\Omega}^g,\gamma],$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0925 (14.1.5)	303	0.892 N +0	$\begin{aligned} & D^{*} \Omega = \tau_e, \quad \& D \Omega \equiv 0, \\ & \frac{1}{2} * (\vartheta \wedge * \Omega^g) = \ell^{*2} (\Sigma_{\text{YM}} + \Sigma_{\text{mat}}), \end{aligned}$	$D^{*} \Omega = \tau_e, \\ D \Omega \equiv 0, \\ \frac{1}{2} * (\vartheta \wedge * \Omega^g) = \ell^{*2} (\Sigma_{\text{YM}} + \Sigma_{\text{mat}}),$	$D^{*} \Omega = \tau_e, \\ D \Omega \equiv 0, \\ \frac{1}{2} * (\vartheta \wedge * \Omega^g) = \ell^{*2} (\Sigma_{\text{YM}} + \Sigma_{\text{mat}}),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0867 (13.2.3)	283	0.893 N +0	$B \cong dA, \quad dB \cong dF \equiv 0,$	$B \cong dA, \quad dB \cong dF \equiv 0,$	$B \cong dA, \quad dB \cong dF \equiv 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0609 (9.2.11)	201	0.894 N +1	$\begin{aligned} & \left(1 + \zeta^2\right) \int_{M^4} L_{\text{SKY}} - (-1)^s \zeta (4\pi)^2 \cdot \chi(M^4) \\ & = \frac{1}{4} \int_{M^4} \left(\Omega^g - \zeta^* \Omega^{g(*)}\right) \wedge^* \left(\Omega^g - \zeta^* \Omega^{g(*)}\right) \geq 0, \end{aligned}$	$\left(1 + \zeta^2\right) \int_{M^4} L_{\text{SKY}} - (-1)^s \zeta (4\pi)^2 \cdot \chi(M^4) \\ = \frac{1}{4} \int_{M^4} \left(\Omega^g - \zeta^* \Omega^{g(*)}\right) \wedge^* \left(\Omega^g - \zeta^* \Omega^{g(*)}\right) \geq 0,$	$\left(1 + \zeta^2\right) \int_{M^4} L_{\text{SKY}} - (-1)^s \zeta (4\pi)^2 \cdot \chi(M^4) \\ = \frac{1}{4} \int_{M^4} (\Omega^g - \zeta^* \Omega^{g(*)}) \wedge^* (\Omega^g - \zeta^* \Omega^{g(*)}) \geq 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0959 (14.1.41)	309	0.894 N +0	$\oint d\delta = 2\pi b_1 = 2\pi v$	$\oint d\delta = 2\pi b_1 = 2\pi v$	$\oint d\delta = 2\pi b_1 = 2\pi v$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0299 (4.5.10)	102	0.896 N +1	$\begin{aligned} & \Delta g_{\alpha\beta} = 2\varepsilon_{(\alpha}{}^\gamma g_{\beta)\gamma}, \quad \Delta \vartheta^\alpha = -\vartheta^\beta \varepsilon_\beta{}^\alpha, \\ & \Delta \Gamma_\alpha{}^\beta = D\varepsilon_\alpha{}^\beta, \quad \Delta \Psi = -\varepsilon_\alpha{}^\beta I^\alpha{}_\beta \Psi. \end{aligned}$	$\delta g_{\alpha\beta} = 2\varepsilon_{(\alpha}{}^\gamma g_{\beta)\gamma}, \quad \delta \vartheta^\alpha = -\vartheta^\beta \varepsilon_\beta{}^\alpha, \\ \delta \Gamma_\alpha{}^\beta = D\varepsilon_\alpha{}^\beta, \quad \delta \Psi = -\varepsilon_\alpha{}^\beta I^\alpha{}_\beta \Psi.$	$\delta g_{\alpha\beta} = 2\varepsilon_{(\alpha}{}^\gamma g_{\beta)\gamma}, \quad \delta \vartheta^\alpha = -\vartheta^\beta \varepsilon_\beta{}^\alpha, \\ \delta \Gamma_\alpha{}^\beta = D\varepsilon_\alpha{}^\beta, \quad \delta \Psi = -\varepsilon_\alpha{}^\beta I^\alpha{}_\beta \Psi.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0236 (4.3.1)	090	0.896 N +0	$\begin{aligned} (\Delta s)^2 &= -c^2(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 \\ &= g_{ij} \Delta x^i \Delta x^j. \end{aligned}$	$\begin{aligned} (\Delta s)^2 &= -c^2(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 \\ &= g_{ij} \Delta x^i \Delta x^j. \end{aligned}$	$\begin{aligned} (\Delta s)^2 &= -c^2(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 \\ &= g_{ij} \Delta x^i \Delta x^j. \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1005 (14.3.18)	323	0.896 N -1	$\begin{aligned} \not{D}^2 &= \operatorname{Tr} \left(\frac{1}{2} \gamma \wedge \gamma \wedge^* \{D, D\} + \frac{1}{2} \gamma \wedge \gamma \wedge^* \tilde{\Omega} - \gamma \wedge \tilde{\Theta} \wedge^* D \right) \\ &= \left(\square^g + \frac{1}{4} R \right) \eta - \vartheta \wedge \Theta \wedge^* D \end{aligned}$	$\begin{aligned} D^2 &= \operatorname{Tr} \left(\frac{1}{2} \gamma \wedge \gamma \wedge^* \{D, D\} + \frac{1}{2} \gamma \wedge \gamma \wedge^* \tilde{\Omega} - \gamma \wedge \tilde{\Theta} \wedge^* D \right) \\ &= \left(\square^g + \frac{1}{4} R \right) \eta - \vartheta \wedge \Theta \wedge^* D \end{aligned}$	$\begin{aligned} \not{D}^2 &= \operatorname{Tr} \left(\frac{1}{2} \gamma \wedge \gamma \wedge^* \{D, D\} + \frac{1}{2} \gamma \wedge \gamma \wedge^* \tilde{\Omega} - \gamma \wedge \tilde{\Theta} \wedge^* D \right) \\ &= (\square^g + \frac{1}{4} R) \eta - \vartheta \wedge \Theta \wedge^* D \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0230 (4.2.23)	088	0.899 N +0	$\stackrel{*}{\sigma} \Theta = \frac{1}{2} T_{\alpha\beta}^c P_c \otimes \vartheta^\alpha \wedge \vartheta^\beta = \frac{1}{2} T_{ij}^c P_c \otimes dx^i \wedge dx^j,$	$\stackrel{*}{\sigma} \Theta = \frac{1}{2} T_{\alpha\beta}^c P_c \otimes \vartheta^\alpha \wedge \vartheta^\beta = \frac{1}{2} T_{ij}^c P_c \otimes dx^i \wedge dx^j,$	$\stackrel{*}{\sigma} \Theta = \frac{1}{2} T_{\alpha\beta}^c P_c \otimes \vartheta^\alpha \wedge \vartheta^\beta = \frac{1}{2} T_{ij}^c P_c \otimes dx^i \wedge dx^j,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0723 (10.10.3)	230	0.900 C +4	$\begin{aligned} \delta_{\text{susy}} \vartheta^\alpha &= i \bar{\sigma} \Psi \gamma^\alpha, \quad \delta_{\text{susy}} \Gamma_\alpha^* = i \bar{\sigma} \gamma_\alpha^* D \Psi + i c \bar{\sigma} (\gamma_\alpha \Psi + e_\alpha]^* \Psi), \\ \delta_{\text{susy}} \Psi &= 2 D \sigma + c \gamma \sigma, \end{aligned}$	$\begin{aligned} \delta_{\text{susy}} \vartheta^\alpha &= i \bar{\sigma} \Psi \gamma^\alpha, \quad \delta_{\text{susy}} \Gamma_\alpha^* = i \bar{\sigma} \gamma_\alpha^* D \Psi + i c \bar{\sigma} (\gamma_\alpha \Psi + e_\alpha]^* \Psi) \\ \delta_{\text{susy}} \Psi &= 2 D \sigma + c \gamma \sigma, \end{aligned}$	$\begin{aligned} \delta_{\text{susy}} \vartheta^\alpha &= i \bar{\sigma} \Psi \gamma^\alpha, \quad \delta_{\text{susy}} \Gamma_\alpha^* = i \bar{\sigma} \gamma_\alpha^* D \Psi + i c \bar{\sigma} (\gamma_\alpha \Psi + e_\alpha]^* \Psi), \\ \delta_{\text{susy}} \Psi &= 2 D \sigma + c \gamma \sigma, \end{aligned} \tag{1}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0216 (4.2.8)	086	0.902 N +0	$\widetilde{\omega} \rightarrow^{A^{-1}} \widetilde{\omega} = A^{-1} \widetilde{\omega} A - A^{-1} dA, \quad A \in \mathcal{A}_p.$	$\widetilde{\omega} \rightarrow^{A^{-1}} \widetilde{\omega} = A^{-1} \widetilde{\omega} A - A^{-1} dA, \quad A \in \mathcal{A}_p.$	$\widetilde{\omega} \rightarrow^{A^{-1}} \widetilde{\omega} = A^{-1} \widetilde{\omega} A - A^{-1} dA, \quad A \in \mathcal{A}_p.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1042 (15.1.19)	340	■ 0.902 ● K +0	$\backslash\mathrm{Gamma}_{\mu\alpha\beta}:\backslashleft(2^{\wedge\{+\}}\backslash\mathrm{right})\backslash\mathrm{oplus}2\backslashleft(1^{\wedge\{-}\}\backslash\mathrm{right})\backslash\mathrm{oplus}\backslashleft(0^{\wedge\{+\}}\backslash\mathrm{right})\backslash\mathrm{oplus}\backslashleft(2^{\wedge\{-}\}\backslash\mathrm{right})\backslash\mathrm{oplus}2\backslashleft(1^{\wedge\{+\}}\backslash\mathrm{right})\backslash\mathrm{oplus}\backslashleft(0^{\wedge\{-}\}\backslash\mathrm{right})\right),$	$\Gamma_{\mu\alpha\beta}:(2^+) \oplus 2(1^-) \oplus (0^+) \oplus (2^-) \oplus 2(1^+) \oplus (0^-),$	$\Gamma_{\mu\alpha\beta}:(2^+) \oplus 2(1^-) \oplus (0^+) \oplus (2^-) \oplus 2(1^+) \oplus (0^-),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0481 (7.5.2)	157	■ 0.903 ● W -1	$\backslash\mathrm{square}^{\wedge\{c\}}g_{ij}=\backslash\mathrm{square}^{\wedge\{c\}}\backslashleft(\backslash\mathrm{Omega}^{\wedge\{2\}}o_{ij}^{\wedge\{\text{Mink}\}}\backslash\mathrm{right})=\backslash\mathrm{Omega}^{\wedge\{6\}}\backslash\mathrm{square}^{\wedge\{c\}}o_{ij}^{\wedge\{\text{Mink}\}}=0$	$\square^c g_{ij} = \square^c \left(\Omega^2 o_{ij}^{\mathrm{Mink}} \right) = \Omega^6 \square^c o_{ij}^{\mathrm{Mink}} = 0$	$\square^c g_{ij} = \square^c (\varOmega^2 o_{ij}^{\mathrm{Mink}}) = \varOmega^6 \square^c o_{ij}^{\mathrm{Mink}} = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0117 (3.3.10)	059	■ 0.905 ● N -1	$L_{\mathrm{Max}}:=\backslash\mathrm{frac}\{1\}\{2\}F\backslash\mathrm{wedge}^{\wedge\{*\}}F=\backslash\mathrm{frac}\{1\}\{2\}\backslashleft(\backslash\mathrm{mathbf}\{E\}^{\wedge\{2\}}-\backslash\mathrm{mathbf}\{B\}^{\wedge\{2\}}\backslash\mathrm{right})\backslash\mathrm{eta},$	$L_{\mathrm{Max}} := -\frac{1}{2}F \wedge^* F = \frac{1}{2}(\mathbf{E}^2 - \mathbf{B}^2)\eta,$	$L_{\mathrm{Max}} := -\frac{1}{2}F \wedge^* F = \frac{1}{2}(\mathbf{E}^2 - \mathbf{B}^2)\eta,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1110 (B.22)	371	■ 0.905 ● K +0	$\backslash\mathrm{square}\alpha=0,$	$\square\alpha=0,$	$\square\alpha=0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0831 (12.1.3)	270	■ 0.907 ● N +0	$\backslash\mathrm{tilde}\{j\}_{5}:=j_{5}+\backslashleft(1\ /\ 96\ \backslash\mathrm{pi}^{\wedge\{2\}}\backslash\mathrm{right})A\backslash\mathrm{wedge}dA$	$\tilde{j}_5 := j_5 + (1/96\pi^2) A \wedge dA$	$\widetilde{j}_5 := j_5 + (1/96\pi^2)A \wedge dA$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00364 (6.2.16)	126	■ 0.912 ● N +0	$D\,D\,\lambda_{\alpha\beta}=-2\,R_{[\alpha\,\mid\beta]}{\,\,}^{\gamma}\,\gamma^{\mid}\wedge\lambda_{\gamma\mid\beta]}=0$	$DD\lambda_{\alpha\beta}=-2R_{[\alpha]}{}^{\gamma}\wedge\lambda_{\gamma \beta]}=0$	$DD\lambda_{\alpha\beta}=-2R_{[\alpha]}{}^{\gamma}\wedge\lambda_{\gamma \beta]}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00840 (12.2.8)	272	■ 0.916 ● N +1	$\begin{aligned} \tau_{\alpha\beta} &:= \frac{\partial L_D}{\partial \Gamma^{\alpha\beta}} = \frac{1}{8} \bar{\psi} \left(\gamma^{\alpha} \sigma_{\alpha\beta} + \sigma_{\alpha\beta} \gamma^{\beta} \right) \psi \\ &= \frac{1}{4} \eta_{\alpha\beta\gamma\delta} \bar{\psi} \gamma^{\delta} \gamma_5 \psi \eta^{\gamma} = \tau_{\alpha\beta\gamma} \eta^{\gamma} = \vartheta_{[\alpha} \wedge \mu_{\beta]} \end{aligned}$	$\tau_{\alpha\beta} := \frac{\partial L_D}{\partial \Gamma^{\alpha\beta}} = \frac{1}{8} \bar{\psi} \left(\gamma^{\alpha} \sigma_{\alpha\beta} + \sigma_{\alpha\beta} \gamma^{\beta} \right) \psi \\ = \frac{1}{4} \eta_{\alpha\beta\gamma\delta} \bar{\psi} \gamma^{\delta} \gamma_5 \psi \eta^{\gamma} = \tau_{\alpha\beta\gamma} \eta^{\gamma} = \vartheta_{[\alpha} \wedge \mu_{\beta]}$	$\tau_{\alpha\beta} := \frac{\partial L_D}{\partial \Gamma^{\alpha\beta}} = \frac{1}{8} \bar{\psi} \left(\gamma^{\alpha} \sigma_{\alpha\beta} + \sigma_{\alpha\beta} \gamma^{\beta} \right) \psi \\ = \frac{1}{4} \eta_{\alpha\beta\gamma\delta} \bar{\psi} \gamma^{\delta} \gamma_5 \psi \eta^{\gamma} = \tau_{\alpha\beta\gamma} \eta^{\gamma} = \vartheta_{[\alpha} \wedge \mu_{\beta]}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00232 (4.2.25)	088	■ 0.917 ● K +0	$\Theta \rightarrow \Theta^A \Theta = G^{-1} \Theta,$	$\Theta \rightarrow \Theta^A \Theta = G^{-1} \Theta,$	$\Theta \rightarrow \Theta^A \Theta = G^{-1} \Theta,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00577 (8.6.15)	185	■ 0.920 ● C +25	$-E_{\alpha}=\frac{\theta_{\mathrm{T}}}{2}R^{\left\{\beta\gamma\right\}}\wedge\eta_{\alpha\beta\gamma}-\theta_{\mathrm{T}}^{\ast}\Lambda_{\mathrm{eff}}\eta_{\alpha}=\ell^2\widehat{\Sigma}_{\alpha}.$	(not rendered)	$-E_{\alpha}=\frac{\theta_{\mathrm{T}}^{\ast}}{2}R^{\left\{\beta\gamma\right\}}\wedge\eta_{\alpha\beta\gamma}-\theta_{\mathrm{T}}^{\ast}\Lambda_{\mathrm{eff}}\eta_{\alpha}=\ell^2\widehat{\Sigma}_{\alpha}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E01031 (15.1.8)	337	■ 0.920 ● W -1	$\begin{aligned} & \frac{1}{2} \left(\vartheta^f \wedge \Omega^f \right) - a \ell^2 \vartheta^f \wedge \left\{ 2 L_{fg} - \frac{1}{2 b \ell^2} \left(\Omega^g \wedge \vartheta^g \wedge \vartheta^g + 2 \Lambda \eta \right) \right\} \\ &= \ell^2 \left\{ \Sigma_{\mathrm{m}} \left(\mathrm{hadrons} \right) + \frac{a}{b} \left(\vartheta^g \wedge \Sigma_{\mathrm{m}} \left(\mathrm{leptons} \right) \right) \wedge \vartheta^f \right\} \end{aligned}$	$\frac{1}{2} \left(\vartheta^f \wedge \Omega^f \right) - a \ell^2 \vartheta^f \wedge \left\{ 2 L_{fg} - \frac{1}{2 b \ell^2} \left(\Omega^g \wedge \vartheta^g \wedge \vartheta^g + 2 \Lambda \eta \right) \right\} \\ = \ell^2 \left\{ \Sigma_{\mathrm{m}} \left(\mathrm{hadrons} \right) + \frac{a}{b} \left(\vartheta^g \wedge \Sigma_{\mathrm{m}} \left(\mathrm{leptons} \right) \right) \wedge \vartheta^f \right\}$	$\frac{1}{2} \left(\vartheta^f \wedge \Omega^f \right) - a \ell^2 \vartheta^f \wedge \left\{ 2 L_{fg} - \frac{1}{2 b \ell^2} \left(\Omega^g \wedge \vartheta^g \wedge \vartheta^g + 2 \Lambda \eta \right) \right\} \\ = \ell^2 \left\{ \Sigma_{\mathrm{m}} \left(\mathrm{hadrons} \right) + \frac{a}{b} \left(\vartheta^g \wedge \Sigma_{\mathrm{m}} \left(\mathrm{leptons} \right) \right) \wedge \vartheta^f \right\}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0473 (7.4.1)	156	■0.921 ● N +1	$\begin{aligned} & \int \mathcal{D}\Gamma \exp \left[\int L_{\text{SKY}}^{(*)} / \hbar \right] \\ &= \int \mathcal{D}\Gamma \exp \left[- \int \left(R_{\alpha\beta} + {}^*R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} + {}^*R^{\alpha\beta(*)} \right) / 4\hbar \right] \end{aligned}$	$\int \mathcal{D}\Gamma \exp \left[\int L_{\text{SKY}}^{(*)} / \hbar \right] \\ = \int \mathcal{D}\Gamma \exp \left[- \int \left(R_{\alpha\beta} + {}^*R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} + {}^*R^{\alpha\beta(*)} \right) / 4\hbar \right]$	$\int \mathcal{D}\Gamma \exp \left[\int L_{\text{SKY}}^{(*)} / \hbar \right] \\ = \int \mathcal{D}\Gamma \exp \left[- \int \left(R_{\alpha\beta} + {}^*R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} + {}^*R^{\alpha\beta(*)} \right) / 4\hbar \right]$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0343 (5.4.1)	115	■0.922 ● N +0	$k \, \partial_k \Gamma_k = \frac{1}{2} \, \text{STr} \left\{ \left[\Gamma_k^{(2)} + R_k \right]^{-1} (k \partial_k R_k) \right\},$	$k \partial_k \Gamma_k = \frac{1}{2} \, \text{STr} \left\{ \left[\Gamma_k^{(2)} + R_k \right]^{-1} (k \partial_k R_k) \right\},$	$k \partial_k \Gamma_k = \frac{1}{2} \text{STr} \left\{ [\Gamma_k^{(2)} + R_k]^{-1} (k \partial_k R_k) \right\},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0829 (12.1.1)	269	■0.922 ● W +0	$\bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \rightarrow \bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \mathbf{P} \exp \left\{ i \int_x^{x+\varepsilon} A \right\}.$	$\bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \rightarrow \bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \mathbf{P} \exp \left\{ i \int_x^{x+\varepsilon} A \right\}.$	$\overline{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \rightarrow \overline{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \mathbf{P} \exp \left\{ i \int_x^{x+\varepsilon} A \right\}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0736 (11.1.3)	238	■0.922 ● N +0	$\gamma^0 := \left(\begin{array}{rr} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{array} \right), \quad \gamma^k := \left(\begin{array}{rr} 0 & \sigma^k \\ -\sigma^k & 0 \end{array} \right), \quad \sigma^k : \text{Pauli matrices.}$	$\gamma^0 := \left(\begin{array}{rr} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{array} \right), \quad \gamma^k := \left(\begin{array}{rr} 0 & \sigma^k \\ -\sigma^k & 0 \end{array} \right), \quad \sigma^k : \text{Pauli matrices.}$	$\gamma^0 := \left(\begin{array}{rr} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{array} \right), \quad \gamma^k := \left(\begin{array}{rr} 0 & \sigma^k \\ -\sigma^k & 0 \end{array} \right), \quad \sigma^k : \text{Pauli matrices.}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0661 (10.3.18)	217	■0.923 ● N +0	$\square H - (-1)^s {}^* D {}^* D H = dt + \frac{\gamma}{\delta \ell^2} d\mathcal{A} + \frac{1}{\delta \ell} dh,$	$\square H - (-1)^s {}^* D {}^* D H = dt + \frac{\gamma}{\delta \ell^2} d\mathcal{A} + \frac{1}{\delta \ell} dh,$	$\square H - (-1)^s {}^* D {}^* D H = dt + \frac{\gamma}{\delta \ell^2} d\mathcal{A} + \frac{1}{\delta \ell} dh,$
Conf. MathPix confidence: ■ ■ ■ — ≥0.90.5–0.9<0.5no value Residual render vs scan (inkdrill): ● ● ● ● ● ● C componentW weakS stableN noiseK cleannot measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__E00685 (10.3.43)	222	■0.923 ●K +0	$C_{\alpha}-E_{\alpha}=\Sigma_{\alpha}-D^{\mathfrak{U}}\mu_{\alpha},$	$C_{\alpha}-E_{\alpha}=\Sigma_{\alpha}-D^{\mathfrak{U}}\mu_{\alpha},$	$C_{\alpha}-E_{\alpha}=\Sigma_{\alpha}-D^{\mathfrak{U}}\mu_{\alpha},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__E00702 (10.6.1)	225	■0.924 ●N -1	$L_{\mathfrak{A}}=\frac{1}{2}\left(F\wedge\right)^{*}F+m_{\text {photon }}F\wedge A\right).$	$L_A=\frac{1}{2}\left(F\wedge^*F+m_{\text{photon}}F\wedge A\right).$	$L_A=\frac{1}{2}(F\wedge^*F+m_{\text{photon}}F\wedge A).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__E00221 (4.2.14)	087	■0.924 ●K +0	$\Omega^{\mathrm{T}}\rightarrow\left\{A^{-1}\right\}\Omega^{\mathrm{T}}=G^{-1}\left(\Omega^{\mathrm{T}}-DDT\right)=G^{-1}\Omega^{\mathrm{T}}-G^{-1}\Omega^{\mathrm{L}}\otimes T.$	$\Omega^{\mathrm{T}}\rightarrow^{A^{-1}}\Omega^{\mathrm{T}}=G^{-1}\left(\Omega^{\mathrm{T}}-DDT\right)=G^{-1}\Omega^{\mathrm{T}}-G^{-1}\Omega^{\mathrm{L}}\otimes T.$	$\Omega^{\mathrm{T}}\rightarrow^{A^{-1}}\Omega^{\mathrm{T}}=G^{-1}(\Omega^{\mathrm{T}}-DDT)=G^{-1}\Omega^{\mathrm{T}}-G^{-1}\Omega^{\mathrm{L}}\otimes T.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__E00443 (6.8.14)	143	■0.924 ●N -1	$H_{\alpha}=-\frac{1}{\ell^2}\ast\left(\sum_{i=1}^3a_i^{(i)}T_{\alpha}\right),$	$H_{\alpha}=-\frac{1}{\ell^2}\ast\left(\sum_{i=1}^3a_i^{(i)}T_{\alpha}\right),$	$H_{\alpha}=-\frac{1}{\ell^2}\ast\left(\sum_{i=1}^3a_i^{(i)}T_{\alpha}\right),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__E00192 (4.1.2)	081	■0.924 ●N +0	$\mathfrak{a}:=\left[\begin{array}{c c}g&\mathfrak{a}\\\hline0&1\end{array}\right]=\left[\begin{array}{c c}1_n&\mathfrak{a}\\\hline0&1\end{array}\right]\left[\begin{array}{c c}g&0\\\hline0&1\end{array}\right]=a^{\mathrm{T}}\cdot a^{\mathrm{L}}\in A(n,\mathbb{R}).$	$a:=\left[\begin{array}{c c}g&\mathfrak{a}\\\hline0&1\end{array}\right]=\left[\begin{array}{c c}1_n&\mathfrak{a}\\\hline0&1\end{array}\right]\left[\begin{array}{c c}g&0\\\hline0&1\end{array}\right]=a^{\mathrm{T}}\cdot a^{\mathrm{L}}\in A(n,\mathbb{R}).$	$a:=\left[\begin{array}{c c}g&\mathfrak{a}\\\hline0&1\end{array}\right]=\left[\begin{array}{c c}1_n&\mathfrak{a}\\\hline0&1\end{array}\right]\left[\begin{array}{c c}g&0\\\hline0&1\end{array}\right]=a^{\mathrm{T}}\cdot a^{\mathrm{L}}\in A(n,\mathbb{R}).$
Conf. MathPix confidence: ■≥0.9 ■0.5–0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__ EQ0857 (12.4.4)	278	■0.925 ●C +3	$\left.\mathrm{bf}\{\mathbb{C}\}_n\ \mathrm{mathscr{G}}^{\mathrm{A}}\ \mathrm{cong}\ n\right\rfloor\left(D\ \mathrm{underline{\tau}}^{\mathrm{A}}\right),$	(not rendered)	$\mathbb{L}_n\mathcal{G}^A\cong n\rfloor(D\underline{\tau}^A),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__ EQ1025 (15.1.1)	336	■0.926 ●W -1	$\begin{aligned} L_{\mathrm{f-g}}= & \frac{1}{\ell^2} L_{\mathrm{W}}(f)+\frac{1}{\ell^2} L_{\mathrm{W}}(g)+L_{fg}+\frac{\Lambda}{\ell^2} \vartheta^g\wedge\vartheta^g\wedge\vartheta^g\wedge\vartheta^g \\ & +L_{\mathrm{m}}\left(\text{hadrons},f\right)+L_{\mathrm{m}}\left(\text{leptons},g\right) . \end{aligned}$	$L_{f-g}=\frac{1}{\ell^2}L_{\mathrm{W}}(f)+\frac{1}{\ell^2}L_{\mathrm{W}}(g)+L_{fg}+\frac{\Lambda}{\ell^2}\vartheta^g\wedge\vartheta^g\wedge\vartheta^g\wedge\vartheta^g+L_{\mathrm{m}}\left(\text{hadrons},f\right)+L_{\mathrm{m}}\left(\text{leptons},g\right).$	$L_{f-g}=\frac{1}{\ell^2}L_{\mathrm{W}}(f)+\frac{1}{\ell^2}L_{\mathrm{W}}(g)+L_{fg}+\frac{\Lambda}{\ell^2}\vartheta^g\wedge\vartheta^g\wedge\vartheta^g\wedge\vartheta^g+L_{\mathrm{m}}(\text{hadrons},f)+L_{\mathrm{m}}(\text{leptons},g).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__ EQ0109 (3.3.10)	058	■0.927 ●N -1	$L_{\mathrm{Max}}=-\frac{1}{2}F\wedge\mathrm{ }^*F,$	$L_{\mathrm{Max}}=-\frac{1}{2}F\wedge\mathrm{ }^*F,$	$L_{\mathrm{Max}}=-\frac{1}{2}F\wedge\mathrm{ }^*F,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__ EQ0587 (9.1.5)	194	■0.928 ●N +0	$d\ r^{*}:=e^{\left(\lambda-v\right)/2}\}d\ r\ .$	$dr^{*}:=e^{(\lambda-v)/2)}dr.$	$dr^{*}:=e^{(\lambda-v)/2)}dr.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__ EQ1101 (B.13)	370	■0.929 ●W -1	$\ast(\underbrace{\vartheta\wedge\cdots\wedge\vartheta}_{\mathrm{p\ products}})=\frac{p!}{(n-p)!}\underbrace{\vartheta\wedge\cdots\wedge\vartheta}_{\mathrm{n-p\ products}}(\star),$	$\ast(\underbrace{\vartheta\wedge\cdots\wedge\vartheta}_{\mathrm{p\ products}})=\frac{p!}{(n-p)!}\underbrace{\vartheta\wedge\cdots\wedge\vartheta}_{\mathrm{n-p\ products}}(\star),$	$\ast(\underbrace{\vartheta\wedge\cdots\wedge\vartheta}_{\mathrm{p\ products}})=\frac{p!}{(n-p)!}\underbrace{\vartheta\wedge\cdots\wedge\vartheta}_{\mathrm{n-p\ products}}(\star),$
Conf. MathPix confidence: ■≥0.9 ■0.5-0.9 ■<0.5 —no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0413 (6.5.1)	137	■0.929 ● N +0	$W(A, \gamma) = \operatorname{Tr} \mathbf{P} \exp \oint A = \operatorname{Tr} \mathbf{P} \exp \oint ds \dot{\gamma}^i(s) A_i^J(\gamma(s)) \lambda_J,$		
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0465 (7.3.4)	153	■0.929 ● N +1	$\left. \left. E_{\alpha} := \partial L_{\text{SKY}} / \partial \vartheta^{\alpha} = \frac{1}{2} (e_{\alpha} \rfloor R^{\mu\nu} \wedge {}^* R_{\mu\nu} - R^{\mu\nu} \wedge e_{\alpha} \rfloor {}^* R_{\mu\nu}) \right = 0 \right.$		
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ1034 (15.1.11)	338	■0.930 ● N +0	$\Lambda_{\text{eff.}}^f = \frac{2a}{\kappa b} \Lambda - 2a \ell^2 {}^* L_{fg}$		
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0512 (7.7.30)	165	■0.931 ● N -1	$\begin{aligned} \Omega^g := d\Gamma + \Gamma \wedge \Gamma &= \frac{i}{4} (d\Gamma_{\alpha}{}^{\beta} - \Gamma_{\alpha}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\beta}) \sigma^{\alpha}{}_{\beta} \\ &= \frac{i}{4} R^{\alpha\beta} \sigma_{\alpha\beta} = \frac{i}{8} R_{ij}{}^{\alpha\beta} \sigma_{\alpha\beta} dx^i \wedge dx^j, \end{aligned}$		
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0854 (12.4.1)	277	■0.931 ● C +4	$\stackrel{\text{rel}}{\text{pm}}\{V\}_{\mathrm{EC}} := V_{\mathrm{EC}} \mp \frac{1}{2} \operatorname{Tr} \left(\left(1 \mp \gamma_5 \right) \Omega \wedge \sigma \right) = - \frac{1}{2 \ell^2} R^{\alpha \beta} \wedge \eta_{\alpha \beta} + \frac{\Lambda}{\ell^2} \eta$		

Conf.

MathPix confidence:

■ ≥0.9

■ 0.5–0.9

■ <0.5

— no value

Residual

render vs scan (inkdrill):

● C component

● W weak

● S stable

● N noise

● K clean

● not measured

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0530 (7.7.50)	169	■0.931 ● N +1	$\left.\right\}^{\ast}\mathrm{R}_{\alpha\,\beta}^{}\equiv(-1)^{\mathrm{sig}}\mathrm{R}_{\alpha\,\beta}+\mathrm{e}_{[\alpha}\!\!\left\lfloor G_{\beta]}+\frac{1}{4}R\eta_{\alpha\beta}+(-1)^{\mathrm{sig}+1}D_{[\alpha}T_{\beta]}$	$*R^{(\star)}_{\alpha\beta}\equiv (-1)^{\mathrm{sig}} R_{\alpha\beta} + e_{[\alpha}\rfloor G_{\beta]} + \frac{1}{4} R \eta_{\alpha\beta} + (-1)^{\mathrm{sig}+1} D_{[\alpha} T_{\beta]}$	$*R^{(\star)}_{\alpha\beta}\equiv (-1)^{\mathrm{sig}} R_{\alpha\beta} + e_{[\alpha}\rfloor G_{\beta]} + \frac{1}{4} R \eta_{\alpha\beta} + (-1)^{\mathrm{sig}+1} D_{[\alpha} T_{\beta]}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0556 (8.4.10)	180	■0.933 ● N +0	$R_{\alpha\,\beta}=\mp\left\}^{\ast}\mathrm{R}_{\alpha\,\beta}^{}\right\}$	$R_{\alpha\beta} = \mp *R^{(\star)}_{\alpha\beta}$	$R_{\alpha\beta} = \mp *R^{(\star)}_{\alpha\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0547 (8.4.1)	178	■0.935 ● N +0	$\begin{aligned} L_{\mathrm{Pontr}} &\&:= d C_{\mathrm{RR}}=\\ \operatorname{Tr}\bigl(\! \omega^g \wedge \Omega^g \\ \bigr) &= \frac{1}{2} \, R_{\alpha}{}^{\beta} \wedge R_{\beta}{}^{\alpha} \\ &= \frac{1}{2} d \Bigl(\Gamma_{\alpha}{}^{\beta} \wedge R_{\beta}{}^{\alpha} - \frac{1}{3} \Gamma_{\alpha}{}^{\beta} \wedge \Gamma_{\beta}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\alpha} \Bigr) \end{aligned}$	$L_{\text{Pontr}} := dC_{\text{RR}} = \text{Tr}\{\Omega^g \wedge \Omega^g\} = \frac{1}{2} R_\alpha^\beta \wedge R_\beta^\alpha$ $= \frac{1}{2}d\left(\Gamma_\alpha^\beta \wedge R_\beta^\alpha - \frac{1}{3}\Gamma_\alpha^\beta \wedge \Gamma_\beta^\gamma \wedge \Gamma_\gamma^\alpha\right)$	$L_{\text{Pontr}} := dC_{\text{RR}} = \text{Tr}\{\varOmega^g \wedge \varOmega^g\} = \frac{1}{2} R_\alpha^\beta \wedge R_\beta^\alpha$ $= \frac{1}{2}d\Bigl(\Gamma_\alpha^\beta \wedge R_\beta^\alpha - \frac{1}{3}\Gamma_\alpha^\beta \wedge \Gamma_\beta^\gamma \wedge \Gamma_\gamma^\alpha\Bigr),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0824 (11.4.38)	259	■0.935 ● W +0	$\left.\left.\right.r_s\right\}\equiv\angle\left \vec{x}\right ^2\right\rangle_{\psi}^{\scriptscriptstyle{1/2}}:=\left(\frac{\int_{H^3}\left \vec{x}\right ^2\bar{\psi}^*\gamma\psi}{\int_{H^3}\bar{\psi}^*\gamma\psi}\right)^{\scriptscriptstyle{1/2}}.$	$r_s \equiv \langle \vec{x} ^2 \rangle_\psi^{1/2} := \left(\frac{\int_{H^3} \vec{x} ^2 \bar{\psi}^* \gamma \psi}{\int_{H^3} \bar{\psi}^* \gamma \psi} \right)^{1/2}.$	$r_s \equiv \langle \vec{x} ^2 \rangle_\psi^{1/2} := \left(\frac{\int_{H^3} \vec{x} ^2 \overline{\psi}^* \gamma \psi}{\int_{H^3} \overline{\psi}^* \gamma \psi} \right)^{1/2}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0288 (4.4.17)	099	■0.936 ● N +0	$\left[\omega^g , \Pi^L \right] + E_{[]} = D \tau_s - \Sigma_{[]} .$	$\left[\Omega^g , \Pi^L \right] + E_{[]} = D \tau_s - \Sigma_{[]} .$	$[\varOmega^g , \varPi^L] + E_{[]} = D \tau_s - \varSigma_{[]} .$

Conf. MathPix confidence:

■≥0.9

■0.5–0.9

■<0.5

— no value

Residual render vs scan (inkdrill):

● C component

● W weak

● S stable

● N noise

● K clean

● not measured

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0135 (3.4.6)	063	0.940 N -1	$\Pi^{\Omega}=\frac{2}{\alpha_g}\left(\frac{\partial L}{\partial \mathcal{F}}^*\Omega+\frac{\partial L}{\partial^*\mathcal{F}}\Omega\right)$	$\Pi^{\Omega}=\frac{2}{\alpha_g}\left(\frac{\partial L}{\partial \mathcal{F}}^*\Omega+\frac{\partial L}{\partial^*\mathcal{F}}\Omega\right)$	$\Pi^{\Omega}=\frac{2}{\alpha_g}\left(\frac{\partial L}{\partial \mathcal{F}}^*\Omega+\frac{\partial L}{\partial^*\mathcal{F}}\Omega\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0772 (11.2.1)	245	0.943 N +0	$L_{\mathrm{D}}=\frac{i}{2}\left(\bar{\psi}\gamma\wedge^*D\psi+(\overline{D\psi})\wedge^*\gamma\psi\right)-m\bar{\psi}\psi\eta.$	$L_{\mathrm{D}}=\frac{i}{2}\left(\bar{\psi}\gamma\wedge^*D\psi+(\overline{D\psi})\wedge^*\gamma\psi\right)-m\bar{\psi}\psi\eta.$	$L_{\mathrm{D}}=\frac{i}{2}\left(\bar{\psi}\gamma\wedge^*D\psi+(\overline{D\psi})\wedge^*\gamma\psi\right)-m\bar{\psi}\psi\eta.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0551 (8.4.5)	179	0.945 W -2	$\begin{aligned} L_{\mathrm{FP}}=& \, -\operatorname{Tr}\left\{\beta\wedge^*\bar{\Omega}^g-\bar{\chi}\wedge^*D\Psi^{(\pm)}-\bar{\chi}\wedge^*\left[C,\bar{\Omega}^{\pm g}\right]\right.\\ & +\bar{\eta}D^*\Psi+\bar{\Phi}D^*D\Phi+\bar{\Phi}[\Psi,*\Psi]+\mathbb{B}d^*\Gamma\\ & \left.-\bar{C}d^*\Psi+\bar{C}d^*(DC)+\frac{1}{2}\rho\beta\wedge^*\beta+\frac{1}{2}\mathbb{B}\wedge^*\mathbb{B}\right\}. \end{aligned}$	$L_{\mathrm{FP}}=-\operatorname{Tr}\left\{\beta\wedge^*\bar{\Omega}^g-\bar{\chi}\wedge^*D\Psi^{(\pm)}-\bar{\chi}\wedge^*\left[C,\bar{\Omega}^{\pm g}\right]+\bar{\eta}D^*\Psi+\bar{\Phi}D^*D\Phi+\bar{\Phi}[\Psi,*\Psi]+\mathbb{B}d^*\Gamma-\bar{C}d^*\Psi+\bar{C}d^*(DC)+\frac{1}{2}\rho\beta\wedge^*\beta+\frac{1}{2}\mathbb{B}\wedge^*\mathbb{B}\right\}.$	$L_{\mathrm{FP}}=-\operatorname{Tr}\left\{\beta\wedge^*\bar{\Omega}^g-\bar{\chi}\wedge^*D\Psi^{(\pm)}-\bar{\chi}\wedge^*\left[C,\bar{\Omega}^{\pm g}\right]+\bar{\eta}D^*\Psi+\bar{\Phi}D^*D\Phi+\bar{\Phi}[\Psi,*\Psi]+\mathbb{B}d^*\Gamma-\bar{C}d^*\Psi+\bar{C}d^*(DC)+\frac{1}{2}\rho\beta\wedge^*\beta+\frac{1}{2}\mathbb{B}\wedge^*\mathbb{B}\right\}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ1069 (16.2.8)	356	0.945 N +0	$V_{\mathrm{eff}}^{\iota}(\rho^*)=\frac{1\pm\rho^2}{\rho^2}\left[\iota(\iota+1)-\left(2-\frac{1}{\beta}\right)\rho^2\right],\,_{(\kappa<0)}^{\kappa>0}.$	$V_{\mathrm{eff}}^{\iota}(\rho^*)=\frac{1\pm\rho^2}{\rho^2}\left[\iota(\iota+1)-\left(2-\frac{1}{\beta}\right)\rho^2\right],\,_{(\kappa<0)}^{\kappa>0}.$	$V_{\mathrm{eff}}^{\iota}(\rho^*)=\frac{1\pm\rho^2}{\rho^2}\left[\iota(\iota+1)-\left(2-\frac{1}{\beta}\right)\rho^2\right],\,_{(\kappa<0)}^{\kappa>0}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0593 (9.1.11)	195	0.945 N +0	$R=\{ \}^*\wedge\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)=-\frac{6\kappa\gamma}{\ell^{*2}}=4\Lambda_{\mathrm{eff}}.$	$R=\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)=-\frac{6\kappa\gamma}{\ell^{*2}}=4\Lambda_{\mathrm{eff}}.$	$R=\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)=-\frac{6\kappa\gamma}{\ell^{*2}}=4\Lambda_{\mathrm{eff}}.$

Conf. MathPix confidence:

0.940

0.5–0.9

<0.5

— no value

Residual render vs scan (inkdrill):

C component

W weak

S stable

N noise

K clean

not measured

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0851 (12.3.4)	276	■ 0.946 ● N +1	$\begin{aligned} & \operatorname{Tr}\left(\gamma_5 K_2\right)=-d^* \mathscr{A} \\ & \operatorname{Tr}\left(\gamma_5 K_4\right)=\frac{1}{6}\left[\operatorname{Tr}\left(R^{\{1} \wedge R^{\{1}\right)}-\frac{1}{4} d \mathscr{A} \wedge d \mathscr{A}+d \mathscr{K}\right], \end{aligned}$	Tr (γ₅ K₂) = −d[*] ℳ Tr (γ₅ K₄) = $\frac{1}{6} \left[\operatorname{Tr} \left(R^{\{1} \wedge R^{\{1\}} \right) - \frac{1}{4} d \mathscr{A} \wedge d \mathscr{A} + d \mathscr{K} \right],$	$Tr(\gamma_5 K_2)=-d^* \mathscr{A},$ $Tr(\gamma_5 K_4)=\frac{1}{6}\left[Tr\left(R^{\{1} \wedge R^{\{1\}}\right)-\frac{1}{4} d \mathscr{A} \wedge d \mathscr{A}+d \mathscr{K}\right],$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0815 (11.4.29)	257	■ 0.946 ● N +2	$\begin{aligned} & H^{\prime} \simeq \frac{1+\omega}{\beta} F \\ & F^{\prime}+\frac{2}{\rho} F \simeq \frac{1-\omega}{\beta} H \end{aligned}$	$H' \simeq \frac{1+\omega}{\beta} F$ $F' + \frac{2}{\rho} F \simeq \frac{1-\omega}{\beta} H$	$H' \simeq \frac{1+\omega}{\beta} F,$ $F' + \frac{2}{\rho} F \simeq \frac{1-\omega}{\beta} H.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1021 (14.4.16)	328	■ 0.947 ● N -1	$S_{\mathrm{A}}:=4 \pi r_{+}^2=16 \pi M_{\mathrm{ir}}^2 \frac{\hbar^2}{M^{* 4} c^2},$	$S_{\mathrm{A}} := 4 \pi r_+^2 = 16 \pi M_{\mathrm{ir}}^2 \frac{\hbar^2}{M^{* 4} c^2},$	$S_{\mathrm{A}}:=4 \pi r_{+}^2=16 \pi M_{\mathrm{ir}}^2 \frac{\hbar^2}{M^{* 4} c^2},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0855 (12.4.2)	277	■ 0.947 ● N +0	$\operatorname{Tr}\left(\gamma_5 K_2\right)=-d^* \mathscr{A}$	$\mathcal{H}_\Lambda \Psi^{(1)}(\underline{\Gamma}) = 0$	$\mathcal{H}_\Lambda \Psi^{(\pm)}(\underline{\Gamma}) = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0452 (7.2.5)	151	■ 0.948 ● W +0	$C_{\cdot \cdot c d}^{\{a b\}}=R_{\cdot \cdot c d}^{\{a b\}}-2 R_{\cdot[c}^{\{a} \delta_{d\}}^{\{b\}}+\frac{R}{6} \delta_{[c}^a \delta_{d]}^b.$	$C_{\cdot \cdot c d}^{ab} = R_{\cdot \cdot c d}^{ab} - 2 R_{\cdot [c}^{[a} \delta_{d]}^{b]} + \frac{R}{6} \delta_{[c}^a \delta_{d]}^b.$	$C_{\cdot \cdot c d}^{ab}=R_{\cdot \cdot c d}^{ab}-2 R_{\cdot[c}^{[a} \delta_{d]}^{\{b\}}+\frac{R}{6} \delta_{[c}^a \delta_{d]}^b.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0061 (2.8.2)	046	■ 0.949 ● N +1	$\left\{ \right\}^*\!\!\!\left(\gamma_k\right)=\frac{(-1)^k}{(2\pi i)^k k!}\operatorname{Tr}(\Omega\wedge\cdots\wedge\Omega).$	$\pi^*(\gamma_k)=\frac{(-1)^k}{(2\pi i)^k k!}\operatorname{Tr}(\Omega\wedge\cdots\wedge\Omega).$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0756 (11.1.23)	242	■ 0.950 ● W +1	$\begin{aligned} S(p) &= \exp \imath \theta_{\alpha\beta}(m) \Lambda^{-1} (L^{\alpha\beta}) = \exp \imath \theta_{\alpha\beta}(m) \sigma^{\alpha\beta} \\ &= \exp \left(-\frac{1}{4} \theta_{\alpha\beta}(m) [\gamma^\alpha, \gamma^\beta] \right). \end{aligned}$	$S(p) = \exp \imath \theta_{\alpha\beta}(m) \Lambda^{-1} (L^{\alpha\beta}) = \exp \imath \theta_{\alpha\beta}(m) \sigma^{\alpha\beta} = \exp \left(-\frac{1}{4} \theta_{\alpha\beta}(m) [\gamma^\alpha, \gamma^\beta] \right).$	$S(p) = \exp \imath \theta_{\alpha\beta}(m) \Lambda^{-1} (L^{\alpha\beta}) = \exp \imath \theta_{\alpha\beta}(m) \sigma^{\alpha\beta} = \exp \left(-\frac{1}{4} \theta_{\alpha\beta}(m) [\gamma^\alpha, \gamma^\beta] \right).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0939 (14.1.20)	305	■ 0.950 ● N +0	$\xi \wedge \xi = 0 \quad (= 2 \vec{E} \cdot \vec{B} \text{ in 3-dimensional vector notation })$	$\xi \wedge \xi = 0 \quad (= 2 \vec{E} \cdot \vec{B} \text{ in 3-dimensional vector notation })$	$\xi \wedge \xi = 0 \quad (= 2 \vec{E} \cdot \vec{B} \text{ in 3-dimensional vector notation })$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0568 (8.6.5)	183	■ 0.950 ● C -4	$L_{\theta} := \theta_T dC_{TT} + \theta_T^* dC_{TT^*} + \theta_L dC_{RR} + \theta_L^* (-1)^{\operatorname{sig}+1} dC_{RR^*} = dC_{\theta}.$	$L_{\theta} := \theta_T dC_{TT} + \theta_T^* dC_{TT^*} + \theta_L dC_{RR} + \theta_L^* (-1)^{\operatorname{sig}+1} dC_{RR^*} = dC_{\theta}.$	$L_{\theta} := \theta_T dC_{TT} + \theta_T^* dC_{TT^*} + \theta_L dC_{RR} + \theta_L^* (-1)^{\operatorname{sig}+1} dC_{RR^*} = dC_{\theta}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0234 (4.2.27)	088	■ 0.950 ● N +0	$\widetilde{\widetilde{D}}\widetilde{\widetilde{\varphi}} = D\widetilde{\widetilde{\varphi}} = 0$	$\widetilde{\widetilde{D}}\widetilde{\widetilde{\varphi}} = D\widetilde{\widetilde{\varphi}} = 0$	$\widetilde{\widetilde{D}}\widetilde{\widetilde{\varphi}} = D\widetilde{\widetilde{\varphi}} = 0$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0302 (4.5.9)	102	■0.950 ●W +0	\begin{aligned} \widehat{D} \Sigma_{\alpha} &\cong (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \Delta^{\beta} \gamma \\ &\quad - \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta \gamma}) \wedge \sigma^{\beta \gamma} \end{aligned}	$\widehat{D} \Sigma_{\alpha} \cong (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \Delta^{\beta} \gamma - \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta \gamma}) \wedge \sigma^{\beta \gamma}$	$\widehat{D} \Sigma_{\alpha} \cong (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \Delta^{\beta} \gamma - \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta \gamma}) \wedge \sigma^{\beta \gamma}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0151 (3.4.22)	066	■0.951 ●C -4	L_{\text {Mills }}\left(\mathrm{F},\left\{ \right\} ^{*}\mathrm{F}\right)=\mathrm{F}\left[1-\left(1+\frac{1}{b^2}\right)\mathrm{F}^2\right]\mathrm{F}^2\right)^{-1}\right]	$L_{\mathrm{Mills}}\left(\mathcal{F},*\mathcal{F}\right)=\mathcal{F}\left[1-\left(1+\frac{1}{b^2}\mathcal{F}^2\right)^{-1}\right]$	$L_{\mathrm{Mills}}(\mathcal{F},*\mathcal{F})=\mathcal{F}\left[1-\left(1+\frac{1}{b^2}\mathcal{F}^2\right)^{-1}\right]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0560 (8.5.3)	181	■0.953 ●N +0	\stackrel{\pm}{\Omega}^g=i\frac{\theta_{\mathrm{T}}^*}{4\ell^2}\sigma	$\stackrel{\pm}{\Omega}^g=i\frac{\theta_{\mathrm{T}}^*}{4\ell^2}\sigma$	$\stackrel{\pm}{\Omega}^g=i\frac{\theta_{\mathrm{T}}^*}{4\ell^2}\sigma$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ1106 (B.18)	370	■0.953 ●N +0	\delta \alpha ^{(p)}:=(-1)^{p\,n+n+1+s}\,d^*\alpha ^{(p)}.	$\delta \alpha ^{(p)}:=(-1)^{pn+n+1+s}d^*\alpha ^{(p)}.$	$\delta \alpha ^{(p)}:=(-1)^{pn+n+1+s}d^*\alpha ^{(p)}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0786 (11.3.5)	248	■0.954 ●W +1	\begin{aligned} L_{\mathrm{NL}} &\wedge \tau_s = 3 \ell^{*2} \bar{\psi}^* \gamma \psi \wedge (\bar{\psi}^* \gamma \psi) \\ &= 3 \ell^{*2} \bar{\psi} \gamma^{n+1} \gamma \psi \wedge (\bar{\psi} \gamma^{n+1} \gamma \psi). \end{aligned}	$\begin{aligned} L_{\mathrm{NL}} &= -3 \ell^{*2} \tau_s \wedge \tau_s = 3 \ell^{*2} \bar{\psi}^* \gamma \psi \wedge (\bar{\psi}^* \gamma \psi) \\ &= 3 \ell^{*2} \bar{\psi} \gamma^{n+1} \gamma \psi \wedge (\bar{\psi} \gamma^{n+1} \gamma \psi). \end{aligned}$	$\begin{aligned} L_{\mathrm{NL}} &= -3 \ell^{*2} \tau_s \wedge \tau_s = 3 \ell^{*2} \bar{\psi}^* \gamma \psi \wedge (\bar{\psi}^* \gamma \psi) \\ &= 3 \ell^{*2} \bar{\psi} \gamma^{n+1} \gamma \psi \wedge (\bar{\psi} \gamma^{n+1} \gamma \psi). \end{aligned}$
Conf. MathPix confidence: ■≥0.9 ■0.5-0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E08727 (10.10.7)	231	■0.955 ●W +2	$\begin{aligned} & \begin{aligned} & i \gamma^\alpha \Psi \wedge \frac{\delta L}{\delta \vartheta^\alpha} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} - 2D \frac{\delta L}{\delta \bar{\Psi}} \\ & \cong i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\text{TL}}}{\ell} R_\alpha^* + \frac{\theta_{\text{T}}}{\ell^2} T_\alpha + \Sigma_\alpha \right) \\ & + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - D \left(i D \Psi + \frac{i}{2} m \gamma \wedge \Psi \right) \\ & = i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\text{TL}}}{\ell} R_\alpha^* + \frac{\theta_{\text{T}}}{\ell^2} T_\alpha \right) + \gamma^\alpha \Psi \wedge \left(\frac{1}{4} m \bar{\Psi} \gamma_\alpha \Psi \right) \\ & + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - i R_\alpha^* \gamma^\alpha \Psi - \frac{i}{2} m T_\alpha \gamma^\alpha \Psi \\ & + \frac{i}{2} m \gamma \wedge D \Psi. \end{aligned} \end{aligned}$	$\begin{aligned} & i \gamma^\alpha \Psi \wedge \frac{\delta L}{\delta \vartheta^\alpha} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} - 2D \frac{\delta L}{\delta \bar{\Psi}} \\ & \cong i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\text{TL}}}{\ell} R_\alpha^* + \frac{\theta_{\text{T}}}{\ell^2} T_\alpha + \Sigma_\alpha \right) \\ & + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - D \left(i D \Psi + \frac{i}{2} m \gamma \wedge \Psi \right) \\ & = i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\text{TL}}}{\ell} R_\alpha^* + \frac{\theta_{\text{T}}}{\ell^2} T_\alpha \right) + \gamma^\alpha \Psi \wedge \left(\frac{1}{4} m \bar{\Psi} \gamma_\alpha \Psi \right) \\ & + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - i R_\alpha^* \gamma^\alpha \Psi - \frac{i}{2} m T_\alpha \gamma^\alpha \Psi \\ & + \frac{i}{2} m \gamma \wedge D \Psi. \end{aligned}$	$\begin{aligned} & i \gamma^\alpha \Psi \wedge \frac{\delta L}{\delta \vartheta^\alpha} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} - 2D \frac{\delta L}{\delta \bar{\Psi}} \\ & \cong i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\text{TL}}}{\ell} R_\alpha^* + \frac{\theta_{\text{T}}}{\ell^2} T_\alpha + \Sigma_\alpha \right) \\ & + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - D \left(i D \Psi + \frac{i}{2} m \gamma \wedge \Psi \right) \\ & = i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\text{TL}}}{\ell} R_\alpha^* + \frac{\theta_{\text{T}}}{\ell^2} T_\alpha \right) + \gamma^\alpha \Psi \wedge \left(\frac{1}{4} m \bar{\Psi} \gamma_\alpha \Psi \right) \\ & + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - i R_\alpha^* \gamma^\alpha \Psi - \frac{i}{2} m T_\alpha \gamma^\alpha \Psi \\ & + \frac{i}{2} m \gamma \wedge D \Psi. \end{aligned} \tag{10.1}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E08768 (11.1.35)	244	■0.956 ●N +0	$\bar{\psi} \longrightarrow S^{-1} \bar{\psi} = \bar{\psi} S,$	$\bar{\psi} \longrightarrow S^{-1} \bar{\psi} = \bar{\psi} S,$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E08973 (14.2.14)	315	■0.956 ●C +4	$\begin{aligned} & \hat{\Theta} := \hat{D} \hat{\vartheta} = \hat{d} \hat{\vartheta} - [\hat{\omega}, \hat{\vartheta}] \\ & = \left[\begin{array}{c c} \Theta - \hat{\ell} \omega \wedge \alpha^{\text{T}} & \hat{\ell} \left(d\omega - \omega^g \wedge \omega - \omega \wedge \overset{\circ}{\omega} \right) + \vartheta \wedge \alpha + \alpha \wedge \overset{\circ}{\vartheta} \\ \hline -\alpha^{\text{T}} \wedge \vartheta - \vartheta \wedge \alpha^{\text{T}} & \overset{\circ}{\Theta} - \hat{\ell} \alpha^{\text{T}} \wedge \omega \end{array} \right]. \end{aligned}$	$\begin{aligned} & \hat{\Theta} := \hat{D} \hat{\vartheta} = \hat{d} \hat{\vartheta} - [\hat{\omega}, \hat{\vartheta}] \\ & = \left[\begin{array}{c c} \Theta - \hat{\ell} \omega \wedge \alpha^{\text{T}} & \hat{\ell} \left(d\omega - \omega^g \wedge \omega - \omega \wedge \overset{\circ}{\omega} \right) + \vartheta \wedge \alpha + \alpha \wedge \overset{\circ}{\vartheta} \\ \hline -\alpha^{\text{T}} \wedge \vartheta - \vartheta \wedge \alpha^{\text{T}} & \overset{\circ}{\Theta} - \hat{\ell} \alpha^{\text{T}} \wedge \omega \end{array} \right]. \end{aligned}$	$\begin{aligned} & \hat{\Theta} := \hat{D} \hat{\vartheta} = \hat{d} \hat{\vartheta} - [\hat{\omega}, \hat{\vartheta}] \\ & = \left[\begin{array}{c c} \Theta - \hat{\ell} \omega \wedge \alpha^{\text{T}} & \hat{\ell} \left(d\omega - \omega^g \wedge \omega - \omega \wedge \overset{\circ}{\omega} \right) + \vartheta \wedge \alpha + \alpha \wedge \overset{\circ}{\vartheta} \\ \hline -\alpha^{\text{T}} \wedge \vartheta - \vartheta \wedge \alpha^{\text{T}} & \overset{\circ}{\Theta} - \hat{\ell} \alpha^{\text{T}} \wedge \omega \end{array} \right]. \end{aligned}$
Conf. MathPix confidence: ■ ■ ■ — ≥0.90.5-0.9<0.5no value Residual render vs scan (inkdrill): ● ● ● ● ● ● C componentW weakS stableN noiseK cleannot measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0761 (11.1.28)	243	■0.957 ● N +0	$\frac{1}{N} \operatorname{Tr}(\gamma \otimes_s \gamma) = ds^2 = g_{ij}(m) dx^i \otimes_s dx^j.$		$\frac{1}{N} \operatorname{Tr}(\gamma \otimes_s \gamma) = ds^2 = g_{ij}(m) dx^i \otimes_s dx^j.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0486 (7.7.2)	160	■0.957 ● N +0	$\begin{aligned} \vartheta^\alpha &\rightarrow \widetilde{\vartheta}^\alpha := \vartheta^\alpha + \ell^2 e_\beta \rfloor^* H^{\alpha\beta}, \\ \Gamma_\alpha{}^\beta &\rightarrow \widetilde{\Gamma}_\alpha{}^\beta := \Gamma_\alpha{}^\beta + \ell^2 e_\alpha \rfloor^* H^\beta. \end{aligned}$	$\vartheta^\alpha \rightarrow \widetilde{\vartheta}^\alpha := \vartheta^\alpha + \ell^2 e_\beta \rfloor^* H^{\alpha\beta},$ $\Gamma_\alpha{}^\beta \rightarrow \widetilde{\Gamma}_\alpha{}^\beta := \Gamma_\alpha{}^\beta + \ell^2 e_\alpha \rfloor^* H^\beta.$	$\vartheta^\alpha \rightarrow \widetilde{\vartheta}^\alpha := \vartheta^\alpha + \ell^2 e_\beta \rfloor^* H^{\alpha\beta},$ $\Gamma_\alpha{}^\beta \rightarrow \widetilde{\Gamma}_\alpha{}^\beta := \Gamma_\alpha{}^\beta + \ell^2 e_\alpha \rfloor^* H^\beta.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1104 (B.16)	370	■0.957 ● N +0	$\begin{aligned} \left\langle \alpha^{(p)}, \beta^{(p)} \right\rangle &:= \alpha^{(p)} \wedge^* \beta^{(p)} = \beta^{(p)} \wedge^* \alpha^{(p)} \\ &= (-1)^{s*} \left(\alpha^{(p)} \wedge^* \beta^{(p)} \right) \eta \\ &= \frac{(-1)^s}{p!} A^{\alpha_1 \cdots \alpha_p} B_{\alpha_1 \cdots \alpha_p} d\mu(M) \in C^\infty(\wedge^n T^*(M)) \end{aligned}$	$\left\langle \alpha^{(p)}, \beta^{(p)} \right\rangle := \alpha^{(p)} \wedge^* \beta^{(p)} = \beta^{(p)} \wedge^* \alpha^{(p)}$ $= (-1)^{s*} \left(\alpha^{(p)} \wedge^* \beta^{(p)} \right) \eta$ $= \frac{(-1)^s}{p!} A^{\alpha_1 \cdots \alpha_p} B_{\alpha_1 \cdots \alpha_p} d\mu(M) \in C^\infty(\wedge^n T^*(M))$	$\langle \alpha^{(p)}, \beta^{(p)} \rangle := \alpha^{(p)} \wedge^* \beta^{(p)} = \beta^{(p)} \wedge^* \alpha^{(p)}$ $= (-1)^{s*} (\alpha^{(p)} \wedge^* \beta^{(p)}) \eta$ $= \frac{(-1)^s}{p!} A^{\alpha_1 \cdots \alpha_p} B_{\alpha_1 \cdots \alpha_p} d\mu(M) \in C^\infty(\wedge^n T^*(M))$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0537 (8.2.4)	176	■0.958 ● W +1	$\begin{array}{ll} s \, \bar{C} = \mathbb{B}, & s \, \mathbb{B} = 0, \\ s \, \bar{\chi} = \beta, & s \, \beta = 0, \\ s \, \bar{\Phi} = \bar{\eta}, & s \, \bar{\eta} = 0. \end{array}$	$s \bar{C} = \mathbb{B}, \quad s \mathbb{B} = 0,$ $s \bar{\chi} = \beta, \quad s \beta = 0,$ $s \bar{\Phi} = \bar{\eta}, \quad s \bar{\eta} = 0.$	$s \bar{C} = \mathbb{B}, \quad s \mathbb{B} = 0,$ $s \bar{\chi} = \beta, \quad s \beta = 0,$ $s \bar{\Phi} = \bar{\eta}, \quad s \bar{\eta} = 0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0269 (4.3.34)	095	■0.958 ● N +0	$R_{\{i \, j\}} := R_{\{\alpha \, i \, j\}}^{\cdots \alpha}$	$R_{ij} := R_{\alpha ij}^{\cdots \alpha}$	$R_{ij} := R_{\alpha ij}^{\cdots \alpha}$
Conf. MathPix confidence: ■≥ 0.9 ■0.5–0.9 ■< 0.5 — no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0926 (14.1.7)	303	■0.960 ● N +0	$D\left(\vartheta\wedge^*\Omega^g\right)\equiv\Theta\wedge^*\Omega^g\text{ .}$	$D\left(\vartheta\wedge^*\Omega^g\right)\equiv\Theta\wedge^*\Omega^g.$	$D(\vartheta\wedge^*\Omega^g)\equiv\Theta\wedge^*\Omega^g.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0248 (4.3.13)	092	■0.961 ● N +0	$D\,\phi^{\left(\circ\right)}=\vartheta\wedge^{\alpha}\nabla_{\alpha}\phi^{\left(\circ\right)}\text{ .}$	$D\phi^{(\circ)}=\vartheta^\alpha\nabla_\alpha\phi^{(\circ)}.$	$D\,\phi^{(\circ)}=\vartheta^\alpha\nabla_\alpha\phi^{(\circ)}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0282 (4.4.11)	098	■0.961 ● N +1	$\begin{aligned} d\,\Theta&\,=\,d\,d\,\vartheta-d\left[\omega^L\{\mathrm{L}\},\,\vartheta\right]\,\,\&=\,-\left[d\,\omega^{\mathrm{L}}\{\mathrm{L}\},\,\vartheta\right]+\left[\omega^L\{\mathrm{L}\},\,d\vartheta\right] \\ &\,=\,-\left[\Omega^L,\vartheta\right]-\left[\omega^L\wedge\omega^L,\vartheta\right]+\left[\omega^L,\Theta\right]+\left[\omega^L,\left[\omega^L,\vartheta\right]\right] \\ &\,=\,\left[\omega^L,\Theta\right]+\left[\vartheta,\Omega^L\right] \end{aligned}$	$\begin{aligned} d\Theta&=dd\vartheta-d\left[\omega^L,\vartheta\right] \\ &=-\left[d\omega^L,\vartheta\right]+\left[\omega^L,d\vartheta\right] \\ &=-\left[\Omega^L,\vartheta\right]-\left[\omega^L\wedge\omega^L,\vartheta\right]+\left[\omega^L,\Theta\right]+\left[\omega^L,\left[\omega^L,\vartheta\right]\right] \\ &=\left[\omega^L,\Theta\right]+\left[\vartheta,\Omega^L\right] \end{aligned}$	$\begin{aligned} d\Theta&=dd\vartheta-d[\omega^L,\vartheta] \\ &=-[d\omega^L,\vartheta]+[\omega^L,d\vartheta] \\ &=-[\Omega^L,\vartheta]-[\omega^L\wedge\omega^L,\vartheta]+[\omega^L,\Theta]+[\omega^L,[\omega^L,\vartheta]] \\ &=[\omega^L,\Theta]+[\vartheta,\Omega^L]. \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0010 (2.2.5)	034	■0.962 ● N +0	$T_{\{p\}^{\{q\}}(M):=\bigcup_{m\in M}T_m^{\{q\}}(M)\otimes^pT_m^*(M)\otimes^qT_m(M)\text{ .}$	$T_p^q(M):=\bigcup_{m\in M}\otimes^pT_m^*(M)\otimes^qT_m(M).$	$T_p^q(M):=\bigcup_{m\in M}\otimes^pT_m^*(M)\otimes^qT_m(M).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0340 (5.3.17)	115	■0.962 ● C +4	$\begin{aligned} G_{\alpha}&\,\&:=\frac{1}{2}\,R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma} \\ &\,=\,G_{\alpha}^{\{\}}+\frac{(-1)^s}{12}\left(e_{\alpha}\rfloor\mathscr{A}\wedge^*\mathscr{A}-\frac{1}{3}\mathscr{A}\wedge e_{\alpha}\rfloor^*\mathscr{A}\right)+\frac{(-1)^s}{6}\vartheta_{\alpha}\wedge d\mathscr{A} \end{aligned}$	$\begin{aligned} G_{\alpha}&:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma} \\ &=G_{\alpha}^{\{\}}+\frac{(-1)^s}{12}\left(e_{\alpha}\rfloor\mathscr{A}\wedge^*\mathscr{A}-\frac{1}{3}\mathscr{A}\wedge e_{\alpha}\rfloor^*\mathscr{A}\right)+\frac{(-1)^s}{6}\vartheta_{\alpha}\wedge d\mathscr{A} \end{aligned}$	$G_{\alpha}:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma}\tag{5.3}$ $=G_{\alpha}^{\{\}}+\frac{(-1)^s}{12}\left(e_{\alpha}\rfloor\mathscr{A}\wedge^*\mathscr{A}-\frac{1}{3}\mathscr{A}\wedge e_{\alpha}\rfloor^*\mathscr{A}\right)+\frac{(-1)^s}{6}\vartheta_{\alpha}\wedge d\mathscr{A}$
Conf. MathPix confidence: ■≥0.9 ■0.5–0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0424 (6.7.6)	140	■ 0.962 ● N +0	\begin{aligned} \frac{1}{2} \kappa \, T^{\gamma} \wedge \eta_{\alpha\beta\gamma} - \frac{D\theta_{\mathrm{T}}}{2\ell^2} \wedge \vartheta_{\alpha} \wedge \vartheta_{\beta} + D\theta_{\mathrm{L}} \wedge R_{\alpha\beta} &= \vartheta_{[\alpha} \wedge \mu_{\beta]} \\ &= \frac{1}{4} \vartheta_{\alpha} \wedge \vartheta_{\beta} \wedge {}^* j_5. \end{aligned}	$\frac{1}{2\kappa} T^{\gamma} \wedge \eta_{\alpha\beta\gamma} - \frac{D\theta_{\mathrm{T}}}{2\ell^2} \wedge \vartheta_{\alpha} \wedge \vartheta_{\beta} + D\theta_{\mathrm{L}} \wedge R_{\alpha\beta} = \vartheta_{[\alpha} \wedge \mu_{\beta]} = \frac{1}{4} \vartheta_{\alpha} \wedge \vartheta_{\beta} \wedge {}^* j_5.$	$\frac{1}{2\kappa} T^{\gamma} \wedge \eta_{\alpha\beta\gamma} - \frac{D\theta_{\mathrm{T}}}{2\ell^2} \wedge \vartheta_{\alpha} \wedge \vartheta_{\beta} + D\theta_{\mathrm{L}} \wedge R_{\alpha\beta} = \vartheta_{[\alpha} \wedge \mu_{\beta]} = \frac{1}{4} \vartheta_{\alpha} \wedge \vartheta_{\beta} \wedge {}^* j_5.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0809 (11.4.21)	255	■ 0.962 ● W +0	i \gamma^0 \vec{\gamma} \cdot \vec{\partial} \left[\begin{array}{cc} i H(\rho) & \chi_{\kappa}^m \\ -F(\rho) & \chi_{-\kappa}^m \end{array} \right] = \frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } \left(\partial_r - \frac{\vec{\sigma} \cdot \vec{L}}{r} \right) \left[\begin{array}{cc} i F(\rho) & \chi_{-\kappa}^m \\ H(\rho) & \chi_{\kappa}^m \end{array} \right].	$i\gamma^0\vec{\gamma}\cdot\vec{\partial}\left[\begin{array}{cc}iH(\rho)&\chi_{\kappa}^m\\-F(\rho)&\chi_{-\kappa}^m\end{array}\right]=\frac{\vec{\sigma}\cdot\vec{x}}{ \vec{x} }\left(\partial_r-\frac{\vec{\sigma}\cdot\vec{L}}{r}\right)\left[\begin{array}{cc}iF(\rho)&\chi_{-\kappa}^m\\H(\rho)&\chi_{\kappa}^m\end{array}\right].$	$i\gamma^0\vec{\gamma}\cdot\vec{\partial}\left[\begin{array}{cc}iH(\rho)&\chi_{\kappa}^m\\-F(\rho)&\chi_{-\kappa}^m\end{array}\right]=\frac{\vec{\sigma}\cdot\vec{x}}{ \vec{x} }\left(\partial_r-\frac{\vec{\sigma}\cdot\vec{L}}{r}\right)\left[\begin{array}{cc}iF(\rho)&\chi_{-\kappa}^m\\H(\rho)&\chi_{\kappa}^m\end{array}\right].$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0475 (7.4.3)	156	■ 0.963 ● N +1	{ }^{\ast} R_{\alpha \beta}^{\wedge (\star)} = - C_{\alpha \beta} + \frac{1}{2} R \, i^{\tau}{}_{\alpha \beta} + \frac{R}{12} \vartheta_{\alpha} \wedge \vartheta_{\beta}.	${}^{\ast} R_{\alpha \beta}^{(\star)} = - C_{\alpha \beta} + \frac{1}{2} R i^{\tau}{}_{\alpha \beta} + \frac{R}{12} \vartheta_{\alpha} \wedge \vartheta_{\beta}.$	${}^{\ast} R_{\alpha \beta}^{(\star)} = - C_{\alpha \beta} + \frac{1}{2} R i^{\tau}{}_{\alpha \beta} + \frac{R}{12} \vartheta_{\alpha} \wedge \vartheta_{\beta}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0485 (7.7.1)	159	■ 0.963 ● W +1	g_{\{i\}j} \rightarrow \widetilde{g}_{\{i\}j} = g_{\{i\}j} + a \operatorname{Ric}_{ij} + b g_{\{i\}j} \operatorname{Ric}_{\{k\}^{\{k\}}}	$g_{ij} \rightarrow \widetilde{g}_{ij} = g_{ij} + a \operatorname{Ric}_{ij} + b g_{ij} \operatorname{Ric}_k^k$	$g_{ij} \rightarrow \widetilde{g}_{ij} = g_{ij} + a \operatorname{Ric}_{ij} + b g_{ij} \operatorname{Ric}_k^k$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ1143 (C.11)	376	■ 0.963 ● N +0	(A, B):=\operatorname{Tr}(\operatorname{Ad} A \operatorname{Ad} B) = c_{\iota k}^i c_{s i}^k A^{\iota} B^s = c_{\iota k}^i c_{s i}^k A^{\iota} B^s	$(A, B) := \operatorname{Tr}(\operatorname{Ad} A \operatorname{Ad} B) = c_{\iota k}^i c_{s i}^k A^{\iota} B^s = c_{\iota k}^i c_{s i}^k A^{\iota} B^s$	$(A, B) := \operatorname{Tr}(\operatorname{Ad} A \operatorname{Ad} B) = c_{\iota k}^i c_{s i}^k A^{\iota} B^s = c_{\iota k}^i c_{s i}^k A^{\iota} B^s$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ1122 (B.34)	372	■ 0.963 ● K +0	$H^{\{p\}\left(S^{\{n\}}\right)=0\quad\text{for}\quad 1\leq p\leq n-1}.$	$H^p(S^n)=0\quad\text{for}\quad 1\leq p\leq n-1.$	$H^p(S^n)=0\quad\text{for}\quad 1\leq p\leq n-1.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0791 (11.4.3)	252	■ 0.964 ● N -1	$\left\{i\gamma^k\nabla_k^{(t)}+\frac{3}{8}\ell^2\left(\bar\psi\psi-a\bar\psi\gamma^5\psi\gamma^5\right)-m\right\}\psi=0$	$\left\{i\gamma^k\nabla_k^{(t)}+\frac{3\varepsilon}{8}\ell^2\left(\bar\psi\psi-a\bar\psi\gamma^5\psi\gamma^5\right)-m\right\}\psi=0$	$\left\{i\gamma^k\nabla_k^{(t)}+\frac{3\varepsilon}{8}\ell^2(\bar\psi\psi-a\bar\psi\gamma^5\psi\gamma^5)-m\right\}\psi=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0575 (8.6.13)	185	■ 0.964 ● N -1	$\left.\left.\left.R:=e_{\beta}\right\rfloor e_{\alpha}\right\rfloor R^{\alpha\beta}=\frac{6\left(\theta_{\mathrm{T}}^{\ast}-a_0\right)}{\ell^2\left(\theta_{\mathrm{L}}^{\ast}-b_6\right)}\right\rfloor R^{\alpha\beta}=\frac{6\left(\theta_{\mathrm{T}}^{\ast}-a_0\right)}{\ell^2\left(\theta_{\mathrm{L}}^{\ast}-b_6\right)}$	$R:=e_{\beta}\rfloor e_{\alpha}\rfloor R^{\alpha\beta}=\frac{6\left(\theta_{\mathrm{T}}^{\ast}-a_0\right)}{\ell^2\left(\theta_{\mathrm{L}}^{\ast}-b_6\right)}$	$R:=e_{\beta}\rfloor e_{\alpha}\rfloor R^{\alpha\beta}=\frac{6(\theta_{\mathrm{T}}^{\ast}-a_0)}{\ell^2(\theta_{\mathrm{L}}^{\ast}-b_6)}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0996 (14.3.9)	321	■ 0.965 ● W -2	$i\widehat{D}\widehat{\psi}:=i\widehat{\gamma}\wedge^*\widehat{D}\widehat{\psi}=i\gamma\wedge^*D\widehat{\psi}+i\mathcal{D}^{(int)}\widehat{\psi}=0,$	$i\widehat{D}\widehat{\psi}:=i\widehat{\gamma}\wedge^*\widehat{D}\widehat{\psi}=i\gamma\wedge^*D\widehat{\psi}+i\mathcal{D}^{(int)}\widehat{\psi}=0,$	$i\widehat{\mathcal{D}}\widehat{\psi}:=i\widehat{\gamma}\wedge^*\widehat{D}\widehat{\psi}=i\gamma\wedge^*D\widehat{\psi}+i\mathcal{D}^{(int)}\widehat{\psi}=0,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0131 (3.4.2)	062	■ 0.966 ● N -1	$\left\{ \right\} ^{\{*\}}\mathrm{Tr}\left\{ \right\} ^{\{*\}}\left(\Omega \wedge \Omega \right) =\frac{1}{4\alpha _g}F_{\alpha \beta }^{j*}F^{\alpha \beta }_{\,j}.$	${}^*\mathcal{F}:=\frac{1}{\alpha_g}\mathrm{Tr}\left\{{}^*(\Omega\wedge\Omega)\right\}=\frac{1}{4\alpha_g}F_{\alpha\beta}{}^{j*}F^{\alpha\beta}{}_j.$	${}^*\mathcal{F}:=\frac{1}{\alpha_g}\mathrm{Tr}\left\{{}^*(\Omega\wedge\Omega)\right\}=\frac{1}{4\alpha_g}F_{\alpha\beta}{}^j{}^*F^{\alpha\beta}{}_j.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1089 (A.16)	366	■0.967 ● N +0	$\begin{aligned} & \left(\widehat{\gamma}^A \right)^{\mathrm{T}} := \left(\widehat{\gamma}^{\mu}, \widehat{\gamma}^{2p+1}, \widehat{\gamma}^{2p+2} \right), \\ & \widehat{\gamma}^{\mu} := \gamma^{\mu} \otimes \left[\bigotimes_{i=1}^{\left[\frac{n-2}{2} \right]} \sigma^1 \right], \\ & \widehat{\gamma}^{2p+1} := i \mathbb{1} \otimes \left(\bigotimes_{i=1}^{p-2} \sigma^0 \right) \otimes \sigma^3 \otimes \left(\bigotimes_{j=1}^{l-p+1} \sigma^1 \right), \\ & \widehat{\gamma}^{2p+2} := i \mathbb{1} \otimes \left(\bigotimes_{i=1}^{p-2} \sigma^0 \right) \otimes \sigma^2 \otimes \left(\bigotimes_{j=1}^{l-p+2} \sigma^1 \right), \end{aligned}$	$\begin{aligned} (\widehat{\gamma}^A)^{\mathrm{T}} &:= (\widehat{\gamma}^{\mu}, \widehat{\gamma}^{2p+1}, \widehat{\gamma}^{2p+2}), \\ \widehat{\gamma}^{\mu} &:= \gamma^{\mu} \otimes \left[\bigotimes_{i=1}^{\left[\frac{n-2}{2} \right]} \sigma^1 \right], \\ \widehat{\gamma}^{2p+1} &:= i \mathbb{1} \otimes \left(\bigotimes_{i=1}^{p-2} \sigma^0 \right) \otimes \sigma^3 \otimes \left(\bigotimes_{j=1}^{l-p+1} \sigma^1 \right), \\ \widehat{\gamma}^{2p+2} &:= i \mathbb{1} \otimes \left(\bigotimes_{i=1}^{p-2} \sigma^0 \right) \otimes \sigma^2 \otimes \left(\bigotimes_{j=1}^{l-p+2} \sigma^1 \right), \end{aligned}$	$(\widehat{\gamma}^A)^{\mathrm{T}} := (\widehat{\gamma}^{\mu}, \widehat{\gamma}^{2p+1}, \widehat{\gamma}^{2p+2}),$ $\widehat{\gamma}^{\mu} := \gamma^{\mu} \otimes \left[\bigotimes_{i=1}^{\left[\frac{n-2}{2} \right]} \sigma^1 \right],$ $\widehat{\gamma}^{2p+1} := i \mathbb{1} \otimes \left(\bigotimes_{i=1}^{p-2} \sigma^0 \right) \otimes \sigma^3 \otimes \left(\bigotimes_{j=1}^{l-p+1} \sigma^1 \right),$ $\widehat{\gamma}^{2p+2} := i \mathbb{1} \otimes \left(\bigotimes_{i=1}^{p-2} \sigma^0 \right) \otimes \sigma^2 \otimes \left(\bigotimes_{j=1}^{l-p+2} \sigma^1 \right),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1090 (A.17)	367	■0.967 ● N +0	$\stackrel{*}{\alpha} \left(e_{\alpha_1}, \dots, e_{\alpha_p} \right) := \alpha \left(\sigma_* e_{\alpha_1}, \dots, \sigma_* e_{\alpha_p} \right).$	$\stackrel{*}{\sigma} \alpha \left(e_{\alpha_1}, \dots, e_{\alpha_p} \right) := \alpha \left(\sigma_* e_{\alpha_1}, \dots, \sigma_* e_{\alpha_p} \right).$	$\stackrel{*}{\sigma} \alpha (e_{\alpha_1}, \dots, e_{\alpha_p}) := \alpha (\sigma_* e_{\alpha_1}, \dots, \sigma_* e_{\alpha_p}).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0592 (9.1.10)	195	■0.968 ● N +0	$\stackrel{+}{\Omega} = \frac{\gamma \kappa}{\ell^{*2}} \vartheta \wedge \vartheta.$	$\stackrel{+}{\Omega} = \frac{\gamma \kappa}{\ell^{*2}} \vartheta \wedge \vartheta.$	$\stackrel{+}{\Omega}^g = \frac{\gamma \kappa}{\ell^{*2}} \vartheta \wedge \vartheta.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0611 (9.3.2)	202	■0.968 ● N +1	$L_{\mathrm{SKY}} := -\frac{1}{2} R_{\alpha\beta} \wedge {}^* R^{\alpha\beta},$	$L_{\mathrm{SKY}} := -\frac{1}{2} R_{\alpha\beta} \wedge {}^* R^{\alpha\beta},$	$L_{\mathrm{SKY}} := -\frac{1}{2} R_{\alpha\beta} \wedge {}^* R^{\alpha\beta},$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0841 (12.2.9)	272	■ 0.968 ● K +0	$\mu_{\alpha}=\frac{1}{4}\vartheta_{\alpha}\wedge^{*}j_5$.	$\mu_{\alpha}=\frac{1}{4}\vartheta_{\alpha}\wedge^{*}j_5$.	$\mu_{\alpha}=\frac{1}{4}\vartheta_{\alpha}\wedge^{*}j_5$.
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0636 (10.2.7)	212	■ 0.970 ● K +0	$T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha},$	$T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha},$	$T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ1137 (C.5)	376	■ 0.970 ● N +0	$3\,c_{[i\mid s]\{ \}^{\{p\}}c_{\mid j\,k]\{ \}^{\{s\}}} = c_{\{i\,s\}\{ \}^{\{p\}}c_{\{j\,k\}\{ \}^{\{s\}}} + c_{\{j\,s\}\{ \}^{\{p\}}c_{\{k\,i\}\{ \}^{\{s\}}} + c_{\{k\,s\}\{ \}^{\{p\}}c_{\{i\,j\}\{ \}^{\{s\}}} \equiv 0$.	$3c_{[i s}{}^pc_{ jk]}{}^s := c_{is}{}^pc_{jk}{}^s + c_{js}{}^pc_{ki}{}^s + c_{ks}{}^pc_{ij}{}^s \equiv 0.$	$3c_{[i s}{}^pc_{ jk]}{}^s := c_{is}{}^pc_{jk}{}^s + c_{js}{}^pc_{ki}{}^s + c_{ks}{}^pc_{ij}{}^s \equiv 0.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0415 (6.5.3)	137	■ 0.970 ● N +0	$\begin{aligned} & \langle \vartheta W(\gamma) \vartheta \rangle = \int_{\mathcal{C} \mathcal{G}} \mathcal{D} A e^{iC} W(A, \gamma) \\ &= \frac{1}{4\pi} \oint ds \oint dt \dot{\gamma}^\mu(s) \dot{\gamma}^\nu(t) \eta_{0\mu\nu\kappa} \frac{\gamma^\kappa(s) - \gamma^\kappa(t)}{ \gamma(s) - \gamma(t) ^3} \end{aligned}$	$\begin{aligned} \langle 0 W(\gamma) 0\rangle^{(1)} &= \int_{\mathcal{C} \mathcal{G}} \mathcal{D} A e^{iC} W(A, \gamma) \\ &= \frac{1}{4\pi} \oint ds \oint dt \dot{\gamma}^\mu(s) \dot{\gamma}^\nu(t) \eta_{0\mu\nu\kappa} \frac{\gamma^\kappa(s) - \gamma^\kappa(t)}{ \gamma(s) - \gamma(t) ^3} \end{aligned}$	$\begin{aligned} \langle 0 W(\gamma) 0\rangle^{(1)} &= \int_{\mathcal{C} \mathcal{G}} \mathcal{D} A e^{iC} W(A, \gamma) \\ &= \frac{1}{4\pi} \oint ds \oint dt \dot{\gamma}^\mu(s) \dot{\gamma}^\nu(t) \eta_{0\mu\nu\kappa} \frac{\gamma^\kappa(s) - \gamma^\kappa(t)}{ \gamma(s) - \gamma(t) ^3} \end{aligned}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0495 (7.7.13)	162	■ 0.971 ● N +0	$\{ \}^{\{*\}}\backslashleft(\backslashvartheta^{\alpha}\backslashwedge\backslashvartheta^{\beta}\backslashwedge\backslashvartheta^{\gamma}\backslashwedge\backslashvartheta^{\delta}\backslashright)=\eta^{\alpha\beta\gamma\delta},\text{ where }\eta^{\alpha\beta\gamma\delta}:=+\delta^{\alpha\beta\gamma\delta},$	$^{*}\left(\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta}\right)=\eta^{\alpha\beta\gamma\delta},\text{ where }\eta_{\alpha\beta\gamma\delta}:=+\delta_{\alpha\beta\gamma\delta}^{0123},$	$^{*}(\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta})=\eta^{\alpha\beta\gamma\delta},\text{ where }\eta_{\alpha\beta\gamma\delta}:=+\delta_{\alpha\beta\gamma\delta}^{0123},$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1147 (C.16)	377	■ 0.971 ● W +1	$\begin{aligned} & \left\{ \left[\lambda_i, \lambda_j \right] \right\} \quad \text{with } \lambda_i = \frac{1}{N} \sum_k \lambda_k^i \quad \text{and } \lambda_j = \frac{1}{N} \sum_k \lambda_k^j \\ & \text{and } \lambda_k^i = \frac{1}{N} \sum_j \lambda_j^i \end{aligned}$	$[\lambda_i, \lambda_j] = 2i f_{ij}^k \lambda_k \quad (c_{ij}^k = 2i f_{ig}^k),$ $\{\lambda_i, \lambda_j\}_+ = \frac{4}{N} \delta_{ij} + 2d_{ij}^k \lambda_k.$	$[\lambda_i, \lambda_j] = 2i f_{ij}^k \lambda_k \quad (c_{ij}^k = 2i f_{ig}^k),$ $\{\lambda_i, \lambda_j\}_+ = \frac{4}{N} \delta_{ij} + 2d_{ij}^k \lambda_k.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0129 (3.3.30)	061	■ 0.972 ● C -6	$\begin{aligned} & j := \frac{\delta L_{\text{matter}}}{\delta A} = \frac{\partial L_{\text{matter}}}{\partial A} + d \frac{\partial L_{\text{matter}}}{\partial dA} \\ & = \Psi \wedge \frac{\partial L_{\text{matter}}}{\partial D\Psi} + D \frac{\partial L_{\text{matter}}}{\partial B}, \end{aligned}$	$j := \frac{\delta L_{\text{matter}}}{\delta A} = \frac{\partial L_{\text{matter}}}{\partial A} + d \frac{\partial L_{\text{matter}}}{\partial dA}$ $= \Psi \wedge \frac{\partial L_{\text{matter}}}{\partial D\Psi} + D \frac{\partial L_{\text{matter}}}{\partial B},$	$j := \frac{\delta L_{\text{matter}}}{\delta A} = \frac{\partial L_{\text{matter}}}{\partial A} + d \frac{\partial L_{\text{matter}}}{\partial dA}$ $= \Psi \wedge \frac{\partial L_{\text{matter}}}{\partial D\Psi} + D \frac{\partial L_{\text{matter}}}{\partial B},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0784 (11.3.3)	248	■ 0.972 ● N +0	$\begin{aligned} & L_{\text{NL}} = \frac{1}{2} \bar{\psi} \left[\tilde{K}, \gamma \right] \psi = -3i^* [\Theta, \vartheta] \wedge (\bar{\psi}^* \gamma \psi) \\ & = -3\tau_s \wedge [\Theta, \vartheta]. \end{aligned}$	$L_{\text{NL}} = \frac{1}{2} \bar{\psi} \left[\tilde{K}, \gamma \right] \psi = -3i^* [\Theta, \vartheta] \wedge (\bar{\psi}^* \gamma \psi)$ $= -3\tau_s \wedge [\Theta, \vartheta].$	$L_{\text{NL}} = \frac{1}{2} \bar{\psi} [\tilde{K}, \gamma] \psi = -3i^* [\Theta, \vartheta] \wedge (\bar{\psi}^* \gamma \psi)$ $= -3\tau_s \wedge [\Theta, \vartheta].$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0666 (10.3.23)	218	■ 0.972 ● N +0	$\square \vartheta_\alpha + \frac{\ell}{\gamma} {}^* D^* (\delta \ell E_\alpha + S_\alpha) = 0,$	$\square \vartheta_\alpha + \frac{\ell}{\gamma} {}^* D^* (\delta \ell E_\alpha + S_\alpha) = 0,$	$\square \vartheta_\alpha + \frac{\ell}{\gamma} {}^* D^* (\delta \ell E_\alpha + S_\alpha) = 0,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0945 (14.1.26)	306	■ 0.972 ● N +0	$\begin{aligned} \Sigma_{\mathrm{YM}}(\Omega) \wedge \Sigma_{\mathrm{YM}}(\Omega) &= \frac{1}{4\alpha_g} \operatorname{Tr} \left\{ -\xi \wedge \xi \wedge (\vartheta \wedge {}^*\xi) \wedge (\vartheta \wedge {}^*\xi) \right. \\ &\quad \left. + \vartheta \wedge \vartheta \wedge {}^*(\xi \wedge {}^*\xi)^2 - \vartheta \wedge {}^*\xi \wedge {}^*(\vartheta \wedge {}^*\xi) \wedge {}^*(\xi \wedge {}^*\xi) \right\} \\ &= \vartheta \wedge \vartheta \wedge {}^*(\Sigma_{\mathrm{YM}}(\xi) \wedge {}^*\Sigma_{\mathrm{YM}}(\xi)) \end{aligned}$	$\Sigma_{\mathrm{YM}}(\Omega) \wedge \Sigma_{\mathrm{YM}}(\Omega) = \frac{1}{4\alpha_g} \operatorname{Tr} \left\{ -\xi \wedge \xi \wedge (\vartheta \wedge {}^*\xi) \wedge (\vartheta \wedge {}^*\xi) \right. \\ \left. + \vartheta \wedge \vartheta \wedge {}^*(\xi \wedge {}^*\xi)^2 - \vartheta \wedge {}^*\xi \wedge {}^*(\vartheta \wedge {}^*\xi) \wedge {}^*(\xi \wedge {}^*\xi) \right\} \\ = \vartheta \wedge \vartheta \wedge {}^*(\Sigma_{\mathrm{YM}}(\xi) \wedge {}^*\Sigma_{\mathrm{YM}}(\xi))$	$\Sigma_{\mathrm{YM}}(\Omega) \wedge \Sigma_{\mathrm{YM}}(\Omega) = \frac{1}{4\alpha_g} \operatorname{Tr} \left\{ -\xi \wedge \xi \wedge (\vartheta \wedge {}^*\xi) \wedge (\vartheta \wedge {}^*\xi) \right. \\ \left. + \vartheta \wedge \vartheta \wedge {}^*(\xi \wedge {}^*\xi)^2 - \vartheta \wedge {}^*\xi \wedge {}^*(\vartheta \wedge {}^*\xi) \wedge {}^*(\xi \wedge {}^*\xi) \right\} \\ = \vartheta \wedge \vartheta \wedge {}^*(\Sigma_{\mathrm{YM}}(\xi) \wedge {}^*\Sigma_{\mathrm{YM}}(\xi))$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0462 (7.3.1)	153	■ 0.973 ● N +1	$L_{\mathrm{SKY}} = \operatorname{Tr}(\Omega^g \wedge {}^*\Omega^g) = -\frac{1}{2} R^{\alpha\beta} \wedge {}^*R_{\alpha\beta},$	$L_{\mathrm{SKY}} = \operatorname{Tr}(\Omega^g \wedge {}^*\Omega^g) = -\frac{1}{2} R^{\alpha\beta} \wedge {}^*R_{\alpha\beta},$	$L_{\mathrm{SKY}} = \operatorname{Tr}(\Omega^g \wedge {}^*\Omega^g) = -\frac{1}{2} R^{\alpha\beta} \wedge {}^*R_{\alpha\beta},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0356 (6.2.7)	124	■ 0.974 ● N +0	$D T^{\alpha} \equiv R_{\beta}{}^{\alpha} \wedge \vartheta^{\beta}$	$DT^{\alpha} \equiv R_{\beta}{}^{\alpha} \wedge \vartheta^{\beta}$	$DT^{\alpha} \equiv R_{\beta}{}^{\alpha} \wedge \vartheta^{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0712 (10.8.1)	228	■ 0.975 ● K +0	$\Psi = \Psi_i dx^i = \Psi_{\alpha} \vartheta^{\alpha}$	$\Psi = \Psi_i dx^i = \Psi_{\alpha} \vartheta^{\alpha}$	$\Psi = \Psi_i dx^i = \Psi_{\alpha} \vartheta^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0971 (14.2.12)	314	■ 0.975 ● N +0	$\hat{d} := \left[\begin{array}{c c} d & 0 \\ 0 & \overset{\circ}{d} \end{array} \right],$	$\hat{d} := \left[\begin{array}{c c} d & 0 \\ 0 & \overset{\circ}{d} \end{array} \right],$	$\hat{d} := \left[\begin{array}{c c} d & 0 \\ 0 & \overset{\circ}{d} \end{array} \right],$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E08557 (8.4.11)	181	■0.975 ● N +0	$\left\langle \bar{\psi} \gamma_5 \psi \right\rangle = 2 i \int d^4 x \left(\frac{1}{48} \epsilon^{\mu \nu \rho \sigma} F_{\mu \nu} F_{\rho \sigma} + \frac{1}{2} \text{Tr} (F_{\mu \nu} F^{\mu \nu}) \right)$	$\langle dj_5 \rangle = 2i \int d^4 x \left(\frac{1}{48 \pi^2} \left(R_{\alpha \beta}^{\{\gamma} \wedge R^{\alpha \beta\}_{\delta}} + \frac{1}{2} d\mathcal{A} \wedge d\mathcal{A} \right) \right),$	$\langle dj_5 \rangle = 2i \int d^4 x \left(\frac{1}{48 \pi^2} \left(R_{\alpha \beta}^{\{\gamma} \wedge R^{\alpha \beta\}_{\delta}} + \frac{1}{2} d\mathcal{A} \wedge d\mathcal{A} \right) \right),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E08487 (7.7.4)	160	■0.977 ● N +0	$\left(\frac{1}{2} \epsilon^{\mu \nu \rho \sigma} F_{\mu \nu} F_{\rho \sigma} + \frac{1}{2} \text{Tr} (F_{\mu \nu} F^{\mu \nu}) \right) = \frac{1}{2} \epsilon^{\mu \nu \rho \sigma} F_{\mu \nu} F_{\rho \sigma} + \frac{1}{2} \text{Tr} (F_{\mu \nu} F^{\mu \nu})$	$\eta_{\alpha \beta} \rightarrow \tilde{\eta}_{\alpha \beta} = \eta_{\alpha \beta} + \ell^2 \eta_{\alpha \beta \gamma} \wedge e_{\mu}^{\gamma} \wedge H^{\mu \nu} + \frac{\ell^4}{2} \eta_{\alpha \beta \gamma \delta} (e_{\mu}^{\gamma} \wedge H^{\mu \nu}) \wedge (e_{\nu}^{\delta} \wedge H^{\delta \nu}).$	$\eta_{\alpha \beta} \rightarrow \tilde{\eta}_{\alpha \beta} = \eta_{\alpha \beta} + \ell^2 \eta_{\alpha \beta \gamma} \wedge e_{\mu}^{\gamma} \wedge H^{\mu \nu} + \frac{\ell^4}{2} \eta_{\alpha \beta \gamma \delta} (e_{\mu}^{\gamma} \wedge H^{\mu \nu}) \wedge (e_{\nu}^{\delta} \wedge H^{\delta \nu}).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E01068 (16.2.7)	355	■0.978 ● N -1	$\left(\frac{1}{2} \epsilon^{\mu \nu \rho \sigma} F_{\mu \nu} F_{\rho \sigma} + \frac{1}{2} \text{Tr} (F_{\mu \nu} F^{\mu \nu}) \right) = \frac{1}{2} \epsilon^{\mu \nu \rho \sigma} F_{\mu \nu} F_{\rho \sigma} + \frac{1}{2} \text{Tr} (F_{\mu \nu} F^{\mu \nu})$	$\left[\partial_{\rho^*}^2 - V_{\text{eff}}^{\iota}(\rho^*) + \frac{\omega^2}{\beta^2} \right] F = 0$	$\left[\partial_{\rho^*}^2 - V_{\text{eff}}^{\iota}(\rho^*) + \frac{\omega^2}{\beta^2} \right] F = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E01114 (B.26)	371	■0.979 ● N +0	$H^p(M) = 0 \text{ for } p > n,$	$H^p(M) = 0 \text{ for } p > n,$	$H^p(M) = 0 \text{ for } p > n,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E08007 (2.2.2)	033	■0.980 ● K +0	$\vartheta = \vartheta^{\alpha} P_{\alpha} \in C^{\infty}(T^*(M)),$	$\vartheta = \vartheta^{\alpha} P_{\alpha} \in C^{\infty}(T^*(M)),$	$\vartheta = \vartheta^{\alpha} P_{\alpha} \in C^{\infty}(T^*(M)),$

Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value

Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library_ EQ1086 (A.12)	366	■0.982 ●W -1	\begin{aligned} & \mathrm{C}(1,1)=\mathrm{C}(2,\theta)= \\ & \mathrm{GL}(2,\mathbb{R}), \quad \& \mathrm{C}(\theta,2)= \\ & \mathbb{H} \quad \text{ (field of quaternions) }, \\ & \end{aligned}	C(1, 1) = C(2, 0) = GL(2, ℝ), C(0, 2) = ℍ (field of quaternions) ,	C(1, 1) = C(2, 0) = GL(2, ℝ), C(0, 2) = ℍ (field of quaternions),
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library_ EQ0229 (4.2.22)	088	■0.982 ●K +0	\Theta:=D \vartheta=d \vartheta-\left[\omega^{\mathrm{L}}, \vartheta\right]	Θ := Dϑ = dϑ − [ω ^L , ϑ]	Θ := Dϑ = dϑ − [ω ^L , ϑ]
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library_ EQ0836 (12.2.4)	271	■0.983 ●N +0	\mathscr{A}:=\frac{1}{4}\{ \ }^{*} \operatorname{Tr}(\gamma \wedge D \gamma)=\{ \ }^{*} \left(\vartheta^{\alpha} \wedge T_{\alpha} \wedge \beta\right)=\frac{1}{2} T^{\left[\alpha \beta \gamma\right]} \eta_{\alpha \beta \gamma}=\mathscr{A}_i d x^i,	ℳ := ¹ / ₄ *Tr(γ ∧ Dγ) = *(ϑ ^α ∧ T _α) = ¹ / ₂ T ^[αβγ] η _{αβγ} = ℳ _i dx ⁱ ,	ℳ := ¹ / ₄ *Tr(γ ∧ Dγ) = *(ϑ ^α ∧ T _α) = ¹ / ₂ T ^[αβγ] η _{αβγ} = ℳ _i dx ⁱ ,
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library_ EQ0511 (7.7.29)	165	■0.983 ●N -1	\begin{aligned} \Theta:=D \gamma=d \gamma+\left[d \vartheta^{\alpha}+\Gamma_{\beta}^{\alpha} \wedge \vartheta^{\beta}\right] \gamma_{\alpha} \\ &=T^{\alpha} \gamma_{\alpha}=\frac{1}{2} T_{ij}^{\alpha} \gamma_{\alpha} d x^i \wedge d x^j \end{aligned}	Θ := Dγ = dγ + [Γ, γ] = (dϑ ^α + Γ _β ^α ∧ ϑ ^β) γ _α = T ^α γ _α = ¹ / ₂ T _{ij} ^α γ _α dx ⁱ ∧ dx ^j	Θ := Dγ = dγ + [Γ, γ] = (dϑ ^α + Γ _β ^α ∧ ϑ ^β) γ _α = T ^α γ _α = ¹ / ₂ T _{ij} ^α γ _α dx ⁱ ∧ dx ^j ,
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library_ EQ0637 (10.2.8)	212	■0.983 ●N +0	\Gamma_{\alpha}^{\star}=\Gamma_{\alpha}^{\{\backslash \backslash \} \star}-K_{\alpha}^{\star}	Γ _α [★] = Γ _α ^{{ }★} − K _α [★]	Γ _α [★] = Γ _α ^{{ }★} − K _α [★]
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0271 (4.3.36)	096	■ 0.983 ● N +0	$\left[D_{\alpha}, D_{\beta}\right]=R_{\alpha\beta}^{c d} L_{c d}-T_{\alpha\beta}^{c} D_c,$	$[D_{\alpha}, D_{\beta}] = R_{\alpha\beta}^{..cd} L_{cd} - T_{\alpha\beta}^{..c} D_c,$	$[D_{\alpha}, D_{\beta}] = R_{\alpha\beta}^{..cd} L_{cd} - T_{\alpha\beta}^{..c} D_c,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0561 (8.5.4)	181	■ 0.984 ● N +0	$\{ \}^{\{*\}} \sigma^{\{(\star)\}}=(-1)^{\{\mathrm{sig}\}} \sigma,$	$^*\sigma^{(\star)} = (-1)^{\mathrm{sig}} \sigma,$	$^*\sigma^{(\star)} = (-1)^{\mathrm{sig}} \sigma,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0097 (3.2.7)	056	■ 0.985 ● K +0	$D \rightarrowtail \{ \}^{G^{-1}} D=G^{-1} D \, G \, .$	$D \rightarrow {}^{G^{-1}} D = G^{-1} D G.$	$D \rightarrow {}^{G^{-1}} D = G^{-1} D G.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0459 (7.2.11)	152	■ 0.985 ● N +0	$\Omega_{\mathrm{C}}=\stackrel{(+)}{\Omega}_{\mathrm{C}}+\stackrel{(-)}{\Omega}_{\mathrm{C}}, \quad \stackrel{(\pm)}{\Omega}_{\mathrm{C}}:=\frac{1}{2}\left(\Omega_{\mathrm{C}}\pm\Omega_{\mathrm{C}}^{(*)}\right)$	$\Omega_{\mathrm{C}}=\stackrel{(+)}{\Omega}_{\mathrm{C}}+\stackrel{(-)}{\Omega}_{\mathrm{C}}, \quad \stackrel{(\pm)}{\Omega}_{\mathrm{C}}:=\frac{1}{2}\left(\Omega_{\mathrm{C}}\pm\Omega_{\mathrm{C}}^{(*)}\right)$	$\Omega_{\mathrm{C}}=\stackrel{(+)}{\Omega}_{\mathrm{C}}+\stackrel{(-)}{\Omega}_{\mathrm{C}}, \quad \stackrel{(\pm)}{\Omega}_{\mathrm{C}}:=\frac{1}{2}\left(\Omega_{\mathrm{C}}\pm\Omega_{\mathrm{C}}^{(*)}\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1051 (15.2.8)	344	■ 0.985 ● N +0	$\begin{aligned} A_i & \& =\frac{1}{2}\left\{ A_i^{(f)j} + \gamma^5 A_i^{(f)j} \right\} \lambda_j^{(f)} \\ & \oplus \frac{1}{2}\left\{ A_i^{(c)j} + \gamma^5 A_i^{(c)j} \right\} \lambda_j^{(c)}, \end{aligned}$	$A_i = \frac{1}{2} \left\{ A_i^{(f)j} + \gamma^5 A_i^{(f)j} \right\} \lambda_j^{(f)} \oplus \frac{1}{2} \left\{ A_i^{(c)j} + \gamma^5 A_i^{(c)j} \right\} \lambda_j^{(c)},$	$A_i = \frac{1}{2} \left\{ A_i^{(f)j} + \gamma^5 A_i^{(f)j} \right\} \lambda_j^{(f)} \oplus \frac{1}{2} \left\{ A_i^{(c)j} + \gamma^5 A_i^{(c)j} \right\} \lambda_j^{(c)},$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ1096 (B.6)	369	■0.985 ● N +0	$d\,d\,\alpha^{\wedge\{p\}}\equiv 0\;.$	$dd\alpha^{(p)}\equiv 0.$	$d\,d\alpha^{(p)}\equiv 0.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0191 (4.1.1)	081	■0.985 ● N +0	$x\rightarrow x^{\prime}=g\,x+\mathrm{a},\quad g\in G\,L(n,\,\mathbb{R}),\,\mathrm{a}\in\mathbb{R}^n\;.$	$x\rightarrow x'=gx+\mathbf{a},\quad g\in GL(n,\mathbb{R}),a\in\mathbb{R}^n.$	$x\rightarrow x'=gx+\mathbf{a},\quad g\in GL(n,\mathbb{R}),\,a\in\mathbb{R}^n.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0694 (10.5.5)	224	■0.985 ● W +0	$\begin{aligned} L_{\mathrm{D}}\,\,\&=\frac{i}{2}\left(\overline{\psi}^{\wedge{*}}\,\gamma\wedge D\psi+D\overline{\psi}\wedge^{*}\gamma\psi\right)-m\overline{\psi}\psi\eta\\ &\gamma\psi\right)-m\overline{\psi}\psi\eta\,\,\&=\overline{\psi}\left(i^{*}\gamma\wedge D-m\eta\right)\psi+d\left(\frac{i}{2}\overline{\psi}^{\wedge{*}}\gamma\psi\right),\\ &\left(\frac{i}{2}\overline{\psi}^{\wedge{*}}\,\gamma\wedge D\psi+D\overline{\psi}\wedge^{*}\gamma\psi\right),\\ &\end{aligned}$	$L_{\mathrm{D}}=\frac{i}{2}\left(\overline{\psi}^{*}\gamma\wedge D\psi+D\overline{\psi}\wedge^{*}\gamma\psi\right)-m\overline{\psi}\psi\eta\\ =\overline{\psi}\left(i^{*}\gamma\wedge D-m\eta\right)\psi+d\left(\frac{i}{2}\overline{\psi}^{*}\gamma\psi\right),$	$L_{\mathrm{D}}=\frac{i}{2}\left(\overline{\psi}^{*}\gamma\wedge D\psi+D\overline{\psi}\wedge^{*}\gamma\psi\right)-m\overline{\psi}\psi\eta\\ =\overline{\psi}\left(i^{*}\gamma\wedge D-m\eta\right)\psi+d\left(\frac{i}{2}\overline{\psi}^{*}\gamma\psi\right),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0554 (8.4.8)	180	■0.986 ● N +0	$D^{\wedge{*}}\,\,\stackrel{\mathrm{rel}}{\sim}\,\,\mathrm{pm}\,\,\{\,\,\omega^{\wedge\{g\}}=0$	$D^{*}\,\,\overset{\pm g}{\Omega}=0$	$D^{*}\,\,\overset{\pm}{\Omega}^g=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0058 (2.7.8)	045	■0.987 ● K +0	$b\rightarrow^G b:=G\,b\quad\mathrm{in}\,\,C^{\wedge\{\infty\}}\left(V^{\wedge\{\rho\}}\right)\;.$	$b\rightarrow^G b:=Gb\quad\in C^{\infty}\left(V^{\rho}\right).$	$b\rightarrow^G b:=Gb\quad\in C^{\infty}(V^{\rho}).$
Conf. MathPix confidence: ■≥0.9 ■0.5–0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Ekehard_W._Mielke_ __Z-Library__ EQ0285 (4.4.14)	099	■0.987 ● N +0	$D\left(\vartheta\wedge^*\Omega^g\right)=\Theta\wedge^*\Omega^g$	$D\left(\vartheta\wedge^*\Omega^g\right)=\Theta\wedge^*\Omega^g$	$D(\vartheta\wedge^*\Omega^g)=\Theta\wedge^*\Omega^g$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Ekehard_W._Mielke_ __Z-Library__ EQ0834 (12.2.2)	271	■0.988 ● N +0	$i^*\gamma\wedge\left(\mathscr{D}+\frac{i}{4}m\gamma-\frac{1}{2}T\right)\psi=0,$	$i^*\gamma\wedge\left(\mathscr{D}+\frac{i}{4}m\gamma-\frac{1}{2}T\right)\psi=0,$	$i^*\gamma\wedge\left(\mathscr{D}+\frac{i}{4}m\gamma-\frac{1}{2}T\right)\psi=0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Ekehard_W._Mielke_ __Z-Library__ EQ0002 (2.1.1)	031	■0.989 ● N +0	$x_{\{2\}}\circ\stackrel{-1}{x}_{\{1\}}:\mathbb{R}\rightarrow\mathbb{R}$	$x_2\circ\stackrel{-1}{x}_1:\mathbb{R}\rightarrow\mathbb{R}$	$x_2\circ\stackrel{-1}{x}_1:\mathbb{R}\rightarrow\mathbb{R}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Ekehard_W._Mielke_ __Z-Library__ EQ0339 (5.3.16)	114	■0.989 ● N +0	$L_{\{\mathrm{EC}\}}\!:=\!\frac{1}{2\,\kappa}\,R^{\alpha\beta}\wedge\eta_{\alpha\beta}\!=\!L_{\{\mathrm{HE}\}}\!+\!\frac{1}{12\,\kappa}\,\mathscr{A}^*\mathscr{A}$	$L_{\mathrm{EC}}:=-\frac{1}{2\kappa}R^{\alpha\beta}\wedge\eta_{\alpha\beta}=L_{\mathrm{HE}}+\frac{1}{12\kappa}\mathscr{A}^*\mathscr{A}$	$L_{\mathrm{EC}}:=-\frac{1}{2\kappa}R^{\alpha\beta}\wedge\eta_{\alpha\beta}=L_{\mathrm{HE}}+\frac{1}{12\kappa}\mathscr{A}\wedge^*\mathscr{A}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Ekehard_W._Mielke_ __Z-Library__ EQ0024 (2.4.6)	039	■0.990 ● N +0	$\left[L_{\{i\}},L_{\{j\}}\right]=c_{\{i\}j}^{\{k\}}L_{\{k\}}.$	$[L_i,L_j]=c_{ij}^{k}L_k.$	$[L_i,L_j]=c_{ij}^{k}L_k.$
Conf. MathPix confidence: ■≥0.9 ■0.5-0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ _Z-Library_ EQ0507 (7.7.25)	164	■ 0.990 ● N +1	$\sigma_{\pm}:=\left(\sigma\pm i^*\sigma\right)/2=\frac{1}{2}\left(1\mp\gamma_5\right)\sigma,\quad\text{with}\quad i^*\sigma_{\pm}=\pm\sigma_{\pm}$		
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ _Z-Library_ EQ0572 (8.6.10)	184	■ 0.990 ● N -2	$\begin{aligned} H_{\alpha}(* *) &= \frac{\theta_{\mathrm{T}}}{2\ell^2} K^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha} = \frac{1}{\ell^2} \left[\theta_{\mathrm{T}}^* K_{\alpha}^{(*)} - \theta_{\mathrm{T}} T_{\alpha} \right] \\ &= \frac{\theta_{\mathrm{T}}^*}{\ell^2} * \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta}] T^{\beta}) - \frac{1}{2} e_{\alpha}] (T^{\beta} \wedge \vartheta_{\beta}) \right] - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha} \\ &\simeq -\partial L_{\theta} / \partial T^{\alpha}. \end{aligned}$	$H_{\alpha}(**) = \frac{\theta_{\mathrm{T}}^*}{2\ell^2} K^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha} = \frac{1}{\ell^2} \left[\theta_{\mathrm{T}}^* K_{\alpha}^{(*)} - \theta_{\mathrm{T}} T_{\alpha} \right]$ $= \frac{\theta_{\mathrm{T}}^*}{\ell^2} * \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta}] T^{\beta}) - \frac{1}{2} e_{\alpha}] (T^{\beta} \wedge \vartheta_{\beta}) \right] - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha}$ $\simeq -\partial L_{\theta} / \partial T^{\alpha}.$	$H_{\alpha}(**) = \frac{\theta_{\mathrm{T}}^*}{2\ell^2} K^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha} = \frac{1}{\ell^2} \left[\theta_{\mathrm{T}}^* K_{\alpha}^{(*)} - \theta_{\mathrm{T}} T_{\alpha} \right]$ $= \frac{\theta_{\mathrm{T}}^*}{\ell^2} * \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta}] T^{\beta}) - \frac{1}{2} e_{\alpha}] (T^{\beta} \wedge \vartheta_{\beta}) \right] - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha}$ $\simeq -\partial L_{\theta} / \partial T^{\alpha}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ _Z-Library_ EQ0110 (3.3.11)	058	■ 0.990 ● C -3	$\begin{aligned} F &:= E \wedge dt + B \\ &= (E_x dx + E_y dy + E_z dz) \wedge dt \\ &\quad + B_x dy \wedge dz + B_y dz \wedge dx + B_z dx \wedge dy. \end{aligned}$	$F := E \wedge dt + B$ $= (E_x dx + E_y dy + E_z dz) \wedge dt$ $+ B_x dy \wedge dz + B_y dz \wedge dx + B_z dx \wedge dy.$	$F := E \wedge dt + B$ $= (E_x dx + E_y dy + E_z dz) \wedge dt$ $+ B_x dy \wedge dz + B_y dz \wedge dx + B_z dx \wedge dy.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ _Z-Library_ EQ0142 (3.4.13)	064	■ 0.990 ● W -2	$\begin{aligned} D\tau &= DD\Pi^{\Omega} = [\Omega, \Pi^{\Omega}] \\ &= \frac{2}{\alpha_g} \left\{ [\Omega, {}^*\Omega] \frac{\partial L}{\partial \mathcal{F}} + (-1)^s [\Omega, \Omega] \frac{\partial L}{\partial {}^*\mathcal{F}} \right\} = 0. \end{aligned}$	$D\tau = DD\Pi^{\Omega} = [\Omega, \Pi^{\Omega}]$ $= \frac{2}{\alpha_g} \left\{ [\Omega, {}^*\Omega] \frac{\partial L}{\partial \mathcal{F}} + (-1)^s [\Omega, \Omega] \frac{\partial L}{\partial {}^*\mathcal{F}} \right\} = 0.$	$D\tau = DD\Pi^{\Omega} = [\Omega, \Pi^{\Omega}]$ $= \frac{2}{\alpha_g} \left\{ [\Omega, {}^*\Omega] \frac{\partial L}{\partial \mathcal{F}} + (-1)^s [\Omega, \Omega] \frac{\partial L}{\partial {}^*\mathcal{F}} \right\} = 0.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0155 (3.5.3)	067	0.990 K +0	$\stackrel{*}{\Omega}=\frac{1}{4}\,F_{\alpha\,\beta}\,\sigma^{\mu\nu}\otimes\vartheta^{\alpha}\wedge\vartheta^{\beta}$	$\stackrel{*}{\sigma}\,\Omega=\frac{1}{4}F_{\alpha\beta\mu\nu}\sigma^{\mu\nu}\otimes\vartheta^{\alpha}\wedge\vartheta^{\beta}$	$\stackrel{*}{\sigma}\Omega=\frac{1}{4}F_{\alpha\beta\mu\nu}\sigma^{\mu\nu}\otimes\vartheta^{\alpha}\wedge\vartheta^{\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0682 (10.3.40)	221	0.990 C -3	$\theta_{\mathrm{L}}C_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}}\tilde{C}_{\mathrm{L}}=L_{\mathrm{MB}}+\frac{\Delta\theta_{\mathrm{TL}}}{\ell}d\left(\Gamma^{\star\alpha}\wedge\vartheta_{\alpha}\right),$	$\theta_{\mathrm{L}}C_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}}\tilde{C}_{\mathrm{L}}=L_{\mathrm{MB}}+\frac{\Delta\theta_{\mathrm{TL}}}{\ell}d\left(\Gamma^{\star\alpha}\wedge\vartheta_{\alpha}\right),$	$\theta_{\mathrm{L}}C_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}}\tilde{C}_{\mathrm{L}}=L_{\mathrm{MB}}+\frac{\Delta\theta_{\mathrm{TL}}}{\ell}d(\Gamma^{\star\alpha}\wedge\vartheta_{\alpha}),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0874 (13.3.2)	284	0.990 W -2	$\left[Z^{\mathrm{A}}_{\mathrm{B}},\,Z^{\mathrm{C}}_{\mathrm{D}}\right]=\delta_{\mathrm{D}}^{\mathrm{A}}Z^{\mathrm{C}}_{\mathrm{B}}-\delta_{\mathrm{B}}^{\mathrm{C}}Z^{\mathrm{A}}_{\mathrm{D}}$	$\left[Z^A{}_B,Z^C{}_D\right]=\delta_D^AZ^C{}_B-\delta_B^CZ^A{}_D$	$[\not{L}^A{}_B,\not{L}^C{}_D]=\delta_D^A\not{L}^C{}_B-\delta_B^C\not{L}^A{}_D$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0042 (2.6.5)	042	0.991 C -3	$F_{\alpha\beta}{}^j=\partial_{\alpha}\Gamma_{\beta}{}^j-\partial_{\beta}\Gamma_{\alpha}{}^j-c_{kl}{}^j\Gamma_{\alpha}{}^k\Gamma_{\beta}{}^l,$	$F_{\alpha\beta}{}^j=\partial_{\alpha}\Gamma_{\beta}{}^j-\partial_{\beta}\Gamma_{\alpha}{}^j-c_{kl}{}^j\Gamma_{\alpha}{}^k\Gamma_{\beta}{}^l,$	$F_{\alpha\beta}{}^j=\partial_{\alpha}\Gamma_{\beta}{}^j-\partial_{\beta}\Gamma_{\alpha}{}^j-c_{kl}{}^j\Gamma_{\alpha}{}^k\Gamma_{\beta}{}^l,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0233 (4.2.26)	088	0.991 K +0	$\Omega^{\mathrm{T}}=f\Theta+df\wedge\vartheta-\Omega^{\mathrm{L}}\otimes\widetilde{\varphi}$	$\Omega^{\mathrm{T}}=f\Theta+df\wedge\vartheta-\Omega^{\mathrm{L}}\otimes\widetilde{\varphi}$	$\Omega^{\mathrm{T}}=f\Theta+df\wedge\vartheta-\Omega^{\mathrm{L}}\otimes\widetilde{\varphi}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0778	246	0.991 N +0	$\langle-\infty D\tau_A \infty\rangle^{\scriptscriptstyle(+)}=\frac{1}{(4.)8\pi^2}\,\mathrm{Tr}\left(\Omega^g\wedge\Omega^{g(*)}\right).$	$\langle-\infty D\tau_A \infty\rangle^{\scriptscriptstyle(+)}=\frac{1}{(4.)8\pi^2}\,\mathrm{Tr}\left(\Omega^g\wedge\Omega^{g(*)}\right).$	$\langle-\infty D\tau_A \infty\rangle^{\scriptscriptstyle(+)}=\frac{1}{(4.)8\pi^2}\mathrm{Tr}(\Omega^g\wedge\Omega^{g(*)}).$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0644 (10.3.1)	214	■0.991 ● N +0	$\widetilde{\Gamma}_{\alpha}^{\star} = \Gamma_{\alpha}^{\star} - (-1)^s \frac{\varepsilon}{2\ell} \vartheta_{\alpha},$	$\widetilde{\Gamma}_{\alpha}^{\star} = \Gamma_{\alpha}^{\star} - (-1)^s \frac{\varepsilon}{2\ell} \vartheta_{\alpha},$	$\widetilde{I}_{\alpha}^{\star} = I_{\alpha}^{\star} - (-1)^s \frac{\varepsilon}{2\ell} \vartheta_{\alpha},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0149 (3.4.20)	065	■0.992 ● C -3	$L\left(\mathrm{F},\{ \}^{\ast}\mathrm{F}\right)=-\mathrm{F}.$	$L\left(\mathcal{F},\ast\mathcal{F}\right)=-\mathcal{F}.$	$L(\mathcal{F},\ast\mathcal{F})=-\mathcal{F}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0368 (6.2.19)	127	■0.992 ● N +0	$\stackrel{\scriptscriptstyle\rm (pm)}{\relax}\{L_{\scriptscriptstyle\rm ()}:=L_{\scriptscriptstyle\rm ()}\pm i\frac{2}{\kappa}\ell^2\}\{\kappa\}dC_{\scriptscriptstyle\rm (TT)}=L_{\scriptscriptstyle\rm (HE)}-L_{\scriptscriptstyle\rm (EC)}\pm i\frac{2}{\kappa}\ell^2\}\{\kappa\}d\stackrel{\scriptscriptstyle\rm (mp)}{\relax}\{C_{\scriptscriptstyle\rm (TT)}\}.$	$\stackrel{\scriptscriptstyle\rm (\pm)}{L}_{\scriptscriptstyle\rm ()}:=L_{\scriptscriptstyle\rm ()}\pm i\frac{2\ell^2}{\kappa}dC_{\scriptscriptstyle\rm TT}=L_{\scriptscriptstyle\rm HE}-L_{\scriptscriptstyle\rm EC}\pm i\frac{2\ell^2}{\kappa}d\stackrel{\scriptscriptstyle\rm (\mp)}{C}_{\scriptscriptstyle\rm TT}.$	$\stackrel{\scriptscriptstyle\rm (\pm)}{L}_{\scriptscriptstyle\rm ()}:=L_{\scriptscriptstyle\rm ()}\pm i\frac{2\ell^2}{\kappa}dC_{\scriptscriptstyle\rm TT}=L_{\scriptscriptstyle\rm HE}-L_{\scriptscriptstyle\rm EC}\pm i\frac{2\ell^2}{\kappa}d\stackrel{\scriptscriptstyle\rm (\mp)}{C}_{\scriptscriptstyle\rm TT}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0449 (7.2.2)	151	■0.992 ● N +0	$\{ \}^{\ast}\Omega^{\scriptscriptstyle\rm (g)}(\star):=\frac{1}{4}\{ \}^{\ast}R_{\scriptscriptstyle\rm (alpha\beta)}^{\scriptscriptstyle\rm (c d)}L_{\scriptscriptstyle\rm (c d)}\otimes\vartheta^{\scriptscriptstyle\rm (alpha)}\wedge\vartheta^{\scriptscriptstyle\rm (beta)},$	$\ast\Omega^{g(\ast)}:=\frac{1}{4}\ast R_{\alpha\beta}^{(\ast)cd}L_{cd}\otimes\vartheta^{\alpha}\wedge\vartheta^{\beta},$	$\ast\Omega^{g(\ast)}:=\frac{1}{4}\ast R_{\alpha\beta}^{(\ast)cd}L_{cd}\otimes\vartheta^{\alpha}\wedge\vartheta^{\beta},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ1050 (15.2.7)	344	■0.992 ● N +0	$\left\{\overline{L}_{\scriptscriptstyle\rm (j)}=\sigma^{\scriptscriptstyle\rm (alpha\beta)}\otimes\lambda_{\scriptscriptstyle\rm (i)},\quad\lambda_{\scriptscriptstyle\rm (i)},\quad\gamma^5\lambda_{\scriptscriptstyle\rm (i)}\mid j=1,\ldots 8N^2-1\right\},$	$\left\{\overline{L}_j=\sigma^{\alpha\beta}\otimes\lambda_i,\quad\lambda_i,\quad i\gamma^5\lambda_i\mid j=1,\ldots 8N^2-1\right\},$	$\left\{\overline{L}_j=\sigma^{\alpha\beta}\otimes\lambda_i,\quad\lambda_i,\quad i\gamma^5\lambda_i\mid j=1,\ldots 8N^2-1\right\},$
Conf. MathPix confidence: ■≥0.9 ■0.5-0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ E09964 (14.2.4)	313	0.992 N +0	$\stackrel{\circ}{\omega}=\stackrel{\circ}{b}\stackrel{\circ}{\vartheta}$.	$\stackrel{\circ}{\omega}=\stackrel{\circ}{b}\stackrel{\circ}{\vartheta}$.	$\stackrel{\circ}{\omega}=\stackrel{\circ}{b}\stackrel{\circ}{\vartheta}$.
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ E08256 (4.3.21)	093	0.992 W +0	$R_{\alpha\beta c}^{\dots d}=2\left\{\partial_{[\alpha}\Gamma_{\beta]c}^{\dots d}+\Gamma_{[\alpha h}^{\dots d}\Gamma_{ \beta]c}^{\dots h}\right\}$	$R_{\alpha\beta c}^{\dots d}=2\left\{\partial_{[\alpha}\Gamma_{\beta]c}^{\dots d}+\Gamma_{[\alpha h}^{\dots d}\Gamma_{ \beta]c}^{\dots h}\right\}$	$R_{\alpha\beta c}^{\dots d}=2\left\{\partial_{[\alpha}\Gamma_{\beta]c}^{\dots d}+\Gamma_{[\alpha h}^{\dots d}\Gamma_{ \beta]c}^{\dots h}\right\}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ E08245 (4.3.10)	091	0.992 N +0	$\vartheta\wedge^*\vartheta=\eta$.	$\vartheta\wedge^*\vartheta=\eta$.	$\vartheta\wedge^*\vartheta=\eta$.
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ E08633 (10.2.4)	212	0.992 N +0	$-(-1)^s\theta_{\mathrm{TL}}T_\alpha-\frac{1}{2\ell}\theta_{\mathrm{T}}\eta_\alpha-\theta_{\mathrm{L}}\ell R_\alpha^\star=\ell\tau_\alpha^\star;$ $\theta_{\mathrm{L}}\ell R_\alpha^\star=\ell\tau_\alpha^\star;$	$-(-1)^s\theta_{\mathrm{TL}}T_\alpha-\frac{1}{2\ell}\theta_{\mathrm{T}}\eta_\alpha-\theta_{\mathrm{L}}\ell R_\alpha^\star=\ell\tau_\alpha^\star;$	$-(-1)^s\theta_{\mathrm{TL}}T_\alpha-\frac{1}{2\ell}\theta_{\mathrm{T}}\eta_\alpha-\theta_{\mathrm{L}}\ell R_\alpha^\star=\ell\tau_\alpha^\star;$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ E08781 (10.5.12)	225	0.992 N +0	$\hat{D}^*\hat{D}\psi-m^2\eta\psi+i^*\sigma\wedge\left(\hat{R}+T^\alpha\hat{D}_\alpha\right)\psi=0,$	$\hat{D}^*\hat{D}\psi-m^2\eta\psi+i^*\sigma\wedge\left(\hat{R}+T^\alpha\hat{D}_\alpha\right)\psi=0,$	$\hat{D}^*\hat{D}\psi-m^2\eta\psi+i^*\sigma\wedge\left(\hat{R}+T^\alpha\hat{D}_\alpha\right)\psi=0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ E08847 (12.2.15)	273	0.992 N +0	$\mathscr{A}_\alpha\mathscr{A}^\alpha\eta=\mathscr{A}\wedge^*\mathscr{A}\cong(\ell^4/4)^*j_5\wedge j_5=0.$	$\mathscr{A}_\alpha\mathscr{A}^\alpha\eta=\mathscr{A}\wedge^*\mathscr{A}\cong(\ell^4/4)^*j_5\wedge j_5=0.$	$\mathscr{A}_\alpha\mathscr{A}^\alpha\eta=\mathscr{A}\wedge^*\mathscr{A}\cong(\ell^4/4)^*j_5\wedge j_5=0.$
Conf. MathPix confidence: ≥0.9 0.5–0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0388 (6.4.3)	130	■0.993 ● N +0	$\backslash\mathrm{left}.\backslash\underline{\backslash\Psi}:=n\backslash\mathrm{rfloor}(d\,t\,\wedge\,\backslash\Psi)=\backslash\mathrm{left}(1-\{ \}^{\wedge}\{\backslash\mathrm{perp}\}\backslash\mathrm{right})\,\backslash\Psi,\,\backslash\mathrm{quad}\,n\backslash\mathrm{right}\backslash\mathrm{rfloor}\backslash\underline{\backslash\Psi}\,\backslash\mathrm{equiv}\,0\,.$	$\underline{\Psi}:=n\rfloor(dt\wedge\Psi)=(1-\perp)\Psi,\quad n\rfloor\underline{\Psi}\equiv 0.$	$\underline{\Psi}:=n\rfloor(dt\wedge\Psi)=(1-\perp)\Psi,\quad n\rfloor\underline{\Psi}\equiv 0.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0228 (4.2.21)	088	■0.993 ● N +0	$\backslash\mathrm{widetilde}\{\backslash\mathrm{widetilde}\{\backslash\mathrm{vartheta}\}\}=f\,\backslash\mathrm{vartheta}\,.$	$\widetilde{\vartheta}=f\vartheta.$	$\widetilde{\vartheta}=f\vartheta.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0287 (4.4.16)	099	■0.993 ● K +0	$D\,D\,\backslash\Pi^{\backslash\mathrm{mathrm}\{L\}}+D\,\backslash\Pi_{[\,]}^{\backslash\mathrm{mathrm}\{T\}}=D\,\backslash\tau_{\{s\}}\,.$	$DD\Pi^{\mathrm{L}}+D\Pi_{\,}^{\mathrm{T}}=D\tau_s.$	$D\,D\Pi^{\mathrm{L}}+D\Pi_{\,}^{\mathrm{T}}=D\tau_s.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0320 (5.2.4)	111	■0.993 ● N +0	$\backslash\mathrm{mathscr}\{A\}:=\frac{1}{4}\{ \}^{\wedge}\{*\}\,\backslash\mathrm{operatorname}\{Tr\}(\backslash\gamma\wedge D\,\backslash\gamma)=\{ \}^{\wedge}\{*\}\backslash\mathrm{left}(\backslash\mathrm{vartheta}^{\backslash\alpha}\wedge T_{\backslash\alpha}\backslash\mathrm{right})=\frac{1}{2}\,T^{\backslash[\alpha\backslash\beta\backslash\gamma]}\,\backslash\mathrm{eta}_{\backslash\alpha\backslash\beta\backslash\gamma}=\backslash\mathrm{mathscr}\{A\}_{\{i\}}\,d\,x^{\{i\}},$	$\mathscr{A}:=\frac{1}{4}{}^*\mathrm{Tr}(\gamma\wedge D\gamma)={}^*(\vartheta^\alpha\wedge T_\alpha)=\frac{1}{2}T^{[\alpha\beta\gamma]}\eta_{\alpha\beta\gamma}=\mathscr{A}_i dx^i,$	$\mathscr{A}:=\frac{1}{4}{}^*Tr(\gamma\wedge D\gamma)={}^*(\vartheta^\alpha\wedge T_\alpha)=\frac{1}{2}T^{[\alpha\beta\gamma]}\eta_{\alpha\beta\gamma}=\mathscr{A}_i\,dx^i\,,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0100 (3.3.1)	056	■0.994 ● N -2	$F:=d\,A=\frac{1}{2}\,F_{\{i\}\{j\}}\,d\,x^{\{i\}}\wedge d\,x^{\{j\}}\,.$	$F:=dA=\frac{1}{2}F_{ij}dx^i\wedge dx^j.$	$F:=dA=\frac{1}{2}F_{ij}dx^i\wedge dx^j\,.$
Conf. MathPix confidence: ■≥0.9 ■0.5–0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ _Z-Library_ EQ1102 (B.14)	370	0.994 N +1	$\begin{aligned} d_{\mu}(M) &:= \sqrt[n]{ \det g_{\alpha\beta} } \vartheta^1 \wedge \cdots \wedge \vartheta^n = \eta \\ &= \sqrt{ \det g_{ij} } dx^1 \wedge \cdots \wedge dx^n = \frac{1}{n} \vartheta \wedge^* \vartheta. \end{aligned}$	$d_{\mu}(M) := \sqrt[n]{ \det g_{\alpha\beta} } \vartheta^1 \wedge \cdots \wedge \vartheta^n = \eta$ $= \sqrt{ \det g_{ij} } dx^1 \wedge \cdots \wedge dx^n = \frac{1}{n} \vartheta \wedge^* \vartheta.$	$d_{\mu}(M) := \sqrt[n]{ \det g_{\alpha\beta} } \vartheta^1 \wedge \cdots \wedge \vartheta^n = \eta$ $= \sqrt{ \det g_{ij} } dx^1 \wedge \cdots \wedge dx^n = \frac{1}{n} \vartheta \wedge^* \vartheta.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ _Z-Library_ EQ0467 (7.3.6)	154	0.994 N +0	$\begin{aligned} L_{\mathrm{SKY}}^{(\star)} &= -\frac{1}{2} R^{\alpha\beta} \wedge^* R_{\alpha\beta} - \frac{(-1)^s}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(\star)} \\ &= -\frac{1}{4} \left(R_{\alpha\beta} +^* R_{\alpha\beta}^{(\star)} \right) \wedge^* \left(R^{\alpha\beta} +^* R^{\alpha\beta(\star)} \right) \\ &= \frac{1}{2} \mathrm{Tr} \left(\bar{\Omega}^g \wedge^* \bar{\Omega}^g \right). \end{aligned}$	$L_{\mathrm{SKY}}^{(\star)} = -\frac{1}{2} R^{\alpha\beta} \wedge^* R_{\alpha\beta} - \frac{(-1)^s}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(\star)}$ $= -\frac{1}{4} \left(R_{\alpha\beta} +^* R_{\alpha\beta}^{(\star)} \right) \wedge^* \left(R^{\alpha\beta} +^* R^{\alpha\beta(\star)} \right)$ $= \frac{1}{2} \mathrm{Tr}(\bar{\Omega}^g \wedge^* \bar{\Omega}^g).$	$L_{\mathrm{SKY}}^{(\star)} = -\frac{1}{2} R^{\alpha\beta} \wedge^* R_{\alpha\beta} - \frac{(-1)^s}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(\star)}$ $= -\frac{1}{4} \left(R_{\alpha\beta} +^* R_{\alpha\beta}^{(\star)} \right) \wedge^* \left(R^{\alpha\beta} +^* R^{\alpha\beta(\star)} \right)$ $= \frac{1}{2} \mathrm{Tr}(\bar{\Omega}^g \wedge^* \bar{\Omega}^g).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ _Z-Library_ EQ0490 (7.7.16)	163	0.994 N +0	$\{ \}^{\star} \Phi := \frac{1}{(4-p)! p!} \sqrt{\det g_{ij}} \varepsilon^{i_1 \cdots i_p}_{j_1 \cdots j_{4-p}} \Phi_{i_1 \cdots i_p} dx^{j_1} \wedge \cdots \wedge dx^{j_{4-p}}$	$^*\Phi := \frac{1}{(4-p)!p!} \sqrt{\det g_{ij}} \varepsilon^{i_1 \cdots i_p}_{j_1 \cdots j_{4-p}} \Phi_{i_1 \cdots i_p} dx^{j_1} \wedge \cdots \wedge dx^{j_{4-p}}$	$^*\Phi := \frac{1}{(4-p)!p!} \sqrt{\det g_{ij}} \varepsilon^{i_1 \cdots i_p}_{j_1 \cdots j_{4-p}} \Phi_{i_1 \cdots i_p} dx^{j_1} \wedge \cdots \wedge dx^{j_{4-p}},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ _Z-Library_ EQ0724 (10.10.4)	230	0.994 N +0	$\delta_{\mathrm{susy}} L = i \bar{\sigma} \Psi \gamma^\alpha \wedge \frac{\delta L}{\delta \vartheta^\alpha} + \delta_{\mathrm{susy}} \Gamma_\alpha^\star \wedge \frac{\delta L}{\delta \Gamma_\alpha^\star} + (2 \overline{D\sigma} + c \bar{\sigma} \gamma) \wedge \frac{\delta L}{\delta \overline{\Psi}},$	$\delta_{\mathrm{susy}} L = i \bar{\sigma} \Psi \gamma^\alpha \wedge \frac{\delta L}{\delta \vartheta^\alpha} + \delta_{\mathrm{susy}} \Gamma_\alpha^\star \wedge \frac{\delta L}{\delta \Gamma_\alpha^\star} + (2 \overline{D\sigma} + c \bar{\sigma} \gamma) \wedge \frac{\delta L}{\delta \overline{\Psi}},$	$\delta_{\mathrm{susy}} L = i \bar{\sigma} \Psi \gamma^\alpha \wedge \frac{\delta L}{\delta \vartheta^\alpha} + \delta_{\mathrm{susy}} \Gamma_\alpha^\star \wedge \frac{\delta L}{\delta \Gamma_\alpha^\star} + (2 \overline{D\sigma} + c \bar{\sigma} \gamma) \wedge \frac{\delta L}{\delta \overline{\Psi}},$
Conf. MathPix confidence: 0.994 0.5-0.9 <0.5 no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0081 (3.1.6)	053	■ 0.994 ● N +0	$\begin{aligned} \delta L &= \delta \varphi \wedge \delta L_\varphi + \delta(d\varphi) \wedge \pi \\ &= \delta \varphi \wedge \delta L_\varphi + d(\delta \varphi) \wedge \pi \\ &= \delta \varphi \wedge (\delta L_\varphi + d\pi) + d(\delta \varphi \wedge \pi), \end{aligned}$	$\begin{aligned} \delta L &= \delta \varphi \wedge \delta L_\varphi + \delta(d\varphi) \wedge \pi \\ &= \delta \varphi \wedge \delta L_\varphi + d(\delta \varphi) \wedge \pi \\ &= \delta \varphi \wedge (\delta L_\varphi + d\pi) + d(\delta \varphi \wedge \pi), \end{aligned}$	$\begin{aligned} \delta L &= \delta \varphi \wedge \delta L_\varphi + \delta(d\varphi) \wedge \pi \\ &= \delta \varphi \wedge \delta L_\varphi + d(\delta \varphi) \wedge \pi \\ &= \delta \varphi \wedge (\delta L_\varphi + d\pi) + d(\delta \varphi \wedge \pi), \end{aligned}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0509 (7.7.27)	165	■ 0.994 ● K +0	$D = d + [\Gamma, \quad]$	$D = d + [\Gamma, \quad]$	$D = d + [\Gamma, \quad]$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0387 (6.4.2)	130	■ 0.994 ● K +0	$\left. \begin{aligned} \Psi_{\perp} &:= d t \wedge \Psi_{\perp}; \quad \Psi_{\perp} := n \rfloor \Psi, \\ \Psi_{\perp} &:= n \rfloor \Psi, \end{aligned} \right\}$	$\left. \begin{aligned} \Psi_{\perp} &:= d t \wedge \Psi_{\perp}; \quad \Psi_{\perp} := n \rfloor \Psi, \\ \Psi_{\perp} &:= n \rfloor \Psi, \end{aligned} \right\}$	$\left. \begin{aligned} \Psi_{\perp} &:= d t \wedge \Psi_{\perp}; \quad \Psi_{\perp} := n \rfloor \Psi, \\ \Psi_{\perp} &:= n \rfloor \Psi, \end{aligned} \right\}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0646 (10.3.3)	214	■ 0.995 ● N +0	$\Gamma_{\alpha}^{\star} = (-1)^s \frac{\varepsilon}{2\ell} \vartheta_{\alpha},$	$\Gamma_{\alpha}^{\star} = (-1)^s \frac{\varepsilon}{2\ell} \vartheta_{\alpha},$	$\Gamma_{\alpha}^{\star} = (-1)^s \frac{\varepsilon}{2\ell} \vartheta_{\alpha},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0806 (11.4.18)	255	■ 0.995 ● W +1	$\overrightarrow{J^2} \chi_{\kappa}^m = j(j+1) \chi_{\kappa}^m$	$\overrightarrow{J^2} \chi_{\kappa}^m = j(j+1) \chi_{\kappa}^m$	$\overrightarrow{J^2} \chi_{\kappa}^m = j(j+1) \chi_{\kappa}^m$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0995 (14.3.8)	321	■ 0.995 ● W -1	$L_{\text{anom.}}=\frac{\widehat{\ell}}{2}\bar{\psi}\gamma\wedge\gamma\wedge{}^*\Omega\psi=\frac{\ell^*}{4\sqrt{\alpha_g}}\bar{\psi}F_{\mu\nu}\gamma^5\left[\gamma^\mu,\gamma^\nu\right]\psi\eta.$	$L_{\text{anom.}}=\frac{\widehat{\ell}}{2}\bar{\psi}\gamma\wedge\gamma\wedge{}^*\Omega\psi=\frac{\ell^*}{4\sqrt{\alpha_g}}\bar{\psi}F_{\mu\nu}\gamma^5\left[\gamma^\mu,\gamma^\nu\right]\psi\eta.$	$L_{\text{anom.}}=\frac{\widehat{\ell}}{2}\bar{\psi}\gamma\wedge\gamma\wedge{}^*\Omega\psi=\frac{\ell^*}{4\sqrt{\alpha_g}}\bar{\psi}F_{\mu\nu}\gamma^5\left[\gamma^\mu,\gamma^\nu\right]\psi\eta.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0098 (3.2.8)	056	■ 0.995 ● K +0	$\Omega\rightarrow{}^{G^{-1}}\Omega=G^{-1}\Omega G.$	$\Omega\rightarrow{}^{G^{-1}}\Omega=G^{-1}\Omega G.$	$\Omega\rightarrow{}^{G^{-1}}\Omega=G^{-1}\Omega G.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0930 (14.1.11)	304	■ 0.995 ● N +0	$\mathrm{R}=\{ \}^*\backslash\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)=0$	$\mathrm{R}=\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)=0$	$\mathrm{R}=\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ1033 (15.1.10)	338	■ 0.995 ● N -1	$\frac{1}{2}\{ \}^*\backslash\left(\vartheta^f\wedge\Omega^f\right)+\Lambda_{\mathrm{eff}}^f\vartheta^f=0$	$\frac{1}{2}\left(\vartheta^f\wedge\Omega^f\right)+\Lambda_{\mathrm{eff}}^f\vartheta^f=0$	$\frac{1}{2}\left(\vartheta^f\wedge\Omega^f\right)+\Lambda_{\mathrm{eff}}^f\vartheta^f=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0207 (4.1.21)	083	■ 0.995 ● N +0	$\delta:\left\{\begin{array}{llll}A(n,\mathbb{R})&\times A(M)&\longrightarrow&A(M)\\ \downarrow&\downarrow&&\downarrow\\ (g,a)&(e(m),x)&=(\mathbf{g.e}(m),g^{-1}(x-a)).\end{array}\right.$	$\delta:\left\{\begin{array}{llll}A(n,\mathbb{R})&\times A(M)&\longrightarrow&A(M)\\ \downarrow&\downarrow&&\downarrow\\ (g,a)&(e(m),x)&=(\mathbf{g.e}(m),g^{-1}(x-a)).\end{array}\right.$	$\delta:\left\{\begin{array}{llll}A(n,\mathbb{R})&\times A(M)&\longrightarrow&A(M)\\ \downarrow&\downarrow&&\downarrow\\ (g,a)&(e(m),x)&=(g.e(m),g^{-1}(x-a)).\end{array}\right.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0658 (10.3.15)	217	■0.995 ● N +0	$\{ \}^{\{ * \}} D \left(\vartheta^{\alpha} \wedge H_{\alpha} \right) - \{ \}^{\{ * \}} \left(T^{\alpha} \wedge H_{\alpha} \right) = \frac{\gamma}{\delta \ell^2} \{ \}^{\{ * \}} (\vartheta^{\alpha} \wedge T_{\alpha}) + \frac{1}{\delta \ell} \{ \}^{\{ * \}} (\eta^{\alpha} \wedge H_{\alpha}) .$	${}^* D \left(\vartheta^{\alpha} \wedge H_{\alpha} \right) - {}^* \left(T^{\alpha} \wedge H_{\alpha} \right) = \frac{\gamma}{\delta \ell^2} {}^* (\vartheta^{\alpha} \wedge T_{\alpha}) + \frac{1}{\delta \ell} {}^* (\eta^{\alpha} \wedge H_{\alpha}) .$	${}^* D (\vartheta^{\alpha} \wedge H_{\alpha}) - {}^* (T^{\alpha} \wedge H_{\alpha}) = \frac{\gamma}{\delta \ell^2} {}^* (\vartheta^{\alpha} \wedge T_{\alpha}) + \frac{1}{\delta \ell} {}^* (\eta^{\alpha} \wedge H_{\alpha}) .$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0154 (3.5.2)	066	■0.995 ● N +0	$\stackrel{\scriptscriptstyle (-)}{\sigma}_{\mu \nu} := \left\{ \begin{array}{l} \sigma^{ij} = \frac{1}{2} \varepsilon^{ijk} \sigma_k \\ \sigma^{i0} = (-) \frac{1}{2} \sigma^i \end{array} \right i, j, k = 1, 2, 3$	$\stackrel{\scriptscriptstyle (-)}{\sigma}_{\mu \nu} := \left\{ \begin{array}{l} \sigma^{ij} = \frac{1}{2} \varepsilon^{ijk} \sigma_k \\ \sigma^{i0} = (-) \frac{1}{2} \sigma^i \end{array} \right i, j, k = 1, 2, 3$	$\stackrel{\scriptscriptstyle (-)}{\sigma}_{\mu \nu} := \left\{ \begin{array}{l} \sigma^{ij} = \frac{1}{2} \varepsilon^{ijk} \sigma_k \\ \sigma^{i0} = (-) \frac{1}{2} \sigma^i \end{array} \right i, j, k = 1, 2, 3$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0928 (14.1.9)	304	■0.995 ● W -1	$\begin{aligned} \Sigma_{\mu \nu}^{\mathrm{YM}} &= \frac{1}{\alpha_g} \mathrm{Tr} \left(F_{\mu \alpha} F_{\nu}^{\alpha} - \frac{1}{4} g_{\mu \nu} F_{\alpha \beta} F^{\alpha \beta} \right) \\ &= \frac{1}{2 \alpha_g} \mathrm{Tr} (F_{\mu \alpha} F_{\nu}^{\alpha} - (-1)^s {}^* F_{\mu \alpha} {}^* F_{\nu}^{\alpha}), \end{aligned}$	$\Sigma_{\mu \nu}^{\mathrm{YM}} = \frac{1}{\alpha_g} \mathrm{Tr} \left(F_{\mu \alpha} F_{\nu}^{\alpha} - \frac{1}{4} g_{\mu \nu} F_{\alpha \beta} F^{\alpha \beta} \right)$ $= \frac{1}{2 \alpha_g} \mathrm{Tr} (F_{\mu \alpha} F_{\nu}^{\alpha} - (-1)^s {}^* F_{\mu \alpha} {}^* F_{\nu}^{\alpha}),$	$\Sigma_{\mu \nu}^{\mathrm{YM}} = \frac{1}{\alpha_g} \mathrm{Tr} (F_{\mu \alpha} F_{\nu}^{\alpha} - \frac{1}{4} g_{\mu \nu} F_{\alpha \beta} F^{\alpha \beta})$ $= \frac{1}{2 \alpha_g} \mathrm{Tr} (F_{\mu \alpha} F_{\nu}^{\alpha} - (-1)^s {}^* F_{\mu \alpha} {}^* F_{\nu}^{\alpha}),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0006 (2.2.1)	033	■0.996 ● K +0	$\left. e_{\beta} \right] \vartheta^{\alpha} = \delta^{\alpha}_{\beta} .$	$e_{\beta} \rfloor \vartheta^{\alpha} = \delta^{\alpha}_{\beta} .$	$e_{\beta} \rfloor \vartheta^{\alpha} = \delta^{\alpha}_{\beta} .$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0284 (4.4.13)	099	■0.996 ● W +2	$\begin{aligned} D[\Theta, \vartheta] &= [D\Theta, \vartheta] + [\Theta, D\vartheta] \\ &= [[\vartheta, \Omega^g], \vartheta] + [\Theta, \Theta] \\ &= \vartheta \wedge \Omega^g \wedge \vartheta - \Omega^g \wedge \vartheta \wedge \vartheta + \vartheta \wedge \vartheta \wedge \Omega^g - \vartheta \wedge \Omega^g \wedge \vartheta \\ &= [\vartheta \wedge \vartheta, \Omega^g] . \end{aligned}$	$D[\Theta, \vartheta] = [D\Theta, \vartheta] + [\Theta, D\vartheta]$ $= [[\vartheta, \Omega^g], \vartheta] + [\Theta, \Theta]$ $= \vartheta \wedge \Omega^g \wedge \vartheta - \Omega^g \wedge \vartheta \wedge \vartheta + \vartheta \wedge \vartheta \wedge \Omega^g - \vartheta \wedge \Omega^g \wedge \vartheta$ $= [\vartheta \wedge \vartheta, \Omega^g] .$	$D[\Theta, \vartheta] = [D\Theta, \vartheta] + [\Theta, D\vartheta]$ $= [[\vartheta, \Omega^g], \vartheta] + [\Theta, \Theta]$ $= \vartheta \wedge \Omega^g \wedge \vartheta - \Omega^g \wedge \vartheta \wedge \vartheta + \vartheta \wedge \vartheta \wedge \Omega^g - \vartheta \wedge \Omega^g \wedge \vartheta$ $= [\vartheta \wedge \vartheta, \Omega^g].$

Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value
Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0220 (4.2.13)	086	■ 0.996 ● K +0	$\Omega^{\mathrm{L}} \rightarrow^{A^{-1}} \Omega^{\mathrm{L}} = G^{-1} \Omega^{\mathrm{L}} G$	$\Omega^{\mathrm{L}} \rightarrow^{A^{-1}} \Omega^{\mathrm{L}} = G^{-1} \Omega^{\mathrm{L}} G$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0460 (7.2.12)	152	■ 0.996 ● W +0	$g_{\mathrm{i\ j}} \rightarrow \bar{g}_{\mathrm{i\ j}} = \phi^{\mathrm{L}} g_{\mathrm{i\ j}},$	$g_{ij} \rightarrow \bar{g}_{ij} = \phi^L g_{ij},$	$g_{ij} \rightarrow \bar{g}_{ij} = \phi^L g_{ij},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0664 (10.3.21)	218	■ 0.996 ● N +0	$\square \vartheta_{\alpha} = \frac{\ell}{\gamma} D^* [\tau_{\alpha}^* - S_{\alpha} - \delta \ell (\Sigma_{\alpha} + E_{\alpha})] + \frac{\ell}{\gamma} D^* (D^* H_{\alpha}^* - \delta \ell D^* H_{\alpha}).$	$\square \vartheta_{\alpha} = \frac{\ell}{\gamma} D^* [\tau_{\alpha}^* - S_{\alpha} - \delta \ell (\Sigma_{\alpha} + E_{\alpha})] + \frac{\ell}{\gamma} D^* (D^* H_{\alpha}^* - \delta \ell D^* H_{\alpha}).$	$\square \vartheta_{\alpha} = \frac{\ell}{\gamma} D^* [\tau_{\alpha}^* - S_{\alpha} - \delta \ell (\Sigma_{\alpha} + E_{\alpha})] + \frac{\ell}{\gamma} D^* (D^* H_{\alpha}^* - \delta \ell D^* H_{\alpha}).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0977 (14.2.18)	315	■ 0.996 ● N +0	$\hat{L}_{\mathrm{L}} = -\frac{1}{2\ell^{*2}} \mathrm{Tr}[\hat{\alpha} \wedge \hat{\alpha}, \quad \hat{\vartheta} \wedge \cdots \wedge \hat{\vartheta}]$	$\hat{L}_{\parallel} = -\frac{1}{2\ell^{*2}} \mathrm{Tr}[\hat{\alpha} \wedge \hat{\alpha}, \quad \hat{\vartheta} \wedge \cdots \wedge \hat{\vartheta}]$	$\hat{L}_{\parallel} = -\frac{1}{2\ell^{*2}} \mathrm{Tr}[\hat{\alpha} \wedge \hat{\alpha}, \quad \hat{\vartheta} \wedge \cdots \wedge \hat{\vartheta}]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0152 (3.4.23)	066	■ 0.996 ● C -4	$L_{\mathrm{BI}}(\mathcal{F}, {}^*\mathcal{F}) = \sqrt{1 - 2\mathcal{F} - ({}^*\mathcal{F})^2} - 1.$	$L_{\mathrm{BI}}(\mathcal{F}, {}^*\mathcal{F}) = \sqrt{1 - 2\mathcal{F} - ({}^*\mathcal{F})^2} - 1.$	$L_{\mathrm{BI}}(\mathcal{F}, {}^*\mathcal{F}) = \sqrt{1 - 2\mathcal{F} - ({}^*\mathcal{F})^2} - 1.$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5-0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0250 (4.3.15)	092	■ 0.996 ● N +0	$\{ \}^{\{ * \}} \omega^{\{ g \}} = \frac{1}{2} \omega^{\{ \alpha \beta \}} L_{\{ \alpha \beta \}} L_{\{ \alpha \beta \}} = \frac{1}{2} \Gamma_{\{ i \}}^{\{ \cdot \}} \omega^{\{ \alpha \beta \}} L_{\alpha \beta} \otimes dx^i,$	${}^* \omega^g = \frac{1}{2} \omega^{\alpha \beta} L_{\alpha \beta} = \frac{1}{2} \Gamma_i^{\cdot \alpha \beta} L_{\alpha \beta} \otimes dx^i,$	${}^* \omega^g = \frac{1}{2} \omega^{\alpha \beta} L_{\alpha \beta} = \frac{1}{2} \Gamma_i^{\cdot \alpha \beta} L_{\alpha \beta} \otimes dx^i,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0376 (6.2.27)	128	■ 0.996 ● N +0	$\stackrel{\{ \}}{\vartheta}^{\alpha} = \pm \frac{i}{2} \ell^2 g^{\alpha \beta} \stackrel{\{ \}}{\Pi}_{\beta},$	$\stackrel{(\pm)}{D} \vartheta^{\alpha} = \pm \frac{i}{2} \ell^2 g^{\alpha \beta} \stackrel{(\pm)}{\Pi}_{\beta},$	$\stackrel{(\pm)}{D} \vartheta^{\alpha} = \pm \frac{i}{2} \ell^2 g^{\alpha \beta} \stackrel{(\pm)}{\Pi}_{\beta},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0367 (6.2.18)	126	■ 0.997 ● N +0	$R_{\{ \alpha \beta \}} = \xi^* R_{\alpha \beta}^{(*)} + (\gamma / \ell^2) [\vartheta_{\alpha} \wedge \vartheta_{\beta} \pm i^* (\vartheta_{\alpha} \wedge \vartheta_{\beta})],$	$R_{\alpha \beta} = \xi^* R_{\alpha \beta}^{(*)} + (\gamma / \ell^2) [\vartheta_{\alpha} \wedge \vartheta_{\beta} \pm i^* (\vartheta_{\alpha} \wedge \vartheta_{\beta})],$	$R_{\alpha \beta} = \xi^* R_{\alpha \beta}^{(*)} + (\gamma / \ell^2) [\vartheta_{\alpha} \wedge \vartheta_{\beta} \pm i^* (\vartheta_{\alpha} \wedge \vartheta_{\beta})],$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0371 (6.2.22)	127	■ 0.997 ● W +0	$\stackrel{\{ + \}}{\Pi}^{\{ \beta \}} \wedge \stackrel{\{ - \}}{\Pi}_{\{ \beta \}} = H^{\{ \beta \}} \wedge H_{\{ \beta \}} + \ell^{-4} T^{\beta} \wedge T_{\beta} = 0.$	$\stackrel{(+)}{\Pi}^{\beta} \wedge \stackrel{(-)}{\Pi}_{\beta} = H^{\beta} \wedge H_{\beta} + \ell^{-4} T^{\beta} \wedge T_{\beta} = 0.$	$\stackrel{(+)}{\Pi}^{\beta} \wedge \stackrel{(-)}{\Pi}_{\beta} = H^{\beta} \wedge H_{\beta} + \ell^{-4} T^{\beta} \wedge T_{\beta} = 0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0602 (9.2.1)	198	■ 0.997 ● W +0	$\begin{aligned} p_1(M^4) &= \frac{1}{2(2\pi)^2} \int_{M^4} \text{Tr}(\Omega^L \wedge \Omega^L) \\ &= \frac{(-1)^{s/2}}{(4\pi)^2} \int_{M^4} R_{\alpha \beta \gamma \delta}^* R^{\alpha \beta \gamma \delta} \sqrt{ g } d^4 x \end{aligned}$	$p_1(M^4) = \frac{1}{2(2\pi)^2} \int_{M^4} \text{Tr}(\Omega^L \wedge \Omega^L) = \frac{(-1)^{s/2}}{(4\pi)^2} \int_{M^4} R_{\alpha \beta \gamma \delta}^* R^{\alpha \beta \gamma \delta} \sqrt{ g } d^4 x$	$p_1(M^4) = \frac{1}{2(2\pi)^2} \int_{M^4} \text{Tr}(\Omega^L \wedge \Omega^L) = \frac{(-1)^{s/2}}{(4\pi)^2} \int_{M^4} R_{\alpha \beta \gamma \delta}^* R^{\alpha \beta \gamma \delta} \sqrt{ g } d^4 x$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0717 (10.9.3)	229	■ 0.997 ● W +0	$\backslash \mathrm{Sigma}_{\{\alpha\}}:=\frac{\delta L_{\Psi}}{\delta \vartheta^{\alpha}}\backslash \mathrm{vartheta}^{\{\alpha\}}=\frac{\partial L_{\Psi}}{\partial \vartheta_{\alpha}}\backslash \mathrm{vartheta}_{\{\alpha\}}+D\frac{\partial L_{\Psi}}{\partial T^{\alpha}}T^{\{\alpha\}},$	$\Sigma_{\alpha}:=\frac{\delta L_{\Psi}}{\delta \vartheta^{\alpha}}=\frac{\partial L_{\Psi}}{\partial \vartheta_{\alpha}}+D\frac{\partial L_{\Psi}}{\partial T^{\alpha}},$	$\Sigma_{\alpha}:=\frac{\delta L_{\Psi}}{\delta \vartheta^{\alpha}}=\frac{\partial L_{\Psi}}{\partial \vartheta_{\alpha}}+D\frac{\partial L_{\Psi}}{\partial T^{\alpha}},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0766 (11.1.33)	243	■ 0.997 ● N +0	$\backslash \mathrm{stackrel{*}{\sigma}}\backslash \mathrm{tilde{\omega}}=\frac{1}{2}\backslash \mathrm{\omega}^{\{\alpha\beta\}}\backslash \mathrm{\sigma}_{\alpha\beta}=\frac{1}{2}\backslash \mathrm{\Gamma}_i^{\alpha\beta}\backslash \mathrm{\sigma}_{\alpha\beta}\otimes dx^i=:\backslash \mathrm{\Gamma}_idx^i.$	$\sigma^*\tilde{\omega}=\frac{1}{2}\omega^{\alpha\beta}\sigma_{\alpha\beta}=\frac{1}{2}\Gamma_i^{\alpha\beta}\sigma_{\alpha\beta}\otimes dx^i=:\Gamma_idx^i.$	$\sigma^*\tilde{\omega}=\frac{1}{2}\omega^{\alpha\beta}\sigma_{\alpha\beta}=\frac{1}{2}\Gamma_i^{\alpha\beta}\sigma_{\alpha\beta}\otimes dx^i=:\Gamma_idx^i.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ1003 (14.3.16)	323	■ 0.997 ● K +0	$D\,D=\frac{1}{2}\backslash \mathrm{D},\,D\backslash +\frac{1}{2}\backslash \mathrm{D},\,D=\frac{1}{2}\backslash \mathrm{D},\,D\backslash +\frac{1}{2}\backslash \mathrm{widetilde{\Omega}}$	$DD=\frac{1}{2}\{D,D\}+\frac{1}{2}[D,D]=\frac{1}{2}\{D,D\}+\frac{1}{2}\tilde{\Omega}$	$D\,D=\frac{1}{2}\{D,D\}+\frac{1}{2}[D,D]=\frac{1}{2}\{D,D\}+\frac{1}{2}\tilde{\Omega}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0225 (4.2.18)	087	■ 0.997 ● N +0	$\backslash \mathrm{\omega}^{\{\mathrm{T}\}}=\backslash \mathrm{widetilde{\widetilde{\vartheta}}}\backslash \mathrm{widetilde{\widetilde{\vartheta}}}-D\backslash \mathrm{widetilde{\widetilde{\varphi}}}$	$\omega^{\mathrm{T}}=\widetilde{\vartheta}-D\widetilde{\varphi}$	$\omega^{\mathrm{T}}=\widetilde{\vartheta}-D\widetilde{\varphi}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0048 (2.6.11)	043	■ 0.997 ● N +0	$D\,D\,D\,b\,\equiv\,0\,.$	$DDDb\equiv 0.$	$DDDb\equiv 0.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0099 (3.2.9)	056	■ 0.997 ● K +0	$\begin{aligned} \{ \}^{\Delta} \omega &:= \{ \}^{G^{-1}} \omega - \omega = G^{-1}[\omega, G] - G^{-1}dG \\ \omega \wedge &= G^{-1}[\omega, G] - G^{-1}dG \quad \& \quad = -G^{-1}D \\ G &= -G^{-1}D \quad . \end{aligned}$	$\Delta \omega := {}^{G^{-1}}\omega - \omega = G^{-1}[\omega, G] - G^{-1}dG \\ = -G^{-1}DG = -{}^{G^{-1}}D.$	${}^{\Delta}\omega := {}^{G^{-1}}\omega - \omega = G^{-1}[\omega, G] - G^{-1}dG \\ = -G^{-1}DG = -{}^{G^{-1}}D.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0219 (4.2.12)	086	■ 0.997 ● N +0	$\widetilde{\omega} \rightarrow {}^{A^{-1}}\widetilde{\omega} = A^{-1}\widetilde{\omega}A$	$\widetilde{\omega} \rightarrow {}^{A^{-1}}\widetilde{\omega} = A^{-1}\widetilde{\omega}A$	$\widetilde{\omega} \rightarrow {}^{A^{-1}}\widetilde{\omega} = A^{-1}\widetilde{\omega}A$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0411 (6.4.31)	136	■ 0.997 ● W +0	$n_{\mathrm{NY}} := \int_{R^4} dC_{\mathrm{TT}} = \int_{S^3_\infty} \underline{C}_{\mathrm{TT}} = 3 \mathrm{Vol}(S^3) k = 6\pi^2 k.$	$n_{\mathrm{NY}} := \int_{R^4} dC_{\mathrm{TT}} = \int_{S^3_\infty} \underline{C}_{\mathrm{TT}} = 3 \mathrm{Vol}(S^3) k = 6\pi^2 k.$	$n_{\mathrm{NY}} := \int_{R^4} dC_{\mathrm{TT}} = \int_{S^3_\infty} \underline{C}_{\mathrm{TT}} = 3 \mathrm{Vol}(S^3) k = 6\pi^2 k.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0492 (7.7.10)	161	■ 0.997 ● N -1	$H_{\alpha\beta} := -\frac{\partial L}{\partial R^{\alpha\beta}} = \tilde{H}_{\alpha\beta} + \frac{1}{2}R^{\mu\nu} \wedge (\partial \tilde{H}_{\mu\nu} / \partial R^{\alpha\beta})$	$H_{\alpha\beta} := -\frac{\partial L}{\partial R^{\alpha\beta}} = \tilde{H}_{\alpha\beta} + \frac{1}{2}R^{\mu\nu} \wedge (\partial \tilde{H}_{\mu\nu} / \partial R^{\alpha\beta})$	$H_{\alpha\beta} := -\frac{\partial L}{\partial R^{\alpha\beta}} = \tilde{H}_{\alpha\beta} + \frac{1}{2}R^{\mu\nu} \wedge (\partial \tilde{H}_{\mu\nu} / \partial R^{\alpha\beta})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0877 (13.3.6)	285	■ 0.997 ● N +0	$\left[P_{\alpha}, P_{\beta} \right] = \frac{1}{\ell^2} L^{\beta}_{\alpha}.$	$\left[P_{\alpha}, P_{\beta} \right] = \frac{1}{\ell^2} L^{\beta}_{\alpha}.$	$\left[P_{\alpha}, P_{\beta} \right] = \frac{1}{\ell^2} L^{\beta}_{\alpha}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0962 (14.2.2)	312	■ 0.997 ● N +0	$\stackrel{\circ}{\vartheta}=\varphi_j d\xi^j,$	$\stackrel{\circ}{\vartheta}=\varphi_j d\xi^j,$	$\stackrel{\circ}{\vartheta}=\varphi_j d\xi^j,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ1060 (16.1.1)	354	■ 0.997 ● N -1	$R_{ij}^f-\frac{1}{2}\mathrm{f}_{ij}\left(R^f-2\Lambda_{\mathrm{eff}}\right)=\ell^2\mathrm{T}_{ij}$	$R_{ij}^f-\frac{1}{2}\mathrm{f}_{ij}\left(R^f-2\Lambda_{\mathrm{eff}}\right)=\ell^2\mathrm{T}_{ij}$	$R_{ij}^f-\frac{1}{2}\mathrm{f}_{ij}(R^f-2\Lambda_{\mathrm{eff}})=\ell^2\mathrm{T}_{ij}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0976 (14.2.17)	315	■ 0.997 ● W -2	$\begin{aligned} \hat{L}_{\mathrm{W}.} &= \frac{1}{2} \operatorname{Tr}(\hat{\Omega} \wedge \underbrace{\hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}}_{2+K \text{ termms}}) \\ &= \hat{L}_{\mathrm{W}.}^g + \hat{L}_{\mathrm{W}.}^G + \ell^{*2} \hat{L}_{\parallel} - \frac{1}{2} d \operatorname{Tr}(\hat{\alpha} \wedge \hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}) \end{aligned}$	$\hat{L}_{\mathrm{W}.} = \frac{1}{2} \operatorname{Tr}(\hat{\Omega} \wedge \underbrace{\hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}}_{2+K \text{ termms}})$ $= \hat{L}_{\mathrm{W}.}^g + \hat{L}_{\mathrm{W}.}^G + \ell^{*2} \hat{L}_{\parallel} - \frac{1}{2} d \operatorname{Tr}(\hat{\alpha} \wedge \hat{\vartheta} \wedge \dots \wedge \hat{\vartheta})$	$\hat{L}_{\mathrm{W}.} = \frac{1}{2} \operatorname{Tr}(\hat{\Omega} \wedge \underbrace{\hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}}_{2+K \text{ termms}})$ $= \hat{L}_{\mathrm{W}.}^g + \hat{L}_{\mathrm{W}.}^G + \ell^{*2} \hat{L}_{\parallel} - \frac{1}{2} d \operatorname{Tr}(\hat{\alpha} \wedge \hat{\vartheta} \wedge \dots \wedge \hat{\vartheta})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0466 (7.3.5)	153	■ 0.997 ● W +0	$G_{\alpha}^{\{\}}:=\frac{1}{2}R^{\{\}\beta\gamma}\wedge\eta_{\alpha\beta\gamma},\quad G_{ij}:=\operatorname{Ric}_{ij}^{\{\}}-\frac{1}{2}g_{ij},$	$G_{\alpha}^{\{\}}:=\frac{1}{2}R^{\{\}\beta\gamma}\wedge\eta_{\alpha\beta\gamma},\quad G_{ij}:=\operatorname{Ric}_{ij}^{\{\}}-\frac{1}{2}g_{ij},$	$G_{\alpha}^{\{\}}:=\frac{1}{2}R^{\{\}\beta\gamma}\wedge\eta_{\alpha\beta\gamma},\quad G_{ij}:=\operatorname{Ric}_{ij}^{\{\}}-\frac{1}{2}g_{ij},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0105 (3.3.6)	058	■ 0.998 ● N -1	$H:=\frac{\partial L}{\partial A}=-\frac{\partial L}{\partial F}\quad\text{and}\quad j:=\frac{\delta L_{\mathrm{mat}}}{\delta A},$	$H:=-\frac{\partial L}{\partial dA}=-\frac{\partial L}{\partial F}\quad\text{and}\quad j:=\frac{\delta L_{\mathrm{mat}}}{\delta A},$	$H:=-\frac{\partial L}{\partial dA}=-\frac{\partial L}{\partial F}\quad\text{and}\quad j:=\frac{\delta L_{\mathrm{mat}}}{\delta A},$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0136 (3.4.7)	063	■0.998 ●N +2	$\begin{aligned} \delta \Omega &= \delta(d\omega - \omega \wedge \omega) = \delta d\omega - \delta\omega \wedge \omega - \omega \wedge \delta\omega \\ &= d\delta\omega - [\omega, \delta\omega] = D\delta\omega. \end{aligned}$	$\delta\Omega = \delta(d\omega - \omega \wedge \omega) = \delta d\omega - \delta\omega \wedge \omega - \omega \wedge \delta\omega$ $= d\delta\omega - [\omega, \delta\omega] = D\delta\omega.$	$\delta\Omega = \delta(d\omega - \omega \wedge \omega) = \delta d\omega - \delta\omega \wedge \omega - \omega \wedge \delta\omega$ $= d\delta\omega - [\omega, \delta\omega] = D\delta\omega.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0390 (6.4.5)	131	■0.998 ●W +1	$\mathscr{H} = n \rfloor L - \ell_n \underline{\Psi} \wedge \left(\frac{\partial L}{\partial d\Psi} \right).$	$\mathcal{H} = n \rfloor L - \ell_n \underline{\Psi} \wedge \left(\frac{\partial L}{\partial d\Psi} \right).$	$\mathcal{H} = n \rfloor L - \ell_n \underline{\Psi} \wedge \left(\frac{\partial L}{\partial d\Psi} \right).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0594 (9.1.12)	195	■0.998 ●K +0	$\bar{\Omega}^{\{\}} = 0.$	$\bar{\Omega}^{\{\}} = 0.$	$\bar{\Omega}^{\{\}} = 0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ1012 (14.4.7)	326	■0.998 ●W -1	$\begin{aligned} S_{\mathrm{A}} &:= 4\pi \left(r_+^2 + \frac{J(J+1)\hbar^2}{M^2 c^2} \right) \\ &= \frac{8\pi\hbar^2}{M^{*4}c^2} \left(M^2 + \frac{\alpha_e}{2} M^{*2} \mathrm{Tr} \bar{Q}^2 + M \sqrt{M^2 - J(J+1) \frac{M^{*4}}{M^2} - \alpha_e M^{*2} \mathrm{Tr} \bar{Q}^2} \right) \end{aligned}$	$S_{\mathrm{A}} := 4\pi \left(r_+^2 + \frac{J(J+1)\hbar^2}{M^2 c^2} \right)$ $= \frac{8\pi\hbar^2}{M^{*4}c^2} \left(M^2 + \frac{\alpha_e}{2} M^{*2} \mathrm{Tr} \bar{Q}^2 + M \sqrt{M^2 - J(J+1) \frac{M^{*4}}{M^2} - \alpha_e M^{*2} \mathrm{Tr} \bar{Q}^2} \right)$	$S_{\mathrm{A}} := 4\pi \left(r_+^2 + \frac{J(J+1)\hbar^2}{M^2 c^2} \right)$ $= \frac{8\pi\hbar^2}{M^{*4}c^2} \left(M^2 + \frac{\alpha_e}{2} M^{*2} \mathrm{Tr} \bar{Q}^2 + M \sqrt{M^2 - J(J+1) \frac{M^{*4}}{M^2} - \alpha_e M^{*2} \mathrm{Tr} \bar{Q}^2} \right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0073 (2.8.16)	049	■0.998 ●N +0	$\begin{aligned} & d \stackrel{\infty}{\omega} - \stackrel{\infty}{\omega} \wedge \stackrel{\infty}{\omega} \\ &= -d \left(G^{-1} \right) \wedge dG - G^{-1} ddG - \left(G^{-1} dG \right) \wedge \left(G^{-1} dG \right) \\ &= -d \left(G^{-1} \right) \wedge dG + G^{-1} G d \left(G^{-1} \right) \wedge dG = 0 \end{aligned}$	$d \stackrel{\infty}{\omega} - \stackrel{\infty}{\omega} \wedge \stackrel{\infty}{\omega}$ $= -d \left(G^{-1} \right) \wedge dG - G^{-1} ddG - \left(G^{-1} dG \right) \wedge \left(G^{-1} dG \right)$ $= -d \left(G^{-1} \right) \wedge dG + G^{-1} G d \left(G^{-1} \right) \wedge dG = 0$	$d \stackrel{\infty}{\omega} - \stackrel{\infty}{\omega} \wedge \stackrel{\infty}{\omega}$ $= -d \left(G^{-1} \right) \wedge dG - G^{-1} ddG - \left(G^{-1} dG \right) \wedge \left(G^{-1} dG \right)$ $= -d \left(G^{-1} \right) \wedge dG + G^{-1} G d \left(G^{-1} \right) \wedge dG = 0$
Conf. MathPix confidence: ■≥0.9 ■0.5–0.9 ■<0.5 —no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0214 (4.2.6)	086	■ 0.998 ● N +0	$\mathrm{mathscr{G}}(n,\mathrm{mathbb{R}}):=C^{\infty}\!\left(A(M)\times_{\mathrm{Ad}}\mathrm{G}\,L(n,\mathrm{mathbb{R}})\right)$	$\mathscr{G}(n,\mathbb{R}):=C^{\infty}(A(M)\times_{\mathrm{Ad}}GL(n,\mathbb{R}))$	$\mathscr{G}(n,\mathbb{R}):=C^{\infty}(A(M)\times_{\mathrm{Ad}}GL(n,\mathbb{R}))$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0656 (10.3.13)	216	■ 0.998 ● N +0	$\left.\left.E_{\alpha}\right _{\star}\right\rfloor L(\star)+(-1)^s\frac{2}{\ell}\left[\left(e_{\alpha}\right\rfloor R^{\star\beta}\right)\wedge\vartheta_{\beta}\right]$	$E_{\alpha}=e_{\alpha}\rfloor L(\star)+(-1)^s\frac{2\gamma}{\ell}\left[\left(e_{\alpha}\rfloor R^{\star\beta}\right)\wedge\vartheta_{\beta}\right]$	$E_{\alpha}=e_{\alpha}\rfloor L(\star)+(-1)^s\frac{2\gamma}{\ell}\left[\left(e_{\alpha}\rfloor R^{\star\beta}\right)\wedge\vartheta_{\beta}\right]$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0813 (11.4.26)	256	■ 0.998 ● K +0	$d\,\rho^{\wedge\{\ast\}}:=e^{\{(\mu-\nu)\,/\,2\}}\,d\,\rho\;.$	$d\rho^{\ast}:=e^{(\mu-\nu)/2}d\rho.$	$d\rho^{\ast}:=e^{(\mu-\nu)/2}d\rho.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0062 (2.8.3)	046	■ 0.998 ● W +1	$\begin{aligned} & \ast\left(d\,\gamma_k\right)=D\,\pi^{\wedge\{\ast\}}\left(\gamma_k\right)=D\,\pi^{\wedge\{\ast\}}\left(\gamma_k\right)\\ & =\frac{(-1)^k}{(2\pi i)^k(k-1)!}\mathrm{Tr}(D\Omega\wedge\Omega\wedge\cdots\wedge\Omega)=0. \end{aligned}$	$\begin{aligned} & \ast(d\gamma_k)=d\pi^{\ast}(\gamma_k)=D\pi^{\ast}(\gamma_k)\\ & =\frac{(-1)^k}{(2\pi i)^k(k-1)!}\mathrm{Tr}(D\Omega\wedge\Omega\wedge\cdots\wedge\Omega)=0. \end{aligned}$	$\begin{aligned} \pi^{\ast}(d\gamma_k)&=d\pi^{\ast}(\gamma_k)=D\pi^{\ast}(\gamma_k)\\ &=\frac{(-1)^k}{(2\pi i)^k(k-1)!}\mathrm{Tr}(D\Omega\wedge\Omega\wedge\cdots\wedge\Omega)=0. \end{aligned}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0426 (6.7.8)	141	■ 0.998 ● K +0	$\mu_{\alpha}=\frac{1}{4}\,\vartheta_{\alpha}\wedge^{\ast}j_5,$	$\mu_{\alpha}=\frac{1}{4}\vartheta_{\alpha}\wedge^{\ast}j_5,$	$\mu_{\alpha}=\frac{1}{4}\vartheta_{\alpha}\wedge^{\ast}j_5,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0069 (2.8.12)	048	■0.998 ● N +0	$\begin{aligned} c_2(M) &= \frac{1}{8\pi^2} \int_M d \operatorname{Tr} \left(\omega \wedge \Omega + \frac{1}{3} \omega \wedge \omega \wedge \omega \right) \\ &= \frac{1}{8\pi^2} \int_{\partial M} \operatorname{Tr} \left(\omega \wedge d\omega - \frac{2}{3} \omega \wedge \omega \wedge \omega \right) . \end{aligned}$	$c_2(M) = \frac{1}{8\pi^2} \int_M d \operatorname{Tr} \left(\omega \wedge \Omega + \frac{1}{3} \omega \wedge \omega \wedge \omega \right) \\ = \frac{1}{8\pi^2} \int_{\partial M} \operatorname{Tr} \left(\omega \wedge d\omega - \frac{2}{3} \omega \wedge \omega \wedge \omega \right) .$	$c_2(M) = \frac{1}{8\pi^2} \int_M d \operatorname{Tr} \left(\omega \wedge \Omega + \frac{1}{3} \omega \wedge \omega \wedge \omega \right) \\ = \frac{1}{8\pi^2} \int_{\partial M} \operatorname{Tr} \left(\omega \wedge d\omega - \frac{2}{3} \omega \wedge \omega \wedge \omega \right) .$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0144 (3.4.15)	064	■0.998 ● N +0	$\iota := \tau + (-1)^{n+s} * \left[\omega, \Pi^{\Omega} \right]$	$\iota := \tau + (-1)^{n+s} * [\omega, \Pi^{\Omega}]$	$\iota := \tau + (-1)^{n+s} * [\omega, \Pi^{\Omega}]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0318 (5.2.2)	110	■0.998 ● N -1	$j := \bar{\psi}^* \gamma \psi = j^\mu \eta_\mu \quad \text{and} \quad j_5 := \bar{\psi}^* \gamma \gamma_5 \psi = j_5^\mu \eta_\mu,$	$j := \bar{\psi}^* \gamma \psi = j^\mu \eta_\mu \quad \text{and} \quad j_5 := \bar{\psi}^* \gamma \gamma_5 \psi = j_5^\mu \eta_\mu,$	$j := \bar{\psi}^* \gamma \psi = j^\mu \eta_\mu \quad \text{and} \quad j_5 := \bar{\psi}^* \gamma \gamma_5 \psi = j_5^\mu \eta_\mu,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0347 (5.4.5)	116	■0.998 ● N +0	$\mu = \frac{4}{3} \kappa \Lambda = \frac{1}{3} (2\kappa)^2 \rho_\Lambda \leq \frac{4}{3} g_{N*} \lambda_* \simeq 0.2$	$\mu = \frac{4}{3} \kappa \Lambda = \frac{1}{3} (2\kappa)^2 \rho_\Lambda \leq \frac{4}{3} g_{N*} \lambda_* \simeq 0.2$	$\mu = \frac{4}{3} \kappa \Lambda = \frac{1}{3} (2\kappa)^2 \rho_\Lambda \leq \frac{4}{3} g_{N*} \lambda_* \simeq 0.2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0500 (7.7.18)	163	■0.998 ● K +0	$\{ \}^{*} \sigma = \frac{1}{2} \sigma_{\alpha \beta} \left(\vartheta^{\alpha} \wedge \vartheta^{\beta} \right) = \frac{1}{2} \sigma_{\alpha \beta} \eta^{\alpha \beta} =: \sigma^{(*)} = i \gamma_5 \sigma .$	$* \sigma = \frac{1}{2} \sigma_{\alpha \beta} * \left(\vartheta^{\alpha} \wedge \vartheta^{\beta} \right) = \frac{1}{2} \sigma_{\alpha \beta} \eta^{\alpha \beta} =: \sigma^{(*)} = i \gamma_5 \sigma .$	$* \sigma = \frac{1}{2} \sigma_{\alpha \beta} * (\vartheta^{\alpha} \wedge \vartheta^{\beta}) = \frac{1}{2} \sigma_{\alpha \beta} \eta^{\alpha \beta} =: \sigma^{(*)} = i \gamma_5 \sigma .$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E08678 (10.3.27)	219	■0.998 ●N +0	$\check{\mathrm{H}}_{\alpha}:=\mathrm{H}_{\alpha}\,\,\mathrm{cong}\,\,\Sigma_{\alpha}\,\,\mathrm{E}_{\alpha} - (-1)^s\,\,\eta_{\alpha\beta\gamma}\,\,\Gamma^{\star\beta}\wedge H^{\gamma}\,\,\Gamma^{\star}\beta\wedge H^{\gamma}$	$\check{\mathcal{E}}_{\alpha}:=dH_{\alpha}\cong\Sigma_{\alpha}+E_{\alpha}-(-1)^s\eta_{\alpha\beta\gamma}\Gamma^{\star\beta}\wedge H^{\gamma}$	$\check{\mathcal{E}}_{\alpha}:=dH_{\alpha}\cong\Sigma_{\alpha}+E_{\alpha}-(-1)^s\eta_{\alpha\beta\gamma}\Gamma^{\star\beta}\wedge H^{\gamma}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E08680 (10.3.37)	221	■0.998 ●W +1	$\theta_{\mathrm{L}}C_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}}\tilde{C}_{\mathrm{L}}=-\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}-\frac{\Lambda}{\ell}\eta+\frac{\theta_{\mathrm{T}}}{2\ell}\vartheta^{\alpha}\wedge T_{\alpha}+\Delta\theta_{\mathrm{L}}C_{\mathrm{L}}+\frac{\Delta\theta_{\mathrm{TL}}}{\ell}d\left(\Gamma^{\star\alpha}\wedge\vartheta_{\alpha}\right)$	$\theta_{\mathrm{L}}C_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}}\tilde{C}_{\mathrm{L}}=-\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}-\frac{\Lambda}{\ell}\eta+\frac{\theta_{\mathrm{T}}}{2\ell}\vartheta^{\alpha}\wedge T_{\alpha}+\Delta\theta_{\mathrm{L}}C_{\mathrm{L}}+\frac{\Delta\theta_{\mathrm{TL}}}{\ell}d\left(\Gamma^{\star\alpha}\wedge\vartheta_{\alpha}\right)$	$\theta_{\mathrm{L}}C_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}}\tilde{C}_{\mathrm{L}}=-\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}-\frac{\Lambda}{\ell}\eta+\frac{\theta_{\mathrm{T}}}{2\ell}\vartheta^{\alpha}\wedge T_{\alpha}+\Delta\theta_{\mathrm{L}}C_{\mathrm{L}}+\frac{\Delta\theta_{\mathrm{TL}}}{\ell}d\left(\Gamma^{\star\alpha}\wedge\vartheta_{\alpha}\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E08700 (10.5.11)	225	■0.998 ●K +0	$[\hat{D},\hat{D}]=\hat{R}+T^{\alpha}\hat{D}_{\alpha}\,.$	$[\hat{D},\hat{D}]=\hat{R}+T^{\alpha}\hat{D}_{\alpha}\,.$	$[\hat{D},\hat{D}]=\hat{R}+T^{\alpha}\hat{D}_{\alpha}\,.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E08820 (11.4.34)	258	■0.998 ●N -1	$f^{\{l\}}(\rho)=\bar{\rho}^{\{l\}}\left(1+\bar{e}^{4l+2}\right)^{-1/2}$	$f^l(\rho)=\bar{\rho}^l\left(1+\bar{e}^{4l+2}\right)^{-1/2}$	$f^l(\rho)=\bar{\rho}^l(1+\bar{e}^{4l+2})^{-1/2}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E01126 (B.38)	373	■0.998 ●N +0	$\partial:C_p(M,\mathbb{R})\rightarrow C_{p-1}(M,\mathbb{R}),$	$\partial:C_p(M,\mathbb{R})\rightarrow C_{p-1}(M,\mathbb{R}),$	$\partial:C_p(M,\mathbb{R})\rightarrow C_{p-1}(M,\mathbb{R}),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E08516 (7.7.34)	166	■0.998 ●N +0	$\widetilde{T}=T-3d\varphi,\quad\mathscr{A}^{\theta}=\mathscr{A}-id\theta.$	$\widetilde{T}=T-3d\varphi,\quad\mathscr{A}^{\theta}=\mathscr{A}-id\theta.$	$\widetilde{T}=T-3d\varphi,\quad\mathscr{A}^{\theta}=\mathscr{A}-id\theta.$
Conf. MathPix confidence: ■≥0.9 ■0.5–0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0967 (14.2.8)	313	■ 0.998 ● N +1	$d \stackrel{\circ}{s}^2 = \stackrel{\circ}{g}_{ij} d\xi^i \otimes_s d\xi^j$		$ds^{\circ 2} = \overset{\circ}{g}_{ij} d\xi^i \otimes_s d\xi^j,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0060 (2.8.1)	046	■ 0.999 ● N +0	$\operatorname{det}\left(1+\frac{i}{2\pi}\Omega\right)=\stackrel{*}{\pi}\left(1+\gamma_1+\cdots+\gamma_m\right)$	$\det\left(1+\frac{i}{2\pi}\Omega\right)=\overset{*}{\pi}\left(1+\gamma_1+\cdots+\gamma_m\right)$	$\det\left(1+\frac{i}{2\pi}\Omega\right)=\overset{*}{\pi}\left(1+\gamma_1+\cdots+\gamma_m\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0071 (2.8.14)	048	■ 0.999 ● K +0	$\stackrel{\infty}{\omega}:= -G^{-1}dG,$	$\overset{\infty}{\omega}:= -G^{-1}dG,$	$\overset{\infty}{\omega}:= -G^{-1}dG,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0235 (4.2.28)	089	■ 0.999 ● K +0	$Q_{\kappa\mu\nu}:= -D_{\kappa}g_{\mu\nu}$	$Q_{\kappa\mu\nu}:= -D_{\kappa}g_{\mu\nu}$	$Q_{\kappa\mu\nu}:= -D_{\kappa}g_{\mu\nu}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0278 (4.4.6)	097	■ 0.999 ● W +0	$\begin{aligned} & \& D\,\Pi^{\mathrm{T}}-E=\Sigma \quad \& D\,\Pi^{\mathrm{L}}+\Pi_{\square}^{\mathrm{T}}=\tau_{\mathrm{s}} \\ & \end{aligned}$	$D\Pi^{\mathrm{T}}-E=\Sigma$ $D\Pi^{\mathrm{L}}+\Pi_{\square}^{\mathrm{T}}=\tau_{\mathrm{s}}$	$D\Pi^{\mathrm{T}}-E=\Sigma$ $D\Pi^{\mathrm{L}}+\Pi_{\square}^{\mathrm{T}}=\tau_{\mathrm{s}}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0313 (4.6.5)	105	■ 0.999 ● N +0	$\backslash \mathrm{Phi}(A,\backslash \gamma)=P\backslash \exp\left[(i/\hbar)\oint A\right],$	$\Phi(A,\gamma)=P\exp[(i/\hbar)\oint A],$	$\Phi(A,\gamma)=P\exp[(i/\hbar)\oint A],$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0358 (6.2.9)	125	■ 0.999 ● N +0	$L_{\parallel}:= -\frac{1}{2\kappa}T^{\alpha}\wedge *\left({}^{(1)}T_{\alpha}-2{}^{(2)}T_{\alpha}-\frac{1}{2}{}^{(3)}T_{\alpha}\right),$	$L_{\parallel}:= -\frac{1}{2\kappa}T^{\alpha}\wedge *\left({}^{(1)}T_{\alpha}-2{}^{(2)}T_{\alpha}-\frac{1}{2}{}^{(3)}T_{\alpha}\right),$	$L_{\parallel}:= -\frac{1}{2\kappa}T^{\alpha}\wedge *\left({}^{(1)}T_{\alpha}-2{}^{(2)}T_{\alpha}-\frac{1}{2}{}^{(3)}T_{\alpha}\right),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0461 (7.2.13)	152	■ 0.999 ● N +0	$\backslash \mathrm{stackrel{\scriptscriptstyle{(\mp)}}{\Omega}}_{\mathrm{C}}=0$	$\stackrel{(\mp)}{\Omega}_{\mathrm{C}}=0$	$\stackrel{(\mp)}{\Omega}_{\mathrm{C}}=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0620 (10.1.8)	209	■ 0.999 ● N +0	$\frac{\delta C_{\mathrm{T}}}{\delta \vartheta^{\alpha}}=\frac{1}{\ell^2}T_{\alpha} \quad \text{and} \quad \frac{\delta C_{\mathrm{T}}}{\delta \Gamma^{\star\alpha}}=\frac{(-1)^s}{\ell^2}\eta_{\alpha},$	$\frac{\delta C_{\mathrm{T}}}{\delta \vartheta^{\alpha}}=\frac{1}{\ell^2}T_{\alpha} \quad \text{and} \quad \frac{\delta C_{\mathrm{T}}}{\delta \Gamma^{\star\alpha}}=\frac{(-1)^s}{\ell^2}\eta_{\alpha},$	$\frac{\delta C_{\mathrm{T}}}{\delta \vartheta^{\alpha}}=\frac{1}{\ell^2}T_{\alpha} \quad \text{and} \quad \frac{\delta C_{\mathrm{T}}}{\delta \Gamma^{\star\alpha}}=\frac{(-1)^s}{\ell^2}\eta_{\alpha},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0667 (10.3.24)	218	■ 0.999 ● N +0	$\left[\backslash \mathrm{square}+(-1)^s\,m^2\right]\backslash \vartheta^{\alpha}\backslash \cong 0,$	$\left[\square+(-1)^sm^2\right]\vartheta^{\alpha}\cong 0,$	$\left[\square+(-1)^sm^2\right]\vartheta^{\alpha}\cong 0,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0678 (10.3.35)	220	■ 0.999 ● N -2	$\check{\mathrm{E}}_{\alpha}^{\mathrm{MB}}-\check{\mathrm{E}}_{\alpha}=\frac{\theta_{\mathrm{TL}}}{\ell}d\Gamma_{\alpha}^{\star}$ d $\Gamma_{\alpha}^{\star}$,	$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}}-\check{\mathcal{E}}_{\alpha}=\frac{\theta_{\mathrm{TL}}}{\ell}d\Gamma_{\alpha}^{\star}$,	$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}}-\check{\mathcal{E}}_{\alpha}=\frac{\theta_{\mathrm{TL}}}{\ell}d\Gamma_{\alpha}^{\star}$,
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0740 (11.1.7)	238	■ 0.999 ● N +0	$\mathrm{XX}=-\mathrm{o}_{\alpha\beta}x^{\alpha}x^{\beta}\mathbb{1}$	$\mathrm{XX}=-\mathrm{o}_{\alpha\beta}x^{\alpha}x^{\beta}\mathbb{1}$	$\mathrm{XX}=-\mathrm{o}_{\alpha\beta}x^{\alpha}x^{\beta}\mathbb{1}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0776 (11.2.5)	246	■ 0.999 ● W +0	$L_{\mathrm{D}}=\frac{i}{2}\left(\bar{\psi}\gamma\wedge^{\ast}d\psi+d\bar{\psi}\wedge^{\ast}\gamma\psi\right)-m\bar{\psi}\psi\eta-\frac{1}{2}\bar{\psi}\left[\tilde{\omega},^{\ast}\gamma\right]\psi,$	$L_{\mathrm{D}}=\frac{i}{2}\left(\bar{\psi}\gamma\wedge^{\ast}d\psi+d\bar{\psi}\wedge^{\ast}\gamma\psi\right)-m\bar{\psi}\psi\eta-\frac{1}{2}\bar{\psi}\left[\tilde{\omega},^{\ast}\gamma\right]\psi,$	$L_{\mathrm{D}}=\frac{i}{2}(\bar{\psi}\gamma\wedge^{\ast}d\psi+d\bar{\psi}\wedge^{\ast}\gamma\psi)-m\bar{\psi}\psi\eta-\frac{1}{2}\bar{\psi}[\tilde{\omega},^{\ast}\gamma]\psi,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0805 (11.4.17)	255	■ 0.999 ● N +1	$\kappa=\mp\left(j+\frac{1}{2}\right)\quad\text{for}\quad j=l\pm\frac{1}{2}.$	$\kappa=\mp\left(j+\frac{1}{2}\right)\quad\text{for}\quad j=l\pm\frac{1}{2}.$	$\kappa=\mp(j+\frac{1}{2})\quad\text{for}\quad j=l\pm\frac{1}{2}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0909 (14.3.2)	320	■ 0.999 ● W +0	$\hat{L}_{\mathrm{D}}=\frac{i}{2}\left(\overline{\hat{\psi}}\hat{\gamma}\wedge^{\ast}\hat{D}\hat{\psi}+(\overline{\hat{D}\hat{\psi}})\wedge^{\ast}\hat{\gamma}\hat{\psi}\right)-\hat{m}\hat{\psi}\hat{\psi}\eta$	$\hat{L}_{\mathrm{D}}=\frac{i}{2}\left(\overline{\hat{\psi}}\hat{\gamma}\wedge^{\ast}\hat{D}\hat{\psi}+(\overline{\hat{D}\hat{\psi}})\wedge^{\ast}\hat{\gamma}\hat{\psi}\right)-\hat{m}\hat{\psi}\hat{\psi}\eta$	$\hat{L}_{\mathrm{D}}=\frac{i}{2}(\bar{\hat{\psi}}\hat{\gamma}\wedge^{\ast}\hat{D}\hat{\psi}+(\overline{\hat{D}\hat{\psi}})\wedge^{\ast}\hat{\gamma}\hat{\psi})-\hat{m}\hat{\psi}\hat{\psi}\eta$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1057 (15.2.14)	346	■ 0.999 ● N +0	i \bar{\gamma} \wedge^* \bar{D}^{\{\}} \psi - 6 \ell^2 \bar{\gamma}^5 \bar{\gamma} \wedge^* (\bar{\psi} \bar{\gamma}^5 \bar{\gamma} \psi) \psi - m \psi \eta = 0	$i \bar{\gamma} \wedge^* \bar{D}^{\{\}} \psi - 6 \ell^2 \bar{\gamma}^5 \bar{\gamma} \wedge^* (\bar{\psi} \bar{\gamma}^5 \bar{\gamma} \psi) \psi - m \psi \eta = 0$	$i \bar{\gamma} \wedge^* \bar{D}^{\{\}} \psi - 6 \ell^2 \bar{\gamma}^5 \bar{\gamma} \wedge^* (\bar{\psi} \bar{\gamma}^5 \bar{\gamma} \psi) \psi - m \psi \eta = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0669 (10.3.26)	219	■ 0.999 ● C -12	m=\frac{\theta_{\mathrm{T}} \theta_{\mathrm{TL}}}{\theta_{\mathrm{TL}}^2 + \theta_{\mathrm{T}} \theta_{\mathrm{L}}} \ell, \quad \Lambda_{\mathrm{ind}} = \frac{\theta_{\mathrm{T}}^2 [-(-1)^s 3 \theta_{\mathrm{TL}}^2 + 8 \theta_{\mathrm{T}} \theta_{\mathrm{L}}]}{4 [(-1)^s \theta_{\mathrm{TL}}^2 - 2 \theta_{\mathrm{T}} \theta_{\mathrm{L}}]^2 \ell^2},	$m = \frac{\theta_{\mathrm{T}} \theta_{\mathrm{TL}}}{2 (\theta_{\mathrm{TL}}^2 + \theta_{\mathrm{T}} \theta_{\mathrm{L}}) \ell}, \quad \Lambda_{\mathrm{ind}} = \frac{\theta_{\mathrm{T}}^2 [-(-1)^s 3 \theta_{\mathrm{TL}}^2 + 8 \theta_{\mathrm{T}} \theta_{\mathrm{L}}]}{4 [(-1)^s \theta_{\mathrm{TL}}^2 - 2 \theta_{\mathrm{T}} \theta_{\mathrm{L}}]^2 \ell^2},$	$m = \frac{\theta_{\mathrm{T}} \theta_{\mathrm{TL}}}{2 (\theta_{\mathrm{TL}}^2 + \theta_{\mathrm{T}} \theta_{\mathrm{L}}) \ell}, \quad \Lambda_{\mathrm{ind}} = \frac{\theta_{\mathrm{T}}^2 [-(-1)^s 3 \theta_{\mathrm{TL}}^2 + 8 \theta_{\mathrm{T}} \theta_{\mathrm{L}}]}{4 [(-1)^s \theta_{\mathrm{TL}}^2 - 2 \theta_{\mathrm{T}} \theta_{\mathrm{L}}]^2 \ell^2},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0929 (14.1.10)	304	■ 0.999 ● N +0	\begin{aligned} & \operatorname{Tr} \left(\Sigma_{\mathrm{YM}} \wedge^* \vartheta \right) = \\ & \operatorname{Tr} \left(\Sigma_{\mathrm{YM}} \right) \wedge^* \vartheta = \frac{1}{2 \alpha_g} \operatorname{Tr} \left\{ \vartheta \wedge \Omega \wedge^* (\vartheta \wedge^* \Omega) + \vartheta \wedge^* \vartheta \wedge^* (\Omega \wedge^* \Omega) \right\} = 0 \end{aligned}	$\operatorname{Tr} (\Sigma_{\mathrm{YM}} \wedge^* \vartheta) = \operatorname{Tr} (\vartheta \wedge^* \Sigma_{\mathrm{YM}})$ $= \frac{1}{2 \alpha_g} \operatorname{Tr} \{ \vartheta \wedge \Omega \wedge^* (\vartheta \wedge^* \Omega) + \vartheta \wedge^* \vartheta \wedge^* (\Omega \wedge^* \Omega) \} = 0$	$\operatorname{Tr} (\Sigma_{\mathrm{YM}} \wedge^* \vartheta) = \operatorname{Tr} (\vartheta \wedge^* \Sigma_{\mathrm{YM}})$ $= \frac{1}{2 \alpha_g} \operatorname{Tr} \{ \vartheta \wedge \Omega \wedge^* (\vartheta \wedge^* \Omega) + \vartheta \wedge^* \vartheta \wedge^* (\Omega \wedge^* \Omega) \} = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0093 (3.2.3)	055	■ 0.999 ● K +0	\omega^{\{\prime\}} \rightarrow G^{\{-1\}} \omega^{\{\prime\}} G + \left(d G^{\{-1\}} \right) G	$\omega' \rightarrow G^{-1} \omega' = G^{-1} \omega' G + (dG^{-1}) G$	$\omega' \rightarrow G^{-1} \omega' = G^{-1} \omega' G + (dG^{-1}) G$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0113 (3.3.14)	059	■ 0.999 ● N +0	H={ }^* F .	$H = {}^* F.$	$H = {}^* F.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0173 (3.7.1)	071	0.999 ● N +0	$\mathrm{L}_{\mathrm{H}}=-\bar{\Theta}^{\mathrm{(1)}}\wedge^*D\phi^0+\bar{\Theta}^{\mathrm{(1)}}\wedge^*\Theta^{\mathrm{(1)}}+\frac{\lambda}{4}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)^2\eta.$	$L_{\mathrm{H}}=-\bar{\Theta}^{\mathrm{(1)}}\wedge^*D\phi^0+\bar{\Theta}^{\mathrm{(1)}}\wedge^*\Theta^{\mathrm{(1)}}+\frac{\lambda}{4}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)^2\eta.$	$L_{\mathrm{H}}=-\bar{\Theta}^{\mathrm{(1)}}\wedge^*D\phi^0+\bar{\Theta}^{\mathrm{(1)}}\wedge^*\Theta^{\mathrm{(1)}}+\frac{\lambda}{4}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)^2\eta.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0201 (4.1.15)	083	0.999 ● N +0	$\delta_g^*\vartheta=g^{-1}\vartheta,\quad g\in GL(n,\mathbb{R}),$	$\delta_g^*\vartheta=g^{-1}\vartheta,\quad g\in GL(n,\mathbb{R}),$	$\delta_g^*\vartheta=g^{-1}\vartheta,\quad g\in GL(n,\mathbb{R}),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0275 (4.4.3)	097	0.999 ● W +1	$\begin{aligned} &=d(\delta\vartheta)-\left[\omega^{\mathrm{L}},\delta\vartheta\right]-\left[\delta\omega^{\mathrm{L}},\vartheta\right] \\ &=D(\delta\vartheta)-\left[\delta\omega^{\mathrm{L}},\vartheta\right]. \end{aligned}$	$\begin{aligned} &=d(\delta\vartheta)-\left[\omega^{\mathrm{L}},\delta\vartheta\right]-\left[\delta\omega^{\mathrm{L}},\vartheta\right] \\ &=D(\delta\vartheta)-\left[\delta\omega^{\mathrm{L}},\vartheta\right]. \end{aligned}$	$\begin{aligned} &=d(\delta\vartheta)-[\omega^{\mathrm{L}},\delta\vartheta]-[\delta\omega^{\mathrm{L}},\vartheta] \\ &=D(\delta\vartheta)-[\delta\omega^{\mathrm{L}},\vartheta]. \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0341 (5.3.18)	115	0.999 ● K +0	$\mathscr{A}=\frac{\kappa}{2}j_5,$	$^*\mathscr{A}=\frac{\kappa}{2}j_5,$	$^*\mathscr{A}=\frac{\kappa}{2}j_5,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0441 (6.8.12)	143	0.999 ● N +0	$D\tau_{\alpha\beta}+\vartheta_{[\alpha}\wedge\Sigma_{\beta]}\cong0,$	$D\tau_{\alpha\beta}+\vartheta_{[\alpha}\wedge\Sigma_{\beta]}\cong0,$	$D\tau_{\alpha\beta}+\vartheta_{[\alpha}\wedge\Sigma_{\beta]}\cong0,$
Conf. MathPix confidence: 0.999 0.5–0.9 <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0476 (7.4.4)	156	■ 0.999 ● K +0	$C_{\alpha\beta}=0 \quad \text{and} \quad R=0,$	$C_{\alpha\beta} = 0 \quad \text{and} \quad R = 0,$	$C_{\alpha\beta} = 0 \quad \text{and} \quad R = 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0519 (7.7.37)	166	■ 0.999 ● N +0	$D \Theta \equiv \left[\Omega^g, \gamma \right], \quad D \Omega^g \equiv 0,$	$D\Theta \equiv [\Omega^g, \gamma], \quad D\Omega^g \equiv 0,$	$D\Theta \equiv [\Omega^g, \gamma], \quad D\Omega^g \equiv 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0567 (8.6.4)	183	■ 0.999 ● N +1	$\begin{aligned} H_{\alpha\beta} &= \theta_L R_{\alpha\beta} + \theta_L^* R_{\alpha\beta}^* + \frac{\theta_T^*}{2\ell^2} \eta_{\alpha\beta} - \frac{\theta_T}{2\ell^2} \vartheta_\alpha \wedge \vartheta_\beta \\ &\simeq -\partial L_\theta / \partial R^{\alpha\beta} \end{aligned}$	$H_{\alpha\beta}^{(**)} = \theta_L R_{\alpha\beta} + \theta_L^* R_{\alpha\beta}^{(*)} + \frac{\theta_T^*}{2\ell^2} \eta_{\alpha\beta} - \frac{\theta_T}{2\ell^2} \vartheta_\alpha \wedge \vartheta_\beta \\ \simeq -\partial L_\theta / \partial R^{\alpha\beta}$	$H_{\alpha\beta}^{(**)} = \theta_L R_{\alpha\beta} + \theta_L^* R_{\alpha\beta}^{(*)} + \frac{\theta_T^*}{2\ell^2} \eta_{\alpha\beta} - \frac{\theta_T}{2\ell^2} \vartheta_\alpha \wedge \vartheta_\beta \\ \simeq -\partial L_\theta / \partial R^{\alpha\beta},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0624 (10.1.12)	210	■ 0.999 ● K +0	$D H_{\alpha} - E_{\alpha} = \Sigma_{\alpha},$	$DH_{\alpha} - E_{\alpha} = \Sigma_{\alpha},$	$DH_{\alpha} - E_{\alpha} = \Sigma_{\alpha},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0673 (10.3.30)	220	■ 0.999 ● C -4	$H_{\alpha} = -\frac{\theta_T}{2\ell^2} \vartheta_{\alpha} - \frac{\theta_{TL}}{\ell} \Gamma_{\alpha}^*, \quad H_{\alpha}^* = -\frac{\theta_L}{2} \Gamma_{\alpha}^*,$	$H_{\alpha} = -\frac{\theta_T}{2\ell^2} \vartheta_{\alpha} - \frac{\theta_{TL}}{\ell} \Gamma_{\alpha}^*, \quad H_{\alpha}^* = -\frac{\theta_L}{2} \Gamma_{\alpha}^*,$	$H_{\alpha} = -\frac{\theta_T}{2\ell^2} \vartheta_{\alpha} - \frac{\theta_{TL}}{\ell} \Gamma_{\alpha}^*, \quad H_{\alpha}^* = -\frac{\theta_L}{2} \Gamma_{\alpha}^*,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00722 (10.10.2)	230	■ 0.999 ● N +0	$\backslash\delta L=\backslash\delta\vartheta^{\alpha}\wedge\frac{\delta L}{\delta\vartheta^{\alpha}}+\delta\Gamma_{\alpha}^{\star}\wedge\frac{\delta L}{\delta\Gamma_{\alpha}^{\star}}+\delta\bar{\Psi}\wedge\frac{\delta L}{\delta\bar{\Psi}},$	$\delta L = \delta \vartheta^{\alpha} \wedge \frac{\delta L}{\delta \vartheta^{\alpha}} + \delta \Gamma_{\alpha}^{\star} \wedge \frac{\delta L}{\delta \Gamma_{\alpha}^{\star}} + \delta \bar{\Psi} \wedge \frac{\delta L}{\delta \bar{\Psi}},$	$\delta L = \delta \vartheta^{\alpha} \wedge \frac{\delta L}{\delta \vartheta^{\alpha}} + \delta \Gamma_{\alpha}^{\star} \wedge \frac{\delta L}{\delta \Gamma_{\alpha}^{\star}} + \delta \bar{\Psi} \wedge \frac{\delta L}{\delta \bar{\Psi}},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00759 (11.1.26)	242	■ 0.999 ● N +0	$\{ \}^{\ast}(\underbrace{\gamma \wedge \cdots \wedge \gamma}_{p})=\frac{p!}{(n-p)!} \gamma^{\wedge n+1} \underbrace{\gamma \wedge \cdots \wedge \gamma}_{n-p}.$	$^*(\underbrace{\gamma \wedge \cdots \wedge \gamma}_p) = \frac{p!}{(n-p)!} \gamma^{n+1} \underbrace{\gamma \wedge \cdots \wedge \gamma}_{n-p}.$	$^*(\underbrace{\gamma \wedge \cdots \wedge \gamma}_p) = \frac{p!}{(n-p)!} \gamma^{n+1} \underbrace{\gamma \wedge \cdots \wedge \gamma}_{n-p}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00780 (11.2.8)	247	■ 0.999 ● K +0	$(\backslash\psi,\backslash\chi):=\backslash\int_{H^{\wedge n-1}}\backslash\tau_{\{s\}}(\backslash\psi,\backslash\chi)=i\backslash\int_{H^{\wedge n-1}}\backslash\psi^*\backslash\gamma\backslash\chi$	$(\psi,\chi):=\int_{H^{n-1}}\tau_s(\psi,\chi)=i\int_{H^{n-1}}\psi^*\gamma\chi$	$(\psi,\chi):=\int_{H^{n-1}}\tau_s(\psi,\chi)=i\int_{H^{n-1}}\psi^*\gamma\chi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00789 (11.4.1)	251	■ 0.999 ● N +0	$E_{\{r\}}=\sqrt{\alpha}r_{\{\circ\}}^2\backslash\left[1+\backslash\left(\frac{r}{r_{\circ}}\right)^4\right]^{\wedge -1/2}\sim\sqrt{\alpha}\frac{1}{r^2}$	$E_r = \sqrt{\alpha} r_{\circ}^2 \left[1 + \left(\frac{r}{r_{\circ}} \right)^4 \right]^{-1/2} \sim \sqrt{\alpha} \frac{1}{r^2}$	$E_r = \sqrt{\alpha} r_{\circ}^2 \left[1 + \left(\frac{r}{r_{\circ}} \right)^4 \right]^{-1/2} \sim \sqrt{\alpha} \frac{1}{r^2}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00825 (11.4.39)	259	■ 0.999 ● N +0	$m_{\{\pi\}}=\sqrt{1-\omega^{\wedge 2}}\ m$	$m_{\pi} = \sqrt{1 - \omega^2} m$	$m_{\pi} = \sqrt{1 - \omega^2} \ m$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0892 (13.4.11)	287	■0.999 ● N +0	$\vartheta \xrightarrow{A^{-1}(x)} \vartheta' = \Lambda^{-1}(x) \vartheta, \quad \theta \xrightarrow{A^{-1}(x)} \theta' = \theta \Lambda^{-1}(x).$	$\vartheta \xrightarrow{A^{-1}(x)} \vartheta' = \Lambda^{-1}(x) \vartheta, \quad \theta \xrightarrow{A^{-1}(x)} \theta' = \theta \Lambda^{-1}(x).$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0932 (14.1.13)	304	■0.999 ● N +0	$e^{\delta} \circ e^{\varepsilon} = e^{*\varepsilon} \circ e^{*\delta} = e^{*(\delta+\varepsilon)},$	$e^{*\delta} \circ e^{*\varepsilon} = e^{*\varepsilon} \circ e^{*\delta} = e^{*(\delta+\varepsilon)},$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0940 (14.1.21)	305	■0.999 ● N +0	$\xi \wedge \xi = \xi \wedge^* \xi = 0$	$\xi \wedge \xi = \xi \wedge^* \xi = 0$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0951 (14.1.33)	307	■0.999 ● W +2	$\begin{aligned} & D^* \xi - (-1)^z \xi \wedge D \delta = \bar{\tau}_e \\ & D \xi + (-1)^z \xi \wedge D \delta = \bar{\tau}_p \end{aligned}$	$D^* \xi - (-1)^z \xi \wedge D \delta = \bar{\tau}_e, \\ D \xi + (-1)^z \xi \wedge D \delta = \bar{\tau}_p.$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0968 (14.2.9)	314	■0.999 ● N +2	$\hat{\omega} = \left[\begin{array}{c c} \omega^g & -\alpha \\ \hline \alpha^T & \hat{\omega} \end{array} \right] =: \omega^g \oplus \hat{\omega} - \hat{\alpha}; \quad \hat{\alpha} = \left[\begin{array}{c c} 0 & \alpha \\ \hline -\alpha^T & 0 \end{array} \right]$	$\hat{\omega} = \left[\begin{array}{c c} \omega^g & -\alpha \\ \hline \alpha^T & \hat{\omega} \end{array} \right] =: \omega^g \oplus \hat{\omega} - \hat{\alpha}; \quad \hat{\alpha} = \left[\begin{array}{c c} 0 & \alpha \\ \hline -\alpha^T & 0 \end{array} \right]$	
Conf. MathPix confidence: ■ ■ ■ — ≥0.9 0.5–0.9 <0.5 no value Residual render vs scan (inkdrill): ● ● ● ● ● ● C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1039 (15.1.16)	339	■ 0.999 ● C +9	$\begin{aligned} ds_{\mathrm{f}}^2= & f_{ij} dx^i \\ & \otimes_s dx^j \\ & = -\frac{3\gamma}{2} e^{-\lambda^f} dt^2 + 2\left[\gamma\left\{1 - \left(1 + \frac{9\gamma}{4}\right) e^{\lambda^f - \lambda^g} + \frac{9\gamma}{4} e^{2(\lambda^f - \lambda^g)}\right\}\right]^{1/2} dt dr \\ & + e^{\lambda^g} \left\{\frac{2}{3} + \frac{3\gamma}{2} \left(1 - e^{\lambda^g - \lambda^f}\right)\right\} dr^2 + \frac{2}{3} r^2 d\Omega^2 \end{aligned}$	$ds_f^2=f_{ij}dx^i\otimes_sdx^j$ $=-\frac{3\gamma}{2}e^{-\lambda^f}dt^2+2\left[\gamma\left\{1-\left(1+\frac{9\gamma}{4}\right)e^{\lambda^f-\lambda^g}+\frac{9\gamma}{4}e^{2(\lambda^f-\lambda^g)}\right\}\right]^{1/2}dtdr$ $+e^{\lambda^g}\left\{\frac{2}{3}+\frac{3\gamma}{2}\left(1-e^{\lambda^g-\lambda^f}\right)\right\}dr^2+\frac{2}{3}r^2d\Omega^2$	$ds_f^2=f_{ij}dx^i\otimes_sdx^j$ $=-\frac{3\gamma}{2}e^{-\lambda^f}dt^2+2\left[\gamma\left\{1-\left(1+\frac{9\gamma}{4}\right)e^{\lambda^f-\lambda^g}+\frac{9\gamma}{4}e^{2(\lambda^f-\lambda^g)}\right\}\right]^{1/2}dtdr$ $+e^{\lambda^g}\left\{\frac{2}{3}+\frac{3\gamma}{2}\left(1-e^{\lambda^g-\lambda^f}\right)\right\}dr^2+\frac{2}{3}r^2d\Omega^2$ (15.1.16)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1088 (A.15)	366	■ 0.999 ● N +0	$C(1, n-1)=\left[\bigotimes_{i=1}^{\left[\frac{n-2}{2}\right]} C(0,2)\right] \otimes C(1,1) \text { . }$	$C(1, n-1)=\left[\bigotimes_{i=1}^{\left[\frac{n-2}{2}\right]} C(0,2)\right] \otimes C(1,1) \text { . }$	$C(1, n-1)=\left[\bigotimes_{i=1}^{\left[\frac{n-2}{2}\right]} C(0,2)\right] \otimes C(1,1) \text { . }$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0434 (6.8.5)	142	■ 0.999 ● N -1	$H_{\alpha}:=\frac{\partial L_g}{\partial d \vartheta^{\alpha}}=-\frac{\partial L_g}{\partial T^{\alpha}}, \quad \text { and } \quad H_{\alpha \beta}:=\frac{\partial L_g}{\partial d \Gamma^{\alpha \beta}}=-\frac{\partial L_g}{\partial R^{\alpha \beta}} \text { . }$	$H_{\alpha}:=\frac{\partial L_g}{\partial d \vartheta^{\alpha}}=-\frac{\partial L_g}{\partial T^{\alpha}}, \quad \text { and } \quad H_{\alpha \beta}:=\frac{\partial L_g}{\partial d \Gamma^{\alpha \beta}}=-\frac{\partial L_g}{\partial R^{\alpha \beta}} \text { . }$	$H_{\alpha}:=\frac{\partial L_g}{\partial d \vartheta^{\alpha}}=-\frac{\partial L_g}{\partial T^{\alpha}}, \quad \text { and } \quad H_{\alpha \beta}:=\frac{\partial L_g}{\partial d \Gamma^{\alpha \beta}}=-\frac{\partial L_g}{\partial R^{\alpha \beta}} \text { . }$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0566 (8.6.3)	183	■ 0.999 ● N +0	$H_{\alpha \beta}:=\frac{\partial L_{\mathrm{qPG}}}{\partial R^{\alpha \beta}}=\frac{a_0}{2 \ell^2} \eta_{\alpha \beta}+\left(\sum_{N=1}^6 b_{(N)}^{(N)} R_{\alpha \beta}\right)$	$H_{\alpha \beta}:=\frac{\partial L_{\mathrm{qPG}}}{\partial R^{\alpha \beta}}=\frac{a_0}{2 \ell^2} \eta_{\alpha \beta}+\left(\sum_{N=1}^6 b_{(N)}^{(N)} R_{\alpha \beta}\right)$	$H_{\alpha \beta}:=\frac{\partial L_{\mathrm{qPG}}}{\partial R^{\alpha \beta}}=\frac{a_0}{2 \ell^2} \eta_{\alpha \beta}+\left(\sum_{N=1}^6 b_{(N)}^{(N)} R_{\alpha \beta}\right)$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0798 (11.4.10)	254	0.999 W +0	$\begin{aligned} & \& \left\{ i \, \bar{\gamma}^k \left(\partial_k + i \bar{\Gamma}_k + \frac{3}{4} \partial_k \mu \right) \right. \\ & \left. + \frac{3\varepsilon}{8} \ell^2 e^{\mu/2} \left(\bar{\psi} \psi - \mathfrak{a} \bar{\psi} \gamma^5 \psi \gamma^5 \right) - e^{\mu/2} m \right\} \psi = 0. \end{aligned}$	$\left\{ i \bar{\gamma}^k \left(\partial_k + i \bar{\Gamma}_k + \frac{3}{4} \partial_k \mu \right) + \frac{3\varepsilon}{8} \ell^2 e^{\mu/2} \left(\bar{\psi} \psi - \mathfrak{a} \bar{\psi} \gamma^5 \psi \gamma^5 \right) - e^{\mu/2} m \right\} \psi = 0.$	$\left\{ i \bar{\gamma}^k (\partial_k + i \bar{\Gamma}_k + \frac{3}{4} \partial_k \mu) + \frac{3\varepsilon}{8} \ell^2 e^{\mu/2} (\bar{\psi} \psi - \mathfrak{a} \bar{\psi} \gamma^5 \psi \gamma^5) - e^{\mu/2} m \right\} \psi = 0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0833 (12.2.1)	270	0.999 N +0	$L_{\mathrm{D}} = L(\gamma, \psi, \mathscr{D}\psi) = \frac{i}{2} \left\{ \bar{\psi}^* \gamma \wedge \mathscr{D}\psi + \overline{\mathscr{D}\psi} \wedge^* \gamma \psi \right\} - m \bar{\psi} \psi \eta,$	$L_{\mathrm{D}} = L(\gamma, \psi, \mathscr{D}\psi) = \frac{i}{2} \left\{ \bar{\psi}^* \gamma \wedge \mathscr{D}\psi + \overline{\mathscr{D}\psi} \wedge^* \gamma \psi \right\} - m \bar{\psi} \psi \eta,$	$L_{\mathrm{D}} = L(\gamma, \psi, \mathscr{D}\psi) = \frac{i}{2} \left\{ \bar{\psi}^* \gamma \wedge \mathscr{D}\psi + \overline{\mathscr{D}\psi} \wedge^* \gamma \psi \right\} - m \bar{\psi} \psi \eta,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0020 (2.4.2)	038	1.000 N +0	$V^{\rho} := V^{\rho} \left(M, \mathbb{C}^N, \rho(G) \subseteq GL(N, \mathbb{C}), P \right),$	$V^{\rho} := V^{\rho} \left(M, \mathbb{C}^N, \rho(G) \subseteq GL(N, \mathbb{C}), P \right),$	$V^{\rho} := V^{\rho}(M, \mathbb{C}^N, \rho(G) \subseteq GL(N, \mathbb{C}), P),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0091 (3.2.1)	055	1.000 W +0	$\mathscr{O} \rightarrow {}^{G^{-1}}\mathscr{O} = G^{-1} \mathscr{O} G.$	$\mathscr{O} \rightarrow {}^{G^{-1}}\mathscr{O} = G^{-1} \mathscr{O} G.$	$\mathscr{O} \rightarrow {}^{G^{-1}}\mathscr{O} = G^{-1} \mathscr{O} G.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0101 (3.3.2)	057	1.000 N ·1	$A \rightarrow A + d\theta(x) \quad \Rightarrow \quad F \rightarrow dA + dd\theta = dA = F.$	$A \rightarrow A + d\theta(x) \quad \Rightarrow \quad F \rightarrow dA + dd\theta = dA = F.$	$A \rightarrow A + d\theta(x) \quad \Rightarrow \quad F \rightarrow dA + dd\theta = dA = F.$
Conf. MathPix confidence: ≥ 0.9 $0.5\text{--}0.9$ < 0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0137 (3.4.8)	063	■ 1.000 ● N +0	$\begin{aligned} \delta L_{\omega} &= \delta \Omega \wedge \Pi^{\Omega} = D(\delta \omega) \wedge \Pi^{\Omega} \\ &= -\delta \omega \wedge D \Pi^{\Omega} + d\left(\delta \omega \wedge \Pi^{\Omega}\right). \end{aligned}$	$\delta L_{\omega} = \delta \Omega \wedge \Pi^{\Omega} = D(\delta \omega) \wedge \Pi^{\Omega} \\ = -\delta \omega \wedge D \Pi^{\Omega} + d\left(\delta \omega \wedge \Pi^{\Omega}\right).$	$\delta L_{\omega} = \delta \Omega \wedge \Pi^{\Omega} = D(\delta \omega) \wedge \Pi^{\Omega} \\ = -\delta \omega \wedge D \Pi^{\Omega} + d(\delta \omega \wedge \Pi^{\Omega}).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0146 (3.4.17)	065	■ 1.000 ● N +0	$Q=Q^{\{i\}}I_i:=\int_{H^3}\iota_{\partial H^3}\Pi^{\Omega},$	$Q=Q^{(i)}I_i:=\int_{H^3}\iota=\int_{\partial H^3}\Pi^{\Omega},$	$Q=Q^{(i)}I_i:=\int_{H^3}\iota=\int_{\partial H^3}\Pi^{\Omega},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0175 (3.7.4)	072	■ 1.000 ● N +0	$\tau_{\mathrm{H}}=-\left[\phi^{\left(0\right)},\left\{ \right\} ^{\ast}\Theta^{\left(1\right)}\right]$	$\tau_{\mathrm{H}}=-\left[\phi^{\left(0\right)},\ast\Theta^{\left(1\right)}\right]$	$\tau_{\mathrm{H}}=-[\phi^{\left(0\right)},\ast\Theta^{\left(1\right)}]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0195 (4.1.5)	082	■ 1.000 ● N +0	$\frac{a}{g}(n,\mathbb{R})=\mathbb{R}^n\otimes\mathfrak{gl}(n,\mathbb{R})$	$\mathfrak{a}(n,\mathbb{R})=\mathbb{R}^n\otimes\mathfrak{gl}(n,\mathbb{R})$	$\mathfrak{a}(n,\mathbb{R})=\mathbb{R}^n\otimes\mathfrak{gl}(n,\mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0208 (4.1.22)	084	■ 1.000 ● N +0	$T_{\mathrm{A}}(M)=\stackrel{\circ}{V}(M,\mathbb{R}^n,A(n,\mathbb{R}),A(M))\subset V^{\mathrm{id}}$	$T_{\mathrm{A}}(M)=\stackrel{\circ}{V}(M,\mathbb{R}^n,A(n,\mathbb{R}),A(M))\subset V^{\mathrm{id}}$	$T_{\mathrm{A}}(M)=\stackrel{\circ}{V}(M,\mathbb{R}^n,A(n,\mathbb{R}),A(M))\subset V^{\mathrm{id}}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0212 (4.2.4)	085	■ 1.000 ● K +0	$\Omega^{\mathrm{T}}:=D\,\omega^{\mathrm{T}}=d\omega^{\mathrm{T}}-\left[\omega^{\mathrm{T}},\omega^{\mathrm{L}}\right],$	$\Omega^{\mathrm{T}}:=D\omega^{\mathrm{T}}=d\omega^{\mathrm{T}}-\left[\omega^{\mathrm{T}},\omega^{\mathrm{L}}\right],$	$\Omega^{\mathrm{T}}:=D\omega^{\mathrm{T}}=d\omega^{\mathrm{T}}-\left[\omega^{\mathrm{T}},\omega^{\mathrm{L}}\right],$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0215 (4.2.7)	086	■ 1.000 ● N +0	$\mathrm{T}(n,\,\mathbb{R}):=C^{\infty}\bigl(\mathrm{A}(M)\,\times_{\mathrm{Ad}}\mathbb{R}^n\bigr)$	$\mathcal{T}(n,\mathbb{R}):=C^{\infty}\bigl(A(M)\times_{\mathrm{Ad}}\mathbb{R}^n\bigr)$	$\mathcal{T}(n,\,\mathbb{R}):=C^{\infty}(A(M)\times_{\mathrm{Ad}}\mathbb{R}^n)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0217 (4.2.9)	086	■ 1.000 ● N +0	$A=\left[\begin{array}{c c} G&T\\\hline 0&1\end{array}\right]\in\mathscr{A}_p,\quad G\in\mathscr{G}_p,T\in\mathscr{T}_p$	$A=\left[\begin{array}{c c} G&T\\\hline 0&1\end{array}\right]\in\mathscr{A}_p,\quad G\in\mathscr{G}_p,T\in\mathscr{T}_p$	$A=\left[\begin{array}{c c} G&T\\\hline 0&1\end{array}\right]\in\mathscr{A}_p,\quad G\in\mathscr{G}_p,T\in\mathscr{T}_p$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0243 (4.3.8)	091	■ 1.000 ● W +2	$\cos\phi=\frac{g_{\mu\nu}dV^{\mu}dW^{\nu}}{\sqrt{g_{\alpha\beta}dV^{\alpha}dV^{\beta}}\sqrt{g_{\alpha\beta}dW^{\alpha}dW^{\beta}}}.$	$\cos\phi=\frac{g_{\mu\nu}dV^{\mu}dW^{\nu}}{\sqrt{g_{\alpha\beta}dV^{\alpha}dV^{\beta}}\sqrt{g_{\alpha\beta}dW^{\alpha}dW^{\beta}}}.$	$\cos\phi=\frac{g_{\mu\nu}dV^{\mu}dW^{\nu}}{\sqrt{g_{\alpha\beta}dV^{\alpha}dV^{\beta}}\sqrt{g_{\alpha\beta}dW^{\alpha}dW^{\beta}}}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0323 (5.2.7)	111	■ 1.000 ● N +0	$\begin{aligned}\Sigma_{\alpha}&\rightarrow\sigma_{\alpha}:=\Sigma_{\alpha}-D\mu_{\alpha}+e_{\beta}](T^{\beta}\wedge\mu_{\alpha}),\\ \tau_{\alpha\beta}&\rightarrow\hat{\tau}_{\alpha\beta}:=\tau_{\alpha\beta}-\vartheta_{[\alpha}\wedge\mu_{\beta]}=0\end{aligned}$	$\begin{aligned}\Sigma_{\alpha}\rightarrow\sigma_{\alpha}&:=\Sigma_{\alpha}-D\mu_{\alpha}+e_{\beta}](T^{\beta}\wedge\mu_{\alpha}),\\ \tau_{\alpha\beta}\rightarrow\hat{\tau}_{\alpha\beta}&:=\tau_{\alpha\beta}-\vartheta_{[\alpha}\wedge\mu_{\beta]}=0\end{aligned}$	$\begin{aligned}\Sigma_{\alpha}\rightarrow\sigma_{\alpha}&:=\Sigma_{\alpha}-D\mu_{\alpha}+e_{\beta}](T^{\beta}\wedge\mu_{\alpha}),\\ \tau_{\alpha\beta}\rightarrow\hat{\tau}_{\alpha\beta}&:=\tau_{\alpha\beta}-\vartheta_{[\alpha}\wedge\mu_{\beta]}=0\end{aligned}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0320 (5.3.5)	112	■ 1.000 ● N +0	$\Theta:=D\,\gamma\quad\text{and}\quad\Omega^g:=d\Gamma+\Gamma\wedge\Gamma.$	$\Theta:=D\gamma\quad\text{and}\quad\Omega^g:=d\Gamma+\Gamma\wedge\Gamma.$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0370 (6.2.21)	127	■ 1.000 ● N +0	$\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\alpha=-\partial\stackrel{\scriptscriptstyle{(\pm)}}{L}/\partial T^\alpha=H_\alpha^\parallel\mp(i/\kappa)T_\alpha$	$\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\alpha=-\partial\stackrel{\scriptscriptstyle{(\pm)}}{L}/\partial T^\alpha=H_\alpha^\parallel\mp(i/\kappa)T_\alpha$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0300 (6.2.31)	129	■ 1.000 ● N +0	$\stackrel{\scriptscriptstyle{(\pm)}}{D}\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\alpha=0,$	$\stackrel{\scriptscriptstyle{(\pm)}}{D}\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\alpha=0,$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0420 (6.7.10)	141	■ 1.000 ● N +0	$\kappa D\theta_L\wedge DT_\beta=-T\wedge\eta_\beta=D\eta_\beta,$	$\kappa D\theta_L\wedge DT_\beta=-T\wedge\eta_\beta=D\eta_\beta,$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0522 (7.7.41)	167	■ 1.000 ● N +1	$\begin{gathered} K^{\scriptscriptstyle{(\star)}}:=\eta_{\alpha\beta\gamma}\wedge K^{\beta\gamma}\gamma^\alpha,\\ \Omega^{g(\star)}:=\frac{i}{8}R_{\alpha\beta}\eta^{\alpha\beta\gamma\delta}\sigma_{\gamma\delta}=-\frac{1}{4}R^{\alpha\beta}\gamma_5\sigma_{\alpha\beta}=i\gamma_5\Omega^g. \end{gathered}$	$K^{\scriptscriptstyle{(\star)}}:=\eta_{\alpha\beta\gamma}\wedge K^{\beta\gamma}\gamma^\alpha,$ $\Omega^{g(\star)}:=\frac{i}{8}R_{\alpha\beta}\eta^{\alpha\beta\gamma\delta}\sigma_{\gamma\delta}=-\frac{1}{4}R^{\alpha\beta}\gamma_5\sigma_{\alpha\beta}=i\gamma_5\Omega^g.$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0526 (7.7.46)	168	■ 1.000 ● N -1	$L_{\mathrm{NY}}:=dC_{\mathrm{TT}}=\frac{1}{2}\ell^2\left(T^{\alpha}\wedge T_{\alpha}+R_{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}\right),$	$L_{\mathrm{NY}}:=dC_{\mathrm{TT}}=\frac{1}{2\ell^2}\left(T^{\alpha}\wedge T_{\alpha}+R_{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}\right),$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0535 (8.2.2)	175	■ 1.000 ● N +0	$\widetilde{\Gamma}:=\Gamma\oplus C,\quad\widetilde{\Omega}^g:=\Omega^g\oplus\Psi\oplus\Phi.$	$\widetilde{\Gamma}:=\Gamma\oplus C,\quad\widetilde{\Omega}^g:=\Omega^g\oplus\Psi\oplus\Phi.$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0538 (8.2.5)	176	■ 1.000 ● K +0	$s_{\mathrm{loc}}(\alpha+\lambda)=-d\lambda,\quad s_{\mathrm{loc}}\lambda=0$	$s_{\mathrm{loc}}(\alpha+\lambda)=-d\lambda,\quad s_{\mathrm{loc}}\lambda=0$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0539 (8.2.6)	176	■ 1.000 ● K +0	$(d\oplus s_{\mathrm{loc}})(\alpha+\lambda)=d\alpha.$	$(d\oplus s_{\mathrm{loc}})(\alpha+\lambda)=d\alpha.$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0573 (8.6.11)	185	■ 1.000 ● W +2	$\begin{aligned} L_{\mathrm{qPG}}(*)&= \frac{\Lambda}{\ell^2}\eta-\frac{a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2}T^{\alpha}\wedge H_{\alpha}(**)-\frac{1}{2}R^{\alpha\beta}\wedge H_{\alpha\beta}(**)\\ &= \frac{\Lambda}{\ell^2}\eta-\frac{\theta_{\mathrm{T}}^*+a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2\ell^2}T^{\alpha}\wedge\left[\theta_{\mathrm{T}}^*K_{\alpha}^{(*)}-\theta_{\mathrm{T}}T_{\alpha}\right]\\ &\quad -\frac{1}{2}R^{\alpha\beta}\wedge\left[\theta_{\mathrm{L}}R_{\alpha\beta}+\theta_{\mathrm{L}}^*R_{\alpha\beta}^{(*)}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}\right]. \end{aligned}$	$L_{\mathrm{qPG}}(**)=\frac{\Lambda}{\ell^2}\eta-\frac{a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2}T^{\alpha}\wedge H_{\alpha}(**)-\frac{1}{2}R^{\alpha\beta}\wedge H_{\alpha\beta}(**)\\ =\frac{\Lambda}{\ell^2}\eta-\frac{\theta_{\mathrm{T}}^*+a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2\ell^2}T^{\alpha}\wedge\left[\theta_{\mathrm{T}}^*K_{\alpha}^{(*)}-\theta_{\mathrm{T}}T_{\alpha}\right]\\ -\frac{1}{2}R^{\alpha\beta}\wedge\left[\theta_{\mathrm{L}}R_{\alpha\beta}+\theta_{\mathrm{L}}^*R_{\alpha\beta}^{(*)}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}\right]. \quad (8.6)$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0580 (8.6.18)	186	■1.000 C +6	$G_{\alpha}^{\{\backslash\ \backslash\}}\text{-}\Lambda_{\text{eff}}\ \eta_{\alpha}=\kappa_{\text{eff}}\ \overparen{\Sigma}_{\alpha}\ .$	$(not\ rendered)$	$G_{\alpha}^{\{\}}-\Lambda_{\text{eff}}\ \eta_{\alpha}=\kappa_{\text{eff}}\ \widehat{\Sigma}_{\alpha}\ .$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0595 (9.1.13)	195	■1.000 N +0	$R^{\{\{\ \backslash\}\}}\text{-}\frac{6\ \kappa\ \gamma}{\ell^{*2}}\ .$	$R^{\{\}}=-\frac{6\kappa\gamma}{\ell^{*2}}.$	$R^{\{\}}=-\frac{6\kappa\gamma}{\ell^{*2}}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0645 (10.3.2)	214	■1.000 K +0	$\widetilde{T}_{\alpha}=T_{\alpha}\text{-}\frac{\varepsilon}{\ell}\{\backslash\text{varepsilonpsilon}\}\{\backslash\ell\}\ \eta_{\alpha},\ \text{quad}\ \widetilde{R}_{\alpha}^{\star}=R_{\alpha}^{\star}\text{-}(-1)^s\frac{\varepsilon}{2\ell}T_{\alpha}+(-1)^s\frac{\varepsilon^2}{4\ell^2}\eta_{\alpha}$	$\tilde{T}_{\alpha}=T_{\alpha}-\frac{\varepsilon}{\ell}\eta_{\alpha},\quad \tilde{R}_{\alpha}^{\star}=R_{\alpha}^{\star}-(-1)^s\frac{\varepsilon}{2\ell}T_{\alpha}+(-1)^s\frac{\varepsilon^2}{4\ell^2}\eta_{\alpha}$	$\tilde{T}_{\alpha}=T_{\alpha}-\frac{\varepsilon}{\ell}\eta_{\alpha},\quad \tilde{R}_{\alpha}^{\star}=R_{\alpha}^{\star}-(-1)^s\frac{\varepsilon}{2\ell}T_{\alpha}+(-1)^s\frac{\varepsilon^2}{4\ell^2}\eta_{\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0676 (10.3.33)	220	■1.000 N -2	$\check{\mathrm{E}}_{\alpha}^{\{\mathrm{T}\}\ \mathrm{T}}\text{-}(-1)^s\frac{\theta_{\mathrm{T}}\theta_{\mathrm{L}}}{2\chi\ell}\left(d\Gamma_{\alpha}^{\star}+\frac{2}{\theta_{\mathrm{L}}}\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}\star}\right),$	$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}}=-(-1)^s\frac{\theta_{\mathrm{T}}\theta_{\mathrm{L}}}{2\chi\ell}\left(d\Gamma_{\alpha}^{\star}+\frac{2}{\theta_{\mathrm{L}}}\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}\star}\right),$	$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}}=-(-1)^s\frac{\theta_{\mathrm{T}}\theta_{\mathrm{L}}}{2\chi\ell}\left(d\Gamma_{\alpha}^{\star}+\frac{2}{\theta_{\mathrm{L}}}\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}\star}\right),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0688 (10.3.46)	222	■1.000 N +0	$G_{\alpha}^{\{\backslash\ \backslash\}}\text{-}\tilde{\Lambda}_{\text{eff}}\ \eta_{\alpha}=\ell\widetilde{\Sigma}_{\alpha},$	$G_{\alpha}^{\{\}}-\tilde{\Lambda}\eta_{\alpha}=\ell\tilde{\Sigma}_{\alpha},$	$G_{\alpha}^{\{\}}-\tilde{\Lambda}\eta_{\alpha}=\ell\ \tilde{\Sigma}_{\alpha},$
Conf. MathPix confidence: ■≥0.9 ■0.5-0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0690 (10.5.1)	223	■ 1.000 ● N +0	$\begin{aligned} \sigma^1 &= \left(\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array} \right), \quad \sigma^2 = \left(\begin{array}{ll} 0 & -i \\ i & 0 \end{array} \right), \quad \sigma^3 = \left(\begin{array}{ll} 1 & 0 \\ 0 & -1 \end{array} \right). \\ \sigma^1 &= \left(\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array} \right), \quad \sigma^2 = \left(\begin{array}{ll} 0 & -i \\ i & 0 \end{array} \right), \quad \sigma^3 = \left(\begin{array}{ll} 1 & 0 \\ 0 & -1 \end{array} \right). \end{aligned}$	$\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$	$\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0710 (10.7.6)	227	■ 1.000 ● W +2	$\begin{aligned} \vartheta^{\hat{0}} &= dt + \ell^2 \sigma \rho^{*2} [1 - \cos(\rho/\rho^*)] d\phi \\ \vartheta^{\hat{1}} &= d\rho, \quad \vartheta^{\hat{2}} = \rho^* \sin(\rho/\rho^*) d\phi \\ \Gamma^{\hat{1}\hat{2}} &= \cos(\rho/\rho^*) d\phi = -\Gamma^{\hat{2}\hat{1}} \end{aligned}$	$\begin{aligned} \vartheta^{\hat{0}} &= dt + \ell^2 \sigma \rho^{*2} [1 - \cos(\rho/\rho^*)] d\phi \\ \vartheta^{\hat{1}} &= d\rho, \quad \vartheta^{\hat{2}} = \rho^* \sin(\rho/\rho^*) d\phi \\ \Gamma^{\hat{1}\hat{2}} &= \cos(\rho/\rho^*) d\phi = -\Gamma^{\hat{2}\hat{1}} \end{aligned}$	$\begin{aligned} \vartheta^{\hat{0}} &= dt + \ell^2 \sigma \rho^{*2} [1 - \cos(\rho/\rho^*)] d\phi \\ \vartheta^{\hat{1}} &= d\rho, \quad \vartheta^{\hat{2}} = \rho^* \sin(\rho/\rho^*) d\phi, \\ \Gamma^{\hat{1}\hat{2}} &= \cos(\rho/\rho^*) d\phi = -\Gamma^{\hat{2}\hat{1}}. \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0716 (10.9.2)	229	■ 1.000 ● K +0	$D \Psi = d \Psi - \frac{1}{2} \gamma_{\alpha} \Gamma^{\star \alpha} \wedge \Psi,$	$D \Psi = d \Psi - \frac{1}{2} \gamma_{\alpha} \Gamma^{\star \alpha} \wedge \Psi,$	$D \Psi = d \Psi - \frac{1}{2} \gamma_{\alpha} \Gamma^{\star \alpha} \wedge \Psi,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0720 (10.9.6)	229	■ 1.000 ● N +0	$\begin{aligned} \tau_{\alpha}^{\star} &:= \frac{1}{2} \eta_{\alpha \beta \gamma} \tau^{\beta \gamma} = \frac{(-1)^s}{2} \frac{\delta L_{\Psi}}{\delta \Gamma_{\alpha}^{\star}} = \frac{(-1)^s}{2} \frac{i}{4} \bar{\Psi} \wedge \gamma_{\alpha} \Psi. \\ \tau_{\alpha}^{\star} &:= \frac{1}{2} \eta_{\alpha \beta \gamma} \tau^{\beta \gamma} = \frac{(-1)^s}{2} \frac{\delta L_{\Psi}}{\delta \Gamma_{\alpha}^{\star}} = \frac{(-1)^s}{2} \frac{i}{4} \bar{\Psi} \wedge \gamma_{\alpha} \Psi. \end{aligned}$	$\tau_{\alpha}^{\star} := \frac{1}{2} \eta_{\alpha \beta \gamma} \tau^{\beta \gamma} = \frac{(-1)^s}{2} \frac{\delta L_{\Psi}}{\delta \Gamma_{\alpha}^{\star}} = \frac{(-1)^s}{2} \frac{i}{4} \bar{\Psi} \wedge \gamma_{\alpha} \Psi.$	$\tau_{\alpha}^{\star} := \frac{1}{2} \eta_{\alpha \beta \gamma} \tau^{\beta \gamma} = \frac{(-1)^s}{2} \frac{\delta L_{\Psi}}{\delta \Gamma_{\alpha}^{\star}} = \frac{(-1)^s}{2} \frac{i}{4} \bar{\Psi} \wedge \gamma_{\alpha} \Psi.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0728 (10.10.8)	231	■ 1.000 ● K +0	$\gamma^{\alpha} \Psi \wedge \bar{\Psi} \gamma_{\alpha} \Psi = 0,$	$\gamma^{\alpha} \Psi \wedge \bar{\Psi} \gamma_{\alpha} \Psi = 0,$	$\gamma^{\alpha} \Psi \wedge \bar{\Psi} \gamma_{\alpha} \Psi = 0,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0770 (11.1.37)	244	■ 1.000 ● N +0	$D \longrightarrow {}^S D = S^{-1} D S \text{ .}$	$D \longrightarrow {}^{S^{-1}} D = S^{-1} D S.$	$D \longrightarrow {}^{S^{-1}} D = S^{-1} D S.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0839 (12.2.7)	272	■ 1.000 ● N +0	$j_{\pm} := \frac{1}{2} \frac{\bar{\psi}}{\psi} \left(1 \pm \gamma_5 \right)^* \gamma \psi = \bar{\psi}_{\mathrm{L},\mathrm{R}}^* \gamma \psi_{\mathrm{L},\mathrm{R}} \text{ .}$	$j_{\pm} := \frac{1}{2} \bar{\psi} \left(1 \pm \gamma_5 \right)^* \gamma \psi = \bar{\psi}_{\mathrm{L},\mathrm{R}}^* \gamma \psi_{\mathrm{L},\mathrm{R}}.$	$j_{\pm} := \frac{1}{2} \bar{\psi} (1 \pm \gamma_5)^* \gamma \psi = \bar{\psi}_{\mathrm{L},\mathrm{R}}^* \gamma \psi_{\mathrm{L},\mathrm{R}}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0873 (13.3.1)	284	■ 1.000 ● N +0	$\nvdash^A{}_B := L^A{}_B - \frac{1}{n} \delta_B^A L^C{}_C \text{ .}$	$\not\vdash^A{}_B := L^A{}_B - \frac{1}{n} \delta_B^A L^C{}_C.$	$\not\vdash^A{}_B := L^A{}_B - \frac{1}{n} \delta_B^A L^C{}_C.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0893 (13.4.12)	287	■ 1.000 ● N +0	$\mathrm{A}_*(n,\mathbb{R})/ \mathrm{GL}(n,\mathbb{R}) \approx \mathrm{GL}(n,\mathbb{R}) \otimes \mathrm{GL}(n,\mathbb{R}) \text{ ,}$	$\mathbf{A}_*(n,\mathbb{R})/\mathrm{GL}(n,\mathbb{R}) \approx \mathbb{R}^n \otimes \mathbb{R}_*^n,$	$\mathbf{A}_*(n,\mathbb{R})/\mathrm{GL}(n,\mathbb{R}) \approx \mathbb{R}^n \otimes \mathbb{R}_*^n,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0907 (13.5.12)	290	■ 1.000 ● N +0	$\tilde{L}_{\mathrm{SSB}} + d\hat{C} \cong \frac{1}{2\mu} \eta_{\alpha\beta\gamma\delta} \left[R^{\alpha\beta} - \frac{\Lambda}{3} \vartheta^\alpha \wedge \vartheta^\beta \right] \wedge \left[R^{\gamma\delta} - \frac{\Lambda}{3} \vartheta^\gamma \wedge \vartheta^\delta \right] \text{ .}$	$\tilde{L}_{\mathrm{SSB}} + d\hat{C} \cong \frac{1}{2\mu} \eta_{\alpha\beta\gamma\delta} \left[R^{\alpha\beta} - \frac{\Lambda}{3} \vartheta^\alpha \wedge \vartheta^\beta \right] \wedge \left[R^{\gamma\delta} - \frac{\Lambda}{3} \vartheta^\gamma \wedge \vartheta^\delta \right] \text{ .}$	$\tilde{L}_{\mathrm{SSB}} + d\hat{C} \cong \frac{1}{2\mu} \eta_{\alpha\beta\gamma\delta} \left[R^{\alpha\beta} - \frac{\Lambda}{3} \vartheta^\alpha \wedge \vartheta^\beta \right] \wedge \left[R^{\gamma\delta} - \frac{\Lambda}{3} \vartheta^\gamma \wedge \vartheta^\delta \right] \text{ .}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0947 (14.1.28)	307	■ 1.000 ● N +0	$R_{i\alpha}^{\{\}}R_{\cdot j}^{\{\}}=\frac{1}{4}g_{ij}R_{\alpha\beta}^{\{\}}R^{\{\}}_{\alpha\beta}$		$R_{i\alpha}^{\{\}}R_{\cdot j}^{\{\}}=\frac{1}{4}g_{ij}R_{\alpha\beta}^{\{\}}R^{\{\}}_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1004 (14.3.17)	323	■ 1.000 ● W +0	$\widetilde{\Theta}:=D\gamma=\frac{1}{2}T_{\alpha\beta}^{\cdot\cdot c}\gamma_c\otimes\vartheta^\alpha\wedge\vartheta^\beta$		$\widetilde{\Theta}:=D\gamma=\frac{1}{2}T_{\alpha\beta}^{\cdot\cdot c}\gamma_c\otimes\vartheta^\alpha\wedge\vartheta^\beta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1013 (14.4.8)	326	■ 1.000 ● N +0	$M_{\mathrm{ir}}^2:=\frac{S_{\mathrm{A}}}{16\pi}\frac{M^{*4}c^2}{\hbar^2}$		$M_{\mathrm{ir}}^2:=\frac{S_{\mathrm{A}}}{16\pi}\frac{M^{*4}c^2}{\hbar^2}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1135 (C.3)	375	■ 1.000 ● N +0	$\left[I_i,I_j\right]=c_{ij}^{k}I_k\;.$	$[I_i,I_j]=c_{ij}^{k}I_k.$	$[I_i,I_j]=c_{ij}^{k}I_k.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0001 (1.1)	024	■ 1.000 ● N +0	$\ell^*:=\sqrt{8\pi\hbar G_N/c^3}\simeq 8\times 10^{-33}\,\mathrm{cm},$	$\ell^*:=\sqrt{8\pi\hbar G_N/c^3}\simeq 8\times 10^{-33}\,\mathrm{cm},$	$\ell^*:=\sqrt{8\pi\hbar G_N/c^3}\simeq 8\times 10^{-33}\mathrm{cm},$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0003 (2.1.2)	032	■1.000 ● N -1	$e^{\mathrm{f}} := \left. \frac{d f(s(t))}{d t} \right _{t_o = s^{-1}(m)} = \frac{\partial f}{\partial x^i} \cdot \frac{d x^i(s(t))}{d t}$	$e^{\mathrm{f}} := \left. \frac{d f(s(t))}{d t} \right _{t_o = s^{-1}(m)} = \frac{\partial f}{\partial x^i} \cdot \frac{d x^i(s(t))}{d t}$	$e^{\mathrm{f}} := \left. \frac{d f(s(t))}{d t} \right _{t_o = \overset{-1}{s}(m)} = \frac{\partial f}{\partial x^i} \cdot \frac{d x^i(s(t))}{d t}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0005 (2.1.4)	033	■1.000 ● N +1	$T(M) = \bigcup_{m \in M} T_m(M)$	$T(M) = \bigcup_{m \in M} T_m(M)$	$T(M) = \bigcup_{m \in M} T_m(M).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0008 (2.2.3)	034	■1.000 ● N -1	$\vartheta^{\alpha} = \mathrm{E}_{\mathrm{j}}^{\alpha}(m) \, d \, x^{\mathrm{j}} \; .$	$\vartheta^{\alpha} = \mathrm{E}_j^{\alpha}(m) dx^j.$	$\vartheta^{\alpha} = \mathrm{E}_j^{\alpha}(m) \, dx^j.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0011 (2.2.6)	034	■1.000 ● W +2	$T = T_{\alpha_1 \cdots \alpha_p}^{\beta_1 \cdots \beta_q}(m) \, \vartheta^{\alpha_1} \cdots \vartheta^{\alpha_p} \otimes e_{\beta_1} \otimes \cdots \otimes e_{\beta_q} \in C^{\infty}(T_p^q(M)) \; .$	$T = T_{\alpha_1 \cdots \alpha_p}^{\beta_1 \cdots \beta_q}(m) \vartheta^{\alpha_1} \otimes \cdots \otimes \vartheta^{\alpha_p} \otimes e_{\beta_1} \otimes \cdots \otimes e_{\beta_q} \in C^{\infty}(T_p^q(M)) \; .$	$T = T_{\alpha_1 \cdots \alpha_p}^{\beta_1 \cdots \beta_q}(m) \vartheta^{\alpha_1} \otimes \cdots \otimes \vartheta^{\alpha_p} \otimes e_{\beta_1} \otimes \cdots \otimes e_{\beta_q} \in C^{\infty}(T_p^q(M)).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0012 (2.2.7)	034	■1.000 ● N -2	$d \, s^{\mathrm{2}} = g_{\alpha \, \beta} \, \vartheta^{\alpha} \, \vartheta^{\beta} =: g_{i \, j} \, d \, x^{\mathrm{i}} \, \vartheta^{\beta} \otimes_s d \, x^{\mathrm{j}} \; .$	$ds^2 = g_{\alpha \beta} \vartheta^{\alpha} \otimes_s \vartheta^{\beta} =: g_{ij} dx^i \otimes_s dx^j.$	$ds^2 = g_{\alpha \beta} \vartheta^{\alpha} \otimes_s \vartheta^{\beta} =: g_{ij} \, dx^i \otimes_s \, dx^j.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0014 (2.2.9)	035	■ 1.000 ● N -1	$\alpha^{\{p\}}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\cdots\vartheta^{\alpha_p}\in C^\infty(\wedge^pT^*(M)).$	$\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty(\wedge^pT^*(M)).$	$\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty(\wedge^pT^*(M)).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0015 (2.3.1)	036	■ 1.000 ● N +0	$\pi\colon F\rightarrow M.$	$\pi:F\rightarrow M.$	$\pi:F\rightarrow M.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0016 (2.3.2)	037	■ 1.000 ● K +0	$\sigma\colon U\rightarrow F\quad\text{where}\quad\pi\circ\sigma=\text{id}$	$\sigma:U\rightarrow F\quad\text{where}\quad\pi\circ\sigma=\text{id}$	$\sigma:U\rightarrow F\quad\text{where}\quad\pi\circ\sigma=\text{id}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0017 (2.3.3)	037	■ 1.000 ● W +0	$\iota\left(m,g_1g_2\right)=\iota\left(m,g_1\right)\cdot g_2,\quad m\in U,g\in G.$	$\iota(m,g_1g_2)=\iota(m,g_1)\cdot g_2,\quad m\in U,g\in G.$	$\iota(m,g_1g_2)=\iota(m,g_1)\cdot g_2,\quad m\in U,g\in G.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0019 (2.4.1)	038	■ 1.000 ● N -2	$\delta_g(p,\zeta):=\left(pg,g^{-1}\zeta\right)\subset P\times F,\quad g\in G.$	$\delta_g(p,\zeta):=(pg,g^{-1}\zeta)\subset P\times F,\quad g\in G.$	$\delta_g(p,\zeta):=(pg,g^{-1}\zeta)\subset P\times F,\quad g\in G.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0022 (2.4.4)	038	■ 1.000 ● N -2	$\varphi^{\{p\}}=\left\{\varphi^{\{p\}A}\mid A=1,\ldots,N\right\}.$	$\varphi^{(p)}=\left\{\varphi^{(p)A}\mid A=1,\ldots,N\right\}.$	$\varphi^{(p)}=\left\{\varphi^{(p)A}\mid A=1,\ldots,N\right\}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0023 (2.4.5)	039	■ 1.000 ● N +0	$L_{\{j\}}:=\left[\left[\frac{\partial \rho_A}{\partial \xi^j}\right]_{g=e}\right]_{\in T_e(\rho(G))}$	$L_j:=\left[\frac{\partial \rho_A^B(g(\xi))}{\partial \xi^j}\right]_{g=e}\in T_e(\rho(G))$	$L_j:=\left[\frac{\partial \rho_A^B(g(\xi))}{\partial \xi^j}\right]_{g=e}\in T_e(\rho(G))$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0025 (2.4.7)	039	■ 1.000 ● W -2	$\varphi_{\{p\}}=\frac{1}{p!}\varphi_{\alpha_1\ldots\alpha_p}L_j\otimes\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^{Ad}).$	$\phi^{(p)}=\frac{1}{p!}\phi_{\alpha_1\ldots\alpha_p}^jL_j\otimes\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^{Ad}).$	$\phi^{(p)}=\frac{1}{p!}\phi_{\alpha_1\ldots\alpha_p}^jL_j\otimes\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^{Ad}).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0026 (2.4.8)	039	■ 1.000 ● C -5	$\left[\varphi^{(p)},\psi^{(q)}\right]:=\varphi^{(p)}\wedge\psi^{(q)}-(-1)^{pq}\psi^{(q)}\wedge\varphi^{(p)}=(-1)^{pq+1}\left[\psi^{(q)},\varphi^{(p)}\right]$	$\left[\phi^{(p)},\psi^{(q)}\right]:=\phi^{(p)}\wedge\psi^{(q)}-(-1)^{pq}\psi^{(q)}\wedge\phi^{(p)}=(-1)^{pq+1}\left[\psi^{(q)},\phi^{(p)}\right]$	$\left[\phi^{(p)},\psi^{(q)}\right]:=\phi^{(p)}\wedge\psi^{(q)}-(-1)^{pq}\psi^{(q)}\wedge\phi^{(p)}=(-1)^{pq+1}\left[\psi^{(q)},\phi^{(p)}\right]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0027 (2.4.9)	039	■ 1.000 ● N +0	$\left[\varphi^{(2k)},\varphi^{(2k)}\right]=-\left[\varphi^{(2k)},\varphi^{(2k)}\right]=0.$	$\left[\phi^{(2k)},\phi^{(2k)}\right]=-\left[\phi^{(2k)},\phi^{(2k)}\right]=0.$	$\left[\phi^{(2k)},\phi^{(2k)}\right]=-\left[\phi^{(2k)},\phi^{(2k)}\right]=0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0029 (2.5.2)	040	■ 1.000 ● N +0	$\omega(e\,g)=g^{-1}\omega(e)\,g,\quad e\in T(P).$	$\omega(eg)=g^{-1}\omega(e)g,\quad e\in T(P).$	$\omega(eg)=g^{-1}\omega(e)g,\quad e\in T(P).$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0030 (2.5.3)	040	1.000 N -1	$\omega(p\,dg)=g^{-1}\,dg\quad\in\quad T_e(G)\approx\mathfrak{g}$	$\omega(pdg)=g^{-1}dg\quad\in\quad T_e(G)\approx\mathfrak{g}$	$\omega(p\,dg)=g^{-1}dg\quad\in\quad T_e(G)\approx\mathfrak{g}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0031 (2.5.4)	040	1.000 N +0	$D:C^{\infty}\left(V^{\rho}\right)\longrightarrow C^{\infty}\left(T_{\mathbb{C}}^{\ast}(M)\otimes V^{\rho}\right)$	$D:C^{\infty}(V^{\rho})\longrightarrow C^{\infty}(T_{\mathbb{C}}^{\ast}(M)\otimes V^{\rho})$	$D:C^{\infty}(V^{\rho})\longrightarrow C^{\infty}(T_{\mathbb{C}}^{\ast}(M)\otimes V^{\rho})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0032 (2.5.6)	041	1.000 N +0	$D\,b=\omega\otimes b,\quad b\in C^{\infty}\left(V^{\rho}\right).$	$Db=\omega\otimes b,\quad b\in C^{\infty}(V^{\rho}).$	$Db=\omega\otimes b,\quad b\in C^{\infty}(V^{\rho}).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0033 (2.5.7)	041	1.000 C -3	$D\left(\varphi^{\left(p\right)}\otimes b\right)=d\,\varphi^{\left(p\right)}\otimes b+(-1)^p\varphi^{\left(p\right)}\otimes Db.$	$D\left(\varphi^{\left(p\right)}\otimes b\right)=d\varphi^{\left(p\right)}\otimes b+(-1)^p\varphi^{\left(p\right)}\otimes Db.$	$D(\varphi^{(p)}\otimes b)=d\varphi^{(p)}\otimes b+(-1)^p\varphi^{(p)}\otimes Db.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0034 (2.5.8)	041	1.000 C -3	$D\left(\varphi^{\left(p\right)}\otimes b\right)=\left\{d\varphi^p+(-1)^p\left[\varphi^{\left(p\right)},\omega\right]\right\}\otimes b.$	$D\left(\varphi^{\left(p\right)}\otimes b\right)=\left\{d\varphi^p+(-1)^p\left[\varphi^{\left(p\right)},\omega\right]\right\}\otimes b.$	$D(\varphi^{(p)}\otimes b)=\left\{d\varphi^p+(-1)^p[\varphi^{(p)},\omega]\right\}\otimes b.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0035 (2.5.9)	041	1.000 K +0	$D\,b_{\{A\}}=\omega_{\{A\}}\{ \}^{\{B\}}\,b_{\{B\}},\,A,\,B=1,\,\ldots,\,N.$	$Db_A=\omega_A{}^Bb_B,A,B=1,\ldots,N.$	$Db_A=\omega_A{}^Bb_B,A,B=1,\ldots,N.$
Conf. MathPix confidence: ≥0.9 0.5-0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0036 (2.5.10)	041	■ 1.000 ● N +0	$\backslash Gamma:=\stackrel{*}{\sigma} \backslash \omega=\backslash Gamma_{\backslash \alpha}\{ \}^{\backslash j} \backslash L_{\backslash j} \backslash \otimes \backslash \vartheta^{\backslash \alpha}, \quad \backslash quad \backslash j=1, \backslash \ldots, \backslash \operatorname{dim} \backslash G,$	$\Gamma := \overset{*}{\sigma} \omega = \Gamma_{\alpha}^j L_j \otimes \vartheta^{\alpha}, \quad j = 1, \dots, \dim G,$	$\Gamma := \overset{*}{\sigma} \omega = \Gamma_{\alpha}^j L_j \otimes \vartheta^{\alpha}, \quad j = 1, \dots, \dim G,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0037 (2.5.11)	041	■ 1.000 ● N -1	$A_{\backslash \alpha}\{ \}^{\backslash j}(m):=\backslash Gamma_{\backslash \alpha}\{ \}^{\backslash j}(m)_{\backslash \mid G=\backslash U(f)}$	$A_{\alpha}^j(m) := \Gamma_{\alpha}^j(m)_{ G=U(f)}$	$A_{\alpha}^j(m) := \Gamma_{\alpha}^j(m)_{ G=U(f)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0038 (2.6.1)	042	■ 1.000 ● N +0	$D \, D \, b =: \backslash \omega \backslash \otimes b \, .$	$DDb =: \Omega \otimes b.$	$DDb =: \Omega \otimes b.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0039 (2.6.2)	042	■ 1.000 ● N +1	$\backslash \begin{aligned} D \, D \, b &= D(\backslash \omega \backslash \otimes b) \, \& \, = d \, \backslash \omega \backslash \otimes b - \backslash \omega \backslash \wedge D \, b \, \backslash \backslash \, \& \, = (d \, \backslash \omega - \backslash \omega \backslash \wedge \backslash \omega) \backslash \otimes b \, \backslash \end{aligned}$	$DDb = D(\omega \otimes b) = d\omega \otimes b - \omega \wedge Db \\ = (d\omega - \omega \wedge \omega) \otimes b$	$DDb = D(\omega \otimes b) = d\omega \otimes b - \omega \wedge Db \\ = (d\omega - \omega \wedge \omega) \otimes b$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0040 (2.6.3)	042	■ 1.000 ● C +3	$\backslash \omega = d \, \backslash \omega - \backslash \omega \backslash \wedge \backslash \omega = d \, \backslash \omega - \backslash \frac{1}{2}[\backslash \omega, \backslash \omega]$	$\Omega = d\omega - \omega \wedge \omega = d\omega - \frac{1}{2}[\omega, \omega]$	$\Omega = d\omega - \omega \wedge \omega = d\omega - \frac{1}{2}[\omega, \omega] \Big .$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0043 (2.6.6)	042	■ 1.000 ● N +0	$F_{ij}=\partial_i A_j-\partial_j A_i$	$F_{ij}=\partial_i A_j-\partial_j A_i$	$F_{ij}=\partial_i A_j-\partial_j A_i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0044 (2.6.7)	042	■ 1.000 ● N +1	$\Delta b=b-b(\circ)=\oint (db)_{\mathrm{hor}}$	$\Delta b=b-b(\circ)=\oint (db)_{\mathrm{hor}}$	$\Delta b=b-b(\circ)=\oint (db)_{\mathrm{hor}},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0047 (2.6.10)	043	■ 1.000 ● N +0	$K=\lim_{U\rightarrow 0}\frac{\int_U\Omega\otimes b}{\int_U1\otimes b}$	$K=\lim_{U\rightarrow 0}\left(\int_U\Omega\otimes b\right)/\left(\int_U1\otimes b\right)$	$K=\lim_{U\rightarrow 0}\left(\int_U\Omega\otimes b\right)/\left(\int_U1\otimes b\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0049 (2.6.12)	043	■ 1.000 ● N +0	$D\,\Omega\equiv 0$	$D\Omega\equiv 0$	$D\Omega\equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0052 (2.7.2)	044	■ 1.000 ● N -1	$G(g\,p)=g\,G(p),\quad g\in G,\quad p\in P\,;$	$G(gp)=gG(p),\quad g\in G,\quad p\in P;$	$G(gp)=g\,G(p),\quad g\in G,\quad p\in P;$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0053 (2.7.3)	044	1.000 K +0	$\pi \circ G = \pi \text{ .}$	$\pi \circ G = \pi \text{ .}$	$\pi \circ G = \pi \text{ .}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0054 (2.7.4)	045	1.000 N +0	$\mathrm{G}_p \approx C^{\infty} (P \times_{\mathrm{Ad}} G) \text{ .}$	$\mathcal{G}_p \approx C^{\infty} (P \times_{\mathrm{Ad}} G) \text{ .}$	$\mathcal{G}_p \approx C^{\infty} (P \times_{\mathrm{Ad}} G) \text{ .}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0055 (2.7.5)	045	1.000 N -1	$G(p) = \exp i \theta^k(m) L_k \in \mathrm{G}_p \text{ .}$	$G(p) = \exp i \theta^k(m) L_k \in \mathcal{G}_p \text{ .}$	$G(p) = \exp i \theta^k(m) L_k \in \mathcal{G}_p \text{ .}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0057 (2.7.7)	045	1.000 K +0	$\varphi \rightarrow^{G^{-1}} \varphi := G^{-1} \varphi \in C^{\infty}(M, \mathbb{C}), G \in \mathcal{G}_V,$	$\varphi \rightarrow^{G^{-1}} \varphi := G^{-1} \varphi \in C^{\infty}(M, \mathbb{C}), G \in \mathcal{G}_V,$	$\varphi \rightarrow^{G^{-1}} \varphi := G^{-1} \varphi \in C^{\infty}(M, \mathbb{C}), G \in \mathcal{G}_V,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0063 (2.8.4)	046	1.000 N +0	$p_k(V) = (-1)^k c_{2k}(V) \text{ .}$	$p_k(V^{\mathbb{R}}) = (-1)^k c_{2k}(V) \text{ .}$	$p_k(V^{\mathbb{R}}) = (-1)^k c_{2k}(V) \text{ .}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0066 (2.8.9)	047	1.000 N +0	$\mathrm{Tr}(\Omega \wedge \Omega) = d \mathrm{Tr} \left(\Omega \wedge \Omega + \frac{1}{3} \Omega \wedge \Omega \wedge \Omega \right)$	$\mathrm{Tr}(\Omega \wedge \Omega) = d \mathrm{Tr} \left(\omega \wedge \Omega + \frac{1}{3} \omega \wedge \omega \wedge \omega \right)$	$\mathrm{Tr}(\Omega \wedge \Omega) = d \mathrm{Tr} \left(\omega \wedge \Omega + \frac{1}{3} \omega \wedge \omega \wedge \omega \right)$
Conf. MathPix confidence: 1.000 0.5-0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00067 (2.8.10)	048	■ 1.000 ● C +7	$\begin{aligned} & \operatorname{Tr}\left\{d\left(\omega\wedge\omega-\frac{2}{3}\omega\wedge\omega\wedge\omega\right)\right\}=\operatorname{Tr}\{d\omega\wedge d\omega-\omega\wedge dd\omega \\ & -\frac{2}{3}d\omega\wedge\omega\wedge\omega+\frac{2}{3}\omega\wedge d\omega\wedge\omega-\frac{2}{3}\omega\wedge\omega\wedge d\omega\} \\ & =\operatorname{Tr}\{d\omega\wedge d\omega-d\omega\wedge\omega\wedge\omega-\omega\wedge\omega\wedge d\omega\} \\ & =\operatorname{Tr}\{(d\omega-\omega\wedge\omega)\wedge(d\omega-\omega\wedge\omega)\}=\operatorname{Tr}(\Omega\wedge\Omega). \end{aligned}$	$\begin{aligned} & \operatorname{Tr}\left\{d\left(\omega\wedge d\omega-\frac{2}{3}\omega\wedge\omega\wedge\omega\right)\right\}=\operatorname{Tr}\{d\omega\wedge d\omega-\omega\wedge dd\omega \\ & -\frac{2}{3}d\omega\wedge\omega\wedge\omega+\frac{2}{3}\omega\wedge d\omega\wedge\omega-\frac{2}{3}\omega\wedge\omega\wedge d\omega\} \\ & =\operatorname{Tr}\{d\omega\wedge d\omega-d\omega\wedge\omega\wedge\omega-\omega\wedge\omega\wedge d\omega\} \\ & =\operatorname{Tr}\{(d\omega-\omega\wedge\omega)\wedge(d\omega-\omega\wedge\omega)\}=\operatorname{Tr}(\Omega\wedge\Omega) \end{aligned}$	$\begin{aligned} & \operatorname{Tr}\left\{d\left(\omega\wedge d\omega-\frac{2}{3}\omega\wedge\omega\wedge\omega\right)\right\}=\operatorname{Tr}\{d\omega\wedge d\omega-\omega\wedge dd\omega \\ & -\frac{2}{3}d\omega\wedge\omega\wedge\omega+\frac{2}{3}\omega\wedge d\omega\wedge\omega-\frac{2}{3}\omega\wedge\omega\wedge d\omega\} \\ & =\operatorname{Tr}\{d\omega\wedge d\omega-d\omega\wedge\omega\wedge\omega-\omega\wedge\omega\wedge d\omega\} \\ & =\operatorname{Tr}\{(d\omega-\omega\wedge\omega)\wedge(d\omega-\omega\wedge\omega)\}=\operatorname{Tr}(\Omega\wedge\Omega). \end{aligned}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00068 (2.8.11)	048	■ 1.000 ● K +0	$\operatorname{Tr}\{\omega\wedge(\omega\wedge\omega\wedge\omega)\}=-\operatorname{Tr}\{(\omega\wedge\omega\wedge\omega)\wedge\omega\}=0.$	$\operatorname{Tr}\{\omega\wedge(\omega\wedge\omega\wedge\omega)\}=-\operatorname{Tr}\{(\omega\wedge\omega\wedge\omega)\wedge\omega\}=0.$	$\operatorname{Tr}\{\omega\wedge(\omega\wedge\omega\wedge\omega)\}=-\operatorname{Tr}\{(\omega\wedge\omega\wedge\omega)\wedge\omega\}=0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00070 (2.8.13)	048	■ 1.000 ● N -1	$S_{\mathrm{YM}}=\int_M L_{\mathrm{YM}}$	$S_{\mathrm{YM}}=\int_M L_{\mathrm{YM}}$	$S_{\mathrm{YM}}=\int_M L_{\mathrm{YM}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00072 (2.8.15)	048	■ 1.000 ● K +0	$\omega\sim\omega$	$\omega\sim\omega$	$\omega\sim\omega$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00074 (2.8.17)	049	■ 1.000 ● K +0	$\Omega(\omega)=0.$	$\Omega(\omega)=0.$	$\Omega(\omega)=0.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0075 (2.8.18)	049	■ 1.000 ● N +0	$c_2(M)=\frac{1}{24\pi^2}\int_{\partial M}\mathrm{Tr}(\stackrel{\infty}{\omega}\wedge\stackrel{\infty}{\omega}\wedge\stackrel{\infty}{\omega}),$	$c_2(M)=\frac{1}{24\pi^2}\int_{\partial M}\mathrm{Tr}(\overset{\infty}{\omega}\wedge\overset{\infty}{\omega}\wedge\overset{\infty}{\omega}),$	$c_2(M)=\frac{1}{24\pi^2}\int_{\partial M}\mathrm{Tr}(\overset{\infty}{\omega}\wedge\overset{\infty}{\omega}\wedge\overset{\infty}{\omega}),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0076 (3.1.1)	052	■ 1.000 ● K +0	$L=L(\varphi,d\varphi)\eta$	$L=L(\varphi,d\varphi)\eta$	$L=L(\varphi,d\varphi)\eta$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0077 (3.1.2)	052	■ 1.000 ● N +0	$\delta\int_M L=0$	$\delta\int_M L=0$	$\delta\int_M L=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0078 (3.1.3)	052	■ 1.000 ● N -1	$\delta\varphi=0\quad\text{at the boundary of }\mathrm{M}$	$\delta\varphi=0\quad\text{at the boundary }\partial\mathrm{M}\text{ of }\mathrm{M}$	$\delta\varphi=0\quad\text{at the boundary }\partial\mathrm{M}\text{ of }\mathrm{M}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0079 (3.1.4)	052	■ 1.000 ● K +0	$\delta(d\varphi):=d(\varphi+\delta\varphi)-d\varphi=d\delta\varphi$	$\delta(d\varphi):=d(\varphi+\delta\varphi)-d\varphi=d\delta\varphi$	$\delta(d\varphi):=d(\varphi+\delta\varphi)-d\varphi=d\delta\varphi$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0080 (3.1.5)	052	■ 1.000 ● K +0	$\pi := \frac{\partial L}{\partial (\partial_\alpha \varphi)} \vartheta^\alpha$	$\pi := \frac{\partial L}{\partial (\partial_\alpha \varphi)} \vartheta^\alpha$	$\pi := \frac{\partial L}{\partial (\partial_\alpha \varphi)} \vartheta^\alpha$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0082 (3.1.7)	053	■ 1.000 ● N +1	$\int_M d(\delta \varphi \wedge \pi) = \int_{\partial M} \delta \varphi \wedge \pi$	$\int_M d(\delta \varphi \wedge \pi) = \int_{\partial M} \delta \varphi \wedge \pi$	$\int_M d(\delta \varphi \wedge \pi) = \int_{\partial M} \delta \varphi \wedge \pi.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0083 (3.1.9)	053	■ 1.000 ● N +0	$\lambda^\varphi := (-1)^s L_\varphi + {}^* d\pi = 0,$	$\lambda^\varphi := (-1)^s L_\varphi + {}^* d\pi = 0,$	$\lambda^\varphi := (-1)^s L_\varphi + {}^* d\pi = 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0084 (3.1.10)	053	■ 1.000 ● K +0	$\frac{\partial L}{\partial \varphi} - \partial_\alpha \frac{\partial L}{\partial (\partial_\alpha \varphi)} = 0.$	$\frac{\partial L}{\partial \varphi} - \partial_\alpha \frac{\partial L}{\partial (\partial_\alpha \varphi)} = 0.$	$\frac{\partial L}{\partial \varphi} - \partial_\alpha \frac{\partial L}{\partial (\partial_\alpha \varphi)} = 0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0085 (3.1.11)	053	■ 1.000 ● K +0	${}^* H := L - d\varphi \wedge \pi$	${}^* H := L - d\varphi \wedge \pi$	${}^* H := L - d\varphi \wedge \pi$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0086 (3.1.12)	053	■ 1.000 ● N +0	$d^{*}\mathrm{H}=\mathrm{d}\,L+\mathrm{d}\,\varphi\wedge L_{\varphi}+\mathrm{d}\,\varphi\wedge d\pi-d\,d\varphi\wedge\pi$	$d^{*}H=dL+d\varphi\wedge^{*}L_{\varphi}+d\varphi\wedge d\pi-dd\varphi\wedge\pi$	$d^{*}H=dL+d\varphi\wedge^{*}L_{\varphi}+d\varphi\wedge d\pi-d\,d\varphi\wedge\pi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0087 (3.1.13)	053	■ 1.000 ● K +0	$d^{*}\mathrm{H}\simeq 0$	$d^{*}H\simeq 0$	$d^{*}H\simeq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0088 (3.1.14)	054	■ 1.000 ● N +0	$\beta\wedge D\psi=m\psi\eta.$	$\beta\wedge^{*}D\psi=m\psi\eta.$	$\beta\wedge^{*}D\psi=m\psi\eta.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0089 (3.1.15)	054	■ 1.000 ● N +0	$\square\varphi+m^2\varphi=0$	$\square\varphi+m^2\varphi=0$	$\square\varphi+m^2\varphi=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0090 (3.1.16)	054	■ 1.000 ● W -1	$\overline{\mathrm{L}}(\overline{\psi}\psi,\overline{\psi}\beta\cdot d\psi)\eta=\overline{\psi}\beta\wedge^{*}d\psi-m\overline{\psi}\psi\eta.$	$\overline{L}_{\mathrm{mat}}=\overline{L}_{\mathrm{mat}}(\overline{\psi}\psi,\overline{\psi}\beta\cdot d\psi)\eta=\overline{\psi}\beta\wedge^{*}d\psi-m\overline{\psi}\psi\eta.$	$\overline{\overline{L}}_{\mathrm{mat}}=\overline{\overline{L}}_{\mathrm{mat}}(\overline{\overline{\psi}}\overline{\overline{\psi}},\overline{\overline{\psi}}\overline{\overline{\beta}}\cdot d\overline{\overline{\psi}})\eta=\overline{\overline{\psi}}\overline{\overline{\beta}}\wedge^{*}d\overline{\overline{\psi}}-m\overline{\overline{\psi}}\overline{\overline{\psi}}\eta.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0092 (3.2.2)	055	■ 1.000 ● N -1	$L(\overline{\varphi}\varphi,\overline{\varphi}\beta\cdot(d+\omega')\varphi)\eta=L(\overline{\varphi}\varphi,\overline{\varphi}\beta\cdot D'\varphi)\eta$	$L_{\mathrm{mat}}=L(\overline{\varphi}\varphi,\overline{\varphi}\beta\cdot(d+\omega')\varphi)\eta=L(\overline{\varphi}\varphi,\overline{\varphi}\beta\cdot D'\varphi)\eta$	$L_{\mathrm{mat}}=L(\overline{\varphi}\varphi,\overline{\varphi}\beta\cdot(d+\omega')\varphi)\eta=L(\overline{\varphi}\varphi,\overline{\varphi}\beta\cdot D'\varphi)\eta$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0094 (3.2.4)	055	■ 1.000 ● K +0	$D\left(\left\{ \right\} ^{G}b\right) = \left\{ \right\} ^{G} \omega \otimes \left\{ \right\} ^{G} b \; .$	$D\left({}^G b\right) = {}^G \omega \otimes {}^G b.$	$D({}^G b) = {}^G \omega \otimes {}^G b.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0095 (3.2.5)	055	■ 1.000 ● W +0	$\begin{aligned} D\left(\left\{ \right\} ^{G}b\right) &= dG \otimes b + G \omega \otimes b + G \omega \otimes b \; \& \; = \left\{ dG G^{-1} + G \omega G^{-1} \right\} \otimes {}^G b \; . \end{aligned}$	$\begin{aligned} D\left({}^G b\right) &= dG \otimes b + G \omega \otimes b \\ &= \left\{ dG G^{-1} + G \omega G^{-1} \right\} \otimes {}^G b. \end{aligned}$	$\begin{aligned} D({}^G b) &= dG \otimes b + G \omega \otimes b \\ &= \left\{ dG G^{-1} + G \omega G^{-1} \right\} \otimes {}^G b. \end{aligned}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0096 (3.2.6)	055	■ 1.000 ● N +0	$\omega \rightarrow \left\{ \right\} ^{G} \omega = G \omega G^{-1} + dG G^{-1} \; .$	$\omega \rightarrow {}^G \omega = G \omega G^{-1} + dG G^{-1}.$	$\omega \rightarrow {}^G \omega = G \omega G^{-1} + dG G^{-1}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0102 (3.3.3)	057	■ 1.000 ● N +1	$\theta = - \frac{e}{\hbar} \int \mathbf{A} \cdot d\mathbf{l}$	$\theta = - \frac{e}{\hbar} \int \mathbf{A} \cdot d\mathbf{l}$	$\theta = - \frac{e}{\hbar} \int \mathbf{A} \cdot d\mathbf{l}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0103 (3.3.4)	057	■ 1.000 ● N -2	$L = L(A, dA) + L_{\mathrm{mat}}(A, dA, \Psi, d\Psi) \; .$	$L = L(A, dA) + L_{\mathrm{mat}}(A, dA, \Psi, d\Psi).$	$L = L(A, dA) + L_{\mathrm{mat}}(A, dA, \Psi, d\Psi).$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0104 (3.3.5)	057	■ 1.000 ● N -1	$\frac{\delta L}{\delta A} := \frac{\partial L}{\partial A} + d \frac{\partial L}{\partial dA} = 0,$	$\frac{\delta L}{\delta A} := \frac{\partial L}{\partial A} + d \frac{\partial L}{\partial dA} = 0,$	$\frac{\delta L}{\delta A} := \frac{\partial L}{\partial A} + d \frac{\partial L}{\partial dA} = 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0106 (3.3.7)	058	■ 1.000 ● N +1	$dF \equiv 0, \quad dH = j$	$dF \equiv 0, \quad dH = j$	$dF \equiv 0, \quad dH = j.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0107 (3.3.8)	058	■ 1.000 ● K +0	$dj \simeq 0.$	$dj \simeq 0.$	$dj \simeq 0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0111 (3.3.12)	058	■ 1.000 ● N -1	$H := -\mathfrak{H} \wedge dt + \mathfrak{D} = \frac{1}{2} H_{ij} dx^i \wedge dx^j$	$H := -\mathfrak{H} \wedge dt + \mathfrak{D} = \frac{1}{2} H_{ij} dx^i \wedge dx^j$	$H := -\mathfrak{H} \wedge dt + \mathfrak{D} = \frac{1}{2} H_{ij} dx^i \wedge dx^j,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0114 (3.3.15)	059	■ 1.000 ● W +1	$H_{mn} = \frac{\sqrt{ g }}{2} \varepsilon_{mnkl} g^{ki} g^{lj} F_{ij}.$	$H_{mn} = \frac{\sqrt{ g }}{2} \varepsilon_{mnkl} g^{ki} g^{lj} F_{ij}.$	$H_{mn} = \frac{\sqrt{ g }}{2} \varepsilon_{mnkl} g^{ki} g^{lj} F_{ij}.$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5-0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0121 (3.3.22)	060	■ 1.000 ● K +0	$L_{\mathrm{BF}}=-B\wedge F=-B\wedge dA\ .$	$L_{\mathrm{BF}}=-B\wedge F=-B\wedge dA.$	$L_{\mathrm{BF}}=-B\wedge F=-B\wedge dA\ .$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0122 (3.3.23)	060	■ 1.000 ● N -1	$\begin{aligned} \tilde{L}_{\mathrm{BF}} &:= -B\wedge dA + \frac{1}{2}B\wedge B \\ &\cong -B\wedge dA + dC \end{aligned}$	$\tilde{L}_{\mathrm{BF}} := -B\wedge dA + \frac{1}{2}B\wedge B \\ \cong -B\wedge dA + dC$	$\tilde{L}_{\mathrm{BF}} := -B\wedge dA + \frac{1}{2}B\wedge B \\ \cong -B\wedge dA + dC\ .$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0123 (3.3.24)	060	■ 1.000 ● N +0	$B\congd A,\quad d B\congd F\equiv 0$	$B\cong dA,\quad dB\cong dF\equiv 0$	$B\cong dA,\quad dB\cong dF\equiv 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0124 (3.3.25)	060	■ 1.000 ● K +0	$C=\frac{1}{2}A\wedge F,\quad d C=\frac{1}{2}F\wedge F,$	$C=\frac{1}{2}A\wedge F,\quad dC=\frac{1}{2}F\wedge F,$	$C=\frac{1}{2}A\wedge F,\quad dC=\frac{1}{2}F\wedge F,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0125 (3.3.26)	061	■ 1.000 ● K +0	$A\rightarrowtail A^{\mathrm{\prime}}=A+d\,\theta(x)\ .$	$A\rightarrowtail A^{\prime}=A+d\theta(x).$	$A\rightarrowtail A^{\prime}=A+d\theta(x).$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0126 (3.3.27)	061	1.000 N +0	$A \rightarrow \widetilde{A} = A + \psi, \quad B \rightarrow \widetilde{B} = B + d\psi,$	$A \rightarrow \widetilde{A} = A + \psi, \quad B \rightarrow \widetilde{B} = B + d\psi,$	$A \rightarrow \widetilde{A} = A + \psi, \quad B \rightarrow \widetilde{B} = B + d\psi,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0127 (3.3.28)	061	1.000 K +0	$L_{\mathrm{Max}} = -B \wedge dA + \frac{1}{2} B \wedge {}^*B + L_{\mathrm{matter}},$	$L_{\mathrm{Max}} = -B \wedge dA + \frac{1}{2} B \wedge {}^*B + L_{\mathrm{matter}},$	$L_{\mathrm{Max}} = -B \wedge dA + \frac{1}{2} B \wedge {}^*B + L_{\mathrm{matter}},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0128 (3.3.29)	061	1.000 N +0	$-d B = d^* {}^* F \cong j,$	$-dB = d^* F \cong j,$	$-dB = d^* F \cong j,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0132 (3.4.3)	063	1.000 N +0	$\alpha_g := g^2 / \hbar c$	$\alpha_g := g^2 / \hbar c$	$\alpha_g := g^2 / \hbar c$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0134 (3.4.5)	063	1.000 K +0	$\delta L_{\omega} =: \delta \Omega \wedge \Pi^{\Omega}.$	$\delta L_{\omega} =: \delta \Omega \wedge \Pi^{\Omega}.$	$\delta L_{\omega} =: \delta \Omega \wedge \Pi^{\Omega}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0138 (3.4.9)	063	1.000 N -1	$L = L_{\mathrm{mat}} + L_{\omega},$	$L = L_{\mathrm{mat}} + L_{\omega},$	$L = L_{\mathrm{mat}} + L_{\omega},$
Conf. MathPix confidence: ≥0.9 0.5-0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0139 (3.4.10)	063	■ 1.000 ● K +0	$\delta L_{\mathrm{mat}} =: \delta \omega \wedge \tau$.	$\delta L_{\mathrm{mat}} =: \delta \omega \wedge \tau$.	$\delta L_{\mathrm{mat}} =: \delta \omega \wedge \tau$.
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0140 (3.4.11)	064	■ 1.000 ● N +2	$D \Pi^{\omega} = \tau$	$D \Pi^{\Omega} = \tau$	$\underline{D \Pi^{\Omega} = \tau}$.
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0141 (3.4.12)	064	■ 1.000 ● N +0	$D \Omega \equiv 0$.	$D \Omega \equiv 0$.	$D \Omega \equiv 0$.
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0145 (3.4.16)	064	■ 1.000 ● N +0	$d \Pi^{\omega} + \left[\omega, \Pi^{\Omega} \right] = \tau$.	$d \Pi^{\Omega} + \left[\omega, \Pi^{\Omega} \right] = \tau$.	$d \Pi^{\Omega} + \left[\omega, \Pi^{\Omega} \right] = \tau$.
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0147 (3.4.18)	065	■ 1.000 ● N +0	$Q_e = \frac{1}{2} Y + I_3$.	$Q_e = \frac{1}{2} Y + I_3$.	$Q_e = \frac{1}{2} Y + I_3$.
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0148 (3.4.19)	065	■ 1.000 ● N +0	$L_{\mathrm{YM}} = - \frac{1}{\alpha_g} \operatorname{Tr} \left(\Omega \wedge {}^* \Omega \right)$.	$L_{\mathrm{YM}} = - \frac{1}{\alpha_g} \operatorname{Tr} (\Omega \wedge {}^* \Omega)$.	$L_{\mathrm{YM}} = - \frac{1}{\alpha_g} \operatorname{Tr} (\Omega \wedge {}^* \Omega)$.
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0150 (3.4.21)	065	■ 1.000 ● N +1	$\backslash\mathrm{Pi}^{\backslash\mathrm{Omega}}=-\frac{2}{\backslash\mathrm{alpha}_{\mathrm{g}}}\backslash\mathrm{Omega}$	$\Pi^{\Omega}=-\frac{2}{\alpha_g}*\Omega$	$\Pi^{\Omega}=-\frac{2}{\alpha_g}*\Omega.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0153 (3.5.1)	066	■ 1.000 ● N +1	$\backslash\mathrm{Pi}^{\backslash\mathrm{Omega}}=\backslash\mathrm{zeta}\ \backslash\mathrm{Omega}\ .$	$\Pi^{\Omega}=\zeta\Omega.$	$\Pi^{\Omega}=\zeta\Omega.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0156 (3.5.4)	067	■ 1.000 ● N +0	$\backslash\mathrm{sigma}^{\backslash\mathrm{mu}\ \backslash\mathrm{nu}}=\backslash\mathrm{sigma}^{\backslash\mathrm{mu}\ \backslash\mathrm{nu}},\ \backslash\mathrm{quad}\backslash\mathrm{sigma}^{\backslash\mathrm{mu}\ \backslash\mathrm{nu}}=-\backslash\mathrm{bar}\{\backslash\mathrm{sigma}\}^{\backslash\mathrm{mu}\ \backslash\mathrm{nu}},$	$*\sigma^{\mu\nu}=\sigma^{\mu\nu},\quad *\bar{\sigma}^{\mu\nu}=-\bar{\sigma}^{\mu\nu},$	$*\sigma^{\mu\nu}=\sigma^{\mu\nu},\quad *\bar{\sigma}^{\mu\nu}=-\bar{\sigma}^{\mu\nu},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0158 (3.5.6)	067	■ 1.000 ● N +0	$A_{\backslash\mathrm{mu}}=i\ \backslash\mathrm{sigma}_{\backslash\mathrm{mu}\ i}\ \backslash\mathrm{partial}^{\backslash\mathrm{i}}\ \backslash\mathrm{ln}\ h\ .$	$A_{\mu}=i\sigma_{\mu i}\partial^i\ln h.$	$A_{\mu}=i\sigma_{\mu i}\partial^i\ln h.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0159 (3.5.7)	067	■ 1.000 ● N +0	$\backslash\mathrm{square}\ h^{-1}\ \backslash\mathrm{partial}_{\backslash\mathrm{i}}\ h\ \backslash\mathrm{partial}^{\backslash\mathrm{i}}\ h=h^3$	$\square h-h^{-1}\partial_i h\partial^i h=h^3$	$\square h-h^{-1}\partial_i h\partial^i h=h^3$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0160 (3.5.8)	067	■ 1.000 ● W +0	$h=\sum_{i=1}^{k+1} \frac{l_{(i)}^2}{\left(x-x_{(i)}\right)^2},$	$h=\sum_{i=1}^{k+1} \frac{l_{(i)}^2}{\left(x-x_{(i)}\right)^2},$	$h=\sum_{i=1}^{k+1} \frac{l_{(i)}^2}{\left(x-x_{(i)}\right)^2},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0161 (3.5.9)	067	■ 1.000 ● K +0	$p_{\{1\}}(M)=k \text{ .}$	$p_1(M)=k.$	$p_1(M)=k.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0162 (3.5.10)	068	■ 1.000 ● N +2	$\langle \infty -\infty \rangle = Z \int_{\mathcal{M}_\omega} d\omega \mu[\omega] \exp i \int_M (L_\omega + L_\infty).$	$\langle \infty -\infty \rangle = Z \int_{\mathcal{M}_\omega} d\omega \mu[\omega] \exp i \int_M (L_\omega + L_\infty)$	$\langle \infty -\infty \rangle = Z \int_{\mathcal{M}_\omega} d\omega \mu[\omega] \exp i \int_M (L_\omega + L_\infty).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0163 (3.5.11)	068	■ 1.000 ● N +0	$L_{\{\infty\}}=(\stackrel{+}{-})\frac{1}{\alpha_g}\operatorname{Tr}(\Omega\wedge\Omega)$	$L_\infty=(\stackrel{+}{-})\frac{1}{\alpha_g}\operatorname{Tr}(\Omega\wedge\Omega)$	$L_\infty=(\stackrel{+}{-})\frac{1}{\alpha_g}\operatorname{Tr}(\Omega\wedge\Omega)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0167 (3.6.2)	070	■ 1.000 ● N +0	$\mathscr{P}_{\{ \pm \}}:=\bar{\psi} \sigma_{\pm} \psi / 2 \text{ m},$	$\mathscr{P}_{\pm}:=\bar{\psi} \sigma_{\pm} \psi / 2 m,$	$\mathscr{P}_{\pm}:=\bar{\psi} \sigma_{\pm} \psi / 2 m,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0168 (3.6.3)	070	■ 1.000 ● N +0	$D_{\{ \pm \}}^{\{ \}} D_{\pm} \psi \mp i \left(F^{\pm} - \frac{1}{2} \bar{\psi} \sigma_{\pm} \psi \right) \wedge \sigma_{\pm} \psi = 0$	$D_{\pm}^* D_{\pm} \psi \mp i \left(F^{\pm} - \frac{1}{2} \bar{\psi} \sigma_{\pm} \psi \right) \wedge \sigma_{\pm} \psi = 0$	$D_{\pm}^* D_{\pm} \psi \mp i \left(F^{\pm} - \frac{1}{2} \bar{\psi} \sigma_{\pm} \psi \right) \wedge \sigma_{\pm} \psi = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0170 (3.6.5)	070	■ 1.000 ● N +0	$D^{\{ \}} D \psi = \left(\nabla^* \nabla + \frac{1}{4} R + \frac{1}{2} F^+ \right) \psi.$	$D^* D \psi = \left(\nabla^* \nabla + \frac{1}{4} R + \frac{1}{2} F^+ \right) \psi.$	$D^* D \psi = \left(\nabla^* \nabla + \frac{1}{4} R + \frac{1}{2} F^+ \right) \psi.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0172 (3.6.7)	070	■ 1.000 ● N +0	$\Omega_{\pm}^g = \frac{1}{2\ell^2} \sigma_{\pm},$	$\Omega_{\pm}^g = \frac{1}{2\ell^2} \sigma_{\pm},$	$\Omega_{\pm}^g = \frac{1}{2\ell^2} \sigma_{\pm},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0176 (3.7.5)	072	■ 1.000 ● N +0	$\begin{aligned} & D^{\{ \}} \Theta^{\{ (1) \} - \frac{\lambda}{2} \left(\frac{\mu^2}{\lambda} - \bar{\varphi} \cdot \varphi \right) \varphi \eta = 0, \\ & D \Pi^{\Omega} + \left[\phi^{(0)}, * \Theta^{(1)} \right] = 0. \end{aligned}$	$D^* \Theta^{(1)} - \frac{\lambda}{2} \left(\frac{\mu^2}{\lambda} - \bar{\varphi} \cdot \varphi \right) \varphi \eta = 0,$ $D \Pi^{\Omega} + \left[\phi^{(0)}, * \Theta^{(1)} \right] = 0.$	$D^* \Theta^{(1)} - \frac{\lambda}{2} \left(\frac{\mu^2}{\lambda} - \bar{\varphi} \cdot \varphi \right) \varphi \eta = 0,$ $D \Pi^{\Omega} + [\phi^{(0)}, * \Theta^{(1)}] = 0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ __Z-Library__ EQ0177 (3.7.7)	072	■ 1.000 ● N +0	$\bar{\varphi} \cdot \varphi := \varphi^{*j} \varphi_j = \frac{\mu^2}{\lambda}.$	$\bar{\varphi} \cdot \varphi := \varphi^{*j} \varphi_j = \frac{\mu^2}{\lambda}.$	$\bar{\varphi} \cdot \varphi := \varphi^{*j} \varphi_j = \frac{\mu^2}{\lambda}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0179 (3.7.9)	072	■ 1.000 ● N -1	$\varphi_{\min }: P \rightarrow C^{\infty }\left(\stackrel{\circ}{V}(M, \rho (G/H), \rho (G), P) \subset V^{\rho }\right).$	$\varphi_{\min }: P \rightarrow C^{\infty }\left(\stackrel{\circ}{V}(M, \rho (G/H), \rho (G), P) \subset V^{\rho }\right).$	$\varphi_{\min }: P \rightarrow C^{\infty }\left(\stackrel{\circ}{V}(M, \rho (G/H), \rho (G), P) \subset V^{\rho }\right).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0180 (3.7.10)	073	■ 1.000 ● N +0	$H:=\left\{h\in G\mid \rho (h)\mathrm{v}_{0}=\mathrm{v}_{0}\right\},$	$H:=\left\{h\in G\mid \rho (h)\mathbf{v}_0=\mathbf{v}_0\right\},$	$H:=\left\{h\in G\mid \rho (h)\mathbf{v}_0=\mathbf{v}_0\right\},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0181 (3.7.11)	073	■ 1.000 ● N -2	$Q=\left\{p\in P\mid \varphi (p)=\sigma \left(\mathrm{v}_{0}\right)\right\}$	$Q=\left\{p\in P\mid \varphi (p)=\sigma \left(\mathbf{v}_0\right)\right\}$	$Q=\left\{p\in P\mid \varphi (p)=\sigma \left(\mathbf{v}_0\right)\right\}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0182 (3.7.12)	073	■ 1.000 ● N +0	$D\,\phi^{\{0\}}=0\;.$	$D\phi^{(0)}=0.$	$D\phi^{(0)}=0.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0183 (3.7.13)	073	■ 1.000 ● K +0	$\begin{aligned}\Theta^{\{\perp\}}&:=\Omega_{\{0\}}\wedge d\Omega_{\{0\}}-d\Omega_{\{0\}}\wedge\Omega_{\{0\}}=d\Omega_{\{0\}}\wedge\Omega_{\{0\}}-d\Omega_{\{0\}}\wedge\Omega_{\{0\}}\\&=d\Omega_{\{0\}}-[\Omega_{\{0\}},\Omega_{\{0\}}]=D\Omega_{\{0\}}\end{aligned}$	$\Theta^{\perp}:=\Omega_0=d\omega_0-(\omega\wedge\omega)_0=d\omega_0-\omega_0\wedge\omega-\omega\wedge\omega_0\\=d\omega_0-[\omega_0,\omega]=D\omega_0$	$\Theta^{\perp}:=\Omega_0=d\omega_0-(\omega\wedge\omega)_0=d\omega_0-\omega_0\wedge\omega-\omega\wedge\omega_0\\=d\omega_0-[\omega_0,\omega]=D\omega_0$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0185 (3.7.15)	074	■ 1.000 ● K +0	$d_{\{0\}}\{ \}^{\{*\}} \backslash \Omega_{\{0\}}=0$.	$d_0^*\Omega_0=0.$	$d_0^*\Omega_0=0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0186 (3.7.16)	074	■ 1.000 ● N +0	$\left(D^{\{*\}} \backslash \Omega_{\mathrm{right}}\right)_{\{0\}}=\mathrm{vec}\{D\}^{\{*\}} \backslash \Omega_{\{0\}}=\mathrm{vec}\{D\}^{\{*\}} \backslash \Theta^{\{\perp\}}=0$	$(D^*\Omega)_0=\vec{D}^*\Omega_0=\vec{D}^*\Theta^\perp=0$	$(D^*\Omega)_0=\vec{D}^*\Omega_0=\vec{D}^*\Theta^\perp=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0188 (3.7.18)	074	■ 1.000 ● N -1	$\omega_{\{0\}}=\varphi^{\{j\}}L_{\{j\}} \backslash \otimes \vartheta_{\{0\}}$.	$\omega_0=\varphi^jL_j\otimes\vartheta_0.$	$\omega_0=\varphi^jL_j\otimes\vartheta_0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0190	080	■ 1.000 ● C +23	$\left[\begin{array}{l} T_{\{G\}}:\\quad T_{\{g\}} \backslash \psi(m)=D\left(g_{\{m\}}^{\{-1\}} g_{g_{\{g^{\{-1\}} m\}}, m_{\{0\}}\right)} \backslash \psi\left(g^{\{-1\}} m\right) \\ . \end{array}\right]$	<i>(not rendered)</i>	$T_G:\quad T_g\psi(m)=D\left(g_m^{-1}gg_{g^{-1}m},m_0\right)\psi(g^{-1}m).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0193 (4.1.3)	081	■ 1.000 ● N +0	$A(n,\mathbb{R})=\mathbb{R}^n\backslash \otimes GL(n,\mathbb{R})$	$A(n,\mathbb{R})=\mathbb{R}^n\otimes GL(n,\mathbb{R})$	$A(n,\mathbb{R})=\mathbb{R}^n\oslash GL(n,\mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ0194 (4.1.4)	081	■ 1.000 ● N +0	$P:=\mathbb{R}^n\backslash \otimes \Theta(1,n-1)\backslash \subset A(n,\mathbb{R}),$	$P:=\mathbb{R}^n\otimes \Theta(1,n-1)\subset A(n,\mathbb{R}),$	$P:=\mathbb{R}^n\oslash \Theta(1,n-1)\subset A(n,\mathbb{R}),$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0197 (4.1.10)	082	■ 1.000 ● N +0	$e_{\{\alpha\}}(m)=e_{\{\alpha\}}^{\{i\}}\,\partial_{\{i\}}$	$e_{\alpha}(m)=e_{\alpha}^i\partial_i$	$e_{\alpha}(m)=e_{\alpha}^i\partial_i$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0199 (4.1.13)	082	■ 1.000 ● N +0	$L(M)=\bigcup_{m\in M}L_{\{m\}}(M)$	$L(M)=\bigcup_{m\in M}L_m(M)$	$L(M)=\bigcup_{m\in M}L_m(M)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0200 (4.1.14)	083	■ 1.000 ● K +0	$L(M)=P(M,\,G\,L(n,\,\mathbb{R}),\,\pi,\,\delta)$	$L(M)=P(M, GL(n,\mathbb{R}),\pi,\delta)$	$L(M)=P(M, GL(n,\mathbb{R}),\pi,\delta)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0202 (4.1.16)	083	■ 1.000 ● K +0	$\vartheta(e)=0\,\Leftrightarrow\,\pi(e)=0,\,e\in T_{\{p\}}(P),$	$\vartheta(e)=0\Leftrightarrow\pi(e)=0,e\in T_p(P),$	$\vartheta(e)=0\Leftrightarrow\pi(e)=0,e\in T_p(P),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0204 (4.1.18)	083	■ 1.000 ● N +0	$T(M)=V\left(M,\,\mathbb{R}^{\{n\}},\,G\,L(n,\,\mathbb{R}),\,L(M)\right)$	$T(M)=V\left(M,\mathbb{R}^n, GL(n,\mathbb{R}), L(M)\right)$	$T(M)=V\left(M,\mathbb{R}^n, GL(n,\mathbb{R}), L(M)\right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0205 (4.1.19)	083	■ 1.000 ● N +0	$A(M):=P(M,\,A(n,\,\mathbb{R}),\,\pi,\,\delta)=L(M)\,\times\,\mathbb{R}^{\{n\}}.$	$A(M):=P(M,A(n,\mathbb{R}),\pi,\delta)=L(M)\times\mathbb{R}^n.$	$A(M):=P(M,A(n,\mathbb{R}),\pi,\delta)=L(M)\times\mathbb{R}^n.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0206 (4.1.20)	083	■ 1.000 ● N +0	$\beta: A(n, \mathbb{R}) \rightarrow GL(n, \mathbb{R}) = A(n, \mathbb{R})/\mathbb{R}^n.$		
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0213 (4.2.5)	086	■ 1.000 ● N +0	$\mathscr{A}(n, \mathbb{R}) := C^\infty(A(M) \times_{\text{Ad}} A(n, \mathbb{R})).$		$\mathscr{A}(n, \mathbb{R}) := C^\infty\Big(A(M) \times_{\text{Ad}} A(n, \mathbb{R})\Big).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0223 (4.2.16)	087	■ 1.000 ● N -2	$Q_A = \left\{p \in A(M) \mid \widetilde{\varphi}(p) = \sigma(0_m)\right\} = L(M)$		$Q_A = \{p \in A(M) \mid \widetilde{\varphi}(p) = \sigma(0_m)\} = L(M)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0237 (4.3.2)	090	■ 1.000 ● W +0	$\ell(\mathscr{C}) := \int_{t_0}^{t_1} ds(t)$		$\ell(\mathscr{C}) := \int_{t_0}^{t_1} ds(t)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0238 (4.3.3)	090	■ 1.000 ● K +0	$\ell(\mathscr{C}_1 + \mathscr{C}_2) = \ell(\mathscr{C}_1) + \ell(\mathscr{C}_2)$		$\ell(\mathscr{C}_1 + \mathscr{C}_2) = \ell(\mathscr{C}_1) + \ell(\mathscr{C}_2)$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0253 (4.3.18)	093	■ 1.000 ● N +0	$\partial_{\{k} g_{\{i j\}} \stackrel{(*)}{=} 0$.	$\partial_k g_{ij} \stackrel{(*)}{=} 0.$	$\partial_k g_{ij} \stackrel{(*)}{=} 0.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0254 (4.3.19)	093	■ 1.000 ● N +1	$\Omega^g = d \omega^g - \omega^g \wedge \omega^g$	$\Omega^g = d\omega^g - \omega^g \wedge \omega^g$	$\Omega^g = d\omega^g - \omega^g \wedge \omega^g$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0257 (4.3.22)	094	■ 1.000 ● K +0	$\omega^g = \omega^{\{ \} } - K$.	$\omega^g = \omega^{\{ \} } - K.$	$\omega^g = \omega^{\{ \} } - K.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0258 (4.3.23)	094	■ 1.000 ● K +0	$d \vartheta - [\omega^{\{ \} }, \vartheta] = 0$	$d\vartheta - \left[\omega^{\{ \} }, \vartheta \right] = 0$	$d\vartheta - [\omega^{\{ \} }, \vartheta] = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0259 (4.3.24)	094	■ 1.000 ● N +0	$\Theta = [K, \vartheta]$	$\Theta = [K, \vartheta]$	$\Theta = [K, \vartheta]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0265 (4.3.30)	095	■ 1.000 ● W +0	$\left\{ \begin{array}{c} \kappa \\ ij \end{array} \right\} = \frac{1}{2} g^{\kappa l} (\partial_i g_{jl} + \partial_j g_{il} - \partial_l g_{ij})$	$\left\{ \begin{array}{c} \kappa \\ ij \end{array} \right\} = \frac{1}{2} g^{\kappa l} (\partial_i g_{jl} + \partial_j g_{il} - \partial_l g_{ij})$	$\left\{ \begin{array}{c} \kappa \\ ij \end{array} \right\} = \frac{1}{2} g^{\kappa l} (\partial_i g_{jl} + \partial_j g_{il} - \partial_l g_{ij})$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0272 (4.4.1)	096	■ 1.000 ● N -1	$L_{\mathrm{g}}=\mathcal{L}\left(g_{\alpha\beta}\left(\vartheta\right);\vartheta,\Theta,\Omega^g\right).$	$L_g=L\left(g_{\alpha\beta};\vartheta,\left(\omega^g\right),\Theta,\Omega^g\right).$	$L_g=L(g_{\alpha\beta};\vartheta,(\omega^g),\Theta,\Omega^g).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0273 (4.4.2)	097	■ 1.000 ● N +0	$\delta L_{\mathrm{g}}=\delta\vartheta\wedge E+\delta\Theta\wedge\Pi^{\mathrm{T}}+\delta\Omega^g\wedge\Pi^{\mathrm{L}},$	$\delta L_g=\delta\vartheta\wedge E+\delta\Theta\wedge\Pi^{\mathrm{T}}+\delta\Omega^g\wedge\Pi^{\mathrm{L}},$	$\delta L_g=\delta\vartheta\wedge E+\delta\Theta\wedge\Pi^{\mathrm{T}}+\delta\Omega^g\wedge\Pi^{\mathrm{L}},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0274 (4.4.7)	097	■ 1.000 ● K +0	$\delta\Theta=\delta\left(d\vartheta-\left[\omega^{\mathrm{L}},\vartheta\right]\right)$	$\delta\Theta=\delta\left(d\vartheta-\left[\omega^{\mathrm{L}},\vartheta\right]\right)$	$\delta\Theta=\delta(d\vartheta-[\omega^{\mathrm{L}},\vartheta])$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0277 (4.4.5)	097	■ 1.000 ● N +0	$S=\int_M\left(L_{\mathrm{g}}+L_{\mathrm{mat}}\right)$	$S=\int_M(L_g+L_{\mathrm{mat}})$	$S=\int_M(L_g+L_{\mathrm{mat}})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0279 (4.4.8)	098	■ 1.000 ● N +0	$\delta L_{\mathrm{matter}}=\delta\vartheta\wedge\Sigma+\delta\omega^g\wedge\tau_s+\delta\varphi\wedge\lambda^\varphi$	$\delta L_{\mathrm{matter}}=\delta\vartheta\wedge\Sigma+\delta\omega^g\wedge\tau_s+\delta\varphi\wedge\lambda^\varphi$	$\delta L_{\mathrm{matter}}=\delta\vartheta\wedge\Sigma+\delta\omega^g\wedge\tau_s+\delta\varphi\wedge\lambda^\varphi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0280 (4.4.9)	098	■ 1.000 ● N +0	$D\Theta\equiv\left[\vartheta,\Omega^{\mathrm{L}}\right]$	$D\Theta\equiv\left[\vartheta,\Omega^{\mathrm{L}}\right]$	$D\Theta\equiv[\vartheta,\Omega^{\mathrm{L}}]$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0281 (4.4.10)	098	■ 1.000 ● N +0	$D\,\Omega^g\equiv 0,$	$D\Omega^g\equiv 0,$	$D\Omega^g\equiv 0,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0283 (4.4.12)	099	■ 1.000 ● N +0	$D[\Theta,\vartheta]\equiv\left[\vartheta\wedge\vartheta,\Omega^g\right]$	$D[\Theta,\vartheta]\equiv[\vartheta\wedge\vartheta,\Omega^g]$	$D[\Theta,\vartheta]\equiv[\vartheta\wedge\vartheta,\Omega^g]$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0289 (4.4.18)	100	■ 1.000 ● N +0	$\left[\Omega^g,\Pi^L\right]+\mathrm{E}_{\square}=0$	$\left[\Omega^g,\Pi^L\right]+\mathrm{E}_{\square}=0$	$[\Omega^g,\Pi^L]+\mathrm{E}_{\square}=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0290 (4.5.1)	100	■ 1.000 ● N +0	$L_{\mathrm{mat}}=L(g_{\alpha\beta},\vartheta^\alpha,\Psi,D\Psi).$	$L_{\mathrm{mat}}=L(g_{\alpha\beta},\vartheta^\alpha,\Psi,D\Psi).$	$L_{\mathrm{mat}}=L(g_{\alpha\beta},\vartheta^\alpha,\Psi,D\Psi).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0291 (4.5.2)	100	■ 1.000 ● W -1	$\Sigma_{\alpha}:=\frac{\partial L}{\partial\vartheta^\alpha},\quad\sigma^{\alpha\beta}:=2\frac{\partial L}{\partial g_{\alpha\beta}},\quad\text{and}\quad\tau_{\alpha\beta}:=\frac{\partial L}{\partial\Gamma^{\alpha\beta}}=I_{\alpha\beta}\Psi\wedge\frac{\partial L}{\partial D\Psi}$	$\Sigma_{\alpha}:=\frac{\partial L}{\partial\vartheta^\alpha},\quad\sigma^{\alpha\beta}:=2\frac{\partial L}{\partial g_{\alpha\beta}},\quad\text{and}\quad\tau_{\alpha\beta}:=\frac{\partial L}{\partial\Gamma^{\alpha\beta}}=I_{\alpha\beta}\Psi\wedge\frac{\partial L}{\partial D\Psi}$	$\Sigma_{\alpha}:=\frac{\partial L}{\partial\vartheta^\alpha},\quad\sigma^{\alpha\beta}:=2\frac{\partial L}{\partial g_{\alpha\beta}},\quad\text{and}\quad\tau_{\alpha\beta}:=\frac{\partial L}{\partial\Gamma^{\alpha\beta}}=I_{\alpha\beta}\Psi\wedge\frac{\partial L}{\partial D\Psi}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0297 (8.10)	101	■1.000 ● N +0	$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha} + \delta \Psi \wedge \frac{\partial L}{\partial \Psi} + \delta(D\Psi) \wedge \frac{\partial L}{\partial D\Psi}.$	$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha} + \delta \Psi \wedge \frac{\partial L}{\partial \Psi} + \delta(D\Psi) \wedge \frac{\partial L}{\partial D\Psi}.$	$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha} + \delta \Psi \wedge \frac{\partial L}{\partial \Psi} + \delta(D\Psi) \wedge \frac{\partial L}{\partial D\Psi}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0300 (4.5.11)	102	■1.000 ● N +0	$\delta L = -\varepsilon_\alpha{}^\beta \left[-\sigma^{\alpha\gamma} g_{\gamma\beta} + \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\beta} + D \frac{\partial L}{\partial \Gamma_\alpha{}^\beta} + I^\alpha{}_\beta \Psi \wedge \frac{\delta L}{\delta \Psi} \right].$	$\delta L = -\varepsilon_\alpha{}^\beta \left[-\sigma^{\alpha\gamma} g_{\gamma\beta} + \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\beta} + D \frac{\partial L}{\partial \Gamma_\alpha{}^\beta} + I^\alpha{}_\beta \Psi \wedge \frac{\delta L}{\delta \Psi} \right].$	$\delta L = -\varepsilon_\alpha{}^\beta \left[-\sigma^{\alpha\gamma} g_{\gamma\beta} + \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\beta} + D \frac{\partial L}{\partial \Gamma_\alpha{}^\beta} + I^\alpha{}_\beta \Psi \wedge \frac{\delta L}{\delta \Psi} \right].$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0301 (4.5.12)	102	■1.000 ● N +1	$D \tau_{\alpha\beta} + \vartheta_{[\alpha} \wedge \Sigma_{\beta]} = -I_{\alpha\beta} \Psi \wedge \frac{\delta L}{\delta \Psi} \simeq 0.$	$D\tau_{\alpha\beta} + \vartheta_{[\alpha} \wedge \Sigma_{\beta]} = -I_{\alpha\beta} \Psi \wedge \frac{\delta L}{\delta \Psi} \simeq 0.$	$D\tau_{\alpha\beta} + \vartheta_{[\alpha} \wedge \Sigma_{\beta]} = -I_{\alpha\beta} \Psi \wedge \frac{\delta L}{\delta \Psi} \simeq 0.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0303 (4.5.13)	102	■1.000 ● W -1	$\begin{aligned} & R_{\alpha}{}^{\beta} = d\Gamma_{\alpha}{}^{\beta} \\ & - \Gamma_{\alpha}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\beta} \end{aligned}$	$R_{\alpha}{}^{\beta} = d\Gamma_{\alpha}{}^{\beta} - \Gamma_{\alpha}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\beta}$	$R_{\alpha}{}^{\beta} = d\Gamma_{\alpha}{}^{\beta} - \Gamma_{\alpha}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0307 (4.5.17)	103	■1.000 ● N +0	$\hat{M} := M^3 / \left[M^2 + \frac{\Lambda}{3} J^2 \right], \quad \hat{J} := JM^2 / \left[M^2 + \frac{\Lambda}{3} J^2 \right],$	$\hat{M} := M^3 / \left[M^2 + \frac{\Lambda}{3} J^2 \right], \quad \hat{J} := JM^2 / \left[M^2 + \frac{\Lambda}{3} J^2 \right],$	$\hat{M} := M^3 / [M^2 + \frac{\Lambda}{3} J^2], \quad \hat{J} := JM^2 / [M^2 + \frac{\Lambda}{3} J^2],$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0317 (5.2.1)	110	■ 1.000 ● W +0	$L_{\mathrm{D}}=L(\gamma,\psi,D\psi)=\frac{i}{2}\left\{\bar{\psi}^*\gamma\wedge D\psi+\overline{D\psi}\wedge^*\gamma\psi\right\}-m\bar{\psi}\psi\eta,$	$L_{\mathrm{D}}=L(\gamma,\psi,D\psi)=\frac{i}{2}\left\{\bar{\psi}^*\gamma\wedge D\psi+\overline{D\psi}\wedge^*\gamma\psi\right\}-m\bar{\psi}\psi\eta,$	$L_{\mathrm{D}}=L(\gamma,\psi,D\psi)=\frac{i}{2}\left\{\bar{\psi}^*\gamma\wedge D\psi+\overline{D\psi}\wedge^*\gamma\psi\right\}-m\bar{\psi}\psi\eta,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0322 (5.2.6)	111	■ 1.000 ● K +0	$\left.d\,U=T^{\alpha}\wedge\mu_{\alpha}-\vartheta^{\alpha}\wedge D\mu_{\alpha}=\vartheta^{\alpha}\wedge[e_{\beta}]\left(T^{\beta}\wedge\mu_{\alpha}\right)-D\mu_{\alpha}\right]$	$dU=T^{\alpha}\wedge\mu_{\alpha}-\vartheta^{\alpha}\wedge D\mu_{\alpha}=\vartheta^{\alpha}\wedge[e_{\beta}]\left(T^{\beta}\wedge\mu_{\alpha}\right)-D\mu_{\alpha}]$	$dU=T^{\alpha}\wedge\mu_{\alpha}-\vartheta^{\alpha}\wedge D\mu_{\alpha}=\vartheta^{\alpha}\wedge[e_{\beta}]\left(T^{\beta}\wedge\mu_{\alpha}\right)-D\mu_{\alpha}]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0324 (5.3.1)	111	■ 1.000 ● K +0	$L_{\mathrm{EC}}:=-\frac{1}{2\,\kappa}\,R^{\alpha\,\beta}\wedge\eta_{\alpha\,\beta},$	$L_{\mathrm{EC}}:=-\frac{1}{2\kappa}R^{\alpha\beta}\wedge\eta_{\alpha\beta},$	$L_{\mathrm{EC}}:=-\frac{1}{2\kappa}R^{\alpha\beta}\wedge\eta_{\alpha\beta},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0325 (5.3.2)	112	■ 1.000 ● K +0	$G_{\alpha}:=\frac{1}{2}\,R^{\beta\,\gamma}\wedge\eta_{\alpha\beta\gamma}=\kappa\,\Sigma_{\alpha},$	$G_{\alpha}:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma}=\kappa\Sigma_{\alpha},$	$G_{\alpha}:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma}=\kappa\Sigma_{\alpha},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0327 (5.3.4)	112	■ 1.000 ● N +0	$D^{\{\backslash\backslash\}}G_{\alpha}^{\{\backslash\backslash\}}\equiv 0$	$D^{\{\}}G_{\alpha}^{\{\}}\equiv 0$	$D^{\{\}}G_{\alpha}^{\{\}}\equiv 0$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0331 (5.3.8)	112	■ 1.000 ● N +0	$D[\gamma, \Theta] \equiv 2i \left[\sigma, \Omega^g \right], \quad D[\gamma, \Omega^g] \equiv [\Theta, \Omega^g],$	$D[\gamma, \Theta] \equiv 2i [\sigma, \Omega^g], \quad D[\gamma, \Omega^g] \equiv [\Theta, \Omega^g],$	$D[\gamma, \Theta] \equiv 2i [\sigma, \Omega^g], \quad D[\gamma, \Omega^g] \equiv [\Theta, \Omega^g],$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0332 (5.3.9)	113	■ 1.000 ● N +0	$DG \equiv i \left[\Theta, \gamma_5 \Omega^g \right]$	$DG \equiv i [\Theta, \gamma_5 \Omega^g]$	$DG \equiv i [\Theta, \gamma_5 \Omega^g]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0333 (5.3.10)	113	■ 1.000 ● K +0	$\eta_{\alpha\beta\gamma} \wedge T^\gamma = 2\kappa \tau_{\alpha\beta}$	$\eta_{\alpha\beta\gamma} \wedge T^\gamma = 2\kappa \tau_{\alpha\beta}$	$\eta_{\alpha\beta\gamma} \wedge T^\gamma = 2\kappa \tau_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0334 (5.3.11)	113	■ 1.000 ● N +0	$\mathrm{T}_{\mu \nu} := \frac{2}{\sqrt{g}} \frac{\delta L}{\delta g_{\mu \nu}}$	$T_{\mu\nu} := \frac{2}{\sqrt{g}} \frac{\delta L}{\delta g_{\mu\nu}}$	$T_{\mu\nu} := \frac{2}{\sqrt{g}} \frac{\delta L}{\delta g_{\mu\nu}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0338 (5.3.15)	114	■ 1.000 ● N +0	$\{ \}^{(3)} T_{\alpha} = \frac{(-1)^s}{3} * (\vartheta_{\alpha} \wedge \mathscr{A})$	$^{(3)}T_{\alpha} = \frac{(-1)^s}{3} * (\vartheta_{\alpha} \wedge \mathscr{A})$	$^{(3)}T_{\alpha} = \frac{(-1)^s}{3} * (\vartheta_{\alpha} \wedge \mathscr{A})$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0342 (5.3.19)	115	■ 1.000 ● N -1	$\Delta \widetilde{\mathrm{L}} := \kappa \left(L_{\mathrm{EC}} - L_{\mathrm{HE}} \right) \simeq \frac{\kappa^2}{48} j_5 \wedge \mathrm{^*} j_5 = 4 f^2 j_5 \wedge \mathrm{^*} j_5,$		$\Delta \widetilde{L} := \kappa (L_{\mathrm{EC}} - L_{\mathrm{HE}}) \simeq \frac{\kappa^2}{48} j_5 \wedge \mathrm{^*} j_5 = 4 f^2 j_5 \wedge \mathrm{^*} j_5,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0344 (5.4.2)	116	■ 1.000 ● N -1	$L_{\mathrm{HE}} := L_{\mathrm{HE}} + \frac{\Lambda}{\kappa} \eta$		$L_{\mathrm{HE}} := L_{\mathrm{HE}} + \frac{\Lambda}{\kappa} \eta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0345 (5.4.3)	116	■ 1.000 ● N +0	$g_{\mathrm{N}} := \kappa k^2, \quad \lambda := \Lambda / k^2,$	$g_{\mathrm{N}} := \kappa k^2, \quad \lambda := \Lambda / k^2,$	$g_{\mathrm{N}} := \kappa k^2, \quad \lambda := \Lambda / k^2,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0346 (5.4.4)	116	■ 1.000 ● N +0	$k \frac{\partial}{\partial k} g_{\mathrm{N}} = \beta_1(g_{\mathrm{N}}, \lambda) = (2 + d_{\mathrm{N}}) g_{\mathrm{N}}, \quad k \frac{\partial}{\partial k} \lambda = \beta_2(g_{\mathrm{N}}, \lambda),$		$k \frac{\partial}{\partial k} g_{\mathrm{N}} = \beta_1(g_{\mathrm{N}}, \lambda) = (2 + d_{\mathrm{N}}) g_{\mathrm{N}}, \quad k \frac{\partial}{\partial k} \lambda = \beta_2(g_{\mathrm{N}}, \lambda),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0348 (5.4.6)	117	■ 1.000 ● N +0	$f^2 = \frac{1}{192} \kappa^2 = 3 \times 2^{-10} \left(\frac{\mu}{\Lambda} \right)^2 = 2^{-8} \frac{\mu}{\rho_{\Lambda}}$		$f^2 = \frac{1}{192} \kappa^2 = 3 \times 2^{-10} \left(\frac{\mu}{\Lambda} \right)^2 = 2^{-8} \frac{\mu}{\rho_{\Lambda}}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0350 (6.2.1)	123	■ 1.000 ● N +0	$C_{\mathrm{TT}}:=\frac{1}{2}\ell^2\left(\vartheta^{\alpha}\wedge T_{\alpha}\right)= -\frac{(-1)^{\mathrm{sig}}}{2\ell^2}\ast\mathscr{A},$		$C_{\mathrm{TT}}:=\frac{1}{2\ell^2}\left(\vartheta^{\alpha}\wedge T_{\alpha}\right)= -\frac{(-1)^{\mathrm{sig}}}{2\ell^2}\ast\mathscr{A},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0351 (6.2.2)	124	■ 1.000 ● W -1	$\begin{aligned} C_{\mathrm{RR}}\ \& \; := -\operatorname{Tr}\left(\Gamma\wedge\Omega-\frac{1}{3}\Gamma\wedge\Gamma\wedge\Gamma\right) \\ & = -\frac{1}{2}\left(\Gamma_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}+\frac{1}{3}\Gamma_{\alpha}{}^{\beta}\wedge\Gamma_{\beta}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\alpha}\right). \end{aligned}$	$C_{\mathrm{RR}}:= -\operatorname{Tr}\left(\Gamma\wedge\Omega-\frac{1}{3}\Gamma\wedge\Gamma\wedge\Gamma\right) \\ = -\frac{1}{2}\left(\Gamma_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}+\frac{1}{3}\Gamma_{\alpha}{}^{\beta}\wedge\Gamma_{\beta}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\alpha}\right).$	$C_{\mathrm{RR}}:= -\operatorname{Tr}\left(\Gamma\wedge\Omega-\frac{1}{3}\Gamma\wedge\Gamma\wedge\Gamma\right) \\ = -\frac{1}{2}\left(\Gamma_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}+\frac{1}{3}\Gamma_{\alpha}{}^{\beta}\wedge\Gamma_{\beta}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\alpha}\right).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0352 (6.2.3)	124	■ 1.000 ● N -1	$\hat{C}=C_{\mathrm{RR}}-2\,C_{\mathrm{TT}}$	$\hat{C}=C_{\mathrm{RR}}-2C_{\mathrm{TT}}$	$\hat{C}=C_{\mathrm{RR}}-2C_{\mathrm{TT}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0353 (6.2.4)	124	■ 1.000 ● N -1	$d\,C_{\mathrm{TT}}\equiv\frac{1}{2}\ell^2\left(T^{\alpha}\wedge T_{\alpha}+R_{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}\right),$	$dC_{\mathrm{TT}}\equiv\frac{1}{2\ell^2}\left(T^{\alpha}\wedge T_{\alpha}+R_{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}\right),$	$dC_{\mathrm{TT}}\equiv\frac{1}{2\ell^2}\left(T^{\alpha}\wedge T_{\alpha}+R_{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}\right),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0354 (6.2.5)	124	■ 1.000 ● N -1	$d\,C_{\mathrm{RR}}=-\operatorname{Tr}(\Omega\wedge\Omega)=\frac{1}{2}R^{\alpha\beta}\wedge R_{\alpha\beta}$	$dC_{\mathrm{RR}}=-\operatorname{Tr}(\Omega\wedge\Omega)=\frac{1}{2}R^{\alpha\beta}\wedge R_{\alpha\beta}$	$dC_{\mathrm{RR}}=-\operatorname{Tr}(\Omega\wedge\Omega)=\frac{1}{2}R^{\alpha\beta}\wedge R_{\alpha\beta}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0355 (6.2.6)	124	■ 1.000 ● N +0	$D\,R_{\{\alpha\,\beta\}}\equiv 0,$	$DR_{\alpha\beta}\equiv 0,$	$DR_{\alpha\beta}\equiv 0,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0359 (6.2.10)	125	■ 1.000 ● N +0	$H_{\{\alpha\}^{\{\}}}:=-\partial L_{\{\}}/\partial T^{\{\}}/\partial T^{\alpha}=(1/\kappa)\,\eta_{\alpha\beta\gamma}\wedge K^{\beta\gamma}_{\{\beta\,\gamma\}}$	$H^{\parallel}_{\alpha}:= -\partial L_{\parallel}/\partial T^{\alpha}=(1/\kappa)\eta_{\alpha\beta\gamma}\wedge K^{\beta\gamma}$	$H^{\parallel}_{\alpha}:= -\partial L_{\parallel}/\partial T^{\alpha}=(1/\kappa)\eta_{\alpha\beta\gamma}\wedge K^{\beta\gamma}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0361 (6.2.12)	126	■ 1.000 ● N +0	$\tilde{L}_{\{\}}=L_{\{\}}-R^{\{\alpha\,\beta\}}\wedge\lambda_{\alpha\beta}\,.$	$\tilde{L}_{\parallel}=L_{\parallel}-R^{\alpha\beta}\wedge\lambda_{\alpha\beta}.$	$\tilde{L}_{\parallel}=L_{\parallel}-R^{\alpha\beta}\wedge\lambda_{\alpha\beta}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0362 (6.2.14)	126	■ 1.000 ● W +0	$\begin{gathered} D\,H_{\{\alpha\}^{\{\}}}-E_{\{\alpha\}^{\{\}}}=\\ \Sigma_{\{\alpha\}},\quad \backslash\,D\,\lambda_{\{\alpha\,\beta\}}+\vartheta_{[\alpha}\wedge H_{\beta]}^{\{\}}=\tau_{\alpha\beta},\\ \end{gathered}$	$DH^{\parallel}_{\alpha}-E^{\parallel}_{\alpha}=\Sigma_{\alpha},$ $D\lambda_{\alpha\beta}+\vartheta_{[\alpha}\wedge H^{\parallel}_{\beta]}=\tau_{\alpha\beta},$	$DH^{\parallel}_{\alpha}-E^{\parallel}_{\alpha}=\Sigma_{\alpha},$ $D\lambda_{\alpha\beta}+\vartheta_{[\alpha}\wedge H^{\parallel}_{\beta]}=\tau_{\alpha\beta},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0363 (6.2.15)	126	■ 1.000 ● N +1	$R^{\{\alpha\,\beta\}}=0\,.$	$R^{\alpha\beta}=0.$	$R^{\alpha\beta}=0.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0365 (6.2.17)	126	■ 1.000 ● N +0	$D\left(\tau_{\alpha\beta}-\vartheta_{[\alpha}\wedge H_{\beta]}^{\parallel}\right)=0$	$D\left(\tau_{\alpha\beta}-\vartheta_{[\alpha}\wedge H_{\beta]}^{\parallel}\right)=0$	$D(\tau_{\alpha\beta}-\vartheta_{[\alpha}\wedge H_{\beta]}^{\parallel})=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0369 (6.2.20)	127	■ 1.000 ● N +1	$\stackrel{\scriptscriptstyle{C}}{\mathrm{TT}}:=\frac{1}{2}\ell^2\left(\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{T}}{\alpha}\right),$	$\stackrel{\scriptscriptstyle{C}}{\mathrm{TT}}:=\frac{1}{2\ell^2}\left(\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{T}}{\alpha}\right),$	$\stackrel{\scriptscriptstyle{C}}{\mathrm{TT}}:=\frac{1}{2\ell^2}\left(\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{T}}{\alpha}\right),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0373 (6.2.24)	127	■ 1.000 ● N +0	$\stackrel{\scriptscriptstyle{D}}{\Pi}_{\alpha}=\Sigma_{\alpha}.$	$\stackrel{\scriptscriptstyle{D}}{\Pi}_{\alpha}=\Sigma_{\alpha}.$	$\stackrel{\scriptscriptstyle{D}}{\Pi}_{\alpha}=\Sigma_{\alpha}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0379 (6.2.30)	128	■ 1.000 ● K +0	$\operatorname{Tr}(\stackrel{\scriptscriptstyle{F}}{\wedge}\stackrel{\scriptscriptstyle{F}}{\wedge})$	$\operatorname{Tr}(\stackrel{\scriptscriptstyle{F}}{\wedge}\stackrel{\scriptscriptstyle{F}}{\wedge})$	$\operatorname{Tr}(\stackrel{\scriptscriptstyle{F}}{\wedge}\stackrel{\scriptscriptstyle{F}}{\wedge})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0383 (6.3.1)	129	■ 1.000 ● N +0	$\check{E}_{\alpha}:=E_{\alpha}+\Gamma_{\alpha}^{\beta}\wedge H_{\beta}+\Sigma_{\alpha}\cong dH_{\alpha},$	$\check{E}_{\alpha}:=E_{\alpha}+\Gamma_{\alpha}^{\beta}\wedge H_{\beta}+\Sigma_{\alpha}\cong dH_{\alpha},$	$\check{E}_{\alpha}:=E_{\alpha}+\Gamma_{\alpha}^{\beta}\wedge H_{\beta}+\Sigma_{\alpha}\cong dH_{\alpha},$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0386 (6.4.1)	130	■ 1.000 ● N +0	$n:=\partial_t-N^A\partial/\partial x^A,\quad A=1,2,3,$	$n:=\partial_t-N^A\partial/\partial x^A,\quad A=1,2,3,$	$n:=\partial_t-N^A\partial/\partial x^A,\quad A=1,2,3,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0395 (6.4.10)	132	■ 1.000 ● W +0	$\Sigma_{\alpha}:=\frac{\partial L}{\partial \vartheta^{\alpha}},\quad \tau_{\alpha\beta}:=\frac{\partial L}{\partial \Gamma^{\alpha\beta}}=\rho(I_{\alpha\beta})\Psi\wedge\frac{\partial L}{\partial D\Psi}$	$\Sigma_{\alpha}:=\frac{\partial L}{\partial \vartheta^{\alpha}},\quad \tau_{\alpha\beta}:=\frac{\partial L}{\partial \Gamma^{\alpha\beta}}=\rho(I_{\alpha\beta})\Psi\wedge\frac{\partial L}{\partial D\Psi}$	$\Sigma_{\alpha}:=\frac{\partial L}{\partial \vartheta^{\alpha}},\quad \tau_{\alpha\beta}:=\frac{\partial L}{\partial \Gamma^{\alpha\beta}}=\rho(I_{\alpha\beta})\Psi\wedge\frac{\partial L}{\partial D\Psi}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0408 (6.4.28)	136	■ 1.000 ● N +0	$d\,s^2=h^2\,d\,r^2+f^2\left[d\,\psi^2+\sin^2\psi\left(d\theta^2+\sin^2\theta\,d\phi^2\right)\right],$	$ds^2=h^2dr^2+f^2\left[d\psi^2+\sin^2\psi\left(d\theta^2+\sin^2\theta d\phi^2\right)\right],$	$ds^2=h^2dr^2+f^2\left[d\psi^2+\sin^2\psi\left(d\theta^2+\sin^2\theta d\phi^2\right)\right],$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0410 (6.4.30)	136	■ 1.000 ● N +1	$\ell^2\underline{C}_{\mathrm{TT}}=\frac{1}{2}\underline{\vartheta}^{\alpha}\wedge d\underline{\vartheta}_{\alpha}=\frac{1}{2}\underline{\vartheta}^A\wedge d\underline{\vartheta}_A=3\underline{\vartheta}^{\hat{1}}\wedge\underline{\vartheta}^{\hat{2}}\wedge\underline{\vartheta}^{\hat{3}}$	$\ell^2\underline{C}_{\mathrm{TT}}=\frac{1}{2}\underline{\vartheta}^{\alpha}\wedge d\underline{\vartheta}_{\alpha}=\frac{1}{2}\underline{\vartheta}^A\wedge d\underline{\vartheta}_A=3\underline{\vartheta}^{\hat{1}}\wedge\underline{\vartheta}^{\hat{2}}\wedge\underline{\vartheta}^{\hat{3}}$	$\ell^2\underline{C}_{\mathrm{TT}}=\frac{1}{2}\underline{\vartheta}^{\alpha}\wedge d\underline{\vartheta}_{\alpha}=\frac{1}{2}\underline{\vartheta}^A\wedge d\underline{\vartheta}_A=3\underline{\vartheta}^{\hat{1}}\wedge\underline{\vartheta}^{\hat{2}}\wedge\underline{\vartheta}^{\hat{3}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0412 (6.4.32)	136	■ 1.000 ● N +1	$f=\frac{a\,r^2}{r^2+c^2}$	$f=\frac{ar^2}{r^2+c^2}$	$f=\frac{ar^2}{r^2+c^2}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0414 (6.5.2)	137	■ 1.000 ● N +0	$\Psi(\gamma)=\int_{\mathcal{C}} \frac{1}{W(A,\gamma)} \Psi(A) \, dA$	$\Psi(\gamma) = \int_{\mathcal{C}/\mathcal{G}} \mathcal{D}A W(A,\gamma) \Psi(A).$	$\Psi(\gamma) = \int_{\mathcal{C}/\mathcal{G}} \mathcal{D}A W(A,\gamma) \Psi(A).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0416 (6.5.4)	137	■ 1.000 ● K +0	$\langle 0 W(\gamma) 0\rangle e^{\Lambda G(\gamma)} V(\gamma),$	$\langle 0 W(\gamma) 0\rangle e^{\Lambda G(\gamma)} V(\gamma),$	$\langle 0 W(\gamma) 0\rangle e^{\Lambda G(\gamma)} V(\gamma),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0417 (6.6.1)	138	■ 1.000 ● N +1	$\int T = \int d\sigma = 2\pi N$	$\int T = \int d\sigma = 2\pi N$	$\int T = \int d\sigma = 2\pi N,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0418 (6.6.2)	139	■ 1.000 ● K +0	$W^N:=S^3\#^N(S^1\times S^2)$	$W^N:=S^3\#^N(S^1\times S^2).$	$W^N:=S^3\#^N(S^1\times S^2).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0419 (6.7.1)	140	■ 1.000 ● K +0	$L:=L_{\mathrm{EC}}+L_{\mathrm{\theta}}+L_{\mathrm{D}},$	$L:=L_{\mathrm{EC}}+L_{\theta}+L_{\mathrm{D}},$	$L:=L_{\mathrm{EC}}+L_{\theta}+L_{\mathrm{D}},$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0420 (6.7.2)	140	■ 1.000 ● N +1	$L_{\theta}=\theta_{\mathrm{T}}dC_{\mathrm{TT}}+\theta_{\mathrm{L}}dC_{\mathrm{RR}}$	$L_{\theta}=\theta_{\mathrm{T}}dC_{\mathrm{TT}}+\theta_{\mathrm{L}}dC_{\mathrm{RR}}$	$L_{\theta}=\theta_{\mathrm{T}}dC_{\mathrm{TT}}+\theta_{\mathrm{L}}dC_{\mathrm{RR}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0421 (6.7.3)	140	■ 1.000 ● K +0	$H_{\alpha}:= -\frac{\partial L}{\partial T^{\alpha}}= -\frac{\theta_{\mathrm{T}}}{\ell^2}T_{\alpha},$	$H_{\alpha}:= -\frac{\partial L}{\partial T^{\alpha}}= -\frac{\theta_{\mathrm{T}}}{\ell^2}T_{\alpha},$	$H_{\alpha}:= -\frac{\partial L}{\partial T^{\alpha}}= -\frac{\theta_{\mathrm{T}}}{\ell^2}T_{\alpha},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0422 (6.7.4)	140	■ 1.000 ● W +0	$H_{\alpha\beta}:= -\frac{\partial L}{\partial R^{\alpha\beta}}= \frac{1}{2\kappa}\eta_{\alpha\beta}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\theta_{\mathrm{L}}R_{\alpha\beta}$	$H_{\alpha\beta}:= -\frac{\partial L}{\partial R^{\alpha\beta}}= \frac{1}{2\kappa}\eta_{\alpha\beta}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\theta_{\mathrm{L}}R_{\alpha\beta}$	$H_{\alpha\beta}:= -\frac{\partial L}{\partial R^{\alpha\beta}}= \frac{1}{2\kappa}\eta_{\alpha\beta}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\theta_{\mathrm{L}}R_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0423 (6.7.5)	140	■ 1.000 ● N +0	$D\theta_{\mathrm{T}}\wedge T_{\alpha}+\theta_{\mathrm{T}}DT_{\alpha}+\ell^2E_{\alpha}=-\ell^2\Sigma_{\alpha}.$	$D\theta_{\mathrm{T}}\wedge T_{\alpha}+\theta_{\mathrm{T}}DT_{\alpha}+\ell^2E_{\alpha}=-\ell^2\Sigma_{\alpha}.$	$D\theta_{\mathrm{T}}\wedge T_{\alpha}+\theta_{\mathrm{T}}DT_{\alpha}+\ell^2E_{\alpha}=-\ell^2\Sigma_{\alpha}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0425 (6.7.7)	140	■ 1.000 ● N +0	$C_{\mathrm{TT}}\cong\frac{\kappa}{4\ell^2}j_5$	$C_{\mathrm{TT}}\cong\frac{\kappa}{4\ell^2}j_5$	$C_{\mathrm{TT}}\cong\frac{\kappa}{4\ell^2}j_5$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0427 (6.7.9)	141	■ 1.000 ● N +0	$T^{\gamma} \wedge \eta_{\gamma\beta} + \kappa D\theta_{\mathrm{L}} \wedge DT_{\beta} = 0$.	$T^{\gamma} \wedge \eta_{\gamma\beta} + \kappa D\theta_{\mathrm{L}} \wedge DT_{\beta} = 0$.	$T^{\gamma} \wedge \eta_{\gamma\beta} + \kappa D\theta_{\mathrm{L}} \wedge DT_{\beta} = 0$.
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0429 (6.7.11)	141	■ 1.000 ● N +0	$\kappa d\theta_{\mathrm{L}} \wedge T_{\beta} = -\eta_{\beta}$	$\kappa d\theta_{\mathrm{L}} \wedge T_{\beta} = -\eta_{\beta}$	$\kappa d\theta_{\mathrm{L}} \wedge T_{\beta} = -\eta_{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0430 (6.7.12)	141	■ 1.000 ● N -1	$\kappa \ell^2 d\theta_{\mathrm{L}} \wedge C_{\mathrm{TT}} = 2\eta$.	$\kappa \ell^2 d\theta_{\mathrm{L}} \wedge C_{\mathrm{TT}} = 2\eta$.	$\kappa \ell^2 d\theta_{\mathrm{L}} \wedge C_{\mathrm{TT}} = 2\eta$.
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0431 (6.7.13)	141	■ 1.000 ● N +0	$D T^{\gamma} \wedge \eta_{\alpha\beta\gamma} + \kappa D\theta_{\mathrm{T}} \wedge T_{[\alpha} \wedge \vartheta_{\beta]} / \ell^2 = 2\kappa (T_{[\alpha} \wedge \mu_{\beta]} - \vartheta_{[\alpha} \wedge D\mu_{\beta]})$.	$DT^{\gamma} \wedge \eta_{\alpha\beta\gamma} + \kappa D\theta_{\mathrm{T}} \wedge T_{[\alpha} \wedge \vartheta_{\beta]} / \ell^2 = 2\kappa (T_{[\alpha} \wedge \mu_{\beta]} - \vartheta_{[\alpha} \wedge D\mu_{\beta]})$.	$DT^{\gamma} \wedge \eta_{\alpha\beta\gamma} + \kappa D\theta_{\mathrm{T}} \wedge T_{[\alpha} \wedge \vartheta_{\beta]} / \ell^2 = 2\kappa (T_{[\alpha} \wedge \mu_{\beta]} - \vartheta_{[\alpha} \wedge D\mu_{\beta]})$.
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0432 (6.8.1)	142	■ 1.000 ● N +0	$W = \int \left[L(\vartheta^{\alpha}, T^{\alpha}, R^{\alpha\beta}, \Psi, D\Psi) + L_g(\vartheta^{\alpha}, T^{\alpha}, R^{\alpha\beta}) \right]$.	$W = \int \left[L(\vartheta^{\alpha}, T^{\alpha}, R^{\alpha\beta}, \Psi, D\Psi) + L_g(\vartheta^{\alpha}, T^{\alpha}, R^{\alpha\beta}) \right]$.	$W = \int \left[L(\vartheta^{\alpha}, T^{\alpha}, R^{\alpha\beta}, \Psi, D\Psi) + L_g(\vartheta^{\alpha}, T^{\alpha}, R^{\alpha\beta}) \right]$.
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0433 (6.8.3)	142	■ 1.000 ● C -3	$\begin{gathered} \frac{\delta L}{\delta \Psi} = \frac{\partial L}{\partial \Psi} - (-1)^p D \frac{\partial L}{\partial D \Psi} = 0, \quad (\text{MATTER}) \\ DH_{\alpha} - E_{\alpha} = \Sigma_{\alpha}, \quad (\text{FIRST}) \\ DH_{\alpha\beta} - E_{\alpha\beta} = \tau_{\alpha\beta}. \quad (\text{SECOND}) \end{gathered}$	$\frac{\delta L}{\delta \Psi} = \frac{\partial L}{\partial \Psi} - (-1)^p D \frac{\partial L}{\partial D \Psi} = 0, \quad (\text{MATTER})$ $DH_{\alpha} - E_{\alpha} = \Sigma_{\alpha}, \quad (\text{FIRST})$ $DH_{\alpha\beta} - E_{\alpha\beta} = \tau_{\alpha\beta}. \quad (\text{SECOND})$	$\frac{\delta L}{\delta \Psi} = \frac{\partial L}{\partial \Psi} - (-1)^p D \frac{\partial L}{\partial D \Psi} = 0, \quad (\text{MATTER})$ $DH_{\alpha} - E_{\alpha} = \Sigma_{\alpha}, \quad (\text{FIRST})$ $DH_{\alpha\beta} - E_{\alpha\beta} = \tau_{\alpha\beta}. \quad (\text{SECOND})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0435 (6.8.6)	142	■ 1.000 ● N +1	$\Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = \frac{\partial L}{\partial \vartheta^{\alpha}} + D \frac{\partial L}{\partial T^{\alpha}}$	$\Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = \frac{\partial L}{\partial \vartheta^{\alpha}} + D \frac{\partial L}{\partial T^{\alpha}}$	$\Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = \frac{\partial L}{\partial \vartheta^{\alpha}} + D \frac{\partial L}{\partial T^{\alpha}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0436 (6.8.7)	142	■ 1.000 ● W +0	$\tau_{\alpha\beta} := \frac{\delta L}{\delta \Gamma^{\alpha\beta}} = \rho(I_{\alpha\beta}) \Psi \wedge \frac{\partial L}{\partial (D \Psi)} + \vartheta_{[\alpha} \wedge \frac{\partial L}{\partial T^{\beta]} + D \frac{\partial L}{\partial R^{\alpha\beta}},$	$\tau_{\alpha\beta} := \frac{\delta L}{\delta \Gamma^{\alpha\beta}} = \rho(I_{\alpha\beta}) \Psi \wedge \frac{\partial L}{\partial (D \Psi)} + \vartheta_{[\alpha} \wedge \frac{\partial L}{\partial T^{\beta]} + D \frac{\partial L}{\partial R^{\alpha\beta}},$	$\tau_{\alpha\beta} := \frac{\delta L}{\delta \Gamma^{\alpha\beta}} = \rho(I_{\alpha\beta}) \Psi \wedge \frac{\partial L}{\partial (D \Psi)} + \vartheta_{[\alpha} \wedge \frac{\partial L}{\partial T^{\beta]} + D \frac{\partial L}{\partial R^{\alpha\beta}},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0438 (6.8.9)	142	■ 1.000 ● N +0	$E_{\alpha\beta} := -\vartheta_{[\alpha} \wedge H_{\beta]}$	$E_{\alpha\beta} := -\vartheta_{[\alpha} \wedge H_{\beta]}$	$E_{\alpha\beta} := -\vartheta_{[\alpha} \wedge H_{\beta]}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0439 (6.8.10)	143	■ 1.000 ● N +0	$\vartheta^{\alpha} \wedge E_{\alpha} = 4L_g + 2T^{\alpha} \wedge H_{\alpha} + 2R^{\alpha\beta} \wedge H_{\alpha\beta}.$	$\vartheta^{\alpha} \wedge E_{\alpha} = 4L_g + 2T^{\alpha} \wedge H_{\alpha} + 2R^{\alpha\beta} \wedge H_{\alpha\beta}.$	$\vartheta^{\alpha} \wedge E_{\alpha} = 4L_g + 2T^{\alpha} \wedge H_{\alpha} + 2R^{\alpha\beta} \wedge H_{\alpha\beta}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0442 (6.8.13)	143	■ 1.000 ● N +0	$L_{\mathrm{qPG}}=-\frac{1}{2}T^{\alpha}\wedge H_{\alpha}-\frac{1}{2}R^{\alpha\beta}\wedge\left(H_{\alpha\beta}-\frac{a_0}{2\ell^2}\eta_{\alpha\beta}\right),$		$L_{\mathrm{qPG}}=-\frac{1}{2}T^{\alpha}\wedge H_{\alpha}-\frac{1}{2}R^{\alpha\beta}\wedge\left(H_{\alpha\beta}-\frac{a_0}{2\ell^2}\eta_{\alpha\beta}\right),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0444 (6.8.15)	143	■ 1.000 ● N +0	$H_{\alpha\beta}=-\frac{a_0}{2\ell^2}\eta_{\alpha\beta}-\frac{1}{\kappa}\ast\left(\sum_{j=1}^6b_j^{(j)}R_{\alpha\beta}\right).$	$H_{\alpha\beta}=-\frac{a_0}{2\ell^2}\eta_{\alpha\beta}-\frac{1}{\kappa}\ast\left(\sum_{j=1}^6b_j^{(j)}R_{\alpha\beta}\right).$	$H_{\alpha\beta}=-\frac{a_0}{2\ell^2}\eta_{\alpha\beta}-\frac{1}{\kappa}\ast\left(\sum_{j=1}^6b_j^{(j)}R_{\alpha\beta}\right).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0447 (6.9.3)	144	■ 1.000 ● N +0	$\vartheta=\{ \}^{\ast}{}_1=\frac{1}{4!}\vartheta_{\alpha\beta\gamma\delta}\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta},$	$\eta=\ast 1=\frac{1}{4!}\eta_{\alpha\beta\gamma\delta}\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta},$	$\eta=\ast 1=\frac{1}{4!}\eta_{\alpha\beta\gamma\delta}\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0448 (6.9.4)	145	■ 1.000 ● N +0	$p=\frac{1}{8\pi^2}\int R_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}.$	$p=\frac{1}{8\pi^2}\int R_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}.$	$p=\frac{1}{8\pi^2}\int R_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ _Z-Library__ EQ0453 (7.2.1)	151	■ 1.000 ● C +14	$\left[L_{ab}^{\left(\star\right)}:=\frac{1}{2}\varepsilon_{abcd}L^{cd}=L_{ab}.\right]$	(not rendered)	$L_{ab}^{(\star)}:=\frac{1}{2}\varepsilon_{abcd}L^{cd}=L_{ab}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0454 (7.2.6)	152	■ 1.000 ● N +0	$\backslash 0\mathrm{mega}^{\mathrm{g}}=\backslash 0\mathrm{mega}_{\mathrm{C}}+\backslash \mathrm{bar}\{\backslash 0\mathrm{mega}\}^{\mathrm{g}}-\backslash \mathrm{frac}\{1\}\{6\}\ \mathrm{vartheta}\ \mathrm{wedge}\ \mathrm{vartheta}\ R\ .$	$\Omega^g=\Omega_{\mathrm{C}}+\bar{\Omega}^g-\frac{1}{6}\vartheta\wedge\vartheta R.$	$\Omega^g=\Omega_{\mathrm{C}}+\bar{\Omega}^g-\frac{1}{6}\vartheta\wedge\vartheta R.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0455 (7.2.7)	152	■ 1.000 ● N +0	$\backslash \mathrm{bar}\{\backslash 0\mathrm{mega}\}^{\mathrm{g}}=\mathrm{vartheta}\ \mathrm{wedge}^{\mathrm{*}}\backslash \mathrm{left}(\mathrm{vartheta}\ \mathrm{wedge}\ \backslash 0\mathrm{mega}^{\mathrm{g}}\backslash \mathrm{right})-\backslash \mathrm{frac}\{1\}\{4\}\ \mathrm{vartheta}\ \mathrm{wedge}\ \mathrm{vartheta}\ R,$	$\bar{\Omega}^g=\vartheta\wedge^*(\vartheta\wedge\Omega^g)-\frac{1}{4}\vartheta\wedge\vartheta R,$	$\bar{\Omega}^g=\vartheta\wedge^*(\vartheta\wedge\Omega^g)-\frac{1}{4}\vartheta\wedge\vartheta R,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0457 (7.2.9)	152	■ 1.000 ● N +0	$\backslash \mathrm{operatorname}\{\mathrm{Tr}\}\backslash \mathrm{left}(\mathrm{vartheta}\ \mathrm{wedge}\ \mathrm{vartheta}\ \mathrm{wedge}\ \backslash 0\mathrm{mega}_{\mathrm{C}}\backslash \mathrm{right})=\backslash \mathrm{operatorname}\{\mathrm{Tr}\}\backslash \mathrm{left}(\mathrm{vartheta}\ \mathrm{wedge}\ \mathrm{vartheta}\ \mathrm{wedge}\ \backslash \mathrm{bar}\{\backslash 0\mathrm{mega}\}^{\mathrm{g}}\backslash \mathrm{right})=0$	$\mathrm{Tr}\left(\vartheta\wedge\vartheta\wedge\Omega_{\mathrm{C}}\right)=\mathrm{Tr}\left(\vartheta\wedge\vartheta\wedge\bar{\Omega}^g\right)=0$	$\mathrm{Tr}(\vartheta\wedge\vartheta\wedge\Omega_{\mathrm{C}})=\mathrm{Tr}(\vartheta\wedge\vartheta\wedge\bar{\Omega}^g)=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0463 (7.3.2)	153	■ 1.000 ● N +0	$D^{\mathrm{*}}\backslash 0\mathrm{mega}^{\mathrm{g}}=0,$	$D^*\Omega^g=0,$	$D^*\Omega^g=0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0464 (7.3.3)	153	■ 1.000 ● K +0	$D^{\mathrm{*}}\ R_{\alpha\beta}=0,$	$D^*R_{\alpha\beta}=0,$	$D^*R_{\alpha\beta}=0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0468 (7.3.7)	154	■ 1.000 ● N +0	$R_{\alpha\beta}=-\mathrm{*}\ R_{\alpha\beta}^{\mathrm{(\star)}},$	$R_{\alpha\beta}=-\mathrm{*}\ R_{\alpha\beta}^{(*)},$	$R_{\alpha\beta}=-\mathrm{*}\ R_{\alpha\beta}^{(*)},$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0469 (7.3.8)	154	■ 1.000 ● N +0	$R_{\alpha\beta} = {}^*R_{\alpha\beta}^{(\star)},$	$R_{\alpha\beta} = {}^*R_{\alpha\beta}^{(\star)},$	$R_{\alpha\beta} = {}^*R_{\alpha\beta}^{(\star)},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0479 (7.4.7)	157	■ 1.000 ● K +0	$G_{\alpha}-\Lambda\eta_{\alpha}=0$	$G_{\alpha}-\Lambda\eta_{\alpha}=0$	$G_{\alpha}-\Lambda\eta_{\alpha}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0480 (7.5.1)	157	■ 1.000 ● N -1	$\square^c:=\square-\frac{1}{6}R,\quad\square:=(1/\sqrt{g})\,\partial_i\left(\sqrt{g}\,g^{ij}\,\partial_j\right).$	$\square^c:=\square-\frac{1}{6}R,\quad\square:=(1/\sqrt{g})\partial_i\left(\sqrt{g}g^{ij}\partial_j\right).$	$\square^c:=\square-\frac{1}{6}R,\quad\square:=(1/\sqrt{g})\partial_i(\sqrt{g}g^{ij}\partial_j).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0482 (7.5.3)	157	■ 1.000 ● N +0	$\left(\square-\frac{2}{3}\Lambda\right)h_{ij}=0$	$\left(\square-\frac{2}{3}\Lambda\right)h_{ij}=0$	$\left(\square-\frac{2}{3}\Lambda\right)h_{ij}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ0483 (7.5.4)	158	■ 1.000 ● N +0	$\partial^i h_{ij}-\partial_j h=0$	$\partial^i h_{ij}-\partial_j h=0$	$\partial^i h_{ij}-\partial_j h=0$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0484 (7.5.5)	158	■ 1.000 ● K +0	$\Delta m=\sqrt{2\Lambda/3}>0$	$\Delta m = \sqrt{2\Lambda/3} > 0$	$\Delta m = \sqrt{2\Lambda/3} > 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0488 (7.7.5)	160	■ 1.000 ● K +0	$L_{\mathrm{EC}}=-\frac{1}{2\kappa^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}$	$L_{\mathrm{EC}}=-\frac{1}{2\kappa^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}$	$L_{\mathrm{EC}}=-\frac{1}{2\kappa^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0489 (7.7.6)	160	■ 1.000 ● N +0	$\widetilde{L}=-\sum_{k=0}^K(1/2k)R^{\alpha\beta}\wedge\stackrel{(2k)}{H}_{\alpha\beta}+L_{\theta},$	$\widetilde{L}=-\sum_{k=0}^K(1/2k)R^{\alpha\beta}\wedge\stackrel{(2k)}{H}_{\alpha\beta}+L_{\theta},$	$\widetilde{L}=-\sum_{k=0}^K(1/2k)R^{\alpha\beta}\wedge\stackrel{(2k)}{H}_{\alpha\beta}+L_{\theta},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0491 (7.7.9)	161	■ 1.000 ● K +0	$\widetilde{L}\rightarrow L=-\frac{1}{2}\left(R^{\alpha\beta}\wedge\widetilde{H}_{\alpha\beta}-\widetilde{L}\right)$	$\widetilde{L}\rightarrow L=-\frac{1}{2}\left(R^{\alpha\beta}\wedge\widetilde{H}_{\alpha\beta}-\widetilde{L}\right)$	$\widetilde{L}\rightarrow L=-\frac{1}{2}\left(R^{\alpha\beta}\wedge\widetilde{H}_{\alpha\beta}-\widetilde{L}\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0493 (7.7.11)	161	■ 1.000 ● N +0	$\widetilde{H}_{\alpha\beta\mu\nu}:=\frac{\partial^2\widetilde{L}}{\partial R^{\mu\nu}\partial R^{\alpha\beta}}=-\frac{\partial\widetilde{H}_{\mu\nu}}{\partial R^{\alpha\beta}}$	$\widetilde{H}_{\alpha\beta\mu\nu}:=\frac{\partial^2\widetilde{L}}{\partial R^{\mu\nu}\partial R^{\alpha\beta}}=-\frac{\partial\widetilde{H}_{\mu\nu}}{\partial R^{\alpha\beta}}$	$\widetilde{H}_{\alpha\beta\mu\nu}:=\frac{\partial^2\widetilde{L}}{\partial R^{\mu\nu}\partial R^{\alpha\beta}}=-\frac{\partial\widetilde{H}_{\mu\nu}}{\partial R^{\alpha\beta}}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0494 (7.7.12)	161	■ 1.000 ● N -1	$\widetilde{\eta}_{\alpha\beta}:=\frac{\theta_{\mathrm{T}}^*}{2}\eta_{\alpha\beta}-\frac{\theta_{\mathrm{T}}}{2}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\ell^2\widetilde{H}_{\alpha\beta}=0$	$\widetilde{\eta}_{\alpha\beta}:=\frac{\theta_{\mathrm{T}}^*}{2}\eta_{\alpha\beta}-\frac{\theta_{\mathrm{T}}}{2}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\ell^2\widetilde{H}_{\alpha\beta}=0$	$\widetilde{\eta}_{\alpha\beta}:=\frac{\theta_{\mathrm{T}}^*}{2}\eta_{\alpha\beta}-\frac{\theta_{\mathrm{T}}}{2}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\ell^2\widetilde{H}_{\alpha\beta}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0496 (7.7.14)	162	■ 1.000 ● K +0	$\eta:=\eta_{\alpha\beta\gamma\delta}\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta}/4!$	$\eta:=\eta_{\alpha\beta\gamma\delta}\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta}/4!$	$\eta:=\eta_{\alpha\beta\gamma\delta}\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta}/4!$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0499 (7.7.17)	163	■ 1.000 ● K +0	$\gamma_{\alpha}\gamma_{\beta}+\gamma_{\beta}\gamma_{\alpha}=2o_{\alpha\beta}\mathbf{1}_4.$	$\gamma_{\alpha}\gamma_{\beta}+\gamma_{\beta}\gamma_{\alpha}=2o_{\alpha\beta}\mathbf{1}_4.$	$\gamma_{\alpha}\gamma_{\beta}+\gamma_{\beta}\gamma_{\alpha}=2o_{\alpha\beta}\mathbf{1}_4.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0501 (7.7.19)	163	■ 1.000 ● K +0	$\gamma:=\gamma^{\alpha}\vartheta_{\alpha}$	$\gamma:=\gamma^{\alpha}\vartheta_{\alpha}$	$\gamma:=\gamma^{\alpha}\vartheta_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0502 (7.7.20)	163	■ 1.000 ● N +0	$\gamma^{\alpha}\eta_{\alpha}=\frac{i}{6}\gamma_5\gamma\wedge\gamma\wedge\gamma,$	$*\gamma=\gamma^{\alpha}\eta_{\alpha}=\frac{i}{6}\gamma_5\gamma\wedge\gamma\wedge\gamma,$	$*\gamma=\gamma^{\alpha}\eta_{\alpha}=\frac{i}{6}\gamma_5\gamma\wedge\gamma\wedge\gamma,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0503 (7.7.21)	164	■ 1.000 ● N +1	$\backslash\sigma:=\frac{1}{2}\backslash\sigma_{\alpha}\backslash\beta}\backslashvartheta^{\alpha}\backslash\alpha}\backslashwedge\backslashvartheta^{\alpha}\backslash\beta}=\frac{1}{2}\backslash\gamma\wedge\gamma,\quad\backslash\sigma=\frac{1}{2}\sigma_{\alpha\beta}\eta^{\alpha\beta}=: \sigma^{(*)}=i\gamma_5\sigma.$	$\sigma:=\frac{1}{2}\sigma_{\alpha\beta}\vartheta^{\alpha}\wedge\vartheta^{\beta}=\frac{i}{2}\gamma\wedge\gamma,\quad\sigma=\frac{1}{2}\sigma_{\alpha\beta}\eta^{\alpha\beta}=: \sigma^{(*)}=i\gamma_5\sigma.$	$\sigma:=\frac{1}{2}\sigma_{\alpha\beta}\vartheta^{\alpha}\wedge\vartheta^{\beta}=\frac{i}{2}\gamma\wedge\gamma,\quad\sigma=\frac{1}{2}\sigma_{\alpha\beta}\eta^{\alpha\beta}=: \sigma^{(*)}=i\gamma_5\sigma.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0504 (7.7.22)	164	■ 1.000 ● N +0	$\backslash\sigma^{\{(*)\}}=:i\backslash\gamma_5\backslash\sigma,\quad\backslash\sigma^{\{(**)\}}=i^2\gamma_5^2\sigma=-\sigma.$	$\sigma^{(*)}:=i\gamma_5\sigma,\quad\sigma^{(**)}=i^2\gamma_5^2\sigma=-\sigma.$	$\sigma^{(*)}:=i\gamma_5\sigma,\quad\sigma^{(**)}=i^2\gamma_5^2\sigma=-\sigma.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0505 (7.7.23)	164	■ 1.000 ● N +0	$\backslash\left[\backslash\gamma,\backslash\sigma^{\{(*)\}}\right]=\backslash\gamma\wedge i\backslash\gamma_5\backslash\sigma-i\backslash\gamma_5\sigma\wedge\gamma=-i\gamma_5(\gamma\wedge\sigma+\sigma\wedge\gamma)=2\gamma_5\gamma\wedge\gamma\wedge\gamma$	$\left[\gamma,\sigma^{(*)}\right]=\gamma\wedge i\gamma_5\sigma-i\gamma_5\sigma\wedge\gamma=-i\gamma_5(\gamma\wedge\sigma+\sigma\wedge\gamma)=2\gamma_5\gamma\wedge\gamma\wedge\gamma$	$[\gamma,\sigma^{(*)}]=\gamma\wedge i\gamma_5\sigma-i\gamma_5\sigma\wedge\gamma=-i\gamma_5(\gamma\wedge\sigma+\sigma\wedge\gamma)=2\gamma_5\gamma\wedge\gamma\wedge\gamma$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0508 (7.7.26)	165	■ 1.000 ● N +0	$\backslash\gamma\rightarrow\backslash\gamma^{\theta}=e^{i\gamma^5\theta}\backslash\gamma e^{-i\gamma^5\theta}$	$\gamma\rightarrow\gamma^{\theta}=e^{i\gamma^5\theta}\gamma e^{-i\gamma^5\theta}$	$\gamma\rightarrow\gamma^{\theta}=e^{i\gamma^5\theta}\gamma e^{-i\gamma^5\theta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0510 (7.7.28)	165	■ 1.000 ● K +0	$\backslash\gamma:=\backslash\vartheta^{\alpha}\backslash\gamma_{\alpha},\quad\backslash\Gamma:=\frac{i}{4}\Gamma^{\alpha\beta}\sigma_{\alpha\beta}=\Gamma^{\{ \}}-K,$	$\gamma:=\vartheta^{\alpha}\gamma_{\alpha},\quad\Gamma:=\frac{i}{4}\Gamma^{\alpha\beta}\sigma_{\alpha\beta}=\Gamma^{\{ \}}-K,$	$\gamma:=\vartheta^{\alpha}\gamma_{\alpha},\quad\Gamma:=\frac{i}{4}\Gamma^{\alpha\beta}\sigma_{\alpha\beta}=\Gamma^{\{ \}}-K,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0513 (7.7.31)	166	■ 1.000 ● N +0	$\Omega^{\{\star\}}=\frac{1}{4}R_{\{\alpha\}\{\beta\}}^{\{\star\}}\sigma^{\alpha\beta},\quad R_{\{\alpha\}\{\beta\}}^{\{\star\}}:=\frac{1}{2}\eta_{\alpha\beta}\gamma_{\delta}R^{\gamma\delta}.$	$\Omega^{(\star)}=\frac{1}{4}R_{\alpha\beta}^{(\star)}\sigma^{\alpha\beta},\quad R_{\alpha\beta}^{(\star)}:=\frac{1}{2}\eta_{\alpha\beta}\gamma_{\delta}R^{\gamma\delta}.$	$\Omega^{(\star)}=\frac{1}{4}R_{\alpha\beta}^{(\star)}\sigma^{\alpha\beta},\quad R_{\alpha\beta}^{(\star)}:=\frac{1}{2}\eta_{\alpha\beta}\gamma_{\delta}R^{\gamma\delta}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0514 (7.7.32)	166	■ 1.000 ● K +0	$D\sigma=\frac{i}{2}[\Theta,\gamma]$	$D\sigma=\frac{i}{2}[\Theta,\gamma]$	$D\sigma=\frac{i}{2}[\Theta,\gamma]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0517 (7.7.35)	166	■ 1.000 ● N +0	$d^*\gamma=0,\quad d^*\Gamma=0$	$d^*\gamma=0,\quad d^*\Gamma=0$	$d^*\gamma=0,\quad d^*\Gamma=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0518 (7.7.36)	166	■ 1.000 ● N +0	$D\Phi=\left[\Omega^g,\Phi\right],$	$DD\Phi=\left[\Omega^g,\Phi\right],$	$DD\Phi=\left[\Omega^g,\Phi\right],$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0520 (7.7.38)	166	■ 1.000 ● C +3	$G:=G_{\{\alpha\}\{\beta\}}^{\{\star\}}\sigma^{\alpha\beta}:=\frac{1}{2}R^{\mu\nu}\wedge\eta_{\mu\nu\lambda}\gamma^\lambda=-i\gamma_5(\Omega^g\wedge\gamma+\gamma\wedge\Omega^g)=i[\gamma,\gamma_5\Omega^g].$	$G:=G_{\alpha}\gamma^{\alpha}:=\frac{1}{2}R^{\mu\nu}\wedge\eta_{\mu\nu\lambda}\gamma^{\lambda}=-i\gamma_5(\Omega^g\wedge\gamma+\gamma\wedge\Omega^g)=i[\gamma,\gamma_5\Omega^g].$	$G:=G_{\alpha}\gamma^{\alpha}:=\frac{1}{2}R^{\mu\nu}\wedge\eta_{\mu\nu\lambda}\gamma^{\lambda}=-i\gamma_5(\Omega^g\wedge\gamma+\gamma\wedge\Omega^g)=i[\gamma,\gamma_5\Omega^g].$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0521 (7.7.39)	167	■ 1.000 ● W +0	$D[\gamma, \Theta] \equiv 2i \left[\sigma, \Omega^g \right], \quad D \left[\gamma, \Omega^g \right] \equiv [\Theta, \Omega^g], \quad DG \equiv i [\Theta, \gamma_5 \Omega^g],$	$D[\gamma, \Theta] \equiv 2i [\sigma, \Omega^g], \quad D [\gamma, \Omega^g] \equiv [\Theta, \Omega^g], \quad DG \equiv i [\Theta, \gamma_5 \Omega^g],$	$D[\gamma, \Theta] \equiv 2i[\sigma, \Omega^g], \quad D[\gamma, \Omega^g] \equiv [\Theta, \Omega^g], \quad DG \equiv i[\Theta, \gamma_5 \Omega^g],$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0523 (7.7.42)	167	■ 1.000 ● N +0	$D R_{\alpha \beta}^{(\star)} \equiv 0,$	$DR_{\alpha \beta}^{(\star)} \equiv 0,$	$DR_{\alpha \beta}^{(\star)} \equiv 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0532 (7.7.52)	169	■ 1.000 ● K +0	$\{ \}^{(6)} R^{\alpha \beta} = -\frac{1}{12} R \vartheta^{\alpha \beta}$	${}^{(6)}R^{\alpha \beta} = -\frac{1}{12} R \vartheta^{\alpha \beta}$	${}^{(6)}R^{\alpha \beta} = -\frac{1}{12} R \vartheta^{\alpha \beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0541 (8.3.2)	177	■ 1.000 ● N +0	$s \gamma = \psi - [C, \gamma]$	$s \gamma = \psi - [C, \gamma]$	$s \gamma = \psi - [C, \gamma]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0542 (8.3.3)	177	■ 1.000 ● K +0	$s \sigma = \frac{i}{2} s (\gamma \wedge \gamma) = \frac{i}{2} ([\psi, \gamma] - [[C, \gamma], \gamma]) = \frac{i}{2} [\psi, \gamma] - [C, \sigma].$	$s \sigma := \frac{i}{2} s (\gamma \wedge \gamma) = \frac{i}{2} ([\psi, \gamma] - [[C, \gamma], \gamma]) = \frac{i}{2} [\psi, \gamma] - [C, \sigma].$	$s \sigma := \frac{i}{2} s (\gamma \wedge \gamma) = \frac{i}{2} ([\psi, \gamma] - [[C, \gamma], \gamma]) = \frac{i}{2} [\psi, \gamma] - [C, \sigma].$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0543 (8.3.4)	177	■ 1.000 ● N +0	$s \bar{c} = b, \quad s b = 0.$	$s \bar{c} = b, \quad s b = 0.$	$s \bar{c} = b, \quad s b = 0.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0546 (8.3.7)	178	■ 1.000 ● N +0	$d\,s^2=\frac{1}{4}\,\operatorname{Tr}\!\left\{\frac{\gamma}{\gamma}\otimes\frac{\gamma}{\gamma}\right\}\,o_{\alpha\beta}\,D\,x^{\alpha}\otimes D\,x^{\beta}$	$ds^2=\frac{1}{4}\operatorname{Tr}\{\gamma\otimes\gamma\}^*=o_{\alpha\beta}D\xi^\alpha\otimes D\xi^\beta$	$ds^2=\frac{1}{4}\operatorname{Tr}\{\gamma\otimes\gamma\}^*=o_{\alpha\beta}D\xi^\alpha\otimes D\xi^\beta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0550 (8.4.4)	179	■ 1.000 ● N -1	$L_{\mathrm{TG}}=d\,C_{\mathrm{RR}}+L_{\mathrm{FP}}=d\,C_{\mathrm{RR}}+s\{\cdots\}$	$L_{\mathrm{TG}}=dC_{\mathrm{RR}}+L_{\mathrm{FP}}=dC_{\mathrm{RR}}+s\{\cdots\}$	$L_{\mathrm{TG}}=dC_{\mathrm{RR}}+L_{\mathrm{FP}}=dC_{\mathrm{RR}}+s\{\cdots\}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0559 (8.5.2)	181	■ 1.000 ● N +0	$\sigma:=\frac{i}{2}\,\gamma\wedge\gamma\,.$	$\sigma:=\frac{i}{2}\gamma\wedge\gamma.$	$\sigma:=\frac{i}{2}\gamma\wedge\gamma.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0563 (8.5.6)	182	■ 1.000 ● N +0	$\Omega^{g(\theta)}:=\sqrt{2}\left(\Omega^g\sin\theta+{}^*\Omega^{g(*)}\cos\theta\right)$	$\Omega^{g(\theta)}:=\sqrt{2}\left(\Omega^g\sin\theta+{}^*\Omega^{g(*)}\cos\theta\right)$	$\Omega^{g(\theta)}:=\sqrt{2}\left(\Omega^g\sin\theta+{}^*\Omega^{g(*)}\cos\theta\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0564 (8.6.1)	183	■ 1.000 ● K +0	$L_{\mathrm{qPG}}=\frac{\Lambda}{\ell^2}\eta-\frac{a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2}T^\alpha\wedge H_\alpha-\frac{1}{2}R^{\alpha\beta}\wedge H_{\alpha\beta}\,.$	$L_{\mathrm{qPG}}=\frac{\Lambda}{\ell^2}\eta-\frac{a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2}T^\alpha\wedge H_\alpha-\frac{1}{2}R^{\alpha\beta}\wedge H_{\alpha\beta}.$	$L_{\mathrm{qPG}}=\frac{\Lambda}{\ell^2}\eta-\frac{a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2}T^\alpha\wedge H_\alpha-\frac{1}{2}R^{\alpha\beta}\wedge H_{\alpha\beta}\,.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0565 (8.6.2)	183	■ 1.000 ● N +1	$H_{\alpha}:= -\partial L_{\mathrm{qPG}}/\partial T^{\alpha} \, / \, \partial T^{\alpha} = \frac{1}{\ell^2} * \left(\sum_{M=1}^3 a_{(M)}^{(M)} T_{\alpha} \right)^{\wedge (M)} T_{[\alpha} \wedge H_{\beta]}$,	$H_{\alpha} := -\partial L_{\mathrm{qPG}}/\partial T^{\alpha} = \frac{1}{\ell^2} * \left(\sum_{M=1}^3 a_{(M)}^{(M)} T_{\alpha} \right),$	$H_{\alpha} := -\partial L_{\mathrm{qPG}}/\partial T^{\alpha} = \frac{1}{\ell^2} * \left(\sum_{M=1}^3 a_{(M)}^{(M)} T_{\alpha} \right),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0569 (8.6.7)	184	■ 1.000 ● N +0	$\begin{aligned} D H_{\alpha} - E_{\alpha} &= \Sigma_{\alpha}, \\ D H_{\alpha \beta} + \vartheta_{[\alpha} \wedge H_{\beta]} &= \tau_{\alpha \beta}, \end{aligned}$	$\begin{aligned} D H_{\alpha} - E_{\alpha} &= \Sigma_{\alpha}, \\ D H_{\alpha \beta} + \vartheta_{[\alpha} \wedge H_{\beta]} &= \tau_{\alpha \beta}, \end{aligned}$	$\begin{aligned} D H_{\alpha} - E_{\alpha} &= \Sigma_{\alpha}, \\ D H_{\alpha \beta} + \vartheta_{[\alpha} \wedge H_{\beta]} &= \tau_{\alpha \beta}, \end{aligned}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0571 (8.6.9)	184	■ 1.000 ● N -1	$\frac{\theta_{\mathrm{T}}^*}{2\ell^2} \eta_{\alpha\beta\gamma} T^{\gamma} + \frac{\theta_{\mathrm{T}}}{\ell^2} \vartheta_{[\alpha} \wedge T_{\beta]} + \vartheta_{[\alpha} \wedge H_{\beta]} (**) = \tau_{\alpha\beta}.$	$\frac{\theta_{\mathrm{T}}^*}{2\ell^2} \eta_{\alpha\beta\gamma} T^{\gamma} + \frac{\theta_{\mathrm{T}}}{\ell^2} \vartheta_{[\alpha} \wedge T_{\beta]} + \vartheta_{[\alpha} \wedge H_{\beta]} (**) = \tau_{\alpha\beta}.$	$\frac{\theta_{\mathrm{T}}^*}{2\ell^2} \eta_{\alpha\beta\gamma} T^{\gamma} + \frac{\theta_{\mathrm{T}}}{\ell^2} \vartheta_{[\alpha} \wedge T_{\beta]} + \vartheta_{[\alpha} \wedge H_{\beta]} (**) = \tau_{\alpha\beta}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0576 (8.6.14)	185	■ 1.000 ● N -1	$\Lambda_{\mathrm{eff}} = \frac{1}{\theta_{\mathrm{T}}^*} \left[\Lambda - \frac{1}{4} (\theta_{\mathrm{T}}^* - a_0) R \right] = \frac{\Lambda}{\theta_{\mathrm{T}}^*} - \Lambda_{\theta}$	$\Lambda_{\mathrm{eff}} = \frac{1}{\theta_{\mathrm{T}}^*} \left[\Lambda - \frac{1}{4} (\theta_{\mathrm{T}}^* - a_0) R \right] = \frac{\Lambda}{\theta_{\mathrm{T}}^*} - \Lambda_{\theta}$	$\Lambda_{\mathrm{eff}} = \frac{1}{\theta_{\mathrm{T}}^*} \left[\Lambda - \frac{1}{4} (\theta_{\mathrm{T}}^* - a_0) R \right] = \frac{\Lambda}{\theta_{\mathrm{T}}^*} - \Lambda_{\theta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0578 (8.6.16)	186	■ 1.000 ● K +0	$\widetilde{L}_{\mathrm{qPG}} = L_{\mathrm{qPG}} + \lambda_{\alpha} \wedge T^{\alpha}.$	$\widetilde{L}_{\mathrm{qPG}} = L_{\mathrm{qPG}} + \lambda_{\alpha} \wedge T^{\alpha}.$	$\widetilde{L}_{\mathrm{qPG}} = L_{\mathrm{qPG}} + \lambda_{\alpha} \wedge T^{\alpha}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E09582 (8.6.20)	188	1.000 N +0	$\Lambda_{\theta}=\frac{3\left(\theta_{\mathrm{T}}-\mathrm{a}_{\mathrm{0}}\right)^2}{2\ell^2\left(\theta_{\mathrm{L}}^{\star}-b_6\right)\theta_{\mathrm{T}}^{\star}}$	$\Lambda_{\theta}=\frac{3\left(\theta_{\mathrm{T}}^{\star}-a_0\right)^2}{2\ell^2\left(\theta_{\mathrm{L}}^{\star}-b_6\right)\theta_{\mathrm{T}}^{\star}}$	$\Lambda_{\theta}=\frac{3(\theta_{\mathrm{T}}^{\star}-a_0)^2}{2\ell^2(\theta_{\mathrm{L}}^{\star}-b_6)\theta_{\mathrm{T}}^{\star}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E09586 (9.1.4)	193	1.000 N +0	$ds^2=\left(\frac{N(r\pm t)}{r}\right)^2\left(-dt^2+dr^2+r^2d\Omega^2\right),$	$ds^2=\left(\frac{N(r\pm t)}{r}\right)^2\left(-dt^2+dr^2+r^2d\Omega^2\right),$	$ds^2=\left(\frac{N(r\pm t)}{r}\right)^2(-dt^2+dr^2+r^2d\Omega^2),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E09588 (9.1.6)	194	1.000 N +0	$u^2-v^2=e^{r^*/4\mu}$	$u^2-v^2=e^{r^*/4\mu}$	$u^2-v^2=e^{r^*/4\mu}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E09589 (9.1.7)	194	1.000 N +0	$(u+v)/(u-v)=e^{t/2\mu},$	$(u+v)/(u-v)=e^{t/2\mu},$	$(u+v)/(u-v)=e^{t/2\mu},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E09590 (9.1.8)	194	1.000 N +0	$ds^2=16\mu^2e^{\nu-r^*/2\mu}\left(-dv^2+du^2\right)+r^2d\Omega^2,$	$ds^2=16\mu^2e^{\nu-r^*/2\mu}\left(-dv^2+du^2\right)+r^2d\Omega^2,$	$ds^2=16\mu^2e^{v-r^*/2\mu}\left(-dv^2+du^2\right)+r^2d\Omega^2,$
Conf. MathPix confidence: ≥0.9 0.5–0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E08591 (9.1.9)	195	■ 1.000 ● N +0	$\Omega^{\{\}}=\Omega_{\mathrm{C}}^{\{\}}^{\{\}}+\bar{\Omega}^{\{\}}\vartheta\wedge\vartheta,$	$\Omega^{\{\}}=\Omega_{\mathrm{C}}^{\{\}}+\bar{\Omega}^{\{\}}-\frac{1}{6}R^{\{\}}\vartheta\wedge\vartheta,$	$\Omega^{\{\}}=\Omega_{\mathrm{C}}^{\{\}}+\bar{\Omega}^{\{\}}-\frac{1}{6}R^{\{\}}\vartheta\wedge\vartheta,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E08688 (9.1.18)	197	■ 1.000 ● W +1	$\begin{aligned}ds_{\mathrm{Sds}}^2&=\left(1-\frac{2}{\mu}\frac{\kappa}{\ell^2}\right)d\tau^2\\&+\left(1-\frac{2}{r}\frac{\kappa}{\ell^2}\right)^{-1}dr^2+r^2d\Omega^2,\end{aligned}$	$ds_{\mathrm{Sds}}^2=\left(1-\frac{2}{r}\frac{\kappa}{\ell^2}\right)d\tau^2+\left(1-\frac{2}{r}\frac{\kappa}{\ell^2}\right)^{-1}dr^2+r^2d\Omega^2$	$ds_{\mathrm{Sds}}^2=\left(1-\frac{2}{r}\frac{\kappa}{\ell^2}\right)d\tau^2+\left(1-\frac{2}{r}\frac{\kappa}{\ell^2}\right)^{-1}dr^2+r^2d\Omega^2,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E08681 (9.1.19)	197	■ 1.000 ● N +0	$\begin{aligned}&f=\frac{2}{r^2}\left(1-\frac{2}{r}\frac{\kappa}{\ell^2}\right)^{-1/2}\\&g=h=k=0\end{aligned}$	$f=\frac{2}{r^2}\left(1-\frac{2}{r}\frac{\kappa}{\ell^2}\right)^{-1/2}$ $g=h=k=0$	$f=\frac{2}{r^2}\left(1-\frac{2}{r}\frac{\kappa}{\ell^2}\right)^{-1/2}$ $g=h=k=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E08683 (9.2.2)	198	■ 1.000 ● K +0	$p_1(M^4)=-3\tau$	$p_1(M^4)=-3\tau$	$p_1(M^4)=-3\tau$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E08687 (9.2.8)	200	■ 1.000 ● W +0	$\begin{aligned}\chi(M\#N)&=\chi(M)+\chi(N)-\chi\left(S^n\right),\\ \chi(M\times N)&=\chi(M)\cdot\chi(N),\\ \chi(M^2)&=2(g-1),\\ \chi(\mathbb{C}P^2)&=3\end{aligned}$	$\chi(M\#N)=\chi(M)+\chi(N)-\chi(S^n),$ $\chi(M\times N)=\chi(M)\cdot\chi(N),$ $\chi(M^2)=2(g-1),$ $\chi(\mathbb{C}P^2)=3$	$\chi(M\#N)=\chi(M)+\chi(N)-\chi(S^n),$ $\chi(M\times N)=\chi(M)\cdot\chi(N),$ $\chi(M^2)=2(g-1),$ $\chi(\mathbb{C}P^2)=3$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0608 (9.2.10)	201	■ 1.000 ● K +0	$\chi\left(M^{\{2\ k+1\}}\right)=0$	$\chi\left(M^{2k+1}\right)=0$	$\chi(M^{2k+1})=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0612 (10.0.1)	208	■ 1.000 ● N +0	$L_{\mathrm{EC}}=-\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}$	$L_{\mathrm{EC}}=-\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}$	$L_{\mathrm{EC}}=-\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0614 (10.1.2)	209	■ 1.000 ● N +0	$T^{\{\alpha\}}:=d\vartheta^{\{\alpha\}}-(-1)^s\vartheta^{\{\alpha}\ \beta\}\wedge\Gamma_{\beta}^{\star}$	$T^{\alpha}:=d\vartheta^{\alpha}-(-1)^s\eta^{\alpha\beta}\wedge\Gamma_{\beta}^{\star}$	$T^{\alpha}:=d\vartheta^{\alpha}-(-1)^s\eta^{\alpha\beta}\wedge\Gamma_{\beta}^{\star}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0615 (10.1.3)	209	■ 1.000 ● W +0	$R_{\{\alpha\}}^{\{\star\}}=\frac{1}{2}\vartheta_{\{\alpha}\ \beta\ \gamma\}\wedge\Gamma_{\beta\ \gamma}^{\star}$ $R^{\{\beta\}\ \gamma\}\wedge\Gamma_{\beta\ \gamma}^{\star}:=d\Gamma_{\alpha}^{\{\star\}}+\frac{1}{2}(-1)^s\vartheta_{\alpha}^{\{\beta\}\ \gamma\}\wedge\Gamma_{\gamma}^{\{\star\}}\wedge\Gamma_{\beta}^{\{\star\}},$	$R_{\alpha}^{\star}=\frac{1}{2}\eta_{\alpha\beta\gamma}R^{\beta\gamma}:=d\Gamma_{\alpha}^{\star}+\frac{(-1)^s}{2}\eta_{\alpha}^{\beta\gamma}\Gamma_{\beta}^{\star}\wedge\Gamma_{\gamma}^{\star},$	$R_{\alpha}^{\star}=\frac{1}{2}\eta_{\alpha\beta\gamma}R^{\beta\gamma}:=d\Gamma_{\alpha}^{\star}+\frac{(-1)^s}{2}\eta_{\alpha}^{\beta\gamma}\Gamma_{\beta}^{\star}\wedge\Gamma_{\gamma}^{\star},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0616 (10.1.4)	209	■ 1.000 ● N +0	$D\ T^{\{\alpha\}}\equiv(-1)^s\vartheta^{\{\alpha}\ \beta\}\wedge R_{\beta}^{\star}$	$DT^{\alpha}\equiv(-1)^s\eta^{\alpha\beta}\wedge R_{\beta}^{\star}$	$DT^{\alpha}\equiv(-1)^s\eta^{\alpha\beta}\wedge R_{\beta}^{\star}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0617 (10.1.5)	209	■1.000 ● N +0	$D\,R_{\alpha}^{\star}\equiv 0\;.$	$DR_{\alpha}^{\star}\equiv 0.$	$DR_{\alpha}^{\star}\equiv 0\,.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0618 (10.1.6)	209	■1.000 ● N +0	$C_{\mathrm{T}}:=\frac{1}{2\,\ell^2}\,\vartheta^{\alpha}\wedge T_{\alpha}=\frac{(-1)^s}{\ell^2}\,\eta^{\alpha}\wedge K_{\alpha}^{\star},$ $C_{\mathrm{T}}:=\frac{1}{2\ell^2}\vartheta^{\alpha}\wedge T_{\alpha}=-\frac{(-1)^s}{\ell^2}\eta^{\alpha}\wedge K_{\alpha}^{\star},$	$C_{\mathrm{T}}:=\frac{1}{2\ell^2}\vartheta^{\alpha}\wedge T_{\alpha}=-\frac{(-1)^s}{\ell^2}\eta^{\alpha}\wedge K_{\alpha}^{\star},$	$C_{\mathrm{T}}:=\frac{1}{2\ell^2}\vartheta^{\alpha}\wedge T_{\alpha}=-\frac{(-1)^s}{\ell^2}\eta^{\alpha}\wedge K_{\alpha}^{\star},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0619 (10.1.7)	209	■1.000 ● K +0	$C_{\mathrm{L}}:=(-1)^s\Gamma^{\star\alpha}\wedge R_{\alpha}^{\star}-\frac{1}{3!}\eta_{\alpha\beta\gamma}\Gamma^{\star\alpha}\wedge\Gamma^{\star\beta}\wedge\Gamma^{\star\gamma}.$	$C_{\mathrm{L}}:=(-1)^s\Gamma^{\star\alpha}\wedge R_{\alpha}^{\star}-\frac{1}{3!}\eta_{\alpha\beta\gamma}\Gamma^{\star\alpha}\wedge\Gamma^{\star\beta}\wedge\Gamma^{\star\gamma}.$	$C_{\mathrm{L}}:=(-1)^s\Gamma^{\star\alpha}\wedge R_{\alpha}^{\star}-\frac{1}{3!}\eta_{\alpha\beta\gamma}\Gamma^{\star\alpha}\wedge\Gamma^{\star\beta}\wedge\Gamma^{\star\gamma}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0621 (10.1.9)	209	■1.000 ● N -2	$\frac{\delta C_{\mathrm{L}}}{\delta\vartheta^{\alpha}}=0\quad\text{and}\quad\frac{\delta C_{\mathrm{L}}}{\delta\Gamma^{\star\alpha}}=(-1)^s2R_{\alpha}^{\star},$	$\frac{\delta C_{\mathrm{L}}}{\delta\vartheta^{\alpha}}=0\quad\text{and}\quad\frac{\delta C_{\mathrm{L}}}{\delta\Gamma^{\star\alpha}}=(-1)^s2R_{\alpha}^{\star},$	$\frac{\delta C_{\mathrm{L}}}{\delta\vartheta^{\alpha}}=0\quad\text{and}\quad\frac{\delta C_{\mathrm{L}}}{\delta\Gamma^{\star\alpha}}=(-1)^s2R_{\alpha}^{\star},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0622 (10.1.10)	210	■1.000 ● N -1	$C_{\mathrm{TL}}:=\frac{1}{\ell}\left(\Gamma^{\star\alpha}\wedge T_{\alpha}-\frac{(-1)^s}{2}\eta_{\alpha\beta\gamma}\Gamma^{\star\alpha}\wedge\Gamma^{\star\beta}\wedge\vartheta^{\gamma}\right).$	$C_{\mathrm{TL}}:=\frac{1}{\ell}\left(\Gamma^{\star\alpha}\wedge T_{\alpha}-\frac{(-1)^s}{2}\eta_{\alpha\beta\gamma}\Gamma^{\star\alpha}\wedge\Gamma^{\star\beta}\wedge\vartheta^{\gamma}\right).$	$C_{\mathrm{TL}}:=\frac{1}{\ell}\left(\Gamma^{\star\alpha}\wedge T_{\alpha}-\frac{(-1)^s}{2}\eta_{\alpha\beta\gamma}\Gamma^{\star\alpha}\wedge\Gamma^{\star\beta}\wedge\vartheta^{\gamma}\right).$
Conf. MathPix confidence: ■≥0.9 ■0.5-0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0625 (10.1.13)	210	■ 1.000 ● N +0	$D\,H_{\alpha}^{\star}-S_{\alpha}=\tau_{\alpha}^{\star},$	$DH_{\alpha}^{\star}-S_{\alpha}=\tau_{\alpha}^{\star},$	$DH_{\alpha}^{\star}-S_{\alpha}=\tau_{\alpha}^{\star},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0626 (10.1.14)	211	■ 1.000 ● K +0	$H_{\alpha}:=\frac{\partial L}{\partial T^{\alpha}},\quad H_{\alpha}^{\star}:=\frac{\partial L}{\partial R^{\star\alpha}},$	$H_{\alpha}:= -\frac{\partial L}{\partial T^{\alpha}},\quad H_{\alpha}^{\star}:= -\frac{(-1)^s}{2}\frac{\partial L}{\partial R^{\star\alpha}},$	$H_{\alpha}:= -\frac{\partial L}{\partial T^{\alpha}},\quad H_{\alpha}^{\star}:= -\frac{(-1)^s}{2}\frac{\partial L}{\partial R^{\star\alpha}},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0628 (10.1.16)	211	■ 1.000 ● N +0	$S_{\alpha}:=\frac{\partial L}{\partial \Gamma^{\star\alpha}}=\frac{1}{2}\eta_{\alpha\beta}\wedge H^{\beta}.$	$S_{\alpha}:=\frac{(-1)^s}{2}\frac{\partial L}{\partial \Gamma^{\star\alpha}}=\frac{1}{2}\eta_{\alpha\beta}\wedge H^{\beta}.$	$S_{\alpha}:=\frac{(-1)^s}{2}\frac{\partial L}{\partial \Gamma^{\star\alpha}}=\frac{1}{2}\eta_{\alpha\beta}\wedge H^{\beta}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0629 (10.1.17)	211	■ 1.000 ● N +0	$\Sigma_{\alpha}:=\frac{\delta L_{\mathrm{MAT}}}{\delta \vartheta^{\alpha}},\quad \tau_{\alpha}^{\star}:=\frac{\delta L_{\mathrm{MAT}}}{\delta \Gamma^{\star\alpha}},$	$\Sigma_{\alpha}:=\frac{\delta L_{\mathrm{MAT}}}{\delta \vartheta^{\alpha}},\quad \tau_{\alpha}^{\star}:=\frac{(-1)^s}{2}\frac{\delta L_{\mathrm{MAT}}}{\delta \Gamma^{\star\alpha}},$	$\Sigma_{\alpha}:=\frac{\delta L_{\mathrm{MAT}}}{\delta \vartheta^{\alpha}},\quad \tau_{\alpha}^{\star}:=\frac{(-1)^s}{2}\frac{\delta L_{\mathrm{MAT}}}{\delta \Gamma^{\star\alpha}},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0630 (10.2.1)	211	■ 1.000 ● N -1	$L_{\mathrm{EC}}:= -\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}\equiv -\chi\,C_{\mathrm{TL}}-\frac{\chi}{\ell}d(\Gamma_{\alpha}^{\star}\wedge\vartheta^{\alpha})$	$L_{\mathrm{EC}}:= -\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}\equiv -\chi\,C_{\mathrm{TL}}-\frac{\chi}{\ell}d(\Gamma_{\alpha}^{\star}\wedge\vartheta^{\alpha})$	$L_{\mathrm{EC}}:= -\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}\equiv -\chi\,C_{\mathrm{TL}}-\frac{\chi}{\ell}d(\Gamma_{\alpha}^{\star}\wedge\vartheta^{\alpha})$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0631 (10.2.2)	211	■1.000 ●C -5	$L_{\mathrm{MB}}(\vartheta^{\alpha}, \Gamma_{\alpha}^{\star}) = \theta_{\mathrm{T}} C_{\mathrm{T}} + \theta_{\mathrm{L}} C_{\mathrm{L}} + \theta_{\mathrm{TL}} C_{\mathrm{TL}},$	$L_{\mathrm{MB}}(\vartheta^{\alpha}, \Gamma_{\alpha}^{\star}) = \theta_{\mathrm{T}} C_{\mathrm{T}} + \theta_{\mathrm{L}} C_{\mathrm{L}} + \theta_{\mathrm{TL}} C_{\mathrm{TL}},$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0632 (10.2.3)	212	■1.000 ●C -3	$-\theta_{\mathrm{TL}} R_{\alpha}^{\star} - \frac{1}{\ell} \theta_{\mathrm{T}} T_{\alpha} = \ell \Sigma_{\alpha}$	$-\theta_{\mathrm{TL}} R_{\alpha}^{\star} - \frac{1}{\ell} \theta_{\mathrm{T}} T_{\alpha} = \ell \Sigma_{\alpha}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0634 (10.2.5)	212	■1.000 ●N -2	$T_{\alpha} - \frac{\varepsilon}{\ell} \eta_{\alpha} = \frac{2}{A} \ell (\theta_{\mathrm{TL}} \tau_{\alpha}^{\star} - \theta_{\mathrm{L}} \ell \Sigma_{\alpha})$	$T_{\alpha} - \frac{\varepsilon}{\ell} \eta_{\alpha} = \frac{2}{A} \ell (\theta_{\mathrm{TL}} \tau_{\alpha}^{\star} - \theta_{\mathrm{L}} \ell \Sigma_{\alpha})$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0635 (10.2.6)	212	■1.000 ●C -3	$R_{\alpha}^{\star} - \frac{\rho}{\ell^2} \eta_{\alpha} = \frac{2}{A} (\theta_{\mathrm{TL}} \ell \Sigma_{\alpha} - \theta_{\mathrm{T}} \tau_{\alpha}^{\star})$	$R_{\alpha}^{\star} - \frac{\rho}{\ell^2} \eta_{\alpha} = \frac{2}{A} (\theta_{\mathrm{TL}} \ell \Sigma_{\alpha} - \theta_{\mathrm{T}} \tau_{\alpha}^{\star})$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0638 (10.2.9)	213	■1.000 ●N +0	$R_{\alpha}^{\star} \equiv R_{\alpha}^{\{\} \star} - DK_{\alpha}^{\star} + \frac{1}{2} \eta_{\alpha \beta \gamma} K^{\star \beta} \wedge K^{\star \gamma}$	$R_{\alpha}^{\star} \equiv R_{\alpha}^{\{\} \star} - DK_{\alpha}^{\star} + \frac{1}{2} \eta_{\alpha \beta \gamma} K^{\star \beta} \wedge K^{\star \gamma}$	
Conf. MathPix confidence: ■≥0.9 ■0.5-0.9 ■<0.5 —no value Residual render vs scan (inkdrill): ●C ●W ●S ●N ●K ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0640 (10.2.11)	213	■ 1.000 ● C -4	$\Lambda_{\mathrm{eff}}=-\left[4\,\rho-\left(1+2(-1)^s\right)\varepsilon^2\right]/4\,\ell^2=\theta_{\mathrm{T}}^2\left[\left(9+2(-1)^s\right)\theta_{\mathrm{TL}}^2+8\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right]/(2A\ell)^2$	$\Lambda_{\mathrm{eff}}=-\left[4\rho-\left(1+2(-1)^s\right)\varepsilon^2\right]/4\ell^2=\theta_{\mathrm{T}}^2\left[\left(9+2(-1)^s\right)\theta_{\mathrm{TL}}^2+8\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right]/(2A\ell)^2$	$\Lambda_{\mathrm{eff}}=-[4\rho-(1+2(-1)^s)\varepsilon^2]/4\ell^2=\theta_{\mathrm{T}}^2[(9+2(-1)^s)\theta_{\mathrm{TL}}^2+8\theta_{\mathrm{T}}\theta_{\mathrm{L}}]/(2A\ell)^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0642 (10.2.13)	213	■ 1.000 ● N +0	$ds^2=-N^2(r)dt^2+N^{-2}(r)dr^2+r^2\left[d\phi+N^\phi(r)dt\right]^2$	$ds^2=-N^2(r)dt^2+N^{-2}(r)dr^2+r^2\left[d\phi+N^\phi(r)dt\right]^2$	$ds^2=-N^2(r)dt^2+N^{-2}(r)dr^2+r^2\left[d\phi+N^\phi(r)dt\right]^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0643 (10.2.14)	213	■ 1.000 ● N -1	$N^2(r)=-M-\Lambda_{\mathrm{eff}}r^2+\frac{J^2}{4r^2},\quad N^\phi(r)=-\frac{J}{2r^2},$	$N^2(r)=-M-\Lambda_{\mathrm{eff}}r^2+\frac{J^2}{4r^2},\quad N^\phi(r)=-\frac{J}{2r^2},$	$N^2(r)=-M-\Lambda_{\mathrm{eff}}r^2+\frac{J^2}{4r^2},\quad N^\phi(r)=-\frac{J}{2r^2},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0647 (10.3.4)	214	■ 1.000 ● K +0	$S_{\alpha}=\frac{\ell}{\varepsilon}E_{\alpha}$	$S_{\alpha}=\frac{\ell}{\varepsilon}E_{\alpha}$	$S_{\alpha}=\frac{\ell}{\varepsilon}E_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0648 (10.3.5)	215	■ 1.000 ● N +0	$H_{\alpha}^{\star}=\frac{\ell}{\varepsilon}H_{\alpha}$	$H_{\alpha}^{\star}=\frac{\ell}{\varepsilon}H_{\alpha}$	$H_{\alpha}^{\star}=\frac{\ell}{\varepsilon}H_{\alpha}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ E00649 (10.3.6)	215	■ 1.000 ● N +1	$T_{\{\alpha\}}=\frac{\varepsilon}{\ell}\,\eta_{\alpha},\quad R_{\{\alpha\}}^{\star}=\frac{\rho}{\ell^2}\,\eta_{\alpha}$	$T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha}$	$T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ E00650 (10.3.7)	215	■ 1.000 ● N +0	$H_{\{\alpha\}}^{\star}=\delta\ell\,\eta_{\alpha}+\frac{\gamma}{\ell}\,\vartheta_{\alpha}$	$H_{\alpha}^{\star}=\delta\ell H_{\alpha}+\frac{\gamma}{\ell}\vartheta_{\alpha}$	$H_{\alpha}^{\star}=\delta\ell H_{\alpha}+\frac{\gamma}{\ell}\vartheta_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ E00651 (10.3.8)	215	■ 1.000 ● N +1	$\delta\ell\,D\,H_{\{\alpha\}}+\frac{\gamma}{\ell}\,T_{\{\alpha\}}-\frac{1}{2}\,\eta_{\alpha\beta}\,\eta_{\beta\gamma}\,\vartheta^{\gamma}=\tau_{\alpha}^{\star}$	$\delta\ell DH_{\alpha}+\frac{\gamma}{\ell}T_{\alpha}-\frac{1}{2}\eta_{\alpha\beta}\wedge H^{\beta}=\tau_{\alpha}^{\star}$	$\delta\ell DH_{\alpha}+\frac{\gamma}{\ell}T_{\alpha}-\frac{1}{2}\eta_{\alpha\beta}\wedge H^{\beta}=\tau_{\alpha}^{\star}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ E00653 (10.3.10)	216	■ 1.000 ● K +0	$H_{\{\alpha\}}^{\star}=\frac{\gamma}{\ell(1+\delta\varepsilon)}\vartheta_{\alpha}=-\frac{\ell}{\varepsilon}H_{\alpha},$	$H_{\alpha}^{\star}=\frac{\gamma}{\ell(1+\delta\varepsilon)}\vartheta_{\alpha}=-\frac{\ell}{\varepsilon}H_{\alpha},$	$H_{\alpha}^{\star}=\frac{\gamma}{\ell(1+\delta\varepsilon)}\vartheta_{\alpha}=-\frac{\ell}{\varepsilon}H_{\alpha},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ E00659 (10.3.16)	217	■ 1.000 ● K +0	$H=\vartheta^{\alpha}\wedge H_{\alpha}$	$H=\vartheta^{\alpha}\wedge H_{\alpha},$	$H=\vartheta^{\alpha}\wedge H_{\alpha},$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0660 (10.3.17)	217	■ 1.000 ● N +0	$t=\{ \}^*\backslash\left(T^{\alpha}\wedge H_{\alpha}\right), \quad \mathscr{A}:=\{ \}^*\backslash\left(\vartheta^{\alpha}\wedge T_{\alpha}\right), \quad h=\{ \}^*\backslash\left(\eta^{\alpha}\wedge H_{\alpha}\right),$	$t=*(T^{\alpha}\wedge H_{\alpha}), \quad \mathscr{A}:=*(\vartheta^{\alpha}\wedge T_{\alpha}), \quad h=*(\eta^{\alpha}\wedge H_{\alpha}),$	$t=*(T^{\alpha}\wedge H_{\alpha}), \quad \mathscr{A}:=*(\vartheta^{\alpha}\wedge T_{\alpha}), \quad h=*(\eta^{\alpha}\wedge H_{\alpha}),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0663 (10.3.20)	217	■ 1.000 ● N +2	$d\,s^2=(-1)^s\,d\,t^2+d\,\mathscr{A}^2+d\,h^2$	$ds^2=(-1)^sdt^2+d\mathscr{A}^2+dh^2$	$ds^2=(-1)^sdt^2+d\mathscr{A}^2+dh^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0665 (10.3.22)	218	■ 1.000 ● N +0	$D^{\{*\}}\vartheta_{\alpha}:=\frac{\ell}{\gamma}\left[D^*H_{\alpha}^{\star}-\delta\ell D^*H_{\alpha}\right]$	$D^*\vartheta_{\alpha}:=\frac{\ell}{\gamma}\left[D^*H_{\alpha}^{\star}-\delta\ell D^*H_{\alpha}\right]$	$D^*\vartheta_{\alpha}:=\frac{\ell}{\gamma}\left[D^*H_{\alpha}^{\star}-\delta\ell D^*H_{\alpha}\right]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0668 (10.3.25)	219	■ 1.000 ● N +0	$D^{\{*\}}\vartheta^{\alpha}\cong 0,$	$D^*\vartheta^{\alpha}\cong 0,$	$D^*\vartheta^{\alpha}\cong 0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0672 (10.3.29)	220	■ 1.000 ● N +0	$P_{\alpha}=\int\check{\mathscr{E}}_{\alpha}, \quad J_{\alpha}=\int\check{\mathscr{J}}_{\alpha}^{\star}.$	$P_{\alpha}=\int\check{\mathscr{E}}_{\alpha}, \quad J_{\alpha}=\int\check{\mathscr{J}}_{\alpha}^{\star}.$	$P_{\alpha}=\int\check{\mathscr{E}}_{\alpha}, \quad J_{\alpha}=\int\check{\mathscr{J}}_{\alpha}^{\star}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0675 (10.3.32)	220	■ 1.000 ● N -1	$\check{\mathrm{J}}_{\alpha}^{\star}=-\frac{\theta_{\mathrm{L}}}{2}\mathrm{d}\,\Gamma_{\alpha}^{\star}$	$\check{\mathcal{J}}_{\alpha}^{\star}=-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$	$\check{\mathcal{J}}_{\alpha}^{\star}=-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0677 (10.3.34)	220	■ 1.000 ● N -1	$\check{\mathrm{J}}_{\alpha}^{\mathrm{MB}\star}=(-1)^s\frac{\chi}{2\ell}d\vartheta_{\alpha}-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$	$\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}\star}=(-1)^s\frac{\chi}{2\ell}d\vartheta_{\alpha}-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$	$\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}\star}=(-1)^s\frac{\chi}{2\ell}d\vartheta_{\alpha}-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0679 (10.3.36)	221	■ 1.000 ● K +0	$\widetilde{\mathrm{C}}_{\alpha}^{\star}=(-1)^s\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{R}_{\alpha}^{\star}-\frac{1}{3!}\eta_{\alpha\beta\gamma}\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{\Gamma}^{\star\beta}\wedge\widetilde{\Gamma}^{\star\gamma},$	$\widetilde{C}_{\mathrm{L}}=(-1)^s\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{R}_{\alpha}^{\star}-\frac{1}{3!}\eta_{\alpha\beta\gamma}\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{\Gamma}^{\star\beta}\wedge\widetilde{\Gamma}^{\star\gamma},$	$\widetilde{C}_{\mathrm{L}}=(-1)^s\widetilde{F}^{\star\alpha}\wedge\widetilde{R}_{\alpha}^{\star}-\frac{1}{3!}\eta_{\alpha\beta\gamma}\widetilde{F}^{\star\alpha}\wedge\widetilde{F}^{\star\beta}\wedge\widetilde{F}^{\star\gamma},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0683 (10.3.41)	221	■ 1.000 ● N -1	$c=12\cdot 4\pi\theta_{\mathrm{L}},\quad\mathrm{t}\tilde{c}=12\cdot 4\pi\tilde{\theta}_{\mathrm{L}},$	$c=12\cdot 4\pi\theta_{\mathrm{L}},\quad\tilde{c}=12\cdot 4\pi\tilde{\theta}_{\mathrm{L}},$	$c=12\cdot 4\pi\theta_{\mathrm{L}},\quad\tilde{c}=12\cdot 4\pi\tilde{\theta}_{\mathrm{L}},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0684 (10.3.42)	222	■ 1.000 ● K +0	$\widetilde{L}=L+\lambda_{\alpha}\wedge T^{\alpha}.$	$\widetilde{L}=L+\lambda_{\alpha}\wedge T^{\alpha}.$	$\widetilde{L}=L+\lambda_{\alpha}\wedge T^{\alpha}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0691 (10.5.2)	223	■ 1.000 ● K +0	$\left[\sigma^{\alpha}, \sigma^{\beta}\right]=2 \mathrm{i} \sigma^{\alpha \beta \gamma} \sigma_{\gamma} .$	$\left[\sigma^{\alpha}, \sigma^{\beta}\right]=2 i \eta^{\alpha \beta \gamma} \sigma_{\gamma} .$	$\left[\sigma^{\alpha}, \sigma^{\beta}\right]=2 i \eta^{\alpha \beta \gamma} \sigma_{\gamma} .$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0692 (10.5.3)	224	■ 1.000 ● K +0	$\gamma_{\{0\}}=\mathrm{i} \sigma^{\{2\}}, \quad \gamma_{\{1\}}=\sigma^{\{1\}}, \quad \gamma_{\{2\}}=\sigma^{\{3\}} .$	$\gamma_0=i \sigma^2, \quad \gamma_1=\sigma^1, \quad \gamma_2=\sigma^3 .$	$\gamma_0=i \sigma^2, \quad \gamma_1=\sigma^1, \quad \gamma_2=\sigma^3 .$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0693 (10.5.4)	224	■ 1.000 ● N +0	$\gamma_{\{\alpha\}} \gamma_{\{\beta\}}=g_{\{\alpha \beta\}} \mathbf{1}+\eta_{\alpha \beta \nu} \gamma^{\nu},$	$\gamma_{\alpha} \gamma_{\beta}=g_{\alpha \beta} \mathbf{1}+\eta_{\alpha \beta \nu} \gamma^{\nu},$	$\gamma_{\alpha} \gamma_{\beta}=g_{\alpha \beta} \mathbf{1}+\eta_{\alpha \beta \nu} \gamma^{\nu},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0695 (10.5.6)	224	■ 1.000 ● N +0	$\mathrm{i}^{\{*\}} \gamma \wedge D \psi-m \psi \eta-\frac{\mathrm{i}}{2}\left(D^{\{*\}} \gamma\right) \psi=0$	$i^* \gamma \wedge D \psi-m \psi \eta-\frac{i}{2}\left(D^* \gamma\right) \psi=0$	$i^* \gamma \wedge D \psi-m \psi \eta-\frac{i}{2}\left(D^* \gamma\right) \psi=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0697 (10.5.8)	224	■ 1.000 ● K +0	$\hat{D}:=D-\frac{1}{2} T,$	$\hat{D}:=D-\frac{1}{2} T,$	$\hat{D}:=D-\frac{1}{2} T,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0698 (10.5.9)	224	■ 1.000 ● N +0	$i^{\ast}\gamma\wedge\hat{D}\psi-m\eta\psi=0,$	$i^{\ast}\gamma\wedge\hat{D}\psi-m\eta\psi=0,$	$i^{\ast}\gamma\wedge\hat{D}\psi-m\eta\psi=0,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0699 (10.5.10)	224	■ 1.000 ● N +0	$\hat{R}=R-\frac{1}{2}\,dT=\frac{i}{4}\,R^{\alpha\beta}\sigma_{\alpha\beta}-\frac{1}{2}\,dT$	$\hat{R}=R-\frac{1}{2}\,dT=\frac{i}{4}\,R^{\alpha\beta}\sigma_{\alpha\beta}-\frac{1}{2}\,dT$	$\hat{R}=R-\frac{1}{2}\,dT=\frac{i}{4}\,R^{\alpha\beta}\sigma_{\alpha\beta}-\frac{1}{2}\,dT$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0703 (10.6.2)	225	■ 1.000 ● N +0	$d^{\ast}F-m_{\text{photon}}F=0$	$d^{\ast}F-m_{\text{photon}}F=0$	$d^{\ast}F-m_{\text{photon}}F=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0705 (10.7.1)	226	■ 1.000 ● N +0	$X:=x_{\alpha}\vartheta^{\alpha},\quad Y:=y_{\alpha}\vartheta^{\alpha}.$	$X:=x_{\alpha}\vartheta^{\alpha},\quad Y:=y_{\alpha}\vartheta^{\alpha}.$	$X:=x_{\alpha}\vartheta^{\alpha},\quad Y:=y_{\alpha}\vartheta^{\alpha}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0706 (10.7.2)	226	■ 1.000 ● N +0	$\Sigma_{\alpha}=\varepsilon x_{\alpha}X\wedge Y,\quad \tau_{\alpha}^{\star}=\sigma y_{\alpha}X\wedge Y,$	$\Sigma_{\alpha}=\varepsilon x_{\alpha}X\wedge Y,\quad \tau_{\alpha}^{\star}=\sigma y_{\alpha}X\wedge Y,$	$\Sigma_{\alpha}=\varepsilon x_{\alpha}X\wedge Y,\quad \tau_{\alpha}^{\star}=\sigma y_{\alpha}X\wedge Y,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0707 (10.7.3)	226	■ 1.000 ● N +0	$\vartheta^{\alpha}\wedge\Sigma_{\alpha}=0,\quad \vartheta^{\alpha}\wedge\tau_{\alpha}^{\star}=0$	$\vartheta^{\alpha}\wedge\Sigma_{\alpha}=0,\quad \vartheta^{\alpha}\wedge\tau_{\alpha}^{\star}=0$	$\vartheta^{\alpha}\wedge\Sigma_{\alpha}=0,\quad \vartheta^{\alpha}\wedge\tau_{\alpha}^{\star}=0$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0708 (10.7.4)	227	■ 1.000 ● N +0	$R^{\alpha\beta}=\varepsilon\ell^2x^{\left[\alpha}y^{\beta\right]}X\wedge Y,$	$R^{\alpha\beta}=\varepsilon\ell^2x^{\left[\alpha}y^{\beta\right]}X\wedge Y,$	$R^{\alpha\beta}=\varepsilon\ell^2x^{\left[\alpha}y^{\beta\right]}X\wedge Y,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0711 (10.7.7)	228	■ 1.000 ● N +0	$\mathrm{\mathscr{A}}:=\left\{\vartheta^{\alpha}\wedge T_{\alpha}\right\} =-(-1)^s\frac{\varepsilon}{\ell^2}$	$\mathscr{A}:=\left(\vartheta^{\alpha}\wedge T_{\alpha}\right)=-(-1)^s\frac{\varepsilon}{\ell^2}$	$\mathscr{A}:=\left(\vartheta^{\alpha}\wedge T_{\alpha}\right)=-(-1)^s\frac{\varepsilon}{\ell^2}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0713 (10.8.2)	228	■ 1.000 ● K +0	$\gamma=\gamma_{\alpha}\vartheta^{\alpha}.$	$\gamma=\gamma_{\alpha}\vartheta^{\alpha}.$	$\gamma=\gamma_{\alpha}\vartheta^{\alpha}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0714	228	■ 1.000 ● K +0	$\bar{\Psi}\wedge\Psi=0,\quad\bar{\Psi}\wedge\gamma_5\Psi=0,\quad\bar{\Psi}\wedge\gamma_5\Psi=0.$	$\bar{\Psi}\wedge\Psi=0,\quad\bar{\Psi}\wedge\gamma_5\gamma^{\alpha}\Psi=0,\quad\bar{\Psi}\wedge\gamma_5\Psi=0.$	$\bar{\Psi}\wedge\Psi=0,\quad\bar{\Psi}\wedge\gamma_5\gamma^{\alpha}\Psi=0,\quad\bar{\Psi}\wedge\gamma_5\Psi=0.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0715 (10.9.1)	229	■ 1.000 ● N -1	$L_{\mathrm{RS}}=\frac{i}{4}(\bar{\Psi}\wedge D\Psi-\Psi\wedge\overline{D\Psi})+\frac{i}{4}m\bar{\Psi}\wedge\gamma\wedge\Psi,$	$L_{\mathrm{RS}}=\frac{i}{4}(\bar{\Psi}\wedge D\Psi-\Psi\wedge\overline{D\Psi})+\frac{i}{4}m\bar{\Psi}\wedge\gamma\wedge\Psi,$	$L_{\mathrm{RS}}=\frac{i}{4}(\overline{\Psi}\wedge D\Psi-\Psi\wedge\overline{D\Psi})+\frac{i}{4}m\overline{\Psi}\wedge\gamma\wedge\Psi,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0719 (10.9.5)	229	■ 1.000 ● N +1	$\Sigma_{\alpha}=-\frac{i}{4}m\bar{\Psi}\wedge\gamma_{\alpha}\Psi=-(-1)^s2m\tau_{\alpha}^{\star}$	$\Sigma_{\alpha}=-\frac{i}{4}m\bar{\Psi}\wedge\gamma_{\alpha}\Psi=-(-1)^s2m\tau_{\alpha}^{\star}$	$\Sigma_{\alpha}=-\frac{i}{4}m\overline{\Psi}\wedge\gamma_{\alpha}\Psi=-(-1)^s2m\tau_{\alpha}^{\star},$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E00721 (10.10.1)	230	■1.000 ● N +0	$L=L_{\infty}\left(\vartheta^{\alpha},\,\Gamma_{\alpha}^{\star},\,\Psi\right)=L_{\mathrm{MB}}+L_{\Psi}$	$L=L_{\infty}\left(\vartheta^{\alpha},\Gamma_{\alpha}^{\star},\Psi\right)=L_{\mathrm{MB}}+L_{\Psi}$	$L=L_{\infty}(\vartheta^{\alpha},\Gamma_{\alpha}^{\star},\Psi)=L_{\mathrm{MB}}+L_{\psi}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E00726 (10.10.6)	230	■1.000 ● N +0	$\frac{2}{i}\frac{\delta L}{\delta \Psi}=\frac{1}{2}m\gamma\wedge\Psi\cong 0$	$\frac{2}{i}\frac{\delta L}{\delta \Psi}=D\Psi+\frac{1}{2}m\gamma\wedge\Psi\cong 0$	$\frac{2}{i}\frac{\delta L}{\delta \Psi}=D\Psi+\frac{1}{2}m\gamma\wedge\Psi\cong 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E00729 (10.10.9)	231	■1.000 ● N +0	$\gamma\wedge\gamma=-2\gamma^{\alpha}\eta_{\alpha}$	$\gamma\wedge\gamma=-2\gamma^{\alpha}\eta_{\alpha}$	$\gamma\wedge\gamma=-2\gamma^{\alpha}\eta_{\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E00730 (10.10.10)	231	■1.000 ● W -2	$i\left[\left(\frac{\theta_{\mathrm{TL}}}{\ell}-1\right)R_{\alpha}^{\star}+\left(\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right)T_{\alpha}+\frac{m^2}{2}\eta_{\alpha}\right]\wedge\gamma^{\alpha}\Psi=0,$	$i\left[\left(\frac{\theta_{\mathrm{TL}}}{\ell}-1\right)R_{\alpha}^{\star}+\left(\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right)T_{\alpha}+\frac{m^2}{2}\eta_{\alpha}\right]\wedge\gamma^{\alpha}\Psi=0,$	$i\left[\left(\frac{\theta_{\mathrm{TL}}}{\ell}-1\right)R_{\alpha}^{\star}+\left(\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right)T_{\alpha}+\frac{m^2}{2}\eta_{\alpha}\right]\wedge\gamma^{\alpha}\Psi=0,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__E00731 (10.10.11)	231	■1.000 ● W +0	$\begin{aligned} & i\left[\left(\frac{\theta_{\mathrm{TL}}}{\ell}-1\right)R_{\alpha}^{\star}+\left(\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right)T_{\alpha}+\frac{m^2}{2}\eta_{\alpha}\right]\wedge\gamma^{\alpha}\Psi=0, \\ & +\tau_{\alpha}^{\star}\wedge\gamma^{\alpha}\Psi\simeq 0 \end{aligned}$	$i\left[\left(\theta_{\mathrm{L}}+\frac{\theta_{\mathrm{TL}}}{\ell}-1\right)R_{\alpha}^{\star}+\left((-1)^s\frac{\theta_{\mathrm{TL}}}{\ell}+\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right)T_{\alpha}+\frac{1}{2}\left(\frac{\theta_{\mathrm{T}}}{\ell^2}+m^2\right)\eta_{\alpha}+\tau_{\alpha}^{\star}\right]\wedge\gamma^{\alpha}\Psi\simeq 0$	$i\left[\left(\theta_{\mathrm{L}}+\frac{\theta_{\mathrm{TL}}}{\ell}-1\right)R_{\alpha}^{\star}+\left((-1)^s\frac{\theta_{\mathrm{TL}}}{\ell}+\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right)T_{\alpha}+\frac{1}{2}\left(\frac{\theta_{\mathrm{T}}}{\ell^2}+m^2\right)\eta_{\alpha}+\tau_{\alpha}^{\star}\right]\wedge\gamma^{\alpha}\Psi\simeq 0,$ (10.10)
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0732 (10.10.12)	232	■ 1.000 ● W +2	$\theta_{\mathrm{T}} \simeq m^2 \ell^2, \theta_{\mathrm{TL}} \simeq \frac{(-1)^s}{2} m(2m+1) \ell, \theta_{\mathrm{L}} \simeq 1 - \frac{\theta_{\mathrm{TL}}}{\ell} = 1 - \frac{(-1)^s}{2} m(2m+1)$	$\theta_{\mathrm{T}} \simeq -m^2 \ell^2, \theta_{\mathrm{TL}} \simeq \frac{(-1)^s}{2} m(2m+1) \ell, \theta_{\mathrm{L}} \simeq 1 - \frac{\theta_{\mathrm{TL}}}{\ell} = 1 - \frac{(-1)^s}{2} m(2m+1)$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0733 (10.10.13)	232	■ 1.000 ● N +0	$\vartheta_{\alpha} = (-1)^s \ell \Gamma_{\alpha}^{\star} + \bar{\sigma} \gamma_{\alpha} \Psi.$	$\vartheta_{\alpha} = (-1)^s \ell \Gamma_{\alpha}^{\star} + \bar{\sigma} \gamma_{\alpha} \Psi.$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0735 (11.1.2)	238	■ 1.000 ● N +0	$\gamma_{\alpha} \gamma_{\beta} + \gamma_{\beta} \gamma_{\alpha} = 2 \mathbf{o}_{\alpha\beta} \mathbb{1}$	$\gamma_{\alpha} \gamma_{\beta} + \gamma_{\beta} \gamma_{\alpha} = 2 \mathbf{o}_{\alpha\beta} \mathbb{1}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0738 (11.1.5)	238	■ 1.000 ● N +0	$\mathrm{X} := \imath x^{\alpha} \gamma_{\alpha} \longleftrightarrow x^{\alpha} = -\frac{l}{N} \mathrm{Tr}(\gamma^{\alpha} X).$	$\mathrm{X} := \imath x^{\alpha} \gamma_{\alpha} \longleftrightarrow x^{\alpha} = -\frac{l}{N} \mathrm{Tr}(\gamma^{\alpha} X).$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0739 (11.1.6)	238	■ 1.000 ● K +0	$\operatorname{det} \mathrm{X} = -\mathrm{o}_{\alpha\beta} x^{\alpha} x^{\beta},$	$\det \mathrm{X} = -\mathbf{o}_{\alpha\beta} x^{\alpha} x^{\beta},$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0741 (11.1.8)	238	■ 1.000 ● N +0	$\mathrm{X} - \mathrm{X}' = SXS^{-1}, \quad S \in \widetilde{SO}_{\circ}(s, n-s).$	$\mathrm{X} - \mathrm{X}' = SXS^{-1}, \quad S \in \widetilde{SO}_{\circ}(s, n-s).$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00743 (11.1.10)	239	1.000 N +0	$\sigma^{\alpha\beta}:=\frac{i}{4}\left[\gamma^{\alpha},\gamma^{\beta}\right],\quad\Lambda\left(\sigma^{\alpha\beta}\right)=L^{\alpha\beta},$	$\sigma^{\alpha\beta}:=\frac{i}{4}\left[\gamma^{\alpha},\gamma^{\beta}\right],\quad\Lambda\left(\sigma^{\alpha\beta}\right)=L^{\alpha\beta},$	$\sigma^{\alpha\beta}:=\frac{i}{4}[\gamma^{\alpha},\gamma^{\beta}],\quad\Lambda(\sigma^{\alpha\beta})=L^{\alpha\beta},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00744 (11.1.11)	239	1.000 N +0	$L_{\circ}^g(M):=P\left(M,S_{0_{\circ}}(s,n-s),\pi,\delta\right)$	$L_{\circ}^g(M):=P\left(M,S_{O_{\circ}}(s,n-s),\pi,\delta\right)$	$L_{\circ}^g(M):=P(M,S_{O_{\circ}}(s,n-s),\pi,\delta)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00745 (11.1.12)	239	1.000 N +0	$\widetilde{L}(M):=P\left(M,\widetilde{S}_{0_{\circ}}(s,n-2),\widetilde{\pi},\widetilde{\delta}\right)\ni\widetilde{p}$	$\widetilde{L}(M):=P\left(M,\widetilde{S}\widetilde{O}_{\circ}(s,n-2),\widetilde{\pi},\widetilde{\delta}\right)\ni\tilde{p}$	$\widetilde{L}(M):=P(M,\widetilde{S}\widetilde{O}_{\circ}(s,n-2),\tilde{\pi},\tilde{\delta})\ni\tilde{p}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00746 (11.1.13)	239	1.000 N +0	$f(\tilde{p}S)=f(\tilde{p})\Lambda(S),S\in\widetilde{S}_{0_{\circ}}(s,n-2).$	$f(\tilde{p}S)=f(\tilde{p})\Lambda(S),S\in\widetilde{S}\widetilde{O}_{\circ}(s,n-2).$	$f(\tilde{p}S)=f(\tilde{p})\Lambda(S),S\in\widetilde{S}\widetilde{O}_{\circ}(s,n-2).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00748 (11.1.15)	240	1.000 N +0	$\tilde{L}\left(M^4\right)=M^4\times S\,L(2,\mathbb{C})$	$\tilde{L}\left(M^4\right)=M^4\times SL(2,\mathbb{C})$	$\tilde{L}(M^4)=M^4\times SL(2,\mathbb{C})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ E00749 (11.1.16)	241	1.000 N +0	$V^{\bar{s}}:=V\left(M,\mathbb{C}^N,\rho^{\bar{s}}\left(\widetilde{S}\widetilde{O}_{\circ}(s,n-s)\right)\subset GL(N,\mathbb{C}),\tilde{L}(M)\right)$	$V^{\bar{s}}:=V\left(M,\mathbb{C}^N,\rho^{\bar{s}}\left(\widetilde{S}\widetilde{O}_{\circ}(s,n-s)\right)\subset GL(N,\mathbb{C}),\tilde{L}(M)\right)$	$V^{\bar{s}}:=V(M,\mathbb{C}^N,\rho^{\bar{s}}(\widetilde{S}\widetilde{O}_{\circ}(s,n-s))\subset GL(N,\mathbb{C}),\tilde{L}(M))$
Conf. MathPix confidence: 1.000 0.5-0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0751 (11.1.18)	241	■ 1.000 ● K +0	$D^{\frac{1}{2}}\{\frac{1}{2}\}:=D^{\left(-\frac{1}{2},\frac{3}{2},\frac{3}{2}\right)}\oplus D^{\left(\frac{1}{2},\frac{3}{2},\frac{3}{2}\right)}\left(\widetilde{SO}_o(1,3)\right)$	$D^{\frac{1}{2}}:=D^{\left(-\frac{1}{2},\frac{3}{2},\frac{3}{2}\right)}\oplus D^{\left(\frac{1}{2},\frac{3}{2},\frac{3}{2}\right)}\left(\widetilde{SO}_o(1,3)\right)$	$D^{\frac{1}{2}}:=D^{\left(-\frac{1}{2},\frac{3}{2},\frac{3}{2}\right)}\oplus D^{\left(\frac{1}{2},\frac{3}{2},\frac{3}{2}\right)}\left(\widetilde{SO}_o(1,3)\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0754 (11.1.21)	241	■ 1.000 ● N +0	$\psi \rightarrow^{S^{-1}} \psi := S^{-1} \psi$	$\psi \rightarrow^{S^{-1}} \psi := S^{-1} \psi$	$\psi \rightarrow^{S^{-1}} \psi := S^{-1} \psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0755 (11.1.22)	242	■ 1.000 ● W +0	$\begin{gathered} S(p) \in \mathscr{G}_S \approx C^\infty\left(\tilde{L}(M) \times_{\text{Ad}} \widetilde{SO}_o(s, n-s)\right) \\ \Lambda(S_{(p)}) = G(p) \in \mathscr{G}_p \end{gathered}$	$S(p) \in \mathscr{G}_S \approx C^\infty\left(\tilde{L}(M) \times_{\text{Ad}} \widetilde{SO}_o(s, n-s)\right) \\ \Lambda(S_{(p)}) = G(p) \in \mathscr{G}_p$	$S(p) \in \mathscr{G}_S \approx C^\infty(\tilde{L}(M) \times_{\text{Ad}} \widetilde{SO}_o(s, n-s)) \\ \Lambda(S_{(p)}) = G(p) \in \mathscr{G}_p$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0757 (11.1.24)	242	■ 1.000 ● N +0	$S^{-1}(p) \gamma_\alpha S(p) = \gamma_\beta G_\alpha{}^\beta(p), \quad [G_\alpha{}^\beta(p)] \in \mathscr{G}_p.$	$S^{-1}(p) \gamma_\alpha S(p) = \gamma_\beta G_\alpha{}^\beta(p), \quad [G_\alpha{}^\beta(p)] \in \mathscr{G}_p.$	$S^{-1}(p) \gamma_\alpha S(p) = \gamma_\beta G_\alpha{}^\beta(p), \quad [G_\alpha{}^\beta(p)] \in \mathscr{G}_p.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0762 (11.1.29)	243	■ 1.000 ● N +0	$\bar{\psi} := \psi^+ \gamma^0,$	$\bar{\psi} := \psi^+ \gamma^0,$	$\bar{\psi} := \psi^+ \gamma^0,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0763 (11.1.30)	243	■ 1.000 ● W -2	$\bar{\sigma}^{\alpha\beta} := \gamma^0 \sigma^{\alpha\beta} + \gamma^0 \sigma^{\alpha\beta}, \quad \bar{\gamma}^\alpha := \gamma^0 \gamma^\alpha + \gamma^0 \gamma^\alpha.$	$\bar{\sigma}^{\alpha\beta} := \gamma^0 \sigma^{\alpha\beta} + \gamma^0 \sigma^{\alpha\beta}, \quad \bar{\gamma}^\alpha := \gamma^0 \gamma^\alpha + \gamma^0 \gamma^\alpha.$	$\bar{\sigma}^{\alpha\beta} := \gamma^0 \sigma^{\alpha\beta} + \gamma^0 \sigma^{\alpha\beta}, \quad \bar{\gamma}^\alpha := \gamma^0 \gamma^\alpha + \gamma^0 \gamma^\alpha.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0764 (11.1.31)	243	■ 1.000 ● N +0	$D \psi = d \psi + i \tilde{\omega} \psi,$	$D \psi = d \psi + i \tilde{\omega} \psi,$	$D \psi = d \psi + i \tilde{\omega} \psi,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0765 (11.1.32)	243	■ 1.000 ● N +0	$\Lambda(i \tilde{\omega}) = \omega^g$	$\Lambda(i \tilde{\omega}) = \omega^g$	$\Lambda(i \tilde{\omega}) = \omega^g$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0771 (11.1.38)	244	■ 1.000 ● K +0	$\gamma \longrightarrow S^{-1} \gamma = S^{-1} \gamma S$	$\gamma \longrightarrow S^{-1} \gamma = S^{-1} \gamma S$	$\gamma \longrightarrow S^{-1} \gamma = S^{-1} \gamma S$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0775 (11.2.4)	246	■ 1.000 ● K +0	$\overline{\widetilde{\omega}} = \gamma^0 \widetilde{\omega}^+ \gamma^0 = \widetilde{\omega}$	$\overline{\widetilde{\omega}} = \gamma^0 \widetilde{\omega}^+ \gamma^0 = \widetilde{\omega}$	$\overline{\widetilde{\omega}} = \gamma^0 \widetilde{\omega}^+ \gamma^0 = \widetilde{\omega}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0777 (11.2.6)	246	■ 1.000 ● N +0	$\tau_s = i \left(\bar{\psi} \gamma \psi \right).$	$\tau_s = i \left(\bar{\psi}^* \gamma \psi \right).$	$\tau_s = i \left(\bar{\psi}^* \gamma \psi \right).$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0779 (11.2.7)	247	■ 1.000 ● K +0	$d\,\tau_s=0$	$d\tau_s=0$	$d\tau_s=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0781 (11.2.9)	247	■ 1.000 ● N +1	$(\psi,\chi)=i\int_{H^{n-1}}\psi^+\chi d^{n-1}x$	$(\psi,\chi)=i\int_{H^{n-1}}\psi^+\chi d^{n-1}x$	$(\psi,\chi)=i\int_{H^{n-1}}\psi^+\chi d^{n-1}x.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0782 (11.3.1)	247	■ 1.000 ● N +0	$\widetilde{\omega}=\widetilde{\omega}^{\{\}}-\widetilde{\omega}^{\{K\}},\quad \Lambda(i\widetilde{K})=K$	$\widetilde{\omega}=\widetilde{\omega}^{\{}}-\widetilde{K},\quad \Lambda(i\widetilde{K})=K$	$\widetilde{\omega}=\widetilde{\omega}^{()}-\widetilde{K},\quad \Lambda(i\widetilde{K})=K$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0785 (11.3.4)	248	■ 1.000 ● N +0	$[\Theta,\vartheta]=\ell^{*2}\tau_s\;.$	$[\Theta,\vartheta]=\ell^{*2}\tau_s.$	$[\Theta,\vartheta]=\ell^{*2}\tau_s.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0790 (11.4.2)	252	■ 1.000 ● W +0	$\begin{aligned} &\left(\overline{\psi}\gamma^5\gamma_\mu\psi\right)\left(\overline{\psi}\gamma^5\gamma^\mu\psi\right)=\left(\overline{\psi}\gamma_\mu\psi\right)\left(\overline{\psi}\gamma^\mu\psi\right)\\ &=\left(\overline{\psi}\psi\right)^2-\left(\overline{\psi}\gamma^5\psi\right)^2. \end{aligned}$	$\left(\overline{\psi}i\gamma^5\gamma_\mu\psi\right)\left(\overline{\psi}i\gamma^5\gamma^\mu\psi\right)=\left(\overline{\psi}\gamma_\mu\psi\right)\left(\overline{\psi}\gamma^\mu\psi\right)\\ =\left(\overline{\psi}\psi\right)^2-\left(\overline{\psi}\gamma^5\psi\right)^2.$	$\left(\overline{\psi}i\gamma^5\gamma_\mu\psi\right)\left(\overline{\psi}i\gamma^5\gamma^\mu\psi\right)=\left(\overline{\psi}\gamma_\mu\psi\right)\left(\overline{\psi}\gamma^\mu\psi\right)\\ =\left(\overline{\psi}\psi\right)^2-\left(\overline{\psi}\gamma^5\psi\right)^2.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0793 (11.4.5)	253	■1.000 ● N +0	$\rho:=\frac{\sqrt{2\pi}}{\ell^*}\ell^*r,$	$\rho := \frac{\sqrt{2\pi}}{\ell^*}r,$	$\rho := \frac{\sqrt{2\pi}}{\ell^*}r,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0794 (11.4.6)	253	■1.000 ● N +0	$\bar{g}_{kl} \longleftarrow g_{kl} = e^{\mu} \bar{g}_{kl}.$	$\bar{g}_{kl} \longleftarrow g_{kl} = e^{\mu} \bar{g}_{kl}.$	$\bar{g}_{kl} \longleftarrow g_{kl} = e^{\mu} \bar{g}_{kl}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0795 (11.4.7)	253	■1.000 ● N -2	$\bar{E}_{\cdot k}^{\alpha} \longleftarrow E_{\cdot k}^{\alpha} = e^{\mu/2} \bar{E}_{\cdot k}^{\alpha}$	$\bar{E}_{\cdot k}^{\alpha} \longleftarrow E_{\cdot k}^{\alpha} = e^{\mu/2} \bar{E}_{\cdot k}^{\alpha}$	$\bar{E}_{\cdot k}^{\alpha} \longleftarrow E_{\cdot k}^{\alpha} = e^{\mu/2} \bar{E}_{\cdot k}^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0796 (11.4.8)	253	■1.000 ● N +0	$\overline{\left\{ \begin{array}{c} \alpha \\ kl \end{array} \right\}} = \left\{ \begin{array}{c} \alpha \\ kl \end{array} \right\} + \frac{1}{2} (\delta_k^{\alpha} \partial_l + \delta_l^{\alpha} \partial_k - g_{kl} \partial^{\alpha}) \mu$	$\left\{ \begin{array}{c} \alpha \\ kl \end{array} \right\} = \overline{\left\{ \begin{array}{c} \alpha \\ kl \end{array} \right\}} + \frac{1}{2} (\delta_k^{\alpha} \partial_l + \delta_l^{\alpha} \partial_k - g_{kl} \partial^{\alpha}) \mu$	$\left\{ \begin{array}{c} \alpha \\ kl \end{array} \right\} = \overline{\left\{ \begin{array}{c} \alpha \\ kl \end{array} \right\}} + \frac{1}{2} (\delta_k^{\alpha} \partial_l + \delta_l^{\alpha} \partial_k - g_{kl} \partial^{\alpha}) \mu$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0797 (11.4.9)	253	■1.000 ● N +0	$\Gamma_k = \bar{\Gamma}_k - \frac{1}{4} \delta_k^{[\alpha} \partial^{\beta]} \mu \sigma_{\alpha\beta}.$	$\Gamma_k = \bar{\Gamma}_k - \frac{1}{4} \delta_k^{[\alpha} \partial^{\beta]} \mu \sigma_{\alpha\beta}.$	$\Gamma_k = \bar{\Gamma}_k - \frac{1}{4} \delta_k^{[\alpha} \partial^{\beta]} \mu \sigma_{\alpha\beta}.$
Conf. MathPix confidence: ■≥0.9 ■0.5–0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0801 (11.4.13)	254	■ 1.000 ● N +1	$\begin{aligned} & \bar{\gamma}_0 = \frac{i}{4} e^{(\nu-\mu)/2} \partial_r (\nu - \mu) \gamma_0 \gamma_1 \\ & \bar{\gamma}_1 = 0, \quad \bar{\gamma}_2 = \frac{i}{2} \gamma_2 \gamma_1 \\ & \bar{\gamma}_3 = \frac{i}{2} (\sin \vartheta \gamma_3 \gamma_1 + \cos \vartheta \gamma_3 \gamma_2) \end{aligned}$	$\bar{\gamma}_0 = \frac{i}{4} e^{(\nu-\mu)/2} \partial_r (\nu - \mu) \gamma_0 \gamma_1$ $\bar{\gamma}_1 = 0, \quad \bar{\gamma}_2 = \frac{i}{2} \gamma_2 \gamma_1$ $\bar{\gamma}_3 = \frac{i}{2} (\sin \vartheta \gamma_3 \gamma_1 + \cos \vartheta \gamma_3 \gamma_2)$	$\bar{\gamma}_0 = \frac{i}{4} e^{(v-\mu)/2} \partial_r (v - \mu) \gamma_0 \gamma_1$ $\bar{\gamma}_1 = 0, \quad \bar{\gamma}_2 = \frac{i}{2} \gamma_2 \gamma_1$ $\bar{\gamma}_3 = \frac{i}{2} (\sin \vartheta \gamma_3 \gamma_1 + \cos \vartheta \gamma_3 \gamma_2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0804 (11.4.16)	255	■ 1.000 ● N +0	$\chi_{\kappa}^m = \sum_{\bar{m}=\pm 1/2} C \left(l_{\frac{1}{2}} j; m - \bar{m}, \bar{m} \right) Y_l^{m-\bar{m}}(\vartheta, \phi) \chi^{\bar{m}}.$	$\chi_{\kappa}^m = \sum_{\bar{m}=\pm 1/2} C \left(l_{\frac{1}{2}} j; m - \bar{m}, \bar{m} \right) Y_l^{m-\bar{m}}(\vartheta, \phi) \chi^{\bar{m}}.$	$\chi_{\kappa}^m = \sum_{\bar{m}=\pm 1/2} C \left(l_{\frac{1}{2}} j; m - \bar{m}, \bar{m} \right) Y_l^{m-\bar{m}}(\vartheta, \phi) \chi^{\bar{m}}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0808 (11.4.20)	255	■ 1.000 ● W +0	$\frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } \chi_{\kappa}^m = -\chi_{-\kappa}^m$	$\frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } \chi_{\kappa}^m = -\chi_{-\kappa}^m$	$\frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } \chi_{\kappa}^m = -\chi_{-\kappa}^m$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0814 (11.4.27)	257	■ 1.000 ● W +1	$\beta = \frac{M^*}{2m}, \quad M^* := \frac{\sqrt{8\pi} \hbar}{c \ell^*}.$	$\beta = \frac{M^*}{2m}, \quad M^* := \frac{\sqrt{8\pi} \hbar}{c \ell^*}.$	$\beta = \frac{M^*}{2m}, \quad M^* := \frac{\sqrt{8\pi} \hbar}{c \ell^*}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0816 (11.4.30)	257	■ 1.000 ● K +0	$F'' + \frac{2}{\rho} F' - \frac{2}{\rho^2} F \simeq \frac{1 - \omega^2}{\beta^2} F$	$F'' + \frac{2}{\rho} F' - \frac{2}{\rho^2} F \simeq \frac{1 - \omega^2}{\beta^2} F$	$F'' + \frac{2}{\rho} F' - \frac{2}{\rho^2} F \simeq \frac{1 - \omega^2}{\beta^2} F$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0817 (11.4.31)	257	■ 1.000 ● N +0	$F \sim C_{\infty} \left(\frac{1}{\bar{\rho}} + \frac{1}{\bar{\rho}^2} \right) e^{-\bar{\rho}}$	$F \sim C_{\infty} \left(\frac{1}{\bar{\rho}} + \frac{1}{\bar{\rho}^2} \right) e^{-\bar{\rho}}$	$F \sim C_{\infty} \left(\frac{1}{\bar{\rho}} + \frac{1}{\bar{\rho}^2} \right) e^{-\bar{\rho}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0818 (11.4.32)	257	■ 1.000 ● N +0	$H \sim C_{\infty} \sqrt{\frac{1+\omega}{1-\omega}} \frac{1}{\bar{\rho}} e^{-\bar{\rho}}$	$H \sim C_{\infty} \sqrt{\frac{1+\omega}{1-\omega}} \frac{1}{\bar{\rho}} e^{-\bar{\rho}}$	$H \sim C_{\infty} \sqrt{\frac{1+\omega}{1-\omega}} \frac{1}{\bar{\rho}} e^{-\bar{\rho}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0819 (11.4.33)	257	■ 1.000 ● N +0	$\bar{\rho} := \frac{1}{\beta} \sqrt{1-\omega^2} \rho$	$\bar{\rho} := \frac{1}{\beta} \sqrt{1-\omega^2} \rho$	$\bar{\rho} := \frac{1}{\beta} \sqrt{1-\omega^2} \rho$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0821 (11.4.35)	258	■ 1.000 ● N +0	$F(\rho) \simeq C_0 f^1(\bar{\rho}) \frac{1}{\cosh \bar{\rho}}$	$F(\rho) \simeq C_0 f^1(\bar{\rho}) \frac{1}{\cosh \bar{\rho}}$	$F(\rho) \simeq C_0 f^1(\bar{\rho}) \frac{1}{\cosh \bar{\rho}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0822 (11.4.36)	258	■ 1.000 ● N +0	$H(\rho) \simeq C_0 \sqrt{\frac{1+\omega}{1-\omega}} f^0(\bar{\rho}) \frac{1}{\cosh \bar{\rho}}$	$H(\rho) \simeq C_0 \sqrt{\frac{1+\omega}{1-\omega}} f^0(\bar{\rho}) \frac{1}{\cosh \bar{\rho}}$	$H(\rho) \simeq C_0 \sqrt{\frac{1+\omega}{1-\omega}} f^0(\bar{\rho}) \frac{1}{\cosh \bar{\rho}}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0823 (11.4.37)	258	■ 1.000 ● K +0	$C_{\{0\}}=(1-\omega)\sqrt{\frac{-2}{\varepsilon(\omega+1)}}$	$C_0 = (1 - \omega) \sqrt{\frac{-2}{\varepsilon(\omega + 1)}}$	$C_0 = (1 - \omega) \sqrt{\frac{-2}{\varepsilon(\omega + 1)}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0826 (11.4.40)	259	■ 1.000 ● N +0	$m_{\{s\}}=\frac{1}{c}\int_{H^3}\Sigma_{\{\mathrm{D}\}}^{\{0\}}$	$m_s = \frac{1}{c} \int_{H^3} \Sigma_{\mathrm{D}}^0$	$m_s = \frac{1}{c} \int_{H^3} \Sigma_{\mathrm{D}}^0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0827 (11.5.1)	262	■ 1.000 ● N +1	$Q:=i\int_{H^3}\bar{\psi}^*\gamma\psi=\sum_{k=0,1,2,\ldots}k\int_{H^3}\bar{\psi}^*\gamma\psi=\sum_{k=0,1,2,\ldots}k\int_{H^3}\bar{\psi}^*\gamma\psi$	$Q := ie \int_{H^3} \bar{\psi}^* \gamma \psi = ke, \quad k = 0, 1, 2, \dots$	$Q := ie \int_{H^3} \bar{\psi}^* \gamma \psi = ke, \quad k = 0, 1, 2, \dots$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0828 (11.5.2)	263	■ 1.000 ● N +0	$i\int_{H^3}\bar{\psi}^*\gamma\psi=\sum_{k=0,1,2,\ldots}k\int_{H^3}\bar{\psi}^*\gamma\psi$	$i\gamma\wedge^*d\psi(x)+e^2\int_{y\in M}\bar{\psi}(y)\gamma\psi(y)D_F(y-x)\wedge^*\gamma\psi(x)=m\psi(x)\eta$	$i\gamma\wedge^*d\psi(x)+e^2\int_{y\in M}\bar{\psi}(y)\gamma\psi(y)D_F(y-x)\wedge^*\gamma\psi(x)=m\psi(x)\eta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0830 (12.1.2)	269	■ 1.000 ● N +0	$\langle dj_5(x) \rangle = 2im\langle P \rangle - (1/96\pi^2) F \wedge F$	$\langle dj_5(x) \rangle = 2im\langle P \rangle - (1/96\pi^2) F \wedge F$	$\langle dj_5(x) \rangle = 2im\langle P \rangle - (1/96\pi^2) F \wedge F$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0838 (12.2.6)	271	■ 1.000 ● N +0	$d\,j_{\{5\}=2\,i\,m\,P=2\,i\,m\,\bar{\psi}\,\gamma_5\,\psi\,\eta}$	$dj_5 = 2imP = 2im\bar{\psi}\gamma_5\psi\eta$	$dj_5 = 2imP = 2im\bar{\psi}\gamma_5\psi\eta$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0843 (12.2.11)	272	■ 1.000 ● N +0	$L_{\{a\,\psi\,\psi\}}=\frac{1}{2}\,d\,\theta\,\wedge\,j_{\{5\}}=\frac{1}{2}\,f_{\{a\}}\,d\,a\,\wedge\,\bar{\psi}\wedge\gamma_5\psi,\,$	$L_{a\psi\psi} = \frac{1}{2}d\theta \wedge j_5 = \frac{1}{2f_a}da \wedge \bar{\psi}^*\gamma\gamma_5\psi,$	$L_{a\psi\psi} = \frac{1}{2}d\theta \wedge j_5 = \frac{1}{2f_a}da \wedge \bar{\psi}^*\gamma\gamma_5\psi,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0846 (12.2.14)	273	■ 1.000 ● N -1	$d\,j_{\{5\}}\,\cong\,4\,d\,C_{\{\mathrm{TT}\}}=\frac{2}{\ell^2}\,\left(T^{\alpha}\wedge T_{\alpha}+R_{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}\right)$	$dj_5 \cong 4dC_{\mathrm{TT}} = \frac{2}{\ell^2} \left(T^{\alpha} \wedge T_{\alpha} + R_{\alpha\beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta} \right)$	$dj_5 \cong 4dC_{\mathrm{TT}} = \frac{2}{\ell^2} \left(T^{\alpha} \wedge T_{\alpha} + R_{\alpha\beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta} \right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0849 (12.3.2)	275	■ 1.000 ● N +0	$K\left(t,\,x,\,\mathrm{not}\,D^{\{2\}}\right)=(4\,\pi)^{-n\,/\,2}\,\sum_{k=0}^{\infty}\,t^{\left(k-n\right)/2}\,K_k\left(x,\,\mathrm{not}\,D^{\{2\}}\right)$	$K\left(t,x,\,D^2\right)=\left(4\pi\right)^{-n/2}\sum_{k=0}^{\infty}t^{\left(k-n\right)/2}K_k\left(x,\,D^2\right)$	$K(t,x,\,D^2)=(4\pi)^{-n/2}\sum_{k=0}^{\infty}t^{(k-n)/2}K_k(x,\,D^2)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0852 (12.3.5)	276	■ 1.000 ● N -1	$-K_{\{2\}}\,/\,t=\left(2\,\ell^2\,/\,t\right)\,d\,C_{\{\mathrm{TT}\}}\,\cong\left(\ell^2\,/\,2\,t\right)\,d\,j_{\{5\}}=\left(i\,m\,\ell^2\,/\,t\right)\,\bar{\psi}\,\gamma_5\,\psi\,\sim\ell^2\,M^2\,m^4\,\rightarrow\,0\,.$	$-K_2/t = (2\ell^2/t)\,dC_{\mathrm{TT}} \cong (\ell^2/2t)\,dj_5 = (im\ell^2/t)\,\bar{\psi}\gamma_5\psi \sim \ell^2M^2m^4 \rightarrow 0.$	$-K_2/t = (2\ell^2/t)dC_{\mathrm{TT}} \cong (\ell^2/2t) dj_5 = (im\ell^2/t)\bar{\psi}\gamma_5\psi \sim \ell^2M^2m^4 \rightarrow 0.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0856 (12.4.3)	277	■ 1.000 ● N +0	$\mathrm{G}^A \cong 0, \quad A=1,2,3$	$\mathcal{G}^A \cong 0, \quad A = 1, 2, 3$	$\mathcal{G}^A \cong 0, \quad A = 1, 2, 3$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0858 (12.4.5)	278	■ 1.000 ● K +0	$\gamma:=\gamma_{\alpha} \vartheta^{\alpha}, \quad \gamma^*=\gamma^{\alpha} \eta_{\alpha} .$ $\gamma=\gamma^{\alpha} \eta_{\alpha} .$	$\gamma:=\gamma_{\alpha} \vartheta^{\alpha}, \quad \gamma^*=\gamma^{\alpha} \eta_{\alpha} .$	$\gamma:=\gamma_{\alpha} \vartheta^{\alpha}, \quad \gamma^*=\gamma^{\alpha} \eta_{\alpha} .$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0859 (12.4.6)	278	■ 1.000 ● N +0	$\Theta:=D \gamma=T^{\alpha} \gamma_{\alpha}, \quad \Omega:=d \Gamma+\Gamma \wedge \Gamma=\frac{i}{4} R^{\alpha \beta} \sigma_{\alpha \beta}$ $\Omega:=d \Gamma+\Gamma \wedge \Gamma=\frac{i}{4} R^{\alpha \beta} \sigma_{\alpha \beta}$	$\Theta:=D \gamma=T^{\alpha} \gamma_{\alpha}, \quad \Omega:=d \Gamma+\Gamma \wedge \Gamma=\frac{i}{4} R^{\alpha \beta} \sigma_{\alpha \beta}$	$\Theta:=D \gamma=T^{\alpha} \gamma_{\alpha}, \quad \Omega:=d \Gamma+\Gamma \wedge \Gamma=\frac{i}{4} R^{\alpha \beta} \sigma_{\alpha \beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0860 (12.4.7)	278	■ 1.000 ● N +1	$C_{\mathrm{RR}}:=\operatorname{Tr}\left(\Gamma \wedge \Omega-\frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma\right) .$	$C_{\mathrm{RR}}:=\operatorname{Tr}\left(\Gamma \wedge \Omega-\frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma\right) .$	$C_{\mathrm{RR}}:=\operatorname{Tr}\left(\Gamma \wedge \Omega-\frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma\right) .$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0862 (12.4.9)	279	■ 1.000 ● N +1	$C_{\mathrm{TT}}:=\frac{1}{8 \ell^2} \operatorname{Tr}\left(\gamma \wedge \Theta\right)=\frac{1}{2 \ell^2} \vartheta^{\alpha} \wedge T_{\alpha} .$	$C_{\mathrm{TT}}:=\frac{1}{8 \ell^2} \operatorname{Tr}\left(\gamma \wedge \Theta\right)=\frac{1}{2 \ell^2} \vartheta^{\alpha} \wedge T_{\alpha} .$	$C_{\mathrm{TT}}:=\frac{1}{8 \ell^2} \operatorname{Tr}\left(\gamma \wedge \Theta\right)=\frac{1}{2 \ell^2} \vartheta^{\alpha} \wedge T_{\alpha} .$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0863 (12.4.10)	279	■ 1.000 ● C +5	$d\,C_{\mathrm{TT}}=\frac{1}{8\ell^2}\mathrm{Tr}(\Theta\wedge\Theta-4i\Omega\wedge\sigma)=\frac{1}{2\ell^2}\left(T^\alpha\wedge T_\alpha+R_{\alpha\beta}\wedge\vartheta^\alpha\wedge\vartheta^\beta\right).$	$dC_{\mathrm{TT}}=\frac{1}{8\ell^2}\mathrm{Tr}(\Theta\wedge\Theta-4i\Omega\wedge\sigma)=\frac{1}{2\ell^2}\left(T^\alpha\wedge T_\alpha+R_{\alpha\beta}\wedge\vartheta^\alpha\wedge\vartheta^\beta\right).$	$dC_{\mathrm{TT}}=\frac{1}{8\ell^2}\mathrm{Tr}\left(\Theta\wedge\Theta-4i\Omega\wedge\sigma\right)=\frac{1}{2\ell^2}\left(T^\alpha\wedge T_\alpha+R_{\alpha\beta}\wedge\vartheta^\alpha\wedge\vartheta^\beta\right).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0864 (12.4.11)	279	■ 1.000 ● N -2	$\hat{C}_{\mathrm{C}}=\mathrm{C}_{\mathrm{RR}}-2\,C_{\mathrm{TT}}$	$\hat{C}_{\mathrm{RR}}=C_{\mathrm{RR}}-2C_{\mathrm{TT}}$	$\hat{C}_{\mathrm{RR}}=C_{\mathrm{RR}}-2C_{\mathrm{TT}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0865 (13.2.1)	283	■ 1.000 ● N -1	$L_{\mathrm{BF}}=-B\wedge F=-B\wedge dA.$	$L_{\mathrm{BF}}=-B\wedge F=-B\wedge dA.$	$L_{\mathrm{BF}}=-B\wedge F=-B\wedge dA.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0866 (13.2.2)	283	■ 1.000 ● N +1	$\tilde{L}_{\mathrm{BF}}:=-B\wedge dA+\frac{1}{2}B\wedge B\cong -B\wedge dA+dC$	$\tilde{L}_{\mathrm{BF}}:=-B\wedge dA+\frac{1}{2}B\wedge B\cong -B\wedge dA+dC$	$\tilde{L}_{\mathrm{BF}}:=-B\wedge dA+\frac{1}{2}B\wedge B\cong -B\wedge dA+dC.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0868 (13.2.4)	283	■ 1.000 ● N +0	$C=\frac{1}{2}A\wedge F,\quad dC=\frac{1}{2}F\wedge F,$	$C=\frac{1}{2}A\wedge F,\quad dC=\frac{1}{2}F\wedge F,$	$C=\frac{1}{2}A\wedge F,\quad dC=\frac{1}{2}F\wedge F,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0869 (13.2.5)	283	■ 1.000 ● K +0	$A \rightarrow A^{\prime}=A+d \theta(x),$	$A \rightarrow A^{\prime}=A+d \theta(x),$	$A \rightarrow A^{\prime}=A+d \theta(x),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0870 (13.2.6)	284	■ 1.000 ● N +0	$A \rightarrow \widetilde{A}=A+\sigma, \quad B \rightarrow \widetilde{B}=B+d \sigma,$	$A \rightarrow \widetilde{A}=A+\sigma, \quad B \rightarrow \widetilde{B}=B+d \sigma,$	$A \rightarrow \widetilde{A}=A+\sigma, \quad B \rightarrow \widetilde{B}=B+d \sigma,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0871 (13.2.7)	284	■ 1.000 ● K +0	$L_{\mathrm{Max}}=-B \wedge d A+\frac{1}{2} B \wedge \mathrm{B} \wedge L_{\mathrm{matter}},$	$L_{\mathrm{Max}}=-B \wedge d A+\frac{1}{2} B \wedge \mathrm{B}+L_{\mathrm{matter}},$	$L_{\mathrm{Max}}=-B \wedge d A+\frac{1}{2} B \wedge \mathrm{B}+L_{\mathrm{matter}},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0872 (13.2.8)	284	■ 1.000 ● N +1	$-d B=d^{\ast} F \cong j,$	$-d B=d^{\ast} F \cong j,$	$-d B=d^{\ast} F \cong j,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ0875 (13.3.3)	284	■ 1.000 ● N +0	$\left[L^{\alpha}{}_{\beta}, L^{\gamma}{}_{\delta}\right]=\delta_{\delta}^{\alpha} L^{\gamma}{}_{\beta}-\delta_{\beta}^{\gamma} L^{\alpha}{}_{\delta},$	$\left[L^{\alpha}{}_{\beta}, L^{\gamma}{}_{\delta}\right]=\delta_{\delta}^{\alpha} L^{\gamma}{}_{\beta}-\delta_{\beta}^{\gamma} L^{\alpha}{}_{\delta},$	$\left[L^{\alpha}{}_{\beta}, L^{\gamma}{}_{\delta}\right]=\delta_{\delta}^{\alpha} L^{\gamma}{}_{\beta}-\delta_{\beta}^{\gamma} L^{\alpha}{}_{\delta},$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5-0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0880 (13.4.1)	285	■ 1.000 ● N +0	$A(x):=\left\{\begin{array}{cc} \Lambda(x) & \& \tau(x) \\ \tau_*(x) & 1 \end{array}\right\},$	$A(x) := \left\{ \begin{array}{cc} \Lambda(x) & \tau(x) \\ \tau_*(x) & 1 \end{array} \right\},$	$A(x) := \left\{ \begin{array}{cc} \Lambda(x) & \tau(x) \\ \tau_*(x) & 1 \end{array} \right\},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0881 (13.4.2)	285	■ 1.000 ● N +0	$\hat{\Gamma}=\left(\begin{array}{cc} \Gamma^{\{L\}} & \& \Gamma^{\{T\}} \\ \Gamma_*^{\{T\}} & 0 \end{array}\right)=\left(\begin{array}{cc} \Gamma_{\alpha}^{\{L\}\beta} L_{\beta}^{\alpha} & \Gamma^{\{T\}\alpha} \ell P_{\alpha} \\ \Gamma_{\beta}^{*(T)} & \ell P_{*}^{\beta} \end{array}\right)$	$\hat{\Gamma} = \left(\begin{array}{cc} \Gamma^{(L)} & \Gamma^{(T)} \\ \Gamma_*^{(T)} & 0 \end{array} \right) = \left(\begin{array}{cc} \Gamma_{\alpha}^{(L)\beta} L_{\beta}^{\alpha} & \Gamma^{(T)\alpha} \ell P_{\alpha} \\ \Gamma_{\beta}^{*(T)} & \ell P_{*}^{\beta} \end{array} \right)$	$\hat{F} = \left(\begin{array}{cc} \Gamma^{(L)} & \Gamma^{(T)} \\ \Gamma_*^{(T)} & 0 \end{array} \right) = \left(\begin{array}{cc} \Gamma_{\alpha}^{(L)\beta} L_{\beta}^{\alpha} & \Gamma^{(T)\alpha} \ell P_{\alpha} \\ \Gamma_{\beta}^{*(T)} & \ell P_{*}^{\beta} \end{array} \right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0882 (13.4.3)	286	■ 1.000 ● N +1	$\hat{\Gamma} \mapsto \hat{\Gamma}' = A^{-1}(x) \hat{\Gamma} A(x) + A^{-1}(x) dA(x).$	$\hat{\Gamma} \xrightarrow{A^{-1}(x)} \hat{\Gamma}' = A^{-1}(x) \hat{\Gamma} A(x) + A^{-1}(x) dA(x).$	$\hat{F} \xrightarrow{A^{-1}(x)} \hat{F}' = A^{-1}(x) \hat{F} A(x) + A^{-1}(x) dA(x).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__EQ0883	286	■ 1.000 ● N +0	$\tau \tau_* = -\Lambda \tau_* \Lambda^{-1} \tau.$	$\tau \tau_* = -\Lambda \tau_* \Lambda^{-1} \tau.$	$\tau \tau_* = -\Lambda \tau_* \Lambda^{-1} \tau.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0884 (13.4.4)	286	■ 1.000 ● W +1	$\hat{R}:=d\hat{\Gamma}+\hat{\Gamma}\wedge\hat{\Gamma}=\left(\begin{array}{cc} R^{\{L\}}+\Gamma^{\{L\}}\wedge\Gamma_*^{\{L\}} & d\Gamma^{\{T\}}+\Gamma^{\{L\}}\wedge\Gamma^{\{T\}} \\ d\Gamma_*^{\{T\}}+\Gamma_*^{\{T\}}\wedge\Gamma^{\{L\}} & 0 \end{array}\right)$	$\hat{R} := d\hat{\Gamma} + \hat{\Gamma} \wedge \hat{\Gamma} = \left(\begin{array}{cc} R^{(L)} + \Gamma^{(L)} \wedge \Gamma_*^{(L)} & d\Gamma^{(T)} + \Gamma^{(L)} \wedge \Gamma^{(T)} \\ d\Gamma_*^{(T)} + \Gamma_*^{(T)} \wedge \Gamma^{(L)} & 0 \end{array} \right)$	$\hat{R} := d\hat{F} + \hat{F} \wedge \hat{F} = \left(\begin{array}{cc} R^{(L)} + \Gamma^{(L)} \wedge \Gamma_*^{(L)} & d\Gamma^{(T)} + \Gamma^{(L)} \wedge \Gamma^{(T)} \\ d\Gamma_*^{(T)} + \Gamma_*^{(T)} \wedge \Gamma^{(L)} & 0 \end{array} \right),$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0885	286	■ 1.000 ● K +0	$R^{\{L\}}:=d\ \Gamma^{\{L\}}+\Gamma^{\{L\}}\ \wedge\ \Gamma^{\{L\}}$	$R^{(L)}:=d\Gamma^{(L)}+\Gamma^{(L)}\wedge\Gamma^{(L)}$	$R^{(L)}:=d\Gamma^{(L)}+\Gamma^{(L)}\wedge\Gamma^{(L)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0886 (13.4.5)	286	■ 1.000 ● N +0	$R^{\{T\}}:=d\ \Gamma^{\{T\}}+\Gamma^{\{L\}}\ \wedge\ \Gamma^{\{T\}}=\left(T^{\beta}-R_{\alpha}^{\beta}\xi^{\alpha}\right)\ell P_{\beta}\ell\!\!\!\ell P_{\{\beta\}}$	$R^{(T)}:=d\Gamma^{(T)}+\Gamma^{(L)}\wedge\Gamma^{(T)}=\left(T^{\beta}-R_{\alpha}^{\beta}\xi^{\alpha}\right)\ell P_{\beta}$	$R^{(T)}:=d\Gamma^{(T)}+\Gamma^{(L)}\wedge\Gamma^{(T)}=\left(T^{\beta}-R_{\alpha}^{\beta}\xi^{\alpha}\right)\ell P_{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0887 (13.4.6)	286	■ 1.000 ● N +1	$\hat{R}\ \xrightarrow{A^{-1}(x)}\ \hat{R}'=A^{-1}(x)\hat{R}A(x)\ .$	$\hat{R}\ \xrightarrow{A^{-1}(x)}\ \hat{R}'=A^{-1}(x)\hat{R}A(x).$	$\hat{R}\ \xrightarrow{A^{-1}(x)}\ \hat{R}'=A^{-1}(x)\hat{R}A(x)\ .$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0888 (13.4.7)	286	■ 1.000 ● N +0	$\Gamma^{\{L\}}\ \xrightarrow{A^{-1}(x)}\ \Gamma^{\{L\}}\prime=\Lambda^{-1}(x)\Gamma^{\{L\}}\Lambda(x)+\Lambda^{-1}(x)d\Lambda(x)$	$\Gamma^{(L)}\ \xrightarrow{A^{-1}(x)}\ \Gamma^{(L)\prime}=\Lambda^{-1}(x)\Gamma^{(L)}\Lambda(x)+\Lambda^{-1}(x)d\Lambda(x)$	$\Gamma^{(L)}\ \xrightarrow{A^{-1}(x)}\ \Gamma^{(L)\prime}=\Lambda^{-1}(x)\ \Gamma^{(L)}\ \Lambda(x)+\Lambda^{-1}(x)d\Lambda(x)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0891 (13.4.10)	287	■ 1.000 ● N +0	$\vartheta:=\Gamma^{\{T\}}+D\xi:=\vartheta^{\alpha}P_{\alpha},\quad \theta:=\Gamma_{*}^{\{T\}}+D\xi_{*}:=\theta_{\beta}P_{*}^{\beta},$	$\vartheta:=\Gamma^{(T)}+D\xi:=\vartheta^{\alpha}P_{\alpha},\quad \theta:=\Gamma_{*}^{\{T\}}+D\xi_{*}:=\theta_{\beta}P_{*}^{\beta},$	$\vartheta:=\Gamma^{(T)}+D\xi:=\vartheta^{\alpha}P_{\alpha},\quad \theta:=\Gamma_{*}^{\{T\}}+D\xi_{*}:=\theta_{\beta}P_{*}^{\beta},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0894 (13.4.13)	287	■ 1.000 ● N +0	$D\ \xi\stackrel{*}{=}0,\quad D\ \xi_{*}\stackrel{*}{=}0,$	$D\xi\stackrel{*}{=}0,\quad D\xi_{*}\stackrel{*}{=}0,$	$D\xi\stackrel{*}{=}0,\quad D\xi_{*}\stackrel{*}{=}0,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0895 (13.4.14)	287	■ 1.000 ● N +1	$\overline{\mathrm{\Gamma}}=\left(\begin{array}{cc}\mathrm{\Gamma}^{\mathrm{(L)}} & \vartheta \\ \theta & 0\end{array}\right),$	$\bar{\Gamma}=\left(\begin{array}{cc}\Gamma^{(L)} & \vartheta \\ \theta & 0\end{array}\right),$	$\bar{\bar{\Gamma}}=\left(\begin{array}{cc}\Gamma^{(L)} & \vartheta \\ \theta & 0\end{array}\right),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0896 (13.5.1)	288	■ 1.000 ● N +0	$\mathrm{\hat{\Gamma}}=\mathrm{\Gamma}_{\alpha}^{\beta}L^{\alpha}_{\beta}+\vartheta^{\alpha}P_{\alpha}+\theta_{\beta}P_{*}^{\beta},$	$\hat{\Gamma}=\Gamma_{\alpha}^{\beta}L^{\alpha}_{\beta}+\vartheta^{\alpha}P_{\alpha}+\theta_{\beta}P_{*}^{\beta},$	$\hat{\hat{\Gamma}}=\Gamma_{\alpha}^{\beta}L^{\alpha}_{\beta}+\vartheta^{\alpha}P_{\alpha}+\theta_{\beta}P_{*}^{\beta},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0899 (13.5.4)	288	■ 1.000 ● K +0	$\mathrm{\tilde{L}}_{\mathrm{\mathrm{SL}}(5,\mathrm{\mathbb{R}})}=-\mathrm{operatorname{Tr}}\{\mathrm{\hat{B}}\wedge\mathrm{\hat{R}}\}-d\hat{C}.$	$\tilde{L}_{\mathrm{SL}(5,\mathbb{R})}=-\mathrm{Tr}\{\hat{B}\wedge\hat{R}\}-d\hat{C}.$	$\tilde{\tilde{L}}_{\mathrm{SL}(5,\mathbb{R})}=-\mathrm{Tr}\{\hat{\hat{B}}\wedge\hat{\hat{R}}\}-d\hat{\hat{C}}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0900 (13.5.5)	289	■ 1.000 ● N +0	$\mathrm{\hat{D}}B_{\mathrm{A}}^{\mathrm{B}}\cong\mathrm{\hat{D}}\mathrm{\hat{R}}_{\mathrm{A}}^{\mathrm{B}}\equiv 0.$	$\hat{D}B_A^B\cong\hat{D}\hat{R}_A^B\equiv 0.$	$\hat{\hat{D}}B_A^B\cong\hat{\hat{D}}\hat{\hat{R}}_A^B\equiv 0.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0901 (13.5.6)	289	■ 1.000 ● N -1	$\mathrm{\tilde{L}}_{\mathrm{\mathrm{SSB}}}=\mathrm{\tilde{L}}_{\mathrm{\mathrm{SL}}(5,\mathrm{\mathbb{R}})}+\frac{1}{2}\eta_{ABCDE}B^{AB}\wedge B^{CD}\Phi^E.$	$\tilde{L}_{\mathrm{SSB}}=\tilde{L}_{\mathrm{SL}(5,\mathbb{R})}+\frac{1}{2}\eta_{ABCDE}B^{AB}\wedge B^{CD}\Phi^E.$	$\tilde{\tilde{L}}_{\mathrm{SSB}}=\tilde{\tilde{L}}_{\mathrm{SL}(5,\mathbb{R})}+\frac{1}{2}\eta_{ABCDE}B^{AB}\wedge B^{CD}\Phi^E.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0902 (13.5.7)	289	■ 1.000 ● N +0	$\hat{g}_{AB} \Phi^A \Phi^B = \mu^2, \quad \Phi_E \hat{D} \Phi^E = 0$	$\hat{g}_{AB} \Phi^A \Phi^B = \mu^2, \quad \Phi_E \hat{D} \Phi^E = 0$	$\hat{g}_{AB} \Phi^A \Phi^B = \mu^2, \quad \Phi_E \hat{D} \Phi^E = 0,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0903 (13.5.8)	289	■ 1.000 ● K +0	$\left\langle \Phi^E \right\rangle = \Phi_0^E = (0, 0, 0, 0, \mu)^T$	$\langle \Phi^E \rangle = \Phi_0^E = (0, 0, 0, 0, \mu)^T$	$\langle \Phi^E \rangle = \Phi_0^E = (0, 0, 0, 0, \mu)^T$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0904 (13.5.9)	289	■ 1.000 ● K +0	$\tilde{L}_{\text{SSB}} = -\text{Tr}\{\hat{B} \wedge \hat{R}\} + \frac{\mu}{2} \eta_{\alpha\beta\gamma\delta} B^{\alpha\beta} \wedge B^{\gamma\delta} - dC_{\text{RR}} + 2dC_{\text{TT}}$	$\tilde{L}_{\text{SSB}} = -\text{Tr}\{\hat{B} \wedge \hat{R}\} + \frac{\mu}{2} \eta_{\alpha\beta\gamma\delta} B^{\alpha\beta} \wedge B^{\gamma\delta} - dC_{\text{RR}} + 2dC_{\text{TT}}$	$\tilde{L}_{\text{SSB}} = -\text{Tr}\{\hat{B} \wedge \hat{R}\} + \frac{\mu}{2} \eta_{\alpha\beta\gamma\delta} B^{\alpha\beta} \wedge B^{\gamma\delta} - dC_{\text{RR}} + 2dC_{\text{TT}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0905 (13.5.10)	290	■ 1.000 ● N +1	$d \hat{C} = dC_{\text{RR}} - 2dC_{\text{TT}},$	$d\hat{C} = dC_{\text{RR}} - 2dC_{\text{TT}},$	$d\hat{C} = dC_{\text{RR}} - 2dC_{\text{TT}},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0906 (13.5.11)	290	■ 1.000 ● K +0	$B_{\alpha\beta} \cong \frac{1}{\mu} \eta_{\alpha\beta\gamma\delta} \left[R^{\gamma\delta} - \frac{\Lambda}{3} \theta^\gamma \wedge \vartheta^\delta \right]$	$B_{\alpha\beta} \cong \frac{1}{\mu} \eta_{\alpha\beta\gamma\delta} \left[R^{\gamma\delta} - \frac{\Lambda}{3} \theta^\gamma \wedge \vartheta^\delta \right]$	$B_{\alpha\beta} \cong \frac{1}{\mu} \eta_{\alpha\beta\gamma\delta} \left[R^{\gamma\delta} - \frac{\Lambda}{3} \theta^\gamma \wedge \vartheta^\delta \right]$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ9909 (13.5.14)	291	■ 1.000 ● N -1	$\tilde{L}_{\mathrm{SSB}} \cong -\frac{1}{2\kappa} R^{\alpha\beta} \wedge \eta_{\alpha\beta} + \frac{\Lambda}{\kappa} \eta$		$\tilde{L}_{\mathrm{SSB}} \cong -\frac{1}{2\kappa} R^{\alpha\beta} \wedge \eta_{\alpha\beta} + \frac{\Lambda}{\kappa} \eta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ9910 (13.5.15)	291	■ 1.000 ● N +0	$\mu=\frac{4}{3}\kappa\Lambda=\frac{1}{3}(2\kappa)^2\rho_\Lambda$		$\mu=\frac{4}{3}\kappa\Lambda=\frac{1}{3}(2\kappa)^2\rho_\Lambda.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ9912 (13.7.1)	293	■ 1.000 ● N +0	$g_{\mathrm{N}}:=\kappa\,k^2,\quad \lambda:=\Lambda/k^2,$	$g_{\mathrm{N}}:=\kappa k^2,\quad \lambda:=\Lambda/k^2,$	$g_{\mathrm{N}}:=\kappa k^2,\quad \lambda:=\Lambda/k^2,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ9913 (13.7.2)	293	■ 1.000 ● W +0	$k\,\frac{\partial}{\partial k}g_{\mathrm{N}}=\beta_1(g_{\mathrm{N}},\lambda)=(2+d_{\mathrm{N}})g_{\mathrm{N}},\quad k\,\frac{\partial}{\partial k}\lambda=\beta_2(g_{\mathrm{N}},\lambda),$		$k\frac{\partial}{\partial k}g_{\mathrm{N}}=\beta_1(g_{\mathrm{N}},\lambda)=(2+d_{\mathrm{N}})g_{\mathrm{N}},\quad k\frac{\partial}{\partial k}\lambda=\beta_2(g_{\mathrm{N}},\lambda),$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ9914 (13.7.3)	293	■ 1.000 ● N +0	$\mu_*\leq\frac{4}{3}g_{\mathrm{N}*}\lambda_*\simeq 0.2$		$\mu_*\leq\frac{4}{3}g_{\mathrm{N}*}\lambda_*\simeq 0.2$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0916 (13.8.2)	295	■ 1.000 ● N -1	$d\,C_{\mathrm{TT}}=\frac{1}{2\,\ell^2}\,\left(T^{\alpha}\wedge T_{\alpha}+R_{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}\right)$		
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0917 (13.8.3)	295	■ 1.000 ● N +0	$d\,C_{\mathrm{RR}}=-\frac{1}{2}\,R_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}$		
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0918 (13.8.4)	295	■ 1.000 ● N +0	$D\,R_{\alpha}{}^{\beta}\equiv 0,$		
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0921 (13.8.7)	296	■ 1.000 ● N +1	$L_{\mathrm{SKY}}:=-\frac{1}{2}\,R_{\alpha\beta}\wedge R^{\alpha\beta},$		
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0922 (14.1.1)	302	■ 1.000 ● N +0	$L=L_g+L_{\omega}+L_{\mathrm{mat}}$		
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0923 (14.1.2)	302	■ 1.000 ● N +1	$\mathrm{L}_{\{\omega\text{-}g\}}=\operatorname{Tr}\left\{\wedge^*\left(\vartheta\wedge\Omega^g\right)\wedge^*\left(\vartheta\wedge\Omega\right)\right\}$	$L_{\omega-g}=\operatorname{Tr}\left\{\wedge^*\left(\vartheta\wedge\Omega^g\right)\wedge^*\left(\vartheta\wedge\Omega\right)\right\}$	$L_{\omega-g}=\operatorname{Tr}\left\{\wedge^*\left(\vartheta\wedge\Omega^g\right)\wedge^*\left(\vartheta\wedge\Omega\right)\right\}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0924 (14.1.3)	303	■ 1.000 ● N -1	$\mathrm{L}_{\{\mathrm{GMD}\}}.=\frac{1}{\ell^{*2}}\mathrm{L}_{\{\mathrm{W}\}}.+L_{\{\mathrm{YM}\}}+L_{\{\mathrm{mat}\}}$	$L_{\mathrm{GMD}.}=\frac{1}{\ell^{*2}}L_{\mathrm{W}.}+L_{\mathrm{YM}}+L_{\mathrm{mat}}$	$L_{\mathrm{GMD}.}=\frac{1}{\ell^{*2}}L_{\mathrm{W}.}+L_{\mathrm{YM}}+L_{\mathrm{mat}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0933 (14.1.14)	305	■ 1.000 ● K +0	$e^{\{*\pi/2\}}=(-1)^{z*},\quad z:=\frac{s-1}{2}.$	$e^{*\pi/2}=(-1)^{z*},\quad z:=\frac{s-1}{2}.$	$e^{*\pi/2}=(-1)^{z*},\quad z:=\frac{s-1}{2}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0934 (14.1.15)	305	■ 1.000 ● K +0	$\xi:=e^{\{*(-\delta)\}}\Omega$	$\xi:=e^{*(-\delta)}\Omega$	$\xi:=e^{*(-\delta)}\Omega$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0935 (14.1.16)	305	■ 1.000 ● N +0	$\Omega\wedge^*\Omega\rightarrow\xi\wedge^*\xi=\Omega\wedge^*\Omega\cos2\delta-(-1)^z\Omega\wedge\Omega\sin2\delta$	$\Omega\wedge^*\Omega\rightarrow\xi\wedge^*\xi=\Omega\wedge^*\Omega\cos2\delta-(-1)^z\Omega\wedge\Omega\sin2\delta$	$\Omega\wedge^*\Omega\rightarrow\xi\wedge^*\xi=\Omega\wedge^*\Omega\cos2\delta-(-1)^z\Omega\wedge\Omega\sin2\delta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0936 (14.1.17)	305	■ 1.000 ● N +0	$\Omega\wedge\Omega\rightarrow\xi\wedge\xi=\Omega\wedge\Omega\cos2\delta-(-1)^z\Omega\wedge^*\Omega\sin2\delta.$	$\Omega\wedge\Omega\rightarrow\xi\wedge\xi=\Omega\wedge\Omega\cos2\delta-(-1)^z\Omega\wedge^*\Omega\sin2\delta.$	$\Omega\wedge\Omega\rightarrow\xi\wedge\xi=\Omega\wedge\Omega\cos2\delta-(-1)^z\Omega\wedge^*\Omega\sin2\delta.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0937 (14.1.18)	305	■ 1.000 ● N +0	$\vec{E}=\left\{E_{\text{j}}:=F_{\text{j } 0} \mid \text{j}=1,2,3\right\}$	$\vec{E}=\left\{E_j:=F_{j0} \mid j=1,2,3\right\}$	$\vec{E}=\left\{E_j:=F_{j0} \mid j=1,2,3\right\}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0938 (14.1.19)	305	■ 1.000 ● N +0	$\vec{B}=\left\{B_{\text{j}}: \left.=\frac{1}{2} \epsilon_{i j k} F^{\text{j } k}\right i, j, k=1,2,3\right\}$	$\vec{B}=\left\{B_j:=\frac{1}{2} \epsilon_{ijk} F^{jk} \mid i, j, k=1,2,3\right\}$	$\vec{B}=\left\{B_j:=\frac{1}{2} \epsilon_{ijk} F^{jk} \mid i, j, k=1,2,3\right\}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0943 (14.1.24)	306	■ 1.000 ● N +1	$\Omega=e^{\ast} \delta \xi$	$\Omega=e^{\ast \delta} \xi$	$\Omega=e^{\ast \delta} \xi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0944 (14.1.25)	306	■ 1.000 ● N -1	$\Sigma_{\mathrm{YM}}(\Omega)=\Sigma_{\mathrm{YM}}(\xi)$	$\Sigma_{\mathrm{YM}}(\Omega)=\Sigma_{\mathrm{YM}}(\xi)$	$\Sigma_{\mathrm{YM}}(\Omega)=\Sigma_{\mathrm{YM}}(\xi)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0950 (14.1.31)	307	■ 1.000 ● K +0	$\tau_{\text{e}}=\bar{\tau}_{\text{e}} \cos \delta-\left(-1\right)^z \bar{\tau}_{\text{p}} \sin \delta$	$\tau_e=\bar{\tau}_e \cos \delta-\left(-1\right)^z \bar{\tau}_p \sin \delta$	$\tau_e=\bar{\tau}_e \cos \delta-\left(-1\right)^z \bar{\tau}_p \sin \delta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ0952 (14.1.34)	307	■ 1.000 ● N +0	$D \delta=d \delta=0,$	$D \delta=d \delta=0,$	$D \delta=d \delta=0,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0955 (14.1.37)	308	■ 1.000 ● N +0	$(-1)^{z}\,\mathrm{d}\,\Sigma_{\mathrm{YM}}(\xi)\wedge\mathrm{d}\,\Delta=D\,\Sigma_{\mathrm{YM}}(\xi)\,\mathrm{d}\,\xi\,.$	$(-1)^z\Sigma_{\mathrm{YM}}(\xi)\wedge d\delta=D\Sigma_{\mathrm{YM}}(\xi).$	$(-1)^z\Sigma_{\mathrm{YM}}(\xi)\wedge d\delta=D\Sigma_{\mathrm{YM}}(\xi).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0958 (14.1.40)	309	■ 1.000 ● N +1	$\Delta=\int_{c_1}\mathrm{d}\,\Delta+\Delta_0$	$\delta=\int_{c_1}d\delta+\delta_0$	$\delta=\int_{c_1}d\delta+\delta_0.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0969 (14.2.10)	314	■ 1.000 ● N +1	$\hat{\omega}^{AB}=-\hat{\omega}^{BA},\quad A,B=0,\ldots,3+K,\ldots,3+K$	$\hat{\omega}^{AB}=-\hat{\omega}^{BA},\quad A,B=0,\ldots,3+K$	$\hat{\omega}^{AB}=-\hat{\omega}^{BA},\quad A,B=0,\ldots,3+K$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0975 (14.2.16)	315	■ 1.000 ● N +1	$\Omega^{\prime}:=\pi\left(d\omega-\omega\wedge\overset{\circ}{\omega}-\omega^g\wedge\omega\right)=\Omega-\omega^g\wedge\omega.$	$\Omega^{\prime}:=\pi\left(d\omega-\omega\wedge\overset{\circ}{\omega}-\omega^g\wedge\omega\right)=\Omega-\omega^g\wedge\omega.$	$\Omega^{\prime}:=\pi(d\omega-\omega\wedge\overset{\circ}{\omega}-\omega^g\wedge\omega)=\Omega-\omega^g\wedge\omega.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__EQ0980 (14.2.21)	316	■ 1.000 ● N +0	$\hat{g}_{AB}\left(x,\xi^j\right)=\sum_{n=-\infty}^{+\infty}g_{AB}^{\left(n\right)}\left(x\right)Y_n\left(\xi^j\right)$	$\hat{g}_{AB}\left(x,\xi^j\right)=\sum_{n=-\infty}^{+\infty}g_{AB}^{\left(n\right)}\left(x\right)Y_n\left(\xi^j\right)$	$\hat{g}_{AB}(x,\xi^j)=\sum_{n=-\infty}^{+\infty}g_{AB}^{(n)}(x)Y_n(\xi^j)$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0981 (14.2.22)	317	■ 1.000 ● N -1	$M^*=\sqrt{\hbar c / G_{\mathrm{N}}}} \,\simeq\, 10^{19} \,\mathrm{GeV} / c^2 \, .$	$M^* = \sqrt{\hbar c / G_{\mathrm{N}}} \simeq 10^{19} \mathrm{GeV} / c^2 .$	$M^* = \sqrt{\hbar c / G_{\mathrm{N}}} \simeq 10^{19} \mathrm{GeV} / c^2 .$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0983 (14.2.24)	318	■ 1.000 ● K +0	$G=S\,U(3)\,\times\,S\,U(2)\,\times\,U(1)$	$G = SU(3) \times SU(2) \times U(1)$	$G = SU(3) \times SU(2) \times U(1)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0984 (14.2.25)	318	■ 1.000 ● N +0	$B=\mathbb{C}P^2\,\times\,S^2\,\times\,S^1,$	$B = \mathbb{C}P^2 \times S^2 \times S^1,$	$B = \mathbb{C}P^2 \times S^2 \times S^1,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0985 (14.2.26)	318	■ 1.000 ● N +1	$G/H=\frac{S\,U(3)\,\times\,S\,U(2)\,\times\,U(1)}{S\,U(2)\,\times\,U(1)\,\times\,U(1)},$	$G/H = \frac{SU(3) \times SU(2) \times U(1)}{SU(2) \times U(1) \times U(1)} ,$	$G/H = \frac{SU(3) \times SU(2) \times U(1)}{SU(2) \times U(1) \times U(1)} ,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0986 (14.2.27)	318	■ 1.000 ● K +0	$\widetilde{S}^7:=\frac{S\,O(5)\,\times\,S\,O(3)}{S\,O(3)\,\times\,S\,O(3)} \, .$	$\widetilde{S}^7 := \frac{SO(5) \times SO(3)}{SO(3) \times SO(3)} .$	$\widetilde{S}^7 := \frac{SO(5) \times SO(3)}{SO(3) \times SO(3)} .$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0988 (14.3.1)	319	■ 1.000 ● K +0	$\varphi \rightarrow \varphi^{\mathrm{C}} := \bar{\varphi}$	$\varphi \rightarrow \varphi^{\mathrm{C}} := \bar{\varphi}$	$\varphi \rightarrow \varphi^{\mathrm{C}} := \bar{\varphi}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0991 (14.3.4)	320	■ 1.000 ● N +0	$\hat{\psi} = \sum_{F=1}^{\operatorname{dim} \rho(G)} \left(\psi_{-}^F(x) \chi_{\rho F}^{-}(\xi) + \psi_{+}^F(x) \chi_{\rho F}^{+}(\xi) \right).$	$\hat{\psi} = \sum_{F=1}^{\operatorname{dim} \rho(G)} \left(\psi_{-}^F(x) \chi_{\rho F}^{-}(\xi) + \psi_{+}^F(x) \chi_{\rho F}^{+}(\xi) \right).$	$\hat{\psi} = \sum_{F=1}^{\operatorname{dim} \rho(G)} (\psi_{-}^F(x) \chi_{\rho F}^{-}(\xi) + \psi_{+}^F(x) \chi_{\rho F}^{+}(\xi)).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0994 (14.3.7)	321	■ 1.000 ● N +0	$D^{g+\omega} := d + i \widetilde{\omega}^g \otimes i \omega$	$D^{g+\omega} := d + i \widetilde{\omega}^g \otimes i \omega$	$D^{g+\omega} := d + i \widetilde{\omega}^g \otimes i \omega$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ0999 (14.3.12)	322	■ 1.000 ● N +0	$\left(\widehat{\gamma}^{n+1}\right)^2 = -1 \rightarrow \widehat{\psi} = \mp i \widehat{\psi}.$	$\left(\widehat{\gamma}^{n+1}\right)^2 = -1 \Rightarrow \widehat{\gamma}^{n+1} \widehat{\psi} = \mp i \widehat{\psi}.$	$(\widehat{\gamma}^{n+1})^2 = -1 \Rightarrow \widehat{\gamma}^{n+1} \widehat{\psi} = \mp i \widehat{\psi}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1000 (14.3.13)	322	■ 1.000 ● N +0	$\left(\widehat{\gamma}^{n+1}\right)^2 = +1 \rightarrow \widehat{\psi} = \mp \widehat{\psi}.$	$\left(\widehat{\gamma}^{n+1}\right)^2 = +1 \Rightarrow \widehat{\gamma}^{n+1} \widehat{\psi} = \mp \widehat{\psi}.$	$(\widehat{\gamma}^{n+1})^2 = +1 \Rightarrow \widehat{\gamma}^{n+1} \widehat{\psi} = \mp \widehat{\psi}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1006 (14.4.1)	324	■ 1.000 ● N +0	$\bar{Q}^{(i)} = Q^{(i)} \cos \delta + P^{(i)} \sin \delta, \quad Q^{(\circ)} \equiv Q_e.$	$\bar{Q}^{(i)} = Q^{(i)} \cos \delta + P^{(i)} \sin \delta, \quad Q^{(\circ)} \equiv Q_e.$	$\bar{Q}^{(i)} = Q^{(i)} \cos \delta + P^{(i)} \sin \delta, \quad Q^{(\circ)} \equiv Q_e.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1009 (14.4.4)	325	■ 1.000 ● N +0	$\rho^{\wedge}{2}:=r^{\wedge}{2}+\frac{J(J+1)}{\hbar^{\wedge}{2}}{M^{\wedge}{2}}{c^{\wedge}{2}}\cos^{\wedge}{2}\vartheta$	$\rho^2:=r^2+\frac{J(J+1)\hbar^2}{M^2c^2}\cos^2\vartheta$	$\rho^2:=r^2+\frac{J(J+1)\hbar^2}{M^2c^2}\cos^2\vartheta$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1010 (14.4.5)	325	■ 1.000 ● W +0	$\Delta:=r^{\wedge}{2}-\frac{2M}{\hbar^{\wedge}{2}}{M^{\wedge}{2}}{c}r+\frac{J(J+1)}{\hbar^{\wedge}{2}}{M^{\wedge}{2}}{c^{\wedge}{2}}+\frac{\alpha_e}{\hbar^{\wedge}{2}}\hbar^{\wedge}{2}{M^{\wedge}{2}}{c^{\wedge}{2}}\operatorname{Tr}\bar{Q}^{\wedge}{2}$	$\Delta:=r^2-\frac{2M\hbar}{M^{*2}c}r+\frac{J(J+1)\hbar^2}{M^2c^2}+\frac{\alpha_e\hbar^2}{M^{*2}c^2}\operatorname{Tr}\bar{Q}^2$	$\Delta:=r^2-\frac{2M\hbar}{M^{*2}c}r+\frac{J(J+1)\hbar^2}{M^2c^2}+\frac{\alpha_e\hbar^2}{M^{*2}c^2}\operatorname{Tr}\bar{Q}^2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1017 (14.4.12)	327	■ 1.000 ● N +0	$d s^{\wedge}{2}=-\frac{\bar{\Delta}}{N}{\hbar^{\wedge}{2}}{\sin^{\wedge}{2}\left(\frac{\vartheta}{2}\right)}d\phi+c d t^{\wedge}{2}+\frac{\bar{\rho}^{\wedge}{2}}{\Delta}\left(d r^{\wedge}{2}+\bar{\Delta} d \Omega^{\wedge}{2}\right),$	$ds^2=-\frac{\bar{\Delta}}{\bar{\rho}^2}\left(\frac{4N\hbar}{Mc}\sin^2\left(\frac{\vartheta}{2}\right)d\phi+cdt\right)^2+\frac{\bar{\rho}^2}{\bar{\Delta}}\left(dr^2+\bar{\Delta}d\Omega^2\right),$	$ds^2=-\frac{\bar{\Delta}}{\bar{\rho}^2}\left(\frac{4N\hbar}{Mc}\sin^2\left(\frac{\vartheta}{2}\right)d\phi+cdt\right)^2+\frac{\bar{\rho}^2}{\bar{\Delta}}(dr^2+\bar{\Delta}d\Omega^2),$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1018 (14.4.13)	328	■ 1.000 ● N +0	$\bar{\Delta}:=r^{\wedge}{2}-\frac{2M}{\hbar^{\wedge}{2}}{M^{\wedge}{2}}{c}r-\frac{N^{\wedge}{2}}{\hbar^{\wedge}{2}}{\hbar^{\wedge}{2}}{M^{\wedge}{2}}{c^{\wedge}{2}}+\frac{\alpha_e}{\hbar^{\wedge}{2}}\hbar^{\wedge}{2}{M^{\wedge}{2}}{c^{\wedge}{2}}\operatorname{Tr}\bar{Q}^{\wedge}{2}$	$\bar{\Delta}:=r^2-\frac{2M\hbar}{M^{*2}c}r-\frac{N^2\hbar^2}{M^2c^2}+\frac{\alpha_e\hbar^2}{M^{*2}c^2}\operatorname{Tr}\bar{Q}^2$	$\bar{\Delta}:=r^2-\frac{2M\hbar}{M^{*2}c}r-\frac{N^2\hbar^2}{M^2c^2}+\frac{\alpha_e\hbar^2}{M^{*2}c^2}\operatorname{Tr}\bar{Q}^2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1019 (14.4.14)	328	■ 1.000 ● N +0	$\bar{\rho}^{\wedge}{2}:=r^{\wedge}{2}+\frac{N^{\wedge}{2}}{\hbar^{\wedge}{2}}{\hbar^{\wedge}{2}}{M^{\wedge}{2}}{c^{\wedge}{2}}$	$\bar{\rho}^2:=r^2+\frac{N^2\hbar^2}{M^2c^2}$	$\bar{\rho}^2:=r^2+\frac{N^2\hbar^2}{M^2c^2}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1020 (14.4.15)	328	■ 1.000 ● W +1	$\begin{aligned} A = & -\frac{1}{\bar{\rho}^2}(\bar{Q}\cos\delta+N\bar{Q}\sin\delta)c\,d t\,\, \& +\frac{1-\cos\vartheta}{\bar{\rho}^2}\bar{Q}\left[-\bar{\rho}^2\sin\delta+2r\frac{N\hbar}{Mc}\cos\delta\right]d\phi. \end{aligned}$		
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1022 (14.4.17)	328	■ 1.000 ● W +0	$M^{\{2\}}/M_{\mathrm{ir}}^{\{2\}}=\frac{\left(1+\frac{1}{4}\left(\alpha_e^2\mathrm{Tr}\,\bar{Q}^2-2N^2\right)\left(M^*/M_{\mathrm{ir}}\right)^4\right)^2}{1-\frac{1}{4}N^2\left(M^*/M_{\mathrm{ir}}\right)^4},$		$M^2/M_{\mathrm{ir}}^2=\frac{\left[1+\frac{1}{4}\left(\alpha_e^2\mathrm{Tr}\,\bar{Q}^2-2N^2\right)\left(M^*/M_{\mathrm{ir}}\right)^4\right]^2}{1-\frac{1}{4}N^2\left(M^*/M_{\mathrm{ir}}\right)^4},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1023 (15.0.1)	335	■ 1.000 ● N +0	$\ell:=2\,\pi\,\hbar/\left(M_P\,c\right)=\ell^*/\sqrt{\kappa}\simeq1.3\,\times\,10^{-13}\,\mathrm{cm}$		$\ell:=2\pi\hbar/(M_Pc)=\ell^*/\sqrt{\kappa}\simeq1.3\times10^{-13}\,\mathrm{cm}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1024 (15.0.2)	335	■ 1.000 ● N +0	$V_{\mathrm{Yu}}(r)\,\sim\,\frac{1}{r}e^{-rm_{\pi}c/\hbar},$	$V_{\mathrm{Yu}}(r)\sim\frac{1}{r}e^{-rm_{\pi}c/\hbar},$	$V_{\mathrm{Yu}}(r)\sim\frac{1}{r}e^{-rm_{\pi}c/\hbar},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1026 (15.1.2)	336	■ 1.000 ● N +0	$L_{fg}=P\left(\vartheta^f,\vartheta^g\right)\eta,$	$L_{fg}=P\left(\vartheta^f,\vartheta^g\right)\eta,$	$L_{fg}=P(\vartheta^f,\vartheta^g)\eta,$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1027 (15.1.3)	337	■ 1.000 ● N +0	$\backslash\mathrm{delta}\ \mathrm{L}_{\mathrm{f}\ \mathrm{g}}=: \backslash\mathrm{delta}\ \mathrm{\vartheta}^{\mathrm{f}}\ \mathrm{\wedge}\ \mathrm{\Sigma}^{\mathrm{f}}+\backslash\mathrm{delta}\ \mathrm{\vartheta}^{\mathrm{g}}\ \mathrm{\wedge}\ \mathrm{\Sigma}^{\mathrm{g}}\ .$	$\delta L_{fg} =: \delta \vartheta^f \wedge \Sigma^f + \delta \vartheta^g \wedge \Sigma^g.$	$\delta L_{fg} =: \delta \vartheta^f \wedge \Sigma^f + \delta \vartheta^g \wedge \Sigma^g.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1028 (15.1.4)	337	■ 1.000 ● N +0	$\backslash\mathrm{frac}\{1\}\{a\}\ \mathrm{\vartheta}^{\mathrm{f}}\ \mathrm{\wedge}\ \mathrm{\Sigma}^{\mathrm{f}}+\backslash\mathrm{frac}\{1\}\{b\}\ \mathrm{\vartheta}^{\mathrm{g}}\ \mathrm{\wedge}\ \mathrm{\Sigma}^{\mathrm{g}}=2\ \mathrm{L}_{\mathrm{f}\ \mathrm{g}}\ .$	$\frac{1}{a} \vartheta^f \wedge \Sigma^f + \frac{1}{b} \vartheta^g \wedge \Sigma^g = 2 L_{fg}.$	$\frac{1}{a} \vartheta^f \wedge \Sigma^f + \frac{1}{b} \vartheta^g \wedge \Sigma^g = 2\ L_{fg}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1029 (15.1.5)	337	■ 1.000 ● W +0	$\mathrm{L}_{\mathrm{f}\ \mathrm{g}}^{\mathrm{\prime}}=\mathrm{frac}\{\mathrm{m}_{\mathrm{f}}^{\mathrm{2}}\}\{\mathrm{ell}^{\mathrm{2}}\}\backslash\mathrm{left}\ \mathrm{operatorname{det}}\ \mathrm{g}_{\mathrm{i}\ \mathrm{j}}\backslash\mathrm{right}\ ^{\mathrm{u}}\backslash\mathrm{left}\ \mathrm{operatorname{det}}\ \mathrm{f}_{\mathrm{i}\ \mathrm{j}}\backslash\mathrm{right}\ ^{\mathrm{1}\ /\ 2-\mathrm{u}}\backslash\mathrm{left}(\mathrm{\vartheta}^{\mathrm{f}}-\mathrm{\vartheta}^{\mathrm{g}}\backslash\mathrm{right})\ \mathrm{\wedge}\ \backslash\mathrm{left}(\mathrm{\vartheta}^{\mathrm{f}}-\mathrm{\vartheta}^{\mathrm{g}}\backslash\mathrm{right})\ \mathrm{\wedge}\ \mathrm{\vartheta}^{\mathrm{g}}\ \mathrm{\wedge}\ \mathrm{\vartheta}^{\mathrm{f}}$	$L'_{fg} = \frac{m_f^2}{\ell^2} \det g_{ij} ^u \det f_{ij} ^{1/2-u} (\vartheta^f - \vartheta^g) \wedge (\vartheta^f - \vartheta^g) \wedge \vartheta^g \wedge \vartheta^f$	$L'_{fg} = \frac{m_f^2}{\ell^2} \det g_{ij} ^u \det f_{ij} ^{1/2-u} (\vartheta^f - \vartheta^g) \wedge (\vartheta^f - \vartheta^g) \wedge \vartheta^g \wedge \vartheta^f$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1035 (15.1.12)	338	■ 1.000 ● N +1	$\mathrm{\vartheta}^{\mathrm{m}_{\mathrm{f}}}:=\mathrm{\vartheta}^{\mathrm{f}}-\mathrm{\vartheta}^{\mathrm{g}}$	$\vartheta^{m_f} := \vartheta^f - \vartheta^g$	$\vartheta^{m_f} := \vartheta^f - \vartheta^g$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1036 (15.1.13)	338	■ 1.000 ● W +1	$\backslash\mathrm{begin}\{\mathrm{aligned}\}\ \mathrm{d}\ \mathrm{x}^{\mathrm{2}}\ \&=\mathrm{frac}\{1\}\{1+\mathrm{\kappa}\}\ \mathrm{g}^{\{\mathrm{\alpha}\ \mathrm{\beta}\}}\backslash\mathrm{left}(\mathrm{\vartheta}_{\mathrm{\alpha}}^{\mathrm{g}}\ \mathrm{\otimes}_s\ \mathrm{\vartheta}_{\mathrm{\beta}}^{\mathrm{g}}+\mathrm{\kappa}\ \mathrm{\vartheta}_{\mathrm{\alpha}}^{\mathrm{f}}\ \mathrm{\otimes}\ \mathrm{\vartheta}_{\mathrm{\beta}}^{\mathrm{f}}\backslash\mathrm{right})+\mathrm{\kappa}\ \mathrm{\vartheta}_{\mathrm{\alpha}}^{\mathrm{f}}\ \mathrm{\otimes}\ \mathrm{\vartheta}_{\mathrm{\beta}}^{\mathrm{g}}\backslash\mathrm{right})\ \backslash\&=\mathrm{frac}\{1\}\{1+\mathrm{\kappa}\}\backslash\mathrm{left}(\mathrm{g}_{\mathrm{i}\ \mathrm{j}}+\mathrm{\kappa}\ \mathrm{f}_{\mathrm{i}\ \mathrm{j}}\backslash\mathrm{right})\ \mathrm{d}\ \mathrm{x}^{\mathrm{i}}\ \mathrm{\otimes}_s\ \mathrm{d}\ \mathrm{x}^{\mathrm{j}}\ \backslash\mathrm{end}\{\mathrm{aligned}\}$	$ds^2 = \frac{1}{1+\kappa} g^{\alpha\beta} \left(\vartheta_{\alpha}^g \otimes_s \vartheta_{\beta}^g + \kappa \vartheta_{\alpha}^f \otimes \vartheta_{\beta}^f \right) = \frac{1}{1+\kappa} (g_{ij} + \kappa f_{ij}) dx^i \otimes_s dx^j$	$ds^2 = \frac{1}{1+\kappa} g^{\alpha\beta} \left(\vartheta_{\alpha}^g \otimes_s \vartheta_{\beta}^g + \kappa \vartheta_{\alpha}^f \otimes \vartheta_{\beta}^f \right) = \frac{1}{1+\kappa} (g_{ij} + \kappa f_{ij}) dx^i \otimes_s dx^j.$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1037 (15.1.14)	338	■ 1.000 ● N +0	$\kappa=\ell^{{*}2}/\ell^2=G_{\mathrm{N}}/G_{\mathrm{S}}=0.38\,\times\,10^{-38}$	$\kappa = \ell^{*2}/\ell^2 = G_{\mathrm{N}}/G_{\mathrm{S}} = 0.38 \times 10^{-38}$	$\kappa = \ell^{*2}/\ell^2 = G_{\mathrm{N}}/G_{\mathrm{S}} = 0.38 \times 10^{-38}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1038 (15.1.15)	339	■ 1.000 ● W +1	$e^{\{-\lambda^{\mathrm{g},\,\mathrm{f}}\}}\!:=\!1\!-\!\frac{2\,\mu^{\mathrm{g},\,\mathrm{f}}}{r}\!-\!\frac{1}{3}\Lambda^{\mathrm{g},\,\mathrm{f}}r^2,$	$e^{-\lambda^{g,f}} := 1 - \frac{2\mu^{g,f}}{r} - \frac{1}{3}\Lambda^{g,f}r^2,$	$e^{-\lambda^{g,f}} := 1 - \frac{2\mu^{g,f}}{r} - \frac{1}{3}\Lambda^{g,f}r^2,$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1040 (15.1.17)	339	■ 1.000 ● N +0	$e^{\{-\lambda^{\mathrm{f}}\}}\,\,\,\simeq\,\,1\!-\!\frac{2\,\mu^{\mathrm{f}}}{r}\,\exp\left(-r\,m_{\mathrm{f}}\,c\,/\,\hbar\right)\,.$	$e^{-\lambda^f} \simeq 1 - \frac{2\mu^f}{r} \exp\left(-rm_f c/\hbar\right).$	$e^{-\lambda^f} \simeq 1 - \frac{2\mu^f}{r} \exp\left(-r\,m_f\,c/\hbar\right).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1041 (15.1.18)	340	■ 1.000 ● K +0	$E_{\{\alpha\,\beta\}}\!:\!\left(2^{+}\right)\,\oplus\,2\!\left(1^{-}\right)\,\oplus\,2\!\left(0^{+}\right)\,\oplus\,2\!\left(1^{+}\right)\,\oplus\,2\!\left(0^{+}\right)\,\oplus\!\left(1^{+}\right)$	$E_{\alpha\beta} : (2^+) \oplus 2\,(1^-) \oplus 2\,(0^+) \oplus (1^+)$	$E_{\alpha\beta} : (2^+) \oplus 2(1^-) \oplus 2(0^+) \oplus (1^+)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1043 (15.1.20)	342	■ 1.000 ● N +0	$\frac{M_{\mathrm{U}}}{M_{\mathrm{P}}}=\frac{\ell_{\mathrm{H}}}{\ell^{*}}\cdot\frac{\ell}{\ell^{*}}=\frac{1}{\kappa}\frac{\ell_{\mathrm{H}}}{\ell}\simeq 10^{80}$	$\frac{M_{\mathrm{U}}}{M_{\mathrm{P}}} = \frac{\ell_{\mathrm{H}}}{\ell^{*}} \cdot \frac{\ell}{\ell^{*}} = \frac{1}{\kappa} \frac{\ell_{\mathrm{H}}}{\ell} \simeq 10^{80}$	$\frac{M_{\mathrm{U}}}{M_{\mathrm{P}}} = \frac{\ell_{\mathrm{H}}}{\ell^{*}} \cdot \frac{\ell}{\ell^{*}} = \frac{1}{\kappa} \frac{\ell_{\mathrm{H}}}{\ell} \simeq 10^{80}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1044 (15.2.1)	343	■ 1.000 ● W +0	$\begin{aligned} \bar{G} &= SL(2f, \mathbb{C}) \otimes SL(2c, \mathbb{C}) \\ &\supset SU(f)_L \otimes SU(f)_R \otimes SU(c)_L \otimes SU(c)_R \end{aligned}$	$\bar{G} = SL(2f, \mathbb{C}) \otimes SL(2c, \mathbb{C})$ $\supset SU(f)_L \otimes SU(f)_R \otimes SU(c)_L \otimes SU(c)_R$	$\bar{G} = SL(2f, \mathbb{C}) \otimes SL(2c, \mathbb{C})$ $\supset SU(f)_L \otimes SU(f)_R \otimes SU(c)_L \otimes SU(c)_R$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1045 (15.2.2)	343	■ 1.000 ● N +0	$\bar{L}(M) := M^4 \times \bar{G} \supset M^4 \times SL(2, \mathbb{C})$	$\bar{L}(M) := M^4 \times \bar{G} \supset M^4 \times SL(2, \mathbb{C})$	$\bar{L}(M) := M^4 \times \bar{G} \supset M^4 \times SL(2, \mathbb{C})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1046 (15.2.3)	344	■ 1.000 ● N -1	$\bar{\gamma} = \bar{E}_j^\alpha(m) \gamma_\alpha \otimes dx^j = \bar{E}_j dx^j$	$\bar{\gamma} = \bar{E}_j^\alpha(m) \gamma_\alpha \otimes dx^j = \bar{E}_j dx^j$	$\bar{\gamma} = \bar{E}_j^\alpha(m) \gamma_\alpha \otimes dx^j = \bar{E}_j dx^j$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1048 (15.2.5)	344	■ 1.000 ● K +0	$\frac{1}{4} \text{Tr}(\bar{\gamma} \otimes_s \bar{\gamma}) = f_{ij} dx^i \otimes_s dx^j$	$\frac{1}{4} \text{Tr}(\bar{\gamma} \otimes_s \bar{\gamma}) = f_{ij} dx^i \otimes_s dx^j$	$\frac{1}{4} \text{Tr}(\bar{\gamma} \otimes_s \bar{\gamma}) = f_{ij} dx^i \otimes_s dx^j$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1052 (15.2.9)	345	■ 1.000 ● N +0	$\bar{\omega}_i = \frac{1}{2} \Gamma_i^{(f) \alpha \beta j} \sigma_{\alpha \beta} \lambda_j^{(f)} \oplus \frac{1}{2} \Gamma_i^{(c) \alpha \beta j} \sigma_{\alpha \beta} \lambda_j^{(c)} \oplus A_i.$	$\bar{\omega}_i = \frac{1}{2} \Gamma_i^{(f) \alpha \beta j} \sigma_{\alpha \beta} \lambda_j^{(f)} \oplus \frac{1}{2} \Gamma_i^{(c) \alpha \beta j} \sigma_{\alpha \beta} \lambda_j^{(c)} \oplus A_i.$	$\bar{\omega}_i = \frac{1}{2} \Gamma_i^{(f) \alpha \beta j} \sigma_{\alpha \beta} \lambda_j^{(f)} \oplus \frac{1}{2} \Gamma_i^{(c) \alpha \beta j} \sigma_{\alpha \beta} \lambda_j^{(c)} \oplus A_i.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ1053 (15.2.10)	345	■ 1.000 ● N +0	$\backslash\mathrm{bar}\{\Theta\}:=\backslash\mathrm{bar}\{D\} \ \backslash\mathrm{bar}\{\gamma\}=d \ \backslash\mathrm{bar}\{\gamma\}-[\backslash\mathrm{bar}\{\omega\}, \ \backslash\mathrm{bar}\{\gamma\}]$	$\bar{\Theta} := \bar{D}\bar{\gamma} = d\bar{\gamma} - [\bar{\omega}, \bar{\gamma}]$	$\overline{\Theta} := \overline{D\gamma} = d\overline{\gamma} - [\overline{\omega}, \overline{\gamma}]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ1054 (15.2.11)	345	■ 1.000 ● W +1	$\backslash\mathrm{bar}\{\Omega\}=d \ \backslash\mathrm{bar}\{\omega\}-\backslash\mathrm{bar}\{\omega\} \ \wedge \ \backslash\mathrm{bar}\{\omega\}$	$\bar{\Omega} = d\bar{\omega} - \bar{\omega} \wedge \bar{\omega}$	$\overline{\Omega} = d\overline{\omega} - \overline{\omega} \wedge \overline{\omega}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ1056 (15.2.13)	346	■ 1.000 ● N +0	$\backslash\mathrm{psi}=\backslash\mathrm{left}\{\backslash\mathrm{psi}^{\backslash\mathrm{left}\{q_{\mathrm{f}}\}, \ q_{\mathrm{c}}\}\backslash\mathrm{right}\} \ \backslash\mathrm{mid} \ q_{\mathrm{f}}=1, \ \backslash\mathrm{ldots}, \ \mathrm{f} \ ; \ q_{\mathrm{c}}=1, \ \backslash\mathrm{ldots}, \ \mathrm{c}\backslash\mathrm{right}\}$.	$\psi = \left\{ \psi^{(q_f, q_c)} \mid q_f = 1, \dots, f; q_c = 1, \dots, c \right\}.$	$\psi = \{\psi^{(q_f, q_c)} q_f = 1, \dots, f; q_c = 1, \dots, c\}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ1061 (16.1.2)	354	■ 1.000 ● C -3	$B=\frac{2 \ \mathrm{hbar} \ c}{\ell^{*2}} \ \Lambda_{\mathrm{eff}}=-\frac{\frac{3}{\kappa}\{\frac{\mathrm{hbar} \ c}{\ell^{*4}}\}}{2}$	$B = \frac{2\hbar c}{\ell^{*2}} \Lambda_{\mathrm{eff}} = -\frac{3\kappa}{2} \frac{\hbar c}{\ell^{*4}}$	$B = \frac{2\hbar c}{\ell^{*2}} \Lambda_{\mathrm{eff}} = -\frac{3\kappa}{2} \frac{\hbar c}{\ell^{*4}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library_ EQ1063 (16.2.2)	355	■ 1.000 ● N +0	$\backslash\mathrm{square}=\frac{1}{\sqrt{ f }} \ \backslash\mathrm{partial}_{\mathrm{i}}\backslash\mathrm{left}\{\mathrm{f}^{\mathrm{i} \ \mathrm{j}} \ \backslash\mathrm{sqrt}\{ f \} \ \backslash\mathrm{partial}_{\mathrm{j}}\backslash\mathrm{right}\}$	$\square = \frac{1}{\sqrt{ f }} \partial_i \left(f^{ij} \sqrt{ f } \partial_j \right)$	$\square = \frac{1}{\sqrt{ f }} \partial_i (f^{ij} \sqrt{ f } \partial_j)$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1064 (16.2.3)	355	1.000 ● N +0	$\varphi = m \frac{c}{\hbar} \frac{1}{\rho} F_{nm}^l(\rho^*) Y_l^m(\vartheta, \phi) e^{-i t \omega m c^2 / \hbar} \cdot \left(\rho^* \right) Y_{\iota}^m(\vartheta, \phi) e^{-i t \omega m c^2 / \hbar} \cdot$	$\varphi = m \frac{c}{\hbar} \frac{1}{\rho} F_{nm}^l(\rho^*) Y_l^m(\vartheta, \phi) e^{-i t \omega m c^2 / \hbar} \cdot$	$\varphi = m \frac{c}{\hbar} \frac{1}{\rho} F_{nm}^l(\rho^*) Y_l^m(\vartheta, \phi) e^{-i t \omega m c^2 / \hbar} \cdot$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1067 (16.2.6)	355	1.000 ● K +0	$\beta = \frac{\sqrt{2\pi\kappa}}{\ell^*} / \frac{mc}{\hbar} = \frac{\ell_m}{\ell} \cdot$	$\beta = \frac{\sqrt{2\pi\kappa}}{\ell^*} / \frac{mc}{\hbar} = \frac{\ell_m}{\ell} \cdot$	$\beta = \frac{\sqrt{2\pi\kappa}}{\ell^*} / \frac{mc}{\hbar} = \frac{\ell_m}{\ell} \cdot$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1072 (16.3.1)	358	1.000 ● N +0	$\operatorname{Tr} \bar{Q}^2 = -\frac{4b}{\alpha_e} Y + I(I+1) - \frac{1}{4} Y^2 \cdot$	$\operatorname{Tr} \bar{Q}^2 = -\frac{4b}{\alpha_e} Y + I(I+1) - \frac{1}{4} Y^2 \cdot$	$\operatorname{Tr} \bar{Q}^2 = -\frac{4b}{\alpha_e} Y + I(I+1) - \frac{1}{4} Y^2 \cdot$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1073 (16.3.2)	358	1.000 ● N +0	$Q_e = I_3 + \frac{1}{2} Y - \frac{2}{3} C \cdot$	$Q_e = I_3 + \frac{1}{2} Y - \frac{2}{3} C \cdot$	$Q_e = I_3 + \frac{1}{2} Y - \frac{2}{3} C \cdot$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1074 (16.3.3)	359	1.000 ● N +0	$\frac{M^2}{M^{*2}} = \left\{ 1 - bY + \frac{\alpha_e}{4} \left[I(I+1) - \frac{1}{4} Y^2 \right] \right\}^2 + \frac{1}{4} J(J+1)$	$\frac{M^2}{M^{*2}} = \left\{ 1 - bY + \frac{\alpha_e}{4} \left[I(I+1) - \frac{1}{4} Y^2 \right] \right\}^2 + \frac{1}{4} J(J+1)$	$\frac{M^2}{M^{*2}} = \left\{ 1 - bY + \frac{\alpha_e}{4} \left[I(I+1) - \frac{1}{4} Y^2 \right] \right\}^2 + \frac{1}{4} J(J+1)$
Conf. MathPix confidence: ≥0.9 0.5–0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1075 (16.3.4)	360	■ 1.000 ● N +0	$T_{\bullet}=\frac{2\hbar c}{S_Ak_B}\left(r_+-\frac{M}{M^{*2}}\frac{\hbar}{c}\right)-\frac{\frac{M}{M^{*2}}\frac{\hbar}{c}}{\frac{2\hbar c}{S_Ak_B}}$.	$T_{\bullet}=\frac{2\hbar c}{S_Ak_B}\left(r_+-\frac{M}{M^{*2}}\frac{\hbar}{c}\right).$	$T_{\bullet}=\frac{2\hbar c}{S_Ak_B}\left(r_+-\frac{M}{M^{*2}}\frac{\hbar}{c}\right).$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1076 (A.1)	364	■ 1.000 ● N +0	$x^{\{A\}}y_{\{A\}}:=\sum_{A=0}^n\sum_{B=0}^nx^{\{A\}}y^{\{B\}}g_{\{A\}B}$.	$x^Ay_A:=\sum_{A=0}^n\sum_{B=0}^nx^Ay^Bg_{AB}.$	$x^Ay_A:=\sum_{A=0}^n\sum_{B=0}^nx^Ay^Bg_{AB}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1077 (A.2)	364	■ 1.000 ● N +0	$c=\hbar=1$.	$c=\hbar=1.$	$c=\hbar=1.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1080 (A.5)	365	■ 1.000 ● W +0	$\left\{\sigma^i,\sigma^j\right\}=g^{ij}$	$\left\{\sigma^i,\sigma^j\right\}=g^{ij}$	$\left\{\sigma^i,\sigma^j\right\}=g^{ij}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1081 (A.6)	365	■ 1.000 ● K +0	$\left(\gamma^{\alpha}\right)^{\mathrm{T}}:=\left(\gamma^{\{0\}},\gamma^{\{k\}}\right),$	$\left(\gamma^{\alpha}\right)^{\mathrm{T}}:=\left(\gamma^0,\gamma^k\right),$	$\left(\gamma^{\alpha}\right)^{\mathrm{T}}:=\left(\gamma^0,\gamma^k\right),$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1083 (A.8)	365	■ 1.000 ● N +0	$\backslash\gamma^{\alpha}\backslash\gamma^{\beta}+\backslash\gamma^{\beta}\backslash\gamma^{\alpha}\backslash\gamma^{\alpha}\backslash\gamma^{\beta}=2\ g^{\alpha}\backslash\gamma^{\beta}$	$\gamma^{\alpha}\gamma^{\beta}+\gamma^{\beta}\gamma^{\alpha}=2g^{\alpha\beta}$	$\gamma^{\alpha}\gamma^{\beta}+\gamma^{\beta}\gamma^{\alpha}=2g^{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1084 (A.9)	365	■ 1.000 ● N +0	$\backslash\gamma^{\{0+\}}=\backslash\gamma^{\{0\}},\quad\backslash\gamma^{\{k+\}}=-\backslash\gamma^{\{k\}},\quad\backslash\gamma^{\{5+\}}=\backslash\gamma^{\{5\}}\ .$	$\gamma^{0+}=\gamma^0,\quad\gamma^{k+}=-\gamma^k,\quad\gamma^{5+}=\gamma^5.$	$\gamma^{0+}=\gamma^0,\quad\gamma^{k+}=-\gamma^k,\quad\gamma^{5+}=\gamma^5.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1087 (A.14)	366	■ 1.000 ● K +0	$C(1,3)=C(0,2)\otimes C(1,1)\ .$	$C(1,3)=C(0,2)\otimes C(1,1).$	$C(1,3)=C(0,2)\otimes C(1,1).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1091 (B.1)	368	■ 1.000 ● N +0	$\backslash\alpha^{\{p\}}=\frac{1}{p!}\ A_{\backslash\alpha_1\{1\}}\ \backslash\cdots\ \backslash\alpha_{\{p\}}\ \backslash\vartheta^{\{\backslash\alpha_1\{1\}\}}\ \backslash\wedge\ \backslash\cdots\ \backslash\wedge\ \backslash\vartheta^{\{\backslash\alpha_p\{p\}\}}\ \backslash\in\ C^{\{\infty\}}\backslash\left(\backslash\wedge^{\{p\}}\ T^{\{*\}}(M)\backslash\right)\ .$	$\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^{\infty}(\wedge^pT^*(M)).$	$\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^{\infty}(\wedge^pT^*(M)).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1092 (B.2)	368	■ 1.000 ● W +0	$\backslash\begin{gathered}\backslash\alpha^{\{p\}}\ \backslash\wedge\ \backslash\beta^{\{(q)\}}:=\frac{1}{(p!)\cdot(q!)}\ A_{\backslash\alpha_1\{1\}}\ \backslash\cdots\ \backslash\alpha_{\{p\}}\ \backslash\cdots\ \backslash\beta_{\{1\}}\ \backslash\cdots\ \backslash\beta_{\{q\}}\ \backslash\vartheta^{\{\backslash\alpha_1\{1\}\}}\ \backslash\wedge\ \backslash\cdots\ \backslash\wedge\ \backslash\vartheta^{\{\backslash\alpha_p\{p\}\}}\ \backslash\backslash\ \backslash\wedge\ \backslash\vartheta^{\{\backslash\beta_1\{1\}\}}\ \backslash\wedge\ \backslash\cdots\ \backslash\wedge\ \backslash\vartheta^{\{\backslash\beta_q\{q\}\}}\ \backslash\end{gathered}$	$\alpha^{(p)}\wedge\beta^{(q)}:=\frac{1}{(p!)\cdot(q!)}A_{\alpha_1\cdots\alpha_p}\cdot B_{\beta_1\cdots\beta_q}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\wedge\vartheta^{\beta_1}\wedge\cdots\wedge\vartheta^{\beta_q}$	$\alpha^{(p)}\wedge\beta^{(q)}:=\frac{1}{(p!)\cdot(q!)}A_{\alpha_1\cdots\alpha_p}\cdot B_{\beta_1\cdots\beta_q}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\wedge\vartheta^{\beta_1}\wedge\cdots\wedge\vartheta^{\beta_q}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1093 (B.3)	368	■ 1.000 ● N +0	$\backslash\alpha^{\{p\}}\wedge\backslash\beta^{\{q\}}=(-1)^{p\ q}\backslash\beta^{\{q\}}\backslash\wedge\backslash\alpha^{\{p\}},$	$\alpha^{(p)}\wedge\beta^{(q)}=(-1)^{pq}\beta^{(q)}\wedge\alpha^{(p)},$	$\alpha^{(p)}\wedge\beta^{(q)}=(-1)^{pq}\beta^{(q)}\wedge\alpha^{(p)},$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1094 (B.4)	368	■ 1.000 ● N +0	$D(M):=\sum_{p=0}^n\backslash\wedge^pT^*(M)$	$D(M):=\sum_{p=0}^n\wedge^pT^*(M)$	$D(M):=\sum_{p=0}^n\wedge^pT^*(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1095 (B.5)	369	■ 1.000 ● N +1	$d\left(\backslash\alpha^{\{p\}}\wedge\backslash\beta^{\{q\}}\right)=d\backslash\alpha^{\{p\}}\backslash\wedge\backslash\beta^{\{q\}}+(-1)^p\backslash\alpha^{\{p\}}\backslash\wedge d\backslash\beta^{\{q\}}$	$d\left(\alpha^{(p)}\wedge\beta^{(q)}\right)=d\alpha^{(p)}\wedge\beta^{(q)}+(-1)^p\alpha^{(p)}\wedge d\beta^{(q)}$	$d(\alpha^{(p)}\wedge\beta^{(q)})=d\alpha^{(p)}\wedge\beta^{(q)}+(-1)^p\alpha^{(p)}\wedge d\beta^{(q)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1097 (B.7)	369	■ 1.000 ● N +0	$\backslash\sigma:M\rightarrowtail N,\quad\backslash\sigma\in\mathscr{D}(M)$	$\sigma:M\rightarrow N,\quad\sigma\in\mathscr{D}(M)$	$\sigma:M\rightarrow N,\quad\sigma\in\mathscr{D}(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1098 (B.8)	369	■ 1.000 ● N +0	$\backslash\sigma_*:\backslash\wedge^pT^*(N)\rightarrowtail\backslash\wedge^pT^*(M)$	$\sigma_*:\wedge^pT^*(N)\rightarrow\wedge^pT^*(M)$	$\sigma_*:\wedge^pT^*(N)\rightarrow\wedge^pT^*(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1103 (B.15)	370	■ 1.000 ● K +0	$\{ \}^{\{* \}}\backslash\alpha^{\{p\}}=(-1)^{p(n+1)+s}\backslash\alpha^{\{p\}}\ .$	$**\alpha^{(p)}=(-1)^{p(n+1)+s}\alpha^{(p)}.$	$**\alpha^{(p)}=(-1)^{p(n+1)+s}\alpha^{(p)}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1105 (B.17)	370	■ 1.000 ● N +0	$(\alpha, \beta) := \int_M \alpha^{(p)} \wedge \beta^{(p)}$	$(\alpha, \beta) := \int_M \alpha^{(p)} \wedge \beta^{(p)}$	$(\alpha, \beta) := \int_M \alpha^{(p)} \wedge \beta^{(p)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1107 (B.19)	370	■ 1.000 ● N +0	$\square \alpha^{(p)} := d \alpha^{(p)} + \delta \alpha^{(p)}$	$\square \alpha^{(p)} := d \alpha^{(p)} + \delta \alpha^{(p)}$	$\square \alpha^{(p)} := d \alpha^{(p)} + \delta \alpha^{(p)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1108 (B.20)	371	■ 1.000 ● N +0	$\begin{aligned} (\alpha, \square \alpha) &= (\alpha, d \alpha) + (\alpha, \delta \alpha) \\ &= (\delta \alpha, \delta \alpha) + (d \alpha, d \alpha) \end{aligned}$	$(\alpha, \square \alpha) = (\alpha, d \alpha) + (\alpha, \delta \alpha)$	$(\alpha, \square \alpha) = (\alpha, d \alpha) + (\alpha, \delta \alpha)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1109 (B.21)	371	■ 1.000 ● K +0	$d \alpha = 0$	$d \alpha = 0$	$d \alpha = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1111 (B.23)	371	■ 1.000 ● N +0	$\beta^{(p)} = d \alpha^{(p-1)} \in C^\infty(B(M))$	$\beta^{(p)} = d \alpha^{(p-1)} \in C^\infty(B(M))$	$\beta^{(p)} = d \alpha^{(p-1)} \in C^\infty(B(M))$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ1112 (B.24)	371	■ 1.000 ● N +0	$H(M) := Z(M) / B(M)$	$H(M) := Z(M)/B(M)$	$H(M) := Z(M)/B(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ1113 (B.25)	371	■ 1.000 ● N +0	$H(M) = \sum_{p=0}^n H^p(M) := \sum_{p=0}^n Z^p(M) / B^p(M) \text{ .}$	$H(M) = \sum_{p=0}^n H^p(M) := \sum_{p=0}^n Z^p(M)/B^p(M).$	$H(M) = \sum_{p=0}^n H^p(M) := \sum_{p=0}^n Z^p(M)/B^p(M).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ1115 (B.27)	371	■ 1.000 ● N +0	$H^{\{n\}}(M) = \wedge^{\{n\}} T^{\{*\}}(M) / B^{\{n\}}(M) \text{ .}$	$H^n(M) = \wedge^n T^*(M)/B^n(M).$	$H^n(M) = \wedge^n T^*(M)/B^n(M).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ1116 (B.28)	371	■ 1.000 ● N +0	$H^{\{0\}}(M) = Z^{\{0\}}(M)$	$H^0(M) = Z^0(M)$	$H^0(M) = Z^0(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ1117 (B.29)	372	■ 1.000 ● N +0	$H^{\{0\}}(M) \approx \mathbb{R}$	$H^0(M) \approx \mathbb{R}$	$H^0(M) \approx \mathbb{R}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ ECKEHARD_W._MIELKE_ _Z-Library_ EQ1118 (B.30)	372	■ 1.000 ● N -1	$b_{\{p\}} := \operatorname{dim} H^{\{p\}}(M)$	$b_p := \dim H^p(M)$	$b_p := \dim H^p(M)$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1119 (B.31)	372	■ 1.000 ● N +1	$\mathrm{f}_{\mathrm{M}}(t)=\sum_{p=0}^n\mathrm{b}_{\mathrm{p}}\mathrm{t}^{\mathrm{p}}$	$\mathrm{f}_{\mathrm{M}}(t)=\sum_{p=0}^nb_pt^p$	$\mathrm{f}_{\mathrm{M}}(t)=\sum_{p=0}^nb_pt^p,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1120 (B.32)	372	■ 1.000 ● N +0	$\chi(M):=\mathrm{f}_{\mathrm{M}}(-1)=\sum_{p=0}^n(-1)^{\mathrm{p}}\mathrm{b}_{\mathrm{p}},$	$\chi(M):=\mathrm{f}_{\mathrm{M}}(-1)=\sum_{p=0}^n(-1)^pb_p,$	$\chi(M):=\mathrm{f}_{\mathrm{M}}(-1)=\sum_{p=0}^n(-1)^pb_p,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1121 (B.33)	372	■ 1.000 ● N +0	$\mathrm{H}^{\{0\}}\left(S^{\mathrm{n}}\right)=\mathrm{H}^{\mathrm{n}}\left(S^{\mathrm{n}}\right)\approx\mathrm{mathbb{R}}$	$\mathrm{H}^0(S^n)=H^n(S^n)\approx\mathbb{R}$	$\mathrm{H}^0(S^n)=H^n(S^n)\approx\mathbb{R}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1123 (B.35)	372	■ 1.000 ● N +0	$\mathrm{f}_{\mathrm{S}^{\mathrm{n}}}(t)=1+\mathrm{t}^{\mathrm{n}},$	$\mathrm{f}_{S^n}(t)=1+t^n,$	$\mathrm{f}_{S^n}(t)=1+t^n,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__ EQ1124 (B.36)	372	■ 1.000 ● N +0	$\chi\left(S^{\mathrm{n}}\right)=\left\{\begin{array}{l}0\text{ for }n=2\mathrm{k}+1,\\2\text{ for }n=2\mathrm{k},\end{array}\right.$	$\chi(S^n)=\left\{\begin{array}{l}0\text{ for }n=2k+1,\\2\text{ for }n=2k,\end{array}\right.$	$\chi(S^n)=\left\{\begin{array}{l}0\text{ for }n=2k+1,\\2\text{ for }n=2k,\end{array}\right.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1125 (B.37)	372	■ 1.000 ● N +0	$\Delta_p := \left\{ x \in E^n \mid x = \sum_{i=0}^p \lambda^i \eta_i, \lambda^i \geq 0, \sum_{i=0}^p \lambda^i = 1 \right\}.$	$\Delta_p := \left\{ x \in E^n \mid x = \sum_{i=0}^p \lambda^i \eta_i, \lambda^i \geq 0, \sum_{i=0}^p \lambda^i = 1 \right\}.$	$\Delta_p := \left\{ x \in E^n \mid x = \sum_{i=0}^p \lambda^i \eta_i, \lambda^i \geq 0, \sum_{i=0}^p \lambda^i = 1 \right\}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1127 (B.39)	373	■ 1.000 ● N +0	$\partial c = \sum_{i=0}^p (-1)^i c \left(\sum_{j=0}^{i-1} \lambda^j \eta_j + \sum_{j=i}^{p-1} \lambda^j \eta_{j+1} \right).$	$\partial c = \sum_{i=0}^p (-1)^i c \left(\sum_{j=0}^{i-1} \lambda^j \eta_j + \sum_{j=i}^{p-1} \lambda^j \eta_{j+1} \right).$	$\partial c = \sum_{i=0}^p (-1)^i c \left(\sum_{j=0}^{i-1} \lambda^j \eta_j + \sum_{j=i}^{p-1} \lambda^j \eta_{j+1} \right).$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1129 (B.41)	373	■ 1.000 ● N +0	$H_{\mathrm{p}}(M, \mathbb{R}) := Z_{\mathrm{p}}(M, \mathbb{R}) / B_{\mathrm{p}}(M, \mathbb{R})$	$H_{\mathrm{p}}(M, \mathbb{R}) := Z_{\mathrm{p}}(M, \mathbb{R}) / B_{\mathrm{p}}(M, \mathbb{R})$	$H_{\mathrm{p}}(M, \mathbb{R}) := Z_{\mathrm{p}}(M, \mathbb{R}) / B_{\mathrm{p}}(M, \mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1130 (B.42)	373	■ 1.000 ● N +1	$\int_{c_{p+1}} d\alpha^{(p)} = \int_{\partial c_{p+1}} \alpha^{(p)}.$	$\int_{c_{p+1}} d\alpha^{(p)} = \int_{\partial c_{p+1}} \alpha^{(p)}.$	$\int_{c_{p+1}} d\alpha^{(p)} = \int_{\partial c_{p+1}} \alpha^{(p)}.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1132 (B.44)	373	■ 1.000 ● K +0	$\langle d\alpha, c \rangle_{\mathrm{M}} = \langle \alpha, \partial c \rangle_{\mathrm{M}}$	$\langle d\alpha, c \rangle_{\mathrm{M}} = \langle \alpha, \partial c \rangle_{\mathrm{M}}$	$\langle d\alpha, c \rangle_{\mathrm{M}} = \langle \alpha, \partial c \rangle_{\mathrm{M}}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1133 (C.1)	375	■ 1.000 ● N +0	$\mathrm{I}_{\{j\}}:=\left.\frac{\partial g(\xi)}{\partial \xi^j}\right _{g=e}\in T_e(G)\approx \mathfrak{g}$	$I_j:=\left.\frac{\partial g(\xi)}{\partial \xi^j}\right _{g=e}\in T_e(G)\approx \mathfrak{g}$	$I_j:=\left.\frac{\partial g(\xi)}{\partial \xi^j}\right _{g=e}\in T_e(G)\approx \mathfrak{g}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1138 (C.6)	376	■ 1.000 ● N +0	$[\mathrm{AB}]:=\mathrm{A}^{\mathrm{i}}\mathrm{B}^{\mathrm{j}}\mathrm{c}_{\mathrm{ij}}^{\mathrm{k}}\mathrm{I}_{\mathrm{k}}.$	$[\mathrm{AB}]:=\mathrm{A}^{\mathrm{i}}\mathrm{B}^{\mathrm{j}}\mathrm{c}_{\mathrm{ij}}^{\mathrm{k}}\mathrm{I}_{\mathrm{k}}.$	$[\mathrm{AB}]:=\mathrm{A}^{\mathrm{i}}\mathrm{B}^{\mathrm{j}}\mathrm{c}_{\mathrm{ij}}^{\mathrm{k}}\mathrm{I}_{\mathrm{k}}.$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1139 (C.7)	376	■ 1.000 ● W +0	$\rho_{\mathrm{A}}^{\mathrm{C}}(g_1)\rho_{\mathrm{C}}^{\mathrm{B}}(g_2)=\rho_{\mathrm{A}}^{\mathrm{B}}(g_1g_2)$	$\rho_{\mathrm{A}}^{\mathrm{C}}(g_1)\rho_{\mathrm{C}}^{\mathrm{B}}(g_2)=\rho_{\mathrm{A}}^{\mathrm{B}}(g_1g_2)$	$\rho_{\mathrm{A}}^{\mathrm{C}}(g_1)\rho_{\mathrm{C}}^{\mathrm{B}}(g_2)=\rho_{\mathrm{A}}^{\mathrm{B}}(g_1g_2)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1140 (C.8)	376	■ 1.000 ● K +0	$[\mathrm{A},\mathrm{B}]\longrightarrow \rho([\mathrm{A},\mathrm{B}])=[\rho(\mathrm{A}),\rho(\mathrm{B})].$	$[\mathrm{A},\mathrm{B}]\longrightarrow \rho([\mathrm{A},\mathrm{B}])=[\rho(\mathrm{A}),\rho(\mathrm{B})].$	$[\mathrm{A},\mathrm{B}]\longrightarrow \rho([\mathrm{A},\mathrm{B}])=[\rho(\mathrm{A}),\rho(\mathrm{B})].$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1141 (C.9)	376	■ 1.000 ● K +0	$\operatorname{Ad} \mathrm{A}(\mathrm{B}):=[\mathrm{A},\mathrm{B}];\quad \mathrm{A},\mathrm{B}\in \mathfrak{g},$	$\operatorname{Ad} \mathrm{A}(\mathrm{B}):=[\mathrm{A},\mathrm{B}];\quad \mathrm{A},\mathrm{B}\in \mathfrak{g},$	$\operatorname{Ad} \mathrm{A}(\mathrm{B}):=[\mathrm{A},\mathrm{B}];\quad \mathrm{A},\mathrm{B}\in \mathfrak{g},$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke____Z-Library__EQ1142 (C.10)	376	■ 1.000 ● N +0	$(\operatorname{Ad} \mathrm{A})_{\{j\}\{ \}^{\mathrm{k}}}=\mathrm{c}_{\mathrm{ij}}^{\mathrm{k}}\mathrm{A}^{\mathrm{i}}.$	$(\operatorname{Ad} \mathrm{A})_j^k=c_{ij}^kA^i.$	$(\operatorname{Ad} \mathrm{A})_j^k=c_{ij}^kA^i.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1144 (C.12)	376	■ 1.000 ● K +0	$(A, B)=2\,N\,\operatorname{Tr}\{A\,B\}-2\,\operatorname{Tr}\{A\}\operatorname{Tr}\{B\}$	$(A, B) = 2N\operatorname{Tr}(AB) - 2\operatorname{Tr}(A)\operatorname{Tr}(B)$	$(A, B) = 2N\operatorname{Tr}(AB) - 2\operatorname{Tr}(A)\operatorname{Tr}(B)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1146 (C.14)	377	■ 1.000 ● N +1	$g=\exp^{\left(-\frac{i}{2}\,\xi^j\,\lambda_j\right)}\in\mathrm{SU}(N),\,j=1,\,\ldots,\,N^2-1\,.$	$g=\exp^{\left(-\frac{i}{2}\xi^j\lambda_j\right)}\in SU(N),j=1,\ldots,N^2-1.$	$g=\exp^{\left(-\frac{i}{2}\xi^j\lambda_j\right)}\in SU(N),j=1,\ldots,N^2-1.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1148 (C.17)	377	■ 1.000 ● W +0	$\lambda_i\,\lambda_j=\frac{2}{N}\,\delta_{ij}+\left(d_{ij}^k+i\,f_{ij}^k\right)\lambda_k\,.$	$\lambda_i\lambda_j=\frac{2}{N}\delta_{ij}+(d_{ij}^k+if_{ij}^k)\lambda_k.$	$\lambda_i\,\lambda_j=\frac{2}{N}\delta_{ij}+(d_{ij}^k+if_{ij}^k)\lambda_k.$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__ EQ1149 (C.18)	377	■ 1.000 ● N +0	$\operatorname{Tr}\{\left(\lambda_i\,\lambda_j\right)\}=2\,\delta_{ij}\,.$	$\operatorname{Tr}(\lambda_i\lambda_j)=2\delta_{ij}.$	$\operatorname{Tr}(\lambda_i\,\lambda_j)=2\delta_{ij}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Inline formulas (first occurrence)

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00001		—	A	A
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00002		—	$\operatorname{SL}(5, \mathbb{R})$	$\operatorname{SL}(5, \mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00003		—	$\mathbb{R}$	$\mathbb{R}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00004		—	β	β
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00005		—	$\mathrm{W}^{\pm}$	$\mathrm{W}^{\pm}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00006		—	Z^{o}	Z^{o}
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00007		—	1×10^{-12}	1×10^{-12}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00008		—	∂_j	∂_j
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00009		—	$D_j := \partial_j + i A_j$	$D_j := \partial_j + i A_j$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00010		—	A_j	A_j
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00011		—	$\psi \rightarrow e^{i\theta} \psi$	$\psi \rightarrow e^{i\theta} \psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00012		—	ψ	ψ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00013		—	$A_j \rightarrow A_j - \partial_j \theta$	$A_j \rightarrow A_j - \partial_j \theta$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00014		—	$\mathrm{n}>4$	$n > 4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00015		—	$\{ \}^{+}$	$+$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00016		—	$\mathrm{A}_{2}, \mathrm{~K}^{*}(1420)$	$A_2, K^*(1420)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00017		—	$\operatorname{SL}(2, \mathbb{C})$	$SL(2, \mathbb{C})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00018		—	$\operatorname{SU}(3)$	$SU(3)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00019		—	$\operatorname{SL}(2, \mathbb{C}) \subset \operatorname{SL}(2, \mathbb{C}) \otimes \operatorname{SU}(3) \subset \operatorname{SL}(6, \mathbb{C})$	$SL(2, \mathbb{C}) \subset SL(2, \mathbb{C}) \otimes SU(3) \subset SL(6, \mathbb{C})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00020		—	$\operatorname{SL}(6, \mathbb{C})$	$SL(6, \mathbb{C})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00021		—	$\ell:=\ell^* / \sqrt{\kappa}$	$\ell := \ell^* / \sqrt{\kappa}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00022		—	$\mathrm{G}_{\mathrm{S}}=\mathrm{G}_{\mathrm{N}} / \kappa$	$G_S = G_N / \kappa$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00023		—	$\sqrt{s}=540 \mathrm{GeV}$	$\sqrt{s} = 540\mathrm{GeV}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00024		—	$\mathrm{x}: \mathrm{M} \rightarrow \mathbb{R}^{\mathrm{n}}$	$\mathbf{x} : \mathbf{M} \rightarrow \mathbb{R}^{\mathbf{n}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00025		—	E^{n}	$E^{\mathbf{n}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00026		—	U_{i}	$\mathbf{U}_i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00027		—	π_i	π_i
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00028		—	$\mathbb{R}^{\mathrm{n}}$	$\mathbb{R}^n$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00029		—	$\mathrm{x}_{i=\pi_i} \circ \mathrm{x}, i=1, \ldots, \mathrm{n}$	$\mathbf{x}_i = \pi_i \circ \mathbf{x}, i = 1, \dots, \mathbf{n}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00030		—	S^{n}	$\mathbf{S}^n$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00031		—	$\mathrm{U}_{\mathrm{i}} \subset \mathrm{M}$	$\mathbf{U}_i \subset \mathbf{M}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00032		—	$\mathscr{D}(\mathrm{M})$	$\mathcal{D}(\mathbf{M})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00033		—	$\left(\mathrm{U}_{\mathrm{i}}, \mathrm{x}_{\mathrm{i}}\right)$	$(\mathbf{U}_i, \mathbf{x}_i)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00034		—	C^{∞}	$\mathbf{C}^\infty$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00035		—	C^{r}	C^{r}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00036		—	$\mathrm{E}^{\mathrm{n}(\mathrm{n}+1) / 2}$	$\mathrm{E}^{\mathrm{n}(\mathrm{n}+1) / 2}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00037		—	$\mathrm{N}=\mathrm{n}(3 \mathrm{n}+11) / 2$	$N=n(3 n+11) / 2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00038		—	$\mathrm{N}=\mathrm{n}(\mathrm{n}+1)(3 \mathrm{n}+11) / 2$	$N=n(n+1)(3 n+11) / 2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00039		—	$\mathbb{E}^4$	$\mathbb{E}^4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00040		—	$\mathrm{T}_{\mathrm{m}}(\mathrm{M})$	$\mathrm{T}_{\mathrm{m}}(\mathrm{M})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00041		—	$\mathrm{m} \in \mathrm{M}$	$\mathrm{m} \in \mathrm{M}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00042		—	$\mathrm{s}(\mathrm{t})$	s(t)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00043		—	$\mathrm{e}(\mathrm{m})$	e(m)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00044		—	$\mathrm{e}_{\alpha}(\mathrm{m}), \alpha=1, \ldots, \mathrm{n}$	$\mathbf{e}_{\alpha}(\mathbf{m}), \alpha = 1, \dots, \mathbf{n}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00045		—	$\partial_{\mathrm{i}}:=\partial / \partial \mathrm{x}^{\mathrm{i}}$	$\partial_i := \partial / \partial \mathbf{x}^i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00046		—	$\left[\mathrm{e}_{\mathrm{i}}(\mathrm{m}), \mathrm{e}_{\mathrm{j}}(\mathrm{m})\right]=\left[\partial_{\mathrm{i}}, \partial_{\mathrm{j}}\right]=0$	$[\mathbf{e}_i(\mathbf{m}), \mathbf{e}_j(\mathbf{m})] = [\partial_i, \partial_j] = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00047		—	$\mathrm{e}_{\alpha}^{\mathrm{i}}(\mathrm{m})$	$\mathbf{e}_{\alpha}^i(m)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00048		—	$\mathrm{T}(\mathrm{M})$	T(M)
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00049		—	$\mathrm{T}^{\ast}(\mathrm{M})=\bigcup_{\mathrm{m}\in\mathrm{M}}\mathrm{T}_{\mathrm{m}}^{\ast}(\mathrm{M})$	$T^*(M) = \bigcup_{m \in M} T_m^*(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00050		—	$\mathrm{T}_{\mathrm{m}}^{\ast}(\mathrm{M})$	$T_m^*(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00051		—	$\vartheta^{\alpha}, \alpha=1, \ldots, n$	$\vartheta^\alpha, \alpha = 1, \dots, n$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00052		—	ϑ^{α}	ϑ^α
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00053		—	$\mathrm{d}x^i$	$\mathrm{d}x^i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00054		—	$\mathrm{d}\vartheta^{\alpha}$	$\mathrm{d}\vartheta^\alpha$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00055		—	$\mathrm{d}x^j$	$\mathrm{d}x^j$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00056		—	$\mathrm{E}_{\cdot}^{\alpha}{}_{\{j\}}$	$E_{\cdot j}^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00057		—	e_{β}^i	e_{β}^i
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00058		—	$\otimes_{\mathrm{s}, \mathrm{a}}$	$\otimes_{s,a}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00059		—	$\vee$	$\vee$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00060		—	$\wedge$	$\wedge$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00061		—	$\mathrm{T}_{\{\alpha_1{}_1\} \cdots \alpha_{\mathrm{p}}\}}^{\beta_1 \cdots \beta_q}(\mathrm{m})$	$T_{\alpha_1 \cdots \alpha_p}^{\beta_1 \cdots \beta_q}(\mathbf{m})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00062		—	$\mathrm{T}_{\{\mathrm{p}\}}^{\{\mathrm{q}\}}(\mathrm{M})$	$T_p^q(\mathbf{M})$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00063		—	$\mathrm{L}(\mathrm{M})$	$L(M)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00064		—	$\mathrm{D}^{\{(\mathrm{p}, \mathrm{q})\}}$	$D^{(p,q)}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00065		—	$\rho^{\{(\mathrm{p}, \mathrm{q})\}}(\mathrm{GL}(\mathrm{n}, \mathbb{R}))$	$\rho^{(p,q)}(GL(n, \mathbb{R}))$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00066		—	$\otimes^{\{\mathrm{p} * \}} \mathbb{R}^{\{\mathrm{n}\}} \otimes^{\{\mathrm{q}\}} \mathbb{R}^{\{\mathrm{n}\}}$	$\otimes^{p*} \mathbb{R}^n \otimes^q \mathbb{R}^n$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00067		—	$\mathrm{g}_{\mu \nu}$	$\mathfrak{g}_{\mu\nu}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00068		—	$(\mathrm{p}, 0)$	$(p, 0)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00069		—	$\wedge := \otimes_{\mathbf{a}}$	$\wedge := \otimes_{\mathbf{a}}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00070		—	(1939,1957)	(1939,1957)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00071		—	H	$\mathcal{H}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00072		—	φ	φ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00073		—	$\varphi(\mathrm{m})$	$\varphi(\mathbf{m})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00074		—	V_{m}	$V_{\mathbf{m}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00075		—	π	π
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00076		—	$\mathrm{F}_{\mathrm{m}}:=-\pi^{-1}(\mathrm{\sim m})$	$\mathbf{F}_{\mathbf{m}} := -\pi^{-1}(\mathbf{m})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00077		—	$\pi^{-1}(\mathrm{U}) \approx \mathrm{U} \times \mathrm{F}_{\mathrm{m}}$	$\pi^{-1}(\mathrm{U}) \approx \mathrm{U} \times \mathrm{F}_{\mathrm{m}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00078		—	$\left(\mathrm{U} \times \mathrm{F}_{\mathrm{m}}, \mathrm{U}, \pi_{\mathrm{u}}\right)$	$(\mathrm{U} \times \mathrm{F}_{\mathrm{m}}, \mathrm{U}, \pi_{\mathrm{u}})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00079		—	$\mathrm{P}(\mathrm{M}, \mathrm{G}, \pi, \delta)$	$\mathrm{P}(\mathrm{M}, \mathrm{G}, \pi, \delta)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00080		—	$(\mathrm{P}, \mathrm{M}, \pi)$	$(\mathrm{P}, \mathrm{M}, \pi)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00081		—	$\mathrm{P}, \mathrm{G}, \delta$	$\mathrm{P}, \mathrm{G}, \delta$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00082		—	$\mathrm{U} \subset$	$\mathrm{U} \subset$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00083		—	$\iota: \mathrm{U} \times \mathrm{G} \rightarrow {}^{-1}(\mathrm{U})$	$\iota: \mathrm{U} \times \mathrm{G} \rightarrow {}^{-1}(\mathrm{U})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00084		—	G	G
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00085		—	$\mathrm{p}_2=\mathrm{g}\ \mathrm{p}_1$	$\mathbf{p}_2 = \mathbf{g}\mathbf{p}_1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00086		—	p_1	$\mathbf{p}_1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00087		—	p_2	$\mathbf{p}_2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00088		—	π, δ	π, δ
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00089		—	$\mathrm{P}\ \times \mathrm{F}$	$\mathbf{P} \times \mathbf{F}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00090		—	$\mathrm{V}=\mathrm{P}\ \times_{\mathrm{G}}\ \mathrm{F}$	$\mathbf{V} = \mathbf{P} \times_{\mathbf{G}} \mathbf{F}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00091		—	$\{ \}^{-1}(\mathrm{U}) \approx \mathrm{U} \times \mathrm{G}$	${}^{-1}(\mathrm{U}) \approx \mathrm{U} \times \mathrm{G}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00092		—	$\{ \}^{-1}\{ \}_{\mathrm{v}}(\mathrm{U}) \approx \mathrm{U} \times \mathrm{F}$	${}^{-1}_{\mathrm{v}}(\mathrm{U}) \approx \mathrm{U} \times \mathrm{F}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00093		—	$\{ \}^{-1} \pi_{\mathrm{v}}(\mathrm{U})$	${}^{-1} \pi_{\mathrm{v}}(\mathrm{U})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00094		—	$\rho: \mathrm{G} \rightarrow \mathrm{GL}(\mathrm{N}, \mathbb{C})$	$\rho: \mathrm{G} \rightarrow \mathrm{GL}(\mathrm{N}, \mathbb{C})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00095		—	$\mathrm{N} \times \mathrm{N}$	$\mathrm{N} \times \mathrm{N}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00096		—	$\mathbb{C}^{\mathrm{N}}$	$\mathbb{C}^{\mathrm{N}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00097		—	$\mathrm{T}_{\mathbb{C}}^*(\mathrm{M})$	$\mathrm{T}_{\mathbb{C}}^*(\mathrm{M})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00098		—	ρ	ρ
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00099		—	$\mathrm{N} \geq 1$	$N \geq 1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00100		—	$\mathrm{b}=\left(\mathrm{b}_{\mathrm{A}}\right): \mathrm{U} \subset \mathrm{M} \rightarrow$	$\mathbf{b} = (\mathbf{b}_A) : \mathcal{U} \subset \mathcal{M} \rightarrow$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00101		—	V^{ρ}	V^{ρ}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00102		—	$\rho: \mathrm{G} \rightarrow \operatorname{GL}(N, \mathbb{C})$	$\rho: \mathcal{G} \rightarrow \operatorname{GL}(N, \mathbb{C})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00103		—	$\rho_{\mathrm{A}}^{\mathrm{B}}(\mathbf{g}(\xi))$	$\rho_A^{\mathbf{B}}(\mathbf{g}(\xi))$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00104		—	$\xi^j(j=1, \ldots, \operatorname{dim} \mathrm{G})$	$\xi^j(j = 1, \dots, \dim \mathcal{G})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00105		—	$\rho=\mathrm{id}$	$\rho = \mathrm{id}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00106		—	I_{j}	I_{j}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00107		—	$\mathfrak{g}$	$\mathfrak{g}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00108		—	$\mathrm{c}_{\mathrm{ij}}\{\}^{\mathrm{k}}$	$\mathrm{c}_{\mathrm{ij}}^{\mathrm{k}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00109		—	V^{Ad}	V^{Ad}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00110		—	$\mathrm{s}(\mathrm{t}) \subset \mathrm{M}$	$\mathrm{s}(\mathrm{t}) \subset \mathrm{M}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00111		—	$\mathrm{T}_{\mathrm{p}}\{\}^{\ast}(\mathrm{P})$	$\mathrm{T}_{\mathrm{p}}^{\ast}(\mathrm{P})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00112		—	$\mathrm{p} \in \mathrm{P}$	$\mathrm{p} \in \mathrm{P}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00113		—	$\mathrm{T}_{\mathrm{p}}^*(\mathrm{P}) = \mathrm{H}_{\mathrm{p}} \oplus \mathrm{V}_{\mathrm{p}}$	$\mathrm{T}_{\mathrm{p}}^*(\mathrm{P}) = \mathrm{H}_{\mathrm{p}} \oplus \mathrm{V}_{\mathrm{p}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00114		—	$\mathrm{F}_{\mathrm{m}}(\mathrm{M})$	$\mathrm{F}_{\mathrm{m}}(\mathrm{M})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00115		—	$\omega(\mathbf{e}) \in \mathcal{C}$	$\omega(\mathbf{e}) \in \mathcal{C}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00116		—	$\mathbf{e}_{\mathrm{v}} = \mathbf{pdg} \in \mathrm{V}_{\mathrm{p}}$	$\mathbf{e}_{\mathrm{v}} = \mathbf{pdg} \in \mathrm{V}_{\mathrm{p}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00117		—	$\bigwedge^{\mathrm{p}} \mathrm{T}_{\mathbb{C}}^*(\mathrm{M}) \otimes V^{\rho}$	$\bigwedge^{\mathrm{p}} \mathrm{T}_{\mathbb{C}}^*(\mathrm{M}) \otimes V^{\rho}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00118		—	$\mathrm{T}_{\mathbb{C}}^*(\mathrm{M}) \otimes V^{\rho}$	$\mathrm{T}_{\mathbb{C}}^*(\mathrm{M}) \otimes V^{\rho}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00119		—	$V^{\{\rho\}}$	V^{ρ}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00120		—	$\mathbf{b}$	b
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00121		—	$\mathrm{b}_{\{\mathrm{A}\}}, \mathrm{A}=1, \ldots, \mathrm{N}$	$\mathbf{b}_A, A = 1, \dots, N$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00122		—	$\mathrm{V}_{\{\mathbf{U}\}^{\rho}}$	V_{U}^{ρ}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00123		—	$\mathrm{b}_{\{\mathrm{A}\}}$	$\mathbf{b}_A$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00124		—	$\left[\omega_{\{\mathrm{A}\}}^{\{\mathrm{B}\}}\right]$	$\left[\omega_{\mathrm{A}}^{\mathrm{B}}\right]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00125		—	$\Gamma_{\alpha}^{\{\mathrm{j}\}}(\mathrm{m})$	$\Gamma_{\alpha}^{\mathbf{j}}(\mathbf{m})$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00126		—	$\mathrm{dd} \equiv 0$	$\mathrm{dd} \equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00127		—	$\partial \partial \equiv 0$	$\partial \partial \equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00128		—	Ω	Ω
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00129		—	$\mathrm{G}=\mathrm{U}(1)$	$\mathrm{G} = \mathrm{U}(1)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00130		—	$\circlearrowleft=\partial \mathrm{U} \subset \mathrm{M}$	$\circ = \partial \mathrm{U} \subset \mathrm{M}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00131		—	$\partial \mathrm{U}$	$\partial \mathrm{U}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00132		—	$\mathrm{e}(\mathrm{m}) \in \mathrm{T}_{\mathrm{m}}(\mathrm{M})$	$\mathrm{e}(\mathrm{m}) \in \mathrm{T}_{\mathrm{m}}(\mathrm{M})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00133		—	$\mathrm{G}(\mathrm{p})$	$G(p)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00134		—	$\mathrm{F}_{\mathrm{m}}=\pi^{-1}(\mathrm{~m})$	$F_m = \pi^{-1}(\text{ m})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00135		—	$\mathscr{G}_{\mathrm{P}}$	$\mathcal{G}_P$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00136		—	$\exp : \mathfrak{g} \rightarrow \mathrm{G}$	$\exp : \mathfrak{g} \rightarrow G$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00137		—	$P \times_{\mathrm{Ad}} \mathrm{G}$	$P \times_{\mathrm{Ad}} G$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00138		—	$\theta^k(m) \in C^{\infty}(M)$	$\theta^k(m) \in C^\infty(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00139		—	$\mathrm{V}^{\rho}\left(\mathrm{M}, \mathbb{C}^{\mathrm{N}}, \mathrm{GL}(\mathrm{N}, \mathbb{C}), \mathrm{P}\right)$	$V^\rho\left(M, \mathbb{C}^N, \mathrm{GL}(N, \mathbb{C}), P\right)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00140		—	$\mathrm{scr}{G}_{\mathrm{V}}$	$\mathcal{G}_V$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00141		—	L^{k}	$\mathbf{L}^k$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00142		—	$\rho: \mathrm{G} \rightarrow \operatorname{GL}(\mathrm{N}, \mathbb{C})$	$\rho: \mathbf{G} \rightarrow \mathrm{GL}(\mathbf{N}, \mathbb{C})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00143		—	$\bar{\mathrm{varphi}}$	$\bar{\varphi}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00144		—	$\bar{\rho}$	$\bar{\rho}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00145		—	$\mathrm{V}\left(\mathrm{M}, \mathbb{C}^{\mathrm{N}}, \mathrm{GL}(\mathrm{N}, \mathbb{C}), \mathrm{P}\right)$	$\mathbf{V}\left(\mathbf{M}, \mathbb{C}^{\mathbf{N}}, \mathrm{GL}(\mathbf{N}, \mathbb{C}), \mathbf{P}\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00146		—	γ_{k}	$\gamma_{\mathbf{k}}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00147		—	$\mathrm{k} \leq n / 2$	$k \leq n/2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00148		—	ω	ω
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00149		—	k^{th}	k^{th}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00150		—	$\mathrm{c}_{\mathrm{k}}(\mathrm{V})$	$c_k(V)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00151		—	$V^{\{\mathbb{R}\}}$	$V^{\mathbb{R}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00152		—	$\mathbb{R}^{\{\mathrm{N}\}}$	$\mathbb{R}^N$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00153		—	$\mathbb{C}^{\{N\}}$	$\mathbb{C}^N$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00154		—	V^{R}	$V^{\mathbb{R}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00155		—	$\mathrm{s}=0$	$s = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00156		—	$\mathrm{P}\backslash\left(\mathrm{S}^{\mathrm{4}},\mathrm{SU}(\mathrm{f}),\pi,\delta\right)$	$P\left(S^4,\mathrm{SU}(\mathrm{f}),\pi,\delta\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00157		—	$\operatorname{det}\mathrm{G}=1$	$\det G = 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00158		—	$\operatorname{Tr}(\backslash 0\mathrm{omega}\wedge\backslash 0\mathrm{omega})$	$\mathrm{Tr}(\Omega\wedge\Omega)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00159		—	$\stackrel{*}{\mathrm{pi}}\backslash\left(\gamma_2\right)$	$\overset{*}{\pi}\left(\gamma_2\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00160		—	$\mathrm{G}=\mathrm{SU}(\mathrm{f})$	$G = \mathrm{SU}(\mathrm{f})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00161		—	$\mathrm{SU}(2)$	$\mathrm{SU}(2)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00162		—	$\widetilde{\mathrm{SO}}(4) \approx \mathrm{SU}(2) \otimes \mathrm{SU}(2)$	$\mathrm{SO}(4) \approx \mathrm{SU}(2) \otimes \mathrm{SU}(2)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00163		—	γ_2	γ_2
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00164		—	$\mathrm{F}_{\alpha\beta}^j$	$\mathrm{F}_{\alpha\beta}^j$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00165		—	$ \mathrm{x} ^{-2}$	$ \mathbf{x} ^{-2}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00166		—	$ x \rightarrow \infty$	$ x \rightarrow \infty$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00167		—	$\stackrel{\infty}{\omega}$	$\stackrel{\infty}{\omega}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00168		—	$\partial \mathrm{M}$	∂M
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00169		—	$\mathrm{G}=\mathrm{SU}(2)$	$G = \mathrm{SU}(2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00170		—	S^3	S^3
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00171		—	$\mathrm{c}_2(\mathrm{M})$	$c_2(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00172		—	$\mathrm{S}^3 \approx \mathrm{SU}(2)$	$S^3 \approx \mathrm{SU}(2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00173		—	n^{th}	n^{th}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00174		—	$\mathbb{Z}$	$\mathbb{Z}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00175		—	$\pi_n\left(S^n\right)=\mathbb{Z}$	$\pi_n\left(S^n\right)=\mathbb{Z}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00176		—	$\mathrm{S}^{\mathrm{n}} \rightarrow \mathrm{S}^{\mathrm{n}}$	$\mathrm{S}^{\mathrm{n}} \rightarrow \mathrm{S}^{\mathrm{n}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00177		—	$\delta \varphi$	$\delta \varphi$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00178		—	$\varphi+\delta \varphi$	$\varphi+\delta \varphi$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00179		—	$(n-1)$	$(n-1)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00180		—	$d \varphi$	$d \varphi$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00181		—	δ	δ
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00182		—	s	s
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00183		—	$d\,d\,\equiv 0$	$dd \equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00184		—	x^i	$\mathbf{x}^i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00185		—	$\varphi \rightarrow \psi := \left(\varphi, \Theta^{\{1\}} := d\,\varphi\right)$	$\varphi \rightarrow \psi := \left(\varphi, \Theta^{(1)} := d\varphi\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00186		—	$\mathscr{D}(M)$	$\mathscr{D}(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00187		—	$\mathscr{G}$	$\mathscr{G}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00188		—	$L_{\{\text{mat}\}}$	L_{mat}
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00189		—	$\bar{\varphi} \, \mathit{\text{scr}}{0} \, \varphi$	$\bar{\varphi} \varphi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00190		—	$\mathit{\text{scr}}{0}$	$\varnothing$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00191		—	$\omega^{\{\mathrm{prime}\}}$	ω'
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00192		—	$\mathrm{GdG}^{\{-1\}}$	GdG^{-1}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00193		—	$\mathrm{G} \, \mathrm{in} \, \mathit{\text{scr}}{G}_{\{\mathrm{V}\}}$	$\mathrm{G} \in \mathcal{B}_V$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00194		—	$\{ \ }^{\{\mathrm{G}\}} \, \omega$	G_{ω}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00195		—	$\{ \ }^{\{\mathrm{G}\}} \, \mathrm{b}$	$\mathrm{G}_{\mathfrak{b}}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00196		—	G^{-1}	G^{-1}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00197		—	$\{ \}_{\sigma_{\alpha}}^{\alpha} \omega$	$\sigma_{\alpha}^* \omega$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00198		—	$G^{-1} \omega$	$G^{-1} \omega$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00199		—	$\bar{\omega}$	$\bar{\omega}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00200		—	$\bar{\Omega} = \mathrm{G}^{-1} \Omega \mathrm{G}$	$\bar{\Omega} = G^{-1} \Omega G$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00201		—	$d^* \omega \simeq 0$	$d^* \omega \simeq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00202		—	$A = A_i dx^i$	$A = A_i dx^i$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00203		—	F	F
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00204		—	$d\,d\,\Phi \equiv 0$	$dd\Phi \equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00205		—	Φ	Φ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00206		—	$d\,F \equiv 0$	$dF \equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00207		—	$F=d\,A$	$F = dA$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00208		—	$U(1)$	$U(1)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00209		—	$D \rightarrowtail \mathscr{D}:=D+i\,e\,A$	$D \rightarrow \mathscr{D}:=D+ieA$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00210		—	L	L
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00211		—	\Psi	Ψ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00212		—	S=\int L	$S = \int L$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00213		—	H	H
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00214		—	d d H \equiv 0	$ddH \equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00215		—	j	j
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00216		—	C:=A \wedge F=A \wedge d A	$C := A \wedge F = A \wedge dA$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00217		—	P	P
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00218		—	$\stackrel{\theta}{L}=-\left(\cos \theta F \wedge * F+\sin \theta F \wedge F\right) / 2$	$\stackrel{\theta}{L}=-\left(\cos \theta F \wedge * F+\sin \theta F \wedge F\right) / 2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00219		—	θ	θ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00220		—	*	*
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00221		—	g	g
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00222		—	η	η
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00223		—	$\partial \mathfrak{D} / \partial t$	$\partial \mathfrak{D} / \partial t$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00224		—	$\mathfrak{D}=\varepsilon \mathbf{E}$	$\mathfrak{D} = \varepsilon \mathbf{E}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00225		—	$\vartheta_{[\alpha} \wedge \Sigma_{\beta]}=0$	$\vartheta_{[\alpha} \wedge \Sigma_{\beta]} = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00226		—	$B=\frac{1}{2} B_{i j} d x^{i} \wedge d x^{j}$	$B = \frac{1}{2} B_{ij} dx^i \wedge dx^j$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00227		—	B	B
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00228		—	$d B \not\cong 0$	$dB \cong 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00229		—	$F:=d A \not\cong 0$	$F := dA \cong 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00230		—	$d C$	dC
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00231		—	$H=d\ B$	$H = dB$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00232		—	$\delta C / \delta A = dF \equiv 0$	$\delta dC / \delta A = dF \equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00233		—	$\psi=\psi_{\mathrm{i}}\,dx^{\mathrm{i}}$	$\psi = \psi_{\mathrm{i}}dx^{\mathrm{i}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00234		—	$s\,\psi=D\,\phi$	$s\psi = D\phi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00235		—	$s\,\phi=0$	$s\phi = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00236		—	$B\,F$	BF
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00237		—	$F=\{ \,\}^{\ast}\,B$	$F = {}^*B$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00238		—	$j:=\delta L_{\text{matter}}/\delta A$	$j := \delta L_{\text{matter}}/\delta A$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00239		—	$d\,j\,\cong 0$	$dj \cong 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00240		—	$d\,A$	dA
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00241		—	$V(B)$	$V(B)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00242		—	$d\,B+j\,\cong 0$	$dB + j \cong 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00243		—	$F\,\cong\,\partial V\,/\,\partial B$	$F \cong \partial V/\partial B$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00244		—	$-B\,\wedge\,F+B\,\wedge\,B\,\wedge\,B\,/\,3$	$-B \wedge F + B \wedge B \wedge B/3$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00245		—	$C:=A \wedge F-A \wedge A \wedge A / 3$	$C := A \wedge F - A \wedge A \wedge A/3$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00246		—	$F=-B \wedge B$	$F = -B \wedge B$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00247		—	$D \, B=D \, F \equiv 0$	$DB = DF \equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00248		—	$\Gamma_{\alpha}^{\mathrm{j}}(\mathrm{m})$	$\Gamma_{\alpha}^{\mathrm{J}}(\mathbf{m})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00249		—	$d \, \omega$	$d\omega$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00250		—	$\mathscr{F}$	$\mathcal{F}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00251		—	$\{ \}^{\ast} \mathscr{F}$	$\ast \mathcal{F}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00252		—	Π^{Ω}	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00253		—	$n-1$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00254		—	τ	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00255		—	$\delta L / \delta \omega$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00256		—	(1978,1980)	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00257		—	H^3	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00258		—	∂H^3	
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00259		—	ι	ι
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00260		—	$\mathrm{SU}(\mathrm{N})$	$\mathrm{SU}(\mathrm{N})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00261		—	$L\left(\mathrm{mathscr{F}},\{ \}^*\mathrm{mathscr{F}}\right)$	$L\left(\mathcal{F},*\mathcal{F}\right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00262		—	$\mathrm{I}^{\mathrm{j}}=\mathrm{i}\,\sigma^{\mathrm{j}}$	$\mathbf{I}^j = \mathrm{i}\sigma^j$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00263		—	$\mathrm{SU}(2)\,\times\,\mathrm{SU}(2)\,\approx$	$\mathrm{SU}(2)\times\mathrm{SU}(2)\approx$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00264		—	$\mathrm{S0}(4)$	$\mathrm{SO}(4)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00265		—	$\mathrm{h}=\mathrm{h}(\mathrm{x})$	$\mathbf{h} = \mathbf{h}(\mathbf{x})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00266		—	M^4	M^4
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00267		—	$\mathrm{k}=1$	$k = 1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00268		—	$5\ \mathrm{k}+4$	$5k + 4$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00269		—	$\mathbb{R}^4\ \cup\ \{\infty\}\ \approx\ \mathrm{S}^4$	$\mathbb{R}^4 \cup \{\infty\} \approx S^4$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00270		—	$ \cdot\infty\rangle$	$ \, -\, \infty\rangle$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00271		—	$\langle\infty $	$\langle\infty $
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00272		—	$\mathscr{M}_{\omega}$	$\mathcal{M}_{\omega}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00273		—	$\mathrm{M}_{\omega}=\mathrm{C} \ / \ \mathrm{G}_{\mathrm{p}}$	$\mathcal{M}_{\omega}=\mathcal{C}/\mathcal{G}_{\mathrm{p}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00274		—	$\mu[\omega]$	$\mu[\omega]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00275		—	S	S
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00276		—	$\operatorname{Spin}^c(4)$	$\operatorname{Spin}^c(4)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00277		—	$\psi=\psi_{\mathrm{L}}+\psi_{\mathrm{R}}=\mathrm{P}_{-}\psi+\mathrm{P}_{+}\psi$	$\psi=\psi_L+\psi_R=P_{-}\psi+P_{+}\psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00278		—	$\bar{\psi}$	$\bar{\psi}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00279		—	R	R
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00280		—	$\psi=0$	$\psi = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00281		—	$F^{\{ \text{ } \pm \}}$	$F^{\pm}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00282		—	$\mathrm{A}_{\alpha}^{\{\alpha\}\{\mathrm{j}\}}$	A_{α}^{J}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00283		—	$\varphi^{\{j\}}$	φ^j
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00284		—	$\Theta^{\{1\} \text{ } j\}$	$\Theta^{(1)j}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00285		—	$\phi^{\{(\emptyset)\}}$	$\phi^{(0)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00286		—	$\Theta^{\{1\}}$	$\Theta^{(1)}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00287		—	$\mathrm{A}_{\alpha}^{\mathrm{j}}(\mathrm{m})$	$A_{\alpha}^j(m)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00288		—	$\delta L_{\mathrm{H}} / \delta \bar{\phi}^{\{0\}}$	$\delta L_H / \delta \bar{\phi}^{(0)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00289		—	$\delta L_{\mathrm{H}} / \delta \bar{\Theta}^{\{1\}}$	$\delta L_H / \delta \bar{\Theta}^{(1)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00290		—	μ	μ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00291		—	$\varphi=0$	$\varphi = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00292		—	$ \varphi $	$ \varphi $
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00293		—	λ^{φ}	λ_{φ}
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00294		—	$\mathrm{\varphi}_{\mathrm{H}}:=\mathrm{\varphi}-\mathrm{\mu} \, / \, \sqrt{\mathrm{\lambda}}$	$\varphi_{\mathrm{H}} := \varphi - \mu / \sqrt{\lambda}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00295		—	$\mathrm{m}_{\mathrm{H}}=\sqrt{2} \, \, \mathrm{\mu}$	$\mathfrak{m}_{\mathrm{H}} = \sqrt{2} \mu$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00296		—	$\mathrm{G} \, / \, \mathrm{H}$	G / H
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00297		—	$\mathrm{v}_{\mathrm{0}} \, \in \, \mathrm{V}^{\mathrm{\rho}}$	$\mathfrak{v}_0 \in V^{\rho}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00298		—	$\stackrel{\circ}{\mathrm{\varphi}}_{\mathrm{min}}:=\mathrm{\sigma}\mathrm{\left(\mathrm{v}_{\mathrm{0}}\right)} \, \in \, \mathrm{C}^{\infty}\left(\mathrm{V}^{\mathrm{\rho}}\right)$	$\overset{\circ}{\varphi}_{\min} := \sigma \left(\mathfrak{v}_0\right) \in \mathrm{C}^{\infty}\left(\mathfrak{V}^{\rho}\right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00299		—	$\mathrm{a} \, \mathrm{g} \, \in \, \mathrm{G}$	$\mathfrak{a} \mathfrak{g} \in \mathrm{G}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00300		—	v_{0}	$\mathfrak{v}_0$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00301		—	$\mathrm{v}=\rho(\mathrm{g})\ \cdot\ \mathrm{v}_{\{0\}}$	$\mathbf{v}=\rho(\mathbf{g})\cdot\mathbf{v}_0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00302		—	$\stackrel{\circ}{\mathrm{rel}}{\mathrm{circ}}{\mathrm{\varphi}}_{\{\mathrm{text\ min}\}}=\sigma\mathrm{\left(\mathrm{v}_{\{0\}\mathrm{right}}\right)}$	$\overset{\circ}{\varphi}_{\min}=\sigma\left(\mathbf{v}_0\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00303		—	$\mathrm{P}(\mathrm{G},\mathrm{M},\pi,\delta)$	$\mathbf{P}(\mathbf{G},\mathbf{M},\pi,\delta)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00304		—	$\mathfrak{h}$	$\mathfrak{h}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00305		—	ω_{H}	ω_{H}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00306		—	Q	$\mathcal{Q}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00307		—	$\omega_{\{0\}}$	ω_0
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00308		—	ϕ	ϕ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00309		—	$\lambda \rightarrow 0$	$\lambda \rightarrow 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00310		—	$\stackrel{*}{\sigma} \vec{\omega}$	$\overset{*}{\sigma} \vec{\omega}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00311		—	(1981,1983)	(1981,1983)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00312		—	σ	σ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00313		—	$\mathbb{R}^2 \times \mathrm{S}^2$	$\mathbb{R}^2 \times \mathrm{S}^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00314		—	S^4	S^4
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00315		—	$\mathrm{N}=2$	$\mathbb{N} = 2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00316		—	$\ell^{\ast}$	ℓ^*
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00317		—	(1980,1981)	(1980,1981)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00318		—	$\mathrm{A}(\mathrm{n},\,\mathbb{R})$	$\mathbf{A}(\mathbf{n},\mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00319		—	A^{n}	$\mathbf{A}^{\mathbf{n}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00320		—	$\mathrm{a}=\left(\mathrm{a}^{\alpha}\right)\,\mathrm{in}\,\mathbb{R}^{\mathrm{n}}$	$\mathbf{a} = \left(\mathbf{a}^{\alpha}\right) \in \mathbb{R}^{\mathbf{n}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00321		—	$\mathrm{x}\,\,\mathrm{in}\,\,\mathrm{A}^{\mathrm{n}}$	$\mathbf{x} \in \mathbf{A}^{\mathbf{n}}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00322		—	$\tilde{x}:=\binom{x}{1} \in \mathbb{R}^{\mathrm{n}+1}$	$\tilde{x} := \binom{x}{1} \in \mathbb{R}^{\mathbf{n}+1}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00323		—	$\mathrm{n}+1$	$\mathbf{n} + 1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00324		—	$\tilde{\mathrm{x}}$	$\tilde{\mathbf{x}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00325		—	$\mathrm{GL}(\mathrm{n}+1, \mathbb{R})$	$\mathrm{GL}(\mathbf{n} + 1, \mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00326		—	$\mathrm{a} \in \mathbb{R}^{\mathrm{n}}$	$\mathbf{a} \in \mathbb{R}^{\mathbf{n}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00327		—	$\mathrm{g} \in \mathrm{GL}(\mathrm{n}, \mathbb{R})$	$\mathbf{g} \in \mathrm{GL}(\mathbf{n}, \mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00328		—	$\theta(1, \mathrm{n}-1) \subset \mathrm{GL}(\mathrm{n}, \mathbb{R})$	$0(1, \mathbf{n} - 1) \subset \mathrm{GL}(\mathbf{n}, \mathbb{R})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00329		—	$\mathfrak{a}(\mathrm{n}, \mathbb{R})$	$\mathfrak{a}(\mathbf{n}, \mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00330		—	$L_{\{\alpha \beta\}} \in \mathfrak{gl}(\mathrm{n}, \mathbb{R})$	$L_{\alpha\beta} \in \mathfrak{gl}(\mathbf{n}, \mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00331		—	$\mathrm{P}_{\{\alpha\}} \in \mathbb{R}^{\mathrm{n}}$	$\mathbf{P}_{\alpha} \in \mathbb{R}^{\mathbf{n}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00332		—	$\mathrm{GL}(\mathrm{n}, \mathbb{R})$	$\mathbf{GL}(\mathbf{n}, \mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00333		—	$\mathrm{L}_{\{\mathrm{m}\}}(\mathrm{M})$	$\mathbf{L}_{\mathbf{m}}(\mathbf{M})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00334		—	$\mathrm{U} \subset \mathrm{M}$	$\mathbf{U} \subset \mathbf{M}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00335		—	$\mathrm{e}_{\{\mathrm{a}\}}(\mathrm{m})$	$\mathbf{e}_{\mathbf{a}}(\mathbf{m})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00336		—	$\mathbb{R}^{\mathrm{n}^2}$	$\mathbb{R}^{\mathrm{n}^2}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00337		—	$\mathrm{e}_{\alpha^j}(\mathrm{~m})$	$\mathbf{e}_{\alpha}^j(\mathrm{~m})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00338		—	$n \times n$	$n \times n$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00339		—	$\left(\mathrm{x}^i, \mathrm{e}_{\alpha^{-j}}\right)$	$\left(\mathbf{x}^i, \mathbf{e}_{\alpha}^{-j}\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00340		—	$\mathrm{U} \times \mathrm{GL}(\mathrm{n}, \mathbb{R})$	$\mathbf{U} \times \mathrm{GL}(\mathbf{n}, \mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00341		—	$\pi^{-1}(\mathrm{U})$	$\pi^{-1}(\mathbf{U})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00342		—	$\operatorname{GL}(\mathrm{n})$	$\mathrm{GL}(\mathbf{n})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00343		—	ϑ	ϑ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00344		—	$\mathrm{n}=\operatorname{dim} \mathrm{M}$	$n = \dim M$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00345		—	$\mathbb{R}^n$	$\mathbb{R}^n$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00346		—	$\mathrm{A}(\mathrm{M}) / \mathbb{R}^n$	$A(M)/\mathbb{R}^n$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00347		—	$\mathrm{e}_{\alpha}(\mathrm{m})$	$e_{\alpha}(m)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00348		—	$\mathrm{T}=\mathbb{R}^n$	$T = \mathbb{R}^n$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00349		—	$\mathrm{A}(\mathrm{n}, \mathbb{R}) / \mathrm{GL}(\mathrm{n}, \mathbb{R})$	$A(n, \mathbb{R})/GL(n, \mathbb{R})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00350		—	$\mathrm{T}=\mathrm{P} \ / \ \mathrm{L}$	T = P/L
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00351		—	$\widetilde{\widetilde{\omega}}$	$\widetilde{\widetilde{\omega}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00352		—	$\chi: \mathrm{L}(\mathrm{L})(\mathrm{M}) \rightarrow \mathrm{0}_{\mathrm{m}} \times \mathrm{L}(\mathrm{M}) \subset \mathrm{A}(\mathrm{M})$	$\chi: \mathrm{L}(\mathrm{M}) \rightarrow 0_{\mathrm{m}} \times \mathrm{L}(\mathrm{M}) \subset \mathrm{A}(\mathrm{M})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00353		—	$\stackrel{*}{\chi} \ \widetilde{\omega}$	$\overset{*}{\chi} \ \widetilde{\omega}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00354		—	ω^{L}	ω^{L}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00355		—	$\mathfrak{gl}(\mathrm{n}, \mathbb{R})$	$\mathfrak{gl}(\mathfrak{n}, \mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00356		—	ω^{T}	ω^{T}
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00357		—	$\mathrm{H} \subset \mathrm{G}$	$\mathrm{H} \subset \mathrm{G}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00358		—	$\widetilde{\widetilde{\Omega}}$	$\widetilde{\widetilde{\Omega}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00359		—	$\left[\omega^{\mathrm{T}}, \omega^{\mathrm{T}}\right]$	$\left[\omega^{\mathrm{T}}, \omega^{\mathrm{T}}\right]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00360		—	$\omega^{\mathrm{T}}, \omega^{\mathrm{L}}$	$\omega^{\mathrm{T}}, \omega^{\mathrm{L}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00361		—	$\mathscr{A}(n, \mathbb{R})$	$\mathscr{A}(n, \mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00362		—	$C^{\{\infty\}}$	C^{∞}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00363		—	$\mathrm{GL}(n, \mathbb{R})$	$GL(n, \mathbb{R})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00364		—	$\mathscr{T}(n, \mathbb{R})$	$\mathcal{T}(n, \mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00365		—	$\operatorname{Diff}(n, \mathbb{R})$	$\operatorname{Diff}(n, \mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00366		—	$(n, \mathbb{R})$	$(n, \mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00367		—	$\left(n+n^2\right)$	$\left(n+n^2\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00368		—	$A(n, \mathbb{R})_{\mathrm{H}}$	$A(n, \mathbb{R})_{\mathrm{H}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00369		—	$\mathrm{P}_{\mathrm{i}}=\partial_{\mathrm{i}}:=\partial/\partial x^{\mathrm{i}}$	$P_i = \partial_i := \partial/\partial x^i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00370		—	$L^{\mathrm{i}}{ }_{\mathrm{j}}=x^{\mathrm{i}} \partial_{\mathrm{j}}$	$L^i{}_j = x^i \partial_j$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00371		—	χ	χ
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00372		—	V^{id}	V^{id}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00373		—	$\stackrel{\circ}{\varphi}=\sigma\left(0_{\mathrm{m}}\right)$	$\overset{\circ}{\tilde{\varphi}}=\sigma\left(0_{\mathrm{m}}\right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00374		—	$\mathrm{H}=\mathrm{GL}\left(\mathrm{n},\mathbb{R}\right)$	$\mathrm{H}=\mathrm{GL}(\mathrm{n},\mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00375		—	$\widetilde{\vartheta}$	$\widetilde{\vartheta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00376		—	$\mathscr{A}_{\mathrm{p}}$	$\mathscr{A}_{\mathrm{p}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00377		—	$\mathrm{D}=(\stackrel{*}{\chi}\widetilde{\mathrm{D}})$	$\mathrm{D}=(\overset{*}{\tilde{\chi}}\widetilde{\mathrm{D}})$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00378		—	G_{p}	$\mathcal{G}_{\mathrm{p}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00379		—	$\widetilde{\mathrm{D}}=\mathrm{D}$	$\widetilde{\mathrm{D}}=\mathrm{D}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00380		—	$\mathrm{C} \subset \mathrm{M}$	$\mathcal{C} \subset \mathrm{M}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00381		—	$\mathrm{F}(\mathrm{m}, \vartheta): \mathrm{M} \times \mathrm{T}^*(\mathrm{M}) \rightarrow \mathbb{R}^+$	$\mathrm{F}(\mathrm{m}, \vartheta): \mathrm{M} \times \mathrm{T}^*(\mathrm{M}) \rightarrow \mathbb{R}^+$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00382		—	C	$\mathcal{C}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00383		—	$(2, \theta)$	$(2, 0)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00384		—	$\mathrm{g}_{ij} \in \mathrm{C}^\infty(\mathrm{M})$	$\mathbf{g}_{ij} \in \mathrm{C}^\infty(\mathrm{M})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00385		—	$\mathrm{e}_{\alpha}^{\cdot i}$	$\mathrm{e}_{\alpha}^{\cdot i}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00386		—	$\mathfrak{0}(\mathrm{s}, \mathrm{n}-\mathrm{s})$	$\mathfrak{D}(\mathrm{s}, \mathrm{n}-\mathrm{s})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00387		—	ω^{g}	ω^{g}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00388		—	$\phi^{(\circ)}$	$\phi^{(\circ)}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00389		—	$\Gamma_{\mathrm{i}}^{\alpha \beta}$	$\Gamma_{\mathrm{i}}^{\alpha \beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00390		—	$L_{\alpha \beta}$	$L_{\alpha \beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00391		—	$\omega^{\alpha \beta}=-\omega^{\beta \alpha}$	$\omega^{\alpha \beta}=-\omega^{\beta \alpha}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00392		—	$\mathrm{m}_{\mathrm{o}} \in \mathrm{M}$	$\mathfrak{m}_o \in M$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00393		—	$\mathfrak{o}_{i\ j}$	$\mathfrak{o}_{ij}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00394		—	$\mathrm{H}=\mathrm{H}_0(\mathrm{s},\ \mathrm{n}-\mathrm{s})$	$H = O(s, n - s)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00395		—	$\mathfrak{o}(\mathrm{s},\ \mathrm{n}-\mathrm{s})$	$\mathfrak{o}(s, n - s)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00396		—	$\mathrm{L}^{\mathrm{g}}(\mathrm{M})$	$L^g(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00397		—	$\mathrm{D} \vartheta = 0$	$D\vartheta = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00398		—	Θ	Θ
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00399		—	ϑ^g	ϑ^g
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00400		—	Ω^g	Ω^g
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00401		—	$\omega^{\{ \}$	$\omega^{\{ \}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00402		—	K	K
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00403		—	U_{n}	U_n
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00404		—	V_{n}	V_n
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00405		—	$R_{i\ j\ c}^{\{ \{ . \ \ldots \ d \}$	$R_{ijc}^{\{ \dots d$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00406		—	$R_{i\ j\ k\ l}^{\{\ \}}$	$R_{ijkl}^{\{\ \}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00407		—	$S\ S\ B$	SSB
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00408		—	$\mathrm{O}(1,\ \mathrm{n}-1)$	$O(1,\ n-1)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00409		—	$\operatorname{GL}(\mathrm{n},\ \mathbb{R})$	$GL(n,\mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00410		—	$\mathrm{R}_{\alpha\beta}^{\{\alpha\ \beta\}^{\{\ .\ .\ \mathrm{cd}\}}(\mathrm{m})$	$R_{\alpha\beta}^{..cd}(m)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00411		—	$\vartheta,\ \Theta,\ \Omega^{\mathrm{g}}$	$\vartheta,\ \Theta,\ \Omega^g$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00412		—	$L_{\{\mathrm{g}\}}$	L_g
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00413		—	t	t
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00414		—	Π^{T}	Π^{T}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00415		—	Π^{L}	Π^{L}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00416		—	E	E
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00417		—	$\Pi_{\varnothing}^{\mathrm{T}}$	$\Pi_{\varnothing}^{\mathrm{T}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00418		—	$\Pi^{\mathrm{T}} := \left[\Pi_{\lceil}^{\mathrm{T}}, \vartheta \right]$	$\Pi^{\mathrm{T}} := \left[\Pi_{\lceil}^{\mathrm{T}}, \vartheta \right]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00419		—	$\delta \vartheta$	$\delta \vartheta$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00420		—	$\delta \omega^{\mathrm{g}}$	$\delta \omega^g$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00421		—	Σ	Σ
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00422		—	$\lambda^{\varphi}=\mathrm{o}$	$\lambda^\varphi = \mathbf{o}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00423		—	τ_{s}	τ_{s}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00424		—	Υ	Υ
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00425		—	$\Gamma^{\alpha\beta}=-\Gamma^{\beta\alpha}$	$\Gamma^{\alpha\beta} = -\Gamma^{\beta\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00426		—	$\delta L=\delta \Gamma^{\alpha\beta} \wedge \left(\partial L / \partial \Gamma^{\alpha\beta}\right. \left.\right)$	$\delta L = \delta \Gamma^{\alpha\beta} \wedge (\partial L / \partial \Gamma^{\alpha\beta})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00427		—	$S=\int_M L_{\text{mat}}$	$S = \int_M L_{\text{mat}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00428		—	$\operatorname{Diff}(M)$	$\operatorname{Diff}(M)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00429		—	$\mathscr{T} \subset \operatorname{Diff}(M)$	$\mathscr{T} \subset \operatorname{Diff}(M)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00430		—	$S_{0(1,3)}$	$SO(1,3)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00431		—	$\mathrm{t}_{\xi}:=\xi \jmath \mathbf{D}+\xi \jmath$	$\mathbb{L}_{\xi}:=\xi \jmath D+D \xi \jmath$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00432		—	M	M
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00433		—	ξ	ξ
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00434		—	$D\,g_{\{\alpha\,\beta\}}=0$	$Dg_{\alpha\beta}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00435		—	$\mathrm{L}_{\xi}L=D\,\xi\,\mathrm{jmath}\,L$	$L_{\xi}L=D\xi jL$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00436		—	$\xi\rightarrow e_{\alpha}$	$\xi\rightarrow e_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00437		—	$\left.F_{\alpha}^{\text{Max}}\right.=\left(e_{\alpha}\right\rfloor F\right)\wedge j$	$F_{\alpha}^{\text{Max}}=\left(e_{\alpha}\rfloor F\right)\wedge j$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00438		—	$\simeq$	$\simeq$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00439		—	$\delta L\,/\,\delta\Psi=0$	$\delta L/\delta\Psi=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00440		—	$\delta D\,\Psi=D\,\delta\Psi+\delta\Gamma_{\alpha}^{\beta}\wedge I^{\alpha}_{\beta}\Psi$	$\delta D\Psi=D\delta\Psi+\delta\Gamma_{\alpha}^{\beta}\wedge I^{\alpha}_{\beta}\Psi$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00441		—	$\varepsilon_{\alpha}^{\beta}(x):=\Lambda_{\alpha}^{\beta}(x)-\delta_{\alpha}^{\beta}$	$\varepsilon_{\alpha}^{\beta}(x):=\Lambda_{\alpha}^{\beta}(x)-\delta_{\alpha}^{\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00442		—	$T^{\alpha}=D\vartheta^{\alpha}$	$T^{\alpha}=D\vartheta^{\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00443		—	$D\,T^{\alpha}\equiv R_{\gamma}^{\alpha}\wedge\vartheta^{\gamma}$	$DT^{\alpha}\equiv R_{\gamma}^{\alpha}\wedge\vartheta^{\gamma}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00444		—	$\begin{aligned} &\widehat{D}\,\Sigma_{\alpha}\,\&\,\left.\cong\right\left(e_{\alpha}\right\rfloor R_{\beta}^{\gamma}\wedge\Delta^{\beta}\gamma\\ &\qquad\qquad\qquad-\frac{1}{2}\left(e_{\alpha}\right\rfloor Q_{\beta\gamma}\right)\wedge\sigma^{\beta\gamma} \end{aligned}$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00445		—	Γ_{α}^{β}	Γ_{α}^{β}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00446		—	$\begin{aligned} &R_{\alpha}^{\beta}=d\Gamma_{\alpha}^{\beta}\\ &\qquad\qquad\qquad-\Gamma_{\alpha}^{\gamma}\wedge\Gamma_{\gamma}^{\beta} \end{aligned}$	$R_{\alpha}^{\beta}=d\Gamma_{\alpha}^{\beta}$ $-\Gamma_{\alpha}^{\gamma}\wedge\Gamma_{\gamma}^{\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00447		—	$D\,R_{\alpha}^{\beta}\equiv 0$	$DR_{\alpha}^{\beta}\equiv 0$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00448		—	$D\,\Delta^{\alpha}_{\beta} + \vartheta^{\alpha} \wedge \Sigma_{\beta} \cong \sigma^{\alpha}_{\beta}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00449		—	$\overparen{D}$	(not rendered)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00450		—	$\left.\overparen{\Gamma}_{\alpha}^{\beta}:=\Gamma_{\alpha}^{\beta}+e_{\alpha}^{\beta}\right\rfloor T^{\beta}$	(not rendered)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00451		—	$\tau_{\alpha\beta}=-\tau_{\beta\alpha}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00452		—	$x_{[\alpha}\wedge\Sigma_{\beta]}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00453		—	x^i	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00454		—	$x^{\alpha}=\delta_i^{\alpha}x^i$	
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00455		—	$\xi=\partial / \partial t$	$\xi = \partial/\partial t$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00456		—	$\xi=\partial / \partial \varphi$	$\xi = \partial/\partial \varphi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00457		—	$\pm(\Lambda / 3) \ r^{\{3\}}$	$\pm(\Lambda/3)r^3$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00458		—	Λ	Λ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00459		—	$\Gamma^{\{T\} \ \alpha}$	$\Gamma^{(T)\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00460		—	$\Gamma_{\alpha}^{\{\alpha\}^{\{L\} \ \beta}}$	$\Gamma_{\alpha}^{(L)\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00461		—	$A(n, \ \mathbb{R}):=\mathbb{R}^{\{n\}} \ \otimes \ G \ L(n, \ \mathbb{R})$	$A(n, \mathbb{R}) := \mathbb{R}^n \otimes GL(n, \mathbb{R})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00469		—	$F:=d\ A=0$	$F:=dA=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00470		—	$F=d\ A+A\ \wedge\ A=0$	$F=dA+A\wedge A=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00471		—	$F\neq 0$	$F\neq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00472		—	$A(4,\ \mathbb{R})$	$A(4,\mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00473		—	$G\ L(4,\ \mathbb{R})$	$GL(4,\mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00474		—	$\Gamma^{\{(T)\ \alpha\}}\stackrel{*}{=}0$	$\Gamma^{(T)\alpha}\stackrel{*}{=}0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00475		—	(1993,1994)	(1993,1994)
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00476		—	P_{α}	P_{α}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00477		—	$2 \pi \hbar / \ell$	$2\pi\hbar/\ell$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00478		—	$\mathscr{M}_n = n M_{\text{Planck}}$	$\mathcal{M}_n = n M_{\text{Planck}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00479		—	$\Gamma^{\alpha\beta} = \Gamma^{\{j\alpha\beta} - K^{\alpha\beta}$	$\Gamma^{\alpha\beta} = \Gamma^{\{j\alpha\beta} - K^{\alpha\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00480		—	$\Gamma^{\{\alpha\beta\}}$	$\Gamma^{\{\alpha\beta\}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00481		—	$K_{\alpha\beta} = -K_{\beta\alpha}$	$K_{\alpha\beta} = -K_{\beta\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00482		—	$K = i K^{\alpha\beta} \sigma_{\alpha\beta} / 4$	$K = i K^{\alpha\beta} \sigma_{\alpha\beta} / 4$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00483		—	$T^{\alpha}=K^{\alpha}{}_{\beta}\wedge\vartheta^{\beta}$	$T^{\alpha}=K^{\alpha}{}_{\beta}\wedge\vartheta^{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00484		—	$\Gamma=\Gamma^{\{\ \ \}}-K$	$\Gamma=\Gamma^{\{\ \ \}}-K$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00485		—	$\Gamma^{\{\ \ \}}$	$\Gamma^{\{\ \ \}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00486		—	$D>2$	$D>2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00487		—	$j_{\{5\}}$	$j_{\{5\}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00488		—	$\bar{\psi}:=\psi^{\dagger}\gamma_0$	$\bar{\psi}:=\psi^{\dagger}\gamma_0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00489		—	$D\psi:=d\psi+\Gamma\wedge\psi$	$D\psi:=d\psi+\Gamma\wedge\psi$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00490		—	$\backslash\gamma:=\backslash\gamma_{\backslash\alpha}\ \backslash\vartheta^{\backslash\alpha}$	$\gamma := \gamma_{\alpha} \vartheta^{\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00491		—	$D\ \backslash\gamma=[\backslash\gamma,\ K]$	$D\gamma = [\gamma, K]$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00492		—	$=\backslash\gamma_{\backslash\alpha}\ T^{\backslash\alpha}$	$= \gamma_{\alpha} T^{\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00493		—	$T^{\backslash\alpha}:=D\ \backslash\vartheta^{\backslash\alpha}$	$T^{\alpha} := D\vartheta^{\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00494		—	$L_{\backslash\mathrm{D}}=\backslash\mathrm{bar}\{L_{\backslash\mathrm{D}}\}=L_{\backslash\mathrm{D}}^{\backslash\dagger}$	$L_D = \bar{L}_D = L_D^{\dagger}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00495		—	$\backslash\tau_{\backslash\alpha\ \backslash\beta\ \backslash\gamma}=\backslash\tau_{[\backslash\alpha\ \backslash\beta\ \backslash\gamma]}$	$\tau_{\alpha\beta\gamma} = \tau_{[\alpha\beta\gamma]}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00496		—	$\backslash\mu_{\backslash\alpha}=\backslash\vartheta_{\backslash\alpha}\ \backslash\wedge{\ }^{\backslash{*}}\ j_{\backslash{5}}\ /\ 4$	$\mu_{\alpha} = \vartheta_{\alpha} \wedge^* j_5/4$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00497		—	$\mathrm{GR}_{\parallel}$	$\mathrm{GR}_{\parallel}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00498		—	$C_{\mathrm{TT}*} := \vartheta^{\alpha} \wedge T_{\alpha}$	$C_{\mathrm{TT}*} := \vartheta^{\alpha} \wedge T_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00499		—	$L_{\mathrm{D}} \rightarrow L_{\mathrm{D}} + dU$	$L_{\mathrm{D}} \rightarrow L_{\mathrm{D}} + dU$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00500		—	$U = \vartheta^{\alpha} \wedge \mu_{\alpha}$	$U = \vartheta^{\alpha} \wedge \mu_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00501		—	$\hat{\tau}_{\alpha\beta}$	$\hat{\tau}_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00502		—	$U = \vartheta^{\alpha} \wedge \vartheta_{\alpha} \wedge j_5 / 4$	$U = \vartheta^{\alpha} \wedge \vartheta_{\alpha} \wedge j_5 / 4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00503		—	$\kappa = 8\pi G_{\mathrm{N}}$	$\kappa = 8\pi G_{\mathrm{N}}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00504		—	Σ_{α}	Σ_{α}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00505		—	L_{EC}	L_{EC}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00506		—	$\left.\left.\widehat{\Gamma}_{\alpha}^{\beta}:=\Gamma_{\alpha}^{\beta}+e_{\alpha}\right\rfloor T^{\beta}\right\rfloor$	$\left.\widehat{\Gamma}_{\alpha}^{\beta}:=\Gamma_{\alpha}^{\beta}+e_{\alpha}\right\rfloor T^{\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00507		—	$\sigma:=\frac{1}{2}\sigma_{\alpha\beta}\vartheta^{\alpha}\wedge\vartheta^{\beta}=\frac{i}{2}\gamma\wedge\gamma$	$\sigma:=\frac{1}{2}\sigma_{\alpha\beta}\vartheta^{\alpha}\wedge\vartheta^{\beta}=\frac{i}{2}\gamma\wedge\gamma$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00508		—	$\Theta=0$	$\Theta=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00509		—	$\Gamma^{\alpha\beta}$	$\Gamma^{\alpha\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00510		—	$\tilde{\mathrm{T}}_{\mu\nu}$	$\tilde{\mathrm{T}}_{\mu\nu}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00511		—	$\tau_{\mathrm{s}}=\mathrm{o}$	$\tau_{\mathrm{s}} = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00512		—	$\ell^{*~4}$	ℓ^{*4}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00513		—	$\ell^{*}=10^{-33}~\mathrm{\sim cm}$	$\ell^{*} = 10^{-33}~\mathrm{cm}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00514		—	$10^{\{80\}}$	10^{80}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00515		—	$R^{\{\backslash\lambda\alpha~\backslash\beta\}}$	$R^{\{\lambda\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00516		—	$\{~\}^{\{i\}}~T_{\backslash\alpha}$	$^{(i)}T_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00517		—	$\mathscr{A}$	$\mathscr{A}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00518		—	L_{HE}	L_{HE}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00519		—	$T^{\alpha} \wedge^* \left(T_{\alpha} - 2 T_{\alpha} - \frac{1}{2} T_{\alpha} \right)$	$T^{\alpha} \wedge^* \left({}^{(1)}T_{\alpha} - 2 {}^{(2)}T_{\alpha} - \frac{1}{2} {}^{(3)}T_{\alpha} \right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00520		—	$d C_{\mathrm{TT}}^*$	$d C_{\mathrm{TT}}^*$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00521		—	$G_{\alpha}^{\{\backslash\}} = G_{\alpha}^{\{\backslash\}} \eta_{\beta}$	$G_{\alpha}^{\{\}} = G_{\alpha}^{\beta} \eta_{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00522		—	$\ell = \sqrt{\kappa}$	$\ell = \sqrt{\kappa}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00523		—	$k \rightarrow \infty$	$k \rightarrow \infty$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00524		—	Γ_k	Γ_k
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00525		—	$k \, \partial_{\mathrm{k}}$	$k\partial_k$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00526		—	k	k
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00527		—	ρ_{Λ}	ρ_{Λ}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00528		—	d_{N}	d_{N}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00529		—	g_{N}	g_{N}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00530		—	$g_{\mathrm{N} \, *}$	$g_{\mathrm{N}*}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00531		—	λ_{*}	λ_{*}
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00532		—	$R^{\{n\}}$	R^n
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00533		—	$f(R)$	$f(R)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00534		—	κ	κ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00535		—	$1 / k^{\{4\}}$	$1/k^4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00536		—	$f_{\{*\}}^{\{2\}}$	f_*^2
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00537		—	$R_{\{\alpha \beta\}} \equiv 0$	$R_{\alpha\beta} \equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00538		—	$S\,L(5,\,\mathbb{R})$	$SL(5,\mathbb{R})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00539		—	$\Delta c / c \, \simeq 10^{-17}$	$\Delta c / c \simeq 10^{-17}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00540		—	10^{-13}	10^{-13}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00541		—	10^{-8}	10^{-8}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00542		—	$\bar{\nu}_e$	$\bar{\nu}_e$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00543		—	ν_e	ν_e
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00544		—	10^{-6}	10^{-6}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00545		—	C P T	CPT
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00546		—	$K^{\{0\}}$	K^0
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00547		—	$\bar{K}^{\{0\}}$	$\bar{K}^0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00548		—	$10^{\{-18\}}$	10^{-18}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00549		—	$(5, \mathbb{R})$	$(5, \mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00550		—	$(1986, 1991)$	$(1986, 1991)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00551		—	$d\,C_{\mathrm{TT}}$	dC_{TT}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00552		—	$\left(\mathrm{GR}_{\parallel}\right)$	$(\mathrm{GR}_{\parallel})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00553		—	$\ell := \sqrt{8 \pi \hbar G_{\mathrm{N}}} / c^{\{3\}} \simeq 10^{\{-33\}} \mathrm{\,cm}$	$\ell := \sqrt{8 \pi \hbar G_{\mathrm{N}} / c^3} \simeq 10^{-33} \text{ cm}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00554		—	q	q
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00555		—	$\mathscr{C} = q^{\{2\}} / 2$	$\mathscr{C} = q^2 / 2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00556		—	$\mathscr{S} = \exp \left(- \mathscr{C} \right) = \exp \left(- q^{\{2\}} / 2 \right)$	$\mathscr{S} = \exp(-\mathscr{C}) = \exp(-q^2 / 2)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00557		—	$\Psi = N \exp \left(i \int \vartheta \, \mathscr{C} \right)$	$\Psi = N \exp \left(i \theta \int \mathscr{C} \right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00558		—	$P := \mathbb{R}^{\{4\}} \otimes S \, 0(1,3)$	$P := \mathbb{R}^4 \otimes SO(1,3)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00559		—	$\mathscr{A} := \{ \, \}^{\{*\}} \left(\vartheta_{\alpha} \wedge T^{\alpha} \right) =$	$\mathscr{A} := {}^*(\vartheta_{\alpha} \wedge T^{\alpha}) =$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00560		—	$\mathrm{d} x^i$	$\mathcal{A}_i dx^i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00561		—	ℓ	ℓ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00562		—	$\mathrm{SO}(1,4)$	$\mathrm{SO}(1,4)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00563		—	$\mathrm{SO}(2,3)$	$\mathrm{SO}(2,3)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00564		—	$g_{\alpha \beta}$	$g_{\alpha \beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00565		—	$T_{\alpha}=g_{\alpha \beta} T^{\beta}$	$T_{\alpha}=g_{\alpha \beta} T^{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00566		—	ϑ_{α}	ϑ_{α}
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00567		—	$R_{\{\alpha \ \beta\}^{\{\ \ \}}}$	$R_{\alpha\beta}^{\{\ \}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00568		—	$\Gamma_{\{\alpha \ \beta\}^{\{\ \ \}}}$	$\Gamma_{\alpha\beta}^{\{\ \}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00569		—	$K_{\{\alpha \ \beta\}}$	$K_{\alpha\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00570		—	$\Gamma_{\{\alpha \ \beta\}}=$	$\Gamma_{\alpha\beta} =$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00571		—	$\backslash\mathrm{left}.\backslash\mathrm{left}.\backslash\mathrm{left}.\backslash\Gamma_{\{\beta \ \alpha\}}=\Gamma_{\{\alpha \ \beta\}^{\{\ \ \}}-\mathrm{K}_{\{\alpha \ \beta\}}=\Gamma_{\{\alpha \ \beta\}^{\{\ \ \}}+e_{\{\alpha\}\mathrm{right}\rfloor}T_{\{\beta\}}+\backslash\mathrm{left}(e_{\{\alpha\}\mathrm{right}\rfloor}e_{\{\beta\}\mathrm{right}\rfloor}T_{\{\gamma\}}\backslash\mathrm{right}\rfloor T_{\{\gamma\}}\wedge\vartheta^{\{\gamma\}}$	$-\Gamma_{\beta\alpha}=\Gamma_{\alpha\beta}^{\{\ \}}-K_{\alpha\beta}=\Gamma_{\alpha\beta}^{\{\ \}}+e_{\alpha}\rfloor T_{\beta}+(e_{\alpha}\rfloor e_{\beta}\rfloor T_{\gamma})\wedge\vartheta^{\gamma}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00572		—	$C_{\{\mathrm{TT}^*\}}:=\vartheta^{\{\alpha\}}\wedge^*T_{\alpha}/2\ell^2$	$C_{\mathrm{TT}^*}:=\vartheta^{\alpha}\wedge^*T_{\alpha}/2\ell^2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00573		—	$R^{\{\alpha \ \beta\}}=0$	$R^{\alpha\beta}=0$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00574		—	$R^{\alpha\beta} \wedge \lambda_{\alpha\beta}$	$R^{\alpha\beta} \wedge \lambda_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00575		—	$\lambda_{\alpha\beta} = -\lambda_{\beta\alpha}$	$\lambda_{\alpha\beta} = -\lambda_{\beta\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00576		—	$\vartheta^{\alpha}, \Gamma^{\alpha\beta}$	$\vartheta^{\alpha}, \Gamma^{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00577		—	$\lambda_{\alpha\beta}$	$\lambda_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00578		—	$L_{\parallel}$	$L_{\parallel}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00579		—	$\lambda_{\alpha\beta} = \left(\gamma/\ell^2\right) \eta_{\alpha\beta}$	$\lambda_{\alpha\beta} = \left(\gamma/\ell^2\right) \eta_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00580		—	$\boldsymbol{G R}_{\parallel}$	$\boldsymbol{G R}_{\parallel}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00581		—	$\sigma = \pm i \gamma$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00582		—	$\theta_{\mathrm{T}} = \pm i$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00583		—	$\stackrel{\scriptscriptstyle (\pm)}{T}^{\alpha} := \frac{1}{2} (T^{\alpha} \pm i^* T^{\alpha})$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00584		—	$\tau^{\lambda\beta}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00585		—	$D^{\tau} \lambda_{\alpha\beta} = \tau_{\alpha\beta}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00586		—	Π^{α}	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00587		—	$\delta L_{\parallel} / \delta \vartheta^{\alpha}$	
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00588		—	$R_{\alpha \beta} = 0$	$R_{\alpha \beta} = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00589		—	$\stackrel{(\pm)}{\Pi}_{\alpha}$	$\stackrel{(\pm)}{\Pi}_{\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00590		—	$\stackrel{(\pm)}{\Sigma}_{\alpha}$	$\stackrel{(\pm)}{\Sigma}_{\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00591		—	$\stackrel{(\pm)}{D}$	$\stackrel{(\pm)}{D}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00592		—	$\stackrel{(\mp)}{A}_{\alpha}^{\beta}$	$\stackrel{(\mp)}{A}_{\alpha}^{\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00593		—	$\underline{\Gamma}_{ }^{\alpha \beta} \rightarrow 0$	$\underline{\Gamma}_{ }^{\alpha \beta} \rightarrow 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00594		—	H_{α}	H_{α}
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00595		—	$\overparen{D} \ \eta_{\{\alpha\}} := D \ \eta_{\{\alpha\}} -$	<i>(not rendered)</i>
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00596		—	$\left(e_{\{\alpha\}} \downarrow T^{\beta}\right) \wedge \eta_{\{\beta\}} \equiv 0$	$\left(e_{\alpha} \downarrow T^{\beta}\right) \wedge \eta_{\beta} \equiv 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00597		—	$\stackrel{\scriptscriptstyle{D}}{\rel{\pm}} \eta_{\{\alpha\}} = \pm (i/2) \ell^2 \eta_{\alpha\beta} \wedge \stackrel{\scriptscriptstyle{\Pi}}{\rel{\pm}}$	$\stackrel{(\pm)}{D} \eta_{\alpha} = \pm (i/2) \ell^2 \eta_{\alpha\beta} \wedge \stackrel{(\pm)}{\Pi}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00598		—	$E_{\{\alpha\}}$	E_{α}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00599		—	$D \ E_{\{\alpha\}} =$	$DE_{\alpha} =$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00600		—	$D \ D \ H_{\{\alpha\}} \equiv R_{\{\alpha\}} \{ \ }^{\{\beta\}} \wedge H_{\{\beta\}} = 0$	$DDH_{\alpha} \equiv -R_{\alpha}^{\beta} \wedge H_{\beta} = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00601		—	$\mathbf{G \ R}_{\{\boldsymbol{\{\}}\}}$	$\mathbf{GR}_{\parallel}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00602		—	Σ_t	Σ_t
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00603		—	$x^0=t$	$x^0 = t$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00604		—	$u:=-N^2\,d\,t,\,\longrightarrow$	$u := -N^2 dt, \Longleftrightarrow$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00605		—	$d\,u\wedge u=0$	$du \wedge u = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00606		—	$u=g(n,$	$u = g(n,$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00607		—	$)$. The lapse function N	$)$.The lapse function N
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00608		—	N^A	N^A
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00609		—	p	p
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00610		—	(3+1)	$(3 + 1)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00611		—	$L=L\left(\vartheta^{\alpha},\,\Psi,\,D\,\Psi\right)$	$L=L\left(\vartheta^{\alpha},\Psi,D\Psi\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00612		—	$\left.\mathscr{K}\right _{L_{\perp}:=n}\right\rfloor L$	$\mathscr{K}=L_{\perp}:=n\rfloor L$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00613		—	$\{ \}^{\bullet}$	$\bullet$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00614		—	$\left.\ell\right _n:=d\left\rfloor n\right\rfloor +n\downarrow d$	$\ell_n:=dn\rfloor +n\downarrow d$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00615		—	n	n
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00616		—	$\left. n^{\alpha} \right\rceil := n \rfloor \vartheta^{\alpha}$	$n^{\alpha} \rceil \vartheta^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00617		—	$\left. n^{\alpha} e_{\alpha} \right\rceil \Psi = n \downarrow \Psi$	$n^{\alpha} e_{\alpha} \rceil \Psi = n \downarrow \Psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00618		—	$\mathrm{L}_n = D \ n \downarrow + n \downarrow D$	$\mathbf{L}_n = D n \downarrow + n \downarrow D$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00619		—	$n \rfloor \Psi = n \downarrow (dt \wedge n \rfloor \Psi) = n \rfloor \Psi$	$n \rfloor \Psi = n \downarrow (dt \wedge n \rfloor \Psi) = n \rfloor \Psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00620		—	$\left. \Gamma_{\perp}^{\alpha\beta} \right\rceil := n \rfloor \Gamma^{\alpha\beta}$	$\Gamma_{\perp}^{\alpha\beta} \rceil := n \rfloor \Gamma^{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00621		—	$\mathscr{G}_{\alpha}$	$\mathcal{G}_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00622		—	$\mathscr{G}_{\alpha\beta}$	$\mathcal{G}_{\alpha\beta}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00623		—	$\left. D_{\alpha} := e_{\alpha} \right\rfloor D$	$D_{\alpha} := e_{\alpha} \rfloor D$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00624		—	$D \, \lambda_{\alpha \, \beta}$	$D \lambda_{\alpha \beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00625		—	$\left. \underline{\Psi} := \Psi - d \, t \wedge n \right\rfloor \Psi$	$\underline{\Psi} := \Psi - dt \wedge n \rfloor \Psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00626		—	$\underline{\lambda_{\alpha \, \beta}} = \left(1 / 2 \, \ell^2 \right) * \left(\vartheta_{\alpha} \wedge \vartheta_{\beta} \right) = - \left(1 / N \ell^2 \right) n_{[\alpha} * \underline{\vartheta_{\beta]}}$	$\underline{\lambda}_{\alpha \beta} = \left(1 / 2 \ell^2 \right) * \left(\vartheta_{\alpha} \wedge \vartheta_{\beta} \right) = - \left(1 / N \ell^2 \right) n_{[\alpha} * \underline{\vartheta}_{\beta]}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00627		—	$d \, \stackrel{\scriptscriptstyle (\mp)}{\text{rel}} \{ C \}_\text{\texttt{TT}}$	$d \, \stackrel{\scriptscriptstyle (\mp)}{C}_{\texttt{TT}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00628		—	$(\alpha = \hat{0})$	$(\alpha = \hat{0})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00629		—	$(\alpha = B)$	$(\alpha = B)$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00630		—	$\stackrel{\scriptscriptstyle{(\pm)}}{\underline{\mathscr{A}}}_\alpha:=\stackrel{\scriptscriptstyle{*}}{\prod}_\alpha \underline{\Pi}_\alpha$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00631		—	$\stackrel{\scriptscriptstyle{*}}{-}_\alpha$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00632		—	$\stackrel{\scriptscriptstyle{**}}{=} +1$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00633		—	$\underline{\vartheta}^0=0$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00634		—	$\stackrel{\scriptscriptstyle{(\pm)}}{\mathscr{A}}_0= d^{(\pm)}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00635		—	$\alpha=\hat{0}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00636		—	$\stackrel{\scriptscriptstyle{*}}{-}\vartheta_\alpha$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00637		—	$\stackrel{\scriptscriptstyle{(\pm)}}{\mathrel{\mathop{\rightarrow}\limits^{\scriptscriptstyle{(\pm)}}}}_{\alpha} \mathscr{A}_{\alpha} := \prod_{\alpha}^*$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00638		—	$\stackrel{\scriptscriptstyle{(\pm)}}{\mathrel{\mathop{\rightarrow}\limits^{\scriptscriptstyle{(\pm)}}}}_{\alpha} \mathscr{A}_{\alpha} := \prod_{\alpha}^*$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00639		—	$\stackrel{\scriptscriptstyle{(\pm)}}{\mathrel{\mathop{\rightarrow}\limits^{\scriptscriptstyle{(\pm)}}}}_{\alpha} \mathscr{A}_{\alpha} := \prod_{\alpha}^*$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00640		—	$\stackrel{\scriptscriptstyle{(\pm)}}{\mathrel{\mathop{\rightarrow}\limits^{\scriptscriptstyle{(\pm)}}}}_{\alpha} \mathscr{A}_{\alpha} := \prod_{\alpha}^*$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00641		—	$\stackrel{\scriptscriptstyle{(\pm)}}{\mathrel{\mathop{\rightarrow}\limits^{\scriptscriptstyle{(\pm)}}}}_{\alpha} \mathscr{A}_{\alpha} := \prod_{\alpha}^*$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00642		—	$\stackrel{\scriptscriptstyle{(\pm)}}{\mathrel{\mathop{\rightarrow}\limits^{\scriptscriptstyle{(\pm)}}}}_{\alpha} \mathscr{A}_{\alpha} := \prod_{\alpha}^*$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00643		—	$\stackrel{\scriptscriptstyle{(\pm)}}{\mathrel{\mathop{\rightarrow}\limits^{\scriptscriptstyle{(\pm)}}}}_{\alpha} \mathscr{A}_{\alpha} := \prod_{\alpha}^*$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00644		—	$B=1,2,3$	$B = 1, 2, 3$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00645		—	$\stackrel{\scriptscriptstyle{(\pm)}}{\underline{T}}^B$	$\stackrel{\scriptscriptstyle{(\pm)}}{\underline{T}}^B$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00646		—	$\underline{\Gamma}_{\parallel}^{\alpha\beta} \stackrel{*}{=} 0$	$\underline{\Gamma}_{\parallel}^{\alpha\beta} \stackrel{*}{=} 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00647		—	$\Psi \wedge^* \Phi = \Phi \wedge^* \Psi$	$\Psi \wedge^* \Phi = \Phi \wedge^* \Psi$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00648		—	$S\,U(2)$	$SU(2)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00649		—	$\mathscr{H}_{\Lambda} \Psi_{\Lambda}(A) = 0$	$\mathscr{H}_{\Lambda} \Psi_{\Lambda}(A) = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F00650		—	$\ell \rightarrow 0$	$\ell \rightarrow 0$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00651		—	$\operatorname{dim} \underline{\vartheta}^B = [\ell]$	$\dim \underline{\vartheta}^B = [\ell]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00652		—	$\underline{T}^{\{(\mp)\}} \{ \}^B = 0$	$\underline{T}^{(\mp)B} = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00653		—	$\stackrel{\pm}{\operatorname{rel}}\{(\pm)\} \{ \}^B = 0$	$\stackrel{(\pm)}{\underline{T}} = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00654		—	$\mathbb{R}^4 \cup \infty = S^4$	$\mathbb{R}^4 \cup \infty = S^4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00655		—	(r, ψ, θ, ϕ)	(r, ψ, θ, ϕ)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00656		—	S^3	S^3
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00657		—	f	f
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00658		—	$d f= \pm 2 \, h \, d r$	$df = \pm 2 h dr$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00659		—	$\eta_{\mu\nu}^A := \eta_{\{0\} \{ \}^{\{A\}}_{\{\} \}^{\{\mu \, \nu\}} + \delta_{\mu}^A \, \delta_{\nu}^{\{0\}} - \delta_{\nu}^A \delta_{\mu}^{\{0\}}$	$\eta_{\mu\nu}^A := \eta_0^A{}_{\mu\nu} + \delta_{\mu}^A \delta_{\nu}^0 - \delta_{\nu}^A \delta_{\mu}^0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00660		—	$r \rightarrow \infty$	$r \rightarrow \infty$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00661		—	$\stackrel{\scriptscriptstyle{B}}{\stackrel{\scriptscriptstyle{B}}{\relax}}\{(\mp)\}^{\scriptscriptstyle{B}}_T=0$	$\stackrel{\scriptscriptstyle{B}}{T}^{\scriptscriptstyle{B}} = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00662		—	$\stackrel{\scriptscriptstyle{C}}{\stackrel{\scriptscriptstyle{C}}{\relax}}\{(\pm)\}^{\scriptscriptstyle{C}}_{\text{TT}}$	$\stackrel{\scriptscriptstyle{C}}{C}_{\text{TT}}^{\scriptscriptstyle{C}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00663		—	$\Psi_{\vartheta}(\vartheta) \propto e^{-C}$	$\Psi_{\parallel}(\vartheta) \propto e^{-C}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00664		—	$\underline{\vartheta}^{\hat{0}}=3\kappa d\,\vartheta_{\mathrm{L}}=h\,d\,r=\pm d f/2$	$\underline{\vartheta}^{\hat{0}}=3\kappa d\theta_{\mathrm{L}}=h dr=\pm df/2$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00665		—	$T^{\hat{0}}=0$	$T^{\hat{0}}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00666		—	$A=A_{\mu}^{\{I}\}\lambda_{\{I\}}$	$A=A_{\mu}^I\lambda_I$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00667		—	$d\,x^{\mu}\in\mathscr{C}$	$dx^{\mu}\in\mathcal{C}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00668		—	$\gamma^{\mu}(s)$	$\gamma^{\mu}(s)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00669		—	$\Psi(A)$	$\Psi(A)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00670		—	$\mathscr{C}/\mathscr{G}$	$\mathcal{C}/\mathcal{G}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00671		—	$G(\gamma)$	$G(\gamma)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00672		—	$V(\gamma)$	$V(\gamma)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00673		—	$ \Psi\rangle$	$ \Psi\rangle$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00674		—	$\mathbb{R}^4$	$\mathbb{R}^4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00675		—	$\left. n^{\alpha} = n \right\rfloor \vartheta^{\alpha}$	$n^{\alpha} = n \rfloor \vartheta^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00676		—	$\left(1 / 16 \pi^2\right)$	$(1/16\pi^2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00677		—	$\int R^{\alpha \beta} \wedge R_{\alpha \beta}^{(\star)}$	$\int R^{\alpha \beta} \wedge R_{\alpha \beta}^{(\star)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00678		—	$\left(1 / 8 \pi^2\right) \int R_{\alpha}\left\{\right\}^{\beta} \wedge R_{\beta}\left\{\right\}^{\alpha}$	$(1/8\pi^2) \int R_{\alpha}{}^{\beta} \wedge R_{\beta}{}^{\alpha}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00679		—	$\backslash\epsilon$	ϵ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00680		—	$0 \leq \epsilon \leq 1$	$0 \leq \epsilon \leq 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00681		—	$\backslash\Gamma_{\alpha\ \beta}(\epsilon)=\backslash\Gamma_{\alpha\ \beta}^{\{\ \ \}}-\epsilon K_{\alpha\ \beta}$	$\Gamma_{\alpha\beta}(\epsilon) = \Gamma_{\alpha\beta}^{\{\}} - \epsilon K_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00682		—	$\backslash\Gamma_{\alpha\ \beta}(0)=\backslash\Gamma_{\alpha\ \beta}^{\{\ \ \}}$	$\Gamma_{\alpha\beta}(0) = \Gamma_{\alpha\beta}^{\{\}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00683		—	$\backslash\Gamma_{\alpha\ \beta}$	$\Gamma_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00684		—	$\backslash\left.T:=e_{\alpha}\right\rceil T^{\alpha}=d\ \sigma$	$T := e_{\alpha}] T^{\alpha} = d\sigma$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00685		—	N	N
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00686		—	$\sigma=$	$\sigma =$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00687		—	$\arcsin(x)$	$\arcsin(x)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00688		—	$\pi_3(SO(4))=$	$\pi_3(SO(4)) =$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00689		—	$\mathbb{Z} \oplus \mathbb{Z}$	$\mathbb{Z} \oplus \mathbb{Z}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00690		—	$\ell \, \underline{\vartheta}^A \rightarrow$	$\ell \underline{\vartheta}^A \rightarrow$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00691		—	$\underline{\Gamma}^{\star A} := \frac{1}{2} \eta^{ABC} \underline{\Gamma}_{BC}$	$\underline{\Gamma}^{\star A} := \frac{1}{2} \eta^{ABC} \underline{\Gamma}_{BC}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00692		—	$\mathfrak{su}(2)$	$\mathfrak{su}(2)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00693		—	$\underline{C}_{\mathrm{TT}} \rightarrow \underline{C}_{\mathrm{RR}} + \Lambda^{\prime} \underline{\eta}$	$\underline{C}_{\mathrm{TT}} \rightarrow \underline{C}_{\mathrm{RR}} + \Lambda^{\prime} \underline{\eta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00694		—	$\mathbb{R} \times W^3$	$\mathbb{R} \times W^3$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00695		—	$d\,j_5$	dj_5
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00696		—	$d\,\mathrm{d}\,\mathscr{A} \wedge d\,\mathrm{d}\,\mathscr{A}$	$d\mathscr{A} \wedge d\mathscr{A}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00697		—	$\theta_{\mathrm{T}}=2\,/\,\gamma$	$\theta_{\mathrm{T}}=2/\gamma$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00698		—	$d\,C_{\mathrm{\text{TT}}}$	dC_{TT}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00699		—	$\vartheta_{[\beta} \wedge H_{\alpha]}$	$\vartheta_{[\beta} \wedge H_{\alpha]}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00700		—	$D\,H_{\alpha\,\beta}$	$DH_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00701		—	μ_{α}	μ_{α}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00702		—	$j_5=\overline{\psi}^*\,\gamma\gamma_5\,\psi$	$j_5=\psi^*\gamma\gamma_5\psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00703		—	$D\,\theta_{\mathrm{L}}\,\neq\,0$	$D\theta_{\mathrm{L}}\neq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00704		—	$\mu_{\alpha}\,\wedge\,\vartheta^{\alpha}=0$	$\mu_{\alpha}\wedge\vartheta^{\alpha}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00705		—	$\left.T=e_{\alpha}\right\rceil\rfloor T^{\alpha}$	$T=e_{\alpha}\rfloor T^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00706		—	$D\,D\,\theta=d\,d\,\theta\,\equiv\,0$	$DD\theta=dd\theta\equiv 0$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00707		—	ϑ^{β}	ϑ^{β}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00708		—	$\mathscr{A} := \left(\vartheta^{\alpha} \wedge T_{\alpha} \right)$	$\mathscr{A} := \left(\vartheta^{\alpha} \wedge T_{\alpha} \right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00709		—	θ_{L}	θ_{L}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00710		—	$T^{\left[\gamma \wedge T^{\mu} \right] \eta_{\alpha \beta \gamma \mu} = 0}$	$T^{\left[\gamma \wedge T^{\mu} \right] \eta_{\alpha \beta \gamma \mu} = 0}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00711		—	$D \ H$	DH
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00712		—	$L_{\{g\}}$	L_g
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00713		—	$\left\{ \right\}^{\left(i \right) } T^{\alpha}$	$\left(i \right) T^{\alpha}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00714		—	$\{ \}^{\{j\}} R^{\alpha \beta}$	${}^{(j)}R^{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00715		—	$\ell \simeq 10^{-32} \mathrm{~cm}$	$\ell \simeq 10^{-32} \text{ cm}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00716		—	$a_{\{0\}}, a_{\{i\}}(i=1,2,3), \kappa$	$a_0, a_i(i=1,2,3), \kappa$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00717		—	$b_{\{j\}}(j=1, \ldots, 6)$	$b_j(j=1, \ldots, 6)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00718		—	$C: Q \rightarrow Q^{\prime}=-Q$	$C: Q \rightarrow Q' = -Q$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00719		—	C	C
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00720		—	T	T
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00721		—	$\stackrel{\wedge}{=}$	$\stackrel{\wedge}{=}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00722		—	$a_{\{0\}}=0, a_{\{i\}}=1$	$a_0 = 0, a_i = 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00723		—	$b_{\{j\}}=1$	$b_j = 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00724		—	θ_{T}	θ_{T}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00725		—	P: $\vartheta^{\mathrm{\prime A}}=-\vartheta^{\mathrm{A}}$	$P : \vartheta^{\prime A} = -\vartheta^A$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00726		—	$A=1,2,3$	$A = 1, 2, 3$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00727		—	$\eta_{\{\alpha \ \beta \gamma \ \delta \}}$	$\eta_{\alpha\beta\gamma\delta}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00728		—	$R_{\{\alpha\}}^{\{\beta\}} \wedge R_{\{\beta\}}^{\{\alpha\}}$	$R_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00729		—	$\langle \text{angle } \theta R^{\{\alpha \beta\}} \theta \rangle = 0$	$\langle 0 R^{\alpha \beta} 0 \rangle = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00730		—	Ω_{C}	Ω_{C}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00731		—	$\operatorname{SL}(2, \mathbb{C}) \approx \widetilde{\operatorname{SO}}(3,1)$	$\operatorname{SL}(2, \mathbb{C}) \approx \widetilde{\operatorname{SO}}(3,1)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00732		—	$\sigma_{\alpha \beta} = \frac{1}{2} \left[\gamma_{\alpha}, \gamma_{\beta} \right]$	$\sigma_{\alpha \beta} = \frac{1}{2} \left[\gamma_{\alpha}, \gamma_{\beta} \right]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00733		—	$\sigma_{\alpha \beta}^{(\star)} = \gamma^5 \sigma_{\alpha \beta}$	$\sigma_{\alpha \beta}^{(\star)} = \gamma^5 \sigma_{\alpha \beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00734		—	$\gamma^{\{5\}} := \frac{1}{4!} \varepsilon_{\mathrm{abcd}} \gamma^{\mathrm{a}} \gamma^{\mathrm{b}} \gamma^{\mathrm{c}} \gamma^{\mathrm{d}} \gamma^{\mathrm{c}} \gamma^{\mathrm{d}}$	$\gamma^5 := \frac{1}{4!} \varepsilon_{\mathrm{abcd}} \gamma^{\mathrm{a}} \gamma^{\mathrm{b}} \gamma^{\mathrm{c}} \gamma^{\mathrm{d}}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00735		—	$s=0$	$s = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00736		—	$R_{\alpha \beta}$	$R_{\alpha \beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00737		—	$\Gamma^{\alpha \beta}_{\alpha \beta} := \Gamma_i^{\alpha \beta} d x^i$	$\Gamma^{\alpha \beta} := \Gamma_i^{\alpha \beta} dx^i$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00738		—	$\vartheta^{\alpha} = E_j^{\alpha} d x^j$	$\vartheta^{\alpha} = E_j^{\alpha} dx^j$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00739		—	e_{α}	e_{α}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00740		—	$G_{i j}$	G_{ij}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00741		—	$\operatorname{Ric}_{\alpha \beta} := (-1)^s * \left(R_{(\alpha} \{ \}^{\delta} \wedge \eta_{\delta \beta)} \right)$	$\operatorname{Ric}_{\alpha \beta} := (-1)^s * \left(R_{(\alpha}{}^{\delta} \wedge \eta_{\delta \beta)} \right)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00742		—	$\tau_{\alpha \beta}=\vartheta_{[\alpha} \wedge$	$\tau_{\alpha\beta}=\vartheta_{[\alpha}\wedge$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00743		—	$\mu_{\beta}]$	$\mu_{\beta}]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00744		—	$\Gamma^{\{\beta\alpha\beta}\{ }\wedge^{\alpha\beta} \propto \partial g$	$\Gamma^{\{\beta\alpha\beta}\propto\partial g$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00745		—	$\{ \}^{\{\backslash star\}}$	$(\star)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00746		—	$L_{\{\text{SKY}\}}^{\{\backslash star\}}=2\,L_{\text{SKY}}$	$L_{\text{SKY}}^{(\star)}=2L_{\text{SKY}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00747		—	$j_5:=\bar{\psi}^{\{\psi\}}\gamma\gamma_5\psi$	$j_5:=\psi^*\gamma\gamma_5\psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00748		—	$D\,\eta_{\alpha\beta}=0$	$D\eta_{\alpha\beta}=0$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00749		—	$\zeta = \pm 1$	$\zeta = \pm 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00750		—	$\zeta = +1$	$\zeta = +1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00751		—	$\zeta = -1$	$\zeta = -1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00752		—	$\mathscr{A} := \left(\vartheta_{\alpha} \wedge T^{\alpha} \right)$	$\mathscr{A} := \left(\vartheta_{\alpha} \wedge T^{\alpha} \right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00753		—	$\left\langle d j_5 \right\rangle = 2 i^* m \left\langle \bar{\psi} \gamma_5 \psi \right\rangle - \left(R_{\alpha \beta}^{\{ } \wedge R^{\} j \alpha \beta} + \frac{1}{2} d \mathscr{A} \wedge d \mathscr{A} \right) /$	$\left\langle d j_5 \right\rangle = 2 i^* m \left\langle \bar{\psi} \gamma_5 \psi \right\rangle - \left(R_{\alpha \beta}^{\{ } \wedge R^{\} j \alpha \beta} + \frac{1}{2} d \mathscr{A} \wedge d \mathscr{A} \right) /$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00754		—	$48 \pi^2$	$48 \pi^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00755		—	$d^4 x$	$d^4 x$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00756		—	$\{ \}^{*} R_{\{\alpha \beta\}}=\theta_{\mathrm{L}}^{\{\star\}} R_{\{\alpha \beta\}^{\{\star\}}}+\left(\theta_{\mathrm{T}}\right) \eta_{\{\alpha \beta\}}$	${}^*R_{\alpha\beta}=\theta_{\mathrm{L}}^{\star}R_{\alpha\beta}^{(\star)}+(\theta_{\mathrm{T}}^*\Lambda/6)\,\eta_{\alpha\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00757		—	$\sigma_{\alpha}:=\Sigma_{\alpha}-D^{\{\}}\mu_{\alpha}$	$\sigma_{\alpha}:=\Sigma_{\alpha}-D^{\{\}}\mu_{\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00758		—	$\left.\left.R:=e_{\{\alpha\}}\right\rfloor e_{\{\beta\}}\right\rfloor R^{\{\alpha \beta\}}$	$R:=e_{\alpha}\rfloor e_{\beta}\rfloor R^{\alpha\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00759		—	$o_{ij}^{\{\text{Mink}\}}$	o_{ij}^{Mink}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00760		—	$D^{\{\}}G_{\{\alpha\}}^{\{\}}\equiv 0$	$D^{\{\}}G_{\alpha}^{\{\}}\equiv 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00761		—	$R=-4\,\Lambda$	$R=-4\Lambda$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00762		—	$\Lambda\simeq 1/\kappa$	$\Lambda\simeq 1/\kappa$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00763		—	$R=0$	$R = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00764		—	$h_{i\ j}:=g_{i\ j}-g_{i\ j}^{\mathrm{dS}}$	$h_{ij} := g_{ij} - g_{ij}^{\mathrm{dS}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00765		—	$g_{i\ j}^{\mathrm{dS}}$	g_{ij}^{dS}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00766		—	$m\ /\ H$	m/H
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00767		—	$m\ \leq 7\ \times 10^{\{-32\}}\ \mathrm{eV}$	$m \leq 7 \times 10^{-32} \mathrm{eV}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00768		—	$H\ \simeq 10^{\{-33\}}\ \mathrm{eV}$	$H \simeq 10^{-33} \mathrm{eV}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00769		—	$\Lambda_{\mathrm{obs}} > 0$	$\Lambda_{\mathrm{obs}} > 0$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00770		—	$1 / g^2$	$1/g^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00771		—	$\operatorname{SU}(2)$	$\mathrm{SU}(2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00772		—	1974)	1974)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00773		—	$\overline{S_0(4)} \cong$	$\overline{SO(4)} \cong$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00774		—	$SU(2) \otimes SU(2)$	$SU(2) \otimes SU(2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00775		—	$\kappa \Lambda_{\text{obs}} \simeq 10^{-123}$	$\kappa \Lambda_{\text{obs}} \simeq 10^{-123}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00776		—	ΔL	ΔL
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00777		—	$\operatorname{Ric}_{\alpha\beta} := \left(R_{(\alpha}{}^{\delta}{}_{\beta)} \wedge \eta_{\delta} \right)$	$\operatorname{Ric}_{\alpha\beta} := {}^*(R_{(\alpha}{}^{\delta}{}_{\beta)} \wedge \eta_{\delta})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00778		—	$H_{\alpha\beta}$	$H_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00779		—	L_{eff}	L_{eff}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00780		—	$n-2$	$n-2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00781		—	$P := \mathbb{R}^4 \otimes$	$P := \mathbb{R}^4 \otimes$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00782		—	$H^{\alpha\beta} \rightarrow \vartheta^{\alpha}$	$H^{\alpha\beta} \rightarrow \vartheta^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00783		—	$H^{\alpha} \rightarrow \Gamma^{\alpha\beta}$	$H^{\alpha} \rightarrow \Gamma^{\alpha\beta}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00784		—	$\{ \}^{\{ * \}} D H_{\alpha} \{ \}^{\beta}$	$*DH_{\alpha}^{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00785		—	$\{ \}^{\{ * \}} \left(H^{\beta} \wedge \vartheta_{\alpha} \right) \equiv e_{\alpha}$	$*\left(H^{\beta} \wedge \vartheta_{\alpha}\right) \equiv e_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00786		—	H^{β}	H^{β}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00787		—	$\{ \}^{\{ * \}} D H^{\alpha} \cong \{ \}^{\{ * \}} E^{\alpha}$	$*DH^{\alpha} \cong *E^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00788		—	$\widetilde{G}$	$\widetilde{G}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00789		—	$\widetilde{H}_{\alpha \beta} := -\partial \widetilde{L} / \partial R^{\alpha \beta}$	$\widetilde{H}_{\alpha \beta} := -\partial \widetilde{L} / \partial R^{\alpha \beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00790		—	$\widetilde{H}_{\alpha \beta} = \stackrel{(2)}{H}_{\alpha \beta} + \stackrel{(4)}{H}_{\alpha \beta} + \cdots$	$\widetilde{H}_{\alpha \beta} = \stackrel{(2)}{H}_{\alpha \beta} + \stackrel{(4)}{H}_{\alpha \beta} + \cdots$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00791		—	$\widetilde{H}_{\alpha} := -\partial \widetilde{L} / \partial T^{\alpha}$	$H_{\alpha} := -\partial \widetilde{L} / \partial T^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00792		—	$T^{\alpha} :=$	$T^{\alpha} :=$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00793		—	$D \vartheta^{\alpha}$	$D \vartheta^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00794		—	$\widetilde{H}_{\alpha\beta\mu\nu}$	$\widetilde{H}_{\alpha\beta\mu\nu}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00795		—	$\partial \widetilde{H}_{\mu\nu} / \partial R^{\alpha\beta} = 0$	$\partial \widetilde{H}_{\mu\nu} / \partial R^{\alpha\beta} = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00796		—	$\theta_{\mathrm{T}} R^{\alpha\beta} \wedge \vartheta_{\alpha} \wedge \vartheta_{\beta}$	$\theta_{\mathrm{T}} R^{\alpha\beta} \wedge \vartheta_{\alpha} \wedge \vartheta_{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00797		—	$\widetilde{L}$	$\widetilde{L}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00798		—	$\eta_{\alpha\beta}$	$\eta_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00799		—	$\widetilde{H}_{\alpha\beta}$	$\widetilde{H}_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00800		—	$\widetilde{L}=L_{\mathrm{EC}}$	$\widetilde{L}=L_{\mathrm{EC}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00801		—	$L=L_{\mathrm{EC}}$	$L=L_{\mathrm{EC}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00802		—	$\theta_{\mathrm{T}}^*=1$	$\theta_{\mathrm{T}}^*=1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00803		—	$\theta_{\mathrm{T}}=0$	$\theta_{\mathrm{T}}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00804		—	$\theta_{\mathrm{T}}=i$	$\theta_{\mathrm{T}}=i$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00805		—	$\left.\Gamma_{\alpha}\right\}^{\beta}\rightarrow\left.\Gamma_{\alpha}\right\}^{\beta}:=\Gamma_{\alpha}^{\beta}\mp\left(i\ell^2/2\right)e_{\alpha}\right\}^{\beta}$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00806		—	$\mathrm{id}C_{\mathrm{TT}}$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00807		—	$\left\{ \right\} ^{*}\backslash \Phi =(-1)^{p(4-p)+s}\backslash \Phi$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00808		—	$s:=\operatorname{sig}=1$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00809		—	$\left\{ \eta,\eta_{\alpha},\eta_{\alpha\beta},\eta_{\alpha\beta\gamma}\eta_{\alpha\beta\gamma\delta}\right\}$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00810		—	$\left.\eta_{\alpha}\right\}:=e_{\alpha}\right\}\eta=*\vartheta_{\alpha},\eta_{\alpha\beta}:=$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00811		—	$\left.\left.\left.\left.e_{\beta}\right\}\right\}\right\}\eta_{\alpha}=\eta_{\alpha\beta\gamma}\vartheta^{\gamma}=e_{\beta}\right\}e_{\alpha}\right\}\eta=*(\vartheta_{\alpha}\wedge\vartheta_{\beta})=\frac{1}{2}\eta_{\alpha\beta\gamma\delta}\vartheta^{\gamma}\wedge\vartheta^{\delta}$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00812		—	$\backslash eta_{\backslash alpha \backslash beta \backslash gamma}:=e_{\backslash gamma}$	$\eta_{\alpha\beta\gamma} := e_{\gamma}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00813		—	$\backslash rfloor \backslash eta_{\backslash alpha \backslash beta}={ }^{\backslash *} \backslash left(\backslash vartheta_{\backslash alpha} \backslash wedge \backslash vartheta_{\backslash beta} \backslash wedge \backslash vartheta_{\backslash gamma} \backslash right)$	$\rfloor \eta_{\alpha\beta} = {}^*(\vartheta_{\alpha} \wedge \vartheta_{\beta} \wedge \vartheta_{\gamma})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00814		—	$o_{\backslash alpha \backslash beta}=e^{\{i\}\{ }\backslash alpha\} e^{\{j\}\{ }\backslash beta\} g_{\{i \ j\}}$	$o_{\alpha\beta} = e^i_{\alpha} e^j_{\beta} g_{ij}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00815		—	$o_{\backslash alpha \backslash beta}$	$o_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00816		—	$\backslash Phi=\frac{1}{p!} \backslash Phi_{\{i_{\{1\}} \backslash cdots i_{\{p\}}\} } d \ x^{\{i_{\{1\}}\} } \backslash wedge \backslash cdots \backslash wedge d \ x^{\{i_{\{p\}}\} }$	$\Phi = \frac{1}{p!} \Phi_{i_1 \cdots i_p} dx^{i_1} \wedge \cdots \wedge dx^{i_p}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00817		—	${ }^{\backslash *} \backslash *} \backslash Phi=(-1)^{p(4-p)+\operatorname{sig}} \backslash Phi$	$^{**} \Phi = (-1)^{p(4-p)+\operatorname{sig}} \Phi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00818		—	$\backslash left\{\backslash mathbf{1}_{\{4\}}, \backslash gamma_{\backslash alpha}, \backslash sigma_{\backslash alpha \backslash beta}, \backslash gamma_{\{5\}}, \backslash gamma_{\{5\}} \backslash gamma_{\backslash alpha}\backslash right\}$	$\{\mathbf{1}_4, \gamma_{\alpha}, \sigma_{\alpha\beta}, \gamma_5, \gamma_5 \gamma_{\alpha}\}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00819		—	$\sigma_{\alpha\beta}:=\frac{i}{2}\left(\gamma_{\alpha}\gamma_{\beta}-\gamma_{\beta}\gamma_{\alpha}\right)$	$\sigma_{\alpha\beta}:=\frac{i}{2}(\gamma_{\alpha}\gamma_{\beta}-\gamma_{\beta}\gamma_{\alpha})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00820		—	$\gamma_5=-i\gamma_{\hat{0}}\gamma_{\hat{1}}\gamma_{\hat{2}}\gamma_{\hat{3}}$	$\gamma_5=-i\gamma_{\hat{0}}\gamma_{\hat{1}}\gamma_{\hat{2}}\gamma_{\hat{3}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00821		—	$\gamma_5^2=(-1)^{\mathrm{sig}+1}\mathbf{1}_4$	$\gamma_5^2=(-1)^{\mathrm{sig}+1}\mathbf{1}_4$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00822		—	$\operatorname{Tr}\left(\gamma_{\alpha}\gamma_{\beta}\right)=4\,o_{\alpha\beta}$	$\operatorname{Tr}\left(\gamma_{\alpha}\gamma_{\beta}\right)=4o_{\alpha\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00823		—	$\operatorname{Tr}\left(\sigma_{\alpha\beta}\sigma^{\gamma\delta}\right)=8\delta_{[\alpha}^{\gamma}\delta_{\beta]}^{\delta}$	$\operatorname{Tr}\left(\sigma_{\alpha\beta}\sigma^{\gamma\delta}\right)=8\delta_{[\alpha}^{\gamma}\delta_{\beta]}^{\delta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00824		—	$[\alpha\beta]=\frac{1}{2}(\alpha\beta-\beta\alpha)$	$[\alpha\beta]=\frac{1}{2}(\alpha\beta-\beta\alpha)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00825		—	Lie	Lie
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00826		—	$\sigma_{\alpha\beta}^{\{\alpha\beta\}^{\star}}:=\frac{1}{2}\sigma^{\{\gamma\delta\}}\eta_{\alpha\beta\gamma\delta}$ $\delta=i\gamma_5\sigma_{\alpha\beta}$	$\sigma_{\alpha\beta}^{(*)}:=\frac{1}{2}\sigma^{\gamma\delta}\eta_{\alpha\beta\gamma\delta}=i\gamma_5\sigma_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00827		—	$\sigma^{\{\star\}}$	$\sigma^{(*)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00828		—	γ_5	γ_5
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00829		—	$12i^*\gamma$	$12i^*\gamma$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00830		—	$P_{\pm}:=\frac{1}{2}\left(1\pm\gamma_5\right)$	$P_{\pm}=\frac{1}{2}\left(1\pm\gamma_5\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00831		—	$P_{\pm}P_{\pm}=P_{\pm}$	$P_{\pm}P_{\pm}=P_{\pm}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00832		—	$\gamma\rightarrow\widetilde{\gamma}=e^{\varphi}\gamma$	$\gamma\rightarrow\widetilde{\gamma}=e^{\varphi}\gamma$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00833		—	$\{ \}^{\{\backslash\# \}}$	$\#$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00834		—	h	h
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00835		—	$\{ \}^{\{* * \}}=-1$	$** = -1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00836		—	$i^{\{ * \}}$	i^*
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00837		—	$\backslash Gamma:=\frac{\{ i \}}{4} \backslash Gamma^{\{ \alpha \} \backslash beta} \backslash sigma_{\{ \alpha \} \backslash beta}$	$\Gamma := \frac{i}{4} \Gamma^{\alpha \beta} \sigma_{\alpha \beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00838		—	$\overline{\{ S \}_\{\circ \}}(1,3) \backslash cong S L(2, C)$	$SO_{\circ}(1,3) \cong SL(2,C)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00839		—	$[\backslash Psi, \backslash Phi]:=\backslash Psi \wedge \backslash Phi - (-1)^{\{ p \} q}$	$[\Psi, \Phi]:=\Psi \wedge \Phi - (-1)^{pq}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00840		—	$\Phi \wedge \Psi$	$\Phi \wedge \Psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00841		—	$K:=\frac{i}{4} K^{\alpha \beta} \sigma_{\alpha \beta}$	$K:=\frac{i}{4} K^{\alpha \beta} \sigma_{\alpha \beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00842		—	$\{ \ }^{\{2\}} \Theta:=$	$^{(2)} \Theta :=$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00843		—	$\frac{1}{3} \gamma \wedge T$	$\frac{1}{3} \gamma \wedge T$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00844		—	$\{ \ }^{\{3\}} \Theta:=-\frac{1}{3} *(\gamma \wedge \mathscr{A})$	$^{(3)} \Theta :=-\frac{1}{3} *(\gamma \wedge \mathscr{A})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00845		—	$\{ \ }^{\{1\}} \Theta:=$	$^{(1)} \Theta :=$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00846		—	$\Theta-\{ \ }^{\{2\}} \Theta-\{ \ }^{\{3\}} \Theta$	$\Theta-^{(2)} \Theta-^{(3)} \Theta$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00847		—	$\widetilde{\mathrm{A}}=\mathrm{A}$	$\mathcal{A} = \mathcal{A}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00848		—	$D\,\Omega^{\mathrm{g}(\star)}\equiv 0$	$D\Omega^{g(\star)}\equiv 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00849		—	$D\,\eta_{\alpha\,\beta\,\gamma\,\delta}=0$	$D\eta_{\alpha\beta\gamma\delta}=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00850		—	$\mathrm{A}:=\{ \ }^*\!\!\left(\vartheta_{\alpha}\,\wedge\,T^{\alpha}\right)=\mathrm{A}_{\mathrm{i}}\,\mathrm{d}\,x^{\mathrm{i}}$	$\mathcal{A}:=*(\vartheta_{\alpha}\wedge T^{\alpha})=\mathcal{A}_i dx^i$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00851		—	$R^{\{\ \ \}}:=(-1)^{\text{sig}+1}\sim*\!\!\left(R^{\{\mathrm{j}\}}\,\wedge\,\eta_{\beta\,\alpha}\right)$	$R^{\{ \}}:=(-1)^{\mathrm{sig}+1}\ast\left(R^{\{j\}^{\beta}}\wedge\eta_{\beta\alpha}\right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00852		—	L_{SKY}	L_{SKY}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00853		—	$\operatorname{Ric}_{\alpha\,\beta}:=(-1)^{\text{sig}\,*}\!\!\left(R_{(\alpha\{ \}^{\delta}\,\wedge\,\eta_{\delta \beta})}(\delta\,\mid\,\beta)\right)$	$\operatorname{Ric}_{\alpha\beta}:=(-1)^{\mathrm{sig}\,*}\left(R_{(\alpha}{}^{\delta}\wedge\eta_{\delta \beta)}\right)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00854		—	$G_{\{\alpha\}}:=R^{\{\beta\}\gamma}\wedge\eta_{\{\alpha\}\beta\gamma}/2$	$G_{\alpha}:=R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma}/2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00855		—	$G^{\{\}\{\}}=G_{\{\alpha\}}^{\{\}\{\}}\eta_{\{\beta\}}^{\{\}\{\}}\gamma^{\{\alpha\}}$	$G^{\{\}}=G_{\alpha}^{\{\}\beta}\eta_{\beta}\gamma^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00856		—	$2+1$	$2+1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00857		—	$\mathrm{m}\rightarrow 0$	$\mathbf{m}\rightarrow 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00858		—	$\mathrm{dS}_{\{4\}}$	dS_4
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00859		—	$\mathrm{AdS}_{\{4\}}$	AdS_4
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00860		—	$5,\mathbb{R}$	$5,\mathbb{R}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00861		—	$R+R^{\wedge}2+Q^{\wedge}2$	$R+R^2+Q^2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00862		—	(2014,2006)	(2014,2006)
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00863		—	$\lambda_{\mathrm{DE}}=\left(\hbar c / \rho_{\mathrm{DE}}\right)^{1 / 4} \simeq 85 \mu \mathrm{m}$	$\lambda_{\mathrm{DE}}=\left(\hbar c / \rho_{\mathrm{DE}}\right)^{1 / 4} \simeq 85 \mu \mathrm{m}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00864		—	$[\hbar]=[p][q]$	$[\hbar]=[p][q]$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00865		—	$\hbar$	$\hbar$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00866		—	$C:=\frac{i}{4} C^{\alpha \beta} \sigma_{\alpha \beta}$	$C:=\frac{i}{4} C^{\alpha \beta} \sigma_{\alpha \beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00867		—	$\Psi:=\frac{i}{4} \Psi_j^{\alpha \beta} \sigma_{\alpha \beta} d x^j$	$\Psi:=\frac{i}{4} \Psi_j^{\alpha \beta} \sigma_{\alpha \beta} d x^j$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00868		—	$\Phi:=\frac{i}{4}\,\Phi^{\alpha\beta}\,\sigma_{\alpha\beta}$	$\Phi := \frac{i}{4}\Phi^{\alpha\beta}\sigma_{\alpha\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00869		—	$\sigma_{\alpha\beta}$	$\sigma_{\alpha\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00870		—	Γ	Γ
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00871		—	Ω^g	Ω^g
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00872		—	$C,\,\Psi$	C,Ψ
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00873		—	$s^2=0$	$s^2 = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00874		—	$^{\{2\}}$	2
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00875		—	$s \, s=0$	$ss = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00876		—	$(d \, \text{\o plus } s)(d \, \text{\o plus } s) \, \text{\equiv } 0$	$(d \oplus s)(d \oplus s) \equiv 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00877		—	$[s, \, d] := s \, d + d \, s = 0$	$[s, d] := sd + ds = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00878		—	$\text{\bar{C}}, \, \text{\bar{\chi}}$	$C, \bar{\chi}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00879		—	$\text{\bar{\Phi}}$	$\bar{\Phi}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00880		—	$[\Psi, \, \Phi] := \Psi \, \text{\wedge } \Phi -$	$[\Psi, \Phi] := \Psi \wedge \Phi -$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00881		—	$(-1)^{\text{\left(p_{\scriptscriptstyle 1}+g_{\scriptscriptstyle 1}\right)}\text{\left(p_{\scriptscriptstyle 2}+g_{\scriptscriptstyle 2}\right)}} \, \Phi \, \text{\wedge } \Psi$	$(-1)^{(p_1+g_1)(p_2+g_2)} \Phi \wedge \Psi$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00882		—	$\bar{\eta}$	$\bar{\eta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00883		—	$\mathbb{B}$	$\mathbb{B}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00884		—	$\alpha=\alpha_{i} \text{ d } x^{i}$	$\alpha=\alpha_i dx^i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00885		—	λ	λ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00886		—	α	α
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00887		—	$C \rightarrow (\alpha+$	$C \rightarrow (\alpha+$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00888		—	$\lambda) C, \Psi \rightarrow (\alpha+\lambda) \Psi$	$\lambda) C, \Psi \rightarrow (\alpha+\lambda) \Psi$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00889		—	$\backslash\mathrm{Phi} \rightarrow (\alpha + \lambda)^2 \backslash\mathrm{Phi}$	$\Phi \rightarrow (\alpha + \lambda)^2 \Phi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00890		—	$\backslash\mathrm{gamma} = \vartheta^{\alpha} \backslash\mathrm{gamma}_{\alpha}$	$\gamma = \vartheta^{\alpha} \gamma_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00891		—	$\backslash\mathrm{widetilde{\Theta}} := \backslash\mathrm{Theta} \oplus \backslash\mathrm{psi}$	$\tilde{\Theta} := \Theta \oplus \psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00892		—	$\backslash\mathrm{psi} := \backslash\mathrm{psi}_{\mathrm{j}}^{\alpha} \backslash\mathrm{gamma}_{\alpha} \mathrm{d} \mathrm{x}^{\mathrm{j}}$	$\psi := \psi_j^{\alpha} \gamma_{\alpha} dx^j$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00893		—	$\mathrm{s} \rightarrow \backslash\mathrm{widetilde{s}} = \mathrm{s} + \mathrm{f}_{\backslash\mathrm{zeta}}$	$s \rightarrow \widetilde{s} = s + \mathfrak{f}_{\zeta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00894		—	$\backslash\mathrm{left}.\backslash\mathrm{left}.\mathrm{mathrm{L}}_{\backslash\mathrm{zeta}} := \backslash\mathrm{zeta} \backslash\mathrm{right}\backslash\mathrm{rcorner} \mathrm{D} \backslash\mathrm{zeta} \backslash\mathrm{right}\backslash\mathrm{rfloor}$	$\mathbb{L}_{\zeta} := \zeta \lrcorner D - D \zeta \rfloor$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00895		—	$\backslash\mathrm{zeta} = \mathrm{zeta}^{\mathrm{i}} \backslash\mathrm{partial}_{\mathrm{i}}$	$\zeta = \zeta^i \partial_i$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00896		—	$\mathfrak{L}_{\zeta}$	$\mathfrak{L}_{\zeta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00897		—	ζ	ζ
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00898		—	$s \ \zeta = \phi + \mathfrak{L}_{\zeta} \ \zeta$	$s\zeta = \phi + \mathfrak{L}_{\zeta}\zeta$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00899		—	$p+1$	$p + 1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00900		—	$\exp \left(\zeta \rfloor \right) \widetilde{\Omega}^g$	$\exp(\zeta \rfloor) \widetilde{\Omega}^g$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00901		—	$\left. \exp \left(\zeta \rfloor \right) \right(\widetilde{\Theta} \oplus \phi)$	$\exp(\zeta \rfloor) \left(\widetilde{\Theta} \oplus \phi \right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00902		—	$\phi := \phi^{\alpha} \gamma_{\alpha}$	$\phi := \phi^{\alpha} \gamma_{\alpha}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00903		—	d	d
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00904		—	$\Gamma^{\mathrm{T}}=(\gamma-D\xi)/\ell$	$\Gamma^{(\mathrm{T})}=(\gamma-D\xi)/\ell$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00905		—	$\Omega^{\mathrm{T}}=$	$\Omega^{(\mathrm{T})}=$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00906		—	$D\,\Gamma^{\mathrm{T}}=(\Theta-DD\xi)/\ell$	$D\Gamma^{(\mathrm{T})}=(\Theta-DD\xi)/\ell$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00907		—	Γ^{T}	$\Gamma^{(\mathrm{T})}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00908		—	$\gamma\rightarrow\widetilde{\gamma}:=\ell\left(\Gamma^{\mathrm{T}}\oplus c\right)$	$\gamma\rightarrow\widetilde{\gamma}:=\ell\left(\Gamma^{(\mathrm{T})}\oplus c\right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00909		—	$\xi:=\xi^{\alpha}\,\gamma_{\alpha}$	$\xi:=\xi^{\alpha}\gamma_{\alpha}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00910		—	$A(4, \mathbb{R}) / GL(4, \mathbb{R}) \approx \mathbb{R}^4$	$A(4, \mathbb{R})/GL(4, \mathbb{R}) \approx \mathbb{R}^4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00911		—	$\Gamma^{\mathrm{T}} \stackrel{*}{=} 0$	$\Gamma^{(\mathrm{T})} \stackrel{*}{=} 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00912		—	$SU(N)$	$SU(N)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00913		—	$SL(4, \mathbb{R})$	$SL(4, \mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00914		—	$\mathrm{SO}(1,3)$	$\mathrm{SO}(1,3)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00915		—	$\scriptstyle =0$	$\scriptstyle = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00916		—	$s\{\cdots\}$	$s\{\cdots\}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00917		—	$d^{\{*\}} \Gamma=0, D^{\{*\}} \Psi=0$	$d^*\Gamma=0, D^*\Psi=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00918		—	$\Omega^{g(\pm)}=0$	$\Omega^{g(\pm)}=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00919		—	$C\ P$	CP
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00920		—	$s\ L_{\text{\text {Euler }}}=$	$sL_{\text{Euler}}=$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00921		—	$(-1)^{\text{\text {sig }+1}} d\left(\Psi^{\alpha\ \beta}\wedge R_{\alpha\ \beta}^{(\star)}\right)$	$(-1)^{\text{sig }+1}d\left(\Psi^{\alpha\beta}\wedge R_{\alpha\beta}^{(*)}\right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00922		—	$\bar{\chi}$	$\bar{\chi}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00923		—	$\widetilde{\Psi}_{\mathrm{F}}:=\bar{C}_{\alpha\ \beta}\left(d^{\{*\}} \Gamma^{\alpha\ \beta}+\frac{1}{2}\{*\}\mathbb{B}^{\alpha\ \beta}\right)$	$\widetilde{\Psi}_{\mathrm{F}}:=\bar{C}_{\alpha\beta}\left(d^*\Gamma^{\alpha\beta}+\frac{1}{2}*\mathbb{B}^{\alpha\beta}\right)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00924		—	$\{\cdots\}$	$\{\cdots\}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00925		—	$d^*\Gamma = \mathbb{B}$	$d^*\Gamma = \mathbb{B}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00926		—	$\rho=0$	$\rho = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00927		—	$\rho=1$	$\rho = 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00928		—	$D\,\Omega^g \equiv 0$	$D\Omega^g \equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00929		—	$T_{\mu \nu} := 2^* \left(\partial L / \partial g_{\mu \nu} \right) = 0$	$T_{\mu\nu} := 2^* (\partial L / \partial g_{\mu\nu}) = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F00930		—	$\mathscr{A} := \left(\vartheta_{\alpha} \wedge \right.$	$\mathscr{A} := \left(\vartheta_{\alpha} \wedge \right.$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00931		—	T^{α}	T^{α}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00932		—	$\gamma:=\vartheta^{\alpha} \gamma_{\alpha}$	$\gamma := \vartheta^{\alpha} \gamma_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00933		—	$d^{*} \gamma=0$	$d^{*} \gamma = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00934		—	$\widetilde{L}_{\mathrm{FP}}:=L_{\mathrm{FP}}+\Delta L_{\mathrm{FP}}$	$\widetilde{L}_{\mathrm{FP}} := L_{\mathrm{FP}} + \Delta L_{\mathrm{FP}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00935		—	i	i
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00936		—	$\stackrel{\pm}{\Omega}^g$	$\stackrel{\pm}{\Omega}^g$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00937		—	$\theta=\pi / 2$	$\theta = \pi / 2$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00938		—	$\stackrel{+}{\Omega}^g$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00939		—	$\theta=3\pi/2$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00940		—	$\bar{\Omega}^g$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00941		—	$\theta_{\mathrm{L}}=\theta_{\mathrm{T}}=0$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00942		—	$\theta_{\mathrm{L}}^{\star}=\mp(-1)^{\text{sig}}$	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00943		—	L_{qPG}	
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00944		—	$a_{\{M\}}$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00945		—	$b_{(N)}$	$b_{(N)}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00946		—	$\theta_{\mathrm{T}}, \theta_{\mathrm{T}}^*$	$\theta_{\mathrm{T}}, \theta_{\mathrm{T}}^*$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00947		—	θ_{L}^*	θ_{L}^*
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00948		—	$a_{\{0\}}=0, b_{(N)}=1$	$a_0 = 0, b_{(N)} = 1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00949		—	$d\,C_{\mathrm{TT}}\{ \}^{\{*\}}:=d\left(\vartheta^{\alpha}\wedge T_{\alpha}\right)/2\,\ell^2$	$dC_{\mathrm{TT}}^* := d\left(\vartheta^{\alpha}\wedge T_{\alpha}\right)/2\ell^2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00950		—	$\frac{1}{2}\,s\,\overline{\chi}^{\alpha\beta}\wedge\left(H_{\alpha\beta}+\partial L_{\theta}/\partial R^{\alpha\beta}\right)R^{\alpha\beta}$	$\frac{1}{2}s\overline{\chi}^{\alpha\beta}\wedge\left(H_{\alpha\beta}+\partial L_{\theta}/\partial R^{\alpha\beta}\right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00951		—	$\widetilde{L}_{\mathrm{FP}}$	$\widetilde{L}_{\mathrm{FP}}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00952		—	D	D
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00953		—	$\vartheta_{\alpha} \wedge \vartheta_{\beta}$	$\vartheta_{\alpha} \wedge \vartheta_{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00954		—	$\tau_{\alpha\beta}=0$	$\tau_{\alpha\beta} = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00955		—	$H_{\alpha}(**)$	$H_{\alpha}(**)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00956		—	$H_{\alpha\beta}(**) \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta}$	$H_{\alpha\beta}(**) \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00957		—	$\theta_{\mathrm{L}} \neq 0$	$\theta_{\mathrm{L}} \neq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00958		—	$R_{\alpha\beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta}=0$	$R_{\alpha\beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta} = 0$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00959		—	$\theta_{\mathrm{L}}=0$	$\theta_{\mathrm{L}}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00960		—	$\theta_{\mathrm{T}} R^{\alpha\beta} \wedge \vartheta_{\alpha} \wedge \vartheta_{\beta} / 4 \ell^2$	$\theta_{\mathrm{T}} R^{\alpha\beta} \wedge \vartheta_{\alpha} \wedge \vartheta_{\beta} / 4 \ell^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00961		—	$T^{\alpha} \simeq 0$	$T^{\alpha} \simeq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00962		—	$T^{\alpha}=0$	$T^{\alpha}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00963		—	λ_{α}	λ_{α}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00964		—	$\Gamma^{\{j}_{\alpha\beta}$	$\Gamma^{\{j}_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00965		—	$\mu_{\alpha}=\frac{1}{4} \vartheta_{\alpha} \wedge ^* j_5$	$\mu_{\alpha}=\frac{1}{4} \vartheta_{\alpha} \wedge ^* j_5$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00966		—	$\mu_{\alpha} \wedge \delta T^{\alpha}$	$\mu_{\alpha} \wedge \delta T^{\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00967		—	$D R_{\alpha \beta} \equiv 0, D R_{\alpha \beta}^{(\star)} \equiv$	$DR_{\alpha \beta} \equiv 0, DR_{\alpha \beta}^{(\star)} \equiv$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00968		—	$\theta, D \eta_{\alpha \beta}=0$	$0, D\eta_{\alpha \beta} = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00969		—	$D\left(\vartheta_{\alpha} \wedge \vartheta_{\beta}\right)=2 T_{[\alpha} \wedge \vartheta_{\beta]}=0$	$D\left(\vartheta_{\alpha} \wedge \vartheta_{\beta}\right)=2 T_{[\alpha} \wedge \vartheta_{\beta]} = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00970		—	$R_{\alpha \beta}^{(\star)}$	$R_{\alpha \beta}^{(\star)}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00971		—	$G^{\{\backslash \ \backslash\}}:=R^{\{\backslash 1\}} \backslash \beta \backslash \gamma \wedge \eta_{\alpha \beta \gamma} \gamma^{\alpha} / 2$	$G^{\{ \}} := R^{\{1\} \beta \gamma} \wedge \eta_{\alpha \beta \gamma} \gamma^{\alpha} / 2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00972		—	$G_{i j}:=\operatorname{Ric}_{i j}^{\{\backslash \ \backslash\}}-\frac{1}{2} g_{i j}$	$G_{ij} := \operatorname{Ric}_{ij}^{\{ \}} - \frac{1}{2} g_{ij}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00973		—	$\kappa_{\text{eff}}=\ell^2/\theta_{\mathrm{T}}^*$	$\kappa_{\text{eff}} = \ell^2/\theta_{\text{T}}^*$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00974		—	θ_{T}^*	θ_{T}^*
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00975		—	$\widehat{\Sigma}_{\alpha}$	$\widehat{\Sigma}_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00976		—	$\kappa_{\text{eff}}=8\,\pi\,G_{\mathrm{N}}/c^4$	$\kappa_{\text{eff}} = 8\pi G_{\text{N}}/c^4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00977		—	L_{qPG}	L_{qPG}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00978		—	$\varphi\propto 1/\ell$	$\langle\varphi\rangle\propto 1/\ell$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00979		—	$\mathfrak{a}$	$\mathfrak{a}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00980		—	$\Phi = a + i f_{\varphi} \exp \left(- \varphi / f_{\varphi} \right)$	$\Phi = a + i f_{\varphi} \exp \left(- \varphi / f_{\varphi} \right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00981		—	$\Lambda_{\text{eff}} > 0$	$\Lambda_{\text{eff}} > 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00982		—	$\mathrm{N} = \mathrm{N}(\mathrm{r} \pm \mathrm{t})$	$\mathrm{N} = \mathrm{N}(\mathrm{r} \pm \mathrm{t})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00983		—	r	r
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00984		—	$\Lambda_{\text{eff}} = 0$	$\Lambda_{\text{eff}} = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00985		—	$\{ \ }^{\{ * \}}$	$*$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00986		—	$\mathrm{u}^{\{ 2 \}} - \mathrm{v}^{\{ 2 \}} = 1$	$\mathrm{u}^2 - \mathrm{v}^2 = 1$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5 - 0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00987		—	$\mathrm{r}=\mathrm{r}^{\ast}=\mathrm{o}$	$\mathbf{r} = \mathbf{r}^{\ast} = \mathbf{0}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00988		—	$\mathrm{\tilde{v}}:=\mathrm{i}v$	$\tilde{\mathbf{v}} := \mathrm{i}v$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00989		—	S^2	$\mathbf{S}^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00990		—	$\mathrm{e}^v=0$	$\mathbf{e}^v = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00991		—	$\tau:=$	$\tau :=$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00992		—	$\mathrm{K}_{_3}$	$\mathbf{K}_3$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F00993		—	ω^g	ω^g
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00994		—	$R\ C$	RC
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00995		—	$\Omega_{\mathrm{C}}^{\{\ \}}$	$\Omega_C^{\{\}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00996		—	$\mathrm{f}_{\mathrm{i}}=\mathrm{o}$	$\mathbf{f}_i = \mathbf{o}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00997		—	$K\ \wedge\ K$	$K \wedge K$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00998		—	$\mu=0$	$\mu = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F00999		—	$m\ u$	μ
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01000		—	$\kappa<0$	$\kappa < 0$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01001		—	$\mathrm{t} \rightarrow \mathrm{i} \tau$	$t \rightarrow -i\tau$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01002		—	$\mathrm{f}=\mathrm{T}_{\hat{0}}^{\hat{0}} \hat{1}^{\hat{1}}$	$\mathbf{f} = \mathbf{T}_{01}^{\cdot 0}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01003		—	$\mathrm{g}=\mathrm{T}_{\hat{1}}^{\hat{1}} \hat{2}^{\hat{2}} \cdots \hat{2}^{\hat{2}}=\mathrm{T}_{\hat{1}}^{\hat{1}} \hat{3}^{\hat{3}} \cdots \hat{3}^{\hat{3}}$	$\mathbf{g} = \mathbf{T}_{12}^{\cdot 2} = \mathbf{T}_{13}^{\cdot 3}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01004		—	$\mathrm{n}=4$	$\mathbf{n} = 4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01005		—	R^6	$\mathbf{R}^6$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01006		—	γ_2^g	γ_2^g
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01007		—	$\mathrm{H}_{\mathrm{P}}(\mathrm{M}, \mathbb{R})$	$\mathbf{H_P(M, \mathbb{R})}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01008		—	$\mathrm{b}:=\operatorname{dim} \mathrm{H}_{\mathrm{p}}(\mathrm{M}, \mathbb{R})$	$\mathrm{b}:=\dim \mathrm{H}_{\mathrm{p}}(\mathrm{M}, \mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01009		—	$\chi(\mathrm{M})$	$\chi(\mathrm{M})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01010		—	$\mathrm{s}=1$	$\mathrm{s}=1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01011		—	$\alpha: \omega^{\mathrm{g}}-\omega^{\mathrm{g}}$	$\alpha: \omega^{\mathrm{g}}-\omega^{\mathrm{g}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01012		—	B^{n}	B^{n}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01013		—	$\partial \mathrm{B}^{\mathrm{n}}=\mathrm{S}^{\mathrm{n}-1}$	$\partial \mathrm{B}^{\mathrm{n}}=\mathrm{S}^{\mathrm{n}-1}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01014		—	$\mathrm{W}^{\mathrm{n}}:=\mathrm{S}^1 \times \mathrm{S}^{\mathrm{n}-1}$	$\mathrm{W}^{\mathrm{n}}:=\mathrm{S}^1 \times \mathrm{S}^{\mathrm{n}-1}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01015		—	$\left(\mathrm{S}^{\mathrm{1}} \times \mathrm{S}^{\mathrm{\{n\}}-1}\right)$	$\left(\mathrm{S}^1 \times \mathrm{S}^{\mathrm{n}-1}\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01016		—	$\mathbb{C} \, \mathrm{P}^{\mathrm{2}}$	$\mathbb{C}\mathbb{P}^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01017		—	E^{2}	E^2
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01018		—	$\mathrm{T}^{\mathrm{4}}=\mathrm{S}^{\mathrm{1}} \times \mathrm{S}^{\mathrm{1}} \times \mathrm{S}^{\mathrm{1}} \times \mathrm{S}^{\mathrm{1}}$	$\mathrm{T}^4=\mathrm{S}^1 \times \mathrm{S}^1 \times \mathrm{S}^1 \times \mathrm{S}^1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01019		—	$\mathscr{S}_{\mathrm{*}}(\mathrm{M}):=\mathscr{M} / \tilde{\mathscr{D}}$	$\mathscr{S}_*(\mathbf{M}):=\mathscr{M}/\tilde{\mathscr{D}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01020		—	S^{2}	S^2
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01021		—	$\mathbb{R} \times$	$\mathbb{R} \times$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01022		—	$S^{\{1\}} \times S^{\{2\}}$	$S^1 \times S^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01023		—	$\mathbf{R}^{\{4\}}$	$\mathbf{R}^4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01024		—	$U_{\{4\}}$	U_4
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01025		—	$\mathrm{R}+\mathrm{R}^{\{2\}}$	$\mathrm{R}+\mathrm{R}^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01026		—	$C_{\{\mathrm{TL}\}}$	C_{TL}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01027		—	$\varepsilon \rightarrow 1/\varepsilon$	$\varepsilon \rightarrow 1/\varepsilon$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01028		—	$\Gamma^{\{\beta \gamma\}}=\Gamma_{\{j\}}^{\{\beta \gamma\}} \, \mathrm{d} \, x^{\{j\}}$	$\Gamma^{\beta \gamma}=\Gamma_j^{\beta \gamma} \, dx^j$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01029		—	$\Gamma_{\alpha}^{\star}$	$\Gamma_{\alpha}^{\star}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01030		—	$\vartheta_{\alpha} / \ell$	$\vartheta_{\alpha} / \ell$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01031		—	$C=\operatorname{Tr}\{A \wedge F\}$	$C = \operatorname{Tr}\{A \wedge F\}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01032		—	T_{α}	T_{α}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01033		—	$R_{\alpha}^{\star}$	$R_{\alpha}^{\star}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01034		—	η_{α}	η_{α}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01035		—	$n^{\{2\}}$	n^2
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01036		—	$n^{\{2\}}(n-1) / 2$	$n^2(n-1)/2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01037		—	$R^{\{\alpha \beta \}}$	$R^{\alpha \beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01038		—	$n^{\{2\}}(n-1)^{\{2\}} / 4$	$n^2(n-1)^2/4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01039		—	$\tau_{\{\alpha \beta \}}$	$\tau_{\alpha \beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01040		—	$n=3$	$n = 3$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01041		—	$\mathrm{SO}\{1,2\}$	$\mathrm{SO}(1,2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01042		—	$9=5 \otimes 3 \otimes 1$	$9 = 5 \otimes 3 \otimes 1$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01043		—	$H=-\partial L / \partial F$	$H = -\partial L / \partial F$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01044		—	$H_{\{\alpha\}}=H_{\{\{\alpha \ \beta\}\} \ \vartheta^{\{\beta\}}}$	$H_{\alpha} = H_{[\alpha\beta]}\vartheta^{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01045		—	$S_{\{\alpha\}}=-\{ \}^{\{*\}} H_{\{\alpha\}}$	$S_{\alpha} = -^*H_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01046		—	$\mathbb{R}^3 \otimes S_{0(1,2)}$	$\mathbb{R}^3 \otimes SO(1,2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01047		—	$\theta_{\mathrm{T}}, \ \theta_{\mathrm{L}}$	$\theta_{\mathrm{T}}, \theta_{\mathrm{L}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01048		—	$\theta_{\mathrm{TL}}=-\chi$	$\theta_{\mathrm{TL}} = -\chi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01049		—	$C_{\mathrm{T}}, \ C_{\mathrm{L}}$	$C_{\mathrm{T}}, C_{\mathrm{L}}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01050		—	C_{T}	C_{T}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01051		—	$\Gamma^{\star\alpha}$	$\Gamma^{\star\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01052		—	$\theta_{\mathrm{T}}/2\ell$	$\theta_{\mathrm{T}}/2\ell$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01053		—	$\varepsilon = \theta_{\mathrm{TL}}\theta_{\mathrm{T}}/A$	$\varepsilon = \theta_{\mathrm{TL}}\theta_{\mathrm{T}}/A$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01054		—	$\rho = -\theta_{\mathrm{T}}^2/A$	$\rho = -\theta_{\mathrm{T}}^2/A$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01055		—	$A := -(-1)^s \theta_{\mathrm{TL}}^2 + 2\theta_{\mathrm{T}}\theta_{\mathrm{L}} \neq 0$	$A := -(-1)^s \theta_{\mathrm{TL}}^2 + 2\theta_{\mathrm{T}}\theta_{\mathrm{L}} \neq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01056		—	$K_{\alpha}^{\star} = -\{ \}^{\star} T_{\alpha} / 2 = -(-1)^s (\varepsilon / 2\ell) \vartheta_{\alpha}$	$K_{\alpha}^{\star} = -^{\star} T_{\alpha} / 2 = -(-1)^s (\varepsilon / 2\ell) \vartheta_{\alpha}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01057		—	$\varepsilon=0$	$\varepsilon = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01058		—	Λ_{eff}	Λ_{eff}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01059		—	J	J
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01060		—	$J^2 \leq M^2$	$J^2 \leq M^2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01061		—	$M=-1$	$M = -1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01062		—	$J=0$	$J = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01063		—	$\Gamma_{\alpha}^{\{\left\{\right\}^{\star}\right\}} \rightarrow \Gamma_{\alpha}^{\{\star\}} = \Gamma_{\alpha}^{\{\star\}} - K_{\alpha}^{\star}$	$\Gamma_{\alpha}^{\{\}^{\star}} \rightarrow \Gamma_{\alpha}^{\star} = \Gamma_{\alpha}^{\{\}^{\star}} - K_{\alpha}^{\star}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01064		—	$\backslash\mathrm{varepsilon}$	ε
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01065		—	$\backslash\Gamma_{\alpha}^{\{\star\}}=\frac{1}{2}\ \eta_{\alpha\ \beta\ \gamma}\ \backslash\Gamma^{\{\beta\}\ \gamma}$	$\Gamma_{\alpha}^{\star}=\frac{1}{2}\eta_{\alpha\beta\gamma}\Gamma^{\beta\gamma}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01066		—	$\backslash\mathrm{widetilde{T}}_{\alpha}=0$	$\widetilde{T}_{\alpha}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01067		—	$\backslash\mathrm{widetilde{\Gamma}}_{\alpha}^{\{\star\}}\ \backslash\mathrm{stackrel{*}{=}}\ 0$	$\widetilde{\Gamma}_{\alpha}^{\star}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01068		—	$\backslash\mathrm{widetilde{R}}_{\alpha}^{\{\star\}}=0$	$\widetilde{R}_{\alpha}^{\star}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01069		—	$1\ /\ \backslash\mathrm{varepsilon}$	$1/\varepsilon$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01070		—	$\backslash\rho=(-1)^{\{\mathrm{text\ sign}\}}\ \backslash\mathrm{varepsilon}^{\{2\}}\ /\ 4$	$\rho=(-1)^{\mathrm{sign}}\varepsilon^2/4$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01071		—	$R_{\alpha}^{\star}=(-1)^{\text{sign}}\,\varepsilon T_{\alpha}/4\,\ell$	$R_{\alpha}^{\star}=(-1)^{\text{sign}}\,\varepsilon T_{\alpha}/4\ell$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01072		—	$\varepsilon \neq 0$	$\varepsilon \neq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01073		—	$\varepsilon \rightarrow -\varepsilon$	$\varepsilon \rightarrow -\varepsilon$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01074		—	$\gamma=(1-\delta\varepsilon)\rightarrow$	$\gamma=(1-\delta\varepsilon)\rightarrow$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01075		—	$\gamma \neq 0$	$\gamma \neq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01076		—	$\varepsilon \rightarrow 0$	$\varepsilon \rightarrow 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01077		—	$H_{\alpha}^{\star}=\gamma\vartheta_{\alpha}/\ell$	$H_{\alpha}^{\star}=\gamma\vartheta_{\alpha}/\ell$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01078		—	$\backslash\mathrm{varepsilon} \rightarrow \infty$	$\varepsilon \rightarrow \infty$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01079		—	$H_{\alpha}=-\gamma \vartheta_{\alpha} / \delta \ell^2$	$H_{\alpha}=-\gamma \vartheta_{\alpha} / \delta \ell^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01080		—	$H_{\alpha}^{\star}=0$	$H_{\alpha}^{\star}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01081		—	$\{H, d t, d \mathscr{A}, d h\}$	$\{H, d t, d \mathscr{A}, d h\}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01082		—	$H=0$	$H=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01083		—	$H_{\alpha} \neq 0$	$H_{\alpha} \neq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01084		—	$d \mathscr{A}$	$d \mathscr{A}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01085		—	$d^{\ast}\vartheta_{\alpha}=0$	$d^{\ast}\vartheta_{\alpha}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01086		—	$\eta_{\alpha}=\vartheta^{\ast}_{\alpha}\vartheta_{\alpha}$	$\eta_{\alpha}=\vartheta^{\ast}_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01087		—	$m=\varepsilon/\ell$	$m=\varepsilon/\ell$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01088		—	$\vartheta^{\alpha}=E_i^{\alpha}dx^i$	$\vartheta^{\alpha}=E_i^{\alpha}dx^i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01089		—	$3\times 3=9$	$3\times 3=9$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01090		—	$S_0(1,2)$	$SO(1,2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01091		—	$\vartheta^{\prime\alpha}=\Lambda_{\beta}^{\alpha}(x)\vartheta^{\beta}$	$\vartheta^{\prime\alpha}=\Lambda_{\beta}^{\alpha}(x)\vartheta^{\beta}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01092		—	$2 / \theta_{\mathrm{L}}$	$2/\theta_{\mathrm{L}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01093		—	$\theta_{\mathrm{T}} \theta_{\mathrm{L}} / 2 \chi=1$	$\theta_{\mathrm{T}}\theta_{\mathrm{L}}/2\chi = 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01094		—	$\theta_{\mathrm{T}} \neq 0$	$\theta_{\mathrm{T}} \neq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01095		—	$\varepsilon \neq \tilde{\varepsilon}$	$\varepsilon \neq \tilde{\varepsilon}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01096		—	$\tilde{\theta}_{\mathrm{L}}$	$\tilde{\theta}_{\mathrm{L}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01097		—	$C^{\mathrm{k} \mathrm{l}}=C^{\mathrm{l} \mathrm{k}}$	$C^{kl} = C^{lk}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01098		—	$\Gamma^{\{\lambda \alpha \beta\}}$	$\Gamma^{\{\lambda \alpha \beta\}}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01099		—	$T_{\alpha}=D\ \vartheta_{\alpha}=0$	$T_{\alpha}=D\vartheta_{\alpha}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01100		—	$H_{\alpha}=0$	$H_{\alpha}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01101		—	$D\ H_{\alpha}^{\star}=\delta\ell\ D\ H_{\alpha}+(\gamma/\ell)\ D\ \vartheta_{\alpha}\ \text{equiv}\ 0$	$DH_{\alpha}^{\star}=\delta\ell DH_{\alpha}+(\gamma/\ell)D\vartheta_{\alpha}\equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01102		—	$D^{\{4\}}\ H_{\{\alpha\ \beta\}}$	$D^{[4]}H_{\{\alpha\beta\}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01103		—	$G_{\alpha}=R_{\alpha}^{\{\left\{\ \right\}^{\}\}}\ \star\right. .\}$	$G_{\alpha}=R_{\alpha}^{\{\}\star}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01104		—	$\widetilde{\Sigma}_{\alpha}:=\Sigma_{\alpha}-D^{\{\ \ \}}\ \mu_{\alpha}$	$\tilde{\Sigma}_{\alpha}:=\Sigma_{\alpha}-D^{\{\}}\mu_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01105		—	$E=p\ c_{\mathrm{F}}$	$E=pc_{\mathrm{F}}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01106		—	c_{F}	c_F
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01107		—	$\mathrm{SO}(3)$	$\mathrm{SO}(3)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01108		—	$s=1$	$s = 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01109		—	$S\,L(2,\,\mathbb{R})$	$SL(2,\mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01110		—	$\mathfrak{sp}(2,\,\mathbb{R})$	$\mathfrak{sp}(2,\mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01111		—	$\gamma_{\alpha}\,\gamma_{\beta}+\gamma_{\beta}\,\gamma_{\alpha}=2\,g_{\alpha\beta}$	$\gamma_{\alpha}\gamma_{\beta}+\gamma_{\beta}\gamma_{\alpha}=2g_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01112		—	$D\,\psi:=d\,\psi-\frac{1}{2}\,\gamma_{\alpha}\,\wedge\,\Gamma^{\star\alpha}\psi$	$D\psi:=d\psi-\frac{1}{2}\gamma_{\alpha}\wedge\Gamma^{\star\alpha}\psi$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01113		—	L_{D}	L_{D}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01114		—	$\bar{\Psi}$	$\bar{\Psi}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01115		—	$\hat{D}$	$\hat{D}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01116		—	$F := d \ A$	$F := dA$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01117		—	$d\left(A \wedge^* F\right) / 2$	$d\left(A \wedge^* F\right) / 2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01118		—	$\square := d^* d^* - * d^* d$	$\square := d^* d^* - * d^* d$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01119		—	$\lambda = \hbar / c \ m_{\text {photon }}$	$\lambda = \hbar / c m_{\text{photon}}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01120		—	x^{α}	x^{α}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01121		—	y^{α}	y^{α}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01122		—	(x, y)	(x, y)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01123		—	n^{α}	n^{α}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01124		—	$n^{\alpha} n_{\alpha}=s$	$n^{\alpha} n_{\alpha} = s$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01125		—	$\tau_{\alpha}^{\star}$	$\tau_{\alpha}^{\star}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01126		—	X	X
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01127		—	Y	Y
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01128		—	$\Sigma_{\hat{0}} \neq 0$	$\Sigma_{\hat{0}} \neq 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01129		—	$\tau_{\hat{1} \hat{2}} =$	$\tau_{\hat{1} \hat{2}} =$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01130		—	$-\tau_{\hat{2} \hat{1}} \neq 0$	$-\tau_{\hat{2} \hat{1}} \neq 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01131		—	$x^{\{\alpha\} \beta} R_{\alpha \beta} = R_{\hat{1} \hat{2}} = -R_{\hat{2} \hat{1}} \neq 0$	$x^{[\alpha} y^{\beta]} R_{\alpha \beta} = R_{\hat{1} \hat{2}} = -R_{\hat{2} \hat{1}} \neq 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01132		—	$\left. N^{\alpha} = n \right\rfloor \vartheta^{\alpha}$	$N^{\alpha} = n \rfloor \vartheta^{\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01133		—	$d(\mathscr{A} \wedge d\mathscr{A})$	$d(\mathscr{A} \wedge d\mathscr{A})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01134		—	$d\,C_{\mathrm{T}}$	dC_{T}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01135		—	$d^{*}\,\mathscr{A}$	$d^{*}\mathscr{A}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01136		—	$\mathscr{N}=1$	$\mathcal{N}=1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01137		—	$\left.\Psi_{\alpha}:=e_{\alpha}\right\rfloor\Psi$	$\Psi_{\alpha}:=e_{\alpha}\rfloor\Psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01138		—	$i,j,k,\,\ldots=0,1,2$	$i,j,k,\ldots=0,1,2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01139		—	$\alpha,\beta,\,\ldots=\hat{0},\,\hat{1},\,\hat{2}$	$\alpha,\beta,\ldots=\hat{0},\hat{1},\hat{2}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01140		—	$\bar{\Psi}:=\Psi^{\dagger}\gamma^0$	$\bar{\Psi}:=\Psi^{\dagger}\gamma^0$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01141		—	$\backslash\Psi:=\backslash\Psi_{\alpha}\ \vartheta^{\alpha}$	$\Psi := \Psi_{\alpha} \vartheta^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01142		—	$\backslash\Psi=C\ \bar{\backslash\Psi}^{\mathrm{T}}$	$\Psi = C \bar{\Psi}^{\mathrm{T}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01143		—	$C=-i\ \gamma_{\{0\}}$	$C = -i\gamma_0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01144		—	$C^{\{\dagger\}}=$	$C^{\dagger} =$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01145		—	$C^{\{-1\}},\ \quad C^{\mathrm{T}}=-C$	$C^{-1},\ \ C^{\mathrm{T}} = -C$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01146		—	$C^{\{-1\}}\ \gamma^{\{\alpha\}}\ C=-\left(\gamma^{\{\alpha\}}\right)^{\mathrm{T}}$	$C^{-1}\gamma^{\alpha}C = -(\gamma^{\alpha})^{\mathrm{T}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01147		—	$\gamma^{\{\alpha\}}$	γ^{α}
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01148		—	L_{RS}	L_{RS}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01149		—	$\vartheta^\alpha, \Gamma_\alpha^\star, \Psi$	$\vartheta^\alpha, \Gamma_\alpha^\star, \Psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01150		—	c	c
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01151		—	$\overline{c \ \gamma \ \sigma} = c \ \overline{\sigma} \ \gamma$	$\overline{c \gamma \sigma} = c \overline{\sigma} \gamma$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01152		—	$\delta L \ / \ \delta \Gamma_\alpha^\star \ \cong \ 0$	$\delta L / \delta \Gamma_\alpha^\star \cong 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01153		—	$C = -m$	$c = -m$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01154		—	$\gamma \wedge D \Psi$	$\gamma \wedge D \Psi$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01155		—	L_{∞}	L_{∞}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01156		—	$\mathrm{N}=2$	$\mathcal{N} = 2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01157		—	$\mathrm{D}=3$	$D = 3$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01158		—	$(2+1)$	$(2 + 1)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01159		—	$1 / 2$	$1/2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01160		—	$D^{\left(-\frac{1}{2}, \frac{3}{2}\right)} \otimes D^{\left(\frac{1}{2}, \frac{3}{2}\right)} \left(\widetilde{SO}_0(1,3)\right)$	$D^{\left(-\frac{1}{2}, \frac{3}{2}\right)} \otimes D^{\left(\frac{1}{2}, \frac{3}{2}\right)} \left(\widetilde{SO}_0(1,3)\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01161		—	$S_0(\mathrm{circ})(1,3)$	$SO_0(1,3)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01162		—	$P_{\circ}:=\mathbb{R}^4\otimes S_{0_{\circ}}(1,3)$	$P_{\circ}:=\mathbb{R}^4\otimes SO_{\circ}(1,3)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01163		—	$\psi_{\infty}:=\psi_{1/2}\otimes\psi_{5/2}\otimes\psi_{9/2}\otimes\ldots$	$\psi_{\infty}:=\psi_{1/2}\otimes\psi_{5/2}\otimes\psi_{9/2}\otimes\ldots$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01164		—	$\mathrm{S0}(s,\,n-s)$	$SO(s,n-s)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01165		—	$s\neq 0$	$s\neq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01166		—	$S_{0_{\circ}}(s,\,n-s)$	$SO_{\circ}(s,n-s)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01167		—	$\widetilde{S_{0_{\circ}}}(s,\,n-s)(=:\operatorname{Spin}(n))$	$\widetilde{SO_{\circ}}(s,n-s)(=:\operatorname{Spin}(n))$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01168		—	$s=0)$	$s=0)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01169		—	$\Lambda: \tilde{\mathrm{G}} \rightarrow \mathrm{G}$	$\Lambda: \tilde{G} \rightarrow G$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01170		—	$C(s, n-s)$	$C(s, n-s)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01171		—	γ_{α}	γ_{α}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01172		—	$N \times N$	$N \times N$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01173		—	$N=2^{\lfloor n/2 \rfloor}$	$N=2^{\lfloor n/2 \rfloor}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01174		—	$\mathbf{E}^n$	$\mathbf{E}^n$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01175		—	$\mathfrak{g} \in \mathfrak{so}(s, n-s)$	$\mathfrak{g} \in \mathfrak{so}(s, n-s)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01176		—	$2^{\mathbb{N}}$	$2^{\mathbb{N}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01177		—	$\widetilde{SO}(1,3) \approx SL(2, \mathbb{C})$	$\widetilde{SO}_o(1,3) \approx SL(2, \mathbb{C})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01178		—	$\widetilde{SO}(s, n-s)$	$\widetilde{SO}_o(s, n-s)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01179		—	$\mathrm{P}_0 := \mathbb{R}^n \otimes \widetilde{SO}$	$\mathrm{P}_0 := \mathbb{R}^n \otimes \widetilde{SO}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01180		—	$(s, n-s)$	$(s, n-s)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01181		—	$L^{\{\mathrm{g}\}}(M)$	$L^g(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01182		—	$L_{\circ}^{\circ\mathrm{g}}(M)$	$L_{\circ}^g(M)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01183		—	$\mathrm{f}: \tilde{L}(M) \rightarrow L_{\circ}^g(M)$	$\mathbf{f}: \tilde{L}(M) \rightarrow L_{\circ}^g(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01184		—	$L_{\circ}^g(M)$	$L_{\circ}^g(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01185		—	$\tilde{L}^{\prime}(M), f^{\prime}$	$\tilde{L}^{\prime}(M), f^{\prime}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01186		—	$f^{\prime}$	$f^{\prime}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01187		—	$\mathrm{w}_{\{2\}} \in \mathrm{H}^{\{2\}}\!\!\left(M, \mathbb{Z}_{\{2\}}\right)$	$\mathbf{w}_2 \in \mathrm{H}^2(M, \mathbb{Z}_2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01188		—	$\mathrm{g}^{\{\text{th}\}}$	$\mathbf{g}^{\text{th}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01189		—	$\mathrm{H}^{\{\mathrm{g}\}}\!\!\left(M, \mathbb{Z}_{\{2\}}\right)$	$\mathrm{H}^{\mathfrak{g}}(M, \mathbb{Z}_2)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01190		—	$\mathrm{H}^1\left(M, \mathbb{Z}_2\right)$	$\tilde{s} = \dim H^1(M, \mathbb{Z}_2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01191		—	$\tilde{s}=2^{\{v\}}$	$\tilde{s} = 2^v$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01192		—	ν	ν
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01193		—	$W^{\{3\}}:=$	$W^3 :=$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01194		—	$\mathrm{w}_{\{1\}}=0$	$w_1 = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01195		—	$\mathrm{H}^{\circ}\left(M, \mathbb{Z}_2\right)$	$H^{\circ}(M, \mathbb{Z}_2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01196		—	$S^{\{7\}}$	S^7
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01197		—	(1968,1970)	(1968,1970)
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01198		—	$T(M)$	$T(M)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01199		—	$S\ U(3)$	$SU(3)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01200		—	$\tilde{\mathrm{L}}(\mathrm{M})$	$\tilde{\mathrm{L}}(\mathrm{M})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01201		—	D^{F}	D^{F}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01202		—	$\widetilde{S\ 0}_{\circ}$	$\widetilde{SO}_{\circ}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01203		—	$\widetilde{S\ 0}_{\circ}(1,3)$	$\widetilde{SO}_{\circ}(1,3)$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01204		—	$S\,L(2,\,\mathbb{C})\,\approx\,\widetilde{SO}_{\circ}(1,3)$	$SL(2,\mathbb{C})\approx SO_{\circ}(1,3)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01205		—	$\left(-\frac{1}{2},\,\frac{3}{2}\right)\oplus\left(\frac{1}{2},\,\frac{3}{2}\right)$	$\left(-\frac{1}{2},\frac{3}{2}\right)\oplus\left(\frac{1}{2},\frac{3}{2}\right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01206		—	$\frac{1}{2}$	$\frac{1}{2}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01207		—	$\widetilde{0}(1,3)$	$\tilde{O}(1,3)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01208		—	$\widetilde{\mathbf{b}}_{\{(1)\}},\,i=1,\,\ldots,\,N$	$\tilde{b}_{(1)},i=1,\ldots,N$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01209		—	$\mathscr{G}_{\{S\}}$	$\mathscr{G}_S$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01210		—	$\mathscr{G}_{\{p\}}$	$\mathscr{G}_p$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01211		—	$\tilde{L}(M)$	$\tilde{L}(M)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01212		—	$\gamma^{\mathrm{n}+1}$	$\gamma^{\mathrm{n}+1}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01213		—	γ^{n+1}	γ^{n+1}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01214		—	$\tilde{\omega}$	$\tilde{\omega}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01215		—	$\Gamma_{i\{ }^{\{\alpha } \beta \}$	$\Gamma_i^{\alpha \beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01216		—	$\Gamma_{i\ j\{ }^{\{\cdots k\}$	$\Gamma_{ij}^{\cdots k}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01217		—	$\mathbb{R}^n \otimes S\,0$	$\mathbb{R}^n \otimes SO$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01218		—	$\mathrm{e}_{\alpha}^{-i}(m)$	$\mathfrak{e}_{\alpha}^{-i}(m)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01219		—	$\mathcal{D}(\mathrm{M})$	$\mathcal{D}(\mathbf{M})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01220		—	$\delta L_{\mathrm{D}} / \delta \bar{\psi}$	$\delta L_{\mathbf{D}} / \delta \psi$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01221		—	$\delta L_{\mathrm{D}} / \delta \psi$	$\delta L_{\mathbf{D}} / \delta \psi$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01222		—	$\mathrm{D} \gamma = 0$	$\mathbf{D} \gamma = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01223		—	$\tau := \delta L / \delta \omega$	$\tau := \delta \mathbf{L} / \delta \omega$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01224		—	$\tau = i \bar{\psi} \gamma \psi$	$\tau = i \psi \gamma \psi$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01225		—	τ_{s}	τ_s
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01226		—	$j_{n+1}:=i\,\overline{\psi}^{\,*}\,\gamma\,\gamma^{n+1}\,\psi$	$j_{n+1} := i\psi^*\gamma\gamma^{n+1}\psi$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01227		—	$\mathrm{H}^{\mathrm{n}-1}$	$\mathrm{H}^{\mathrm{n}-1}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01228		—	$\tau_{\mathrm{s}}^{\mathrm{0}}$	τ_s^0
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01229		—	$\tilde{\omega}^{\{\ \ \}}$	$\tilde{\omega}^{\{ \}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01230		—	$\mathrm{D}^{\{\ \ \}}$	$\mathrm{D}^{\{ \}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01231		—	$\Theta=[K,\,\vartheta]$	$\Theta = [K, \vartheta]$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01232		—	$[\Theta, \vartheta]$	$[\Theta, \vartheta]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01233		—	$\delta L_D / \delta \psi$	$\delta L_D / \delta \psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01234		—	$D=\vartheta^\alpha \nabla_\alpha$	$D = \vartheta^\alpha \nabla_\alpha$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01235		—	(1938,1957)	(1938,1957)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01236		—	P_C	PC
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01237		—	$\psi := \binom{\psi_p}{\psi_N}$	$\psi := \binom{\psi_p}{\psi_N}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01238		—	$SL(2, \mathbb{C})$	$SL(2, \mathbb{C})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01239		—	$\overline{\mathrm{G}}=S\,L(2\,\mathrm{f},\,\mathbb{C})$	$\overline{G} = SL(2f, \mathbb{C})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01240		—	$\overline{\mathrm{G}}$	$\overline{G}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01241		—	$1\,/\,r^{\mathrm{2}}$	$1/r^2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01242		—	$r\!:=\! \overrightarrow{\mathrm{x}} $	$r := \overrightarrow{\mathbf{x}} $
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01243		—	$(1934,1935)$	$(1934,1935)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01244		—	$n\!=\!4$	$n = 4$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01245		—	$\varepsilon=\,\mathrm{pm}\,1,\,\mathrm{ell}$	$\varepsilon = \pm 1, \ell$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01246		—	$v=v(\rho)$	$v = v(\rho)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01247		—	$\mu=\mu(\rho)$	$\mu = \mu(\rho)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01248		—	$\mathrm{r}:= \overrightarrow{\mathrm{x}} , \vartheta$	$\mathbf{r} := \overrightarrow{\mathbf{x}} , \vartheta$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01249		—	$\overrightarrow{\mathrm{x}}$	$\overrightarrow{\mathbf{x}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01250		—	$g_{\mathrm{k\ l}}$	g_{kl}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01251		—	$\Gamma_{\kappa}^{f\cdot\alpha\beta}$	$\Gamma_{\kappa}^{f\cdot\alpha\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01252		—	$\mathrm{R}_{\circ}$	$\mathrm{R}_{\circ}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01253		—	$\lambda, \mathbf{y}$	$\lambda, \mathbf{y}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01254		—	$S_{0\circ}(1,4)$	$SO_{\circ}(1,4)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01255		—	$\bar{\Gamma}_k$	Γ_k
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01256		—	$i \, \vec{\gamma} \cdot \vec{\partial}$	$i\vec{\gamma} \cdot \vec{\partial}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01257		—	$\exp(-\mu/2 - \nu/4)$	$\exp(-\mu/2 - \nu/4)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01258		—	χ_k^m	χ_k^m
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01259		—	$\mathrm{P}=(-1)^l$	$P=(-1)^l$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01260		—	$\mathbb{L}$	$\mathbb{L}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01261		—	$\mathrm{C}\left(\mathbb{L} \frac{1}{2} \mathrm{j} ; \mathrm{m}-\overline{\mathrm{m}}, \overline{\mathrm{m}}\right)$	$\mathrm{C}\left(\frac{1}{2} \mathrm{j} ; \mathrm{m}-\overline{\mathrm{m}}, \overline{\mathrm{m}}\right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01262		—	$\mathrm{Y}_{\mathbb{L}}^m(\vartheta, \phi)$	$\mathrm{Y}_l^m(\vartheta, \phi)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01263		—	$\vec{\mathbb{J}}$	$\vec{J}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01264		—	$\vec{\mathbb{L}}$	$\vec{L}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01265		—	$ \kappa =1$	$ \kappa =1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01266		—	$\mathrm{j}=\frac{1}{2}, \quad \mathbb{L}=0,1$	$j=\frac{1}{2}, l=0,1$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01267		—	$\mathrm{m}=\pm\frac{1}{2}$	$m = \pm \frac{1}{2}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01268		—	$\int L_{\mathrm{D}^{\prime}}$	$\int L_{\mathrm{D}'}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01269		—	$\rho^{\ast}$	ρ^*
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01270		—	$M^{\ast}$	M^*
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01271		—	$j=\frac{1}{2}$	$j = \frac{1}{2}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01272		—	$\kappa=-1$	$\kappa = -1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01273		—	$H(\rho)$	$H(\rho)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01274		—	$l=0$	$l = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01275		—	$l=1$	$l = 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01276		—	$F(\rho)$	$F(\rho)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01277		—	$ \varphi ^6$	$ \varphi ^6$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01278		—	$\mathrm{Y}_{\mathrm{l}}^{\mathrm{m}}(\vartheta, \phi)$	$Y_l^m(\vartheta, \phi)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01279		—	$ \omega <1$	$ \omega < 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01280		—	$ \omega >1$	$ \omega > 1$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01281		—	$ \psi \sim e^{-m_{\pi} r}$	$ \psi \sim e^{-m_{\pi} r}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01282		—	m_{s}	m_{s}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01283		—	Σ_{D}	Σ_{D}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01284		—	m_{s}	m_s
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01285		—	(1951,1956)	(1951,1956)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01286		—	$a=0$	$a = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01287		—	$a=1$	$a = 1$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01288		—	$\varepsilon=+1$	$\varepsilon = +1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01289		—	$S_{1 / 2}$	$S_{1/2}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01290		—	$Q=e$	$Q = e$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01291		—	$D_{\mathrm{F}}(x)$	$D_{\mathrm{F}}(x)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01292		—	$(1957,1958)$	$(1957,1958)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01293		—	$\ell^*=8.1 \times 10^{-33} \mathrm{~cm}$	$\ell^* = 8.1 \times 10^{-33} \mathrm{~cm}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01294		—	$\ell=10^{-13}$	$\ell = 10^{-13}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01295		—	$\ell=10^{-13} \mathrm{~cm}$	$\ell = 10^{-13} \text{ cm}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01296		—	$\langle \text{angle } d \text{ } j \rangle = 0$	$\langle dj \rangle = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01297		—	$\langle \text{angle } d \text{ } j_5 \rangle \neq 0$	$\langle dj_5 \rangle \neq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01298		—	$j_5(x \text{ ; } \varepsilon) := \overline{\psi}(x) \gamma_5 \gamma \psi(x + \varepsilon)$	$j_5(x; \varepsilon) := \psi(x) \gamma_5^* \gamma \psi(x + \varepsilon)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01299		—	$\delta / \delta A$	$\delta / \delta A$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01300		—	$j_5(x \text{ ; } \varepsilon)$	$j_5(x; \varepsilon)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01301		—	$x^i \rightarrow x^i + \varepsilon^i$	$x^i \rightarrow x^i + \varepsilon^i$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01302		—	$P=\bar{\psi} \gamma_5 \psi$	$P = \psi \gamma_5 \psi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01303		—	$C:=A \wedge d A$	$C := A \wedge dA$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01304		—	$\mathbf{A} \cdot \mathbf{B}$	$\mathbf{A} \cdot \mathbf{B}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01305		—	$\mathscr{D}:=D+i A \wedge$	$\mathscr{D} := D + iA\wedge$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01306		—	$D \psi:=d \psi+\Gamma \wedge$	$D\psi := d\psi + \Gamma\wedge$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01307		—	$\Gamma^{\alpha \beta}=\Gamma_{i}^{\alpha \beta} d x^i$	$\Gamma^{\alpha \beta} = \Gamma_i^{\alpha \beta} dx^i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01308		—	$\Theta:=D \gamma$	$\Theta := D\gamma$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01309		—	$\left.\left.T:=\frac{1}{4}\operatorname{Tr}(\check{\gamma}\rfloor\Theta)\right)=e_{\alpha}\right\lrcorner T^{\alpha}$	$T:=\frac{1}{4}\operatorname{Tr}(\check{\gamma}\rfloor\Theta)=e_{\alpha}\lrcorner T^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01310		—	$K=\frac{i}{4}K^{\alpha\beta}\sigma_{\alpha\beta}$	$K=\frac{i}{4}K^{\alpha\beta}\sigma_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01311		—	$D\gamma=[\gamma,K]=\gamma_{\alpha}T^{\alpha}$	$D\gamma=[\gamma,K]=\gamma_{\alpha}T^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01312		—	$D^{\{t\}}\gamma=0$	$D^{(t)}\gamma=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01313		—	$\gamma\rightarrow\gamma^{\beta}=$	$\gamma\rightarrow\gamma^{\beta}=$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01314		—	$e^{i\gamma^5}\gamma e^{-i\gamma^5}$	$e^{i\gamma^5}\gamma e^{-i\gamma^5}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01315		—	$P:\vartheta^B\rightarrow-\vartheta^B$	$P:\vartheta^B\rightarrow-\vartheta^B$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01316		—	$B=$	$B =$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01317		—	$1,2,3$	$1,2,3$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01318		—	$d\,j\,\simeq 0$	$dj \simeq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01319		—	$m\rightarrow 0$	$m \rightarrow 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01320		—	$d\,j_{\{5\}}=0$	$dj_5 = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01321		—	$\mathscr{A}=2\,d\,\theta$	$\mathscr{A} = 2d\theta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01322		—	$a=\theta f_a$	$a = \theta f_a$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01323		—	$U(1)_{\mathrm{PQ}}$	$U(1)_{\mathrm{PQ}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01324		—	$\eta^{\alpha\beta} := \left(\vartheta^\alpha \wedge \vartheta^\beta \right)$	$\eta^{\alpha\beta} := \left(\vartheta^\alpha \wedge \vartheta^\beta \right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01325		—	$\left\langle dj_5 \right\rangle$	$\langle dj_5 \rangle$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01326		—	$d\,C_{\mathrm{TT}} \cong (1/4)\,dj_5 \rightarrow 0$	$dC_{\mathrm{TT}} \cong (1/4) dj_5 \rightarrow 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01327		—	$\{ \}^* j_5$	$^* j_5$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01328		—	$n=4\,k+2$	$n = 4k + 2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01329		—	$d^* \mathscr{A} \sim dC_{\mathrm{TT}}$	$d^* \mathscr{A} \sim dC_{\mathrm{TT}}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01330		—	$d \, \mathrm{d} \wedge \mathrm{d} = -2 \, \mathrm{d} \, \mathrm{d} \cdot \mathrm{d} \, \mathrm{d}$	$d\mathcal{A} \wedge d\mathcal{A} = -2\mathcal{E} \cdot \mathcal{B}\eta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01331		—	$F \wedge F = dA \wedge dA$	$F \wedge F = dA \wedge dA$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01332		—	$d^* \, \mathrm{d} \wedge \mathrm{d} = 4 \, \mathrm{d} \, \mathrm{d} \cdot \mathrm{d} \, \mathrm{d} \wedge \mathrm{d} = 4 \, \mathrm{d} \, \mathrm{d} \cdot \mathrm{d} \, \mathrm{d} \wedge \mathrm{d}$	$d^* \mathcal{A} \wedge (d^* \mathcal{A}) = 4\ell^4 V_{\mathrm{NY}} \wedge^* V_{\mathrm{NY}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01333		—	$d^* \, \mathrm{d} = 2 \, \mathrm{d} \, \mathrm{d} \cdot \mathrm{d} \, \mathrm{d} \wedge \mathrm{d} = 2 \, \mathrm{d} \, \mathrm{d} \cdot \mathrm{d} \, \mathrm{d} \wedge \mathrm{d}$	$d^* \mathcal{A} = 2\ell^2 dC_{\mathrm{TT}} = 2\ell^2 V_{\mathrm{NY}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01334		—	Z	Z
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01335		—	$\Psi = \Psi_i \, dx^i$	$\Psi = \Psi_i dx^i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01336		—	$J_5 := i \, \bar{\Psi} \wedge \gamma \wedge \Psi$	$J_5 := i\Psi \wedge \gamma \wedge \Psi$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01337		—	t \rightarrow +0	$t \rightarrow +0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01338		—	$\mathbb{A} :=$	$\mathbb{A} :=$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01339		—	$\check{\gamma} \rfloor A = \gamma^\alpha e_\alpha \downarrow A = (-1)^{s+1} [* \gamma \wedge A]$	$\tilde{\gamma} \rfloor A = \gamma^\alpha e_\alpha \downarrow A = (-1)^{s+1} [* \gamma \wedge A]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01340		—	$\square :=$	$\square :=$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01341		—	$\frac{\partial_i \left(\sqrt{ g } g^{ij} \partial_j \right)}{\sqrt{ g }}$	$\partial_i \left(\sqrt{ g } g^{ij} \partial_j \right) / \sqrt{ g }$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01342		—	$K_k \left(x, \not{D}^2 \right)$	$K_k \left(x, \not{D}^2 \right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01343		—	$\not{D}^2$	$\not{D}^2$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01344		—	$\mathrm{o}_{\alpha\beta} =$	$\mathrm{o}_{\alpha\beta} =$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01345		—	$(-1, \ldots, -1)$	$(-1, \ldots, -1)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01346		—	$k=1,3, \ldots$	$k=1,3,\ldots$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01347		—	$d\,\mathrm{scr}{K}=d^{\wedge{*}}\,\stackrel{\smile}{D}\,\wedge^{\wedge{*}}\,\breve{D}^{\wedge{*}}\,\mathrm{scr}{A}$	$d\mathscr{K} = d^*\widetilde{D} \wedge^* \breve{D}^* \mathscr{A}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01348		—	$D=D^{\wedge\{\backslash\backslash\}}+i\,\mathrm{scr}{A}\,\gamma_5\,/\,4$	$D = D^{\mathfrak{U}} + i\mathscr{A}\gamma_5/4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01349		—	$K_{\{2\}}$	K_2
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01350		—	$K_{\{4\}}$	K_4
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01351		—	$k=4$	$k = 4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01352		—	$t^{\{(k-4) / 2\}}=1$	$t^{(k-4)/2} = 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01353		—	$K_{\{2\}} \sim$	$\bar{K}_2 \sim$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01354		—	$d^{\{*\}} \backslash \mathrm{mathscr{A}}=2 \backslash \mathrm{ell}^{\{2\}} \mathrm{d} \mathrm{C}_{\{\mathrm{TT}\}}$	$d^* \mathcal{A} = 2 \ell^2 d C_{\mathrm{TT}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01355		—	$[\backslash \mathrm{text} \{ \mathrm{length} \}]^{\{-3 / 2\}}$	$[\mathrm{length}]^{-3/2}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01356		—	$\backslash \mathrm{psi} \sim m^{\{3 / 2\}}$	$\psi \sim m^{3/2}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01357		—	$t=1 / M^{\{2\}}$	$t = 1/M^2$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01358		—	$M \rightarrow \infty$	$M \rightarrow \infty$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01359		—	$m \neq 0$	$m \neq 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01360		—	$\vartheta^\alpha \rightarrow \widetilde{\vartheta}^\alpha = M \vartheta^\alpha$	$\vartheta^\alpha \rightarrow \widetilde{\vartheta}^\alpha = M \vartheta^\alpha$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01361		—	$[\hbar]$	$[\hbar]$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01362		—	$(n=2)$	$(n = 2)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01363		—	$\sigma_{\alpha} := \Sigma_{\alpha} - D \mu_{\alpha}$	$\sigma_{\alpha} := \Sigma_{\alpha} - D \mu_{\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01364		—	$\stackrel{\scriptscriptstyle (\pm)}{\Gamma}(\pm)$	$\stackrel{\scriptscriptstyle (\pm)}{\Gamma}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01365		—	$d\,C_{\mathrm{RR}}^{\{\}}^{\backslash\{ \backslash\}}$	$dC_{\mathrm{RR}}^{\{\}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01366		—	$\Lambda \neq 0$	$\Lambda \neq 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01367		—	$\underline{C}_{\mathrm{RR}}^{\{\}}^{\backslash\{ \backslash\}}$	$\underline{C}_{\mathrm{RR}}^{\{\}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01368		—	$\stackrel{(\pm)}{\mathrm{rel}}\{(\ \mathrm{pm})\}\{\underline{\Gamma}\}$	$\stackrel{(\pm)}{\underline{\Gamma}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01369		—	$\mathrm{mathscr{C}} := \Gamma / \mathrm{mathscr{G}}$	$\mathcal{C} := \Gamma / \mathcal{G}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01370		—	$\mathrm{t}_n := n \downarrow D + D\ n$	$\mathbb{L}_n := n \downarrow D + Dn$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01371		—	τ^A	τ^A
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01372		—	$\tau^A := (1/2) \eta^A{}_{\beta\gamma} \tau^{\beta\gamma}$	$\tau^A := (1/2)\eta^{0A\beta\gamma}\tau_{\beta\gamma}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01373		—	$\gamma_\alpha\gamma_\beta + \gamma_\beta\gamma_\alpha = 2\mathrm{o}_{\alpha\beta}$	$\gamma_\alpha\gamma_\beta + \gamma_\beta\gamma_\alpha = 2\mathbf{o}_{\alpha\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01374		—	$\eta^\alpha :=$	$\eta^\alpha :=$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01375		—	$\{ \}^* \vartheta^\alpha$	$^*\vartheta^\alpha$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01376		—	$\Gamma := \frac{i}{4} \Gamma_i^{\alpha\beta} \sigma_{\alpha\beta} dx^i$	$\Gamma := \frac{i}{4}\Gamma_i^{\alpha\beta}\sigma_{\alpha\beta}dx^i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01377		—	$D = d + \Gamma \wedge$	$D = d + \Gamma \wedge$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01378		—	$\sigma_{\alpha\beta} = \frac{i}{2} \left(\gamma_\alpha\gamma_\beta - \gamma_\beta\gamma_\alpha \right)$	$\sigma_{\alpha\beta} = \frac{i}{2} \left(\gamma_\alpha\gamma_\beta - \gamma_\beta\gamma_\alpha \right)$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01379		—	$\sigma:=\frac{i}{2}\,\gamma\wedge\gamma=\frac{1}{2}\sigma_{\alpha\beta}\vartheta^\alpha\wedge\vartheta^\beta$	$\sigma:=\frac{i}{2}\gamma\wedge\gamma=\frac{1}{2}\sigma_{\alpha\beta}\vartheta^\alpha\wedge\vartheta^\beta$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01380		—	$R:=\left(R^\alpha{}_\beta\wedge\eta_{\beta\alpha}\right)$	$R:=\left(R^{\alpha\beta}\wedge\eta_{\beta\alpha}\right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01381		—	$\mathfrak{sl}(5,\mathbb{R})$	$\mathfrak{sl}(5,\mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01382		—	$\hat{\Gamma}=\frac{1}{\ell}\left(\vartheta^\alpha L^4{}_\alpha+\vartheta_\beta L^\beta{}_4\right)$	$\hat{\Gamma}=\Gamma+\left(\vartheta^\alpha L^4{}_\alpha+\vartheta_\beta L^\beta{}_4\right)/\ell$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01383		—	$\hat{C}_{\mathrm{RR}}$	$\hat{C}_{\mathrm{RR}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01384		—	$\pi^0\rightarrow 2\,\gamma$	$\pi^0\rightarrow 2\gamma$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01385		—	$\mathbb{R}^4\otimes\mathrm{SO}(1,3)$	$\mathbb{R}^4\otimes\mathrm{SO}(1,3)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01386		—	$\mathrm{d} B=0$	$dB = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01387		—	$\sigma=\sigma_{\mathrm{i}}\,\mathrm{d} x^{\mathrm{i}}$	$\sigma = \sigma_i dx^i$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01388		—	$\mathrm{SL}(5,\,\mathbb{R})$	$\mathrm{SL}(5,\mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01389		—	$\operatorname{SL}(n+1,\,\mathbb{R})$	$\mathrm{SL}(n+1,\mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01390		—	$\operatorname{det} g=1$	$\det g = 1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01391		—	$n(n+2)$	$n(n+2)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01392		—	$\mathfrak{gl}(n,\,\mathbb{R})$	$\mathfrak{gl}(n,\mathbb{R})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01393		—	$\backslash\alpha, \backslash\ldots, \backslash\delta=0, \backslash\ldots, n-1$	$\alpha, \dots, \delta = 0, \dots, n - 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01394		—	$\backslash\ell\ell\ P_{\backslash\alpha}:=\mathfrak{L}^{\backslash n+1}\{ \ }_{\backslash\alpha}=\mathfrak{L}^{\backslash n+1}\{ \ }_{\backslash\alpha}$	$\ell P_{\alpha} := \mathfrak{L}^{n+1}_{\alpha} = L^{n+1}_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01395		—	$\backslash\ell\ell\ P_{\{*\}^{\backslash\beta}}:=\swarrow^{\backslash\beta}\{ \ }_{\backslash n+1}=\mathfrak{L}^{\backslash\beta}\{ \ }_{\backslash n+1}$	$\ell P_{*}^{\beta} := \swarrow^{\beta}_{n+1} = L^{\beta}_{n+1}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01396		—	$P_{\{*\}^{\backslash\alpha}}$	P_{*}^{α}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01397		—	$P_{\backslash\beta}$	P_{β}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01398		—	$\backslash\operatorname{GL}\{n, \backslash\mathbb{R}\}$	$\operatorname{GL}(n, \mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01399		—	$\backslash\mathrm{A}_{\{*\}}(n, \backslash\mathbb{R})$	$\mathbf{A}_{*}(n, \mathbb{R})$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01400		—	$\mathrm{SO}(n+1)$	$\mathrm{SO}(n+1)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01401		—	$\hat{g}_{AB}$	$\hat{g}_{AB}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01402		—	$n(n+1) / 2$	$n(n+1)/2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01403		—	$A(x) \in \mathrm{A}_*(n, R)$	$A(x) \in \mathbf{A}_*(n, R)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01404		—	$\Lambda(x) \in \mathrm{GL}(n, \mathbb{R})$	$\Lambda(x) \in \mathrm{GL}(n, \mathbb{R})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01405		—	$\tau(x) := \exp \left[\xi^{\alpha} P_{\alpha} \right] \in \mathrm{mathscr{T}}(n, R)$	$\tau(x) := \exp \left[\xi^{\alpha} P_{\alpha} \right] \in \mathcal{T}(n, R)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01406		—	$\tau_{*}(x) := \exp \left[\xi^{\alpha} P_{\alpha} \right] \in \mathrm{mathscr{T}}_{*}(n, R)$	$\tau_{*}(x) := \exp \left[\xi^{\alpha} P_{\alpha} \right] \in \mathcal{T}_{*}(n, R)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01407		—	$\ell \rightarrow \infty$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01408		—	$\operatorname{Ad} A(B) = [A, B]$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01409		—	$\hat{\Gamma}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01410		—	$\Gamma^{(L)}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01411		—	$\mathrm{GL}(n, \mathbb{R})$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01412		—	$D \tau(x) := d \tau(x) + \Gamma^{(L)} \tau(x)$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01413		—	$\Gamma^{(T)}$	
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01414		—	$\Gamma_*^{(T)}$	$\Gamma_*^{(T)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01415		—	$\xi=\xi^{\alpha} P_{\alpha}$	$\xi = \xi^{\alpha} P_{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01416		—	$\xi_*=\xi_{\beta}^* P_{\beta}^*$	$\xi_* = \xi_{\beta}^* P_{\beta}^*$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01417		—	$\mathscr{T}(n, R) \otimes \mathscr{T}_*(n, R)$	$\mathcal{T}(n, R) \otimes \mathcal{T}_*(n, R)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01418		—	$\xi=\xi^*=0$	$\xi = \xi^* = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01419		—	$\theta_{\alpha}=g_{\alpha\beta} \vartheta^{\beta}$	$\theta_{\alpha} = g_{\alpha\beta} \vartheta^{\beta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01420		—	$D \, \xi^{\alpha} \stackrel{*}{=} \vartheta^{\alpha}$	$D \xi^{\alpha} \stackrel{*}{=} \vartheta^{\alpha}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01421		—	ξ^{α}	ξ^{α}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01422		—	$1/[$	$1/[$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01423		—	$]$	$]$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01424		—	P_*^{β}	P_*^{β}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01425		—	$\vartheta^{\alpha}=\ell\ \Gamma_{\hat{4}}^{\alpha}$	$\vartheta^{\alpha}=\ell\Gamma_4^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01426		—	$\theta_{\beta}=\ell\Gamma_{\beta}^{\hat{4}}$	$\theta_{\beta}=\ell\Gamma_{\beta}^4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01427		—	$\Lambda=3\ /\ \ell^2$	$\Lambda=3/\ell^2$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01428		—	$d\,\hat{C}$	$d\hat{C}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01429		—	$\Gamma_{A\{ \} }^{\{ \} B}$	Γ_A^{B}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01430		—	$\hat{B}$	$\hat{B}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01431		—	$\hat{R}=0$	$\hat{R}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01432		—	$\eta_{A\,B\,C\,D\,E}:=\sqrt{\hat{g}}\,\,\,\varepsilon_{A\,B\,C\,D\,E}$	$\eta_{ABCDE}:=\sqrt{\hat{g}}\varepsilon_{ABCDE}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01433		—	$\varepsilon_{A\,B\,C\,D\,E}$	ε_{ABCDE}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01434		—	$\hat{g}$	$\hat{g}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01435		—	$B^{\{A\ B\}}:=\hat{g}^{\{g\}^{\{A\ C\}}\ B_{\{C\}\{ \}^{\{B\}}}$	$B^{AB}:=\hat{g}^{AC}B_C{}^B$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01436		—	$\operatorname{SO}(2,3)$	$\operatorname{SO}(2,3)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01437		—	$\eta_{\{A\ B\ C\ D\ E\}}$	η_{ABCDE}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01438		—	$\Phi^{\{A\}=\left\{\phi^{\{\alpha\}},\ \phi^{\{\hat{4}\}}\right\}}$	$\Phi^A=\left\{\phi^\alpha,\phi^{\hat{4}}\right\}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01439		—	$U\left(\Phi^{\{B\}}\ \Phi_{\{B\}}\right)$	$U\left(\Phi^B\Phi_B\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01440		—	$\Phi=\Re\ \Phi+i\ \Im\ \Phi$	$\Phi=\Re\Phi+i\Im\Phi$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01441		—	$d\ C_{\{\mathrm{RR}\}}$	dC_{RR}
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01442		—	$\stackrel{\circ}{\Gamma}^{\alpha\beta} =$	$\stackrel{\circ}{\Gamma}^{\alpha\beta} =$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01443		—	$-\stackrel{\circ}{\Gamma}^{\beta\alpha}$	$-\stackrel{\circ}{\Gamma}^{\beta\alpha}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01444		—	$B^{\alpha\beta}$	$B^{\alpha\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01445		—	$\mu \neq 0$	$\mu \neq 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01446		—	$\vartheta^{\alpha} = g^{\alpha\beta} \theta_{\beta}$	$\vartheta^{\alpha} = g^{\alpha\beta} \theta_{\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01447		—	$\kappa =$	$\kappa =$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01448		—	$8 \pi G_{\mathrm{N}} = 8 \pi \ell_{\mathrm{Planck}}^2$	$8 \pi G_{\mathrm{N}} = 8 \pi \ell_{\mathrm{Planck}}^2$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01449		—	$\rho_{\Lambda}=\Lambda / \kappa=\Omega_{\Lambda} \rho_{\text {crit }} < \rho_{\text {crit }}$	$\rho_{\Lambda}=\Lambda / \kappa=\Omega_{\Lambda} \rho_{\text {crit }} < \rho_{\text {crit }}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01450		—	$\kappa=\mu \ell ^{2} / 4$	$\kappa=\mu \ell ^2 / 4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01451		—	$\Lambda =12 \kappa / \mu \ell ^{4}=3 / \ell ^2$	$\Lambda =12 \kappa / \mu \ell ^4=3 / \ell ^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01452		—	$\mu \simeq 10 ^{-123}$	$\mu \simeq 10 ^{-123}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01453		—	$\hat{B} \wedge \hat{B}$	$\hat{B} \wedge \hat{B}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01454		—	G_{N}	G_{N}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01455		—	$O(5)$	$O(5)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01456		—	$\operatorname{SO}(1,3)$	SO(1,3)
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01457		—	g_{ij}	g_{ij}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01458		—	$\Gamma_{\{4\}}^{\alpha} \otimes_{\mathrm{s}} \Gamma_{\alpha}^4$	$\Gamma_4^{\alpha} \otimes_{\mathrm{s}} \Gamma_{\alpha}^4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01459		—	$\ell^2/\mu^2=\left(\kappa/4\,\Lambda_{\mathrm{H}}\right)$	$\ell^2/\mu^2=(\kappa/4\Lambda_{\mathrm{H}})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01460		—	$W^{\{+\}}$	W^+
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01461		—	$W^{\{-\}}$	W^-
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01462		—	$\mu \rightarrow 0$	$\mu \rightarrow 0$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01463		—	$d\,s$	ds
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01464		—	$1/\alpha :=$	$1/\alpha :=$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01465		—	$\hbar c/e^2 \simeq 137$	$\hbar c/e^2 \simeq 137$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01466		—	$1/\mu \simeq 10^{123}$	$1/\mu \simeq 10^{123}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01467		—	C_{TT}	C_{TT}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01468		—	$\mathfrak{gl}(4,\mathbb{R})$	$\mathfrak{gl}(4,\mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01469		—	$\operatorname{Ric}_{\alpha\beta}:=(-1)^{\text{sig}}*$	$\operatorname{Ric}_{\alpha\beta}:=(-1)^{\text{sig}}*$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01470		—	$\left(R_{\{\alpha\}}^{\{\delta\}}\wedge\eta_{\{\delta\mid\beta\}}\right)$	$\left(R_{\alpha}^{\delta}\wedge\eta_{\delta\mid\beta}\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01471		—	$\mathscr{S}:=\mathscr{M} \,/\, \mathscr{D}$	$\mathcal{S}:=\mathcal{M}\big/\mathcal{D}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01472		—	$\mathscr{D}$	$\mathcal{D}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01473		—	M^{3}	$\mathbf{M}^3$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01474		—	$\mathrm{F}_{\mathrm{i} \, \mathrm{j}}$	$\mathbf{F}_{ij}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01475		—	$\mathbf{g}_{\{\mathrm{i} \, \mathrm{j}\}} \, \mathbf{g}^{\{\mathrm{i} \, \mathrm{j}\}}$	$g_{[ij]}g^{[ij]}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01476		—	$\mathrm{g}_{\{\mathrm{i} \, \mathrm{j}\}}:=\ell^{\ast \, 2} \, \mathrm{\sim F}_{\mathrm{i} \, \mathrm{j}}$	$\mathbf{g}_{[ij]}:=\ell^{\ast 2} \, \mathbf{F}_{ij}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01477		—	$m \in M$	$m \in M$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01478		—	Σ_{ω}	Σ_{ω}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01479		—	$\Sigma_{\text{mat}}=0$	$\Sigma_{\text{mat}} = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01480		—	$\delta=\pi / 2$	$\delta = \pi/2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01481		—	Σ_{YM}	Σ_{YM}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01482		—	$\cos ^{2} \delta+$	$\cos ^2 \delta+$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01483		—	$\sin ^{2} \delta=1$	$\sin ^2 \delta = 1$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01484		—	$R_{ij}^{\{\}}$	$R_{ij}^{\{\}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01485		—	$\bar{\tau}_e$	$\bar{\tau}_e$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01486		—	$\bar{\tau}_p$	$\bar{\tau}_p$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01487		—	$\delta=0$	$\delta=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01488		—	$\xi \wedge \vartheta$	$\xi \wedge \vartheta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01489		—	$\{ \}^{\{ * \}} \xi \wedge \vartheta$	$^* \xi \wedge \vartheta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01490		—	$d \delta$	$d \delta$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01491		—	$\{ \}^{\ast} \ \Sigma_{\mathrm{YM}}(\xi)$	$\ast\Sigma_{\mathrm{YM}}(\xi)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01492		—	$d \ d \ \delta \ \equiv \ 0$	$dd\delta \equiv 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01493		—	$b_{\{1\}}=\operatorname{dim} \ H^{\{1\}}(M)=\nu$	$b_1 = \dim H^1(M) = \nu$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01494		—	$\mathbf{v}$	$\boldsymbol{v}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01495		—	$(1962,1969)$	$(1962,1969)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01496		—	$\mathrm{n}=4+\mathrm{K}$	$\mathbf{n} = 4 + \mathbf{K}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01497		—	$(1926,1928)$	$(1926,1928)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01498		—	M^{4+K}	M^{4+K}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01499		—	$\hat{L}\left(M^{4+K}\right)$	$\hat{L}\left(M^{4+K}\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01500		—	$\hat{\vartheta}$	$\hat{\vartheta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01501		—	$\hat{E} \approx M^4 \times B$	$\hat{E} \approx M^4 \times B$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01502		—	G / H	G/H
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01503		—	$\mathrm{K}=\operatorname{dim} \mathrm{G}$	$K = \dim G$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01504		—	$P\left(M^4, G, \pi, \delta\right)$	$P\left(M^4, G, \pi, \delta\right)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01505		—	$P\left(M^4, N / H, \pi, \delta\right)$	$P\left(M^4, N / H, \pi, \delta\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01506		—	$\hat{E}$	$\hat{E}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01507		—	$M^4 \times G / H$	$M^4 \times G / H$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01508		—	$\hat{E} \backslash \left(M^4, F=G / H, N / H, P\right)$	$\hat{E}\left(M^4, F=G / H, N / H, P\right)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01509		—	N / H	N / H
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01510		—	$\omega=\omega^j_{\left\{\alpha\right\} L_j} \otimes \vartheta^{\alpha}$	$\omega=\omega^j_{\alpha} L_j \otimes \vartheta^{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01511		—	$\stackrel{\circ}{\vartheta}$	$\stackrel{\circ}{\vartheta}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01512		—	$T_{\{e\}\ }^{\{*\}}(G)$	$T_e^*(G)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01513		—	ξ^j	ξ^j
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01514		—	φ_j	φ_j
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01515		—	$d\,\xi^j$	$d\xi^j$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01516		—	$I_{\{j\}}$	I_j
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01517		—	$V{=}{V^{\{j\}}\ I_{\{j\}}}$, $\stackrel{\circ}{\relax}\circ{\circ}\{\vartheta\}$	$V = V^j I_j, \overset{\circ}{\vartheta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01518		—	$\stackrel{\circ}{\relax}\circ{\circ}\{\omega\}$	$\overset{\circ}{\omega}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01519		—	$\stackrel{\circ}{\circ}\rel{\circ}{\Theta}$	$\overset{\circ}{\Theta}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01520		—	$\stackrel{\circ}{\circ}\rel{\circ}{\Omega}$	$\overset{\circ}{\Omega}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01521		—	$b=1 / 2$	$b = 1/2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01522		—	$\stackrel{\circ}{\circ}\rel{\circ}{R}=1 / 4$	$\overset{\circ}{R}= 1/4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01523		—	$b=0, 1$	$b = 0, 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01524		—	$\operatorname{Tr}(\stackrel{\circ}{\circ}\rel{\circ}{\vartheta} \ \otimes \ \stackrel{\circ}{\circ}\rel{\circ}{\vartheta})$	$\operatorname{Tr}(\overset{\circ}{\vartheta} \otimes \overset{\circ}{\vartheta})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01525		—	$\stackrel{\circ}{\circ}\rel{\circ}{g}_{i \ j}$	$\overset{\circ}{g}_{ij}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01526		—	$\operatorname{Tr}(\omega \otimes \stackrel{\circ}{\vartheta})$	$\operatorname{Tr}(\omega \otimes \stackrel{\circ}{\vartheta})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01527		—	$\operatorname{Tr}(\omega \otimes \omega)$	$\operatorname{Tr}(\omega \otimes \omega)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01528		—	$d \, \hat{s}^2$	$d\hat{s}^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01529		—	$\mathrm{G} \, \times \, \mathrm{G} / \mathrm{G}_{\mathrm{D}}$	$\mathrm{G} \times \mathrm{G} / \mathrm{G}_{\mathrm{D}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01530		—	G_{D}	G_{D}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01531		—	$\hat{\mathrm{L}}(\hat{\mathrm{E}})$	$\hat{L}(\hat{E})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01532		—	$\mathrm{L}\left(\mathrm{M}^4\right), \stackrel{\circ}{\omega}$	$L\left(M^4\right), \stackrel{\circ}{\omega}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01533		—	$\hat{\alpha}$	$\hat{\alpha}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01534		—	$4+\mathrm{K}$	$4 + \mathbb{K}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01535		—	$\hat{\Omega}$	$\hat{\Omega}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01536		—	$(4+\mathrm{K})$	$(4 + \mathbb{K})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01537		—	$\hat{\mathrm{d}}$	$\hat{\mathrm{d}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01538		—	$M^{\{4\}}$	M^4
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01539		—	$\Omega^{\prime}$	Ω'
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01540		—	$\alpha_g:=g^2/\hbar c$	$\alpha_g := g^2/\hbar c$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01541		—	$\hat{L}_{\mathrm{W}}^G$	$\hat{L}_{\mathrm{W}}^G$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01542		—	$\stackrel{\circ}{\operatorname{R}}$	$\stackrel{\circ}{R}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01543		—	$\sqrt{\operatorname{det} \stackrel{\circ}{\operatorname{g}}}$	$\sqrt{\det \stackrel{\circ}{g}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01544		—	$M^{4+K} \approx M^4 \times B$	$M^{4+K} \approx M^4 \times B$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01545		—	$R_{\{4\}} \gg R_{\{K\}}$	$R_4 \gg R_K$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01546		—	$\mathscr{D}(\mathcal{M}^{4+K})$	$\mathscr{D}(M^{4+K})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01547		—	$R_{\{\circ\}}$	$R_{\circ}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01548		—	$Y_{\{n\}}(B)$	$Y_n(B)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01549		—	$B=G \ / \ H$	$B = G/H$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01550		—	$M^{\{4\}} \ \times \ S^{\{1\}}$	$M^4 \times S^1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01551		—	$R_{\{\mathrm{o}\}}$	$R_{\circ}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01552		—	$S^{\{1\}}$	S^1
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01553		—	$S \ U(2) \ \times \ U(1)$	$SU(2) \times U(1)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01554		—	$\mathrm{C} \ P^2 = S \ U(3) \ / \ U(2)$	$\mathbb{C}P^2 = SU(3)/U(2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01555		—	$S^{\{n\}}$	S^n
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01556		—	$M^{\{4\}} \ \times \ B$	$M^4 \times B$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01557		—	$\mathrm{n}=11$	$\mathfrak{n} = 11$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01558		—	$S^{\{7\}} = S \ 0(8) \ / \ S \ 0(7)$	$S^{\prime} = SO(8)/SO(7)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01559		—	$\mathrm{K}=4+2+1=7$	$\mathbb{K} = 4 + 2 + 1 = 7$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01560		—	$\mathrm{n}=4+7=11$	$\mathfrak{n} = 4 + 7 = 11$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01561		—	$(\mathrm{scr}{N}=1)$	$(\mathcal{N} = 1)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01562		—	$\overline{\mathrm{s}} \leq 3 / 2$	$\bar{s} \leq 3/2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01563		—	$\overline{\mathrm{s}}=2$	$\bar{s} = 2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01564		—	$A_{\text{ IJK }}$	$A_{\mathfrak{I}\mathfrak{J}\mathfrak{K}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01565		—	$M^{11}=(\mathrm{AdS})^{\wedge 4} \times S^{\wedge 7}$	$M^{11} = (\mathrm{AdS})^4 \times S^7$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01566		—	$\tilde{\mathrm{s}}^{\wedge 7}$	$\bar{S}^7$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01567		—	$M^{\{5\}}=\mathrm{R}^{\wedge 4} \times S^{\wedge 1}$	$M^5 = \mathbb{R}^4 \times S^1$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01568		—	$\mathrm{E}_{2,4}$	$E_{2,4}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01569		—	$\hat{\omega}$	$\hat{\omega}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01570		—	$\hat{\psi}$	$\hat{\psi}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01571		—	$2^{\{[2+\mathrm{K}]/2\}}$	$2^{[2+K/2]}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01572		—	$\chi_{\rho F}^{\wedge}(\xi)$	$\chi_{\rho F}^{\pm}(\xi)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01573		—	$\stackrel{\circ}{\gamma}^{\mathrm{K}+1}$	$\overset{\circ}{\gamma}^{K+1}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01574		—	$\tilde{\omega}^{\mathrm{g}}$	$\tilde{\omega}^g$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01582		—	$\widetilde{S^0}(1,3+K)$	$SO(1,3+K)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01583		—	$\hat{\gamma}^{n+1}$	$\hat{\gamma}^{n+1}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01584		—	$\overparen{S^0}(1,3) \otimes \widetilde{S^0}(K)$	<i>(not rendered)</i>
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01585		—	$n=2 \text{ } p+1$	$n = 2p + 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01586		—	$\hat{\gamma}^A \rightarrow \hat{\gamma}^A$	$\hat{\gamma}^A \rightarrow -\hat{\gamma}^A$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01587		—	$\hat{\sigma}^{A \text{ } B}$	$\hat{\sigma}^{AB}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01588		—	$\hat{\gamma}^{n+1} \rightarrow \hat{\gamma}^{n+1}$	$\hat{\gamma}^{n+1} \rightarrow -\hat{\gamma}^{n+1}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01589		—	$n=4 \ p$	$n = 4p$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01590		—	$n=4 \ p+2$	$n = 4p + 2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01591		—	$i \ \not D^{\{(\text{int })\}}$	$i \ \not D^{(\text{int })}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01592		—	$\mathrm{D}^{\wedge\{*\}}=\{ \ }^{\wedge\{*\}} \ \mathrm{D}$	$\mathbf{D}^* = *\mathbf{D}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01593		—	$\stackrel{\circ}{\rel{\circ}}{\Theta}=0$	$\overset{\circ}{\Theta}=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01594		—	$n>2$	$n > 2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01595		—	$n=11$	$n = 11$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01596		—	$G=\mathrm{SO}(32)$	$G = \mathrm{SO}(32)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01597		—	$\mathrm{E}_{\!8} \times \mathrm{E}_{\!8}$	$\mathrm{E}_8 \times \mathrm{E}_8$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01598		—	$n=10+\operatorname{dim} G=506$	$n = 10 + \dim G = 506$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01599		—	$\{ \}^{\!10} J_{\!3} \hbar$	$^{10} J_3 \hbar$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01600		—	$\sqrt{J(J+1)} \hbar$	$\sqrt{J(J+1)} \hbar$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01601		—	$Q^{\{i\}}$	$Q^{(i)}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01602		—	$P^{\{i\}}$	$P^{(i)}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01603		—	$J_{\{3\}} \hat{=} J$	$J_3 \hat{=} J$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01604		—	$r^2 e^v$	$r^2 e^v$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01605		—	$\bar{Q}^{\{\circ\}}$	$\bar{Q}^{(\circ)}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01606		—	$J_{\{3\}} \hbar$	$J_3 \hbar$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01607		—	$\mu_m = \frac{Q_{\{e\}}}{M} J$	$\mu_m = \frac{Q_e}{M} J$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01608		—	$\mathrm{M}_{\{\mathrm{ir}\}}$	$\mathbf{M}_{\mathrm{ir}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01609		—	$\mathrm{M}_{\{\mathrm{ir}\}}, \bar{Q}^{\{\circ\}}$	$\mathbf{M}_{\mathrm{ir}}, \bar{Q}^{(\circ)}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01610		—	$J\,\hat{=}\,J_3=0$	$J\hat{=}\,J_3=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01611		—	λ_3	λ_3
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01612		—	σ_3	σ_3
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01613		—	Δ	Δ
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01614		—	$r=r_{\mathrm{o}}$	$r=r_{\mathrm{o}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01615		—	$r=0$	$r=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01616		—	$Q=-\bar{Q}\,\cos\,\delta$	$Q=-Q\cos\delta$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01617		—	$P=-\bar{Q} \sin \delta$	$P = -Q \sin \delta$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01618		—	S_{A}	S_A
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01619		—	$\bar{Q}$	Q
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01620		—	$1 / \kappa$	$1/\kappa$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01621		—	$\mathrm{d}=10$	$d = 10$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01622		—	$1 / \alpha_e$	$1/\alpha_e$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01623		—	$m_{\pi}=2\ m_e / \alpha_e$	$m_{\pi} = 2m_e/\alpha_e$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01624		—	$S\,U(3)^c$	$SU(3)^c$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01625		—	$f_{\mathbf{i}\,\mathbf{j}}$	f_{ij}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01626		—	ϑ^f	ϑ^f
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01627		—	ϑ^g	ϑ^g
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01628		—	$L_{\mathbf{f}\,\mathbf{g}}$	L_{fg}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01629		—	ϑ^{f}	ϑ^f
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01630		—	$\mathbf{b-4}$	$b-4$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01631		—	$\sum^f$	$\sum^f$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01632		—	$\sum^g$	$\sum^g$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01633		—	G_{N}	G_N
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01634		—	G_{S}	G_S
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01635		—	$=10^{-13} \mathrm{~cm}$	$= 10^{-13} \text{ cm}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01636		—	f^0	$\mathbf{f}^0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01637		—	μ^g, μ^f	μ^g, μ^f
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01638		—	$\mathrm{u}=1/2$	$u = 1/2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01639		—	$\mu^g=\Lambda=0$	$\mu^g = \Lambda = 0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01640		—	$m_{\rm f}=1-2\,\mathrm{GeV}/c^2$	$m_f = 1 - 2\mathrm{GeV}/c^2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01641		—	E_j^β	E_j^β
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01642		—	$\Gamma_i^{\alpha\beta}$	$\Gamma_i^{\alpha\beta}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01643		—	$\left(\frac{1}{2},\frac{1}{2}\right)$	$\left(\frac{1}{2},\frac{1}{2}\right)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01644		—	$(1,0)\oplus$	$(1,0)\oplus$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01645		—	$(0,1)$	$(0,1)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01646		—	$\operatorname{spin} \bar{s}_a=1$	$\operatorname{spin} \bar{s}_a = 1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01647		—	$\bar{s}=2$	$\bar{s} = 2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01648		—	$\Lambda_{\text{eff}} \neq 0$	$\Lambda_{\text{eff}} \neq 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01649		—	$\ell=\ell^* / \sqrt{\kappa}$	$\ell = \ell^* / \sqrt{\kappa}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01650		—	$1 / \sqrt{\kappa}$	$1/\sqrt{\kappa}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01651		—	$10^{13} \mathrm{\sim cm}$	10^{13} cm
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01652		—	ℓ^2	ℓ^2
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01653		—	ℓ^{*2}	ℓ^{*2}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01654		—	$\kappa=0,38 \times 10^{-38}$	$\kappa = 0,38 \times 10^{-38}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01655		—	10^{20}	10^{20}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01656		—	M_{U}	M_{U}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01657		—	M_{P}	M_{P}
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01658		—	ℓ_{H}	ℓ_{H}
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01659		—	$SU(f)$	$SU(f)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01660		—	$SU(4)$	$SU(4)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01661		—	$SU(5)$	$SU(5)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01662		—	$SO(3) \approx SU(2)$	$SO(3) \approx SU(2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01663		—	$SU(4) \approx SO(6) \supset$	$SU(4) \approx SO(6) \supset$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01664		—	$SO(4) \approx SU(2) \times SU(2)$	$SO(4) \approx SU(2) \times SU(2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01665		—	$SU(6)$	$SU(6)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01666		—	$J=1 / 2$	$J = 1/2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01667		—	$J=3 / 2$	$J = 3/2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01668		—	$S L(6, \mathbb{C}) \supset S L(2, \mathbb{C}) \times S U(3)$	$SL(6, \mathbb{C}) \supset SL(2, \mathbb{C}) \times SU(3)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01669		—	$\mathrm{SU}(3)$	$SU(3)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01670		—	$S L(6, \mathbb{C})$	$SL(6, \mathbb{C})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01671		—	$\mathrm{P}(\mathrm{M}, \overline{\mathrm{G}}, \pi, \delta)$	$P(M, \overline{G}, \pi, \delta)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01672		—	$[S U(4) \otimes]^4$	$[SU(4) \otimes]^4$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01673		—	f=c=4	$f = c = 4$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01674		—	$\overline{\mathrm{L}}(\mathrm{M})$	$\overline{\mathbf{L}}(\mathbf{M})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01675		—	A_{2}	$\mathbf{A}_2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01676		—	$S\,L(2\,\mathbb{N},\,\mathbb{C})\supset S\,L(2,\,\mathbb{C})\times U(N)$	$SL(2N,\mathbb{C})\supset SL(2,\mathbb{C})\times U(N)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01677		—	λ_i	λ_i
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01678		—	$\mathrm{N}^{\wedge 2}-1$	$\mathbf{N}^2-1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories_ Eckehard_W._Mielke_ __Z-Library__F01679		—	$\operatorname{SU}(N)$	$\mathrm{SU}(N)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01680		—	A_{i}	A_i
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01681		—	$\bar{\gamma}$	$\bar{\gamma}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01682		—	$L_{\mathrm{W}}(\mathrm{f})$	$L_{\mathrm{W}}(\mathfrak{f})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01683		—	$\mathfrak{f}=\mathfrak{c}=\emptyset$	$f = c = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01684		—	$\mathfrak{f}=3$	$f = 3$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01685		—	$\mathfrak{c}=\emptyset$	$c = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01686		—	$\bar{L}_{\mathrm{L}}(\mathrm{fg})$	$L_{\mathfrak{fg}}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01687		—	$G=S\ U(2)$	$G = SU(2)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01688		—	$S\ U(4)^{\{f\}} \otimes S\ U(4)^{\{c\}}$	$SU(4)^f \otimes SU(4)^c$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01689		—	$S\ L(8,\ \mathbb{C})^{\{f\}} \otimes S\ U(8,\ \mathbb{C})^{\{c\}}$	$SL(8,\mathbb{C})^f \otimes SU(8,\mathbb{C})^c$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01690		—	$M=\mathbb{R}\ \times\ S^{\{1\}}\ \times\ S^{\{2\}}$	$M = \mathbb{R} \times S^1 \times S^2$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01691		—	$G\ L(8,\ \mathbb{C})^{\{f\}} \otimes$	$GL(8,\mathbb{C})^f \otimes$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01692		—	$G\ L(8,\ \mathbb{C})^{\{c\}}$	$GL(8,\mathbb{C})^c$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01693		—	$U(5)$	$U(5)$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01694		—	$S\ 0(1,9)$	$SO(1,9)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01695		—	$S\ 0(10)$	$SO(10)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01696		—	$S\ 0(1,13)$	$SO(1,13)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01697		—	$S\ 0(4,\ \mathbb{C})$	$SO(4,\mathbb{C})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01698		—	$E(8)$	$E(8)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01699		—	$E(10)$	$E(10)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01700		—	$(5,\ \mathbb{C})$	$(5,\mathbb{C})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01701		—	$\operatorname{SL}(2\,\mathrm{n},\,\mathbb{C})\,\otimes\,\operatorname{SL}(2,\,\mathbb{C})$	$\mathrm{SL}(2\mathrm{n},\mathbb{C})\otimes\mathrm{SL}(2,\mathbb{C})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01702		—	$1000\,\mathrm{~m}_{\mathrm{~}\mathrm{W}}$	1000 m _W
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01703		—	$2\,\mathrm{n},\,\mathbb{C}$	$2\mathrm{n},\mathbb{C}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01704		—	$2^{\{+\}}$	2^+
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01705		—	$\mathrm{R}-\mathrm{R}_{\mathrm{~}\mathrm{o}}$	$\mathrm{R}-\mathrm{R}_{\mathrm{o}}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01706		—	$(1966,1967)$	$(1966,1967)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01707		—	$\kappa>0$	$\kappa>0$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01708		—	$2\,\mu\,/\,\mathrm{r}$	$2\mu/\mathrm{r}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01709		—	ℓ_{m}	ℓ_m
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01710		—	$\mathrm{F}=\mathrm{F}_{\mathrm{nm}}^{\mathrm{iota}}\left(\rho^*\right)$	$\mathbf{F}=\mathbf{F}_{\mathrm{nm}}^{\iota}(\rho^*)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01711		—	$\mathrm{iota}=0$	$\iota=0$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01712		—	$\rho^*=\frac{\pi}{2}$	$\rho^*=\frac{\pi}{2}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01713		—	$\mathrm{beta}^2<2$	$\beta^2<2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke___Z-Library__F01714		—	$\mathrm{V}(\mathrm{r})=2\,\mu\,/\,\mathrm{r}+\mathrm{cr}$	$\mathbf{V}(\mathbf{r})=2\mu/\mathrm{r}+\mathbf{cr}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01715		—	$\overline{\mathrm{Q}}$	Q
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01716		—	Q_{e}	Q _e
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01717		—	$\mathrm{M}_{\text{\text{ir}}}$	M _{ir}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01718		—	$\mathrm{M}^* := \hbar \sqrt{8 \pi} / \ell^* \mathrm{c}$	$M^* := \hbar \sqrt{8 \pi} / \ell^* c$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01719		—	$\mathrm{M}_{\text{\text{ir}}} = \mathrm{M}^* = 1 \mathrm{GeV} / \mathrm{c}^2$	M _{ir} = M* = 1GeV/c ²
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01720		—	M, J	M, J
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01721		—	I	I
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01722		—	$\mathrm{J}=0$	$J = 0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01723		—	$\mathrm{b}=\frac{1}{5}$	$\mathbf{b} = \frac{1}{5}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01724		—	$\Lambda(1116)$	$\Lambda(1116)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01725		—	$k_{\mathrm{B}}=8.6 \times 10^{-5} \mathrm{eV} / \mathrm{K}$	$k_B = 8.6 \times 10^{-5} \mathrm{eV/K}$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01726		—	$\mathrm{S\ U}(8)$	$SU(8)$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01727		—	$(6, \mathbb{C})$	$(6, \mathbb{C})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01728		—	equiv	$\equiv$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01729		—	$\backslash\mathrm{approx}$	$\approx$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01730		—	$\backslash\mathrm{G}_{\backslash\mathrm{circ}}$	$\mathrm{G}_{\circ}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01731		—	$[\backslash\mathrm{A}, \backslash\mathrm{B}]:=\mathrm{AB}-\mathrm{BA}$	$[A, B] := AB - BA$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01732		—	$\backslash\{\backslash\mathrm{A}, \backslash\mathrm{B}\backslash\}:=\mathrm{AB}+\mathrm{BA}$	$\{A, B\} := AB + BA$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01733		—	$\backslash\mathrm{A}^{\backslash+}=\mathrm{A}^{\backslash*} \backslash\mathrm{T}$	$\mathrm{A}^{+} = \mathrm{A}^{*\mathrm{T}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01734		—	$\backslash\mathrm{pi}^{\backslash-1}(\backslash\mathrm{N})$	$\pi^{-1}(\mathrm{N})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01735		—	$\backslash\mathrm{pi}: \backslash\mathrm{M} \rightarrow \backslash\mathrm{N}$	$\pi : \mathrm{M} \rightarrow \mathrm{N}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01736		—	$\mathrm{T}_{\{(\mu \ \nu)\}}:=\frac{1}{2}\left(\mathrm{T}_{\sim\{\mu \ \nu\}}+\mathrm{T}_{\{\nu \ \mu\}}\right)$	$T_{(\mu\nu)} := \frac{1}{2} (\ T_{\mu\nu} + T_{\nu\mu})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01737		—	$\mathrm{T}_{\{[\mu \ \nu]\}}:=\frac{1}{2}\left(\mathrm{T}_{\sim\{\mu \ \nu\}}-\mathrm{T}_{\{\nu \ \mu\}}\right)$	$T_{[\mu\nu]} := \frac{1}{2} (\ T_{\mu\nu} - T_{\nu\mu})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01738		—	2×2	2×2
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01739		—	$\mathrm{C}\{(\boldsymbol{0},2)=\mathbb{H}$	$\mathbf{C}(0,2) = \mathbb{H}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01740		—	4×4	4×4
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01741		—	$4<2 \ \mathrm{p}+1<2 \ \mathrm{p}+2 \ \leq n=2 \ l+2$	$4 < 2\mathbf{p} + 1 < 2\mathbf{p} + 2 \leq n = 2l + 2$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01742		—	$\emptyset$	$\emptyset$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01743		—	T_{1}	T_1
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01744		—	$\mathrm{m}_{1}, \mathrm{~m}_{2} \in \mathrm{X}$	$m_1, m_2 \in X$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01745		—	$\mathrm{U}_{\mathrm{i}} \ni \mathrm{m}_{\mathrm{i}}$	$U_i \ni m_i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01746		—	T_{2}	T_2
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01747		—	$\mathrm{U}_{1} \bigcap \mathrm{U}_{2}=\varnothing$	$U_1 \bigcap U_2 = \emptyset$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01748		—	$\sigma: \mathrm{M} \rightarrow \mathrm{N}$	$\sigma: M \rightarrow N$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01749		—	$\mathrm{M} \approx$	$M \approx$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01750		—	$\mathrm{N} \approx \mathbb{R}^{\mathrm{n}}$	$\mathbb{N} \approx \mathbb{R}^{\mathrm{n}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01751		—	$\{ \}^{-1}$	-1
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01752		—	$\sigma_{*}: \mathrm{T}_{\mathrm{m}}(\mathrm{M}) \rightarrow \mathrm{T}_{\sigma(\mathrm{m})}(\mathrm{N})$	$\sigma_{*}: \mathrm{T}_{\mathrm{m}}(\mathrm{M}) \rightarrow \mathrm{T}_{\sigma(\mathrm{m})}(\mathrm{N})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01753		—	$\sigma(\mathrm{s}(\mathrm{t}))$	$\sigma(\mathrm{s}(\mathrm{t}))$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01754		—	e^{f}	e^f
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01755		—	σ_{*}	σ_{*}
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01756		—	$\sigma_{*}^{\mathrm{T}}: T_{\sigma(\mathrm{m})}^{*}(\mathrm{N}) \rightarrow \mathrm{T}_{\mathrm{M}}^{*}(\mathrm{M})$	$\sigma_{*}^{\mathrm{T}}: T_{\sigma(\mathrm{m})}^{*}(\mathrm{N}) \rightarrow \mathrm{T}_{\mathrm{M}}^{*}(\mathrm{M})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01757		—	$\{ \}^{\ast} \ \alpha$	$\ast \alpha$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01758		—	$(p, \ 0)$	$(p, 0)$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01759		—	$\{ \}^{\{1\}} \ \mathrm{d}f\!=\!\vartheta^{\alpha}\partial_{\alpha} \ \mathrm{f}$	$\mathrm{d}f = \vartheta^{\alpha}\partial_{\alpha}f$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01760		—	$\sigma_{\ast}^{\ast} \ \alpha^{\{(\emptyset)\}}\!:=\!\alpha^{\{(\emptyset)\}} \ \circ \sigma$	$\sigma_{\ast}\alpha^{(0)} := \alpha^{(0)} \circ \sigma$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01761		—	$\varepsilon_{\alpha_1\{1\} \ \ldots \ \alpha_n\}$	$\varepsilon_{\alpha_1 \ldots \alpha_n}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01762		—	$\varepsilon_{\hat{1} \ldots \tilde{n}}\!=\!+1$	$\varepsilon_{\hat{1} \ldots \tilde{n}} = +1$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01763		—	$g_{\{\mathrm{i}j\}}$	g_{ij}
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\!-\!0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01764		—	$\mathrm{d}f=\left(\partial\mathrm{f}/\partial x^i\right)\mathrm{d}x^i$	$\mathrm{d}f=\left(\partial\mathrm{f}/\partial x^i\right)dx^i$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01765		—	$\delta\alpha=0$	$\delta\alpha=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01766		—	$\mathrm{B}(\mathrm{M}):=\mathrm{d}\mathrm{D}(\mathrm{M})$	$\mathrm{B}(\mathrm{M}):=\mathrm{d}\mathrm{D}(\mathrm{M})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01767		—	$\mathrm{B}^{\{0\}}(\mathrm{M})=0$	$\mathrm{B}^0(\mathrm{M})=0$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01768		—	$\mathrm{H}^{\{p\}}(\mathrm{M})$	$\mathrm{H}^p(\mathrm{M})$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01769		—	$\mathrm{t}=-1$	$\mathrm{t}=-1$
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01770		—	η_{i}	η_{i}
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5\text{--}0.9$ ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01771		—	$\mathrm{c}: \Delta_{\mathrm{p}} \rightarrow \mathrm{M}$	$\mathbf{c} : \Delta_{\mathbf{p}} \rightarrow \mathbf{M}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01772		—	$\mathrm{C}_{\mathrm{p}}(\mathrm{M}, \mathbb{R})$	$\mathbf{C}_{\mathbf{p}}(\mathbf{M}, \mathbb{R})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01773		—	$\partial \mathrm{c}$	$\partial \mathbf{c}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01774		—	$\mathrm{Z}_{\mathrm{p}}(\mathrm{M}, \mathbb{R}) = (\operatorname{Ker} \partial)_{\mathrm{p}}, \mathrm{B}_{\mathrm{p}}(\mathrm{M}, \mathbb{R}) := (\operatorname{Im} \partial)_{\mathrm{p}}$	$\mathbf{Z}_{\mathbf{p}}(\mathbf{M}, \mathbb{R}) = (\operatorname{Ker} \partial)_{\mathbf{p}}, \mathbf{B}_{\mathbf{p}}(\mathbf{M}, \mathbb{R}) := (\operatorname{Im} \partial)_{\mathbf{p}}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01775		—	$\mathrm{N}_{\mathrm{p}}(\mathrm{M}) \subset \mathrm{M}$	$\mathbf{N}_{(\mathbf{p})} \subset \mathbf{M}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01776		—	$\mathrm{N}_{\mathrm{p}}(\mathrm{M})$	$\mathbf{N}_{(\mathbf{p})}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01777		—	$\mathrm{c}_{\mathrm{p}} \in \mathrm{C}_{\mathrm{p}}(\mathrm{M}, \mathbb{R})$	$\mathbf{c}_{\mathbf{p}} \in \mathbf{C}_{\mathbf{p}}(\mathbf{M}, \mathbb{R})$
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01778		—	$\mathrm{H}^{\mathrm{P}}(\mathrm{M}) \times \mathrm{H}_{\mathrm{p}}(\mathrm{M}, \mathbb{R}) \rightarrow \mathbb{R}$ $\rightarrow \mathbb{R}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01779		—	$\mathrm{g} \in \mathrm{G}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01780		—	$\mathrm{G} \times \mathrm{G} \rightarrow \mathrm{G}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01781		—	$\left(\mathrm{g}_1, \mathrm{~g}_2\right) \rightarrow \mathrm{g}_1 \mathrm{~g}_2^{-1}$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01782		—	$\mathrm{G}=\mathrm{GL}(\mathrm{N}, \mathbb{C})$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01783		—	$\mathrm{T}_{\mathrm{e}}(\mathrm{G})$	
Geometrodynamics_ of_Gauge_Fields_ On_the_Geometry_ of_Yang-Mills_ and_Gravitational_ Gauge_Theories__ Eckehard_W._Mielke_ __Z-Library__F01784		—	$\mathrm{e} \in \mathrm{G}$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01785		—	$\xi=\left\{\xi^{\mathrm{j}},\ \mathrm{j}=1,\ \ldots,\ \mathrm{~K}\right\}$	$\xi=\left\{\xi^j,j=1,\ldots,\ K\right\}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01786		—	$\mathrm{c}_{\mathrm{ij}}^{\mathrm{~k}}$	c_{ij}^k
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01787		—	$\mathrm{A}=\mathrm{A}^{\mathrm{i}}\ \mathrm{I}_{\mathrm{i}}$	$A=A^iI_i$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01788		—	$\mathrm{B}=\mathrm{B}^{\mathrm{j}}\ \mathrm{I}_{\mathrm{j}}$	$B=B^jI_j$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01789		—	$\stackrel{\circ}{g}_{ls}$	$\overset{\circ}{g}_{ls}$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01790		—	$\mathfrak{gl}(\mathrm{N},\ \mathbb{C})$	$\mathfrak{gl}(\mathbf{N},\mathbb{C})$
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01791		—	$\mathrm{G}=\exp^{\mathrm{~Trg}}$	$G=\exp^{\mathrm{Trg}}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				



Identifier	Page	Conf.	LaTeX source	Rendered
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01792		—	ξ^{j}	ξ^j
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__F01793		—	λ_{j}	λ_j
Geometrodynamics_of_Gauge_Fields_On_the_Geometry_of_Yang-Mills_and_Gravitational_Gauge_Theories__Eckehard_W._Mielke__Z-Library__FOX_499d85f9b6		—	$O(s, n - s$	$O(s, n - s$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5-0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				

Tables




Identifier	Page	Content (LaTeX source if any)	Scan image
Geometrodynamics_of_Gauge_Fields_in_the_Geometry_of_Yang-Mills_Theories_1075	075	(no LaTeX source: 1111 × 569 px region — see tables.html)	—
Geometrodynamics_of_Gauge_Fields_in_the_Geometry_of_Yang-Mills_Theories_1102	102	(no LaTeX source: 902 × 202 px region — see tables.html)	—
Geometrodynamics_of_Gauge_Fields_in_the_Geometry_of_Yang-Mills_Theories_1210	210	(no LaTeX source: 1159 × 360 px region — see tables.html)	—
Geometrodynamics_of_Gauge_Fields_in_the_Geometry_of_Yang-Mills_Theories_1275	275	(no LaTeX source: 441 × 116 px region — see tables.html)	—
Geometrodynamics_of_Gauge_Fields_in_the_Geometry_of_Yang-Mills_Theories_1364	364	(no LaTeX source: 1058 × 670 px region — see tables.html)	—





Unrecovered image regions — TikZ / table / failed-math candidates




Regions MathPix left as images (no LaTeX). Reconstruct with pdfdrill vision (LLM classify + transcribe to TikZ/tabular/math); verify an LLM result against the real ink with inkdrill.





Identifier	Page	Image	Source
	258		tex.zip (local)
	256		tex.zip (local)

Identifier	Page	Image	Source
Geometrodynamics_of_Gauge_Fields_on_	359		tex.zip (local)
Geometrodynamics_of_Gauge_Fields_on_	025		tex.zip (local)

Identifier	Page	Image	Source
	031	 Diagram illustrating a 2D plane E^2 with a grid and a sphere $M^2 \approx S^2$. The plane has axes x_1 and x_2, and a transformation $x_2 \circ x_1^{-1}$. The sphere has regions U_1 and U_2, and a point s.	tex.zip (local)
	032	 Diagram illustrating a sphere U with a dashed curve $O = \partial U$. A tangent plane $T_m(M)$ is shown at point m, with axes $\frac{\partial}{\partial \theta}$ and $\frac{\partial}{\partial \phi}$. A vector e is shown at the top, with a small triangle Δe.	tex.zip (local)
	036	 Diagram illustrating a torus $U \subset M \approx S^1$. A point m is marked on the torus, and a vertical line segment is labeled π.	tex.zip (local)

Identifier	Page	Image	Source
Geometrodynamics_of_Gauge_Fields_on_	040		tex.zip (local)
Geometrodynamics_of_Gauge_Fields_on_	057		tex.zip (local)
Geometrodynamics_of_Gauge_Fields_on_	073		tex.zip (local)
Geometrodynamics_of_Gauge_Fields_on_	084		tex.zip (local)

Identifier	Page	Image	Source
Geometrodynamics_of_Gauge_Fields_On_	131		tex.zip (local)
Geometrodynamics_of_Gauge_Fields_On_	200		tex.zip (local)
Geometrodynamics_of_Gauge_Fields_On_	226		tex.zip (local)

Identifier	Page	Image	Source
	227		tex.zip (local)
	288		tex.zip (local)
	290		tex.zip (local)
	347		tex.zip (local)