

mielke-geometroynamics: equation evidence

62 rows show a *REFINEMENT* below, not the OCR reading: 62 verified against the census. Each is marked [refined] beside its identifier. The model is unchanged — the original reading is still what ‘latex’ holds; this report chose the refinement for them. Rows: mielke-geometroynamics_EQ0006, mielke-geometroynamics_EQ0119, mielke-geometroynamics_EQ0120, mielke-geometroynamics_EQ0292, mielke-geometroynamics_EQ0293, mielke-geometroynamics_EQ0294 ...

Display equations

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometroynamics_EQ0966 (14.2.7) [conf 0.024]	313	■ 0.024	$\begin{aligned} d\hat{s}^2 &= \frac{1}{4+K} \operatorname{Tr} \left(\hat{\vartheta} \otimes_s \hat{\vartheta}^T \right) = \hat{g}_{AB} \hat{\vartheta}^A \otimes_s \hat{\vartheta}^B; \quad A, B, = 0, \dots, 3+K \\ \hat{\vartheta} \otimes_s \hat{\vartheta}^T &= \hat{g}_{AB} \hat{\vartheta}^A \otimes_s \hat{\vartheta}^B; \quad A, B, = 0, \dots, 3+K \\ \hat{\vartheta} \otimes_s \hat{\vartheta}^T &= \hat{g}_{AB} \hat{\vartheta}^A \otimes_s \hat{\vartheta}^B; \quad A, B, = 0, \dots, 3+K \end{aligned}$	$d\hat{s}^2 = \frac{1}{4+K} \operatorname{Tr} \left(\hat{\vartheta} \otimes_s \hat{\vartheta}^T \right) = \hat{g}_{AB} \hat{\vartheta}^A \otimes_s \hat{\vartheta}^B; \quad A, B, = 0, \dots, 3+K$ $= \frac{1}{4+K} \operatorname{Tr} \left[\begin{array}{c c} \vartheta \otimes \vartheta + \hat{\ell}^2 \omega \otimes \omega & \hat{\ell} \omega \otimes \overset{\circ}{\vartheta} \\ \hline \hat{\ell} \vartheta \otimes \omega & \ddots \\ \hline \vartheta \otimes & \end{array} \right].$	$d\hat{s}^2 = \frac{1}{4+K} \operatorname{Tr}(\hat{\vartheta} \otimes_s \hat{\vartheta}^T) = \hat{g}_{AB} \hat{\vartheta}^A \otimes_s \hat{\vartheta}^B; \quad A, B, = 0, \dots, 3+K$ $= \frac{1}{4+K} \operatorname{Tr} \left[\frac{\vartheta \otimes \vartheta + \hat{\ell}^2 \omega \otimes \omega}{\hat{\ell} \vartheta \otimes \omega} \middle \frac{\hat{\ell} \omega \otimes \overset{\circ}{\vartheta}}{\vartheta \otimes \overset{\circ}{\vartheta}} \right]. \quad (1)$
mielke-geometroynamics_EQ0403 (6.4.23) [conf 0.028]	134	■ 0.028	$\Psi_{\parallel}(\vartheta) = \exp \left(- \int_{M_3} \overset{(\pm)}{C}_{\text{TT}} \right) = \exp \left(\frac{-1}{2\ell^2} \int_{M_3} \left[\overset{*}{+} \underline{\vartheta}_B \wedge \overset{*}{T}_T^B \right] \right)$	$\Psi_{\parallel}(\vartheta) = \exp \left(- \int_{M_3} \overset{(\pm)}{C}_{\text{TT}} \right) = \exp \left(\frac{-1}{2\ell^2} \int_{M_3} \left[\overset{*}{+} \underline{\vartheta}_B \wedge \overset{*}{T}_T^B \right] \right)$	$\Psi_{\parallel}(\vartheta) = \exp \left(- \int_{M_3} \overset{(\pm)}{C}_{\text{TT}} \right) = \exp \left(\frac{-1}{2\ell^2} \int_{M_3} \left[\overset{*}{+} \underline{\vartheta}_B \wedge \overset{*}{T}_T^B \right] \right)$
mielke-geometroynamics_EQ0850 (12.3.3) [conf 0.032]	275	■ 0.032	$\begin{aligned} \not{D}^2 &= -\frac{1}{2} \gamma^\alpha \gamma^\beta \left(\{ D_\alpha^{\{\}}, D_\beta^{\{\}} \} + [D_\alpha^{\{\}}, D_\beta^{\{\}}] \right) - 2im \not{D}^{\{\}} \\ &\cong -\square - \frac{1}{8} \sigma^{\alpha\beta} R_{\alpha\beta\mu\nu}^{\{\}} \sigma^{\mu\nu} \\ &= -\frac{1}{4} \gamma_5 \left(\not{D}^{\{\}} \not{\not{D}} \right) + \frac{1}{2} \gamma_5 \sigma^{\alpha\beta} \mathcal{A}_\alpha D_\beta^{\{\}} + m^2 - \frac{1}{2} m \gamma_5 \not{\not{D}} - \frac{1}{16} \not{\not{D}} \not{\not{D}} \\ &\cong -\square - \frac{1}{8} \sigma^{\alpha\beta} R_{\alpha\beta\mu\nu}^{\{\}} \sigma^{\mu\nu} \\ &= -\frac{1}{4} \gamma_5 \left(\not{D}^{\{\}} \not{\not{D}} \right) + \frac{1}{2} \gamma_5 \sigma^{\alpha\beta} \mathcal{A}_\alpha D_\beta^{\{\}} - \frac{1}{16} \mathcal{A}_\alpha \mathcal{A}^\alpha - m^2 \end{aligned}$	$D^2 = -\frac{1}{2} \gamma^\alpha \gamma^\beta \left(\{ D_\alpha^{\{\}}, D_\beta^{\{\}} \} + [D_\alpha^{\{\}}, D_\beta^{\{\}}] \right) - 2im \not{D}^{\{\}}$ $= -\frac{1}{4} \gamma_5 \left(\not{D}^{\{\}} \not{\not{D}} \right) + \frac{1}{2} \gamma_5 \sigma^{\alpha\beta} \mathcal{A}_\alpha D_\beta^{\{\}} + m^2 - \frac{1}{2} m \gamma_5 \not{\not{D}} - \frac{1}{16} \not{\not{D}} \not{\not{D}}$ $\cong -\square - \frac{1}{8} \sigma^{\alpha\beta} R_{\alpha\beta\mu\nu}^{\{\}} \sigma^{\mu\nu}$ $= -\frac{1}{4} \gamma_5 \left(\not{D}^{\{\}} \not{\not{D}} \right) + \frac{1}{2} \gamma_5 \sigma^{\alpha\beta} \mathcal{A}_\alpha D_\beta^{\{\}} - \frac{1}{16} \mathcal{A}_\alpha \mathcal{A}^\alpha - m^2$	$\not{D}^2 = -\frac{1}{2} \gamma^\alpha \gamma^\beta \left(\{ D_\alpha^{\{\}}, D_\beta^{\{\}} \} + [D_\alpha^{\{\}}, D_\beta^{\{\}}] \right) - 2im \not{D}^{\{\}}$ $= -\frac{1}{4} \gamma_5 (\not{D}^{\{\}} \not{\not{D}}) + \frac{1}{2} \gamma_5 \sigma^{\alpha\beta} \mathcal{A}_\alpha D_\beta^{\{\}} + m^2 - \frac{1}{2} m \gamma_5 \not{\not{D}} - \frac{1}{16} \not{\not{D}} \not{\not{D}}$ $\cong -\square - \frac{1}{8} \sigma^{\alpha\beta} R_{\alpha\beta\mu\nu}^{\{\}} \sigma^{\mu\nu}$ $= -\frac{1}{4} \gamma_5 (\not{D}^{\{\}} \not{\not{D}}) + \frac{1}{2} \gamma_5 \sigma^{\alpha\beta} \mathcal{A}_\alpha D_\beta^{\{\}} - \frac{1}{16} \mathcal{A}_\alpha \mathcal{A}^\alpha - m^2$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_134 EQ0402 (6.4.22) [conf 0.042]	134	■ 0.042	$\begin{array}{ll} \text{q: } & \underline{*}^{\vartheta\beta} \Psi_{\parallel}(\vartheta) = \underline{*}'^{\vartheta\beta} \Psi_{\parallel}(\vartheta), \\ \text{p: } & \underline{*}^{\vartheta\beta} \Psi_{\parallel}(\vartheta) = -i\ell^2 \frac{\delta}{\delta \underline{*}_{\alpha}} \Psi_{\parallel}(\vartheta), \end{array}$		$q : \quad \underline{*}^{\vartheta\beta} \Psi_{\parallel}(\vartheta) = \underline{*}^{\vartheta\beta} \Psi_{\parallel}(\vartheta),$ $\underline{p}: \quad \underline{*}^{\vartheta\beta} \Psi_{\parallel}(\vartheta) = -i\ell^2 \frac{\delta}{\delta \underline{*}_{\alpha}} \Psi_{\parallel}(\vartheta),$
mielke-geometrodynamics_101 EQ0293 (4.5.4) [conf 0.052] [refined: census]	101	■ 0.052	$\lfloor L \equiv \left(\lfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + (\lfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\lfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi}.$		$\lfloor L \equiv (\lfloor \vartheta^{\alpha}) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + (\lfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\lfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi}.$
mielke-geometrodynamics_47 EQ0064 (2.8.6) [conf 0.054]	47	■ 0.054	$\begin{aligned} & \stackrel{*}{\pi}(\gamma_1) = -\frac{1}{2\pi i} \text{Tr}(\Omega) \quad (=0) \\ & \stackrel{*}{\pi}(\gamma_2) = -\frac{1}{8\pi^2} \{ \text{Tr}(\Omega \wedge \Omega) - \text{Tr}(\Omega) \wedge \text{Tr}(\Omega) \} \\ & \quad \left(= -\frac{1}{32\pi^2} \epsilon^{\alpha\beta\mu\nu} F_{\alpha\beta}^j F_{\mu\nu j} \sqrt{ g } d^4x \right) \\ & \stackrel{*}{\pi}(\gamma_k) = 0 \quad \text{for } k \geq 3. \end{aligned}$		$\stackrel{*}{\pi}(\gamma_1) = -\frac{1}{2\pi i} \text{Tr} \Omega \quad (=0)$ $\stackrel{*}{\pi}(\gamma_2) = -\frac{1}{8\pi^2} \{ \text{Tr}(\Omega \wedge \Omega) - \text{Tr}(\Omega) \wedge \text{Tr}(\Omega) \}$ $\left(= -\frac{1}{32\pi^2} \epsilon^{\alpha\beta\mu\nu} F_{\alpha\beta}^j F_{\mu\nu j} \sqrt{ g } d^4x \right)$ $\stackrel{*}{\pi}(\gamma_k) = 0 \quad \text{for } k \geq 3.$
mielke-geometrodynamics_133 EQ0400 (6.4.19) [conf 0.060]	133	■ 0.060	$\stackrel{(\pm)}{\mathcal{G}}_{\alpha} := \underline{D}^{\underline{(\pm)}} \underline{\Pi}_{\alpha} = \underline{D}^{\underline{(\pm)}} \underline{A}_{\alpha}^{\underline{(\pm)}} = 0,$		$\stackrel{(\pm)}{\mathcal{G}}_{\alpha} := \underline{D}^{\underline{(\pm)}} \underline{\Pi}_{\alpha} = \underline{D}^{\underline{(\pm)}} \underline{\mathcal{A}}_{\alpha}^{\underline{(\pm)}} = 0,$
mielke-geometrodynamics_229 EQ0718 (10.9.4) [conf 0.066] [refined: census]	229	■ 0.066	$\begin{aligned} & \Sigma_{\alpha} := e_{\alpha} \rfloor L_{\Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L_{\Psi}}{\partial \Psi} - (e_{\alpha} \rfloor \bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial \bar{\Psi}} \\ & \quad - (e_{\alpha} \rfloor D\Psi) \wedge \frac{\partial L_{\Psi}}{\partial D\Psi} - (e_{\alpha} \rfloor D\bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial D\bar{\Psi}}; \end{aligned}$		$\Sigma_{\alpha} := e_{\alpha} \rfloor L_{\Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L_{\Psi}}{\partial \Psi} - (e_{\alpha} \rfloor \bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial \bar{\Psi}} - (e_{\alpha} \rfloor D\Psi) \wedge \frac{\partial L_{\Psi}}{\partial D\Psi} - (e_{\alpha} \rfloor D\bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial D\bar{\Psi}};$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_131 EQ0391 (6.4.6) [conf 0.066] [refined: census]	131	■ 0.066	$\begin{aligned} \Sigma_{\alpha} &:= \frac{\delta L}{\delta \vartheta^{\alpha}} = e_{\alpha} \rfloor L - (e_{\alpha} \rfloor D \Psi) \wedge \frac{\partial L}{\partial D \Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} \\ &\quad - (e_{\alpha} \rfloor T^{\beta}) \wedge \frac{\partial L}{\partial T^{\beta}} + D \frac{\partial L}{\partial T^{\alpha}} - \left(e_{\alpha} \rfloor R_{\beta}^{\gamma} \right) \wedge \frac{\partial L}{\partial R_{\beta}^{\gamma}}. \end{aligned}$	$\Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = e_{\alpha} \rfloor L - (e_{\alpha} \rfloor D \Psi) \wedge \frac{\partial L}{\partial D \Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} - (e_{\alpha} \rfloor T^{\beta}) \wedge \frac{\partial L}{\partial T^{\beta}} + D \frac{\partial L}{\partial T^{\alpha}} - \left(e_{\alpha} \rfloor R_{\beta}^{\gamma} \right) \wedge \frac{\partial L}{\partial R_{\beta}^{\gamma}}.$	$\Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = e_{\alpha} \rfloor L - (e_{\alpha} \rfloor D \Psi) \wedge \frac{\partial L}{\partial D \Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} - (e_{\alpha} \rfloor T^{\beta}) \wedge \frac{\partial L}{\partial T^{\beta}} + D \frac{\partial L}{\partial T^{\alpha}} - (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \frac{\partial L}{\partial R_{\beta}^{\gamma}}.$
mielke-geometrodynamics_168 EQ0528 (7.7.48) [conf 0.078] [refined: census]	168	■ 0.078	$\begin{aligned} 2 \ell^2 d C_{\mathrm{T}} &:= 2 d \left(\vartheta^{\alpha} \wedge * T_{\alpha} \right) \\ &= R^{\{f\alpha\beta} \wedge \eta_{\alpha\beta} - R^{\alpha\beta} \wedge \eta_{\alpha\beta} \\ &\quad - T^{\alpha} * \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta} \rfloor T^{\beta}) - \frac{1}{2} e_{\alpha} \rfloor (T^{\beta} \wedge \vartheta_{\beta}) \right] \\ &= R^{\{f\alpha\beta} \wedge \eta_{\alpha\beta} - R^{\alpha\beta} \wedge \eta_{\alpha\beta} - T^{\alpha} \wedge * \left(- {}^{(1)}T_{\alpha} + 2 {}^{(2)}T_{\alpha} + \frac{1}{2} {}^{(3)}T_{\alpha} \right), \end{aligned}$	$2\ell^2 d C_{\mathrm{T}} := 2d(\vartheta^{\alpha} \wedge * T_{\alpha}) = R^{\{f\alpha\beta} \wedge \eta_{\alpha\beta} - R^{\alpha\beta} \wedge \eta_{\alpha\beta} - T^{\alpha} * \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta} \rfloor T^{\beta}) - \frac{1}{2} e_{\alpha} \rfloor (T^{\beta} \wedge \vartheta_{\beta}) \right] = R^{\{f\alpha\beta} \wedge \eta_{\alpha\beta} - R^{\alpha\beta} \wedge \eta_{\alpha\beta} - T^{\alpha} \wedge * \left(- {}^{(1)}T_{\alpha} + 2 {}^{(2)}T_{\alpha} + \frac{1}{2} {}^{(3)}T_{\alpha} \right),$	$2\ell^2 d C_{\mathrm{T}} := 2d(\vartheta^{\alpha} \wedge * T_{\alpha}) = R^{\{f\alpha\beta} \wedge \eta_{\alpha\beta} - R^{\alpha\beta} \wedge \eta_{\alpha\beta} - T^{\alpha} * \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta} \rfloor T^{\beta}) - \frac{1}{2} e_{\alpha} \rfloor (T^{\beta} \wedge \vartheta_{\beta}) \right] = R^{\{f\alpha\beta} \wedge \eta_{\alpha\beta} - R^{\alpha\beta} \wedge \eta_{\alpha\beta} - T^{\alpha} \wedge * \left(- {}^{(1)}T_{\alpha} + 2 {}^{(2)}T_{\alpha} + \frac{1}{2} {}^{(3)}T_{\alpha} \right),$
mielke-geometrodynamics_129 EQ0302 (6.2.33) [conf 0.087] [refined: census]	129	■ 0.087	$\stackrel{\mathrm{rel}}{\Gamma}_{\alpha}^{\beta} := \Gamma_{\alpha}^{\beta} + i \stackrel{\mathrm{rel}}{A}^{\beta}, \quad \stackrel{\mathrm{rel}}{A}^{\beta} := \pm \frac{1}{2} \ell^2 e_{\alpha} \rfloor \stackrel{\mathrm{rel}}{\Pi}^{\beta},$	$\stackrel{(\pm)}{\Gamma}_{\alpha}^{\beta} := \Gamma_{\alpha}^{\beta} + i \stackrel{(\mp)}{A}^{\beta}, \quad \stackrel{(\pm)}{A}^{\beta} := \pm \frac{1}{2} \ell^2 e_{\alpha} \rfloor \stackrel{(\pm)}{\Pi}^{\beta},$	$\stackrel{(\pm)}{\Gamma}_{\alpha}^{\beta} := \Gamma_{\alpha}^{\beta} + i \stackrel{(\mp)}{A}_{\alpha}^{\beta}, \quad \stackrel{(\pm)}{A}_{\alpha}^{\beta} := \pm \frac{1}{2} \ell^2 e_{\alpha} \rfloor \stackrel{(\pm)}{\Pi}^{\beta},$
mielke-geometrodynamics_135 EQ0405 (6.4.25) [conf 0.098]	135	■ 0.098	$\underline{D}^{\alpha} \stackrel{\mathrm{rel}}{=} \underline{D}^{\alpha} + i \underline{A}^{\alpha}, \quad \underline{A}^{\alpha} := \pm \frac{1}{2} \ell^2 e_{\alpha} \rfloor \underline{\Pi}^{\alpha},$	$-\ell^2 \underline{D}^{\alpha} \stackrel{\mathrm{rel}}{=} \underline{D}^{\alpha} + i \underline{A}^{\alpha}, \quad \underline{A}^{\alpha} := \pm \frac{1}{2} \ell^2 e_{\alpha} \rfloor \underline{\Pi}^{\alpha},$	$-\ell^2 \underline{D}^{\alpha} \stackrel{\mathrm{rel}}{=} \underline{D}^{\alpha} + i \underline{A}^{\alpha}, \quad \underline{A}^{\alpha} := \pm \frac{1}{2} \ell^2 e_{\alpha} \rfloor \underline{\Pi}^{\alpha},$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_168 E08529 (7.7.49) [conf 0.099]	168	■ 0.099	$\begin{aligned} L_{\text{Euler}} &:= (-1)^{\text{sig}+1} \text{Tr} \left\{ \Omega \wedge \Omega^{(*)} \right\} = \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= dC_{\text{RR}^{(*)}} = \frac{(-1)^{\text{sig}}}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta^{(*)}} - \frac{1}{3} \Gamma_{\alpha}^{\beta^{(*)}} \wedge \Gamma_{\beta}^{\gamma} \wedge \Gamma_{\gamma}^{\alpha} \right) \\ &\equiv \frac{1}{2} R_{\alpha\beta} \wedge {}^* R^{\alpha\beta} - 2 \text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} + \frac{1}{2} \text{Ric}_{\alpha}^{\alpha} \wedge {}^* \text{Ric}_{\beta}^{\beta} \\ &\equiv -L_{\text{SKY}} \\ &\quad - 2 \left(\text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} - \frac{1}{4} \text{Ric}_{\alpha}^{\alpha} \wedge {}^* \text{Ric}_{\beta}^{\beta} \right) \end{aligned}$	$\begin{aligned} L_{\text{Euler}} &:= (-1)^{\text{sig}+1} \text{Tr} \left\{ \Omega \wedge \Omega^{(*)} \right\} = \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= dC_{\text{RR}^{(*)}} = \frac{(-1)^{\text{sig}}}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta^{(*)}} - \frac{1}{3} \Gamma_{\alpha}^{\beta^{(*)}} \wedge \Gamma_{\beta}^{\gamma} \wedge \Gamma_{\gamma}^{\alpha} \right) \\ &\equiv \frac{1}{2} R_{\alpha\beta} \wedge {}^* R^{\alpha\beta} - 2 \text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} + \frac{1}{2} \text{Ric}_{\alpha}^{\alpha} \wedge {}^* \text{Ric}_{\beta}^{\beta} \\ &\equiv -L_{\text{SKY}} \\ &\quad - 2 \left(\text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} - \frac{1}{4} \text{Ric}_{\alpha}^{\alpha} \wedge {}^* \text{Ric}_{\beta}^{\beta} \right) \end{aligned}$	$L_{\text{Euler}} := (-1)^{\text{sig}+1} \text{Tr} \{ \Omega \wedge \Omega^{(*)} \} = \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)}$ $= dC_{\text{RR}^{(*)}} = \frac{(-1)^{\text{sig}}}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta^{(*)}} - \frac{1}{3} \Gamma_{\alpha}^{\beta^{(*)}} \wedge \Gamma_{\beta}^{\gamma} \wedge \Gamma_{\gamma}^{\alpha} \right)$ $\equiv \frac{1}{2} R_{\alpha\beta} \wedge {}^* R^{\alpha\beta} - 2 \text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} + \frac{1}{2} \text{Ric}_{\alpha}^{\alpha} \wedge {}^* \text{Ric}_{\beta}^{\beta}$ $\equiv -L_{\text{SKY}}$ $- 2 \left(\text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} - \frac{1}{4} \text{Ric}_{\alpha}^{\alpha} \wedge {}^* \text{Ric}_{\beta}^{\beta} \right)$
mielke-geometrodynamics_315 E09974 (14.2.15) [conf 0.099]	315	■ 0.099	$\Theta^{\perp} := \pi(\hat{\Theta} - [\hat{\alpha}, \hat{\vartheta}]) = \left[\frac{\Theta}{0} \middle \frac{\hat{\ell}\Omega'}{\circ\Theta} = 0 \right],$	$\Theta^{\perp} := \pi(\hat{\Theta} - [\hat{\alpha}, \hat{\vartheta}]) = \left[\frac{\Theta}{0} \middle \frac{\hat{\ell}\Omega'}{\circ\Theta} = 0 \right],$	$\Theta^{\perp} := \pi(\hat{\Theta} - [\hat{\alpha}, \hat{\vartheta}]) = \left[\frac{\Theta}{0} \middle \frac{\hat{\ell}\Omega'}{\pi\hat{\Theta} = 0} \right],$
mielke-geometrodynamics_180 E08553 (8.4.7)	180	■ 0.100	$\begin{aligned} L_{\text{SKY}}^{(*)} &= \frac{1}{2} \text{Tr} \left(\overset{\pm}{\Omega}^g \wedge {}^* \overset{\pm}{\Omega}^g \right) \\ &= -\frac{1}{2} R^{\alpha\beta} \wedge {}^* R_{\alpha\beta} \mp \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= -\frac{1}{4} \left(R_{\alpha\beta} \pm {}^* R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} \pm {}^* R^{\alpha\beta^{(*)}} \right). \end{aligned}$	$\begin{aligned} L_{\text{SKY}}^{(*)} &= \frac{1}{2} \text{Tr} \left(\overset{\pm}{\Omega}^g \wedge {}^* \overset{\pm}{\Omega}^g \right) \\ &= -\frac{1}{2} R^{\alpha\beta} \wedge {}^* R_{\alpha\beta} \mp \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= -\frac{1}{4} \left(R_{\alpha\beta} \pm {}^* R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} \pm {}^* R^{\alpha\beta^{(*)}} \right). \end{aligned}$	$\begin{aligned} L_{\text{SKY}}^{(*)} &= \frac{1}{2} \text{Tr} \left(\overset{\pm}{\Omega}^g \wedge {}^* \overset{\pm}{\Omega}^g \right) \\ &= -\frac{1}{2} R^{\alpha\beta} \wedge {}^* R_{\alpha\beta} \mp \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= -\frac{1}{4} \left(R_{\alpha\beta} \pm {}^* R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} \pm {}^* R^{\alpha\beta^{(*)}} \right). \end{aligned}$
mielke-geometrodynamics_104 E08311 (4.6.3) [refined: census]	104	■ 0.111	$\begin{aligned} \widetilde{\leftarrow}_{\widetilde{y}} \widetilde{D} \widetilde{\xi}^{\alpha} &= \widetilde{y} \rfloor \left(\widetilde{D} \widetilde{D} \widetilde{\xi}^{\alpha} \right) + \widetilde{D}(\widetilde{y} \rfloor \widetilde{D} \widetilde{\xi}^{\alpha}) \\ &= y \rfloor \left(DD \xi^{\alpha} + R^{(T)\alpha} \right) = 0, \end{aligned}$	$\begin{aligned} \widetilde{\leftarrow}_{\widetilde{y}} \widetilde{D} \widetilde{\xi}^{\alpha} &= \widetilde{y} \rfloor \left(\widetilde{D} \widetilde{D} \widetilde{\xi}^{\alpha} \right) + \widetilde{D}(\widetilde{y} \rfloor \widetilde{D} \widetilde{\xi}^{\alpha}) \\ &= y \rfloor (DD \xi^{\alpha} + R^{(T)\alpha}) = 0, \end{aligned}$	$\begin{aligned} \widetilde{\leftarrow}_{\widetilde{y}} \widetilde{D} \widetilde{\xi}^{\alpha} &= \widetilde{y} \rfloor (\widetilde{D} \widetilde{D} \widetilde{\xi}^{\alpha}) + \widetilde{D}(\widetilde{y} \rfloor \widetilde{D} \widetilde{\xi}^{\alpha}) \\ &= y \rfloor (DD \xi^{\alpha} + R^{(T)\alpha}) = 0, \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_ E00378 (6.2.29) [re-fined: census] 128 0.136			$D\,D\,\vartheta^{\alpha}=R_{\beta}^{\alpha}\wedge\vartheta^{\beta}=0,\,D\,D\,\Pi_{\alpha}=-\,R_{\alpha}^{\beta}\wedge\Pi_{\beta}=\pm\frac{i}{2}\ell^2\left(e_{\alpha}\lrcorner\Pi^{\beta}\right)\wedge\Sigma_{\beta}\cong0.$		$DD\vartheta^{\alpha}=R_{\beta}^{\alpha}\wedge\vartheta^{\beta}=0,\,D\,D\,\Pi_{\alpha}=-\,R_{\alpha}^{\beta}\wedge\Pi_{\beta}=\pm\frac{i}{2}\ell^2\left(e_{\alpha}\lrcorner\Pi^{\beta}\right)\wedge\Sigma_{\beta}\cong0.$
mielke-geometrodynamics_ E01059 (15.2.16) 346 0.147			$\begin{aligned} \psi = & \left[\begin{array}{ccc c} \mathcal{P}_a & \mathcal{P}_b & \mathcal{P}_c & \nu_e \\ \eta_a & \eta_b & \eta_c & e^- \\ \lambda_a & \lambda_b & \lambda_c & \mu^- \\ \hline c_a & c_b & c_c & \nu_{\mu} \end{array} \right] \begin{array}{c} \text{up} \\ \text{down} \\ \text{strange} \\ \text{charm} \end{array} \begin{array}{c} \uparrow \\ \text{flavors} \\ \downarrow \end{array} \\ & \text{red} \quad \text{yellow} \quad \text{blue} \quad \text{lilac} \end{aligned}$		$\psi = \left[\begin{array}{ccc c} \mathcal{P}_a & \mathcal{P}_b & \mathcal{P}_c & \nu_e \\ \eta_a & \eta_b & \eta_c & e^- \\ \lambda_a & \lambda_b & \lambda_c & \mu^- \\ \hline c_a & c_b & c_c & \nu_{\mu} \end{array} \right] \begin{array}{c} \text{up} \\ \text{down} \\ \text{strange} \\ \text{charm} \end{array} \begin{array}{c} \uparrow \\ \text{flavors} \\ \downarrow \end{array}$
mielke-geometrodynamics_ E00972 (14.2.13) 314 0.153			$\stackrel{\circ}{D}\,\alpha:=\stackrel{\circ}{d}\,\alpha-\omega^g\wedge\alpha-\alpha\wedge\stackrel{\circ}{\omega}$		$\stackrel{\circ}{D}\alpha:=\stackrel{\circ}{d}\alpha-\omega^g\wedge\alpha-\alpha\wedge\stackrel{\circ}{\omega}$
mielke-geometrodynamics_ E00264 (4.3.29) 95 0.154			$\Gamma_{ij}'=\left\{\begin{array}{c}\kappa\\ij\end{array}\right\}-K_{ij}^{\cdot\kappa},$		$\Gamma_{ij}^{\cdot\kappa}=\left\{\begin{array}{c}\kappa\\ij\end{array}\right\}-K_{ij}^{\cdot\kappa},$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_ EQ0292 (4.5.3) [re- fined: census]	100	0.165	$\begin{aligned} \mathfrak{f}_{\xi} L &= \left(\mathfrak{f}_{\xi} g_{\alpha\beta} \right) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \left(\mathfrak{f}_{\xi} \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} \\ &+ \left(\mathfrak{f}_{\xi} \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\mathfrak{f}_{\xi} D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \\ &= D \left[\left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \right] \\ &- \left(\xi \rfloor \vartheta^{\alpha} \right) D \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor T^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor R_{\beta}^{\gamma} \right) \wedge I^{\beta}{}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} \\ &+ \left(\xi \rfloor D\Psi \right) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p \left(\xi \rfloor \Psi \right) \wedge D \frac{\delta L}{\delta \Psi} \end{aligned}$	$\mathfrak{f}_{\xi} L = \left(\mathfrak{f}_{\xi} g_{\alpha\beta} \right) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \left(\mathfrak{f}_{\xi} \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\mathfrak{f}_{\xi} \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\mathfrak{f}_{\xi} D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} = D \left[\left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \right] - \left(\xi \rfloor \vartheta^{\alpha} \right) D \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor T^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor R_{\beta}^{\gamma} \right) \wedge I^{\beta}{}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p \left(\xi \rfloor \Psi \right) \wedge D \frac{\delta L}{\delta \Psi}$	$\mathfrak{f}_{\xi} L = \left(\mathfrak{f}_{\xi} g_{\alpha\beta} \right) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \left(\mathfrak{f}_{\xi} \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\mathfrak{f}_{\xi} \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\mathfrak{f}_{\xi} D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} = D \left[\left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \right] - \left(\xi \rfloor \vartheta^{\alpha} \right) D \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor T^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor R_{\beta}^{\gamma} \right) \wedge I^{\beta}{}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p \left(\xi \rfloor \Psi \right) \wedge D \frac{\delta L}{\delta \Psi}.$
mielke-geometrodynamics_ EQ0734 (11.1.1)	237	0.220	$\pi_1 \left(SO_{\circ}(s, n-s), e \right) = \begin{cases} \mathbb{Z}_2 & n = 2k \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2 & n = 4k \\ \mathbb{Z}_4 & n = 2(2k+1) \end{cases}$	$\pi_1 \left(SO_{\circ}(s, n-s), e \right) = \begin{cases} \mathbb{Z}_2 & n = 2k \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2 & n = 4k \\ \mathbb{Z}_4 & n = 2(2k+1) \end{cases}$	$\pi_1(SO_{\circ}(s, n-s), e) = \begin{cases} \mathbb{Z}_2 & n = 2k \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2 & n = 4k \\ \mathbb{Z}_4 & n = 2(2k+1). \end{cases}$
mielke-geometrodynamics_ EQ0961 (14.2.1)	312	0.220	$\hat{\vartheta} = \hat{\vartheta}(m, g) = \underbrace{\left[\begin{array}{c c} \vartheta & \hat{\ell}\omega \\ \hline 0 & \vartheta \end{array} \right]}_4 \}^4 K.$	$\hat{\vartheta} = \hat{\vartheta}(m, g) = \left[\begin{array}{c c} \vartheta & \hat{\ell}\omega \\ \hline 0 & \vartheta \end{array} \right] \}^4 \overset{\sim}{4} \tilde{K}.$	$\hat{\vartheta} = \hat{\vartheta}(m, g) = \left[\begin{array}{c c} \vartheta & \hat{\ell}\omega \\ \hline 0 & \vartheta \end{array} \right] \}^4 \overset{\sim}{4} \tilde{K}.$
mielke-geometrodynamics_ EQ0211 (4.2.3)	85	0.223	$\begin{aligned} * \widetilde{\Omega} &= \chi \left(d\widetilde{\omega} - \frac{1}{2} [\widetilde{\omega}, \widetilde{\omega}] \right) \\ &= d\omega^L - \omega^L \wedge \omega^L + d\omega^T - \omega^T \wedge \omega^L - \omega^L \wedge \omega^T + \frac{1}{2} [\omega^T, \omega^T] \\ &= \Omega^L + \Omega^T \end{aligned}$	$\begin{aligned} {}^*\widetilde{\Omega} &= {}^*\chi \left(d\widetilde{\omega} - \frac{1}{2} [\widetilde{\omega}, \widetilde{\omega}] \right) \\ &= d\omega^L - \omega^L \wedge \omega^L + d\omega^T - \omega^T \wedge \omega^L - \omega^L \wedge \omega^T + \frac{1}{2} [\omega^T, \omega^T] \\ &= \Omega^L + \Omega^T \end{aligned}$	$\begin{aligned} {}^*\widetilde{\Omega} &= {}^*\chi (d\widetilde{\omega} - \frac{1}{2} [\widetilde{\omega}, \widetilde{\omega}]) \\ &= d\omega^L - \omega^L \wedge \omega^L + d\omega^T - \omega^T \wedge \omega^L - \omega^L \wedge \omega^T + \frac{1}{2} [\omega^T, \omega^T] \\ &= \Omega^L + \Omega^T \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geotrodynamics_E00266 (4.3.31)	95	■ 0.227	$R_{\{i\ j\}c}^{\{::d\}} = R_{\{i\ j\}c}^{\{!\ldots d\}} - 2D_{[i}K_{j]c}^{\cdot d} + 2K_{[i c}^{\cdot h}K_{ j]h}^{\cdot d}$	$R_{ijc}^{::d} = R_{ijc}^{\{!\ldots d\}} - 2D_{[i}K_{j]c}^{\cdot d} + 2K_{[i c}^{\cdot h}K_{ j]h}^{\cdot d}$	$R_{ijc}^{..d} = R_{ijc}^{\{!\ldots d\}} - 2D_{[i}K_{j]c}^{\cdot d} + 2K_{[i c}^{\cdot h}K_{ j]h}^{\cdot d}$
mielke-geotrodynamics_E00286 (4.4.15)	99	■ 0.229	$D\ \Pi_{[]}^{\mathrm{T}} - \mathrm{E}_{[]} = \Sigma_{[]}$	$D\Pi_{[]}^{\mathrm{T}} - \mathrm{E}_{[]} = \Sigma_{[]}$	$D\Pi_{[]}^{\mathrm{T}} - \mathrm{E}_{[]} = \Sigma_{[]}$
mielke-geotrodynamics_E00337 (5.3.14)	114	■ 0.239	$\begin{aligned} R^{\{\{\alpha\ \beta\}\}} \wedge \eta_{\alpha\beta} &\equiv R^{\{\alpha\ \beta\}} \wedge \eta_{\alpha\beta} - K^{\alpha\mu} \wedge K_{\mu}^{\beta} \wedge \eta_{\alpha\beta} + K^{\alpha\beta} \wedge T^{\gamma} \wedge \eta_{\alpha\beta\gamma} \\ &\quad + d(K^{\alpha\beta} \wedge \eta_{\alpha\beta}) \\ &= R^{\alpha\beta} \wedge \eta_{\alpha\beta} + T^{\alpha} \wedge \left(-^{(1)}T_{\alpha} + 2^{(2)}T_{\alpha} + \frac{1}{2}^{(3)}T_{\alpha} \right) + 2dC_{\mathrm{TT}} \end{aligned}$	$\begin{aligned} R^{\{\alpha\beta\beta\}} \wedge \eta_{\alpha\beta} &\equiv R^{\alpha\beta} \wedge \eta_{\alpha\beta} - K^{\alpha\mu} \wedge K_{\mu}^{\beta} \wedge \eta_{\alpha\beta} + K^{\alpha\beta} \wedge T^{\gamma} \wedge \eta_{\alpha\beta\gamma} \\ &\quad + d(K^{\alpha\beta} \wedge \eta_{\alpha\beta}) \\ &= R^{\alpha\beta} \wedge \eta_{\alpha\beta} + T^{\alpha} \wedge \left(-^{(1)}T_{\alpha} + 2^{(2)}T_{\alpha} + \frac{1}{2}^{(3)}T_{\alpha} \right) + 2dC_{\mathrm{TT}} \end{aligned}$	$\begin{aligned} R^{\{\alpha\beta\}} \wedge \eta_{\alpha\beta} &\equiv R^{\alpha\beta} \wedge \eta_{\alpha\beta} - K^{\alpha\mu} \wedge K_{\mu}^{\beta} \wedge \eta_{\alpha\beta} + K^{\alpha\beta} \wedge T^{\gamma} \wedge \eta_{\alpha\beta\gamma} \\ &\quad + d(K^{\alpha\beta} \wedge \eta_{\alpha\beta}) \\ &= R^{\alpha\beta} \wedge \eta_{\alpha\beta} + T^{\alpha} \wedge \left(-^{(1)}T_{\alpha} + 2^{(2)}T_{\alpha} + \frac{1}{2}^{(3)}T_{\alpha} \right) + 2dC_{\mathrm{TT}} \end{aligned} \quad (5.3)$
mielke-geotrodynamics_E00835 (12.2.3)	271	■ 0.247	$\begin{aligned} L_{\mathrm{D}} &= L\left(\gamma, \psi, D^{(f)}\psi\right) - \frac{i}{2}\bar{\psi}\left(*\gamma \wedge K - K \wedge *\gamma\right)\psi + A \wedge j \\ &= L\left(\gamma, \psi, D^{(\dagger)}\psi\right) + \frac{1}{4}\mathscr{A} \wedge j_5 + A \wedge j \\ &= L\left(\gamma, \psi, D^{(i)}\psi\right) - T^{\alpha} \wedge \mu_{\alpha} + A \wedge j. \end{aligned}$	$\begin{aligned} L_{\mathrm{D}} &= L\left(\gamma, \psi, D^{(f)}\psi\right) - \frac{i}{2}\bar{\psi}\left(*\gamma \wedge K - K \wedge *\gamma\right)\psi + A \wedge j \\ &= L\left(\gamma, \psi, D^{(\dagger)}\psi\right) + \frac{1}{4}\mathscr{A} \wedge j_5 + A \wedge j \\ &= L\left(\gamma, \psi, D^{(i)}\psi\right) - T^{\alpha} \wedge \mu_{\alpha} + A \wedge j. \end{aligned}$	$\begin{aligned} L_{\mathrm{D}} &= L(\gamma, \psi, D^{(1)}\psi) - \frac{i}{2}\bar{\psi}\left(*\gamma \wedge K - K \wedge *\gamma\right)\psi + A \wedge j \\ &= L(\gamma, \psi, D^{(1)}\psi) + \frac{1}{4}\mathscr{A} \wedge j_5 + A \wedge j \\ &= L(\gamma, \psi, D^{(1)}\psi) - T^{\alpha} \wedge \mu_{\alpha} + A \wedge j. \end{aligned}$
mielke-geotrodynamics_E00787 (11.3.6)	248	■ 0.258	$i\ \gamma \wedge *\ D^{[1}\psi - 6\ell^{*2}\gamma^{n+1}\gamma \wedge *(\bar{\psi}\gamma^{n+1}\gamma\psi)\psi - m\psi\eta = 0$	$i\gamma \wedge * D^{[1}\psi - 6\ell^{*2}\gamma^{n+1}\gamma \wedge *(\bar{\psi}\gamma^{n+1}\gamma\psi)\psi - m\psi\eta = 0$	$i\gamma \wedge * D^{(1)}\psi - 6\ell^{*2}\gamma^{n+1}\gamma \wedge *(\bar{\psi}\gamma^{n+1}\gamma\psi)\psi - m\psi\eta = 0 \Big ,$
mielke-geotrodynamics_E01078 (A.3)	365	■ 0.258	$\begin{aligned} &\& \left(\sigma^a\right)^{\mathrm{T}} := \left(\begin{array}{l} \sigma^0 \\ \sigma^1 \\ \sigma^2 \end{array}\right) \\ &\& \sigma^0 := \left(\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array}\right) \equiv \mathbb{1}, \quad \sigma^1 := \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right) \quad \text{..} \\ &\& \sigma^2 := \left(\begin{array}{cc} 0 & -i \\ i & 0 \end{array}\right), \quad \sigma^3 := \left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right). \end{aligned}$	$(\sigma^a)^{\mathrm{T}} := \begin{pmatrix} \sigma^0, \sigma^k \\ \sigma^0 := \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \equiv \mathbb{1}, \quad \sigma^1 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{..} \\ \sigma^2 := \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \end{pmatrix}$	$\begin{aligned} (\sigma^a)^{\mathrm{T}} &:= (\sigma^0, \sigma^k) \quad k = 1, 2, 3. \\ \sigma^0 &:= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \equiv \mathbb{1}, \quad \sigma^1 := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \\ \sigma^2 &:= \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \end{aligned}$
mielke-geotrodynamics_E00010 (2.3.4)	37	■ 0.261	$\delta: \left\{ \begin{array}{l} G \times P \rightarrow P \\ w \quad \psi \quad \psi \\ g_p \quad p_{\circ} = p \end{array} \middle e \cdot p_{\circ} = p_{\circ} \in P \right\}.$	$\delta: \left\{ \begin{array}{l} G \times P \rightarrow P \\ \psi \quad \psi \quad \psi \\ g_p \cdot p_{\circ} = p \end{array} \middle e \cdot p_{\circ} = p_{\circ} \in P \right\}.$	$\delta: \left\{ \begin{array}{l} G \times P \rightarrow P \\ \psi \quad \psi \quad \psi \\ g_p \cdot p_{\circ} = p \end{array} \middle e \cdot p_{\circ} = p_{\circ} \in P \right\}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_177 EQ0544 (8.3.5) [re-fined: census]	177	■ 0.261	$\exp(\lfloor\zeta\rfloor):=\mathbf{1}+\lfloor\zeta\rfloor+\frac{1}{2!}\lfloor\zeta\rfloor\lfloor\zeta\rfloor+\frac{1}{3!}\lfloor\zeta\rfloor\lfloor\zeta\rfloor\lfloor\zeta\rfloor+\cdots.$	$\exp(\zeta\rfloor) := \mathbf{1} + \zeta\rfloor + \frac{1}{2!}\zeta\rfloor\zeta\rfloor + \frac{1}{3!}\zeta\rfloor\zeta\rfloor\zeta\rfloor + \cdots.$	$\exp(\zeta\rfloor) := \mathbf{1} + \zeta\rfloor + \frac{1}{2!}\zeta\rfloor\zeta\rfloor + \frac{1}{3!}\zeta\rfloor\zeta\rfloor\zeta\rfloor + \cdots.$
mielke-geometrodynamics_199 EQ0605 (9.2.4)	199	■ 0.263	$\left.\chi(M):=\sum_{p=0}^n(-1)^pb_p=\int_M\pi^*(\gamma_2^g)\right\}\pi^*(\gamma_2^g)$	$\chi(M):=\sum_{p=0}^n(-1)^pb_p=\int_M\pi^*(\gamma_2^g)$	$\chi(M):=\sum_{p=0}^n(-1)^pb_p=\int_M\pi^*(\gamma_2^g)$
mielke-geometrodynamics_211 EQ0627 (10.1.15) [re-fined: census]	211	■ 0.271	$E_{\alpha}:=\frac{\partial L}{\partial\vartheta^{\alpha}}=e_{\alpha}\rfloor L+\left(e_{\alpha}\rfloor T^{\beta}\right)\wedge H_{\beta}+2(-1)^s\left(e_{\alpha}\rfloor R^{*\beta}\right)\wedge H_{\beta}^{\star},$	$E_{\alpha}:=\frac{\partial L}{\partial\vartheta^{\alpha}}=e_{\alpha}\rfloor L+\left(e_{\alpha}\rfloor T^{\beta}\right)\wedge H_{\beta}+2(-1)^s\left(e_{\alpha}\rfloor R^{*\beta}\right)\wedge H_{\beta}^{\star},$	$E_{\alpha}:=\frac{\partial L}{\partial\vartheta^{\alpha}}=e_{\alpha}\rfloor L+\left(e_{\alpha}\rfloor T^{\beta}\right)\wedge H_{\beta}+2(-1)^s\left(e_{\alpha}\rfloor R^{*\beta}\right)\wedge H_{\beta}^{\star},$
mielke-geometrodynamics_316 EQ0979 (14.2.20)	316	■ 0.272	$\hat{\ell}=\left(\begin{smallmatrix}+\\-\end{smallmatrix}\right)\ell^*/\sqrt{\alpha_g}=\left(\begin{smallmatrix}+\\-\end{smallmatrix}\right)\ell^*\frac{\sqrt{\hbar c}}{g}$	$\hat{\ell}=\left(\begin{smallmatrix}+\\-\end{smallmatrix}\right)\ell^*/\sqrt{\alpha_g}=\left(\begin{smallmatrix}+\\-\end{smallmatrix}\right)\ell^*\frac{\sqrt{\hbar c}}{g}$	$\hat{\ell}=\left(\begin{smallmatrix}+\\-\end{smallmatrix}\right)\ell^*/\sqrt{\alpha_g}=\left(\begin{smallmatrix}+\\-\end{smallmatrix}\right)\ell^*\frac{\sqrt{\hbar c}}{g}$
mielke-geometrodynamics_216 EQ0654 (10.3.11) [re-fined: census]	216	■ 0.274	$E_{\alpha}=e_{\alpha}\rfloor L+\left[\left(e_{\alpha}\rfloor T^{\beta}\right)+(-1)^s2\delta\ell\left(e_{\alpha}\rfloor R^{*\beta}\right)\right]\wedge H_{\beta}+(-1)^s\frac{2\gamma}{\ell}\left(e_{\alpha}\rfloor R^{*\beta}\right)\wedge\vartheta_{\beta},$	$E_{\alpha}=e_{\alpha}\rfloor L+\left[\left(e_{\alpha}\rfloor T^{\beta}\right)+(-1)^s2\delta\ell\left(e_{\alpha}\rfloor R^{*\beta}\right)\right]\wedge H_{\beta}+(-1)^s\frac{2\gamma}{\ell}\left(e_{\alpha}\rfloor R^{*\beta}\right)\wedge\vartheta_{\beta},$	$E_{\alpha}=e_{\alpha}\rfloor L+\left[\left(e_{\alpha}\rfloor T^{\beta}\right)+(-1)^s2\delta\ell\left(e_{\alpha}\rfloor R^{*\beta}\right)\right]\wedge H_{\beta}+(-1)^s\frac{2\gamma}{\ell}\left(e_{\alpha}\rfloor R^{*\beta}\right)\wedge\vartheta_{\beta},$
mielke-geometrodynamics_95 EQ0268 (4.3.33)	95	■ 0.276	$R_{i[j\kappa l]}^{\{1\}}\equiv 0$	$R_{i[j\kappa l]}^{\{1\}}\equiv 0$	$R_{i[j\kappa l]}^{\{1\}}\equiv 0$
mielke-geometrodynamics_213 EQ0639 (10.2.10)	213	■ 0.278	$R_{\alpha}^{\{i\star}=-\Lambda_{\mathrm{eff}}\,\eta_{\alpha},$	$R_{\alpha}^{\{i\star}=-\Lambda_{\mathrm{eff}}\,\eta_{\alpha},$	$R_{\alpha}^{\{1\star}=-\Lambda_{\mathrm{eff}}\,\eta_{\alpha},$
mielke-geometrodynamics_168 EQ0527 (7.7.47)	168	■ 0.282	$\begin{aligned} L_{\mathrm{Pontr}}:=&\,d\,C_{\mathrm{RR}}=\frac{1}{2}R_{\alpha}^{\beta}\wedge R_{\beta}^{\alpha}=-\frac{1}{2}R_{\alpha\beta}^{\{1\}}\wedge R^{\{j\alpha\beta}\\ &-\frac{1}{12}d\left[\ast\mathscr{A}\wedge R^{\{1\}}-\frac{1}{3}\mathscr{A}\wedge d\mathscr{A}+\frac{1}{9}\ast\mathscr{A}\wedge\ast(\mathscr{A}\wedge\ast\mathscr{A})\right]. \end{aligned}$	$L_{\mathrm{Pontr}}:=dC_{\mathrm{RR}}=\frac{1}{2}R_{\alpha}^{\beta}\wedge R_{\beta}^{\alpha}=-\frac{1}{2}R_{\alpha\beta}^{\{1\}}\wedge R^{\{j\alpha\beta}\\ -\frac{1}{12}d\left[\ast\mathscr{A}\wedge R^{\{1\}}-\frac{1}{3}\mathscr{A}\wedge d\mathscr{A}+\frac{1}{9}\ast\mathscr{A}\wedge\ast(\mathscr{A}\wedge\ast\mathscr{A})\right].$	$L_{\mathrm{Pontr}}:=dC_{\mathrm{RR}}=\frac{1}{2}R_{\alpha}^{\beta}\wedge R_{\beta}^{\alpha}=-\frac{1}{2}R_{\alpha\beta}^{\{1\}}\wedge R^{\{1\}\alpha\beta}\\ -\frac{1}{12}d\left[\ast\mathscr{A}\wedge R^{\{1\}}-\frac{1}{3}\mathscr{A}\wedge d\mathscr{A}+\frac{1}{9}\ast\mathscr{A}\wedge\ast(\mathscr{A}\wedge\ast\mathscr{A})\right].$
mielke-geometrodynamics_306 EQ0942 (14.1.23)	306	■ 0.287	$\xi\wedge\ast\xi=\pm\frac{\Omega\wedge\ast\Omega\,\mathrm{Tr}^*(\Omega\wedge\ast\Omega)-(-1)^{2z}\Omega\wedge\Omega\,\mathrm{Tr}^*(\Omega\wedge\Omega)}{\sqrt{(\mathrm{Tr}^*(\Omega\wedge\ast\Omega))^2+(-1)^{2z}(\mathrm{Tr}^*(\Omega\wedge\Omega))^2}}.$	$\xi\wedge\ast\xi=\pm\frac{\Omega\wedge\ast\Omega\,\mathrm{Tr}^*(\Omega\wedge\ast\Omega)-(-1)^{2z}\Omega\wedge\Omega\,\mathrm{Tr}^*(\Omega\wedge\Omega)}{\sqrt{(\mathrm{Tr}^*(\Omega\wedge\ast\Omega))^2+(-1)^{2z}(\mathrm{Tr}^*(\Omega\wedge\Omega))^2}}.$	$\xi\wedge\ast\xi=\pm\frac{\Omega\wedge\ast\Omega\,\mathrm{Tr}^*(\Omega\wedge\ast\Omega)-(-1)^{2z}\Omega\wedge\Omega\,\mathrm{Tr}^*(\Omega\wedge\Omega)}{\sqrt{(\mathrm{Tr}^*(\Omega\wedge\ast\Omega))^2+(-1)^{2z}(\mathrm{Tr}^*(\Omega\wedge\Omega))^2}}.$
mielke-geometrodynamics_143 EQ0440 (6.8.11) [re-fined: census]	143	■ 0.288	$D\Sigma_{\alpha}\cong(e_{\alpha}\rfloor T^{\gamma})\wedge\Sigma_{\gamma}+(e_{\alpha}\rfloor R^{\gamma\delta})\wedge\tau_{\gamma\delta}$	$D\Sigma_{\alpha}\cong(e_{\alpha}\rfloor T^{\gamma})\wedge\Sigma_{\gamma}+(e_{\alpha}\rfloor R^{\gamma\delta})\wedge\tau_{\gamma\delta}$	$D\Sigma_{\alpha}\cong(e_{\alpha}\rfloor T^{\gamma})\wedge\Sigma_{\gamma}+(e_{\alpha}\rfloor R^{\gamma\delta})\wedge\tau_{\gamma\delta}$
mielke-geometrodynamics_105 EQ0315 (4.6.7)	105	■ 0.292	$\Delta\Phi(\widetilde{\Gamma},\gamma)\simeq\frac{i}{\hbar}\int_S\Big(R^{(T)\alpha}P_{\alpha}+R_{\alpha}^{(L)\beta}L^{\alpha}_{\beta}\Big)$	$\Delta\Phi(\widetilde{\Gamma},\gamma)\simeq\frac{i}{\hbar}\int_S\Big(R^{(T)\alpha}P_{\alpha}+R_{\alpha}^{(L)\beta}L^{\alpha}_{\beta}\Big)$	$\Delta\Phi(\widetilde{\Gamma},\gamma)\simeq\frac{i}{\hbar}\int_S(R^{(T)\alpha}P_{\alpha}+R_{\alpha}^{(L)\beta}L^{\alpha}_{\beta}).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_365 EQ1082 (A.7)		■ 0.326	$\begin{aligned} & \gamma^0 := \left(\begin{array}{cc} \sigma^0 & 0 \\ 0 & \sigma^0 \end{array} \right), & \gamma^k := \left(\begin{array}{cc} 0 & \sigma^k \\ -\sigma^k & 0 \end{array} \right); \\ & \gamma^5 := i\gamma^0\gamma^1\gamma^2\gamma^3 = \left(\begin{array}{cc} 0 & \sigma^0 \\ \sigma^0 & 0 \end{array} \right). \end{aligned}$	$\gamma^0 := \left(\begin{array}{cc} \sigma^0 & 0 \\ 0 & \sigma^0 \end{array} \right), \quad \gamma^k := \left(\begin{array}{cc} 0 & \sigma^k \\ -\sigma^k & 0 \end{array} \right);$ $\gamma^5 := i\gamma^0\gamma^1\gamma^2\gamma^3 = \left(\begin{array}{cc} 0 & \sigma^0 \\ \sigma^0 & 0 \end{array} \right).$	$\gamma^0 := \left(\begin{array}{cc} \sigma^0 & 0 \\ 0 & \sigma^0 \end{array} \right), \quad \gamma^k := \left(\begin{array}{cc} 0 & \sigma^k \\ -\sigma^k & 0 \end{array} \right);$ $\gamma^5 := i\gamma^0\gamma^1\gamma^2\gamma^3 = \left(\begin{array}{cc} 0 & \sigma^0 \\ \sigma^0 & 0 \end{array} \right).$
mielke-geometrodynamics_316 EQ0978 (14.2.19)		■ 0.331	$\begin{aligned} & \pi \hat{L}_{ } = -\frac{\hat{\ell}^2}{\ell^{*2}} \text{Tr}(\Omega' \wedge {}^*\Omega' \wedge \underbrace{\overset{\circ}{\vartheta} \wedge \cdots \wedge \overset{\circ}{\vartheta}}_{\text{K terms}}) \\ & = -\frac{1}{\alpha_g} \text{Tr}(\Omega' \wedge {}^*\Omega') \sqrt{\det \overset{\circ}{g}} = L'_{\text{YM}} d_\mu(G) \end{aligned}$	$\pi \hat{L}_{ } = -\frac{\hat{\ell}^2}{\ell^{*2}} \text{Tr}(\Omega' \wedge {}^*\Omega' \wedge \underbrace{\overset{\circ}{\vartheta} \wedge \cdots \wedge \overset{\circ}{\vartheta}}_{\text{K terms}})$ $= -\frac{1}{\alpha_g} \text{Tr}(\Omega' \wedge {}^*\Omega') \sqrt{\det \overset{\circ}{g}} = L'_{\text{YM}} d_\mu(G)$	$\pi \hat{L}_{ } = -\frac{\hat{\ell}^2}{\ell^{*2}} \text{Tr}(\Omega' \wedge {}^*\Omega' \wedge \underbrace{\overset{\circ}{\vartheta} \wedge \cdots \wedge \overset{\circ}{\vartheta}}_{\text{K terms}})$ $= -\frac{1}{\alpha_g} \text{Tr}(\Omega' \wedge {}^*\Omega') \sqrt{\det \overset{\circ}{g}} = L'_{\text{YM}} d_\mu(G)$
mielke-geometrodynamics_186 EQ0579 (8.6.17) [re-fined: census]		■ 0.351	$2 \, D \left(e^\beta \rfloor DH_{\alpha\beta} - \frac{1}{4} \vartheta_\alpha \wedge e^\gamma \rfloor e^\delta \rfloor DH_{\gamma\delta} \right) - E_\alpha = \Sigma_\alpha - D\mu_\alpha,$	$2D \left(e^\beta \rfloor DH_{\alpha\beta} - \frac{1}{4} \vartheta_\alpha \wedge e^\gamma \rfloor e^\delta \rfloor DH_{\gamma\delta} \right) - E_\alpha = \Sigma_\alpha - D\mu_\alpha,$	$2D \left(e^\beta \rfloor DH_{\alpha\beta} - \frac{1}{4} \vartheta_\alpha \wedge e^\gamma \rfloor e^\delta \rfloor DH_{\gamma\delta} \right) - E_\alpha = \Sigma_\alpha - D\mu_\alpha,$
mielke-geometrodynamics_309 EQ0956 (14.1.38)		■ 0.353	$\begin{aligned} & d \, \delta = (-1)^{-z} {}^*\Sigma_{\text{YM}}(\Omega) \wedge D\Sigma_{\text{YM}}(\Omega) / {}^*(\Sigma_{\text{YM}} \wedge \Sigma_{\text{YM}}) \\ & = (-1)^{-z} {}^*(\vartheta \wedge {}^*\Omega^g) \wedge D(\vartheta \wedge {}^*\Omega^g) / {}^*(\vartheta \wedge {}^*\Omega^g \wedge {}^*(\vartheta \wedge {}^*\Omega^g)). \end{aligned}$	$d\delta = (-1)^{-z} {}^*\Sigma_{\text{YM}}(\Omega) \wedge D\Sigma_{\text{YM}}(\Omega) / {}^*(\Sigma_{\text{YM}} \wedge \Sigma_{\text{YM}})$ $= (-1)^{-z} {}^*(\vartheta \wedge {}^*\Omega^g) \wedge D(\vartheta \wedge {}^*\Omega^g) / {}^*(\vartheta \wedge {}^*\Omega^g \wedge {}^*(\vartheta \wedge {}^*\Omega^g)).$	$d\delta = (-1)^{-z} {}^*\Sigma_{\text{YM}}(\Omega) \wedge D\Sigma_{\text{YM}}(\Omega) / {}^*(\Sigma_{\text{YM}} \wedge \Sigma_{\text{YM}})$ $= (-1)^{-z} {}^*(\vartheta \wedge {}^*\Omega^g) \wedge D(\vartheta \wedge {}^*\Omega^g) / {}^*(\vartheta \wedge {}^*\Omega^g \wedge {}^*(\vartheta \wedge {}^*\Omega^g)).$
mielke-geometrodynamics_180 EQ0555 (8.4.9) [re-fined: census]		■ 0.371	$E_\alpha = \frac{1}{2} \text{Tr} \left(\overset{\pm}{\Omega}^g \wedge e_\alpha \rfloor {}^*\overset{\pm}{\Omega}^g - e_\alpha \rfloor \overset{\pm}{\Omega}^g \wedge {}^*\overset{\pm}{\Omega}^g \right) = 0,$	$E_\alpha = \frac{1}{2} \text{Tr} \left(\overset{\pm}{\Omega}^g \wedge e_\alpha \rfloor {}^*\overset{\pm}{\Omega}^g - e_\alpha \rfloor \overset{\pm}{\Omega}^g \wedge {}^*\overset{\pm}{\Omega}^g \right) = 0,$	$E_\alpha = \frac{1}{2} \text{Tr} \left(\overset{\pm}{\Omega}^g \wedge e_\alpha \rfloor {}^*\overset{\pm}{\Omega}^g - e_\alpha \rfloor \overset{\pm}{\Omega}^g \wedge {}^*\overset{\pm}{\Omega}^g \right) = 0,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_182 E09562 (8.5.5)		■ 0.377	$\begin{aligned} \widetilde{L}_{\mathrm{SKY}}^{\left(\star\right)} &= \frac{1}{2} \operatorname{Tr} \left[\left(\left(\frac{\pm}{\varOmega}{}^g - i \frac{\theta_{\mathrm{T}}^{\star}}{4 \ell^2} \sigma \right) \wedge^* \left(\frac{\pm}{\varOmega}{}^g - i \frac{\theta_{\mathrm{T}}^{\star}}{4 \ell^2} \sigma \right) \right) \right. \\ &\quad \left. - \frac{\theta_{\mathrm{T}}^{\star}}{2 \ell^2} R^{\alpha \beta} \wedge \eta_{\alpha \beta} + \theta_{\mathrm{T}}^{\star} \Lambda_{\mathrm{eff}} \eta - \frac{1}{2} R^{\alpha \beta} \wedge^* R_{\alpha \beta} \mp \frac{\left(-1\right)^{\mathrm{sig}}}{2} R^{\alpha \beta} \wedge R_{\alpha \beta}^{\left(\star\right)} \right] \end{aligned}$	$\widetilde{L}_{\mathrm{SKY}}^{\left(\star\right)} = \frac{1}{2} \operatorname{Tr} \left[\left(\left(\otimes^g - i \frac{\theta_{\mathrm{T}}^{\star}}{4 \ell^2} \sigma \right) \wedge^* \left(\frac{\pm}{\varOmega}{}^g - i \frac{\theta_{\mathrm{T}}^{\star}}{4 \ell^2} \sigma \right) \right) \right. \\ \left. = - \frac{\theta_{\mathrm{T}}^{\star}}{2 \ell^2} R^{\alpha \beta} \wedge \eta_{\alpha \beta} + \theta_{\mathrm{T}}^{\star} \Lambda_{\mathrm{eff}} \eta - \frac{1}{2} R^{\alpha \beta} \wedge^* R_{\alpha \beta} \mp \frac{\left(-1\right)^{\mathrm{sig}}}{2} R^{\alpha \beta} \wedge R_{\alpha \beta}^{\left(\star\right)} \right]$	$\widetilde{L}_{\mathrm{SKY}}^{\left(\star\right)} = \frac{1}{2} \operatorname{Tr} \left[\left(\left(\frac{\pm}{\varOmega}{}^g - i \frac{\theta_{\mathrm{T}}^{\star}}{4 \ell^2} \sigma \right) \wedge^* \left(\frac{\pm}{\varOmega}{}^g - i \frac{\theta_{\mathrm{T}}^{\star}}{4 \ell^2} \sigma \right) \right) \right] \quad (8.5)$ $= - \frac{\theta_{\mathrm{T}}^{\star}}{2 \ell^2} R^{\alpha \beta} \wedge \eta_{\alpha \beta} + \theta_{\mathrm{T}}^{\star} \Lambda_{\mathrm{eff}} \eta - \frac{1}{2} R^{\alpha \beta} \wedge^* R_{\alpha \beta} \mp \frac{\left(-1\right)^{\mathrm{sig}}}{2} R^{\alpha \beta} \wedge R_{\alpha \beta}^{\left(\star\right)}.$
mielke-geometrodynamics_132 E09393 (6.4.8) [re-fined: census]		■ 0.382	$\begin{aligned} n^{\alpha} \Sigma_{\alpha} &= n \rfloor L - \left(\mathfrak{L}_n \Psi \right) \wedge \frac{\partial L}{\partial D \Psi} - \left(n \rfloor \Psi \right) \wedge \frac{\delta L}{\delta \Psi} + D \left[\left(n \rfloor \Psi \right) \wedge \frac{\partial L}{\partial D \Psi} \right] \\ &\cong n \rfloor L - \left(\ell_n \Psi + \Gamma_{\perp}^{\alpha \beta} \rho \left(I_{\alpha \beta} \right) \Psi \right) \wedge \frac{\partial L}{\partial D \Psi} + D \left[\left(n \rfloor \Psi \right) \wedge \frac{\partial L}{\partial D \Psi} \right]. \end{aligned}$	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - \left(\mathfrak{L}_n \Psi \right) \wedge \frac{\partial L}{\partial D \Psi} - \left(n \rfloor \Psi \right) \wedge \frac{\delta L}{\delta \Psi} + D \left[\left(n \rfloor \Psi \right) \wedge \frac{\partial L}{\partial D \Psi} \right]$ $\cong n \rfloor L - \left(\ell_n \Psi + \Gamma_{\perp}^{\alpha \beta} \rho \left(I_{\alpha \beta} \right) \Psi \right) \wedge \frac{\partial L}{\partial D \Psi} + D \left[\left(n \rfloor \Psi \right) \wedge \frac{\partial L}{\partial D \Psi} \right].$	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - \left(\mathfrak{L}_n \Psi \right) \wedge \frac{\partial L}{\partial D \Psi} - \left(n \rfloor \Psi \right) \wedge \frac{\delta L}{\delta \Psi} + D \left[\left(n \rfloor \Psi \right) \wedge \frac{\partial L}{\partial D \Psi} \right]$ $\cong n \rfloor L - \left(\ell_n \Psi + \Gamma_{\perp}^{\alpha \beta} \rho \left(I_{\alpha \beta} \right) \Psi \right) \wedge \frac{\partial L}{\partial D \Psi} + D \left[\left(n \rfloor \Psi \right) \wedge \frac{\partial L}{\partial D \Psi} \right].$
mielke-geometrodynamics_94 E09261 (4.3.26)		■ 0.394	$K_{\cdot \alpha \beta}^c = - \frac{1}{2} \left(T_{\cdot \alpha \beta}^c - T_{\alpha \cdot \beta}^{\cdot c} + T_{\beta \cdot \alpha}^{\cdot c} \right) = - K_{\cdot \beta \alpha}^c.$	$K_{\cdot \alpha \beta}^c = - \frac{1}{2} \left(T_{\cdot \alpha \beta}^c - T_{\alpha \cdot \beta}^{\cdot c} + T_{\beta \cdot \alpha}^{\cdot c} \right) = - K_{\cdot \beta \alpha}^c.$	$K_{\cdot \alpha \beta}^c = - \frac{1}{2} \left(T_{\cdot \alpha \beta}^c - T_{\alpha \cdot \beta}^{\cdot c} + T_{\beta \cdot \alpha}^{\cdot c} \right) = - K_{\cdot \beta \alpha}^c.$
mielke-geometrodynamics_128 E09377 (6.2.28)		■ 0.396	$\stackrel{\scriptscriptstyle{(\pm)}}{L}_{\parallel} = - \frac{1}{2} \left[\vartheta^{\alpha} \wedge \stackrel{\scriptscriptstyle{(\pm)}}{D}^{\scriptscriptstyle{(\pm)}}_{\alpha} + d \left(\vartheta^{\alpha} \wedge \stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha} \right) \right] = - \frac{1}{2} \left(\stackrel{\scriptscriptstyle{(\pm)}}{D} \vartheta^{\alpha} \wedge \stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}, \right.$	$\stackrel{\scriptscriptstyle{(\pm)}}{L}_{\parallel} = - \frac{1}{2} \left[\vartheta^{\alpha} \wedge \stackrel{\scriptscriptstyle{(\pm)}}{D}^{\scriptscriptstyle{(\pm)}}_{\alpha} + d \left(\vartheta^{\alpha} \wedge \stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha} \right) \right] = - \frac{1}{2} \left(\stackrel{\scriptscriptstyle{(\pm)}}{D} \vartheta^{\alpha} \wedge \stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}, \right.$	$\stackrel{\scriptscriptstyle{(\pm)}}{L}_{\parallel} = - \frac{1}{2} \left[\vartheta^{\alpha} \wedge \stackrel{\scriptscriptstyle{(\pm)}}{D}^{\scriptscriptstyle{(\pm)}}_{\alpha} + d \left(\vartheta^{\alpha} \wedge \stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha} \right) \right] = - \frac{1}{2} \stackrel{\scriptscriptstyle{(\pm)}}{D} \vartheta^{\alpha} \wedge \stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},$
mielke-geometrodynamics_94 E09262 (4.3.27)		■ 0.399	$\begin{aligned} \Omega^g &= \Omega^{\{\}} - dK + K \wedge \omega^{\{\}} + \omega^{\{\}} \wedge K - K \wedge K \\ &= \Omega^{\{\}} - D^{\{\}} K - K \wedge K = \Omega^{\{\}} - DK + K \wedge K \end{aligned}$	$\Omega^g = \Omega^{\{\}} - dK + K \wedge \omega^{\{\}} + \omega^{\{\}} \wedge K - K \wedge K$ $= \Omega^{\{\}} - D^{\{\}} K - K \wedge K = \Omega^{\{\}} - DK + K \wedge K$	$\Omega^g = \Omega^{\{\}} - dK + K \wedge \omega^{\{\}} + \omega^{\{\}} \wedge K - K \wedge K$ $= \Omega^{\{\}} - D^{\{\}} K - K \wedge K = \Omega^{\{\}} - DK + K \wedge K$
mielke-geometrodynamics_184 E09579 (8.6.8) [re-fined: census]		■ 0.403	$E_{\alpha} := \partial L / \partial \vartheta^{\alpha} = e_{\alpha} \rfloor L + \left(e_{\alpha} \rfloor T^{\beta} \right) \wedge H_{\beta} + \left(e_{\alpha} \rfloor R^{\beta \gamma} \right) \wedge H_{\beta \gamma}$	$E_{\alpha} := \partial L / \partial \vartheta^{\alpha} = e_{\alpha} \rfloor L + \left(e_{\alpha} \rfloor T^{\beta} \right) \wedge H_{\beta} + \left(e_{\alpha} \rfloor R^{\beta \gamma} \right) \wedge H_{\beta \gamma}$	$E_{\alpha} := \partial L / \partial \vartheta^{\alpha} = e_{\alpha} \rfloor L + \left(e_{\alpha} \rfloor T^{\beta} \right) \wedge H_{\beta} + \left(e_{\alpha} \rfloor R^{\beta \gamma} \right) \wedge H_{\beta \gamma}$
mielke-geometrodynamics_103 E09396 (4.5.16) [re-fined: census]		■ 0.406	$\widetilde{\varepsilon}_{\mathrm{MA}} := \varepsilon_{\mathrm{RC}} + \xi \rfloor L \cong - \left(\mathcal{L}_{\xi} \Gamma_{\beta}^{\gamma} \right) \wedge H^{\beta}{}_{\gamma} + dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}} := \varepsilon_{\mathrm{RC}} + \xi \rfloor L \cong - \left(\mathcal{L}_{\xi} \Gamma_{\beta}^{\gamma} \right) \wedge H^{\beta}{}_{\gamma} + dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}} := \varepsilon_{\mathrm{RC}} + \xi \rfloor L \cong - \left(\mathcal{L}_{\xi} \Gamma_{\beta}^{\gamma} \right) \wedge H^{\beta}{}_{\gamma} + dH_{\infty}$
mielke-geometrodynamics_114 E09336 (5.3.13)		■ 0.407	$R^{\alpha \beta} = R^{\{\} j \alpha \beta} - D^{\{\}} K^{\alpha \beta} - K^{\alpha}{}_{\mu} \wedge K^{\mu \beta}$	$R^{\alpha \beta} = R^{\{\} j \alpha \beta} - D^{\{\}} K^{\alpha \beta} - K^{\alpha}{}_{\mu} \wedge K^{\mu \beta}$	$R^{\alpha \beta} = R^{\{\} \alpha \beta} - D^{\{\}} K^{\alpha \beta} - K^{\alpha}{}_{\mu} \wedge K^{\mu \beta}$
mielke-geometrodynamics_375 EQ1136 (C.4)		■ 0.409	$c_{ij}^k = - c_{ji}^k$	$c_{ij}^k = - c_{ji}^k$	$c_{ij}^k = - c_{ji}^k$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_279 EQ0861 (12.4.8)		■ 0.420	$\begin{aligned} d\,C_{\mathrm{RR}} &= -\operatorname{Tr}(\Omega \wedge \Omega) \\ &= \frac{1}{2}R_{\alpha\beta}^{\{\}} \wedge R^{\{\beta\alpha\}} \\ &+ \frac{1}{12}d\left[{}^*\mathcal{A} \wedge R^{\{\}} - \frac{1}{3}\mathcal{A} \wedge d\mathcal{A} + \frac{1}{9}{}^*\mathcal{A} \wedge {}^*(\mathcal{A} \wedge {}^*\mathcal{A})\right]. \end{aligned}$	$dC_{\mathrm{RR}} = -\operatorname{Tr}(\Omega \wedge \Omega)$ $= \frac{1}{2}R_{\alpha\beta}^{\{\}} \wedge R^{\{\beta\alpha\}}$ $+ \frac{1}{12}d\left[{}^*\mathcal{A} \wedge R^{\{\}} - \frac{1}{3}\mathcal{A} \wedge d\mathcal{A} + \frac{1}{9}{}^*\mathcal{A} \wedge {}^*(\mathcal{A} \wedge {}^*\mathcal{A})\right].$	$dC_{\mathrm{RR}} = -\operatorname{Tr}(\Omega \wedge \Omega)$ $= \frac{1}{2}R_{\alpha\beta}^{\{\}} \wedge R^{\{\}\alpha\beta}$ $+ \frac{1}{12}d\left[{}^*\mathcal{A} \wedge R^{\{\}} - \frac{1}{3}\mathcal{A} \wedge d\mathcal{A} + \frac{1}{9}{}^*\mathcal{A} \wedge {}^*(\mathcal{A} \wedge {}^*\mathcal{A})\right].$
mielke-geometrodynamics_151 EQ0451 (7.2.4)		■ 0.424	$\stackrel{+}{\Omega} := \frac{1}{2}\left(\Omega^g \pm {}^*\Omega^{g(*)}\right),$	$\stackrel{+}{\Omega} := \frac{1}{2}\left(\Omega^g \pm {}^*\Omega^{g(*)}\right),$	$\stackrel{\pm}{\Omega}^g := \frac{1}{2}\left(\Omega^g \pm {}^*\Omega^{g(*)}\right),$
mielke-geometrodynamics_127 EQ0372 (6.2.23) [re-fined: census]		■ 0.437	$\stackrel{(\pm)}{D}\stackrel{(\pm)}{\Pi}_\alpha \mp \frac{i}{4}\ell^2 e_\alpha \Bigg[\stackrel{(+)}{\Pi}^\beta \wedge \stackrel{(-)}{\Pi}_\beta \Bigg] = \Sigma_\alpha,$	$\stackrel{(\pm)}{D}\stackrel{(\pm)}{\Pi}_\alpha \mp \frac{i}{4}\ell^2 e_\alpha \Bigg[\stackrel{(+)}{\Pi}^\beta \wedge \stackrel{(-)}{\Pi}_\beta \Bigg] = \Sigma_\alpha,$	$\stackrel{(\pm)}{D}\stackrel{(\pm)}{\Pi}_\alpha \mp \frac{i}{4}\ell^2 e_\alpha \Bigg[\stackrel{(+)}{\Pi}^\beta \wedge \stackrel{(-)}{\Pi}_\beta \Bigg] = \Sigma_\alpha,$
mielke-geometrodynamics_322 EQ0997 (14.3.10) [re-fined: census]		■ 0.442	$\cancel{D}^{\text{int}} = \stackrel{\circ}{D} \wedge {}^*\stackrel{\circ}{D} = \stackrel{\circ}{D} \wedge {}^*(\stackrel{\circ}{d} + i\stackrel{\circ}{\tilde{\omega}})$	$\cancel{D}^{(\text{int})} = \stackrel{\circ}{D} \wedge {}^*\stackrel{\circ}{D} = \stackrel{\circ}{D} \wedge {}^*(\stackrel{\circ}{d} + i\stackrel{\circ}{\tilde{\omega}})$	$\cancel{D}^{(int)} = \stackrel{\circ}{D} \wedge {}^*\stackrel{\circ}{D} = \stackrel{\circ}{D} \wedge {}^*(\stackrel{\circ}{d} + i\stackrel{\circ}{\tilde{\omega}})$
mielke-geometrodynamics_156 EQ0474 (7.4.2)		■ 0.444	$R_{\alpha\beta} \equiv C_{\alpha\beta} + \frac{1}{2}Ri^\tau_{\alpha\beta} - \frac{R}{12}\vartheta_\alpha \wedge \vartheta_\beta,$	$R_{\alpha\beta} \equiv C_{\alpha\beta} + \frac{1}{2}Ri^\tau_{\alpha\beta} - \frac{R}{12}\vartheta_\alpha \wedge \vartheta_\beta,$	$R_{\alpha\beta} \equiv C_{\alpha\beta} + \frac{1}{2}Ri\check{\epsilon}_{\alpha\beta} - \frac{R}{12}\vartheta_\alpha \wedge \vartheta_\beta,$
mielke-geometrodynamics_221 EQ0681 (10.3.38)		■ 0.452	$\begin{aligned} \chi &= \theta_{\mathrm{L}}\varepsilon - \tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}, \quad \Lambda = \frac{(-1)^s}{4\ell^2}\left(\theta_{\mathrm{L}}\varepsilon^3 - \tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^3\right), \quad \theta_{\mathrm{T}} = \frac{(-1)^s}{2}\left(\theta_{\mathrm{L}}\varepsilon^2 - \tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^2\right), \\ \Delta\theta_{\mathrm{L}} &= \theta_{\mathrm{L}} - \tilde{\theta}_{\mathrm{L}}, \quad \Delta\theta_{\mathrm{TL}} = \frac{1}{2}\left(\theta_{\mathrm{L}}\varepsilon - \tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}\right). \end{aligned}$	$\chi = \theta_{\mathrm{L}}\varepsilon - \tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}, \quad \Lambda = \frac{(-1)^s}{4\ell^2}\left(\theta_{\mathrm{L}}\varepsilon^3 - \tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^3\right), \quad \theta_{\mathrm{T}} = \frac{(-1)^s}{2}\left(\theta_{\mathrm{L}}\varepsilon^2 - \tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^2\right),$ $\Delta\theta_{\mathrm{L}} = \theta_{\mathrm{L}} - \tilde{\theta}_{\mathrm{L}}, \quad \Delta\theta_{\mathrm{TL}} = \frac{1}{2}\left(\theta_{\mathrm{L}}\varepsilon - \tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}\right).$	$\chi = \theta_{\mathrm{L}}\varepsilon - \tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}, \quad \Lambda = \frac{(-1)^s}{4\ell^2}(\theta_{\mathrm{L}}\varepsilon^3 - \tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^3), \quad \theta_{\mathrm{T}} = \frac{(-1)^s}{2}(\theta_{\mathrm{L}}\varepsilon^2 - \tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^2),$ $\Delta\theta_{\mathrm{L}} = \theta_{\mathrm{L}} - \tilde{\theta}_{\mathrm{L}}, \quad \Delta\theta_{\mathrm{TL}} = \frac{1}{2}(\theta_{\mathrm{L}}\varepsilon - \tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}).$
mielke-geometrodynamics_273 EQ0845 (12.2.13)		■ 0.457	$C_{\mathrm{TT}} \cong \frac{1}{4}j_5, \quad \mathscr{A} = 2\ell^2{}^*C_{\mathrm{TT}} = (\ell^2/2)\bar{\psi}\gamma_5\gamma\psi.$	$C_{\mathrm{TT}} \cong \frac{1}{4}j_5, \quad \mathscr{A} = 2\ell^2{}^*C_{\mathrm{TT}} = (\ell^2/2)\bar{\psi}\gamma_5\gamma\psi.$	$C_{\mathrm{TT}} \cong \frac{1}{4}j_5, \quad \mathscr{A} = 2\ell^2{}^*C_{\mathrm{TT}} = (\ell^2/2)\bar{\psi}\gamma_5\gamma\psi.$
mielke-geometrodynamics_128 EQ0375 (6.2.26)		■ 0.466	$\stackrel{(\pm)}{L}_\parallel = \mp \frac{i}{4}\ell^2 \stackrel{(\pm)}{\Pi}^\alpha \wedge \stackrel{(\pm)}{\Pi}_\alpha, \quad \overline{\stackrel{(+)}{L}_\parallel} = \stackrel{(-)}{L}_\parallel,$	$\stackrel{(\pm)}{L}_\parallel = \mp \frac{i}{4}\ell^2 \stackrel{(\pm)}{\Pi}^\alpha \wedge \stackrel{(\pm)}{\Pi}_\alpha, \quad \overline{\stackrel{(+)}{L}_\parallel} = \stackrel{(-)}{L}_\parallel,$	$\stackrel{(\pm)}{L}_\parallel = \mp \frac{i}{4}\ell^2 \stackrel{(\pm)}{\Pi}^\alpha \wedge \stackrel{(\pm)}{\Pi}_\alpha, \quad \overline{\stackrel{(+)}{L}_\parallel} = \stackrel{(-)}{L}_\parallel,$
mielke-geometrodynamics_87 EQ0222 (4.2.15)		■ 0.466	$\widetilde{\varphi}: A(M) \rightarrow C^\infty(\overset{\circ}{V})$	$\widetilde{\varphi}: A(M) \rightarrow C^\infty(\overset{\circ}{V})$	$\widetilde{\varphi}: A(M) \rightarrow C^\infty(\overset{\circ}{V})$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometroynamics_199 EQ0604 (9.2.3)		■ 0.474	$\begin{aligned} L_{\text{Euler}} &= \frac{1}{(4\pi)^2} \text{Tr} \left(\Omega^g \wedge \Omega^{g(*)} \right) \\ &= \frac{(-1)^{s/4}}{32\pi^2} R^{\alpha\beta\gamma\delta} R^*_{\alpha\beta\gamma\delta} \sqrt{ g } d^4x. \end{aligned}$	$L_{\text{Euler}} = \frac{1}{(4\pi)^2} \text{Tr} \left(\Omega^g \wedge \Omega^{g(*)} \right) = \frac{(-1)^{s/4}}{32\pi^2} R^{\alpha\beta\gamma\delta} R^*_{\alpha\beta\gamma\delta} \sqrt{ g } d^4x.$	$L_{\text{Euler}} = \pi^* (\gamma_2^g) = \frac{1}{(4\pi)^2} \text{Tr} (\Omega^g \wedge \Omega^{g(*)}) = \frac{(-1)^{s/4}}{32\pi^2} R^{\alpha\beta\gamma\delta} R^*_{\alpha\beta\gamma\delta} \sqrt{ g } d^4x.$
mielke-geometroynamics_133 EQ0401 (6.4.20)		■ 0.482	$\mathscr{H}_{\parallel} = \underline{D} - \underline{\mathscr{A}}_{\hat{0}} = 0,$	$\mathscr{H}_{\parallel} = \underline{D} - \underline{\mathscr{A}}_{\hat{0}} = 0,$	$\mathscr{H}_{\parallel} = \underline{D} - \underline{\mathscr{A}}_{\hat{0}} = 0,$
mielke-geometroynamics_241 EQ0750 (11.1.17)		■ 0.484	$D_{\left[\overparen{S_0}(1,3)\right]}^{\mathrm{F}} = \underbrace{D^{\frac{1}{2}} \times \cdots \times D^{\frac{1}{2}}}_{(n-4)/4} \times D^{\frac{1}{2}}.$	$D_{\widehat{SO}(1,3)}^{\mathrm{F}} = \underbrace{D^{\frac{1}{2}} \times \cdots \times D^{\frac{1}{2}}}_{(n-4)/4}.$	$D_{\widehat{SO}(1,3)}^{\mathrm{F}} = \underbrace{D^{\frac{1}{2}} \times \cdots \times D^{\frac{1}{2}}}_{(n-4)/4}.$
mielke-geometroynamics_222 EQ0686 (10.3.44) [re-fined: census]		■ 0.497	$C_l = E_l^\alpha * \left\{ D^{\{\}} \left[2 \left(e^\beta \rfloor D^{\{\}} H_{[\alpha\beta]} \right) + \frac{1}{2} \vartheta_\alpha \wedge \left(e^\beta \rfloor e^\gamma \rfloor D^{\{\}} H_{[\beta\gamma]} \right) \right] \right\}$	$C_l = E_l^\alpha * \left\{ D^{\{\}} \left[2 \left(e^\beta \rfloor D^{\{\}} H_{[\alpha\beta]} \right) + \frac{1}{2} \vartheta_\alpha \wedge \left(e^\beta \rfloor e^\gamma \rfloor D^{\{\}} H_{[\beta\gamma]} \right) \right] \right\}$	$C_l = E_l^\alpha * \left\{ D^{\{\}} \left[2 \left(e^\beta \rfloor D^{\{\}} H_{[\alpha\beta]} \right) + \frac{1}{2} \vartheta_\alpha \wedge \left(e^\beta \rfloor e^\gamma \rfloor D^{\{\}} H_{[\beta\gamma]} \right) \right] \right\}$
mielke-geometroynamics_166 EQ0515 (7.7.33)		■ 0.497	$\left. \left. T := \frac{1}{4} \operatorname{Tr}(\check{\gamma} \rfloor \Theta) = e_\alpha \rfloor T^\alpha, \quad \mathscr{A} := \frac{1}{4} {}^* \operatorname{Tr}(\gamma \wedge \Theta) = {}^*(\vartheta_\alpha \wedge T^\alpha) \right. \right.$	$T := \frac{1}{4} \operatorname{Tr}(\check{\gamma} \rfloor \Theta) = e_\alpha \rfloor T^\alpha, \quad \mathscr{A} := \frac{1}{4} {}^* \operatorname{Tr}(\gamma \wedge \Theta) = {}^*(\vartheta_\alpha \wedge T^\alpha).$	$T := \frac{1}{4} \operatorname{Tr}(\check{\gamma} \rfloor \Theta) = e_\alpha \rfloor T^\alpha, \quad \mathscr{A} := \frac{1}{4} {}^* \operatorname{Tr}(\gamma \wedge \Theta) = {}^*(\vartheta_\alpha \wedge T^\alpha).$
mielke-geometroynamics_254 EQ0799 (11.4.11)		■ 0.505	$ds^2 = \frac{1}{\lambda^2} \left(R_\circ^2 d\lambda^2 - \circ_{ij} dy^i \otimes_s dy^j \right).$	$ds^2 = \frac{1}{\lambda^2} \left(R_\circ^2 d\lambda^2 - \circ_{ij} dy^i \otimes_s dy^j \right).$	$ds^2 = \frac{1}{\lambda^2} \left(R_\circ^2 d\lambda^2 - \circ_{ij} dy^i \otimes_s dy^j \right).$
mielke-geometroynamics_309 EQ0960 (14.1.42)		■ 0.508	$\xi_{\mu\nu} \xi_{\rho\sigma} = -\frac{1}{2} \bar{R}_{\mu\nu\rho\sigma}^{\{\}} - \frac{1}{2} \frac{\bar{R}_{\mu\nu\gamma\delta}^{\{\}} \bar{R}_{\rho\sigma}^{\{\}\gamma\delta}}{R_{\alpha\beta}^{\{\}} R^{\{\}j\alpha\beta}}.$	$\xi_{\mu\nu} \xi_{\rho\sigma} = -\frac{1}{2} \bar{R}_{\mu\nu\rho\sigma}^{\{\}} - \frac{1}{2} \frac{\bar{R}_{\mu\nu\gamma\delta}^{\{\}} \bar{R}_{\rho\sigma}^{\{\}\gamma\delta}}{R_{\alpha\beta}^{\{\}} R^{\{\}j\alpha\beta}}.$	$\xi_{\mu\nu} \xi_{\rho\sigma} = -\frac{1}{2} \bar{R}_{\mu\nu\rho\sigma}^{\{\}} - \frac{1}{2} \frac{\bar{R}_{\mu\nu\gamma\delta}^{\{\}} \bar{R}_{\rho\sigma}^{\{\}\gamma\delta}}{R_{\alpha\beta}^{\{\}} R^{\{\}j\alpha\beta}}.$
mielke-geometroynamics_373 EQ1131 (B.43)		■ 0.509	$\langle \cdot, \cdot \rangle_M : \begin{cases} \bigwedge^p T^*(M) \times C_p(M, \mathbb{R}) \rightarrow \mathbb{R} \\ \langle \alpha^{(p)}, c_p \rangle_M \rightarrow \int_{c_p} \alpha^{(p)} \end{cases}$	$\langle \cdot, \cdot \rangle_M : \begin{cases} \bigwedge^p T^*(M) \times C_p(M, \mathbb{R}) \rightarrow \mathbb{R} \\ \langle \alpha^{(p)}, c_p \rangle_M \rightarrow \int_{c_p} \alpha^{(p)} \end{cases}$	$\langle \cdot, \cdot \rangle_M : \begin{cases} \bigwedge^p T^*(M) \times C_p(M, \mathbb{R}) \rightarrow \mathbb{R} \\ \langle \alpha^{(p)}, c_p \rangle_M \rightarrow \int_{c_p} \alpha^{(p)} \end{cases}$
mielke-geometroynamics_101 EQ0294 (4.5.5) [re-fined: census]		■ 0.510	$\Sigma_\alpha = e_\alpha \rfloor L - (e_\alpha \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_\alpha \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$	$\Sigma_\alpha = e_\alpha \rfloor L - (e_\alpha \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_\alpha \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$	$\Sigma_\alpha = e_\alpha \rfloor L - (e_\alpha \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_\alpha \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$
mielke-geometroynamics_59 EQ0118 (3.3.19)		■ 0.513	$L_{\mathrm{Pontr}} := -\frac{1}{2} F \wedge F = \mathbf{E} \cdot \mathbf{B} \eta.$	$L_{\mathrm{Pontr}} := -\frac{1}{2} F \wedge F = \mathbf{E} \cdot \mathbf{B} \eta.$	$L_{\mathrm{Pontr}} := -\frac{1}{2} F \wedge F = \mathbf{E} \cdot \mathbf{B} \eta.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_169 E00533 (7.7.53) [re-fined: census]		0.513	$\begin{aligned} G &:= \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} \gamma^\alpha \\ &= G^{(\beta)} + \frac{(-1)^{\text{sig}}}{12} \left(e_\alpha \rfloor \mathcal{A} \wedge {}^* \mathcal{A} - \frac{1}{3} \mathcal{A} \wedge e_\alpha \rfloor {}^* \mathcal{A} \right) \gamma^\alpha + \frac{(-1)^{\text{sig}}}{6} \gamma \wedge d\mathcal{A} \end{aligned}$	$G := \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} \gamma^\alpha$ $= G^{(\beta)} + \frac{(-1)^{\text{sig}}}{12} \left(e_\alpha \rfloor \mathcal{A} \wedge {}^* \mathcal{A} - \frac{1}{3} \mathcal{A} \wedge e_\alpha \rfloor {}^* \mathcal{A} \right) \gamma^\alpha + \frac{(-1)^{\text{sig}}}{6} \gamma \wedge d\mathcal{A}$	$G := \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} \gamma^\alpha \quad (7.7.5)$ $= G^{(\beta)} + \frac{(-1)^{\text{sig}}}{12} \left(e_\alpha \rfloor \mathcal{A} \wedge {}^* \mathcal{A} - \frac{1}{3} \mathcal{A} \wedge e_\alpha \rfloor {}^* \mathcal{A} \right) \gamma^\alpha + \frac{(-1)^{\text{sig}}}{6} \gamma \wedge d\mathcal{A}$
mielke-geometrodynamics_288 E00898 (13.5.3)		0.515	$\begin{aligned} \hat{B} &:= B_A^{BAA} B = \left(b_\alpha P_*^\alpha + \hat{b}^\alpha P_\alpha \right) \ell^2 + B_\alpha{}^\beta L^\alpha{}_\beta \\ \ell^2 &= \left(\frac{1}{2} \left(\hat{P}_\alpha \wedge \hat{P}_\beta \right) + \frac{1}{2} \left(\hat{P}_\alpha \wedge \hat{P}_\beta \right) \right) \end{aligned}$	$\hat{B} := B_A^{BAA} B = \left(b_\alpha P_*^\alpha + \hat{b}^\alpha P_\alpha \right) \ell^2 + B_\alpha{}^\beta L^\alpha{}_\beta$	$\hat{B} := B_A^B \mathcal{E}^A{}_B = (b_\alpha P_*^\alpha + \hat{b}^\alpha P_\alpha) \ell^2 + B_\alpha{}^\beta L^\alpha{}_\beta$
mielke-geometrodynamics_43 E00045 (2.6.8)		0.516	$(db)_{\text{hor}} = Db = \omega \otimes b$	$(db)_{\text{hor}} = Db = \omega \otimes b$	$(db)_{\text{hor}} = Db = \omega \otimes b$
mielke-geometrodynamics_285 E00876 (13.3.5)		0.518	$\begin{aligned} &\left[P_\alpha, P_\beta \right] = 0, \quad \left[P_\alpha, P_\beta \right] = 0, \\ &\left[L_\beta^\alpha, P_\gamma \right] = \delta_\gamma^\alpha P_\beta, \quad \left[L_\beta^\alpha, P_\gamma \right] = -\delta_\beta^\gamma P_\alpha^*. \end{aligned}$	$\begin{aligned} [P_\alpha, P_\beta] &= 0, [P_\alpha^*, P_\beta^*] = 0, \\ [L_\beta^\alpha, P_\gamma] &= \delta_\gamma^\alpha P_\beta, [L_\beta^\alpha, P_\gamma^*] = -\delta_\beta^\gamma P_\alpha^*. \end{aligned}$	$\begin{aligned} [P_\alpha, P_\beta] &= 0, \quad [P_\alpha^*, P_\beta^*] = 0, \\ [L_\beta^\alpha, P_\gamma] &= \delta_\gamma^\alpha P_\beta, \quad [L_\beta^\alpha, P_\gamma^*] = -\delta_\beta^\gamma P_\alpha^*. \end{aligned}$
mielke-geometrodynamics_320 E00990 (14.3.3)		0.526	$\begin{aligned} &\left(\overline{\hat{\psi}} \hat{\gamma} \wedge {}^* \hat{D}^{g+\circ} \hat{\psi} + \left(\overline{\hat{D}^{g+\circ} \hat{\psi}} \right) \wedge {}^* \hat{\gamma} \hat{\psi} \right) - \hat{m} \overline{\hat{\psi}} \hat{\psi} \eta \\ &+ \frac{1}{2} \overline{\hat{\psi}} [\hat{\alpha}, {}^* \hat{\gamma}] \hat{\psi}. \end{aligned}$	$\hat{L}_D = \frac{i}{2} \left(\overline{\hat{\psi}} \hat{\gamma} \wedge {}^* \hat{D}^{g+\circ} \hat{\psi} + \left(\overline{\hat{D}^{g+\circ} \hat{\psi}} \right) \wedge {}^* \hat{\gamma} \hat{\psi} \right) - \hat{m} \overline{\hat{\psi}} \hat{\psi} \eta$ $+ \frac{1}{2} \overline{\hat{\psi}} [\hat{\alpha}, {}^* \hat{\gamma}] \hat{\psi}.$	$\hat{L}_D = \frac{i}{2} (\overline{\hat{\psi}} \hat{\gamma} \wedge {}^* \hat{D}^{g+\circ} \hat{\psi} + (\overline{\hat{D}^{g+\circ} \hat{\psi}}) \wedge {}^* \hat{\gamma} \hat{\psi}) - \hat{m} \overline{\hat{\psi}} \hat{\psi} \eta$ $+ \frac{1}{2} \overline{\hat{\psi}} [\hat{\alpha}, {}^* \hat{\gamma}] \hat{\psi}.$
mielke-geometrodynamics_152 E00456 (7.2.8) [re-fined: census]		0.527	$\begin{aligned} \bar{R}_{..cd}^{ab} &= 2 R_{..[c}^{[a} \delta_{d]}^{b]}; \quad \mathcal{R}_{..b}^a := \mathcal{R}_{..b}^a - \frac{1}{4} \delta_b^a R, \\ &= \frac{1}{4} \delta_b^a R, \end{aligned}$	$\bar{R}_{..cd}^{ab} = 2 R_{..[c}^{[a} \delta_{d]}^{b]}; \quad \mathcal{R}_{..b}^a := \mathcal{R}_{..b}^a - \frac{1}{4} \delta_b^a R,$	$\bar{R}_{..cd}^{ab} = 2 \mathcal{R}_{..[c}^{[a} \delta_{d]}^{b]}; \quad \mathcal{R}_{..b}^a := \mathcal{R}_{..b}^a - \frac{1}{4} \delta_b^a R,$
mielke-geometrodynamics_125 E00360 (6.2.11)		0.531	$L_{\parallel} \equiv L_{\text{HE}} + \frac{1}{2\kappa} R_{\alpha\beta} \wedge {}^* (\vartheta^\alpha \wedge \vartheta^\beta) + \frac{2\ell^2}{\kappa} dC_{\text{TT}}^*,$	$L_{\parallel} \equiv L_{\text{HE}} + \frac{1}{2\kappa} R_{\alpha\beta} \wedge {}^* (\vartheta^\alpha \wedge \vartheta^\beta) + \frac{2\ell^2}{\kappa} dC_{\text{TT}}^*,$	$L_{\parallel} \equiv L_{\text{HE}} + \frac{1}{2\kappa} R_{\alpha\beta} \wedge {}^* (\vartheta^\alpha \wedge \vartheta^\beta) + \frac{2\ell^2}{\kappa} dC_{\text{TT}}^*,$
mielke-geometrodynamics_142 E00437 (6.8.8) [re-fined: census]		0.532	$E_\alpha := \frac{\partial L_g}{\partial \vartheta^\alpha} = e_\alpha \rfloor L_g + (e_\alpha \rfloor T^\beta) \wedge H_\beta + (e_\alpha \rfloor R^{\beta\gamma}) \wedge H_{\beta\gamma},$	$E_\alpha := \frac{\partial L_g}{\partial \vartheta^\alpha} = e_\alpha \rfloor L_g + (e_\alpha \rfloor T^\beta) \wedge H_\beta + (e_\alpha \rfloor R^{\beta\gamma}) \wedge H_{\beta\gamma},$	$E_\alpha := \frac{\partial L_g}{\partial \vartheta^\alpha} = e_\alpha \rfloor L_g + (e_\alpha \rfloor T^\beta) \wedge H_\beta + (e_\alpha \rfloor R^{\beta\gamma}) \wedge H_{\beta\gamma},$
mielke-geometrodynamics_82 E00198 (4.1.11)		0.533	$\begin{aligned} e_\alpha(m) &\rightarrow e_\alpha'(m) = g_\alpha{}^\beta e_\beta(m), \\ g &= [g_\alpha{}^\beta] \in GL(n, \mathbb{R}). \end{aligned}$	$e_\alpha(m) \rightarrow e_\alpha'(m) = g_\alpha{}^\beta e_\beta(m),$ $g = [g_\alpha{}^\beta] \in GL(n, \mathbb{R}).$	$e_\alpha(m) \rightarrow e_\alpha'(m) = g_\alpha{}^\beta e_\beta(m),$ $g = [g_\alpha{}^\beta] \in GL(n, \mathbb{R}).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_59 EQ0116 (3.3.17)	59	■ 0.585	$\mathbf{H}=\left[\begin{array}{l} \boldsymbol{\nabla} \times \mathbf{B}-\frac{\partial \mathbf{E}}{c^2 \partial t}=\mu \mathbf{J} \\ \boldsymbol{\nabla} \cdot \mathbf{E}=\rho / \varepsilon . \end{array}\right]$	$dH=j\left\{\begin{array}{l} \boldsymbol{\nabla} \times \mathbf{B}-\frac{\partial \mathbf{E}}{c^2 \partial t}=\mu \mathbf{J} \\ \boldsymbol{\nabla} \cdot \mathbf{E}=\rho / \varepsilon . \end{array}\right.$	$dH=j\left\{\begin{array}{l} \boldsymbol{\nabla} \times \mathbf{B}-\frac{\partial \mathbf{E}}{c^2 \partial t}=\mu \mathbf{J} \\ \boldsymbol{\nabla} \cdot \mathbf{E}=\rho / \varepsilon . \end{array}\right.$
mielke-geometrodynamics_129 EQ0381 (6.2.32) [re-fined: census]	129	■ 0.588	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha}:=D \stackrel{(\pm)}{\Pi}_{\alpha} \pm \frac{i}{2} \ell^2\left(e_{\alpha} \stackrel{(\mp)}{\Pi}\right) \wedge \stackrel{(\pm)}{\Pi}_{\beta},$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha}:=D \stackrel{(\pm)}{\Pi}_{\alpha} \pm \frac{i}{2} \ell^2\left(e_{\alpha} \stackrel{(\mp)}{\Pi}\right) \wedge \stackrel{(\pm)}{\Pi}_{\beta},$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha}:=D \stackrel{(\pm)}{\Pi}_{\alpha} \pm \frac{i}{2} \ell^2\left(e_{\alpha} \stackrel{(\mp)}{\Pi}^{\beta}\right) \wedge \stackrel{(\pm)}{\Pi}_{\beta},$
mielke-geometrodynamics_45 EQ0056 (2.7.6)	45	■ 0.592	$\phi=\varphi \otimes b=G^{-1} \varphi \otimes^G b \in C^{\infty}\left(V^{\rho}\right) .$	$\phi=\varphi \otimes b=G^{-1} \varphi \otimes^G b \in C^{\infty}\left(V^{\rho}\right) .$	$\phi=\varphi \otimes b=G^{-1} \varphi \otimes^G b \in C^{\infty}\left(V^{\rho}\right) .$
mielke-geometrodynamics_74 EQ0187 (3.7.17)	74	■ 0.593	$\begin{aligned} \left(D^* \Omega\right) &=\vec{D}^* \vec{\Omega}+\left[\omega_0, \Omega_0\right] \\ &=\vec{D}^* \vec{\Omega}+\left[\omega_0, \Theta^{\perp}\right]=0 . \end{aligned}$	$\begin{aligned} \left(D^* \Omega\right) &=\vec{D}^* \vec{\Omega}+\left[\omega_0, \Omega_0\right] \\ &=\vec{D}^* \vec{\Omega}+\left[\omega_0, \Theta^{\perp}\right]=0 . \end{aligned}$	$\begin{aligned} \left(D^* \Omega\right) &=\vec{D}^* \vec{\Omega}+\left[\omega_0, \Omega_0\right] \\ &=\vec{D}^* \vec{\Omega}+\left[\omega_0, \Theta^{\perp}\right]=0 . \end{aligned}$
mielke-geometrodynamics_186 EQ0581 (8.6.19)	186	■ 0.593	$\widehat{\Sigma}_{\alpha}:=\Sigma_{\alpha}-D^{\{\}} \mu_{\alpha},$	$\widehat{\Sigma}_{\alpha}:=\Sigma_{\alpha}-D^{\{\}} \mu_{\alpha},$	$\widehat{\Sigma}_{\alpha}:=\Sigma_{\alpha}-D^{\{\}} \mu_{\alpha},$
mielke-geometrodynamics_307 EQ0948 (14.1.29)	307	■ 0.593	$\begin{aligned} D^* \Omega &=\left(D^* \xi-(-1)^z \xi \wedge D \delta\right) \cos \delta \\ &-(-1)^z\left(D \xi+(-1)^z * \xi \wedge D \delta\right) \sin \delta=\tau_e \end{aligned}$	$\begin{aligned} D^* \Omega &=\left(D^* \xi-(-1)^z \xi \wedge D \delta\right) \cos \delta \\ &-(-1)^z\left(D \xi+(-1)^z * \xi \wedge D \delta\right) \sin \delta=\tau_e \end{aligned}$	$\begin{aligned} D^* \Omega &=\left(D^* \xi-(-1)^z \xi \wedge D \delta\right) \cos \delta \\ &-(-1)^z\left(D \xi+(-1)^z * \xi \wedge D \delta\right) \sin \delta=\tau_e \end{aligned}$
mielke-geometrodynamics_125 EQ0357 (6.2.8)	125	■ 0.600	$L_{\mathrm{HE}}=-\frac{1}{2 \kappa} R_{\alpha \beta}^{\{\}} \wedge^*\left(\vartheta^{\alpha} \wedge \vartheta^{\beta}\right)$	$L_{\mathrm{HE}}=-\frac{1}{2 \kappa} R_{\alpha \beta}^{\{\}} \wedge^*\left(\vartheta^{\alpha} \wedge \vartheta^{\beta}\right)$	$L_{\mathrm{HE}}=-\frac{1}{2 \kappa} R_{\alpha \beta}^{\{\}} \wedge^*\left(\vartheta^{\alpha} \wedge \vartheta^{\beta}\right)$
mielke-geometrodynamics_225 EQ0704 (10.6.3)	225	■ 0.601	$\square F-m_{\text {photon }}^2 F=0,$	$\square F-m_{\text {photon }}^2 F=0,$	$\square F-m_{\text {photon }}^2 F=0,$
mielke-geometrodynamics_303 EQ0927 (14.1.8)	303	■ 0.604	$\begin{aligned} \Sigma_{\mathrm{YM}} &=\frac{(-1)^s}{2 \alpha_g} \operatorname{Tr}\left\{\left(\Omega \wedge^*\left(\vartheta \wedge^* \Omega\right)+* \vartheta \wedge^*\left(\Omega \wedge^* \Omega\right)\right)\right. \\ &=\frac{(-1)^s}{4 \alpha_g} \operatorname{Tr}\left\{\Omega \wedge^*\left(\vartheta \wedge^* \Omega\right)-* \Omega \wedge^*\left(\vartheta \wedge \Omega\right)\right\} . \end{aligned}$	$\begin{aligned} \Sigma_{\mathrm{YM}} &=\frac{(-1)^s}{2 \alpha_g} \operatorname{Tr}\left\{\Omega \wedge^*\left(\vartheta \wedge^* \Omega\right)+* \vartheta \wedge^*\left(\Omega \wedge^* \Omega\right)\right\} \\ &=\frac{(-1)^s}{4 \alpha_g} \operatorname{Tr}\left\{\Omega \wedge^*\left(\vartheta \wedge^* \Omega\right)-* \Omega \wedge^*\left(\vartheta \wedge \Omega\right)\right\} . \end{aligned}$	$\begin{aligned} \Sigma_{\mathrm{YM}} &=\frac{(-1)^s}{2 \alpha_g} \operatorname{Tr}\left\{\Omega \wedge^*\left(\vartheta \wedge^* \Omega\right)+* \vartheta \wedge^*\left(\Omega \wedge^* \Omega\right)\right\} \\ &=\frac{(-1)^s}{4 \alpha_g} \operatorname{Tr}\left\{\Omega \wedge^*\left(\vartheta \wedge^* \Omega\right)-* \Omega \wedge^*\left(\vartheta \wedge \Omega\right)\right\} . \end{aligned}$
mielke-geometrodynamics_112 EQ0326 (5.3.3) [re-fined: census]	112	■ 0.613	$\widehat{D} G_{\alpha} \equiv \frac{1}{2}\left(e_{\alpha} \rfloor R^{\beta \gamma}\right) \wedge \eta_{\beta \gamma \mu} \wedge T^{\mu}$	$\widehat{D} G_{\alpha} \equiv \frac{1}{2}\left(e_{\alpha} \rfloor R^{\beta \gamma}\right) \wedge \eta_{\beta \gamma \mu} \wedge T^{\mu}$	$\widehat{D} G_{\alpha} \equiv \frac{1}{2}\left(e_{\alpha} \rfloor R^{\beta \gamma}\right) \wedge \eta_{\beta \gamma \mu} \wedge T^{\mu}$
mielke-geometrodynamics_242 EQ0758 (11.1.25)	242	■ 0.615	$\gamma=E_{\cdot j}^{\alpha}(m) \gamma_{\alpha} \otimes d x^j=\gamma_j d x^j \in C^{\infty}\left(T_{\mathbb{C}}^*(M) \times C V\right)$	$\gamma=E_{\cdot j}^{\alpha}(m) \gamma_{\alpha} \otimes d x^j=\gamma_j d x^j \in C^{\infty}\left(T_{\mathbb{C}}^*(M) \times C V\right)$	$\gamma=E_{\cdot j}^{\alpha}(m) \gamma_{\alpha} \otimes d x^j=\gamma_j d x^j \in C^{\infty}\left(T_{\mathbb{C}}^*(M) \times C V\right)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_308 EQ0953 (14.1.35)	308	■ 0.616	$\begin{aligned} & (-1)^z \operatorname{Tr} \left\{ \left(\vartheta \wedge \xi \wedge \xi - \vartheta \wedge \xi \wedge \xi \right) \wedge d\delta \right\} = \operatorname{Tr} \{ \xi \wedge \vartheta \wedge D^* \xi + \xi \wedge \vartheta \wedge D \xi \} \\ & = \frac{1}{2} D \operatorname{Tr} \{ (\xi \wedge (\vartheta \wedge \xi) - \xi \wedge (\vartheta \wedge \xi)) \} \\ & = (-1)^s 2\alpha_g D \Sigma_{\text{YM}} = \frac{\alpha_g}{\ell^{*2}} D (\vartheta \wedge \Omega^g) \equiv 2 \frac{\alpha_g}{\ell^{*2}} \Theta \wedge \Omega^g. \end{aligned}$	$\begin{aligned} & (-1)^z \operatorname{Tr} \{ \vartheta \wedge \xi \wedge \xi - \vartheta \wedge \xi \wedge \xi \} \wedge d\delta = \operatorname{Tr} \{ \xi \wedge \vartheta \wedge D^* \xi + \xi \wedge \vartheta \wedge D \xi \} \\ & = \frac{1}{2} D \operatorname{Tr} \{ (\xi \wedge (\vartheta \wedge \xi) - \xi \wedge (\vartheta \wedge \xi)) \} \\ & = (-1)^s 2\alpha_g D \Sigma_{\text{YM}} = \frac{\alpha_g}{\ell^{*2}} D (\vartheta \wedge \Omega^g) \equiv 2 \frac{\alpha_g}{\ell^{*2}} \Theta \wedge \Omega^g. \end{aligned}$	$\begin{aligned} & (-1)^z \operatorname{Tr} \{ \vartheta \wedge \xi \wedge \xi - \vartheta \wedge \xi \wedge \xi \} \wedge d\delta = \operatorname{Tr} \{ \xi \wedge \vartheta \wedge D^* \xi + \xi \wedge \vartheta \wedge D \xi \} \\ & = \frac{1}{2} D \operatorname{Tr} \{ (\xi \wedge (\vartheta \wedge \xi) - \xi \wedge (\vartheta \wedge \xi)) \} \\ & = (-1)^s 2\alpha_g D \Sigma_{\text{YM}} = \frac{\alpha_g}{\ell^{*2}} D (\vartheta \wedge \Omega^g) \equiv 2 \frac{\alpha_g}{\ell^{*2}} \Theta \wedge \Omega^g. \end{aligned}$
mielke-geometrodynamics_322 EQ0998 (14.3.11)	322	■ 0.619	$\begin{array}{l} \psi_{\text{L}} : \quad \gamma^5 = +1, \quad \overset{\circ}{\gamma}^{K+1} = +\widehat{\gamma}^{n+1} \\ \psi_{\text{R}} : \quad \gamma^5 = -1, \quad \overset{\circ}{\gamma}^{K+1} = -\widehat{\gamma}^{n+1} \end{array}$	$\begin{array}{l} \psi_{\text{L}} : \quad \gamma^5 = +1, \quad \overset{\circ}{\gamma}^{K+1} = +\widehat{\gamma}^{n+1} \\ \psi_{\text{R}} : \quad \gamma^5 = -1, \quad \overset{\circ}{\gamma}^{K+1} = -\widehat{\gamma}^{n+1} \end{array}$	$\begin{array}{l} \psi_{\text{L}} : \quad \gamma^5 = +1, \quad \overset{\circ}{\gamma}^{K+1} = +\widehat{\gamma}^{n+1} \\ \psi_{\text{R}} : \quad \gamma^5 = -1, \quad \overset{\circ}{\gamma}^{K+1} = -\widehat{\gamma}^{n+1} \end{array}$
mielke-geometrodynamics_167 EQ0525 (7.7.44)	167	■ 0.623	$\begin{aligned} & C_{\text{TT}} := \frac{1}{8\ell^2} \operatorname{Tr} (\gamma \wedge \Theta) = \frac{1}{2\ell^2} \vartheta^\alpha \wedge T_\alpha = -\frac{(-1)^{\text{sig}}}{2\ell^2} * \mathcal{A}, \\ & C_{\text{RR}} := \operatorname{Tr} \left(\Gamma \wedge \Omega^g - \frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma \right), \end{aligned}$	$\begin{aligned} & C_{\text{TT}} := \frac{1}{8\ell^2} \operatorname{Tr} (\gamma \wedge \Theta) = \frac{1}{2\ell^2} \vartheta^\alpha \wedge T_\alpha = -\frac{(-1)^{\text{sig}}}{2\ell^2} * \mathcal{A}, \\ & C_{\text{RR}} := \operatorname{Tr} \left(\Gamma \wedge \Omega^g - \frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma \right), \end{aligned}$	$\begin{aligned} & C_{\text{TT}} := \frac{1}{8\ell^2} \operatorname{Tr} (\gamma \wedge \Theta) = \frac{1}{2\ell^2} \vartheta^\alpha \wedge T_\alpha = -\frac{(-1)^{\text{sig}}}{2\ell^2} * \mathcal{A}, \\ & C_{\text{RR}} := \operatorname{Tr} \left(\Gamma \wedge \Omega^g - \frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma \right), \end{aligned}$
mielke-geometrodynamics_44 EQ0050 (2.6.13)	44	■ 0.626	$\begin{aligned} & D D D b = & D (\Omega \otimes b) = (d\Omega + [\Omega, \omega]) \otimes b \\ & = (d\Omega + \Omega \wedge \omega - \omega \wedge \Omega) \otimes b \\ & = (dd\omega - d\omega \wedge \omega + \omega \wedge d\omega \\ & \quad + d\omega \wedge \omega - \omega \wedge \omega \wedge \omega \\ & \quad + \omega \wedge \omega \wedge \omega - \omega \wedge d\omega) \otimes b \equiv 0. \end{aligned}$	$\begin{aligned} & D D D b = D (\Omega \otimes b) = (d\Omega + [\Omega, \omega]) \otimes b \\ & = (d\Omega + \Omega \wedge \omega - \omega \wedge \Omega) \otimes b \\ & = (dd\omega - d\omega \wedge \omega + \omega \wedge d\omega \\ & \quad + d\omega \wedge \omega - \omega \wedge \omega \wedge \omega \\ & \quad + \omega \wedge \omega \wedge \omega - \omega \wedge d\omega) \otimes b \equiv 0. \end{aligned}$	$\begin{aligned} & D D D b = D (\Omega \otimes b) = (d\Omega + [\Omega, \omega]) \otimes b \\ & = (d\Omega + \Omega \wedge \omega - \omega \wedge \Omega) \otimes b \\ & = (dd\omega - d\omega \wedge \omega + \omega \wedge d\omega \\ & \quad + d\omega \wedge \omega - \omega \wedge \omega \wedge \omega \\ & \quad + \omega \wedge \omega \wedge \omega - \omega \wedge d\omega) \otimes b \equiv 0. \end{aligned}$
mielke-geometrodynamics_59 EQ0115 (3.3.16)	59	■ 0.630	$dF \equiv 0 \quad \Leftrightarrow \quad \left\{ \begin{array}{l} \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} \equiv 0 \\ \nabla \cdot \mathbf{B} \equiv 0. \end{array} \right.$	$dF \equiv 0 \quad \left\{ \begin{array}{l} \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} \equiv 0 \\ \nabla \cdot \mathbf{B} \equiv 0 \end{array} \right.$	$dF \equiv 0 \quad \left\{ \begin{array}{l} \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} \equiv 0 \\ \nabla \cdot \mathbf{B} \equiv 0. \end{array} \right.$
mielke-geometrodynamics_70 EQ0171 (3.6.6)	70	■ 0.630	$i^* \gamma \wedge D_\pm \psi = 0 \quad \text{and} \quad F^\pm = \frac{1}{2} \bar{\psi} \sigma_\pm \psi = m \mathcal{P}_\pm,$	$i^* \gamma \wedge D_\pm \psi = 0 \quad \text{and} \quad F^\pm = \frac{1}{2} \bar{\psi} \sigma_\pm \psi = m \mathcal{P}_\pm,$	$i^* \gamma \wedge D_\pm \psi = 0 \quad \text{and} \quad F^\pm = \frac{1}{2} \bar{\psi} \sigma_\pm \psi = m \mathcal{P}_\pm,$
mielke-geometrodynamics_74 EQ0189 (3.7.19)	74	■ 0.631	$\{ \}^* \Omega = (\stackrel{+}{-}) \Omega.$	$^* \Omega = (\stackrel{+}{-}) \Omega.$	$^* \Omega = (\stackrel{+}{-}) \Omega.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_ EQ0788 (11.3.7)	249	■ 0.635	$\left\{i\gamma^{\kappa}\nabla_{\kappa}^{()}-\frac{6}{n^2}\ell^{*2}\gamma^{n+1}\gamma^{\mu}\bar{\psi}\gamma^{n+1}\gamma_{\mu}\psi-m\right\}\psi=0.$	$\left\{i\gamma^{\kappa}\nabla_{\kappa}^{()} - \frac{6}{n^2}\ell^{*2}\gamma^{n+1}\gamma^{\mu}\bar{\psi}\gamma^{n+1}\gamma_{\mu}\psi - m\right\}\psi = 0.$	$\left\{i\gamma^{\kappa}\nabla_{\kappa}^{()} - \frac{6}{n^2}\ell^{*2}\gamma^{n+1}\gamma^{\mu}\bar{\psi}\gamma^{n+1}\gamma_{\mu}\psi - m\right\}\psi = 0.$
mielke-geometrodynamics_ EQ0164 (3.5.12)	68	■ 0.636	$L_{\mathrm{YM}}+L_{\infty}=(\stackrel{+}{-})\frac{1}{2\alpha_g}\mathrm{Tr}\left(*\Omega\stackrel{-}{+}\Omega\right)^2\geq 0$	$L_{\mathrm{YM}}+L_{\infty}=(\stackrel{+}{-})\frac{1}{2\alpha_g}\mathrm{Tr}\left(*\Omega\stackrel{-}{+}\Omega\right)^2\geq 0$	$L_{\mathrm{YM}}+L_{\infty}=(\stackrel{+}{-})\frac{1}{2\alpha_g}\mathrm{Tr}(*\Omega\stackrel{-}{+}\Omega)^2\geq 0$
mielke-geometrodynamics_ EQ0549 (8.4.3)	179	■ 0.637	$\begin{aligned} L_{\mathrm{FP}} & \text{ \& } := -s \operatorname{Tr} \left\{ \bar{\chi} \wedge * \Omega^g + \bar{\Phi} D^* \Psi + \frac{1}{2} \rho \bar{\chi} \wedge * \beta + \bar{C} d^* \Gamma + \frac{1}{2} \bar{C} \wedge * \mathbb{B} \right\} \\ & = \frac{1}{2} s \left[\bar{\chi}^{\alpha\beta} \wedge * R_{\alpha\beta}^{(\pm)} + \bar{\Phi}_{\alpha\beta} D^* \Psi^{\alpha\beta} + \frac{1}{2} \rho \bar{\chi}^{\alpha\beta} \wedge * b_{\alpha\beta} + \tilde{\Psi}_{\mathrm{F}} \right] \end{aligned}$	$L_{\mathrm{FP}} := -s \operatorname{Tr} \left\{ \bar{\chi} \wedge * \Omega^g + \bar{\Phi} D^* \Psi + \frac{1}{2} \rho \bar{\chi} \wedge * \beta + \bar{C} d^* \Gamma + \frac{1}{2} \bar{C} \wedge * \mathbb{B} \right\} \\ = \frac{1}{2} s \left[\bar{\chi}^{\alpha\beta} \wedge * R_{\alpha\beta}^{(\pm)} + \bar{\Phi}_{\alpha\beta} D^* \Psi^{\alpha\beta} + \frac{1}{2} \rho \bar{\chi}^{\alpha\beta} \wedge * b_{\alpha\beta} + \tilde{\Psi}_{\mathrm{F}} \right]$	$L_{\mathrm{FP}} := -s \operatorname{Tr} \left\{ \bar{\chi} \wedge * \overset{\pm}{\Omega}^g + \bar{\Phi} D^* \Psi + \frac{1}{2} \rho \bar{\chi} \wedge * \beta + \bar{C} d^* \Gamma + \frac{1}{2} \bar{C} \wedge * \mathbb{B} \right\} \\ = \frac{1}{2} s \left[\bar{\chi}^{\alpha\beta} \wedge * R_{\alpha\beta}^{(\pm)} + \bar{\Phi}_{\alpha\beta} D^* \Psi^{\alpha\beta} + \frac{1}{2} \rho \bar{\chi}^{\alpha\beta} \wedge * b_{\alpha\beta} + \tilde{\Psi}_{\mathrm{F}} \right]. \quad (8)$
mielke-geometrodynamics_ EQ1058 (15.2.15)	346	■ 0.638	$\bar{D}^{\{\}} := d + i \bar{\omega}^{\{\}}$	$\bar{D}^{\{\}} := d + i \bar{\omega}^{\{\}}$	$\bar{D}^{\{\}} := d + i \bar{\omega}^{\{\}}$
mielke-geometrodynamics_ EQ1008 (14.4.3)	325	■ 0.640	$\begin{aligned} ds^2 & = -\frac{\Delta}{\rho^2} \left[c dt - J_3 \frac{\hbar}{M} \sin^2 \vartheta d\phi \right]^2 + \frac{\rho^2}{\Delta} dr + \rho^2 d\vartheta^2 \\ & + \frac{\sin^2 \vartheta}{\rho^2} \left\{ \left[r^2 + J(J+1) \frac{\hbar^2}{M^2 c^2} \right] d\phi - J_3 \frac{\hbar}{M} dt \right\}^2 \end{aligned}$	$ds^2 = -\frac{\Delta}{\rho^2} \left[cdt - J_3 \frac{\hbar}{M} \sin^2 \vartheta d\phi \right]^2 + \frac{\rho^2}{\Delta} dr + \rho^2 d\vartheta^2 \\ + \frac{\sin^2 \vartheta}{\rho^2} \left\{ \left[r^2 + J(J+1) \frac{\hbar^2}{M^2 c^2} \right] d\phi - J_3 \frac{\hbar}{M} dt \right\}^2$	$ds^2 = -\frac{\Delta}{\rho^2} \left[cdt - J_3 \frac{\hbar}{M} \sin^2 \vartheta d\phi \right]^2 + \frac{\rho^2}{\Delta} dr + \rho^2 d\vartheta^2 \\ + \frac{\sin^2 \vartheta}{\rho^2} \left\{ \left[r^2 + J(J+1) \frac{\hbar^2}{M^2 c^2} \right] d\phi - J_3 \frac{\hbar}{M} dt \right\}^2.$
mielke-geometrodynamics_ EQ0255 (4.3.20)	93	■ 0.642	$\stackrel{*}{\sigma}\Omega^g=\frac{1}{4}R_{\alpha\beta}^{\cdot cd}L_{cd}\otimes\vartheta^\alpha\wedge\vartheta^\beta.$	$\stackrel{*}{\sigma}\Omega^g=\frac{1}{4}R_{\alpha\beta}^{\cdot cd}L_{cd}\otimes\vartheta^\alpha\wedge\vartheta^\beta.$	$\stackrel{*}{\sigma}\Omega^g=\frac{1}{4}R_{\alpha\beta}^{\cdot cd}L_{cd}\otimes\vartheta^\alpha\wedge\vartheta^\beta.$
mielke-geometrodynamics_ EQ1015 (14.4.10)	327	■ 0.646	$A=\{ \ }^*\}\omega=A^{\scriptscriptstyle{(\circ)}}\lambda_3.$	$A = *\omega = A^{(\circ)}\lambda_3.$	$A = *\omega = A^{(\circ)}\lambda_3.$
mielke-geometrodynamics_ EQ0119 (3.3.20) [re-fined: census]	59	■ 0.646	$\begin{aligned} \Sigma_{\alpha} & \text{ \& } := e_{\alpha} \rfloor L - (e_{\alpha} \rfloor dA) \wedge \frac{\partial L}{\partial (dA)} \\ & = \frac{1}{2} [(e_{\alpha} \rfloor F) \wedge H - F \wedge (e_{\alpha} \rfloor H)] \end{aligned}$	$\Sigma_{\alpha} := e_{\alpha} \rfloor L - (e_{\alpha} \rfloor dA) \wedge \frac{\partial L}{\partial (dA)} \\ = \frac{1}{2} [(e_{\alpha} \rfloor F) \wedge H - F \wedge (e_{\alpha} \rfloor H)]$	$\Sigma_{\alpha} := e_{\alpha} \rfloor L - (e_{\alpha} \rfloor dA) \wedge \frac{\partial L}{\partial (dA)} \\ = \frac{1}{2} [(e_{\alpha} \rfloor F) \wedge H - F \wedge (e_{\alpha} \rfloor H)]$
mielke-geometrodynamics_ EQ0120 (3.3.21) [re-fined: census]	60	■ 0.651	$d\Sigma_{\alpha}\simeq(e_{\alpha}\rfloor F)\wedge j$	$d\Sigma_{\alpha}\simeq(e_{\alpha}\rfloor F)\wedge j$	$d\Sigma_{\alpha}\simeq(e_{\alpha}\rfloor F)\wedge j$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00497 (7.7.15)	163	■ 0.653	$ds^2 = o_{\alpha\beta} \vartheta^\alpha \otimes \vartheta^\beta = g_{ij} dx^i \otimes dx^j, \quad o_{\alpha\beta} := \text{diag}\{1, \underbrace{-1, \dots, -1}_{\text{sig}}\},$	$ds^2 = o_{\alpha\beta} \vartheta^\alpha \otimes \vartheta^\beta = g_{ij} dx^i \otimes dx^j, \quad o_{\alpha\beta} := \text{diag}\{1, \underbrace{-1, \dots, -1}_{\text{sig}}\},$	$ds^2 = o_{\alpha\beta} \vartheta^\alpha \otimes \vartheta^\beta = g_{ij} dx^i \otimes dx^j, \quad o_{\alpha\beta} := \text{diag}\{1, \underbrace{-1, \dots, -1}_{\text{sig}}\},$
mielke-geometrodynamics_E00800 (11.4.12)	254	■ 0.658	$\left(i \gamma^k \partial_k - \frac{3i}{2R_o \lambda} \gamma^0 - \frac{m}{\lambda} \right) \psi = 0,$	$\left\{ i \gamma^k \partial_k - \frac{3i}{2R_o \lambda} \gamma^0 - \frac{m}{\lambda} \right\} \psi = 0,$	$\left\{ i \gamma^k \partial_k - \frac{3i}{2R_o \lambda} \gamma^0 - \frac{m}{\lambda} \right\} \psi = 0,$
mielke-geometrodynamics_E00316 (4.6.8)	105	■ 0.664	$\Phi(\widetilde{\Gamma}, \gamma) = (2\pi \hbar \mathcal{M}_n G / \hbar \ell c^2) = 2\pi n$	$\Phi(\widetilde{\Gamma}, \gamma) = (2\pi \hbar \mathcal{M}_n G / \hbar \ell c^2) = 2\pi n$	$\Phi(\widetilde{\Gamma}, \gamma) = (2\pi \hbar \mathcal{M}_n G / \hbar \ell c^2) = 2\pi n,$
mielke-geometrodynamics_E00335 (5.3.12)	113	■ 0.665	$\begin{aligned} & \widetilde{T}_{\mu\nu} = T_{\mu\nu} + \ell^{*2} \left\{ -4 \tau_{\mu \cdot [\lambda]}^{\cdot \kappa} \tau_{\nu \cdot [\kappa]}^{\cdot \lambda} - 2 \tau_{\mu}^{\cdot \kappa \lambda} \tau_{\nu \kappa \lambda} + \tau_{\cdot, \mu}^{\kappa \lambda} \tau_{\kappa \lambda \nu} \right. \\ & \quad \left. + \frac{1}{2} g_{\mu\nu} \left(4 \tau_{\delta \cdot [\lambda]}^{\cdot \kappa} \tau_{\cdot, \cdot \kappa]}^{\delta \lambda} + \tau^{\delta \kappa \lambda} \tau_{\delta \kappa \lambda} \right) \right\}. \end{aligned}$	$\widetilde{T}_{\mu\nu} = T_{\mu\nu} + \ell^{*2} \left\{ -4 \tau_{\mu \cdot [\lambda]}^{\cdot \kappa} \tau_{\nu \cdot [\kappa]}^{\cdot \lambda} - 2 \tau_{\mu}^{\cdot \kappa \lambda} \tau_{\nu \kappa \lambda} + \tau_{\cdot, \mu}^{\kappa \lambda} \tau_{\kappa \lambda \nu} \right. \\ \quad \left. + \frac{1}{2} g_{\mu\nu} \left(4 \tau_{\delta \cdot [\lambda]}^{\cdot \kappa} \tau_{\cdot, \cdot \kappa]}^{\delta \lambda} + \tau^{\delta \kappa \lambda} \tau_{\delta \kappa \lambda} \right) \right\}.$	$\widetilde{T}_{\mu\nu} = T_{\mu\nu} + \ell^{*2} \left\{ -4 \tau_{\mu \cdot [\lambda]}^{\cdot \kappa} \tau_{\nu \cdot [\kappa]}^{\cdot \lambda} - 2 \tau_{\mu}^{\cdot \kappa \lambda} \tau_{\nu \kappa \lambda} + \tau_{\cdot, \mu}^{\kappa \lambda} \tau_{\kappa \lambda \nu} \right. \\ \quad \left. + \frac{1}{2} g_{\mu\nu} (4 \tau_{\delta \cdot [\lambda]}^{\cdot \kappa} \tau_{\cdot, \cdot \kappa]}^{\delta \lambda} + \tau^{\delta \kappa \lambda} \tau_{\delta \kappa \lambda}) \right\}.$
mielke-geometrodynamics_E01055 (15.2.12)	345	■ 0.665	$\bar{L}_{\mathbf{w}}(f) = \frac{i}{8} \text{Tr}(\bar{\gamma} \wedge \bar{\gamma} \wedge {}^* \bar{\Omega}).$	$\bar{L}_{\mathbf{w}}(f) = \frac{i}{8} \text{Tr}(\bar{\gamma} \wedge \bar{\gamma} \wedge {}^* \bar{\Omega}).$	$\bar{L}_{\mathbf{w}}(f) = \frac{i}{8} \text{Tr}(\bar{\gamma} \wedge \bar{\gamma} \wedge {}^* \bar{\Omega}).$
mielke-geometrodynamics_E01145 (C.13)	377	■ 0.667	$\stackrel{\circ}{g}_{lm} := c_{lk}{}^i c_{mi}{}^k$	$\stackrel{\circ}{g}_{lm} := c_{lk}{}^i c_{mi}{}^k$	$\stackrel{\circ}{g}_{lm} := c_{lk}{}^i c_{mi}{}^k$
mielke-geometrodynamics_E00450 (7.2.3)	151	■ 0.676	$\Omega^g = \stackrel{+}{\Omega}^g + \bar{\Omega}^g$	$\Omega^g = \stackrel{+}{\Omega}^g + \bar{\Omega}^g$	$\Omega^g = \stackrel{+}{\Omega}^g + \bar{\Omega}^g$
mielke-geometrodynamics_E00143 (3.4.14)	64	■ 0.679	$\operatorname{div} \iota := (-1)^{1+s} d^* \iota = 0$	$\operatorname{div} \iota := (-1)^{1+s} d^* \iota = 0$	$\operatorname{div} \iota := (-1)^{1+s} d^* \iota = 0$
mielke-geometrodynamics_E00472 (7.3.11) [re-fined: census]	155	■ 0.682	$\stackrel{+}{\Omega} = 0 \iff R_{ij}^{\cancel{\{}} = 0 \iff R_{ij}^{\{}} = \Lambda' g_{ij}.$	$\stackrel{+}{\Omega} = 0 \iff R_{ij}^{\cancel{\{}} = 0 \iff R_{ij}^{\{}} = \Lambda' g_{ij}.$	$\stackrel{+}{\Omega}^{\{]} = 0 \iff \mathcal{R}_{ij}^{\{]} = 0 \iff R_{ij}^{\{]} = \Lambda' g_{ij}.$
mielke-geometrodynamics_E00812 (11.4.24)	256	■ 0.686	$\begin{aligned} & \partial_{\rho^*} H + \frac{1+\kappa}{\rho} e^{(\nu-\mu)/2} H = \frac{1}{\beta} \left[\omega + e^{\nu/2} - \varepsilon e^{-\mu} \left(H^2 - F^2 - \frac{2}{3} a H^2 \right) \right] F, \\ & \partial_{\rho^*} F + \frac{1-\kappa}{\rho} e^{(\nu-\mu)/2} F = \frac{1}{\beta} \left[-\omega + e^{\nu/2} - \varepsilon e^{-\mu} \left(H^2 - F^2 + \frac{2}{3} a F^2 \right) \right] H. \end{aligned}$	$\begin{aligned} & \partial_{\rho^*} H + \frac{1+\kappa}{\rho} e^{(\nu-\mu)/2} H = \frac{1}{\beta} \left[\omega + e^{\nu/2} - \varepsilon e^{-\mu} \left(H^2 - F^2 - \frac{2}{3} a H^2 \right) \right] F, \\ & \partial_{\rho^*} F + \frac{1-\kappa}{\rho} e^{(\nu-\mu)/2} F = \frac{1}{\beta} \left[-\omega + e^{\nu/2} - \varepsilon e^{-\mu} \left(H^2 - F^2 + \frac{2}{3} a F^2 \right) \right] H. \end{aligned}$	$\begin{aligned} & \partial_{\rho^*} H + \frac{1+\kappa}{\rho} e^{(\nu-\mu)/2} H = \frac{1}{\beta} \left[\omega + e^{\nu/2} - \varepsilon e^{-\mu} \left(H^2 - F^2 - \frac{2}{3} a H^2 \right) \right] F, \\ & \partial_{\rho^*} F + \frac{1-\kappa}{\rho} e^{(\nu-\mu)/2} F = \frac{1}{\beta} \left[-\omega + e^{\nu/2} - \varepsilon e^{-\mu} \left(H^2 - F^2 + \frac{2}{3} a F^2 \right) \right] H. \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_111 EQ0321 (5.2.5)		■ 0.686	$\tau_{\alpha\beta}:=\frac{\partial L_{\mathrm{D}}}{\partial \Gamma^{\alpha\beta}}=\frac{1}{8}\bar{\Psi}\left(\gamma^{\sigma}\sigma_{\alpha\beta}+\sigma_{\alpha\beta}\gamma^{\sigma}\right)\Psi=\frac{1}{4}\eta_{\alpha\beta\gamma\delta}\bar{\Psi}\gamma^{\delta}\gamma_5\Psi\eta^{\gamma}=\tau_{\alpha\beta\gamma}\eta^{\gamma}$	$\tau_{\alpha\beta}:=\frac{\partial L_{\mathrm{D}}}{\partial \Gamma^{\alpha\beta}}=\frac{1}{8}\bar{\Psi}\left(\gamma^{\sigma}\sigma_{\alpha\beta}+\sigma_{\alpha\beta}\gamma^{\sigma}\right)\Psi=\frac{1}{4}\eta_{\alpha\beta\gamma\delta}\bar{\Psi}\gamma^{\delta}\gamma_5\Psi\eta^{\gamma}=\tau_{\alpha\beta\gamma}\eta^{\gamma}$	$\tau_{\alpha\beta}:=\frac{\partial L_{\mathrm{D}}}{\partial \Gamma^{\alpha\beta}}=\frac{1}{8}\bar{\Psi}\left(\gamma^{\sigma}\sigma_{\alpha\beta}+\sigma_{\alpha\beta}\gamma^{\sigma}\right)\Psi=\frac{1}{4}\eta_{\alpha\beta\gamma\delta}\bar{\Psi}\gamma^{\delta}\gamma_5\Psi\eta^{\gamma}=\tau_{\alpha\beta\gamma}\eta^{\gamma}$
mielke-geometrodynamics_131 EQ0389 (6.4.4)		■ 0.688	$\mathrm{H}=\mathrm{K}-\dot{\Psi}\wedge\left(\frac{\partial\mathrm{K}}{\partial\dot{\Psi}}\right),$	$\mathcal{H}=\mathcal{K}-\dot{\underline{\Psi}}\wedge\left(\frac{\partial\mathcal{K}}{\partial\dot{\underline{\Psi}}}\right),$	$\mathcal{H}=\mathcal{K}-\dot{\underline{\Psi}}\wedge\left(\frac{\partial\mathcal{K}}{\partial\dot{\underline{\Psi}}}\right),$
mielke-geometrodynamics_307 EQ0949 (14.1.30)		■ 0.688	$\begin{aligned} D\,\Omega\,\&=\left(D\,\xi+(-1)^z*\xi\wedge D\delta\right)\cos\delta\\ &+(-1)^z\left(D^*\xi-(-1)^z\xi\wedge D\delta\right)\sin\delta=0. \end{aligned}$	$D\Omega=(D\xi+(-1)^z*\xi\wedge D\delta)\cos\delta+(-1)^z(D^*\xi-(-1)^z\xi\wedge D\delta)\sin\delta=0.$	$D\Omega=(D\xi+(-1)^z*\xi\wedge D\delta)\cos\delta+(-1)^z(D^*\xi-(-1)^z\xi\wedge D\delta)\sin\delta=0.$
mielke-geometrodynamics_224 EQ0696 (10.5.7) [re-fined: census]		■ 0.694	$D^*\gamma=\gamma\wedge T,\quad T:=e_\alpha\rfloor T^\alpha,$	$D^*\gamma=\gamma\wedge T,\quad T:=e_\alpha\rfloor T^\alpha,$	$D^*\gamma=\gamma\wedge T,\quad T:=e_\alpha\rfloor T^\alpha,$
mielke-geometrodynamics_175 EQ0534 (8.2.1)		■ 0.699	$\begin{aligned} s\,\Gamma\,\&=\Psi-D\,C\,\,\,\,s\,C\,\&=\Phi-\frac{1}{2}[C,C]\,\,\,s\,\Omega^g\,\&=-D\Psi-[C,\Omega^g]\,\,\,s\Psi\,\&=-D\Phi-[C,\Psi]\,\,\,s\Phi\,\&=-[C,\Phi] \end{aligned}$	$\begin{aligned} s\Gamma &= \Psi - DC \\ sC &= \Phi - \frac{1}{2}[C,C] \\ s\Omega^g &= -D\Psi - [C,\Omega^g] \\ s\Psi &= -D\Phi - [C,\Psi] \\ s\Phi &= -[C,\Phi] \end{aligned}$	$\begin{aligned} s\Gamma &= \Psi - DC, \\ sC &= \Phi - \frac{1}{2}[C,C], \\ s\Omega^g &= -D\Psi - [C,\Omega^g], \\ s\Psi &= -D\Phi - [C,\Psi], \\ s\Phi &= -[C,\Phi]. \end{aligned}$
mielke-geometrodynamics_112 EQ0329 (5.3.6)		■ 0.699	$D\,\Theta\,\equiv\left[\Omega^g,\gamma\right],\quad D\Omega^g\equiv 0,$	$D\Theta\equiv[\Omega^g,\gamma],\quad D\Omega^g\equiv 0,$	$D\Theta\equiv[\Omega^g,\gamma],\quad D\Omega^g\equiv 0,$
mielke-geometrodynamics_155 EQ0471 (7.3.10)		■ 0.706	$\bar{\Omega}^{\{\}}=0\Longleftrightarrow\Omega_{\mathrm{C}}^{\{\}}=0\text{ and }R^{\{\}}=0.$	$\bar{\Omega}^{\{\}}=0\Longleftrightarrow\Omega_{\mathrm{C}}^{\{\}}=0\text{ and }R^{\{\}}=0.$	$\overline{\Omega}^{\{\}}=0\Longleftrightarrow\Omega_{\mathrm{C}}^{\{\}}=0\text{ and }R^{\{\}}=0.$
mielke-geometrodynamics_240 EQ0747 (11.1.14)		■ 0.706	$M=\mathbb{R}\times S^3\#^vW^3.$	$M=\mathbb{R}\times S^3\#^vW^3.$	$M=\mathbb{R}\times S^3\#^vW^3.$
mielke-geometrodynamics_105 EQ0314 (4.6.6)		■ 0.710	$\Phi(\stackrel{\sim}{\Gamma},\gamma)=P\exp\left[(i/\hbar)\oint\left(\Gamma^{(T)\alpha}P_\alpha+\Gamma_\alpha^{(L)\beta}L^\alpha{}_\beta\right)\right]$	$\Phi(\tilde{\Gamma},\gamma)=P\exp\left[(i/\hbar)\oint\left(\Gamma^{(T)\alpha}P_\alpha+\Gamma_\alpha^{(L)\beta}L^\alpha{}_\beta\right)\right]$	$\Phi(\tilde{\Gamma},\gamma)=P\exp\left[(i/\hbar)\oint\left(\Gamma^{(T)\alpha}P_\alpha+\Gamma_\alpha^{(L)\beta}L^\alpha{}_\beta\right)\right]$
mielke-geometrodynamics_58 EQ0108 (3.3.9)		■ 0.711	$L_{\mathrm{Pontr}}=-\frac{1}{2}F\wedge F=-\frac{1}{2}dC,$	$L_{\mathrm{Pontr}}=-\frac{1}{2}F\wedge F=-\frac{1}{2}dC,$	$L_{\mathrm{Pontr}}=-\frac{1}{2}F\wedge F=-\frac{1}{2}dC,$
mielke-geometrodynamics_177 EQ0545 (8.3.6)		■ 0.711	$d\rightarrow\exp(\zeta\rfloor)d\exp(-\zeta\rfloor).$	$d\rightarrow\exp(\zeta\rfloor)d\exp(-\zeta\rfloor).$	$d\rightarrow\exp(\zeta\rfloor)d\exp(-\zeta\rfloor).$
mielke-geometrodynamics_192 EQ0583 (9.1.1)		■ 0.712	$ds^2=-e^vdt^2+e^\lambda dr^2+r^2d\Omega^{n-2}$	$ds^2=-e^vdt^2+e^\lambda dr^2+r^2d\Omega^{n-2}$	$ds^2=-e^vdt^2+e^\lambda dr^2+r^2d\Omega^{n-2}$
mielke-geometrodynamics_94 EQ0260 (4.3.25)		■ 0.718	$\stackrel{*}{K}=\frac{1}{2}K_\alpha^{\beta c}L_{\beta c}\otimes\vartheta^\alpha.$	$\overset{*}{\sigma}K=\frac{1}{2}K_\alpha^{\beta c}L_{\beta c}\otimes\vartheta^\alpha.$	$\overset{*}{\sigma}K=\frac{1}{2}K_\alpha^{\beta c}L_{\beta c}\otimes\vartheta^\alpha.$
mielke-geometrodynamics_285 EQ0878 (13.3.7)		■ 0.722	$\mathrm{SL}(n+1,\mathbb{R})\approx\mathrm{A}_{*}(n,\mathbb{R}):=\mathbb{R}^n\otimes\mathrm{GL}(n,\mathbb{R})\otimes\mathbb{R}_{*}^n.$	$\mathrm{SL}(n+1,\mathbb{R})\approx\mathrm{A}_{*}(n,\mathbb{R}):=\mathbb{R}^n\otimes\mathrm{GL}(n,\mathbb{R})\otimes\mathbb{R}_{*}^n.$	$\mathrm{SL}(n+1,\mathbb{R})\approx\mathrm{A}_{*}(n,\mathbb{R}):=\mathbb{R}^n\otimes\mathrm{GL}(n,\mathbb{R})\otimes\mathbb{R}_{*}^n.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00041 (2.6.4)	42	■ 0.725	$\stackrel{*}{\sigma} \Omega = \frac{1}{2} F_{\alpha} \beta^j L_j \otimes \vartheta^{\alpha} \wedge \vartheta^{\beta}$	$\sigma^* \Omega = \frac{1}{2} F_{\alpha} \beta^j L_j \otimes \vartheta^{\alpha} \wedge \vartheta^{\beta}$	$\sigma^* \Omega = \frac{1}{2} F_{\alpha} \beta^j L_j \otimes \vartheta^{\alpha} \wedge \vartheta^{\beta}$
mielke-geometrodynamics_E00574 (8.6.12)	185	■ 0.725	$L_{\mathrm{eff}} := L_{\mathrm{qPG}}(**) - dC_{\theta} = -\frac{\theta_{\mathrm{T}}^*}{2\ell^2} R^{\{\alpha\beta} \wedge \eta_{\alpha\beta} + \frac{\theta_{\mathrm{T}}^* \Lambda_{\mathrm{eff}}}{\ell^2} \eta.$	$L_{\mathrm{eff}} := L_{\mathrm{qPG}}(**) - dC_{\theta} = -\frac{\theta_{\mathrm{T}}^*}{2\ell^2} R^{\{\alpha\beta} \wedge \eta_{\alpha\beta} + \frac{\theta_{\mathrm{T}}^* \Lambda_{\mathrm{eff}}}{\ell^2} \eta.$	$L_{\mathrm{eff}} := L_{\mathrm{qPG}}(**) - dC_{\theta} = -\frac{\theta_{\mathrm{T}}^*}{2\ell^2} R^{\{\alpha\beta} \wedge \eta_{\alpha\beta} + \frac{\theta_{\mathrm{T}}^* \Lambda_{\mathrm{eff}}}{\ell^2} \eta.$
mielke-geometrodynamics_E00366 (6.2.13)	126	■ 0.725	$\stackrel{(\pm)}{L}_{\mathrm{EC}} := L_{\mathrm{EC}} \pm idC_{\mathrm{TT}} = -\frac{1}{2\ell^2} \stackrel{(\pm)}{R}_{\alpha\beta} \wedge *(\vartheta^{\alpha} \wedge \vartheta^{\beta}) + \frac{\Lambda}{\ell^2} \eta.$	$\stackrel{(\pm)}{L}_{\mathrm{EC}} := L_{\mathrm{EC}} \pm idC_{\mathrm{TT}} = -\frac{1}{2\ell^2} \stackrel{(\pm)}{R}_{\alpha\beta} \wedge *(\vartheta^{\alpha} \wedge \vartheta^{\beta}) + \frac{\Lambda}{\ell^2} \eta.$	$\stackrel{(\pm)}{L}_{\mathrm{EC}} := L_{\mathrm{EC}} \pm idC_{\mathrm{TT}} = -\frac{1}{2\ell^2} \stackrel{(\pm)}{R}_{\alpha\beta} \wedge *(\vartheta^{\alpha} \wedge \vartheta^{\beta}) + \frac{\Lambda}{\ell^2} \eta.$
mielke-geometrodynamics_E00585 (9.1.3)	193	■ 0.725	$ds^2 = \frac{n-2}{2\Lambda_{\mathrm{eff}}} (-dt^2 + dr^2) + \frac{(n-2)(n-3)}{2\Lambda_{\mathrm{eff}}} d\Omega^{n-2}$	$ds^2 = \frac{n-2}{2\Lambda_{\mathrm{eff}}} (-dt^2 + dr^2) + \frac{(n-2)(n-3)}{2\Lambda_{\mathrm{eff}}} d\Omega^{n-2}$	$ds^2 = \frac{n-2}{2\Lambda_{\mathrm{eff}}} (-dt^2 + dr^2) + \frac{(n-2)(n-3)}{2\Lambda_{\mathrm{eff}}} d\Omega^{n-2}$
mielke-geometrodynamics_E00021 (2.4.3)	38	■ 0.726	$\phi^{(p)} := \varphi^p \otimes b \in C^{\infty}(\wedge^p T_{\mathbb{C}}^*(M) \otimes V^p), \quad b \in C^{\infty}(V^p).$	$\phi^{(p)} := \varphi^p \otimes b \in C^{\infty}(\wedge^p T_{\mathbb{C}}^*(M) \otimes V^p), \quad b \in C^{\infty}(V^p).$	$\phi^{(p)} := \varphi^p \otimes b \in C^{\infty}(\wedge^p T_{\mathbb{C}}^*(M) \otimes V^p), \quad b \in C^{\infty}(V^p).$
mielke-geometrodynamics_E00209 (4.2.1)	85	■ 0.728	$\stackrel{*}{\chi} \widetilde{\omega} = \omega^{\mathrm{L}} \otimes \omega^{\mathrm{T}},$	$\stackrel{*}{\chi} \widetilde{\omega} = \omega^{\mathrm{L}} \otimes \omega^{\mathrm{T}},$	$\stackrel{*}{\chi} \widetilde{\omega} = \omega^{\mathrm{L}} \otimes \omega^{\mathrm{T}},$
mielke-geometrodynamics_E00309 (4.6.1)	104	■ 0.728	$\widetilde{\Gamma} := \Gamma^{(T)\alpha} P_{\alpha} + \Gamma_{\alpha}^{(L)\beta} L^{\alpha}_{\beta}$	$\widetilde{\Gamma} := \Gamma^{(T)\alpha} P_{\alpha} + \Gamma_{\alpha}^{(L)\beta} L^{\alpha}_{\beta}$	$\widetilde{\Gamma} := \Gamma^{(T)\alpha} P_{\alpha} + \Gamma_{\alpha}^{(L)\beta} L^{\alpha}_{\beta}$
mielke-geometrodynamics_E00009 (2.2.4)	34	■ 0.729	$E_i^{\alpha} e^i_{\beta} = \delta^{\alpha}_{\beta}.$	$E_i^{\alpha} e^i_{\beta} = \delta^{\alpha}_{\beta}.$	$E_i^{\alpha} e^i_{\beta} = \delta^{\alpha}_{\beta}.$
mielke-geometrodynamics_E00384 (6.3.2)	130	■ 0.730	$H_{\alpha}^{\parallel} = \frac{1}{\ell^2} \left[\vartheta^{\beta} \wedge * (d\vartheta_{\beta} \wedge \vartheta_{\alpha}) - \frac{1}{2} \vartheta_{\alpha} \wedge * (d\vartheta^{\beta} \wedge \vartheta_{\beta}) \right] = \frac{1}{2\ell^2} \Gamma^{\{\beta\gamma} \wedge \eta_{\alpha\beta\gamma}$	$H_{\alpha}^{\parallel} = \frac{1}{\ell^2} \left[\vartheta^{\beta} \wedge * (d\vartheta_{\beta} \wedge \vartheta_{\alpha}) - \frac{1}{2} \vartheta_{\alpha} \wedge * (d\vartheta^{\beta} \wedge \vartheta_{\beta}) \right] = \frac{1}{2\ell^2} \Gamma^{\{\beta\gamma} \wedge \eta_{\alpha\beta\gamma}$	$H_{\alpha}^{\parallel} = \frac{1}{\ell^2} \left[\vartheta^{\beta} \wedge * (d\vartheta_{\beta} \wedge \vartheta_{\alpha}) - \frac{1}{2} \vartheta_{\alpha} \wedge * (d\vartheta^{\beta} \wedge \vartheta_{\beta}) \right] = \frac{1}{2\ell^2} \Gamma^{\{\beta\gamma} \wedge \eta_{\alpha\beta\gamma}$
mielke-geometrodynamics_E00609 (10.3.47)	222	■ 0.731	$\tilde{\Lambda} = (-1)^{s*} L(\star) + \frac{1}{3\ell} [4\gamma + (-1)^s \ell] * \left(R^{\{\star\alpha} \wedge \vartheta_{\alpha} \right),$	$\tilde{\Lambda} = (-1)^{s*} L(\star) + \frac{1}{3\ell} [4\gamma + (-1)^s \ell] * \left(R^{\{\star\alpha} \wedge \vartheta_{\alpha} \right),$	$\tilde{\Lambda} = (-1)^s * L(\star) + \frac{1}{3\ell} [4\gamma + (-1)^s \ell] * \left(R^{\{\star\alpha} \wedge \vartheta_{\alpha} \right),$
mielke-geometrodynamics_E00662 (10.3.19)	217	■ 0.733	$\square := (-1)^{pn+s} [*D^*D + (-1)^n D^*D^*]$	$\square := (-1)^{pn+s} [*D^*D + (-1)^n D^*D^*]$	$\square := (-1)^{pn+s} [*D^*D + (-1)^n D^*D^*]$
mielke-geometrodynamics_E01128 (B.40)	373	■ 0.735	$\text{\texttt{d}} \, \partial \, c \equiv 0,$	$\text{\texttt{d}} \, \partial \, c \equiv 0,$	$\partial \, \partial \, c \equiv 0,$
mielke-geometrodynamics_E00374 (6.2.25)	127	■ 0.742	$\begin{aligned} \stackrel{(\pm)}{L} = & -\frac{i}{4} \left[\stackrel{(+)}{\Pi}^{\alpha} - \stackrel{(-)}{\Pi}^{\alpha} \right] \wedge \stackrel{(\pm)}{\Pi}_{\alpha} \\ & - \frac{i}{4} \left[\stackrel{(+)}{\Pi}^{\alpha\beta} - \stackrel{(-)}{\Pi}^{\alpha\beta} \right] \wedge \left(\stackrel{(\pm)}{\Pi}_{\alpha\beta} \mp \frac{i}{\ell^2} \vartheta_{\alpha} \wedge \vartheta_{\beta} - \frac{a_0}{2\ell^2} \eta_{\alpha\beta} \right) \end{aligned}$	$\begin{aligned} \stackrel{(\pm)}{L} = & -\frac{i}{4} \left[\stackrel{(+)}{\Pi}^{\alpha} - \stackrel{(-)}{\Pi}^{\alpha} \right] \wedge \stackrel{(\pm)}{\Pi}_{\alpha} \\ & - \frac{i}{4} \left[\stackrel{(+)}{\Pi}^{\alpha\beta} - \stackrel{(-)}{\Pi}^{\alpha\beta} \right] \wedge \left(\stackrel{(\pm)}{\Pi}_{\alpha\beta} \mp \frac{i}{\ell^2} \vartheta_{\alpha} \wedge \vartheta_{\beta} - \frac{a_0}{2\ell^2} \eta_{\alpha\beta} \right) \end{aligned}$	$\begin{aligned} \stackrel{(\pm)}{L} = & -\frac{i}{4} \ell^2 \left[\stackrel{(+)}{\Pi}^{\alpha} - \stackrel{(-)}{\Pi}^{\alpha} \right] \wedge \stackrel{(\pm)}{\Pi}_{\alpha} \\ & - \frac{i}{4} \left[\stackrel{(+)}{\Pi}^{\alpha\beta} - \stackrel{(-)}{\Pi}^{\alpha\beta} \right] \wedge \left(\stackrel{(\pm)}{\Pi}_{\alpha\beta} \mp \frac{i}{\ell^2} \vartheta_{\alpha} \wedge \vartheta_{\beta} - \frac{a_0}{2\ell^2} \eta_{\alpha\beta} \right) \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_167 EQ8524 (7.7.43)	167	■ 0.743	$\Theta^{\pm}\{\pm\}:=\frac{1}{2}\left(\Theta\pm\Theta^{\pm}\right),\quad\Omega^g\{\pm\}:=\frac{1}{2}\left(\Omega^g\pm\Omega^g\right),\quad\Omega^{\left(\pm\right)}:=\frac{1}{2}\left(\Omega^g\pm\Omega^{g\left(\pm\right)}\right),$	$\Theta^{\pm}:=\frac{1}{2}\left(\Theta\pm\Theta^{\pm}\right),\quad\Omega^{g\pm}:=\frac{1}{2}\left(\Omega^g\pm\Omega^g\right),\quad\Omega^{\left(\pm\right)}:=\frac{1}{2}\left(\Omega^g\pm\Omega^{g\left(\pm\right)}\right),$	$\Theta^{\pm}:=\frac{1}{2}\left(\Theta\pm\Theta^{\pm}\right),\quad\Omega^{g\pm}:=\frac{1}{2}\left(\Omega^g\pm\Omega^g\right),\quad\Omega^{\left(\pm\right)}:=\frac{1}{2}\left(\Omega^g\pm\Omega^{g\left(\pm\right)}\right),$
mielke-geometrodynamics_67 EQ8157 (3.5.5)	67	■ 0.746	$\{ \}^* F_{\alpha\beta\mu\nu}=\left(\stackrel{+}{-}\right)F_{\alpha\beta\mu\nu}$	$*F_{\alpha\beta\mu\nu}=\left(\stackrel{+}{-}\right)F_{\alpha\beta\mu\nu}$	$*F_{\alpha\beta\mu\nu}=\left(\stackrel{+}{-}\right)F_{\alpha\beta\mu\nu}$
mielke-geometrodynamics_369 EQ1899 (B.10)	369	■ 0.746	$\begin{aligned} \sigma_*\left(\alpha^{(p)}+\beta^{(p)}\right) &= \sigma_*\left(\alpha^{(p)}\right)+\sigma_*\left(\beta^{(p)}\right), \\ \sigma_*\left(\alpha^{(p)}\wedge\beta^{(q)}\right) &= \sigma_*\left(\alpha^{(p)}\right)\wedge\sigma_*\left(\beta^{(q)}\right), \\ d\left(\sigma_*\alpha^{(p)}\right) &= \sigma_*\left(d\alpha^{(p)}\right). \end{aligned}$	$\sigma_*\left(\alpha^{(p)}+\beta^{(p)}\right)=\sigma_*\left(\alpha^{(p)}\right)+\sigma_*\left(\beta^{(p)}\right)$ $\sigma_*\left(\alpha^{(p)}\wedge\beta^{(q)}\right)=\sigma_*\left(\alpha^{(p)}\right)\wedge\sigma_*\left(\beta^{(q)}\right)$ $d\left(\sigma_*\alpha^{(p)}\right)=\sigma_*\left(d\alpha^{(p)}\right)$	$\sigma_*(\alpha^{(p)}+\beta^{(p)})=\sigma_*(\alpha^{(p)})+\sigma_*(\beta^{(p)}),$ $\sigma_*(\alpha^{(p)}\wedge\beta^{(q)})=\sigma_*(\alpha^{(p)})\wedge\sigma_*(\beta^{(q)}),$ $d(\sigma_*\alpha^{(p)})=\sigma_*(d\alpha^{(p)}).$
mielke-geometrodynamics_216 EQ8655 (10.3.12) [re-fined: census]	216	■ 0.746	$E_{\alpha}=e_{\alpha}\rfloor L(\star)-\frac{2}{\ell^3}\frac{\gamma}{(1+\delta\varepsilon)}\eta_{\alpha}+(-1)^s\frac{2}{\ell}\frac{\gamma}{(1+\delta\varepsilon)}\left[\left(e_{\alpha}\rfloor R^{\star\beta}\right)\wedge\vartheta_{\beta}\right],$	$E_{\alpha}=e_{\alpha}\rfloor L(\star)-\frac{2\gamma\varepsilon^2}{\ell^3(1+\delta\varepsilon)}\eta_{\alpha}+(-1)^s\frac{2\gamma}{\ell(1+\delta\varepsilon)}\left[\left(e_{\alpha}\rfloor R^{\star\beta}\right)\wedge\vartheta_{\beta}\right],$	$E_{\alpha}=e_{\alpha}\rfloor L(\star)-\frac{2\gamma\varepsilon^2}{\ell^3(1+\delta\varepsilon)}\eta_{\alpha}+(-1)^s\frac{2\gamma}{\ell(1+\delta\varepsilon)}\left[\left(e_{\alpha}\rfloor R^{\star\beta}\right)\wedge\vartheta_{\beta}\right],$
mielke-geometrodynamics_156 EQ8477 (7.4.5)	156	■ 0.748	$g_{ij}=\Omega^2o_{ij}^{\mathrm{Mink}}$	$g_{ij}=\Omega^2o_{ij}^{\mathrm{Mink}}$	$g_{ij}=\Omega^2o_{ij}^{\mathrm{Mink}},$
mielke-geometrodynamics_312 EQ9963 (14.2.3)	312	■ 0.750	$\stackrel{\circ}{d}\mathcal{U}=\stackrel{\circ}{\vartheta}\wedge\stackrel{\circ}{\vartheta}.$	$\stackrel{\circ}{d}\mathcal{U}=\stackrel{\circ}{\vartheta}\wedge\stackrel{\circ}{\vartheta}.$	$\stackrel{\circ}{d}\vartheta=\stackrel{\circ}{\vartheta}\wedge\stackrel{\circ}{\vartheta}.$
mielke-geometrodynamics_222 EQ8687 (10.3.45) [re-fined: census]	222	■ 0.753	$-E_{\alpha}=-e_{\alpha}\rfloor L(\star)-(-1)^s\frac{2}{\ell}\left(e_{\alpha}\rfloor R^{\{\star\beta}\right)\wedge\vartheta_{\beta}=\ell\widetilde{\Sigma}_{\alpha}$	$-E_{\alpha}=-e_{\alpha}\rfloor L(\star)-(-1)^s\frac{2\gamma}{\ell}\left(e_{\alpha}\rfloor R^{\{\star\beta}\right)\wedge\vartheta_{\beta}=\ell\widetilde{\Sigma}_{\alpha}$	$-E_{\alpha}=-e_{\alpha}\rfloor L(\star)-(-1)^s\frac{2\gamma}{\ell}(e_{\alpha}\rfloor R^{\{\star\beta}\right)\wedge\vartheta_{\beta}=\ell\widetilde{\Sigma}_{\alpha}$
mielke-geometrodynamics_199 EQ8686 (9.2.5)	199	■ 0.753	$\chi(M)=\int_M\left(\gamma_2^g\right)-\frac{1}{(4\pi)^2}\int_{\partial M}\left\{\alpha\wedge\Omega^{g(\star)}+\frac{2}{3}\alpha\wedge(\alpha\wedge\alpha)^{(\star)}\right\}.$	$\chi(M)=\int_M\left(\gamma_2^g\right)-\frac{1}{(4\pi)^2}\int_{\partial M}\left\{\alpha\wedge\Omega^{g(\star)}+\frac{2}{3}\alpha\wedge(\alpha\wedge\alpha)^{(\star)}\right\}.$	$\chi(M)=\int_M\left(\gamma_2^g\right)-\frac{1}{(4\pi)^2}\int_{\partial M}\left\{\alpha\wedge\Omega^{g(\star)}+\frac{2}{3}\alpha\wedge(\alpha\wedge\alpha)^{(\star)}\right\}.$
mielke-geometrodynamics_320 EQ9992 (14.3.5)	320	■ 0.754	$\stackrel{\circ}{\gamma}^{K+1}\chi_{\rho F}^{\pm}(\xi)=\pm\chi_{\rho F}^{\pm}(\xi)$	$\stackrel{\circ}{\gamma}^{K+1}\chi_{\rho F}^{\pm}(\xi)=\pm\chi_{\rho F}^{\pm}(\xi)$	$\stackrel{\circ}{\gamma}^{K+1}\chi_{\rho F}^{\pm}(\xi)=\pm\chi_{\rho F}^{\pm}(\xi)$
mielke-geometrodynamics_92 EQ8249 (4.3.14)	92	■ 0.755	$D_kg_{ij}=E_k^{\alpha}\nabla_{\alpha}g_{ij}=0$	$D_kg_{ij}=E_k^{\alpha}\nabla_{\alpha}g_{ij}=0$	$D_kg_{ij}=E_k^{\alpha}\nabla_{\alpha}g_{ij}=0$
mielke-geometrodynamics_133 EQ8399 (6.4.17)	133	■ 0.755	$\begin{aligned} \left[D_{\alpha},D_{\beta}\right] &= -T_{\alpha\beta}{}^{\gamma}(x)D_{\gamma}+R_{\alpha\beta\gamma}{}^{\delta}(x)L^{\gamma}{}_{\delta}, \\ \left[L^{\alpha}{}_{\beta},D_{\gamma}\right] &= \delta_{\gamma}^{\alpha}D_{\beta}, \\ \left[L^{\alpha}{}_{\beta},L^{\gamma}{}_{\delta}\right] &= \delta_{\delta}^{\alpha}L^{\gamma}{}_{\beta}-\delta_{\beta}^{\gamma}L^{\alpha}{}_{\delta}, \end{aligned}$	$\left[D_{\alpha},D_{\beta}\right]=-T_{\alpha\beta}{}^{\gamma}(x)D_{\gamma}+R_{\alpha\beta\gamma}{}^{\delta}(x)L^{\gamma}{}_{\delta},$ $\left[L^{\alpha}{}_{\beta},D_{\gamma}\right]=\delta_{\gamma}^{\alpha}D_{\beta},$ $\left[L^{\alpha}{}_{\beta},L^{\gamma}{}_{\delta}\right]=\delta_{\delta}^{\alpha}L^{\gamma}{}_{\beta}-\delta_{\beta}^{\gamma}L^{\alpha}{}_{\delta},$	$\left[D_{\alpha},D_{\beta}\right]=-T_{\alpha\beta}{}^{\gamma}(x)D_{\gamma}+R_{\alpha\beta\gamma}{}^{\delta}(x)L^{\gamma}{}_{\delta},$ $\left[L^{\alpha}{}_{\beta},D_{\gamma}\right]=\delta_{\gamma}^{\alpha}D_{\beta},$ $\left[L^{\alpha}{}_{\beta},L^{\gamma}{}_{\delta}\right]=\delta_{\delta}^{\alpha}L^{\gamma}{}_{\beta}-\delta_{\beta}^{\gamma}L^{\alpha}{}_{\delta},$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_00957 (14.1.39)	309	■ 0.756	$\partial_{\kappa}\delta=\sqrt{ g }\varepsilon_{\kappa\lambda\rho\sigma}R^{\{\}\sigma}{}_{\cdot\nu}\nabla^{\rho}R^{\{\}\nu\lambda}/\left(R^{\{\}\alpha\beta}_{\alpha\beta}R^{\{\}\alpha\beta}\right).$	$\partial_{\kappa}\delta=\sqrt{ g }\varepsilon_{\kappa\lambda\rho\sigma}R^{\{\}\sigma}{}_{\cdot\nu}\nabla^{\rho}R^{\{\}\nu\lambda}/\left(R^{\{\}\alpha\beta}_{\alpha\beta}R^{\{\}\alpha\beta}\right).$	
mielke-geometrodynamics_00970 (14.2.11)	314	■ 0.758	$\hat{\omega}=\hat{d}\hat{\omega}-\hat{\omega}\wedge\hat{\omega}=\left[\begin{array}{l l}\Omega^g+\alpha\wedge\alpha^T&-D\alpha\\\hline\circ&\overset{\circ}{\Omega}+\alpha^T\wedge\alpha\end{array}\right].$	$\hat{\Omega}=\hat{d}\hat{\omega}-\hat{\omega}\wedge\hat{\omega}=\left[\begin{array}{l l}\Omega^g+\alpha\wedge\alpha^T&-D\alpha\\\hline\circ&\overset{\circ}{\Omega}+\alpha^T\wedge\alpha\end{array}\right].$	
mielke-geometrodynamics_01016 (14.4.11)	327	■ 0.758	$d\,s^2=-\left(1-\left(\frac{r_{\circ}}{r}\right)^4\right)\frac{\Delta}{r^2}dt^2+\frac{r^2}{\Delta}dr^2+r^2d\Omega^2.$	$ds^2=-\left(1-\left(\frac{r_{\circ}}{r}\right)^4\right)\frac{\Delta}{r^2}dt^2+\frac{r^2}{\Delta}dr^2+r^2d\Omega^2.$	$ds^2=-\left(1-\left(\frac{r_{\circ}}{r}\right)^4\right)\frac{\Delta}{r^2}dt^2+\frac{r^2}{\Delta}dr^2+r^2d\Omega^2.$
mielke-geometrodynamics_009310 (4.6.2)	104	■ 0.760	$\begin{aligned} \widetilde{\xi}^{\alpha}&=d\xi^{\alpha}+\Gamma_{\beta}^{(L)\alpha}\wedge\xi^{\beta}+\Gamma^{(T)\alpha}\\ &=D\xi^{\alpha}+\Gamma^{(T)\alpha}=0 \end{aligned}$	$\widetilde{D}\widetilde{\xi}^{\alpha}=d\xi^{\alpha}+\Gamma_{\beta}^{(L)\alpha}\wedge\xi^{\beta}+\Gamma^{(T)\alpha}=D\xi^{\alpha}+\Gamma^{(T)\alpha}=0$	$\widetilde{D}\widetilde{\xi}^{\alpha}=d\xi^{\alpha}+\Gamma_{\beta}^{(L)\alpha}\wedge\xi^{\beta}+\Gamma^{(T)\alpha}=D\xi^{\alpha}+\Gamma^{(T)\alpha}=0.$
mielke-geometrodynamics_00832 (12.1.4)	270	■ 0.763	$\left\langle d\widetilde{j}_5\right\rangle =0,$	$\left\langle d\widetilde{j}_5\right\rangle =0,$	$\langle d\widetilde{j}_5\rangle =0,$
mielke-geometrodynamics_00231 (4.2.24)	88	■ 0.766	$T_{\alpha\beta}^{\cdot\cdot c}=2D_{[\alpha}E_{\cdot \beta]}^c=2\left(\partial_{[\alpha}E_{\cdot \beta]}^c+\Gamma_{[\alpha d}^{\cdot c}E_{\cdot \beta]}^d\right).$	$T_{\alpha\beta}^{\cdot\cdot c}=2D_{[\alpha}E_{\cdot \beta]}^c=2\left(\partial_{[\alpha}E_{\cdot \beta]}^c+\Gamma_{[\alpha d}^{\cdot c}E_{\cdot \beta]}^d\right).$	$T_{\alpha\beta}^{\cdot\cdot c}=2D_{[\alpha}E_{\cdot \beta]}^c=2\left(\partial_{[\alpha}E_{\cdot \beta]}^c+\Gamma_{[\alpha d}^{\cdot c}E_{\cdot \beta]}^d\right).$
mielke-geometrodynamics_01079 (A.4)	365	■ 0.766	$\left[\sigma^i,\sigma^j\right]=\varepsilon^{ij}{}_k\sigma^k$	$\left[\sigma^i,\sigma^j\right]=\varepsilon^{ij}{}_k\sigma^k$	$[\sigma^i,\sigma^j]=\varepsilon^{ij}{}_k\sigma^k$
mielke-geometrodynamics_00276 (4.4.4)	97	■ 0.767	$\begin{aligned} \delta L_g&=\delta\vartheta\wedge E-\delta\vartheta\wedge D\Pi^T+d\left(\delta\vartheta\wedge\Pi^T\right)\\ &-\left[\delta\omega^g,\vartheta\right]\wedge\left[\Pi_{[]}^T,\vartheta\right]-\delta\omega^g\wedge D\Pi^L+d\left(\delta\omega^g\wedge\Pi^L\right), \end{aligned}$	$\delta L_g=\delta\vartheta\wedge E-\delta\vartheta\wedge D\Pi^T+d\left(\delta\vartheta\wedge\Pi^T\right)-\left[\delta\omega^g,\vartheta\right]\wedge\left[\Pi_{[]}^T,\vartheta\right]-\delta\omega^g\wedge D\Pi^L+d\left(\delta\omega^g\wedge\Pi^L\right),$	$\delta L_g=\delta\vartheta\wedge E-\delta\vartheta\wedge D\Pi^T+d\left(\delta\vartheta\wedge\Pi^T\right)-\left[\delta\omega^g,\vartheta\right]\wedge\left[\Pi_{[]}^T,\vartheta\right]-\delta\omega^g\wedge D\Pi^L+d\left(\delta\omega^g\wedge\Pi^L\right),$
mielke-geometrodynamics_00613 (10.1.1)	209	■ 0.769	$\vartheta^{\alpha}=E_i^{\alpha}dx^i\quad\text{and}\quad\Gamma_{\alpha}^{\star}:=\frac{1}{2}\eta_{\alpha\beta\gamma}\Gamma^{\beta\gamma}.$	$\vartheta^{\alpha}=E_i^{\alpha}dx^i\quad\text{and}\quad\Gamma_{\alpha}^{\star}:=\frac{1}{2}\eta_{\alpha\beta\gamma}\Gamma^{\beta\gamma}.$	$\vartheta^{\alpha}=E_i^{\alpha}dx^i\quad\text{and}\quad\Gamma_{\alpha}^{\star}:=\frac{1}{2}\eta_{\alpha\beta\gamma}\Gamma^{\beta\gamma}.$
mielke-geometrodynamics_01066 (16.2.5)	355	■ 0.770	$\rho^*=\int_o^{\rho}\frac{dx}{1\pm x^2}=\left\{\begin{array}{l} \operatorname{artan}\rho,\kappa>0\\ \operatorname{Arth}\rho,\kappa<0\end{array}\right\}.$	$\rho^*=\int_o^{\rho}\frac{dx}{1\pm x^2}=\left\{\begin{array}{l} \operatorname{artan}\rho,\kappa>0\\ \operatorname{Arth}\rho,\kappa<0\end{array}\right\}.$	$\rho^*=\int_o^{\rho}\frac{dx}{1\pm x^2}=\left\{\begin{array}{l} \operatorname{artan}\rho,\kappa>0\\ \operatorname{Arth}\rho,\kappa<0\end{array}\right\}.$
mielke-geometrodynamics_01030 (15.1.6)	337	■ 0.771	$\begin{aligned} \frac{1}{2}\vartheta^g\wedge^*\Omega^g+\Lambda^*\vartheta_g&=\ell^{*2}\left\{\Sigma_{\mathrm{m}}(\text{leptons})+\Sigma^{\mathrm{g}}\right\},\\ \frac{1}{2}\vartheta^f\wedge^*\Omega^f&=\ell^2\left\{\Sigma_{\mathrm{m}}(\text{hadrons})+\Sigma^{\mathrm{f}}\right\}. \end{aligned}$	$\frac{1}{2}\vartheta^g\wedge^*\Omega^g+\Lambda^*\vartheta_g=\ell^{*2}\left\{\Sigma_{\mathrm{m}}(\text{leptons})+\Sigma^{\mathrm{g}}\right\},$ $\frac{1}{2}\vartheta^f\wedge^*\Omega^f=\ell^2\left\{\Sigma_{\mathrm{m}}(\text{hadrons})+\Sigma^{\mathrm{f}}\right\}.$	$\frac{1}{2}\vartheta^g\wedge^*\Omega^g+\Lambda^*\vartheta_g=\ell^{*2}\left\{\Sigma_{\mathrm{m}}(\text{leptons})+\Sigma^{\mathrm{g}}\right\},$ $\frac{1}{2}\vartheta^f\wedge^*\Omega^f=\ell^2\left\{\Sigma_{\mathrm{m}}(\text{hadrons})+\Sigma^{\mathrm{f}}\right\}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_ E08112 (3.3.13)	59	■ 0.772	$\begin{aligned} j := & \mathscr{J} \wedge dt + \hat{\rho} \\ & = -(J_x dy \wedge dz + J_y dz \wedge dx + J_z dx \wedge dy) \wedge dt \\ & \quad + \rho dx \wedge dy \wedge dz \end{aligned}$	$j := -\mathscr{J} \wedge dt + \hat{\rho} = -(J_x dy \wedge dz + J_y dz \wedge dx + J_z dx \wedge dy) \wedge dt + \rho dx \wedge dy \wedge dz$	$j := -\mathscr{J} \wedge dt + \hat{\rho} = -(J_x dy \wedge dz + J_y dz \wedge dx + J_z dx \wedge dy) \wedge dt + \rho dx \wedge dy \wedge dz$
mielke-geometrodynamics_ E08138 (3.4.1)	62	■ 0.780	$\mathscr{F} := \frac{1}{\alpha_g} \operatorname{Tr} \left\{ \left(\Omega \wedge \right) \right\} = \frac{1}{4\alpha_g} F_{\alpha\beta}{}^j F^{\alpha\beta}{}_j$	$\mathscr{F} := \frac{1}{\alpha_g} \operatorname{Tr} \left\{ \left(\Omega \wedge \right) \right\} = \frac{1}{4\alpha_g} F_{\alpha\beta}{}^j F^{\alpha\beta}{}_j$	$\mathscr{F} := \frac{1}{\alpha_g} \operatorname{Tr} \left\{ \left(\Omega \wedge \right) \right\} = \frac{1}{4\alpha_g} F_{\alpha\beta}{}^j F^{\alpha\beta}{}_j$
mielke-geometrodynamics_ E08478 (7.3.9)	155	■ 0.782	$\left\{ \right\}^{\ast} \Omega^g = \zeta \Omega^g,$	$\left\{ \right\}^{\ast} \Omega^g = \zeta \Omega^g,$	$\left\{ \right\}^{\ast} \Omega^g = \zeta \Omega^g,$
mielke-geometrodynamics_ E08846 (2.6.9)	43	■ 0.783	$\begin{aligned} \Delta b &= \int_U d(\omega \otimes b)_{\mathrm{hor}} = \int_U D(\omega \otimes b) \\ &= \int_U \Omega \otimes b \simeq \Omega \otimes b \end{aligned}$	$\Delta b = \int_{\partial U} \omega \otimes b = \int_U d(\omega \otimes b)_{\mathrm{hor}} = \int_U D(\omega \otimes b) = \int_U \Omega \otimes b \simeq \Omega \otimes b$	$\Delta b = \int_{\partial U} \omega \otimes b = \int_U d(\omega \otimes b)_{\mathrm{hor}} = \int_U D(\omega \otimes b) = \int_U \Omega \otimes b \simeq \Omega \otimes b$
mielke-geometrodynamics_ E08849 (15.2.6)	344	■ 0.785	$\left\{ \right\}^{\ast} \bar{\omega} = \bar{\omega}_i^j \bar{L}_j \otimes dx^i = \bar{\omega}_i dx^i$	$\left\{ \right\}^{\ast} \bar{\omega} = \bar{\omega}_i^j \bar{L}_j \otimes dx^i = \bar{\omega}_i dx^i$	$\left\{ \right\}^{\ast} \bar{\omega} = \bar{\omega}_i^j \bar{L}_j \otimes dx^i = \bar{\omega}_i dx^i$
mielke-geometrodynamics_ E08133 (3.4.4)	63	■ 0.785	$L_{\omega} = L(\mathscr{F}, \ast \mathscr{F}) \eta.$	$L_{\omega} = L(\mathscr{F}, \ast \mathscr{F}) \eta.$	$L_{\omega} = L(\mathscr{F}, \ast \mathscr{F}) \eta.$
mielke-geometrodynamics_ E08312 (4.6.4) [re-fined: census]	104	■ 0.786	$\begin{aligned} \Delta \xi^{\alpha} &= - \oint_C y \rfloor (DD \xi^{\alpha}) = \oint_C y \rfloor R^{(T)\alpha} = \int_S R^{(T)\alpha} \\ &\simeq \frac{1}{2} (T_{ij}^{\alpha} - R_{ij\beta}^{\alpha} \xi^{\beta}) \int_S dy^i \wedge dy^j \end{aligned}$	$\Delta \xi^{\alpha} = - \oint_C y \rfloor (DD \xi^{\alpha}) = \oint_C y \rfloor R^{(T)\alpha} = \int_S R^{(T)\alpha} \simeq \frac{1}{2} (T_{ij}^{\alpha} - R_{ij\beta}^{\alpha} \xi^{\beta}) \int_S dy^i \wedge dy^j$	$\Delta \xi^{\alpha} = - \oint_C y \rfloor (DD \xi^{\alpha}) = \oint_C y \rfloor R^{(T)\alpha} = \int_S R^{(T)\alpha} \simeq \frac{1}{2} (T_{ij}^{\alpha} - R_{ij\beta}^{\alpha} \xi^{\beta}) \int_S dy^i \wedge dy^j.$
mielke-geometrodynamics_ E08224 (4.2.17)	87	■ 0.787	$\widetilde{\varphi} \rightarrow A^{-1} \widetilde{\varphi} = G^{-1} (\widetilde{\varphi} - T).$	$\widetilde{\varphi} \rightarrow A^{-1} \widetilde{\varphi} = G^{-1} (\widetilde{\varphi} - T).$	$\widetilde{\varphi} \rightarrow A^{-1} \widetilde{\varphi} = G^{-1} (\widetilde{\varphi} - T).$
mielke-geometrodynamics_ E08941 (14.1.22)	306	■ 0.788	$\tan 2\delta = (-1)^z \frac{\operatorname{Tr}^*(\Omega \wedge \Omega)}{\operatorname{Tr}^*(\Omega \wedge \ast \Omega)}.$	$\tan 2\delta = (-1)^z \frac{\operatorname{Tr}^*(\Omega \wedge \Omega)}{\operatorname{Tr}^*(\Omega \wedge \ast \Omega)}.$	$\tan 2\delta = (-1)^z \frac{\operatorname{Tr}^*(\Omega \wedge \Omega)}{\operatorname{Tr}^*(\Omega \wedge \ast \Omega)}.$
mielke-geometrodynamics_ E08768 (11.1.27)	242	■ 0.791	$\left(\gamma^{n+1} \right)^2 = \mp \mathbb{1}.$	$\left(\gamma^{n+1} \right)^2 = \mp \mathbb{1}.$	$\left(\gamma^{n+1} \right)^2 = \mp \mathbb{1}.$
mielke-geometrodynamics_ E08725 (10.10.5)	230	■ 0.793	$\delta_{\mathrm{susy}} L \cong \bar{\sigma} \left(i \gamma^{\alpha} \Psi \wedge \frac{\delta L}{\delta \vartheta^{\alpha}} - 2 D \frac{\delta L}{\delta \bar{\Psi}} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} \right) + 2 d \left(\bar{\sigma} \wedge \frac{\delta L}{\delta \bar{\Psi}} \right).$	$\delta_{\mathrm{susy}} L \cong \bar{\sigma} \left(i \gamma^{\alpha} \Psi \wedge \frac{\delta L}{\delta \vartheta^{\alpha}} - 2 D \frac{\delta L}{\delta \bar{\Psi}} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} \right) + 2 d \left(\bar{\sigma} \wedge \frac{\delta L}{\delta \bar{\Psi}} \right).$	$\delta_{\mathrm{susy}} L \cong \bar{\sigma} \left(i \gamma^{\alpha} \Psi \wedge \frac{\delta L}{\delta \vartheta^{\alpha}} - 2 D \frac{\delta L}{\delta \bar{\Psi}} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} \right) + 2 d \left(\bar{\sigma} \wedge \frac{\delta L}{\delta \bar{\Psi}} \right).$
mielke-geometrodynamics_ E08818 (11.4.22)	256	■ 0.793	$\frac{3}{8} \ell^2 \bar{\psi} \psi = 4 \pi m e^{-\mu - \nu/2} (H^2 - F^2) \left Y_{ \kappa -1}^{ \kappa -1}(\vartheta, \phi) \right ^2$	$\frac{3}{8} \ell^2 \bar{\psi} \psi = 4 \pi m e^{-\mu - \nu/2} (H^2 - F^2) \left Y_{ \kappa -1}^{ \kappa -1}(\vartheta, \phi) \right ^2$	$\frac{3}{8} \ell^2 \bar{\psi} \psi = 4 \pi m e^{-\mu - \nu/2} (H^2 - F^2) \left Y_{ \kappa -1}^{ \kappa -1}(\vartheta, \phi) \right ^2$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E01032 (15.1.9)	338	■ 0.793	$\frac{1}{2}*\left(\vartheta^g\wedge^*\Omega^g\right)+\Lambda\vartheta^g=0,$	$\frac{1}{2}*(\vartheta^g\wedge^*\Omega^g)+\Lambda\vartheta^g=0,$	$\frac{1}{2}*(\vartheta^g\wedge^*\Omega^g)+\Lambda\vartheta^g=0,$
mielke-geometrodynamics_E00305 (4.5.15) [re-fined: census]	103	■ 0.793	$\mathrm{varepsilon}_{\mathrm{RC}}:=\mathrm{x}^{\mathrm{\alpha}}\mathrm{\Sigma}_{\mathrm{\alpha}}+\left(e_{\beta}\rfloor\overparen{\mathrm{D}\xi^{\gamma}}\right)\tau^{\beta}_{\gamma},$	$\varepsilon_{\mathrm{RC}}:=\xi^{\alpha}\Sigma_{\alpha}+\left(e_{\beta}\rfloor\widehat{D\xi^{\gamma}}\right)\tau^{\beta}_{\gamma},$	$\varepsilon_{\mathrm{RC}}:=\xi^{\alpha}\Sigma_{\alpha}+(e_{\beta}\rfloor\widehat{D}\xi^{\gamma})\tau^{\beta}_{\gamma},$
mielke-geometrodynamics_E00848 (12.3.1)	274	■ 0.794	$\left\langle d j_5\right\rangle =2 i m\left\langle \bar{\psi} \gamma_5 \psi\right\rangle \eta-\frac{1}{4 \pi^2} \operatorname{Tr}(G \wedge G)-\frac{1}{96 \pi^2}\left[2 R_{\alpha \beta}^{\{\}} \wedge R^{\{\beta \alpha \beta}+d \mathscr{A} \wedge d \mathscr{A}\right]$	$\langle dj_5\rangle=2im\langle\bar{\psi}\gamma_5\psi\rangle\eta-\frac{1}{4\pi^2}\operatorname{Tr}(G\wedge G)-\frac{1}{96\pi^2}\left[2R_{\alpha\beta}^{\{\}}\wedge R^{\{\beta\alpha\beta}+d\mathscr{A}\wedge d\mathscr{A}\right]$	$\langle dj_5\rangle=2im\langle\bar{\psi}\gamma_5\psi\rangle\eta-\frac{1}{4\pi^2}Tr(G\wedge G)-\frac{1}{96\pi^2}\left[2R_{\alpha\beta}^{\{\}}\wedge R^{\{\}\alpha\beta}+d\mathscr{A}\wedge d\mathscr{A}\right]$
mielke-geometrodynamics_E00319 (5.2.3)	110	■ 0.795	$\begin{aligned} L_{\mathrm{D}} &= L\left(\gamma, \psi, D^{\{\}} \psi\right)-\frac{i}{2} \overline{\psi}\left(* \gamma \wedge K-K \wedge * \gamma\right) \psi \\ &= L\left(\gamma, \psi, D^{\{\}} \psi\right)+\frac{1}{4} \mathscr{A} \wedge j_5 . \end{aligned}$	$L_{\mathrm{D}}=L\left(\gamma, \psi, D^{\{\}} \psi\right)-\frac{i}{2} \bar{\psi}\left(* \gamma \wedge K-K \wedge * \gamma\right) \psi \\ =L\left(\gamma, \psi, D^{\{\}} \psi\right)+\frac{1}{4} \mathscr{A} \wedge j_5 .$	$L_{\mathrm{D}}=L(\gamma, \psi, D^{\{\}} \psi)-\frac{i}{2} \overline{\psi}\left(* \gamma \wedge K-K \wedge * \gamma\right) \psi \\ =L(\gamma, \psi, D^{\{\}} \psi)+\frac{1}{4} \mathscr{A} \wedge j_5 .$
mielke-geometrodynamics_E00552 (8.4.6)	179	■ 0.797	$\stackrel{+g}{\Omega}:=\frac{1}{2}\left(\Omega^g \pm * \Omega^{g(\star)}\right)=0$	$\stackrel{+g}{\Omega}:=\frac{1}{2}\left(\Omega^g\pm*\Omega^{g(\star)}\right)=0$	$\stackrel{\pm}{\Omega}^g:=\frac{1}{2}\left(\Omega^g\pm*\Omega^{g(\star)}\right)=0$
mielke-geometrodynamics_E01001 (14.3.14)	323	■ 0.798	$(i\not{D})^2:=-\operatorname{Tr}\left\{\left(\gamma \wedge^* D^*\right)\left(\gamma \wedge^* D\right)\right\} .$	$(i\not{D})^2:=-\operatorname{Tr}\left\{\gamma \wedge^* D^*\left(\gamma \wedge^* D\right)\right\} .$	$(i\not{D})^2:=-\operatorname{Tr}\left\{\gamma \wedge^* D^*\left(\gamma \wedge^* D\right)\right\} .$
mielke-geometrodynamics_E00065 (2.8.8)	47	■ 0.803	$c_2(M)=-\int_M \pi\left(\gamma_2\right)=\frac{1}{8 \pi^2} \int_M \operatorname{Tr}(\Omega \wedge \Omega) .$	$c_2(M)=-\int_M^*\pi\left(\gamma_2\right)=\frac{1}{8 \pi^2} \int_M \operatorname{Tr}(\Omega \wedge \Omega) .$	$c_2(M)=-\int_M^*\pi\left(\gamma_2\right)=\frac{1}{8 \pi^2} \int_M \operatorname{Tr}(\Omega \wedge \Omega) .$
mielke-geometrodynamics_E00652 (10.3.9)	216	■ 0.803	$H_{\alpha}=-\frac{\gamma \varepsilon}{\ell^2\left(1+\delta \varepsilon\right)} \vartheta_{\alpha} \quad \Leftrightarrow \quad T_{\alpha}=\frac{\varepsilon}{\ell} \eta_{\alpha},$	$H_{\alpha}=-\frac{\gamma\varepsilon}{\ell^2(1+\delta\varepsilon)}\vartheta_{\alpha}\quad\Leftrightarrow\quad T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},$	$H_{\alpha}=-\frac{\gamma\varepsilon}{\ell^2(1+\delta\varepsilon)}\vartheta_{\alpha}\quad\Leftrightarrow\quad T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},$
mielke-geometrodynamics_E01065 (16.2.4)	355	■ 0.803	$\rho:=\sqrt{\frac{1}{3} \Lambda_{\text {eff }}} r, \quad r= \mathbf{x} $	$\rho:=\sqrt{\frac{1}{3} \Lambda_{\text{eff}}} r, \quad r= \mathbf{x} $	$\rho:=\sqrt{\frac{1}{3} \Lambda_{\text{eff}}} r, \quad r= \mathbf{x} $
mielke-geometrodynamics_E00166 (3.6.1)	69	■ 0.806	$\begin{aligned} L_{\mathrm{SW}} &= \frac{1}{2} \overline{D_{\pm} \psi} \wedge^* D_{\pm} \psi \mp i\left(F^{\pm}-\frac{1}{2} \bar{\psi} \sigma_{\pm} \psi\right)^2 \\ &= \mp i \operatorname{Tr}\left(F^{\pm} \wedge F^{\pm}\right)+\frac{1}{2} \overline{D_{\pm} \psi} \wedge^* D_{\pm} \psi \\ &\quad \pm i \bar{\psi} \sigma_{\pm} \psi \wedge F^{\pm} \mp \frac{i}{4} \bar{\psi} \sigma_{\pm} \psi \wedge \bar{\psi} \sigma_{\pm} \psi ; \end{aligned}$	$L_{\mathrm{SW}}=\frac{1}{2}\overline{D_{\pm}\psi}\wedge^*D_{\pm}\psi\mp i\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)^2 \\ =\mp i\operatorname{Tr}\left(F^{\pm}\wedge F^{\pm}\right)+\frac{1}{2}\overline{D_{\pm}\psi}\wedge^*D_{\pm}\psi \\ \pm i\bar{\psi}\sigma_{\pm}\psi\wedge F^{\pm}\mp\frac{i}{4}\bar{\psi}\sigma_{\pm}\psi\wedge\bar{\psi}\sigma_{\pm}\psi;$	$L_{\mathrm{SW}}=\frac{1}{2}\overline{D_{\pm}\psi}\wedge^*D_{\pm}\psi\mp i\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)^2 \\ =\mp i\operatorname{Tr}\left(F^{\pm}\wedge F^{\pm}\right)+\frac{1}{2}\overline{D_{\pm}\psi}\wedge^*D_{\pm}\psi \\ \pm i\bar{\psi}\sigma_{\pm}\psi\wedge F^{\pm}\mp\frac{i}{4}\bar{\psi}\sigma_{\pm}\psi\wedge\bar{\psi}\sigma_{\pm}\psi;$
mielke-geometrodynamics_E01070 (16.2.9)	356	■ 0.806	$V_{\mathrm{eff}}^o\left(\rho^*\right)=\left(\frac{1}{\beta^2}-2\right)\left[1+\tan ^2 \rho^*\right]$	$V_{\mathrm{eff}}^o\left(\rho^*\right)=\left(\frac{1}{\beta^2}-2\right)\left[1+\tan ^2 \rho^*\right]$	$V_{\mathrm{eff}}^o\left(\rho^*\right)=\left(\frac{1}{\beta^2}-2\right)\left[1+\tan ^2 \rho^*\right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_144 EQ0446 (6.9.2)	144	■ 0.806	$\begin{aligned} \stackrel{\theta}{L} &= \frac{1}{2} \left[T^\alpha \wedge {}^* T_\alpha + \theta_{\mathrm{T}} \left(T^\alpha \wedge T_\alpha + R_{\alpha\beta} \wedge \vartheta^\alpha \wedge \vartheta^\beta \right) \right. \\ &\quad \left. + \frac{1}{2} \left[R^{\alpha\beta} \wedge {}^* R_{\alpha\beta} + \theta_{\mathrm{L}} R^{\alpha\beta} \wedge R_{\alpha\beta} \right] \right]. \end{aligned}$	$\stackrel{\theta}{L} = \frac{1}{2\ell} \left[T^\alpha \wedge {}^* T_\alpha + \theta_{\mathrm{T}} \left(T^\alpha \wedge T_\alpha + R_{\alpha\beta} \wedge \vartheta^\alpha \wedge \vartheta^\beta \right) \right. \\ \left. + \frac{1}{2} \left[R^{\alpha\beta} \wedge {}^* R_{\alpha\beta} + \theta_{\mathrm{L}} R^{\alpha\beta} \wedge R_{\alpha\beta} \right] \right].$	$\stackrel{\theta}{L} = \frac{1}{2\ell} \left[T^\alpha \wedge {}^* T_\alpha + \theta_{\mathrm{T}} \left(T^\alpha \wedge T_\alpha + R_{\alpha\beta} \wedge \vartheta^\alpha \wedge \vartheta^\beta \right) \right. \\ \left. + \frac{1}{2} \left[R^{\alpha\beta} \wedge {}^* R_{\alpha\beta} + \theta_{\mathrm{L}} R^{\alpha\beta} \wedge R_{\alpha\beta} \right] \right].$
mielke-geometrodynamics_202 EQ0610 (9.3.1)	202	■ 0.806	$\begin{aligned} d\,C_{\mathrm{RR}^{(\star)}} &\&:= \\ &\frac{1}{2}\,d\left(\Gamma_{\alpha\beta}\wedge R^{\alpha\beta(\star)}-\frac{1}{3}\,\Gamma_{\alpha}{}^{\beta(\star)}\wedge\Gamma_{\beta}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\alpha}\right) \\ &\equiv -L_{\mathrm{SKY}}-2\,\mathrm{Ric}_{\alpha\beta}\wedge{}^*\mathrm{Ric}^{\alpha\beta}+\frac{1}{2}\,\mathrm{Ric}_{\alpha}{}^{\alpha}\wedge{}^*\mathrm{Ric}_{\beta}{}^{\beta} \end{aligned}$	$dC_{\mathrm{RR}^{(\star)}} := \frac{1}{2}d\left(\Gamma_{\alpha\beta}\wedge R^{\alpha\beta(\star)}-\frac{1}{3}\Gamma_{\alpha}{}^{\beta(\star)}\wedge\Gamma_{\beta}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\alpha}\right) \\ \equiv -L_{\mathrm{SKY}}-2\,\mathrm{Ric}_{\alpha\beta}\wedge{}^*\mathrm{Ric}^{\alpha\beta}+\frac{1}{2}\,\mathrm{Ric}_{\alpha}{}^{\alpha}\wedge{}^*\mathrm{Ric}_{\beta}{}^{\beta}$	$dC_{\mathrm{RR}^{(\star)}} := \frac{1}{2}d\left(\Gamma_{\alpha\beta}\wedge R^{\alpha\beta(\star)}-\frac{1}{3}\Gamma_{\alpha}{}^{\beta(\star)}\wedge\Gamma_{\beta}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\alpha}\right) \\ \equiv -L_{\mathrm{SKY}}-2\mathrm{Ric}_{\alpha\beta}\wedge{}^*\mathrm{Ric}^{\alpha\beta}+\frac{1}{2}\mathrm{Ric}_{\alpha}{}^{\alpha}\wedge{}^*\mathrm{Ric}_{\beta}{}^{\beta}$
mielke-geometrodynamics_244 EQ0767 (11.1.34)	244	■ 0.807	$\psi \longrightarrow S^{-1}\psi = S^{-1}\psi$	$\psi \longrightarrow S^{-1}\psi = S^{-1}\psi$	$\psi \longrightarrow S^{-1}\psi = S^{-1}\psi$
mielke-geometrodynamics_95 EQ0267 (4.3.32)	95	■ 0.808	$\begin{aligned} &R_{(ij)\kappa l} = R_{ij(\kappa l)} = 0 \\ &R_{ij\kappa l} = R_{\kappa lij}. \end{aligned}$	$R_{(ij)\kappa l} = R_{ij(\kappa l)} = 0 \\ R_{ij\kappa l} = R_{\kappa lij}.$	$R_{(ij)\kappa l} = R_{ij(\kappa l)} = 0 \\ R_{ij\kappa l} = R_{\kappa lij}.$
mielke-geometrodynamics_101 EQ0298 (4.5.8)	101	■ 0.811	$\begin{aligned} \delta L &= \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha} \\ &\quad + \delta \Gamma_\alpha{}^\beta \wedge I^\alpha{}_\beta \Psi \wedge \frac{\partial L}{\partial D\Psi} + \delta \Psi \wedge \frac{\delta L}{\delta \Psi} + D\left(\delta \Psi \wedge \frac{\partial L}{\partial D\Psi}\right). \end{aligned}$	$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha} \\ + \delta \Gamma_\alpha{}^\beta \wedge I^\alpha{}_\beta \Psi \wedge \frac{\partial L}{\partial D\Psi} + \delta \Psi \wedge \frac{\delta L}{\delta \Psi} + D\left(\delta \Psi \wedge \frac{\partial L}{\partial D\Psi}\right)$	$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha} \\ + \delta \Gamma_\alpha{}^\beta \wedge I^\alpha{}_\beta \Psi \wedge \frac{\partial L}{\partial D\Psi} + \delta \Psi \wedge \frac{\delta L}{\delta \Psi} + D\left(\delta \Psi \wedge \frac{\partial L}{\partial D\Psi}\right).$
mielke-geometrodynamics_295 EQ0919 (13.8.5)	295	■ 0.811	$D\,T^\alpha \wedge R_{\beta}{}^\alpha \wedge \vartheta^\beta$	$DT^\alpha \equiv R_\beta{}^\alpha \wedge \vartheta^\beta$	$DT^\alpha \equiv R_\beta{}^\alpha \wedge \vartheta^\beta$
mielke-geometrodynamics_254 EQ0802 (11.4.14)	254	■ 0.812	$\begin{aligned} &\left\{ i\gamma^0\partial_0 - e^{(\nu-\mu)/2} \left[i\vec{\gamma}\cdot\vec{\partial} + i\gamma^1\partial_r \left(\frac{\mu}{2} + \frac{\nu}{4} \right) \right] \right. \\ &\quad \left. + \frac{3\varepsilon}{8}\ell^2 e^{\nu/2} \left(\bar{\psi}\psi - \mathrm{a}\bar{\psi}\gamma^5\psi\gamma^5 \right) - e^{\nu/2}m \right\} \psi = 0 \end{aligned}$	$\left\{ i\gamma^0\partial_0 - e^{(\nu-\mu)/2} \left[i\vec{\gamma}\cdot\vec{\partial} + i\gamma^1\partial_r \left(\frac{\mu}{2} + \frac{\nu}{4} \right) \right] \right. \\ \left. + \frac{3\varepsilon}{8}\ell^2 e^{\nu/2} \left(\bar{\psi}\psi - \mathrm{a}\bar{\psi}\gamma^5\psi\gamma^5 \right) - e^{\nu/2}m \right\} \psi = 0$	$\left\{ i\gamma^0\partial_0 - e^{(\nu-\mu)/2} \left[i\vec{\gamma}\cdot\vec{\partial} + i\gamma^1\partial_r \left(\frac{\mu}{2} + \frac{\nu}{4} \right) \right] \right. \\ \left. + \frac{3\varepsilon}{8}\ell^2 e^{\nu/2} (\bar{\psi}\psi - \mathrm{a}\bar{\psi}\gamma^5\psi\gamma^5) - e^{\nu/2}m \right\} \psi = 0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E08887 (11.4.19)	255	■ 0.812	$\vec{\sigma} \cdot \vec{L} \chi_{\kappa}^m = -(\kappa + 1) \chi_{\kappa}^m$.	$\vec{\sigma} \cdot \vec{L} \chi_{\kappa}^m = -(\kappa + 1) \chi_{\kappa}^m$.	$\vec{\sigma} \cdot \vec{L} \chi_{\kappa}^m = -(\kappa + 1) \chi_{\kappa}^m$.
mielke-geometrodynamics_E08397 (6.4.12)	132	■ 0.813	$\mathscr{H} \cong n^{\alpha} \mathscr{G}_{\alpha} + \Gamma_{\perp}^{\alpha\beta} \mathscr{G}_{\alpha\beta}$,	$\mathscr{H} \cong n^{\alpha} \mathscr{G}_{\alpha} + \Gamma_{\perp}^{\alpha\beta} \mathscr{G}_{\alpha\beta}$,	$\mathscr{H} \cong n^{\alpha} \mathscr{G}_{\alpha} + \Gamma_{\perp}^{\alpha\beta} \mathscr{G}_{\alpha\beta}$,
mielke-geometrodynamics_E08789 (10.7.5)	227	■ 0.813	$R_{\beta}{}^{\alpha} \wedge \vartheta^{\beta} = \frac{\ell^2}{2} \varepsilon(x^{\alpha} Y \wedge X \wedge Y - y^{\alpha} X \wedge X \wedge Y) = 0$.	$R_{\beta}{}^{\alpha} \wedge \vartheta^{\beta} = \frac{\ell^2}{2} \varepsilon(x^{\alpha} Y \wedge X \wedge Y - y^{\alpha} X \wedge X \wedge Y) = 0$.	$R_{\beta}{}^{\alpha} \wedge \vartheta^{\beta} = \frac{\ell^2}{2} \varepsilon(x^{\alpha} Y \wedge X \wedge Y - y^{\alpha} X \wedge X \wedge Y) = 0$.
mielke-geometrodynamics_E08218 (4.2.2)	85	■ 0.814	$\stackrel{*}{\chi} \approx \left[\begin{array}{c c} \omega^L & \omega^T \\ \hline 0 & 0 \end{array} \right]$.	$\stackrel{*}{\chi} \approx \left[\begin{array}{c c} \omega^L & \omega^T \\ \hline 0 & 0 \end{array} \right]$.	$\stackrel{*}{\chi} \omega \approx \left[\begin{array}{c c} \omega^L & \omega^T \\ \hline 0 & 0 \end{array} \right]$.
mielke-geometrodynamics_E08769 (11.1.36)	244	■ 0.817	$\widetilde{\omega} \longrightarrow {}^S \widetilde{\omega} = S \widetilde{\omega} S^{-1} + (dS) S^{-1}$	$\widetilde{\omega} \longrightarrow {}^S \widetilde{\omega} = S \widetilde{\omega} S^{-1} + (dS) S^{-1}$	$\widetilde{\omega} \longrightarrow {}^S \widetilde{\omega} = S \widetilde{\omega} S^{-1} + (dS) S^{-1}$
mielke-geometrodynamics_E08398 (6.4.14)	133	■ 0.820	$\begin{aligned} & \left(\mathscr{G}_{\alpha}(t, \mathbf{x}), \mathscr{G}_{\beta}(t, \mathbf{y}) \right) = \left(-T_{\alpha\beta}{}^{\gamma} \mathscr{G}_{\gamma} + R_{\alpha\beta\gamma}{}^{\delta} \mathscr{G}^{\gamma}_{\delta} \right) \cdot \delta(\mathbf{x} - \mathbf{y}), \\ & \left(\mathscr{G}^{\alpha}_{\beta}(t, \mathbf{x}), \mathscr{G}_{\gamma}(t, \mathbf{y}) \right) = \delta_{\beta}^{\gamma} \mathscr{G}_{\beta} \delta(\mathbf{x} - \mathbf{y}), \\ & \left(\mathscr{G}^{\alpha}_{\beta}(t, \mathbf{x}), \mathscr{G}^{\gamma}_{\delta}(t, \mathbf{y}) \right) = \left(\delta_{\delta}^{\alpha} \mathscr{G}^{\gamma}_{\beta} - \delta_{\beta}^{\gamma} \mathscr{G}^{\alpha}_{\delta} \right) \cdot \delta(\mathbf{x} - \mathbf{y}). \end{aligned}$	$\begin{aligned} & \left(\mathscr{G}_{\alpha}(t, \mathbf{x}), \mathscr{G}_{\beta}(t, \mathbf{y}) \right) = \left(-T_{\alpha\beta}{}^{\gamma} \mathscr{G}_{\gamma} + R_{\alpha\beta\gamma}{}^{\delta} \mathscr{G}^{\gamma}_{\delta} \right) \cdot \delta(\mathbf{x} - \mathbf{y}), \\ & \left(\mathscr{G}^{\alpha}_{\beta}(t, \mathbf{x}), \mathscr{G}_{\gamma}(t, \mathbf{y}) \right) = \delta_{\beta}^{\gamma} \mathscr{G}_{\beta} \delta(\mathbf{x} - \mathbf{y}), \\ & \left(\mathscr{G}^{\alpha}_{\beta}(t, \mathbf{x}), \mathscr{G}^{\gamma}_{\delta}(t, \mathbf{y}) \right) = \left(\delta_{\delta}^{\alpha} \mathscr{G}^{\gamma}_{\beta} - \delta_{\beta}^{\gamma} \mathscr{G}^{\alpha}_{\delta} \right) \cdot \delta(\mathbf{x} - \mathbf{y}). \end{aligned}$	$\begin{aligned} & \left(\mathscr{G}_{\alpha}(t, \mathbf{x}), \mathscr{G}_{\beta}(t, \mathbf{y}) \right) = \left(-T_{\alpha\beta}{}^{\gamma} \mathscr{G}_{\gamma} + R_{\alpha\beta\gamma}{}^{\delta} \mathscr{G}^{\gamma}_{\delta} \right) \cdot \delta(\mathbf{x} - \mathbf{y}), \\ & \left(\mathscr{G}^{\alpha}_{\beta}(t, \mathbf{x}), \mathscr{G}_{\gamma}(t, \mathbf{y}) \right) = \delta_{\beta}^{\gamma} \mathscr{G}_{\beta} \delta(\mathbf{x} - \mathbf{y}), \\ & \left(\mathscr{G}^{\alpha}_{\beta}(t, \mathbf{x}), \mathscr{G}^{\gamma}_{\delta}(t, \mathbf{y}) \right) = \left(\delta_{\delta}^{\alpha} \mathscr{G}^{\gamma}_{\beta} - \delta_{\beta}^{\gamma} \mathscr{G}^{\alpha}_{\delta} \right) \cdot \delta(\mathbf{x} - \mathbf{y}). \end{aligned}$
mielke-geometrodynamics_E08659 (2.7.9)	45	■ 0.820	$\bar{\varphi} \rightarrow {}^{G^{-1}} \bar{\varphi} := \bar{\varphi} G \in C^{\infty}(V^{\bar{s}})$.	$\bar{\varphi} \rightarrow {}^{G^{-1}} \bar{\varphi} := \bar{\varphi} G \in C^{\infty}(V^{\bar{s}})$.	$\bar{\varphi} \rightarrow {}^{G^{-1}} \bar{\varphi} := \bar{\varphi} G \in C^{\infty}(V^{\bar{s}})$.
mielke-geometrodynamics_E08993 (14.3.6)	320	■ 0.822	$\int_G \pi \hat{L}_{\text{D}} = \frac{i}{2} \left(\bar{\psi} \gamma \wedge {}^* D^{g+\omega} \psi + \left(\overline{D^{g+\omega} \psi} \right) \wedge {}^* \gamma \psi \right) + \frac{\hat{\ell}}{2} \bar{\psi} \gamma \wedge \gamma \wedge {}^* \Omega \psi - \widehat{m} \bar{\psi} \psi \eta,$	$\int_G \pi \hat{L}_{\text{D}} = \frac{i}{2} \left(\bar{\psi} \gamma \wedge {}^* D^{g+\omega} \psi + \left(\overline{D^{g+\omega} \psi} \right) \wedge {}^* \gamma \psi \right) + \frac{\hat{\ell}}{2} \bar{\psi} \gamma \wedge \gamma \wedge {}^* \Omega \psi - \widehat{m} \bar{\psi} \psi \eta,$	$\int_G \pi \hat{L}_{\text{D}} = \frac{i}{2} \left(\bar{\psi} \gamma \wedge {}^* D^{g+\omega} \psi + \left(\overline{D^{g+\omega} \psi} \right) \wedge {}^* \gamma \psi \right) + \frac{\hat{\ell}}{2} \bar{\psi} \gamma \wedge \gamma \wedge {}^* \Omega \psi - \widehat{m} \bar{\psi} \psi \eta,$
mielke-geometrodynamics_E08844 (12.2.12)	273	■ 0.831	$\mathrm{L} = \frac{1}{2\ell^2} \mathrm{Tr}(\Omega \wedge {}^* \sigma) + \mathrm{L}_{\text{D}} = \frac{1}{2\ell^2} \mathrm{R}^{\alpha\beta} \wedge \eta_{\alpha\beta} + \mathrm{L}_{\text{D}}$,	$\mathrm{L} = \frac{1}{2\ell^2} \mathrm{Tr}(\Omega \wedge {}^* \sigma) + \mathrm{L}_{\text{D}} = \frac{1}{2\ell^2} \mathrm{R}^{\alpha\beta} \wedge \eta_{\alpha\beta} + \mathrm{L}_{\text{D}}$,	$\mathrm{L} = \frac{1}{2\ell^2} \mathrm{Tr}(\Omega \wedge {}^* \sigma) + \mathrm{L}_{\text{D}} = \frac{1}{2\ell^2} \mathrm{R}^{\alpha\beta} \wedge \eta_{\alpha\beta} + \mathrm{L}_{\text{D}}$,

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00218 (4.2.10)	86	■ 0.836	$\begin{aligned} & \omega^{\mathrm{L}} \rightarrow {}^{A^{-1}}\omega^{\mathrm{L}} = G^{-1}\omega^{\mathrm{L}}G - G^{-1}dG, \\ & \omega^{\mathrm{T}} \rightarrow {}^{A^{-1}}\omega^{\mathrm{T}} = G^{-1}\omega^{\mathrm{T}} - G^{-1}DT. \end{aligned}$	$\omega^{\mathrm{L}} \rightarrow {}^{A^{-1}}\omega^{\mathrm{L}} = G^{-1}\omega^{\mathrm{L}}G - G^{-1}dG,$ $\omega^{\mathrm{T}} \rightarrow {}^{A^{-1}}\omega^{\mathrm{T}} = G^{-1}\omega^{\mathrm{T}} - G^{-1}DT.$	$\omega^{\mathrm{L}} \rightarrow {}^{A^{-1}}\omega^{\mathrm{L}} = G^{-1}\omega^{\mathrm{L}}G - G^{-1}dG,$ $\omega^{\mathrm{T}} \rightarrow {}^{A^{-1}}\omega^{\mathrm{T}} = G^{-1}\omega^{\mathrm{T}} - G^{-1}DT.$
mielke-geometrodynamics_E00240 (4.3.5)	90	■ 0.836	$\begin{aligned} & F(m, f \, \vartheta) = f \, F(m, \, \vartheta), \\ & F(m, \vartheta) > 0 \quad \text{if } \vartheta \neq 0. \end{aligned}$	$F(m, f \vartheta) = f F(m, \vartheta),$ $F(m, \vartheta) > 0 \quad \text{if } \vartheta \neq 0.$	$F(m, f \vartheta) = f F(m, \vartheta),$ $F(m, \vartheta) > 0 \quad \text{if } \vartheta \neq 0.$
mielke-geometrodynamics_E00445 (6.9.1)	144	■ 0.837	$ds^2 = -\left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right)^{-1} dr^2 + r^2 d\Omega^2,$	$ds^2 = -\left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right)^{-1} dr^2 + r^2 d\Omega^2,$	$ds^2 = -\left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2}\right)^{-1} dr^2 + r^2 d\Omega^2,$
mielke-geometrodynamics_E00227 (4.2.20)	88	■ 0.839	$\widetilde{\vartheta} \rightarrow {}^{A^{-1}}\widetilde{\vartheta} = G^{-1}\widetilde{\vartheta}.$	$\widetilde{\vartheta} \rightarrow {}^{A^{-1}}\widetilde{\vartheta} = G^{-1}\widetilde{\vartheta}.$	$\widetilde{\vartheta} \rightarrow {}^{A^{-1}}\widetilde{\vartheta} = G^{-1}\widetilde{\vartheta}.$
mielke-geometrodynamics_E00773 (11.2.2)	245	■ 0.840	$i \gamma \wedge {}^* D \psi - \frac{i}{2} (D \wedge {}^* \gamma) \psi - m \psi \eta = 0$	$i \gamma \wedge {}^* D \psi - \frac{i}{2} (D \wedge {}^* \gamma) \psi - m \psi \eta = 0$	$i \gamma \wedge {}^* D \psi - \frac{i}{2} (D \wedge {}^* \gamma) \psi - m \psi \eta = 0$
mielke-geometrodynamics_E00811 (11.4.23)	256	■ 0.840	$\frac{3}{8} \ell^2 \bar{\psi} \gamma^5 \psi = 4 \pi m e^{-\mu - v/2} \frac{2}{3} H F \left Y_{ \kappa - 1}^{ \kappa - 1}(\vartheta, \phi) \right ^2$	$\frac{3}{8} \ell^2 \bar{\psi} \gamma^5 \psi = 4 \pi m e^{-\mu - v/2} \frac{2}{3} H F \left Y_{ \kappa - 1}^{ \kappa - 1}(\vartheta, \phi) \right ^2$	$\frac{3}{8} \ell^2 \bar{\psi} \gamma^5 \psi = 4 \pi m e^{-\mu - v/2} \frac{2}{3} H F \left Y_{ \kappa - 1}^{ \kappa - 1}(\vartheta, \phi) \right ^2$
mielke-geometrodynamics_E00584 (9.1.2)	192	■ 0.843	$e^\nu = e^{-\lambda} = 1 - \left(\frac{2\mu}{r}\right)^{n-3} - \frac{2\Lambda_{\text{eff.}}}{(n-1)(n-2)} r^2$ $\mu = \frac{M}{m^{*2}}; \quad d\Omega^2 := d\vartheta^2 + \sin^2 \vartheta d\phi^2$	$e^\nu = e^{-\lambda} = 1 - \left(\frac{2\mu}{r}\right)^{n-3} - \frac{2\Lambda_{\text{eff.}}}{(n-1)(n-2)} r^2$ $\mu = \frac{M}{m^{*2}}; \quad d\Omega^2 := d\vartheta^2 + \sin^2 \vartheta d\phi^2$	$e^\nu = e^{-\lambda} = 1 - \left(\frac{2\mu}{r}\right)^{n-3} - \frac{2\Lambda_{\text{eff.}}}{(n-1)(n-2)} r^2$ $\mu = \frac{M}{m^{*2}}; \quad d\Omega^2 := d\vartheta^2 + \sin^2 \vartheta d\phi^2$
mielke-geometrodynamics_E00349 (5.4.7)	117	■ 0.845	$\langle dj_5 \rangle = 2im \langle \bar{\psi} \gamma_5 \psi \rangle \eta - \frac{1}{96\pi^2} \left[2R_{\alpha\beta}^{\{\} \} \wedge R^{\{\beta\alpha\beta\}} + d\mathcal{A} \wedge d\mathcal{A} \right]$	$\langle dj_5 \rangle = 2im \langle \bar{\psi} \gamma_5 \psi \rangle \eta - \frac{1}{96\pi^2} \left[2R_{\alpha\beta}^{\{\} \} \wedge R^{\{\beta\alpha\beta\}} + d\mathcal{A} \wedge d\mathcal{A} \right]$	$\langle dj_5 \rangle = 2im \langle \bar{\psi} \gamma_5 \psi \rangle \eta - \frac{1}{96\pi^2} \left[2R_{\alpha\beta}^{\{\} \} \wedge R^{\{\beta\alpha\beta\}} + d\mathcal{A} \wedge d\mathcal{A} \right]$
mielke-geometrodynamics_E00792 (11.4.4)	253	■ 0.845	$ds^2 = g_{kl} dx^k \otimes_s dx^l = e^\nu dt^2 - e^\mu (dr^2 + r^2 d\vartheta^2 + r^2 \sin^2 \vartheta d\phi^2).$	$ds^2 = g_{kl} dx^k \otimes_s dx^l = e^\nu dt^2 - e^\mu (dr^2 + r^2 d\vartheta^2 + r^2 \sin^2 \vartheta d\phi^2).$	$ds^2 = g_{kl} dx^k \otimes_s dx^l = e^\nu dt^2 - e^\mu (dr^2 + r^2 d\vartheta^2 + r^2 \sin^2 \vartheta d\phi^2).$
mielke-geometrodynamics_E00742 (11.1.9)	238	■ 0.846	$\Lambda : \begin{cases} \widetilde{SO}_\circ(s, n-s) & \longrightarrow SO_\circ(s, n-s), \\ \stackrel{\Psi}{\Psi} & \longrightarrow g^\alpha{}_\beta = -\frac{i}{N} \text{Tr} \left(\gamma^\alpha \text{Si} \gamma_\beta S^{-1} \right), \end{cases}$	$\Lambda : \begin{cases} \widetilde{SO}_\circ(s, n-s) & \longrightarrow SO_\circ(s, n-s), \\ \stackrel{\Psi}{\Psi} & \longrightarrow g^\alpha{}_\beta = -\frac{i}{N} \text{Tr} \left(\gamma^\alpha \text{Si} \gamma_\beta S^{-1} \right), \end{cases}$	$\Lambda : \begin{cases} \widetilde{SO}_\circ(s, n-s) & \longrightarrow SO_\circ(s, n-s), \\ \stackrel{\Psi}{\Psi} & \longrightarrow g^\alpha{}_\beta = -\frac{i}{N} \text{Tr} \left(\gamma^\alpha \text{Si} \gamma_\beta S^{-1} \right), \end{cases}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E0897 (13.5.2)	288	■ 0.846	$\begin{aligned} \hat{R} &:= R_{A}{}^B E^A{}_B := [d\Gamma_A{}^B - \Gamma_A{}^C \wedge \Gamma_C{}^B] E^A{}_B \\ &= \left[R_{\alpha}{}^{\beta} - \frac{\Lambda}{3} \theta_{\alpha} \wedge \vartheta^{\beta} \right] L^{\alpha}{}_{\beta} + T^{\beta} P_{\beta} + D\theta_{\alpha} P^{\alpha}_{*} \end{aligned}$	$\hat{R} := R_A{}^B E^A{}_B := [d\Gamma_A{}^B - \Gamma_A{}^C \wedge \Gamma_C{}^B] E^A{}_B = \left[R_{\alpha}{}^{\beta} - \frac{\Lambda}{3} \theta_{\alpha} \wedge \vartheta^{\beta} \right] L^{\alpha}{}_{\beta} + T^{\beta} P_{\beta} + D\theta_{\alpha} P^{\alpha}_{*}$	$\hat{R} := R_A{}^B \not{E}^A{}_B := \left[d\Gamma_A{}^B - \Gamma_A{}^C \wedge \Gamma_C{}^B \right] \not{E}^A{}_B = \left[R_{\alpha}{}^{\beta} - \frac{\Lambda}{3} \theta_{\alpha} \wedge \vartheta^{\beta} \right] L^{\alpha}{}_{\beta} + T^{\beta} P_{\beta} + D\theta_{\alpha} P^{\alpha}_{*},$
mielke-geometrodynamics_E08296 (4.5.7) [re-fined: census]	101	■ 0.847	$F_{\{\alpha\}} = \left(e_{\{\alpha\}} \rfloor D \Psi \right) \frac{\delta L}{\delta \Psi} + (-1)^p \left(e_{\{\alpha\}} \rfloor \Psi \right) \wedge D \frac{\delta L}{\delta \Psi}$	$F_{\alpha} = (e_{\alpha} \rfloor D\Psi) \frac{\delta L}{\delta \Psi} + (-1)^p (e_{\alpha} \rfloor \Psi) \wedge D \frac{\delta L}{\delta \Psi}$	$F_{\alpha} = (e_{\alpha} \rfloor D\Psi) \frac{\delta L}{\delta \Psi} + (-1)^p (e_{\alpha} \rfloor \Psi) \wedge D \frac{\delta L}{\delta \Psi}$
mielke-geometrodynamics_E08241 (4.3.6)	91	■ 0.848	$g_{\{\alpha\} \{\beta\}}(m, \vartheta) := \frac{1}{2} \frac{\partial^2 F^2(m, \vartheta)}{\partial \vartheta^{\alpha} \otimes_s \partial \vartheta^{\beta}}.$	$g_{\alpha\beta}(m, \vartheta) := \frac{1}{2} \frac{\partial^2 F^2(m, \vartheta)}{\partial \vartheta^{\alpha} \otimes_s \partial \vartheta^{\beta}}.$	$g_{\alpha\beta}(m, \vartheta) := \frac{1}{2} \frac{\partial^2 F^2(m, \vartheta)}{\partial \vartheta^{\alpha} \otimes_s \partial \vartheta^{\beta}}.$
mielke-geometrodynamics_E08478 (7.4.6) [re-fined: census]	157	■ 0.849	$\operatorname{R}\cancel{\mathrm{i}}_{\alpha\beta} := \operatorname{Ric}_{\alpha\beta} - \frac{1}{4} \operatorname{Rg}_{\alpha\beta} = 0.$	$R\cancel{c}_{\alpha\beta} := \operatorname{Ric}_{\alpha\beta} - \frac{1}{4} \operatorname{Rg}_{\alpha\beta} = 0.$	$Ri\cancel{c}_{\alpha\beta} := \operatorname{Ric}_{\alpha\beta} - \frac{1}{4} Rg_{\alpha\beta} = 0.$
mielke-geometrodynamics_E08295 (4.5.6) [re-fined: census]	101	■ 0.850	$\begin{aligned} D \Sigma_{\{\alpha\}} &\&= \left(e_{\{\alpha\}} \rfloor T^{\beta} \right) \wedge \Sigma_{\beta} + \left(e_{\{\alpha\}} \rfloor R^{\beta\gamma} \right) \wedge \tau_{\beta\gamma} + F_{\alpha} \\ &\simeq \left(e_{\alpha} \rfloor T^{\beta} \right) \wedge \Sigma_{\beta} + \left(e_{\alpha} \rfloor R^{\beta\gamma} \right) \wedge \tau_{\beta\gamma}. \end{aligned}$	$D\Sigma_{\alpha} = (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma} + F_{\alpha} \simeq (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma}.$	$D\Sigma_{\alpha} = (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma} + F_{\alpha} \simeq (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma}.$
mielke-geometrodynamics_E08842 (12.2.10) [re-fined: census]	272	■ 0.852	$\mu := e_{\{\alpha\}} \rfloor \mu^{\alpha} = \frac{3}{4} j_5$	$\mu := e_{\alpha} \rfloor \mu^{\alpha} = \frac{3}{4} j_5$	$\mu := e_{\alpha} \rfloor \mu^{\alpha} = \frac{3}{4} j_5$
mielke-geometrodynamics_E08586 (7.7.24)	164	■ 0.852	$\gamma_5 := (i/4!)^*(\gamma \wedge \gamma \wedge \gamma \wedge \gamma)$	$\gamma_5 := (i/4!)*(\gamma \wedge \gamma \wedge \gamma \wedge \gamma)$	$\gamma_5 := (i/4!)*(\gamma \wedge \gamma \wedge \gamma \wedge \gamma)$
mielke-geometrodynamics_E01814 (14.4.9)	326	■ 0.856	$M^{\{2\}} = \left(M_{\mathrm{ir}} + \frac{\alpha_e M^{*2}}{4 M_{\mathrm{ir}}} \operatorname{Tr} \bar{Q}^2 \right)^2 + \frac{1}{4} J(J+1) \frac{M^{*4}}{M_{\mathrm{ir}}^2},$	$M^2 = \left(M_{\mathrm{ir}} + \frac{\alpha_e M^{*2}}{4 M_{\mathrm{ir}}} \operatorname{Tr} \bar{Q}^2 \right)^2 + \frac{1}{4} J(J+1) \frac{M^{*4}}{M_{\mathrm{ir}}^2},$	$M^2 = \left(M_{\mathrm{ir}} + \frac{\alpha_e M^{*2}}{4 M_{\mathrm{ir}}} \operatorname{Tr} \bar{Q}^2 \right)^2 + \frac{1}{4} J(J+1) \frac{M^{*4}}{M_{\mathrm{ir}}^2},$
mielke-geometrodynamics_E08753 (11.1.20)	241	■ 0.859	$\hat{\Psi}_{ \widehat{SO}(1,3)} = \begin{bmatrix} \Psi_1 \\ \Psi_2 \\ \vdots \\ \Psi_{\lfloor \frac{n}{4} \rfloor} \end{bmatrix}.$	$\hat{\Psi}_{ \widehat{SO}(1,3)} = \begin{bmatrix} \Psi_1 \\ \Psi_2 \\ \vdots \\ \Psi_{\lfloor \frac{n}{4} \rfloor} \end{bmatrix}.$	$\hat{\Psi}_{ \widehat{SO}_s(1,3)} = \begin{bmatrix} \Psi_1 \\ \Psi_2 \\ \vdots \\ \Psi_{\lfloor \frac{n}{4} \rfloor} \end{bmatrix}.$
mielke-geometrodynamics_E08783 (11.3.2)	247	■ 0.862	$\begin{aligned} L_{\{\mathrm{D}\}} &\&= \frac{i}{2} \left(\bar{\psi} \gamma \wedge^* D^{\{\}} \psi + \left(\overline{D^{\{\}} \psi} \right) \wedge^* \gamma \psi \right) - m \bar{\psi} \psi \eta \\ &+ \frac{1}{2} \bar{\psi} \left[\tilde{K}, * \gamma \right] \psi =: L_{\mathrm{D}}^{\{\}} + L_{\mathrm{NL}}. \end{aligned}$	$L_{\mathrm{D}} = \frac{i}{2} \left(\bar{\psi} \gamma \wedge^* D^{\{\}} \psi + \left(\overline{D^{\{\}} \psi} \right) \wedge^* \gamma \psi \right) - m \bar{\psi} \psi \eta + \frac{1}{2} \bar{\psi} \left[\tilde{K}, * \gamma \right] \psi =: L_{\mathrm{D}}^{\{\}} + L_{\mathrm{NL}}.$	$L_{\mathrm{D}} = \frac{i}{2} \left(\bar{\psi} \gamma \wedge^* D^{\{\}} \psi + \left(\overline{D^{\{\}} \psi} \right) \wedge^* \gamma \psi \right) - m \bar{\psi} \psi \eta + \frac{1}{2} \bar{\psi} [\tilde{K}, * \gamma] \psi =: L_{\mathrm{D}}^{\{\}} + L_{\mathrm{NL}}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_369 EQ1100 (B.12)		■ 0.864	$\begin{aligned} \{ \}^{[*]} \alpha^{\{p\}} &:= \frac{1}{(n-p)! \cdot (p!)} \sqrt{ \det g_{ij} } \varepsilon_{\alpha_1 \dots \alpha_p \beta_1 \dots \beta_{n-p}} \\ &\quad \times A^{\alpha_1 \dots \alpha_p} \vartheta^{\beta_1} \wedge \dots \wedge \vartheta^{\beta_{n-p}}. \end{aligned}$	${}^* \alpha^{(p)} := \frac{1}{(n-p)! \cdot (p!)} \sqrt{ \det g_{ij} } \varepsilon_{\alpha_1 \dots \alpha_p \beta_1 \dots \beta_{n-p}} \times A^{\alpha_1 \dots \alpha_p} \vartheta^{\beta_1} \wedge \dots \wedge \vartheta^{\beta_{n-p}}.$	${}^* \alpha^{(p)} := \frac{1}{(n-p)! \cdot (p!)} \sqrt{ \det g_{ij} } \varepsilon_{\alpha_1 \dots \alpha_p \beta_1 \dots \beta_{n-p}} \times A^{\alpha_1 \dots \alpha_p} \vartheta^{\beta_1} \wedge \dots \wedge \vartheta^{\beta_{n-p}}.$
mielke-geometrodynamics_74 EQ0184 (3.7.14)		■ 0.864	$\vec{\omega} = d \vec{\omega} - (\overrightarrow{\omega \wedge})$	$\vec{\Omega} = d\vec{\omega} - (\overrightarrow{\omega \wedge})$	$\vec{\Omega} = d\vec{\omega} - (\overrightarrow{\omega \wedge}) = d\vec{\omega} - \vec{\omega} \wedge \vec{\omega}.$
mielke-geometrodynamics_69 EQ0165 (3.5.13)		■ 0.865	$\{ \}^{[*]} \omega = (\stackrel{+}{\omega}) \omega ;$	${}^* \Omega = (\stackrel{+}{-}) \Omega ;$	${}^* \Omega = (\stackrel{+}{-}) \Omega ;$
mielke-geometrodynamics_238 EQ0737 (11.1.4)		■ 0.867	$x^{\{\alpha\}} \rightarrow x^{\{\prime \alpha\}} = g_{\beta}^{\alpha} \quad x^{\beta}, [g_{\beta}^{\alpha}] \in SO(s, n-s), \quad x^{\alpha} \in \mathbf{E}^n.$	$x^{\alpha} \rightarrow x^{\prime \alpha} = g_{\beta}^{\alpha} \quad x^{\beta}, [g_{\beta}^{\alpha}] \in SO(s, n-s), \quad x^{\alpha} \in \mathbf{E}^n.$	$x^{\alpha} \rightarrow x^{\prime \alpha} = g_{\beta}^{\alpha} \quad x^{\beta}, [g_{\beta}^{\alpha}] \in SO(s, n-s), \quad x^{\alpha} \in \mathbf{E}^n.$
mielke-geometrodynamics_276 EQ0853 (12.3.6)		■ 0.868	$\left\langle \vartheta^{\alpha} \wedge \sigma_{\alpha} \right\rangle = -\frac{1}{3\pi^2} \left[\text{Tr} (G \wedge {}^* G) + \frac{1}{24} \left(2R^{\alpha\beta\gamma\delta} \wedge R_{\alpha\beta}^{\{\gamma\delta\}} + d\mathcal{A} \wedge {}^* d\mathcal{A} \right) \right],$	$\langle \vartheta^{\alpha} \wedge \sigma_{\alpha} \rangle = -\frac{1}{3\pi^2} \left[\text{Tr} (G \wedge {}^* G) + \frac{1}{24} \left(2R^{\alpha\beta\gamma\delta} \wedge R_{\alpha\beta}^{\{\gamma\delta\}} + d\mathcal{A} \wedge {}^* d\mathcal{A} \right) \right],$	$\langle \vartheta^{\alpha} \wedge \sigma_{\alpha} \rangle = -\frac{1}{3\pi^2} \left[\text{Tr} (G \wedge {}^* G) + \frac{1}{24} \left(2R^{\alpha\beta\gamma\delta} \wedge R_{\alpha\beta}^{\{\gamma\delta\}} + d\mathcal{A} \wedge {}^* d\mathcal{A} \right) \right],$
mielke-geometrodynamics_245 EQ0774 (11.2.3)		■ 0.868	$i(\overline{D\psi}) \wedge {}^* \gamma + \frac{i}{2} \bar{\psi} D \wedge {}^* \gamma + m \bar{\psi} \eta = 0,$	$i(\overline{D\psi}) \wedge {}^* \gamma + \frac{i}{2} \bar{\psi} D \wedge {}^* \gamma + m \bar{\psi} \eta = 0,$	$i(\overline{D\psi}) \wedge {}^* \gamma + \frac{i}{2} \bar{\psi} D \wedge {}^* \gamma + m \bar{\psi} \eta = 0,$
mielke-geometrodynamics_178 EQ0548 (8.4.2)		■ 0.871	$s_{L_{\text{Pontr}}} = -2 d \operatorname{Tr} \left(\Psi \wedge \Omega^g \right) = -d \left(\Psi_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha} \right).$	$s_{L_{\text{Pontr}}} = -2d \operatorname{Tr} \{ \Psi \wedge \Omega^g \} = -d \left(\Psi_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha} \right).$	$s_{L_{\text{Pontr}}} = -2d \operatorname{Tr} \{ \Psi \wedge \Omega^g \} = -d \left(\Psi_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha} \right).$
mielke-geometrodynamics_91 EQ0244 (4.3.9)		■ 0.873	$g_{\{i\}j} e_{\alpha}^{\{i\}} \cdot e_{\beta}^{\{j\}} = g_{\alpha\beta}.$	$g_{ij} e_{\alpha}^i e_{\beta}^j = g_{\alpha\beta}.$	$g_{ij} e_{\alpha}^i e_{\beta}^j = g_{\alpha\beta}.$
mielke-geometrodynamics_287 EQ0890 (13.4.9)		■ 0.876	$\hat{\xi} = \begin{pmatrix} \xi \\ 1 \end{pmatrix} = \begin{pmatrix} \xi^{\alpha} P_{\alpha} \\ 1 \end{pmatrix}, \quad \hat{\xi}_{*} = (\xi_{*}, 1) = (\xi_{\beta}^{*} P_{*}^{\beta}, 1)$	$\hat{\xi} = \begin{pmatrix} \xi \\ 1 \end{pmatrix} = \begin{pmatrix} \xi^{\alpha} P_{\alpha} \\ 1 \end{pmatrix}, \quad \hat{\xi}_{*} = (\xi_{*}, 1) = (\xi_{\beta}^{*} P_{*}^{\beta}, 1)$	$\hat{\xi} = \begin{pmatrix} \xi \\ 1 \end{pmatrix} = \begin{pmatrix} \xi^{\alpha} P_{\alpha} \\ 1 \end{pmatrix}, \quad \hat{\xi}_{*} = (\xi_{*}, 1) = (\xi_{\beta}^{*} P_{*}^{\beta}, 1)$
mielke-geometrodynamics_325 EQ1007 (14.4.2)		■ 0.877	$\begin{aligned} A^{(\circ)} &= \sigma^{(\circ)} \omega^{(\circ)} = -\frac{r^2}{\rho^2} \left[dt - J_3 \frac{\hbar}{Mc} \sin^2 \vartheta d\phi \right] Q^{(\circ)} \cos \delta \\ &\quad - \frac{\cos \vartheta}{\rho^2} \left[\left(r^2 + J(J+1) \frac{\hbar^2}{M^2 c^2} \right) d\phi - J_3 \frac{\hbar}{M} dt \right] P^{(\circ)} \sin \delta. \end{aligned}$	$A^{(\circ)} = \sigma^{(\circ)} \omega^{(\circ)} = -\frac{r^2}{\rho^2} \left[dt - J_3 \frac{\hbar}{Mc} \sin^2 \vartheta d\phi \right] Q^{(\circ)} \cos \delta - \frac{\cos \vartheta}{\rho^2} \left[\left(r^2 + J(J+1) \frac{\hbar^2}{M^2 c^2} \right) d\phi - J_3 \frac{\hbar}{M} dt \right] P^{(\circ)} \sin \delta$	$A^{(\circ)} = \sigma^{(\circ)} \omega^{(\circ)} = -\frac{r^2}{\rho^2} \left[dt - J_3 \frac{\hbar}{Mc} \sin^2 \vartheta d\phi \right] Q^{(\circ)} \cos \delta - \frac{\cos \vartheta}{\rho^2} \left[\left(r^2 + J(J+1) \frac{\hbar^2}{M^2 c^2} \right) d\phi - J_3 \frac{\hbar}{M} dt \right] P^{(\circ)} \sin \delta.$
mielke-geometrodynamics_355 EQ1062 (16.2.1)		■ 0.877	$\left[\square - (mc/\hbar)^2 \right] \varphi = 0$	$\left[\square - (mc/\hbar)^2 \right] \varphi = 0$	$\left[\square - (mc/\hbar)^2 \right] \varphi = 0$
mielke-geometrodynamics_44 EQ0051 (2.7.1)		■ 0.881	$G(p) : P \rightarrow P, \quad G(p) \in \mathcal{G}_p$	$G(p) : P \rightarrow P, \quad G(p) \in \mathcal{G}_p$	$G(p) : P \rightarrow P, \quad G(p) \in \mathcal{G}_p$
mielke-geometrodynamics_72 EQ0178 (3.7.8)		■ 0.881	$\left\langle O \right ^{\lambda} \varphi \left O \right\rangle = \mu / \sqrt{\lambda}$	$\left\langle O \right ^{\lambda} \varphi \left O \right\rangle = \mu / \sqrt{\lambda}$	$\left\langle O \right ^{\lambda} \varphi \left O \right\rangle = \mu / \sqrt{\lambda}$
mielke-geometrodynamics_40 EQ0028 (2.5.1)		■ 0.885	$V_{\{p\}}^{[*]} = T_{\{p\}}^{[*]} \left(F_{\{m\}}(M) \right),$	$V_p^{*} = T_p^{*} (F_m(M)),$	$V_p^{*} = T_p^{*} (F_m(M)),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_295 E09926 (13.8.6)		■ 0.888	$\begin{aligned} & d\,C_{\mathrm{RR}(\star)} \, \& := \frac{1}{2} \\ & d\left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta(\star)} - \frac{1}{3}\Gamma_{\alpha}{}^{\beta(\star)} \wedge \Gamma_{\beta}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\alpha}\right) \\ & \equiv -L_{\mathrm{SKY}} - 2\,\mathrm{Ric}_{\alpha\beta} \wedge {}^*\mathrm{Ric}^{\alpha\beta} + \frac{1}{2}\,\mathrm{Ric}_{\alpha}{}^{\alpha} \wedge {}^*\mathrm{Ric}_{\beta}{}^{\beta} \end{aligned}$	$dC_{\mathrm{RR}(\star)} := \frac{1}{2}d\left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta(\star)} - \frac{1}{3}\Gamma_{\alpha}{}^{\beta(\star)} \wedge \Gamma_{\beta}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\alpha}\right) \\ \equiv -L_{\mathrm{SKY}} - 2\,\mathrm{Ric}_{\alpha\beta} \wedge {}^*\mathrm{Ric}^{\alpha\beta} + \frac{1}{2}\,\mathrm{Ric}_{\alpha}{}^{\alpha} \wedge {}^*\mathrm{Ric}_{\beta}{}^{\beta}$	$dC_{\mathrm{RR}^{(\star)}} := \frac{1}{2}d\left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta(\star)} - \frac{1}{3}\Gamma_{\alpha}{}^{\beta(\star)} \wedge \Gamma_{\beta}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\alpha}\right) \\ \equiv -L_{\mathrm{SKY}} - 2\mathrm{Ric}_{\alpha\beta} \wedge {}^*\mathrm{Ric}^{\alpha\beta} + \frac{1}{2}\mathrm{Ric}_{\alpha}{}^{\alpha} \wedge {}^*\mathrm{Ric}_{\beta}{}^{\beta}$
mielke-geometrodynamics_220 E09674 (10.3.31)		■ 0.890	$\check{\mathrm{E}}_{\alpha} = -\frac{\theta_{\mathrm{T}}}{2\ell^2} \left(\frac{1}{\theta_{\mathrm{TL}}} d\vartheta_{\alpha} - \frac{4\ell}{\theta_{\mathrm{T}}\theta_{\mathrm{L}}} \check{\mathcal{J}}_{\alpha}^{\star} \right),$	$\check{\mathcal{E}}_{\alpha} = -\frac{\theta_{\mathrm{T}}\theta_{\mathrm{TL}}}{2\ell^2} \left(\frac{1}{\theta_{\mathrm{TL}}} d\vartheta_{\alpha} - \frac{4\ell}{\theta_{\mathrm{T}}\theta_{\mathrm{L}}} \check{\mathcal{J}}_{\alpha}^{\star} \right),$	$\check{\mathcal{E}}_{\alpha} = -\frac{\theta_{\mathrm{T}}\theta_{\mathrm{TL}}}{2\ell^2} \left(\frac{1}{\theta_{\mathrm{TL}}} d\vartheta_{\alpha} - \frac{4\ell}{\theta_{\mathrm{T}}\theta_{\mathrm{L}}} \check{\mathcal{J}}_{\alpha}^{\star} \right),$
mielke-geometrodynamics_169 E09531 (7.7.51) [re-fined: census]		■ 0.890	$R := e_{\beta} \rfloor e_{\alpha} \rfloor R^{\alpha\beta}, \quad {}^*R^{\alpha\beta} \wedge \vartheta_{\alpha} \wedge \vartheta_{\beta} \equiv R^{\alpha\beta} \wedge \eta_{\alpha\beta} \equiv -R\eta,$	$R := e_{\beta} \rfloor e_{\alpha} \rfloor R^{\alpha\beta}, \quad {}^*R^{\alpha\beta} \wedge \vartheta_{\alpha} \wedge \vartheta_{\beta} \equiv R^{\alpha\beta} \wedge \eta_{\alpha\beta} \equiv -R\eta,$	$R := e_{\beta} \rfloor e_{\alpha} \rfloor R^{\alpha\beta}, \quad {}^*R^{\alpha\beta} \wedge \vartheta_{\alpha} \wedge \vartheta_{\beta} \equiv R^{\alpha\beta} \wedge \eta_{\alpha\beta} \equiv -R\eta,$
mielke-geometrodynamics_176 E09540 (8.3.1)		■ 0.891	$\begin{aligned} & (d \oplus s) \gamma + [\widetilde{\Gamma}, \gamma] = \widetilde{\Theta} \\ & (d \oplus s) \widetilde{\Theta} + [\widetilde{\Gamma}, \widetilde{\Theta}] \equiv [\widetilde{\Omega}^g, \gamma] \end{aligned}$	$\begin{aligned} & (d \oplus s) \gamma + [\widetilde{\Gamma}, \gamma] = \widetilde{\Theta} \\ & (d \oplus s) \widetilde{\Theta} + [\widetilde{\Gamma}, \widetilde{\Theta}] \equiv [\widetilde{\Omega}^g, \gamma] \end{aligned}$	$\begin{aligned} & (d \oplus s) \gamma + [\widetilde{\Gamma}, \gamma] = \widetilde{\Theta}, \\ & (d \oplus s) \widetilde{\Theta} + [\widetilde{\Gamma}, \widetilde{\Theta}] \equiv [\widetilde{\Omega}^g, \gamma], \end{aligned}$
mielke-geometrodynamics_303 E09925 (14.1.5)		■ 0.892	$\begin{aligned} & D^{\star} \Omega = \tau_e, \\ & D\Omega \equiv 0, \\ & \frac{1}{2} \star (\vartheta \wedge \star \Omega^g) = \ell^{\star 2} (\Sigma_{\mathrm{YM}} + \Sigma_{\mathrm{mat}}), \end{aligned}$	$\begin{aligned} & D^{\star} \Omega = \tau_e, \\ & D\Omega \equiv 0, \\ & \frac{1}{2} \star (\vartheta \wedge \star \Omega^g) = \ell^{\star 2} (\Sigma_{\mathrm{YM}} + \Sigma_{\mathrm{mat}}), \end{aligned}$	$\begin{aligned} & D^{\star} \Omega = \tau_e, \\ & D\Omega \equiv 0, \\ & \frac{1}{2} \star (\vartheta \wedge \star \Omega^g) = \ell^{\star 2} (\Sigma_{\mathrm{YM}} + \Sigma_{\mathrm{mat}}), \end{aligned}$
mielke-geometrodynamics_283 E09867 (13.2.3)		■ 0.893	$B \cong dA, \quad dB \cong dF \equiv 0,$	$B \cong dA, \quad dB \cong dF \equiv 0,$	$B \cong dA, \quad dB \cong dF \equiv 0,$
mielke-geometrodynamics_201 E09609 (9.2.11)		■ 0.894	$\begin{aligned} & \left(1 + \zeta^2\right) \int_{M^4} L_{\mathrm{SKY}} - (-1)^s \zeta (4\pi)^2 \cdot \chi(M^4) \\ & = \frac{1}{4} \int_{M^4} \left(\Omega^g - \zeta^{\star} \Omega^{g(\star)}\right) \wedge {}^{\star} \left(\Omega^g - \zeta^{\star} \Omega^{g(\star)}\right) \geq 0, \end{aligned}$	$\begin{aligned} & \left(1 + \zeta^2\right) \int_{M^4} L_{\mathrm{SKY}} - (-1)^s \zeta (4\pi)^2 \cdot \chi(M^4) \\ & = \frac{1}{4} \int_{M^4} \left(\Omega^g - \zeta^{\star} \Omega^{g(\star)}\right) \wedge {}^{\star} \left(\Omega^g - \zeta^{\star} \Omega^{g(\star)}\right) \geq 0, \end{aligned}$	$\begin{aligned} & (1 + \zeta^2) \int_{M^4} L_{\mathrm{SKY}} - (-1)^s \zeta (4\pi)^2 \cdot \chi(M^4) \\ & = \frac{1}{4} \int_{M^4} (\Omega^g - \zeta^{\star} \Omega^{g(\star)}) \wedge {}^{\star} (\Omega^g - \zeta^{\star} \Omega^{g(\star)}) \geq 0, \end{aligned}$
mielke-geometrodynamics_309 E09959 (14.1.41)		■ 0.894	$\oint d\delta = 2\pi b_1 = 2\pi v$	$\oint d\delta = 2\pi b_1 = 2\pi v$	$\oint d\delta = 2\pi b_1 = 2\pi v$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_102 E00299 (4.5.10)	102	0.896	$\begin{aligned} & \& \delta g_{\alpha\beta} = 2\varepsilon_{(\alpha}{}^{\gamma} g_{\beta)\gamma}, \quad \delta\vartheta^{\alpha} = -\vartheta^{\beta}\varepsilon_{\beta}{}^{\alpha}, \\ & \delta\Gamma_{\alpha}{}^{\beta} = D\varepsilon_{\alpha}{}^{\beta}, \quad \delta\Psi = -\varepsilon_{\alpha}{}^{\beta} I^{\alpha}{}_{\beta}\Psi. \end{aligned}$		
mielke-geometrodynamics_90 E00236 (4.3.1)	90	0.896	$\begin{aligned} (\Delta s)^2 &= -c^2(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 \\ &= g_{ij}\Delta x^i\Delta x^j. \end{aligned}$		
mielke-geometrodynamics_323 E01005 (14.3.18)	323	0.896	$\begin{aligned} \not{D}^2 &= \operatorname{Tr}\left(\frac{1}{2}\gamma\wedge\gamma\wedge^*\{D,D\} + \frac{1}{2}\gamma\wedge\gamma\wedge^*\tilde{\Omega} - \gamma\wedge\tilde{\Theta}\wedge^*D\right) \\ &= \left(\Box^g + \frac{1}{4}R\right)\eta - \vartheta\wedge\Theta\wedge^*D \end{aligned}$		
mielke-geometrodynamics_88 E00230 (4.2.23)	88	0.899	$\stackrel{*}{\sigma}\Theta = \frac{1}{2}T_{\alpha\beta}^cP_c\otimes\vartheta^{\alpha}\wedge\vartheta^{\beta} = \frac{1}{2}T_{ij}^cP_c\otimes dx^i\wedge dx^j,$		
mielke-geometrodynamics_230 E00723 (10.10.3) [re-fined: census]	230	0.900	$\begin{aligned} \delta_{\text{susy}}\vartheta^{\alpha} &= i\bar{\sigma}\Psi\gamma^{\alpha}, \quad \delta_{\text{susy}}\Gamma_{\alpha}^{\star} = i\bar{\sigma}\gamma_{\alpha}^{\star}D\Psi + ic\bar{\sigma}(\gamma_{\alpha}\Psi + e_{\alpha}\lceil^*\Psi) \\ \delta_{\text{susy}}\Psi &= 2D\sigma + c\gamma\sigma, \end{aligned}$		
mielke-geometrodynamics_86 E00216 (4.2.8)	86	0.902	$\widetilde{\omega} \rightarrow^{A^{-1}} \widetilde{\bar{\omega}} = A^{-1}\widetilde{\omega}A - A^{-1}dA, \quad A \in \mathscr{A}_p.$		
mielke-geometrodynamics_340 E01042 (15.1.19)	340	0.902	$\Gamma_{\mu\alpha\beta}:(2^+)\oplus 2(1^-)\oplus(0^+)\oplus(2^-)\oplus 2(1^+)\oplus(0^-),$		
mielke-geometrodynamics_157 E00481 (7.5.2)	157	0.903	$\square^c g_{ij} = \square^c \left(\Omega^2 o_{ij}^{\text{Mink}} \right) = \Omega^6 \square^c o_{ij}^{\text{Mink}} = 0$		
mielke-geometrodynamics_59 E00117 (3.3.18)	59	0.905	$L_{\text{Max}} := -\frac{1}{2}F\wedge^*F = \frac{1}{2}(\mathbf{E}^2 - \mathbf{B}^2)\eta,$		
mielke-geometrodynamics_371 E01110 (B.22)	371	0.905	$\Box\alpha = 0,$		
mielke-geometrodynamics_270 E00831 (12.1.3)	270	0.907	$\tilde{j}_5 := j_5 + (1/96\pi^2)A\wedge dA$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_136 EQ0409 (6.4.29)		■ 0.907	$T^A=\frac{1}{f}\left(d\,f\wedge\vartheta^A-2\,\eta^A\wedge\vartheta^0\wedge\vartheta^A\wedge\vartheta^0\wedge\vartheta^A\right)=-\frac{2}{f}\,\eta^A\wedge\vartheta^0\wedge\vartheta^A\wedge\vartheta^0\wedge\vartheta^A,$	$T^A=\frac{1}{f}\left(df\wedge\vartheta^A-2\eta^{0A\beta\gamma}\vartheta_\beta\wedge\vartheta_\gamma\right)=-\frac{2}{f}\eta^A{}_{\mu\nu}\vartheta^\mu\wedge\vartheta^\nu=\pm^*T^A,\quad T^{\hat{0}}=0,$	$T^A=\frac{1}{f}\left(df\wedge\vartheta^A-2\eta^{0A\beta\gamma}\vartheta_\beta\wedge\vartheta_\gamma\right)=-\frac{2}{f}\eta^A{}_{\mu\nu}\vartheta^\mu\wedge\vartheta^\nu=\pm^*T^A,\quad T^{\hat{0}}=0,$
mielke-geometrodynamics_217 EQ0657 (10.3.14) [re-fined: census]		■ 0.907	$E_{\alpha}=e_{\alpha}\rfloor L(\star)-\frac{2}{\delta\ell^2}\Gamma_{\alpha},$	$E_{\alpha}=e_{\alpha}\rfloor L(\star)-\frac{2\gamma}{\delta\ell^2}T_{\alpha},$	$E_{\alpha}=e_{\alpha}\rfloor L(\star)-\frac{2\gamma}{\delta\ell^2}T_{\alpha},$
mielke-geometrodynamics_132 EQ0396 (6.4.11)		■ 0.909	$\begin{aligned} \mathcal{H} &\cong n^{\alpha}\frac{\delta L}{\delta\vartheta^{\alpha}}+\Gamma_{\perp}^{\alpha\beta}\frac{\delta L}{\delta\Gamma^{\alpha\beta}} \\ &\quad -d\left[(n\rfloor\Psi)\wedge\frac{\partial L}{\partial D\Psi}+n^{\alpha}\frac{\partial L}{\partial T^{\alpha}}+\Gamma_{\perp}^{\alpha\beta}\frac{\partial L}{\partial R^{\alpha\beta}}\right]. \end{aligned}$	$\mathcal{H}\cong n^{\alpha}\frac{\delta L}{\delta\vartheta^{\alpha}}+\Gamma_{\perp}^{\alpha\beta}\frac{\delta L}{\delta\Gamma^{\alpha\beta}}-d\left[(n\rfloor\Psi)\wedge\frac{\partial L}{\partial D\Psi}+n^{\alpha}\frac{\partial L}{\partial T^{\alpha}}+\Gamma_{\perp}^{\alpha\beta}\frac{\partial L}{\partial R^{\alpha\beta}}\right].$	$\mathcal{H}\cong n^{\alpha}\frac{\delta L}{\delta\vartheta^{\alpha}}+\Gamma_{\perp}^{\alpha\beta}\frac{\delta L}{\delta\Gamma^{\alpha\beta}}-d\left[(n\rfloor\Psi)\wedge\frac{\partial L}{\partial D\Psi}+n^{\alpha}\frac{\partial L}{\partial T^{\alpha}}+\Gamma_{\perp}^{\alpha\beta}\frac{\partial L}{\partial R^{\alpha\beta}}\right].$
mielke-geometrodynamics_213 EQ0641 (10.2.12)		■ 0.909	$ds^2=-\left(1-\Lambda_{\text{eff}}\right)r^2dt^2+\left(1-\Lambda_{\text{eff}}\right)r^2dr^2+r^2d\phi^2$	$ds^2=-\left(1-\Lambda_{\text{eff}}\,r^2\right)dt^2+\left(1-\Lambda_{\text{eff}}\,r^2\right)^{-1}dr^2+r^2d\phi^2$	$ds^2=-\left(1-\Lambda_{\text{eff}}\,r^2\right)dt^2+\left(1-\Lambda_{\text{eff}}\,r^2\right)^{-1}dr^2+r^2d\phi^2$
mielke-geometrodynamics_94 EQ0263 (4.3.28)		■ 0.912	$\Gamma_{ij}^{\cdot k}=e_b^{\kappa}\Gamma_{i\alpha}^{\cdot b}E_j^{\alpha}+e_{\beta}^{\kappa}\partial_iE_{\cdot j}^{\beta}$	$\Gamma_{ij}^{\cdot k}=e_b^{\kappa}\Gamma_{i\alpha}^{\cdot b}E_j^{\alpha}+e_{\beta}^{\kappa}\partial_iE_{\cdot j}^{\beta}$	$\Gamma_{ij}^{\cdot \kappa}=e_b^{\kappa}\Gamma_{i\alpha}^{\cdot b}E_j^{\alpha}+e_{\beta}^{\kappa}\partial_iE_{\cdot j}^{\beta}$
mielke-geometrodynamics_126 EQ0364 (6.2.16)		■ 0.912	$D\,D\,\lambda_{\alpha\beta}=-2\,R_{[\alpha}\wedge\lambda_{\beta]}=0$	$DD\lambda_{\alpha\beta}=-2R_{[\alpha]}{}^{\gamma}\wedge\lambda_{\gamma \beta]}=0$	$DD\lambda_{\alpha\beta}=-2R_{[\alpha]}{}^{\gamma}\wedge\lambda_{\gamma \beta]}=0$
mielke-geometrodynamics_272 EQ0040 (12.2.8)		■ 0.916	$\begin{aligned} \tau_{\alpha\beta} &:=\frac{\partial L_D}{\partial\Gamma^{\alpha\beta}}=\frac{1}{8}\bar{\psi}\left(\gamma^{\alpha}\sigma_{\alpha\beta}+\sigma_{\alpha\beta}\gamma^{\alpha}\right)\psi \\ &=\frac{1}{4}\eta_{\alpha\beta\gamma\delta}\bar{\psi}\gamma^{\delta}\gamma_5\psi\eta^{\gamma}=\tau_{\alpha\beta\gamma}\eta^{\gamma}=\vartheta_{[\alpha}\wedge\mu_{\beta]} \end{aligned}$	$\tau_{\alpha\beta}:=\frac{\partial L_D}{\partial\Gamma^{\alpha\beta}}=\frac{1}{8}\bar{\psi}\left(\gamma^{\alpha}\sigma_{\alpha\beta}+\sigma_{\alpha\beta}\gamma^{\alpha}\right)\psi=\frac{1}{4}\eta_{\alpha\beta\gamma\delta}\bar{\psi}\gamma^{\delta}\gamma_5\psi\eta^{\gamma}=\tau_{\alpha\beta\gamma}\eta^{\gamma}=\vartheta_{[\alpha}\wedge\mu_{\beta]}$	$\tau_{\alpha\beta}:=\frac{\partial L_D}{\partial\Gamma^{\alpha\beta}}=\frac{1}{8}\bar{\psi}\left(\gamma^{\alpha}\sigma_{\alpha\beta}+\sigma_{\alpha\beta}\gamma^{\alpha}\right)\psi=\frac{1}{4}\eta_{\alpha\beta\gamma\delta}\bar{\psi}\gamma^{\delta}\gamma_5\psi\eta^{\gamma}=\tau_{\alpha\beta\gamma}\eta^{\gamma}=\vartheta_{[\alpha}\wedge\mu_{\beta]}.$
mielke-geometrodynamics_88 EQ0232 (4.2.25)		■ 0.917	$\Theta\rightarrow^{A^{-1}}\Theta=G^{-1}\Theta,$	$\Theta\rightarrow^{A^{-1}}\Theta=G^{-1}\Theta,$	$\Theta\rightarrow^{A^{-1}}\Theta=G^{-1}\Theta,$
mielke-geometrodynamics_185 EQ0577 (8.6.15)		■ 0.920	$-E_{\alpha}=\frac{\theta_{\mathrm{T}}^*}{2}R^{\{\beta\gamma}\wedge\eta_{\alpha\beta\gamma}-\theta_{\mathrm{T}}^*\Lambda_{\mathrm{eff}}\eta_{\alpha}=\ell^2\widehat{\Sigma}_{\alpha}.$	$-E_{\alpha}=\frac{\theta_{\mathrm{T}}^*}{2}R^{\{\beta\gamma}\wedge\eta_{\alpha\beta\gamma}-\theta_{\mathrm{T}}^*\Lambda_{\mathrm{eff}}\eta_{\alpha}=\ell^2\widehat{\Sigma}_{\alpha}.$	$-E_{\alpha}=\frac{\theta_{\mathrm{T}}^*}{2}R^{\{\beta\gamma}\wedge\eta_{\alpha\beta\gamma}-\theta_{\mathrm{T}}^*\Lambda_{\mathrm{eff}}\eta_{\alpha}=\ell^2\widehat{\Sigma}_{\alpha}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_337 EQ1031 (15.1.8)		■ 0.920	$\begin{aligned} & \frac{1}{2} * \left(\vartheta^f \wedge * \Omega^f \right) - a \ell^2 \vartheta^f \wedge * \left\{ 2L_{fg} - \frac{1}{2b\ell^{*2}} (\Omega^g \wedge \vartheta^g \wedge \vartheta^g + 2\Lambda\eta) \right\} \\ & = \ell^2 * \left\{ \Sigma_{\text{m}}(\text{ hadrons }) + \frac{a}{b} * (\vartheta^g \wedge * \Sigma_{\text{m}}(\text{ leptons })) \wedge \vartheta^f \right\} \end{aligned}$	$\frac{1}{2} * \left(\vartheta^f \wedge * \Omega^f \right) - a \ell^2 \vartheta^f \wedge * \left\{ 2L_{fg} - \frac{1}{2b\ell^{*2}} (\Omega^g \wedge \vartheta^g \wedge \vartheta^g + 2\Lambda\eta) \right\} \\ = \ell^2 * \left\{ \Sigma_{\text{m}}(\text{ hadrons }) + \frac{a}{b} * (\vartheta^g \wedge * \Sigma_{\text{m}}(\text{ leptons })) \wedge \vartheta^f \right\}$	$\frac{1}{2} * (\vartheta^f \wedge * \Omega^f) - a \ell^2 \vartheta^f \wedge * \left\{ 2L_{fg} - \frac{1}{2b\ell^{*2}} (\Omega^g \wedge \vartheta^g \wedge \vartheta^g + 2\Lambda\eta) \right\} \\ = \ell^2 * \left\{ \Sigma_{\text{m}}(\text{hadrons}) + \frac{a}{b} * (\vartheta^g \wedge * \Sigma_{\text{m}}(\text{leptons})) \wedge \vartheta^f \right\}$
mielke-geometrodynamics_156 EQ0473 (7.4.1)		■ 0.921	$\begin{aligned} & \int \mathcal{D}\Gamma \exp \left[\int L_{\text{SKY}}^{(*)} / \hbar \right] \\ & = \int \mathcal{D}\Gamma \exp \left[- \int \left(R_{\alpha\beta} + *R_{\alpha\beta}^{(*)} \right) \wedge * \left(R^{\alpha\beta} + *R^{\alpha\beta(*)} \right) / 4\hbar \right] \end{aligned}$	$\int \mathcal{D}\Gamma \exp \left[\int L_{\text{SKY}}^{(*)} / \hbar \right] \\ = \int \mathcal{D}\Gamma \exp \left[- \int \left(R_{\alpha\beta} + *R_{\alpha\beta}^{(*)} \right) \wedge * \left(R^{\alpha\beta} + *R^{\alpha\beta(*)} \right) / 4\hbar \right]$	$\int \mathcal{D}\Gamma \exp \left[\int L_{\text{SKY}}^{(*)} / \hbar \right] \\ = \int \mathcal{D}\Gamma \exp \left[- \int \left(R_{\alpha\beta} + *R_{\alpha\beta}^{(*)} \right) \wedge * \left(R^{\alpha\beta} + *R^{\alpha\beta(*)} \right) / 4\hbar \right]$
mielke-geometrodynamics_115 EQ0343 (5.4.1)		■ 0.922	$k \partial_k \Gamma_k = \frac{1}{2} \text{S Tr} \left\{ \left[\Gamma_k^{(2)} + R_k \right]^{-1} (k \partial_k R_k) \right\},$	$k \partial_k \Gamma_k = \frac{1}{2} \text{S Tr} \left\{ \left[\Gamma_k^{(2)} + R_k \right]^{-1} (k \partial_k R_k) \right\},$	$k \partial_k \Gamma_k = \frac{1}{2} \text{STr} \left\{ [\Gamma_k^{(2)} + R_k]^{-1} (k \partial_k R_k) \right\},$
mielke-geometrodynamics_269 EQ0829 (12.1.1)		■ 0.922	$\bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \rightarrow \bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \mathbf{P} \exp \left\{ i \int_x^{x+\varepsilon} A \right\}.$	$\bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \rightarrow \bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \mathbf{P} \exp \left\{ i \int_x^{x+\varepsilon} A \right\}.$	$\bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \rightarrow \bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \mathbf{P} \exp \left\{ i \int_x^{x+\varepsilon} A \right\}.$
mielke-geometrodynamics_238 EQ0736 (11.1.3)		■ 0.922	$\gamma^0 := \begin{pmatrix} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{pmatrix}, \quad \gamma^k := \begin{pmatrix} 0 & \sigma^k \\ -\sigma^k & 0 \end{pmatrix}, \quad \sigma^k : \text{Pauli matrices.}$	$\gamma^0 := \begin{pmatrix} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{pmatrix}, \quad \gamma^k := \begin{pmatrix} 0 & \sigma^k \\ -\sigma^k & 0 \end{pmatrix}, \quad \sigma^k : \text{Pauli matrices.}$	$\gamma^0 := \begin{pmatrix} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{pmatrix}, \quad \gamma^k := \begin{pmatrix} 0 & \sigma^k \\ -\sigma^k & 0 \end{pmatrix}, \quad \sigma^k : \text{Pauli matrices.}$
mielke-geometrodynamics_217 EQ0661 (10.3.18)		■ 0.923	$\square H - (-1)^s * D^* D H = dt + \frac{\gamma}{\delta \ell^2} d\mathcal{A} + \frac{1}{\delta \ell} dh,$	$\square H - (-1)^s * D^* D H = dt + \frac{\gamma}{\delta \ell^2} d\mathcal{A} + \frac{1}{\delta \ell} dh,$	$\square H - (-1)^s * D^* D H = dt + \frac{\gamma}{\delta \ell^2} d\mathcal{A} + \frac{1}{\delta \ell} dh,$
mielke-geometrodynamics_222 EQ0685 (10.3.43)		■ 0.923	$C_\alpha - E_\alpha = \Sigma_\alpha - D^{\{1\}} \mu_\alpha,$	$C_\alpha - E_\alpha = \Sigma_\alpha - D^{\{1\}} \mu_\alpha,$	$C_\alpha - E_\alpha = \Sigma_\alpha - D^{\{1\}} \mu_\alpha,$
mielke-geometrodynamics_225 EQ0702 (10.6.1)		■ 0.924	$L_A = \frac{1}{2} (F \wedge *F + m_{\text{photon}} F \wedge A).$	$L_A = \frac{1}{2} (F \wedge *F + m_{\text{photon}} F \wedge A).$	$L_A = \frac{1}{2} (F \wedge *F + m_{\text{photon}} F \wedge A).$
mielke-geometrodynamics_87 EQ0221 (4.2.14)		■ 0.924	$\Omega^{\text{T}} \rightarrow {}^{A^{-1}} \Omega^{\text{T}} = G^{-1} (\Omega^{\text{T}} - D D T) = G^{-1} \Omega^{\text{T}} - G^{-1} \Omega^{\text{L}} \otimes T.$	$\Omega^{\text{T}} \rightarrow {}^{A^{-1}} \Omega^{\text{T}} = G^{-1} (\Omega^{\text{T}} - D D T) = G^{-1} \Omega^{\text{T}} - G^{-1} \Omega^{\text{L}} \otimes T.$	$\Omega^{\text{T}} \rightarrow {}^{A^{-1}} \Omega^{\text{T}} = G^{-1} (\Omega^{\text{T}} - D D T) = G^{-1} \Omega^{\text{T}} - G^{-1} \Omega^{\text{L}} \otimes T.$
mielke-geometrodynamics_143 EQ0443 (6.8.14)		■ 0.924	$H_\alpha = -\frac{1}{\ell^2} * \left(\sum_{i=1}^3 a_i^{(i)} T_\alpha \right),$	$H_\alpha = -\frac{1}{\ell^2} * \left(\sum_{i=1}^3 a_i^{(i)} T_\alpha \right),$	$H_\alpha = -\frac{1}{\ell^2} * \left(\sum_{i=1}^3 a_i^{(i)} T_\alpha \right),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_81 EQ0192 (4.1.2)	81	■ 0.924	$\mathbf{a}:=\left[\begin{array}{c c} g & \mathrm{a} \\ \hline 0 & 1 \end{array}\right] \in \mathbb{R}^{n \times 2}$	$a := \left[\begin{array}{c c} g & \mathbf{a} \\ \hline 0 & 1 \end{array}\right] = \left[\begin{array}{c c} \mathbb{1}_n & \mathbf{a} \\ \hline 0 & 1 \end{array}\right] \left[\begin{array}{c c} g & 0 \\ \hline 0 & 1 \end{array}\right] = a^{\mathrm{T}} \cdot a^{\mathrm{L}} \in A(n, \mathbb{R}).$	$a := \left[\begin{array}{c c} g & \mathbf{a} \\ \hline 0 & 1 \end{array}\right] = \left[\begin{array}{c c} \mathbb{1}_n & \mathbf{a} \\ \hline 0 & 1 \end{array}\right] \left[\begin{array}{c c} g & 0 \\ \hline 0 & 1 \end{array}\right] = a^{\mathrm{T}} \cdot a^{\mathrm{L}} \in A(n, \mathbb{R}).$
mielke-geometrodynamics_278 EQ0857 (12.4.4) [re-fined: census]	278	■ 0.925	$\mathbf{t}_n \in \mathbb{R}^n \text{ such that } \mathbf{t}_n \in \mathbb{R}^n \text{ and } \mathbf{t}_n \in \mathbb{R}^n$	$\mathbb{L}_n \mathcal{G}^A \cong n \left(D_{\mathbb{T}}^A \right),$	$\mathbb{L}_n \mathcal{G}^A \cong n \left(D_{\mathbb{T}}^A \right),$
mielke-geometrodynamics_336 EQ1025 (15.1.1)	336	■ 0.926	$\begin{aligned} L_{f-g} &= \frac{1}{\ell^2} L_{\mathrm{W}}(f) + \frac{1}{\ell^2} L_{\mathrm{W}}(g) + L_{fg} + \frac{\Lambda}{\ell^2} \vartheta^g \wedge \vartheta^g \wedge \vartheta^g \wedge \vartheta^g \\ &\quad + L_{\mathrm{m}}(\mathrm{hadrons}, f) + L_{\mathrm{m}}(\mathrm{leptons}, g). \end{aligned}$	$L_{f-g} = \frac{1}{\ell^2} L_{\mathrm{W}}(f) + \frac{1}{\ell^2} L_{\mathrm{W}}(g) + L_{fg} + \frac{\Lambda}{\ell^2} \vartheta^g \wedge \vartheta^g \wedge \vartheta^g \wedge \vartheta^g + L_{\mathrm{m}}(\mathrm{hadrons}, f) + L_{\mathrm{m}}(\mathrm{leptons}, g).$	$L_{f-g} = \frac{1}{\ell^2} L_{\mathrm{W}}(f) + \frac{1}{\ell^2} L_{\mathrm{W}}(g) + L_{fg} + \frac{\Lambda}{\ell^2} \vartheta^g \wedge \vartheta^g \wedge \vartheta^g \wedge \vartheta^g + L_{\mathrm{m}}(\mathrm{hadrons}, f) + L_{\mathrm{m}}(\mathrm{leptons}, g).$
mielke-geometrodynamics_58 EQ0109 (3.3.10)	58	■ 0.927	$L_{\mathrm{Max}} = -\frac{1}{2} F \wedge {}^*F,$	$L_{\mathrm{Max}} = -\frac{1}{2} F \wedge {}^*F,$	$L_{\mathrm{Max}} = -\frac{1}{2} F \wedge {}^*F,$
mielke-geometrodynamics_194 EQ0587 (9.1.5)	194	■ 0.928	$d r^* := e^{(\lambda - v)/2} d r.$	$d r^* := e^{(\lambda - v)/2} d r.$	$d r^* := e^{(\lambda - v)/2} d r.$
mielke-geometrodynamics_370 EQ1101 (B.13)	370	■ 0.929	$\underbrace{(\vartheta \wedge \cdots \wedge \vartheta)}_{p \text{ products}} = \frac{p!}{(n-p)!} \underbrace{\vartheta \wedge \cdots \wedge \vartheta}_{n-p \text{ products}}(\star),$	$\underbrace{(\vartheta \wedge \cdots \wedge \vartheta)}_{p \text{ products}} = \frac{p!}{(n-p)!} \underbrace{\vartheta \wedge \cdots \wedge \vartheta}_{n-p \text{ products}}(\star),$	$\underbrace{(\vartheta \wedge \cdots \wedge \vartheta)}_{p \text{ products}} = \frac{p!}{(n-p)!} \underbrace{\vartheta \wedge \cdots \wedge \vartheta}_{n-p \text{ products}}(\star),$
mielke-geometrodynamics_137 EQ0413 (6.5.1)	137	■ 0.929	$W(A, \gamma) = \mathrm{Tr} \mathbf{P} \exp \oint A = \mathrm{Tr} \mathbf{P} \exp \oint ds \dot{\gamma}^i(s) A_i^J(\gamma(s)) \lambda_J,$	$W(A, \gamma) = \mathrm{Tr} \mathbf{P} \exp \oint A = \mathrm{Tr} \mathbf{P} \exp \oint ds \dot{\gamma}^i(s) A_i^J(\gamma(s)) \lambda_J,$	$W(A, \gamma) = \mathrm{Tr} \mathbf{P} \exp \oint A = \mathrm{Tr} \mathbf{P} \exp \oint ds \dot{\gamma}^i(s) A_i^J(\gamma(s)) \lambda_J,$
mielke-geometrodynamics_153 EQ0465 (7.3.4) [re-fined: census]	153	■ 0.929	$E_{\alpha} := \partial L_{\mathrm{SKY}} / \partial \vartheta^{\alpha} = \frac{1}{2} (e_{\alpha} \rfloor R^{\mu\nu} \wedge {}^* R_{\mu\nu} - R^{\mu\nu} \wedge e_{\alpha} \rfloor {}^* R_{\mu\nu}) = 0$	$E_{\alpha} := \partial L_{\mathrm{SKY}} / \partial \vartheta^{\alpha} = \frac{1}{2} (e_{\alpha} \rfloor R^{\mu\nu} \wedge {}^* R_{\mu\nu} - R^{\mu\nu} \wedge e_{\alpha} \rfloor {}^* R_{\mu\nu}) = 0$	$E_{\alpha} := \partial L_{\mathrm{SKY}} / \partial \vartheta^{\alpha} = \frac{1}{2} (e_{\alpha} \rfloor R^{\mu\nu} \wedge {}^* R_{\mu\nu} - R^{\mu\nu} \wedge e_{\alpha} \rfloor {}^* R_{\mu\nu}) = 0$
mielke-geometrodynamics_338 EQ1034 (15.1.11)	338	■ 0.930	$\Lambda_{\mathrm{eff.}}^f = \frac{2a}{\kappa b} \Lambda - 2a \ell^2 {}^* L_{fg}$	$\Lambda_{\mathrm{eff.}}^f = \frac{2a}{\kappa b} \Lambda - 2a \ell^2 {}^* L_{fg}$	$\Lambda_{\mathrm{eff.}}^f = \frac{2a}{\kappa b} \Lambda - 2a \ell^2 {}^* L_{fg}$
mielke-geometrodynamics_165 EQ0512 (7.7.30)	165	■ 0.931	$\begin{aligned} \Omega^g &:= d\Gamma + \Gamma \wedge \Gamma = \frac{i}{4} (d\Gamma_{\alpha}^{\beta} - \Gamma_{\alpha}^{\gamma} \wedge \Gamma_{\gamma}^{\beta}) \sigma^{\alpha}_{\beta} \\ &= \frac{i}{4} R^{\alpha\beta} \sigma_{\alpha\beta} = \frac{i}{8} R_{ij}{}^{\alpha\beta} \sigma_{\alpha\beta} dx^i \wedge dx^j, \end{aligned}$	$\Omega^g := d\Gamma + \Gamma \wedge \Gamma = \frac{i}{4} (d\Gamma_{\alpha}^{\beta} - \Gamma_{\alpha}^{\gamma} \wedge \Gamma_{\gamma}^{\beta}) \sigma^{\alpha}_{\beta} = \frac{i}{4} R^{\alpha\beta} \sigma_{\alpha\beta} = \frac{i}{8} R_{ij}{}^{\alpha\beta} \sigma_{\alpha\beta} dx^i \wedge dx^j,$	$\Omega^g := d\Gamma + \Gamma \wedge \Gamma = \frac{i}{4} (d\Gamma_{\alpha}^{\beta} - \Gamma_{\alpha}^{\gamma} \wedge \Gamma_{\gamma}^{\beta}) \sigma^{\alpha}_{\beta} = \frac{i}{4} R^{\alpha\beta} \sigma_{\alpha\beta} = \frac{i}{8} R_{ij}{}^{\alpha\beta} \sigma_{\alpha\beta} dx^i \wedge dx^j,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_00772 (11.2.1)	245	■ 0.943	$L_{\mathrm{D}}=\frac{i}{2}\left(\bar{\psi}\gamma\wedge^*\psi+\overline{(D\psi)}\wedge^*\gamma\psi\right)-m\bar{\psi}\psi\eta.$	$L_{\mathrm{D}}=\frac{i}{2}\left(\bar{\psi}\gamma\wedge^*D\psi+\overline{(D\psi)}\wedge^*\gamma\psi\right)-m\bar{\psi}\psi\eta.$	$L_{\mathrm{D}}=\frac{i}{2}\left(\bar{\psi}\gamma\wedge^*D\psi+\overline{(D\psi)}\wedge^*\gamma\psi\right)-m\bar{\psi}\psi\eta.$
mielke-geometrodynamics_00551 (8.4.5)	179	■ 0.945	$\begin{aligned} L_{\mathrm{FP}}=& \operatorname{Tr}\left\{\beta\wedge^*\bar{\Omega}^g-\bar{\chi}\wedge^*D\Psi^{(\pm)}-\bar{\chi}\wedge^*\left[C,\bar{\Omega}^g\right]\right.\\ & \left.+\bar{\eta}D^*\Psi+\bar{\Phi}D^*D\Phi+\bar{\Phi}\left[\Psi,^*\Psi\right]+\mathbb{B}d^*\Gamma\right.\\ & \left.-\bar{C}d^*\Psi+\bar{C}d^*(DC)+\frac{1}{2}\rho\beta\wedge^*\beta+\frac{1}{2}\mathbb{B}\wedge^*\mathbb{B}\right\}. \end{aligned}$	$L_{\mathrm{FP}}=-\operatorname{Tr}\left\{\beta\wedge^*\bar{\Omega}^g-\bar{\chi}\wedge^*D\Psi^{(\pm)}-\bar{\chi}\wedge^*\left[C,\bar{\Omega}^g\right]\right.\\ \left.+\bar{\eta}D^*\Psi+\bar{\Phi}D^*D\Phi+\bar{\Phi}\left[\Psi,^*\Psi\right]+\mathbb{B}d^*\Gamma\right.\\ \left.-\bar{C}d^*\Psi+\bar{C}d^*(DC)+\frac{1}{2}\rho\beta\wedge^*\beta+\frac{1}{2}\mathbb{B}\wedge^*\mathbb{B}\right\}.$	$L_{\mathrm{FP}}=-\operatorname{Tr}\left\{\beta\wedge^*\bar{\Omega}^g-\bar{\chi}\wedge^*D\Psi^{(\pm)}-\bar{\chi}\wedge^*\left[C,\bar{\Omega}^g\right]\right.\\ \left.+\bar{\eta}D^*\Psi+\bar{\Phi}D^*D\Phi+\bar{\Phi}\left[\Psi,^*\Psi\right]+\mathbb{B}d^*\Gamma\right.\\ \left.-\bar{C}d^*\Psi+\bar{C}d^*(DC)+\frac{1}{2}\rho\beta\wedge^*\beta+\frac{1}{2}\mathbb{B}\wedge^*\mathbb{B}\right\}.$
mielke-geometrodynamics_01069 (16.2.8)	356	■ 0.945	$V_{\mathrm{eff}}^{\iota}(\rho^*)=\frac{1\pm\rho^2}{\rho^2}\left[\iota(\iota+1)-\left(2-\frac{1}{\beta}\right)\rho^2\right],\kappa>0\\ \&(\kappa<0).$	$V_{\mathrm{eff}}^{\iota}(\rho^*)=\frac{1\pm\rho^2}{\rho^2}\left[\iota(\iota+1)-\left(2-\frac{1}{\beta}\right)\rho^2\right],\kappa>0\\ (\kappa<0).$	$V_{\mathrm{eff}}^{\iota}(\rho^*)=\frac{1\pm\rho^2}{\rho^2}\left[\iota(\iota+1)-\left(2-\frac{1}{\beta}\right)\rho^2\right],\kappa>0\\ (\kappa<0).$
mielke-geometrodynamics_00593 (9.1.11)	195	■ 0.945	$R=\left\{\right\}\wedge^*\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)=-\frac{6\kappa\gamma}{\ell^{*2}}=4\Lambda_{\mathrm{eff}}.$	$R=\left\{\right\}\wedge^*\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)=-\frac{6\kappa\gamma}{\ell^{*2}}=4\Lambda_{\mathrm{eff}}.$	$R=\left\{\right\}\wedge^*\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)=-\frac{6\kappa\gamma}{\ell^{*2}}=4\Lambda_{\mathrm{eff}}.$
mielke-geometrodynamics_00851 (12.3.4)	276	■ 0.946	$\begin{aligned} & \operatorname{Tr}\left(\gamma_5K_2\right)=-d^*\mathscr{A},\\ & \operatorname{Tr}\left(\gamma_5K_4\right)=\frac{1}{6}\left[\operatorname{Tr}\left(R^{\lrcorner}\wedge R^{\lrcorner}\right)-\frac{1}{4}d\mathscr{A}\wedge d\mathscr{A}+d\mathscr{K}\right], \end{aligned}$	$\operatorname{Tr}\left(\gamma_5K_2\right)=-d^*\mathscr{A}$ $\operatorname{Tr}\left(\gamma_5K_4\right)=\frac{1}{6}\left[\operatorname{Tr}\left(R^{\lrcorner}\wedge R^{\lrcorner}\right)-\frac{1}{4}d\mathscr{A}\wedge d\mathscr{A}+d\mathscr{K}\right],$	$\operatorname{Tr}\left(\gamma_5K_2\right)=-d^*\mathscr{A},$ $\operatorname{Tr}\left(\gamma_5K_4\right)=\frac{1}{6}\left[\operatorname{Tr}\left(R^{\lrcorner}\wedge R^{\lrcorner}\right)-\frac{1}{4}d\mathscr{A}\wedge d\mathscr{A}+d\mathscr{K}\right],$
mielke-geometrodynamics_00815 (11.4.29)	257	■ 0.946	$\begin{aligned} & H^{\prime}\simeq\frac{1+\omega}{\beta}F,\\ & F^{\prime}+\frac{2}{\rho}F\simeq\frac{1-\omega}{\beta}H \end{aligned}$	$H^{\prime}\simeq\frac{1+\omega}{\beta}F$ $F^{\prime}+\frac{2}{\rho}F\simeq\frac{1-\omega}{\beta}H$	$H^{\prime}\simeq\frac{1+\omega}{\beta}F,$ $F^{\prime}+\frac{2}{\rho}F\simeq\frac{1-\omega}{\beta}H.$
mielke-geometrodynamics_01021 (14.4.16)	328	■ 0.947	$S_{\mathrm{A}}:=4\pi r_+^2=16\pi M_{\mathrm{ir}}^2\frac{\hbar^2}{M^{*4}c^2},$	$S_{\mathrm{A}}:=4\pi r_+^2=16\pi M_{\mathrm{ir}}^2\frac{\hbar^2}{M^{*4}c^2},$	$S_{\mathrm{A}}:=4\pi r_+^2=16\pi M_{\mathrm{ir}}^2\frac{\hbar^2}{M^{*4}c^2},$
mielke-geometrodynamics_00855 (12.4.2)	277	■ 0.947	$\mathscr{H}_{\Lambda}\Psi(\underline{\Gamma})=0$	$\mathscr{H}_{\Lambda}\Psi^{(1)}(\underline{\Gamma})=0$	$\mathscr{H}_{\Lambda}\Psi^{(\pm)}(\underline{\Gamma})=0$
mielke-geometrodynamics_00452 (7.2.5)	151	■ 0.948	$C_{\cdot\cdot cd}^{\cdot\cdot cd}=\frac{R_{\cdot\cdot cd}^{\cdot\cdot cd}}{2}+\frac{R_{\cdot\cdot cd}^{\cdot\cdot cd}}{6}\delta_{[c}^a\delta_{d]}^b.$	$C_{\cdot\cdot cd}^{\cdot\cdot cd}=\frac{R_{\cdot\cdot cd}^{\cdot\cdot cd}}{2}+\frac{R_{\cdot\cdot cd}^{\cdot\cdot cd}}{6}\delta_{[c}^a\delta_{d]}^b.$	$C_{\cdot\cdot cd}^{\cdot\cdot cd}=\frac{R_{\cdot\cdot cd}^{\cdot\cdot cd}}{2}+\frac{R_{\cdot\cdot cd}^{\cdot\cdot cd}}{6}\delta_{[c}^a\delta_{d]}^b.$
mielke-geometrodynamics_00061 (2.8.2)	46	■ 0.949	$\left\{\right\}\wedge^*\left(\gamma_k\right)=\frac{(-1)^k}{(2\pi i)^kk!}\operatorname{Tr}(\Omega\wedge\cdots\wedge\Omega).$	$\left\{\right\}\wedge^*(\gamma_k)=\frac{(-1)^k}{(2\pi i)^kk!}\operatorname{Tr}(\Omega\wedge\cdots\wedge\Omega).$	$\left\{\right\}\wedge^*(\gamma_k)=\frac{(-1)^k}{(2\pi i)^kk!}\operatorname{Tr}(\Omega\wedge\cdots\wedge\Omega).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00756 (11.1.23)	242	■ 0.950	$\begin{aligned} S(p) &= \exp \imath \theta_{\alpha\beta}(m) \Lambda^{-1} (L^{\alpha\beta}) = \exp \imath \theta_{\alpha\beta}(m) \sigma^{\alpha\beta} \\ &= \exp \left(-\frac{1}{4} \theta_{\alpha\beta}(m) [\gamma^\alpha, \gamma^\beta] \right). \end{aligned}$	$S(p) = \exp \imath \theta_{\alpha\beta}(m) \Lambda^{-1} (L^{\alpha\beta}) = \exp \imath \theta_{\alpha\beta}(m) \sigma^{\alpha\beta} \\ = \exp \left(-\frac{1}{4} \theta_{\alpha\beta}(m) [\gamma^\alpha, \gamma^\beta] \right).$	$S(p) = \exp \imath \theta_{\alpha\beta}(m) \Lambda^{-1} (L^{\alpha\beta}) = \exp \imath \theta_{\alpha\beta}(m) \sigma^{\alpha\beta} \\ = \exp \left(-\frac{1}{4} \theta_{\alpha\beta}(m) [\gamma^\alpha, \gamma^\beta] \right).$
mielke-geometrodynamics_E00939 (14.1.20)	305	■ 0.950	$\xi \wedge \xi = 0 \quad (= 2 \vec{E} \cdot \vec{B} \text{ in 3-dimensional vector notation })$	$\xi \wedge \xi = 0 \quad (= 2 \vec{E} \cdot \vec{B} \text{ in 3-dimensional vector notation })$	$\xi \wedge \xi = 0 \quad (= 2 \vec{E} \cdot \vec{B} \text{ in 3-dimensional vector notation })$
mielke-geometrodynamics_E00568 (8.6.5)	183	■ 0.950	$L_{\theta} := \theta_T dC_{TT} + \theta_T^* dC_{TT^*} + \theta_L dC_{RR} + \theta_L^* (-1)^{\text{sig}+1} dC_{RR^*} = dC_{\theta}.$	$L_{\theta} := \theta_T dC_{TT} + \theta_T^* dC_{TT^*} + \theta_L dC_{RR} + \theta_L^* (-1)^{\text{sig}+1} dC_{RR^*} = dC_{\theta}.$	$L_{\theta} := \theta_T dC_{TT} + \theta_T^* dC_{TT^*} + \theta_L dC_{RR} + \theta_L^* (-1)^{\text{sig}+1} dC_{RR^*} = dC_{\theta}.$
mielke-geometrodynamics_E00234 (4.2.27)	88	■ 0.950	$\widetilde{D}\widetilde{\varphi} = D\widetilde{\varphi} = 0$	$\widetilde{D}\widetilde{\varphi} = D\widetilde{\varphi} = 0$	$\widetilde{D}\widetilde{\varphi} = D\widetilde{\varphi} = 0$
mielke-geometrodynamics_E00302 (4.5.9) [re-fined: census]	102	■ 0.950	$\begin{aligned} \widehat{D} \Sigma_{\alpha} &\cong \left(e_{\alpha} \rfloor R_{\beta}^{\gamma} \right) \wedge \Delta^{\beta} \gamma \\ &\quad - \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta\gamma}) \wedge \sigma^{\beta\gamma} \end{aligned}$	$\widehat{D} \Sigma_{\alpha} \cong \left(e_{\alpha} \rfloor R_{\beta}^{\gamma} \right) \wedge \Delta^{\beta} \gamma \\ - \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta\gamma}) \wedge \sigma^{\beta\gamma}$	$\widehat{D} \Sigma_{\alpha} \cong (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \Delta^{\beta}_{\gamma} \\ - \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta\gamma}) \wedge \sigma^{\beta\gamma}$
mielke-geometrodynamics_E00151 (3.4.22)	66	■ 0.951	$L_{\text{Mills}}(\mathcal{F}, * \mathcal{F}) = \mathcal{F} \left[1 - \left(1 + \frac{1}{b^2} \mathcal{F}^2 \right)^{-1} \right]$	$L_{\text{Mills}}(\mathcal{F}, * \mathcal{F}) = \mathcal{F} \left[1 - \left(1 + \frac{1}{b^2} \mathcal{F}^2 \right)^{-1} \right]$	$L_{\text{Mills}}(\mathcal{F}, * \mathcal{F}) = \mathcal{F} \left[1 - \left(1 + \frac{1}{b^2} \mathcal{F}^2 \right)^{-1} \right]$
mielke-geometrodynamics_E00560 (8.5.3)	181	■ 0.953	$\stackrel{\pm}{\Omega}^g = i \frac{\theta_T^*}{4\ell^2} \sigma$	$\stackrel{\pm}{\Omega}^g = i \frac{\theta_T^*}{4\ell^2} \sigma$	$\stackrel{\pm}{\Omega}^g = i \frac{\theta_T^*}{4\ell^2} \sigma$
mielke-geometrodynamics_E01106 (B.18)	370	■ 0.953	$\delta \alpha^{(p)} := (-1)^{pn+n+1+s} d^* \alpha^{(p)}.$	$\delta \alpha^{(p)} := (-1)^{pn+n+1+s} d^* \alpha^{(p)}.$	$\delta \alpha^{(p)} := (-1)^{pn+n+1+s} d^* \alpha^{(p)}.$
mielke-geometrodynamics_E00786 (11.3.5)	248	■ 0.954	$\begin{aligned} L_{\text{NL}} &= -3\ell^{*2} \tau_s \wedge^* \tau_s = 3\ell^{*2} \bar{\psi}^* \gamma \psi \wedge^* (\bar{\psi}^* \gamma \psi) \\ &= 3\ell^{*2} \bar{\psi} \gamma^{n+1} \gamma \psi \wedge^* (\bar{\psi} \gamma^{n+1} \gamma \psi). \end{aligned}$	$L_{\text{NL}} = -3\ell^{*2} \tau_s \wedge^* \tau_s = 3\ell^{*2} \bar{\psi}^* \gamma \psi \wedge^* (\bar{\psi}^* \gamma \psi) \\ = 3\ell^{*2} \bar{\psi} \gamma^{n+1} \gamma \psi \wedge^* (\bar{\psi} \gamma^{n+1} \gamma \psi).$	$L_{\text{NL}} = -3\ell^{*2} \tau_s \wedge^* \tau_s = 3\ell^{*2} \bar{\psi}^* \gamma \psi \wedge^* (\bar{\psi}^* \gamma \psi) \\ = 3\ell^{*2} \bar{\psi} \gamma^{n+1} \gamma \psi \wedge^* (\bar{\psi} \gamma^{n+1} \gamma \psi).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_00727 (10.10.7)	231	■ 0.955	$\begin{aligned} & i \gamma^\alpha \Psi \wedge \frac{\delta L}{\delta \vartheta^\alpha} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} - 2D \frac{\delta L}{\delta \bar{\Psi}} \\ & \cong i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\text{TL}}}{\ell} R_\alpha^\star + \frac{\theta_{\text{T}}}{\ell^2} T_\alpha + \Sigma_\alpha \right) \\ & + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - D \left(i D \Psi + \frac{i}{2} m \gamma \wedge \Psi \right) \\ & = i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\text{TL}}}{\ell} R_\alpha^\star + \frac{\theta_{\text{T}}}{\ell^2} T_\alpha \right) + \gamma^\alpha \Psi \wedge \left(\frac{1}{4} m \bar{\Psi} \gamma_\alpha \Psi \right) \\ & + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - i R_\alpha^\star \gamma^\alpha \Psi - \frac{i}{2} m T_\alpha \gamma^\alpha \Psi \\ & + \frac{i}{2} m \gamma \wedge D \Psi. \end{aligned} \tag{10.1}$	$i \gamma^\alpha \Psi \wedge \frac{\delta L}{\delta \vartheta^\alpha} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} - 2D \frac{\delta L}{\delta \bar{\Psi}} \\ \cong i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\text{TL}}}{\ell} R_\alpha^\star + \frac{\theta_{\text{T}}}{\ell^2} T_\alpha + \Sigma_\alpha \right) \\ + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - D \left(i D \Psi + \frac{i}{2} m \gamma \wedge \Psi \right) \\ = i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\text{TL}}}{\ell} R_\alpha^\star + \frac{\theta_{\text{T}}}{\ell^2} T_\alpha \right) + \gamma^\alpha \Psi \wedge \left(\frac{1}{4} m \bar{\Psi} \gamma_\alpha \Psi \right) \\ + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - i R_\alpha^\star \gamma^\alpha \Psi - \frac{i}{2} m T_\alpha \gamma^\alpha \Psi \\ + \frac{i}{2} m \gamma \wedge D \Psi.$	$i \gamma^\alpha \Psi \wedge \frac{\delta L}{\delta \vartheta^\alpha} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} - 2D \frac{\delta L}{\delta \bar{\Psi}} \\ \cong i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\text{TL}}}{\ell} R_\alpha^\star + \frac{\theta_{\text{T}}}{\ell^2} T_\alpha + \Sigma_\alpha \right) \\ + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - D \left(i D \Psi + \frac{i}{2} m \gamma \wedge \Psi \right) \\ = i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\text{TL}}}{\ell} R_\alpha^\star + \frac{\theta_{\text{T}}}{\ell^2} T_\alpha \right) + \gamma^\alpha \Psi \wedge \left(\frac{1}{4} m \bar{\Psi} \gamma_\alpha \Psi \right) \\ + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - i R_\alpha^\star \gamma^\alpha \Psi - \frac{i}{2} m T_\alpha \gamma^\alpha \Psi \\ + \frac{i}{2} m \gamma \wedge D \Psi. \tag{10.1}$
mielke-geometrodynamics_00768 (11.1.35)	244	■ 0.956	$\bar{\psi} \longrightarrow \bar{\psi} S,$	$\bar{\psi} \longrightarrow S^{-1} \bar{\psi} = \bar{\psi} S,$	$\bar{\psi} \longrightarrow S^{-1} \bar{\psi} = \bar{\psi} S,$
mielke-geometrodynamics_00973 (14.2.14)	315	■ 0.956	$\begin{aligned} & \hat{\Theta} \wedge \hat{\vartheta} = \hat{D} \hat{\vartheta} - [\hat{\omega}, \hat{\vartheta}] \\ & = \left[\begin{array}{c c} \Theta - \hat{\ell} \omega \wedge \alpha^\text{T} & \hat{\ell} (d\omega - \omega^g \wedge \omega - \omega \wedge \mathring{\omega}) + \vartheta \wedge \alpha + \alpha \wedge \mathring{\vartheta} \\ \hline -\alpha^\text{T} \wedge \vartheta - \vartheta \wedge \alpha^\text{T} & \mathring{\Theta} - \hat{\ell} \alpha^\text{T} \wedge \omega \end{array} \right]. \end{aligned}$	$\begin{aligned} \hat{\Theta} &:= \hat{D} \hat{\vartheta} = \hat{d} \hat{\vartheta} - [\hat{\omega}, \hat{\vartheta}] \\ &= \left[\begin{array}{c c} \Theta - \hat{\ell} \omega \wedge \alpha^\text{T} & \hat{\ell} (d\omega - \omega^g \wedge \omega - \omega \wedge \mathring{\omega}) + \vartheta \wedge \alpha + \alpha \wedge \mathring{\vartheta} \\ \hline -\alpha^\text{T} \wedge \vartheta - \vartheta \wedge \alpha^\text{T} & \mathring{\Theta} - \hat{\ell} \alpha^\text{T} \wedge \omega \end{array} \right]. \end{aligned}$	$\begin{aligned} \hat{\Theta} &:= \hat{D} \hat{\vartheta} = \hat{d} \hat{\vartheta} - [\hat{\omega}, \hat{\vartheta}] \\ &= \left[\begin{array}{c c} \Theta - \hat{\ell} \omega \wedge \alpha^\text{T} & \hat{\ell} (d\omega - \omega^g \wedge \omega - \omega \wedge \mathring{\omega}) + \vartheta \wedge \alpha + \alpha \wedge \mathring{\vartheta} \\ \hline -\alpha^\text{T} \wedge \vartheta - \vartheta \wedge \alpha^\text{T} & \mathring{\Theta} - \hat{\ell} \alpha^\text{T} \wedge \omega \end{array} \right]. \end{aligned}$
mielke-geometrodynamics_00761 (11.1.28)	243	■ 0.957	$\frac{1}{N} \text{Tr}(\gamma \otimes_s \gamma) = ds^2 = g_{ij}(m) dx^i \otimes_s dx^j.$	$\frac{1}{N} \text{Tr}(\gamma \otimes_s \gamma) = ds^2 = g_{ij}(m) dx^i \otimes_s dx^j.$	$\frac{1}{N} \text{Tr}(\gamma \otimes_s \gamma) = ds^2 = g_{ij}(m) dx^i \otimes_s dx^j.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_160 E00486 (7.7.2) [re-fined: census]	160	■ 0.957	$\begin{aligned} & \vartheta^{\alpha} \rightarrow \widetilde{\vartheta}^{\alpha} := \vartheta^{\alpha} + \ell^2 e_{\beta} \rfloor^* H^{\alpha\beta}, \\ & \Gamma_{\alpha}{}^{\beta} \rightarrow \widetilde{\Gamma}_{\alpha}{}^{\beta} := \Gamma_{\alpha}{}^{\beta} + \ell^2 e_{\alpha} \rfloor^* H^{\beta}. \end{aligned}$	$\vartheta^{\alpha} \rightarrow \widetilde{\vartheta}^{\alpha} := \vartheta^{\alpha} + \ell^2 e_{\beta} \rfloor^* H^{\alpha\beta},$ $\Gamma_{\alpha}{}^{\beta} \rightarrow \widetilde{\Gamma}_{\alpha}{}^{\beta} := \Gamma_{\alpha}{}^{\beta} + \ell^2 e_{\alpha} \rfloor^* H^{\beta}.$	$\vartheta^{\alpha} \rightarrow \widetilde{\vartheta}^{\alpha} := \vartheta^{\alpha} + \ell^2 e_{\beta} \rfloor^* H^{\alpha\beta},$ $\Gamma_{\alpha}{}^{\beta} \rightarrow \widetilde{\Gamma}_{\alpha}{}^{\beta} := \Gamma_{\alpha}{}^{\beta} + \ell^2 e_{\alpha} \rfloor^* H^{\beta}.$
mielke-geometrodynamics_370 E01104 (B.16)	370	■ 0.957	$\begin{aligned} & \left\langle \alpha^{(p)}, \beta^{(p)} \right\rangle := \alpha^{(p)} \wedge^* \beta^{(p)} = \beta^{(p)} \wedge^* \alpha^{(p)} \\ & = (-1)^{s*} \left(\alpha^{(p)} \wedge^* \beta^{(p)} \right) \eta \\ & = \frac{(-1)^s}{p!} A^{\alpha_1 \cdots \alpha_p} B_{\alpha_1 \cdots \alpha_p} d\mu(M) \in C^{\infty}(\wedge^n T^*(M)) \end{aligned}$	$\left\langle \alpha^{(p)}, \beta^{(p)} \right\rangle := \alpha^{(p)} \wedge^* \beta^{(p)} = \beta^{(p)} \wedge^* \alpha^{(p)}$ $= (-1)^{s*} \left(\alpha^{(p)} \wedge^* \beta^{(p)} \right) \eta$ $= \frac{(-1)^s}{p!} A^{\alpha_1 \cdots \alpha_p} B_{\alpha_1 \cdots \alpha_p} d\mu(M) \in C^{\infty}(\wedge^n T^*(M))$	$\langle \alpha^{(p)}, \beta^{(p)} \rangle := \alpha^{(p)} \wedge^* \beta^{(p)} = \beta^{(p)} \wedge^* \alpha^{(p)}$ $= (-1)^{s*} (\alpha^{(p)} \wedge^* \beta^{(p)}) \eta$ $= \frac{(-1)^s}{p!} A^{\alpha_1 \cdots \alpha_p} B_{\alpha_1 \cdots \alpha_p} d\mu(M) \in C^{\infty}(\wedge^n T^*(M))$
mielke-geometrodynamics_176 E00537 (8.2.4)	176	■ 0.958	$\begin{array}{l} s\,\bar{C}=\mathbb{B},\quad s\,\mathbb{B}=0,\\ s\,\bar{\chi}=\beta,\quad s\,\beta=0,\\ s\,\bar{\Phi}=\bar{\eta},\quad s\,\bar{\eta}=0. \end{array}$	$\begin{array}{l} s\bar{C}=\mathbb{B},\quad s\mathbb{B}=0,\\ s\bar{\chi}=\beta,\quad s\beta=0,\\ s\bar{\Phi}=\bar{\eta},\quad s\bar{\eta}=0. \end{array}$	$\begin{array}{l} s\bar{C}=\mathbb{B},\quad s\mathbb{B}=0,\\ s\bar{\chi}=\beta,\quad s\beta=0,\\ s\bar{\Phi}=\bar{\eta},\quad s\bar{\eta}=0. \end{array}$
mielke-geometrodynamics_95 E00269 (4.3.34)	95	■ 0.958	$R_{\{i\}j}:=R_{\{\alpha\}i\}^j\cdots\alpha\}$	$R_{ij}:=R_{\alpha ij}^{\cdots\alpha}$	$R_{ij}:=R_{\alpha ij}^{\cdots\alpha}$
mielke-geometrodynamics_132 E00394 (6.4.9)	132	■ 0.958	$\left.\mathscr{H}\right.\cong n^{\alpha}\underline{\Sigma}_{\alpha}+\Gamma_{\perp}^{\alpha\beta}\rho\left(I_{\alpha\beta}\right)\underline{\Psi}\wedge\frac{\partial L}{\partial D\Psi}-\underline{d}\left[\left(n\right]\Psi\right)\wedge\frac{\partial L}{\partial D\Psi}\Big].$	$\mathcal{H}\cong n^{\alpha}\underline{\Sigma}_{\alpha}+\Gamma_{\perp}^{\alpha\beta}\rho\left(I_{\alpha\beta}\right)\underline{\Psi}\wedge\frac{\partial L}{\partial D\Psi}-\underline{d}\left[\left(n\right]\Psi\right)\wedge\frac{\partial L}{\partial D\Psi}\Big].$	$\mathcal{H}\cong n^{\alpha}\underline{\Sigma}_{\alpha}+\Gamma_{\perp}^{\alpha\beta}\rho\left(I_{\alpha\beta}\right)\underline{\Psi}\wedge\frac{\partial L}{\partial D\Psi}-\underline{d}\left[\left(n\right]\Psi\right)\wedge\frac{\partial L}{\partial D\Psi}\Big].$
mielke-geometrodynamics_135 E00406 (6.4.26)	135	■ 0.959	$\stackrel{\scriptscriptstyle\pm}{\operatorname{D}}\stackrel{\scriptscriptstyle\pm}{\operatorname{T}}^B=0$	$\stackrel{(\pm)}{\underline{D}}\stackrel{(\pm)}{\underline{T}}^B=0$	$\stackrel{(\pm)}{\underline{D}}\stackrel{(\pm)}{\underline{T}}^B=0$
mielke-geometrodynamics_152 E00458 (7.2.10)	152	■ 0.959	$\mathfrak{s}\mathfrak{o}(4)\approx\mathfrak{s}\mathfrak{o}(3)\times\mathfrak{s}\mathfrak{o}(3)\approx\mathfrak{s}\mathfrak{u}(2)\times\mathfrak{s}\mathfrak{u}(2).$	$\mathfrak{so}(4)\approx\mathfrak{so}(3)\times\mathfrak{so}(3)\approx\mathfrak{su}(2)\times\mathfrak{su}(2).$	$\mathfrak{so}(4)\approx\mathfrak{so}(3)\times\mathfrak{so}(3)\approx\mathfrak{su}(2)\times\mathfrak{su}(2).$
mielke-geometrodynamics_308 E00954 (14.1.36)	308	■ 0.959	$\begin{aligned} & \left. \operatorname{Tr}_{\{\xi\}} \left(\xi \wedge \vartheta \wedge \xi - \xi \wedge \vartheta \wedge \xi \right) \wedge d\delta \right. \\ & = \operatorname{Tr}_{\{\xi\}} \left(\xi \wedge \vartheta \wedge D^* \xi + \xi \wedge \vartheta \wedge D \xi \right) \\ & = -\frac{1}{2} D \operatorname{Tr}_{\{\xi\}} \left(\xi \wedge \vartheta \wedge \xi - \xi \wedge \vartheta \wedge \xi \right). \end{aligned}$	$\begin{aligned} & (-1)^z \operatorname{Tr}_{\{\xi\}} \left(\xi \wedge \vartheta \wedge \xi - \xi \wedge \vartheta \wedge \xi \right) \wedge d\delta \\ & = \operatorname{Tr}_{\{\xi\}} \left(\xi \wedge \vartheta \wedge D^* \xi + \xi \wedge \vartheta \wedge D \xi \right) \\ & = -\frac{1}{2} D \operatorname{Tr}_{\{\xi\}} \left(\xi \wedge \vartheta \wedge \xi - \xi \wedge \vartheta \wedge \xi \right). \end{aligned}$	$\begin{aligned} & (-1)^z \operatorname{Tr}_{\{\xi\}} \left(\xi \wedge \vartheta \wedge \xi - \xi \wedge \vartheta \wedge \xi \right) \wedge d\delta \\ & = \operatorname{Tr}_{\{\xi\}} \left(\xi \wedge \vartheta \wedge D^* \xi + \xi \wedge \vartheta \wedge D \xi \right) \\ & = -\frac{1}{2} D \operatorname{Tr}_{\{\xi\}} \left(\xi \wedge \vartheta \wedge \xi - \xi \wedge \vartheta \wedge \xi \right). \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_112 E08338 (5.3.7)	112	■ 0.960	$G:=G_{\alpha}\gamma^{\alpha}\wedge\gamma^{\alpha}:=\frac{1}{2}R^{\mu\nu}\wedge\eta_{\mu\nu\lambda}\gamma^{\lambda}=-i\gamma_5\left(\Omega^g\wedge\gamma+\gamma\wedge\Omega^g\right)=i\left[\gamma,\gamma_5\Omega^g\right].$	$G:=G_{\alpha}\gamma^{\alpha}:=\frac{1}{2}R^{\mu\nu}\wedge\eta_{\mu\nu\lambda}\gamma^{\lambda}=-i\gamma_5\left(\Omega^g\wedge\gamma+\gamma\wedge\Omega^g\right)=i\left[\gamma,\gamma_5\Omega^g\right].$	$G:=G_{\alpha}\gamma^{\alpha}:=\frac{1}{2}R^{\mu\nu}\wedge\eta_{\mu\nu\lambda}\gamma^{\lambda}=-i\gamma_5(\Omega^g\wedge\gamma+\gamma\wedge\Omega^g)=i[\gamma,\gamma_5\Omega^g].$
mielke-geometrodynamics_303 E09926 (14.1.7)	303	■ 0.960	$D\left(\vartheta\wedge\Omega^g\right)\equiv\Theta\wedge\Omega^g.$	$D\left(\vartheta\wedge\Omega^g\right)\equiv\Theta\wedge\Omega^g.$	$D(\vartheta\wedge\Omega^g)\equiv\Theta\wedge\Omega^g.$
mielke-geometrodynamics_92 E08248 (4.3.13)	92	■ 0.961	$D\phi^{(\circ)}=\vartheta^{\alpha}\nabla_{\alpha}\phi^{(\circ)}.$	$D\phi^{(\circ)}=\vartheta^{\alpha}\nabla_{\alpha}\phi^{(\circ)}.$	$D\phi^{(\circ)}=\vartheta^{\alpha}\nabla_{\alpha}\phi^{(\circ)}.$
mielke-geometrodynamics_98 E08282 (4.4.11)	98	■ 0.961	$\begin{aligned} d\Theta&=dd\vartheta-d[\omega^L,\vartheta] \\ &=-[d\omega^L,\vartheta]+[\omega^L,d\vartheta] \\ &=-[\Omega^L,\vartheta]-[\omega^L\wedge\omega^L,\vartheta]+[\omega^L,\Theta]+[\omega^L,[\omega^L,\vartheta]] \\ &=[\omega^L,\Theta]+[\vartheta,\Omega^L] \end{aligned}$	$\begin{aligned} d\Theta&=dd\vartheta-d[\omega^L,\vartheta] \\ &=-[d\omega^L,\vartheta]+[\omega^L,d\vartheta] \\ &=-[\Omega^L,\vartheta]-[\omega^L\wedge\omega^L,\vartheta]+[\omega^L,\Theta]+[\omega^L,[\omega^L,\vartheta]] \\ &=[\omega^L,\Theta]+[\vartheta,\Omega^L] \end{aligned}$	$\begin{aligned} d\Theta&=dd\vartheta-d[\omega^L,\vartheta] \\ &=-[d\omega^L,\vartheta]+[\omega^L,d\vartheta] \\ &=-[\Omega^L,\vartheta]-[\omega^L\wedge\omega^L,\vartheta]+[\omega^L,\Theta]+[\omega^L,[\omega^L,\vartheta]] \\ &=[\omega^L,\Theta]+[\vartheta,\Omega^L]. \end{aligned}$
mielke-geometrodynamics_34 E09016 (2.2.5)	34	■ 0.962	$T_{\{p\}^q}(M):=\bigcup_{m\in M}T_m^*(M)\otimes^qT_m(M).$	$T_p^q(M):=\bigcup_{m\in M}\otimes^pT_m^*(M)\otimes^qT_m(M).$	$T_p^q(M):=\bigcup_{m\in M}\otimes^pT_m^*(M)\otimes^qT_m(M).$
mielke-geometrodynamics_115 E08348 (5.3.17) [re-fined; census]	115	■ 0.962	$\begin{aligned} G_{\alpha}&:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma} \\ &=G_{\alpha}^{\{\}}+\frac{(-1)^s}{12}\left(e_{\alpha}\rfloor\mathcal{A}\wedge{}^*\mathcal{A}-\frac{1}{3}\mathcal{A}\wedge e_{\alpha}\rfloor{}^*\mathcal{A}\right)+\frac{(-1)^s}{6}\vartheta_{\alpha}\wedge d\mathcal{A} \end{aligned}$	$\begin{aligned} G_{\alpha}&:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma} \\ &=G_{\alpha}^{\{\}}+\frac{(-1)^s}{12}\left(e_{\alpha}\rfloor\mathcal{A}\wedge{}^*\mathcal{A}-\frac{1}{3}\mathcal{A}\wedge e_{\alpha}\rfloor{}^*\mathcal{A}\right)+\frac{(-1)^s}{6}\vartheta_{\alpha}\wedge d\mathcal{A} \end{aligned}$	$\begin{aligned} G_{\alpha}&:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma} \\ &=G_{\alpha}^{\{\}}+\frac{(-1)^s}{12}\left(e_{\alpha}\rfloor\mathcal{A}\wedge{}^*\mathcal{A}-\frac{1}{3}\mathcal{A}\wedge e_{\alpha}\rfloor{}^*\mathcal{A}\right)+\frac{(-1)^s}{6}\vartheta_{\alpha}\wedge d\mathcal{A} \end{aligned} \tag{5.3}$
mielke-geometrodynamics_140 E08424 (6.7.6)	140	■ 0.962	$\begin{aligned} \frac{1}{2\kappa}T^{\gamma}\wedge\eta_{\alpha\beta\gamma}-\frac{D\theta_{\mathrm{T}}}{2\ell^2}\wedge\vartheta_{\alpha}\wedge\vartheta_{\beta}+D\theta_{\mathrm{L}}\wedge R_{\alpha\beta}&=\vartheta_{[\alpha}\wedge\mu_{\beta]} \\ &=\frac{1}{4}\vartheta_{\alpha}\wedge\vartheta_{\beta}\wedge{}^*j_5. \end{aligned}$	$\begin{aligned} \frac{1}{2\kappa}T^{\gamma}\wedge\eta_{\alpha\beta\gamma}-\frac{D\theta_{\mathrm{T}}}{2\ell^2}\wedge\vartheta_{\alpha}\wedge\vartheta_{\beta}+D\theta_{\mathrm{L}}\wedge R_{\alpha\beta}&=\vartheta_{[\alpha}\wedge\mu_{\beta]} \\ &=\frac{1}{4}\vartheta_{\alpha}\wedge\vartheta_{\beta}\wedge{}^*j_5. \end{aligned}$	$\begin{aligned} \frac{1}{2\kappa}T^{\gamma}\wedge\eta_{\alpha\beta\gamma}-\frac{D\theta_{\mathrm{T}}}{2\ell^2}\wedge\vartheta_{\alpha}\wedge\vartheta_{\beta}+D\theta_{\mathrm{L}}\wedge R_{\alpha\beta}&=\vartheta_{[\alpha}\wedge\mu_{\beta]} \\ &=\frac{1}{4}\vartheta_{\alpha}\wedge\vartheta_{\beta}\wedge{}^*j_5. \end{aligned}$
mielke-geometrodynamics_255 E08889 (11.4.21)	255	■ 0.962	$i\gamma^0\vec{\gamma}\cdot\vec{\partial}\left[\begin{array}{cc} iH(\rho) & \chi_{\kappa}^m \\ -F(\rho) & \chi_{-\kappa}^m \end{array}\right]=\frac{\vec{\sigma}\cdot\vec{x}}{ \vec{x} }\left(\partial_r-\frac{\vec{\sigma}\cdot\vec{L}}{r}\right)\left[\begin{array}{cc} iF(\rho) & \chi_{-\kappa}^m \\ H(\rho) & \chi_{\kappa}^m \end{array}\right].$	$i\gamma^0\vec{\gamma}\cdot\vec{\partial}\left[\begin{array}{cc} iH(\rho) & \chi_{\kappa}^m \\ -F(\rho) & \chi_{-\kappa}^m \end{array}\right]=\frac{\vec{\sigma}\cdot\vec{x}}{ \vec{x} }\left(\partial_r-\frac{\vec{\sigma}\cdot\vec{L}}{r}\right)\left[\begin{array}{cc} iF(\rho) & \chi_{-\kappa}^m \\ H(\rho) & \chi_{\kappa}^m \end{array}\right].$	$i\gamma^0\vec{\gamma}\cdot\vec{\partial}\left[\begin{array}{cc} iH(\rho) & \chi_{\kappa}^m \\ -F(\rho) & \chi_{-\kappa}^m \end{array}\right]=\frac{\vec{\sigma}\cdot\vec{x}}{ \vec{x} }\left(\partial_r-\frac{\vec{\sigma}\cdot\vec{L}}{r}\right)\left[\begin{array}{cc} iF(\rho) & \chi_{-\kappa}^m \\ H(\rho) & \chi_{\kappa}^m \end{array}\right].$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00592 (9.1.10)	195	■ 0.968	$\stackrel{+}{\Omega}=\frac{\gamma}{\ell^{*2}}\vartheta\wedge\vartheta.$	$\stackrel{+}{\Omega}=\frac{\gamma}{\ell^{*2}}\vartheta\wedge\vartheta.$	$\stackrel{+}{\Omega}=\frac{\gamma}{\ell^{*2}}\vartheta\wedge\vartheta.$
mielke-geometrodynamics_E00611 (9.3.2)	202	■ 0.968	$L_{\mathrm{SKY}}:=-\frac{1}{2}R_{\alpha\beta}\wedge{}^*R^{\alpha\beta},$	$L_{\mathrm{SKY}}:=-\frac{1}{2}R_{\alpha\beta}\wedge{}^*R^{\alpha\beta},$	$L_{\mathrm{SKY}}:=-\frac{1}{2}R_{\alpha\beta}\wedge{}^*R^{\alpha\beta},$
mielke-geometrodynamics_E00841 (12.2.9)	272	■ 0.968	$\mu_{\alpha}=\frac{1}{4}\vartheta_{\alpha}\wedge{}^*j_5.$	$\mu_{\alpha}=\frac{1}{4}\vartheta_{\alpha}\wedge{}^*j_5.$	$\mu_{\alpha}=\frac{1}{4}\vartheta_{\alpha}\wedge{}^*j_5.$
mielke-geometrodynamics_E00636 (10.2.7)	212	■ 0.970	$T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha},$	$T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha},$	$T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha},$
mielke-geometrodynamics_E01137 (C.5)	376	■ 0.970	$3\,c_{[i}{}^pc_{j]k]}{}^s:=c_{is}{}^pc_{jk}{}^s+c_{js}{}^pc_{ki}{}^s+c_{ks}{}^pc_{ij}{}^s\equiv 0.$	$3c_{[i}{}^pc_{j]k]}{}^s:=c_{is}{}^pc_{jk}{}^s+c_{js}{}^pc_{ki}{}^s+c_{ks}{}^pc_{ij}{}^s\equiv 0.$	$3c_{[i}{}^pc_{j]k]}{}^s:=c_{is}{}^pc_{jk}{}^s+c_{js}{}^pc_{ki}{}^s+c_{ks}{}^pc_{ij}{}^s\equiv 0.$
mielke-geometrodynamics_E00415 (6.5.3)	137	■ 0.970	$\begin{aligned} \left\langle 0\right W(\gamma)\left 0\right\rangle ^{(1)}&= \int_{\mathcal{C} \mathcal{G}}\mathcal{D}A\,e^{iC}W(A,\gamma)\\ &= \frac{1}{4\pi}\oint ds\oint dt\dot{\gamma}^{\mu}(s)\dot{\gamma}^{\nu}(t)\eta_{0\mu\nu\kappa}\frac{\gamma^{\kappa}(s)-\gamma^{\kappa}(t)}{ \gamma(s)-\gamma(t) ^3} \end{aligned}$	$\begin{aligned} \left\langle 0\right W(\gamma)\left 0\right\rangle ^{(1)}&= \int_{\mathcal{C} \mathcal{G}}\mathcal{D}A\,e^{iC}W(A,\gamma)\\ &= \frac{1}{4\pi}\oint ds\oint dt\dot{\gamma}^{\mu}(s)\dot{\gamma}^{\nu}(t)\eta_{0\mu\nu\kappa}\frac{\gamma^{\kappa}(s)-\gamma^{\kappa}(t)}{ \gamma(s)-\gamma(t) ^3} \end{aligned}$	$\begin{aligned} \left\langle 0\right W(\gamma)\left 0\right\rangle ^{(1)}&= \int_{\mathcal{C} \mathcal{G}}\mathcal{D}A\,e^{iC}W(A,\gamma)\\ &= \frac{1}{4\pi}\oint ds\oint dt\dot{\gamma}^{\mu}(s)\dot{\gamma}^{\nu}(t)\eta_{0\mu\nu\kappa}\frac{\gamma^{\kappa}(s)-\gamma^{\kappa}(t)}{ \gamma(s)-\gamma(t) ^3} \end{aligned}$
mielke-geometrodynamics_E00495 (7.7.13)	162	■ 0.971	$\{ \}^*\left(\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta}\right)=\eta^{\alpha\beta\gamma\delta},$ where $\eta_{\alpha\beta\gamma\delta}:=+\delta_{\alpha\beta\gamma\delta}^{0123},$	$\left(\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta}\right)=\eta^{\alpha\beta\gamma\delta},$ where $\eta_{\alpha\beta\gamma\delta}:=+\delta_{\alpha\beta\gamma\delta}^{0123},$	$\left(\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta}\right)=\eta^{\alpha\beta\gamma\delta},$ where $\eta_{\alpha\beta\gamma\delta}:=+\delta_{\alpha\beta\gamma\delta}^{0123},$
mielke-geometrodynamics_E01147 (C.16)	377	■ 0.971	$\begin{aligned} \left[\lambda_i,\lambda_j\right]&=2if_{ij}^k\lambda_k\quad\left(c_{ij}^k=2if_{ig}^k\right),\\ \left\{\lambda_i,\lambda_j\right\}_+&=\frac{4}{N}\delta_{ij}+2d_{ij}^k\lambda_k. \end{aligned}$	$\begin{aligned} \left[\lambda_i,\lambda_j\right]&=2if_{ij}^k\lambda_k\quad\left(c_{ij}^k=2if_{ig}^k\right),\\ \left\{\lambda_i,\lambda_j\right\}_+&=\frac{4}{N}\delta_{ij}+2d_{ij}^k\lambda_k. \end{aligned}$	$\begin{aligned} \left[\lambda_i,\lambda_j\right]&=2if_{ij}^k\lambda_k\quad\left(c_{ij}^k=2if_{ig}^k\right),\\ \left\{\lambda_i,\lambda_j\right\}_+&=\frac{4}{N}\delta_{ij}+2d_{ij}^k\lambda_k. \end{aligned}$
mielke-geometrodynamics_E00129 (3.3.30)	61	■ 0.972	$\begin{aligned} j:=\frac{\delta L_{\mathrm{matter}}}{\delta A}&= \frac{\partial L_{\mathrm{matter}}}{\partial A}+d\frac{\partial L_{\mathrm{matter}}}{\partial dA}\\ &= \Psi\wedge\frac{\partial L_{\mathrm{matter}}}{\partial D\Psi}+D\frac{\partial L_{\mathrm{matter}}}{\partial B}, \end{aligned}$	$\begin{aligned} j:=\frac{\delta L_{\mathrm{matter}}}{\delta A}&= \frac{\partial L_{\mathrm{matter}}}{\partial A}+d\frac{\partial L_{\mathrm{matter}}}{\partial dA}\\ &= \Psi\wedge\frac{\partial L_{\mathrm{matter}}}{\partial D\Psi}+D\frac{\partial L_{\mathrm{matter}}}{\partial B}, \end{aligned}$	$\begin{aligned} j:=\frac{\delta L_{\mathrm{matter}}}{\delta A}&= \frac{\partial L_{\mathrm{matter}}}{\partial A}+d\frac{\partial L_{\mathrm{matter}}}{\partial dA}\\ &= \Psi\wedge\frac{\partial L_{\mathrm{matter}}}{\partial D\Psi}+D\frac{\partial L_{\mathrm{matter}}}{\partial B}, \end{aligned}$
mielke-geometrodynamics_E00784 (11.3.3)	248	■ 0.972	$\begin{aligned} L_{\mathrm{NL}}&=\frac{1}{2}\bar{\psi}\left[\widetilde{K},{}^*\gamma\right]\psi=-3i^*[\Theta,\vartheta]\wedge(\bar{\psi}{}^*\gamma\psi)\\ &=-3\tau_s\wedge[\Theta,\vartheta]. \end{aligned}$	$\begin{aligned} L_{\mathrm{NL}}&=\frac{1}{2}\bar{\psi}\left[\widetilde{K},{}^*\gamma\right]\psi=-3i^*[\Theta,\vartheta]\wedge(\bar{\psi}{}^*\gamma\psi)\\ &=-3\tau_s\wedge[\Theta,\vartheta]. \end{aligned}$	$\begin{aligned} L_{\mathrm{NL}}&=\frac{1}{2}\bar{\psi}\left[\widetilde{K},{}^*\gamma\right]\psi=-3i^*[\Theta,\vartheta]\wedge(\bar{\psi}{}^*\gamma\psi)\\ &=-3\tau_s\wedge[\Theta,\vartheta]. \end{aligned}$
mielke-geometrodynamics_E00666 (10.3.23)	218	■ 0.972	$\square\vartheta_{\alpha}+\frac{\ell}{\gamma}{}^*D^*(\delta\ell E_{\alpha}+S_{\alpha})=0,$	$\square\vartheta_{\alpha}+\frac{\ell}{\gamma}{}^*D^*(\delta\ell E_{\alpha}+S_{\alpha})=0,$	$\square\vartheta_{\alpha}+\frac{\ell}{\gamma}{}^*D^*(\delta\ell E_{\alpha}+S_{\alpha})=0,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E08945 (14.1.26)	306	■ 0.972	$\begin{aligned} & \Sigma_{\mathrm{YM}}(\Omega) \wedge \Sigma_{\mathrm{YM}}(\Omega) = \frac{1}{4\alpha_g} \mathrm{Tr} \left\{ -\xi \wedge \xi \wedge (\vartheta \wedge {}^*\xi) \wedge (\vartheta \wedge {}^*\xi) \right. \\ & \left. + \vartheta \wedge \vartheta \wedge {}^*(\xi \wedge {}^*\xi)^2 - \vartheta \wedge {}^*\xi \wedge {}^*(\vartheta \wedge {}^*\xi) \wedge {}^*(\xi \wedge {}^*\xi) \right\} \\ & = \vartheta \wedge \vartheta \wedge {}^*(\Sigma_{\mathrm{YM}}(\xi) \wedge {}^*\Sigma_{\mathrm{YM}}(\xi)) \end{aligned}$	$\Sigma_{\mathrm{YM}}(\Omega) \wedge \Sigma_{\mathrm{YM}}(\Omega) = \frac{1}{4\alpha_g} \mathrm{Tr} \left\{ -\xi \wedge \xi \wedge (\vartheta \wedge {}^*\xi) \wedge (\vartheta \wedge {}^*\xi) \right. \\ \left. + \vartheta \wedge \vartheta \wedge {}^*(\xi \wedge {}^*\xi)^2 - \vartheta \wedge {}^*\xi \wedge {}^*(\vartheta \wedge {}^*\xi) \wedge {}^*(\xi \wedge {}^*\xi) \right\} \\ = \vartheta \wedge \vartheta \wedge {}^*(\Sigma_{\mathrm{YM}}(\xi) \wedge {}^*\Sigma_{\mathrm{YM}}(\xi))$	$\Sigma_{\mathrm{YM}}(\Omega) \wedge \Sigma_{\mathrm{YM}}(\Omega) = \frac{1}{4\alpha_g} \mathrm{Tr} \left\{ -\xi \wedge \xi \wedge (\vartheta \wedge {}^*\xi) \wedge (\vartheta \wedge {}^*\xi) \right. \\ \left. + \vartheta \wedge \vartheta \wedge {}^*(\xi \wedge {}^*\xi)^2 - \vartheta \wedge {}^*\xi \wedge {}^*(\vartheta \wedge {}^*\xi) \wedge {}^*(\xi \wedge {}^*\xi) \right\} \\ = \vartheta \wedge \vartheta \wedge {}^*(\Sigma_{\mathrm{YM}}(\xi) \wedge {}^*\Sigma_{\mathrm{YM}}(\xi))$
mielke-geometrodynamics_E08462 (7.3.1)	153	■ 0.973	$L_{\mathrm{SKY}} = \mathrm{Tr}(\Omega^g \wedge {}^*\Omega^g) = -\frac{1}{2} R^{\alpha\beta} \wedge {}^*R_{\alpha\beta},$	$L_{\mathrm{SKY}} = \mathrm{Tr}(\Omega^g \wedge {}^*\Omega^g) = -\frac{1}{2} R^{\alpha\beta} \wedge {}^*R_{\alpha\beta},$	$L_{\mathrm{SKY}} = \mathrm{Tr}(\Omega^g \wedge {}^*\Omega^g) = -\frac{1}{2} R^{\alpha\beta} \wedge {}^*R_{\alpha\beta},$
mielke-geometrodynamics_E08356 (6.2.7)	124	■ 0.974	$D T^{\alpha} \equiv R_{\beta}^{\alpha} \wedge \vartheta^{\beta}$	$DT^{\alpha} \equiv R_{\beta}^{\alpha} \wedge \vartheta^{\beta}$	$DT^{\alpha} \equiv R_{\beta}^{\alpha} \wedge \vartheta^{\beta}$
mielke-geometrodynamics_E08712 (10.8.1)	228	■ 0.975	$\Psi = \Psi_i dx^i = \Psi_{\alpha} \vartheta^{\alpha}$	$\Psi = \Psi_i dx^i = \Psi_{\alpha} \vartheta^{\alpha}$	$\Psi = \Psi_i dx^i = \Psi_{\alpha} \vartheta^{\alpha}$
mielke-geometrodynamics_E08971 (14.2.12)	314	■ 0.975	$\hat{d} := \left[\begin{array}{c c} d & 0 \\ \hline 0 & \overset{\circ}{d} \end{array} \right],$	$\hat{d} := \left[\begin{array}{c c} d & 0 \\ \hline 0 & \overset{\circ}{d} \end{array} \right],$	$\hat{d} := \left[\begin{array}{c c} d & 0 \\ \hline 0 & \overset{\circ}{d} \end{array} \right],$
mielke-geometrodynamics_E08557 (8.4.11)	181	■ 0.975	$\langle dj_5 \rangle = 2i {}^*m \langle \bar{\psi} \gamma_5 \psi \rangle - \frac{1}{48\pi^2} \left(R_{\alpha\beta}^{\{ \} } \wedge R^{\{\beta\alpha\beta} + \frac{1}{2} d\mathscr{A} \wedge d\mathscr{A} \right),$	$\langle dj_5 \rangle = 2i {}^*m \langle \bar{\psi} \gamma_5 \psi \rangle - \frac{1}{48\pi^2} \left(R_{\alpha\beta}^{\{ \} } \wedge R^{\{\beta\alpha\beta} + \frac{1}{2} d\mathscr{A} \wedge d\mathscr{A} \right),$	$\langle dj_5 \rangle = 2i {}^*m \langle \bar{\psi} \gamma_5 \psi \rangle - \frac{1}{48\pi^2} \left(R_{\alpha\beta}^{\{ \} } \wedge R^{\{\beta\alpha\beta} + \frac{1}{2} d\mathscr{A} \wedge d\mathscr{A} \right),$
mielke-geometrodynamics_E08487 (7.7.4) [re-fined: census]	160	■ 0.977	$\eta_{\alpha\beta} \rightarrow \tilde{\eta}_{\alpha\beta} = \eta_{\alpha\beta} + \ell^2 \eta_{\alpha\beta\gamma} \wedge e_{\mu} {}^*H^{\mu\gamma} + \frac{\ell^4}{2} \eta_{\alpha\beta\gamma\delta} (e_{\mu} {}^*H^{\gamma\mu}) \wedge (e_{\nu} {}^*H^{\delta\nu}).$	$\eta_{\alpha\beta} \rightarrow \tilde{\eta}_{\alpha\beta} = \eta_{\alpha\beta} + \ell^2 \eta_{\alpha\beta\gamma} \wedge e_{\mu} {}^*H^{\mu\gamma} + \frac{\ell^4}{2} \eta_{\alpha\beta\gamma\delta} (e_{\mu} {}^*H^{\gamma\mu}) \wedge (e_{\nu} {}^*H^{\delta\nu}).$	$\eta_{\alpha\beta} \rightarrow \tilde{\eta}_{\alpha\beta} = \eta_{\alpha\beta} + \ell^2 \eta_{\alpha\beta\gamma} \wedge e_{\mu} {}^*H^{\mu\gamma} + \frac{\ell^4}{2} \eta_{\alpha\beta\gamma\delta} (e_{\mu} {}^*H^{\gamma\mu}) \wedge (e_{\nu} {}^*H^{\delta\nu}).$
mielke-geometrodynamics_E01868 (16.2.7)	355	■ 0.978	$\left[\partial_{\rho^*}^2 - V_{\mathrm{eff}}^{\iota}(\rho^*) + \frac{\omega^2}{\beta^2} \right] F = 0$	$\left[\partial_{\rho^*}^2 - V_{\mathrm{eff}}^{\iota}(\rho^*) + \frac{\omega^2}{\beta^2} \right] F = 0$	$\left[\partial_{\rho^*}^2 - V_{\mathrm{eff}}^{\iota}(\rho^*) + \frac{\omega^2}{\beta^2} \right] F = 0$
mielke-geometrodynamics_E01114 (B.26)	371	■ 0.979	$H^{\{p\}}(M) = 0 \text{ for } p > n,$	$H^p(M) = 0 \text{ for } p > n,$	$H^p(M) = 0 \text{ for } p > n,$
mielke-geometrodynamics_E08807 (2.2.2)	33	■ 0.980	$\vartheta = \vartheta^{\alpha} P_{\alpha} \in C^{\infty}(T^*(M)),$	$\vartheta = \vartheta^{\alpha} P_{\alpha} \in C^{\infty}(T^*(M)),$	$\vartheta = \vartheta^{\alpha} P_{\alpha} \in C^{\infty}(T^*(M)),$
mielke-geometrodynamics_E08988 (13.5.13)	291	■ 0.980	$D g_{\alpha\beta} = 0,$	$Dg_{\alpha\beta} = 0,$	$Dg_{\alpha\beta} = 0,$
mielke-geometrodynamics_E08536 (8.2.3)	175	■ 0.980	$\begin{aligned} & (d \oplus s) \tilde{\Gamma} + \frac{1}{2} [\tilde{\Gamma}, \tilde{\Gamma}] = \tilde{\Omega}^g \\ & (d \oplus s) \tilde{\Omega}^g + [\tilde{\Gamma}, \tilde{\Omega}^g] \equiv 0 \end{aligned}$	$\begin{aligned} & (d \oplus s) \tilde{\Gamma} + \frac{1}{2} [\tilde{\Gamma}, \tilde{\Gamma}] = \tilde{\Omega}^g \\ & (d \oplus s) \tilde{\Omega}^g + [\tilde{\Gamma}, \tilde{\Omega}^g] \equiv 0 \end{aligned}$	$\begin{aligned} & (d \oplus s) \tilde{\Gamma} + \frac{1}{2} [\tilde{\Gamma}, \tilde{\Gamma}] = \tilde{\Omega}^g, \\ & (d \oplus s) \tilde{\Omega}^g + [\tilde{\Gamma}, \tilde{\Omega}^g] \equiv 0, \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_292 EQ0911 (13.6.1)	292	■ 0.980	$\begin{aligned} ds^2 &:= \frac{\ell^2}{\mu^2} \left\langle \hat{D}\Phi^A \otimes_s \hat{D}\Phi_A \right\rangle = \frac{\ell^2}{\mu^2} \hat{D}\Phi_0^A \otimes_s \hat{D}\Phi_{A0} \\ &= \ell^2 \Gamma_4^\alpha \otimes_s \Gamma_\alpha^4 = (\vartheta^\alpha - D\xi^\alpha) \otimes_s (\theta_\alpha - D\xi_\alpha^*) \\ &\stackrel{*}{=} \vartheta^\alpha \otimes_s \theta_\alpha = g_{ij} dx^i \otimes_s dx^j \end{aligned}$	$ds^2 := \frac{\ell^2}{\mu^2} \left\langle \hat{D}\Phi^A \otimes_s \hat{D}\Phi_A \right\rangle = \frac{\ell^2}{\mu^2} \hat{D}\Phi_0^A \otimes_s \hat{D}\Phi_{A0}$ $= \ell^2 \Gamma_4^\alpha \otimes_s \Gamma_\alpha^4 = (\vartheta^\alpha - D\xi^\alpha) \otimes_s (\theta_\alpha - D\xi_\alpha^*)$ $\stackrel{*}{=} \vartheta^\alpha \otimes_s \theta_\alpha = g_{ij} dx^i \otimes_s dx^j$	$ds^2 := \frac{\ell^2}{\mu^2} \langle \hat{D}\Phi^A \otimes_s \hat{D}\Phi_A \rangle = \frac{\ell^2}{\mu^2} \hat{D}\Phi_0^A \otimes_s \hat{D}\Phi_{A0}$ $= \ell^2 \Gamma_4^\alpha \otimes_s \Gamma_\alpha^4 = (\vartheta^\alpha - D\xi^\alpha) \otimes_s (\theta_\alpha - D\xi_\alpha^*)$ $\stackrel{*}{=} \vartheta^\alpha \otimes_s \theta_\alpha = g_{ij} dx^i \otimes_s dx^j .$
mielke-geometrodynamics_304 EQ0931 (14.1.12)	304	■ 0.981	$\Omega \rightarrow e^{*\delta} \Omega := \Omega \cos \delta + (-1)^{z*} \Omega \sin \delta.$	$\Omega \rightarrow e^{*\delta} \Omega := \Omega \cos \delta + (-1)^{z*} \Omega \sin \delta.$	$\Omega \rightarrow e^{*\delta} \Omega := \Omega \cos \delta + (-1)^z {}^* \Omega \sin \delta.$
mielke-geometrodynamics_219 EQ0671 (10.3.28)	219	■ 0.982	$\check{\mathscr{H}}_\alpha^* := dH_\alpha^* \cong \tau_\alpha^* + E_\alpha^* - (-1)^s \eta_{\alpha\beta\gamma} \Gamma^{*\beta} \wedge H^{*\gamma},$	$\check{\mathscr{H}}_\alpha^* := dH_\alpha^* \cong \tau_\alpha^* + E_\alpha^* - (-1)^s \eta_{\alpha\beta\gamma} \Gamma^{*\beta} \wedge H^{*\gamma},$	$\check{\mathscr{H}}_\alpha^* := dH_\alpha^* \cong \tau_\alpha^* + E_\alpha^* - (-1)^s \eta_{\alpha\beta\gamma} \Gamma^{*\beta} \wedge H^{*\gamma},$
mielke-geometrodynamics_366 EQ1086 (A.12)	366	■ 0.982	$\begin{aligned} \mathrm{C}(1,1) &= \mathrm{C}(2,0) = \mathrm{GL}(2, \mathbb{R}), \\ \mathrm{C}(0,2) &= \mathbb{H} \quad (\text{field of quaternions}), \end{aligned}$	$\begin{aligned} \mathrm{C}(1,1) &= \mathrm{C}(2,0) = \mathrm{GL}(2, \mathbb{R}), \\ \mathrm{C}(0,2) &= \mathbb{H} \quad (\text{field of quaternions}), \end{aligned}$	$\begin{aligned} \mathrm{C}(1,1) &= \mathrm{C}(2,0) = \mathrm{GL}(2, \mathbb{R}), \\ \mathrm{C}(0,2) &= \mathbb{H} \quad (\text{field of quaternions}), \end{aligned}$
mielke-geometrodynamics_88 EQ0229 (4.2.22)	88	■ 0.982	$\Theta := D\vartheta = d\vartheta - [\omega^L, \vartheta]$	$\Theta := D\vartheta = d\vartheta - [\omega^L, \vartheta]$	$\Theta := D\vartheta = d\vartheta - [\omega^L, \vartheta]$
mielke-geometrodynamics_271 EQ0836 (12.2.4)	271	■ 0.983	$\mathscr{A} := \frac{1}{4} {}^* \mathrm{Tr}(\gamma \wedge D\gamma) = {}^*(\vartheta^\alpha \wedge T_\alpha) = \frac{1}{2} T^{[\alpha\beta\gamma]} \eta_{\alpha\beta\gamma} = \mathscr{A}_i dx^i,$	$\mathscr{A} := \frac{1}{4} {}^* \mathrm{Tr}(\gamma \wedge D\gamma) = {}^*(\vartheta^\alpha \wedge T_\alpha) = \frac{1}{2} T^{[\alpha\beta\gamma]} \eta_{\alpha\beta\gamma} = \mathscr{A}_i dx^i,$	$\mathscr{A} := \frac{1}{4} {}^* Tr(\gamma \wedge D\gamma) = {}^*(\vartheta^\alpha \wedge T_\alpha) = \frac{1}{2} T^{[\alpha\beta\gamma]} \eta_{\alpha\beta\gamma} = \mathscr{A}_i dx^i ,$
mielke-geometrodynamics_165 EQ0511 (7.7.29)	165	■ 0.983	$\begin{aligned} \Theta &:= D\gamma = d\gamma + [\Gamma, \gamma] = (d\vartheta^\alpha + \Gamma_\beta^\alpha \wedge \vartheta^\beta) \gamma_\alpha \\ &= T^\alpha \gamma_\alpha = \frac{1}{2} T_{ij}^\alpha \gamma_\alpha dx^i \wedge dx^j \end{aligned}$	$\begin{aligned} \Theta &:= D\gamma = d\gamma + [\Gamma, \gamma] = (d\vartheta^\alpha + \Gamma_\beta^\alpha \wedge \vartheta^\beta) \gamma_\alpha \\ &= T^\alpha \gamma_\alpha = \frac{1}{2} T_{ij}^\alpha \gamma_\alpha dx^i \wedge dx^j \end{aligned}$	$\begin{aligned} \Theta &:= D\gamma = d\gamma + [\Gamma, \gamma] = (d\vartheta^\alpha + \Gamma_\beta^\alpha \wedge \vartheta^\beta) \gamma_\alpha \\ &= T^\alpha \gamma_\alpha = \frac{1}{2} T_{ij}^\alpha \gamma_\alpha dx^i \wedge dx^j , \end{aligned}$
mielke-geometrodynamics_212 EQ0637 (10.2.8)	212	■ 0.983	$\Gamma_\alpha^* = \Gamma_\alpha^{\{\}}{}^* - K_\alpha^*$	$\Gamma_\alpha^* = \Gamma_\alpha^{\{\}}{}^* - K_\alpha^*$	$\Gamma_\alpha^* = \Gamma_\alpha^{\{\}}{}^* - K_\alpha^*$
mielke-geometrodynamics_96 EQ0271 (4.3.36)	96	■ 0.983	$[D_\alpha, D_\beta] = R_{\alpha\beta}^{..cd} L_{cd} - T_{\alpha\beta}^{..c} D_c,$	$[D_\alpha, D_\beta] = R_{\alpha\beta}^{..cd} L_{cd} - T_{\alpha\beta}^{..c} D_c,$	$[D_\alpha, D_\beta] = R_{\alpha\beta}^{..cd} L_{cd} - T_{\alpha\beta}^{..c} D_c,$
mielke-geometrodynamics_181 EQ0561 (8.5.4)	181	■ 0.984	$\{\}^* \sigma^{\{\}} = (-1)^{\mathrm{sig}} \sigma,$	${}^* \sigma^{(*)} = (-1)^{\mathrm{sig}} \sigma,$	${}^* \sigma^{(*)} = (-1)^{\mathrm{sig}} \sigma ,$
mielke-geometrodynamics_56 EQ0097 (3.2.7)	56	■ 0.985	$D \rightarrow {}^{G^{-1}} D = G^{-1} D G.$	$D \rightarrow {}^{G^{-1}} D = G^{-1} D G.$	$D \rightarrow {}^{G^{-1}} D = G^{-1} D G.$
mielke-geometrodynamics_152 EQ0459 (7.2.11)	152	■ 0.985	$\Omega_{\mathrm{C}} = \stackrel{(+)}{\Omega}_{\mathrm{C}} + \stackrel{(-)}{\Omega}_{\mathrm{C}}, \quad \stackrel{(\pm)}{\Omega}_{\mathrm{C}} := \frac{1}{2} \left(\Omega_{\mathrm{C}} \pm \Omega_{\mathrm{C}}^{(*)} \right)$	$\Omega_{\mathrm{C}} = \stackrel{(+)}{\Omega}_{\mathrm{C}} + \stackrel{(-)}{\Omega}_{\mathrm{C}}, \quad \stackrel{(\pm)}{\Omega}_{\mathrm{C}} := \frac{1}{2} \left(\Omega_{\mathrm{C}} \pm \Omega_{\mathrm{C}}^{(*)} \right)$	$\Omega_{\mathrm{C}} = \stackrel{(+)}{\Omega}_{\mathrm{C}} + \stackrel{(-)}{\Omega}_{\mathrm{C}}, \quad \stackrel{(\pm)}{\Omega}_{\mathrm{C}} := \frac{1}{2} \left(\Omega_{\mathrm{C}} \pm \Omega_{\mathrm{C}}^{(*)} \right)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometroynamics_344 EQ1051 (15.2.8)		■ 0.985	$\begin{aligned} A_i &= \frac{1}{2} \left\{ A_i^{(f)j} + {}^* A_i^{(f)j} \gamma^5 \right\} \lambda_j^{(f)} \\ &\oplus \frac{1}{2} \left\{ A_i^{(c)j} + {}^* A_i^{(c)j} \gamma^5 \right\} \lambda_j^{(c)}, \end{aligned}$	$A_i = \frac{1}{2} \left\{ A_i^{(f)j} + {}^* A_i^{(f)j} \gamma^5 \right\} \lambda_j^{(f)} \\ \oplus \frac{1}{2} \left\{ A_i^{(c)j} + {}^* A_i^{(c)j} \gamma^5 \right\} \lambda_j^{(c)},$	$A_i = \frac{1}{2} \left\{ A_i^{(f)j} + {}^* A_i^{(f)j} \gamma^5 \right\} \lambda_j^{(f)} \\ \oplus \frac{1}{2} \left\{ A_i^{(c)j} + {}^* A_i^{(c)j} \gamma^5 \right\} \lambda_j^{(c)},$
mielke-geometroynamics_369 EQ1096 (B.6)		■ 0.985	$d \, d \, \alpha^{(p)} \equiv 0.$	$dd\alpha^{(p)} \equiv 0.$	$d \, d \alpha^{(p)} \equiv 0.$
mielke-geometroynamics_81 EQ0191 (4.1.1)		■ 0.985	$x \mapsto x' = g x + a, \quad g \in GL(n, \mathbb{R}), \quad a \in \mathbb{R}^n.$	$x \rightarrow x' = gx + a, \quad g \in GL(n, \mathbb{R}), a \in \mathbb{R}^n.$	$x \rightarrow x' = gx + a, \quad g \in GL(n, \mathbb{R}), \quad a \in \mathbb{R}^n.$
mielke-geometroynamics_224 EQ0694 (10.5.5)		■ 0.985	$\begin{aligned} L_{\mathrm{D}} &= \frac{i}{2} \left(\bar{\psi}^* \gamma \wedge D \psi + D \bar{\psi} \wedge {}^* \gamma \psi \right) - m \bar{\psi} \psi \eta \\ &= \bar{\psi} \left(i^* \gamma \wedge D - m \eta \right) \psi + d \left(\frac{i}{2} \bar{\psi}^* \gamma \psi \right), \end{aligned}$	$L_{\mathrm{D}} = \frac{i}{2} \left(\bar{\psi}^* \gamma \wedge D \psi + D \bar{\psi} \wedge {}^* \gamma \psi \right) - m \bar{\psi} \psi \eta \\ = \bar{\psi} \left(i^* \gamma \wedge D - m \eta \right) \psi + d \left(\frac{i}{2} \bar{\psi}^* \gamma \psi \right),$	$L_{\mathrm{D}} = \frac{i}{2} \left(\bar{\psi}^* \gamma \wedge D \psi + D \bar{\psi} \wedge {}^* \gamma \psi \right) - m \bar{\psi} \psi \eta \\ = \bar{\psi} \left(i^* \gamma \wedge D - m \eta \right) \psi + d \left(\frac{i}{2} \bar{\psi}^* \gamma \psi \right),$
mielke-geometroynamics_180 EQ0554 (8.4.8)		■ 0.986	$D^* \Omega = 0$	$D^* \Omega^{\pm g} = 0$	$D^* \Omega^{\pm g} = 0$
mielke-geometroynamics_45 EQ0058 (2.7.8)		■ 0.987	$b \mapsto {}^G b := G b \quad \in C^\infty(V^\rho).$	$b \rightarrow {}^G b := G b \quad \in C^\infty(V^\rho).$	$b \rightarrow {}^G b := G b \quad \in C^\infty(V^\rho).$
mielke-geometroynamics_99 EQ0285 (4.4.14)		■ 0.987	$D \left(\vartheta \wedge {}^* \Omega^g \right) = \Theta \wedge {}^* \Omega^g$	$D \left(\vartheta \wedge {}^* \Omega^g \right) = \Theta \wedge {}^* \Omega^g$	$D(\vartheta \wedge {}^* \Omega^g) = \Theta \wedge {}^* \Omega^g$
mielke-geometroynamics_271 EQ0834 (12.2.2)		■ 0.988	$i^* \gamma \wedge \left(\mathscr{D} + \frac{i}{4} m \gamma - \frac{1}{2} T \right) \psi = 0,$	$i^* \gamma \wedge \left(\mathscr{D} + \frac{i}{4} m \gamma - \frac{1}{2} T \right) \psi = 0,$	$i^* \gamma \wedge \left(\mathscr{D} + \frac{i}{4} m \gamma - \frac{1}{2} T \right) \psi = 0,$
mielke-geometroynamics_31 EQ0002 (2.1.1)		■ 0.989	$x_2 \circ \bar{x}_1 : \mathbb{R} \rightarrow \mathbb{R}$	$x_2 \circ \bar{x}_1 : \mathbb{R} \rightarrow \mathbb{R}$	$x_2 \circ \bar{x}_1 : \mathbb{R} \rightarrow \mathbb{R}$
mielke-geometroynamics_114 EQ0339 (5.3.16)		■ 0.989	$L_{\mathrm{EC}} := -\frac{1}{2\kappa} R^{\alpha\beta} \wedge \eta_{\alpha\beta} = L_{\mathrm{HE}} + \frac{1}{12\kappa} \mathscr{A}^* \mathscr{A}$	$L_{\mathrm{EC}} := -\frac{1}{2\kappa} R^{\alpha\beta} \wedge \eta_{\alpha\beta} = L_{\mathrm{HE}} + \frac{1}{12\kappa} \mathscr{A}^* \mathscr{A}$	$L_{\mathrm{EC}} := -\frac{1}{2\kappa} R^{\alpha\beta} \wedge \eta_{\alpha\beta} = L_{\mathrm{HE}} + \frac{1}{12\kappa} \mathscr{A} \wedge {}^* \mathscr{A}$
mielke-geometroynamics_39 EQ0024 (2.4.6)		■ 0.990	$[L_i, L_j] = c_{ij}^k L_k.$	$[L_i, L_j] = c_{ij}^k L_k.$	$[L_i, L_j] = c_{ij}^k L_k.$
mielke-geometroynamics_164 EQ0507 (7.7.25)		■ 0.990	$\sigma_{\pm} := (\sigma \pm i^* \sigma) / 2 = \frac{1}{2} (1 \mp \gamma_5) \sigma, \quad \text{with} \quad i^* \sigma_{\pm} = \pm \sigma_{\pm}$	$\sigma_{\pm} := (\sigma \pm i^* \sigma) / 2 = \frac{1}{2} (1 \mp \gamma_5) \sigma, \quad \text{with} \quad i^* \sigma_{\pm} = \pm \sigma_{\pm}$	$\sigma_{\pm} := (\sigma \pm i^* \sigma) / 2 = \frac{1}{2} (1 \mp \gamma_5) \sigma, \quad \text{with} \quad i^* \sigma_{\pm} = \pm \sigma_{\pm}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_008572 (8.6.10) [re-fined: census]	184	■ 0.990	$\begin{aligned} H_{\{\alpha\}}(* *) &= \frac{\theta_{\{\alpha\}}}{\ell^2} K^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} - \frac{\theta_{\{\alpha\}}}{\ell^2} T_{\alpha} = \frac{1}{\ell^2} \left[\theta_{\{\alpha\}}^* K_{\alpha}^{(*)} - \theta_{\{\alpha\}} T_{\alpha} \right] \\ &= \frac{\theta_{\{\alpha\}}^*}{\ell^2} * \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta}] T^{\beta}) - \frac{1}{2} e_{\alpha}] (T^{\beta} \wedge \vartheta_{\beta}) \right] - \frac{\theta_{\{\alpha\}}}{\ell^2} T_{\alpha} \\ &\simeq -\partial L_{\theta} / \partial T^{\alpha}. \end{aligned}$	$H_{\alpha}(**) = \frac{\theta_{\mathrm{T}}^*}{2\ell^2} K^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha} = \frac{1}{\ell^2} \left[\theta_{\mathrm{T}}^* K_{\alpha}^{(*)} - \theta_{\mathrm{T}} T_{\alpha} \right]$ $= \frac{\theta_{\mathrm{T}}^*}{\ell^2} * \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta}] T^{\beta}) - \frac{1}{2} e_{\alpha}] (T^{\beta} \wedge \vartheta_{\beta}) \right] - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha}$ $\simeq -\partial L_{\theta} / \partial T^{\alpha}.$	$H_{\alpha}(**) = \frac{\theta_{\mathrm{T}}^*}{2\ell^2} K^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha} = \frac{1}{\ell^2} \left[\theta_{\mathrm{T}}^* K_{\alpha}^{(*)} - \theta_{\mathrm{T}} T_{\alpha} \right]$ $= \frac{\theta_{\mathrm{T}}^*}{\ell^2} * \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta}] T^{\beta}) - \frac{1}{2} e_{\alpha}] (T^{\beta} \wedge \vartheta_{\beta}) \right] - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha}$ $\simeq -\partial L_{\theta} / \partial T^{\alpha}.$
mielke-geometrodynamics_00110 (3.3.11)	58	■ 0.990	$\begin{aligned} F := & E \wedge dt + B \\ &= (E_x dx + E_y dy + E_z dz) \wedge dt \\ &+ B_x dy \wedge dz + B_y dz \wedge dx + B_z dx \wedge dy. \end{aligned}$	$F := E \wedge dt + B$ $= (E_x dx + E_y dy + E_z dz) \wedge dt$ $+ B_x dy \wedge dz + B_y dz \wedge dx + B_z dx \wedge dy.$	$F := E \wedge dt + B$ $= (E_x dx + E_y dy + E_z dz) \wedge dt$ $+ B_x dy \wedge dz + B_y dz \wedge dx + B_z dx \wedge dy.$
mielke-geometrodynamics_00142 (3.4.13)	64	■ 0.990	$\begin{aligned} D \tau &= D D \Pi^{\Omega} = [\Omega, \Pi^{\Omega}] \\ &= \frac{2}{\alpha_g} \left\{ [\Omega, * \Omega] \frac{\partial L}{\partial \mathcal{F}} + (-1)^s [\Omega, \Omega] \frac{\partial L}{\partial * \mathcal{F}} \right\} = 0. \end{aligned}$	$D\tau = DD\Pi^{\Omega} = [\Omega, \Pi^{\Omega}]$ $= \frac{2}{\alpha_g} \left\{ [\Omega, * \Omega] \frac{\partial L}{\partial \mathcal{F}} + (-1)^s [\Omega, \Omega] \frac{\partial L}{\partial * \mathcal{F}} \right\} = 0.$	$D\tau = DD\Pi^{\Omega} = [\Omega, \Pi^{\Omega}]$ $= \frac{2}{\alpha_g} \left\{ [\Omega, * \Omega] \frac{\partial L}{\partial \mathcal{F}} + (-1)^s [\Omega, \Omega] \frac{\partial L}{\partial * \mathcal{F}} \right\} = 0.$
mielke-geometrodynamics_00155 (3.5.3)	67	■ 0.990	$\stackrel{*}{\sigma} \Omega = \frac{1}{4} F_{\alpha\beta\mu\nu} \sigma^{\mu\nu} \otimes \vartheta^{\alpha} \wedge \vartheta^{\beta}$	$\stackrel{*}{\sigma} \Omega = \frac{1}{4} F_{\alpha\beta\mu\nu} \sigma^{\mu\nu} \otimes \vartheta^{\alpha} \wedge \vartheta^{\beta}$	$\stackrel{*}{\sigma} \Omega = \frac{1}{4} F_{\alpha\beta\mu\nu} \sigma^{\mu\nu} \otimes \vartheta^{\alpha} \wedge \vartheta^{\beta}$
mielke-geometrodynamics_00682 (10.3.40)	221	■ 0.990	$\theta_{\mathrm{L}} C_{\mathrm{L}} - \tilde{\theta}_{\mathrm{L}} \tilde{C}_{\mathrm{L}} = L_{\mathrm{MB}} + \frac{\Delta \theta_{\mathrm{TL}}}{\ell} d(\Gamma^{*\alpha} \wedge \vartheta_{\alpha}),$	$\theta_{\mathrm{L}} C_{\mathrm{L}} - \tilde{\theta}_{\mathrm{L}} \tilde{C}_{\mathrm{L}} = L_{\mathrm{MB}} + \frac{\Delta \theta_{\mathrm{TL}}}{\ell} d(\Gamma^{*\alpha} \wedge \vartheta_{\alpha}),$	$\theta_{\mathrm{L}} C_{\mathrm{L}} - \tilde{\theta}_{\mathrm{L}} \tilde{C}_{\mathrm{L}} = L_{\mathrm{MB}} + \frac{\Delta \theta_{\mathrm{TL}}}{\ell} d(\Gamma^{*\alpha} \wedge \vartheta_{\alpha}),$
mielke-geometrodynamics_00874 (13.3.2) [re-fined: census]	284	■ 0.990	$\cancel{\{Z^A\}_{\{B\}} \cancel{\{Z^C\}_{\{D\}}} = \delta_{\{D\}}^A \{Z^C\}_{\{B\}} - \delta_{\{B\}}^C \{Z^A\}_{\{D\}}$	$\not\!\! [Z^A{}_B \not\!\! Z^C{}_D] = \delta_D^A Z^C{}_B - \delta_B^C Z^A{}_D$	$[\not\!\! E^A{}_B, \not\!\! E^C{}_D] = \delta_D^A \not\!\! E^C{}_B - \delta_B^C \not\!\! E^A{}_D$
mielke-geometrodynamics_00842 (2.6.5)	42	■ 0.991	$F_{\{\alpha\}\{\beta\}} = \partial_{\alpha} \Gamma_{\beta}^j - \partial_{\beta} \Gamma_{\alpha}^j - c_{kl}^j \Gamma_{\alpha}^k \Gamma_{\beta}^l,$	$F_{\alpha\beta}{}^j = \partial_{\alpha} \Gamma_{\beta}{}^j - \partial_{\beta} \Gamma_{\alpha}{}^j - c_{kl}{}^j \Gamma_{\alpha}{}^k \Gamma_{\beta}{}^l,$	$F_{\alpha\beta}{}^j = \partial_{\alpha} \Gamma_{\beta}{}^j - \partial_{\beta} \Gamma_{\alpha}{}^j - c_{kl}{}^j \Gamma_{\alpha}{}^k \Gamma_{\beta}{}^l,$
mielke-geometrodynamics_00233 (4.2.26)	88	■ 0.991	$\Omega^{\mathrm{T}} = f \Theta + df \wedge \vartheta - \Omega^{\mathrm{L}} \otimes \widetilde{\varphi}$	$\Omega^{\mathrm{T}} = f \Theta + df \wedge \vartheta - \Omega^{\mathrm{L}} \otimes \widetilde{\varphi}$	$\Omega^{\mathrm{T}} = f \Theta + df \wedge \vartheta - \Omega^{\mathrm{L}} \otimes \widetilde{\varphi}$
mielke-geometrodynamics_00778	246	■ 0.991	$\langle -\infty D \tau_A \infty \rangle^{(+)} = \frac{1}{(4.)8\pi^2} \mathrm{Tr} \left(\Omega^g \wedge \Omega^{g(*)} \right).$	$\langle -\infty D \tau_A \infty \rangle^{(+)} = \frac{1}{(4.)8\pi^2} \mathrm{Tr} \left(\Omega^g \wedge \Omega^{g(*)} \right).$	$\langle -\infty D \tau_A \infty \rangle^{(+)} = \frac{1}{(4.)8\pi^2} \mathrm{Tr} (\Omega^g \wedge \Omega^{g(*)}).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00644 (10.3.1)	214	■ 0.991	$\widetilde{\Gamma}_{\alpha}^{\star}=\Gamma_{\alpha}^{\star}-(-1)^s\frac{\varepsilon}{2\ell}\vartheta_{\alpha},$	$\widetilde{\Gamma}_{\alpha}^{\star}=\Gamma_{\alpha}^{\star}-(-1)^s\frac{\varepsilon}{2\ell}\vartheta_{\alpha},$	$\widetilde{\Gamma}_{\alpha}^{\star}=\Gamma_{\alpha}^{\star}-(-1)^s\frac{\varepsilon}{2\ell}\vartheta_{\alpha},$
mielke-geometrodynamics_E00149 (3.4.20)	65	■ 0.992	$L(\mathscr{F},\{ \}^{\ast}\mathscr{F}\backslash\mathrm{right})=-\mathscr{F}.$	$L(\mathscr{F},\ast\mathscr{F})=-\mathscr{F}.$	$L(\mathscr{F},\ast\mathscr{F})=-\mathscr{F}.$
mielke-geometrodynamics_E00368 (6.2.19)	127	■ 0.992	$\stackrel{\scriptscriptstyle\pm}{L}_{\parallel}:=L_{\parallel}\pm i\frac{2\ell^2}{\kappa}dC_{\mathrm{TT}}=L_{\mathrm{HE}}-L_{\mathrm{EC}}\pm i\frac{2\ell^2}{\kappa}d\stackrel{\scriptscriptstyle(\mp)}{C}_{\mathrm{TT}}.$	$\stackrel{\scriptscriptstyle(\pm)}{L}_{\parallel}:=L_{\parallel}\pm i\frac{2\ell^2}{\kappa}dC_{\mathrm{TT}}=L_{\mathrm{HE}}-L_{\mathrm{EC}}\pm i\frac{2\ell^2}{\kappa}d\stackrel{\scriptscriptstyle(\mp)}{C}_{\mathrm{TT}}.$	$\stackrel{\scriptscriptstyle(\pm)}{L}_{\parallel}:=L_{\parallel}\pm i\frac{2\ell^2}{\kappa}dC_{\mathrm{TT}}=L_{\mathrm{HE}}-L_{\mathrm{EC}}\pm i\frac{2\ell^2}{\kappa}d\stackrel{\scriptscriptstyle(\mp)}{C}_{\mathrm{TT}}.$
mielke-geometrodynamics_E00449 (7.2.2)	151	■ 0.992	$\{ \}^{\ast}\Omega^{\mathrm{g}(\star)}:=\frac{1}{4}\{ \}^{\ast}R_{\alpha\beta}^{\left(\star\right)cd}L_{cd}\otimes\vartheta^{\alpha}\wedge\vartheta^{\beta},$	$\ast\Omega^{g(\ast)}:=\frac{1}{4}\ast R_{\alpha\beta}^{\left(\star\right)cd}L_{cd}\otimes\vartheta^{\alpha}\wedge\vartheta^{\beta},$	$\ast\Omega^{g(\ast)}:=\frac{1}{4}\ast R_{\alpha\beta}^{\left(\star\right)cd}L_{cd}\otimes\vartheta^{\alpha}\wedge\vartheta^{\beta},$
mielke-geometrodynamics_E01050 (15.2.7)	344	■ 0.992	$\left\{\bar{L}_j=\sigma^{\alpha\beta}\otimes\lambda_i,\quad\lambda_i,\quad i\gamma^5\lambda_i\mid j=1,\dots 8N^2-1\right\},$	$\left\{\bar{L}_j=\sigma^{\alpha\beta}\otimes\lambda_i,\quad\lambda_i,\quad i\gamma^5\lambda_i\mid j=1,\dots 8N^2-1\right\},$	$\left\{\bar{L}_j=\sigma^{\alpha\beta}\otimes\lambda_i,\quad\lambda_i,\quad i\gamma^5\lambda_i\mid j=1,\dots 8N^2-1\right\},$
mielke-geometrodynamics_E00964 (14.2.4)	313	■ 0.992	$\stackrel{\scriptscriptstyle\circ}{\omega}=\stackrel{\scriptscriptstyle\circ}{b}\stackrel{\scriptscriptstyle\circ}{\vartheta}.$	$\stackrel{\scriptscriptstyle\circ}{\omega}=\stackrel{\scriptscriptstyle\circ}{b}\stackrel{\scriptscriptstyle\circ}{\vartheta}.$	$\stackrel{\scriptscriptstyle\circ}{\omega}=\stackrel{\scriptscriptstyle\circ}{b}\stackrel{\scriptscriptstyle\circ}{\vartheta}.$
mielke-geometrodynamics_E00256 (4.3.21)	93	■ 0.992	$R_{\alpha\beta c}^{\cdots d}=2\left\{\partial_{[\alpha}\Gamma_{\beta]c}^{\cdots d}+\Gamma_{[\alpha h}^{\cdots d}\Gamma_{ \beta]c}^{\cdots h}\right\}$	$R_{\alpha\beta c}^{\cdots d}=2\left\{\partial_{[\alpha}\Gamma_{\beta]c}^{\cdots d}+\Gamma_{[\alpha h}^{\cdots d}\Gamma_{ \beta]c}^{\cdots h}\right\}$	$R_{\alpha\beta c}^{\cdots d}=2\left\{\partial_{[\alpha}\Gamma_{\beta]c}^{\cdots d}+\Gamma_{[\alpha h}^{\cdots d}\Gamma_{ \beta]c}^{\cdots h}\right\}$
mielke-geometrodynamics_E00245 (4.3.10)	91	■ 0.992	$\vartheta\wedge^*\vartheta=\eta.$	$\vartheta\wedge^*\vartheta=\eta.$	$\vartheta\wedge^*\vartheta=\eta.$
mielke-geometrodynamics_E00633 (10.2.4)	212	■ 0.992	$-(-1)^s\theta_{\mathrm{TL}}T_{\alpha}-\frac{1}{2\ell}\theta_{\mathrm{T}}\eta_{\alpha}-\theta_{\mathrm{L}}\ell R_{\alpha}^{\star}=\ell\tau_{\alpha}^{\star};$	$-(-1)^s\theta_{\mathrm{TL}}T_{\alpha}-\frac{1}{2\ell}\theta_{\mathrm{T}}\eta_{\alpha}-\theta_{\mathrm{L}}\ell R_{\alpha}^{\star}=\ell\tau_{\alpha}^{\star};$	$-(-1)^s\theta_{\mathrm{TL}}T_{\alpha}-\frac{1}{2\ell}\theta_{\mathrm{T}}\eta_{\alpha}-\theta_{\mathrm{L}}\ell R_{\alpha}^{\star}=\ell\tau_{\alpha}^{\star};$
mielke-geometrodynamics_E00701 (10.5.12)	225	■ 0.992	$\hat{D}^*\hat{D}\psi-m^2\eta\psi+i^*\sigma\wedge\left(\hat{R}+T^{\alpha}\hat{D}_{\alpha}\right)\psi=0,$	$\hat{D}^*\hat{D}\psi-m^2\eta\psi+i^*\sigma\wedge\left(\hat{R}+T^{\alpha}\hat{D}_{\alpha}\right)\psi=0,$	$\hat{D}^*\hat{D}\psi-m^2\eta\psi+i^*\sigma\wedge\left(\hat{R}+T^{\alpha}\hat{D}_{\alpha}\right)\psi=0,$
mielke-geometrodynamics_E00847 (12.2.15)	273	■ 0.992	$\mathscr{A}_{\alpha}\mathscr{A}^{\alpha}\eta=\mathscr{A}\wedge^*\mathscr{A}\cong(\ell^4/4)^*j_5\wedge j_5=0.$	$\mathscr{A}_{\alpha}\mathscr{A}^{\alpha}\eta=\mathscr{A}\wedge^*\mathscr{A}\cong(\ell^4/4)^*j_5\wedge j_5=0.$	$\mathscr{A}_{\alpha}\mathscr{A}^{\alpha}\eta=\mathscr{A}\wedge^*\mathscr{A}\cong(\ell^4/4)^*j_5\wedge j_5=0.$
mielke-geometrodynamics_E00388 (6.4.3) [re-fined: census]	130	■ 0.993	$\underline{\Psi}:=n\rfloor(dt\wedge\Psi)=\left(1-\stackrel{\scriptscriptstyle\perp}{}\right)\Psi,\quad n\rfloor\underline{\Psi}\equiv 0.$	$\underline{\Psi}:=n\rfloor(dt\wedge\Psi)=\left(1-\stackrel{\scriptscriptstyle\perp}{}\right)\Psi,\quad n\rfloor\underline{\Psi}\equiv 0.$	$\underline{\Psi}:=n\rfloor(dt\wedge\Psi)=\left(1-\stackrel{\scriptscriptstyle\perp}{}\right)\Psi,\quad n\rfloor\underline{\Psi}\equiv 0.$
mielke-geometrodynamics_E00228 (4.2.21)	88	■ 0.993	$\widetilde{\vartheta}=f\vartheta.$	$\widetilde{\vartheta}=f\vartheta.$	$\widetilde{\vartheta}=f\vartheta.$
mielke-geometrodynamics_E00287 (4.4.16)	99	■ 0.993	$DD\Pi^{\mathrm{L}}+D\Pi_{\square}^{\mathrm{T}}=D\tau_s.$	$DD\Pi^{\mathrm{L}}+D\Pi_{\square}^{\mathrm{T}}=D\tau_s.$	$D\,D\Pi^{\mathrm{L}}+D\Pi_{\square}^{\mathrm{T}}=D\tau_s.$
mielke-geometrodynamics_E00320 (5.2.4)	111	■ 0.993	$\mathscr{A}:=\frac{1}{4}\ast\mathrm{Tr}(\gamma\wedge D\gamma)=\ast(\vartheta^{\alpha}\wedge T_{\alpha})=\frac{1}{2}T^{[\alpha\beta\gamma]}\eta_{\alpha\beta\gamma}=\mathscr{A}_idx^i,$	$\mathscr{A}:=\frac{1}{4}\ast\mathrm{Tr}(\gamma\wedge D\gamma)=\ast(\vartheta^{\alpha}\wedge T_{\alpha})=\frac{1}{2}T^{[\alpha\beta\gamma]}\eta_{\alpha\beta\gamma}=\mathscr{A}_idx^i,$	$\mathscr{A}:=\frac{1}{4}\ast Tr(\gamma\wedge D\gamma)=\ast(\vartheta^{\alpha}\wedge T_{\alpha})=\frac{1}{2}T^{[\alpha\beta\gamma]}\eta_{\alpha\beta\gamma}=\mathscr{A}_idx^i,$
mielke-geometrodynamics_E00100 (3.3.1)	56	■ 0.994	$F:=dA=\frac{1}{2}F_{ij}dx^i\wedge dx^j.$	$F:=dA=\frac{1}{2}F_{ij}dx^i\wedge dx^j.$	$F:=dA=\frac{1}{2}F_{ij}dx^i\wedge dx^j.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_321 EQ0995 (14.3.8)		■ 0.995	$L_{\text{anom.}}=\frac{\widehat{\ell}}{2}\bar{\psi}\gamma\wedge\gamma\wedge{}^*\Omega\psi=\frac{\ell^*}{4\sqrt{\alpha_g}}\bar{\psi}F_{\mu\nu}\gamma^5[\gamma^\mu,\gamma^\nu]\psi\eta.$	$L_{\text{anom.}}=\frac{\widehat{\ell}}{2}\bar{\psi}\gamma\wedge\gamma\wedge{}^*\Omega\psi=\frac{\ell^*}{4\sqrt{\alpha_g}}\bar{\psi}F_{\mu\nu}\gamma^5[\gamma^\mu,\gamma^\nu]\psi\eta.$	
mielke-geometrodynamics_56 EQ0098 (3.2.8)		■ 0.995	$\Omega\rightarrow{}^{G^{-1}}\Omega=G^{-1}\Omega G.$	$\Omega\rightarrow{}^{G^{-1}}\Omega=G^{-1}\Omega G.$	
mielke-geometrodynamics_304 EQ0930 (14.1.11)		■ 0.995	$\mathrm{R}=\{ \wedge^*\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)\wedge\vartheta\wedge\vartheta\right)=0$	$\mathrm{R}=\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)=0$	
mielke-geometrodynamics_338 EQ1033 (15.1.10)		■ 0.995	$\frac{1}{2}\{ \wedge^*\left(\vartheta^f\wedge\Omega^f\right)+\Lambda_{\mathrm{eff}}^f\vartheta^f\wedge\vartheta^f\}=0$	$\frac{1}{2}\left(\vartheta^f\wedge\Omega^f\right)+\Lambda_{\mathrm{eff}}^f\vartheta^f=0$	$\frac{1}{2}\left(\vartheta^f\wedge\Omega^f\right)+\Lambda_{\mathrm{eff}}^f\vartheta^f=0$
mielke-geometrodynamics_217 EQ0658 (10.3.15)		■ 0.995	$\{ \wedge^*\mathrm{D}\left(\vartheta^\alpha\wedge H_\alpha\right)-\{ \wedge^*\left(T^\alpha\wedge H_\alpha\right)=\frac{\gamma}{\delta\ell^2}\left(\vartheta^\alpha\wedge T_\alpha\right)+\frac{1}{\delta\ell}\left(\eta^\alpha\wedge H_\alpha\right).$	$\left\{ \wedge^*\mathrm{D}\left(\vartheta^\alpha\wedge H_\alpha\right)-\left\{ \wedge^*\left(T^\alpha\wedge H_\alpha\right)=\frac{\gamma}{\delta\ell^2}\left(\vartheta^\alpha\wedge T_\alpha\right)+\frac{1}{\delta\ell}\left(\eta^\alpha\wedge H_\alpha\right).$	$\left\{ \wedge^*\mathrm{D}\left(\vartheta^\alpha\wedge H_\alpha\right)-\left\{ \wedge^*\left(T^\alpha\wedge H_\alpha\right)=\frac{\gamma}{\delta\ell^2}\left(\vartheta^\alpha\wedge T_\alpha\right)+\frac{1}{\delta\ell}\left(\eta^\alpha\wedge H_\alpha\right).$
mielke-geometrodynamics_83 EQ0207 (4.1.21)		■ 0.995	$\delta:\left\{ \begin{array}{l} A(n,\mathbb{R})\times A(M)\longrightarrow A(M)\\ \downarrow\qquad\qquad\downarrow\qquad\qquad\qquad\downarrow\\ (g,a)\qquad\qquad(e(m),x)\qquad=\left(\mathbf{g.e}\left(m\right),g^{-1}(x-a)\right).\end{array}\right.$	$\delta:\left\{ \begin{array}{l} A(n,\mathbb{R})\times A(M)\longrightarrow A(M)\\ \downarrow\qquad\qquad\downarrow\qquad\qquad\qquad\downarrow\\ (g,a)\qquad\qquad(e(m),x)\qquad=\left(\mathbf{g.e}\left(m\right),g^{-1}(x-a)\right).\end{array}\right.$	$\delta:\left\{ \begin{array}{l} A(n,\mathbb{R})\times A(M)\longrightarrow A(M)\\ \downarrow\qquad\qquad\downarrow\qquad\qquad\qquad\downarrow\\ (g,a)\qquad\qquad(e(m),x)\qquad=\left(g.e(m),g^{-1}(x-a)\right).\end{array}\right.$
mielke-geometrodynamics_66 EQ0154 (3.5.2)		■ 0.995	$\stackrel{(-)}{\sigma}_{\mu\nu}:=\left\{ \begin{array}{l} \sigma^{ij}=\frac{1}{2}\varepsilon^{ijk}\sigma_k\\ \sigma^{i0}=\left(-\right)\frac{1}{2}\sigma^i\end{array}\right \left.i,j,k=1,2,3\right.$	$\stackrel{(-)}{\sigma}_{\mu\nu}:=\left\{ \begin{array}{l} \sigma^{ij}=\frac{1}{2}\varepsilon^{ijk}\sigma_k\\ \sigma^{i0}=\left(-\right)\frac{1}{2}\sigma^i\end{array}\right \left.i,j,k=1,2,3\right.$	$\stackrel{(-)}{\sigma}_{\mu\nu}:=\left\{ \begin{array}{l} \sigma^{ij}=\frac{1}{2}\varepsilon^{ijk}\sigma_k\\ \sigma^{i0}=\left(-\right)\frac{1}{2}\sigma^i\end{array}\right \left.i,j,k=1,2,3\right.$
mielke-geometrodynamics_304 EQ0928 (14.1.9)		■ 0.995	$\begin{aligned} \Sigma_{\mu\nu}^{\mathrm{YM}}&=\frac{1}{\alpha_g}\mathrm{Tr}\left(F_{\mu\alpha}F_{\nu}^{\cdot\alpha}-\frac{1}{4}g_{\mu\nu}F_{\alpha\beta}F^{\alpha\beta}\right)\\ &=\frac{1}{2\alpha_g}\mathrm{Tr}\left(F_{\mu\alpha}F_{\nu}^{\cdot\alpha}-(-1)^s{}^*F_{\mu\alpha}{}^*F_{\nu}^{\cdot\alpha}\right),\end{aligned}$	$\Sigma_{\mu\nu}^{\mathrm{YM}}=\frac{1}{\alpha_g}\mathrm{Tr}\left(F_{\mu\alpha}F_{\nu}^{\cdot\alpha}-\frac{1}{4}g_{\mu\nu}F_{\alpha\beta}F^{\alpha\beta}\right)\\ =\frac{1}{2\alpha_g}\mathrm{Tr}\left(F_{\mu\alpha}F_{\nu}^{\cdot\alpha}-(-1)^s{}^*F_{\mu\alpha}{}^*F_{\nu}^{\cdot\alpha}\right),$	$\Sigma_{\mu\nu}^{\mathrm{YM}}=\frac{1}{\alpha_g}\mathrm{Tr}\left(F_{\mu\alpha}F_{\nu}^{\cdot\alpha}-\frac{1}{4}g_{\mu\nu}F_{\alpha\beta}F^{\alpha\beta}\right)\\ =\frac{1}{2\alpha_g}\mathrm{Tr}\left(F_{\mu\alpha}F_{\nu}^{\cdot\alpha}-(-1)^s{}^*F_{\mu\alpha}{}^*F_{\nu}^{\cdot\alpha}\right),$
mielke-geometrodynamics_33 EQ0006 (2.2.1) [re-fined: census]		■ 0.996	$e_{\beta}\rfloor\vartheta^\alpha=\delta^\alpha{}_\beta.$	$e_{\beta}\rfloor\vartheta^\alpha=\delta^\alpha{}_\beta.$	$e_{\beta}\rfloor\vartheta^\alpha=\delta^\alpha{}_\beta.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_99 EQ0284 (4.4.13)	99	■ 0.996	$\begin{aligned} D[\Theta, \vartheta] &= [D\Theta, \vartheta] + [\Theta, D\vartheta] \\ &= [[\vartheta, \Omega^g], \vartheta] + [\Theta, \Theta] \\ &= \vartheta \wedge \Omega^g \wedge \vartheta - \Omega^g \wedge \vartheta \wedge \vartheta + \vartheta \wedge \vartheta \wedge \Omega^g - \vartheta \wedge \Omega^g \wedge \vartheta \\ &= [\vartheta \wedge \vartheta, \Omega^g]. \end{aligned}$		$\begin{aligned} D[\Theta, \vartheta] &= [D\Theta, \vartheta] + [\Theta, D\vartheta] \\ &= [[\vartheta, \Omega^g], \vartheta] + [\Theta, \Theta] \\ &= \vartheta \wedge \Omega^g \wedge \vartheta - \Omega^g \wedge \vartheta \wedge \vartheta + \vartheta \wedge \vartheta \wedge \Omega^g - \vartheta \wedge \Omega^g \wedge \vartheta \\ &= [\vartheta \wedge \vartheta, \Omega^g]. \end{aligned} \quad (4)$
mielke-geometrodynamics_103 EQ0308 (4.5.18)	103	■ 0.996	$C_{\mathrm{AdS}} = \hat{M}^2 + \frac{\Lambda}{3} \hat{J}^2$	$C_{\mathrm{AdS}} = \hat{M}^2 + \frac{\Lambda}{3} \hat{J}^2$	$C_{\mathrm{AdS}} = \hat{M}^2 + \frac{\Lambda}{3} \hat{J}^2$
mielke-geometrodynamics_210 EQ0623 (10.1.11)	210	■ 0.996	$\frac{\delta C_{\mathrm{TL}}}{\delta \vartheta^\alpha} = \frac{1}{\ell} R_\alpha^* \quad \text{and} \quad \frac{\delta C_{\mathrm{TL}}}{\delta \Gamma^{\star\alpha}} = \frac{1}{\ell} T_\alpha,$	$\frac{\delta C_{\mathrm{TL}}}{\delta \vartheta^\alpha} = \frac{1}{\ell} R_\alpha^* \quad \text{and} \quad \frac{\delta C_{\mathrm{TL}}}{\delta \Gamma^{\star\alpha}} = \frac{1}{\ell} T_\alpha,$	$\frac{\delta C_{\mathrm{TL}}}{\delta \vartheta^\alpha} = \frac{1}{\ell} R_\alpha^* \quad \text{and} \quad \frac{\delta C_{\mathrm{TL}}}{\delta \Gamma^{\star\alpha}} = \frac{1}{\ell} T_\alpha,$
mielke-geometrodynamics_295 EQ0915 (13.8.1)	295	■ 0.996	$\hat{C} := -\frac{1}{2} \left(\Gamma_A{}^B \wedge d\Gamma_B{}^A - \frac{2}{3} \Gamma_A{}^B \wedge \Gamma_B{}^C \wedge \Gamma_C{}^A \right)$	$\hat{C} := -\frac{1}{2} \left(\Gamma_A{}^B \wedge d\Gamma_B{}^A - \frac{2}{3} \Gamma_A{}^B \wedge \Gamma_B{}^C \wedge \Gamma_C{}^A \right)$	$\hat{C} := -\frac{1}{2} \left(\Gamma_A{}^B \wedge d\Gamma_B{}^A - \frac{2}{3} \Gamma_A{}^B \wedge \Gamma_B{}^C \wedge \Gamma_C{}^A \right)$
mielke-geometrodynamics_357 EQ1071 (16.2.10)	357	■ 0.996	$E_n = \beta m c^2 \left(2n + \iota + \frac{3}{2} + \sqrt{\frac{g}{4} + \frac{1}{\beta^2}} \right), n = 0, 1, 2, \dots$	$E_n = \beta m c^2 \left(2n + \iota + \frac{3}{2} + \sqrt{\frac{g}{4} + \frac{1}{\beta^2}} \right), n = 0, 1, 2, \dots$	$E_n = \beta m c^2 \left(2n + \iota + \frac{3}{2} + \sqrt{\frac{g}{4} + \frac{1}{\beta^2}} \right), n = 0, 1, 2, \dots$
mielke-geometrodynamics_161 EQ0490 (7.7.7) [re-fined: census]	161	■ 0.996	$\begin{aligned} \widetilde{E}_\alpha &:= -e_\alpha \rfloor \widetilde{L} - (e_\alpha \rfloor R^{\beta\gamma}) \wedge \widetilde{H}_{\beta\gamma} = \Sigma_\alpha, \\ D\widetilde{H}_{\alpha\beta} &= \tau_{\alpha\beta}. \end{aligned}$	$\begin{aligned} -\widetilde{E}_\alpha &:= -e_\alpha \rfloor \widetilde{L} - (e_\alpha \rfloor R^{\beta\gamma}) \wedge \widetilde{H}_{\beta\gamma} = \Sigma_\alpha, \\ D\widetilde{H}_{\alpha\beta} &= \tau_{\alpha\beta}. \end{aligned}$	$\begin{aligned} -\widetilde{E}_\alpha &:= -e_\alpha \rfloor \widetilde{L} - (e_\alpha \rfloor R^{\beta\gamma}) \wedge \widetilde{H}_{\beta\gamma} = \Sigma_\alpha, \\ D\widetilde{H}_{\alpha\beta} &= \tau_{\alpha\beta}. \end{aligned}$
mielke-geometrodynamics_86 EQ0220 (4.2.13)	86	■ 0.996	$\Omega^L \xrightarrow{A^{-1}} \Omega^L = G^{-1} \Omega^L G$	$\Omega^L \xrightarrow{A^{-1}} \Omega^L = G^{-1} \Omega^L G$	$\Omega^L \xrightarrow{A^{-1}} \Omega^L = G^{-1} \Omega^L G$
mielke-geometrodynamics_152 EQ0460 (7.2.12)	152	■ 0.996	$g_{ij} \rightarrow \bar{g}_{ij} = \phi^L g_{ij},$	$g_{ij} \rightarrow \bar{g}_{ij} = \phi^L g_{ij},$	$g_{ij} \rightarrow \bar{g}_{ij} = \phi^L g_{ij},$
mielke-geometrodynamics_218 EQ0664 (10.3.21)	218	■ 0.996	$\square \vartheta_\alpha = \frac{\ell}{\gamma} {}^* D^* [\tau_\alpha^* - S_\alpha - \delta \ell (\Sigma_\alpha + E_\alpha)] + \frac{\ell}{\gamma} D^* (D^* H_\alpha^* - \delta \ell D^* H_\alpha).$	$\square \vartheta_\alpha = \frac{\ell}{\gamma} {}^* D^* [\tau_\alpha^* - S_\alpha - \delta \ell (\Sigma_\alpha + E_\alpha)] + \frac{\ell}{\gamma} D^* (D^* H_\alpha^* - \delta \ell D^* H_\alpha).$	$\square \vartheta_\alpha = \frac{\ell}{\gamma} {}^* D^* [\tau_\alpha^* - S_\alpha - \delta \ell (\Sigma_\alpha + E_\alpha)] + \frac{\ell}{\gamma} D^* (D^* H_\alpha^* - \delta \ell D^* H_\alpha).$
mielke-geometrodynamics_315 EQ0977 (14.2.18)	315	■ 0.996	$\hat{L}_\parallel = -\frac{1}{2\ell^{*2}} \mathrm{Tr}[\hat{\alpha} \wedge \hat{\alpha}, \quad \hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}]$	$\hat{L}_\parallel = -\frac{1}{2\ell^{*2}} \mathrm{Tr}[\hat{\alpha} \wedge \hat{\alpha}, \quad \hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}]$	$\hat{L}_\parallel = -\frac{1}{2\ell^{*2}} \mathrm{Tr}[\hat{\alpha} \wedge \hat{\alpha}, \quad \hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}]$
mielke-geometrodynamics_92 EQ0250 (4.3.15)	92	■ 0.996	${}^* \omega^g = \frac{1}{2} \omega^{\alpha\beta} L_{\alpha\beta} = \frac{1}{2} \Gamma_i^{\alpha\beta} L_{\alpha\beta} \otimes dx^i,$	${}^* \omega^g = \frac{1}{2} \omega^{\alpha\beta} L_{\alpha\beta} = \frac{1}{2} \Gamma_i^{\alpha\beta} L_{\alpha\beta} \otimes dx^i,$	${}^* \omega^g = \frac{1}{2} \omega^{\alpha\beta} L_{\alpha\beta} = \frac{1}{2} \Gamma_i^{\alpha\beta} L_{\alpha\beta} \otimes dx^i,$
mielke-geometrodynamics_128 EQ0376 (6.2.27)	128	■ 0.996	$\stackrel{(\pm)}{D} \vartheta^\alpha = \pm \frac{i}{2} \ell^2 g^{\alpha\beta} \stackrel{(\pm)}{\Pi}_\beta,$	$\stackrel{(\pm)}{D} \vartheta^\alpha = \pm \frac{i}{2} \ell^2 g^{\alpha\beta} \stackrel{(\pm)}{\Pi}_\beta,$	$\stackrel{(\pm)}{D} \vartheta^\alpha = \pm \frac{i}{2} \ell^2 g^{\alpha\beta} \stackrel{(\pm)}{\Pi}_\beta,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_152 (3.4.23)	66	■ 0.996	$L_{\mathrm{BI}}(\mathscr{F},*\mathscr{F})=\sqrt{1-2\mathscr{F}-(\ast\mathscr{F})^2}-1.$	$L_{\mathrm{BI}}(\mathscr{F},*\mathscr{F})=\sqrt{1-2\mathscr{F}-(\ast\mathscr{F})^2}-1.$	$L_{\mathrm{BI}}(\mathscr{F},*\mathscr{F})=\sqrt{1-2\mathscr{F}-(\ast\mathscr{F})^2}-1.$
mielke-geometrodynamics_367 (6.2.18)	126	■ 0.997	$R_{\alpha\beta}=\xi^{\ast}R_{\alpha\beta}^{(\ast)}+(\gamma/\ell^2)\left[\vartheta_{\alpha}\wedge\vartheta_{\beta}\pm i^{\ast}(\vartheta_{\alpha}\wedge\vartheta_{\beta})\right],$	$R_{\alpha\beta}=\xi^{\ast}R_{\alpha\beta}^{(\ast)}+(\gamma/\ell^2)\left[\vartheta_{\alpha}\wedge\vartheta_{\beta}\pm i^{\ast}(\vartheta_{\alpha}\wedge\vartheta_{\beta})\right],$	$R_{\alpha\beta}=\xi^{\ast}R_{\alpha\beta}^{(\ast)}+(\gamma/\ell^2)\left[\vartheta_{\alpha}\wedge\vartheta_{\beta}\pm i^{\ast}(\vartheta_{\alpha}\wedge\vartheta_{\beta})\right],$
mielke-geometrodynamics_371 (6.2.22)	127	■ 0.997	$\stackrel{(+)}{\Pi}\wedge\stackrel{(-)}{\Pi}_{\beta}=H^{\beta}\wedge H_{\beta}+\ell^{-4}T^{\beta}\wedge T_{\beta}=0.$	$\stackrel{(+)}{\Pi}\wedge\stackrel{(-)}{\Pi}_{\beta}=H^{\beta}\wedge H_{\beta}+\ell^{-4}T^{\beta}\wedge T_{\beta}=0.$	$\stackrel{(+)}{\Pi}^{\beta}\wedge\stackrel{(-)}{\Pi}_{\beta}=H^{\beta}\wedge H_{\beta}+\ell^{-4}T^{\beta}\wedge T_{\beta}=0.$
mielke-geometrodynamics_602 (9.2.1)	198	■ 0.997	$\begin{aligned} p_1(M^4) &= \frac{1}{2(2\pi)^2} \int_{M^4} \mathrm{Tr}(\Omega^L \wedge \Omega^L) \\ &= \frac{(-1)^{s/2}}{(4\pi)^2} \int_{M^4} R_{\alpha\beta\gamma\delta}^{\ast} R^{\alpha\beta\gamma\delta} \sqrt{ g } d^4x \end{aligned}$	$\begin{aligned} p_1(M^4) &= \frac{1}{2(2\pi)^2} \int_{M^4} \mathrm{Tr}(\Omega^L \wedge \Omega^L) \\ &= \frac{(-1)^{s/2}}{(4\pi)^2} \int_{M^4} R_{\alpha\beta\gamma\delta}^{\ast} R^{\alpha\beta\gamma\delta} \sqrt{ g } d^4x \end{aligned}$	$\begin{aligned} p_1(M^4) &= \frac{1}{2(2\pi)^2} \int_{M^4} \mathrm{Tr}(\Omega^L \wedge \Omega^L) \\ &= \frac{(-1)^{s/2}}{(4\pi)^2} \int_{M^4} R_{\alpha\beta\gamma\delta}^{\ast} R^{\alpha\beta\gamma\delta} \sqrt{ g } d^4x \end{aligned}$
mielke-geometrodynamics_717 (10.9.3)	229	■ 0.997	$\Sigma_{\alpha}:=\frac{\delta L_{\Psi}}{\delta \vartheta^{\alpha}}=\frac{\partial L_{\Psi}}{\partial \vartheta_{\alpha}}+D\frac{\partial L_{\Psi}}{\partial T^{\alpha}},$	$\Sigma_{\alpha}:=\frac{\delta L_{\Psi}}{\delta \vartheta^{\alpha}}=\frac{\partial L_{\Psi}}{\partial \vartheta_{\alpha}}+D\frac{\partial L_{\Psi}}{\partial T^{\alpha}},$	$\Sigma_{\alpha}:=\frac{\delta L_{\Psi}}{\delta \vartheta^{\alpha}}=\frac{\partial L_{\Psi}}{\partial \vartheta_{\alpha}}+D\frac{\partial L_{\Psi}}{\partial T^{\alpha}},$
mielke-geometrodynamics_766 (11.1.33)	243	■ 0.997	$\stackrel{\ast}{\sigma}\tilde{\omega}=\frac{1}{2}\omega^{\alpha\beta}\sigma_{\alpha\beta}=\frac{1}{2}\Gamma_i^{\cdot\alpha\beta}\sigma_{\alpha\beta}\otimes dx^i=:\Gamma_idx^i.$	$\stackrel{\ast}{\sigma}\tilde{\omega}=\frac{1}{2}\omega^{\alpha\beta}\sigma_{\alpha\beta}=\frac{1}{2}\Gamma_i^{\cdot\alpha\beta}\sigma_{\alpha\beta}\otimes dx^i=:\Gamma_idx^i.$	$\stackrel{\ast}{\sigma}\tilde{\omega}=\frac{1}{2}\omega^{\alpha\beta}\sigma_{\alpha\beta}=\frac{1}{2}\Gamma_i^{\cdot\alpha\beta}\sigma_{\alpha\beta}\otimes dx^i=:\Gamma_idx^i.$
mielke-geometrodynamics_1003 (14.3.16)	323	■ 0.997	$D\,D=\frac{1}{2}\{D,D\}+\frac{1}{2}[D,D]=\frac{1}{2}\{D,D\}+\frac{1}{2}\widetilde{\Omega}$	$D\,D=\frac{1}{2}\{D,D\}+\frac{1}{2}[D,D]=\frac{1}{2}\{D,D\}+\frac{1}{2}\widetilde{\Omega}$	$D\,D=\frac{1}{2}\{D,D\}+\frac{1}{2}[D,D]=\frac{1}{2}\{D,D\}+\frac{1}{2}\widetilde{\Omega}$
mielke-geometrodynamics_225 (4.2.18)	87	■ 0.997	$\omega^{\mathrm{T}}=\widetilde{\vartheta}-D\widetilde{\varphi}$	$\omega^{\mathrm{T}}=\widetilde{\vartheta}-D\widetilde{\varphi}$	$\omega^{\mathrm{T}}=\widetilde{\vartheta}-D\widetilde{\varphi}$
mielke-geometrodynamics_048 (2.6.11)	43	■ 0.997	$D\,D\,D\,b\equiv 0.$	$D\,D\,D\,b\equiv 0.$	$D\,D\,D\,b\equiv 0.$
mielke-geometrodynamics_099 (3.2.9)	56	■ 0.997	$\begin{aligned} \{ \ }^{\Delta}\omega &:= G^{-1}\omega-\omega=G^{-1}[\omega,G]-G^{-1}dG \\ &=-G^{-1}DG=-G^{-1}D. \end{aligned}$	$\begin{aligned} \Delta\omega &:= G^{-1}\omega-\omega=G^{-1}[\omega,G]-G^{-1}dG \\ &=-G^{-1}DG=-G^{-1}D. \end{aligned}$	$\begin{aligned} \Delta\omega &:= G^{-1}\omega-\omega=G^{-1}[\omega,G]-G^{-1}dG \\ &=-G^{-1}DG=-G^{-1}D. \end{aligned}$
mielke-geometrodynamics_219 (4.2.12)	86	■ 0.997	$\widetilde{\widetilde{\Omega}}\rightarrow^{A^{-1}}\widetilde{\widetilde{\Omega}}=A^{-1}\widetilde{\widetilde{\Omega}}A$	$\widetilde{\widetilde{\Omega}}\rightarrow^{A^{-1}}\widetilde{\widetilde{\Omega}}=A^{-1}\widetilde{\widetilde{\Omega}}A$	$\widetilde{\widetilde{\Omega}}\rightarrow^{A^{-1}}\widetilde{\widetilde{\Omega}}=A^{-1}\widetilde{\widetilde{\Omega}}A$
mielke-geometrodynamics_411 (6.4.31)	136	■ 0.997	$n_{\mathrm{NY}}:=\int_{R^4}dC_{\mathrm{TT}}=\int_{S^3_{\infty}}\underline{C}_{\mathrm{TT}}=3\,\mathrm{Vol}\left(S^3\right)k=6\pi^2k.$	$n_{\mathrm{NY}}:=\int_{R^4}dC_{\mathrm{TT}}=\int_{S^3_{\infty}}\underline{C}_{\mathrm{TT}}=3\,\mathrm{Vol}\left(S^3\right)k=6\pi^2k.$	$n_{\mathrm{NY}}:=\int_{R^4}dC_{\mathrm{TT}}=\int_{S^3_{\infty}}\underline{C}_{\mathrm{TT}}=3\,\mathrm{Vol}\left(S^3\right)k=6\pi^2k.$
mielke-geometrodynamics_492 (7.7.10)	161	■ 0.997	$H_{\alpha\beta}:=-\frac{\partial L}{\partial R^{\alpha\beta}}=\widetilde{H}_{\alpha\beta}+\frac{1}{2}R^{\mu\nu}\wedge\left(\partial\widetilde{H}_{\mu\nu}/\partial R^{\alpha\beta}\right)$	$H_{\alpha\beta}:=-\frac{\partial L}{\partial R^{\alpha\beta}}=\widetilde{H}_{\alpha\beta}+\frac{1}{2}R^{\mu\nu}\wedge\left(\partial\widetilde{H}_{\mu\nu}/\partial R^{\alpha\beta}\right)$	$H_{\alpha\beta}:=-\frac{\partial L}{\partial R^{\alpha\beta}}=\widetilde{H}_{\alpha\beta}+\frac{1}{2}R^{\mu\nu}\wedge\left(\partial\widetilde{H}_{\mu\nu}/\partial R^{\alpha\beta}\right)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_ E08877 (13.3.6)	285	■ 0.997	$\left[P_{\alpha}, P_{*}^{\beta}\right]=\frac{1}{\ell^2} L^{\beta}{}_{\alpha} L^{\beta}{}_{\alpha} .$	$\left[P_{\alpha}, P_{*}^{\beta}\right]=\frac{1}{\ell^2} L^{\beta}{}_{\alpha} .$	$\left[P_{\alpha}, P_{*}^{\beta}\right]=\frac{1}{\ell^2} L^{\beta}{}_{\alpha} .$
mielke-geometrodynamics_ E09962 (14.2.2)	312	■ 0.997	$\stackrel{\circ}{\vartheta}=\varphi_j d \xi^j,$	$\stackrel{\circ}{\vartheta}=\varphi_j d \xi^j,$	$\stackrel{\circ}{\vartheta}=\varphi_j d \xi^j,$
mielke-geometrodynamics_ E01060 (16.1.1)	354	■ 0.997	$R_{i j}^f-\frac{1}{2} f_{i j}\left(R^f-2 \Lambda_{\text {eff }}\right)=\ell^2 T_{i j}$	$R_{i j}^f-\frac{1}{2} f_{i j}\left(R^f-2 \Lambda_{\text {eff }}\right)=\ell^2 T_{i j}$	$R_{i j}^f-\frac{1}{2} f_{i j}\left(R^f-2 \Lambda_{\text {eff }}\right)=\ell^2 T_{i j}$
mielke-geometrodynamics_ E0976 (14.2.17)	315	■ 0.997	$\begin{aligned} \hat{L}_{\mathrm{W}} &= \frac{1}{2} \operatorname{Tr}\left(\hat{\Omega} \wedge \underbrace{\hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}}_{2+\mathrm{K} \text { termms }}\right) \\ &= \hat{L}_{\mathrm{W}}^g + \hat{L}_{\mathrm{W}}^G + \ell^{* 2} \hat{L}_{\parallel} - \frac{1}{2} d \operatorname{Tr}\left(\hat{\alpha} \wedge \hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}\right) \end{aligned}$	$\hat{L}_{\mathrm{W}} = \frac{1}{2} \operatorname{Tr}\left(\hat{\Omega} \wedge \underbrace{\hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}}_{2+\mathrm{K} \text { termms }}\right) \\ = \hat{L}_{\mathrm{W}}^g + \hat{L}_{\mathrm{W}}^G + \ell^{* 2} \hat{L}_{\parallel} - \frac{1}{2} d \operatorname{Tr}\left(\hat{\alpha} \wedge \hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}\right)$	$\hat{L}_{\mathrm{W}} = \frac{1}{2} \operatorname{Tr}\left(\hat{\Omega} \wedge \underbrace{\hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}}_{2+\mathrm{K} \text { termms }}\right) \\ = \hat{L}_{\mathrm{W}}^g + \hat{L}_{\mathrm{W}}^G + \ell^{* 2} \hat{L}_{\parallel} - \frac{1}{2} d \operatorname{Tr}\left(\hat{\alpha} \wedge \hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}\right)$
mielke-geometrodynamics_ E09466 (7.3.5)	153	■ 0.997	$G_{\alpha}^{\{\}}:=\frac{1}{2} R^{\{\beta \gamma} \wedge \eta_{\alpha \beta \gamma}, \quad G_{i j}:=\operatorname{Ric}_{i j}^{\{\}}-\frac{1}{2} g_{i j},$	$G_{\alpha}^{\{\}}:=\frac{1}{2} R^{\{\beta \gamma} \wedge \eta_{\alpha \beta \gamma}, \quad G_{i j}:=\operatorname{Ric}_{i j}^{\{\}}-\frac{1}{2} g_{i j},$	$G_{\alpha}^{\{\}}:=\frac{1}{2} R^{\{\beta \gamma} \wedge \eta_{\alpha \beta \gamma}, \quad G_{i j}:=\operatorname{Ric}_{i j}^{\{\}}-\frac{1}{2} g_{i j},$
mielke-geometrodynamics_ E08105 (3.3.6)	58	■ 0.998	$H:=-\frac{\partial L}{\partial d A}=-\frac{\partial L}{\partial F} \quad \text { and } \quad j:=\frac{\delta L_{\text {mat }}}{\delta A},$	$H:=-\frac{\partial L}{\partial d A}=-\frac{\partial L}{\partial F} \quad \text { and } \quad j:=\frac{\delta L_{\text {mat }}}{\delta A},$	$H:=-\frac{\partial L}{\partial d A}=-\frac{\partial L}{\partial F} \quad \text { and } \quad j:=\frac{\delta L_{\text {mat }}}{\delta A},$
mielke-geometrodynamics_ E08136 (3.4.7)	63	■ 0.998	$\begin{aligned} \delta \Omega &= \delta(d \omega-\omega \wedge \omega)=\delta d \omega-\delta \omega \wedge \omega-\omega \wedge \delta \omega \\ &= d \delta \omega-[\omega, \delta \omega]=D \delta \omega . \end{aligned}$	$\delta \Omega = \delta(d \omega-\omega \wedge \omega)=\delta d \omega-\delta \omega \wedge \omega-\omega \wedge \delta \omega \\ = d \delta \omega-[\omega, \delta \omega]=D \delta \omega .$	$\delta \Omega = \delta(d \omega-\omega \wedge \omega)=\delta d \omega-\delta \omega \wedge \omega-\omega \wedge \delta \omega \\ = d \delta \omega-[\omega, \delta \omega]=D \delta \omega .$
mielke-geometrodynamics_ E08390 (6.4.5)	131	■ 0.998	$\mathscr{H}=n \rfloor L-\ell_n \underline{\Psi} \wedge \left(\frac{\partial L}{\partial d \Psi}\right) .$	$\mathscr{H}=n \rfloor L-\ell_n \underline{\Psi} \wedge \left(\frac{\partial L}{\partial d \Psi}\right) .$	$\mathscr{H}=n \rfloor L-\ell_n \underline{\Psi} \wedge \left(\frac{\partial L}{\partial d \Psi}\right) .$
mielke-geometrodynamics_ E08594 (9.1.12)	195	■ 0.998	$\bar{\Omega}^{\{\}}=0 .$	$\bar{\Omega}^{\{\}}=0 .$	$\bar{\Omega}^{\{\}}=0 .$
mielke-geometrodynamics_ E01012 (14.4.7)	326	■ 0.998	$\begin{aligned} S_{\mathrm{A}} &:= 4 \pi\left(r_+^2+\frac{J(J+1) \hbar^2}{M^2 c^2}\right) \\ &= \frac{8 \pi \hbar^2}{M^{* 4} c^2}\left(M^2+\frac{\alpha_e}{2} M^{* 2} \operatorname{Tr} \bar{Q}^2+M \sqrt{M^2-J(J+1) \frac{M^{* 4}}{M^2}-\alpha_e M^{* 2} \operatorname{Tr} \bar{Q}^2}\right) \end{aligned}$	$S_{\mathrm{A}}:=4 \pi\left(r_+^2+\frac{J(J+1) \hbar^2}{M^2 c^2}\right) \\ = \frac{8 \pi \hbar^2}{M^{* 4} c^2}\left(M^2+\frac{\alpha_e}{2} M^{* 2} \operatorname{Tr} \bar{Q}^2+M \sqrt{M^2-J(J+1) \frac{M^{* 4}}{M^2}-\alpha_e M^{* 2} \operatorname{Tr} \bar{Q}^2}\right)$	$S_{\mathrm{A}}:=4 \pi\left(r_+^2+\frac{J(J+1) \hbar^2}{M^2 c^2}\right) \\ = \frac{8 \pi \hbar^2}{M^{* 4} c^2}\left(M^2+\frac{\alpha_e}{2} M^{* 2} \operatorname{Tr} \bar{Q}^2+M \sqrt{M^2-J(J+1) \frac{M^{* 4}}{M^2}-\alpha_e M^{* 2} \operatorname{Tr} \bar{Q}^2}\right)$
mielke-geometrodynamics_ E08813 (11.4.26)	256	■ 0.998	$d \rho^*:=e^{(\mu-\nu) / 2} d \rho .$	$d \rho^*:=e^{(\mu-\nu) / 2} d \rho .$	$d \rho^*:=e^{(\mu-\nu) / 2} d \rho .$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00073 (2.8.16)	49	■ 0.998	$\begin{aligned} & \mathrm{d} \stackrel{\infty}{\omega} - \stackrel{\infty}{\omega} \wedge \stackrel{\infty}{\omega} \\ & = -d \left(G^{-1} \right) \wedge dG - G^{-1} ddG - \left(G^{-1} dG \right) \wedge \left(G^{-1} dG \right) \\ & = -d \left(G^{-1} \right) \wedge dG + G^{-1} G d \left(G^{-1} \right) \wedge dG = 0 \end{aligned}$	$\begin{aligned} & d \stackrel{\infty}{\omega} - \stackrel{\infty}{\omega} \wedge \stackrel{\infty}{\omega} \\ & = -d \left(G^{-1} \right) \wedge dG - G^{-1} ddG - \left(G^{-1} dG \right) \wedge \left(G^{-1} dG \right) \\ & = -d \left(G^{-1} \right) \wedge dG + G^{-1} G d \left(G^{-1} \right) \wedge dG = 0 \end{aligned}$	$\begin{aligned} & d \stackrel{\infty}{\omega} - \stackrel{\infty}{\omega} \wedge \stackrel{\infty}{\omega} \\ & = -d \left(G^{-1} \right) \wedge dG - G^{-1} ddG - \left(G^{-1} dG \right) \wedge \left(G^{-1} dG \right) \\ & = -d \left(G^{-1} \right) \wedge dG + G^{-1} G d \left(G^{-1} \right) \wedge dG = 0 \end{aligned}$
mielke-geometrodynamics_E00214 (4.2.6)	86	■ 0.998	$\mathrm{mathscr{G}}(n, \mathbb{R}) := C^\infty \left(A(M) \times_{\mathrm{Ad}} GL(n, \mathbb{R}) \right)$	$\mathcal{G}(n, \mathbb{R}) := C^\infty \left(A(M) \times_{\mathrm{Ad}} GL(n, \mathbb{R}) \right)$	$\mathcal{G}(n, \mathbb{R}) := C^\infty \left(A(M) \times_{\mathrm{Ad}} GL(n, \mathbb{R}) \right)$
mielke-geometrodynamics_E00656 (10.3.13) [re-fined: census]	216	■ 0.998	$E_{\alpha} = e_{\alpha} \rfloor L(\star) + (-1)^s \frac{2\gamma}{\ell} \left[(e_{\alpha} \rfloor R^{\star\beta}) \wedge \vartheta_{\beta} \right]$	$E_{\alpha} = e_{\alpha} \rfloor L(\star) + (-1)^s \frac{2\gamma}{\ell} \left[(e_{\alpha} \rfloor R^{\star\beta}) \wedge \vartheta_{\beta} \right]$	$E_{\alpha} = e_{\alpha} \rfloor L(\star) + (-1)^s \frac{2\gamma}{\ell} \left[(e_{\alpha} \rfloor R^{\star\beta}) \wedge \vartheta_{\beta} \right]$
mielke-geometrodynamics_E00062 (2.8.3)	46	■ 0.998	$\begin{aligned} & \ast \left(d \gamma_k \right) = d \pi^{\ast} \left(\gamma_k \right) = D \pi^{\ast} \left(\gamma_k \right) \\ & = \frac{(-1)^k}{(2\pi i)^k (k-1)!} \mathrm{Tr} \left(D \Omega \wedge \Omega \wedge \cdots \wedge \Omega \right) = 0. \end{aligned}$	$\begin{aligned} & \ast \left(d \gamma_k \right) = d \pi^{\ast} \left(\gamma_k \right) = D \pi^{\ast} \left(\gamma_k \right) \\ & = \frac{(-1)^k}{(2\pi i)^k (k-1)!} \mathrm{Tr} \left(D \Omega \wedge \Omega \wedge \cdots \wedge \Omega \right) = 0. \end{aligned}$	$\begin{aligned} & \pi^{\ast} \left(d \gamma_k \right) = d \pi^{\ast} \left(\gamma_k \right) = D \pi^{\ast} \left(\gamma_k \right) \\ & = \frac{(-1)^k}{(2\pi i)^k (k-1)!} \mathrm{Tr} \left(D \Omega \wedge \Omega \wedge \cdots \wedge \Omega \right) = 0. \end{aligned}$
mielke-geometrodynamics_E00426 (6.7.8)	141	■ 0.998	$\mu_{\alpha} = \frac{1}{4} \vartheta_{\alpha} \wedge \ast j_5,$	$\mu_{\alpha} = \frac{1}{4} \vartheta_{\alpha} \wedge \ast j_5,$	$\mu_{\alpha} = \frac{1}{4} \vartheta_{\alpha} \wedge \ast j_5,$
mielke-geometrodynamics_E00069 (2.8.12)	48	■ 0.998	$\begin{aligned} & c_2(M) = \frac{1}{8\pi^2} \int_M d \mathrm{Tr} \left(\omega \wedge \Omega + \frac{1}{3} \omega \wedge \omega \wedge \omega \right) \\ & = \frac{1}{8\pi^2} \int_{\partial M} \mathrm{Tr} \left(\omega \wedge d\omega - \frac{2}{3} \omega \wedge \omega \wedge \omega \right). \end{aligned}$	$\begin{aligned} & c_2(M) = \frac{1}{8\pi^2} \int_M d \mathrm{Tr} \left(\omega \wedge \Omega + \frac{1}{3} \omega \wedge \omega \wedge \omega \right) \\ & = \frac{1}{8\pi^2} \int_{\partial M} \mathrm{Tr} \left(\omega \wedge d\omega - \frac{2}{3} \omega \wedge \omega \wedge \omega \right). \end{aligned}$	$\begin{aligned} & c_2(M) = \frac{1}{8\pi^2} \int_M d \mathrm{Tr} \left(\omega \wedge \Omega + \frac{1}{3} \omega \wedge \omega \wedge \omega \right) \\ & = \frac{1}{8\pi^2} \int_{\partial M} \mathrm{Tr} \left(\omega \wedge d\omega - \frac{2}{3} \omega \wedge \omega \wedge \omega \right). \end{aligned}$
mielke-geometrodynamics_E00144 (3.4.15)	64	■ 0.998	$\iota := \tau + (-1)^{n+s} \ast \left[\omega, \Pi^{\Omega} \right]$	$\iota := \tau + (-1)^{n+s} \ast \left[\omega, \Pi^{\Omega} \right]$	$\iota := \tau + (-1)^{n+s} \ast \left[\omega, \Pi^{\Omega} \right]$
mielke-geometrodynamics_E00318 (5.2.2)	110	■ 0.998	$j := \bar{\psi}^{\ast} \gamma \psi = j^{\mu} \eta_{\mu} \quad \text{and} \quad j_5 := \bar{\psi}^{\ast} \gamma \gamma_5 \psi = j_5^{\mu} \eta_{\mu},$	$j := \bar{\psi}^{\ast} \gamma \psi = j^{\mu} \eta_{\mu} \quad \text{and} \quad j_5 := \bar{\psi}^{\ast} \gamma \gamma_5 \psi = j_5^{\mu} \eta_{\mu},$	$j := \bar{\psi}^{\ast} \gamma \psi = j^{\mu} \eta_{\mu} \quad \text{and} \quad j_5 := \bar{\psi}^{\ast} \gamma \gamma_5 \psi = j_5^{\mu} \eta_{\mu},$
mielke-geometrodynamics_E00347 (5.4.5)	116	■ 0.998	$\mu = \frac{4}{3} \kappa \Lambda = \frac{1}{3} (2\kappa)^2 \rho_{\Lambda} \leq \frac{4}{3} g_{\mathrm{N}\ast} \lambda_{\ast} \simeq 0.2$	$\mu = \frac{4}{3} \kappa \Lambda = \frac{1}{3} (2\kappa)^2 \rho_{\Lambda} \leq \frac{4}{3} g_{\mathrm{N}\ast} \lambda_{\ast} \simeq 0.2$	$\mu = \frac{4}{3} \kappa \Lambda = \frac{1}{3} (2\kappa)^2 \rho_{\Lambda} \leq \frac{4}{3} g_{\mathrm{N}\ast} \lambda_{\ast} \simeq 0.2$
mielke-geometrodynamics_E00500 (7.7.18)	163	■ 0.998	$\ast \sigma = \frac{1}{2} \sigma_{\alpha\beta}^{\ast} \left(\vartheta^{\alpha} \wedge \vartheta^{\beta} \right) = \frac{1}{2} \sigma_{\alpha\beta} \eta^{\alpha\beta} =: \sigma^{(\ast)} = i \gamma_5 \sigma.$	$\ast \sigma = \frac{1}{2} \sigma_{\alpha\beta}^{\ast} \left(\vartheta^{\alpha} \wedge \vartheta^{\beta} \right) = \frac{1}{2} \sigma_{\alpha\beta} \eta^{\alpha\beta} =: \sigma^{(\ast)} = i \gamma_5 \sigma.$	$\ast \sigma = \frac{1}{2} \sigma_{\alpha\beta}^{\ast} \left(\vartheta^{\alpha} \wedge \vartheta^{\beta} \right) = \frac{1}{2} \sigma_{\alpha\beta} \eta^{\alpha\beta} =: \sigma^{(\ast)} = i \gamma_5 \sigma.$
mielke-geometrodynamics_E00670 (10.3.27)	219	■ 0.998	$\check{\mathcal{E}}_{\alpha} := dH_{\alpha} \cong \Sigma_{\alpha} + E_{\alpha} - (-1)^s \eta_{\alpha\beta\gamma} \Gamma^{\star\beta} \wedge H^{\gamma}$	$\check{\mathcal{E}}_{\alpha} := dH_{\alpha} \cong \Sigma_{\alpha} + E_{\alpha} - (-1)^s \eta_{\alpha\beta\gamma} \Gamma^{\star\beta} \wedge H^{\gamma}$	$\check{\mathcal{E}}_{\alpha} := dH_{\alpha} \cong \Sigma_{\alpha} + E_{\alpha} - (-1)^s \eta_{\alpha\beta\gamma} \Gamma^{\star\beta} \wedge H^{\gamma}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_221 EQ0680 (10.3.37)	221	■ 0.998	$\theta_{\mathrm{L}}\ C_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}}\tilde{C}_{\mathrm{L}}=-\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}-\frac{\Lambda}{\ell}\eta+\frac{\theta_{\mathrm{T}}}{2\ell}\vartheta^{\alpha}\wedge T_{\alpha}+\Delta\theta_{\mathrm{L}}C_{\mathrm{L}}+\frac{\Delta\theta_{\mathrm{TL}}}{\ell}d(\Gamma^{\star\alpha}\wedge\vartheta_{\alpha})$	$\theta_{\mathrm{L}}C_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}}\tilde{C}_{\mathrm{L}}=-\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}-\frac{\Lambda}{\ell}\eta+\frac{\theta_{\mathrm{T}}}{2\ell}\vartheta^{\alpha}\wedge T_{\alpha}+\Delta\theta_{\mathrm{L}}C_{\mathrm{L}}+\frac{\Delta\theta_{\mathrm{TL}}}{\ell}d(\Gamma^{\star\alpha}\wedge\vartheta_{\alpha})$	$\theta_{\mathrm{L}}C_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}}\tilde{C}_{\mathrm{L}}=-\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}-\frac{\Lambda}{\ell}\eta+\frac{\theta_{\mathrm{T}}}{2\ell}\vartheta^{\alpha}\wedge T_{\alpha}+\Delta\theta_{\mathrm{L}}C_{\mathrm{L}}+\frac{\Delta\theta_{\mathrm{TL}}}{\ell}d(\Gamma^{\star\alpha}\wedge\vartheta_{\alpha})$
mielke-geometrodynamics_225 EQ0700 (10.5.11)	225	■ 0.998	$[\hat{D},\hat{D}]=\hat{R}+T^{\alpha}\hat{D}_{\alpha}$	$[\hat{D},\hat{D}]=\hat{R}+T^{\alpha}\hat{D}_{\alpha}$	$[\hat{D},\hat{D}]=\hat{R}+T^{\alpha}\hat{D}_{\alpha}$
mielke-geometrodynamics_258 EQ0820 (11.4.34)	258	■ 0.998	$f^l(\rho)=\bar{\rho}^l\left(1+\bar{e}^{4l+2}\right)^{-1/2}$	$f^l(\rho)=\bar{\rho}^l\left(1+\bar{e}^{4l+2}\right)^{-1/2}$	$f^l(\rho)=\bar{\rho}^l(1+\bar{e}^{4l+2})^{-1/2}$
mielke-geometrodynamics_373 EQ1126 (B.38)	373	■ 0.998	$\partial:C_p(M,\mathbb{R})\rightarrow C_{p-1}(M,\mathbb{R}),$	$\partial:C_p(M,\mathbb{R})\rightarrow C_{p-1}(M,\mathbb{R}),$	$\partial:C_p(M,\mathbb{R})\rightarrow C_{p-1}(M,\mathbb{R}),$
mielke-geometrodynamics_166 EQ0516 (7.7.34)	166	■ 0.998	$\widetilde{T}=T-3d\varphi,\quad\mathscr{A}^{\theta}=\mathscr{A}-id\theta.$	$\widetilde{T}=T-3d\varphi,\quad\mathscr{A}^{\theta}=\mathscr{A}-id\theta.$	$\widetilde{T}=T-3d\varphi,\quad\mathscr{A}^{\theta}=\mathscr{A}-id\theta.$
mielke-geometrodynamics_313 EQ0967 (14.2.8)	313	■ 0.998	$d\stackrel{\circ}{s}=g_{ij}\,d\xi^i\otimes_s d\xi^j$	$d\stackrel{\circ}{s}=g_{ij}\,d\xi^i\otimes_s d\xi^j$	$d\stackrel{\circ}{s}^2=\stackrel{\circ}{g}_{ij}d\xi^i\otimes_s d\xi^j,$
mielke-geometrodynamics_46 EQ0060 (2.8.1)	46	■ 0.999	$\operatorname{det}\left(1+\frac{i}{2\pi}\Omega\right)=\stackrel{*}{\operatorname{det}}\left(1+\gamma_1+\cdots+\gamma_m\right)$	$\det\left(1+\frac{i}{2\pi}\Omega\right)=\stackrel{*}{\pi}(1+\gamma_1+\cdots+\gamma_m)$	$\det\left(1+\frac{i}{2\pi}\Omega\right)=\stackrel{*}{\pi}(1+\gamma_1+\cdots+\gamma_m)$
mielke-geometrodynamics_48 EQ0071 (2.8.14)	48	■ 0.999	$\stackrel{\infty}{\omega}:=-G^{-1}dG,$	$\stackrel{\infty}{\omega}:=-G^{-1}dG,$	$\stackrel{\infty}{\omega}:=-G^{-1}dG,$
mielke-geometrodynamics_89 EQ0235 (4.2.28)	89	■ 0.999	$Q_{\kappa\mu\nu}:=-D_{\kappa}g_{\mu\nu}$	$Q_{\kappa\mu\nu}:=-D_{\kappa}g_{\mu\nu}$	$Q_{\kappa\mu\nu}:=-D_{\kappa}g_{\mu\nu}$
mielke-geometrodynamics_97 EQ0278 (4.4.6)	97	■ 0.999	$\begin{aligned} &\& D\,\Pi^{\mathrm{T}}-E=\Sigma \\ & D\Pi^{\mathrm{L}}+\Pi_{\square}^{\mathrm{T}}=\tau_{\mathrm{s}} \end{aligned}$	$\begin{aligned} &D\Pi^{\mathrm{T}}-E=\Sigma \\ &D\Pi^{\mathrm{L}}+\Pi_{\square}^{\mathrm{T}}=\tau_{\mathrm{s}} \end{aligned}$	$\begin{aligned} &D\Pi^{\mathrm{T}}-E=\Sigma \\ &D\Pi^{\mathrm{L}}+\Pi_{\square}^{\mathrm{T}}=\tau_{\mathrm{s}} \end{aligned}$
mielke-geometrodynamics_105 EQ0313 (4.6.5)	105	■ 0.999	$\Phi(A,\gamma)=P\exp\left[(i/\hbar)\oint A\right],$	$\Phi(A,\gamma)=P\exp\left[(i/\hbar)\oint A\right],$	$\Phi(A,\gamma)=P\exp\left[(i/\hbar)\oint A\right],$
mielke-geometrodynamics_125 EQ0350 (6.2.9)	125	■ 0.999	$L_{\parallel}:=-\frac{1}{2\kappa}T^{\alpha}\wedge\left(^{(1)}T_{\alpha}-2^{(2)}T_{\alpha}-\frac{1}{2}^{(3)}T_{\alpha}\right),$	$L_{\parallel}:=-\frac{1}{2\kappa}T^{\alpha}\wedge\left(^{(1)}T_{\alpha}-2^{(2)}T_{\alpha}-\frac{1}{2}^{(3)}T_{\alpha}\right),$	$L_{\parallel}:=-\frac{1}{2\kappa}T^{\alpha}\wedge\left(^{(1)}T_{\alpha}-2^{(2)}T_{\alpha}-\frac{1}{2}^{(3)}T_{\alpha}\right),$
mielke-geometrodynamics_152 EQ0461 (7.2.13)	152	■ 0.999	$\stackrel{(\mp)}{\Omega}_{\mathrm{C}}=0$	$\stackrel{(\mp)}{\Omega}_{\mathrm{C}}=0$	$\stackrel{(\mp)}{\Omega}_{\mathrm{C}}=0$
mielke-geometrodynamics_209 EQ0620 (10.1.8)	209	■ 0.999	$\frac{\delta C_{\mathrm{T}}}{\delta\vartheta^{\alpha}}=\frac{1}{\ell^2}T_{\alpha}\quad\text{and}\quad\frac{\delta C_{\mathrm{T}}}{\delta\Gamma^{\star\alpha}}=\frac{(-1)^s}{\ell^2}\eta_{\alpha},$	$\frac{\delta C_{\mathrm{T}}}{\delta\vartheta^{\alpha}}=\frac{1}{\ell^2}T_{\alpha}\quad\text{and}\quad\frac{\delta C_{\mathrm{T}}}{\delta\Gamma^{\star\alpha}}=\frac{(-1)^s}{\ell^2}\eta_{\alpha},$	$\frac{\delta C_{\mathrm{T}}}{\delta\vartheta^{\alpha}}=\frac{1}{\ell^2}T_{\alpha}\quad\text{and}\quad\frac{\delta C_{\mathrm{T}}}{\delta\Gamma^{\star\alpha}}=\frac{(-1)^s}{\ell^2}\eta_{\alpha},$
mielke-geometrodynamics_218 EQ0667 (10.3.24)	218	■ 0.999	$\left[\square+(-1)^sm^2\right]\vartheta^{\alpha}\cong0,$	$\left[\square+(-1)^sm^2\right]\vartheta^{\alpha}\cong0,$	$\left[\square+(-1)^sm^2\right]\vartheta^{\alpha}\cong0,$
mielke-geometrodynamics_220 EQ0678 (10.3.35)	220	■ 0.999	$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}}-\check{\mathcal{E}}_{\alpha}=\frac{\theta_{\mathrm{TL}}}{\ell}d\Gamma_{\alpha}^{\star},$	$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}}-\check{\mathcal{E}}_{\alpha}=\frac{\theta_{\mathrm{TL}}}{\ell}d\Gamma_{\alpha}^{\star},$	$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}}-\check{\mathcal{E}}_{\alpha}=\frac{\theta_{\mathrm{TL}}}{\ell}d\Gamma_{\alpha}^{\star},$
mielke-geometrodynamics_238 EQ0740 (11.1.7)	238	■ 0.999	$\mathrm{XX}=-\mathrm{o}_{\alpha\beta}x^{\alpha}x^{\beta}\mathbb{1}$	$\mathrm{XX}=-\mathrm{o}_{\alpha\beta}x^{\alpha}x^{\beta}\mathbb{1}$	$\mathrm{XX}=-\mathrm{o}_{\alpha\beta}x^{\alpha}x^{\beta}\mathbb{1}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_00776 (11.2.5)	246	■ 0.999	$L_{\mathrm{D}}=\frac{i}{2}\left(\bar{\psi}\gamma\wedge d\psi+d\bar{\psi}\wedge\gamma\psi\right)-m\bar{\psi}\psi\eta-\frac{1}{2}\bar{\psi}\left[\widetilde{\omega},*\gamma\right]\psi,$	$L_{\mathrm{D}}=\frac{i}{2}\left(\bar{\psi}\gamma\wedge^*d\psi+d\bar{\psi}\wedge^*\gamma\psi\right)-m\bar{\psi}\psi\eta-\frac{1}{2}\bar{\psi}\left[\widetilde{\omega},*\gamma\right]\psi,$	$L_{\mathrm{D}}=\frac{i}{2}(\bar{\psi}\gamma\wedge^*d\psi+d\bar{\psi}\wedge^*\gamma\psi)-m\bar{\psi}\psi\eta-\frac{1}{2}\bar{\psi}[\widetilde{\omega},*\gamma]\psi,$
mielke-geometrodynamics_00805 (11.4.17)	255	■ 0.999	$\kappa=\mp\left(j+\frac{1}{2}\right)\quad\text{for}\quad j=l\pm\frac{1}{2}.$	$\kappa=\mp\left(j+\frac{1}{2}\right)\quad\text{for}\quad j=l\pm\frac{1}{2}.$	$\kappa=\mp(j+\frac{1}{2})\quad\text{for}\quad j=l\pm\frac{1}{2}.$
mielke-geometrodynamics_00989 (14.3.2)	320	■ 0.999	$\hat{L}_{\mathrm{D}}=\frac{i}{2}\left(\overline{\hat{\psi}\hat{\gamma}}\wedge^*\hat{D}\hat{\psi}+(\overline{\hat{D}\hat{\psi}})\wedge^*\hat{\gamma}\hat{\psi}\right)-\hat{m}\hat{\psi}\hat{\psi}\eta$	$\hat{L}_{\mathrm{D}}=\frac{i}{2}\left(\bar{\hat{\psi}}\hat{\gamma}\wedge^*\hat{D}\hat{\psi}+(\overline{\hat{D}\hat{\psi}})\wedge^*\hat{\gamma}\hat{\psi}\right)-\hat{m}\bar{\hat{\psi}}\hat{\psi}\eta$	$\hat{L}_{\mathrm{D}}=\frac{i}{2}(\bar{\hat{\psi}}\hat{\gamma}\wedge^*\hat{D}\hat{\psi}+(\overline{\hat{D}\hat{\psi}})\wedge^*\hat{\gamma}\hat{\psi})-\hat{m}\bar{\hat{\psi}}\hat{\psi}\eta$
mielke-geometrodynamics_01057 (15.2.14)	346	■ 0.999	$i\,\bar{\gamma}\wedge^*\bar{D}^{\{\}}\psi-6\ell^2\bar{\gamma}^5\bar{\gamma}\wedge^*(\bar{\psi}\bar{\gamma}^5\bar{\gamma}\psi)\,\psi-m\psi\eta=0$	$i\bar{\gamma}\wedge^*\bar{D}^{\{\}}\psi-6\ell^2\bar{\gamma}^5\bar{\gamma}\wedge^*(\bar{\psi}\bar{\gamma}^5\bar{\gamma}\psi)\,\psi-m\psi\eta=0$	$i\,\overline{\gamma}\wedge^*\overline{D}^{\{\}}\psi-6\,\ell^2\,\overline{\gamma}^5\,\overline{\gamma}\wedge^*(\overline{\psi}\,\overline{\gamma}^5\,\overline{\gamma}\,\psi)\psi-m\psi\eta=0$
mielke-geometrodynamics_00669 (10.3.26)	219	■ 0.999	$m=\frac{\theta_{\mathrm{T}}\theta_{\mathrm{TL}}}{2\left(\theta_{\mathrm{TL}}^2+\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right)\ell},\quad\Lambda_{\mathrm{ind}}=\frac{\theta_{\mathrm{T}}^2\left[-(-1)^s3\theta_{\mathrm{TL}}^2+8\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right]}{4\left[(-1)^s\theta_{\mathrm{TL}}^2-2\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right]^2\ell^2},$	$m=\frac{\theta_{\mathrm{T}}\theta_{\mathrm{TL}}}{2\left(\theta_{\mathrm{TL}}^2+\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right)\ell},\quad\Lambda_{\mathrm{ind}}=\frac{\theta_{\mathrm{T}}^2\left[-(-1)^s3\theta_{\mathrm{TL}}^2+8\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right]}{4\left[(-1)^s\theta_{\mathrm{TL}}^2-2\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right]^2\ell^2},$	$m=\frac{\theta_{\mathrm{T}}\theta_{\mathrm{TL}}}{2(\theta_{\mathrm{TL}}^2+\theta_{\mathrm{T}}\theta_{\mathrm{L}})\ell},\quad\Lambda_{\mathrm{ind}}=\frac{\theta_{\mathrm{T}}^2[-(-1)^s3\theta_{\mathrm{TL}}^2+8\theta_{\mathrm{T}}\theta_{\mathrm{L}}]}{4[(-1)^s\theta_{\mathrm{TL}}^2-2\theta_{\mathrm{T}}\theta_{\mathrm{L}}]^2\ell^2},$
mielke-geometrodynamics_00929 (14.1.10)	304	■ 0.999	$\begin{aligned} &\operatorname{Tr}\left(\Sigma_{\mathrm{YM}}\wedge^*\vartheta\right)=\operatorname{Tr}\left(\vartheta\wedge^*\Sigma_{\mathrm{YM}}\right)\\ &=\frac{1}{2\alpha_g}\operatorname{Tr}\left\{\vartheta\wedge\Omega\wedge^*(\vartheta\wedge^*\Omega)+\vartheta\wedge^*\vartheta\wedge^*(\Omega\wedge^*\Omega)\right\}=0 \end{aligned}$	$\begin{aligned} \operatorname{Tr}\left(\Sigma_{\mathrm{YM}}\wedge^*\vartheta\right)&=\operatorname{Tr}\left(\vartheta\wedge^*\Sigma_{\mathrm{YM}}\right)\\ &=\frac{1}{2\alpha_g}\operatorname{Tr}\left\{\vartheta\wedge\Omega\wedge^*(\vartheta\wedge^*\Omega)+\vartheta\wedge^*\vartheta\wedge^*(\Omega\wedge^*\Omega)\right\}=0 \end{aligned}$	$\begin{aligned} \operatorname{Tr}(\Sigma_{\mathrm{YM}}\wedge^*\vartheta)&=\operatorname{Tr}(\vartheta\wedge^*\Sigma_{\mathrm{YM}})\\ &=\frac{1}{2\alpha_g}\operatorname{Tr}\left\{\vartheta\wedge\Omega\wedge^*(\vartheta\wedge^*\Omega)+\vartheta\wedge^*\vartheta\wedge^*(\Omega\wedge^*\Omega)\right\}=0 \end{aligned}$
mielke-geometrodynamics_00093 (3.2.3)	55	■ 0.999	$\omega^{\prime}\rightarrow^{G^{-1}}\omega^{\prime}=G^{-1}\omega^{\prime}G+(dG^{-1})\,G$	$\omega'\rightarrow^{G^{-1}}\omega'=G^{-1}\omega'G+(dG^{-1})\,G$	$\omega'\rightarrow^{G^{-1}}\omega'=G^{-1}\omega'G+(dG^{-1})G$
mielke-geometrodynamics_00113 (3.3.14)	59	■ 0.999	$H=\left\langle\right\rangle^*F\;.$	$H=\left\langle\right\rangle^*F.$	$H=\left\langle\right\rangle^*F\;.$
mielke-geometrodynamics_00173 (3.7.1)	71	■ 0.999	$L_{\mathrm{H}}=-\bar{\Theta}^{(1)}\wedge^*D\phi^0+\bar{\Theta}^{(1)}\wedge^*\Theta^{(1)}+\frac{\lambda}{4}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)^2\eta.$	$L_{\mathrm{H}}=-\bar{\Theta}^{(1)}\wedge^*D\phi^0+\bar{\Theta}^{(1)}\wedge^*\Theta^{(1)}+\frac{\lambda}{4}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)^2\eta.$	$L_{\mathrm{H}}=-\bar{\Theta}^{(1)}\wedge^*D\phi^0+\bar{\Theta}^{(1)}\wedge^*\Theta^{(1)}+\frac{\lambda}{4}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)^2\eta.$
mielke-geometrodynamics_00201 (4.1.15)	83	■ 0.999	$\delta_g^*\vartheta=g^{-1}\vartheta,\quad g\in GL(n,\mathbb{R}),$	$\delta_g^*\vartheta=g^{-1}\vartheta,\quad g\in GL(n,\mathbb{R}),$	$\delta_g^*\vartheta=g^{-1}\vartheta,\quad g\in GL(n,\mathbb{R}),$
mielke-geometrodynamics_00275 (4.4.3)	97	■ 0.999	$\begin{aligned} &\delta(\delta\vartheta)-[\omega^{\mathrm{L}},\delta\vartheta]-[\delta\omega^{\mathrm{L}},\vartheta] \\ &=D(\delta\vartheta)-[\delta\omega^{\mathrm{L}},\vartheta]. \end{aligned}$	$\begin{aligned} &=d(\delta\vartheta)-[\omega^{\mathrm{L}},\delta\vartheta]-[\delta\omega^{\mathrm{L}},\vartheta] \\ &=D(\delta\vartheta)-[\delta\omega^{\mathrm{L}},\vartheta]. \end{aligned}$	$\begin{aligned} &=d(\delta\vartheta)-[\omega^{\mathrm{L}},\delta\vartheta]-[\delta\omega^{\mathrm{L}},\vartheta] \\ &=D(\delta\vartheta)-[\delta\omega^{\mathrm{L}},\vartheta]. \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00341 (5.3.18)	115	■ 0.999	$\{ \}^{*} \ \mathrm{mathscr{A}}=\frac{\kappa}{2} \ j_5,$	${}^*\mathscr{A}=\frac{\kappa}{2}j_5,$	${}^*\mathscr{A}=\frac{\kappa}{2}j_5,$
mielke-geometrodynamics_E00441 (6.8.12)	143	■ 0.999	$D \ \tau_{\alpha \beta}+\vartheta_{[\alpha} \ \Sigma_{\beta]} \cong 0,$	$D\tau_{\alpha\beta}+\vartheta_{[\alpha}\wedge\Sigma_{\beta]}\cong 0,$	$D\tau_{\alpha\beta}+\vartheta_{[\alpha}\wedge\Sigma_{\beta]}\cong 0,$
mielke-geometrodynamics_E00476 (7.4.4)	156	■ 0.999	$C_{\alpha \beta}=0 \ \text{and} \ R=0,$	$C_{\alpha\beta}=0 \quad \text{and} \quad R=0,$	$C_{\alpha\beta}=0 \quad \text{and} \quad R=0,$
mielke-geometrodynamics_E00519 (7.7.37)	166	■ 0.999	$D \ \Theta \equiv \left[\Omega^g, \gamma \right], \quad D \ \Omega^g \equiv 0,$	$D\Theta\equiv[\Omega^g,\gamma], \quad D\Omega^g\equiv 0,$	$D\Theta\equiv[\Omega^g,\gamma], \quad D\Omega^g\equiv 0,$
mielke-geometrodynamics_E00567 (8.6.4)	183	■ 0.999	$\begin{aligned} H_{\alpha \beta}(\ast \ast) \ \&= \vartheta_{\alpha} \ R_{\beta} + \vartheta_{\beta} \ R_{\alpha} + \frac{\vartheta_{\alpha} \ \vartheta_{\beta}}{\ell^2} \ \eta_{\alpha \beta} - \frac{\vartheta_{\alpha} \ \vartheta_{\beta}}{\ell^2} \ \vartheta_{\alpha} \wedge \vartheta_{\beta} \\ &\simeq -\partial L_{\theta} / \partial R^{\alpha \beta}, \end{aligned}$	$H_{\alpha\beta}(\ast\ast)=\theta_{\mathrm{L}}R_{\alpha\beta}+\theta_{\mathrm{L}}^{\star}R_{\alpha\beta}^{(\ast)}+\frac{\theta_{\mathrm{T}}^{\star}}{2\ell^2}\eta_{\alpha\beta}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}$ $\simeq -\partial L_{\theta}/\partial R^{\alpha\beta}$	$H_{\alpha\beta}(\ast\ast)=\theta_{\mathrm{L}}R_{\alpha\beta}+\theta_{\mathrm{L}}^{\star}R_{\alpha\beta}^{(\ast)}+\frac{\theta_{\mathrm{T}}^{\star}}{2\ell^2}\eta_{\alpha\beta}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}$ $\simeq -\partial L_{\theta}/\partial R^{\alpha\beta},$
mielke-geometrodynamics_E00624 (10.1.12)	210	■ 0.999	$D \ H_{\alpha} - E_{\alpha} = \Sigma_{\alpha},$	$DH_{\alpha}-E_{\alpha}=\Sigma_{\alpha},$	$DH_{\alpha}-E_{\alpha}=\Sigma_{\alpha},$
mielke-geometrodynamics_E00673 (10.3.30)	220	■ 0.999	$H_{\alpha}=-\frac{\vartheta_{\alpha}}{\ell^2} \ \vartheta_{\alpha} - \frac{\vartheta_{\alpha}}{\ell} \ \Gamma_{\alpha}^{\star}, \quad H_{\alpha}^{\star}=-\frac{\vartheta_{\alpha}}{2} \ \Gamma_{\alpha}^{\star},$	$H_{\alpha}=-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}-\frac{\theta_{\mathrm{TL}}}{\ell}\Gamma_{\alpha}^{\star}, \quad H_{\alpha}^{\star}=-\frac{\theta_{\mathrm{L}}}{2}\Gamma_{\alpha}^{\star},$	$H_{\alpha}=-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}-\frac{\theta_{\mathrm{TL}}}{\ell}\Gamma_{\alpha}^{\star}, \quad H_{\alpha}^{\star}=-\frac{\theta_{\mathrm{L}}}{2}\Gamma_{\alpha}^{\star},$
mielke-geometrodynamics_E00722 (10.10.2)	230	■ 0.999	$\delta L = \delta \vartheta^{\alpha} \wedge \frac{\delta L}{\delta \vartheta^{\alpha}} + \delta \Gamma_{\alpha}^{\star} \wedge \frac{\delta L}{\delta \Gamma_{\alpha}^{\star}} + \delta \bar{\Psi} \wedge \frac{\delta L}{\delta \bar{\Psi}},$	$\delta L=\delta\vartheta^{\alpha}\wedge\frac{\delta L}{\delta\vartheta^{\alpha}}+\delta\Gamma_{\alpha}^{\star}\wedge\frac{\delta L}{\delta\Gamma_{\alpha}^{\star}}+\delta\bar{\Psi}\wedge\frac{\delta L}{\delta\bar{\Psi}},$	$\delta L=\delta\vartheta^{\alpha}\wedge\frac{\delta L}{\delta\vartheta^{\alpha}}+\delta\Gamma_{\alpha}^{\star}\wedge\frac{\delta L}{\delta\Gamma_{\alpha}^{\star}}+\delta\bar{\Psi}\wedge\frac{\delta L}{\delta\bar{\Psi}},$
mielke-geometrodynamics_E00759 (11.1.26)	242	■ 0.999	$\{ \}^{*}(\underbrace{\gamma \wedge \cdots \wedge \gamma}_p)=\frac{p!}{(n-p)!} \ \gamma^{\wedge n+1} \ \underbrace{\gamma \wedge \cdots \wedge \gamma}_{n-p}.$	${}^*(\underbrace{\gamma\wedge\cdots\wedge\gamma}_p)=\frac{p!}{(n-p)!}\gamma^{n+1}\underbrace{\gamma\wedge\cdots\wedge\gamma}_{n-p}.$	${}^*(\underbrace{\gamma\wedge\cdots\wedge\gamma}_p)=\frac{p!}{(n-p)!}\gamma^{n+1}\underbrace{\gamma\wedge\cdots\wedge\gamma}_{n-p}.$
mielke-geometrodynamics_E00780 (11.2.8)	247	■ 0.999	$(\psi, \chi) := \int_{H^{n-1}} \tau_s(\psi, \chi) = i \int_{H^{n-1}} \psi^* \gamma \chi$	$(\psi,\chi):=\int_{H^{n-1}}\tau_s(\psi,\chi)=i\int_{H^{n-1}}\psi^*\gamma\chi$	$(\psi,\chi):=\int_{H^{n-1}}\tau_s(\psi,\chi)=i\int_{H^{n-1}}\psi^*\gamma\chi$
mielke-geometrodynamics_E00789 (11.4.1)	251	■ 0.999	$E_r = \sqrt{\alpha} r_{\circ}^2 \left[1 + \left(\frac{r}{r_{\circ}} \right)^4 \right]^{-1/2} \sim \sqrt{\alpha} \frac{1}{r^2}$	$E_r=\sqrt{\alpha}r_{\circ}^2\left[1+\left(\frac{r}{r_{\circ}}\right)^4\right]^{-1/2}\sim\sqrt{\alpha}\frac{1}{r^2}$	$E_r=\sqrt{\alpha}r_{\circ}^2\left[1+\left(\frac{r}{r_{\circ}}\right)^4\right]^{-1/2}\sim\sqrt{\alpha}\frac{1}{r^2}$
mielke-geometrodynamics_E00825 (11.4.39)	259	■ 0.999	$m_{\pi}=\sqrt{1-\omega^2} \ m$	$m_{\pi}=\sqrt{1-\omega^2}m$	$m_{\pi}=\sqrt{1-\omega^2} \ m$
mielke-geometrodynamics_E00892 (13.4.11)	287	■ 0.999	$\vartheta \xrightarrow{A^{-1}(x)} \vartheta' = \Lambda^{-1}(x) \vartheta, \quad \theta \xrightarrow{A^{-1}(x)} \theta' = \theta \Lambda^{-1}(x).$	$\vartheta \xrightarrow{A^{-1}(x)} \vartheta' = \Lambda^{-1}(x) \vartheta, \quad \theta \xrightarrow{A^{-1}(x)} \theta' = \theta \Lambda^{-1}(x).$	$\vartheta \xrightarrow{A^{-1}(x)} \vartheta' = \Lambda^{-1}(x) \vartheta, \quad \theta \xrightarrow{A^{-1}(x)} \theta' = \theta \Lambda^{-1}(x).$
mielke-geometrodynamics_E00932 (14.1.13)	304	■ 0.999	$e^{\delta} \circ e^{\varepsilon} = e^{\delta + \varepsilon},$	$e^{*\delta}\circ e^{*\varepsilon}=e^{*\varepsilon}\circ e^{*\delta}=e^{*(\delta+\varepsilon)},$	$e^{*\delta}\circ e^{*\varepsilon}=e^{*\varepsilon}\circ e^{*\delta}=e^{*(\delta+\varepsilon)},$
mielke-geometrodynamics_E00940 (14.1.21)	305	■ 0.999	$\xi \wedge \xi = \xi \wedge^* \xi = 0$	$\xi\wedge\xi=\xi\wedge^*\xi=0$	$\xi\wedge\xi=\xi\wedge^*\xi=0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_00833 (12.2.1)	270	■ 0.999	$L_{\mathrm{D}}=L(\gamma,\psi,\mathscr{D}\psi)=\frac{i}{2}\left\{\bar{\psi}\gamma\wedge\mathscr{D}\psi+\overline{\mathscr{D}\psi}\wedge\gamma\psi\right\}-m\bar{\psi}\psi\eta,$	$L_{\mathrm{D}}=L(\gamma,\psi,\mathscr{D}\psi)=\frac{i}{2}\left\{\bar{\psi}^*\gamma\wedge\mathscr{D}\psi+\overline{\mathscr{D}\psi}\wedge^*\gamma\psi\right\}-m\bar{\psi}\psi\eta,$	$L_{\mathrm{D}}=L(\gamma,\psi,\mathscr{D}\psi)=\frac{i}{2}\left\{\overline{\psi}^*\gamma\wedge\mathscr{D}\psi+\overline{\mathscr{D}\psi}\wedge^*\gamma\psi\right\}-m\bar{\psi}\psi\eta,$
mielke-geometrodynamics_00826 (2.4.2)	38	■ 1.000	$V^{\rho}:=V^{\rho}\left(M,\mathbb{C}^N,\rho(G)\subseteq GL(N,\mathbb{C}),P\right),$	$V^{\rho}:=V^{\rho}\left(M,\mathbb{C}^N,\rho(G)\subseteq GL(N,\mathbb{C}),P\right),$	$V^{\rho}:=V^{\rho}(M,\mathbb{C}^N,\rho(G)\subseteq GL(N,\mathbb{C}),P),$
mielke-geometrodynamics_00891 (3.2.1)	55	■ 1.000	$\mathscr{0}\rightarrow{}^{G^{-1}}\mathscr{0}=G^{-1}\mathscr{O}G.$	$\mathscr{O}\rightarrow{}^{G^{-1}}\mathscr{O}=G^{-1}\mathscr{O}G.$	$\mathscr{O}\rightarrow{}^{G^{-1}}\mathscr{O}=G^{-1}\mathscr{O}G.$
mielke-geometrodynamics_00181 (3.3.2)	57	■ 1.000	$A\rightarrowtail A+d\,\theta(x)\quad\quad\quad\rightarrowtail\quad\quad\quad F\rightarrowtail d\,A+d\,\theta=A=F.$	$A\rightarrow A+d\theta(x)\quad\Rightarrow\quad F\rightarrow dA+dd\theta=dA=F.$	$A\rightarrow A+d\theta(x)\quad\Rightarrow\quad F\rightarrow dA+dd\theta=dA=F.$
mielke-geometrodynamics_00137 (3.4.8)	63	■ 1.000	$\begin{aligned} \delta L_{\omega} &= \delta \Omega \wedge \Pi^{\Omega} = D(\delta \omega) \wedge \Pi^{\Omega} \\ &= -\delta \omega \wedge D \Pi^{\Omega} + d\left(\delta \omega \wedge \Pi^{\Omega}\right). \end{aligned}$	$\delta L_{\omega} = \delta \Omega \wedge \Pi^{\Omega} = D(\delta \omega) \wedge \Pi^{\Omega} \\ = -\delta \omega \wedge D \Pi^{\Omega} + d\left(\delta \omega \wedge \Pi^{\Omega}\right).$	$\delta L_{\omega} = \delta \Omega \wedge \Pi^{\Omega} = D(\delta \omega) \wedge \Pi^{\Omega} \\ = -\delta \omega \wedge D \Pi^{\Omega} + d(\delta \omega \wedge \Pi^{\Omega}).$
mielke-geometrodynamics_00146 (3.4.17)	65	■ 1.000	$Q=Q^{\left(i\right)}I_i:=\int_{H^3}\iota=\int_{\partial H^3}\Pi^{\Omega},$	$Q=Q^{\left(i\right)}I_i:=\int_{H^3}\iota=\int_{\partial H^3}\Pi^{\Omega},$	$Q=Q^{\left(i\right)}I_i:=\int\limits_{H^3}\iota=\int\limits_{\partial H^3}\Pi^{\Omega},$
mielke-geometrodynamics_00175 (3.7.4)	72	■ 1.000	$\tau_{\mathrm{H}}=-\left[\phi^{(0)},*\Theta^{(1)}\right]$	$\tau_{\mathrm{H}}=-\left[\phi^{(0)},*\Theta^{(1)}\right]$	$\tau_{\mathrm{H}}=-[\phi^{(0)},*\Theta^{(1)}]$
mielke-geometrodynamics_00195 (4.1.5)	82	■ 1.000	$\mathfrak{a}(n,\mathbb{R})=\mathbb{R}^n\otimes\mathfrak{gl}(n,\mathbb{R})$	$\mathfrak{a}(n,\mathbb{R})=\mathbb{R}^n\otimes\mathfrak{gl}(n,\mathbb{R})$	$\mathfrak{a}(n,\mathbb{R})=\mathbb{R}^n\otimes\mathfrak{gl}(n,\mathbb{R})$
mielke-geometrodynamics_00208 (4.1.22)	84	■ 1.000	$T_{\mathrm{A}}(\mathrm{M})=\stackrel{\circ}{V}(M,\mathbb{R}^n,A(n,\mathbb{R}),A(M))\subset V^{\mathrm{id}}$	$T_{\mathrm{A}}(M)=\stackrel{\circ}{V}(M,\mathbb{R}^n,A(n,\mathbb{R}),A(M))\subset V^{\mathrm{id}}$	$T_{\mathrm{A}}(M)=\stackrel{\circ}{V}(M,\mathbb{R}^n,A(n,\mathbb{R}),A(M))\subset V^{\mathrm{id}}$
mielke-geometrodynamics_00212 (4.2.4)	85	■ 1.000	$\Omega^{\mathrm{T}}:=D\omega^{\mathrm{T}}=d\omega^{\mathrm{T}}-[\omega^{\mathrm{T}},\omega^{\mathrm{L}}],$	$\Omega^{\mathrm{T}}:=D\omega^{\mathrm{T}}=d\omega^{\mathrm{T}}-[\omega^{\mathrm{T}},\omega^{\mathrm{L}}],$	$\Omega^{\mathrm{T}}:=D\omega^{\mathrm{T}}=d\omega^{\mathrm{T}}-[\omega^{\mathrm{T}},\omega^{\mathrm{L}}],$
mielke-geometrodynamics_00215 (4.2.7)	86	■ 1.000	$\mathscr{T}(n,\mathbb{R}):=C^{\infty}\left(A(M)\times_{\mathrm{Ad}}\mathbb{R}^n\right)$	$\mathscr{T}(n,\mathbb{R}):=C^{\infty}\left(A(M)\times_{\mathrm{Ad}}\mathbb{R}^n\right)$	$\mathscr{T}(n,\mathbb{R}):=C^{\infty}(A(M)\times_{\mathrm{Ad}}\mathbb{R}^n)$
mielke-geometrodynamics_00217 (4.2.9)	86	■ 1.000	$A=\left[\begin{array}{c c} G&T\\ \hline 0&1\end{array}\right]\in\mathscr{A}_p,\quad G\in\mathscr{G}_p,T\in\mathscr{T}_p$	$A=\left[\begin{array}{c c} G&T\\ \hline 0&1\end{array}\right]\in\mathscr{A}_p,\quad G\in\mathscr{G}_p,T\in\mathscr{T}_p$	$A=\left[\begin{array}{c c} G&T\\ \hline 0&1\end{array}\right]\in\mathscr{A}_p,\quad G\in\mathscr{G}_p,T\in\mathscr{T}_p$
mielke-geometrodynamics_00243 (4.3.8)	91	■ 1.000	$\cos\phi=\frac{g_{\mu\nu}dV^{\mu}dW^{\nu}}{\sqrt{g_{\alpha\beta}dV^{\alpha}dV^{\beta}}\sqrt{g_{\alpha\beta}dW^{\alpha}dW^{\beta}}}.$	$\cos\phi=\frac{g_{\mu\nu}dV^{\mu}dW^{\nu}}{\sqrt{g_{\alpha\beta}dV^{\alpha}dV^{\beta}}\sqrt{g_{\alpha\beta}dW^{\alpha}dW^{\beta}}}.$	$\cos\phi=\frac{g_{\mu\nu}dV^{\mu}dW^{\nu}}{\sqrt{g_{\alpha\beta}dV^{\alpha}dV^{\beta}}\sqrt{g_{\alpha\beta}dW^{\alpha}dW^{\beta}}}.$
mielke-geometrodynamics_00323 (5.2.7) [re-fined: census]	111	■ 1.000	$\begin{aligned} \Sigma_{\alpha}&\rightarrow\sigma_{\alpha}:=\Sigma_{\alpha}-D\mu_{\alpha}+e_{\beta}\rfloor(T^{\beta}\wedge\mu_{\alpha}),\\ \tau_{\alpha\beta}&\rightarrow\hat{\tau}_{\alpha\beta}:=\tau_{\alpha\beta}-\vartheta_{[\alpha}\wedge\mu_{\beta]}=0 \end{aligned}$	$\begin{aligned} \Sigma_{\alpha}&\rightarrow\sigma_{\alpha}:=\Sigma_{\alpha}-D\mu_{\alpha}+e_{\beta}\rfloor\left(T^{\beta}\wedge\mu_{\alpha}\right),\\ \tau_{\alpha\beta}&\rightarrow\hat{\tau}_{\alpha\beta}:=\tau_{\alpha\beta}-\vartheta_{[\alpha}\wedge\mu_{\beta]}=0 \end{aligned}$	$\begin{aligned} \Sigma_{\alpha}&\rightarrow\sigma_{\alpha}:=\Sigma_{\alpha}-D\mu_{\alpha}+e_{\beta}\rfloor(T^{\beta}\wedge\mu_{\alpha}),\\ \tau_{\alpha\beta}&\rightarrow\hat{\tau}_{\alpha\beta}:=\tau_{\alpha\beta}-\vartheta_{[\alpha}\wedge\mu_{\beta]}=0 \end{aligned}$
mielke-geometrodynamics_00328 (5.3.5)	112	■ 1.000	$\Theta:=D\gamma\quad\text{and}\quad\Omega^g:=d\Gamma+\Gamma\wedge\Gamma.$	$\Theta:=D\gamma\quad\text{and}\quad\Omega^g:=d\Gamma+\Gamma\wedge\Gamma.$	$\Theta:=D\gamma\quad\text{and}\quad\Omega^g:=d\Gamma+\Gamma\wedge\Gamma.$
mielke-geometrodynamics_00370 (6.2.21)	127	■ 1.000	$\stackrel{(\pm)}{\Pi}_{\alpha}=-\partial\stackrel{(\pm)}{L}/\partial T^{\alpha}=H_{\alpha}^{\parallel}\mp(i/\kappa)T_{\alpha}$	$\stackrel{(\pm)}{\Pi}_{\alpha}=-\partial\stackrel{(\pm)}{L}/\partial T^{\alpha}=H_{\alpha}^{\parallel}\mp(i/\kappa)T_{\alpha}$	$\stackrel{(\pm)}{P}_{\alpha}=-\partial\stackrel{(\pm)}{L}/\partial T^{\alpha}=H_{\alpha}^{\parallel}\mp(i/\kappa)T_{\alpha}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_220 EQ0676 (10.3.33)		■ 1.000	$\check{\mathrm{E}}_{\alpha}^{\mathrm{MB}} = -(-1)^s \frac{\theta_{\mathrm{T}} \theta_{\mathrm{L}}}{2 \chi \ell} \left(d \Gamma_{\alpha}^{\star} + \frac{2}{\theta_{\mathrm{L}}} \check{\mathscr{J}}_{\alpha}^{\mathrm{MB}\star} \right),$		$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}} = -(-1)^s \frac{\theta_{\mathrm{T}} \theta_{\mathrm{L}}}{2 \chi \ell} \left(d \Gamma_{\alpha}^{\star} + \frac{2}{\theta_{\mathrm{L}}} \check{\mathscr{J}}_{\alpha}^{\mathrm{MB}\star} \right),$
mielke-geometrodynamics_222 EQ0688 (10.3.46)		■ 1.000	$G_{\alpha}^{\{\}} - \tilde{\Lambda} \eta_{\alpha} = \ell \tilde{\Sigma}_{\alpha},$		$G_{\alpha}^{\{\}} - \tilde{\Lambda} \eta_{\alpha} = \ell \tilde{\Sigma}_{\alpha},$
mielke-geometrodynamics_223 EQ0690 (10.5.1)		■ 1.000	$\sigma^1 = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix} \right), \quad \sigma^2 = \left(\begin{smallmatrix} 0 & -i \\ i & 0 \end{smallmatrix} \right), \quad \sigma^3 = \left(\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix} \right).$		$\sigma^1 = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix} \right), \quad \sigma^2 = \left(\begin{smallmatrix} 0 & -i \\ i & 0 \end{smallmatrix} \right), \quad \sigma^3 = \left(\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix} \right).$
mielke-geometrodynamics_227 EQ0710 (10.7.6)		■ 1.000	$\begin{aligned} \vartheta^{\hat{0}} &= dt + \ell^2 \sigma \rho^{*2} [1 - \cos(\rho/\rho^*)] d\phi \\ \vartheta^{\hat{1}} &= d\rho, \quad \vartheta^{\hat{2}} = \rho^* \sin(\rho/\rho^*) d\phi, \\ \Gamma^{\hat{1}\hat{2}} &= \cos(\rho/\rho^*) d\phi = -\Gamma^{\hat{2}\hat{1}} \end{aligned}$		$\begin{aligned} \vartheta^{\hat{0}} &= dt + \ell^2 \sigma \rho^{*2} [1 - \cos(\rho/\rho^*)] d\phi \\ \vartheta^{\hat{1}} &= d\rho, \quad \vartheta^{\hat{2}} = \rho^* \sin(\rho/\rho^*) d\phi, \\ \Gamma^{\hat{1}\hat{2}} &= \cos(\rho/\rho^*) d\phi = -\Gamma^{\hat{2}\hat{1}}. \end{aligned}$
mielke-geometrodynamics_229 EQ0716 (10.9.2)		■ 1.000	$D \Psi = d \Psi - \frac{1}{2} \gamma_{\alpha} \Gamma^{\star \alpha} \wedge \Psi,$		$D \Psi = d \Psi - \frac{1}{2} \gamma_{\alpha} \Gamma^{\star \alpha} \wedge \Psi,$
mielke-geometrodynamics_229 EQ0720 (10.9.6)		■ 1.000	$\tau_{\alpha}^{\star} := \frac{1}{2} \eta_{\alpha \beta \gamma} \tau^{\beta \gamma} = \frac{(-1)^s}{2} \frac{\delta L_{\Psi}}{\delta \Gamma_{\alpha}^{\star}} = \frac{(-1)^s}{2} \frac{i}{4} \bar{\Psi} \wedge \gamma_{\alpha} \Psi.$		$\tau_{\alpha}^{\star} := \frac{1}{2} \eta_{\alpha \beta \gamma} \tau^{\beta \gamma} = \frac{(-1)^s}{2} \frac{\delta L_{\Psi}}{\delta \Gamma_{\alpha}^{\star}} = \frac{(-1)^s}{2} \frac{i}{4} \bar{\Psi} \wedge \gamma_{\alpha} \Psi.$
mielke-geometrodynamics_231 EQ0728 (10.10.8)		■ 1.000	$\gamma^{\alpha} \Psi \wedge \bar{\Psi} \gamma_{\alpha} \Psi = 0,$		$\gamma^{\alpha} \Psi \wedge \bar{\Psi} \gamma_{\alpha} \Psi = 0,$
mielke-geometrodynamics_244 EQ0770 (11.1.37)		■ 1.000	$D \longrightarrow^{S^{-1}} D = S^{-1} D S.$		$D \longrightarrow^{S^{-1}} D = S^{-1} D S.$
mielke-geometrodynamics_272 EQ0839 (12.2.7)		■ 1.000	$j_{\pm} := \frac{1}{2} \bar{\psi} (1 \pm \gamma_5)^* \gamma \psi = \bar{\psi}_{\mathrm{L}, \mathrm{R}}^* \gamma \psi_{\mathrm{L}, \mathrm{R}}.$		$j_{\pm} := \frac{1}{2} \bar{\psi} (1 \pm \gamma_5)^* \gamma \psi = \bar{\psi}_{\mathrm{L}, \mathrm{R}}^* \gamma \psi_{\mathrm{L}, \mathrm{R}}.$
mielke-geometrodynamics_284 EQ0873 (13.3.1)		■ 1.000	$\lrcorner^A{}_B := L^A{}_B - \frac{1}{n} \delta_B^A L^C{}_C.$		$\lrcorner^A{}_B := L^A{}_B - \frac{1}{n} \delta_B^A L^C{}_C.$
mielke-geometrodynamics_287 EQ0893 (13.4.12)		■ 1.000	$A_{\star}(n, \mathbb{R})/\mathrm{GL}(n, \mathbb{R}) \approx \mathbb{R}^n \otimes \mathbb{R}_{\star}^n,$		$A_{\star}(n, \mathbb{R})/\mathrm{GL}(n, \mathbb{R}) \approx \mathbb{R}^n \otimes \mathbb{R}_{\star}^n,$
mielke-geometrodynamics_290 EQ0907 (13.5.12)		■ 1.000	$\tilde{L}_{\mathrm{SSB}} + d\hat{C} \cong \frac{1}{2\mu} \eta_{\alpha \beta \gamma \delta} \left[R^{\alpha \beta} - \frac{\Lambda}{3} \vartheta^{\alpha} \wedge \vartheta^{\beta} \right] \wedge \left[R^{\gamma \delta} - \frac{\Lambda}{3} \vartheta^{\gamma} \wedge \vartheta^{\delta} \right].$		$\tilde{L}_{\mathrm{SSB}} + d\hat{C} \cong \frac{1}{2\mu} \eta_{\alpha \beta \gamma \delta} \left[R^{\alpha \beta} - \frac{\Lambda}{3} \vartheta^{\alpha} \wedge \vartheta^{\beta} \right] \wedge \left[R^{\gamma \delta} - \frac{\Lambda}{3} \vartheta^{\gamma} \wedge \vartheta^{\delta} \right].$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0947 (14.1.28)	307	■ 1.000	$R_{i\alpha}^{\{\}}R_{j\beta}^{\{\}\alpha}=\frac{1}{4}g_{ij}R_{\alpha\beta}^{\{\}}R^{\{\}\alpha\beta}_{\alpha\beta}$	$R_{i\alpha}^{\{\}}R_{j\beta}^{\{\}\alpha}=\frac{1}{4}g_{ij}R_{\alpha\beta}^{\{\}}R^{\{\}\alpha\beta}$	$R_{i\alpha}^{\{\}}R_{j\beta}^{\{\}\alpha}=\frac{1}{4}g_{ij}R_{\alpha\beta}^{\{\}}R^{\{\}\alpha\beta}$
mielke-geometrodynamics_EQ1004 (14.3.17)	323	■ 1.000	$\widetilde{\Theta}:=D\gamma=\frac{1}{2}T_{\alpha\beta}^{\cdot\cdot c}\gamma_c\otimes\vartheta^\alpha\wedge\vartheta^\beta$	$\tilde{\Theta}:=D\gamma=\frac{1}{2}T_{\alpha\beta}^{\cdot\cdot c}\gamma_c\otimes\vartheta^\alpha\wedge\vartheta^\beta$	$\tilde{\Theta}:=D\gamma=\frac{1}{2}T_{\alpha\beta}^{\cdot\cdot c}\gamma_c\otimes\vartheta^\alpha\wedge\vartheta^\beta$
mielke-geometrodynamics_EQ1013 (14.4.8)	326	■ 1.000	$M_{\mathrm{ir}}^2:=\frac{S_{\mathrm{A}}}{16\pi}\frac{M^{*4}c^2}{\hbar^2}$	$M_{\mathrm{ir}}^2:=\frac{S_{\mathrm{A}}}{16\pi}\frac{M^{*4}c^2}{\hbar^2}$	$M_{\mathrm{ir}}^2:=\frac{S_{\mathrm{A}}}{16\pi}\frac{M^{*4}c^2}{\hbar^2}$
mielke-geometrodynamics_EQ1135 (C.3)	375	■ 1.000	$\left[I_i,I_j\right]=c_{ij}^{k}I_k\;.$	$[I_i,I_j]=c_{ij}^{k}I_k.$	$[I_i,I_j]=c_{ij}^{k}I_k.$
mielke-geometrodynamics_EQ0001 (1.1)	24	■ 1.000	$\ell^*:=\sqrt{8\pi\hbar G_N/c^3}\simeq 8\times 10^{-33}\,\mathrm{cm},$	$\ell^*:=\sqrt{8\pi\hbar G_N/c^3}\simeq 8\times 10^{-33}\,\mathrm{cm},$	$\ell^*:=\sqrt{8\pi\hbar G_N/c^3}\simeq 8\times 10^{-33}\mathrm{cm},$
mielke-geometrodynamics_EQ0003 (2.1.2)	32	■ 1.000	$e^{\mathrm{f}}:=\left.\frac{df(s(t))}{dt}\right _{t_o=s^{-1}(m)}=\frac{\partial f}{\partial x^i}\cdot\frac{dx^i(s(t))}{dt}$	$e^{\mathrm{f}}:=\left.\frac{df(s(t))}{dt}\right _{t_o=s^{-1}(m)}=\frac{\partial f}{\partial x^i}\cdot\frac{dx^i(s(t))}{dt}$	$e^{\mathrm{f}}:=\left.\frac{df(s(t))}{dt}\right _{t_o=\bar{s}^{-1}(m)}=\frac{\partial f}{\partial x^i}\cdot\frac{dx^i(s(t))}{dt}$
mielke-geometrodynamics_EQ0005 (2.1.4)	33	■ 1.000	$T(M)=\bigcup_{m\in M}T_m(M)$	$T(M)=\bigcup_{m\in M}T_m(M)$	$T(M)=\bigcup_{m\in M}T_m(M).$
mielke-geometrodynamics_EQ0008 (2.2.3)	34	■ 1.000	$\vartheta^\alpha=\mathrm{E}_{j}^\alpha(m)\,dx^j\;.$	$\vartheta^\alpha=\mathrm{E}_j^\alpha(m)dx^j.$	$\vartheta^\alpha=\mathrm{E}_j^\alpha(m)\,dx^j.$
mielke-geometrodynamics_EQ0011 (2.2.6)	34	■ 1.000	$T=T_{\alpha_1\cdots\alpha_p}^{\beta_1\cdots\beta_q}(m)\vartheta^{\alpha_1}\otimes\cdots\otimes\vartheta^{\alpha_p}\otimes e_{\beta_1}\otimes\cdots\otimes e_{\beta_q}\in C^\infty\left(T_p^q(M)\right).$	$T=T_{\alpha_1\cdots\alpha_p}^{\beta_1\cdots\beta_q}(m)\vartheta^{\alpha_1}\otimes\cdots\otimes\vartheta^{\alpha_p}\otimes e_{\beta_1}\otimes\cdots\otimes e_{\beta_q}\in C^\infty\left(T_p^q(M)\right).$	$T=T_{\alpha_1\cdots\alpha_p}^{\beta_1\cdots\beta_q}(m)\vartheta^{\alpha_1}\otimes\cdots\otimes\vartheta^{\alpha_p}\otimes e_{\beta_1}\otimes\cdots\otimes e_{\beta_q}\in C^\infty(T_p^q(M)).$
mielke-geometrodynamics_EQ0012 (2.2.7)	34	■ 1.000	$ds^2=g_{\alpha\beta}\vartheta^\alpha\otimes_s\vartheta^\beta=:\,g_{ij}dx^i\otimes_sdx^j.$	$ds^2=g_{\alpha\beta}\vartheta^\alpha\otimes_s\vartheta^\beta=:\,g_{ij}dx^i\otimes_sdx^j.$	$ds^2=g_{\alpha\beta}\vartheta^\alpha\otimes_s\vartheta^\beta=:\,g_{ij}dx^i\otimes_sdx^j.$
mielke-geometrodynamics_EQ0014 (2.2.9)	35	■ 1.000	$\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty\left(\wedge^pT^*(M)\right).$	$\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty\left(\wedge^pT^*(M)\right).$	$\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty(\wedge^pT^*(M)).$
mielke-geometrodynamics_EQ0015 (2.3.1)	36	■ 1.000	$\pi\colon F\rightarrowtail M\;.$	$\pi:F\rightarrowtail M.$	$\pi:F\rightarrowtail M.$
mielke-geometrodynamics_EQ0016 (2.3.2)	37	■ 1.000	$\sigma\colon U\rightarrowtail F\quad\text{where}\quad\pi\circ\sigma=\mathrm{id}$	$\sigma:U\rightarrowtail F\quad\text{where}\quad\pi\circ\sigma=\mathrm{id}$	$\sigma:U\rightarrowtail F\quad\text{where}\quad\pi\circ\sigma=\mathrm{id}$
mielke-geometrodynamics_EQ0017 (2.3.3)	37	■ 1.000	$\iota\!\left(m,g_1g_2\right)=\iota\!\left(m,g_1\right)\cdot g_2,\quad m\in U,g\in G\;.$	$\iota\left(m,g_1g_2\right)=\iota\left(m,g_1\right)\cdot g_2,\quad m\in U,g\in G.$	$\iota(m,g_1g_2)=\iota(m,g_1)\cdot g_2,\quad m\in U,g\in G.$
mielke-geometrodynamics_EQ0019 (2.4.1)	38	■ 1.000	$\delta_g(p,\zeta):=\left(pg,g^{-1}\zeta\right)\subset P\times F,\quad g\in G.$	$\delta_g(p,\zeta):=\left(pg,g^{-1}\zeta\right)\subset P\times F,\quad g\in G.$	$\delta_g(p,\zeta):=\left(pg,g^{-1}\zeta\right)\subset P\times F,\quad g\in G.$
mielke-geometrodynamics_EQ0022 (2.4.4)	38	■ 1.000	$\varphi^{(p)}=\left\{\varphi^{(p)A}\mid A=1,\ldots,N\right\}.$	$\varphi^{(p)}=\left\{\varphi^{(p)A}\mid A=1,\ldots,N\right\}.$	$\varphi^{(p)}=\left\{\varphi^{(p)A} A=1,\ldots,N\right\}.$
mielke-geometrodynamics_EQ0023 (2.4.5)	39	■ 1.000	$L_j:=\left[\left.\frac{\partial\rho_A^B(g(\xi))}{\partial\xi^j}\right _{g=e}\right]\in T_e(\rho(G))$	$L_j:=\left[\left.\frac{\partial\rho_A^B(g(\xi))}{\partial\xi^j}\right _{g=e}\right]\in T_e(\rho(G))$	$L_j:=\left[\left.\frac{\partial\rho_A^B(g(\xi))}{\partial\xi^j}\right _{g=e}\right]\in T_e(\rho(G))$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_39 EQ0025 (2.4.7)	39	■ 1.000	$\phi^{\{p\}}=\frac{1}{p!}\phi_{\alpha_1\ldots\alpha_p}\phi_{\alpha_1}\ldots\phi_{\alpha_p}\wedge\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty\left(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^{Ad}\right).$	$\phi^{(p)}=\frac{1}{p!}\phi_{\alpha_1\ldots\alpha_p}^jL_j\otimes\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty\left(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^{Ad}\right).$	$\phi^{(p)}=\frac{1}{p!}\phi_{\alpha_1\ldots\alpha_p}^jL_j\otimes\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty\left(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^{Ad}\right).$
mielke-geometrodynamics_39 EQ0026 (2.4.8)	39	■ 1.000	$\left[\phi^{\{p\}},\psi^{\{q\}}\right]:=\phi^{\{p\}}\wedge\psi^{\{q\}}-(-1)^{pq}\psi^{\{q\}}\wedge\phi^{\{p\}}=(-1)^{pq+1}\left[\psi^{\{q\}},\phi^{\{p\}}\right]$	$\left[\phi^{(p)},\psi^{(q)}\right]:=\phi^{(p)}\wedge\psi^{(q)}-(-1)^{pq}\psi^{(q)}\wedge\phi^{(p)}=(-1)^{pq+1}\left[\psi^{(q)},\phi^{(p)}\right]$	$\left[\phi^{(p)},\psi^{(q)}\right]:=\phi^{(p)}\wedge\psi^{(q)}-(-1)^{pq}\psi^{(q)}\wedge\phi^{(p)}=(-1)^{pq+1}\left[\psi^{(q)},\phi^{(p)}\right]$
mielke-geometrodynamics_39 EQ0027 (2.4.9)	39	■ 1.000	$\left[\phi^{\{2k\}},\phi^{\{2k\}}\right]=-\left[\phi^{\{2k\}},\phi^{\{2k\}}\right]=0.$	$\left[\phi^{(2k)},\phi^{(2k)}\right]=-\left[\phi^{(2k)},\phi^{(2k)}\right]=0.$	$\left[\phi^{(2k)},\phi^{(2k)}\right]=-\left[\phi^{(2k)},\phi^{(2k)}\right]=0.$
mielke-geometrodynamics_40 EQ0029 (2.5.2)	40	■ 1.000	$\omega(e\,g)=g^{-1}\omega(e)\,g,\quad e\in T(P).$	$\omega(eg)=g^{-1}\omega(e)g,\quad e\in T(P).$	$\omega(eg)=g^{-1}\omega(e)g,\quad e\in T(P).$
mielke-geometrodynamics_40 EQ0030 (2.5.3)	40	■ 1.000	$\omega(p\,dg)=g^{-1}dg\in T_e(G)\approx\mathfrak{g}$	$\omega(pdg)=g^{-1}dg\in T_e(G)\approx\mathfrak{g}$	$\omega(p\,dg)=g^{-1}dg\in T_e(G)\approx\mathfrak{g}$
mielke-geometrodynamics_40 EQ0031 (2.5.4)	40	■ 1.000	$D:C^\infty(V^\rho)\longrightarrow C^\infty(T_{\mathbb{C}}^*(M)\otimes V^\rho)$	$D:C^\infty(V^\rho)\longrightarrow C^\infty(T_{\mathbb{C}}^*(M)\otimes V^\rho)$	$D:C^\infty(V^\rho)\longrightarrow C^\infty(T_{\mathbb{C}}^*(M)\otimes V^\rho)$
mielke-geometrodynamics_41 EQ0032 (2.5.6)	41	■ 1.000	$Db=\omega\otimes b,\quad b\in C^\infty(V^\rho).$	$Db=\omega\otimes b,\quad b\in C^\infty(V^\rho).$	$Db=\omega\otimes b,\quad b\in C^\infty(V^\rho).$
mielke-geometrodynamics_41 EQ0033 (2.5.7)	41	■ 1.000	$D(\varphi^{\{p\}}\otimes b)=d\varphi^{\{p\}}\otimes b+(-1)^p\varphi^{\{p\}}\otimes Db.$	$D\left(\varphi^{(p)}\otimes b\right)=d\varphi^{(p)}\otimes b+(-1)^p\varphi^{(p)}\otimes Db.$	$D(\varphi^{(p)}\otimes b)=d\varphi^{(p)}\otimes b+(-1)^p\varphi^{(p)}\otimes Db.$
mielke-geometrodynamics_41 EQ0034 (2.5.8)	41	■ 1.000	$D(\varphi^{\{p\}}\otimes b)=\left\{d\varphi^p+(-1)^p\left[\varphi^{\{p\}},\omega\right]\right\}\otimes b.$	$D\left(\varphi^{(p)}\otimes b\right)=\left\{d\varphi^p+(-1)^p\left[\varphi^{(p)},\omega\right]\right\}\otimes b.$	$D(\varphi^{(p)}\otimes b)=\left\{d\varphi^p+(-1)^p[\varphi^{(p)},\omega]\right\}\otimes b.$
mielke-geometrodynamics_41 EQ0035 (2.5.9)	41	■ 1.000	$Db_A=\omega_A{}^Bb_B,\,A,B=1,\ldots,N.$	$Db_A=\omega_A{}^Bb_B,\,A,B=1,\ldots,N.$	$Db_A=\omega_A{}^Bb_B,\,A,B=1,\ldots,N.$
mielke-geometrodynamics_41 EQ0036 (2.5.10)	41	■ 1.000	$\Gamma:=\stackrel{*}{\sigma}\omega=\Gamma_\alpha{}^jL_j\otimes\vartheta^\alpha,\quad j=1,\ldots,\dim G,$	$\Gamma:=\stackrel{*}{\sigma}\omega=\Gamma_\alpha{}^jL_j\otimes\vartheta^\alpha,\quad j=1,\ldots,\dim G,$	$\Gamma:=\stackrel{*}{\sigma}\omega=\Gamma_\alpha{}^jL_j\otimes\vartheta^\alpha,\quad j=1,\ldots,\dim G,$
mielke-geometrodynamics_41 EQ0037 (2.5.11)	41	■ 1.000	$A_\alpha{}^j(m):=\Gamma_\alpha{}^j(m) _{G=U(f)}$	$A_\alpha{}^j(m):=\Gamma_\alpha{}^j(m) _{G=U(f)}$	$A_\alpha{}^j(m):=\Gamma_\alpha{}^j(m) _{G=U(f)}$
mielke-geometrodynamics_42 EQ0038 (2.6.1)	42	■ 1.000	$DDb:=\Omega\otimes b.$	$DDb:=\Omega\otimes b.$	$DDb:=\Omega\otimes b.$
mielke-geometrodynamics_42 EQ0039 (2.6.2)	42	■ 1.000	$DDb=D(\omega\otimes b)=d\omega\otimes b-\omega\wedge Db=(d\omega-\omega\wedge\omega)\otimes b$	$DDb=D(\omega\otimes b)=d\omega\otimes b-\omega\wedge Db=(d\omega-\omega\wedge\omega)\otimes b$	$DDb=D(\omega\otimes b)=d\omega\otimes b-\omega\wedge Db=(d\omega-\omega\wedge\omega)\otimes b$
mielke-geometrodynamics_42 EQ0040 (2.6.3)	42	■ 1.000	$\Omega=d\omega-\omega\wedge\omega=d\omega-\frac{1}{2}[\omega,\omega]$	$\Omega=d\omega-\omega\wedge\omega=d\omega-\frac{1}{2}[\omega,\omega]$	$\Omega=d\omega-\omega\wedge\omega=d\omega-\frac{1}{2}[\omega,\omega]$
mielke-geometrodynamics_42 EQ0043 (2.6.6)	42	■ 1.000	$F_{ij}=\partial_iA_j-\partial_jA_i$	$F_{ij}=\partial_iA_j-\partial_jA_i$	$F_{ij}=\partial_iA_j-\partial_jA_i$
mielke-geometrodynamics_42 EQ0044 (2.6.7)	42	■ 1.000	$\Delta b=b-b(\circ)=\oint (db)_{\mathrm{hor}}$	$\Delta b=b-b(\circ)=\oint (db)_{\mathrm{hor}}$	$\Delta b=b-b(\circ)=\oint (db)_{\mathrm{hor}},$
mielke-geometrodynamics_43 EQ0047 (2.6.10)	43	■ 1.000	$K=\lim_{U\rightarrow 0}\left(\int_U\Omega\otimes b\right)/\left(\int_U1\otimes b\right)$	$K=\lim_{U\rightarrow 0}\left(\int_U\Omega\otimes b\right)/\left(\int_U1\otimes b\right)$	$K=\lim_{U\rightarrow 0}\left(\int_U\Omega\otimes b\right)/\left(\int_U1\otimes b\right)$
mielke-geometrodynamics_43 EQ0049 (2.6.12)	43	■ 1.000	$D\Omega\equiv 0$	$D\Omega\equiv 0$	$D\Omega\equiv 0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_52 EQ0078 (3.1.3)	52	■ 1.000	$\delta \varphi = 0$ at the boundary ∂M of M	$\delta \varphi = 0$ at the boundary ∂M of M	$\delta \varphi = 0$ at the boundary ∂M of M
mielke-geometrodynamics_52 EQ0079 (3.1.4)	52	■ 1.000	$\delta(d\varphi) := d(\varphi + \delta\varphi) - d\varphi = d\delta\varphi$	$\delta(d\varphi) := d(\varphi + \delta\varphi) - d\varphi = d\delta\varphi$	$\delta(d\varphi) := d(\varphi + \delta\varphi) - d\varphi = d\delta\varphi$
mielke-geometrodynamics_52 EQ0080 (3.1.5)	52	■ 1.000	$\pi := \frac{\partial L}{\partial(\partial_\alpha \varphi)} \vartheta^\alpha$	$\pi := \frac{\partial L}{\partial(\partial_\alpha \varphi)} \vartheta^\alpha$	$\pi := \frac{\partial L}{\partial(\partial_\alpha \varphi)} \vartheta^\alpha$
mielke-geometrodynamics_53 EQ0082 (3.1.7)	53	■ 1.000	$\int_M d(\delta\varphi \wedge \pi) = \int_{\partial M} \delta\varphi \wedge \pi$	$\int_M d(\delta\varphi \wedge \pi) = \int_{\partial M} \delta\varphi \wedge \pi$	$\int_M d(\delta\varphi \wedge \pi) = \int_{\partial M} \delta\varphi \wedge \pi.$
mielke-geometrodynamics_53 EQ0083 (3.1.9)	53	■ 1.000	$\lambda^\varphi := (-1)^s L_\varphi + {}^*d\pi = 0,$	$\lambda^\varphi := (-1)^s L_\varphi + {}^*d\pi = 0,$	$\lambda^\varphi := (-1)^s L_\varphi + {}^*d\pi = 0,$
mielke-geometrodynamics_53 EQ0084 (3.1.10)	53	■ 1.000	$\frac{\partial L}{\partial \varphi} - \partial_\alpha \frac{\partial L}{\partial(\partial_\alpha \varphi)} = 0.$	$\frac{\partial L}{\partial \varphi} - \partial_\alpha \frac{\partial L}{\partial(\partial_\alpha \varphi)} = 0.$	$\frac{\partial L}{\partial \varphi} - \partial_\alpha \frac{\partial L}{\partial(\partial_\alpha \varphi)} = 0.$
mielke-geometrodynamics_53 EQ0085 (3.1.11)	53	■ 1.000	${}^*H := L - d\varphi \wedge \pi$	${}^*H := L - d\varphi \wedge \pi$	${}^*H := L - d\varphi \wedge \pi$
mielke-geometrodynamics_53 EQ0086 (3.1.12)	53	■ 1.000	$d{}^*H = dL + d\varphi \wedge {}^*L_\varphi + d\varphi \wedge d\pi - dd\varphi \wedge \pi$	$d{}^*H = dL + d\varphi \wedge {}^*L_\varphi + d\varphi \wedge d\pi - dd\varphi \wedge \pi$	$d{}^*H = dL + d\varphi \wedge {}^*L_\varphi + d\varphi \wedge d\pi - d\,d\varphi \wedge \pi$
mielke-geometrodynamics_53 EQ0087 (3.1.13)	53	■ 1.000	$d{}^*H \simeq 0$	$d{}^*H \simeq 0$	$d{}^*H \simeq 0$
mielke-geometrodynamics_54 EQ0088 (3.1.14)	54	■ 1.000	$\beta \wedge {}^*D\psi = m\psi\eta.$	$\beta \wedge {}^*D\psi = m\psi\eta.$	$\beta \wedge {}^*D\psi = m\psi\eta.$
mielke-geometrodynamics_54 EQ0089 (3.1.15)	54	■ 1.000	$\square\varphi + m^2\varphi = 0$	$\square\varphi + m^2\varphi = 0$	$\square\varphi + m^2\varphi = 0$
mielke-geometrodynamics_54 EQ0090 (3.1.16)	54	■ 1.000	$\overline{L}_{\text{mat}} = \overline{L}_{\text{mat}}(\bar{\psi}\psi, \bar{\psi}\beta \cdot d\psi)\eta = \bar{\psi}\beta \wedge {}^*d\psi - m\bar{\psi}\psi\eta.$	$\overline{L}_{\text{mat}} = \overline{L}_{\text{mat}}(\bar{\psi}\psi, \bar{\psi}\beta \cdot d\psi)\eta = \bar{\psi}\beta \wedge {}^*d\psi - m\bar{\psi}\psi\eta.$	$\overline{L}_{\text{mat}} = \overline{L}_{\text{mat}}(\bar{\psi}\psi, \bar{\psi}\beta \cdot d\psi)\eta = \bar{\psi}\beta \wedge {}^*d\psi - m\bar{\psi}\psi\eta.$
mielke-geometrodynamics_55 EQ0092 (3.2.2)	55	■ 1.000	$L_{\text{mat}} = L(\bar{\varphi}\varphi, \bar{\varphi}\beta \cdot (d + \omega')\varphi)\eta = L(\bar{\varphi}\varphi, \bar{\varphi}\beta \cdot D'\varphi)\eta$	$L_{\text{mat}} = L(\bar{\varphi}\varphi, \bar{\varphi}\beta \cdot (d + \omega')\varphi)\eta = L(\bar{\varphi}\varphi, \bar{\varphi}\beta \cdot D'\varphi)\eta$	$L_{\text{mat}} = L(\bar{\varphi}\varphi, \bar{\varphi}\beta \cdot (d + \omega')\varphi)\eta = L(\bar{\varphi}\varphi, \bar{\varphi}\beta \cdot D'\varphi)\eta$
mielke-geometrodynamics_55 EQ0094 (3.2.4)	55	■ 1.000	$D\left({}^G b\right) = {}^G\omega \otimes {}^G b.$	$D\left({}^G b\right) = {}^G\omega \otimes {}^G b.$	$D({}^G b) = {}^G\omega \otimes {}^G b.$
mielke-geometrodynamics_55 EQ0095 (3.2.5)	55	■ 1.000	$D\left({}^G b\right) = dG \otimes b + G\omega \otimes b = \left\{dGG^{-1} + G\omega G^{-1}\right\} \otimes {}^G b.$	$D\left({}^G b\right) = dG \otimes b + G\omega \otimes b = \left\{dGG^{-1} + G\omega G^{-1}\right\} \otimes {}^G b.$	$D({}^G b) = dG \otimes b + G\omega \otimes b = \left\{dGG^{-1} + G\omega G^{-1}\right\} \otimes {}^G b.$
mielke-geometrodynamics_55 EQ0096 (3.2.6)	55	■ 1.000	$\omega \rightarrow {}^G\omega = G\omega G^{-1} + dGG^{-1}.$	$\omega \rightarrow {}^G\omega = G\omega G^{-1} + dGG^{-1}.$	$\omega \rightarrow {}^G\omega = G\omega G^{-1} + dG\,G^{-1}.$
mielke-geometrodynamics_57 EQ0102 (3.3.3)	57	■ 1.000	$\theta = -\frac{e}{\hbar} \int \mathbf{A} \cdot d\mathbf{l}$	$\theta = -\frac{e}{\hbar} \int \mathbf{A} \cdot d\mathbf{l}$	$\theta = -\frac{e}{\hbar} \int \mathbf{A} \cdot d\mathbf{l}.$
mielke-geometrodynamics_57 EQ0103 (3.3.4)	57	■ 1.000	$L = L(A, dA) + L_{\text{mat}}(A, dA, \Psi, d\Psi).$	$L = L(A, dA) + L_{\text{mat}}(A, dA, \Psi, d\Psi).$	$L = L(A, dA) + L_{\text{mat}}(A, dA, \Psi, d\Psi).$
mielke-geometrodynamics_57 EQ0104 (3.3.5)	57	■ 1.000	$\frac{\delta L}{\delta A} := \frac{\partial L}{\partial A} + d\frac{\partial L}{\partial dA} = 0,$	$\frac{\delta L}{\delta A} := \frac{\partial L}{\partial A} + d\frac{\partial L}{\partial dA} = 0,$	$\frac{\delta L}{\delta A} := \frac{\partial L}{\partial A} + d\frac{\partial L}{\partial dA} = 0,$
mielke-geometrodynamics_58 EQ0106 (3.3.7)	58	■ 1.000	$dF \equiv 0, \quad dH = j$	$dF \equiv 0, \quad dH = j$	$dF \equiv 0, \quad dH = j.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_58 EQ0107 (3.3.8)	58	■ 1.000	$d\,j\,\simeq\,0\;.$	$dj \simeq 0.$	$dj \simeq 0\,.$
mielke-geometrodynamics_58 EQ0111 (3.3.12)	58	■ 1.000	$H\!:=\!\frac{H}{\wedge dt+\frac{D}{\wedge}}\frac{1}{2}\,H_{ij}\,dx^i\wedge dx^j$	$H := -\mathfrak{H} \wedge dt + \mathfrak{D} = \frac{1}{2}H_{ij}dx^i \wedge dx^j$	$H := -\mathfrak{H} \wedge dt + \mathfrak{D} = \frac{1}{2}H_{ij}dx^i \wedge dx^j\,,$
mielke-geometrodynamics_59 EQ0114 (3.3.15)	59	■ 1.000	$H_{mn}=\frac{\sqrt{ g }}{2}\,\varepsilon_{mnkl}\,g^{ki}\,g^{lj}\,F_{ij}\;.$	$H_{mn} = \frac{\sqrt{ g }}{2}\varepsilon_{mnkl}g^{ki}g^{lj}F_{ij}.$	$H_{mn} = \frac{\sqrt{ g }}{2}\varepsilon_{mnkl}g^{ki}g^{lj}F_{ij}\,.$
mielke-geometrodynamics_60 EQ0121 (3.3.22)	60	■ 1.000	$L_{\mathrm{BF}}\!=\!-B\,\wedge F\!=\!-B\,\wedge dA\;.$	$L_{\mathrm{BF}} = -B \wedge F = -B \wedge dA.$	$L_{\mathrm{BF}} = -B \wedge F = -B \wedge dA\;.$
mielke-geometrodynamics_60 EQ0122 (3.3.23)	60	■ 1.000	$\begin{aligned} \tilde{L}_{\mathrm{BF}} & \& := -B \wedge dA + \frac{1}{2} B \wedge B \\ & \cong -B \wedge dA + dC \end{aligned}$	$\begin{aligned} \tilde{L}_{\mathrm{BF}} & := -B \wedge dA + \frac{1}{2} B \wedge B \\ & \cong -B \wedge dA + dC \end{aligned}$	$\begin{aligned} \tilde{L}_{\mathrm{BF}} & := -B \wedge dA + \frac{1}{2} B \wedge B \\ & \cong -B \wedge dA + dC\;.\end{aligned}$
mielke-geometrodynamics_60 EQ0123 (3.3.24)	60	■ 1.000	$B\,\congd A,\,\,\,\quad d\,B\,\congd d\,F\,\equiv\,0$	$B \cong dA,\quad dB \cong dF \equiv 0$	$B \cong dA\,,\quad dB \cong dF \equiv 0\,,$
mielke-geometrodynamics_60 EQ0124 (3.3.25)	60	■ 1.000	$C\!=\!\frac{1}{2}\,A\,\wedge F,\,\,\,\quad d\,C\!=\!\frac{1}{2}\,F\,\wedge F,$	$C = \frac{1}{2}A \wedge F,\quad dC = \frac{1}{2}F \wedge F,$	$C = \frac{1}{2}A \wedge F,\quad dC = \frac{1}{2}F \wedge F\,,$
mielke-geometrodynamics_61 EQ0125 (3.3.26)	61	■ 1.000	$A\rightarrow A^{\prime}\!=\!A\!+\!d\,\theta(x)\;.$	$A \rightarrow A' = A + d\theta(x).$	$A \rightarrow A' = A + d\theta(x).$
mielke-geometrodynamics_61 EQ0126 (3.3.27)	61	■ 1.000	$A\rightarrow\widetilde{A}\!=\!A\!+\!\psi,\,\,\,\quad B\rightarrow\widetilde{B}\!=\!B\!+\!d\psi,$	$A \rightarrow \widetilde{A} = A + \psi,\quad B \rightarrow \widetilde{B} = B + d\psi,$	$A \rightarrow \widetilde{A} = A + \psi,\quad B \rightarrow \widetilde{B} = B + d\psi\,,$
mielke-geometrodynamics_61 EQ0127 (3.3.28)	61	■ 1.000	$L_{\mathrm{Max}}\!=\!-B\,\wedge dA\!+\!\frac{1}{2}\,B\,\wedge\{ \}^{\ast} B\!+\!L_{\mathrm{matter}},$	$L_{\mathrm{Max}} = -B \wedge dA + \frac{1}{2}B \wedge {}^*B + L_{\mathrm{matter}},$	$L_{\mathrm{Max}} = -B \wedge dA + \frac{1}{2}B \wedge {}^*B + L_{\mathrm{matter}},$
mielke-geometrodynamics_61 EQ0128 (3.3.29)	61	■ 1.000	$-d\,B\!=\!d^{\ast}\,F\,\congd j,$	$-dB = d^*F \cong j,$	$-dB = d^*F \cong j,$
mielke-geometrodynamics_63 EQ0132 (3.4.3)	63	■ 1.000	$\alpha_g\!:=\!g^2\,/\,\hbar\,c$	$\alpha_g := g^2/\hbar c$	$\alpha_g := g^2/\hbar c$
mielke-geometrodynamics_63 EQ0134 (3.4.5)	63	■ 1.000	$\delta L_{\omega}\!:=\,\delta\Omega\,\wedge\,\Pi^{\Omega}\;.$	$\delta L_{\omega} =: \delta\Omega \wedge \Pi^{\Omega}.$	$\delta L_{\omega} =: \delta\Omega \wedge \Pi^{\Omega}.$
mielke-geometrodynamics_63 EQ0138 (3.4.9)	63	■ 1.000	$L\!=\!L_{\mathrm{mat}}\!+\!L_{\omega},$	$L = L_{\mathrm{mat}} + L_{\omega},$	$L = L_{\mathrm{mat}} + L_{\omega},$
mielke-geometrodynamics_63 EQ0139 (3.4.10)	63	■ 1.000	$\delta L_{\mathrm{mat}}\!:=\,\delta\omega\,\wedge\,\tau\;.$	$\delta L_{\mathrm{mat}} =: \delta\omega \wedge \tau.$	$\delta L_{\mathrm{mat}} =: \delta\omega \wedge \tau.$
mielke-geometrodynamics_64 EQ0140 (3.4.11)	64	■ 1.000	$D\,\Pi^{\Omega}\!=\!\tau$	$D\Pi^{\Omega} = \tau$	$\underline{D\Pi^{\Omega} = \tau}.$
mielke-geometrodynamics_64 EQ0141 (3.4.12)	64	■ 1.000	$D\,\Omega\,\equiv\,0\;.$	$D\Omega \equiv 0.$	$D\Omega \equiv 0.$
mielke-geometrodynamics_64 EQ0145 (3.4.16)	64	■ 1.000	$d\,\Pi^{\Omega}\!+\![\omega,\Pi^{\Omega}]\!=\!\tau\;.$	$d\Pi^{\Omega} + [\omega, \Pi^{\Omega}] = \tau.$	$d\,\Pi^{\Omega} + [\omega, \Pi^{\Omega}] = \tau.$
mielke-geometrodynamics_65 EQ0147 (3.4.18)	65	■ 1.000	$Q_e\!=\!\frac{1}{2}\,Y\!+\!I_3\;.$	$Q_e = \frac{1}{2}Y + I_3.$	$Q_e = \frac{1}{2}Y + I_3.$
mielke-geometrodynamics_65 EQ0148 (3.4.19)	65	■ 1.000	$L_{\mathrm{YM}}\!=\!-\frac{1}{\alpha_g}\,\mathrm{Tr}\left(\Omega\,\wedge^{\ast}\,\Omega\right)\;.$	$L_{\mathrm{YM}} = -\frac{1}{\alpha_g}\mathrm{Tr}\left(\Omega \wedge {}^*\Omega\right).$	$L_{\mathrm{YM}} = -\frac{1}{\alpha_g}\mathrm{Tr}(\Omega \wedge {}^*\Omega).$
mielke-geometrodynamics_65 EQ0150 (3.4.21)	65	■ 1.000	$\Pi^{\Omega}\!=\!-\frac{2}{\alpha_g}\,\ast\Omega$	$\Pi^{\Omega} = -\frac{2}{\alpha_g}{}^*\Omega$	$\Pi^{\Omega} = -\frac{2}{\alpha_g}{}^*\Omega.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0153 (3.5.1)	66	■ 1.000	$\backslash\mathrm{Pi}^{\backslash\mathrm{Omega}}=\backslash\mathrm{zeta} \ \backslash\mathrm{Omega} \ .$	$\Pi^\Omega = \zeta \Omega.$	$\varPi^\varOmega = \zeta \varOmega.$
mielke-geometrodynamics_EQ0156 (3.5.4)	67	■ 1.000	$\{ \ }^{\{ * \} } \backslash\mathrm{sigma}^{\backslash\mathrm{mu} \ \backslash\mathrm{nu}}=\backslash\mathrm{sigma}^{\backslash\mathrm{mu} \ \backslash\mathrm{nu}} , \ \backslash\mathrm{quad} \{ \ }^{\{ * \} } \backslash\mathrm{bar} \{ \backslash\mathrm{sigma} \} ^{\backslash\mathrm{mu} \ \backslash\mathrm{nu}}=-\backslash\mathrm{bar} \{ \backslash\mathrm{sigma} \} ^{\backslash\mathrm{mu} \ \backslash\mathrm{nu}} ,$	${}^*\sigma^{\mu\nu}=\sigma^{\mu\nu}, \quad {}^*\bar{\sigma}^{\mu\nu}=-\bar{\sigma}^{\mu\nu},$	${}^*\sigma^{\mu\nu}=\sigma^{\mu\nu}, \quad {}^*\bar{\sigma}^{\mu\nu}=-\bar{\sigma}^{\mu\nu},$
mielke-geometrodynamics_EQ0158 (3.5.6)	67	■ 1.000	$A_{\backslash\mathrm{mu}}=\mathrm{i} \ \backslash\mathrm{sigma}_{\backslash\mathrm{mu} \ i} \ \backslash\mathrm{partial}^{\backslash\mathrm{i}} \ \backslash\mathrm{ln} \ h \ .$	$A_\mu=i\sigma_{\mu i}\partial^i\ln h.$	$A_\mu=i\sigma_{\mu i}\partial^i\ln h.$
mielke-geometrodynamics_EQ0159 (3.5.7)	67	■ 1.000	$\backslash\mathrm{square} \ h\text{-}h^{\backslash\mathrm{-}1} \ \backslash\mathrm{partial}_{\backslash\mathrm{i}} \ h \ \backslash\mathrm{partial}^{\backslash\mathrm{i}} \ h\text{=}\mathrm{h}^{\backslash\mathrm{h}^{\backslash\mathrm{3}}}$	$\square h-h^{-1}\partial_i h\partial^i h=h^3$	$\square h-h^{-1}\partial_i h\partial^i h=h^3$
mielke-geometrodynamics_EQ0160 (3.5.8)	67	■ 1.000	$\mathrm{h}=\sum_{\mathrm{i}=1}^{\mathrm{k}+1} \ \backslash\mathrm{frac} \{ \mathrm{l}_{\backslash\mathrm{(i)}}^2 \} {\backslash\mathrm{left} (\mathrm{x}\text{-}\mathrm{x}_{\backslash\mathrm{(i)}} \ \backslash\mathrm{right})^{\backslash\mathrm{2}} \} ,$	$h=\sum_{i=1}^{k+1}\frac{l_{(i)}^2}{\left(x-x_{(i)}\right)^2},$	$h=\sum_{i=1}^{k+1}\frac{l_{(i)}^2}{(x-x_{(i)})^2},$
mielke-geometrodynamics_EQ0161 (3.5.9)	67	■ 1.000	$\mathrm{p}_{\backslash\mathrm{1}}(\mathrm{M})\text{=}\mathrm{k} \ .$	$p_1(M)=k.$	$p_1(M)=k.$
mielke-geometrodynamics_EQ0162 (3.5.10)	68	■ 1.000	$\backslash\mathrm{rangle}\backslash\mathrm{infty} \ \backslash\mathrm{mid}\text{-}\backslash\mathrm{infty}\backslash\mathrm{rangle}=\mathrm{Z} \ \backslash\mathrm{int}_{\backslash\mathrm{mathscr}\{\mathrm{M}\}_{\backslash\mathrm{omega}} \} \ \mathrm{d} \ \backslash\mathrm{omega} \ \backslash\mathrm{mu} [\backslash\mathrm{omega}] \ \backslash\mathrm{exp} \ \mathrm{i} \ \backslash\mathrm{int}_{\backslash\mathrm{M}} \backslash\mathrm{left} (\mathrm{L}_{\backslash\mathrm{omega}} \text{+}\mathrm{L}_{\backslash\mathrm{infty}} \backslash\mathrm{right})$	$\langle\infty\mid-\infty\rangle=Z\int_{\mathscr{M}_\omega}d\omega\mu[\omega]\exp i\int_M(L_\omega+L_\infty)$	$\langle\infty\mid-\infty\rangle=Z\int_{\mathscr{M}_\omega}d\omega\mu[\omega]\exp i\int_M(L_\omega+L_\infty).$
mielke-geometrodynamics_EQ0163 (3.5.11)	68	■ 1.000	$\mathrm{L}_{\backslash\mathrm{infty}}\text{=}\left(\backslash\mathrm{stackrel{\mathrm{+}}{\mathrm{-}}}\right)\backslash\mathrm{frac} \{ 1 \} {\backslash\mathrm{alpha}_{\backslash\mathrm{g}}} \ \backslash\mathrm{operatorname}\{\mathrm{Tr}\} (\backslash\mathrm{Omega} \ \backslash\mathrm{wedge} \ \backslash\mathrm{Omega})$	$L_\infty=\left(\overset{+}{-}\right)\frac{1}{\alpha_g}\mathrm{Tr}(\Omega\wedge\Omega)$	$L_\infty=\left(\overset{+}{-}\right)\frac{1}{\alpha_g}\mathrm{Tr}(\varOmega\wedge\varOmega)$
mielke-geometrodynamics_EQ0167 (3.6.2)	70	■ 1.000	$\backslash\mathrm{mathscr}\{\mathrm{P}\}_{\backslash\mathrm{pm}}\text{:}=\backslash\mathrm{bar} \{ \backslash\mathrm{psi} \} \ \backslash\mathrm{sigma}_{\backslash\mathrm{pm}} \ \backslash\mathrm{psi} \ / \ 2 \ \mathrm{m} ,$	$\mathscr{P}_\pm:=\bar{\psi}\sigma_\pm\psi/2m,$	$\mathscr{P}_\pm:=\bar{\psi}\sigma_\pm\psi/2m,$
mielke-geometrodynamics_EQ0168 (3.6.3)	70	■ 1.000	$\mathrm{D}_{\backslash\mathrm{pm}} \{ \ }^{\{ * \} } \mathrm{D}_{\backslash\mathrm{pm}} \ \backslash\mathrm{psi} \ \backslash\mathrm{mp} \ \mathrm{i} \backslash\mathrm{left} (\mathrm{F}^{\backslash\mathrm{pm}} \text{-}\backslash\mathrm{frac} \{ 1 \} {2} \ \backslash\mathrm{bar} \{ \backslash\mathrm{psi} \} \ \backslash\mathrm{sigma}_{\backslash\mathrm{pm}} \ \backslash\mathrm{psi} \backslash\mathrm{right}) \ \backslash\mathrm{wedge} \ \backslash\mathrm{sigma}_{\backslash\mathrm{pm}} \ \backslash\mathrm{psi} \text{=}\mathrm{0}$	$D_\pm{}^*D_\pm\psi\mp i\left(F^\pm-\frac{1}{2}\bar{\psi}\sigma_\pm\psi\right)\wedge\sigma_\pm\psi=0$	$D_\pm{}^*D_\pm\psi\mp i\left(F^\pm-\frac{1}{2}\bar{\psi}\sigma_\pm\psi\right)\wedge\sigma_\pm\psi=0$
mielke-geometrodynamics_EQ0170 (3.6.5)	70	■ 1.000	$\mathrm{D}^{\{ * \} } \mathrm{D} \ \backslash\mathrm{psi} \text{=}\backslash\mathrm{left} (\backslash\mathrm{nabla}^{\{ * \} } \ \backslash\mathrm{nabla} \text{+}\backslash\mathrm{frac} \{ 1 \} {4} \ \mathrm{R} \text{+}\backslash\mathrm{frac} \{ 1 \} {2} \ \mathrm{F}^{\{ + \} } \backslash\mathrm{right}) \ \backslash\mathrm{psi} \ .$	$D^*D\psi=\left(\nabla^*\nabla+\frac{1}{4}R+\frac{1}{2}F^+\right)\psi.$	$D^*D\psi=\left(\nabla^*\nabla+\frac{1}{4}R+\frac{1}{2}F^+\right)\psi.$
mielke-geometrodynamics_EQ0172 (3.6.7)	70	■ 1.000	$\backslash\mathrm{Omega}_{\backslash\mathrm{pm}}^{\backslash\mathrm{g}}\text{=}\backslash\mathrm{frac} \{ 1 \} {2 \ \mathrm{ell}^{\backslash\mathrm{2}} \} \ \backslash\mathrm{sigma}_{\backslash\mathrm{pm}} ,$	$\Omega_\pm^g=\frac{1}{2\ell^2}\sigma_\pm,$	$\varOmega_\pm^g=\frac{1}{2\ell^2}\sigma_\pm,$
mielke-geometrodynamics_EQ0176 (3.7.5)	72	■ 1.000	$\backslash\mathrm{begin}\{\mathrm{aligned}\} \ \& \ \mathrm{D}^{\{ * \} } \ \backslash\mathrm{Theta}^{\backslash\mathrm{(1)}}\text{-}\backslash\mathrm{frac} \{ \backslash\mathrm{lambda} \} {2} \ \backslash\mathrm{left} (\backslash\mathrm{frac} \{ \backslash\mathrm{mu}^{\backslash\mathrm{2}} \} {\backslash\mathrm{lambda}} \text{-}\backslash\mathrm{bar} \{ \backslash\mathrm{varphi} \} \ \backslash\mathrm{cdot} \ \backslash\mathrm{varphi} \backslash\mathrm{right}) \ \backslash\mathrm{varphi} \ \backslash\mathrm{eta} \text{=}\mathrm{0} , \ \backslash\backslash \ \& \ \mathrm{D} \ \backslash\mathrm{Pi}^{\backslash\mathrm{Omega}}\text{+}\backslash\mathrm{left} [\ \backslash\mathrm{phi}^{\backslash\mathrm{(0)}} \} , \{ \ }^{\{ * \} } \ \backslash\mathrm{Theta}^{\backslash\mathrm{(1)}} \} \backslash\mathrm{right}] \text{=}\mathrm{0} \ . \ \backslash\mathrm{end}\{\mathrm{aligned}\}$	$D^*\Theta^{(1)}-\frac{\lambda}{2}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)\varphi\eta=0,$ $D\Pi^\Omega+\left[\phi^{(0)},{}^*\Theta^{(1)}\right]=0.$	$D^*\Theta^{(1)}-\frac{\lambda}{2}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)\varphi\eta=0,$ $D\Pi^\varOmega+\left[\phi^{(0)},{}^*\Theta^{(1)}\right]=0.$
mielke-geometrodynamics_EQ0177 (3.7.7)	72	■ 1.000	$\backslash\mathrm{bar} \{ \backslash\mathrm{varphi} \} \ \backslash\mathrm{cdot} \ \backslash\mathrm{varphi} \text{:}=\backslash\mathrm{varphi}^{\{ * \} \ j} \ \backslash\mathrm{varphi}_{\backslash\mathrm{j}} \text{=}\backslash\mathrm{frac} \{ \backslash\mathrm{mu}^{\backslash\mathrm{2}} \} {\backslash\mathrm{lambda}} \ .$	$\bar{\varphi}\cdot\varphi:=\varphi^{*j}\varphi_j=\frac{\mu^2}{\lambda}.$	$\bar{\varphi}\cdot\varphi:=\varphi^{*j}\varphi_j=\frac{\mu^2}{\lambda}.$
mielke-geometrodynamics_EQ0179 (3.7.9)	72	■ 1.000	$\backslash\mathrm{varphi}_{\backslash\mathrm{min}} \text{:} \ \mathrm{P} \ \backslash\mathrm{rightarrow} \ \mathrm{C}^{\backslash\mathrm{infty}} \backslash\mathrm{left} (\backslash\mathrm{stackrel{\mathrm{c}}{\mathrm{circ}} \} {\mathrm{V}} (\mathrm{M} , \ \mathrm{rho} (\mathrm{G} \ / \ \mathrm{H}) , \ \mathrm{rho} (\mathrm{G}) , \ \mathrm{P}) \ \backslash\mathrm{subset} \ \mathrm{V}^{\backslash\mathrm{rho}} \ \backslash\mathrm{right}) \ .$	$\varphi_{\min}:P\rightarrow C^\infty\left(\overset{\circ}{V}\left(M,\rho(G/H),\rho(G),P\right)\subset V^\rho\right).$	$\varphi_{\min}:P\rightarrow C^\infty\left(\overset{\circ}{V}\left(M,\rho(G/H),\rho(G),P\right)\subset V^\rho\right).$
mielke-geometrodynamics_EQ0180 (3.7.10)	73	■ 1.000	$\mathrm{H}\text{:}=\backslash\mathrm{left} \{ \ h \ \backslash\mathrm{in} \ \mathrm{G} \ \backslash\mathrm{mid} \ \mathrm{rho} (h) \ \mathrm{mathrm}\{ \mathrm{v} \}_{\backslash\mathrm{0}} \text{=}\backslash\mathrm{mathrm}\{ \mathrm{v} \}_{\backslash\mathrm{0}} \} \backslash\mathrm{right} \backslash ,$	$H:=\{h\in G\mid\rho(h)\mathfrak{v}_0=\mathfrak{v}_0\},$	$H:=\{h\in G \rho(h)\mathfrak{v}_0=\mathfrak{v}_0\},$
mielke-geometrodynamics_EQ0181 (3.7.11)	73	■ 1.000	$\mathrm{Q}\text{=}\backslash\mathrm{left} \{ \{ p \ \backslash\mathrm{in} \ \mathrm{P} \ \backslash\mathrm{mid} \ \backslash\mathrm{varphi} (p) \text{=}\backslash\mathrm{sigma} \backslash\mathrm{left} (\ \mathrm{mathrm}\{ \mathrm{v} \}_{\backslash\mathrm{0}} \} \backslash\mathrm{right}) \backslash\mathrm{right} \} \backslash$	$Q=\{p\in P\mid\varphi(p)=\sigma(\mathfrak{v}_0)\}$	$\mathcal{Q}=\{p\in P \varphi(p)=\sigma(\mathfrak{v}_0)\}$
mielke-geometrodynamics_EQ0182 (3.7.12)	73	■ 1.000	$\mathrm{D} \ \backslash\mathrm{phi}^{\backslash\mathrm{(0)}}\text{=}\mathrm{0} \ .$	$D\phi^{(0)}=0.$	$D\phi^{(0)}=0.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_73 EQ0183 (3.7.13)	73	■ 1.000	$\begin{aligned} \Theta^{\perp} &:= \Omega_0 = d\omega_0 - (\omega \wedge \omega)_0 = d\omega_0 - \omega_0 \wedge \omega - \omega \wedge \omega_0 \\ &= d\omega_0 - [\omega_0, \omega] = D\omega_0 \end{aligned}$	$\Theta^{\perp} := \Omega_0 = d\omega_0 - (\omega \wedge \omega)_0 = d\omega_0 - \omega_0 \wedge \omega - \omega \wedge \omega_0 \\ = d\omega_0 - [\omega_0, \omega] = D\omega_0$	$\Theta^{\perp} := \Omega_0 = d\omega_0 - (\omega \wedge \omega)_0 = d\omega_0 - \omega_0 \wedge \omega - \omega \wedge \omega_0 \\ = d\omega_0 - [\omega_0, \omega] = D\omega_0$
mielke-geometrodynamics_74 EQ0185 (3.7.15)	74	■ 1.000	$d_{\{0\}} \Theta^{\perp} = 0.$	$d_0^* \Omega_0 = 0.$	$d_0^* \Omega_0 = 0.$
mielke-geometrodynamics_74 EQ0186 (3.7.16)	74	■ 1.000	$\left(D^* \Omega \right)_0 = \vec{D}^* \Omega_0 = \vec{D}^* \Theta^{\perp} = 0$	$(D^* \Omega)_0 = \vec{D}^* \Omega_0 = \vec{D}^* \Theta^{\perp} = 0$	$(D^* \Omega)_0 = \vec{D}^* \Omega_0 = \vec{D}^* \Theta^{\perp} = 0$
mielke-geometrodynamics_74 EQ0188 (3.7.18)	74	■ 1.000	$\omega_0 = \varphi^j L_j \otimes \vartheta_0.$	$\omega_0 = \varphi^j L_j \otimes \vartheta_0.$	$\omega_0 = \varphi^j L_j \otimes \vartheta_0.$
mielke-geometrodynamics_80 EQ0190	80	■ 1.000	$\left[T_G : \psi(m) = D \left(g_m^{-1} g g_{g^{-1}m}, m_0 \right) \psi \left(g^{-1}m \right) \right]$	$T_G : \quad T_g \psi(m) = D \left(g_m^{-1} g g_{g^{-1}m}, m_0 \right) \psi \left(g^{-1}m \right).$	$T_G : \quad T_g \psi(m) = D \left(g_m^{-1} g g_{g^{-1}m}, m_0 \right) \psi \left(g^{-1}m \right).$
mielke-geometrodynamics_81 EQ0193 (4.1.3)	81	■ 1.000	$A(n, \mathbb{R}) = \mathbb{R}^n \otimes GL(n, \mathbb{R})$	$A(n, \mathbb{R}) = \mathbb{R}^n \otimes GL(n, \mathbb{R})$	$A(n, \mathbb{R}) = \mathbb{R}^n \otimes GL(n, \mathbb{R})$
mielke-geometrodynamics_81 EQ0194 (4.1.4)	81	■ 1.000	$P := \mathbb{R}^n \otimes 0(1, n-1) \subset A(n, \mathbb{R}),$	$P := \mathbb{R}^n \otimes 0(1, n-1) \subset A(n, \mathbb{R}),$	$P := \mathbb{R}^n \otimes 0(1, n-1) \subset A(n, \mathbb{R}),$
mielke-geometrodynamics_82 EQ0197 (4.1.10)	82	■ 1.000	$e_{\alpha}(m) = e_{\alpha}^i \partial_i$	$e_{\alpha}(m) = e_{\alpha}^i \partial_i$	$e_{\alpha}(m) = e_{\alpha}^i \partial_i$
mielke-geometrodynamics_82 EQ0199 (4.1.13)	82	■ 1.000	$L(M) = \bigcup_{m \in M} L_m(M)$	$L(M) = \bigcup_{m \in M} L_m(M)$	$L(M) = \bigcup_{m \in M} L_m(M)$
mielke-geometrodynamics_83 EQ0200 (4.1.14)	83	■ 1.000	$L(M) = P(M, GL(n, \mathbb{R}), \pi, \delta)$	$L(M) = P(M, GL(n, \mathbb{R}), \pi, \delta)$	$L(M) = P(M, GL(n, \mathbb{R}), \pi, \delta)$
mielke-geometrodynamics_83 EQ0202 (4.1.16)	83	■ 1.000	$\vartheta(e) = 0 \Leftrightarrow \pi(e) = 0, e \in T_p(P),$	$\vartheta(e) = 0 \Leftrightarrow \pi(e) = 0, e \in T_p(P),$	$\vartheta(e) = 0 \Leftrightarrow \pi(e) = 0, e \in T_p(P),$
mielke-geometrodynamics_83 EQ0204 (4.1.18)	83	■ 1.000	$T(M) = V(M, \mathbb{R}^n, GL(n, \mathbb{R}), L(M))$	$T(M) = V(M, \mathbb{R}^n, GL(n, \mathbb{R}), L(M))$	$T(M) = V(M, \mathbb{R}^n, GL(n, \mathbb{R}), L(M))$
mielke-geometrodynamics_83 EQ0205 (4.1.19)	83	■ 1.000	$A(M) := P(M, A(n, \mathbb{R}), \pi, \delta) = L(M) \times \mathbb{R}^n.$	$A(M) := P(M, A(n, \mathbb{R}), \pi, \delta) = L(M) \times \mathbb{R}^n.$	$A(M) := P(M, A(n, \mathbb{R}), \pi, \delta) = L(M) \times \mathbb{R}^n.$
mielke-geometrodynamics_83 EQ0206 (4.1.20)	83	■ 1.000	$\beta : A(n, \mathbb{R}) \rightarrow GL(n, \mathbb{R}) = A(n, \mathbb{R}) / \mathbb{R}^n.$	$\beta : A(n, \mathbb{R}) \rightarrow GL(n, \mathbb{R}) = A(n, \mathbb{R}) / \mathbb{R}^n.$	$\beta : A(n, \mathbb{R}) \rightarrow GL(n, \mathbb{R}) = A(n, \mathbb{R}) / \mathbb{R}^n.$
mielke-geometrodynamics_86 EQ0213 (4.2.5)	86	■ 1.000	$\mathcal{A}(n, \mathbb{R}) := C^{\infty}(A(M) \times_{\text{Ad}} A(n, \mathbb{R})).$	$\mathcal{A}(n, \mathbb{R}) := C^{\infty}(A(M) \times_{\text{Ad}} A(n, \mathbb{R})).$	$\mathcal{A}(n, \mathbb{R}) := C^{\infty}(A(M) \times_{\text{Ad}} A(n, \mathbb{R})).$
mielke-geometrodynamics_87 EQ0223 (4.2.16)	87	■ 1.000	$Q_A = \{p \in A(M) \mid \widetilde{\varphi}(p) = \sigma(0_m)\} = L(M)$	$Q_A = \{p \in A(M) \mid \widetilde{\varphi}(p) = \sigma(0_m)\} = L(M)$	$Q_A = \{p \in A(M) \mid \widetilde{\varphi}(p) = \sigma(0_m)\} = L(M)$
mielke-geometrodynamics_90 EQ0237 (4.3.2)	90	■ 1.000	$\ell(\mathcal{C}) := \int_{t_0}^{t_1} ds(t)$	$\ell(\mathcal{C}) := \int_{t_0}^{t_1} ds(t)$	$\ell(\mathcal{C}) := \int_{t_0}^{t_1} ds(t)$
mielke-geometrodynamics_90 EQ0238 (4.3.3)	90	■ 1.000	$\ell(\mathcal{C}_1 + \mathcal{C}_2) = \ell(\mathcal{C}_1) + \ell(\mathcal{C}_2)$	$\ell(\mathcal{C}_1 + \mathcal{C}_2) = \ell(\mathcal{C}_1) + \ell(\mathcal{C}_2)$	$\ell(\mathcal{C}_1 + \mathcal{C}_2) = \ell(\mathcal{C}_1) + \ell(\mathcal{C}_2)$
mielke-geometrodynamics_90 EQ0239 (4.3.4)	90	■ 1.000	$ds = F(m, \vartheta) dt,$	$ds = F(m, \vartheta) dt,$	$ds = F(m, \vartheta) dt,$
mielke-geometrodynamics_91 EQ0242 (4.3.7)	91	■ 1.000	$ds^2 = (F(m, \vartheta) dt)^2 = g_{\alpha\beta} \vartheta^{\alpha} \otimes \vartheta^{\beta} = g_{ij}(m) dx^i \otimes_s dx^j;$	$ds^2 = (F(m, \vartheta) dt)^2 = g_{\alpha\beta} \vartheta^{\alpha} \otimes \vartheta^{\beta} = g_{ij}(m) dx^i \otimes_s dx^j;$	$ds^2 = (F(m, \vartheta) dt)^2 = g_{\alpha\beta} \vartheta^{\alpha} \otimes \vartheta^{\beta} = g_{ij}(m) dx^i \otimes_s dx^j;$
mielke-geometrodynamics_92 EQ0246 (4.3.11)	92	■ 1.000	$L^g(M) := P(M, O(s, n-s), \pi, \delta).$	$L^g(M) := P(M, O(s, n-s), \pi, \delta).$	$L^g(M) := P(M, O(s, n-s), \pi, \delta).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_92 EQ0247 (4.3.12)	92	■ 1.000	$D\,e_{\alpha}^{\cdot\{\alpha\}\{\cdot\}\{i\}}=0$	$De_{\alpha}^{\cdot i}=0$	$D\,e_{\alpha}^{\cdot i}=0$
mielke-geometrodynamics_92 EQ0251 (4.3.16)	92	■ 1.000	$\Gamma_i^{\cdot\{\alpha\}\{\beta\}\{\cdot\}}=-\Gamma_i^{\cdot\{\beta\}\{\alpha\}\{\cdot\}}$	$\Gamma_i^{\cdot\alpha\beta}=-\Gamma_i^{\cdot\beta\alpha}$	$\Gamma_i^{\cdot\alpha\beta}=-\Gamma_i^{\cdot\beta\alpha}$
mielke-geometrodynamics_93 EQ0252 (4.3.17)	93	■ 1.000	$g_{\{i\}\{j\}}\stackrel{(*)}{=}o_{\{i\}\{j\}}$	$g_{ij}\stackrel{(*)}{=}o_{ij}$	$g_{ij}\stackrel{(*)}{=}o_{ij}$
mielke-geometrodynamics_93 EQ0253 (4.3.18)	93	■ 1.000	$\partial_k g_{\{i\}\{j\}}\stackrel{(*)}{=}0$.	$\partial_k g_{ij}\stackrel{(*)}{=}0$.	$\partial_k g_{ij}\stackrel{(*)}{=}0$.
mielke-geometrodynamics_93 EQ0254 (4.3.19)	93	■ 1.000	$\Omega^g=d\,\omega^g-\omega^g\wedge\omega^g$	$\Omega^g=d\omega^g-\omega^g\wedge\omega^g$	$\Omega^g=d\omega^g-\omega^g\wedge\omega^g$
mielke-geometrodynamics_94 EQ0257 (4.3.22)	94	■ 1.000	$\omega^g=\omega^{\{\}}-K$.	$\omega^g=\omega^{\{\}}-K$.	$\omega^g=\omega^{\{\}}-K$.
mielke-geometrodynamics_94 EQ0258 (4.3.23)	94	■ 1.000	$d\,\vartheta-\left[\omega^{\{\}},\vartheta\right]=0$	$d\vartheta-\left[\omega^{\{\}},\vartheta\right]=0$	$d\vartheta-\left[\omega^{\{\}},\vartheta\right]=0$
mielke-geometrodynamics_94 EQ0259 (4.3.24)	94	■ 1.000	$\Theta=[K,\vartheta]$	$\Theta=[K,\vartheta]$	$\Theta=[K,\vartheta]$
mielke-geometrodynamics_95 EQ0265 (4.3.30)	95	■ 1.000	$\left\{\kappa_{ij}\right\}=\frac{1}{2}g^{\kappa l}\left(\partial_i g_{jl}+\partial_j g_{il}-\partial_l g_{ij}\right)$	$\left\{\begin{array}{c}\kappa\\ij\end{array}\right\}=\frac{1}{2}g^{\kappa l}\left(\partial_i g_{jl}+\partial_j g_{il}-\partial_l g_{ij}\right)$	$\left\{\begin{array}{c}\kappa\\ij\end{array}\right\}=\frac{1}{2}g^{\kappa l}\left(\partial_i g_{jl}+\partial_j g_{il}-\partial_l g_{ij}\right)$
mielke-geometrodynamics_96 EQ0272 (4.4.1)	96	■ 1.000	$L_{\{g\}}=L\left(g_{\alpha\beta};\vartheta,\left(\omega^g\right),\Theta,\Omega^g\right)$	$L_g=L\left(g_{\alpha\beta};\vartheta,\left(\omega^g\right),\Theta,\Omega^g\right)$.	$L_g=L\left(g_{\alpha\beta};\vartheta,\left(\omega^g\right),\Theta,\Omega^g\right)$.
mielke-geometrodynamics_97 EQ0273 (4.4.2)	97	■ 1.000	$\delta L_g=:\delta\vartheta\wedge E+\delta\Theta\wedge\Pi^T+\delta\Omega^g\wedge\Pi^L,$	$\delta L_g=:\delta\vartheta\wedge E+\delta\Theta\wedge\Pi^T+\delta\Omega^g\wedge\Pi^L,$	$\delta L_g=:\delta\vartheta\wedge E+\delta\Theta\wedge\Pi^T+\delta\Omega^g\wedge\Pi^L,$
mielke-geometrodynamics_97 EQ0274 (4.4.7)	97	■ 1.000	$\delta\Theta=\delta\left(d\vartheta-\left[\omega^L,\vartheta\right]\right)$	$\delta\Theta=\delta\left(d\vartheta-\left[\omega^L,\vartheta\right]\right)$	$\delta\Theta=\delta\left(d\vartheta-\left[\omega^L,\vartheta\right]\right)$
mielke-geometrodynamics_97 EQ0277 (4.4.5)	97	■ 1.000	$S=\int_M\left(L_g+L_{\mathrm{mat}}\right)$	$S=\int_M\left(L_g+L_{\mathrm{mat}}\right)$	$S=\int_M\left(L_g+L_{\mathrm{mat}}\right)$
mielke-geometrodynamics_98 EQ0279 (4.4.8)	98	■ 1.000	$\delta L_{\mathrm{matter}}=:\delta\vartheta\wedge\Sigma+\delta\omega^g\wedge\tau_s+\delta_{\varphi}\wedge\lambda^{\varphi}$	$\delta L_{\mathrm{matter}}=:\delta\vartheta\wedge\Sigma+\delta\omega^g\wedge\tau_s+\delta_{\varphi}\wedge\lambda^{\varphi}$	$\delta L_{\mathrm{matter}}=:\delta\vartheta\wedge\Sigma+\delta\omega^g\wedge\tau_s+\delta_{\varphi}\wedge\lambda^{\varphi}$
mielke-geometrodynamics_98 EQ0280 (4.4.9)	98	■ 1.000	$D\,\Theta\equiv\left[\vartheta,\Omega^L\right]$	$D\Theta\equiv\left[\vartheta,\Omega^L\right]$	$D\Theta\equiv\left[\vartheta,\Omega^L\right]$
mielke-geometrodynamics_98 EQ0281 (4.4.10)	98	■ 1.000	$D\,\Omega^g\equiv 0,$	$D\Omega^g\equiv 0,$	$D\Omega^g\equiv 0,$
mielke-geometrodynamics_99 EQ0283 (4.4.12)	99	■ 1.000	$D\left[\Theta,\vartheta\right]\equiv\left[\vartheta\wedge\vartheta,\Omega^g\right]$	$D\left[\Theta,\vartheta\right]\equiv\left[\vartheta\wedge\vartheta,\Omega^g\right]$	$D\left[\Theta,\vartheta\right]\equiv\left[\vartheta\wedge\vartheta,\Omega^g\right]$
mielke-geometrodynamics_100 EQ0289 (4.4.18)	100	■ 1.000	$\left[\Omega^g,\Pi^L\right]+E_{\square}=0$	$\left[\Omega^g,\Pi^L\right]+E_{\square}=0$	$\left[\Omega^g,\Pi^L\right]+E_{\square}=0$
mielke-geometrodynamics_100 EQ0290 (4.5.1)	100	■ 1.000	$L_{\mathrm{mat}}=L\left(g_{\alpha\beta},\vartheta^{\alpha},\Psi,D\Psi\right)$.	$L_{\mathrm{mat}}=L\left(g_{\alpha\beta},\vartheta^{\alpha},\Psi,D\Psi\right)$.	$L_{\mathrm{mat}}=L\left(g_{\alpha\beta},\vartheta^{\alpha},\Psi,D\Psi\right)$.
mielke-geometrodynamics_100 EQ0291 (4.5.2)	100	■ 1.000	$\Sigma_{\alpha}:=\frac{\partial L}{\partial\vartheta^{\alpha}},\quad\sigma^{\alpha\beta}:=2\frac{\partial L}{\partial g_{\alpha\beta}},\quad\text{and}\quad\tau_{\alpha\beta}:=\frac{\partial L}{\partial\Gamma^{\alpha\beta}}=I_{\alpha\beta}\Psi\wedge\frac{\partial L}{\partial D\Psi}$	$\Sigma_{\alpha}:=\frac{\partial L}{\partial\vartheta^{\alpha}},\quad\sigma^{\alpha\beta}:=2\frac{\partial L}{\partial g_{\alpha\beta}},\quad\text{and}\quad\tau_{\alpha\beta}:=\frac{\partial L}{\partial\Gamma^{\alpha\beta}}=I_{\alpha\beta}\Psi\wedge\frac{\partial L}{\partial D\Psi}$	$\Sigma_{\alpha}:=\frac{\partial L}{\partial\vartheta^{\alpha}},\quad\sigma^{\alpha\beta}:=2\frac{\partial L}{\partial g_{\alpha\beta}},\quad\text{and}\quad\tau_{\alpha\beta}:=\frac{\partial L}{\partial\Gamma^{\alpha\beta}}=I_{\alpha\beta}\Psi\wedge\frac{\partial L}{\partial D\Psi}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometroynamics_101 EQ0297 (8.10)		■ 1.000	$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha} + \delta \Psi \wedge \frac{\partial L}{\partial \Psi} + \delta(D\Psi) \wedge \frac{\partial L}{\partial D\Psi}.$		$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha} + \delta \Psi \wedge \frac{\partial L}{\partial \Psi} + \delta(D\Psi) \wedge \frac{\partial L}{\partial D\Psi}.$
mielke-geometroynamics_102 EQ0300 (4.5.11)		■ 1.000	$\delta L = -\varepsilon_\alpha{}^\beta \left[-\sigma^{\alpha\gamma} g_{\gamma\beta} + \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\beta} + D \frac{\partial L}{\partial \Gamma_\alpha{}^\beta} + I^\alpha{}_\beta \Psi \wedge \frac{\delta L}{\delta \Psi} \right].$		$\delta L = -\varepsilon_\alpha{}^\beta \left[-\sigma^{\alpha\gamma} g_{\gamma\beta} + \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\beta} + D \frac{\partial L}{\partial \Gamma_\alpha{}^\beta} + I^\alpha{}_\beta \Psi \wedge \frac{\delta L}{\delta \Psi} \right].$
mielke-geometroynamics_102 EQ0301 (4.5.12)		■ 1.000	$D \tau_{\alpha\beta} + \vartheta_{[\alpha} \wedge \Sigma_{\beta]} = -I_{\alpha\beta} \Psi \wedge \frac{\delta L}{\delta \Psi} \simeq 0.$		$D \tau_{\alpha\beta} + \vartheta_{[\alpha} \wedge \Sigma_{\beta]} = -I_{\alpha\beta} \Psi \wedge \frac{\delta L}{\delta \Psi} \simeq 0.$
mielke-geometroynamics_102 EQ0303 (4.5.13)		■ 1.000	$\begin{aligned} & \& R_{\alpha}{}^\beta = d\Gamma_{\alpha}{}^\beta \\ & - \Gamma_{\alpha}{}^\gamma \wedge \Gamma_{\gamma}{}^\beta \end{aligned}$		$\begin{aligned} & R_{\alpha}{}^\beta = d\Gamma_{\alpha}{}^\beta \\ & - \Gamma_{\alpha}{}^\gamma \wedge \Gamma_{\gamma}{}^\beta \end{aligned}$
mielke-geometroynamics_103 EQ0307 (4.5.17)		■ 1.000	$\hat{M} := M^3 / \left[M^2 + \frac{\Lambda}{3} J^2 \right], \quad \hat{J} := J M^2 / \left[M^2 + \frac{\Lambda}{3} J^2 \right],$		$\hat{M} := M^3 / [M^2 + \frac{\Lambda}{3} J^2], \quad \hat{J} := J M^2 / [M^2 + \frac{\Lambda}{3} J^2],$
mielke-geometroynamics_110 EQ0317 (5.2.1)		■ 1.000	$L_{\mathrm{D}} = L(\gamma, \psi, D\psi) = \frac{i}{2} \{ \bar{\psi}^* \gamma \wedge D\psi + \overline{D\psi} \wedge \gamma \psi \} - m \bar{\psi} \psi \eta,$		$L_{\mathrm{D}} = L(\gamma, \psi, D\psi) = \frac{i}{2} \{ \bar{\psi}^* \gamma \wedge D\psi + \overline{D\psi} \wedge \gamma \psi \} - m \bar{\psi} \psi \eta,$
mielke-geometroynamics_111 EQ0322 (5.2.6) [re-fined; census]		■ 1.000	$dU = T^\alpha \wedge \mu_\alpha - \vartheta^\alpha \wedge D\mu_\alpha = \vartheta^\alpha \wedge [e_\beta] (T^\beta \wedge \mu_\alpha) - D\mu_\alpha$		$dU = T^\alpha \wedge \mu_\alpha - \vartheta^\alpha \wedge D\mu_\alpha = \vartheta^\alpha \wedge [e_\beta] (T^\beta \wedge \mu_\alpha) - D\mu_\alpha$
mielke-geometroynamics_111 EQ0324 (5.3.1)		■ 1.000	$L_{\mathrm{EC}} := -\frac{1}{2\kappa} R^{\alpha\beta} \wedge \eta_{\alpha\beta},$		$L_{\mathrm{EC}} := -\frac{1}{2\kappa} R^{\alpha\beta} \wedge \eta_{\alpha\beta},$
mielke-geometroynamics_112 EQ0325 (5.3.2)		■ 1.000	$G_\alpha := \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} = \kappa \Sigma_\alpha,$		$G_\alpha := \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} = \kappa \Sigma_\alpha,$
mielke-geometroynamics_112 EQ0327 (5.3.4)		■ 1.000	$D^{\{\}} G_\alpha^{\{\}} \equiv 0$		$D^{\{\}} G_\alpha^{\{\}} \equiv 0$
mielke-geometroynamics_112 EQ0331 (5.3.8)		■ 1.000	$D[\gamma, \Theta] \equiv 2i[\sigma, \Omega^g], \quad D[\gamma, \Omega^g] \equiv [\Theta, \Omega^g],$		$D[\gamma, \Theta] \equiv 2i[\sigma, \Omega^g], \quad D[\gamma, \Omega^g] \equiv [\Theta, \Omega^g],$
mielke-geometroynamics_113 EQ0332 (5.3.9)		■ 1.000	$DG \equiv i[\Theta, \gamma_5 \Omega^g]$		$DG \equiv i[\Theta, \gamma_5 \Omega^g]$
mielke-geometroynamics_113 EQ0333 (5.3.10)		■ 1.000	$\eta_{\alpha\beta\gamma} \wedge T^\gamma = 2\kappa \tau_{\alpha\beta}$		$\eta_{\alpha\beta\gamma} \wedge T^\gamma = 2\kappa \tau_{\alpha\beta}$
mielke-geometroynamics_113 EQ0334 (5.3.11)		■ 1.000	$T_{\mu\nu} := \frac{2}{\sqrt{g}} \frac{\delta L}{\delta g_{\mu\nu}}$		$T_{\mu\nu} := \frac{2}{\sqrt{g}} \frac{\delta L}{\delta g_{\mu\nu}}$
mielke-geometroynamics_114 EQ0338 (5.3.15)		■ 1.000	${}^{(3)}T_\alpha = \frac{(-1)^s}{3} * (\vartheta_\alpha \wedge \mathscr{A})$		${}^{(3)}T_\alpha = \frac{(-1)^s}{3} * (\vartheta_\alpha \wedge \mathscr{A})$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_115 EQ0342 (5.3.19)	115	■ 1.000	$\Delta\widetilde{L}:=\kappa\mathrm{left}(L_{\mathrm{EC}})-L_{\mathrm{HE}}\right)\simeq\frac{\kappa^2}{48}j_5\wedge^*j_5=4f^2j_5\wedge^*j_5,$	$\Delta\widetilde{L}:=\kappa(L_{\mathrm{EC}}-L_{\mathrm{HE}})\simeq\frac{\kappa^2}{48}j_5\wedge^*j_5=4f^2j_5\wedge^*j_5,$	$\Delta\widetilde{L}:=\kappa(L_{\mathrm{EC}}-L_{\mathrm{HE}})\simeq\frac{\kappa^2}{48}j_5\wedge^*j_5=4f^2j_5\wedge^*j_5,$
mielke-geometrodynamics_116 EQ0344 (5.4.2)	116	■ 1.000	$L_{\mathrm{HE}}:=L_{\mathrm{HE}}+\frac{\Lambda}{\kappa}\eta$	$L_{\mathrm{HE}}:=L_{\mathrm{HE}}+\frac{\Lambda}{\kappa}\eta$	$L_{\mathrm{HE}}:=L_{\mathrm{HE}}+\frac{\Lambda}{\kappa}\eta$
mielke-geometrodynamics_116 EQ0345 (5.4.3)	116	■ 1.000	$g_{\mathrm{N}}:=\kappa k^2,$ $\quad\quad\quad\lambda:=\Lambda/k^2,$	$g_{\mathrm{N}}:=\kappa k^2,$ $\quad\quad\quad\lambda:=\Lambda/k^2,$	$g_{\mathrm{N}}:=\kappa k^2,$ $\quad\quad\quad\lambda:=\Lambda/k^2,$
mielke-geometrodynamics_116 EQ0346 (5.4.4)	116	■ 1.000	$k\frac{\partial}{\partial k}g_{\mathrm{N}}=\beta_1(g_{\mathrm{N}},\lambda)=(2+d_{\mathrm{N}})g_{\mathrm{N}},\quad k\frac{\partial}{\partial k}\lambda=\beta_2(g_{\mathrm{N}},\lambda),$	$k\frac{\partial}{\partial k}g_{\mathrm{N}}=\beta_1(g_{\mathrm{N}},\lambda)=(2+d_{\mathrm{N}})g_{\mathrm{N}},\quad k\frac{\partial}{\partial k}\lambda=\beta_2(g_{\mathrm{N}},\lambda),$	$k\frac{\partial}{\partial k}g_{\mathrm{N}}=\beta_1(g_{\mathrm{N}},\lambda)=(2+d_{\mathrm{N}})g_{\mathrm{N}},\quad k\frac{\partial}{\partial k}\lambda=\beta_2(g_{\mathrm{N}},\lambda),$
mielke-geometrodynamics_117 EQ0348 (5.4.6)	117	■ 1.000	$f^2=\frac{1}{192}\kappa^2=3\times 2^{-10}\left(\frac{\mu}{\Lambda}\right)^2=2^{-8}\frac{\mu}{\rho_{\Lambda}}$	$f^2=\frac{1}{192}\kappa^2=3\times 2^{-10}\left(\frac{\mu}{\Lambda}\right)^2=2^{-8}\frac{\mu}{\rho_{\Lambda}}$	$f^2=\frac{1}{192}\kappa^2=3\times 2^{-10}\left(\frac{\mu}{\Lambda}\right)^2=2^{-8}\frac{\mu}{\rho_{\Lambda}}$
mielke-geometrodynamics_123 EQ0350 (6.2.1)	123	■ 1.000	$C_{\mathrm{TT}}:=\frac{1}{2\ell^2}\left(\vartheta^\alpha\wedge T_\alpha\right)=-\frac{(-1)^{\mathrm{sig}}}{2\ell^2}*\mathscr{A},$	$C_{\mathrm{TT}}:=\frac{1}{2\ell^2}\left(\vartheta^\alpha\wedge T_\alpha\right)=-\frac{(-1)^{\mathrm{sig}}}{2\ell^2}*\mathscr{A},$	$C_{\mathrm{TT}}:=\frac{1}{2\ell^2}\left(\vartheta^\alpha\wedge T_\alpha\right)=-\frac{(-1)^{\mathrm{sig}}}{2\ell^2}*\mathscr{A},$
mielke-geometrodynamics_124 EQ0351 (6.2.2)	124	■ 1.000	$\begin{aligned} C_{\mathrm{RR}}&\&:=-\operatorname{Tr}\left(\Gamma\wedge\Omega-\frac{1}{3}\Gamma\wedge\Gamma\wedge\Gamma\right)\\ &=-\frac{1}{2}\left(\Gamma_\alpha{}^\beta\wedge R_\beta{}^\alpha+\frac{1}{3}\Gamma_\alpha{}^\beta\wedge\Gamma_\beta{}^\gamma\wedge\Gamma_\gamma{}^\alpha\right). \end{aligned}$	$C_{\mathrm{RR}}:=-\operatorname{Tr}\left(\Gamma\wedge\Omega-\frac{1}{3}\Gamma\wedge\Gamma\wedge\Gamma\right)\\ =-\frac{1}{2}\left(\Gamma_\alpha{}^\beta\wedge R_\beta{}^\alpha+\frac{1}{3}\Gamma_\alpha{}^\beta\wedge\Gamma_\beta{}^\gamma\wedge\Gamma_\gamma{}^\alpha\right).$	$C_{\mathrm{RR}}:=-\operatorname{Tr}\left(\Gamma\wedge\Omega-\frac{1}{3}\Gamma\wedge\Gamma\wedge\Gamma\right)\\ =-\frac{1}{2}\left(\Gamma_\alpha{}^\beta\wedge R_\beta{}^\alpha+\frac{1}{3}\Gamma_\alpha{}^\beta\wedge\Gamma_\beta{}^\gamma\wedge\Gamma_\gamma{}^\alpha\right).$
mielke-geometrodynamics_124 EQ0352 (6.2.3)	124	■ 1.000	$\hat{C}=C_{\mathrm{RR}}-2C_{\mathrm{TT}}$	$\hat{C}=C_{\mathrm{RR}}-2C_{\mathrm{TT}}$	$\hat{C}=C_{\mathrm{RR}}-2C_{\mathrm{TT}}$
mielke-geometrodynamics_124 EQ0353 (6.2.4)	124	■ 1.000	$dC_{\mathrm{TT}}\equiv\frac{1}{2\ell^2}\left(T^\alpha\wedge T_\alpha+R_{\alpha\beta}\wedge\vartheta^\alpha\wedge\vartheta^\beta\right),$	$dC_{\mathrm{TT}}\equiv\frac{1}{2\ell^2}\left(T^\alpha\wedge T_\alpha+R_{\alpha\beta}\wedge\vartheta^\alpha\wedge\vartheta^\beta\right),$	$dC_{\mathrm{TT}}\equiv\frac{1}{2\ell^2}\left(T^\alpha\wedge T_\alpha+R_{\alpha\beta}\wedge\vartheta^\alpha\wedge\vartheta^\beta\right),$
mielke-geometrodynamics_124 EQ0354 (6.2.5)	124	■ 1.000	$dC_{\mathrm{RR}}=-\operatorname{Tr}(\Omega\wedge\Omega)=\frac{1}{2}R^{\alpha\beta}\wedge R_{\alpha\beta}$	$dC_{\mathrm{RR}}=-\operatorname{Tr}(\Omega\wedge\Omega)=\frac{1}{2}R^{\alpha\beta}\wedge R_{\alpha\beta}$	$dC_{\mathrm{RR}}=-\operatorname{Tr}(\Omega\wedge\Omega)=\frac{1}{2}R^{\alpha\beta}\wedge R_{\alpha\beta}$
mielke-geometrodynamics_124 EQ0355 (6.2.6)	124	■ 1.000	$DR_{\alpha\beta}\equiv 0,$	$DR_{\alpha\beta}\equiv 0,$	$DR_{\alpha\beta}\equiv 0,$
mielke-geometrodynamics_125 EQ0359 (6.2.10)	125	■ 1.000	$H_\alpha^\parallel:=-\partial L_\parallel/\partial T^\alpha=(1/\kappa)\eta_{\alpha\beta\gamma}\wedge K^{\beta\gamma}$	$H_\alpha^\parallel:=-\partial L_\parallel/\partial T^\alpha=(1/\kappa)\eta_{\alpha\beta\gamma}\wedge K^{\beta\gamma}$	$H_\alpha^\parallel:=-\partial L_\parallel/\partial T^\alpha=(1/\kappa)\eta_{\alpha\beta\gamma}\wedge K^{\beta\gamma}$
mielke-geometrodynamics_126 EQ0361 (6.2.12)	126	■ 1.000	$\tilde{L}_\parallel=L_\parallel-R^{\alpha\beta}\wedge\lambda_{\alpha\beta}.$	$\tilde{L}_\parallel=L_\parallel-R^{\alpha\beta}\wedge\lambda_{\alpha\beta}.$	$\tilde{L}_\parallel=L_\parallel-R^{\alpha\beta}\wedge\lambda_{\alpha\beta}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_126 EQ0362 (6.2.14)	126	■ 1.000	$\begin{gathered} D\,H_{\{\alpha\}}^{\{\}}-E_{\{\alpha\}}^{\{\}}=\\ \Sigma_{\{\alpha\}},\quad D\,\lambda_{\alpha\beta}+\vartheta_{[\alpha}\wedge H_{\beta]}^{\parallel}=\tau_{\alpha\beta},\\ \end{gathered}$	$DH_{\alpha}^{\parallel}-E_{\alpha}^{\parallel}=\Sigma_{\alpha},$ $D\lambda_{\alpha\beta}+\vartheta_{[\alpha}\wedge H_{\beta]}^{\parallel}=\tau_{\alpha\beta},$	$DH_{\alpha}^{\parallel}-E_{\alpha}^{\parallel}=\Sigma_{\alpha},$ $D\lambda_{\alpha\beta}+\vartheta_{[\alpha}\wedge H_{\beta]}^{\parallel}=\tau_{\alpha\beta},$
mielke-geometrodynamics_126 EQ0363 (6.2.15)	126	■ 1.000	$R^{\{\alpha\}\{\beta\}}=0\quad.$	$R^{\alpha\beta}=0.$	$R^{\alpha\beta}=0.$
mielke-geometrodynamics_126 EQ0365 (6.2.17)	126	■ 1.000	$D\left(\tau_{\alpha\beta}-\vartheta_{[\alpha}\wedge H_{\beta]}^{\parallel}\right)=0$	$D\left(\tau_{\alpha\beta}-\vartheta_{[\alpha}\wedge H_{\beta]}^{\parallel}\right)=0$	$D(\tau_{\alpha\beta}-\vartheta_{[\alpha}\wedge H_{\beta]}^{\parallel})=0$
mielke-geometrodynamics_127 EQ0369 (6.2.20)	127	■ 1.000	$\stackrel{\scriptscriptstyle{(\pm)}}{C}_{\mathrm{TT}}:=\frac{1}{2\ell^2}\left(\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{T}_{\alpha}\right),$	$\stackrel{\scriptscriptstyle{(\pm)}}{C}_{\mathrm{TT}}:=\frac{1}{2\ell^2}\left(\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{T}_{\alpha}\right),$	$\stackrel{\scriptscriptstyle{(\pm)}}{C}_{\mathrm{TT}}:=\frac{1}{2\ell^2}\left(\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{T}_{\alpha}\right),$
mielke-geometrodynamics_127 EQ0373 (6.2.24)	127	■ 1.000	$\stackrel{\scriptscriptstyle{(\pm)}}{D}\Pi_{\alpha}=\Sigma_{\alpha}\quad.$	$\stackrel{\scriptscriptstyle{(\pm)}}{D}\Pi_{\alpha}=\Sigma_{\alpha}.$	$\stackrel{\scriptscriptstyle{(\pm)}}{D}\Pi_{\alpha}=\Sigma_{\alpha}.$
mielke-geometrodynamics_128 EQ0379 (6.2.30)	128	■ 1.000	$\stackrel{\scriptscriptstyle{(\pm)}}{L}_{\mathrm{YM}}:=\mp(1/4)\mathrm{Tr}\left(\stackrel{\scriptscriptstyle{(\pm)}}{F}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{F}\right)$	$\stackrel{\scriptscriptstyle{(\pm)}}{L}_{\mathrm{YM}}=\mp(1/4)\mathrm{Tr}\left(\stackrel{\scriptscriptstyle{(\pm)}}{F}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{F}\right)$	$\stackrel{\scriptscriptstyle{(\pm)}}{L}_{\mathrm{YM}}=\mp(1/4)\mathrm{Tr}\left(\stackrel{\scriptscriptstyle{(\pm)}}{F}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{F}\right)$
mielke-geometrodynamics_129 EQ0383 (6.3.1)	129	■ 1.000	$\check{E}_{\alpha}:=E_{\alpha}+\Gamma_{\alpha}^{\beta}\wedge H_{\beta}+\Sigma_{\alpha}\cong dH_{\alpha},$ $\wedge H_{\{\beta\}}+\Sigma_{\{\alpha\}}\congd H_{\{\alpha\}},$	$\check{E}_{\alpha}:=E_{\alpha}+\Gamma_{\alpha}^{\beta}\wedge H_{\beta}+\Sigma_{\alpha}\cong dH_{\alpha},$	$\check{E}_{\alpha}:=E_{\alpha}+\Gamma_{\alpha}^{\beta}\wedge H_{\beta}+\Sigma_{\alpha}\cong dH_{\alpha},$
mielke-geometrodynamics_130 EQ0386 (6.4.1)	130	■ 1.000	$n:=\partial_t-N^A\partial/\partial x^A,\quad\quad\quad A=1,2,3,$	$n:=\partial_t-N^A\partial/\partial x^A,\quad\quad\quad A=1,2,3,$	$n:=\partial_t-N^A\partial/\partial x^A,\quad\quad\quad A=1,2,3,$
mielke-geometrodynamics_132 EQ0395 (6.4.10)	132	■ 1.000	$\Sigma_{\{\alpha\}}:=\frac{\partial L}{\partial\vartheta^{\alpha}},\quad\tau_{\alpha\beta}:=\frac{\partial L}{\partial\Gamma^{\alpha\beta}}=\rho(I_{\alpha\beta})\Psi\wedge\frac{\partial L}{\partial D\Psi}$	$\Sigma_{\alpha}:=\frac{\partial L}{\partial\vartheta^{\alpha}},\quad\tau_{\alpha\beta}:=\frac{\partial L}{\partial\Gamma^{\alpha\beta}}=\rho(I_{\alpha\beta})\Psi\wedge\frac{\partial L}{\partial D\Psi}$	$\Sigma_{\alpha}:=\frac{\partial L}{\partial\vartheta^{\alpha}},\quad\tau_{\alpha\beta}:=\frac{\partial L}{\partial\Gamma^{\alpha\beta}}=\rho(I_{\alpha\beta})\Psi\wedge\frac{\partial L}{\partial D\Psi}$
mielke-geometrodynamics_136 EQ0408 (6.4.28)	136	■ 1.000	$ds^2=h^2dr^2+f^2\left[d\psi^2+\sin^2\psi\left(d\theta^2+\sin^2\theta d\phi^2\right)\right],$	$ds^2=h^2dr^2+f^2\left[d\psi^2+\sin^2\psi\left(d\theta^2+\sin^2\theta d\phi^2\right)\right],$	$ds^2=h^2dr^2+f^2\left[d\psi^2+\sin^2\psi\left(d\theta^2+\sin^2\theta d\phi^2\right)\right],$
mielke-geometrodynamics_136 EQ0410 (6.4.30)	136	■ 1.000	$\ell^2\underline{C}_{\mathrm{TT}}=\frac{1}{2}\underline{\vartheta}^{\alpha}\wedge d\underline{\vartheta}_{\alpha}=\frac{1}{2}\underline{\vartheta}^A\wedge d\underline{\vartheta}_A=3\underline{\vartheta}^{\hat{1}}\wedge\underline{\vartheta}^{\hat{2}}\wedge\underline{\vartheta}^{\hat{3}}$	$\ell^2\underline{C}_{\mathrm{TT}}=\frac{1}{2}\underline{\vartheta}^{\alpha}\wedge d\underline{\vartheta}_{\alpha}=\frac{1}{2}\underline{\vartheta}^A\wedge d\underline{\vartheta}_A=3\underline{\vartheta}^{\hat{1}}\wedge\underline{\vartheta}^{\hat{2}}\wedge\underline{\vartheta}^{\hat{3}}$	$\ell^2\underline{C}_{\mathrm{TT}}=\frac{1}{2}\underline{\vartheta}^{\alpha}\wedge d\underline{\vartheta}_{\alpha}=\frac{1}{2}\underline{\vartheta}^A\wedge d\underline{\vartheta}_A=3\underline{\vartheta}^{\hat{1}}\wedge\underline{\vartheta}^{\hat{2}}\wedge\underline{\vartheta}^{\hat{3}}$
mielke-geometrodynamics_136 EQ0412 (6.4.32)	136	■ 1.000	$f=\frac{ar^2}{r^2+c^2}$	$f=\frac{ar^2}{r^2+c^2}$	$f=\frac{ar^2}{r^2+c^2}$
mielke-geometrodynamics_137 EQ0414 (6.5.2)	137	■ 1.000	$\Psi(\gamma)=\int_{\mathscr{C}/\mathscr{G}}\mathscr{D}A\,W(A,\gamma)\Psi(A).$	$\Psi(\gamma)=\int_{\mathscr{C}/\mathscr{G}}\mathscr{D}A\,W(A,\gamma)\Psi(A).$	$\Psi(\gamma)=\int_{\mathscr{C}/\mathscr{G}}\mathscr{D}A\,W(A,\gamma)\Psi(A).$
mielke-geometrodynamics_137 EQ0416 (6.5.4)	137	■ 1.000	$\langle 0 W(\gamma) 0\rangle e^{\Lambda G(\gamma)}V(\gamma),$	$\langle 0 W(\gamma) 0\rangle e^{\Lambda G(\gamma)}V(\gamma),$	$\langle 0 W(\gamma) 0\rangle e^{\Lambda G(\gamma)}V(\gamma),$
mielke-geometrodynamics_138 EQ0417 (6.6.1)	138	■ 1.000	$\int T=\int d\sigma=2\pi N$	$\int T=\int d\sigma=2\pi N$	$\int T=\int d\sigma=2\pi N,$
mielke-geometrodynamics_139 EQ0418 (6.6.2)	139	■ 1.000	$W^{\{N\}}:=S^{\wedge\{3\}}\,\,\#\^{\{N\}}\!\!\left(S^{\wedge\{1\}}\times S^{\wedge\{2\}}\right)\quad.$	$W^N:=S^3\#^N\left(S^1\times S^2\right).$	$W^N:=S^3\#^N(S^1\times S^2).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_140 EQ0419 (6.7.1)	140	■ 1.000	$L := L_{\mathrm{EC}} + L_{\theta} + L_{\mathrm{D}},$	$L := L_{\mathrm{EC}} + L_{\theta} + L_{\mathrm{D}},$	$L := L_{\mathrm{EC}} + L_{\theta} + L_{\mathrm{D}},$
mielke-geometrodynamics_140 EQ0420 (6.7.2)	140	■ 1.000	$L_{\theta} = \theta_{\mathrm{T}} d C_{\mathrm{TT}} + \theta_{\mathrm{L}} d C_{\mathrm{RR}}$	$L_{\theta} = \theta_{\mathrm{T}} d C_{\mathrm{TT}} + \theta_{\mathrm{L}} d C_{\mathrm{RR}}$	$L_{\theta} = \theta_{\mathrm{T}} d C_{\mathrm{TT}} + \theta_{\mathrm{L}} d C_{\mathrm{RR}}$
mielke-geometrodynamics_140 EQ0421 (6.7.3)	140	■ 1.000	$H_{\alpha} := -\frac{\partial L}{\partial T^{\alpha}} = -\frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha},$	$H_{\alpha} := -\frac{\partial L}{\partial T^{\alpha}} = -\frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha},$	$H_{\alpha} := -\frac{\partial L}{\partial T^{\alpha}} = -\frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha},$
mielke-geometrodynamics_140 EQ0422 (6.7.4)	140	■ 1.000	$H_{\alpha \beta} := -\frac{\partial L}{\partial R^{\alpha \beta}} = \frac{1}{2\kappa} \eta_{\alpha \beta} - \frac{\theta_{\mathrm{T}}}{2\ell^2} \vartheta_{\alpha} \wedge \vartheta_{\beta} - \theta_{\mathrm{L}} R_{\alpha \beta}$	$H_{\alpha \beta} := -\frac{\partial L}{\partial R^{\alpha \beta}} = \frac{1}{2\kappa} \eta_{\alpha \beta} - \frac{\theta_{\mathrm{T}}}{2\ell^2} \vartheta_{\alpha} \wedge \vartheta_{\beta} - \theta_{\mathrm{L}} R_{\alpha \beta}$	$H_{\alpha \beta} := -\frac{\partial L}{\partial R^{\alpha \beta}} = \frac{1}{2\kappa} \eta_{\alpha \beta} - \frac{\theta_{\mathrm{T}}}{2\ell^2} \vartheta_{\alpha} \wedge \vartheta_{\beta} - \theta_{\mathrm{L}} R_{\alpha \beta}$
mielke-geometrodynamics_140 EQ0423 (6.7.5)	140	■ 1.000	$D \theta_{\mathrm{T}} \wedge T_{\alpha} + \theta_{\mathrm{T}} D T_{\alpha} + \ell^2 E_{\alpha} = -\ell^2 \Sigma_{\alpha}.$	$D \theta_{\mathrm{T}} \wedge T_{\alpha} + \theta_{\mathrm{T}} D T_{\alpha} + \ell^2 E_{\alpha} = -\ell^2 \Sigma_{\alpha}.$	$D \theta_{\mathrm{T}} \wedge T_{\alpha} + \theta_{\mathrm{T}} D T_{\alpha} + \ell^2 E_{\alpha} = -\ell^2 \Sigma_{\alpha}.$
mielke-geometrodynamics_140 EQ0425 (6.7.7)	140	■ 1.000	$C_{\mathrm{TT}} \cong \frac{\kappa}{4\ell^2} j_5$	$C_{\mathrm{TT}} \cong \frac{\kappa}{4\ell^2} j_5$	$C_{\mathrm{TT}} \cong \frac{\kappa}{4\ell^2} j_5$
mielke-geometrodynamics_141 EQ0427 (6.7.9)	141	■ 1.000	$T^{\gamma} \wedge \eta_{\gamma \beta} + \kappa D \theta_{\mathrm{L}} \wedge D T_{\beta} = 0.$	$T^{\gamma} \wedge \eta_{\gamma \beta} + \kappa D \theta_{\mathrm{L}} \wedge D T_{\beta} = 0.$	$T^{\gamma} \wedge \eta_{\gamma \beta} + \kappa D \theta_{\mathrm{L}} \wedge D T_{\beta} = 0.$
mielke-geometrodynamics_141 EQ0429 (6.7.11)	141	■ 1.000	$\kappa d \theta_{\mathrm{L}} \wedge T_{\beta} = -\eta_{\beta}$	$\kappa d \theta_{\mathrm{L}} \wedge T_{\beta} = -\eta_{\beta}$	$\kappa d \theta_{\mathrm{L}} \wedge T_{\beta} = -\eta_{\beta}$
mielke-geometrodynamics_141 EQ0430 (6.7.12)	141	■ 1.000	$\kappa \ell^2 d \theta_{\mathrm{L}} \wedge C_{\mathrm{TT}} = 2\eta.$	$\kappa \ell^2 d \theta_{\mathrm{L}} \wedge C_{\mathrm{TT}} = 2\eta.$	$\kappa \ell^2 d \theta_{\mathrm{L}} \wedge C_{\mathrm{TT}} = 2\eta.$
mielke-geometrodynamics_141 EQ0431 (6.7.13)	141	■ 1.000	$D T^{\gamma} \wedge \eta_{\alpha \beta \gamma} + \kappa D \theta_{\mathrm{T}} \wedge T_{[\alpha} \wedge \vartheta_{\beta]} / \ell^2 = 2\kappa (T_{[\alpha} \wedge \mu_{\beta]} - \vartheta_{[\alpha} \wedge D \mu_{\beta]}).$	$D T^{\gamma} \wedge \eta_{\alpha \beta \gamma} + \kappa D \theta_{\mathrm{T}} \wedge T_{[\alpha} \wedge \vartheta_{\beta]} / \ell^2 = 2\kappa (T_{[\alpha} \wedge \mu_{\beta]} - \vartheta_{[\alpha} \wedge D \mu_{\beta]}).$	$D T^{\gamma} \wedge \eta_{\alpha \beta \gamma} + \kappa D \theta_{\mathrm{T}} \wedge T_{[\alpha} \wedge \vartheta_{\beta]} / \ell^2 = 2\kappa (T_{[\alpha} \wedge \mu_{\beta]} - \vartheta_{[\alpha} \wedge D \mu_{\beta]}).$
mielke-geometrodynamics_142 EQ0432 (6.8.1)	142	■ 1.000	$W = \int \left[L(\vartheta^{\alpha}, T^{\alpha}, R^{\alpha \beta}, \Psi, D \Psi) + L_g(\vartheta^{\alpha}, T^{\alpha}, R^{\alpha \beta}) \right].$	$W = \int \left[L(\vartheta^{\alpha}, T^{\alpha}, R^{\alpha \beta}, \Psi, D \Psi) + L_g(\vartheta^{\alpha}, T^{\alpha}, R^{\alpha \beta}) \right].$	$W = \int \left[L(\vartheta^{\alpha}, T^{\alpha}, R^{\alpha \beta}, \Psi, D \Psi) + L_g(\vartheta^{\alpha}, T^{\alpha}, R^{\alpha \beta}) \right].$
mielke-geometrodynamics_142 EQ0433 (6.8.3)	142	■ 1.000	$\frac{\delta L}{\delta \Psi} = \frac{\partial L}{\partial \Psi} - (-1)^p D \frac{\partial L}{\partial D \Psi} = 0, \quad (\text{MATTER})$ $D H_{\alpha} - E_{\alpha} = \Sigma_{\alpha}, \quad (\text{FIRST})$ $D H_{\alpha \beta} - E_{\alpha \beta} = \tau_{\alpha \beta}. \quad (\text{SECOND})$	$\frac{\delta L}{\delta \Psi} = \frac{\partial L}{\partial \Psi} - (-1)^p D \frac{\partial L}{\partial D \Psi} = 0, \quad (\text{MATTER})$ $D H_{\alpha} - E_{\alpha} = \Sigma_{\alpha}, \quad (\text{FIRST})$ $D H_{\alpha \beta} - E_{\alpha \beta} = \tau_{\alpha \beta}. \quad (\text{SECOND})$	$\frac{\delta L}{\delta \Psi} = \frac{\partial L}{\partial \Psi} - (-1)^p D \frac{\partial L}{\partial D \Psi} = 0, \quad (\text{MATTER})$ $D H_{\alpha} - E_{\alpha} = \Sigma_{\alpha}, \quad (\text{FIRST})$ $D H_{\alpha \beta} - E_{\alpha \beta} = \tau_{\alpha \beta}. \quad (\text{SECOND})$
mielke-geometrodynamics_142 EQ0435 (6.8.6)	142	■ 1.000	$\Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = \frac{\partial L}{\partial \vartheta^{\alpha}} + D \frac{\partial L}{\partial T^{\alpha}}$	$\Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = \frac{\partial L}{\partial \vartheta^{\alpha}} + D \frac{\partial L}{\partial T^{\alpha}}$	$\Sigma_{\alpha} := \frac{\delta L}{\delta \vartheta^{\alpha}} = \frac{\partial L}{\partial \vartheta^{\alpha}} + D \frac{\partial L}{\partial T^{\alpha}}$
mielke-geometrodynamics_142 EQ0436 (6.8.7)	142	■ 1.000	$\tau_{\alpha \beta} := \frac{\delta L}{\delta \Gamma^{\alpha \beta}} = \rho(I_{\alpha \beta}) \Psi \wedge \frac{\partial L}{\partial (D \Psi)} + \vartheta_{[\alpha} \wedge \frac{\partial L}{\partial T^{\beta]} + D \frac{\partial L}{\partial R^{\alpha \beta}},$	$\tau_{\alpha \beta} := \frac{\delta L}{\delta \Gamma^{\alpha \beta}} = \rho(I_{\alpha \beta}) \Psi \wedge \frac{\partial L}{\partial (D \Psi)} + \vartheta_{[\alpha} \wedge \frac{\partial L}{\partial T^{\beta]} + D \frac{\partial L}{\partial R^{\alpha \beta}},$	$\tau_{\alpha \beta} := \frac{\delta L}{\delta \Gamma^{\alpha \beta}} = \rho(I_{\alpha \beta}) \Psi \wedge \frac{\partial L}{\partial (D \Psi)} + \vartheta_{[\alpha} \wedge \frac{\partial L}{\partial T^{\beta]} + D \frac{\partial L}{\partial R^{\alpha \beta}},$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_142 EQ0438 (6.8.9)	142	■ 1.000	$E_{\alpha\beta} := -\vartheta_{[\alpha} \wedge H_{\beta]}$	$E_{\alpha\beta} := -\vartheta_{[\alpha} \wedge H_{\beta]}$	$E_{\alpha\beta} := -\vartheta_{[\alpha} \wedge H_{\beta]}$
mielke-geometrodynamics_143 EQ0439 (6.8.10)	143	■ 1.000	$\vartheta^\alpha \wedge E_\alpha = 4L_g + 2T^\alpha \wedge H_\alpha + 2R^{\alpha\beta} \wedge H_{\alpha\beta}.$	$\vartheta^\alpha \wedge E_\alpha = 4L_g + 2T^\alpha \wedge H_\alpha + 2R^{\alpha\beta} \wedge H_{\alpha\beta}.$	$\vartheta^\alpha \wedge E_\alpha = 4L_g + 2T^\alpha \wedge H_\alpha + 2R^{\alpha\beta} \wedge H_{\alpha\beta}.$
mielke-geometrodynamics_143 EQ0442 (6.8.13)	143	■ 1.000	$L_{\mathrm{qPG}} := -\frac{1}{2} T^\alpha \wedge H_\alpha - \frac{1}{2} R^{\alpha\beta} \wedge \left(H_{\alpha\beta} - \frac{a_0}{2\ell^2} \eta_{\alpha\beta} \right),$	$L_{\mathrm{qPG}} = -\frac{1}{2} T^\alpha \wedge H_\alpha - \frac{1}{2} R^{\alpha\beta} \wedge \left(H_{\alpha\beta} - \frac{a_0}{2\ell^2} \eta_{\alpha\beta} \right),$	$L_{\mathrm{qPG}} = -\frac{1}{2} T^\alpha \wedge H_\alpha - \frac{1}{2} R^{\alpha\beta} \wedge \left(H_{\alpha\beta} - \frac{a_0}{2\ell^2} \eta_{\alpha\beta} \right),$
mielke-geometrodynamics_143 EQ0444 (6.8.15)	143	■ 1.000	$H_{\alpha\beta} := -\frac{a_0}{2\ell^2} \eta_{\alpha\beta} - \frac{1}{\kappa} * \left(\sum_{j=1}^6 b_j^{(j)} R_{\alpha\beta} \right).$	$H_{\alpha\beta} = -\frac{a_0}{2\ell^2} \eta_{\alpha\beta} - \frac{1}{\kappa} * \left(\sum_{j=1}^6 b_j^{(j)} R_{\alpha\beta} \right).$	$H_{\alpha\beta} = -\frac{a_0}{2\ell^2} \eta_{\alpha\beta} - \frac{1}{\kappa} * \left(\sum_{j=1}^6 b_j^{(j)} R_{\alpha\beta} \right).$
mielke-geometrodynamics_144 EQ0447 (6.9.3)	144	■ 1.000	$\eta = *1 = \frac{1}{4!} \eta_{\alpha\beta\gamma\delta} \vartheta^\alpha \wedge \vartheta^\beta \wedge \vartheta^\gamma \wedge \vartheta^\delta,$	$\eta = *1 = \frac{1}{4!} \eta_{\alpha\beta\gamma\delta} \vartheta^\alpha \wedge \vartheta^\beta \wedge \vartheta^\gamma \wedge \vartheta^\delta,$	$\eta = *1 = \frac{1}{4!} \eta_{\alpha\beta\gamma\delta} \vartheta^\alpha \wedge \vartheta^\beta \wedge \vartheta^\gamma \wedge \vartheta^\delta,$
mielke-geometrodynamics_145 EQ0448 (6.9.4)	145	■ 1.000	$p = \frac{1}{8\pi^2} \int R_\alpha{}^\beta \wedge R_\beta{}^\alpha.$	$p = \frac{1}{8\pi^2} \int R_\alpha{}^\beta \wedge R_\beta{}^\alpha.$	$p = \frac{1}{8\pi^2} \int R_\alpha{}^\beta \wedge R_\beta{}^\alpha.$
mielke-geometrodynamics_151 EQ0453 (7.2.1)	151	■ 1.000	$L_{ab}^{(*)} := \frac{1}{2} \varepsilon_{abcd} L^{cd} = L_{ab}.$	$L_{ab}^{(*)} := \frac{1}{2} \varepsilon_{abcd} L^{cd} = L_{ab}.$	$L_{ab}^{(*)} := \frac{1}{2} \varepsilon_{abcd} L^{cd} = L_{ab}.$
mielke-geometrodynamics_152 EQ0454 (7.2.6)	152	■ 1.000	$\Omega^g = \Omega_C + \bar{\Omega}^g - \frac{1}{6} \vartheta \wedge \vartheta R.$	$\Omega^g = \Omega_C + \bar{\Omega}^g - \frac{1}{6} \vartheta \wedge \vartheta R.$	$\Omega^g = \Omega_C + \bar{\Omega}^g - \frac{1}{6} \vartheta \wedge \vartheta R.$
mielke-geometrodynamics_152 EQ0455 (7.2.7)	152	■ 1.000	$\bar{\Omega}^g = \vartheta \wedge *(\vartheta \wedge \Omega^g) - \frac{1}{4} \vartheta \wedge \vartheta R,$	$\bar{\Omega}^g = \vartheta \wedge *(\vartheta \wedge \Omega^g) - \frac{1}{4} \vartheta \wedge \vartheta R,$	$\bar{\Omega}^g = \vartheta \wedge *(\vartheta \wedge \Omega^g) - \frac{1}{4} \vartheta \wedge \vartheta R,$
mielke-geometrodynamics_152 EQ0457 (7.2.9)	152	■ 1.000	$\mathrm{Tr}(\vartheta \wedge \vartheta \wedge \Omega_C) = \mathrm{Tr}(\vartheta \wedge \vartheta \wedge \bar{\Omega}^g) = 0$	$\mathrm{Tr}(\vartheta \wedge \vartheta \wedge \Omega_C) = \mathrm{Tr}(\vartheta \wedge \vartheta \wedge \bar{\Omega}^g) = 0$	$\mathrm{Tr}(\vartheta \wedge \vartheta \wedge \Omega_C) = \mathrm{Tr}(\vartheta \wedge \vartheta \wedge \bar{\Omega}^g) = 0$
mielke-geometrodynamics_153 EQ0463 (7.3.2)	153	■ 1.000	$D^* \Omega^g = 0,$	$D^* \Omega^g = 0,$	$D^* \Omega^g = 0,$
mielke-geometrodynamics_153 EQ0464 (7.3.3)	153	■ 1.000	$D^* R_{\alpha\beta} = 0,$	$D^* R_{\alpha\beta} = 0,$	$D^* R_{\alpha\beta} = 0,$
mielke-geometrodynamics_154 EQ0468 (7.3.7)	154	■ 1.000	$R_{\alpha\beta} = -{}^* R_{\alpha\beta}^{(*)},$	$R_{\alpha\beta} = -{}^* R_{\alpha\beta}^{(*)},$	$R_{\alpha\beta} = -{}^* R_{\alpha\beta}^{(*)},$
mielke-geometrodynamics_154 EQ0469 (7.3.8)	154	■ 1.000	$R_{\alpha\beta} = {}^* R_{\alpha\beta}^{(*)},$	$R_{\alpha\beta} = {}^* R_{\alpha\beta}^{(*)},$	$R_{\alpha\beta} = {}^* R_{\alpha\beta}^{(*)},$
mielke-geometrodynamics_157 EQ0479 (7.4.7)	157	■ 1.000	$G_\alpha - \Lambda \eta_\alpha = 0$	$G_\alpha - \Lambda \eta_\alpha = 0$	$G_\alpha - \Lambda \eta_\alpha = 0$
mielke-geometrodynamics_157 EQ0480 (7.5.1)	157	■ 1.000	$\square^c := \square - \frac{1}{6} R, \quad \square := (1/\sqrt{g}) \partial_i (\sqrt{g} g^{ij} \partial_j).$	$\square^c := \square - \frac{1}{6} R, \quad \square := (1/\sqrt{g}) \partial_i (\sqrt{g} g^{ij} \partial_j).$	$\square^c := \square - \frac{1}{6} R, \quad \square := (1/\sqrt{g}) \partial_i (\sqrt{g} g^{ij} \partial_j).$
mielke-geometrodynamics_157 EQ0482 (7.5.3)	157	■ 1.000	$\left(\square - \frac{2}{3} \Lambda \right) h_{ij} = 0$	$\left(\square - \frac{2}{3} \Lambda \right) h_{ij} = 0$	$\left(\square - \frac{2}{3} \Lambda \right) h_{ij} = 0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_158 EQ0483 (7.5.4)	158	■ 1.000	$\partial^i h_{ij} - \partial_j h = 0$	$\partial^i h_{ij} - \partial_j h = 0$	$\partial^i h_{ij} - \partial_j h = 0$
mielke-geometrodynamics_158 EQ0484 (7.5.5)	158	■ 1.000	$\Delta m = \sqrt{2 \Lambda / 3} > 0$	$\Delta m = \sqrt{2 \Lambda / 3} > 0$	$\Delta m = \sqrt{2 \Lambda / 3} > 0$
mielke-geometrodynamics_160 EQ0488 (7.7.5)	160	■ 1.000	$L_{\mathrm{EC}} = -\frac{1}{2\kappa^2} R^{\alpha\beta} \wedge \eta_{\alpha\beta}$	$L_{\mathrm{EC}} = -\frac{1}{2\kappa^2} R^{\alpha\beta} \wedge \eta_{\alpha\beta}$	$L_{\mathrm{EC}} = -\frac{1}{2\kappa^2} R^{\alpha\beta} \wedge \eta_{\alpha\beta}$
mielke-geometrodynamics_160 EQ0489 (7.7.6)	160	■ 1.000	$\widetilde{L} = -\sum_{k=0}^K (1/2k) R^{\alpha\beta} \wedge \stackrel{(2k)}{H}_{\alpha\beta} + L_\theta,$	$\widetilde{L} = -\sum_{k=0}^K (1/2k) R^{\alpha\beta} \wedge \stackrel{(2k)}{H}_{\alpha\beta} + L_\theta,$	$\widetilde{L} = -\sum_{k=0}^K (1/2k) R^{\alpha\beta} \wedge \stackrel{(2k)}{H}_{\alpha\beta} + L_\theta,$
mielke-geometrodynamics_161 EQ0491 (7.7.9)	161	■ 1.000	$\widetilde{L} \rightarrow L = -\frac{1}{2} \left(R^{\alpha\beta} \wedge \widetilde{H}_{\alpha\beta} - \widetilde{L} \right)$	$\widetilde{L} \rightarrow L = -\frac{1}{2} \left(R^{\alpha\beta} \wedge \widetilde{H}_{\alpha\beta} - \widetilde{L} \right)$	$\widetilde{L} \rightarrow L = -\frac{1}{2} \left(R^{\alpha\beta} \wedge \widetilde{H}_{\alpha\beta} - \widetilde{L} \right)$
mielke-geometrodynamics_161 EQ0493 (7.7.11)	161	■ 1.000	$\widetilde{H}_{\alpha\beta\mu\nu} := \frac{\partial^2 \widetilde{L}}{\partial R^{\mu\nu} \partial R^{\alpha\beta}} = -\frac{\partial \widetilde{H}_{\mu\nu}}{\partial R^{\alpha\beta}}$	$\widetilde{H}_{\alpha\beta\mu\nu} := \frac{\partial^2 \widetilde{L}}{\partial R^{\mu\nu} \partial R^{\alpha\beta}} = -\frac{\partial \widetilde{H}_{\mu\nu}}{\partial R^{\alpha\beta}}$	$\widetilde{H}_{\alpha\beta\mu\nu} := \frac{\partial^2 \widetilde{L}}{\partial R^{\mu\nu} \partial R^{\alpha\beta}} = -\frac{\partial \widetilde{H}_{\mu\nu}}{\partial R^{\alpha\beta}}$
mielke-geometrodynamics_161 EQ0494 (7.7.12)	161	■ 1.000	$\widetilde{\eta}_{\alpha\beta} := \frac{\theta_{\mathrm{T}}^*}{2} \eta_{\alpha\beta} - \frac{\theta_{\mathrm{T}}}{2} \vartheta_\alpha \wedge \vartheta_\beta - \ell^2 \widetilde{H}_{\alpha\beta} = 0$	$\widetilde{\eta}_{\alpha\beta} := \frac{\theta_{\mathrm{T}}^*}{2} \eta_{\alpha\beta} - \frac{\theta_{\mathrm{T}}}{2} \vartheta_\alpha \wedge \vartheta_\beta - \ell^2 \widetilde{H}_{\alpha\beta} = 0$	$\widetilde{\eta}_{\alpha\beta} := \frac{\theta_{\mathrm{T}}^*}{2} \eta_{\alpha\beta} - \frac{\theta_{\mathrm{T}}}{2} \vartheta_\alpha \wedge \vartheta_\beta - \ell^2 \widetilde{H}_{\alpha\beta} = 0$
mielke-geometrodynamics_162 EQ0496 (7.7.14)	162	■ 1.000	$\eta := \eta_{\alpha\beta\gamma\delta} \vartheta^\alpha \wedge \vartheta^\beta \wedge \vartheta^\gamma \wedge \vartheta^\delta / 4!$	$\eta := \eta_{\alpha\beta\gamma\delta} \vartheta^\alpha \wedge \vartheta^\beta \wedge \vartheta^\gamma \wedge \vartheta^\delta / 4!$	$\eta := \eta_{\alpha\beta\gamma\delta} \vartheta^\alpha \wedge \vartheta^\beta \wedge \vartheta^\gamma \wedge \vartheta^\delta / 4!$
mielke-geometrodynamics_163 EQ0499 (7.7.17)	163	■ 1.000	$\gamma_\alpha \gamma_\beta + \gamma_\beta \gamma_\alpha = 2 o_{\alpha\beta} \mathbf{1}_4.$	$\gamma_\alpha \gamma_\beta + \gamma_\beta \gamma_\alpha = 2 o_{\alpha\beta} \mathbf{1}_4.$	$\gamma_\alpha \gamma_\beta + \gamma_\beta \gamma_\alpha = 2 o_{\alpha\beta} \mathbf{1}_4.$
mielke-geometrodynamics_163 EQ0501 (7.7.19)	163	■ 1.000	$\gamma := \gamma^\alpha \vartheta_\alpha$	$\gamma := \gamma^\alpha \vartheta_\alpha$	$\gamma := \gamma^\alpha \vartheta_\alpha$
mielke-geometrodynamics_163 EQ0502 (7.7.20)	163	■ 1.000	${}^* \gamma = \gamma^\alpha \eta_\alpha = \frac{i}{6} \gamma_5 \gamma \wedge \gamma \wedge \gamma,$	${}^* \gamma = \gamma^\alpha \eta_\alpha = \frac{i}{6} \gamma_5 \gamma \wedge \gamma \wedge \gamma,$	${}^* \gamma = \gamma^\alpha \eta_\alpha = \frac{i}{6} \gamma_5 \gamma \wedge \gamma \wedge \gamma,$
mielke-geometrodynamics_164 EQ0503 (7.7.21)	164	■ 1.000	$\sigma := \frac{1}{2} \sigma_{\alpha\beta} \vartheta^\alpha \wedge \vartheta^\beta = \frac{i}{2} \gamma \wedge \gamma, \quad {}^* \sigma = \frac{1}{2} \sigma_{\alpha\beta} \eta^{\alpha\beta} =: \sigma^{(*)} = i \gamma_5 \sigma.$	$\sigma := \frac{1}{2} \sigma_{\alpha\beta} \vartheta^\alpha \wedge \vartheta^\beta = \frac{i}{2} \gamma \wedge \gamma, \quad {}^* \sigma = \frac{1}{2} \sigma_{\alpha\beta} \eta^{\alpha\beta} =: \sigma^{(*)} = i \gamma_5 \sigma.$	$\sigma := \frac{1}{2} \sigma_{\alpha\beta} \vartheta^\alpha \wedge \vartheta^\beta = \frac{i}{2} \gamma \wedge \gamma, \quad {}^* \sigma = \frac{1}{2} \sigma_{\alpha\beta} \eta^{\alpha\beta} =: \sigma^{(*)} = i \gamma_5 \sigma.$
mielke-geometrodynamics_164 EQ0504 (7.7.22)	164	■ 1.000	$\sigma^{(*)} := i \gamma_5 \sigma, \quad \sigma^{(**)} = i^2 \gamma_5^2 \sigma = -\sigma.$	$\sigma^{(*)} := i \gamma_5 \sigma, \quad \sigma^{(**)} = i^2 \gamma_5^2 \sigma = -\sigma.$	$\sigma^{(*)} := i \gamma_5 \sigma, \quad \sigma^{(**)} = i^2 \gamma_5^2 \sigma = -\sigma.$
mielke-geometrodynamics_164 EQ0505 (7.7.23)	164	■ 1.000	$[\gamma, \sigma^{(*)}] = \gamma \wedge i \gamma_5 \sigma - i \gamma_5 \sigma \wedge \gamma = -i \gamma_5 (\gamma \wedge \sigma + \sigma \wedge \gamma) = 2 \gamma_5 \gamma \wedge \gamma \wedge \gamma$	$[\gamma, \sigma^{(*)}] = \gamma \wedge i \gamma_5 \sigma - i \gamma_5 \sigma \wedge \gamma = -i \gamma_5 (\gamma \wedge \sigma + \sigma \wedge \gamma) = 2 \gamma_5 \gamma \wedge \gamma \wedge \gamma$	$[\gamma, \sigma^{(*)}] = \gamma \wedge i \gamma_5 \sigma - i \gamma_5 \sigma \wedge \gamma = -i \gamma_5 (\gamma \wedge \sigma + \sigma \wedge \gamma) = 2 \gamma_5 \gamma \wedge \gamma \wedge \gamma$
mielke-geometrodynamics_165 EQ0508 (7.7.26)	165	■ 1.000	$\gamma \rightarrow \gamma^\theta = e^{i \gamma^5 \theta} \gamma e^{-i \gamma^5 \theta}$	$\gamma \rightarrow \gamma^\theta = e^{i \gamma^5 \theta} \gamma e^{-i \gamma^5 \theta}$	$\gamma \rightarrow \gamma^\theta = e^{i \gamma^5 \theta} \gamma e^{-i \gamma^5 \theta}$
mielke-geometrodynamics_165 EQ0510 (7.7.28)	165	■ 1.000	$\gamma := \vartheta^\alpha \gamma_\alpha, \quad \Gamma := \frac{i}{4} \Gamma^{\alpha\beta} \sigma_{\alpha\beta} = \Gamma^{\{\}} - K,$	$\gamma := \vartheta^\alpha \gamma_\alpha, \quad \Gamma := \frac{i}{4} \Gamma^{\alpha\beta} \sigma_{\alpha\beta} = \Gamma^{\{\}} - K,$	$\gamma := \vartheta^\alpha \gamma_\alpha, \quad \Gamma := \frac{i}{4} \Gamma^{\alpha\beta} \sigma_{\alpha\beta} = \Gamma^{\{\}} - K,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_184 EQ0569 (8.6.7)	184	■ 1.000	$\begin{aligned} D H_{\alpha}-E_{\alpha} &=\Sigma_{\alpha}, \\ D H_{\alpha \beta}+\vartheta_{[\alpha} \wedge H_{\beta]} &=\tau_{\alpha \beta}, \end{aligned}$	$D H_{\alpha}-E_{\alpha}=\Sigma_{\alpha},$ $D H_{\alpha \beta}+\vartheta_{[\alpha} \wedge H_{\beta]}=\tau_{\alpha \beta},$	$D H_{\alpha}-E_{\alpha}=\Sigma_{\alpha},$ $D H_{\alpha \beta}+\vartheta_{[\alpha} \wedge H_{\beta]}=\tau_{\alpha \beta},$
mielke-geometrodynamics_184 EQ0571 (8.6.9)	184	■ 1.000	$\frac{\theta_{\mathrm{T}}^*}{2 \ell^2} \eta_{\alpha \beta \gamma} T^{\gamma}+\frac{\theta_{\mathrm{T}}}{\ell^2} \vartheta_{[\alpha} \wedge T_{\beta]}+\vartheta_{[\alpha} \wedge H_{\beta]}(**)=\tau_{\alpha \beta}.$	$\frac{\theta_{\mathrm{T}}^*}{2 \ell^2} \eta_{\alpha \beta \gamma} T^{\gamma}+\frac{\theta_{\mathrm{T}}}{\ell^2} \vartheta_{[\alpha} \wedge T_{\beta]}+\vartheta_{[\alpha} \wedge H_{\beta]}(**)=\tau_{\alpha \beta}.$	$\frac{\theta_{\mathrm{T}}^*}{2 \ell^2} \eta_{\alpha \beta \gamma} T^{\gamma}+\frac{\theta_{\mathrm{T}}}{\ell^2} \vartheta_{[\alpha} \wedge T_{\beta]}+\vartheta_{[\alpha} \wedge H_{\beta]}(**)=\tau_{\alpha \beta}.$
mielke-geometrodynamics_185 EQ0576 (8.6.14)	185	■ 1.000	$\Lambda_{\mathrm{eff}}=\frac{1}{\theta_{\mathrm{T}}^*}\left[\Lambda-\frac{1}{4}\left(\theta_{\mathrm{T}}^*-a_0\right) R\right]=\frac{\Lambda}{\theta_{\mathrm{T}}^*}-\Lambda_{\theta}$	$\Lambda_{\mathrm{eff}}=\frac{1}{\theta_{\mathrm{T}}^*}\left[\Lambda-\frac{1}{4}\left(\theta_{\mathrm{T}}^*-a_0\right) R\right]=\frac{\Lambda}{\theta_{\mathrm{T}}^*}-\Lambda_{\theta}$	$\Lambda_{\mathrm{eff}}=\frac{1}{\theta_{\mathrm{T}}^*}\left[\Lambda-\frac{1}{4}\left(\theta_{\mathrm{T}}^*-a_0\right) R\right]=\frac{\Lambda}{\theta_{\mathrm{T}}^*}-\Lambda_{\theta}$
mielke-geometrodynamics_186 EQ0578 (8.6.16)	186	■ 1.000	$\widetilde{L}_{\mathrm{qPG}}=L_{\mathrm{qPG}}+\lambda_{\alpha} \wedge T^{\alpha}.$	$\widetilde{L}_{\mathrm{qPG}}=L_{\mathrm{qPG}}+\lambda_{\alpha} \wedge T^{\alpha}.$	$\widetilde{L}_{\mathrm{qPG}}=L_{\mathrm{qPG}}+\lambda_{\alpha} \wedge T^{\alpha}.$
mielke-geometrodynamics_188 EQ0582 (8.6.20)	188	■ 1.000	$\Lambda_{\theta}=\frac{3 \left(\theta_{\mathrm{T}}^*-a_0\right)^2}{2 \ell^2\left(\theta_{\mathrm{L}}^*-b_6\right) \theta_{\mathrm{T}}^*}$	$\Lambda_{\theta}=\frac{3 \left(\theta_{\mathrm{T}}^*-a_0\right)^2}{2 \ell^2\left(\theta_{\mathrm{L}}^*-b_6\right) \theta_{\mathrm{T}}^*}$	$\Lambda_{\theta}=\frac{3 \left(\theta_{\mathrm{T}}^*-a_0\right)^2}{2 \ell^2\left(\theta_{\mathrm{L}}^*-b_6\right) \theta_{\mathrm{T}}^*}$
mielke-geometrodynamics_193 EQ0586 (9.1.4)	193	■ 1.000	$ds^2=\left(\frac{N(r \pm t)}{r}\right)^2\left(-dt^2+dr^2+r^2d\Omega^2\right),$	$ds^2=\left(\frac{N(r \pm t)}{r}\right)^2\left(-dt^2+dr^2+r^2d\Omega^2\right),$	$ds^2=\left(\frac{N(r \pm t)}{r}\right)^2\left(-dt^2+dr^2+r^2d\Omega^2\right),$
mielke-geometrodynamics_194 EQ0588 (9.1.6)	194	■ 1.000	$u^2-v^2=e^{r^*/4\mu}$	$u^2-v^2=e^{r^*/4\mu}$	$u^2-v^2=e^{r^*/4\mu}$
mielke-geometrodynamics_194 EQ0589 (9.1.7)	194	■ 1.000	$(u+v)/(u-v)=e^{t/2\mu},$	$(u+v)/(u-v)=e^{t/2\mu},$	$(u+v)/(u-v)=e^{t/2\mu},$
mielke-geometrodynamics_194 EQ0590 (9.1.8) [re-fined: census]	194	■ 1.000	$ds^2=16\,\mu u^2\,e^{\nu-r^*\cancel{\mu}}\cancel{\mu}\left(-d\nu^2+du^2\right)+r^2d\Omega^2,$	$ds^2=16\mu^2e^{\nu-r^*\cancel{\mu}}\left(-dv^2+du^2\right)+r^2d\Omega^2,$	$ds^2=16\mu^2e^{\nu-r^*/2\mu}\left(-dv^2+du^2\right)+r^2d\Omega^2,$
mielke-geometrodynamics_195 EQ0591 (9.1.9)	195	■ 1.000	$\Omega^{\{\}}=\Omega_{\mathrm{C}}^{\{\}}+\bar{\Omega}^{\{\}}-\frac{1}{6}R^{\{\}}\vartheta\wedge\vartheta,$	$\Omega^{\{\}}=\Omega_{\mathrm{C}}^{\{\}}+\bar{\Omega}^{\{\}}-\frac{1}{6}R^{\{\}}\vartheta\wedge\vartheta,$	$\Omega^{\{\}}=\Omega_{\mathrm{C}}^{\{\}}+\bar{\Omega}^{\{\}}-\frac{1}{6}R^{\{\}}\vartheta\wedge\vartheta,$
mielke-geometrodynamics_197 EQ0600 (9.1.18)	197	■ 1.000	$\begin{aligned} ds_{\mathrm{Sds}}^2&=\left(1-\frac{2\mu}{r}+\frac{\kappa}{4\ell^{*2}}\right)d\tau^2 \\ &+\left(1-\frac{2\mu}{r}+\frac{\kappa}{4\ell^{*2}}\right)^{-1}dr^2+r^2d\Omega^2 \end{aligned}$	$ds_{\mathrm{Sds}}^2=\left(1-\frac{2\mu}{r}+\frac{\kappa}{4\ell^{*2}}\right)d\tau^2$ $+\left(1-\frac{2\mu}{r}+\frac{\kappa}{4\ell^{*2}}\right)^{-1}dr^2+r^2d\Omega^2$	$ds_{\mathrm{Sds}}^2=\left(1-\frac{2\mu}{r}+\frac{\kappa}{4\ell^{*2}}\right)d\tau^2$ $+\left(1-\frac{2\mu}{r}+\frac{\kappa}{4\ell^{*2}}\right)^{-1}dr^2+r^2d\Omega^2,$
mielke-geometrodynamics_197 EQ0601 (9.1.19)	197	■ 1.000	$\begin{aligned} f&=\frac{2\mu}{r^2}\left(1-\frac{2\mu}{r}+\frac{\kappa}{4\ell^{*2}}\right)^{-1/2} \\ g&=h=k=0 \end{aligned}$	$f=\frac{2\mu}{r^2}\left(1-\frac{2\mu}{r}+\frac{\kappa}{4\ell^{*2}}\right)^{-1/2}$ $g=h=k=0$	$f=\frac{2\mu}{r^2}\left(1-\frac{2\mu}{r}+\frac{\kappa}{4\ell^{*2}}\right)^{-1/2}$ $g=h=k=0$
mielke-geometrodynamics_198 EQ0603 (9.2.2)	198	■ 1.000	$p_1(M^4)=-3\,\tau$	$p_1(M^4)=-3\tau$	$p_1(M^4)=-3\tau$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_200 EQ0607 (9.2.8)	200	■ 1.000	$\begin{aligned} \chi(M \# N) &= \chi(M) + \chi(N) - \chi \\ &\left(S^n\right), \quad \chi(M \times N) = \chi(M) \cdot \chi(N), \\ \chi(M^2) &= 2(g-1), \quad \chi \\ &\left(\mathbb{C}P^2\right) = 3 \end{aligned}$	$\begin{aligned} \chi(M \# N) &= \chi(M) + \chi(N) - \chi(S^n), \\ \chi(M \times N) &= \chi(M) \cdot \chi(N), \\ \chi(M^2) &= 2(g-1), \\ \chi(\mathbb{C}P^2) &= 3 \end{aligned}$	$\begin{aligned} \chi(M \# N) &= \chi(M) + \chi(N) - \chi(S^n), \\ \chi(M \times N) &= \chi(M) \cdot \chi(N), \\ \chi(M^2) &= 2(g-1), \\ \chi(\mathbb{C}P^2) &= 3 \end{aligned}$
mielke-geometrodynamics_201 EQ0608 (9.2.10)	201	■ 1.000	$\chi\left(M^{2k+1}\right)=0$	$\chi\left(M^{2k+1}\right)=0$	$\chi\left(M^{2k+1}\right)=0$
mielke-geometrodynamics_208 EQ0612 (10.0.1)	208	■ 1.000	$L_{\mathrm{EC}}=-\frac{\chi}{\ell}\vartheta^\alpha\wedge R_\alpha^\star$	$L_{\mathrm{EC}}=-\frac{\chi}{\ell}\vartheta^\alpha\wedge R_\alpha^\star$	$L_{\mathrm{EC}}=-\frac{\chi}{\ell}\vartheta^\alpha\wedge R_\alpha^\star$
mielke-geometrodynamics_209 EQ0614 (10.1.2)	209	■ 1.000	$T^\alpha:=d\vartheta^\alpha-(-1)^s\eta^{\alpha\beta}\wedge\Gamma_\beta^\star$	$T^\alpha:=d\vartheta^\alpha-(-1)^s\eta^{\alpha\beta}\wedge\Gamma_\beta^\star$	$T^\alpha:=d\vartheta^\alpha-(-1)^s\eta^{\alpha\beta}\wedge\Gamma_\beta^\star$
mielke-geometrodynamics_209 EQ0615 (10.1.3)	209	■ 1.000	$R_\alpha^\star=\frac{1}{2}\eta_{\alpha\beta\gamma}R^{\beta\gamma}:=d\Gamma_\alpha^\star+\frac{(-1)^s}{2}\eta_\alpha^{\beta\gamma}\Gamma_\beta^\star\wedge\Gamma_\gamma^\star,$	$R_\alpha^\star=\frac{1}{2}\eta_{\alpha\beta\gamma}R^{\beta\gamma}:=d\Gamma_\alpha^\star+\frac{(-1)^s}{2}\eta_\alpha^{\beta\gamma}\Gamma_\beta^\star\wedge\Gamma_\gamma^\star,$	$R_\alpha^\star=\frac{1}{2}\eta_{\alpha\beta\gamma}R^{\beta\gamma}:=d\Gamma_\alpha^\star+\frac{(-1)^s}{2}\eta_\alpha^{\beta\gamma}\Gamma_\beta^\star\wedge\Gamma_\gamma^\star,$
mielke-geometrodynamics_209 EQ0616 (10.1.4)	209	■ 1.000	$D\,T^\alpha\equiv(-1)^s\eta^{\alpha\beta}\wedge R_\beta^\star$	$DT^\alpha\equiv(-1)^s\eta^{\alpha\beta}\wedge R_\beta^\star$	$DT^\alpha\equiv(-1)^s\eta^{\alpha\beta}\wedge R_\beta^\star$
mielke-geometrodynamics_209 EQ0617 (10.1.5)	209	■ 1.000	$D\,R_\alpha^\star\equiv 0.$	$DR_\alpha^\star\equiv 0.$	$DR_\alpha^\star\equiv 0.$
mielke-geometrodynamics_209 EQ0618 (10.1.6)	209	■ 1.000	$C_{\mathrm{T}}:=\frac{1}{2\ell^2}\vartheta^\alpha\wedge T_\alpha=-\frac{(-1)^s}{\ell^2}\eta^\alpha\wedge K_\alpha^\star,$	$C_{\mathrm{T}}:=\frac{1}{2\ell^2}\vartheta^\alpha\wedge T_\alpha=-\frac{(-1)^s}{\ell^2}\eta^\alpha\wedge K_\alpha^\star,$	$C_{\mathrm{T}}:=\frac{1}{2\ell^2}\vartheta^\alpha\wedge T_\alpha=-\frac{(-1)^s}{\ell^2}\eta^\alpha\wedge K_\alpha^\star,$
mielke-geometrodynamics_209 EQ0619 (10.1.7)	209	■ 1.000	$C_{\mathrm{L}}:=(-1)^s\Gamma^{\star\alpha}\wedge R_\alpha^\star-\frac{1}{3!}\eta_{\alpha\beta\gamma}\Gamma^{\star\alpha}\wedge\Gamma^{\star\beta}\wedge\Gamma^{\star\gamma}.$	$C_{\mathrm{L}}:=(-1)^s\Gamma^{\star\alpha}\wedge R_\alpha^\star-\frac{1}{3!}\eta_{\alpha\beta\gamma}\Gamma^{\star\alpha}\wedge\Gamma^{\star\beta}\wedge\Gamma^{\star\gamma}.$	$C_{\mathrm{L}}:=(-1)^s\Gamma^{\star\alpha}\wedge R_\alpha^\star-\frac{1}{3!}\eta_{\alpha\beta\gamma}\Gamma^{\star\alpha}\wedge\Gamma^{\star\beta}\wedge\Gamma^{\star\gamma}.$
mielke-geometrodynamics_209 EQ0621 (10.1.9)	209	■ 1.000	$\frac{\delta C_{\mathrm{L}}}{\delta\vartheta^\alpha}=0\quad\text{and}\quad\frac{\delta C_{\mathrm{L}}}{\delta\Gamma^{\star\alpha}}=(-1)^s2R_\alpha^\star,$	$\frac{\delta C_{\mathrm{L}}}{\delta\vartheta^\alpha}=0\quad\text{and}\quad\frac{\delta C_{\mathrm{L}}}{\delta\Gamma^{\star\alpha}}=(-1)^s2R_\alpha^\star,$	$\frac{\delta C_{\mathrm{L}}}{\delta\vartheta^\alpha}=0\quad\text{and}\quad\frac{\delta C_{\mathrm{L}}}{\delta\Gamma^{\star\alpha}}=(-1)^s2R_\alpha^\star,$
mielke-geometrodynamics_210 EQ0622 (10.1.10)	210	■ 1.000	$C_{\mathrm{TL}}:=\frac{1}{\ell}\left(\Gamma^{\star\alpha}\wedge T_\alpha-\frac{(-1)^s}{2}\eta_{\alpha\beta\gamma}\Gamma^{\star\alpha}\wedge\Gamma^{\star\beta}\wedge\vartheta^\gamma\right).$	$C_{\mathrm{TL}}:=\frac{1}{\ell}\left(\Gamma^{\star\alpha}\wedge T_\alpha-\frac{(-1)^s}{2}\eta_{\alpha\beta\gamma}\Gamma^{\star\alpha}\wedge\Gamma^{\star\beta}\wedge\vartheta^\gamma\right).$	$C_{\mathrm{TL}}:=\frac{1}{\ell}\left(\Gamma^{\star\alpha}\wedge T_\alpha-\frac{(-1)^s}{2}\eta_{\alpha\beta\gamma}\Gamma^{\star\alpha}\wedge\Gamma^{\star\beta}\wedge\vartheta^\gamma\right).$
mielke-geometrodynamics_210 EQ0625 (10.1.13)	210	■ 1.000	$D\,H_\alpha^\star-S_\alpha=\tau_\alpha^\star,$	$DH_\alpha^\star-S_\alpha=\tau_\alpha^\star,$	$DH_\alpha^\star-S_\alpha=\tau_\alpha^\star,$
mielke-geometrodynamics_211 EQ0626 (10.1.14)	211	■ 1.000	$H_\alpha:=-\frac{\partial L}{\partial T^\alpha},\quad H_\alpha^\star:=-\frac{(-1)^s}{2}\frac{\partial L}{\partial R^{\star\alpha}},$	$H_\alpha:=-\frac{\partial L}{\partial T^\alpha},\quad H_\alpha^\star:=-\frac{(-1)^s}{2}\frac{\partial L}{\partial R^{\star\alpha}},$	$H_\alpha:=-\frac{\partial L}{\partial T^\alpha},\quad H_\alpha^\star:=-\frac{(-1)^s}{2}\frac{\partial L}{\partial R^{\star\alpha}},$
mielke-geometrodynamics_211 EQ0628 (10.1.16)	211	■ 1.000	$S_\alpha:=\frac{(-1)^s}{2}\frac{\partial L}{\partial\Gamma^{\star\alpha}}=\frac{1}{2}\eta_{\alpha\beta}\wedge H^\beta.$	$S_\alpha:=\frac{(-1)^s}{2}\frac{\partial L}{\partial\Gamma^{\star\alpha}}=\frac{1}{2}\eta_{\alpha\beta}\wedge H^\beta.$	$S_\alpha:=\frac{(-1)^s}{2}\frac{\partial L}{\partial\Gamma^{\star\alpha}}=\frac{1}{2}\eta_{\alpha\beta}\wedge H^\beta.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_211 E0629 (10.1.17)	211	■ 1.000	$\Sigma_{\alpha}:=\frac{\delta L_{\mathrm{MAT}}}{\delta \vartheta^{\alpha}}, \quad \tau_{\alpha}^{\star}:=\frac{(-1)^s}{2} \frac{\delta L_{\mathrm{MAT}}}{\delta \Gamma^{\star \alpha}},$	$\Sigma_{\alpha}:=\frac{\delta L_{\mathrm{MAT}}}{\delta \vartheta^{\alpha}}, \quad \tau_{\alpha}^{\star}:=\frac{(-1)^s}{2} \frac{\delta L_{\mathrm{MAT}}}{\delta \Gamma^{\star \alpha}},$	
mielke-geometrodynamics_211 E0630 (10.2.1)	211	■ 1.000	$L_{\mathrm{EC}}:=-\frac{\chi}{\ell} \vartheta^{\alpha} \wedge R_{\alpha}^{\star} \equiv-\chi C_{\mathrm{TL}}-\frac{\chi}{\ell} d\left(\Gamma_{\alpha}^{\star} \wedge \vartheta^{\alpha}\right)$	$L_{\mathrm{EC}}:=-\frac{\chi}{\ell} \vartheta^{\alpha} \wedge R_{\alpha}^{\star} \equiv-\chi C_{\mathrm{TL}}-\frac{\chi}{\ell} d\left(\Gamma_{\alpha}^{\star} \wedge \vartheta^{\alpha}\right)$	
mielke-geometrodynamics_211 E0631 (10.2.2)	211	■ 1.000	$L_{\mathrm{MB}}\left(\vartheta^{\alpha}, \Gamma_{\alpha}^{\star}\right)=\theta_{\mathrm{T}} C_{\mathrm{T}}+\theta_{\mathrm{L}} C_{\mathrm{L}}+\theta_{\mathrm{TL}} C_{\mathrm{TL}},$	$L_{\mathrm{MB}}\left(\vartheta^{\alpha}, \Gamma_{\alpha}^{\star}\right)=\theta_{\mathrm{T}} C_{\mathrm{T}}+\theta_{\mathrm{L}} C_{\mathrm{L}}+\theta_{\mathrm{TL}} C_{\mathrm{TL}},$	
mielke-geometrodynamics_212 E0632 (10.2.3)	212	■ 1.000	$-\theta_{\mathrm{TL}} R_{\alpha}^{\star}-\frac{1}{\ell} \theta_{\mathrm{T}} T_{\alpha}=\ell \Sigma_{\alpha}$	$-\theta_{\mathrm{TL}} R_{\alpha}^{\star}-\frac{1}{\ell} \theta_{\mathrm{T}} T_{\alpha}=\ell \Sigma_{\alpha}$	
mielke-geometrodynamics_212 E0634 (10.2.5)	212	■ 1.000	$T_{\alpha}-\frac{\varepsilon}{\ell} \eta_{\alpha}=\frac{2}{A} \ell\left(\theta_{\mathrm{TL}} \tau_{\alpha}^{\star}-\theta_{\mathrm{L}} \ell \Sigma_{\alpha}\right)$	$T_{\alpha}-\frac{\varepsilon}{\ell} \eta_{\alpha}=\frac{2}{A} \ell\left(\theta_{\mathrm{TL}} \tau_{\alpha}^{\star}-\theta_{\mathrm{L}} \ell \Sigma_{\alpha}\right)$	
mielke-geometrodynamics_212 E0635 (10.2.6)	212	■ 1.000	$R_{\alpha}^{\star}-\frac{\rho}{\ell^2} \eta_{\alpha}=\frac{2}{A}\left(\theta_{\mathrm{TL}} \ell \Sigma_{\alpha}-\theta_{\mathrm{T}} \tau_{\alpha}^{\star}\right)$	$R_{\alpha}^{\star}-\frac{\rho}{\ell^2} \eta_{\alpha}=\frac{2}{A}\left(\theta_{\mathrm{TL}} \ell \Sigma_{\alpha}-\theta_{\mathrm{T}} \tau_{\alpha}^{\star}\right)$	
mielke-geometrodynamics_213 E0638 (10.2.9)	213	■ 1.000	$R_{\alpha}^{\star} \equiv R_{\alpha}^{\{\} \star}-D K_{\alpha}^{\star}+\frac{1}{2} \eta_{\alpha \beta \gamma} K^{\star \beta} \wedge K^{\star \gamma}$	$R_{\alpha}^{\star} \equiv R_{\alpha}^{\{\} \star}-D K_{\alpha}^{\star}+\frac{1}{2} \eta_{\alpha \beta \gamma} K^{\star \beta} \wedge K^{\star \gamma}$	
mielke-geometrodynamics_213 E0640 (10.2.11)	213	■ 1.000	$\Lambda_{\mathrm{eff}}=-\left[4 \rho-\left(1+2(-1)^s\right) \varepsilon^2\right] / 4 \ell^2=\theta_{\mathrm{T}}^2\left[\left(9+2(-1)^s\right) \theta_{\mathrm{TL}}^2+8 \theta_{\mathrm{T}} \theta_{\mathrm{L}}\right] /(2 A \ell)^2$	$\Lambda_{\mathrm{eff}}=-\left[4 \rho-\left(1+2(-1)^s\right) \varepsilon^2\right] / 4 \ell^2=\theta_{\mathrm{T}}^2\left[\left(9+2(-1)^s\right) \theta_{\mathrm{TL}}^2+8 \theta_{\mathrm{T}} \theta_{\mathrm{L}}\right] /(2 A \ell)^2$	
mielke-geometrodynamics_213 E0642 (10.2.13)	213	■ 1.000	$d s^2=-N^2(r) d t^2+N^{-2}(r) d r^2+r^2\left[d \phi+N^{\phi}(r) d t\right]^2$	$d s^2=-N^2(r) d t^2+N^{-2}(r) d r^2+r^2\left[d \phi+N^{\phi}(r) d t\right]^2$	
mielke-geometrodynamics_213 E0643 (10.2.14)	213	■ 1.000	$N^2(r)=-M-\Lambda_{\mathrm{eff}} r^2+\frac{J^2}{4 r^2}, \quad N^{\phi}(r)=-\frac{J}{2 r^2},$	$N^2(r)=-M-\Lambda_{\mathrm{eff}} r^2+\frac{J^2}{4 r^2}, \quad N^{\phi}(r)=-\frac{J}{2 r^2},$	
mielke-geometrodynamics_214 E0647 (10.3.4)	214	■ 1.000	$S_{\alpha}=\frac{\ell}{\varepsilon} E_{\alpha}$	$S_{\alpha}=\frac{\ell}{\varepsilon} E_{\alpha}$	
mielke-geometrodynamics_215 E0648 (10.3.5)	215	■ 1.000	$H_{\alpha}^{\star}=\frac{\ell}{\varepsilon} H_{\alpha}$	$H_{\alpha}^{\star}=\frac{\ell}{\varepsilon} H_{\alpha}$	
mielke-geometrodynamics_215 E0649 (10.3.6)	215	■ 1.000	$T_{\alpha}=\frac{\varepsilon}{\ell} \eta_{\alpha}, \quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2} \eta_{\alpha}$	$T_{\alpha}=\frac{\varepsilon}{\ell} \eta_{\alpha}, \quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2} \eta_{\alpha},$	
mielke-geometrodynamics_215 E0650 (10.3.7)	215	■ 1.000	$H_{\alpha}^{\star}=\delta \ell H_{\alpha}+\frac{\gamma}{\ell} \vartheta_{\alpha}$	$H_{\alpha}^{\star}=\delta \ell H_{\alpha}+\frac{\gamma}{\ell} \vartheta_{\alpha}$	
mielke-geometrodynamics_215 E0651 (10.3.8)	215	■ 1.000	$\delta \ell D H_{\alpha}+\frac{\gamma}{\ell} T_{\alpha}-\frac{1}{2} \eta_{\alpha \beta} \wedge H^{\beta}=\tau_{\alpha}^{\star}$	$\delta \ell D H_{\alpha}+\frac{\gamma}{\ell} T_{\alpha}-\frac{1}{2} \eta_{\alpha \beta} \wedge H^{\beta}=\tau_{\alpha}^{\star}$	

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E0653 (10.3.10)	216	■ 1.000	$H_{\{\alpha\}^{\star}}=\frac{\gamma}{\ell(1+\delta\varepsilon)}\vartheta_{\alpha}=-\frac{\ell}{\varepsilon}H_{\alpha},$	$H_{\alpha}^{\star}=\frac{\gamma}{\ell(1+\delta\varepsilon)}\vartheta_{\alpha}=-\frac{\ell}{\varepsilon}H_{\alpha},$	$H_{\alpha}^{\star}=\frac{\gamma}{\ell(1+\delta\varepsilon)}\vartheta_{\alpha}=-\frac{\ell}{\varepsilon}H_{\alpha},$
mielke-geometrodynamics_E0659 (10.3.16)	217	■ 1.000	$H=\left\{ \right\}^{\star}\left(\vartheta^{\alpha}\wedge H_{\alpha}\right),$	$H=\left\{ \right\}^{\star}\left(\vartheta^{\alpha}\wedge H_{\alpha}\right),$	$H=\left\{ \right\}^{\star}\left(\vartheta^{\alpha}\wedge H_{\alpha}\right),$
mielke-geometrodynamics_E0660 (10.3.17)	217	■ 1.000	$t=\left\{ \right\}^{\star}\left(T^{\alpha}\wedge H_{\alpha}\right),\quad\mathscr{A}:=\left\{ \right\}^{\star}\left(\vartheta^{\alpha}\wedge T_{\alpha}\right),\quad h=\left\{ \right\}^{\star}\left(\eta^{\alpha}\wedge H_{\alpha}\right),$	$t=\left\{ \right\}^{\star}\left(T^{\alpha}\wedge H_{\alpha}\right),\quad\mathscr{A}:=\left\{ \right\}^{\star}\left(\vartheta^{\alpha}\wedge T_{\alpha}\right),\quad h=\left\{ \right\}^{\star}\left(\eta^{\alpha}\wedge H_{\alpha}\right),$	$t=\left\{ \right\}^{\star}\left(T^{\alpha}\wedge H_{\alpha}\right),\quad\mathscr{A}:=\left\{ \right\}^{\star}\left(\vartheta^{\alpha}\wedge T_{\alpha}\right),\quad h=\left\{ \right\}^{\star}\left(\eta^{\alpha}\wedge H_{\alpha}\right),$
mielke-geometrodynamics_E0663 (10.3.20)	217	■ 1.000	$ds^2=(-1)^sdt^2+d\mathscr{A}^2+dh^2$	$ds^2=(-1)^sdt^2+d\mathscr{A}^2+dh^2$	$ds^2=(-1)^sdt^2+d\mathscr{A}^2+dh^2$
mielke-geometrodynamics_E0665 (10.3.22)	218	■ 1.000	$D^{\star}\vartheta_{\alpha}:=\frac{\ell}{\gamma}\left[D^{\star}H_{\alpha}^{\star}-\delta\ell D^{\star}H_{\alpha}\right]$	$D^{\star}\vartheta_{\alpha}:=\frac{\ell}{\gamma}\left[D^{\star}H_{\alpha}^{\star}-\delta\ell D^{\star}H_{\alpha}\right]$	$D^{\star}\vartheta_{\alpha}:=\frac{\ell}{\gamma}\left[D^{\star}H_{\alpha}^{\star}-\delta\ell D^{\star}H_{\alpha}\right]$
mielke-geometrodynamics_E0668 (10.3.25)	219	■ 1.000	$D^{\star}\vartheta^{\alpha}\cong 0,$	$D^{\star}\vartheta^{\alpha}\cong 0,$	$D^{\star}\vartheta^{\alpha}\cong 0,$
mielke-geometrodynamics_E0672 (10.3.29)	220	■ 1.000	$P_{\alpha}=\int\check{\mathcal{E}}_{\alpha},\quad J_{\alpha}=\int\check{\mathcal{J}}_{\alpha}^{\star}.$	$P_{\alpha}=\int\check{\mathcal{E}}_{\alpha},\quad J_{\alpha}=\int\check{\mathcal{J}}_{\alpha}^{\star}.$	$P_{\alpha}=\int\check{\mathcal{E}}_{\alpha},\quad J_{\alpha}=\int\check{\mathcal{J}}_{\alpha}^{\star}.$
mielke-geometrodynamics_E0675 (10.3.32)	220	■ 1.000	$\check{\mathcal{J}}_{\alpha}^{\star}=-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$	$\check{\mathcal{J}}_{\alpha}^{\star}=-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$	$\check{\mathcal{J}}_{\alpha}^{\star}=-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$
mielke-geometrodynamics_E0677 (10.3.34)	220	■ 1.000	$\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}\star}=(-1)^s\frac{\chi}{2\ell}d\vartheta_{\alpha}-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$	$\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}\star}=(-1)^s\frac{\chi}{2\ell}d\vartheta_{\alpha}-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$	$\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}\star}=(-1)^s\frac{\chi}{2\ell}d\vartheta_{\alpha}-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$
mielke-geometrodynamics_E0679 (10.3.36)	221	■ 1.000	$\widetilde{C}_{\mathrm{L}}=(-1)^s\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{R}_{\alpha}^{\star}-\frac{1}{3!}\eta_{\alpha\beta\gamma}\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{\Gamma}^{\star\beta}\wedge\widetilde{\Gamma}^{\star\gamma},$	$\widetilde{C}_{\mathrm{L}}=(-1)^s\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{R}_{\alpha}^{\star}-\frac{1}{3!}\eta_{\alpha\beta\gamma}\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{\Gamma}^{\star\beta}\wedge\widetilde{\Gamma}^{\star\gamma},$	$\widetilde{C}_{\mathrm{L}}=(-1)^s\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{R}_{\alpha}^{\star}-\frac{1}{3!}\eta_{\alpha\beta\gamma}\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{\Gamma}^{\star\beta}\wedge\widetilde{\Gamma}^{\star\gamma},$
mielke-geometrodynamics_E0683 (10.3.41)	221	■ 1.000	$c=12\cdot 4\pi\theta_{\mathrm{L}},\quad\tilde{c}=12\cdot 4\pi\tilde{\theta}_{\mathrm{L}},$	$c=12\cdot 4\pi\theta_{\mathrm{L}},\quad\tilde{c}=12\cdot 4\pi\tilde{\theta}_{\mathrm{L}},$	$c=12\cdot 4\pi\theta_{\mathrm{L}},\quad\tilde{c}=12\cdot 4\pi\tilde{\theta}_{\mathrm{L}},$
mielke-geometrodynamics_E0684 (10.3.42)	222	■ 1.000	$\widetilde{L}=L+\lambda_{\alpha}\wedge T^{\alpha}.$	$\widetilde{L}=L+\lambda_{\alpha}\wedge T^{\alpha}.$	$\widetilde{L}=L+\lambda_{\alpha}\wedge T^{\alpha}.$
mielke-geometrodynamics_E0691 (10.5.2)	223	■ 1.000	$\left[\sigma^{\alpha},\sigma^{\beta}\right]=2i\eta^{\alpha\beta\gamma}\sigma_{\gamma}.$	$\left[\sigma^{\alpha},\sigma^{\beta}\right]=2i\eta^{\alpha\beta\gamma}\sigma_{\gamma}.$	$\left[\sigma^{\alpha},\sigma^{\beta}\right]=2i\eta^{\alpha\beta\gamma}\sigma_{\gamma}.$
mielke-geometrodynamics_E0692 (10.5.3)	224	■ 1.000	$\gamma_0=i\sigma^2,\quad\gamma_1=\sigma^1,\quad\gamma_2=\sigma^3.$	$\gamma_0=i\sigma^2,\quad\gamma_1=\sigma^1,\quad\gamma_2=\sigma^3.$	$\gamma_0=i\sigma^2,\quad\gamma_1=\sigma^1,\quad\gamma_2=\sigma^3.$
mielke-geometrodynamics_E0693 (10.5.4)	224	■ 1.000	$\gamma_{\alpha}\gamma_{\beta}=g_{\alpha\beta}\mathbf{1}+\eta_{\alpha\beta\nu}\gamma^{\nu},$	$\gamma_{\alpha}\gamma_{\beta}=g_{\alpha\beta}\mathbf{1}+\eta_{\alpha\beta\nu}\gamma^{\nu},$	$\gamma_{\alpha}\gamma_{\beta}=g_{\alpha\beta}\mathbf{1}+\eta_{\alpha\beta\nu}\gamma^{\nu},$
mielke-geometrodynamics_E0695 (10.5.6)	224	■ 1.000	$i^{\star}\gamma\wedge D\psi-m\psi\eta-\frac{i}{2}\left(D^{\star}\gamma\right)\psi=0$	$i^{\star}\gamma\wedge D\psi-m\psi\eta-\frac{i}{2}\left(D^{\star}\gamma\right)\psi=0$	$i^{\star}\gamma\wedge D\psi-m\psi\eta-\frac{i}{2}\left(D^{\star}\gamma\right)\psi=0$
mielke-geometrodynamics_E0697 (10.5.8)	224	■ 1.000	$\hat{D}:=D-\frac{1}{2}T,$	$\hat{D}:=D-\frac{1}{2}T,$	$\hat{D}:=D-\frac{1}{2}T,$
mielke-geometrodynamics_E0698 (10.5.9)	224	■ 1.000	$i^{\star}\gamma\wedge\hat{D}\psi-m\eta\psi=0,$	$i^{\star}\gamma\wedge\hat{D}\psi-m\eta\psi=0,$	$i^{\star}\gamma\wedge\hat{D}\psi-m\eta\psi=0,$
mielke-geometrodynamics_E0699 (10.5.10)	224	■ 1.000	$\hat{R}=R-\frac{1}{2}dT=\frac{i}{4}R^{\alpha\beta}\sigma_{\alpha\beta}-\frac{1}{2}dT$	$\hat{R}=R-\frac{1}{2}dT=\frac{i}{4}R^{\alpha\beta}\sigma_{\alpha\beta}-\frac{1}{2}dT$	$\hat{R}=R-\frac{1}{2}dT=\frac{i}{4}R^{\alpha\beta}\sigma_{\alpha\beta}-\frac{1}{2}dT$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_225 EQ0703 (10.6.2)	225	■ 1.000	$d^{*}\{*\} F\text{-}\mathrm{m}_{\{\text{photon}\}} F=0$	$d^{*}F-m_{\mathrm{photon}}F=0$	$d^{*}F-m_{\mathrm{photon}}F=0$
mielke-geometrodynamics_226 EQ0705 (10.7.1)	226	■ 1.000	$X\text{:}=\mathrm{x}_{\{\alpha\}}\vartheta^{\{\alpha\}},\quad Y\text{:}=\mathrm{y}_{\{\alpha\}}\vartheta^{\{\alpha\}}.$	$X:=x_{\alpha}\vartheta^{\alpha},\quad Y:=y_{\alpha}\vartheta^{\alpha}.$	$X:=x_{\alpha}\vartheta^{\alpha},\quad Y:=y_{\alpha}\vartheta^{\alpha}.$
mielke-geometrodynamics_226 EQ0706 (10.7.2)	226	■ 1.000	$\Sigma_{\{\alpha\}}=\varepsilon\mathrm{x}_{\{\alpha\}}X\wedge Y,\quad \tau_{\{\alpha\}}^{\star}=\sigma\mathrm{y}_{\{\alpha\}}X\wedge Y,$	$\Sigma_{\alpha}=\varepsilon x_{\alpha}X\wedge Y,\quad \tau_{\alpha}^{\star}=\sigma y_{\alpha}X\wedge Y,$	$\Sigma_{\alpha}=\varepsilon x_{\alpha}X\wedge Y,\quad \tau_{\alpha}^{\star}=\sigma y_{\alpha}X\wedge Y,$
mielke-geometrodynamics_226 EQ0707 (10.7.3)	226	■ 1.000	$\vartheta^{\{\alpha\}}\wedge\Sigma_{\{\alpha\}}=0,\quad \vartheta^{\{\alpha\}}\wedge\tau_{\{\alpha\}}^{\star}=0$	$\vartheta^{\alpha}\wedge\Sigma_{\alpha}=0,\quad \vartheta^{\alpha}\wedge\tau_{\alpha}^{\star}=0$	$\vartheta^{\alpha}\wedge\Sigma_{\alpha}=0,\quad \vartheta^{\alpha}\wedge\tau_{\alpha}^{\star}=0$
mielke-geometrodynamics_227 EQ0708 (10.7.4)	227	■ 1.000	$R^{\{\alpha\}\{\beta\}}=\varepsilon\ell^2\mathrm{x}^{[\alpha}\mathrm{y}^{\beta]}\wedge X\wedge Y,$	$R^{\alpha\beta}=\varepsilon\ell^2x^{[\alpha}y^{\beta]}X\wedge Y,$	$R^{\alpha\beta}=\varepsilon\ell^2x^{[\alpha}y^{\beta]}X\wedge Y,$
mielke-geometrodynamics_228 EQ0711 (10.7.7)	228	■ 1.000	$\mathscr{A}\text{:}=\{*\}\left(\vartheta^{\{\alpha\}}\wedge\mathrm{T}_{\{\alpha\}}\right)\text{-}(-1)^{\{s\}}\frac{\varepsilon}{\ell^2}$	$\mathscr{A}:=*(\vartheta^{\alpha}\wedge T_{\alpha})=-(-1)^s\frac{\varepsilon}{\ell^2}$	$\mathscr{A}:=*(\vartheta^{\alpha}\wedge T_{\alpha})=-(-1)^s\frac{\varepsilon}{\ell^2}$
mielke-geometrodynamics_228 EQ0713 (10.8.2)	228	■ 1.000	$\gamma=\gamma_{\{\alpha\}}\vartheta^{\{\alpha\}}.$	$\gamma=\gamma_{\alpha}\vartheta^{\alpha}.$	$\gamma=\gamma_{\alpha}\vartheta^{\alpha}.$
mielke-geometrodynamics_228 EQ0714	228	■ 1.000	$\bar{\Psi}\wedge\Psi=0,\quad \bar{\Psi}\wedge\gamma_5\gamma^{\alpha}\Psi=0,\quad \bar{\Psi}\wedge\gamma_5\Psi=0.$	$\bar{\Psi}\wedge\Psi=0,\quad \bar{\Psi}\wedge\gamma_5\gamma^{\alpha}\Psi=0,\quad \bar{\Psi}\wedge\gamma_5\Psi=0.$	$\bar{\Psi}\wedge\Psi=0,\quad \bar{\Psi}\wedge\gamma_5\gamma^{\alpha}\Psi=0,\quad \bar{\Psi}\wedge\gamma_5\Psi=0.$
mielke-geometrodynamics_229 EQ0715 (10.9.1)	229	■ 1.000	$L_{\mathrm{RS}}=\frac{i}{4}(\bar{\Psi}\wedge D\Psi-\Psi\wedge\overline{D\Psi})+\frac{i}{4}m\bar{\Psi}\wedge\gamma\wedge\Psi,$	$L_{\mathrm{RS}}=\frac{i}{4}(\bar{\Psi}\wedge D\Psi-\Psi\wedge\overline{D\Psi})+\frac{i}{4}m\bar{\Psi}\wedge\gamma\wedge\Psi,$	$L_{\mathrm{RS}}=\frac{i}{4}(\bar{\Psi}\wedge D\Psi-\Psi\wedge\overline{D\Psi})+\frac{i}{4}m\bar{\Psi}\wedge\gamma\wedge\Psi,$
mielke-geometrodynamics_229 EQ0719 (10.9.5)	229	■ 1.000	$\Sigma_{\{\alpha\}}=-\frac{i}{4}m\bar{\Psi}\wedge\gamma_{\alpha}\Psi\text{-}(-1)^{\{s\}}2m\tau_{\{\alpha\}}^{\star}$	$\Sigma_{\alpha}=-\frac{i}{4}m\bar{\Psi}\wedge\gamma_{\alpha}\Psi=-(-1)^s2m\tau_{\alpha}^{\star}$	$\Sigma_{\alpha}=-\frac{i}{4}m\bar{\Psi}\wedge\gamma_{\alpha}\Psi=-(-1)^s2m\tau_{\alpha}^{\star},$
mielke-geometrodynamics_230 EQ0721 (10.10.1)	230	■ 1.000	$L\text{=}\mathrm{L}_{\{\infty\}}\left(\vartheta^{\{\alpha\}},\Gamma_{\{\alpha\}}^{\star},\Psi\right)\text{=}\mathrm{L}_{\{\mathrm{MB}\}}\text{+}\mathrm{L}_{\{\Psi\}}$	$L=L_{\infty}(\vartheta^{\alpha},\Gamma_{\alpha}^{\star},\Psi)=L_{\mathrm{MB}}+L_{\Psi}$	$L=L_{\infty}(\vartheta^{\alpha},\Gamma_{\alpha}^{\star},\Psi)=L_{\mathrm{MB}}+L_{\Psi}$
mielke-geometrodynamics_230 EQ0726 (10.10.6)	230	■ 1.000	$\frac{2}{i}\frac{\delta L}{\delta\Psi}=\mathrm{D}\Psi+\frac{1}{2}m\gamma\wedge\Psi\cong 0$	$\frac{2}{i}\frac{\delta L}{\delta\Psi}=D\Psi+\frac{1}{2}m\gamma\wedge\Psi\cong 0$	$\frac{2}{i}\frac{\delta L}{\delta\Psi}=D\Psi+\frac{1}{2}m\gamma\wedge\Psi\cong 0$
mielke-geometrodynamics_231 EQ0729 (10.10.9)	231	■ 1.000	$\gamma\wedge\gamma=-2\gamma^{\alpha}\eta_{\alpha}$	$\gamma\wedge\gamma=-2\gamma^{\alpha}\eta_{\alpha}$	$\gamma\wedge\gamma=-2\gamma^{\alpha}\eta_{\alpha}$
mielke-geometrodynamics_231 EQ0730 (10.10.10)	231	■ 1.000	$i\left[\left(\frac{\theta_{\mathrm{TL}}}{\ell}-1\right)R_{\alpha}^{\star}+\left(\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right)T_{\alpha}+\frac{m^2}{2}\eta_{\alpha}\right]\wedge\gamma^{\alpha}\Psi=0,$	$i\left[\left(\frac{\theta_{\mathrm{TL}}}{\ell}-1\right)R_{\alpha}^{\star}+\left(\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right)T_{\alpha}+\frac{m^2}{2}\eta_{\alpha}\right]\wedge\gamma^{\alpha}\Psi=0,$	$i\left[\left(\frac{\theta_{\mathrm{TL}}}{\ell}-1\right)R_{\alpha}^{\star}+\left(\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right)T_{\alpha}+\frac{m^2}{2}\eta_{\alpha}\right]\wedge\gamma^{\alpha}\Psi=0,$
mielke-geometrodynamics_231 EQ0731 (10.10.11)	231	■ 1.000	$\begin{aligned} & i\left[\left(\theta_{\mathrm{L}}+\frac{\theta_{\mathrm{TL}}}{\ell}-1\right)R_{\alpha}^{\star}+\left((-1)^s\frac{\theta_{\mathrm{TL}}}{\ell}+\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right)T_{\alpha}+\frac{1}{2}\left(\frac{\theta_{\mathrm{T}}}{\ell^2}+m^2\right)\eta_{\alpha}+\tau_{\alpha}^{\star}\right]\wedge\gamma^{\alpha}\Psi\simeq 0 \\ & \quad +\tau_{\alpha}^{\star}\wedge\gamma^{\alpha}\Psi\simeq 0 \end{aligned}$	$i\left[\left(\theta_{\mathrm{L}}+\frac{\theta_{\mathrm{TL}}}{\ell}-1\right)R_{\alpha}^{\star}+\left((-1)^s\frac{\theta_{\mathrm{TL}}}{\ell}+\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right)T_{\alpha}+\frac{1}{2}\left(\frac{\theta_{\mathrm{T}}}{\ell^2}+m^2\right)\eta_{\alpha}+\tau_{\alpha}^{\star}\right]\wedge\gamma^{\alpha}\Psi\simeq 0$	$i\left[\left(\theta_{\mathrm{L}}+\frac{\theta_{\mathrm{TL}}}{\ell}-1\right)R_{\alpha}^{\star}+\left((-1)^s\frac{\theta_{\mathrm{TL}}}{\ell}+\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right)T_{\alpha}+\frac{1}{2}\left(\frac{\theta_{\mathrm{T}}}{\ell^2}+m^2\right)\eta_{\alpha}+\tau_{\alpha}^{\star}\right]\wedge\gamma^{\alpha}\Psi\simeq 0,\quad (10.10)$
mielke-geometrodynamics_232 EQ0732 (10.10.12)	232	■ 1.000	$\theta_{\mathrm{T}}\simeq -m^2\ell^2,\theta_{\mathrm{TL}}\simeq\frac{(-1)^s}{2}m(2m+1)\ell,\theta_{\mathrm{L}}\simeq 1-\frac{\theta_{\mathrm{TL}}}{\ell}=1-\frac{(-1)^s}{2}m(2m+1)$	$\theta_{\mathrm{T}}\simeq -m^2\ell^2,\theta_{\mathrm{TL}}\simeq\frac{(-1)^s}{2}m(2m+1)\ell,\theta_{\mathrm{L}}\simeq 1-\frac{\theta_{\mathrm{TL}}}{\ell}=1-\frac{(-1)^s}{2}m(2m+1)$	$\theta_{\mathrm{T}}\simeq -m^2\ell^2,\theta_{\mathrm{TL}}\simeq\frac{(-1)^s}{2}m(2m+1)\ell,\theta_{\mathrm{L}}\simeq 1-\frac{\theta_{\mathrm{TL}}}{\ell}=1-\frac{(-1)^s}{2}m(2m+1)$
mielke-geometrodynamics_232 EQ0733 (10.10.13)	232	■ 1.000	$\vartheta_{\alpha}=\left(-1\right)^s\ell\Gamma_{\alpha}^{\star}+\overline{\sigma}\gamma_{\alpha}\Psi.$	$\vartheta_{\alpha}=(-1)^s\ell\Gamma_{\alpha}^{\star}+\bar{\sigma}\gamma_{\alpha}\Psi.$	$\vartheta_{\alpha}=(-1)^s\ell\Gamma_{\alpha}^{\star}+\bar{\sigma}\gamma_{\alpha}\Psi.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometroynamics_EQ0735 (11.1.2)	238	■ 1.000	$\gamma_{\alpha}\gamma_{\beta}+\gamma_{\beta}\gamma_{\alpha}=2\mathrm{o}_{\alpha\beta}\mathbb{1}$	$\gamma_{\alpha}\gamma_{\beta}+\gamma_{\beta}\gamma_{\alpha}=2\mathrm{o}_{\alpha\beta}\mathbb{1}$	$\gamma_{\alpha}\gamma_{\beta}+\gamma_{\beta}\gamma_{\alpha}=2\mathrm{o}_{\alpha\beta}\mathbb{1}$
mielke-geometroynamics_EQ0738 (11.1.5)	238	■ 1.000	$\mathrm{X}:=\mathrm{i}\mathrm{m}\mathrm{a}\mathrm{t}\mathrm{h}\ x^{\alpha}\gamma_{\alpha}\longleftarrow x^{\alpha}=-\frac{l}{N}\mathrm{Tr}\left(\gamma^{\alpha}X\right).$ $\mathrm{operatorname{Tr}}\left(\gamma^{\alpha}\right)\left(\gamma^{\alpha}\right)$.	$X:=\imath x^{\alpha}\gamma_{\alpha}\longleftrightarrow x^{\alpha}=-\frac{l}{N}\mathrm{Tr}\left(\gamma^{\alpha}X\right).$	$X:=\imath x^{\alpha}\gamma_{\alpha}\longleftrightarrow x^{\alpha}=-\frac{l}{N}\mathrm{Tr}(\gamma^{\alpha}X).$
mielke-geometroynamics_EQ0739 (11.1.6)	238	■ 1.000	$\mathrm{operatorname{det}}\left(\mathrm{X}\right)=-\mathrm{operatorname{o}}_{\alpha\beta}\gamma^{\alpha}\gamma^{\beta},$	$\det X=-\mathrm{o}_{\alpha\beta}x^{\alpha}x^{\beta},$	$\det X=-\mathrm{o}_{\alpha\beta}x^{\alpha}x^{\beta},$
mielke-geometroynamics_EQ0741 (11.1.8)	238	■ 1.000	$\mathrm{X}-\mathrm{X}'=\mathrm{S}\mathrm{X}\mathrm{S}^{-1},\quad\mathrm{S}\in\widetilde{\mathrm{SO}}_{\mathrm{o}}(s,n-s).$	$X-X'=SXS^{-1},\quad S\in\widetilde{SO}_{\mathrm{o}}(s,n-s).$	$X-X'=SXS^{-1},\quad S\in\widetilde{SO}_{\mathrm{o}}(s,n-s).$
mielke-geometroynamics_EQ0743 (11.1.10)	239	■ 1.000	$\sigma^{\alpha\beta}:=\frac{i}{4}\left[\gamma^{\alpha},\gamma^{\beta}\right],\quad\Lambda\left(\sigma^{\alpha\beta}\right)=L^{\alpha\beta},$	$\sigma^{\alpha\beta}:=\frac{i}{4}\left[\gamma^{\alpha},\gamma^{\beta}\right],\quad\Lambda\left(\sigma^{\alpha\beta}\right)=L^{\alpha\beta},$	$\sigma^{\alpha\beta}:=\frac{i}{4}[\gamma^{\alpha},\gamma^{\beta}],\quad\Lambda(\sigma^{\alpha\beta})=L^{\alpha\beta},$
mielke-geometroynamics_EQ0744 (11.1.11)	239	■ 1.000	$\mathrm{L}_{\mathrm{circ}}^{\mathrm{g}}(\mathrm{M}):=\mathrm{P}\left(\mathrm{M},\mathrm{S}_{\mathrm{O}_{\mathrm{circ}}}(s,n-s),\pi,\delta\right)$	$L_{\mathrm{o}}^g(M):=P\left(M,SO_{\mathrm{o}}(s,n-s),\pi,\delta\right)$	$L_{\mathrm{o}}^g(M):=P(M,SO_{\mathrm{o}}(s,n-s),\pi,\delta)$
mielke-geometroynamics_EQ0745 (11.1.12)	239	■ 1.000	$\widetilde{\mathrm{L}}(\mathrm{M}):=\mathrm{P}\left(\mathrm{M},\widetilde{\mathrm{S}}_{\mathrm{O}_{\mathrm{circ}}}(s,n-2),\widetilde{\pi},\widetilde{\delta}\right)\ni\widetilde{p}$	$\widetilde{L}(M):=P\left(M,\widetilde{SO}_{\mathrm{o}}(s,n-2),\widetilde{\pi},\widetilde{\delta}\right)\ni\tilde{p}$	$\widetilde{L}(M):=P(M,\widetilde{SO}_{\mathrm{o}}(s,n-2),\widetilde{\pi},\widetilde{\delta})\ni\tilde{p}$
mielke-geometroynamics_EQ0746 (11.1.13)	239	■ 1.000	$f(\widetilde{p}S)=f(\widetilde{p})\Lambda(S),S\in\widetilde{\mathrm{SO}}_{\mathrm{o}}(s,n-2).$	$f(\tilde{p}S)=f(\tilde{p})\Lambda(S),S\in\widetilde{SO}_{\mathrm{o}}(s,n-2).$	$f(\tilde{p}S)=f(\tilde{p})\Lambda(S),S\in\widetilde{SO}_{\mathrm{o}}(s,n-2).$
mielke-geometroynamics_EQ0748 (11.1.15)	240	■ 1.000	$\widetilde{\mathrm{L}}(\mathrm{M}^4)=\mathrm{M}^4\times\mathrm{SL}(2,\mathbb{C})$	$\tilde{L}\left(M^4\right)=M^4\times SL(2,\mathbb{C})$	$\tilde{L}(M^4)=M^4\times SL(2,\mathbb{C})$
mielke-geometroynamics_EQ0749 (11.1.16)	241	■ 1.000	$V^{\bar{s}}:=V\left(\mathrm{M},\mathbb{C}^N,\rho^{\bar{s}}\left(\widetilde{\mathrm{SO}}_{\mathrm{o}}(s,n-s)\right)\subset\mathrm{GL}(N,\mathbb{C}),\tilde{L}(\mathrm{M})\right)$	$V^{\bar{s}}:=V\left(M,\mathbb{C}^N,\rho^{\bar{s}}\left(\widetilde{SO}_{\mathrm{o}}(s,n-s)\right)\subset GL(N,\mathbb{C}),\tilde{L}(M)\right)$	$V^{\bar{s}}:=V(M,\mathbb{C}^N,\rho^{\bar{s}}(\widetilde{SO}_{\mathrm{o}}(s,n-s))\subset GL(N,\mathbb{C}),\tilde{L}(M))$
mielke-geometroynamics_EQ0751 (11.1.18)	241	■ 1.000	$D^{\frac{1}{2}}:=D^{\left(-\frac{1}{2},\frac{3}{2}\right)}\oplus D^{\left(\frac{1}{2},\frac{3}{2}\right)}\left(\widetilde{\mathrm{SO}}_{\mathrm{o}}(1,3)\right)$	$D^{\frac{1}{2}}:=D^{\left(-\frac{1}{2},\frac{3}{2}\right)}\oplus D^{\left(\frac{1}{2},\frac{3}{2}\right)}\left(\widetilde{SO}_{\mathrm{o}}(1,3)\right)$	$D^{\frac{1}{2}}:=D^{\left(-\frac{1}{2},\frac{3}{2}\right)}\oplus D^{\left(\frac{1}{2},\frac{3}{2}\right)}\left(\widetilde{SO}_{\mathrm{o}}(1,3)\right)$
mielke-geometroynamics_EQ0754 (11.1.21)	241	■ 1.000	$\psi\rightarrow^{S^{-1}}\psi:=S^{-1}\psi$	$\psi\rightarrow^{S^{-1}}\psi:=S^{-1}\psi$	$\psi\rightarrow^{S^{-1}}\psi:=S^{-1}\psi$
mielke-geometroynamics_EQ0755 (11.1.22)	242	■ 1.000	$\begin{gathered} S(p)\in\mathscr{G}_S\approx C^{\infty}\left(\tilde{L}(M)\times_{\mathrm{Ad}}\widetilde{SO}_{\mathrm{o}}(s,n-s)\right)\\ \Lambda\left(S_{(p)}\right)=G(p)\in\mathscr{G}_p \end{gathered}$	$S(p)\in\mathscr{G}_S\approx C^{\infty}\left(\tilde{L}(M)\times_{\mathrm{Ad}}\widetilde{SO}_{\mathrm{o}}(s,n-s)\right)\\ \Lambda\left(S_{(p)}\right)=G(p)\in\mathscr{G}_p$	$S(p)\in\mathscr{G}_S\approx C^{\infty}(\tilde{L}(M)\times_{\mathrm{Ad}}\widetilde{SO}_{\mathrm{o}}(s,n-s))\\ \Lambda(S_{(p)})=G(p)\in\mathscr{G}_p$
mielke-geometroynamics_EQ0757 (11.1.24)	242	■ 1.000	$S^{-1}(p)\gamma_{\alpha}S(p)=\gamma_{\beta}G_{\alpha}^{\beta}(p),\quad[G_{\alpha}^{\beta}(p)]\in\mathscr{G}_p.$	$S^{-1}(p)\gamma_{\alpha}S(p)=\gamma_{\beta}G_{\alpha}^{\beta}(p),\quad[G_{\alpha}^{\beta}(p)]\in\mathscr{G}_p.$	$S^{-1}(p)\gamma_{\alpha}S(p)=\gamma_{\beta}G_{\alpha}^{\beta}(p),\quad[G_{\alpha}^{\beta}(p)]\in\mathscr{G}_p.$
mielke-geometroynamics_EQ0762 (11.1.29)	243	■ 1.000	$\bar{\psi}:=\psi^{+}\gamma^0,$	$\bar{\psi}:=\psi^{+}\gamma^0,$	$\overline{\psi}:=\psi^{+}\gamma^0,$
mielke-geometroynamics_EQ0763 (11.1.30)	243	■ 1.000	$\bar{\sigma}^{\alpha\beta}:=\gamma^0\sigma^{\alpha\beta}+\gamma^0=\sigma^{\alpha\beta},\quad\bar{\gamma}^{\alpha}:=\gamma^0\gamma^{\alpha}+\gamma^0=\gamma^{\alpha}.$	$\bar{\sigma}^{\alpha\beta}:=\gamma^0\sigma^{\alpha\beta}+\gamma^0=\sigma^{\alpha\beta},\quad\bar{\gamma}^{\alpha}:=\gamma^0\gamma^{\alpha}+\gamma^0=\gamma^{\alpha}.$	$\overline{\sigma}^{\alpha\beta}:=\gamma^0\sigma^{\alpha\beta}+\gamma^0=\sigma^{\alpha\beta},\quad\overline{\gamma}^{\alpha}:=\gamma^0\gamma^{\alpha}+\gamma^0=\gamma^{\alpha}.$
mielke-geometroynamics_EQ0764 (11.1.31)	243	■ 1.000	$D\psi=d\psi+i\widetilde{\omega}\psi,$	$D\psi=d\psi+i\widetilde{\omega}\psi,$	$D\psi=d\psi+i\widetilde{\omega}\psi,$
mielke-geometroynamics_EQ0765 (11.1.32)	243	■ 1.000	$\Lambda(i\widetilde{\omega})=\omega^{\mathfrak{g}}$	$\Lambda(i\widetilde{\omega})=\omega^{\mathfrak{g}}$	$\Lambda(i\widetilde{\omega})=\omega^{\mathfrak{g}}$
mielke-geometroynamics_EQ0771 (11.1.38)	244	■ 1.000	$\gamma\longrightarrow^{S^{-1}}\gamma=S^{-1}\gamma S$	$\gamma\longrightarrow^{S^{-1}}\gamma=S^{-1}\gamma S$	$\gamma\longrightarrow^{S^{-1}}\gamma=S^{-1}\gamma S$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00775 (11.2.4)	246	■ 1.000	$\overline{\widetilde{\omega}}=\gamma^0\widetilde{\omega}^+\gamma^0=\widetilde{\omega}$	$\bar{\tilde{\omega}}=\gamma^0\tilde{\omega}^+\gamma^0=\tilde{\omega}$	$\bar{\tilde{\omega}}=\gamma^0\tilde{\omega}^+\gamma^0=\tilde{\omega}$
mielke-geometrodynamics_E00777 (11.2.6)	246	■ 1.000	$\tau_s=i\left(\bar{\psi}^*\gamma\psi\right)$.	$\tau_s=i\left(\bar{\psi}^*\gamma\psi\right)$.	$\tau_s=i(\bar{\psi}^*\gamma\psi)$.
mielke-geometrodynamics_E00779 (11.2.7)	247	■ 1.000	$d\tau_s=0$	$d\tau_s=0$	$d\tau_s=0$
mielke-geometrodynamics_E00781 (11.2.9)	247	■ 1.000	$(\psi,\chi)=i\int_{H^{n-1}}\psi^+\chi d^{n-1}x$	$(\psi,\chi)=i\int_{H^{n-1}}\psi^+\chi d^{n-1}x$	$(\psi,\chi)=i\int_{H^{n-1}}\psi^+\chi d^{n-1}x$.
mielke-geometrodynamics_E00782 (11.3.1)	247	■ 1.000	$\widetilde{\omega}=\widetilde{\omega}^{\{\}}-\widetilde{K},\quad\Lambda(i\widetilde{K})=K$	$\tilde{\omega}=\tilde{\omega}^{\{\}}-\tilde{K},\quad\Lambda(i\tilde{K})=K$	$\tilde{\omega}=\tilde{\omega}^{(\cdot)}-\tilde{K},\quad\Lambda(i\tilde{K})=K$
mielke-geometrodynamics_E00785 (11.3.4)	248	■ 1.000	$[\Theta,\vartheta]=\ell^{*2}\tau_s$.	$[\Theta,\vartheta]=\ell^{*2}\tau_s$.	$[\Theta,\vartheta]=\ell^{*2}\tau_s$.
mielke-geometrodynamics_E00790 (11.4.2)	252	■ 1.000	$\begin{aligned} \left(\bar{\psi}i\gamma^5\gamma_\mu\psi\right)\left(\bar{\psi}i\gamma^5\gamma^\mu\psi\right)&=\left(\bar{\psi}\gamma_\mu\psi\right)\left(\bar{\psi}\gamma^\mu\psi\right)\\ &=\left(\bar{\psi}\psi\right)^2-\left(\bar{\psi}\gamma^5\psi\right)^2. \end{aligned}$	$(\bar{\psi}i\gamma^5\gamma_\mu\psi)(\bar{\psi}i\gamma^5\gamma^\mu\psi)=(\bar{\psi}\gamma_\mu\psi)(\bar{\psi}\gamma^\mu\psi)= (\bar{\psi}\psi)^2-(\bar{\psi}\gamma^5\psi)^2$.	$(\bar{\psi}i\gamma^5\gamma_\mu\psi)(\bar{\psi}i\gamma^5\gamma^\mu\psi)=(\bar{\psi}\gamma_\mu\psi)(\bar{\psi}\gamma^\mu\psi)= (\bar{\psi}\psi)^2-(\bar{\psi}\gamma^5\psi)^2$.
mielke-geometrodynamics_E00793 (11.4.5)	253	■ 1.000	$\rho:=\frac{\sqrt{2\pi}}{\ell^*}\,r,$	$\rho:=\frac{\sqrt{2\pi}}{\ell^*}\,r,$	$\rho:=\frac{\sqrt{2\pi}}{\ell^*}\,r,$
mielke-geometrodynamics_E00794 (11.4.6)	253	■ 1.000	$\bar{g}_{kl}\longleftarrow g_{kl}=e^{\mu}\bar{g}_{kl}$.	$\bar{g}_{kl}\longleftarrow g_{kl}=e^{\mu}\bar{g}_{kl}$.	$\bar{g}_{kl}\longleftarrow g_{kl}=e^{\mu}\bar{g}_{kl}$.
mielke-geometrodynamics_E00795 (11.4.7)	253	■ 1.000	$\bar{E}_{\cdot k}^{\alpha}\longleftarrow E_{\cdot k}^{\alpha}=e^{\mu/2}\bar{E}_{\cdot k}^{\alpha}$	$\bar{E}_{\cdot k}^{\alpha}\longleftarrow E_{\cdot k}^{\alpha}=e^{\mu/2}\bar{E}_{\cdot k}^{\alpha}$	$\bar{E}_{\cdot k}^{\alpha}\longleftarrow E_{\cdot k}^{\alpha}=e^{\mu/2}\bar{E}_{\cdot k}^{\alpha}$
mielke-geometrodynamics_E00796 (11.4.8)	253	■ 1.000	$\overline{\left\{\begin{array}{c}\alpha\\kl\end{array}\right\}}+\frac{1}{2}\left(\delta_k^{\alpha}\partial_l+\delta_l^{\alpha}\partial_k-g_{kl}\partial^{\alpha}\right)\mu$	$\left\{\begin{array}{c}\alpha\\kl\end{array}\right\}=\overline{\left\{\begin{array}{c}\alpha\\kl\end{array}\right\}}+\frac{1}{2}\left(\delta_k^{\alpha}\partial_l+\delta_l^{\alpha}\partial_k-g_{kl}\partial^{\alpha}\right)\mu$	$\left\{\begin{array}{c}\alpha\\kl\end{array}\right\}=\overline{\left\{\begin{array}{c}\alpha\\kl\end{array}\right\}}+\frac{1}{2}\left(\delta_k^{\alpha}\partial_l+\delta_l^{\alpha}\partial_k-g_{kl}\partial^{\alpha}\right)\mu$
mielke-geometrodynamics_E00797 (11.4.9)	253	■ 1.000	$\Gamma_k=\bar{\Gamma}_k-\frac{1}{4}\delta_k^{[\alpha}\partial^{\beta]}\mu\sigma_{\alpha\beta}$.	$\Gamma_k=\bar{\Gamma}_k-\frac{1}{4}\delta_k^{[\alpha}\partial^{\beta]}\mu\sigma_{\alpha\beta}$.	$\Gamma_k=\bar{\Gamma}_k-\frac{1}{4}\delta_k^{[\alpha}\partial^{\beta]}\mu\sigma_{\alpha\beta}$.
mielke-geometrodynamics_E00801 (11.4.13)	254	■ 1.000	$\begin{aligned} \bar{\Gamma}_0&=\frac{i}{4}e^{(\nu-\mu)/2}\partial_r(\nu-\mu)\gamma_0\gamma_1\\ \bar{\Gamma}_1&=0,\quad\bar{\Gamma}_2=\frac{i}{2}\gamma_2\gamma_1\\ \bar{\Gamma}_3&=\frac{i}{2}(\sin\vartheta\gamma_3\gamma_1+\cos\vartheta\gamma_3\gamma_2) \end{aligned}$	$\begin{aligned} \bar{\Gamma}_0&=\frac{i}{4}e^{(\nu-\mu)/2}\partial_r(\nu-\mu)\gamma_0\gamma_1\\ \bar{\Gamma}_1&=0,\quad\bar{\Gamma}_2=\frac{i}{2}\gamma_2\gamma_1\\ \bar{\Gamma}_3&=\frac{i}{2}(\sin\vartheta\gamma_3\gamma_1+\cos\vartheta\gamma_3\gamma_2) \end{aligned}$	$\begin{aligned} \bar{\Gamma}_0&=\frac{i}{4}e^{(v-\mu)/2}\partial_r(v-\mu)\gamma_0\gamma_1\\ \bar{\Gamma}_1&=0,\quad\bar{\Gamma}_2=\frac{i}{2}\gamma_2\gamma_1\\ \bar{\Gamma}_3&=\frac{i}{2}(\sin\vartheta\gamma_3\gamma_1+\cos\vartheta\gamma_3\gamma_2) \end{aligned}$
mielke-geometrodynamics_E00804 (11.4.16)	255	■ 1.000	$\chi_{\kappa}^m=\sum_{\bar{m}=\pm 1/2}C\left(l\frac{1}{2}j;m-\bar{m},\bar{m}\right)Y_l^{m-\bar{m}}(\vartheta,\phi)\chi^{\bar{m}}$.	$\chi_{\kappa}^m=\sum_{\bar{m}=\pm 1/2}C\left(l\frac{1}{2}j;m-\bar{m},\bar{m}\right)Y_l^{m-\bar{m}}(\vartheta,\phi)\chi^{\bar{m}}$.	$\chi_{\kappa}^m=\sum_{\bar{m}=\pm 1/2}C\left(l\frac{1}{2}j;m-\bar{m},\bar{m}\right)Y_l^{m-\bar{m}}(\vartheta,\phi)\chi^{\bar{m}}$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E09808 (11.4.20)	255	■ 1.000	$\frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } \chi_{\kappa}^m = -\chi_{-\kappa}^m$	$\frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } \chi_{\kappa}^m = -\chi_{-\kappa}^m$	$\frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } \chi_{\kappa}^m = -\chi_{-\kappa}^m$
mielke-geometrodynamics_E09814 (11.4.27)	257	■ 1.000	$\beta = \frac{M^*}{2m}, \quad M^* := \frac{\sqrt{8\pi} \hbar}{c \ell^*}.$	$\beta = \frac{M^*}{2m}, \quad M^* := \frac{\sqrt{8\pi} \hbar}{c \ell^*}.$	$\beta = \frac{M^*}{2m}, \quad M^* := \frac{\sqrt{8\pi} \hbar}{c \ell^*}.$
mielke-geometrodynamics_E09816 (11.4.30)	257	■ 1.000	$F^{\prime} + \frac{2}{\rho} F^{\prime} - \frac{2}{\rho^2} F \simeq \frac{1 - \omega^2}{\beta^2} F$	$F'' + \frac{2}{\rho} F' - \frac{2}{\rho^2} F \simeq \frac{1 - \omega^2}{\beta^2} F$	$F'' + \frac{2}{\rho} F' - \frac{2}{\rho^2} F \simeq \frac{1 - \omega^2}{\beta^2} F$
mielke-geometrodynamics_E09817 (11.4.31)	257	■ 1.000	$F \sim C_{\infty} \left(\frac{1}{\bar{\rho}} + \frac{1}{\bar{\rho}^2} \right) e^{-\bar{\rho}}$	$F \sim C_{\infty} \left(\frac{1}{\bar{\rho}} + \frac{1}{\bar{\rho}^2} \right) e^{-\bar{\rho}}$	$F \sim C_{\infty} \left(\frac{1}{\bar{\rho}} + \frac{1}{\bar{\rho}^2} \right) e^{-\bar{\rho}}$
mielke-geometrodynamics_E09818 (11.4.32)	257	■ 1.000	$H \sim C_{\infty} \sqrt{\frac{1 + \omega}{1 - \omega}} \frac{1}{\bar{\rho}} e^{-\bar{\rho}}$	$H \sim C_{\infty} \sqrt{\frac{1 + \omega}{1 - \omega}} \frac{1}{\bar{\rho}} e^{-\bar{\rho}}$	$H \sim C_{\infty} \sqrt{\frac{1 + \omega}{1 - \omega}} \frac{1}{\bar{\rho}} e^{-\bar{\rho}}$
mielke-geometrodynamics_E09819 (11.4.33)	257	■ 1.000	$\bar{\rho} := \frac{1}{\beta} \sqrt{1 - \omega^2} \rho$	$\bar{\rho} := \frac{1}{\beta} \sqrt{1 - \omega^2} \rho$	$\bar{\rho} := \frac{1}{\beta} \sqrt{1 - \omega^2} \rho$
mielke-geometrodynamics_E09821 (11.4.35)	258	■ 1.000	$F(\rho) \simeq C_0 f^1(\bar{\rho}) \frac{1}{\cosh \bar{\rho}}$	$F(\rho) \simeq C_0 f^1(\bar{\rho}) \frac{1}{\cosh \bar{\rho}}$	$F(\rho) \simeq C_0 f^1(\bar{\rho}) \frac{1}{\cosh \bar{\rho}}$
mielke-geometrodynamics_E09822 (11.4.36)	258	■ 1.000	$H(\rho) \simeq C_0 \sqrt{\frac{1 + \omega}{1 - \omega}} f^0(\bar{\rho}) \frac{1}{\cosh \bar{\rho}}$	$H(\rho) \simeq C_0 \sqrt{\frac{1 + \omega}{1 - \omega}} f^0(\bar{\rho}) \frac{1}{\cosh \bar{\rho}}$	$H(\rho) \simeq C_0 \sqrt{\frac{1 + \omega}{1 - \omega}} f^0(\bar{\rho}) \frac{1}{\cosh \bar{\rho}}$
mielke-geometrodynamics_E09823 (11.4.37)	258	■ 1.000	$C_0 = (1 - \omega) \sqrt{\frac{-2}{\varepsilon(\omega + 1)}}$	$C_0 = (1 - \omega) \sqrt{\frac{-2}{\varepsilon(\omega + 1)}}$	$C_0 = (1 - \omega) \sqrt{\frac{-2}{\varepsilon(\omega + 1)}}$
mielke-geometrodynamics_E09826 (11.4.40)	259	■ 1.000	$m_s = \frac{1}{c} \int_{H^3} \Sigma_{\mathrm{D}}^0$	$m_s = \frac{1}{c} \int_{H^3} \Sigma_{\mathrm{D}}^0$	$m_s = \frac{1}{c} \int_{H^3} \Sigma_{\mathrm{D}}^0$
mielke-geometrodynamics_E09827 (11.5.1)	262	■ 1.000	$Q := i e \int_{H^3} \bar{\psi}^* \gamma \psi = k e, \quad k = 0, 1, 2, \dots$	$Q := i e \int_{H^3} \bar{\psi}^* \gamma \psi = k e, \quad k = 0, 1, 2, \dots$	$Q := i e \int_{H^3} \bar{\psi}^* \gamma \psi = k e, \quad k = 0, 1, 2, \dots$
mielke-geometrodynamics_E09828 (11.5.2)	263	■ 1.000	$i \gamma \wedge^* d\psi(x) + e^2 \int_{y \in M} \bar{\psi}(y) \gamma \psi(y) D_{\mathrm{F}}(y - x) \wedge^* \gamma \psi(x) = m \psi(x) \eta$	$i \gamma \wedge^* d\psi(x) + e^2 \int_{y \in M} \bar{\psi}(y) \gamma \psi(y) D_{\mathrm{F}}(y - x) \wedge^* \gamma \psi(x) = m \psi(x) \eta$	$i \gamma \wedge^* d\psi(x) + e^2 \int_{y \in M} \bar{\psi}(y) \gamma \psi(y) D_{\mathrm{F}}(y - x) \wedge^* \gamma \psi(x) = m \psi(x) \eta$
mielke-geometrodynamics_E09830 (12.1.2)	269	■ 1.000	$\langle dj_5(x) \rangle = 2im\langle P \rangle - (1/96\pi^2) F \wedge F$	$\langle dj_5(x) \rangle = 2im\langle P \rangle - (1/96\pi^2) F \wedge F$	$\langle dj_5(x) \rangle = 2im\langle P \rangle - (1/96\pi^2) F \wedge F$
mielke-geometrodynamics_E09838 (12.2.6)	271	■ 1.000	$dj_5 = 2imP = 2im\bar{\psi}\gamma_5\psi\eta$	$dj_5 = 2imP = 2im\bar{\psi}\gamma_5\psi\eta$	$dj_5 = 2imP = 2im\bar{\psi}\gamma_5\psi\eta$
mielke-geometrodynamics_E09843 (12.2.11)	272	■ 1.000	$L_{a\psi\psi} = \frac{1}{2} d\theta \wedge j_5 = \frac{1}{2f_a} da \wedge \bar{\psi}^* \gamma \gamma_5 \psi,$	$L_{a\psi\psi} = \frac{1}{2} d\theta \wedge j_5 = \frac{1}{2f_a} da \wedge \bar{\psi}^* \gamma \gamma_5 \psi,$	$L_{a\psi\psi} = \frac{1}{2} d\theta \wedge j_5 = \frac{1}{2f_a} da \wedge \bar{\psi}^* \gamma \gamma_5 \psi,$
mielke-geometrodynamics_E09846 (12.2.14)	273	■ 1.000	$dj_5 \cong 4dC_{\mathrm{TT}} = \frac{2}{\ell^2} (T^\alpha \wedge T_\alpha + R_{\alpha\beta} \wedge \vartheta^\alpha \wedge \vartheta^\beta)$	$dj_5 \cong 4dC_{\mathrm{TT}} = \frac{2}{\ell^2} (T^\alpha \wedge T_\alpha + R_{\alpha\beta} \wedge \vartheta^\alpha \wedge \vartheta^\beta)$	$dj_5 \cong 4dC_{\mathrm{TT}} = \frac{2}{\ell^2} (T^\alpha \wedge T_\alpha + R_{\alpha\beta} \wedge \vartheta^\alpha \wedge \vartheta^\beta)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E08849 (12.3.2)	275	■ 1.000	$K\left(t, x, \not{D}^2\right)=(4 \pi)^{-n / 2} \sum_{k=0}^{\infty} t^{(k-n) / 2} K_k(x, \not{D}^2)$	$K\left(t, x, \not{D}^2\right)=(4 \pi)^{-n / 2} \sum_{k=0}^{\infty} t^{(k-n) / 2} K_k\left(x, \not{D}^2\right)$	$K\left(t, x, \not{D}^2\right)=(4 \pi)^{-n / 2} \sum_{k=0}^{\infty} t^{(k-n) / 2} K_k\left(x, \not{D}^2\right)$
mielke-geometrodynamics_E08852 (12.3.5)	276	■ 1.000	$-K_2 / t=\left(2 \ell^2 / t\right) d C_{\mathrm{TT}} \cong\left(\ell^2 / 2 t\right) d j_5=\left(i m \ell^2 / t\right) \bar{\psi} \gamma_5 \psi \sim \ell^2 M^2 m^4 \rightarrow 0 .$	$-K_2 / t=\left(2 \ell^2 / t\right) d C_{\mathrm{TT}} \cong\left(\ell^2 / 2 t\right) d j_5=\left(i m \ell^2 / t\right) \bar{\psi} \gamma_5 \psi \sim \ell^2 M^2 m^4 \rightarrow 0 .$	$-K_2 / t=\left(2 \ell^2 / t\right) d C_{\mathrm{TT}} \cong\left(\ell^2 / 2 t\right) d j_5=\left(i m \ell^2 / t\right) \bar{\psi} \gamma_5 \psi \sim \ell^2 M^2 m^4 \rightarrow 0 .$
mielke-geometrodynamics_E08856 (12.4.3)	277	■ 1.000	$\mathscr{G}^A \cong 0, \quad A=1,2,3$	$\mathscr{G}^A \cong 0, \quad A=1,2,3$	$\mathscr{G}^A \cong 0, \quad A=1,2,3$
mielke-geometrodynamics_E08858 (12.4.5)	278	■ 1.000	$\gamma:=\gamma_{\alpha} \vartheta^{\alpha}, \quad * \gamma=\gamma^{\alpha} \eta_{\alpha} .$ $\gamma=\gamma^{\alpha} \eta_{\alpha} \quad \eta_{\alpha}=\gamma_{\alpha} \vartheta^{\alpha} .$	$\gamma:=\gamma_{\alpha} \vartheta^{\alpha}, \quad * \gamma=\gamma^{\alpha} \eta_{\alpha} .$	$\gamma:=\gamma_{\alpha} \vartheta^{\alpha}, \quad * \gamma=\gamma^{\alpha} \eta_{\alpha} .$
mielke-geometrodynamics_E08859 (12.4.6)	278	■ 1.000	$\Theta:=D \gamma=T^{\alpha} \gamma_{\alpha}, \quad \Omega:=d \Gamma+\Gamma \wedge \Gamma=\frac{i}{4} R^{\alpha \beta} \sigma_{\alpha \beta}$	$\Theta:=D \gamma=T^{\alpha} \gamma_{\alpha}, \quad \Omega:=d \Gamma+\Gamma \wedge \Gamma=\frac{i}{4} R^{\alpha \beta} \sigma_{\alpha \beta}$	$\Theta:=D \gamma=T^{\alpha} \gamma_{\alpha}, \quad \Omega:=d \Gamma+\Gamma \wedge \Gamma=\frac{i}{4} R^{\alpha \beta} \sigma_{\alpha \beta}$
mielke-geometrodynamics_E08860 (12.4.7)	278	■ 1.000	$C_{\mathrm{RR}}:=-\operatorname{Tr}\left(\Gamma \wedge \Omega-\frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma\right) .$	$C_{\mathrm{RR}}:=-\operatorname{Tr}\left(\Gamma \wedge \Omega-\frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma\right) .$	$C_{\mathrm{RR}}:=-\operatorname{Tr}\left(\Gamma \wedge \Omega-\frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma\right) .$
mielke-geometrodynamics_E08862 (12.4.9)	279	■ 1.000	$C_{\mathrm{TT}}:=\frac{1}{8 \ell^2} \operatorname{Tr}\left(\gamma \wedge \Theta\right)=\frac{1}{2 \ell^2} \vartheta^{\alpha} \wedge T_{\alpha} .$	$C_{\mathrm{TT}}:=\frac{1}{8 \ell^2} \operatorname{Tr}\left(\gamma \wedge \Theta\right)=\frac{1}{2 \ell^2} \vartheta^{\alpha} \wedge T_{\alpha} .$	$C_{\mathrm{TT}}:=\frac{1}{8 \ell^2} \operatorname{Tr}\left(\gamma \wedge \Theta\right)=\frac{1}{2 \ell^2} \vartheta^{\alpha} \wedge T_{\alpha} .$
mielke-geometrodynamics_E08863 (12.4.10)	279	■ 1.000	$d C_{\mathrm{TT}}=\frac{1}{8 \ell^2} \operatorname{Tr}\left(\Theta \wedge \Theta-4 i \Omega \wedge \sigma\right)=\frac{1}{2 \ell^2}\left(T^{\alpha} \wedge T_{\alpha}+R_{\alpha \beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta}\right) .$	$d C_{\mathrm{TT}}=\frac{1}{8 \ell^2} \operatorname{Tr}\left(\Theta \wedge \Theta-4 i \Omega \wedge \sigma\right)=\frac{1}{2 \ell^2}\left(T^{\alpha} \wedge T_{\alpha}+R_{\alpha \beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta}\right) .$	$d C_{\mathrm{TT}}=\frac{1}{8 \ell^2} \operatorname{Tr}\left(\Theta \wedge \Theta-4 i \Omega \wedge \sigma\right)=\frac{1}{2 \ell^2}\left(T^{\alpha} \wedge T_{\alpha}+R_{\alpha \beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta}\right) .$
mielke-geometrodynamics_E08864 (12.4.11)	279	■ 1.000	$\hat{C}_{\mathrm{RR}}=C_{\mathrm{RR}}-2 C_{\mathrm{TT}}$	$\hat{C}_{\mathrm{RR}}=C_{\mathrm{RR}}-2 C_{\mathrm{TT}}$	$\hat{C}_{\mathrm{RR}}=C_{\mathrm{RR}}-2 C_{\mathrm{TT}}$
mielke-geometrodynamics_E08865 (13.2.1)	283	■ 1.000	$L_{\mathrm{BF}}=-B \wedge F=-B \wedge d A .$	$L_{\mathrm{BF}}=-B \wedge F=-B \wedge d A .$	$L_{\mathrm{BF}}=-B \wedge F=-B \wedge d A .$
mielke-geometrodynamics_E08866 (13.2.2)	283	■ 1.000	$\tilde{L}_{\mathrm{BF}}:=-B \wedge d A+\frac{1}{2} B \wedge B \cong-B \wedge d A+d C$	$\tilde{L}_{\mathrm{BF}}:=-B \wedge d A+\frac{1}{2} B \wedge B \cong-B \wedge d A+d C$	$\tilde{L}_{\mathrm{BF}}:=-B \wedge d A+\frac{1}{2} B \wedge B \cong-B \wedge d A+d C .$
mielke-geometrodynamics_E08868 (13.2.4)	283	■ 1.000	$C=\frac{1}{2} A \wedge F, \quad d C=\frac{1}{2} F \wedge F,$	$C=\frac{1}{2} A \wedge F, \quad d C=\frac{1}{2} F \wedge F,$	$C=\frac{1}{2} A \wedge F, \quad d C=\frac{1}{2} F \wedge F,$
mielke-geometrodynamics_E08869 (13.2.5)	283	■ 1.000	$A \rightarrow A^{\prime}=A+d \theta(x),$	$A \rightarrow A^{\prime}=A+d \theta(x),$	$A \rightarrow A^{\prime}=A+d \theta(x),$
mielke-geometrodynamics_E08870 (13.2.6)	284	■ 1.000	$A \rightarrow \widetilde{A}=A+\sigma, \quad B \rightarrow \widetilde{B}=B+d \sigma,$	$A \rightarrow \widetilde{A}=A+\sigma, \quad B \rightarrow \widetilde{B}=B+d \sigma,$	$A \rightarrow \widetilde{A}=A+\sigma, \quad B \rightarrow \widetilde{B}=B+d \sigma,$
mielke-geometrodynamics_E08871 (13.2.7)	284	■ 1.000	$L_{\mathrm{Max}}=-B \wedge d A+\frac{1}{2} B \wedge * B+L_{\mathrm{matter}},$	$L_{\mathrm{Max}}=-B \wedge d A+\frac{1}{2} B \wedge * B+L_{\mathrm{matter}},$	$L_{\mathrm{Max}}=-B \wedge d A+\frac{1}{2} B \wedge * B+L_{\mathrm{matter}},$
mielke-geometrodynamics_E08872 (13.2.8)	284	■ 1.000	$-d B=d^* F \cong j,$	$-d B=d^* F \cong j,$	$-d B=d^* F \cong j,$
mielke-geometrodynamics_E08875 (13.3.3)	284	■ 1.000	$\left[L^{\alpha}{}_{\beta}, L^{\gamma}{}_{\delta}\right]=\delta_{\delta}^{\alpha} L^{\gamma}{}_{\beta}-\delta_{\beta}^{\gamma} L^{\alpha}{}_{\delta},$	$\left[L^{\alpha}{}_{\beta}, L^{\gamma}{}_{\delta}\right]=\delta_{\delta}^{\alpha} L^{\gamma}{}_{\beta}-\delta_{\beta}^{\gamma} L^{\alpha}{}_{\delta},$	$\left[L^{\alpha}{}_{\beta}, L^{\gamma}{}_{\delta}\right]=\delta_{\delta}^{\alpha} L^{\gamma}{}_{\beta}-\delta_{\beta}^{\gamma} L^{\alpha}{}_{\delta},$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_285 EQ0880 (13.4.1)	285	■ 1.000	$A(x):=\left\{\begin{array}{cc} \Lambda(x) & \tau(x) \\ \tau_*(x) & 1 \end{array}\right\},$	$A(x) := \left\{ \begin{array}{cc} \Lambda(x) & \tau(x) \\ \tau_*(x) & 1 \end{array} \right\},$	$A(x) := \left\{ \begin{array}{cc} \Lambda(x) & \tau(x) \\ \tau_*(x) & 1 \end{array} \right\},$
mielke-geometrodynamics_285 EQ0881 (13.4.2)	285	■ 1.000	$\hat{\Gamma}=\left(\begin{array}{cc} \Gamma^{(L)} & \Gamma^{(T)} \\ \Gamma_*^{(T)} & 0 \end{array}\right)=\left(\begin{array}{cc} \Gamma_\alpha^{(L)\beta} L_\beta^\alpha & \Gamma^{(T)\alpha} \ell P_\alpha \\ \Gamma_\beta^{*(T)} & \ell P_*^\beta \end{array}\right)$	$\hat{\Gamma} = \left(\begin{array}{cc} \Gamma^{(L)} & \Gamma^{(T)} \\ \Gamma_*^{(T)} & 0 \end{array} \right) = \left(\begin{array}{cc} \Gamma_\alpha^{(L)\beta} L_\beta^\alpha & \Gamma^{(T)\alpha} \ell P_\alpha \\ \Gamma_\beta^{*(T)} & \ell P_*^\beta \end{array} \right)$	$\hat{T} = \left(\begin{array}{cc} \Gamma^{(L)} & \Gamma^{(T)} \\ \Gamma_*^{(T)} & 0 \end{array} \right) = \left(\begin{array}{cc} \Gamma_\alpha^{(L)\beta} L_\beta^\alpha & \Gamma^{(T)\alpha} \ell P_\alpha \\ \Gamma_\beta^{*(T)} & \ell P_*^\beta \end{array} \right)$
mielke-geometrodynamics_286 EQ0882 (13.4.3)	286	■ 1.000	$\hat{\Gamma} \rightarrow \hat{\Gamma}' = A^{-1}(x) \hat{\Gamma} A(x) + A^{-1}(x) dA(x).$	$\hat{\Gamma} \xrightarrow{A^{-1}(x)} \hat{\Gamma}' = A^{-1}(x) \hat{\Gamma} A(x) + A^{-1}(x) dA(x).$	$\hat{T} \xrightarrow{A^{-1}(x)} \hat{T}' = A^{-1}(x) \hat{T} A(x) + A^{-1}(x) dA(x).$
mielke-geometrodynamics_286 EQ0883	286	■ 1.000	$\tau \tau_* = -\Lambda \tau_* \Lambda^{-1} \tau.$	$\tau \tau_* = -\Lambda \tau_* \Lambda^{-1} \tau.$	$\tau \tau_* = -\Lambda \tau_* \Lambda^{-1} \tau.$
mielke-geometrodynamics_286 EQ0884 (13.4.4)	286	■ 1.000	$\hat{R}:=d\hat{\Gamma}+\hat{\Gamma}\wedge\hat{\Gamma}=\left(\begin{array}{cc} R^{(L)}+\Gamma^{(L)}\wedge\Gamma_*^{(L)} & d\Gamma^{(T)}+\Gamma^{(L)}\wedge\Gamma^{(T)} \\ d\Gamma_*^{(T)}+\Gamma_*^{(T)}\wedge\Gamma^{(L)} & 0 \end{array}\right)$	$\hat{R} := d\hat{\Gamma} + \hat{\Gamma} \wedge \hat{\Gamma} = \left(\begin{array}{cc} R^{(L)} + \Gamma^{(L)} \wedge \Gamma_*^{(L)} & d\Gamma^{(T)} + \Gamma^{(L)} \wedge \Gamma^{(T)} \\ d\Gamma_*^{(T)} + \Gamma_*^{(T)} \wedge \Gamma^{(L)} & 0 \end{array} \right)$	$\hat{R} := d\hat{T} + \hat{T} \wedge \hat{T} = \left(\begin{array}{cc} R^{(L)} + \Gamma^{(L)} \wedge \Gamma_*^{(L)} & d\Gamma^{(T)} + \Gamma^{(L)} \wedge \Gamma^{(T)} \\ d\Gamma_*^{(T)} + \Gamma_*^{(T)} \wedge \Gamma^{(L)} & 0 \end{array} \right),$
mielke-geometrodynamics_286 EQ0885	286	■ 1.000	$R^{(L)}:=d\Gamma^{(L)}+\Gamma^{(L)}\wedge\Gamma^{(L)}$	$R^{(L)} := d\Gamma^{(L)} + \Gamma^{(L)} \wedge \Gamma^{(L)}$	$R^{(L)} := d\Gamma^{(L)} + \Gamma^{(L)} \wedge \Gamma^{(L)}$
mielke-geometrodynamics_286 EQ0886 (13.4.5)	286	■ 1.000	$R^{(T)}:=d\Gamma^{(T)}+\Gamma^{(L)}\wedge\Gamma^{(T)}=\left(T^\beta-R_\alpha{}^\beta\xi^\alpha\right)\ell P_\beta$	$R^{(T)} := d\Gamma^{(T)} + \Gamma^{(L)} \wedge \Gamma^{(T)} = \left(T^\beta - R_\alpha{}^\beta \xi^\alpha\right) \ell P_\beta$	$R^{(T)} := d\Gamma^{(T)} + \Gamma^{(L)} \wedge \Gamma^{(T)} = \left(T^\beta - R_\alpha{}^\beta \xi^\alpha\right) \ell P_\beta$
mielke-geometrodynamics_286 EQ0887 (13.4.6)	286	■ 1.000	$\hat{R} \xrightarrow{A^{-1}(x)} \hat{R}' = A^{-1}(x) \hat{R} A(x).$	$\hat{R} \xrightarrow{A^{-1}(x)} \hat{R}' = A^{-1}(x) \hat{R} A(x).$	$\hat{R} \xrightarrow{A^{-1}(x)} \hat{R}' = A^{-1}(x) \hat{R} A(x).$
mielke-geometrodynamics_286 EQ0888 (13.4.7)	286	■ 1.000	$\Gamma^{(L)} \xrightarrow{A^{-1}(x)} \Gamma^{(L)'} = \Lambda^{-1}(x) \Gamma^{(L)} \Lambda(x) + \Lambda^{-1}(x) d\Lambda(x)$	$\Gamma^{(L)} \xrightarrow{A^{-1}(x)} \Gamma^{(L)'} = \Lambda^{-1}(x) \Gamma^{(L)} \Lambda(x) + \Lambda^{-1}(x) d\Lambda(x)$	$\Gamma^{(L)} \xrightarrow{A^{-1}(x)} \Gamma^{(L)'} = \Lambda^{-1}(x) \Gamma^{(L)} \Lambda(x) + \Lambda^{-1}(x) d\Lambda(x)$
mielke-geometrodynamics_287 EQ0891 (13.4.10)	287	■ 1.000	$\vartheta:=\Gamma^{(T)}+D\xi:=\vartheta^\alpha P_\alpha,\quad\theta:=\Gamma_*^{(T)}+D\xi_*:=\theta_\beta P_*^\beta,$	$\vartheta := \Gamma^{(T)} + D\xi := \vartheta^\alpha P_\alpha, \quad \theta := \Gamma_*^{(T)} + D\xi_* := \theta_\beta P_*^\beta,$	$\vartheta := \Gamma^{(T)} + D\xi := \vartheta^\alpha P_\alpha, \quad \theta := \Gamma_*^{(T)} + D\xi_* := \theta_\beta P_*^\beta,$
mielke-geometrodynamics_287 EQ0894 (13.4.13)	287	■ 1.000	$D\xi^*=0,\quad D\xi_*^*=0,$	$D\xi^*=0,\quad D\xi_*^*=0,$	$D\xi^*=0,\quad D\xi_*^*=0,$
mielke-geometrodynamics_287 EQ0895 (13.4.14)	287	■ 1.000	$\overline{\Gamma}=\left(\begin{array}{cc} \Gamma^{(L)} & \vartheta \\ \theta & 0 \end{array}\right),$	$\overline{\Gamma} = \left(\begin{array}{cc} \Gamma^{(L)} & \vartheta \\ \theta & 0 \end{array} \right),$	$\overline{T} = \left(\begin{array}{cc} \Gamma^{(L)} & \vartheta \\ \theta & 0 \end{array} \right),$
mielke-geometrodynamics_288 EQ0896 (13.5.1)	288	■ 1.000	$\hat{\Gamma}=\Gamma_\alpha{}^\beta L^\alpha{}_\beta+\vartheta^\alpha P_\alpha+\theta_\beta P_*^\beta,$	$\hat{\Gamma} = \Gamma_\alpha{}^\beta L^\alpha{}_\beta + \vartheta^\alpha P_\alpha + \theta_\beta P_*^\beta,$	$\hat{T} = \Gamma_\alpha{}^\beta L^\alpha{}_\beta + \vartheta^\alpha P_\alpha + \theta_\beta P_*^\beta,$
mielke-geometrodynamics_288 EQ0899 (13.5.4)	288	■ 1.000	$\tilde{L}_{\mathrm{SL}(5,\mathbb{R})}=-\mathrm{Tr}\{\hat{B}\wedge\hat{R}\}-d\hat{C}.$	$\tilde{L}_{\mathrm{SL}(5,\mathbb{R})} = -\mathrm{Tr}\{\hat{B} \wedge \hat{R}\} - d\hat{C}.$	$\tilde{L}_{\mathrm{SL}(5,\mathbb{R})} = -\mathrm{Tr}\{\hat{B} \wedge \hat{R}\} - d\hat{C}.$
mielke-geometrodynamics_289 EQ9900 (13.5.5)	289	■ 1.000	$\hat{D}B_A{}^B\cong\hat{D}\hat{R}_A{}^B\equiv 0.$	$\hat{D}B_A{}^B \cong \hat{D}\hat{R}_A{}^B \equiv 0.$	$\hat{D}B_A{}^B \cong \hat{D}\hat{R}_A{}^B \equiv 0.$
mielke-geometrodynamics_289 EQ9901 (13.5.6)	289	■ 1.000	$\tilde{L}_{\mathrm{SSB}}=\tilde{L}_{\mathrm{SL}(5,\mathbb{R})}+\frac{1}{2}\eta_{ABCDE}B^{AB}\wedge B^{CD}\Phi^E.$	$\tilde{L}_{\mathrm{SSB}} = \tilde{L}_{\mathrm{SL}(5,\mathbb{R})} + \frac{1}{2} \eta_{ABCDE} B^{AB} \wedge B^{CD} \Phi^E.$	$\tilde{L}_{\mathrm{SSB}} = \tilde{L}_{\mathrm{SL}(5,\mathbb{R})} + \frac{1}{2} \eta_{ABCDE} B^{AB} \wedge B^{CD} \Phi^E.$
mielke-geometrodynamics_289 EQ9902 (13.5.7)	289	■ 1.000	$\hat{g}_{AB}\Phi^A\Phi^B=\mu^2,\quad\Phi_E\hat{D}\Phi^E=0$	$\hat{g}_{AB} \Phi^A \Phi^B = \mu^2, \quad \Phi_E \hat{D} \Phi^E = 0$	$\hat{g}_{AB} \Phi^A \Phi^B = \mu^2, \quad \Phi_E \hat{D} \Phi^E = 0,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E09903 (13.5.8)	289	■ 1.000	$\left\langle \text{angle}\Phi^{\text{E}}\right\rangle \text{angle}=\Phi_{\text{0}}^{\text{E}}=(\theta,0,0,0,\mu)^{\text{T}}$	$\langle \Phi^E \rangle = \Phi_0^E = (0, 0, 0, 0, \mu)^T$	$\langle \Phi^E \rangle = \Phi_0^E = (0, 0, 0, 0, \mu)^T$
mielke-geometrodynamics_E09904 (13.5.9)	289	■ 1.000	$\tilde{\text{L}}_{\text{SSB}}=-\operatorname{Tr}\{\hat{\text{B}}\wedge\hat{\text{R}}\}+\frac{\mu}{2}\eta_{\alpha\beta\gamma\delta}B^{\alpha\beta}\wedge B^{\gamma\delta}-dC_{\text{RR}}+2dC_{\text{TT}}$	$\tilde{L}_{\text{SSB}} = -\text{Tr}\{\hat{B} \wedge \hat{R}\} + \frac{\mu}{2}\eta_{\alpha\beta\gamma\delta}B^{\alpha\beta} \wedge B^{\gamma\delta} - dC_{\text{RR}} + 2dC_{\text{TT}}$	$\tilde{L}_{\text{SSB}} = -\text{Tr}\{\hat{B} \wedge \hat{R}\} + \frac{\mu}{2}\eta_{\alpha\beta\gamma\delta}B^{\alpha\beta} \wedge B^{\gamma\delta} - dC_{\text{RR}} + 2dC_{\text{TT}}$
mielke-geometrodynamics_E09905 (13.5.10)	290	■ 1.000	$d\text{C}=\text{d}C_{\text{RR}}-2\text{d}C_{\text{TT}},$	$d\hat{C} = dC_{\text{RR}} - 2dC_{\text{TT}},$	$d\hat{C} = dC_{\text{RR}} - 2dC_{\text{TT}},$
mielke-geometrodynamics_E09906 (13.5.11)	290	■ 1.000	$B_{\alpha\beta}\cong\frac{1}{\mu}\eta_{\alpha\beta\gamma\delta}\left[R^{\gamma\delta}-\frac{\Lambda}{3}\theta^{\gamma}\wedge\vartheta^{\delta}\right]$	$B_{\alpha\beta} \cong \frac{1}{\mu}\eta_{\alpha\beta\gamma\delta} \left[R^{\gamma\delta} - \frac{\Lambda}{3}\theta^{\gamma} \wedge \vartheta^{\delta} \right]$	$B_{\alpha\beta} \cong \frac{1}{\mu}\eta_{\alpha\beta\gamma\delta} \left[R^{\gamma\delta} - \frac{\Lambda}{3}\theta^{\gamma} \wedge \vartheta^{\delta} \right]$
mielke-geometrodynamics_E09909 (13.5.14)	291	■ 1.000	$\tilde{\text{L}}_{\text{SSB}}\cong-\frac{1}{2\kappa}R^{\alpha\beta}\wedge\eta_{\alpha\beta}+\frac{\Lambda}{\kappa}\eta$	$\tilde{L}_{\text{SSB}} \cong -\frac{1}{2\kappa}R^{\alpha\beta} \wedge \eta_{\alpha\beta} + \frac{\Lambda}{\kappa}\eta$	$\tilde{L}_{\text{SSB}} \cong -\frac{1}{2\kappa}R^{\alpha\beta} \wedge \eta_{\alpha\beta} + \frac{\Lambda}{\kappa}\eta$
mielke-geometrodynamics_E09910 (13.5.15)	291	■ 1.000	$\mu=\frac{4}{3}\kappa\Lambda=\frac{1}{3}(2\kappa)^2\rho_{\Lambda}.$	$\mu = \frac{4}{3}\kappa\Lambda = \frac{1}{3}(2\kappa)^2\rho_{\Lambda}.$	$\mu = \frac{4}{3}\kappa\Lambda = \frac{1}{3}(2\kappa)^2\rho_{\Lambda}.$
mielke-geometrodynamics_E09912 (13.7.1)	293	■ 1.000	$g_{\text{N}}:=\kappa k^2, \quad \lambda:=\Lambda/k^2,$	$g_{\text{N}} := \kappa k^2, \quad \lambda := \Lambda/k^2,$	$g_{\text{N}} := \kappa k^2, \quad \lambda := \Lambda/k^2,$
mielke-geometrodynamics_E09913 (13.7.2)	293	■ 1.000	$k\frac{\partial}{\partial k}g_{\text{N}}=\beta_1(g_{\text{N}},\lambda)=(2+d_{\text{N}})g_{\text{N}}, \quad k\frac{\partial}{\partial k}\lambda=\beta_2(g_{\text{N}},\lambda),$	$k\frac{\partial}{\partial k}g_{\text{N}} = \beta_1(g_{\text{N}},\lambda) = (2 + d_{\text{N}})g_{\text{N}}, \quad k\frac{\partial}{\partial k}\lambda = \beta_2(g_{\text{N}},\lambda),$	$k\frac{\partial}{\partial k}g_{\text{N}} = \beta_1(g_{\text{N}},\lambda) = (2 + d_{\text{N}})g_{\text{N}}, \quad k\frac{\partial}{\partial k}\lambda = \beta_2(g_{\text{N}},\lambda),$
mielke-geometrodynamics_E09914 (13.7.3)	293	■ 1.000	$\mu_*\leq\frac{4}{3}g_{\text{N}*}\lambda_*\simeq0.2$	$\mu_* \leq \frac{4}{3}g_{\text{N}*}\lambda_* \simeq 0.2$	$\mu_* \leq \frac{4}{3}g_{\text{N}*}\lambda_* \simeq 0.2$
mielke-geometrodynamics_E09916 (13.8.2)	295	■ 1.000	$dC_{\text{TT}}=\frac{1}{2\ell^2}\left(T^{\alpha}\wedge T_{\alpha}+R_{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}\right)$	$dC_{\text{TT}} = \frac{1}{2\ell^2} \left(T^{\alpha} \wedge T_{\alpha} + R_{\alpha\beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta} \right)$	$dC_{\text{TT}} = \frac{1}{2\ell^2} \left(T^{\alpha} \wedge T_{\alpha} + R_{\alpha\beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta} \right)$
mielke-geometrodynamics_E09917 (13.8.3)	295	■ 1.000	$dC_{\text{RR}}=-\frac{1}{2}R_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}$	$dC_{\text{RR}} = -\frac{1}{2}R_{\alpha}{}^{\beta} \wedge R_{\beta}{}^{\alpha}$	$dC_{\text{RR}} = -\frac{1}{2}R_{\alpha}{}^{\beta} \wedge R_{\beta}{}^{\alpha}$
mielke-geometrodynamics_E09918 (13.8.4)	295	■ 1.000	$\text{D}R_{\alpha}{}^{\beta}\wedge\vartheta^{\alpha}\equiv0,$	$DR_{\alpha}{}^{\beta} \equiv 0,$	$DR_{\alpha}{}^{\beta} \equiv 0,$
mielke-geometrodynamics_E09921 (13.8.7)	296	■ 1.000	$L_{\text{SKY}}:=-\frac{1}{2}R_{\alpha\beta}\wedge{}^*R^{\alpha\beta},$	$L_{\text{SKY}} := -\frac{1}{2}R_{\alpha\beta} \wedge {}^*R^{\alpha\beta},$	$L_{\text{SKY}} := -\frac{1}{2}R_{\alpha\beta} \wedge {}^*R^{\alpha\beta},$
mielke-geometrodynamics_E09922 (14.1.1)	302	■ 1.000	$L=L_{\text{g}}+L_{\omega}+L_{\text{mat}}$	$L = L_g + L_{\omega} + L_{\text{mat}}$	$L = L_g + L_{\omega} + L_{\text{mat}}$
mielke-geometrodynamics_E09923 (14.1.2)	302	■ 1.000	$L_{\omega-g}=\operatorname{Tr}\left\{\wedge^*\left(\vartheta\wedge\Omega^g\right)\wedge^*\left(\vartheta\wedge\Omega\right)\right\}$	$L_{\omega-g} = \text{Tr} \left\{ \wedge^* (\vartheta \wedge \Omega^g) \wedge^* (\vartheta \wedge \Omega) \right\}$	$L_{\omega-g} = \text{Tr} \left\{ \wedge^* (\vartheta \wedge \Omega^g) \wedge^* (\vartheta \wedge \Omega) \right\}$
mielke-geometrodynamics_E09924 (14.1.3)	303	■ 1.000	$L_{\text{GMD}}.=\frac{1}{\ell^*2}L_{\text{W.}}+L_{\text{YM}}+L_{\text{mat}}$	$L_{\text{GMD.}} = \frac{1}{\ell^{*2}}L_{\text{W.}} + L_{\text{YM}} + L_{\text{mat}}$	$L_{\text{GMD.}} = \frac{1}{\ell^{*2}}L_{\text{W.}} + L_{\text{YM}} + L_{\text{mat}}$
mielke-geometrodynamics_E09933 (14.1.14)	305	■ 1.000	$e^{*\pi/2}=(-1)^z*, \quad z:=\frac{s-1}{2}.$	$e^{*\pi/2} = (-1)^z*, \quad z := \frac{s-1}{2}.$	$e^{*\pi/2} = (-1)^z*, \quad z := \frac{s-1}{2}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E09934 (14.1.15)	305	■ 1.000	$\backslash \mathrm{xi}:=\mathrm{e}^{\ast}\left(-\backslash \mathrm{delta}\right)\}\backslash 0\mathrm{omega}$	$\xi:=e^{\ast(-\delta)}\Omega$	$\xi:=e^{\ast(-\delta)}\varOmega$
mielke-geometrodynamics_E09935 (14.1.16)	305	■ 1.000	$\backslash 0\mathrm{omega}\backslash \mathrm{wedge}^{\ast}\}\backslash 0\mathrm{omega}\backslash \mathrm{rightarrow}\backslash \mathrm{xi}\backslash \mathrm{wedge}^{\ast}\}\backslash \mathrm{xi}=\backslash 0\mathrm{omega}\backslash \mathrm{wedge}^{\ast}\}\}\backslash 0\mathrm{omega}\backslash \cos^2\backslash \mathrm{delta}-\left(-1\right)^{\mathrm{z}}\}\backslash 0\mathrm{omega}\backslash \mathrm{wedge}\backslash 0\mathrm{omega}\backslash \sin^2\backslash \mathrm{delta}$	$\Omega\wedge^*\Omega\rightarrow\xi\wedge^*\xi=\Omega\wedge^*\Omega\cos2\delta-(-1)^z\Omega\wedge\Omega\sin2\delta$	$\varOmega\wedge^*\varOmega\rightarrow\xi\wedge^*\xi=\varOmega\wedge^*\varOmega\cos2\delta-(-1)^z\varOmega\wedge\varOmega\sin2\delta$
mielke-geometrodynamics_E09936 (14.1.17)	305	■ 1.000	$\backslash 0\mathrm{omega}\backslash \mathrm{wedge}\backslash 0\mathrm{omega}\backslash \mathrm{rightarrow}\backslash \mathrm{xi}\backslash \mathrm{wedge}\backslash \mathrm{xi}=\backslash 0\mathrm{omega}\backslash \mathrm{wedge}\backslash 0\mathrm{omega}\backslash \cos^2\backslash \mathrm{delta}-\left(-1\right)^{\mathrm{z}}\}\backslash 0\mathrm{omega}\backslash \mathrm{wedge}^{\ast}\}\}\backslash 0\mathrm{omega}\backslash \sin^2\backslash \mathrm{delta}\text{.}$	$\Omega\wedge\Omega\rightarrow\xi\wedge\xi=\Omega\wedge\Omega\cos2\delta-(-1)^z\Omega\wedge^*\Omega\sin2\delta.$	$\varOmega\wedge\varOmega\rightarrow\xi\wedge\xi=\varOmega\wedge\varOmega\cos2\delta-(-1)^z\varOmega\wedge^*\varOmega\sin2\delta.$
mielke-geometrodynamics_E09937 (14.1.18)	305	■ 1.000	$\backslash \mathrm{vec}\{\mathrm{E}\}=\backslash \mathrm{left}\{\backslash \mathrm{E}_{\mathrm{j}}\}:=\mathrm{F}_{\mathrm{j}\text{ }0}\}\backslash \mathrm{mid}\text{ }j=1,2,3\backslash \mathrm{right}\}$	$\vec{E}=\{E_j:=F_{j0}\mid j=1,2,3\}$	$\vec{E}=\{E_j:=F_{j0} j=1,2,3\}$
mielke-geometrodynamics_E09938 (14.1.19)	305	■ 1.000	$\backslash \mathrm{vec}\{\mathrm{B}\}=\backslash \mathrm{left}\{\backslash \mathrm{B}_{\mathrm{j}}\}:\backslash \mathrm{left}.=\frac{1}{2}\}\backslash \mathrm{epsilon}_{\mathrm{i}\text{ }j\text{ }k}\}\mathrm{F}^{\mathrm{j}\text{ }k}\}\backslash \mathrm{right}\backslash \mathrm{rvert}\backslash \text{, i, j, k=1,2,3}\backslash \mathrm{right}\}$	$\vec{B}=\left\{B_j:=\frac{1}{2}\epsilon_{ijk}F^{jk}\right i,j,k=1,2,3\}$	$\vec{B}=\{B_j:=\frac{1}{2}\epsilon_{ijk}F^{jk} i,j,k=1,2,3\}$
mielke-geometrodynamics_E09943 (14.1.24)	306	■ 1.000	$\backslash 0\mathrm{omega}=\mathrm{e}^{\ast}\backslash \mathrm{delta}\}\backslash \mathrm{xi}\text{.}$	$\Omega=e^{*\delta}\xi.$	$\varOmega=e^{*\delta}\xi.$
mielke-geometrodynamics_E09944 (14.1.25)	306	■ 1.000	$\backslash \mathrm{Sigma}_{\backslash \mathrm{mathrm}\{\mathrm{YM}\}}\}\left(\backslash 0\mathrm{omega}\right)=\backslash \mathrm{Sigma}_{\backslash \mathrm{mathrm}\{\mathrm{YM}\}}\}\}\left(\backslash \mathrm{xi}\right)$	$\Sigma_{\mathrm{YM}}(\Omega)=\Sigma_{\mathrm{YM}}(\xi)$	$\varSigma_{\mathrm{YM}}(\varOmega)=\varSigma_{\mathrm{YM}}(\xi)$
mielke-geometrodynamics_E09950 (14.1.31)	307	■ 1.000	$\backslash \mathrm{tau}_{\mathrm{e}}=\backslash \mathrm{bar}\{\backslash \mathrm{tau}\}_{\mathrm{e}}\}\backslash \cos\backslash \mathrm{delta}-\left(-1\right)^{\mathrm{z}}\}\backslash \mathrm{bar}\{\backslash \mathrm{tau}\}_{\mathrm{p}}\}\backslash \sin\backslash \mathrm{delta}$	$\tau_e=\bar{\tau}_e\cos\delta-(-1)^z\bar{\tau}_p\sin\delta$	$\tau_e=\bar{\tau}_e\cos\delta-(-1)^z\bar{\tau}_p\sin\delta$
mielke-geometrodynamics_E09952 (14.1.34)	307	■ 1.000	$D\backslash \mathrm{delta}=\mathrm{d}\backslash \mathrm{delta}=0,$	$D\delta=d\delta=0,$	$D\delta=d\delta=0,$
mielke-geometrodynamics_E09955 (14.1.37)	308	■ 1.000	$\left(-1\right)^{\mathrm{z}}\}\backslash \mathrm{Sigma}_{\backslash \mathrm{mathrm}\{\mathrm{YM}\}}\}\}\left(\backslash \mathrm{xi}\right)\backslash \mathrm{wedge}\mathrm{d}\backslash \mathrm{delta}=\mathrm{D}\backslash \mathrm{Sigma}_{\backslash \mathrm{mathrm}\{\mathrm{YM}\}}\}\}\left(\backslash \mathrm{xi}\right)\text{.}$	$(-1)^z\Sigma_{\mathrm{YM}}(\xi)\wedge d\delta=D\Sigma_{\mathrm{YM}}(\xi).$	$(-1)^z\varSigma_{\mathrm{YM}}(\xi)\wedge d\delta=D\varSigma_{\mathrm{YM}}(\xi).$
mielke-geometrodynamics_E09958 (14.1.40)	309	■ 1.000	$\backslash \mathrm{delta}=\backslash \mathrm{int}_{\mathrm{c}_{\mathrm{1}}}\}\mathrm{d}\backslash \mathrm{delta}+\backslash \mathrm{delta}_{\mathrm{0}}\}$	$\delta=\int_{c_1}d\delta+\delta_0$	$\delta=\int_{c_1}d\delta+\delta_0.$
mielke-geometrodynamics_E09969 (14.2.10)	314	■ 1.000	$\backslash \mathrm{hat}\{\backslash \mathrm{omega}\}^{\mathrm{A}\text{ }B}=-\backslash \mathrm{hat}\{\backslash \mathrm{omega}\}^{\mathrm{B}\text{ }A},\backslash \mathrm{quad}\text{ }A,\text{ }B=0,\backslash \mathrm{dots},\text{ }3+K$	$\hat{\omega}^{AB}=-\hat{\omega}^{BA},\quad A,B=0,\dots,3+K$	$\hat{\omega}^{AB}=-\hat{\omega}^{BA},\quad A,B=0,\dots,3+K$
mielke-geometrodynamics_E09975 (14.2.16)	315	■ 1.000	$\backslash 0\mathrm{omega}^{\backslash \mathrm{prime}}:=\backslash \mathrm{pi}\backslash \mathrm{left}(\mathrm{d}\backslash \mathrm{omega}-\backslash \mathrm{omega}\backslash \mathrm{wedge}\backslash \mathrm{stackrel{\circ}{\mathrm{c}}}\{\backslash \mathrm{omega}\}-\backslash \mathrm{omega}^{\mathrm{g}}\}\backslash \mathrm{wedge}\backslash \mathrm{omega}\backslash \mathrm{right})=\backslash 0\mathrm{omega}-\backslash \mathrm{omega}^{\mathrm{g}}\}\backslash \mathrm{wedge}\backslash \mathrm{omega}\text{.}$	$\Omega':=\pi\left(d\omega-\omega\wedge\overset{\circ}{\omega}-\omega^g\wedge\omega\right)=\Omega-\omega^g\wedge\omega.$	$\varOmega':=\pi(d\omega-\omega\wedge\overset{\circ}{\omega}-\omega^g\wedge\omega)=\varOmega-\omega^g\wedge\omega.$
mielke-geometrodynamics_E09980 (14.2.21)	316	■ 1.000	$\backslash \mathrm{hat}\{\mathrm{g}\}_{\mathrm{A}\text{ }B}\}\backslash \mathrm{left}(x,\backslash \mathrm{xi}^{\mathrm{j}}\}\backslash \mathrm{right})=\backslash \mathrm{sum}_{n=-\backslash \mathrm{infty}}^{\backslash \mathrm{infty}}\}\mathrm{g}_{\mathrm{A}\text{ }B}^{\mathrm{j}}\}\left(n\right)\}\left(x\right)\mathrm{Y}_{\mathrm{n}}\}\backslash \mathrm{left}(\backslash \mathrm{xi}^{\mathrm{j}}\}\backslash \mathrm{right})$	$\hat{g}_{AB}\left(x,\xi^j\right)=\sum_{n=-\infty}^{+\infty}g_{AB}^{(n)}(x)Y_n\left(\xi^j\right)$	$\hat{g}_{AB}(x,\xi^j)=\sum_{n=-\infty}^{+\infty}g_{AB}^{(n)}(x)Y_n(\xi^j)$
mielke-geometrodynamics_E09981 (14.2.22)	317	■ 1.000	$M^{\ast}=\sqrt{\hbar c/\mathrm{G}_{\backslash \mathrm{mathrm}\{\mathrm{N}\}}}\}\backslash \mathrm{simseq}\text{ }10^{\mathrm{19}}\}\backslash \mathrm{mathrm}\{\mathrm{GeV}\}\text{ }/\text{ }c^{\mathrm{2}}\}\text{.}$	$M^*=\sqrt{\hbar c/G_{\mathrm{N}}}\simeq10^{19}\mathrm{GeV}/c^2.$	$M^*=\sqrt{\hbar c/G_{\mathrm{N}}}\simeq10^{19}\mathrm{GeV}/c^2.$
mielke-geometrodynamics_E09983 (14.2.24)	318	■ 1.000	$\mathrm{G}=\mathrm{S}\text{ }U(3)\backslash \mathrm{times}\text{ }S\text{ }U(2)\backslash \mathrm{times}\text{ }U(1)$	$G=SU(3)\times SU(2)\times U(1)$	$G=SU(3)\times SU(2)\times U(1)$
mielke-geometrodynamics_E09984 (14.2.25)	318	■ 1.000	$\mathrm{B}=\backslash \mathrm{mathbb}\{\mathrm{C}\}\text{ }P^{\mathrm{2}}\}\backslash \mathrm{times}\text{ }S^{\mathrm{2}}\}\backslash \mathrm{times}\text{ }S^{\mathrm{1}},$	$B=\mathbb{C}P^2\times S^2\times S^1,$	$B=\mathbb{C}P^2\times S^2\times S^1,$
mielke-geometrodynamics_E09985 (14.2.26)	318	■ 1.000	$\mathrm{G}\text{ }/\text{ }H=\frac{\mathrm{S}\text{ }U(3)\backslash \mathrm{times}\text{ }S\text{ }U(2)\backslash \mathrm{times}\text{ }U(1)}{\mathrm{S}\text{ }U(2)\backslash \mathrm{times}\text{ }U(1)\backslash \mathrm{times}\text{ }U(1)},$	$G/H=\frac{SU(3)\times SU(2)\times U(1)}{SU(2)\times U(1)\times U(1)},$	$G/H=\frac{SU(3)\times SU(2)\times U(1)}{SU(2)\times U(1)\times U(1)},$
mielke-geometrodynamics_E09986 (14.2.27)	318	■ 1.000	$\backslash \mathrm{widetilde}\{\mathrm{S}\}^{\mathrm{7}}:=\frac{\mathrm{S}\text{ }0(5)\backslash \mathrm{times}\text{ }S\text{ }0(3)}{\mathrm{S}\text{ }0(3)}\}\backslash \mathrm{times}\text{ }S\text{ }0(3)\}\text{.}$	$\widetilde{S}^7:=\frac{SO(5)\times SO(3)}{SO(3)\times SO(3)}.$	$\widetilde{S}^7:=\frac{SO(5)\times SO(3)}{SO(3)\times SO(3)}.$
mielke-geometrodynamics_E09988 (14.3.1)	319	■ 1.000	$\backslash \mathrm{varphi}\backslash \mathrm{rightarrow}\backslash \mathrm{varphi}^{\backslash \mathrm{mathrm}\{\mathrm{C}\}}:=\backslash \mathrm{bar}\{\backslash \mathrm{varphi}\}$	$\varphi\rightarrow\varphi^{\mathrm{C}}:=\bar{\varphi}$	$\varphi\rightarrow\varphi^{\mathrm{C}}:=\bar{\varphi}$
mielke-geometrodynamics_E09991 (14.3.4)	320	■ 1.000	$\backslash \mathrm{hat}\{\backslash \mathrm{psi}\}=\backslash \mathrm{sum}_{F=1}^{\backslash \mathrm{operatorname}\{\mathrm{dim}\}\text{ }\backslash \mathrm{rho}\}\left(\mathrm{G}\right)\}\backslash \mathrm{left}(\backslash \mathrm{psi}_{\mathrm{-}}\}\backslash \mathrm{sup}\{\mathrm{F}\}\}\left(x\right)\backslash \mathrm{chi}_{\backslash \mathrm{rho}\text{ }F}\}\backslash \mathrm{sup}\{-\}\}\left(\backslash \mathrm{xi}\right)+\backslash \mathrm{psi}_{\mathrm{+}}\}\backslash \mathrm{sup}\{\mathrm{F}\}\}\left(x\right)\backslash \mathrm{chi}_{\backslash \mathrm{rho}\text{ }F}\}\backslash \mathrm{sup}\{+\}\}\left(\backslash \mathrm{xi}\right)\}\backslash \mathrm{right})\text{.}$	$\hat{\psi}=\sum_{F=1}^{\dim\rho(G)}\left(\psi_{-}^F(x)\chi_{\rho F}^{-}(\xi)+\psi_{+}^F(x)\chi_{\rho F}^{+}(\xi)\right).$	$\hat{\psi}=\sum_{F=1}^{\dim\rho(G)}\left(\psi_{-}^F(x)\chi_{\rho F}^{-}(\xi)+\psi_{+}^F(x)\chi_{\rho F}^{+}(\xi)\right).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0994 (14.3.7)	321	■ 1.000	$D^{\mathrm{g}+\omega}:=d+i\,\widetilde{\omega}^g\otimes i\,\omega$	$D^{g+\omega}:=d+i\widetilde{\omega}^g\otimes i\omega$	$D^{g+\omega}:=d+i\widetilde{\omega}^g\otimes i\omega$
mielke-geometrodynamics_EQ0999 (14.3.12)	322	■ 1.000	$\left(\widehat{\gamma}^{n+1}\right)^2=-1\,\,\,\text{\rm Rightarrow}\,\,\,\widehat{\gamma}^{n+1}\widehat{\psi}=\mp i\widehat{\psi}.$	$\left(\widehat{\gamma}^{n+1}\right)^2=-1\Rightarrow\widehat{\gamma}^{n+1}\widehat{\psi}=\mp i\widehat{\psi}.$	$(\widehat{\gamma}^{n+1})^2=-1\quad\Rightarrow\quad\widehat{\gamma}^{n+1}\widehat{\psi}=\mp i\widehat{\psi}.$
mielke-geometrodynamics_EQ1000 (14.3.13)	322	■ 1.000	$\left(\widehat{\gamma}^{n+1}\right)^2=+1\,\,\,\text{\rm Rightarrow}\,\,\,\widehat{\gamma}^{n+1}\widehat{\psi}=\mp\widehat{\psi}.$	$\left(\widehat{\gamma}^{n+1}\right)^2=+1\Rightarrow\widehat{\gamma}^{n+1}\widehat{\psi}=\mp\widehat{\psi}.$	$(\widehat{\gamma}^{n+1})^2=+1\quad\Rightarrow\quad\widehat{\gamma}^{n+1}\widehat{\psi}=\mp\widehat{\psi}.$
mielke-geometrodynamics_EQ1006 (14.4.1)	324	■ 1.000	$\bar{Q}^{(i)}=Q^{(i)}\cos\delta+P^{(i)}\sin\delta,\quad Q^{(\circ)}\equiv Q_e.$	$\bar{Q}^{(i)}=Q^{(i)}\cos\delta+P^{(i)}\sin\delta,\quad Q^{(\circ)}\equiv Q_e.$	$\bar{Q}^{(i)}=Q^{(i)}\cos\delta+P^{(i)}\sin\delta,\quad Q^{(\circ)}\equiv Q_e.$
mielke-geometrodynamics_EQ1009 (14.4.4)	325	■ 1.000	$\rho^2:=r^2+\frac{J(J+1)\hbar^2}{M^2c^2}\cos^2\vartheta$	$\rho^2:=r^2+\frac{J(J+1)\hbar^2}{M^2c^2}\cos^2\vartheta$	$\rho^2:=r^2+\frac{J(J+1)\hbar^2}{M^2c^2}\cos^2\vartheta$
mielke-geometrodynamics_EQ1010 (14.4.5)	325	■ 1.000	$\Delta:=r^2-\frac{2M\hbar}{M^{*2}c}r+\frac{J(J+1)\hbar^2}{M^2c^2}+\frac{\alpha_e\hbar^2}{M^{*2}c^2}\mathrm{Tr}\,\bar{Q}^2$	$\Delta:=r^2-\frac{2M\hbar}{M^{*2}c}r+\frac{J(J+1)\hbar^2}{M^2c^2}+\frac{\alpha_e\hbar^2}{M^{*2}c^2}\mathrm{Tr}\,\bar{Q}^2$	$\Delta:=r^2-\frac{2M\hbar}{M^{*2}c}r+\frac{J(J+1)\hbar^2}{M^2c^2}+\frac{\alpha_e\hbar^2}{M^{*2}c^2}\mathrm{Tr}\bar{Q}^2$
mielke-geometrodynamics_EQ1017 (14.4.12)	327	■ 1.000	$ds^2=-\frac{\bar{\Delta}}{\bar{\rho}^2}\left(\frac{4N\hbar}{Mc}\sin^2\left(\frac{\vartheta}{2}\right)d\phi+c\,dt\right)^2+\frac{\bar{\rho}^2}{\bar{\Delta}}(dr^2+\bar{\Delta}d\Omega^2),$	$ds^2=-\frac{\bar{\Delta}}{\bar{\rho}^2}\left(\frac{4N\hbar}{Mc}\sin^2\left(\frac{\vartheta}{2}\right)d\phi+c\,dt\right)^2+\frac{\bar{\rho}^2}{\bar{\Delta}}(dr^2+\bar{\Delta}d\Omega^2),$	$ds^2=-\frac{\bar{\Delta}}{\bar{\rho}^2}\left(\frac{4N\hbar}{Mc}\sin^2\left(\frac{\vartheta}{2}\right)d\phi+c\,dt\right)^2+\frac{\bar{\rho}^2}{\bar{\Delta}}(dr^2+\bar{\Delta}d\Omega^2),$
mielke-geometrodynamics_EQ1018 (14.4.13)	328	■ 1.000	$\bar{\Delta}:=r^2-\frac{2M\hbar}{M^{*2}c}r-\frac{N^2\hbar^2}{M^2c^2}+\frac{\alpha_e\hbar^2}{M^{*2}c^2}\mathrm{Tr}\,\bar{Q}^2$	$\bar{\Delta}:=r^2-\frac{2M\hbar}{M^{*2}c}r-\frac{N^2\hbar^2}{M^2c^2}+\frac{\alpha_e\hbar^2}{M^{*2}c^2}\mathrm{Tr}\,\bar{Q}^2$	$\bar{\Delta}:=r^2-\frac{2M\hbar}{M^{*2}c}r-\frac{N^2\hbar^2}{M^2c^2}+\frac{\alpha_e\hbar^2}{M^{*2}c^2}\mathrm{Tr}\bar{Q}^2$
mielke-geometrodynamics_EQ1019 (14.4.14)	328	■ 1.000	$\bar{\rho}^2:=r^2+\frac{N^2\hbar^2}{M^2c^2}$	$\bar{\rho}^2:=r^2+\frac{N^2\hbar^2}{M^2c^2}$	$\bar{\rho}^2:=r^2+\frac{N^2\hbar^2}{M^2c^2}$
mielke-geometrodynamics_EQ1020 (14.4.15)	328	■ 1.000	$A=-\frac{1}{\bar{\rho}^2}(\bar{Q}\cos\delta+N\bar{Q}\sin\delta)c\,dt+\frac{1-\cos\vartheta}{\bar{\rho}^2}\bar{Q}\left[-\bar{\rho}^2\sin\delta+2r\frac{N\hbar}{Mc}\cos\delta\right]d\phi.$	$A=-\frac{1}{\bar{\rho}^2}(\bar{Q}\cos\delta+N\bar{Q}\sin\delta)c\,dt+\frac{1-\cos\vartheta}{\bar{\rho}^2}\bar{Q}\left[-\bar{\rho}^2\sin\delta+2r\frac{N\hbar}{Mc}\cos\delta\right]d\phi$	$A=-\frac{1}{\bar{\rho}^2}(\bar{Q}\cos\delta+N\bar{Q}\sin\delta)c\,dt+\frac{1-\cos\vartheta}{\bar{\rho}^2}\bar{Q}\left[-\bar{\rho}^2\sin\delta+2r\frac{N\hbar}{Mc}\cos\delta\right]d\phi.$
mielke-geometrodynamics_EQ1022 (14.4.17)	328	■ 1.000	$M^2/M_{\mathrm{ir}}^2=\frac{\left[1+\frac{1}{4}(\alpha_e^2\mathrm{Tr}\,\bar{Q}^2-2N^2)(M^*/M_{\mathrm{ir}})^4\right]^2}{1-\frac{1}{4}N^2(M^*/M_{\mathrm{ir}})^4},$	$M^2/M_{\mathrm{ir}}^2=\frac{\left[1+\frac{1}{4}(\alpha_e^2\mathrm{Tr}\,\bar{Q}^2-2N^2)(M^*/M_{\mathrm{ir}})^4\right]^2}{1-\frac{1}{4}N^2(M^*/M_{\mathrm{ir}})^4},$	$M^2/M_{\mathrm{ir}}^2=\frac{\left[1+\frac{1}{4}(\alpha_e^2\mathrm{Tr}\,\bar{Q}^2-2N^2)(M^*/M_{\mathrm{ir}})^4\right]^2}{1-\frac{1}{4}N^2(M^*/M_{\mathrm{ir}})^4},$
mielke-geometrodynamics_EQ1023 (15.0.1)	335	■ 1.000	$\ell:=2\pi\hbar/(M_{\mathrm{P}}c)=\ell^*/\sqrt{\kappa}\simeq1.3\times10^{-13}\,\mathrm{cm}$	$\ell:=2\pi\hbar/(M_{\mathrm{P}}c)=\ell^*/\sqrt{\kappa}\simeq1.3\times10^{-13}\,\mathrm{cm}$	$\ell:=2\pi\hbar/(M_{\mathrm{P}}c)=\ell^*/\sqrt{\kappa}\simeq1.3\times10^{-13}\,\mathrm{cm}$
mielke-geometrodynamics_EQ1024 (15.0.2)	335	■ 1.000	$V_{\mathrm{Yu}}(r)\sim\frac{1}{r}e^{-rm_{\pi}c/\hbar},$	$V_{\mathrm{Yu}}(r)\sim\frac{1}{r}e^{-rm_{\pi}c/\hbar},$	$V_{\mathrm{Yu}}(r)\sim\frac{1}{r}e^{-rm_{\pi}c/\hbar},$
mielke-geometrodynamics_EQ1026 (15.1.2)	336	■ 1.000	$L_{fg}=P\left(\vartheta^f,\vartheta^g\right)\eta,$	$L_{fg}=P\left(\vartheta^f,\vartheta^g\right)\eta,$	$L_{fg}=P(\vartheta^f,\vartheta^g)\eta,$
mielke-geometrodynamics_EQ1027 (15.1.3)	337	■ 1.000	$\delta L_{fg}=: \delta\vartheta^f\wedge\Sigma^f+\delta\vartheta^g\wedge\Sigma^g.$	$\delta L_{fg}=: \delta\vartheta^f\wedge\Sigma^f+\delta\vartheta^g\wedge\Sigma^g.$	$\delta L_{fg}=: \delta\vartheta^f\wedge\Sigma^f+\delta\vartheta^g\wedge\Sigma^g.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_337 EQ1028 (15.1.4)	337	■ 1.000	$\frac{1}{a} \vartheta^f \wedge \Sigma^f + \frac{1}{b} \vartheta^g \wedge \Sigma^g = 2 L_{fg} \vartheta^f \wedge \Sigma^g$	$\frac{1}{a} \vartheta^f \wedge \Sigma^f + \frac{1}{b} \vartheta^g \wedge \Sigma^g = 2 L_{fg}$	$\frac{1}{a} \vartheta^f \wedge \Sigma^f + \frac{1}{b} \vartheta^g \wedge \Sigma^g = 2 L_{fg}$
mielke-geometrodynamics_337 EQ1029 (15.1.5)	337	■ 1.000	$L_{fg}^{\prime} = \frac{m_f^2}{\ell^2} \det g_{ij} ^u \det f_{ij} ^{1/2-u} (\vartheta^f - \vartheta^g) \wedge (\vartheta^f - \vartheta^g) \wedge \vartheta^g \wedge \vartheta^g$	$L_{fg}^{\prime} = \frac{m_f^2}{\ell^2} \det g_{ij} ^u \det f_{ij} ^{1/2-u} (\vartheta^f - \vartheta^g) \wedge (\vartheta^f - \vartheta^g) \wedge \vartheta^g \wedge \vartheta^g$	$L_{fg}^{\prime} = \frac{m_f^2}{\ell^2} \det g_{ij} ^u \det f_{ij} ^{1/2-u} (\vartheta^f - \vartheta^g) \wedge (\vartheta^f - \vartheta^g) \wedge \vartheta^g \wedge \vartheta^g$
mielke-geometrodynamics_338 EQ1035 (15.1.12)	338	■ 1.000	$\vartheta^{m_f} := \vartheta^f - \vartheta^g$	$\vartheta^{m_f} := \vartheta^f - \vartheta^g$	$\vartheta^{m_f} := \vartheta^f - \vartheta^g$
mielke-geometrodynamics_338 EQ1036 (15.1.13)	338	■ 1.000	$\begin{aligned} ds^2 &= \frac{1}{1+\kappa} g^{\alpha\beta} \left(\vartheta_{\alpha}^g \otimes_s \vartheta_{\beta}^g + \kappa \vartheta_{\alpha}^f \otimes \vartheta_{\beta}^f \right) \\ &= \frac{1}{1+\kappa} (g_{ij} + \kappa f_{ij}) dx^i \otimes_s dx^j \end{aligned}$	$\begin{aligned} ds^2 &= \frac{1}{1+\kappa} g^{\alpha\beta} \left(\vartheta_{\alpha}^g \otimes_s \vartheta_{\beta}^g + \kappa \vartheta_{\alpha}^f \otimes \vartheta_{\beta}^f \right) \\ &= \frac{1}{1+\kappa} (g_{ij} + \kappa f_{ij}) dx^i \otimes_s dx^j \end{aligned}$	$\begin{aligned} ds^2 &= \frac{1}{1+\kappa} g^{\alpha\beta} \left(\vartheta_{\alpha}^g \otimes_s \vartheta_{\beta}^g + \kappa \vartheta_{\alpha}^f \otimes \vartheta_{\beta}^f \right) \\ &= \frac{1}{1+\kappa} (g_{ij} + \kappa f_{ij}) dx^i \otimes_s dx^j. \end{aligned}$
mielke-geometrodynamics_338 EQ1037 (15.1.14)	338	■ 1.000	$\kappa = \ell^{*2} / \ell^2 = G_N / G_S = 0.38 \times 10^{-38}$	$\kappa = \ell^{*2} / \ell^2 = G_N / G_S = 0.38 \times 10^{-38}$	$\kappa = \ell^{*2} / \ell^2 = G_N / G_S = 0.38 \times 10^{-38}$
mielke-geometrodynamics_339 EQ1038 (15.1.15)	339	■ 1.000	$e^{-\lambda^{g,f}} := 1 - \frac{2\mu^{g,f}}{r} - \frac{1}{3} \Lambda^{g,f} r^2,$	$e^{-\lambda^{g,f}} := 1 - \frac{2\mu^{g,f}}{r} - \frac{1}{3} \Lambda^{g,f} r^2,$	$e^{-\lambda^{g,f}} := 1 - \frac{2\mu^{g,f}}{r} - \frac{1}{3} \Lambda^{g,f} r^2,$
mielke-geometrodynamics_339 EQ1040 (15.1.17)	339	■ 1.000	$e^{-\lambda^f} \simeq 1 - \frac{2\mu^f}{r} \exp(-r m_f c / \hbar).$	$e^{-\lambda^f} \simeq 1 - \frac{2\mu^f}{r} \exp(-r m_f c / \hbar).$	$e^{-\lambda^f} \simeq 1 - \frac{2\mu^f}{r} \exp(-r m_f c / \hbar).$
mielke-geometrodynamics_340 EQ1041 (15.1.18)	340	■ 1.000	$E_{\alpha\beta} : (2^+) \oplus 2(1^-) \oplus 2(0^+) \oplus (1^+)$	$E_{\alpha\beta} : (2^+) \oplus 2(1^-) \oplus 2(0^+) \oplus (1^+)$	$E_{\alpha\beta} : (2^+) \oplus 2(1^-) \oplus 2(0^+) \oplus (1^+)$
mielke-geometrodynamics_342 EQ1043 (15.1.20)	342	■ 1.000	$\frac{M_U}{M_P} = \frac{\ell_H}{\ell^*} \cdot \frac{\ell}{\ell^*} = \frac{1}{\kappa} \frac{\ell_H}{\ell} \simeq 10^{80}$	$\frac{M_U}{M_P} = \frac{\ell_H}{\ell^*} \cdot \frac{\ell}{\ell^*} = \frac{1}{\kappa} \frac{\ell_H}{\ell} \simeq 10^{80}$	$\frac{M_U}{M_P} = \frac{\ell_H}{\ell^*} \cdot \frac{\ell}{\ell^*} = \frac{1}{\kappa} \frac{\ell_H}{\ell} \simeq 10^{80}$
mielke-geometrodynamics_343 EQ1044 (15.2.1)	343	■ 1.000	$\bar{G} = SL(2f, \mathbb{C}) \otimes SL(2c, \mathbb{C}) \supset SU(f)_L \otimes SU(f)_R \otimes SU(c)_L \otimes SU(c)_R$	$\bar{G} = SL(2f, \mathbb{C}) \otimes SL(2c, \mathbb{C}) \supset SU(f)_L \otimes SU(f)_R \otimes SU(c)_L \otimes SU(c)_R$	$\bar{G} = SL(2f, \mathbb{C}) \otimes SL(2c, \mathbb{C}) \supset SU(f)_L \otimes SU(f)_R \otimes SU(c)_L \otimes SU(c)_R$
mielke-geometrodynamics_343 EQ1045 (15.2.2)	343	■ 1.000	$\bar{L}(M) := M^4 \times \bar{G} \supset M^4 \times SL(2, \mathbb{C})$	$\bar{L}(M) := M^4 \times \bar{G} \supset M^4 \times SL(2, \mathbb{C})$	$\bar{L}(M) := M^4 \times \bar{G} \supset M^4 \times SL(2, \mathbb{C})$
mielke-geometrodynamics_344 EQ1046 (15.2.3)	344	■ 1.000	$\bar{\gamma} = \bar{E}_j^{\alpha}(m) \gamma_{\alpha} \otimes dx^j = \bar{E}_j dx^j$	$\bar{\gamma} = \bar{E}_j^{\alpha}(m) \gamma_{\alpha} \otimes dx^j = \bar{E}_j dx^j$	$\bar{\gamma} = \bar{E}_j^{\alpha}(m) \gamma_{\alpha} \otimes dx^j = \bar{E}_j dx^j$
mielke-geometrodynamics_344 EQ1048 (15.2.5)	344	■ 1.000	$\frac{1}{4} \text{Tr}(\bar{\gamma} \otimes_s \bar{\gamma}) = f_{ij} dx^i \otimes_s dx^j$	$\frac{1}{4} \text{Tr}(\bar{\gamma} \otimes_s \bar{\gamma}) = f_{ij} dx^i \otimes_s dx^j$	$\frac{1}{4} \text{Tr}(\bar{\gamma} \otimes_s \bar{\gamma}) = f_{ij} dx^i \otimes_s dx^j$
mielke-geometrodynamics_345 EQ1052 (15.2.9)	345	■ 1.000	$\bar{\omega}_i = \frac{1}{2} \Gamma_i^{(f)\alpha\beta j} \sigma_{\alpha\beta} \lambda_j^{(f)} \oplus \frac{1}{2} \Gamma_i^{(c)\alpha\beta j} \sigma_{\alpha\beta} \lambda_j^{(c)} \oplus A_i.$	$\bar{\omega}_i = \frac{1}{2} \Gamma_i^{(f)\alpha\beta j} \sigma_{\alpha\beta} \lambda_j^{(f)} \oplus \frac{1}{2} \Gamma_i^{(c)\alpha\beta j} \sigma_{\alpha\beta} \lambda_j^{(c)} \oplus A_i.$	$\bar{\omega}_i = \frac{1}{2} \Gamma_i^{(f)\alpha\beta j} \sigma_{\alpha\beta} \lambda_j^{(f)} \oplus \frac{1}{2} \Gamma_i^{(c)\alpha\beta j} \sigma_{\alpha\beta} \lambda_j^{(c)} \oplus A_i.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_345 EQ1053 (15.2.10)	345	■ 1.000	$\bar{\Theta}:=\bar{D}\bar{\gamma}=\bar{d}\bar{\gamma}-[\bar{\omega},\bar{\gamma}]$ $\bar{\omega}:=\bar{D}\bar{\gamma}=\bar{d}\bar{\gamma}-[\bar{\omega},\bar{\gamma}]$	$\bar{\Theta}:=\bar{D}\bar{\gamma}=\bar{d}\bar{\gamma}-[\bar{\omega},\bar{\gamma}]$	$\bar{\Theta}:=\bar{D}\bar{\gamma}=\bar{d}\bar{\gamma}-[\bar{\omega},\bar{\gamma}]$
mielke-geometrodynamics_345 EQ1054 (15.2.11)	345	■ 1.000	$\bar{\Omega} = d\bar{\omega} - \bar{\omega} \wedge \bar{\omega}$ $\bar{\Omega} = d\bar{\omega} - \bar{\omega} \wedge \bar{\omega}$	$\bar{\Omega} = d\bar{\omega} - \bar{\omega} \wedge \bar{\omega}$	$\bar{\Omega} = d\bar{\omega} - \bar{\omega} \wedge \bar{\omega}$
mielke-geometrodynamics_346 EQ1056 (15.2.13)	346	■ 1.000	$\psi = \left\{ \psi^{(q_f,q_c)} \mid q_f = 1, \dots, f; q_c = 1, \dots, c \right\}.$ $\psi = \{ \psi^{(q_f,q_c)} q_f = 1, \dots, f; q_c = 1, \dots, c \}.$	$\psi = \left\{ \psi^{(q_f,q_c)} \mid q_f = 1, \dots, f; q_c = 1, \dots, c \right\}.$	$\psi = \{ \psi^{(q_f,q_c)} q_f = 1, \dots, f; q_c = 1, \dots, c \}.$
mielke-geometrodynamics_354 EQ1061 (16.1.2)	354	■ 1.000	$B=\frac{2\hbar c}{\ell^{*2}}\Lambda_{\mathrm{eff}}=-\frac{3\kappa}{2}\frac{\hbar c}{\ell^{*4}}$ $B = \frac{2\hbar c}{\ell^{*2}}\Lambda_{\mathrm{eff}} = -\frac{3\kappa}{2}\frac{\hbar c}{\ell^{*4}}$	$B = \frac{2\hbar c}{\ell^{*2}}\Lambda_{\mathrm{eff}} = -\frac{3\kappa}{2}\frac{\hbar c}{\ell^{*4}}$	$B = \frac{2\hbar c}{\ell^{*2}}\Lambda_{\mathrm{eff}} = -\frac{3\kappa}{2}\frac{\hbar c}{\ell^{*4}}$
mielke-geometrodynamics_355 EQ1063 (16.2.2)	355	■ 1.000	$\square = \frac{1}{\sqrt{ f }}\partial_i\left(f^{ij}\sqrt{ f }\partial_j\right)$ $\square = \frac{1}{\sqrt{ f }}\partial_i\left(f^{ij}\sqrt{ f }\partial_j\right)$	$\square = \frac{1}{\sqrt{ f }}\partial_i\left(f^{ij}\sqrt{ f }\partial_j\right)$	$\square = \frac{1}{\sqrt{ f }}\partial_i(f^{ij}\sqrt{ f }\partial_j)$
mielke-geometrodynamics_355 EQ1064 (16.2.3) [re-fined: census]	355	■ 1.000	$\varphi = m\frac{c}{\hbar}\frac{1}{\rho}F_{nm}^t(\rho^*)Y_t^m(\vartheta,\phi)e^{-it\omega mc^2/\hbar}.$ $\varphi = m\frac{c}{\hbar}\frac{1}{\rho}F_{nm}^t(\rho^*)Y_t^m(\vartheta,\phi)e^{-it\omega mc^2/\hbar}.$	$\varphi = m\frac{c}{\hbar}\frac{1}{\rho}F_{nm}^t(\rho^*)Y_t^m(\vartheta,\phi)e^{-it\omega mc^2/\hbar}.$	$\varphi = m\frac{c}{\hbar}\frac{1}{\rho}F_{nm}^t(\rho^*)Y_t^m(\vartheta,\phi)e^{-it\omega mc^2/\hbar}.$
mielke-geometrodynamics_355 EQ1067 (16.2.6)	355	■ 1.000	$\beta = \frac{\sqrt{2\pi\kappa}}{\ell^*}/\frac{mc}{\hbar} = \frac{\ell_m}{\ell}.$ $\beta = \frac{\sqrt{2\pi\kappa}}{\ell^*}/\frac{mc}{\hbar} = \frac{\ell_m}{\ell}.$	$\beta = \frac{\sqrt{2\pi\kappa}}{\ell^*}/\frac{mc}{\hbar} = \frac{\ell_m}{\ell}.$	$\beta = \frac{\sqrt{2\pi\kappa}}{\ell^*}/\frac{mc}{\hbar} = \frac{\ell_m}{\ell}.$
mielke-geometrodynamics_358 EQ1072 (16.3.1)	358	■ 1.000	$\mathrm{Tr}\,\bar{Q}^2=-\frac{4b}{\alpha_e}Y+I(I+1)-\frac{1}{4}Y^2.$ $\mathrm{Tr}\,\bar{Q}^2=-\frac{4b}{\alpha_e}Y+I(I+1)-\frac{1}{4}Y^2.$	$\mathrm{Tr}\,\bar{Q}^2=-\frac{4b}{\alpha_e}Y+I(I+1)-\frac{1}{4}Y^2.$	$\mathrm{Tr}\,\bar{Q}^2=-\frac{4b}{\alpha_e}Y+I(I+1)-\frac{1}{4}Y^2.$
mielke-geometrodynamics_358 EQ1073 (16.3.2)	358	■ 1.000	$Q_e=I_3+\frac{1}{2}Y-\frac{2}{3}C.$ $Q_e = I_3 + \frac{1}{2}Y - \frac{2}{3}C.$	$Q_e = I_3 + \frac{1}{2}Y - \frac{2}{3}C.$	$Q_e = I_3 + \frac{1}{2}Y - \frac{2}{3}C.$
mielke-geometrodynamics_359 EQ1074 (16.3.3)	359	■ 1.000	$\frac{M^2}{M^{*2}}=\left\{1-bY+\frac{\alpha_e}{4}\left[I(I+1)-\frac{1}{4}Y^2\right]\right\}^2+\frac{1}{4}J(J+1)$ $\frac{M^2}{M^{*2}}=\left\{1-bY+\frac{\alpha_e}{4}\left[I(I+1)-\frac{1}{4}Y^2\right]\right\}^2+\frac{1}{4}J(J+1)$	$\frac{M^2}{M^{*2}}=\left\{1-bY+\frac{\alpha_e}{4}\left[I(I+1)-\frac{1}{4}Y^2\right]\right\}^2+\frac{1}{4}J(J+1)$	$\frac{M^2}{M^{*2}}=\left\{1-bY+\frac{\alpha_e}{4}[I(I+1)-\frac{1}{4}Y^2]\right\}^2+\frac{1}{4}J(J+1)$
mielke-geometrodynamics_360 EQ1075 (16.3.4)	360	■ 1.000	$T_{\bullet}=\frac{2\hbar c}{S_Ak_B}\left(r_+-\frac{M}{M^{*2}}\frac{\hbar}{c}\right).$ $T_{\bullet} = \frac{2\hbar c}{S_Ak_B}\left(r_+-\frac{M}{M^{*2}}\frac{\hbar}{c}\right).$	$T_{\bullet}=\frac{2\hbar c}{S_Ak_B}\left(r_+-\frac{M}{M^{*2}}\frac{\hbar}{c}\right).$	$T_{\bullet}=\frac{2\hbar c}{S_Ak_B}\left(r_+-\frac{M}{M^{*2}}\frac{\hbar}{c}\right).$
mielke-geometrodynamics_364 EQ1076 (A.1)	364	■ 1.000	$x^Ay_A:=\sum_{A=0}^n\sum_{B=0}^nx^Ay^Bg_{AB}.$ $x^Ay_A:=\sum_{A=0}^n\sum_{B=0}^nx^Ay^Bg_{AB}.$	$x^Ay_A:=\sum_{A=0}^n\sum_{B=0}^nx^Ay^Bg_{AB}.$	$x^Ay_A:=\sum_{A=0}^n\sum_{B=0}^nx^Ay^Bg_{AB}.$
mielke-geometrodynamics_364 EQ1077 (A.2)	364	■ 1.000	$c=\hbar=1.$ $c = \hbar = 1.$	$c = \hbar = 1.$	$c = \hbar = 1.$
mielke-geometrodynamics_365 EQ1080 (A.5)	365	■ 1.000	$\{\sigma^i,\sigma^j\}=g^{ij}$ $\{\sigma^i,\sigma^j\}=g^{ij}$	$\{\sigma^i,\sigma^j\}=g^{ij}$	$\{\sigma^i,\sigma^j\}=g^{ij}$
mielke-geometrodynamics_365 EQ1081 (A.6)	365	■ 1.000	$(\gamma^\alpha)^{\mathrm{T}}:=(\gamma^0,\gamma^k),$ $(\gamma^\alpha)^{\mathrm{T}}:=(\gamma^0,\gamma^k),$	$(\gamma^\alpha)^{\mathrm{T}}:=(\gamma^0,\gamma^k),$	$(\gamma^\alpha)^{\mathrm{T}}:=(\gamma^0,\gamma^k),$
mielke-geometrodynamics_365 EQ1083 (A.8)	365	■ 1.000	$\gamma^\alpha\gamma^\beta+\gamma^\beta\gamma^\alpha=2g^{\alpha\beta}$ $\gamma^\alpha\gamma^\beta+\gamma^\beta\gamma^\alpha=2g^{\alpha\beta}$	$\gamma^\alpha\gamma^\beta+\gamma^\beta\gamma^\alpha=2g^{\alpha\beta}$	$\gamma^\alpha\gamma^\beta+\gamma^\beta\gamma^\alpha=2g^{\alpha\beta}$
mielke-geometrodynamics_365 EQ1084 (A.9)	365	■ 1.000	$\gamma^{0+}=\gamma^0,\quad\gamma^{k+}=-\gamma^k,\quad\gamma^{5+}=\gamma^5.$ $\gamma^{0+}=\gamma^0,\quad\gamma^{k+}=-\gamma^k,\quad\gamma^{5+}=\gamma^5.$	$\gamma^{0+}=\gamma^0,\quad\gamma^{k+}=-\gamma^k,\quad\gamma^{5+}=\gamma^5.$	$\gamma^{0+}=\gamma^0,\quad\gamma^{k+}=-\gamma^k,\quad\gamma^{5+}=\gamma^5.$
mielke-geometrodynamics_366 EQ1087 (A.14)	366	■ 1.000	$C(1,3)=C(0,2)\otimes C(1,1).$ $C(1,3)=C(0,2)\otimes C(1,1).$	$C(1,3)=C(0,2)\otimes C(1,1).$	$C(1,3)=C(0,2)\otimes C(1,1).$
mielke-geometrodynamics_368 EQ1091 (B.1)	368	■ 1.000	$\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty(\wedge^pT^*(M)).$ $\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty(\wedge^pT^*(M)).$	$\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty(\wedge^pT^*(M)).$	$\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^\infty(\wedge^pT^*(M)).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_368 EQ1092 (B.2)		■ 1.000	$\begin{gathered} \alpha^{\{p\}} \wedge \beta^{\{q\}} := \\ \frac{1}{(p!) \cdot (q!)} A_{\alpha_1 \dots \alpha_p} \cdot B_{\beta_1 \dots \beta_q} \vartheta^{\alpha_1} \wedge \dots \wedge \vartheta^{\alpha_p} \\ \wedge \vartheta^{\beta_1} \wedge \dots \wedge \vartheta^{\beta_q} \end{gathered}$	$\alpha^{(p)} \wedge \beta^{(q)} := \frac{1}{(p!) \cdot (q!)} A_{\alpha_1 \dots \alpha_p} \cdot B_{\beta_1 \dots \beta_q} \vartheta^{\alpha_1} \wedge \dots \wedge \vartheta^{\alpha_p} \\ \wedge \vartheta^{\beta_1} \wedge \dots \wedge \vartheta^{\beta_q}$	$\alpha^{(p)} \wedge \beta^{(q)} := \frac{1}{(p!) \cdot (q!)} A_{\alpha_1 \dots \alpha_p} \cdot B_{\beta_1 \dots \beta_q} \vartheta^{\alpha_1} \wedge \dots \wedge \vartheta^{\alpha_p} \\ \wedge \vartheta^{\beta_1} \wedge \dots \wedge \vartheta^{\beta_q}$
mielke-geometrodynamics_368 EQ1093 (B.3)		■ 1.000	$\alpha^{\{p\}} \wedge \beta^{\{q\}} = (-1)^{pq} \beta^{\{q\}} \wedge \alpha^{\{p\}},$	$\alpha^{(p)} \wedge \beta^{(q)} = (-1)^{pq} \beta^{(q)} \wedge \alpha^{(p)},$	$\alpha^{(p)} \wedge \beta^{(q)} = (-1)^{pq} \beta^{(q)} \wedge \alpha^{(p)},$
mielke-geometrodynamics_368 EQ1094 (B.4)		■ 1.000	$D(M) := \sum_{p=0}^n \wedge^p T^*(M)$	$D(M) := \sum_{p=0}^n \wedge^p T^*(M)$	$D(M) := \sum_{p=0}^n \wedge^p T^*(M)$
mielke-geometrodynamics_369 EQ1095 (B.5)		■ 1.000	$d(\alpha^{\{p\}} \wedge \beta^{\{q\}}) = d\alpha^{\{p\}} \wedge \beta^{\{q\}} + (-1)^p \alpha^{\{p\}} \wedge d\beta^{\{q\}}$	$d\left(\alpha^{(p)} \wedge \beta^{(q)}\right) = d\alpha^{(p)} \wedge \beta^{(q)} + (-1)^p \alpha^{(p)} \wedge d\beta^{(q)}$	$d(\alpha^{(p)} \wedge \beta^{(q)}) = d\alpha^{(p)} \wedge \beta^{(q)} + (-1)^p \alpha^{(p)} \wedge d\beta^{(q)}$
mielke-geometrodynamics_369 EQ1097 (B.7)		■ 1.000	$\sigma: M \rightarrow N, \quad \sigma \in \mathcal{D}(M)$	$\sigma: M \rightarrow N, \quad \sigma \in \mathcal{D}(M)$	$\sigma: M \rightarrow N, \quad \sigma \in \mathcal{D}(M)$
mielke-geometrodynamics_369 EQ1098 (B.8)		■ 1.000	$\sigma_*: \wedge^p T^*(N) \rightarrow \wedge^p T^*(M)$	$\sigma_*: \wedge^p T^*(N) \rightarrow \wedge^p T^*(M)$	$\sigma_*: \wedge^p T^*(N) \rightarrow \wedge^p T^*(M)$
mielke-geometrodynamics_370 EQ1103 (B.15)		■ 1.000	$\alpha^{\{p\}} = (-1)^{p(n+1)+s} \alpha^{\{p\}}.$	$\alpha^{(p)} = (-1)^{p(n+1)+s} \alpha^{(p)}.$	$\alpha^{(p)} = (-1)^{p(n+1)+s} \alpha^{(p)}.$
mielke-geometrodynamics_370 EQ1105 (B.17)		■ 1.000	$(\alpha, \beta) := \int_M \alpha^{\{p\}} \wedge \beta^{\{p\}} = (\beta, \alpha)$	$(\alpha, \beta) := \int_M \alpha^{(p)} \wedge \beta^{(p)} = (\beta, \alpha)$	$(\alpha, \beta) := \int_M \alpha^{(p)} \wedge \beta^{(p)} = (\beta, \alpha)$
mielke-geometrodynamics_370 EQ1107 (B.19)		■ 1.000	$\square \alpha^{\{p\}} := d\delta \alpha^{\{p\}} + \delta d\alpha^{\{p\}}.$	$\square \alpha^{(p)} := d\delta \alpha^{(p)} + \delta d\alpha^{(p)}.$	$\square \alpha^{(p)} := d\delta \alpha^{(p)} + \delta d\alpha^{(p)}.$
mielke-geometrodynamics_371 EQ1108 (B.20)		■ 1.000	$(\alpha, \square \alpha) = (\alpha, d\delta \alpha) + (\alpha, \delta d\alpha) = (\delta \alpha, \delta \alpha) + (d\alpha, d\alpha).$	$(\alpha, \square \alpha) = (\alpha, d\delta \alpha) + (\alpha, \delta d\alpha) = (\delta \alpha, \delta \alpha) + (d\alpha, d\alpha).$	$(\alpha, \square \alpha) = (\alpha, d\delta \alpha) + (\alpha, \delta d\alpha) = (\delta \alpha, \delta \alpha) + (d\alpha, d\alpha).$
mielke-geometrodynamics_371 EQ1109 (B.21)		■ 1.000	$d\alpha = 0.$	$d\alpha = 0.$	$d\alpha = 0.$
mielke-geometrodynamics_371 EQ1111 (B.23)		■ 1.000	$\beta^{\{p\}} = d\alpha^{\{p-1\}} \in C^\infty(B(M))$	$\beta^{(p)} = d\alpha^{(p-1)} \in C^\infty(B(M))$	$\beta^{(p)} = d\alpha^{(p-1)} \in C^\infty(B(M))$
mielke-geometrodynamics_371 EQ1112 (B.24)		■ 1.000	$H(M) := Z(M) / B(M)$	$H(M) := Z(M) / B(M)$	$H(M) := Z(M) / B(M)$
mielke-geometrodynamics_371 EQ1113 (B.25)		■ 1.000	$H(M) = \sum_{p=0}^n H^p(M) := \sum_{p=0}^n Z^p(M) / B^p(M).$	$H(M) = \sum_{p=0}^n H^p(M) := \sum_{p=0}^n Z^p(M) / B^p(M).$	$H(M) = \sum_{p=0}^n H^p(M) := \sum_{p=0}^n Z^p(M) / B^p(M).$
mielke-geometrodynamics_371 EQ1115 (B.27)		■ 1.000	$H^n(M) = \wedge^n T^*(M) / B^n(M).$	$H^n(M) = \wedge^n T^*(M) / B^n(M).$	$H^n(M) = \wedge^n T^*(M) / B^n(M).$
mielke-geometrodynamics_371 EQ1116 (B.28)		■ 1.000	$H^0(M) = Z^0(M)$	$H^0(M) = Z^0(M)$	$H^0(M) = Z^0(M)$
mielke-geometrodynamics_372 EQ1117 (B.29)		■ 1.000	$H^0(M) \approx \mathbb{R}$	$H^0(M) \approx \mathbb{R}$	$H^0(M) \approx \mathbb{R}$
mielke-geometrodynamics_372 EQ1118 (B.30)		■ 1.000	$b_p := \dim H^p(M)$	$b_p := \dim H^p(M)$	$b_p := \dim H^p(M)$
mielke-geometrodynamics_372 EQ1119 (B.31)		■ 1.000	$f_M(t) = \sum_{p=0}^n b_p t^p$	$f_M(t) = \sum_{p=0}^n b_p t^p$	$f_M(t) = \sum_{p=0}^n b_p t^p,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_372 EQ1120 (B.32)	372	■ 1.000	$\chi(M):=\mathrm{f}_{\mathrm{M}}(-1)=\sum_{p=0}^n(-1)^pb_p,$	$\chi(M):=\mathrm{f}_{\mathrm{M}}(-1)=\sum_{p=0}^n(-1)^pb_p,$	$\chi(M):=\mathrm{f}_{\mathrm{M}}(-1)=\sum_{p=0}^n(-1)^pb_p,$
mielke-geometrodynamics_372 EQ1121 (B.33)	372	■ 1.000	$\mathrm{H}^{\{0\}}\left(S^n\right)=\mathrm{H}^n\left(S^n\right) \approx \mathbb{R}$	$H^0(S^n)=H^n(S^n)\approx\mathbb{R}$	$H^0(S^n)=H^n(S^n)\approx\mathbb{R}$
mielke-geometrodynamics_372 EQ1123 (B.35)	372	■ 1.000	$\mathrm{f}_{S^n}(t)=1+t^n,$	$\mathrm{f}_{S^n}(t)=1+t^n,$	$\mathrm{f}_{S^n}(t)=1+t^n,$
mielke-geometrodynamics_372 EQ1124 (B.36)	372	■ 1.000	$\chi\left(S^n\right)=\left\{\begin{array}{l} 0 \text { for } n=2 k+1, \\ 2 \text { for } n=2 k, \end{array}\right.$	$\chi(S^n)=\left\{\begin{array}{l} 0 \text { for } n=2 k+1, \\ 2 \text { for } n=2 k, \end{array}\right.$	$\chi(S^n)=\left\{\begin{array}{l} 0 \text { for } n=2 k+1, \\ 2 \text { for } n=2 k, \end{array}\right.$
mielke-geometrodynamics_372 EQ1125 (B.37)	372	■ 1.000	$\Delta_p:=\left\{x \in E^n \mid x=\sum_{i=0}^p \lambda^i \eta_i, \lambda^i \geq 0, \sum_{i=0}^p \lambda^i=1\right\}.$	$\Delta_p:=\left\{x \in E^n \mid x=\sum_{i=0}^p \lambda^i \eta_i, \lambda^i \geq 0, \sum_{i=0}^p \lambda^i=1\right\}.$	$\Delta_p:=\left\{x \in E^n \mid x=\sum_{i=0}^p \lambda^i \eta_i, \lambda^i \geq 0, \sum_{i=0}^p \lambda^i=1\right\}.$
mielke-geometrodynamics_373 EQ1127 (B.39)	373	■ 1.000	$\partial c=\sum_{i=0}^p(-1)^ic\left(\sum_{j=0}^{i-1} \lambda^j \eta_j+\sum_{j=i}^{p-1} \lambda^j \eta_{j+1}\right).$	$\partial c=\sum_{i=0}^p(-1)^ic\left(\sum_{j=0}^{i-1} \lambda^j \eta_j+\sum_{j=i}^{p-1} \lambda^j \eta_{j+1}\right).$	$\partial c=\sum_{i=0}^p(-1)^ic\left(\sum_{j=0}^{i-1} \lambda^j \eta_j+\sum_{j=i}^{p-1} \lambda^j \eta_{j+1}\right).$
mielke-geometrodynamics_373 EQ1129 (B.41)	373	■ 1.000	$H_{\{\mathrm{p}\}}(M, \mathbb{R}):=Z_{\{\mathrm{p}\}}(M, \mathbb{R}) / B_{\{\mathrm{p}\}}(M, \mathbb{R})$	$H_{\mathrm{p}}(M, \mathbb{R}):=Z_{\mathrm{p}}(M, \mathbb{R}) / B_{\mathrm{p}}(M, \mathbb{R})$	$H_{\mathrm{p}}(M, \mathbb{R}):=Z_{\mathrm{p}}(M, \mathbb{R}) / B_{\mathrm{p}}(M, \mathbb{R})$
mielke-geometrodynamics_373 EQ1130 (B.42)	373	■ 1.000	$\int_{c_{p+1}} d \alpha^{(p)}=\int_{\partial c_{p+1}} \alpha^{(p)}.$	$\int_{c_{p+1}} d \alpha^{(p)}=\int_{\partial c_{p+1}} \alpha^{(p)}.$	$\int_{c_{p+1}} d \alpha^{(p)}=\int_{\partial c_{p+1}} \alpha^{(p)}.$
mielke-geometrodynamics_373 EQ1132 (B.44)	373	■ 1.000	$\langle d \alpha, c\rangle_{\mathrm{M}}=\langle \alpha, \partial c\rangle_{\mathrm{M}}$	$\langle d \alpha, c\rangle_{\mathrm{M}}=\langle \alpha, \partial c\rangle_{\mathrm{M}}$	$\langle d \alpha, c\rangle_{\mathrm{M}}=\langle \alpha, \partial c\rangle_{\mathrm{M}}$
mielke-geometrodynamics_375 EQ1133 (C.1)	375	■ 1.000	$I_j:=\left.\frac{\partial g(\xi)}{\partial \xi^j}\right _{g=e} \in T_e(G) \approx \mathfrak{g}$	$I_j:=\left.\frac{\partial g(\xi)}{\partial \xi^j}\right _{g=e} \in T_e(G) \approx \mathfrak{g}$	$I_j:=\left.\frac{\partial g(\xi)}{\partial \xi^j}\right _{g=e} \in T_e(G) \approx \mathfrak{g}$
mielke-geometrodynamics_376 EQ1138 (C.6)	376	■ 1.000	$[\mathrm{AB}]:=\mathrm{A}^{\mathrm{i}} \mathrm{B}^{\mathrm{j}} \mathrm{c}_{\mathrm{ij}}^{\mathrm{k}} \mathrm{I}_{\mathrm{k}}.$	$[\mathrm{AB}]:=\mathrm{A}^{\mathrm{i}} \mathrm{B}^{\mathrm{j}} \mathrm{c}_{\mathrm{ij}}^{\mathrm{k}} \mathrm{I}_{\mathrm{k}}.$	$[\mathrm{AB}]:=\mathrm{A}^{\mathrm{i}} \mathrm{B}^{\mathrm{j}} \mathrm{c}_{\mathrm{ij}}^{\mathrm{k}} \mathrm{I}_{\mathrm{k}}.$
mielke-geometrodynamics_376 EQ1139 (C.7)	376	■ 1.000	$\rho_A^C\left(g_1\right) \rho_C^B\left(g_2\right)=\rho_A^B\left(g_1 g_2\right)$	$\rho_A^C(g_1)\rho_C^B(g_2)=\rho_A^B(g_1g_2)$	$\rho_A^C(g_1)\rho_C^B(g_2)=\rho_A^B(g_1g_2)$
mielke-geometrodynamics_376 EQ1140 (C.8)	376	■ 1.000	$[A, B] \longrightarrow \rho([A, B])=[\rho(A), \rho(B)].$	$[A, B] \longrightarrow \rho([A, B])=[\rho(A), \rho(B)].$	$[A, B] \longrightarrow \rho([A, B])=[\rho(A), \rho(B)].$
mielke-geometrodynamics_376 EQ1141 (C.9)	376	■ 1.000	$\operatorname{Ad} A(B):=[A, B] ; \quad A, B \in \mathfrak{g},$	$\operatorname{Ad} A(B):=[A, B]; \quad A, B \in \mathfrak{g},$	$\operatorname{Ad} A(B):=[A, B]; \quad A, B \in \mathfrak{g},$
mielke-geometrodynamics_376 EQ1142 (C.10)	376	■ 1.000	$(\operatorname{Ad} A)_j^k=c_{ij}^k A^i.$	$(\operatorname{Ad} A)_j^k=c_{ij}^k A^i.$	$(\operatorname{Ad} A)_j^k=c_{ij}^k A^i.$
mielke-geometrodynamics_376 EQ1144 (C.12)	376	■ 1.000	$(A, B)=2 N \operatorname{Tr}(A B)-2 \operatorname{Tr}(A) \operatorname{Tr}(B)$	$(A, B)=2 N \operatorname{Tr}(A B)-2 \operatorname{Tr}(A) \operatorname{Tr}(B)$	$(A, B)=2 N \operatorname{Tr}(A B)-2 \operatorname{Tr}(A) \operatorname{Tr}(B)$
mielke-geometrodynamics_377 EQ1146 (C.14)	377	■ 1.000	$g=\exp ^{\left(-\frac{i}{2} \xi^j \lambda_j\right)} \in S U(N), j=1, \ldots, N^2-1.$	$g=\exp ^{\left(-\frac{i}{2} \xi^j \lambda_j\right)} \in S U(N), j=1, \ldots, N^2-1.$	$g=\exp ^{\left(-\frac{i}{2} \xi^j \lambda_j\right)} \in S U(N), j=1, \ldots, N^2-1.$
mielke-geometrodynamics_377 EQ1148 (C.17)	377	■ 1.000	$\lambda_i \lambda_j=\frac{2}{N} \delta_{ij}+\left(d_{ij}^k+i f_{ij}^k\right) \lambda_k.$	$\lambda_i \lambda_j=\frac{2}{N} \delta_{ij}+\left(d_{ij}^k+i f_{ij}^k\right) \lambda_k.$	$\lambda_i \lambda_j=\frac{2}{N} \delta_{ij}+\left(d_{ij}^k+i f_{ij}^k\right) \lambda_k.$
mielke-geometrodynamics_377 EQ1149 (C.18)	377	■ 1.000	$\operatorname{Tr}\left(\lambda_i \lambda_j\right)=2 \delta_{ij}.$	$\operatorname{Tr}\left(\lambda_i \lambda_j\right)=2 \delta_{ij}.$	$\operatorname{Tr}\left(\lambda_i \lambda_j\right)=2 \delta_{ij}.$