

mielke-geometrodynamics: residuals

equations measured 2026-09-15T09:47:50Z against the model of 2026-09-15

Corrected (86)

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_EQ0006 (was)	33	■ 0.996	$\left.e_{\beta}\right\rceil\vartheta^{\alpha}=\delta^{\alpha}_{\beta}$.	$e_{\beta}\rfloor\vartheta^{\alpha}=\delta^{\alpha}_{\beta}$.	$e_{\beta}\rfloor\vartheta^{\alpha}=\delta^{\alpha}_{\beta}$.
mielke-geometrodynamics_EQ0006 (now)	33	—	$e_{\beta}\rfloor\vartheta^{\alpha}=\delta^{\alpha}_{\beta}$.	$e_{\beta}\rfloor\vartheta^{\alpha}=\delta^{\alpha}_{\beta}$.	$e_{\beta}\rfloor\vartheta^{\alpha}=\delta^{\alpha}_{\beta}$.
basis: census / census+ink					
mielke-geometrodynamics_EQ0119 (was)	59	■ 0.646	$\begin{aligned} \Sigma_{\alpha} &:= e_{\alpha}\rfloor L - \left(e_{\alpha}\rfloor dA\right) \wedge \frac{\partial L}{\partial(dA)} \\ &= \frac{1}{2}[(e_{\alpha}\rfloor F) \wedge H - F \wedge (e_{\alpha}\rfloor H)] \end{aligned}$	$\Sigma_{\alpha} := e_{\alpha}\rfloor L - (e_{\alpha}\rfloor dA) \wedge \frac{\partial L}{\partial(dA)} \\ = \frac{1}{2}[(e_{\alpha}\rfloor F) \wedge H - F \wedge (e_{\alpha}\rfloor H)]$	$\Sigma_{\alpha} := e_{\alpha}\rfloor L - (e_{\alpha}\rfloor dA) \wedge \frac{\partial L}{\partial(dA)} \\ = \frac{1}{2}[(e_{\alpha}\rfloor F) \wedge H - F \wedge (e_{\alpha}\rfloor H)]$
mielke-geometrodynamics_EQ0119 (now)	59	—	$\begin{aligned} \Sigma_{\alpha} &:= e_{\alpha}\rfloor L - \left(e_{\alpha}\rfloor dA\right) \wedge \frac{\partial L}{\partial(dA)} \\ &= \frac{1}{2}[(e_{\alpha}\rfloor F) \wedge H - F \wedge (e_{\alpha}\rfloor H)] \end{aligned}$	$\Sigma_{\alpha} := e_{\alpha}\rfloor L - (e_{\alpha}\rfloor dA) \wedge \frac{\partial L}{\partial(dA)} \\ = \frac{1}{2}[(e_{\alpha}\rfloor F) \wedge H - F \wedge (e_{\alpha}\rfloor H)]$	$\Sigma_{\alpha} := e_{\alpha}\rfloor L - (e_{\alpha}\rfloor dA) \wedge \frac{\partial L}{\partial(dA)} \\ = \frac{1}{2}[(e_{\alpha}\rfloor F) \wedge H - F \wedge (e_{\alpha}\rfloor H)]$
basis: census / census+ink					
mielke-geometrodynamics_EQ0120 (was)	60	■ 0.651	$d\Sigma_{\alpha} \simeq (e_{\alpha}\downarrow F) \wedge j$	$d\Sigma_{\alpha} \simeq (e_{\alpha}\downarrow F) \wedge j$	$d\Sigma_{\alpha} \simeq (e_{\alpha}\rfloor F) \wedge j$
mielke-geometrodynamics_EQ0120 (now)	60	—	$d\Sigma_{\alpha} \simeq (e_{\alpha}\rfloor F) \wedge j$	$d\Sigma_{\alpha} \simeq (e_{\alpha}\rfloor F) \wedge j$	$d\Sigma_{\alpha} \simeq (e_{\alpha}\rfloor F) \wedge j$
basis: census / census					
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_100 E00292 (was)		■ 0.165	$\begin{aligned} \mathfrak{f}_{\xi} L &= (\mathfrak{f}_{\xi} g_{\alpha\beta}) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + (\mathfrak{f}_{\xi} \vartheta^{\alpha}) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} \\ &+ (\mathfrak{f}_{\xi} \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\mathfrak{f}_{\xi} D\Psi) \wedge \frac{\partial L}{\partial D\Psi} \\ &= D \left[(\xi \rfloor \vartheta^{\alpha}) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + (\xi \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\xi \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} \right] \\ &- (\xi \rfloor \vartheta^{\alpha}) D \frac{\partial L}{\partial \vartheta^{\alpha}} + (\xi \rfloor T^{\alpha}) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + (\xi \lrcorner R_{\beta}^{\gamma}) \wedge I^{\beta}{}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} \\ &+ (\xi \rfloor D\Psi) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p (\xi \rfloor \Psi) \wedge D \frac{\delta L}{\delta \Psi} . \end{aligned}$	$\begin{aligned} \mathfrak{f}_{\xi} L &= (\mathfrak{f}_{\xi} g_{\alpha\beta}) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + (\mathfrak{f}_{\xi} \vartheta^{\alpha}) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} \\ &+ (\mathfrak{f}_{\xi} \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\mathfrak{f}_{\xi} D\Psi) \wedge \frac{\partial L}{\partial D\Psi} \\ &= D \left[(\xi \rfloor \vartheta^{\alpha}) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + (\xi \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\xi \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} \right] \\ &- (\xi \rfloor \vartheta^{\alpha}) D \frac{\partial L}{\partial \vartheta^{\alpha}} + (\xi \rfloor T^{\alpha}) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + (\xi \lrcorner R_{\beta}^{\gamma}) \wedge I^{\beta}{}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} \\ &+ (\xi \rfloor D\Psi) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p (\xi \rfloor \Psi) \wedge D \frac{\delta L}{\delta \Psi} . \end{aligned}$	
Conf. MathPix confidence: ■ ≥ 0.9 ■ $0.5-0.9$ ■ < 0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_E00292 (now)	100	—	$\begin{aligned} \mathfrak{f}_{\xi} L &= \left(\mathfrak{f}_{\xi} g_{\alpha\beta} \right) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \left(\mathfrak{f}_{\xi} \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} \\ &+ \left(\mathfrak{f}_{\xi} \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\mathfrak{f}_{\xi} D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \\ &= D \left[\left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \right] \\ &- \left(\xi \rfloor \vartheta^{\alpha} \right) D \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor T^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor R_{\beta}^{\gamma} \right) \wedge I^{\beta}{}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} \\ &+ \left(\xi \rfloor D\Psi \right) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p \left(\xi \rfloor \Psi \right) \wedge D \frac{\delta L}{\delta \Psi} . \end{aligned}$	$\begin{aligned} \mathfrak{f}_{\xi} L &= \left(\mathfrak{f}_{\xi} g_{\alpha\beta} \right) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \left(\mathfrak{f}_{\xi} \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} \\ &+ \left(\mathfrak{f}_{\xi} \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\mathfrak{f}_{\xi} D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \\ &= D \left[\left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \right] \\ &- \left(\xi \rfloor \vartheta^{\alpha} \right) D \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor T^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor R_{\beta}^{\gamma} \right) \wedge I^{\beta}{}_{\gamma} \Psi \wedge \frac{\partial L}{\partial D\Psi} \\ &+ \left(\xi \rfloor D\Psi \right) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p \left(\xi \rfloor \Psi \right) \wedge D \frac{\delta L}{\delta \Psi} . \end{aligned}$	
basis: census / census					
mielke-geometrodynamics_E00293 (was)	101	■0.052	$\xi \downharpoonleft L \equiv \left(\xi \downharpoonleft \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \downharpoonleft \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \downharpoonleft D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} .$	$\xi \downharpoonleft L \equiv \left(\xi \downharpoonleft \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \downharpoonleft \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \downharpoonleft D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} .$	
mielke-geometrodynamics_E00293 (now)	101	—	$\xi \rfloor L \equiv \left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} .$	$\xi \rfloor L \equiv \left(\xi \rfloor \vartheta^{\alpha} \right) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} .$	
basis: census / census					
mielke-geometrodynamics_E00294 (was)	101	■0.510	$\Sigma_{\alpha} = e_{\alpha} \downharpoonleft L - \left(e_{\alpha} \downharpoonleft D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} - \left(e_{\alpha} \downharpoonleft \Psi \right) \wedge \frac{\partial L}{\partial \Psi} .$	$\Sigma_{\alpha} = e_{\alpha} \downharpoonleft L - \left(e_{\alpha} \downharpoonleft D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} - \left(e_{\alpha} \downharpoonleft \Psi \right) \wedge \frac{\partial L}{\partial \Psi} .$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_101 E08294 (now)	101	—	$\Sigma_{\alpha} = e_{\alpha} \rfloor L - (e_{\alpha} \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$		$\Sigma_{\alpha} = e_{\alpha} \rfloor L - (e_{\alpha} \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} - (e_{\alpha} \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$
basis: census / census+ink					
mielke-geometrodynamics_101 E08295 (was)	101	■0.850	$\begin{aligned} D \Sigma_{\alpha} &\wedge \left(e_{\alpha} \rfloor T^{\beta} \right) \wedge \left(e_{\alpha} \rfloor R^{\beta\gamma} \right) \wedge \tau_{\beta\gamma} + F_{\alpha} \\ &\simeq (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma}. \end{aligned}$		$\begin{aligned} D \Sigma_{\alpha} &= (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma} + F_{\alpha} \\ &\simeq (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma}. \end{aligned}$
mielke-geometrodynamics_101 E08295 (now)	101	—	$\begin{aligned} D \Sigma_{\alpha} &\wedge \left(e_{\alpha} \rfloor T^{\beta} \right) \wedge \left(e_{\alpha} \rfloor R^{\beta\gamma} \right) \wedge \tau_{\beta\gamma} + F_{\alpha} \\ &\simeq (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma}. \end{aligned}$		$\begin{aligned} D \Sigma_{\alpha} &= (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma} + F_{\alpha} \\ &\simeq (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma}. \end{aligned}$
basis: census / census+ink					
mielke-geometrodynamics_101 E08296 (was)	101	■0.847	$F_{\alpha} = (e_{\alpha} \rfloor D\Psi) \frac{\delta L}{\delta \Psi} + (-1)^p (e_{\alpha} \rfloor \Psi) \wedge D \frac{\delta L}{\delta \Psi}$		$F_{\alpha} = (e_{\alpha} \rfloor D\Psi) \frac{\delta L}{\delta \Psi} + (-1)^p (e_{\alpha} \rfloor \Psi) \wedge D \frac{\delta L}{\delta \Psi}$
mielke-geometrodynamics_101 E08296 (now)	101	—	$F_{\alpha} = (e_{\alpha} \rfloor D\Psi) \frac{\delta L}{\delta \Psi} + (-1)^p (e_{\alpha} \rfloor \Psi) \wedge D \frac{\delta L}{\delta \Psi}$		$F_{\alpha} = (e_{\alpha} \rfloor D\Psi) \frac{\delta L}{\delta \Psi} + (-1)^p (e_{\alpha} \rfloor \Psi) \wedge D \frac{\delta L}{\delta \Psi}$
basis: census / census					
mielke-geometrodynamics_102 E08302 (was)	102	■0.950	$\begin{aligned} \widehat{D} \Sigma_{\alpha} &\wedge \left(e_{\alpha} \rfloor R_{\beta}^{\gamma} \right) \wedge \Delta^{\beta\gamma} \\ &- \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta\gamma}) \wedge \sigma^{\beta\gamma} \end{aligned}$		$\begin{aligned} \widehat{D} \Sigma_{\alpha} &\cong (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \Delta^{\beta\gamma} \\ &- \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta\gamma}) \wedge \sigma^{\beta\gamma} \end{aligned}$
mielke-geometrodynamics_102 E08302 (now)	102	—	$\begin{aligned} \widehat{D} \Sigma_{\alpha} &\cong (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \Delta^{\beta\gamma} \\ &- \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta\gamma}) \wedge \sigma^{\beta\gamma} \end{aligned}$		$\begin{aligned} \widehat{D} \Sigma_{\alpha} &\cong (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \Delta^{\beta\gamma} \\ &- \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta\gamma}) \wedge \sigma^{\beta\gamma} \end{aligned}$
basis: census / census+ink					
mielke-geometrodynamics_103 E08305 (was)	103	■0.793	$\varepsilon_{\mathrm{RC}} := \xi^{\alpha} \Sigma_{\alpha} + (e_{\beta} \rfloor \widehat{D} \xi^{\gamma}) \tau^{\beta}_{\gamma},$		$\varepsilon_{\mathrm{RC}} := \xi^{\alpha} \Sigma_{\alpha} + (e_{\beta} \rfloor \widehat{D} \xi^{\gamma}) \tau^{\beta}_{\gamma},$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_E00305 (now)	103	—	$\backslash\mathrm{varepsilon}_{\mathrm{RC}}:=\xi^{\alpha}\backslash\mathrm{Sigma}_{\alpha}+\backslash\mathrm{left}(e_{\beta}\backslash\mathrm{rfloor}\backslash\mathrm{overparen{D}}\backslash\xi^{\gamma}\backslash\mathrm{right})\backslash\mathrm{tau}^{\beta}_{\beta}\backslash\mathrm{gamma},$	$\varepsilon_{\mathrm{RC}}:=\xi^{\alpha}\Sigma_{\alpha}+\left(e_{\beta}\rfloor\widehat{D}\xi^{\gamma}\right)\tau^{\beta}_{\gamma},$	$\varepsilon_{\mathrm{RC}}:=\xi^{\alpha}\Sigma_{\alpha}+(e_{\beta}\rfloor\widehat{D}\xi^{\gamma})\tau^{\beta}_{\gamma},$
basis: census / census					
mielke-geometrodynamics_E00306 (was)	103	■ 0.406	$\backslash\mathrm{left}.\backslash\mathrm{widetilde{\varepsilon}}_{\mathrm{MA}}:=\backslash\mathrm{varepsilon}_{\mathrm{RC}}+\backslash\xi\backslash\mathrm{right}\backslash\mathrm{rcorner}L\backslash\mathrm{cong}-\backslash\mathrm{left}(\backslash\mathrm{mathscr{L}}_{\xi}\backslash\mathrm{Gamma}_{\beta})^{\gamma}\backslash\mathrm{right})\backslash\mathrm{wedge}H^{\beta}_{\gamma}+dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}}:=\varepsilon_{\mathrm{RC}}+\xi\lrcorner L\cong-(\mathcal{L}_{\xi}\Gamma_{\beta}^{\gamma})\wedge H^{\beta}_{\gamma}+dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}}:=\varepsilon_{\mathrm{RC}}+\xi\rfloor L\cong-(\mathcal{L}_{\xi}\Gamma_{\beta}^{\gamma})\wedge H^{\beta}_{\gamma}+dH_{\infty}$
mielke-geometrodynamics_E00306 (now)	103	—	$\backslash\mathrm{widetilde{\varepsilon}}_{\mathrm{MA}}:=\backslash\mathrm{varepsilon}_{\mathrm{RC}}+\backslash\xi\backslash\mathrm{rfloor}L\backslash\mathrm{cong}-\backslash\mathrm{left}(\backslash\mathrm{mathscr{L}}_{\xi}\backslash\mathrm{Gamma}_{\beta})^{\gamma}\backslash\mathrm{right})\backslash\mathrm{wedge}H^{\beta}_{\gamma}+dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}}:=\varepsilon_{\mathrm{RC}}+\xi\rfloor L\cong-(\mathcal{L}_{\xi}\Gamma_{\beta}^{\gamma})\wedge H^{\beta}_{\gamma}+dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}}:=\varepsilon_{\mathrm{RC}}+\xi\rfloor L\cong-(\mathcal{L}_{\xi}\Gamma_{\beta}^{\gamma})\wedge H^{\beta}_{\gamma}+dH_{\infty}$
basis: census / census+ink					
mielke-geometrodynamics_E00311 (was)	104	■ 0.111	$\backslash\mathrm{begin}{aligned}\backslash\mathrm{widetilde{\leftarrow}}_{\mathrm{y}}\backslash\mathrm{widetilde{D}}\backslash\mathrm{xi}^{\alpha}\backslash\mathrm{right}\backslash\mathrm{rfloor}\backslash\mathrm{left}(\backslash\mathrm{widetilde{D}}\backslash\mathrm{widetilde{D}}\backslash\mathrm{xi}^{\alpha}\backslash\mathrm{right})\backslash\mathrm{wedge}H^{\beta}_{\gamma}+dH_{\infty}$	$\overleftarrow{\widetilde{y}}\widetilde{D}\widetilde{\xi}^{\alpha}=\widetilde{y}\rfloor\left(\widetilde{D}\widetilde{D}\widetilde{\xi}^{\alpha}\right)+\widetilde{D}(\widetilde{y}\rfloor\widetilde{D}\widetilde{\xi}^{\alpha})=yJ\left(DD\xi^{\alpha}+R^{(T)\alpha}\right)=0,$	$\widetilde{\mathfrak{L}}_{\widetilde{y}}\widetilde{D}\widetilde{\xi}^{\alpha}=\widetilde{y}\rfloor\left(\widetilde{D}\widetilde{D}\widetilde{\xi}^{\alpha}\right)+\widetilde{D}(\widetilde{y}\rfloor\widetilde{D}\widetilde{\xi}^{\alpha})=y\rfloor(DD\xi^{\alpha}+R^{(T)\alpha})=0,$
mielke-geometrodynamics_E00311 (now)	104	—	$\backslash\mathrm{begin}{aligned}\backslash\mathrm{widetilde{\leftarrow}}_{\mathrm{y}}\backslash\mathrm{widetilde{D}}\backslash\mathrm{xi}^{\alpha}\backslash\mathrm{right}\backslash\mathrm{rfloor}\backslash\mathrm{left}(\backslash\mathrm{widetilde{D}}\backslash\mathrm{widetilde{D}}\backslash\mathrm{xi}^{\alpha}\backslash\mathrm{right})\backslash\mathrm{wedge}H^{\beta}_{\gamma}+dH_{\infty}$	$\overleftarrow{\widetilde{y}}\widetilde{D}\widetilde{\xi}^{\alpha}=\widetilde{y}\rfloor\left(\widetilde{D}\widetilde{D}\widetilde{\xi}^{\alpha}\right)+\widetilde{D}(\widetilde{y}\rfloor\widetilde{D}\widetilde{\xi}^{\alpha})=y\rfloor\left(DD\xi^{\alpha}+R^{(T)\alpha}\right)=0,$	$\widetilde{\mathfrak{L}}_{\widetilde{y}}\widetilde{D}\widetilde{\xi}^{\alpha}=\widetilde{y}\rfloor\left(\widetilde{D}\widetilde{D}\widetilde{\xi}^{\alpha}\right)+\widetilde{D}(\widetilde{y}\rfloor\widetilde{D}\widetilde{\xi}^{\alpha})=y\rfloor(DD\xi^{\alpha}+R^{(T)\alpha})=0,$
basis: census / census					
mielke-geometrodynamics_E00312 (was)	104	■ 0.786	$\backslash\mathrm{begin}{aligned}\Delta\xi^{\alpha}&=\backslash\mathrm{left}.\backslash\mathrm{left}.\backslash\mathrm{oint}_{\mathrm{C}}\backslash\mathrm{y}\backslash\mathrm{right}\backslash\mathrm{rfloor}\backslash\mathrm{left}(D\backslash\mathrm{D}\backslash\mathrm{xi}^{\alpha}\backslash\mathrm{right})=\backslash\mathrm{oint}_{\mathrm{C}}\backslash\mathrm{y}\backslash\mathrm{right}\backslash\mathrm{rfloor}R^{\alpha}(\mathrm{T})\backslash\mathrm{alpha}=\backslash\mathrm{int}_{\mathrm{S}}R^{\alpha}(\mathrm{T})\backslash\mathrm{alpha}\backslash\mathrm{right})\backslash\mathrm{wedge}d\mathrm{y}^{\mathrm{j}}\backslash\mathrm{right})\backslash\mathrm{int}_{\mathrm{S}}d\mathrm{y}^{\mathrm{i}}\backslash\mathrm{wedge}d\mathrm{y}^{\mathrm{j}}\backslash\mathrm{end}{aligned}$	$\Delta\xi^{\alpha}=-\oint_C y\rfloor(DD\xi^{\alpha})=\oint_C y\rfloor R^{(T)\alpha}=\int_S R^{(T)\alpha}\simeq\frac{1}{2}(T_{ij}^{\alpha}-R_{ij\beta}^{\alpha}\xi^{\beta})\int_S dy^i\wedge dy^j$	$\Delta\xi^{\alpha}=-\oint_C y\rfloor(DD\xi^{\alpha})=\oint_C y\rfloor R^{(T)\alpha}=\int_S R^{(T)\alpha}\simeq\frac{1}{2}(T_{ij}^{\alpha}-R_{ij\beta}^{\alpha}\xi^{\beta})\int_S dy^i\wedge dy^j.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_115 EQ0340 (was)	115	■ 0.962	$\begin{aligned} G_{\alpha} &:= \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} \\ &= G_{\alpha}^{\{\}} + \frac{(-1)^s}{12} (e_{\alpha} \rfloor \mathcal{A} \wedge {}^* \mathcal{A} - \frac{1}{3} \mathcal{A} \wedge e_{\alpha} \rfloor {}^* \mathcal{A}) + \frac{(-1)^s}{6} \vartheta_{\alpha} \wedge d\mathcal{A} \end{aligned}$	$G_{\alpha} := \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma}$ $= G_{\alpha}^{\{\}} + \frac{(-1)^s}{12} \left(e_{\alpha} \rfloor \mathcal{A} \wedge {}^* \mathcal{A} - \frac{1}{3} \mathcal{A} \wedge e_{\alpha} \rfloor {}^* \mathcal{A} \right) + \frac{(-1)^s}{6} \vartheta_{\alpha} \wedge d\mathcal{A}$	$G_{\alpha} := \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} \quad (5.3)$ $= G_{\alpha}^{\{\}} + \frac{(-1)^s}{12} \left(e_{\alpha} \rfloor \mathcal{A} \wedge {}^* \mathcal{A} - \frac{1}{3} \mathcal{A} \wedge e_{\alpha} \rfloor {}^* \mathcal{A} \right) + \frac{(-1)^s}{6} \vartheta_{\alpha} \wedge d\mathcal{A}$
mielke-geometrodynamics_115 EQ0340 (now)	115	—	$\begin{aligned} G_{\alpha} &:= \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} \\ &= G_{\alpha}^{\{\}} + \frac{(-1)^s}{12} \left(e_{\alpha} \rfloor \mathcal{A} \wedge {}^* \mathcal{A} - \frac{1}{3} \mathcal{A} \wedge e_{\alpha} \rfloor {}^* \mathcal{A} \right) + \frac{(-1)^s}{6} \vartheta_{\alpha} \wedge d\mathcal{A} \end{aligned}$	$G_{\alpha} := \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma}$ $= G_{\alpha}^{\{\}} + \frac{(-1)^s}{12} \left(e_{\alpha} \rfloor \mathcal{A} \wedge {}^* \mathcal{A} - \frac{1}{3} \mathcal{A} \wedge e_{\alpha} \rfloor {}^* \mathcal{A} \right) + \frac{(-1)^s}{6} \vartheta_{\alpha} \wedge d\mathcal{A}$	$G_{\alpha} := \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} \quad (5.3)$ $= G_{\alpha}^{\{\}} + \frac{(-1)^s}{12} \left(e_{\alpha} \rfloor \mathcal{A} \wedge {}^* \mathcal{A} - \frac{1}{3} \mathcal{A} \wedge e_{\alpha} \rfloor {}^* \mathcal{A} \right) + \frac{(-1)^s}{6} \vartheta_{\alpha} \wedge d\mathcal{A}$
basis: census / census+ink					
mielke-geometrodynamics_127 EQ0372 (was)	127	■ 0.437	$\left. \left(\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} \right) \stackrel{(-)}{\Pi}_{\alpha} \right] \left(\stackrel{(+)}{\Pi}^{\beta} \wedge \stackrel{(-)}{\Pi}_{\beta} \right) = \Sigma_{\alpha},$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} \mp \frac{i}{4} \ell^2 e_{\alpha} \rfloor \left(\stackrel{(+)}{\Pi}^{\beta} \wedge \stackrel{(-)}{\Pi}_{\beta} \right) = \Sigma_{\alpha},$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} \mp \frac{i}{4} \ell^2 e_{\alpha} \rfloor \left(\stackrel{(+)}{\Pi}^{\beta} \wedge \stackrel{(-)}{\Pi}_{\beta} \right) = \Sigma_{\alpha},$
mielke-geometrodynamics_127 EQ0372 (now)	127	—	$\left. \left(\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} \right) \stackrel{(-)}{\Pi}_{\alpha} \right] \left(\stackrel{(+)}{\Pi}^{\beta} \wedge \stackrel{(-)}{\Pi}_{\beta} \right) = \Sigma_{\alpha},$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} \mp \frac{i}{4} \ell^2 e_{\alpha} \rfloor \left(\stackrel{(+)}{\Pi}^{\beta} \wedge \stackrel{(-)}{\Pi}_{\beta} \right) = \Sigma_{\alpha},$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} \mp \frac{i}{4} \ell^2 e_{\alpha} \rfloor \left(\stackrel{(+)}{\Pi}^{\beta} \wedge \stackrel{(-)}{\Pi}_{\beta} \right) = \Sigma_{\alpha},$
basis: census / census+ink					
mielke-geometrodynamics_128 EQ0378 (was)	128	■ 0.136	$D \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} \wedge \vartheta^{\beta} = 0, \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} = - \stackrel{(\pm)}{R}_{\alpha}{}^{\beta} \wedge \stackrel{(\pm)}{\Pi}_{\beta} = \pm \frac{i}{2} \ell^2 \left(e_{\alpha} \rfloor \stackrel{(\pm)}{\Pi}^{\beta} \right) \wedge \Sigma_{\beta} \cong 0.$	$DD\vartheta^{\alpha} = R_{\beta}{}^{\alpha} \wedge \vartheta^{\beta} = 0, \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} = - \stackrel{(\pm)}{R}_{\alpha}{}^{\beta} \wedge \stackrel{(\pm)}{\Pi}_{\beta} = \pm \frac{i}{2} \ell^2 \left(e_{\alpha} \rfloor \stackrel{(\pm)}{\Pi}^{\beta} \right) \wedge \Sigma_{\beta} \cong 0.$	$DD\vartheta^{\alpha} = R_{\beta}{}^{\alpha} \wedge \vartheta^{\beta} = 0, \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} = - \stackrel{(\pm)}{R}_{\alpha}{}^{\beta} \wedge \stackrel{(\pm)}{\Pi}_{\beta} = \pm \frac{i}{2} \ell^2 (e_{\alpha} \rfloor \stackrel{(\pm)}{\Pi}^{\beta}) \wedge \Sigma_{\beta} \cong 0.$
mielke-geometrodynamics_128 EQ0378 (now)	128	—	$D \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} \wedge \vartheta^{\beta} = 0, \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} = - \stackrel{(\pm)}{R}_{\alpha}{}^{\beta} \wedge \stackrel{(\pm)}{\Pi}_{\beta} = \pm \frac{i}{2} \ell^2 \left(e_{\alpha} \rfloor \stackrel{(\pm)}{\Pi}^{\beta} \right) \wedge \Sigma_{\beta} \cong 0.$	$DD\vartheta^{\alpha} = R_{\beta}{}^{\alpha} \wedge \vartheta^{\beta} = 0, \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} = - \stackrel{(\pm)}{R}_{\alpha}{}^{\beta} \wedge \stackrel{(\pm)}{\Pi}_{\beta} = \pm \frac{i}{2} \ell^2 \left(e_{\alpha} \rfloor \stackrel{(\pm)}{\Pi}^{\beta} \right) \wedge \Sigma_{\beta} \cong 0.$	$DD\vartheta^{\alpha} = R_{\beta}{}^{\alpha} \wedge \vartheta^{\beta} = 0, \stackrel{(\pm)}{D} \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} = - \stackrel{(\pm)}{R}_{\alpha}{}^{\beta} \wedge \stackrel{(\pm)}{\Pi}_{\beta} = \pm \frac{i}{2} \ell^2 (e_{\alpha} \rfloor \stackrel{(\pm)}{\Pi}^{\beta}) \wedge \Sigma_{\beta} \cong 0.$
basis: census / census					
mielke-geometrodynamics_129 EQ0381 (was)	129	■ 0.588	$\left. \left(\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} \right) \stackrel{(\mp)}{\Pi}_{\alpha} \right] \left(e_{\alpha} \rfloor \stackrel{(\mp)}{\Pi}^{\beta} \right) \wedge \stackrel{(\pm)}{\Pi}_{\beta},$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} := \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} \pm \frac{i}{2} \ell^2 \left(e_{\alpha} \rfloor \stackrel{(\mp)}{\Pi}^{\beta} \right) \wedge \stackrel{(\pm)}{\Pi}_{\beta},$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} := \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} \pm \frac{i}{2} \ell^2 (e_{\alpha} \rfloor \stackrel{(\mp)}{\Pi}^{\beta}) \wedge \stackrel{(\pm)}{\Pi}_{\beta},$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
mielke-geometrodynamics_129 EQ0381 (now)	129	—	$\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\alpha:=D\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\alpha\pm\frac{i}{2}\ell^2\left(e_\alpha\downarrow\stackrel{\scriptscriptstyle{(\mp)}}{\Pi}\right)\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_\beta,$	$\stackrel{(\pm)}{D}\stackrel{(\pm)}{\Pi}_\alpha:=D\stackrel{(\pm)}{\Pi}_\alpha\pm\frac{i}{2}\ell^2\left(e_\alpha\downarrow\stackrel{(\mp)}{\Pi}\right)\wedge\stackrel{(\pm)}{\Pi}_\beta,$	$\stackrel{(\pm)}{D}\stackrel{(\pm)}{\Pi}_\alpha:=D\stackrel{(\pm)}{\Pi}_\alpha\pm\frac{i}{2}\ell^2\left(e_\alpha\downarrow\stackrel{(\mp)}{\Pi}^\beta\right)\wedge\stackrel{(\pm)}{\Pi}_\beta,$
basis: census / census					
mielke-geometrodynamics_129 EQ0382 (was)	129	■ 0.087	$\stackrel{\scriptscriptstyle{(\pm)}}{\Gamma}_\alpha{}^\beta:=\Gamma_\alpha{}^\beta+i\stackrel{\scriptscriptstyle{(\mp)}}{A}^\beta{}_\beta,\quad\stackrel{\scriptscriptstyle{(\pm)}}{A}^\beta{}_\beta:=\pm\frac{1}{2}\ell^2e_\alpha\downarrow\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}^\beta,$	$\stackrel{(\pm)}{\Gamma}_\alpha{}^\beta:=\Gamma_\alpha{}^\beta+i\stackrel{(\mp)}{A}^\beta{}_\beta,\quad\stackrel{(\pm)}{A}^\beta{}_\beta:=\pm\frac{1}{2}\ell^2e_\alpha\downarrow\stackrel{(\pm)}{\Pi}^\beta,$	$\stackrel{(\pm)}{\Gamma}_\alpha{}^\beta:=\Gamma_\alpha{}^\beta+i\stackrel{(\mp)}{A}_\alpha{}^\beta,\quad\stackrel{(\pm)}{A}_\alpha{}^\beta:=\pm\frac{1}{2}\ell^2e_\alpha\downarrow\stackrel{(\pm)}{\Pi}^\beta,$
mielke-geometrodynamics_129 EQ0382 (now)	129	—	$\stackrel{\scriptscriptstyle{(\pm)}}{\Gamma}_\alpha{}^\beta:=\Gamma_\alpha{}^\beta+i\stackrel{\scriptscriptstyle{(\mp)}}{A}^\beta{}_\beta,\quad\stackrel{\scriptscriptstyle{(\pm)}}{A}^\beta{}_\beta:=\pm\frac{1}{2}\ell^2e_\alpha\downarrow\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}^\beta,$	$\stackrel{(\pm)}{\Gamma}_\alpha{}^\beta:=\Gamma_\alpha{}^\beta+i\stackrel{(\mp)}{A}^\beta{}_\beta,\quad\stackrel{(\pm)}{A}^\beta{}_\beta:=\pm\frac{1}{2}\ell^2e_\alpha\downarrow\stackrel{(\pm)}{\Pi}^\beta,$	$\stackrel{(\pm)}{\Gamma}_\alpha{}^\beta:=\Gamma_\alpha{}^\beta+i\stackrel{(\mp)}{A}_\alpha{}^\beta,\quad\stackrel{(\pm)}{A}_\alpha{}^\beta:=\pm\frac{1}{2}\ell^2e_\alpha\downarrow\stackrel{(\pm)}{\Pi}^\beta,$
basis: census / census					
mielke-geometrodynamics_130 EQ0387 (was)	130	■ 0.994	$\left.\stackrel{\scriptscriptstyle{(\pm)}}{\Psi}\right :=\left.\left(dt\wedge\Psi_\perp\right)\right ,\quad\left.\stackrel{\scriptscriptstyle{(\pm)}}{\Psi}_\perp\right :=\left.\left(dt\wedge\Psi\right)\right ,$	$\stackrel{\scriptscriptstyle{(\pm)}}{\Psi}:=dt\wedge\Psi_\perp;\quad\Psi_\perp:=n\downarrow\Psi,$	$\stackrel{\scriptscriptstyle{(\pm)}}{\Psi}:=dt\wedge\Psi_\perp;\quad\Psi_\perp:=n\downarrow\Psi,$
mielke-geometrodynamics_130 EQ0387 (now)	130	—	$\left.\stackrel{\scriptscriptstyle{(\pm)}}{\Psi}\right :=\left.\left(dt\wedge\Psi_\perp\right)\right ,\quad\left.\stackrel{\scriptscriptstyle{(\pm)}}{\Psi}_\perp\right :=\left.\left(dt\wedge\Psi\right)\right ,$	$\stackrel{\scriptscriptstyle{(\pm)}}{\Psi}:=dt\wedge\Psi_\perp;\quad\Psi_\perp:=n\downarrow\Psi,$	$\stackrel{\scriptscriptstyle{(\pm)}}{\Psi}:=dt\wedge\Psi_\perp;\quad\Psi_\perp:=n\downarrow\Psi,$
basis: census / census+ink					
mielke-geometrodynamics_130 EQ0388 (was)	130	■ 0.993	$\left.\underline{\stackrel{\scriptscriptstyle{(\pm)}}{\Psi}}\right :=\left.\left(dt\wedge\Psi\right)\right ,\quad\left.\underline{\stackrel{\scriptscriptstyle{(\pm)}}{\Psi}}_\perp\right :=\left.\left(dt\wedge\Psi_\perp\right)\right ,$	$\underline{\Psi}:=n\downarrow(dt\wedge\Psi)=(1-\stackrel{\scriptscriptstyle{(\pm)}}{\Psi})\Psi,\quad n\downarrow\underline{\Psi}\equiv 0.$	$\underline{\Psi}:=n\downarrow(dt\wedge\Psi)=(1-\stackrel{\scriptscriptstyle{(\pm)}}{\Psi})\Psi,\quad n\downarrow\underline{\Psi}\equiv 0.$
mielke-geometrodynamics_130 EQ0388 (now)	130	—	$\underline{\stackrel{\scriptscriptstyle{(\pm)}}{\Psi}}:=\left.\left(dt\wedge\Psi\right)\right ,\quad\left.\underline{\stackrel{\scriptscriptstyle{(\pm)}}{\Psi}}_\perp\right :=\left.\left(dt\wedge\Psi_\perp\right)\right ,$	$\underline{\Psi}:=n\downarrow(dt\wedge\Psi)=(1-\stackrel{\scriptscriptstyle{(\pm)}}{\Psi})\Psi,\quad n\downarrow\underline{\Psi}\equiv 0.$	$\underline{\Psi}:=n\downarrow(dt\wedge\Psi)=(1-\stackrel{\scriptscriptstyle{(\pm)}}{\Psi})\Psi,\quad n\downarrow\underline{\Psi}\equiv 0.$
basis: census / census+ink					
mielke-geometrodynamics_131 EQ0391 (was)	131	■ 0.066	$\begin{aligned}\Sigma_\alpha&:=\frac{\delta L}{\delta\vartheta^\alpha}=e_\alpha\downarrow L-(e_\alpha\downarrow D\Psi)\wedge\frac{\partial L}{\partial D\Psi}-(e_\alpha\downarrow\Psi)\wedge\frac{\partial L}{\partial\Psi}\\&\quad-(e_\alpha\downarrow T^\beta)\wedge\frac{\partial L}{\partial T^\beta}+D\frac{\partial L}{\partial T^\alpha}-(e_\alpha\downarrow R_\beta^\gamma)\wedge\frac{\partial L}{\partial R_\beta^\gamma}.\end{aligned}$	$\Sigma_\alpha:=\frac{\delta L}{\delta\vartheta^\alpha}=e_\alpha\downarrow L-(e_\alpha\downarrow D\Psi)\wedge\frac{\partial L}{\partial D\Psi}-(e_\alpha\downarrow\Psi)\wedge\frac{\partial L}{\partial\Psi}\\-(e_\alpha\downarrow T^\beta)\wedge\frac{\partial L}{\partial T^\beta}+D\frac{\partial L}{\partial T^\alpha}-(e_\alpha\downarrow R_\beta^\gamma)\wedge\frac{\partial L}{\partial R_\beta^\gamma}.$	$\Sigma_\alpha:=\frac{\delta L}{\delta\vartheta^\alpha}=e_\alpha\downarrow L-(e_\alpha\downarrow D\Psi)\wedge\frac{\partial L}{\partial D\Psi}-(e_\alpha\downarrow\Psi)\wedge\frac{\partial L}{\partial\Psi}\\-(e_\alpha\downarrow T^\beta)\wedge\frac{\partial L}{\partial T^\beta}+D\frac{\partial L}{\partial T^\alpha}-(e_\alpha\downarrow R_\beta^\gamma)\wedge\frac{\partial L}{\partial R_\beta^\gamma}.$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value LaTeX source MathPix’s reading, unchanged; Rendered the same reading, with a display environment (align/alignat/flalign/eqnarray/gather/multline/split) mapped to its in-math form so it can typeset in this cell — the two may legitimately differ Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured [refined: <i>basis</i>] beside an identifier: the LaTeX-source and Rendered cells both show a VERIFIED refinement, not MathPix’s own reading — <i>basis</i> names what verified it (e.g. ink, source). The original reading is unchanged in the model but is not the one drawn here.					

