

BradleyGastaldiTerilla2023: equation evidence

Display equations

Identifier	Page	Conf.	LaTeX source	Rendered	Image
BradleyGastaldiTerilla2023EQ0008 (4)	7	0.841	$\begin{array}{r} X=\{-, /, \odot, 1, 2, 3, 4, 5, 6, 7, 8, 9, =, a, b, c, d, e, f, \\ g, h, i, j, k, l, m, n, o, p, q, r, s, t, u, v, w, x, y, z, e \} . \end{array}$	$X = \{-, /, \odot, 1, 2, 3, 4, 5, 6, 7, 8, 9, =, a, b, c, d, e, f, g, h, i, j, k, l, m, n, o, p, q, r, s, t, u, v, w, x, y, z, e\}.$	$X = \{-, /, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, =, a, b, c, d, e, f, g, h, i, j, k, l, m, n, o, p, q, r, s, t, u, v, w, x, y, z, \acute{e}\}.$
BradleyGastaldiTerilla2023EQ0004	4	0.992	$\text{aardvark} \mapsto (0.632, 0.370, -0.620, \dots, -0.475).$	$\text{aardvark} \mapsto (0.632, 0.370, -0.620, \dots, -0.475).$	$\text{aardvark} \mapsto (0.632, 0.370, -0.620, \dots, -0.475).$
BradleyGastaldiTerilla2023EQ0001 (1)	3	0.999	$\begin{aligned} & \& f: \mathbb{R}^n \xrightarrow{f_1} \mathbb{R}^{n_1} \xrightarrow{f_2} \mathbb{R}^{n_2} \xrightarrow{f_3} \dots \\ & \mathbb{R}^{n_1} \xrightarrow{f_2} \mathbb{R}^{n_2} \xrightarrow{f_3} \dots \\ & \mathbb{R}^{n_2} \xrightarrow{f_3} \mathbb{R}^{n_3} \xrightarrow{f_4} \dots \\ & \& \dots \xrightarrow{f_K} \mathbb{R}^{n_K} \xrightarrow{g} \mathbb{R}^m \\ & f_i(\mathbf{x}) = a(\mathbf{M}_i \mathbf{x} + b_i), \end{aligned}$	$\begin{aligned} f: \mathbb{R}^n &\xrightarrow{f_1} \mathbb{R}^{n_1} \xrightarrow{f_2} \mathbb{R}^{n_2} \xrightarrow{f_3} \dots \\ &\dots \xrightarrow{f_K} \mathbb{R}^{n_K} \xrightarrow{g} \mathbb{R}^m \\ f_i(\mathbf{x}) &= a(\mathbf{M}_i \mathbf{x} + b_i), \end{aligned}$	$\begin{aligned} f: \mathbb{R}^n &\xrightarrow{f_1} \mathbb{R}^{n_1} \xrightarrow{f_2} \mathbb{R}^{n_2} \xrightarrow{f_3} \dots \\ &\dots \xrightarrow{f_K} \mathbb{R}^{n_K} \xrightarrow{g} \mathbb{R}^m \\ f_i(\mathbf{x}) &= a(\mathbf{M}_i \mathbf{x} + b_i), \end{aligned}$
BradleyGastaldiTerilla2023EQ0002	4	0.999	$\begin{gathered} \text{\texttt{aardvark}} \mapsto (1, 0, 0, 0, \dots, 0, 0, \\ \dots, 0, 0) \text{ and } \text{\texttt{aardwolf}} \mapsto (0, 1, 0, 0, \\ \dots, 0, 0) \text{ and } \text{\texttt{zyzzyva}} \mapsto (0, 0, 0, 0, \\ \dots, 0, 1) . \end{gathered}$	$\begin{aligned} \text{aardvark} &\mapsto (1, 0, 0, 0, \dots, 0, 0) \\ \text{aardwolf} &\mapsto (0, 1, 0, 0, \dots, 0, 0) \\ &\vdots \\ \text{zyzzyva} &\mapsto (0, 0, 0, 0, \dots, 0, 1). \end{aligned}$	$\begin{aligned} \text{aardvark} &\mapsto (1, 0, 0, 0, \dots, 0, 0) \\ \text{aardwolf} &\mapsto (0, 1, 0, 0, \dots, 0, 0) \\ &\vdots \\ \text{zyzzyva} &\mapsto (0, 0, 0, 0, \dots, 0, 1). \end{aligned}$
BradleyGastaldiTerilla2023EQ0003 (3)	4	1.000	$D \hookrightarrow \mathbb{R}^D \xrightarrow{\sigma} \mathbb{R}^{n_1},$	$D \hookrightarrow \mathbb{R}^D \xrightarrow{\sigma} \mathbb{R}^{n_1},$	$D \hookrightarrow \mathbb{R}^D \xrightarrow{\sigma} \mathbb{R}^{n_1},$
BradleyGastaldiTerilla2023EQ0005	5	1.000	$M^* u_j = \sigma_j v_j \text{ and } M v_j = \sigma_j u_j.$	$M^* u_j = \sigma_j v_j \text{ and } M v_j = \sigma_j u_j.$	$M^* u_j = \sigma_j v_j \text{ and } M v_j = \sigma_j u_j.$
BradleyGastaldiTerilla2023EQ0006	6	1.000	$F^* c_i \cong d_i \text{ and } F_* d_i \cong c_i.$	$F^* c_i \cong d_i \text{ and } F_* d_i \cong c_i.$	$F^* c_i \cong d_i \text{ and } F_* d_i \cong c_i.$
BradleyGastaldiTerilla2023EQ0007	6	1.000	$R^*(A_i) = B_i \text{ and } R_*(B_i) = A_i$	$R^*(A_i) = B_i \text{ and } R_*(B_i) = A_i$	$R^*(A_i) = B_i \text{ and } R_*(B_i) = A_i$