

cardona-qft-methods: residuals

equations measured 2026-09-05T11:34:14Z against the model of 2026-09-03

Unresolved

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E09654 (4)	261	■ 0.167 ●	$\begin{aligned} &\phi_{\theta}(x)\phi_{\theta}(y)=e^{\{-\frac{1}{2}\frac{\partial}{\partial x^{\mu}}\frac{\partial}{\partial y^{\nu}}\theta^{\mu\nu}\}}\phi_0(x)\phi_0(y)e^{\frac{1}{2}\left(\frac{\overleftarrow{\partial}}{\partial x^{\mu}}+\frac{\overrightarrow{\partial}}{\partial y^{\mu}}\right)\theta^{\mu\nu}P_{\nu}}\neq\phi_0(x)\phi_0(y). \end{aligned}$	(not rendered)	$\phi_{\theta}(x)\phi_{\theta}(y)=e^{-\frac{1}{2}\frac{\partial}{\partial x^{\mu}}\theta^{\mu\nu}\frac{\partial}{\partial y^{\nu}}}\phi_0(x)\phi_0(y)e^{\frac{1}{2}\left(\frac{\overleftarrow{\partial}}{\partial x^{\mu}}+\frac{\overrightarrow{\partial}}{\partial y^{\mu}}\right)\theta^{\mu\nu}P_{\nu}}\neq\phi_0(x)\phi_0(y).$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Flagged, not acted on

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0677	46	■ 0.150 ● C +44	\left.\begin{array}{cccccc} 1 & \rightarrow & \operatorname{Inn}(\mathcal{A}) & \rightarrow & \operatorname{Aut}(\mathcal{A}) & \rightarrow & \operatorname{Aut}(\mathcal{A})/\operatorname{Inn}(\mathcal{A}) & \rightarrow & 1, \\ 1 & \rightarrow & \mathcal{G} & \rightarrow & \mathcal{G} \rtimes \operatorname{Diff}(M) & \rightarrow & \operatorname{Diff}(M) & \rightarrow & 1. \end{array}\right\}	(not rendered)	$\begin{array}{lcl} AdS_5 : & -R^2 = -Z_0^2 + Z_1^2 + Z_2^2 + Z_3^2 + Z_4^2 - Z_5^2, \\ S^5 : & R^2 = Y_1^2 + Y_2^2 + Y_3^2 + Y_4^2 + Y_5^2 + Y_6^2. \end{array}$
cardona-qft-methods_ EQ0748	284	■ 0.547 ● C -27	\begin{aligned} A d S_{\{5\}} & : & \& -R^{\{2\}} & \& \\ =-Z_{\{0\}}^{\{2\}}+Z_{\{1\}}^{\{2\}}+Z_{\{2\}}^{\{2\}}+Z_{\{3\}}^{\{2\}}+Z_{\{4\}}^{\{2\}}-Z_{\{5\}}^{\{2\}} & S^{\{5\}} : & R^2 = Y_1^2 + Y_2^2 + Y_3^2 + Y_4^2 + Y_5^2 + Y_6^2 \\ \& S^{\{5\}} & : & \& R^{\{2\}} & =Y_{\{1\}}^{\{2\}}+Y_{\{2\}}^{\{2\}}+Y_{\{3\}}^{\{2\}}+Y_{\{4\}}^{\{2\}}+Y_{\{5\}}^{\{2\}}+Y_{\{6\}}^{\{2\}} \end{aligned}	$\begin{array}{lcl} AdS_5 : & -R^2 = -Z_0^2 + Z_1^2 + Z_2^2 + Z_3^2 + Z_4^2 - Z_5^2, \\ S^5 : & R^2 = Y_1^2 + Y_2^2 + Y_3^2 + Y_4^2 + Y_5^2 + Y_6^2. \end{array}$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

88 more flagged rows below the band, stated as a count: 18 component, 70 weak; confidence high 60, mid 14, low 14, none 0

Low confidence

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ9908 [conf 0.080]	350	■ 0.080 ● U +5	<pre>\begin{array}{rlc} P_{\mathrm{g}} M\!:\!{}_1\times{}_0 P_{\mathrm{h}} M & \longrightarrow & P_{\mathrm{g}} M\times P_{\mathrm{h}} M \\ & \downarrow & \downarrow \\ & M & \downarrow \\ & \downarrow & \downarrow \times M \end{array}</pre>	$\begin{array}{ccc} P_g M : {}_1 \times {}_0 P_h M & \longrightarrow & P_g M \times P_h M \\ \downarrow & & \downarrow \\ & e_0 \circ \pi_1 & \\ M & & \downarrow \\ \downarrow & & \downarrow \times M \end{array}$	$\begin{array}{ccc} P_g M : {}_1 \times {}_0 P_h M & \longrightarrow & P_g M \times P_h M \\ \downarrow e_0 \circ \pi_1 & & \downarrow e_0 \times e_0 \\ M & \xrightarrow{\Delta_g} & M \times M \end{array}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					