

kohlhase-omdoc: residuals

equations measured 2026-09-05T12:56:03Z against the model of 2026-09-03

Unresolved

Identifier	Page	Conf.	LaTeX source	Rendered	Image
kohlhase-omdoc_ E00005 [conf 0.092]	309	■ 0.092 ● C +105	<pre>\begin{figure} \[\begin{aligned} \left.\mathcal{R}\right. \\ \frac{\mathbf{A}:\mathbf{B} \rightarrow \mathbf{C} \quad \mathbf{B}:\mathbb{B}}{\mathbf{AB}:\mathbb{C}} \Bigg) &= \frac{\frac{\forall X.\mathbb{B}(X) \Rightarrow \mathbb{C}(\mathbf{AX})}{\mathbb{B}(\mathbf{B}) \Rightarrow \mathbb{C}(\mathbf{AB})} \quad \mathbb{B}(\mathbf{B})}{\mathbb{C}(\mathbf{AB})} \\ \\ \mathcal{R} \frac{\forall X_{\mathbb{B}}.\mathbf{A} \quad \mathbf{B}:\mathbb{B}}{[\mathbf{B}/X]\mathbf{A}} \Bigg) &= \frac{\frac{\forall X.\mathbb{B}(X) \Rightarrow \mathcal{R}(\mathbf{A})}{\mathbb{B}(\mathcal{R}(\mathbf{B})) \Rightarrow \mathcal{R}([\mathbf{B}/X]\mathbf{A})} \quad \mathbb{B}(\mathbf{B})}{\mathcal{R}([\mathbf{B}/X]\mathbf{A})}\end{aligned}</pre> \begin{figure} \[\begin{aligned} \left.\mathcal{R}\right. \\ \frac{\mathbf{A}:\mathbf{B} \rightarrow \mathbf{C} \quad \mathbf{B}:\mathbb{B}}{\mathbf{AB}:\mathbb{C}} \Bigg) &= \frac{\frac{\forall X.\mathbb{B}(X) \Rightarrow \mathbb{C}(\mathbf{AX})}{\mathbb{B}(\mathbf{B}) \Rightarrow \mathbb{C}(\mathbf{AB})} \quad \mathbb{B}(\mathbf{B})}{\mathbb{C}(\mathbf{AB})} \\ \\ \mathcal{R} \frac{\forall X_{\mathbb{B}}.\mathbf{A} \quad \mathbf{B}:\mathbb{B}}{[\mathbf{B}/X]\mathbf{A}} \Bigg) &= \frac{\frac{\forall X.\mathbb{B}(X) \Rightarrow \mathcal{R}(\mathbf{A})}{\mathbb{B}(\mathcal{R}(\mathbf{B})) \Rightarrow \mathcal{R}([\mathbf{B}/X]\mathbf{A})} \quad \mathbb{B}(\mathbf{B})}{\mathcal{R}([\mathbf{B}/X]\mathbf{A})}\end{aligned} \] \captionsetup{labelformat=empty} \caption{Fig. 26.26. Relativization Morphism on Proofs} \end{figure}	(not rendered)	
kohlhase-omdoc_ F00323	215	■ 0.405	$\{\,p\mid p\mid p+2\text{ are prime}\,\}$	(not rendered)	kin) within math formulas. Of course it may be best to display a formula like this instead of breaking it between lines
kohlhase-omdoc_ F00317	215	■ 0.891	$\{\,p\mid\hbox{p+2 are prime}\,\}$	(not rendered)	English description: for example consider 'p p and p+2 are pri how would do the job? /n
kohlhase-omdoc_ F00319	215	■ 0.891	are prime}\,,}	(not rendered)	English description: for example consider 'p p and p+2 are pri how would do the job? /n
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Flagged, not acted on

Identifier	Page	Conf.	LaTeX source	Rendered	Image
kohlhase-omdoc_ EQ0001	60	■ 0.863 ● C +38	\forall M . \operatorname{magma}(M) \Leftrightarrow \exists E, F . M=(E, F) \wedge \text { law_of_composition (E, F)}	(not rendered)	$\forall M.magma(M) \Leftrightarrow \exists E, F.M=(E, F) \wedge law_of_composition(E, F)$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value					
Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

1 more flagged rows below the band, stated as a count: 1 component, 0 weak; confidence high 0, mid 0, low 1, none 0

Low confidence

Identifier	Page	Conf.	LaTeX source	Rendered	Image
kohlhase-omdoc_ EQ0004 (26.1) [conf 0.035]	297	■ 0.035 ● U +44	<code>\backslash \text { CSum\{k\}1\infty\{\Cexp\{x\}k\}</code> instead of <code>\backslash \text { sum_\{k = 1\}^ }\{ }_{1}</code> <code>\text { infty x^k }</code>	<i>(not rendered)</i>	<code>\CSum{k}1\infty{\Cexp{x}k}</code> instead of <code>\sum_{k = 1}^{\infty} x^k</code>
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					