

voloshin-hypergraph: residuals

equations measured 2026-09-05T13:35:48Z against the model of 2026-09-01

Corrected (3)

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
voloshin-hypergraph_ EQ0133 (was)	170	■ 0.340	$(\mathcal{H})_2=(X,\mathcal{E})$ \text { where } \left\{ x_{\{i\}}, x_{\{j\}} \right\} \in \mathcal{E} \Leftrightarrow \mathcal{D}(x_i) \cap \mathcal{D}(x_j) \neq \emptyset .	$(\mathcal{H})_2=(X,\mathcal{E})$ where $\{x_i,x_j\} \in \mathcal{E} \Leftrightarrow \mathcal{D}(x_i) \cap \mathcal{D}(x_j) \neq \emptyset .$	—
voloshin-hypergraph_ EQ0133 (now)	170	—	$(\mathcal{H})_2=(X,\mathcal{E})$ \text { where } \left\{ x_{\{i\}}, x_{\{j\}} \right\} \in \mathcal{E} \Leftrightarrow \mathcal{D}(x_i) \cap \mathcal{D}(x_j) \neq \emptyset .	$(\mathcal{H})_2=(X,\mathcal{E})$ where $\{x_i,x_j\} \in \mathcal{E} \Leftrightarrow \mathcal{D}(x_i) \cap \mathcal{D}(x_j) \neq \emptyset .$	—
basis: inferred / ink					
voloshin-hypergraph_ EQ0141 (was)	185	■ 0.473	$\tau(\mathcal{H})=\nu(\mathcal{H})$.	$\tau(\mathcal{H})=\nu(\mathcal{H})$.	—
voloshin-hypergraph_ EQ0141 (now)	185	—	$\tau(\mathcal{H})=\nu(\mathcal{H})$. a hypergraph, draw	$\tau(\mathcal{H})=\nu(\mathcal{H}).ahypergraph, draw$	—
basis: inferred / ink					
voloshin-hypergraph_ EQ0197 (was)	220	■ 0.500	$\chi(\mathcal{H})=\bar{\chi}(\mathcal{H})=0$, \quad $R(\mathcal{H})=(0,0,\dots,0)$, \quad $P(\mathcal{H},\lambda)=0$.	$\chi(\mathcal{H})=\bar{\chi}(\mathcal{H})=0$, $R(\mathcal{H})=(0,0,\dots,0)$, $P(\mathcal{H},\lambda)=0$.	—
voloshin-hypergraph_ EQ0197 (now)	220	—	en the incidence matrix of a hypergraph, construct the edge list.	<i>entheincidencematrixofahypergraph,constructtheedgelist.</i>	—
basis: inferred / ink					
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					