

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_ E08517 [conf 0.002]	085	■ 0.002 ● C +43	$\begin{aligned} & \frac{1}{2}(-\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 0 \quad 0 \quad 1 \quad 0 \quad 1 \quad 0 \\ & \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 1 \quad 2 \quad 3 \quad 1 \quad 2 \quad 1 \\ & \frac{1}{2}(-\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 0 \quad 1 \quad 2 \quad 1 \quad 2 \quad 1 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 1 \quad 1 \quad 2 \quad 1 \quad 2 \quad 1 \\ & \frac{1}{2}(\lambda_0 + \lambda_1 - \lambda_2 + \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 1 \quad 2 \quad 2 \quad 1 \quad 2 \quad 1 \\ & \frac{1}{2}(-\lambda_0 - \lambda_1 - \lambda_2 + \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 0 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 - \lambda_2 - \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 1 \quad 1 \quad 1 \quad 0 \quad 1 \quad 0 \\ & \frac{1}{2}(-\lambda_0 + \lambda_1 - \lambda_2 - \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 0 \quad 1 \quad 1 \quad 0 \quad 1 \quad 0 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 1 \quad 1 \quad 2 \quad 1 \quad 1 \quad 0 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 - \lambda_2 + \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 0 \\ & \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3) + \frac{1}{2}(\mu_1 - \mu_2) = 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 1 \end{aligned}$	$\begin{aligned} & \frac{1}{2}(-\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 0 \quad 0 \quad 1 \quad 0 \quad 1 \\ & \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 1 \quad 2 \quad 3 \quad 1 \quad 1 \quad 2 \\ & \frac{1}{2}(-\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 0 \quad 1 \quad 2 \quad 1 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 1 \quad 1 \quad 2 \quad 2 \quad 1 \quad 2 \\ & \frac{1}{2}(\lambda_0 + \lambda_1 - \lambda_2 + \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 1 \quad 2 \quad 2 \quad 1 \\ & \frac{1}{2}(-\lambda_0 - \lambda_1 - \lambda_2 + \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 0 \quad 0, 0 \quad 0 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 - \lambda_2 - \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 1 \quad 1 \quad 1 \quad 1 \quad 0 \quad 1 \\ & \frac{1}{2}(-\lambda_0 + \lambda_1 - \lambda_2 - \lambda_3) + \frac{1}{2}(\mu_3 - \mu_1) = 0 \quad 1 \quad 1 \quad 0 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 1 \quad 1 \quad 2 \quad 1 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 - \lambda_2 + \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 1 \quad 1 \quad 1 \end{aligned}$ $\frac{1}{1} \quad 1 \quad 0$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_ EQ1026 [conf 0.018]	171	■0.018 ●C +25	$\left.\begin{array}{l} \lambda_0 - \lambda_1 = 2 \quad 3 \quad 4 \quad 3 \quad 2 \quad 1 \quad 0 \quad 2 \quad \lambda_0 + \lambda_1 = 2 \quad 4 \quad 6 \quad 5 \quad 4 \quad 3 \quad 2 \quad 3 \\ \lambda_0 - \lambda_2 = 2 \quad 3 \quad 4 \quad 3 \quad 2 \quad 1 \quad 1 \quad 2 \quad \lambda_0 + \lambda_2 = 2 \quad 4 \quad 6 \quad 5 \quad 4 \quad 3 \quad 1 \quad 3 \\ \lambda_0 - \lambda_3 = 2 \quad 3 \quad 4 \quad 3 \quad 2 \quad 2 \quad 1 \quad 2 \quad \lambda_0 + \lambda_3 = 2 \quad 4 \quad 6 \quad 5 \quad 4 \quad 2 \quad 1 \quad 3 \\ \lambda_1 - \lambda_2 = 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 0 \quad \lambda_1 + \lambda_2 = 0 \quad 1 \quad 2 \quad 2 \quad 2 \quad 2 \quad 1 \quad 1 \\ \lambda_1 - \lambda_3 = 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 1 \quad 0 \quad \lambda_1 + \lambda_3 = 0 \quad 1 \quad 2 \quad 2 \quad 2 \quad 1 \quad 1 \quad 1 \\ \lambda_2 - \lambda_3 = 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 0 \quad 0 \quad \lambda_2 + \lambda_3 = 0 \quad 1 \quad 2 \quad 2 \quad 2 \quad 1 \quad 0 \quad 1 \end{array}\right]$	$\left.\begin{array}{l} \lambda_0 - \lambda_1 = 2 \quad 3 \quad 4 \quad 3 \quad 2 \quad 1 \quad 0 \quad 2 \quad \lambda_0 + \lambda_1 = 2 \quad 4 \quad 6 \quad 5 \quad 4 \quad 3 \quad 2 \quad 3 \\ \lambda_0 - \lambda_2 = 2 \quad 3 \quad 4 \quad 3 \quad 2 \quad 1 \quad 1 \quad 2 \quad \lambda_0 + \lambda_2 = 2 \quad 4 \quad 6 \quad 5 \quad 4 \quad 3 \quad 1 \quad 3 \\ \lambda_1 - \lambda_2 = 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 0 \quad 4 \quad \lambda_1 + \lambda_2 = 0 \quad 1 \quad 2 \quad 2 \quad 2 \quad 2 \quad 1 \quad 1 \\ \lambda_1 - \lambda_3 = 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 1 \quad 0 \quad \lambda_1 + \lambda_3 = 0 \quad 1 \quad 2 \quad 2 \quad 2 \quad 1 \quad 1 \quad 1 \\ \lambda_2 - \lambda_3 = 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 0 \quad 0 \quad \lambda_2 + \lambda_3 = 0 \quad 1 \quad 2 \quad 2 \quad 2 \quad 1 \quad 0 \quad 1 \end{array}\right]$	$\begin{array}{l} \lambda_0 - \lambda_1 = 2 \quad 3 \quad 4 \quad 3 \quad 2 \quad 1 \quad 0 \quad 2 \quad \lambda_0 + \lambda_1 = 2 \quad 4 \quad 6 \quad 5 \quad 4 \quad 3 \quad 2 \quad 3 \\ \lambda_0 - \lambda_2 = 2 \quad 3 \quad 4 \quad 3 \quad 2 \quad 1 \quad 1 \quad 2 \quad \lambda_0 + \lambda_2 = 2 \quad 4 \quad 6 \quad 5 \quad 4 \quad 3 \quad 1 \quad 3 \\ \lambda_1 - \lambda_2 = 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 0 \quad 4 \quad \lambda_1 + \lambda_2 = 0 \quad 1 \quad 2 \quad 2 \quad 2 \quad 2 \quad 1 \quad 1 \\ \lambda_1 - \lambda_3 = 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 1 \quad 0 \quad \lambda_1 + \lambda_3 = 0 \quad 1 \quad 2 \quad 2 \quad 2 \quad 1 \quad 1 \quad 1 \\ \lambda_2 - \lambda_3 = 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 1 \quad 0 \quad 0 \quad \lambda_2 + \lambda_3 = 0 \quad 1 \quad 2 \quad 2 \quad 2 \quad 1 \quad 0 \quad 1 \end{array}$
0902.0431_ EQ0518 [conf 0.021]	086	■0.021 ●C +6	$\begin{array}{l} \frac{1}{2}(\lambda_0 + \lambda_1 - \lambda_2 - \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 1 \quad 2 \quad 2 \quad 2 \quad 1 \quad 1 \quad c \\ \frac{1}{2}(-\lambda_0 + \lambda_1 - \lambda_2 + \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 0 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \\ \frac{1}{2}(-\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 0 \quad 1 \quad 2 \quad 1 \quad 1 \quad 1 \\ \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 1 \quad 2 \quad 3 \quad 2 \\ \frac{1}{2}(-\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 0 \quad 0 \end{array}$	$\begin{array}{l} \frac{1}{2}(\lambda_0 + \lambda_1 - \lambda_2 - \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 1 \quad 2 \quad 2 \quad 2 \quad 1 \quad 1 \quad c \\ \frac{1}{2}(-\lambda_0 + \lambda_1 - \lambda_2 + \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 0 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \\ \frac{1}{2}(-\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 0 \quad 1 \quad 2 \quad 1 \quad 1 \quad 1 \\ \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 1 \quad 2 \quad 3 \quad 2 \\ \frac{1}{2}(-\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 0 \quad 0 \end{array}$	$\begin{array}{l} \frac{1}{2}(\lambda_0 + \lambda_1 - \lambda_2 - \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 1 \quad 2 \quad 2 \quad 1 \quad 1 \quad 0 \\ \frac{1}{2}(-\lambda_0 + \lambda_1 - \lambda_2 + \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 0 \quad 1 \quad 1 \quad 1 \quad 1 \quad 0 \\ \frac{1}{2}(-\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 0 \quad 1 \quad 2 \quad 1 \quad 1 \quad 0 \\ \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 1 \quad 2 \quad 3 \quad 2 \quad 2 \quad 1 \\ \frac{1}{2}(-\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3) - \frac{1}{2}(\mu_1 - \mu_2) = 0 \quad 0 \quad 1 \quad 1 \quad 1 \quad 0. \end{array}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_ EQ1632 [conf 0.022]	173	■ 0.022 ● C +579	$\begin{aligned} & \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 + \lambda_1 - \lambda_2 + \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 - \lambda_2 - \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3) - \frac{1}{2}\mu_2 + \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 + \lambda_1 - \lambda_2 + \lambda_3) - \frac{1}{2}\mu_2 + \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3) - \frac{1}{2}\mu_2 - \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 - \lambda_2 - \lambda_3) - \frac{1}{2}\mu_2 + \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3) + \frac{1}{2}\mu_3 - \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 + \lambda_1 - \lambda_2 - \lambda_3) + \frac{1}{2}\mu_3 - \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3) + \frac{1}{2}\mu_3 - \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 - \lambda_2 + \lambda_3) + \frac{1}{2}\mu_3 - \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3) - \frac{1}{2}\mu_3 + \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 + \lambda_1 - \lambda_2 - \lambda_3) - \frac{1}{2}\mu_3 + \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3) - \frac{1}{2}\mu_3 + \frac{2}{3}\nu = 1 \\ & \frac{1}{2}(\lambda_0 - \lambda_1 - \lambda_2 + \lambda_3) - \frac{1}{2}\mu_3 + \frac{2}{3}\nu = 1 \end{aligned}$	(not rendered)	$\begin{aligned} & 2 \quad 3 \quad 3 \quad 2 \quad 2 \quad 1 \quad 1 \\ & 2 \quad 3 \quad 3 \quad 2 \quad 1 \quad 1 \quad 1 \\ & 2 \quad 3 \quad 3 \quad 2 \quad 1 \quad 0 \quad 1 \\ & 1 \quad 1 \quad 1 \quad 0 \quad 0 \quad 0 \quad 0 \\ & 2 \quad 2 \quad 2 \quad 2 \quad 2 \quad 1 \quad 2 \\ & 2 \quad 2 \quad 1 \quad 1 \quad 2 \\ & 2 \quad 2 \quad 1 \quad 0 \quad 2 \\ & 1 \quad 1 \quad 0 \quad 0 \quad 0 \quad 0 \quad 1 \\ & 2 \quad 3 \quad 3 \quad 3 \quad 2 \quad 1 \quad 1 \\ & 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 0 \\ & 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 0 \quad 0 \\ & 1 \quad 1 \quad 1 \quad 1 \quad 0 \quad 0 \quad 0 \\ & 3 \quad 5 \quad 4 \quad 3 \quad 2 \quad 1 \quad 3 \\ & 2 \quad 3 \quad 2 \quad 1 \quad 1 \quad 1 \quad 2 \\ & 2 \quad 3 \quad 2 \quad 1 \quad 1 \quad 0 \quad 2 \\ & 2 \quad 3 \quad 2 \quad 1 \quad 0 \quad 0 \quad 2 \end{aligned}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_ E01031 [conf 0.036]	173	■ 0.036 ● C +5	$\begin{aligned} & \backslash \text{frac}\{2\}{3} \backslash \text{nu} = 0 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 0 \quad \backslash \text{quad} 0 \quad \backslash \backslash \quad \& \quad \backslash \text{lambda}_{2} - \backslash \text{frac}\{1\}{2} \quad \backslash \text{mu}_{1} + \backslash \text{frac}\{2\}{3} \backslash \text{nu} = 0 \quad \backslash \text{quad} 0 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 0 \quad \backslash \backslash \quad \& \quad \backslash \text{lambda}_{3} + \backslash \text{frac}\{1\}{2} \quad \backslash \text{mu}_{1} - \backslash \text{frac}\{2\}{3} \backslash \text{nu} = 0 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \backslash \quad \& \quad \backslash \text{lambda}_{3} - \backslash \text{frac}\{1\}{2} \quad \backslash \text{mu}_{1} + \backslash \text{frac}\{2\}{3} \backslash \text{nu} = 0 \quad \backslash \text{quad} 0 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 0 \quad \backslash \backslash \quad \& \quad 0 \quad \end{aligned}$	$\begin{aligned} \lambda_2 + \frac{1}{2}\mu_1 - \frac{2}{3}\nu &= 0 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ \lambda_2 - \frac{1}{2}\mu_1 + \frac{2}{3}\nu &= 0 & 0 & 1 & 1 & 1 & 1 & 0 & \\ \lambda_3 + \frac{1}{2}\mu_1 - \frac{2}{3}\nu &= 0 & 1 & 1 & 1 & 1 & & & \\ \lambda_3 - \frac{1}{2}\mu_1 + \frac{2}{3}\nu &= 0 & 0 & 1 & 1 & 1 & 0 & & \\ 0 & & & & & & & & \end{aligned}$	$\begin{aligned} \lambda_2 + \frac{1}{2}\mu_1 - \frac{2}{3}\nu &= 0 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ \lambda_2 - \frac{1}{2}\mu_1 + \frac{2}{3}\nu &= 0 & 0 & 1 & 1 & 1 & 1 & 0 & 1 \\ \lambda_3 + \frac{1}{2}\mu_1 - \frac{2}{3}\nu &= 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ \lambda_3 - \frac{1}{2}\mu_1 + \frac{2}{3}\nu &= 0 & 0 & 1 & 1 & 1 & 0 & 0 & 1 \end{aligned}$
0902.0431_ E00761 [conf 0.036]	127	■ 0.036 ● C +497	$\begin{aligned} & \backslash \text{frac}\{1\}{2} \backslash \text{left}(-\backslash \text{lambda}_{0} - \backslash \text{lambda}_{1} + \backslash \text{lambda}_{2} - \backslash \text{lambda}_{3}) + \backslash \text{frac}\{1\}{2} \backslash \text{mu}_{2} - \backslash \text{frac}\{2\}{3} \backslash \text{nu} = 0 \quad \backslash \text{quad} 0 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} [\textbf{U+0930}] [\textbf{U+094D}] 0 \quad \backslash \text{quad} 0 \quad \backslash \backslash \quad \& \quad \backslash \text{frac}\{1\}{2} \backslash \text{left}(\backslash \text{lambda}_{0} + \backslash \text{lambda}_{1} + \backslash \text{lambda}_{2} + \backslash \text{lambda}_{3}) - \backslash \text{frac}\{1\}{2} \backslash \text{mu}_{2} - \backslash \text{frac}\{2\}{3} \backslash \text{nu} = 0 \quad \backslash \text{quad} 0 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 0 \quad \backslash \text{quad} 0 \quad \backslash \backslash \quad \& \quad \backslash \text{frac}\{1\}{2} \backslash \text{left}(\backslash \text{lambda}_{0} + \backslash \text{lambda}_{1} + \backslash \text{lambda}_{2} + \backslash \text{lambda}_{3}) + \backslash \text{frac}\{1\}{2} \backslash \text{mu}_{2} - \backslash \text{frac}\{2\}{3} \backslash \text{nu} = 1 \quad \backslash \text{quad} 2 \quad \backslash \text{quad} 3 \quad \backslash \text{quad} 3 \quad \backslash \text{quad} 2 \quad \backslash \text{end}\{aligned\} \quad \backslash \text{quad} 1 \quad \backslash \text{quad} 2 \quad . \end{aligned}$	$\begin{aligned} \frac{1}{2}(-\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu &= 0 & 0 & 1 & 1 & [\textbf{U+0930}][\textbf{U+094D}] & 0 & 0 & \\ \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu &= 1 & 2 & 3 & 3 & 2 & 1 & 2 & \end{aligned}$	$\begin{aligned} \frac{1}{2}(-\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu &= 0 & 0 & 1 & 1 & 0 & 0 & 1 & \\ \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu &= 1 & 2 & 3 & 3 & 2 & 1 & 2 & \\ \frac{1}{2}(-\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu &= 0 & 1 & 2 & 3 & 2 & 1 & 2 & \\ \frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu &= 1 & 1 & 2 & 3 & 2 & 1 & 2 & \\ \frac{1}{2}(\lambda_0 + \lambda_1 - \lambda_2 + \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu &= 1 & 2 & 2 & 3 & 2 & 1 & 2 & \\ \frac{1}{2}(-\lambda_0 - \lambda_1 - \lambda_2 + \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu &= 0 & 0 & 0 & 1 & 0 & 0 & 1 & \\ \frac{1}{2}(\lambda_0 - \lambda_1 - \lambda_2 - \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu &= 1 & 1 & 1 & 1 & 0 & 0 & 1 & \\ \frac{1}{2}(-\lambda_0 + \lambda_1 - \lambda_2 - \lambda_3) + \frac{1}{2}\mu_2 - \frac{2}{3}\nu &= 0 & 1 & 1 & 1 & 0 & 0 & 1 & \\ \frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3) + \frac{1}{2}\mu_3 - \frac{2}{3}\nu &= 1 & 1 & 2 & 2 & 1 & 0 & 1 & \\ \frac{1}{2}(\lambda_0 - \lambda_1 - \lambda_2 + \lambda_3) + \frac{1}{2}\mu_3 + \frac{2}{3}\nu &= 1 & 1 & 1 & 2 & 1 & 0 & 1 & \\ \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3) - \frac{1}{2}\mu_3 - \frac{2}{3}\nu &= 0 & 0 & 0 & 0 & 1 & 1 & 0 & \\ \frac{1}{2}(\lambda_0 + \lambda_1 - \lambda_2 - \lambda_3) + \frac{1}{2}\mu_3 - \frac{2}{3}\nu &= 1 & 2 & 2 & 2 & 1 & 0 & 1 & \\ \frac{1}{2}(-\lambda_0 + \lambda_1 - \lambda_2 + \lambda_3) + \frac{1}{2}\mu_3 - \frac{2}{3}\nu &= 0 & 1 & 1 & 2 & 1 & 0 & 1 & \\ \frac{1}{2}(-\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3) + \frac{1}{2}\mu_3 - \frac{2}{3}\nu &= 0 & 1 & 2 & 2 & 1 & 0 & 1 & \\ \frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3) + \frac{1}{2}\mu_3 - \frac{2}{3}\nu &= 1 & 2 & 3 & 4 & 3 & 1 & 2 & \\ \frac{1}{2}(-\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3) + \frac{1}{2}\mu_3 - \frac{2}{3}\nu &= 0 & 0 & 1 & 2 & 1 & 0 & 1. & \end{aligned}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_ E00516 [conf 0.041]	085	■ 0.041 ● C +12	$\begin{aligned} & \& \lambda_0 + \frac{1}{2} \left(\mu_2 - \mu_3 \right) = 1 \quad 1 \quad 1 \quad 1 \quad a \quad 0 \\ & \& \lambda_1 + \frac{1}{2} \left(\mu_2 - \mu_3 \right) = 0 \quad 1 \quad 1 \quad 1 \quad a \\ & \& \lambda_2 + \frac{1}{2} \left(\mu_2 - \mu_3 \right) = 0 \quad 0 \quad 1 \quad 1 \\ & \& \lambda_3 + \frac{1}{2} \left(\mu_2 - \mu_3 \right) = 0 \quad 0 \quad 0 \\ & \& \lambda_0 - \frac{1}{2} \left(\mu_2 - \mu_3 \right) = 1 \quad 1 \quad 1 \quad 1 \quad 0 \\ & \& \lambda_1 - \frac{1}{2} \left(\mu_2 - \mu_3 \right) = 0 \quad 1 \quad 1 \quad 1 \quad 0 \\ & \& \lambda_2 - \frac{1}{2} \left(\mu_2 - \mu_3 \right) = 0 \quad 0 \quad 1 \quad 1 \\ & \& \lambda_3 - \frac{1}{2} \left(\mu_2 - \mu_3 \right) = 0 \quad 0 \quad 0 \quad 0 \end{aligned}$	$\begin{aligned} & \lambda_0 + \frac{1}{2} (\mu_2 - \mu_3) = 1 \quad 1 \quad 1 \quad 1 \quad a \quad 0 \\ & \lambda_1 + \frac{1}{2} (\mu_2 - \mu_3) = 0 \quad 1 \quad 1 \quad 1 \quad a \\ & \lambda_2 + \frac{1}{2} (\mu_2 - \mu_3) = 0 \quad 0 \quad 1 \quad 1 \\ & \lambda_3 + \frac{1}{2} (\mu_2 - \mu_3) = 0 \quad 0 \quad 0 \\ & \lambda_0 - \frac{1}{2} (\mu_2 - \mu_3) = 1 \quad 1 \quad 1 \quad 1 \quad 0 \\ & \lambda_1 - \frac{1}{2} (\mu_2 - \mu_3) = 0 \quad 1 \quad 1 \quad 1 \quad 0 \\ & \lambda_2 - \frac{1}{2} (\mu_2 - \mu_3) = 0 \quad 0 \quad 1 \quad 1 \\ & \lambda_3 - \frac{1}{2} (\mu_2 - \mu_3) = 0 \quad 0 \quad 0 \quad 0 \end{aligned}$	
0902.0431_ E00425 [conf 0.082]	070	■ 0.082 ● W -2	$\begin{aligned} & \& A \times A^* \times A \times A^* = A(X \times Y) A^* \\ & \& (PMA^*)^*(PNA^*) = AM^*P^*PNA = A(M^*N)A, \\ & \& (PMA^*)(AYA^*) = P(MY)A^*, \\ & \& \overline{(PMA^* \times PNA^*)} = \overline{{}^t\tilde{P}(M \times N){}^t\tilde{A}^*} = \overline{PM} \times \overline{NA^*}, \text{ etc.} \end{aligned}$	$\begin{aligned} & AXA^* \times AYA^* = A(X \times Y)A^* \\ & (PMA^*)^*(PNA^*) = AM^*P^*PNA = A(M^*N)A, \\ & (PMA^*)(AYA^*) = P(MY)A^*, \\ & \overline{PMA^* \times PNA^*} = \overline{{}^t\tilde{P}(M \times N){}^t\tilde{A}^*} = \overline{PM} \times \overline{NA^*}, \text{ etc.} \end{aligned}$	$\begin{aligned} & AXA^* \times AYA^* = A(X \times Y)A^*, \\ & (PMA^*)^*(PNA^*) = AM^*P^*PNA = A(M^*N)A, \\ & (PMA^*)(AYA^*) = P(MY)A^*, \\ & \overline{PMA^* \times PNA^*} = \overline{{}^t\tilde{P}(M \times N){}^t\tilde{A}^*} = \overline{PM} \times \overline{NA^*}, \text{ etc.} \end{aligned}$
0902.0431_ E01059 [conf 0.093]	180	■ 0.093 ● C +7	$\begin{aligned} & \& \alpha \in (E_8^C)_{1,1-,1-} \mid \tau \tilde{\lambda} \alpha = \alpha \tau \tilde{\lambda} \} \\ & \& \alpha \in E_7^C \mid \tau \tilde{\lambda} \alpha = \alpha \tau \tilde{\lambda} \} \text{ (Proposition 5.7.1)} \\ & \& \alpha \in E_7^C \mid \tau \lambda \alpha = \alpha \tau \lambda \} \text{ (by the correspondence to Proposition 5.7.1)} \\ & \& = E_7 \text{ (Lemma 4.3.3.(4)).} \end{aligned}$	$\begin{aligned} & \alpha \in (E_8^C)_{1,1-,1-} \mid \tau \tilde{\lambda} \alpha = \alpha \tau \tilde{\lambda} \} \\ & \alpha \in E_7^C \mid \tau \tilde{\lambda} \alpha = \alpha \tau \tilde{\lambda} \} \text{ (Proposition 5.7.1)} \\ & \alpha \in E_7^C \mid \tau \lambda \alpha = \alpha \tau \lambda \} \text{ (by the correspondence to Proposition 5.7.1)} \\ & = E_7 \text{ (Lemma 4.3.3.(4)).} \end{aligned}$	$\begin{aligned} & = \{ \alpha \in (E_8^C)_{1,1-,1-} \mid \tau \tilde{\lambda} \alpha = \alpha \tau \tilde{\lambda} \} \\ & = \{ \alpha \in E_7^C \mid \tau \tilde{\lambda} \alpha = \alpha \tau \tilde{\lambda} \} \text{ (Proposition 5.7.1)} \\ & = \{ \alpha \in E_7^C \mid \tau \lambda \alpha = \alpha \tau \lambda \} \text{ (by the correspondence to Proposition 5.7.1)} \\ & = E_7 \text{ (Lemma 4.3.3.(4)).} \end{aligned}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_ E00757 [conf 0.095]	126	■0.095 ●C +4	$\begin{aligned} & \backslash \text{begin}\{\text{aligned}\} \ \& \ \backslash \text{lambda}_{\{0\}}-\backslash \text{frac}\{1\}{2}\backslash \text{left}(\backslash \mu_{\{2\}}-\backslash \mu_{\{3\}}\backslash \text{right})=1 \ \backslash \text{quad} \ 1 \ \backslash \text{quad} \ 1 \ \backslash \text{quad} \ 1 \ \backslash \text{quad} \ 1 \ \backslash \text{quad} \ 1 \ \backslash \text{quad} \ 0 \ \backslash \text{quad} \ 0 \ \backslash \backslash \ \& \ \backslash \text{lambda}_{\{1\}}-\backslash \text{frac}\{1\}{2}\backslash \text{left}(\backslash \mu_{\{2\}}-\backslash \mu_{\{3\}}\backslash \text{right})=0 \ \backslash \text{quad} \ 1 \ \backslash \text{quad} \ 1 \ \backslash \text{quad} \ 1 \ \backslash \text{quad} \ 1 \ \backslash \text{quad} \ 1 \ \backslash \text{quad} \ a \ \backslash \backslash \ \& \ \backslash \text{lambda}_{\{2\}}-\backslash \text{frac}\{1\}{2}\backslash \text{left}(\backslash \mu_{\{2\}}-\backslash \mu_{\{3\}}\backslash \text{right})=0 \ \backslash \text{quad} \ 0 \ \backslash \text{quad} \ 1 \ \backslash \text{quad} \ 1 \ \backslash \text{quad} \ 1 \ \backslash \backslash \ \& \ \backslash \text{lambda}_{\{3\}}-\backslash \text{frac}\{1\}{2}\backslash \text{left}(\backslash \mu_{\{2\}}-\backslash \mu_{\{3\}}\backslash \text{right})=0 \ \backslash \text{quad} \ 0 \ \backslash \text{quad} \ 0 \ \backslash \backslash \ \& \ 0 \ \text{end}\{\text{aligned}\} \end{aligned}$	$\begin{aligned} \lambda_0 - \frac{1}{2}(\mu_2 - \mu_3) &= 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 0 \quad 0 \\ \lambda_1 - \frac{1}{2}(\mu_2 - \mu_3) &= 0 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad a \\ \lambda_2 - \frac{1}{2}(\mu_2 - \mu_3) &= 0 \quad 0 \quad 1 \quad 1 \quad 1 \\ \lambda_3 - \frac{1}{2}(\mu_2 - \mu_3) &= 0 \quad 0 \quad 0 \\ 0 \end{aligned}$	$\begin{aligned} \lambda_0 - \frac{1}{2}(\mu_2 - \mu_3) &= 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 0 \quad 0 \\ \lambda_1 - \frac{1}{2}(\mu_2 - \mu_3) &= 0 \quad 1 \quad 1 \quad 1 \quad 1 \quad 0 \quad 0 \\ \lambda_2 - \frac{1}{2}(\mu_2 - \mu_3) &= 0 \quad 0 \quad 1 \quad 1 \quad 1 \quad 0 \quad 0 \\ \lambda_3 - \frac{1}{2}(\mu_2 - \mu_3) &= 0 \quad 0 \quad 0 \quad 1 \quad 1 \quad 0 \quad 0 \end{aligned}$
0902.0431_E00922	153	■0.102 ●C +237	$\begin{aligned} & f^{\{-1\}}\backslash \text{left}(\backslash \text{sum}_{\{i<j<k\}} \ x_{\{i \ j \ k\}} \ \boldsymbol{e}_{\{i\}} \\ & \ \wedge \ \boldsymbol{e}_{\{j\}} \ \wedge \ \boldsymbol{e}_{\{k\}} \\ & \ \text{right})=\backslash \text{left}(\backslash \text{begin}\{\text{array}\}\{\text{c}\} \ \backslash \text{left}(\backslash \text{begin}\{\text{array}\}\{\text{ccc}\} \\ & \ h\backslash \text{left}(x_{\{156\}}\backslash \text{right}) \ \& \ h\backslash \text{left}(x_{\{164\}}, \ \bar{x}_{\{256\}} \\ & \ \backslash \text{right}) \ \& \ h\backslash \text{left}(x_{\{145\}}, \ \bar{x}_{\{356\}}\backslash \text{right}) \ \backslash \backslash \ h \\ & \ \backslash \text{left}(x_{\{256\}}, \ \bar{x}_{\{164\}}\backslash \text{right}) \ \& \ h\backslash \text{left}(x_{\{264\}} \\ & \ \backslash \text{right}) \ \& \ h\backslash \text{left}(x_{\{245\}}, \ \bar{x}_{\{364\}}\backslash \text{right}) \ \backslash \backslash \ h \\ & \ \backslash \text{left}(x_{\{356\}}, \ \bar{x}_{\{145\}}\backslash \text{right}) \ \& \ h\backslash \text{left}(x_{\{364\}}, \\ & \ \bar{x}_{\{245\}}\backslash \text{right}) \ \& \ h\backslash \text{left}(x_{\{345\}}\backslash \text{right}) \ \text{end}\{\text{array}\} \\ & \backslash \text{right}) \ \backslash \backslash \ \backslash \text{left}(\backslash \text{begin}\{\text{array}\}\{\text{ccc}\} \ h\backslash \text{left}(x_{\{423\}}\backslash \text{right}) \\ & \ \& \ h\backslash \text{left}(x_{\{431\}}, \ \bar{x}_{\{523\}}\backslash \text{right}) \ \& \ h\backslash \text{left}(x_{\{412\}}, \\ & \ \bar{x}_{\{623\}}\backslash \text{right}) \ \backslash \backslash \ h\backslash \text{left}(x_{\{523\}}, \ \bar{x}_{\{431\}} \\ & \ \backslash \text{right}) \ \& \ h\backslash \text{left}(x_{\{531\}}\backslash \text{right}) \ \& \ h\backslash \text{left}(x_{\{512\}}, \\ & \ \bar{x}_{\{631\}}\backslash \text{right}) \ \backslash \backslash \ h\backslash \text{left}(x_{\{623\}}, \ \bar{x}_{\{412\}} \\ & \ \backslash \text{right}) \ \& \ h\backslash \text{left}(x_{\{631\}}, \ \bar{x}_{\{512\}}\backslash \text{right}) \ \& \ h \\ & \ \backslash \text{left}(x_{\{612\}}\backslash \text{right}) \ \text{end}\{\text{array}\}\backslash \text{right}) \ \backslash \backslash \backslash \backslash \backslash \backslash \\ & \ \backslash \text{left}(\backslash \text{begin}\{\text{array}\}\{\text{cc}\} \ h\backslash \text{left}(x_{\{123\}}\backslash \text{right}) \ \& \ \backslash \backslash \ h \\ & \ \backslash \text{left}(x_{\{456\}}\backslash \text{right}) \ \text{end}\{\text{array}\}\backslash \text{right}. \ \text{end}\{\text{array}\} \\ & \backslash \text{right}), \end{aligned}$	(not rendered)	$f^{-1}\Big(\sum_{i<j<k}x_{ijk}\boldsymbol{e}_i\wedge\boldsymbol{e}_j\wedge\boldsymbol{e}_k\Big)=\left(\begin{array}{ccc}h(x_{156}) & h(x_{164},\overline{x}_{256}) & h(x_{145},\overline{x}_{356}) \\ h(x_{256},\overline{x}_{164}) & h(x_{264}) & h(x_{245},\overline{x}_{364}) \\ h(x_{356},\overline{x}_{145}) & h(x_{364},\overline{x}_{245}) & h(x_{345}) \end{array}\right)\left(\begin{array}{ccc}h(x_{423}) & h(x_{431},\overline{x}_{523}) & h(x_{412},\overline{x}_{623}) \\ h(x_{523},\overline{x}_{431}) & h(x_{531}) & h(x_{512},\overline{x}_{631}) \\ h(x_{623},\overline{x}_{412}) & h(x_{631},\overline{x}_{512}) & h(x_{612}) \end{array}\right),$ $\begin{array}{c} h(x_{123}) \\ h(x_{456}) \end{array}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 —no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_E01117	189	■ 0.108 ● C +18	$\begin{aligned} & (\boldsymbol{X}, \boldsymbol{Y}) = (X_1, Y_1) + (X_2, Y_2) + (X_3, Y_3) \in C, \\ & \langle \boldsymbol{X}, \boldsymbol{Y} \rangle = \langle X_1, Y_1 \rangle + \langle X_2, Y_2 \rangle + \langle X_3, Y_3 \rangle \in C, \\ & \boldsymbol{X} \times \boldsymbol{Y} = \begin{pmatrix} X_2 \times Y_3 - Y_2 \times X_3 \\ X_3 \times Y_1 - Y_3 \times X_1 \\ X_1 \times Y_2 - Y_1 \times X_2 \end{pmatrix} \in (\mathfrak{J}^C)^3, \\ & \boldsymbol{X} \cdot \boldsymbol{Y} = \begin{pmatrix} (X_1, Y_1) & (X_1, Y_2) \\ (X_2, Y_1) & (X_2, Y_2) \\ (X_3, Y_1) & (X_2, Y_3) \\ X \vee \boldsymbol{Y} & (X_3, Y_3) \end{pmatrix} - \frac{1}{3}(\boldsymbol{X}, \boldsymbol{Y})E \in \mathfrak{sl}(3, C), \\ & \boldsymbol{X} \vee \boldsymbol{Y} = X_1 \vee Y_1 + X_2 \vee Y_2 + X_3 \vee Y_3 \in \mathfrak{e}_6^C, \end{aligned}$	$\begin{aligned} & (\boldsymbol{X}, \boldsymbol{Y}) = (X_1, Y_1) + (X_2, Y_2) + (X_3, Y_3) \in C, \\ & \langle \boldsymbol{X}, \boldsymbol{Y} \rangle = \langle X_1, Y_1 \rangle + \langle X_2, Y_2 \rangle + \langle X_3, Y_3 \rangle \in C, \\ & \boldsymbol{X} \times \boldsymbol{Y} = \begin{pmatrix} X_2 \times Y_3 - Y_2 \times X_3 \\ X_3 \times Y_1 - Y_3 \times X_1 \\ X_1 \times Y_2 - Y_1 \times X_2 \end{pmatrix} \in (\mathfrak{J}^C)^3, \\ & \boldsymbol{X} \cdot \boldsymbol{Y} = \begin{pmatrix} (X_1, Y_1) & (X_1, Y_2) \\ (X_2, Y_1) & (X_2, Y_2) \\ (X_3, Y_1) & (X_2, Y_3) \\ X \vee \boldsymbol{Y} & (X_3, Y_3) \end{pmatrix} - \frac{1}{3}(\boldsymbol{X}, \boldsymbol{Y})E \in \mathfrak{sl}(3, C), \\ & \boldsymbol{X} Y_1 + X_2 \vee Y_2 + X_3 \vee Y_3 \in \mathfrak{e}_6^C, \end{aligned}$	$\begin{aligned} & (\boldsymbol{X}, \boldsymbol{Y}) = (X_1, Y_1) + (X_2, Y_2) + (X_3, Y_3) \in C, \\ & \langle \boldsymbol{X}, \boldsymbol{Y} \rangle = \langle X_1, Y_1 \rangle + \langle X_2, Y_2 \rangle + \langle X_3, Y_3 \rangle \in C, \\ & \boldsymbol{X} \times \boldsymbol{Y} = \begin{pmatrix} X_2 \times Y_3 - Y_2 \times X_3 \\ X_3 \times Y_1 - Y_3 \times X_1 \\ X_1 \times Y_2 - Y_1 \times X_2 \end{pmatrix} \in (\mathfrak{J}^C)^3, \\ & \boldsymbol{X} \cdot \boldsymbol{Y} = \begin{pmatrix} (X_1, Y_1) & (X_1, Y_2) & (X_1, Y_3) \\ (X_2, Y_1) & (X_2, Y_2) & (X_2, Y_3) \\ (X_3, Y_1) & (X_3, Y_2) & (X_3, Y_3) \end{pmatrix} - \frac{1}{3}(\boldsymbol{X}, \boldsymbol{Y})E \in \mathfrak{sl}(3, C), \\ & \boldsymbol{X} \vee \boldsymbol{Y} = X_1 \vee Y_1 + X_2 \vee Y_2 + X_3 \vee Y_3 \in \mathfrak{e}_6^C, \end{aligned}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
9902.0431_E00756	125	■ 0.119 ● C +10	$\begin{array}{l} \backslash begin{array}{llllllllll} \lambda_0 - \lambda_1 = 1 \\ \& 0 \& 0 \& 0 \& 0 \& 0 \& 0 \& 0 \& \lambda_0 + \lambda_1 = 1 \\ 2 \& 2 \& 2 \& 2 \& 2 \& 1 \backslash \backslash \lambda_0 - \lambda_2 = 1 \& 1 \& \\ 0 \& 0 \& 0 \& 0 \& 0 \& 0 \& \lambda_0 + \lambda_2 = 1 \& 1 \& 2 \\ \& 2 \& 2 \& 2 \& 1 \backslash \backslash \lambda_0 - \lambda_3 = 1 \& 1 \& 1 \& 0 \\ \& 0 \& 0 \& 0 \& 0 \& \lambda_0 + \lambda_3 = 1 \& 1 \& 1 \& 2 \& \\ 2 \& 1 \backslash \backslash \lambda_1 - \lambda_2 = 0 \& 1 \& 0 \& 0 \& 0 \& 0 \& \\ 0 \& 0 \& \lambda_1 + \lambda_2 = 0 \& 1 \& 2 \& 2 \& 2 \& 1 \backslash \backslash \\ \lambda_1 - \lambda_3 = 0 \& 1 \& 1 \& 0 \& 0 \& 0 \& 0 \& 0 \& \\ \& \lambda_1 + \lambda_3 = 0 \& 1 \& 1 \& 2 \& 2 \& 1 \backslash \backslash \\ \lambda_2 - \lambda_3 = 0 \& 0 \& 1 \& 0 \& 0 \& 0 \& 0 \& \lambda_2 + \lambda_3 = 0 \& 0 \& 1 \& 2 \& 2 \& 1 \& 1 \\ \\ \lambda_0 + \frac{1}{2}(\mu_2 - \mu_3) = 1 \& 1 \& 1 \& 1 \& 1 \& 1 \& 1 \\ \lambda_1 + \frac{1}{2}(\mu_2 - \mu_3) = 0 \& 1 \& 1 \& 1 \& 1 \& 1 \& 1 \\ \lambda_2 + \frac{1}{2}(\mu_2 - \mu_3) = 0 \& 0 \& 1 \& 1 \& 1 \& 1 \& 1 \\ +\frac{1}{2}(\mu_2 - \mu_3) = 0 \& 0 \& 1 \& 1 \& 1 \& 1 \& 1 \end{array}$	$\begin{array}{l} \lambda_0 - \lambda_1 = 1 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad \lambda_0 + \lambda_1 = 1 \quad 2 \quad 2 \quad 2 \quad 2 \quad 1 \quad 1 \\ \lambda_0 - \lambda_2 = 1 \quad 1 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad \lambda_0 + \lambda_2 = 1 \quad 1 \quad 2 \quad 2 \quad 2 \quad 1 \quad 1 \\ \lambda_0 - \lambda_3 = 1 \quad 1 \quad 1 \quad 0 \quad 0 \quad 0 \quad 0 \quad \lambda_0 + \lambda_3 = 1 \quad 1 \quad 1 \quad 2 \quad 2 \quad 1 \quad 1 \\ \lambda_1 - \lambda_2 = 0 \quad 1 \quad 0 \quad 0 \quad 0 \quad 0 \quad 0 \quad \lambda_1 + \lambda_2 = 0 \quad 1 \quad 2 \quad 2 \quad 2 \quad 1 \quad 1 \\ \lambda_1 - \lambda_3 = 0 \quad 1 \quad 1 \quad 0 \quad 0 \quad 0 \quad 0 \quad \lambda_1 + \lambda_3 = 0 \quad 1 \quad 1 \quad 2 \quad 2 \quad 1 \quad 1 \\ \lambda_2 - \lambda_3 = 0 \quad 0 \quad 1 \quad 0 \quad 0 \quad 0 \quad 0 \quad \lambda_2 + \lambda_3 = 0 \quad 0 \quad 1 \quad 2 \quad 2 \quad 1 \quad 1 \\ \\ \lambda_0 + \frac{1}{2}(\mu_2 - \mu_3) = 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \\ \lambda_1 + \frac{1}{2}(\mu_2 - \mu_3) = 0 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \\ \lambda_2 + \frac{1}{2}(\mu_2 - \mu_3) = 0 \quad 0 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \\ +\frac{1}{2}(\mu_2 - \mu_3) = 0 \quad 0 \quad 0 \quad 1 \quad 1 \quad 1 \quad 1 \end{array}$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_EQ1187	200	■ 0.183 ● C +279	\begin{aligned} &\&=(\boldsymbol{v}, \boldsymbol{w})(\boldsymbol{a}, *(\boldsymbol{b} \wedge \boldsymbol{x})) - (\boldsymbol{a}_1, \boldsymbol{w})(\boldsymbol{w} \wedge \boldsymbol{a}_2, *(\boldsymbol{b} \wedge \boldsymbol{x})) \\ &\quad + (\boldsymbol{a}_2, \boldsymbol{w})(\boldsymbol{w} \wedge \boldsymbol{a}_1, *(\boldsymbol{b} \wedge \boldsymbol{x})) - \frac{3}{5}(*(\boldsymbol{a} \wedge \boldsymbol{b}), \boldsymbol{x})(\boldsymbol{v}, \boldsymbol{w}) \\ &= \frac{2}{5}(*(\boldsymbol{a} \wedge \boldsymbol{b}), \boldsymbol{x})(\boldsymbol{v}, \boldsymbol{w}) - (\boldsymbol{a}_1, \boldsymbol{w})(\boldsymbol{b} \wedge \boldsymbol{x} \wedge \boldsymbol{w} \wedge \boldsymbol{a}_2, \boldsymbol{e}_1 \wedge \cdots \wedge \boldsymbol{e}_5) \\ &\quad + (\boldsymbol{a}_2, \boldsymbol{w})(\boldsymbol{b} \wedge \boldsymbol{x} \wedge \boldsymbol{w} \wedge \boldsymbol{a}_1, \boldsymbol{e}_1 \wedge \cdots \wedge \boldsymbol{e}_5), \\ &((\boldsymbol{b} \times *(\boldsymbol{a} \wedge \boldsymbol{x}))\boldsymbol{v}, \boldsymbol{w}) \\ &= (\boldsymbol{x} \wedge \boldsymbol{a}, \boldsymbol{w} \wedge *(\boldsymbol{b} \wedge \boldsymbol{v})) - \frac{3}{5}(*(\boldsymbol{a} \wedge \boldsymbol{b}), \boldsymbol{x})(\boldsymbol{v}, \boldsymbol{w}) \\ &= (\boldsymbol{v}, \boldsymbol{w})(*(\boldsymbol{a} \wedge \boldsymbol{b}), \boldsymbol{v}) + (\boldsymbol{a}_1, \boldsymbol{w})(\boldsymbol{b} \wedge \boldsymbol{x} \wedge \boldsymbol{w} \wedge \boldsymbol{a}_2, \boldsymbol{e}_1 \wedge \cdots \wedge \boldsymbol{e}_5) \\ &\quad - (\boldsymbol{a}_2, \boldsymbol{w})(\boldsymbol{b} \wedge \boldsymbol{x} \wedge \boldsymbol{w} \wedge \boldsymbol{a}_1, \boldsymbol{e}_1 \wedge \cdots \wedge \boldsymbol{e}_5) - \frac{3}{5}(*(\boldsymbol{a} \wedge \boldsymbol{b}), \boldsymbol{x})(\boldsymbol{v}, \boldsymbol{w}). \end{aligned}	(not rendered)	$ \begin{aligned} &= (v, w)(a, *(b \wedge x)) - (a_1, w)(w \wedge a_2, *(b \wedge x)) \\ &\quad + (a_2, w)(w \wedge a_1, *(b \wedge x)) - \frac{3}{5}(*(a \wedge b), x)(v, w) \\ &= \frac{2}{5}(*(a \wedge b), x)(v, w) - (a_1, w)(b \wedge x \wedge w \wedge a_2, e_1 \wedge \cdots \wedge e_5) \\ &\quad + (a_2, w)(b \wedge x \wedge w \wedge a_1, e_1 \wedge \cdots \wedge e_5), \\ &((b \times *(a \wedge x))v, w) \\ &= (x \wedge a, w \wedge *(b \wedge v)) - \frac{3}{5}(*(a \wedge b), x)(v, w) \\ &= (v, w)(*(a \wedge b), v) + (a_1, w)(b \wedge x \wedge w \wedge a_2, e_1 \wedge \cdots \wedge e_5) \\ &\quad - (a_2, w)(b \wedge x \wedge w \wedge a_1, e_1 \wedge \cdots \wedge e_5) - \frac{3}{5}(*(a \wedge b), x)(v, w). \end{aligned} $
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_E00830	138	0.186 W +0	$\begin{aligned} \kappa(X, Y, \xi, \eta) = & \kappa \left(\begin{array}{ccc} \xi_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \xi_2 & x_1 \\ x_2 & \bar{x}_1 & \xi_3 \end{array} \right), \left(\begin{array}{ccc} \eta_1 & y_3 & \bar{y}_2 \\ \bar{y}_3 & \eta_2 & y_1 \\ y_2 & \bar{y}_1 & \eta_3 \end{array} \right), \xi, \eta \right) \\ = & \left(\begin{array}{ccc} -\xi_1 & 0 & 0 \\ 0 & \xi_2 & x_1 \\ 0 & \bar{x}_1 & \xi_3 \end{array} \right), \left(\begin{array}{ccc} \eta_1 & 0 & 0 \\ 0 & -\eta_2 & -y_1 \\ 0 & -\bar{y}_1 & -\eta_3 \end{array} \right), -\xi, \eta \right), \\ \mu(X, Y, \xi, \eta) = & \left(\begin{array}{ccc} \eta & 0 & 0 \\ 0 & \eta_3 & -y_1 \\ 0 & -\bar{y}_1 & \eta_2 \end{array} \right), \left(\begin{array}{ccc} \xi & 0 & 0 \\ 0 & \xi_3 & -x_1 \\ 0 & -\bar{x}_1 & \xi_2 \end{array} \right), \eta_1, \xi_1 \right). \end{aligned}$	$\begin{aligned} \kappa(X, Y, \xi, \eta) &= \kappa \left(\begin{pmatrix} \xi_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \xi_2 & x_1 \\ x_2 & \bar{x}_1 & \xi_3 \end{pmatrix}, \begin{pmatrix} \eta_1 & y_3 & \bar{y}_2 \\ \bar{y}_3 & \eta_2 & y_1 \\ y_2 & \bar{y}_1 & \eta_3 \end{pmatrix}, \xi, \eta \right) \\ &= \left(\begin{pmatrix} -\xi_1 & 0 & 0 \\ 0 & \xi_2 & x_1 \\ 0 & \bar{x}_1 & \xi_3 \end{pmatrix}, \begin{pmatrix} \eta_1 & 0 & 0 \\ 0 & -\eta_2 & -y_1 \\ 0 & -\bar{y}_1 & -\eta_3 \end{pmatrix}, -\xi, \eta \right), \\ \mu(X, Y, \xi, \eta) &= \left(\begin{pmatrix} \eta & 0 & 0 \\ 0 & \eta_3 & -y_1 \\ 0 & -\bar{y}_1 & \eta_2 \end{pmatrix}, \begin{pmatrix} \xi & 0 & 0 \\ 0 & \xi_3 & -x_1 \\ 0 & -\bar{x}_1 & \xi_2 \end{pmatrix}, \eta_1, \xi_1 \right). \end{aligned}$	
0902.0431_E01041	177	0.204 C +4	$\begin{aligned} & \left(\alpha_i, \alpha_i \right) = \frac{1}{30}, \quad i=1,2,3,4,5,6,7,8 \quad \& \quad \left(\alpha_i, \alpha_{i+1} \right) = -\frac{1}{60}, \quad i=1,2,\cdots,6, \quad \left(\alpha_3, \alpha_8 \right) = -\frac{1}{60}, \\ & \left(\alpha_i, \alpha_j \right) = -\frac{1}{60}, \quad i=1,2, \cdots, 6, \quad \text{otherwise} \\ & \left(-\mu, -\mu \right) = \frac{1}{30}, \quad \left(-\mu, \alpha_7 \right) = -\frac{1}{60}, \quad \left(-\mu, \alpha_i \right) = 0, \quad i=1,2,3,4,5,6,8, \end{aligned}$	$\begin{aligned} & \left(\alpha_i, \alpha_i \right) = \frac{1}{30}, \quad i=1,2,3,4,5,6,7,8 \\ & \left(\alpha_i, \alpha_{i+1} \right) = -\frac{1}{60}, \quad i=1,2,\cdots,6, \quad \left(\alpha_3, \alpha_8 \right) = -\frac{1}{60} \\ & \left(\alpha_i, \alpha_j \right) = 0, \quad \text{otherwise} \\ & \left(-\mu, -\mu \right) = \frac{1}{30}, \quad \left(-\mu, \alpha_7 \right) = -\frac{1}{60}, \quad \left(-\mu, \alpha_i \right) = 0, \quad i=1,2,3,4,5,6,8, \end{aligned}$	
0902.0431_E00340	054	0.226 W +0	$\begin{aligned} \lambda_0 &= \alpha_1 + \alpha_2 + \alpha_3 \\ \lambda_1 &= \alpha_2 + \alpha_3 \\ \lambda_3 &= \alpha_3 \\ \lambda_3 &= \alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4, \end{aligned}$	$\begin{aligned} \lambda_0 &= \alpha_1 + \alpha_2 + \alpha_3 \\ \lambda_1 &= \alpha_2 + \alpha_3 \\ \lambda_3 &= \alpha_3 \\ \lambda_3 &= \alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4, \end{aligned}$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_EQ0206	035	■ 0.294 ● W +0	$\begin{aligned} & \kappa\left(\alpha_1, \alpha_2, \alpha_3\right)=\left(\kappa \alpha_1, \kappa \alpha_3, \kappa \alpha_2\right), \\ & \pi\left(\alpha_1, \alpha_2, \alpha_3\right)=\left(\kappa \alpha_3, \kappa \alpha_2, \kappa \alpha_1\right), \\ & \nu\left(\alpha_1, \alpha_2, \alpha_3\right)=\left(\alpha_2, \alpha_3, \alpha_1\right), \end{aligned}$	$\begin{aligned} & \kappa\left(\alpha_1, \alpha_2, \alpha_3\right)=\left(\kappa \alpha_1, \kappa \alpha_3, \kappa \alpha_2\right), \\ & \pi\left(\alpha_1, \alpha_2, \alpha_3\right)=\left(\kappa \alpha_3, \kappa \alpha_2, \kappa \alpha_1\right), \\ & \nu\left(\alpha_1, \alpha_2, \alpha_3\right)=\left(\alpha_2, \alpha_3, \alpha_1\right), \end{aligned}$	$\begin{aligned} & \kappa\left(\alpha_1, \alpha_2, \alpha_3\right)=\left(\kappa \alpha_1, \kappa \alpha_3, \kappa \alpha_2\right), \\ & \pi\left(\alpha_1, \alpha_2, \alpha_3\right)=\left(\kappa \alpha_3, \kappa \alpha_2, \kappa \alpha_1\right), \\ & \nu\left(\alpha_1, \alpha_2, \alpha_3\right)=\left(\alpha_2, \alpha_3, \alpha_1\right), \end{aligned}$
0902.0431_EQ0655	108	■ 0.299 ● C +4	$\begin{aligned} & \left(P M \tau h(A, B)^*\right)^*\left(P N \tau h(A, B)^*\right)=\tau h(A, B) M^* P^* P N \tau h(A, B)^* \\ & =\tau\left(h(A, B) \tau\left(M^* N\right) h(A, B)^*\right), \\ & \frac{\left(P M \tau h(A, B)^*\right)\left(h(A, B) Y h(A, B)^*\right)}{\left(P M \tau h(A, B)^*\right) \times\left(P N \tau h(A, B)^*\right)}=\frac{\tau\left(P \tau(M Y) \tau h(A, B)^*\right)}{i \tilde{P}(M \times N) \tau^t\left(h(A, B)^*\right)} \\ & =P(\overline{M} \times \overline{N}) h(A, B)^*=\tau\left(P(\overline{M} \times \overline{N}) \tau h(A, B)^*\right) . \end{aligned}$	$\begin{aligned} & \left(P M \tau h(A, B)^*\right)^*\left(P N \tau h(A, B)^*\right)=\tau h(A, B) M^* P^* P N \tau h(A, B)^* \\ & =\tau\left(h(A, B) \tau\left(M^* N\right) h(A, B)^*\right), \\ & \frac{\left(P M \tau h(A, B)^*\right)\left(h(A, B) Y h(A, B)^*\right)}{\left(P M \tau h(A, B)^*\right) \times\left(P N \tau h(A, B)^*\right)}=\frac{\tau\left(P \tau(M Y) \tau h(A, B)^*\right)}{i \tilde{P}(M \times N) \tau^t\left(h(A, B)^*\right)} \\ & =P(\overline{M} \times \overline{N}) h(A, B)^*=\tau\left(P(\overline{M} \times \overline{N}) \tau h(A, B)^*\right) . \end{aligned}$	$\begin{aligned} & \left(P M \tau h(A, B)^*\right)^*\left(P N \tau h(A, B)^*\right)=\tau h(A, B) M^* P^* P N \tau h(A, B)^* \\ & =\tau\left(h(A, B) \tau\left(M^* N\right) h(A, B)^*\right), \\ & \frac{\left(P M \tau h(A, B)^*\right)\left(h(A, B) Y h(A, B)^*\right)}{\left(P M \tau h(A, B)^*\right) \times\left(P N \tau h(A, B)^*\right)}=\frac{\tau\left(P \tau(M Y) \tau h(A, B)^*\right)}{i \tilde{P}(M \times N) \tau^t\left(h(A, B)^*\right)} \\ & =P(\overline{M} \times \overline{N}) h(A, B)^*=\tau\left(P(\overline{M} \times \overline{N}) \tau h(A, B)^*\right) . \end{aligned}$
0902.0431_EQ0101	020	■ 0.306 ● W +1	$\begin{aligned} & \lambda_1-\lambda_2=\alpha_1, \quad \lambda_1-\lambda_3=2 \alpha_1+3 \alpha_2, \quad \lambda_2-\lambda_3=\alpha_1+3 \alpha_2, \\ & \lambda_1=\alpha_1+\alpha_2, \quad \lambda_2=\alpha_2, \quad \quad \quad =\lambda_3+2 \alpha_2 \end{aligned}$	$\begin{aligned} & \lambda_1-\lambda_2=\alpha_1, \quad \lambda_1-\lambda_3=2 \alpha_1+3 \alpha_2, \quad \lambda_2-\lambda_3=\alpha_1+3 \alpha_2, \\ & \lambda_1=\alpha_1+\alpha_2, \quad \lambda_2=\alpha_2, \quad \quad \quad =\lambda_3+2 \alpha_2 \end{aligned}$	$\begin{aligned} & \lambda_1-\lambda_2=\alpha_1, \quad \lambda_1-\lambda_3=2 \alpha_1+3 \alpha_2, \quad \lambda_2-\lambda_3=\alpha_1+3 \alpha_2, \\ & \lambda_1=\alpha_1+\alpha_2, \quad \lambda_2=\alpha_2, \quad \quad \quad -\lambda_3=\alpha_1+2 \alpha_2 . \end{aligned}$
0902.0431_EQ1154	195	■ 0.318 ● N +0	$\begin{aligned} & +\left(\boldsymbol{v}, \boldsymbol{w}\right)\left(\boldsymbol{u}, \boldsymbol{a}\right)-\sum_{i=1}^3 \sum_{j=1}^3\left(\boldsymbol{u}_i \wedge \boldsymbol{v}_j \wedge \boldsymbol{v}_{j+1}, \boldsymbol{w}\right)\left(\boldsymbol{u}_{i+1} \wedge \boldsymbol{u}_{i+2} \wedge \boldsymbol{v}_{j+2}, \boldsymbol{a}\right) \\ & =-\left(\boldsymbol{u} \wedge \boldsymbol{v}, \boldsymbol{w} \wedge \boldsymbol{a}\right)=-\left(*\left(*\left(\boldsymbol{u} \wedge \boldsymbol{v}\right) \wedge \boldsymbol{w}\right), \boldsymbol{a}\right) \end{aligned}$	$\begin{aligned} & +\left(\boldsymbol{v}, \boldsymbol{w}\right)\left(\boldsymbol{u}, \boldsymbol{a}\right)-\sum_{i=1}^3 \sum_{j=1}^3\left(\boldsymbol{u}_i \wedge \boldsymbol{v}_j \wedge \boldsymbol{v}_{j+1}, \boldsymbol{w}\right)\left(\boldsymbol{u}_{i+1} \wedge \boldsymbol{u}_{i+2} \wedge \boldsymbol{v}_{j+2}, \boldsymbol{a}\right) \\ & =-\left(\boldsymbol{u} \wedge \boldsymbol{v}, \boldsymbol{w} \wedge \boldsymbol{a}\right)=-\left(*\left(*\left(\boldsymbol{u} \wedge \boldsymbol{v}\right) \wedge \boldsymbol{w}\right), \boldsymbol{a}\right) \end{aligned}$	$\begin{aligned} & +\left(\boldsymbol{v}, \boldsymbol{w}\right)\left(\boldsymbol{u}, \boldsymbol{a}\right)-\sum_{i=1}^3 \sum_{j=1}^3\left(\boldsymbol{u}_i \wedge \boldsymbol{v}_j \wedge \boldsymbol{v}_{j+1}, \boldsymbol{w}\right)\left(\boldsymbol{u}_{i+1} \wedge \boldsymbol{u}_{i+2} \wedge \boldsymbol{v}_{j+2}, \boldsymbol{a}\right) \\ & =-\left(\boldsymbol{u} \wedge \boldsymbol{v}, \boldsymbol{w} \wedge \boldsymbol{a}\right)=-\left(*\left(*\left(\boldsymbol{u} \wedge \boldsymbol{v}\right) \wedge \boldsymbol{w}\right), \boldsymbol{a}\right) . \end{aligned}$
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
9902.0431_E00802	134	0.338 W +2	$\left(\begin{array}{c} \left(\begin{array}{ccc} \cos a_1 a_2\frac{\sin a_2 }{ a_2 }a_3\frac{\sin a_3 }{ a_3 } & 0 & 0 \\ 0 & a_1\frac{\sin a_1 }{ a_1 }\cos a_2 a_3\frac{\sin a_3 }{ a_3 } & 0 \\ 0 & 0 & a_1\frac{\sin a_1 }{ a_1 }a_2\frac{\sin a_2 }{ a_2 }\cos a_3 \end{array}\right) \\ \left(\begin{array}{ccc} a_1\frac{\sin a_1 }{ a_1 }\cos a_2 \cos a_3 & 0 & 0 \\ 0 & \cos a_1 a_2\frac{\sin a_2 }{ a_2 }\cos a_3 & 0 \\ 0 & 0 & \cos a_1 \cos a_2 a_3\frac{\sin a_3 }{ a_3 } \end{array}\right) \\ \cos a_1 \cos a_2 \cos a_3 \\ a_1\frac{\sin a_1 }{ a_1 }a_2\frac{\sin a_2 }{ a_2 }a_3\frac{\sin a_3 }{ a_3 } \end{array}\right)$	(not rendered)	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_E00875	145	■ 0.339 ● N +1	$\begin{aligned} & \begin{array}{l} \begin{aligned} & \varphi_2(A) \left(\begin{pmatrix} \xi_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \xi_2 & x_1 \\ x_2 & \bar{x}_1 & \xi_3 \end{pmatrix}, \begin{pmatrix} \eta_1 & y_3 & \bar{y}_2 \\ \bar{y}_3 & \eta_2 & y_1 \\ y_2 & \bar{y}_1 & \eta_3 \end{pmatrix}, \xi, \eta \right) \\ &= \left(\begin{pmatrix} \xi'_1 & x'_3 & \bar{x}'_2 \\ \bar{x}'_3 & \xi'_2 & x'_1 \\ x'_2 & \bar{x}'_1 & \xi'_3 \end{pmatrix}, \begin{pmatrix} \eta'_1 & y'_3 & \bar{y}'_2 \\ \bar{y}'_3 & \eta'_2 & y'_1 \\ y'_2 & \bar{y}'_1 & \eta'_3 \end{pmatrix}, \xi', \eta' \right) \end{aligned} \end{array} \end{aligned}$	$\varphi_2(A) \left(\begin{pmatrix} \xi_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \xi_2 & x_1 \\ x_2 & \bar{x}_1 & \xi_3 \end{pmatrix}, \begin{pmatrix} \eta_1 & y_3 & \bar{y}_2 \\ \bar{y}_3 & \eta_2 & y_1 \\ y_2 & \bar{y}_1 & \eta_3 \end{pmatrix}, \xi, \eta \right) \\ = \left(\begin{pmatrix} \xi'_1 & x'_3 & \bar{x}'_2 \\ \bar{x}'_3 & \xi'_2 & x'_1 \\ x'_2 & \bar{x}'_1 & \xi'_3 \end{pmatrix}, \begin{pmatrix} \eta'_1 & y'_3 & \bar{y}'_2 \\ \bar{y}'_3 & \eta'_2 & y'_1 \\ y'_2 & \bar{y}'_1 & \eta'_3 \end{pmatrix}, \xi', \eta' \right),$	
0902.0431_E00400	066	■ 0.351 ● N +1	$\begin{aligned} & \beta_1 m = \text{prm} \bar{r}, \quad \beta_2 m = r m \bar{r} q, \quad \beta_3 m = \bar{q} r m \bar{r} \bar{p}, \quad m \in \mathbf{H}. \\ & \beta_1 m = \text{prm} \bar{r}, \quad \beta_2 m = r m \bar{r} q, \quad \beta_3 m = \bar{q} r m \bar{r} \bar{p}, \quad m \in \mathbf{H}. \end{aligned}$	$\beta_1 m = \text{prm} \bar{r}, \quad \beta_2 m = r m \bar{r} q, \quad \beta_3 m = \bar{q} r m \bar{r} \bar{p}, \quad m \in \mathbf{H}.$	
0902.0431_E00860	143	■ 0.354 ● W +1	$\begin{aligned} & 1 \longrightarrow \text{Spin}(10) \longrightarrow ((E_7)^{\kappa, \mu})_{(0, E_1, 0, 1)} \longrightarrow S^{10} \longrightarrow * \\ & \quad \quad \quad \downarrow p' \quad \quad \quad \downarrow p \quad \quad \quad \downarrow = \\ & 1 \longrightarrow SO(10) \longrightarrow SO(11) \longrightarrow S^{10} \longrightarrow * \end{aligned}$	$\begin{aligned} & 1 \longrightarrow \text{Spin}(10) \longrightarrow ((E_7)^{\kappa, \mu})_{(0, E_1, 0, 1)} \longrightarrow S^{10} \longrightarrow * \\ & \quad \quad \quad \downarrow p' \quad \quad \quad \downarrow p \quad \quad \quad \downarrow = \\ & 1 \longrightarrow SO(10) \longrightarrow SO(11) \longrightarrow S^{10} \longrightarrow * \end{aligned}$	
0902.0431_E00876	145	■ 0.358 ● C +76	$\begin{aligned} & \binom{\xi'_1}{\eta'} = A \binom{\xi_1}{\eta}, \quad \binom{\xi'_2}{\eta'_1} = A \binom{\xi}{\eta_1}, \quad \binom{\eta'_2}{\xi'_3} = A \binom{\eta_2}{\xi_3}, \quad \binom{\eta'_3}{\xi'_2} = A \binom{\eta_3}{\xi_2} \\ & \binom{x'_1}{y'_1} = \tau A \binom{x_1}{y_1}, \quad \binom{x'_2}{y'_2} = \binom{x_2}{y_2}, \quad \binom{x'_3}{y'_3} = \binom{x_3}{y_3}. \end{aligned}$	$\begin{aligned} & \binom{\xi'_1}{\eta'} = A \binom{\xi_1}{\eta}, \quad \binom{\xi'_2}{\eta'_1} = A \binom{\xi}{\eta_1}, \quad \binom{\eta'_2}{\xi'_3} = A \binom{\eta_2}{\xi_3}, \quad \binom{\eta'_3}{\xi'_2} = A \binom{\eta_3}{\xi_2}, \\ & \binom{x'_1}{y'_1} = \tau A \binom{x_1}{y_1}, \quad \binom{x'_2}{y'_2} = \binom{x_2}{y_2}, \quad \binom{x'_3}{y'_3} = \binom{x_3}{y_3}. \end{aligned}$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5–0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902_0431_EQ0941	156	0.372 W +2	$\begin{aligned} & \text{\texttt{\begin{aligned} & P=\left(F_1(1), 0,0,0\right)} \\ & \text{\texttt{\& \xrightarrow{f} \boldsymbol{e}_2 \wedge \boldsymbol{e}_4 \wedge \boldsymbol{e}_5 + \boldsymbol{e}_3 \wedge \boldsymbol{e}_6 \wedge \boldsymbol{e}_4}} \\ & \text{\texttt{\& \xrightarrow{D} (De_2 \wedge e_4 \wedge e_5 + e_2 \wedge De_4 \wedge e_5 + e_2 \wedge e_4 \wedge De_5)}} \\ & \text{\texttt{+ (De_3 \wedge e_6 \wedge e_4 + e_3 \wedge De_6 \wedge e_4 + e_3 \wedge e_6 \wedge De_4)}} \\ & \text{\texttt{= \left(b_{12}e_1 \wedge e_4 \wedge e_5 + \left(b_{22} + \frac{\nu}{3}\right)e_2 \wedge e_4 \wedge e_5 - \bar{b}_{23}e_3 \wedge e_4 \wedge e_5 - \bar{l}_{23}e_6 \wedge e_4 \wedge e_5}} \\ & \text{\texttt{+ l_{11}e_2 \wedge e_1 \wedge e_5 + l_{31}e_2 \wedge e_3 \wedge e_5 + \left(c_{11} - \frac{\nu}{3}\right)e_2 \wedge e_4 \wedge e_5 - \bar{c}_{13}e_2 \wedge e_6 \wedge e_5}} \\ & \text{\texttt{+ l_{12}e_2 \wedge e_4 \wedge e_1 + l_{32}e_2 \wedge e_4 \wedge e_3 + \left(c_{22} - \frac{\nu}{3}\right)e_3 \wedge e_6 \wedge e_4 - \bar{c}_{23}e_2 \wedge e_4 \wedge e_6}} \\ & \text{\texttt{+ \left(b_{13}e_1 \wedge e_6 \wedge e_4 + b_{23}e_2 \wedge e_6 \wedge e_4 + \left(b_{33} + \frac{\nu}{3}\right)e_3 \wedge e_6 \wedge e_4 - \bar{l}_{32}e_5 \wedge e_6 \wedge e_4}} \\ & \text{\texttt{+ l_{13}e_3 \wedge e_1 \wedge e_4 + l_{23}e_3 \wedge e_2 \wedge e_4 + c_{23}e_3 \wedge e_5 \wedge e_4 + \left(c_{33} - \frac{\nu}{3}\right)e_3 \wedge e_6 \wedge e_4}} \\ & \text{\texttt{+ l_{11}e_3 \wedge e_6 \wedge e_1 + l_{21}e_3 \wedge e_6 \wedge e_2 + \left(c_{11} - \frac{\nu}{3}\right)e_3 \wedge e_6 \wedge e_4 - \bar{c}_{12}e_3 \wedge e_6 \wedge e_5}} \\ & \text{\texttt{\left(\begin{pmatrix} 0 & h(b_{13}, c_{13}) & 0 \\ * & h(b_{23} + \bar{c}_{23}) & h(b_{22} - c_{33} - \frac{\nu}{3}, -b_{33} + c_{33} + \frac{\nu}{3}) \\ h(\bar{c}_{12}, \bar{b}_{12}) & * & -h(\bar{b}_{23} + c_{23}) \end{pmatrix}\right)}} \\ & \text{\texttt{\left(\begin{pmatrix} -h(l_{23} + l_{32}) & h(l_{13}, \bar{l}_{31}) & * \\ * & 0 & -h(l_{11}) \\ h(l_{21}, \bar{l}_{12}) & * & 0 \end{pmatrix}\right)}} \\ & \text{\texttt{\left(\begin{pmatrix} 0 & h(b_{13}, c_{13}) & 0 \\ * & h(b_{23} + \bar{c}_{23}) & h(b_{22} - c_{33} - \frac{\nu}{3}, -b_{33} + c_{33} + \frac{\nu}{3}) \\ * & * & -h(b_{23} + c_{23}) \end{pmatrix}\right)}} \end{aligned}} \end{aligned}$	$P = (F_1(1), 0, 0, 0)$ $\xrightarrow{f} \boldsymbol{e}_2 \wedge \boldsymbol{e}_4 \wedge \boldsymbol{e}_5 + \boldsymbol{e}_3 \wedge \boldsymbol{e}_6 \wedge \boldsymbol{e}_4$ $\xrightarrow{D} (De_2 \wedge e_4 \wedge e_5 + e_2 \wedge De_4 \wedge e_5 + e_2 \wedge e_4 \wedge De_5)$ $+ (De_3 \wedge e_6 \wedge e_4 + e_3 \wedge De_6 \wedge e_4 + e_3 \wedge e_6 \wedge De_4)$ $= \left(b_{12}e_1 \wedge e_4 \wedge e_5 + \left(b_{22} + \frac{\nu}{3}\right)e_2 \wedge e_4 \wedge e_5 - \bar{b}_{23}e_3 \wedge e_4 \wedge e_5 - \bar{l}_{23}e_6 \wedge e_4 \wedge e_5\right.$ $+ l_{11}e_2 \wedge e_1 \wedge e_5 + l_{31}e_2 \wedge e_3 \wedge e_5 + \left(c_{11} - \frac{\nu}{3}\right)e_2 \wedge e_4 \wedge e_5 - \bar{c}_{13}e_2 \wedge e_6 \wedge e_5$ $+ l_{12}e_2 \wedge e_4 \wedge e_1 + l_{32}e_2 \wedge e_4 \wedge e_3 + \left(c_{22} - \frac{\nu}{3}\right)e_3 \wedge e_6 \wedge e_4 - \bar{c}_{23}e_2 \wedge e_4 \wedge e_6$ $+ \left(b_{13}e_1 \wedge e_6 \wedge e_4 + b_{23}e_2 \wedge e_6 \wedge e_4 + \left(b_{33} + \frac{\nu}{3}\right)e_3 \wedge e_6 \wedge e_4 - \bar{l}_{32}e_5 \wedge e_6 \wedge e_4$ $+ l_{13}e_3 \wedge e_1 \wedge e_4 + l_{23}e_3 \wedge e_2 \wedge e_4 + c_{23}e_3 \wedge e_5 \wedge e_4 + \left(c_{33} - \frac{\nu}{3}\right)e_3 \wedge e_6 \wedge e_4$ $+ l_{11}e_3 \wedge e_6 \wedge e_1 + l_{21}e_3 \wedge e_6 \wedge e_2 + \left(c_{11} - \frac{\nu}{3}\right)e_3 \wedge e_6 \wedge e_4 - \bar{c}_{12}e_3 \wedge e_6 \wedge e_5$ $\xrightarrow{f^{-1}} \left(\begin{array}{ccc} 0 & h(b_{13}, c_{13}) & 0 \\ * & h(b_{23} + \bar{c}_{23}) & h(b_{22} - c_{33} - \frac{\nu}{3}, -b_{33} + c_{33} + \frac{\nu}{3}) \\ * & * & -h(b_{23} + c_{23}) \end{array}\right)$	$P = (F_1(1), 0, 0, 0)$ $\xrightarrow{f} \boldsymbol{e}_2 \wedge \boldsymbol{e}_4 \wedge \boldsymbol{e}_5 + \boldsymbol{e}_3 \wedge \boldsymbol{e}_6 \wedge \boldsymbol{e}_4$ $\xrightarrow{D} (De_2 \wedge e_4 \wedge e_5 + e_2 \wedge De_4 \wedge e_5 + e_2 \wedge e_4 \wedge De_5)$ $+ (De_3 \wedge e_6 \wedge e_4 + e_3 \wedge De_6 \wedge e_4 + e_3 \wedge e_6 \wedge De_4)$ $= \left(b_{12}e_1 \wedge e_4 \wedge e_5 + \left(b_{22} + \frac{\nu}{3}\right)e_2 \wedge e_4 \wedge e_5 - \bar{b}_{23}e_3 \wedge e_4 \wedge e_5 - \bar{l}_{23}e_6 \wedge e_4 \wedge e_5\right.$ $+ l_{11}e_2 \wedge e_1 \wedge e_5 + l_{31}e_2 \wedge e_3 \wedge e_5 + \left(c_{11} - \frac{\nu}{3}\right)e_2 \wedge e_4 \wedge e_5 - \bar{c}_{13}e_2 \wedge e_6 \wedge e_5$ $+ l_{12}e_2 \wedge e_4 \wedge e_1 + l_{32}e_2 \wedge e_4 \wedge e_3 + \left(c_{22} - \frac{\nu}{3}\right)e_3 \wedge e_6 \wedge e_4 - \bar{c}_{23}e_2 \wedge e_4 \wedge e_6$ $+ \left(b_{13}e_1 \wedge e_6 \wedge e_4 + b_{23}e_2 \wedge e_6 \wedge e_4 + \left(b_{33} + \frac{\nu}{3}\right)e_3 \wedge e_6 \wedge e_4 - \bar{l}_{32}e_5 \wedge e_6 \wedge e_4$ $+ l_{13}e_3 \wedge e_1 \wedge e_4 + l_{23}e_3 \wedge e_2 \wedge e_4 + c_{23}e_3 \wedge e_5 \wedge e_4 + \left(c_{33} - \frac{\nu}{3}\right)e_3 \wedge e_6 \wedge e_4$ $+ l_{11}e_3 \wedge e_6 \wedge e_1 + l_{21}e_3 \wedge e_6 \wedge e_2 + \left(c_{11} - \frac{\nu}{3}\right)e_3 \wedge e_6 \wedge e_4 - \bar{c}_{12}e_3 \wedge e_6 \wedge e_5$ $\xrightarrow{f^{-1}} \left(\begin{array}{ccc} 0 & h(b_{13}, c_{13}) & 0 \\ * & h(b_{23} + \bar{c}_{23}) & h(b_{22} - c_{33} - \frac{\nu}{3}, -b_{33} + c_{33} + \frac{\nu}{3}) \\ * & * & -h(b_{23} + c_{23}) \end{array}\right)$

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_E00093	019	0.382 W +1	$\begin{aligned} H_{\{1\}} H_{\{1\}} e_{\{2\}} &= H_{\{1\}} e_{\{3\}} = -e_{\{2\}}, \\ &\quad \end{aligned}$	$H_1 H_1 e_2 = H_1 e_3 = -e_2,$	$H_1 H_1 e_2 = H_1 e_3 = -e_2, \quad H_1 H_1 e_3 = -H_1 e_2 = -e_3,$ $H_1 H_1 e_4 = -H_1 e_5 = -e_4, \quad H_1 H_1 e_5 = H_1 e_4 = -e,$ $H_1 H_1 e_i = 0 \text{ otherwise.}$
0902.0431_E00371	061	0.388 C +8	$\begin{aligned} &\operatorname{tr}(X) = \operatorname{tr}(\alpha X) \text{ (Lemma 2.2.1.(2))} = \sum_{i=1}^3 \xi_i, \\ &(X, X) = (\alpha X, \alpha X) \text{ (Lemma 2.2.4)} = \sum_{i=1}^3 \xi_i^2, \\ &\operatorname{tr}(X, X, X) = \operatorname{tr}(\alpha X, \alpha X, \alpha X) \text{ (Lemma 2.2.4)} = \sum_{i=1}^3 \xi_i^3. \end{aligned}$	$\operatorname{tr}(X) = \operatorname{tr}(\alpha X) \text{ (Lemma 2.2.1.(2))} = \sum_{i=1}^3 \xi_i,$ $(X, X) = (\alpha X, \alpha X) \text{ (Lemma 2.2.4)} = \sum_{i=1}^3 \xi_i^2,$ $\operatorname{tr}(X, X, X) = \operatorname{tr}(\alpha X, \alpha X, \alpha X) \text{ (Lemma 2.2.4)} = \sum_{i=1}^3 \xi_i^3.$	$\operatorname{tr}(X) = \operatorname{tr}(\alpha X) \text{ (Lemma 2.2.1.(2))} = \sum_{i=1}^3 \xi_i,$ $(X, X) = (\alpha X, \alpha X) \text{ (Lemma 2.2.4)} = \sum_{i=1}^3 \xi_i^2,$ $\operatorname{tr}(X, X, X) = \operatorname{tr}(\alpha X, \alpha X, \alpha X) \text{ (Lemma 2.2.4)} = \sum_{i=1}^3 \xi_i^3.$
0902.0431_E00602	100	0.396 K +0	$\begin{aligned} &\left. \left. \begin{array}{l} \left(\mathfrak{J}^C \right)_{\tau \gamma} = \left\{ X \in \mathfrak{J}^C \mid \tau \gamma X = X \right\} \right. \right. \\ &\quad = \left\{ \left(\begin{array}{ccc} \mu_1 & m_3 & \bar{m}_2 \\ \bar{m}_3 & \mu_2 & m_1 \\ m_2 & \bar{m}_1 & \mu_3 \end{array} \right) + i \left(\begin{array}{ccc} 0 & a_3 e_4 & -a_2 e_4 \\ -a_3 e_4 & 0 & a_1 e_4 \\ a_2 e_4 & -a_1 e_4 & 0 \end{array} \right) \right\} \left \begin{array}{l} \mu_i \in \mathbf{R} \\ m_i, a_i \in \mathbf{H} \end{array} \right\} \end{aligned}$	$(\mathfrak{J}^C)_{\tau \gamma} = \{X \in \mathfrak{J}^C \mid \tau \gamma X = X\}$ $= \left\{ \left(\begin{array}{ccc} \mu_1 & m_3 & \bar{m}_2 \\ \bar{m}_3 & \mu_2 & m_1 \\ m_2 & \bar{m}_1 & \mu_3 \end{array} \right) + i \left(\begin{array}{ccc} 0 & a_3 e_4 & -a_2 e_4 \\ -a_3 e_4 & 0 & a_1 e_4 \\ a_2 e_4 & -a_1 e_4 & 0 \end{array} \right) \right\} \left \begin{array}{l} \mu_i \in \mathbf{R} \\ m_i, a_i \in \mathbf{H} \end{array} \right\}$	$(\mathfrak{J}^C)_{\tau \gamma} = \{X \in \mathfrak{J}^C \mid \tau \gamma X = X\}$ $= \left\{ \left(\begin{array}{ccc} \mu_1 & m_3 & \bar{m}_2 \\ \bar{m}_3 & \mu_2 & m_1 \\ m_2 & \bar{m}_1 & \mu_3 \end{array} \right) + i \left(\begin{array}{ccc} 0 & a_3 e_4 & -a_2 e_4 \\ -a_3 e_4 & 0 & a_1 e_4 \\ a_2 e_4 & -a_1 e_4 & 0 \end{array} \right) \right\} \left \begin{array}{l} \mu_i \in \mathbf{R} \\ m_i, a_i \in \mathbf{H} \end{array} \right\}$
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5–0.9 ■ < 0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_EQ0900	149	■0.397 ●C +9	$\begin{aligned} P = & (X, Y, \xi, \eta) \\ \xrightarrow{X} & k\left(gX - \frac{\xi}{2}E\right)J + e_1k\left(g(\gamma Y) - \frac{\eta}{2}E\right)J \\ \longrightarrow & e_1k(T)k\left(gX - \frac{\xi}{2}E\right)J - k(T)k\left(g(\gamma Y) - \frac{\eta}{2}E\right)J \\ & + e_1k\left(gX - \frac{\xi}{2}E\right)J^tk(T) - k\left(g(\gamma Y) - \frac{\eta}{2}E\right)J^tk(T) \\ = & k(-Tg(\gamma Y) - g(\gamma Y)T + \eta T)J + e_1k(T(gX) + (gX)T - \xi T)J \\ = & k(-2gA \circ g(\gamma Y) + \eta gA)J + e_1k(2gA \circ gX - \xi gA)J \\ = & k\left(-2g(\gamma A \times Y) - \frac{1}{2}(A, Y)E + \eta gA\right)J \\ & + e_1k\left(2g(\gamma A \times \gamma X) + \frac{1}{2}(\gamma A, X)E - \xi gA\right)J \text{ (Lemma 3.12.1)} \\ = & \chi\left(\begin{array}{c} -2\gamma A \times Y + \eta A \\ 2A \times X - \xi \gamma A \\ (A, Y) \\ (-\gamma A, X) \end{array}\right) = \chi\left(\left(\begin{array}{cccc} 0 & -2\gamma A & 0 & A \\ 2A & 0 & -\gamma A & 0 \\ 0 & A & 0 & 0 \\ -\gamma A & 0 & 0 & 0 \end{array}\right)\left(\begin{array}{c} X \\ Y \\ \xi \\ \eta \end{array}\right)\right) \\ = & \chi(\Phi(0, A, -\gamma A, 0)P). \end{aligned}$	$\begin{aligned} P = & (X, Y, \xi, \eta) \\ \xrightarrow{X} & k\left(gX - \frac{\xi}{2}E\right)J + e_1k\left(g(\gamma Y) - \frac{\eta}{2}E\right)J \\ \longrightarrow & e_1k(T)k\left(gX - \frac{\xi}{2}E\right)J - k(T)k\left(g(\gamma Y) - \frac{\eta}{2}E\right)J \\ & + e_1k\left(gX - \frac{\xi}{2}E\right)J^tk(T) - k\left(g(\gamma Y) - \frac{\eta}{2}E\right)J^tk(T) \\ = & k(-Tg(\gamma Y) - g(\gamma Y)T + \eta T)J + e_1k(T(gX) + (gX)T - \xi T)J \\ = & k(-2gA \circ g(\gamma Y) + \eta gA)J + e_1k(2gA \circ gX - \xi gA)J \\ = & k\left(-2g(\gamma A \times Y) - \frac{1}{2}(A, Y)E + \eta gA\right)J \\ & + e_1k\left(2g(\gamma A \times \gamma X) + \frac{1}{2}(\gamma A, X)E - \xi gA\right)J \text{ (Lemma 3.12.1)} \\ = & \chi\left(\begin{array}{c} -2\gamma A \times Y + \eta A \\ 2A \times X - \xi \gamma A \\ (A, Y) \end{array}\right) = \chi\left(\left(\begin{array}{cccc} 0 & -2\gamma A & 0 & A \\ 2A & 0 & -\gamma A & 0 \\ 0 & A & 0 & 0 \\ -\gamma A & 0 & 0 & 0 \end{array}\right)\left(\begin{array}{c} X \\ Y \\ \xi \\ \eta \end{array}\right)\right) \\ = & \chi(\Phi(0, A, -\gamma A, 0)P) \end{aligned}$	$\begin{aligned} P = & (X, Y, \xi, \eta) \\ \xrightarrow{X} & k\left(gX - \frac{\xi}{2}E\right)J + e_1k\left(g(\gamma Y) - \frac{\eta}{2}E\right)J \\ \longrightarrow & e_1k(T)k\left(gX - \frac{\xi}{2}E\right)J - k(T)k\left(g(\gamma Y) - \frac{\eta}{2}E\right)J \\ & + e_1k\left(gX - \frac{\xi}{2}E\right)J^tk(T) - k\left(g(\gamma Y) - \frac{\eta}{2}E\right)J^tk(T) \\ = & k(-Tg(\gamma Y) - g(\gamma Y)T + \eta T)J + e_1k(T(gX) + (gX)T - \xi T)J \\ = & k(-2gA \circ g(\gamma Y) + \eta gA)J + e_1k(2gA \circ gX - \xi gA)J \\ = & k\left(-2g(\gamma A \times Y) - \frac{1}{2}(A, Y)E + \eta gA\right)J \\ & + e_1k\left(2g(\gamma A \times \gamma X) + \frac{1}{2}(\gamma A, X)E - \xi gA\right)J \text{ (Lemma 3.12.1)} \\ = & \chi\left(\begin{array}{c} -2\gamma A \times Y + \eta A \\ 2A \times X - \xi \gamma A \\ (A, Y) \\ (-\gamma A, X) \end{array}\right) = \chi\left(\left(\begin{array}{cccc} 0 & -2\gamma A & 0 & A \\ 2A & 0 & -\gamma A & 0 \\ 0 & A & 0 & 0 \\ -\gamma A & 0 & 0 & 0 \end{array}\right)\left(\begin{array}{c} X \\ Y \\ \xi \\ \eta \end{array}\right)\right) \\ = & \chi(\Phi(0, A, -\gamma A, 0)P). \end{aligned}$
0902.0431_EQ1000	183	■0.403 ●N +0	$S_{\{1\}} \tau S_{\{2\}} - S_{\{2\}} \tau S_{\{1\}} - \frac{1}{8} (S_{\{1\}} \tau S_{\{2\}} - S_{\{2\}} \tau S_{\{1\}}) \in \mathfrak{su}(8),$	$S_1 \tau S_2 - S_2 \tau S_1 - \frac{1}{8} (S_1 \tau S_2 - S_2 \tau S_1) \in \mathfrak{su}(8),$	$S_1 \tau S_2 - S_2 \tau S_1 - \frac{1}{8} (S_1 \tau S_2 - S_2 \tau S_1) \in \mathfrak{su}(8),$
Conf. MathPix confidence: ■ ■ ■ — ■≥0.9 ■0.5–0.9 ■<0.5 —no value Residual render vs scan (inkdrill): ● ● ● ● ● ● ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_E01030	173	0.410 W -1	$\begin{aligned} & \begin{array}{l} \lambda_0 + \frac{1}{2}\mu_1 - \frac{2}{3}\nu = 2 \quad 4 \quad 5 \quad 4 \quad 3 \quad 2 \\ \lambda_0 - \frac{1}{2}\mu_1 + \frac{2}{3}\nu = 2 \quad 3 \quad 5 \quad 4 \\ \lambda_1 + \frac{1}{2}\mu_1 - \frac{2}{3}\nu = 0 \quad 1 \quad 1 \quad 1 \quad 2 \\ \lambda_1 - \frac{1}{2}\mu_1 + \frac{2}{3}\nu = 0 \quad 0 \quad 1 \quad 1 \quad 1 \quad 1 \quad 1 \end{array} \\ & \begin{array}{l} \begin{$		

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_E00558	094	■0.435 ●C +5	$\begin{array}{l} \rightarrow \text{Spin}(9) \rightarrow (E_6)_{E_1} \rightarrow S^9 \rightarrow * \\ \downarrow p' \quad \downarrow p \quad \downarrow = \\ \rightarrow SO(9) \rightarrow SO(10) \rightarrow S^9 \rightarrow * \end{array}$	$\begin{array}{l} \rightarrow \text{Spin}(9) \rightarrow (E_6)_{E_1} \rightarrow S^9 \rightarrow * \\ \downarrow p' \quad \downarrow p \quad \downarrow = \\ \rightarrow SO(9) \rightarrow SO(10) \rightarrow S^9 \rightarrow * \end{array}$	$\begin{array}{ccccccc} 1 & \longrightarrow & Spin(9) & \longrightarrow & (E_6)_{E_1} & \longrightarrow & S^9 \longrightarrow * \\ & & \downarrow p' & & \downarrow p & & \downarrow = \\ 1 & \longrightarrow & SO(9) & \longrightarrow & SO(10) & \longrightarrow & S^9 \longrightarrow * \end{array}$
0902.0431_E00332	053	■0.438 ●W +0	$\begin{aligned} & \pm(\lambda_k - \lambda_l), \pm(\lambda_k + \lambda_l), \quad 0 \leq k < l \leq 3, \\ & \pm\lambda_0, \pm\lambda_1, \pm\lambda_2, \pm\lambda_3, \\ & \pm\frac{1}{2}(-\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3), \pm\frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3), \\ & \pm\frac{1}{2}(-\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3), \pm\frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3), \\ & \pm\frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3), \pm\frac{1}{2}(-\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3), \\ & \pm\frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3), \pm\frac{1}{2}(-\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3). \end{aligned}$	$\begin{aligned} & \pm(\lambda_k - \lambda_l), \pm(\lambda_k + \lambda_l), \quad 0 \leq k < l \leq 3, \\ & \pm\lambda_0, \pm\lambda_1, \pm\lambda_2, \pm\lambda_3, \\ & \pm\frac{1}{2}(-\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3), \pm\frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3), \\ & \pm\frac{1}{2}(-\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3), \pm\frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3), \\ & \pm\frac{1}{2}(\lambda_0 - \lambda_1 + \lambda_2 - \lambda_3), \pm\frac{1}{2}(-\lambda_0 + \lambda_1 + \lambda_2 - \lambda_3), \\ & \pm\frac{1}{2}(\lambda_0 + \lambda_1 + \lambda_2 + \lambda_3), \pm\frac{1}{2}(-\lambda_0 - \lambda_1 + \lambda_2 + \lambda_3). \end{aligned}$	
Conf. MathPix confidence: ■ ≥0.9 ■ 0.5-0.9 ■ <0.5 — no value Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_E01211	203	0.448 C +195	$\begin{aligned} & \begin{aligned} & \exp(\operatorname{ad}(Z, 0))(C, D, \boldsymbol{x} \otimes \boldsymbol{a}, \boldsymbol{b} \otimes \boldsymbol{y}, \boldsymbol{c} \otimes \boldsymbol{z}, \boldsymbol{w} \otimes \boldsymbol{d}) \\ &= (\exp(\operatorname{ad}(Z))C, D, ((\exp Z)\boldsymbol{x}) \otimes \boldsymbol{a}, \\ & \quad ((\exp Z)\boldsymbol{b}) \otimes \boldsymbol{y}, ((\exp(-^t Z))\boldsymbol{c} \otimes \boldsymbol{z}, ((\exp(-^t Z))\boldsymbol{w}) \otimes \boldsymbol{d}) \\ &= (\operatorname{Ad}(\exp Z)C, D, ((\exp Z)\boldsymbol{x}) \otimes \boldsymbol{a}, \\ & \quad ((\exp Z)\boldsymbol{b}) \otimes \boldsymbol{y}, (^t(\exp Z)^{-1}\boldsymbol{c}) \otimes \boldsymbol{z}, (^t(\exp Z)^{-1}\boldsymbol{w}) \otimes \boldsymbol{d}) \\ &= \varphi_1(\exp Z)(C, D, \boldsymbol{x} \otimes \boldsymbol{a}, \boldsymbol{b} \otimes \boldsymbol{y}, \boldsymbol{c} \otimes \boldsymbol{z}, \boldsymbol{w} \otimes \boldsymbol{d}). \end{aligned} \end{aligned}$	$\begin{aligned} & \exp(\operatorname{ad}(Z, 0))(C, D, \boldsymbol{x} \otimes \boldsymbol{a}, \boldsymbol{b} \otimes \boldsymbol{y}, \boldsymbol{c} \otimes \boldsymbol{z}, \boldsymbol{w} \otimes \boldsymbol{d}) \\ &= (\exp(\operatorname{ad}(Z))C, D, ((\exp Z)\boldsymbol{x}) \otimes \boldsymbol{a}, \\ & \quad ((\exp Z)\boldsymbol{b}) \otimes \boldsymbol{y}, ((\exp(-^t Z))\boldsymbol{c} \otimes \boldsymbol{z}, ((\exp(-^t Z))\boldsymbol{w}) \otimes \boldsymbol{d}) \\ &= (\operatorname{Ad}(\exp Z)C, D, ((\exp Z)\boldsymbol{x}) \otimes \boldsymbol{a}, \\ & \quad ((\exp Z)\boldsymbol{b}) \otimes \boldsymbol{y}, (^t(\exp Z)^{-1}\boldsymbol{c}) \otimes \boldsymbol{z}, (^t(\exp Z)^{-1}\boldsymbol{w}) \otimes \boldsymbol{d}) \\ &= \varphi_1(\exp Z)(C, D, \boldsymbol{x} \otimes \boldsymbol{a}, \boldsymbol{b} \otimes \boldsymbol{y}, \boldsymbol{c} \otimes \boldsymbol{z}, \boldsymbol{w} \otimes \boldsymbol{d}). \end{aligned}$	$\begin{aligned} & \exp(\operatorname{ad}(Z, 0))(C, D, \boldsymbol{x} \otimes \boldsymbol{a}, \boldsymbol{b} \otimes \boldsymbol{y}, \boldsymbol{c} \otimes \boldsymbol{z}, \boldsymbol{w} \otimes \boldsymbol{d}) \\ &= (\exp(\operatorname{ad}(Z))C, D, ((\exp Z)\boldsymbol{x}) \otimes \boldsymbol{a}, \\ & \quad ((\exp Z)\boldsymbol{b}) \otimes \boldsymbol{y}, ((\exp(-^t Z))\boldsymbol{c} \otimes \boldsymbol{z}, ((\exp(-^t Z))\boldsymbol{w}) \otimes \boldsymbol{d}) \\ &= (\operatorname{Ad}(\exp Z)C, D, ((\exp Z)\boldsymbol{x}) \otimes \boldsymbol{a}, \\ & \quad ((\exp Z)\boldsymbol{b}) \otimes \boldsymbol{y}, (^t(\exp Z)^{-1}\boldsymbol{c}) \otimes \boldsymbol{z}, (^t(\exp Z)^{-1}\boldsymbol{w}) \otimes \boldsymbol{d}) \\ &= \varphi_1(\exp Z)(C, D, \boldsymbol{x} \otimes \boldsymbol{a}, \boldsymbol{b} \otimes \boldsymbol{y}, \boldsymbol{c} \otimes \boldsymbol{z}, \boldsymbol{w} \otimes \boldsymbol{d}). \end{aligned}$
0902.0431_E01039	177	0.452 W +2	$\begin{aligned} & \begin{aligned} & H_{\alpha_3} = \left(0, 0, 0, \frac{1}{60}, 0, 0\right), \\ & H_{\alpha_4} = \left(\Phi\left(\frac{1}{90}(-E_1 + 2E_2 - E_3)^\sim, 0, 0, -\frac{1}{120}\right), 0, 0, -\frac{1}{120}, 0, 0\right), \\ & H_{\alpha_5} = \left(\Phi\left(\frac{1}{60}(H_3 - (E_2 - E_3)^\sim), 0, 0, 0\right), 0, 0, 0, 0, 0\right), \\ & H_{\alpha_6} = \left(\Phi\left(\frac{1}{60}(H_2 - H_3), 0, 0, 0\right), 0, 0, 0, 0, 0\right), \\ & H_{\alpha_7} = \left(\Phi\left(\frac{1}{60}(H_1 - H_2), 0, 0, 0\right), 0, 0, 0, 0, 0\right), \\ & H_{\alpha_8} = \left(\Phi\left(0, 0, 0, \frac{1}{40}\right), 0, 0, -\frac{1}{120}, 0, 0\right). \end{aligned} \end{aligned}$	$\begin{aligned} & H_{\alpha_3} = \left(0, 0, 0, \frac{1}{60}, 0, 0\right), \\ & H_{\alpha_4} = \left(\Phi\left(\frac{1}{90}(-E_1 + 2E_2 - E_3)^\sim, 0, 0, -\frac{1}{120}\right), 0, 0, -\frac{1}{120}, 0, 0\right), \\ & H_{\alpha_5} = \left(\Phi\left(\frac{1}{60}(H_3 - (E_2 - E_3)^\sim), 0, 0, 0\right), 0, 0, 0, 0, 0\right), \\ & H_{\alpha_6} = \left(\Phi\left(\frac{1}{60}(H_2 - H_3), 0, 0, 0\right), 0, 0, 0, 0, 0\right), \\ & H_{\alpha_7} = \left(\Phi\left(\frac{1}{60}(H_1 - H_2), 0, 0, 0\right), 0, 0, 0, 0, 0\right), \\ & H_{\alpha_8} = \left(\Phi\left(0, 0, 0, \frac{1}{40}\right), 0, 0, -\frac{1}{120}, 0, 0\right). \end{aligned}$	$\begin{aligned} & H_{\alpha_3} = \left(0, 0, 0, \frac{1}{60}, 0, 0\right), \\ & H_{\alpha_4} = \left(\Phi\left(\frac{1}{90}(-E_1 + 2E_2 - E_3)^\sim, 0, 0, -\frac{1}{120}\right), 0, 0, -\frac{1}{120}, 0, 0\right), \\ & H_{\alpha_5} = \left(\Phi\left(\frac{1}{60}(H_3 - (E_2 - E_3)^\sim), 0, 0, 0\right), 0, 0, 0, 0, 0\right), \\ & H_{\alpha_6} = \left(\Phi\left(\frac{1}{60}(H_2 - H_3), 0, 0, 0\right), 0, 0, 0, 0, 0\right), \\ & H_{\alpha_7} = \left(\Phi\left(\frac{1}{60}(H_1 - H_2), 0, 0, 0\right), 0, 0, 0, 0, 0\right), \\ & H_{\alpha_8} = \left(\Phi\left(0, 0, 0, \frac{1}{40}\right), 0, 0, -\frac{1}{120}, 0, 0\right). \end{aligned}$
Conf. MathPix confidence: ≥0.9 0.5-0.9 <0.5 — no value Residual render vs scan (inkdrill): • C component • W weak • S stable • N noise • K clean • not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_E00939	156	■0.471 ●W +0	$\left.\begin{array}{l} \xrightarrow{\hspace{1cm}} \\ \longrightarrow \end{array}\right\} \left(\begin{array}{ccc} h\left(b_{11}-c_{11}-\frac{\nu}{3}\right) & -h\left(c_{12}, b_{12}\right) & -h\left(c_{13}, b_{13}\right) \\ -h\left(\bar{b}_{12}, \bar{c}_{12}\right) & 0 & 0 \\ -h\left(\bar{b}_{13}, \bar{c}_{13}\right) & 0 & 0 \end{array}\right) \\ \begin{array}{ccc} 0 & 0 & 0 \\ 0 & h\left(l_{33}\right) & -h\left(l_{23}, \bar{l}_{32}\right) \\ 0 & -h\left(l_{32}, \bar{l}_{23}\right) & h\left(l_{22}\right) \end{array} \\ 0 \\ -h\left(\bar{l}_{11}\right) \end{array} \\ =\left(\begin{array}{c} \phi_C(B, C) E_1+\frac{i v e_1}{3} E_1 \\ 2 h(L) \times E_1 \\ 0 \\ -(\tau h(L), E_1) \end{array}\right)=\Phi\left(\phi_C(B, C), h(L),-\tau h(L),-i v e_1\right)\left(\begin{array}{c} E_1 \\ 0 \\ 0 \\ 0 \end{array}\right) .$	(not rendered)	
0902.0431_E01196	201	■0.477 ●K +0	$\begin{aligned} \mathfrak{g}_{01} &= \{(C, 0) \in \mathfrak{g}_0 \mid C \in \mathfrak{sl}(5, C)\} \cong \mathfrak{sl}(5, C) \\ \mathfrak{g}_{02} &= \{(0, D) \in \mathfrak{g}_0 \mid D \in \mathfrak{sl}(5, C)\} \cong \mathfrak{sl}(5, C) \\ \mathfrak{q} &= \mathfrak{g}_1 \oplus \mathfrak{g}_2 \oplus \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} . \end{aligned}$	$\begin{aligned} \mathfrak{g}_{01} &= \{(C, 0) \in \mathfrak{g}_0 \mid C \in \mathfrak{sl}(5, C)\} \cong \mathfrak{sl}(5, C) \\ \mathfrak{g}_{02} &= \{(0, D) \in \mathfrak{g}_0 \mid D \in \mathfrak{sl}(5, C)\} \cong \mathfrak{sl}(5, C) \\ \mathfrak{q} &= \mathfrak{g}_1 \oplus \mathfrak{g}_2 \oplus \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} . \end{aligned}$	$\begin{aligned} \mathfrak{g}_{01} &= \{(C, 0) \in \mathfrak{g}_0 \mid C \in \mathfrak{sl}(5, C)\} \cong \mathfrak{sl}(5, C), \\ \mathfrak{g}_{02} &= \{(0, D) \in \mathfrak{g}_0 \mid D \in \mathfrak{sl}(5, C)\} \cong \mathfrak{sl}(5, C), \\ \mathfrak{q} &= \mathfrak{g}_1 \oplus \mathfrak{g}_2 \oplus \mathfrak{g}_{-2} \oplus \mathfrak{g}_{-1} . \end{aligned}$
0902.0431_E01195	201	■0.485 ●N +0	$\begin{aligned} & \left(- * (y \wedge * (x \wedge a)), b \right) = - (x \wedge a, y \wedge b) \\ & = (y, a_1) (x \wedge a_2, b) - (y, a_2) (x \wedge a_1, b) - (x, y) (a, b) . \end{aligned}$	$\begin{aligned} & \left(- * (y \wedge * (x \wedge a)), b \right) = - (x \wedge a, y \wedge b) \\ & = (y, a_1) (x \wedge a_2, b) - (y, a_2) (x \wedge a_1, b) - (x, y) (a, b) . \end{aligned}$	$\begin{aligned} & \left(- * (\boldsymbol{y} \wedge * (\boldsymbol{x} \wedge \boldsymbol{a})), \boldsymbol{b} \right) = - (\boldsymbol{x} \wedge \boldsymbol{a}, \boldsymbol{y} \wedge \boldsymbol{b}) \\ & = (\boldsymbol{y}, \boldsymbol{a}_1) (\boldsymbol{x} \wedge \boldsymbol{a}_2, \boldsymbol{b}) - (\boldsymbol{y}, \boldsymbol{a}_2) (\boldsymbol{x} \wedge \boldsymbol{a}_1, \boldsymbol{b}) - (\boldsymbol{x}, \boldsymbol{y}) (\boldsymbol{a}, \boldsymbol{b}) . \end{aligned}$
0902.0431_E01104	187	■0.490 ●W +0	$\varphi(R) R_1 = [R, R_1], \quad R \in (\mathfrak{e}_{8(8)})^{\tilde{\lambda}_\gamma}, R_1 \in (\mathfrak{e}_{8(8)})_{-\tilde{\lambda}_\gamma} .$	$\varphi(R) R_1 = [R, R_1], \quad R \in (\mathfrak{e}_{8(8)})^{\tilde{\lambda}_\gamma}, R_1 \in (\mathfrak{e}_{8(8)})_{-\tilde{\lambda}_\gamma} .$	
0902.0431_E01010	168	■0.494 ●	$\begin{aligned} \mathfrak{e}_8 &= \left\{ R \in \mathfrak{e}_8^C \mid \tau \tilde{\lambda} R = R \right\} \\ &= \left\{ (\Phi, P, -\tau \lambda P, r, s, -\tau s) \in \mathfrak{e}_8^C \mid \Phi \in \mathfrak{e}_7, P \in \mathfrak{P}^C, r \in i \boldsymbol{R}, s \in C \right\} . \end{aligned}$	$\begin{aligned} \mathfrak{e}_8 &= \left\{ R \in \mathfrak{e}_8^C \mid \tau \tilde{\lambda} R = R \right\} \\ &= \left\{ (\Phi, P, -\tau \lambda P, r, s, -\tau s) \in \mathfrak{e}_8^C \mid \Phi \in \mathfrak{e}_7, P \in \mathfrak{P}^C, r \in i \boldsymbol{R}, s \in C \right\} . \end{aligned}$	$\begin{aligned} \mathfrak{e}_8 &= \{ R \in \mathfrak{e}_8^C \mid \tau \tilde{\lambda} R = R \} \\ &= \{ (\Phi, P, -\tau \lambda P, r, s, -\tau s) \in \mathfrak{e}_8^C \mid \Phi \in \mathfrak{e}_7, P \in \mathfrak{P}^C, r \in i \boldsymbol{R}, s \in C \} . \end{aligned}$
Conf. MathPix confidence: ■≥0.9 ■0.5–0.9 ■<0.5 — no value Residual render vs scan (inkdrill): ●C component ●W weak ●S stable ●N noise ●K clean ●not measured					

Identifier	Page	Conf.	LaTeX source	Rendered	Scan image
0902.0431_E00952	158	0.496	$\begin{aligned} D &= \left(\begin{array}{cc} B & L \\ -L^* & C \end{array} \right) + \frac{\nu}{3} \left(\begin{array}{cc} E & 0 \\ 0 & -E \end{array} \right) \in \mathfrak{su}(6), \\ \widetilde{M} &= \left(\begin{array}{cc} -N_2 - N_1 e_1 & M_2 + M_1 e_1 \\ M_2 - M_1 e_1 & N_2 - N_1 e_1 \end{array} \right), \quad M_i, N_i \in M(3, \boldsymbol{C}), \\ M &= M_1 + iM_2, N = N_1 + iN_2. \end{aligned}$	$D = \left(\begin{array}{cc} B & L \\ -L^* & C \end{array} \right) + \frac{\nu}{3} \left(\begin{array}{cc} E & 0 \\ 0 & -E \end{array} \right) \in \mathfrak{su}(6),$ $\widetilde{M} = \left(\begin{array}{cc} -N_2 - N_1 e_1 & M_2 + M_1 e_1 \\ M_2 - M_1 e_1 & N_2 - N_1 e_1 \end{array} \right), \quad M_i, N_i \in M(3, \boldsymbol{C}),$ $M = M_1 + iM_2, N = N_1 + iN_2.$	$D = \left(\begin{array}{cc} B & L \\ -L^* & C \end{array} \right) + \frac{\nu}{3} \left(\begin{array}{cc} E & 0 \\ 0 & -E \end{array} \right) \in \mathfrak{su}(6),$ $\widetilde{M} = \left(\begin{array}{cc} -N_2 - N_1 e_1 & M_2 + M_1 e_1 \\ M_2 - M_1 e_1 & N_2 - N_1 e_1 \end{array} \right), \quad M_i, N_i \in M(3, \boldsymbol{C}),$ $M = M_1 + iM_2, N = N_1 + iN_2.$
Conf. MathPix confidence: ≥0.9 0.5-0.9 <0.5 — no value Residual render vs scan (inkdrill): C component W weak S stable N noise K clean not measured					

Inline formulas (first occurrence)

Identifier	Page	Conf.	LaTeX source	Rendered
Conf. MathPix confidence: ■ ≥ 0.9 ■ 0.5-0.9 ■ < 0.5 — no value				
Residual render vs scan (inkdrill): ● C component ● W weak ● S stable ● N noise ● K clean ● not measured				