

cardona-qft-methods: equation evidence

Display equations

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E00717 [conf 0.003]	275	■ 0.003	$\begin{aligned} & \hat{T} = \operatorname{tr}_a M_a(u) \\ & = ^L_L \cdots \\ & \cdots \cdots \cdots a. \\ & \cdots 2 \\ & L \end{aligned}$	$\hat{T} = \operatorname{tr}_a M_a(u)$ $= ^L_L \cdots \cdots a.$	$\hat{T} = \operatorname{tr}_a M_a(u)$ $= \left(\begin{array}{cccc} L & \cdots & 2 & 1 \\ \diagdown & & \diagup & \diagup \\ L & \cdots & 2 & 1 \end{array} \right) a.$
cardona-qft-methods_ E00889 [conf 0.077]	345	■ 0.077	$\begin{array}{rcl} \operatorname{Map}(8, M) = L M \\ \times_M L M & \longrightarrow & L M \times L M \\ \downarrow & \Delta & \downarrow \\ \downarrow^{ev \times ev} M & \downarrow & \downarrow^{cM \times M} \end{array}$	$\operatorname{Map}(8, M) = L M \times_M L M \longrightarrow L M \times L M$ $\downarrow \Delta \downarrow$ $\downarrow^{ev \times ev} M \downarrow \downarrow^{cM \times M}$	$\operatorname{Map}(8, M) = L M \times_M L M \longrightarrow L M \times L M$ $\downarrow \Delta \downarrow^{ev \times ev} M \xrightarrow{\Delta} M \times M \downarrow^{ev \times ev}$
cardona-qft-methods_ E00908 [conf 0.080]	350	■ 0.080	$\begin{array}{rcl} P_g M : {}_1 \times {}_0 P_h M & \longrightarrow & P_g M \times P_h M \\ \downarrow & & \downarrow \\ \downarrow e_0 \circ \pi_1 M & & \downarrow \\ \downarrow & & \downarrow \times M \end{array}$	$P_g M : {}_1 \times {}_0 P_h M \longrightarrow P_g M \times P_h M$ $\downarrow \downarrow e_0 \circ \pi_1 M \downarrow \downarrow \downarrow \times M$	$P_g M : {}_1 \times {}_0 P_h M \longrightarrow P_g M \times P_h M$ $\downarrow e_0 \circ \pi_1 \downarrow e_0 \times e_0 M \xrightarrow{\Delta_g} M \times M$
cardona-qft-methods_ E00077	46	■ 0.150	$\left. \begin{array}{l} 1 \\ 1 \end{array} \right\} \longrightarrow \operatorname{Inn}(\mathcal{A}) \longrightarrow \operatorname{Aut}(\mathcal{A}) \longrightarrow \operatorname{Aut}(\mathcal{A}) / \operatorname{Inn}(\mathcal{A}) \longrightarrow 1,$ $\left. \begin{array}{l} 1 \\ 1 \end{array} \right\} \longrightarrow \mathcal{G} \longrightarrow \mathcal{G} \rtimes \operatorname{Diff}(M) \longrightarrow \operatorname{Diff}(M) \longrightarrow 1.$	$(not\ rendered)$	$1 \rightarrow \operatorname{Inn}(\mathcal{A}) \rightarrow \operatorname{Aut}(\mathcal{A}) \rightarrow \operatorname{Aut}(\mathcal{A}) / \operatorname{Inn}(\mathcal{A}) \rightarrow 1,$ $1 \rightarrow \mathcal{G} \rightarrow \mathcal{G} \rtimes \operatorname{Diff}(M) \rightarrow \operatorname{Diff}(M) \rightarrow 1.$
cardona-qft-methods_ E00716	275	■ 0.155	$\begin{aligned} & M_a(u) = L_{a,L}(u) L_{a,L-1}(u) \cdots L_{a,2}(u) L_{a,1}(u) \\ & = a ^L_L \cdots \\ & \cdots \cdots \cdots L \end{aligned}$	$M_a(u) = L_{a,L}(u) L_{a,L-1}(u) \cdots L_{a,2}(u) L_{a,1}(u)$ $= a ^L_L \cdots \cdots L$	$M_a(u) = L_{a,L}(u) L_{a,L-1}(u) \cdots L_{a,2}(u) L_{a,1}(u)$ $= a \frac{L \cdots 2 1}{L \cdots 2 1} a.$
cardona-qft-methods_ E00654 (4)	261	■ 0.167	$\phi_{\theta}(x) \phi_{\theta}(y) = e^{-\frac{1}{2} \frac{\partial}{\partial x^{\mu}} \theta^{\mu \nu} \frac{\partial}{\partial y^{\nu}} \phi_0(x) \phi_0(y)} e^{\frac{1}{2} \left(\frac{\partial}{\partial x^{\mu}} + \frac{\partial}{\partial y^{\mu}} \right) \theta^{\mu \nu} P_{\nu}} \neq \phi_0(x) \phi_0(y).$	$(not\ rendered)$	$\phi_{\theta}(x) \phi_{\theta}(y) = e^{-\frac{1}{2} \frac{\partial}{\partial x^{\mu}} \theta^{\mu \nu} \frac{\partial}{\partial y^{\nu}} \phi_0(x) \phi_0(y)} e^{\frac{1}{2} \left(\frac{\partial}{\partial x^{\mu}} + \frac{\partial}{\partial y^{\mu}} \right) \theta^{\mu \nu} P_{\nu}} \neq \phi_0(x) \phi_0(y).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0681	269	■ 0.304	\begin{gathered} " \text{ { Group" } } \approx e^{\text{i} \text{ { \text{ { "Algebra" } } } }}, \backslash S U(2) \approx e^{\text{i} \mathfrak{s} u(2)} \end{gathered}	" Group" $\approx e^i$ "Algebra" , $SU(2) \approx e^{isu(2)}$	"Group" $\approx e^i$ "Algebra" , $SU(2) \approx e^{isu(2)}$.
cardona-qft-methods_ EQ0158 (4)	87	■ 0.311	\mathrm{t}-\operatorname{Ind}=i_{!}^{-1}j_{!}:K\left(T^{*}M\right) \rightarrow K(\{p t\}) \simeq \mathbb{Z} .	$t - \text{Ind} = i_!^{-1} j_! : K(T^*M) \rightarrow K(\{pt\}) \simeq \mathbb{Z}.$	$t\text{-Ind} = i_!^{-1} j_! : K(T^*M) \rightarrow K(\{pt\}) \simeq \mathbb{Z}.$
cardona-qft-methods_ EQ0501	226	■ 0.333	\left.\sum_{\text{generations}}\left(m_{\nu}^2+m_e^2+3 m_u^2+3 m_d^2\right)\right _{\Lambda=\Lambda_{u n i f}}=\left.\left.8 M_W^2\right _{\Lambda=\Lambda_{u n i f}}\right _{\Lambda=\Lambda_{u n i f}}	$\sum_{\text{generations}}\left(m_{\nu}^2+m_e^2+3 m_u^2+3 m_d^2\right)\Bigg _{\Lambda=\Lambda_{unif}}=8 M_W^2\Big _{\Lambda=\Lambda_{unif}}$	$\sum_{generations}\left(m_{\nu}^2+m_e^2+3 m_u^2+3 m_d^2\right) _{\Lambda=\Lambda_{unif}}=8 M_W^2 _{\Lambda=\Lambda_{unif}}$
cardona-qft-methods_ EQ0049 (14)	35	■ 0.362	\left.\mathcal{H}=\mathcal{L}^2\left(\left(M, g\right), S\right)\right)=\left.\left\{\psi \in \Gamma^{\infty}(M, S) \mid \int_M\langle\psi, \psi\rangle_x d \operatorname{vol}_g(x)<\infty\right\}\right	$\mathcal{H}=L^2((M,g),S))=\left\{\psi\in\Gamma^\infty(M,S)\mid\int_M\langle\psi,\psi\rangle_xd\operatorname{vol}_g(x)<\infty\right\}$	$\mathcal{H}=L^2((M,g),S))=\left\{\psi\in\Gamma^\infty(M,S)\mid\int_M\langle\psi,\psi\rangle_xd\operatorname{vol}_g(x)<\infty\right\}$
cardona-qft-methods_ EQ0795 (36)	301	■ 0.396	\begin{aligned} & \operatorname{type} \text{ { IIB }} \operatorname{ { \mathrm{NS} }}-\operatorname{ { \mathrm{R} }}: \mathbf{8}_v \otimes \mathbf{8}_s=\mathbf{8}_s \oplus \mathbf{56}_s=\lambda^1 \oplus \Psi^{1 M}, \\ & \mathbf{8}_s \otimes \mathbf{8}_v=\mathbf{8}_s \oplus \mathbf{56}_s=\lambda^2 \oplus \Psi^{2 M}, \\ & \operatorname{type} \text{ { IIA }} \operatorname{ { \mathrm{NS} }}-\operatorname{ { \mathrm{R} }}: \mathbf{8}_v \otimes \mathbf{8}_c=\mathbf{8}_c \oplus \mathbf{56}_c=\lambda^1 \oplus \Psi^{1 M}, \\ & \mathbf{8}_s \otimes \mathbf{8}_v=\mathbf{8}_s \oplus \mathbf{56}_s=\lambda^2 \oplus \Psi^{2 M} . \end{aligned}	type IIB NS – R : $\mathbf{8}_v \otimes \mathbf{8}_s = \mathbf{8}_s \oplus \mathbf{56}_s = \lambda^1 \oplus \Psi^{1M}$, R – NS : $\mathbf{8}_s \otimes \mathbf{8}_v = \mathbf{8}_s \oplus \mathbf{56}_s = \lambda^2 \oplus \Psi^{2M}$, type IIA NS – R : $\mathbf{8}_v \otimes \mathbf{8}_c = \mathbf{8}_c \oplus \mathbf{56}_c = \lambda^1 \oplus \Psi^{1M}$, R – NS : $\mathbf{8}_s \otimes \mathbf{8}_v = \mathbf{8}_s \oplus \mathbf{56}_s = \lambda^2 \oplus \Psi^{2M}$.	type IIB NS – R : $\mathbf{8}_v \otimes \mathbf{8}_s = \mathbf{8}_s \oplus \mathbf{56}_s = \lambda^1 \oplus \Psi^{1M}$, R – NS : $\mathbf{8}_s \otimes \mathbf{8}_v = \mathbf{8}_s \oplus \mathbf{56}_s = \lambda^2 \oplus \Psi^{2M}$, type IIA NS – R : $\mathbf{8}_v \otimes \mathbf{8}_c = \mathbf{8}_c \oplus \mathbf{56}_c = \lambda^1 \oplus \Psi^{1M}$, R – NS : $\mathbf{8}_s \otimes \mathbf{8}_v = \mathbf{8}_s \oplus \mathbf{56}_s = \lambda^2 \oplus \Psi^{2M}$.
cardona-qft-methods_ EQ0028	27	■ 0.431	\begin{aligned} & W R e s \left((1+\Delta)^{-d / 2}\right) =\int_M d x \int_{\mathbb{S}^{d-1}}\left\ \xi\right\ _x^{-d} d \xi=\int_M d x \sqrt{\operatorname{det} g_x} \operatorname{Vol}\left(\mathbb{S}^{d-1}\right) \\ & =\operatorname{Vol}\left(\mathbb{S}^{d-1}\right) \int_M\left \operatorname{dvol}_g\right =\operatorname{Vol}\left(\mathbb{S}^{d-1}\right) \end{aligned}	$WRes\left(\left(1+\Delta\right)^{-d / 2}\right)=\int_M d x \int_{\mathbb{S}^{d-1}}\left\ \xi\right\ _x^{-d} d \xi=\int_M d x \sqrt{\operatorname{det} g_x} \operatorname{Vol}\left(\mathbb{S}^{d-1}\right)$ $=\operatorname{Vol}\left(\mathbb{S}^{d-1}\right) \int_M\left \operatorname{dvol}_g\right =\operatorname{Vol}\left(\mathbb{S}^{d-1}\right)$	$WRes\left(\left(1+\Delta\right)^{-d / 2}\right)=\int_M d x \int_{\mathbb{S}^{d-1}}\left\ \xi\right\ _x^{-d} d \xi=\int_M d x \sqrt{\operatorname{det} g_x} \operatorname{Vol}\left(\mathbb{S}^{d-1}\right)$ $=\operatorname{Vol}\left(\mathbb{S}^{d-1}\right) \int_M\left \operatorname{dvol}_g\right =\operatorname{Vol}\left(\mathbb{S}^{d-1}\right) . \quad \square$
cardona-qft-methods_ EQ0148 (1)	83	■ 0.450	\left.\mathcal{D}\right\rightrightarrows \sigma_L(\mathcal{D}) \rightsquigarrow \text { an element of } K\text {-theory } \rightsquigarrow \mathrm{t}-\operatorname{Ind}\left(\sigma_L(\mathcal{D})\right) \in \mathbb{Z} .	$\mathcal{D} \rightsquigarrow \sigma_L(\mathcal{D}) \rightsquigarrow \text{ an element of } K\text{-theory} \rightsquigarrow t - \text{Ind}(\sigma_L(\mathcal{D})) \in \mathbb{Z}.$	$\mathcal{D} \rightsquigarrow \sigma_L(\mathcal{D}) \rightsquigarrow \text{ an element of } K\text{-theory} \rightsquigarrow t\text{-Ind}(\sigma_L(\mathcal{D})) \in \mathbb{Z}.$
cardona-qft-methods_ EQ0515 (1)	236	■ 0.478	\lambda_C=\frac{\hbar}{M c} \leq L \text { or } M \geq \frac{\hbar}{L c} \simeq 10^{19} \mathrm{GeV} \text { (Planck mass) } \simeq 10^{-5} \text { grams } .	$\lambda_C=\frac{\hbar}{Mc} \leq L \text{ or } M \geq \frac{\hbar}{Lc} \simeq 10^{19}\text{GeV (Planck mass)} \simeq 10^{-5} \text{ grams} .$	$\lambda_C=\frac{\hbar}{Mc} \leq L \text{ or } M \geq \frac{\hbar}{Lc} \simeq 10^{19}\text{GeV (Planck mass)} \simeq 10^{-5}\text{grams}.$
cardona-qft-methods_ EQ0019 (7)	24	■ 0.490	W \operatorname{Res} P=\operatorname{Res} \operatorname{Tr}\left(P \mid \mathcal{D} \mid^{-s}\right) .	$W \operatorname{Res} P=\operatorname{Res} \operatorname{Tr}_{s=0}\left(P \mid \mathcal{D} \mid^{-s}\right) .$	$WResP=\operatorname{Res} \operatorname{Tr}_{s=0}\left(P \mid \mathcal{D} \mid^{-s}\right) .$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0324 (10)	161	■ 0.495	$\begin{aligned} \phi_{\parallel}(x) - \phi^a(x) &= d\phi^a(x) + \omega_b^a(x)\phi^b(x) = D\phi^a(x) \\ \phi^a(x) + \omega_b^a(x)\phi^b(x) &= D\phi^a(x) \\ \omega^{\prime a}(x) &= \Lambda_c^a(x)\Lambda_b^d(x)\omega_d^c(x) + \Lambda_c^a(x)d\Lambda_b^c(x), \\ \Lambda_b^d(x)\omega_d^c(x) + \Lambda_c^a(x)d\Lambda_b^c(x) &= D\Lambda_b^c(x) \end{aligned}$	$\phi_{\parallel}^a(x) - \phi^a(x) = d\phi^a(x) + \omega_b^a(x)\phi^b(x) = D\phi^a(x)$ $\omega^{\prime a}(x) = \Lambda_c^a(x)\Lambda_b^d(x)\omega_d^c(x) + \Lambda_c^a(x)d\Lambda_b^c(x)$	$\phi_{\parallel}^a(x) - \phi^a(x) = d\phi^a(x) + \omega_b^a(x)\phi^b(x) = D\phi^a(x)$ $\omega^{\prime a}_b(x) = \Lambda_c^a(x)\Lambda_b^d(x)\omega_d^c(x) + \Lambda_c^a(x)d\Lambda_b^c(x),$
cardona-qft-methods_ EQ0481	219	■ 0.498	$Pf(\frac{A}{\xi}) = \int e^{-\frac{1}{2}\mathfrak{A}(\tilde{\xi})} D[\tilde{\xi}],$	$Pf(\mathfrak{A}) = \int e^{-\frac{1}{2}\mathfrak{A}(\tilde{\xi})} D[\tilde{\xi}],$	$Pf(\mathfrak{A}) = \int e^{-\frac{1}{2}\mathfrak{A}(\tilde{\xi})} D[\tilde{\xi}],$
cardona-qft-methods_ EQ1001	375	■ 0.508	$\begin{aligned} X(a \circ b) &= (a \circ b) \circ (X^\dagger(\cdot) \otimes \text{Id} + \text{Id} \otimes X^\dagger(\cdot)) \circ \Delta \\ &= (Xa \otimes b + a \otimes Xb) \circ \Delta = (Xa)b + a(Xb). \end{aligned}$	$X(ab) = (a \otimes b) \circ (X^\dagger(\cdot) \otimes \text{Id} + \text{Id} \otimes X^\dagger(\cdot)) \circ \Delta$ $= (Xa \otimes b + a \otimes Xb) \circ \Delta = (Xa)b + a(Xb).$	$X(ab) = (a \otimes b) \circ (X^\dagger(\cdot) \otimes \text{Id} + \text{Id} \otimes X^\dagger(\cdot)) \circ \Delta$ $= (Xa \otimes b + a \otimes Xb) \circ \Delta = (Xa)b + a(Xb).$
cardona-qft-methods_ EQ0712	274	■ 0.516	$R_{\{a, b\}}(u) = u1_{\{a, b\}} + iP_{\{a, b\}} = \begin{pmatrix} u+i & 0 & 0 & 0 \\ 0 & u & i & 0 \\ 0 & i & u & 0 \\ 0 & 0 & 0 & u+i \end{pmatrix},$	$R_{a,b}(u) = u1_{a,b} + iP_{a,b} = \begin{pmatrix} u+i & 0 & 0 & 0 \\ 0 & u & i & 0 \\ 0 & i & u & 0 \\ 0 & 0 & 0 & u+i \end{pmatrix},$	$R_{a,b}(u) = u1_{a,b} + iP_{a,b} = \begin{pmatrix} u+i & 0 & 0 & 0 \\ 0 & u & i & 0 \\ 0 & i & u & 0 \\ 0 & 0 & 0 & u+i \end{pmatrix},$
cardona-qft-methods_ EQ0890	345	■ 0.517	$\begin{aligned} &H_*(L M) \otimes H_*(L M) \rightarrow H_*(L M \times L M) \\ &\xrightarrow{\tau_{\Delta*}} H_*(\text{Map}(8, M)^{ev^*(TM)}) \xrightarrow{\tau_{\Delta*}} H_{*-n}(\text{Map}(8, M)) \\ &\downarrow \\ &H_{*-n}(L M) \end{aligned}$	$H_*(LM) \otimes H_*(LM) \rightarrow H_*(LM \times LM)$ $H_*\left(\text{Map}(8, M)^{ev^*(TM)}\right) \xrightarrow{\tau_{\Delta*}} H_{*-n}(\text{Map}(8, M))$ $\downarrow$ $H_{*-n}(LM)$	$H_*(LM) \otimes H_*(LM) \rightarrow H_*(LM \times LM)$ $\begin{array}{c} \tau_{\Delta*} \downarrow \\ H_*(\text{Map}(8, M)^{ev^*(TM)}) \xrightarrow{Thom} H_{*-n}(\text{Map}(8, M)) \\ \downarrow \\ H_{*-n}(LM) \end{array}$
cardona-qft-methods_ EQ0012	20	■ 0.522	$\sigma_m^P(x, \xi) = \lim_{t \rightarrow \infty} t^{-m} e^{-i \text{th}(x)} P e^{i \text{th}(x)},$	$\sigma_m^P(x, \xi) = \lim_{t \rightarrow \infty} t^{-m} e^{-i \text{th}(x)} P e^{i \text{th}(x)},$	$\sigma_m^P(x, \xi) = \lim_{t \rightarrow \infty} t^{-m} e^{-i \text{th}(x)} P e^{i \text{th}(x)},$
cardona-qft-methods_ EQ0201	96	■ 0.522	$\langle d\alpha, \beta \rangle = \langle \alpha, d^*\beta \rangle, \quad \text{for all } \alpha, \beta \in \Omega^*(M, E).$	$\langle d\alpha, \beta \rangle = \langle \alpha, d^*\beta \rangle, \quad \text{for all } \alpha, \beta \in \Omega^*(M, E).$	$\langle d\alpha, \beta \rangle = \langle \alpha, d^*\beta \rangle, \quad \text{for all } \alpha, \beta \in \Omega^*(M, E).$
cardona-qft-methods_ EQ0095	346	■ 0.523	$LM^{ev^*(-TM)} \wedge LM^{ev^*(-TM)} \rightarrow \text{Map}(8, M)^{ev^*(-TM)}.$	$LM^{ev^*(-TM)} \wedge LM^{ev^*(-TM)} \rightarrow \text{Map}(8, M)^{ev^*(-TM)}.$	$LM^{ev^*(-TM)} \wedge LM^{ev^*(-TM)} \rightarrow \text{Map}(8, M)^{ev^*(-TM)}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E00099	57	■ 0.523	$\begin{aligned} & \operatorname{Tr}(\frac{\mathcal{D}^2}{\Lambda^2} / \Lambda^2) \sim \frac{1}{(4\pi)^2} \left[\operatorname{rk}(\mathcal{D}) \int_0^\infty x f(x) dx \Lambda^4 + b_2(\mathcal{D}^2) \int_0^\infty f(x) dx \Lambda^2 \right. \\ & \quad \left. + \sum_{m=0}^\infty (-1)^m f^{(m)}(0) b_{2m+4}(\mathcal{D}^2) \Lambda^{-2m} \right], \quad \Lambda \rightarrow \infty \end{aligned}$	$\operatorname{Tr}(f(\mathcal{D}^2/\Lambda^2)) \sim \frac{1}{(4\pi)^2} \left[\operatorname{rk}(\mathcal{D}) \int_0^\infty x f(x) dx \Lambda^4 + b_2(\mathcal{D}^2) \int_0^\infty f(x) dx \Lambda^2 \right. \\ \left. + \sum_{m=0}^\infty (-1)^m f^{(m)}(0) b_{2m+4}(\mathcal{D}^2) \Lambda^{-2m} \right], \quad \Lambda \rightarrow \infty$	$\operatorname{Tr}(f(\mathcal{D}^2/\Lambda^2)) \sim \frac{1}{(4\pi)^2} \left[\operatorname{rk}(\mathcal{D}) \int_0^\infty x f(x) dx \Lambda^4 + b_2(\mathcal{D}^2) \int_0^\infty f(x) dx \Lambda^2 \right. \\ \left. + \sum_{m=0}^\infty (-1)^m f^{(m)}(0) b_{2m+4}(\mathcal{D}^2) \Lambda^{-2m} \right], \quad \Lambda \rightarrow \infty$
cardona-qft-methods_ E00154 (3)	85	■ 0.523	$\sigma_L(\mathcal{D}): \pi^*E \rightarrow \pi^*F.$	$\sigma_L(\mathcal{D}): \pi^*E \rightarrow \pi^*F.$	$\sigma_L(\mathcal{D}): \pi^*E \rightarrow \pi^*F.$
cardona-qft-methods_ E00919	352	■ 0.527	$C_*(P_G M^{-TM}) \xrightarrow{\sim} \mathcal{RH}om_{C^*(M)^e}(C^*(M), C^*(M) \# G)$	$C_*(P_G M^{-TM}) \xrightarrow{\sim} \mathcal{RH}om_{C^*(M)^e}(C^*(M), C^*(M) \# G)$	$C_*(P_G M^{-TM}) \xrightarrow{\sim} \mathcal{RH}om_{C^*(M)^e}(C^*(M), C^*(M) \# G)$
cardona-qft-methods_ E00356 (25)	171	■ 0.527	$\begin{aligned} & \mathbf{P}_8 = R_{a_2}^{a_1} R_{a_3}^{a_2} \dots R_{a_1}^{a_4}, \\ & (\mathbf{P}_4)^2 = (R_b^a R_a^b)^2, \end{aligned}$	$\begin{aligned} & \mathbf{P}_8 = R_{a_2}^{a_1} R_{a_3}^{a_2} \dots R_{a_1}^{a_4}, \\ & (\mathbf{P}_4)^2 = (R_b^a R_a^b)^2, \end{aligned}$	$\begin{aligned} & \mathbf{P}_8 = R_{a_2}^{a_1} R_{a_3}^{a_2} \dots R_{a_1}^{a_4}, \\ & (\mathbf{P}_4)^2 = (R_b^a R_a^b)^2, \end{aligned}$
cardona-qft-methods_ E00424	202	■ 0.529	$\begin{aligned} & C_{\lambda\mu\nu\kappa} = R_{\lambda\mu\nu\kappa} - \frac{1}{2} (g_{\lambda\nu} R_{\mu\kappa} - g_{\lambda\kappa} R_{\mu\nu} - g_{\mu\nu} R_{\lambda\kappa} + g_{\mu\kappa} R_{\lambda\nu}) \\ & \quad + \frac{1}{6} R_{\alpha}^{\alpha} (g_{\lambda\nu} g_{\mu\kappa} - g_{\lambda\kappa} g_{\mu\nu}). \end{aligned}$	$\begin{aligned} & C_{\lambda\mu\nu\kappa} = R_{\lambda\mu\nu\kappa} - \frac{1}{2} (g_{\lambda\nu} R_{\mu\kappa} - g_{\lambda\kappa} R_{\mu\nu} - g_{\mu\nu} R_{\lambda\kappa} + g_{\mu\kappa} R_{\lambda\nu}) \\ & \quad + \frac{1}{6} R_{\alpha}^{\alpha} (g_{\lambda\nu} g_{\mu\kappa} - g_{\lambda\kappa} g_{\mu\nu}). \end{aligned}$	$\begin{aligned} & C_{\lambda\mu\nu\kappa} = R_{\lambda\mu\nu\kappa} - \frac{1}{2} (g_{\lambda\nu} R_{\mu\kappa} - g_{\lambda\kappa} R_{\mu\nu} - g_{\mu\nu} R_{\lambda\kappa} + g_{\mu\kappa} R_{\lambda\nu}) \\ & \quad + \frac{1}{6} R_{\alpha}^{\alpha} (g_{\lambda\nu} g_{\mu\kappa} - g_{\lambda\kappa} g_{\mu\nu}). \end{aligned}$
cardona-qft-methods_ E00026	27	■ 0.531	$T(P) = \mu(c_P(x)) = c \int_M c_P(x) dx = c WRes(P)$	$T(P) = \mu(c_P(x)) = c \int_M c_P(x) dx = c WRes(P)$	$T(P) = \mu(c_P(x)) = c \int_M c_P(x) dx = c WRes(P).$
cardona-qft-methods_ E00747	283	■ 0.535	$\mathcal{D} = \mathcal{D}_0 + \frac{g_{\mathrm{YM}}^2}{16\pi^2} \hat{H}_{XXX} + O(g_{\mathrm{YM}}^4).$	$\mathcal{D} = \mathcal{D}_0 + \frac{g_{\mathrm{YM}}^2}{16\pi^2} \hat{H}_{XXX} + O(g_{\mathrm{YM}}^4).$	$\mathcal{D} = \mathcal{D}_0 + \frac{g_{\mathrm{YM}}^2}{16\pi^2} \hat{H}_{XXX} + O(g_{\mathrm{YM}}^4).$
cardona-qft-methods_ E00831 (10)	314	■ 0.539	$\begin{aligned} & \epsilon_{\mathrm{IIA}}^1 = \xi_+^1 \otimes \eta_+^1 + \xi_-^1 \otimes \eta_-^1, \\ & \epsilon_{\mathrm{IIA}}^2 = \xi_+^2 \otimes \eta_-^2 + \xi_-^2 \otimes \eta_+^2, \end{aligned}$	$\begin{aligned} & \epsilon_{\mathrm{IIA}}^1 = \xi_+^1 \otimes \eta_+^1 + \xi_-^1 \otimes \eta_-^1, \\ & \epsilon_{\mathrm{IIA}}^2 = \xi_+^2 \otimes \eta_-^2 + \xi_-^2 \otimes \eta_+^2, \end{aligned}$	$\begin{aligned} & \epsilon_{\mathrm{IIA}}^1 = \xi_+^1 \otimes \eta_+^1 + \xi_-^1 \otimes \eta_-^1, \\ & \epsilon_{\mathrm{IIA}}^2 = \xi_+^2 \otimes \eta_-^2 + \xi_-^2 \otimes \eta_+^2, \end{aligned}$
cardona-qft-methods_ E00834 (13)	317	■ 0.540	$\begin{aligned} & e^{-2A+\phi} (d + H \wedge) (e^{2A-\phi} \Phi_+) = 0, \\ & e^{-2A+\phi} (d + H \wedge) (e^{2A-\phi} \Phi_-) = dA \wedge \bar{\Phi}_- + \frac{i}{16} e^{\phi+A} * F_{\mathrm{IIA}} + \end{aligned}$	$\begin{aligned} & e^{-2A+\phi} (d + H \wedge) (e^{2A-\phi} \Phi_+) = 0, \\ & e^{-2A+\phi} (d + H \wedge) (e^{2A-\phi} \Phi_-) = dA \wedge \bar{\Phi}_- + \frac{i}{16} e^{\phi+A} * F_{\mathrm{IIA}} + \end{aligned}$	$\begin{aligned} & e^{-2A+\phi} (d + H \wedge) (e^{2A-\phi} \Phi_+) = 0, \\ & e^{-2A+\phi} (d + H \wedge) (e^{2A-\phi} \Phi_-) = dA \wedge \bar{\Phi}_- + \frac{i}{16} e^{\phi+A} * F_{\mathrm{IIA}} + \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0773 (14)	295	■ 0.590	$\begin{aligned} \xi_t &\equiv \frac{1}{D} \eta^{MN} \xi_{MN} \\ \xi_{MN}^s &= \frac{1}{2} (\xi_{MN} + \xi_{NM} - 2\xi_t \eta_{MN}) , \\ \xi_{MN}^a &= \frac{1}{2} (\xi_{MN} - \xi_{NM}) . \end{aligned}$	$\xi_t \equiv \frac{1}{D} \eta^{MN} \xi_{MN}$ $\xi_{MN}^s = \frac{1}{2} (\xi_{MN} + \xi_{NM} - 2\xi_t \eta_{MN}) ,$ $\xi_{MN}^a = \frac{1}{2} (\xi_{MN} - \xi_{NM}) .$	$\xi_t \equiv \frac{1}{D} \eta^{MN} \xi_{MN} ,$ $\xi_{MN}^s = \frac{1}{2} (\xi_{MN} + \xi_{NM} - 2\xi_t \eta_{MN}) ,$ $\xi_{MN}^a = \frac{1}{2} (\xi_{MN} - \xi_{NM}) .$
cardona-qft-methods_ EQ0399	182	■ 0.599	$\begin{aligned} &\& \mathbf{N}_{\{4\}} \mathbf{P}_{\{4\}}^{\text{Lor}} \\ &= \left(T^a T_a - e^a e^b R_{ab} \right) R^c{}_d R^d{}_c \end{aligned}$	$\mathbf{N}_4 \mathbf{P}_4^{\text{Lor}} =$ $(T^a T_a - e^a e^b R_{ab}) R^c{}_d R^d{}_c$	$\mathbf{N}_4 \mathbf{P}_4^{\text{Lor}} =$ $(T^a T_a - e^a e^b R_{ab}) R^c{}_d R^d{}_c$
cardona-qft-methods_ EQ0611 (19)	254	■ 0.600	$\begin{array}{r} P_{\mu} c_{p_1}^{\dagger} c_{p_2}^{\dagger} \dots c_{p_N}^{\dagger} 0\rangle e_{p_1} \otimes e_{p_2} \otimes \dots e_{p_N} + \\ c_{p_1}^{\dagger} c_{p_2}^{\dagger} \dots c_{p_N}^{\dagger} 0\rangle (-\sum_i P_{i\mu}) e_{p_1} \otimes e_{p_2} \otimes \dots e_{p_N} = 0, \end{array}$	$P_{\mu} c_{p_1}^{\dagger} c_{p_2}^{\dagger} \dots c_{p_N}^{\dagger} 0\rangle e_{p_1} \otimes e_{p_2} \otimes \dots e_{p_N} +$ $c_{p_1}^{\dagger} c_{p_2}^{\dagger} \dots c_{p_N}^{\dagger} 0\rangle (-\sum_i P_{i\mu}) e_{p_1} \otimes e_{p_2} \otimes \dots e_{p_N} = 0$	$P_{\mu} c_{p_1}^{\dagger} c_{p_2}^{\dagger} \dots c_{p_N}^{\dagger} 0\rangle e_{p_1} \otimes e_{p_2} \otimes \dots e_{p_N} +$ $c_{p_1}^{\dagger} c_{p_2}^{\dagger} \dots c_{p_N}^{\dagger} 0\rangle (-\sum_i P_{i\mu}) e_{p_1} \otimes e_{p_2} \otimes \dots e_{p_N} = 0,$
cardona-qft-methods_ EQ0478	219	■ 0.600	$S = \text{Tr} \left(f \left(D_A / \Lambda \right) / \Lambda \right) + \frac{1}{2} \left\langle J \tilde{\xi}, D_A \tilde{\xi} \right\rangle, \quad \tilde{\xi} \in \mathcal{H}_{cl}^+,$	$S = \text{Tr} \left(f \left(D_A / \Lambda \right) \right) + \frac{1}{2} \left\langle J \tilde{\xi}, D_A \tilde{\xi} \right\rangle, \quad \tilde{\xi} \in \mathcal{H}_{cl}^+,$	$S = \text{Tr} \left(f \left(D_A / \Lambda \right) \right) + \frac{1}{2} \left\langle J \tilde{\xi}, D_A \tilde{\xi} \right\rangle, \quad \tilde{\xi} \in \mathcal{H}_{cl}^+,$
cardona-qft-methods_ EQ0809 (9)	308	■ 0.604	$\mathcal{H} = \left(\begin{array}{cc} g - B g^{-1} B & B g^{-1} \\ -g^{-1} B & g^{-1} \end{array} \right) .$	$\mathcal{H} = \left(\begin{array}{cc} g - B g^{-1} B & B g^{-1} \\ -g^{-1} B & g^{-1} \end{array} \right) .$	$\mathcal{H} = \left(\begin{array}{cc} g - B g^{-1} B & B g^{-1} \\ -g^{-1} B & g^{-1} \end{array} \right) .$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0339 (7)	166	0.644	$\Omega_n \int_{M^{2n}} \mathbf{E}_{2n} \in \mathbb{Z}, \quad \tilde{\Omega}_k \int_{M^{4k}} \mathbf{P}_{4k} \in \mathbb{Z},$		$\Omega_n \int_{M^{2n}} \mathbf{E}_{2n} \in \mathbb{Z}, \quad \tilde{\Omega}_k \int_{M^{4k}} \mathbf{P}_{4k} \in \mathbb{Z},$
cardona-qft-methods_ EQ0846 (2)	329	0.651	$\begin{aligned} & \& \frac{\delta \mathcal{L}}{\delta X} = dX + (\Pi^\# \circ X)\eta = 0 \\ & \& \frac{\delta \mathcal{L}}{\delta \eta} = d\eta + \frac{1}{2} \langle (\partial \Pi^\# \circ X)\eta, \eta \rangle = 0. \end{aligned}$		$\frac{\delta \mathcal{L}}{\delta X} = dX + (\Pi^\# \circ X)\eta = 0$ $\frac{\delta \mathcal{L}}{\delta \eta} = d\eta + \frac{1}{2} \langle (\partial \Pi^\# \circ X)\eta, \eta \rangle = 0.$
cardona-qft-methods_ EQ0104	62	0.660	$\begin{aligned} & \& \gamma^{\alpha_1 \alpha_2} = \frac{1}{2} (\gamma^{\alpha_1} \gamma^{\alpha_2} - \gamma^{\alpha_2} \gamma^{\alpha_1}) \\ & \& \Omega_{\alpha_1 \alpha_2} = [\delta_{\alpha_1} + \tilde{A}_{\alpha_1}, \delta_{\alpha_2} + \tilde{A}_{\alpha_2}] = L(F_{\alpha_1 \alpha_2}) - R(F_{\alpha_1 \alpha_2}) \\ & \& F_{\alpha_1 \alpha_2} = \delta_{\alpha_1}(A_{\alpha_2}) - \delta_{\alpha_2}(A_{\alpha_1}) + [A_{\alpha_1}, A_{\alpha_2}] \end{aligned}$		$\gamma^{\alpha_1 \alpha_2} = \frac{1}{2} (\gamma^{\alpha_1} \gamma^{\alpha_2} - \gamma^{\alpha_2} \gamma^{\alpha_1}),$ $\Omega_{\alpha_1 \alpha_2} = [\delta_{\alpha_1} + \tilde{A}_{\alpha_1}, \delta_{\alpha_2} + \tilde{A}_{\alpha_2}] = L(F_{\alpha_1 \alpha_2}) - R(F_{\alpha_1 \alpha_2})$ $F_{\alpha_1 \alpha_2} = \delta_{\alpha_1}(A_{\alpha_2}) - \delta_{\alpha_2}(A_{\alpha_1}) + [A_{\alpha_1}, A_{\alpha_2}].$
cardona-qft-methods_ EQ0659 (9)	262	0.662	$S = T \exp \left(-i \int d^4 x H_I(x) \right).$		$S = T \exp \left(-i \int d^4 x H_I(x) \right).$
cardona-qft-methods_ EQ0896	346	0.664	$\begin{aligned} & \& LM^{ev^*(-TM)} \wedge LM^{ev^*(-TM)} = (LM \times LM)^{ev^*(-2TM)} \\ & \& \downarrow \\ & \& Map(8, M)^{ev^*(-TM)} \rightarrow LM^{ev^*(-TM)} \end{aligned}$		$LM^{ev^*(-TM)} \wedge LM^{ev^*(-TM)} = (LM \times LM)^{ev^*(-2TM)}$ $\downarrow$ $Map(8, M)^{ev^*(-TM)} \rightarrow LM^{ev^*(-TM)}$
cardona-qft-methods_ EQ0467	217	0.689	$\begin{aligned} & \& D_A^2 = (D^{1,0})^2 + 1_4 \otimes (D^{0,1})^2 - \gamma_5 [D^{1,0}, 1_4 \otimes D^{0,1}] \\ & \& [D^{1,0}, 1_4 \otimes D^{0,1}] = \sqrt{-1} \gamma^\mu [(\nabla_\mu^s + \mathbb{A}_\mu), 1_4 \otimes D^{0,1}] \end{aligned}$		$D_A^2 = (D^{1,0})^2 + 1_4 \otimes (D^{0,1})^2 - \gamma_5 [D^{1,0}, 1_4 \otimes D^{0,1}]$ $[D^{1,0}, 1_4 \otimes D^{0,1}] = \sqrt{-1} \gamma^\mu [(\nabla_\mu^s + \mathbb{A}_\mu), 1_4 \otimes D^{0,1}]$
cardona-qft-methods_ EQ0808 (8)	307	0.696	$[X + \zeta, Y + \eta]_C = [X, Y] + \mathcal{L}_X \eta - \mathcal{L}_Y \zeta - \frac{1}{2} d(\iota_X \eta - \iota_Y \zeta).$		$[X + \zeta, Y + \eta]_C = [X, Y] + \mathcal{L}_X \eta - \mathcal{L}_Y \zeta - \frac{1}{2} d(\iota_X \eta - \iota_Y \zeta).$
cardona-qft-methods_ EQ0330 (17)	162	0.706	$R^{\{a\}b} = \frac{1}{2} e_\alpha^a e_\beta^b R_{\mu\nu}^{\alpha\beta} dx^\mu \wedge dx^\nu.$		$R^{ab} = \frac{1}{2} e_\alpha^a e_\beta^b R_{\mu\nu}^{\alpha\beta} dx^\mu \wedge dx^\nu.$
cardona-qft-methods_ EQ0070 (21)	42	0.716	$a_2(\mathcal{D}^2) = -\frac{1}{12(4\pi)^{d/2}} \int_M s(x) dvol_g(x).$		$a_2(\mathcal{D}^2) = -\frac{1}{12(4\pi)^{d/2}} \int_M s(x) dvol_g(x).$
cardona-qft-methods_ EQ0233	109	0.718	$\chi(Y \cup Z) = \chi(Y) + \chi(Z) - \chi(Y \cap Z).$		$\chi(Y \cup Z) = \chi(Y) + \chi(Z) - \chi(Y \cap Z).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0017 (5)	22	0.722	$c_P(x)=\frac{1}{(2\pi)^d}\int_{\mathbb{S}^{d-1}}\mathrm{tr}\left(\sigma_{-d}^P(x,\xi)\right)d\xi.$	$c_P(x)=\frac{1}{(2\pi)^d}\int_{\mathbb{S}^{d-1}}\mathrm{tr}\left(\sigma_{-d}^P(x,\xi)\right)d\xi.$	$c_P(x)=\frac{1}{(2\pi)^d}\int_{\mathbb{S}^{d-1}}\mathrm{tr}\left(\sigma_{-d}^P(x,\xi)\right)d\xi.$
cardona-qft-methods_ EQ0043	32	0.725	$\sigma_1^d(x,\xi)=\lim_{t\rightarrow\infty}\frac{1}{t}\left(e^{-ith(x)}de^{ith(x)}\right)(x)=\lim_{t\rightarrow\infty}\frac{1}{t}itd_xh=id_xh=i\xi,$	$\sigma_1^d(x,\xi)=\lim_{t\rightarrow\infty}\frac{1}{t}\left(e^{-ith(x)}de^{ith(x)}\right)(x)=\lim_{t\rightarrow\infty}\frac{1}{t}itd_xh=id_xh=i\xi,$	$\sigma_1^d(x,\xi)=\lim_{t\rightarrow\infty}\frac{1}{t}\left(e^{-ith(x)}de^{ith(x)}\right)(x)=\lim_{t\rightarrow\infty}\frac{1}{t}itd_xh=id_xh=i\xi,$
cardona-qft-methods_ EQ0227	102	0.728	$e^{it}\cdot(z,\nu):=\left(e^{it}z,\nu\right),\quad z\in M,\,t,\nu\in\mathbb{R}.$	$e^{it}\cdot(z,\nu):=\left(e^{it}z,\nu\right),\quad z\in M,\,t,\nu\in\mathbb{R}.$	$e^{it}\cdot(z,\nu):=\left(e^{it}z,\nu\right),\quad z\in M,\,t,\nu\in\mathbb{R}.$
cardona-qft-methods_ EQ0855	331	0.737	$G\times_{(s,t)}G\stackrel{\mu}{\rightarrow}G\stackrel{i}{\rightarrow}G\stackrel{s}{\stackrel{\varepsilon}{\rightleftarrows}}_tM$	$G\times_{(s,t)}G\stackrel{\mu}{\rightarrow}G\stackrel{i}{\rightarrow}G\stackrel{s}{\stackrel{\varepsilon}{\rightleftarrows}}_tM$	$G\times_{(s,t)}G\stackrel{\mu}{\rightarrow}G\stackrel{i}{\rightarrow}G\stackrel{s}{\stackrel{\varepsilon}{\rightleftarrows}}_tM$
cardona-qft-methods_ EQ0794 (35)	301	0.738	$\begin{aligned} &\text{type IIB R - R : } \mathbf{8}_s\otimes\mathbf{8}_s=\mathbf{1}\oplus\mathbf{28}\oplus\mathbf{35}_+=C_0\oplus C_2\oplus C_{4+}\\ &\text{type IIA R - R : } \mathbf{8}_s\otimes\mathbf{8}_c=\mathbf{8}_v\oplus\mathbf{56}_t=C_1\oplus C_3. \end{aligned}$	$\begin{aligned} &\text{type IIB R - R : } \mathbf{8}_s\otimes\mathbf{8}_s=\mathbf{1}\oplus\mathbf{28}\oplus\mathbf{35}_+=C_0\oplus C_2\oplus C_{4+}\\ &\text{type IIA R - R : } \mathbf{8}_s\otimes\mathbf{8}_c=\mathbf{8}_v\oplus\mathbf{56}_t=C_1\oplus C_3. \end{aligned}$	$\begin{aligned} &\text{type IIB R - R : } \mathbf{8}_s\otimes\mathbf{8}_s=\mathbf{1}\oplus\mathbf{28}\oplus\mathbf{35}_+=C_0\oplus C_2\oplus C_{4+}\\ &\text{type IIA R - R : } \mathbf{8}_s\otimes\mathbf{8}_c=\mathbf{8}_v\oplus\mathbf{56}_t=C_1\oplus C_3. \end{aligned}$
cardona-qft-methods_ EQ0069	42	0.751	$\mathrm{Tr}_\mathrm{Dix}\left(D^{-d}\right)=\mathrm{Tr}_\omega\left(D^{-d}\right)=\frac{a_0\left(D^2\right)}{\Gamma(d/2+1)}=\frac{1}{d}W\mathrm{Res}\left(D^{-d}\right).$	$\mathrm{Tr}_\mathrm{Dix}\left(D^{-d}\right)=\mathrm{Tr}_\omega\left(D^{-d}\right)=\frac{a_0\left(D^2\right)}{\Gamma(d/2+1)}=\frac{1}{d}W\mathrm{Res}\left(D^{-d}\right).$	$\mathrm{Tr}_{Dix}\left(D^{-d}\right)=\mathrm{Tr}_\omega\left(D^{-d}\right)=\frac{a_0\left(D^2\right)}{\Gamma(d/2+1)}=\frac{1}{d}W\mathrm{Res}\left(D^{-d}\right).$
cardona-qft-methods_ EQ0687	269	0.753	$\begin{array}{ll} S^{+} \uparrow \rangle = 0, & S^{-} \uparrow \rangle = \downarrow \rangle, \\ S^{+} \downarrow \rangle = \uparrow \rangle, & S^{-} \downarrow \rangle = 0. \end{array}$	$\begin{array}{ll} S^{+} \uparrow \rangle = 0, & S^{-} \uparrow \rangle = \downarrow \rangle, \\ S^{+} \downarrow \rangle = \uparrow \rangle, & S^{-} \downarrow \rangle = 0. \end{array}$	$\begin{array}{ll} S^{+} \uparrow \rangle = 0, & S^{-} \uparrow \rangle = \downarrow \rangle, \\ S^{+} \downarrow \rangle = \uparrow \rangle, & S^{-} \downarrow \rangle = 0. \end{array}$
cardona-qft-methods_ EQ0282	137	0.755	$\underline{t}\in Y_G\Longleftrightarrow \Psi_G(\underline{t})\neq 0\Longleftrightarrow \Psi_{G/e}(\underline{t}')\neq -t_e\Psi_{G\setminus e}(\underline{t}'),$	$\underline{t}\in Y_G\Longleftrightarrow \Psi_G(\underline{t})\neq 0\Longleftrightarrow \Psi_{G/e}(\underline{t}')\neq -t_e\Psi_{G\setminus e}(\underline{t}'),$	$\underline{t}\in Y_G\Longleftrightarrow \Psi_G(\underline{t})\neq 0\Longleftrightarrow \Psi_{G/e}(\underline{t}')\neq -t_e\Psi_{G\setminus e}(\underline{t}'),$
cardona-qft-methods_ EQ0652 (2)	261	0.755	$\exp\left(-\beta V_{STAT}\left(x_1,x_2\right)\right)=\left\langle x_1,x_2\right e^{-\beta H}\left x_1,x_2\right\rangle ,$	$\exp\left(-\beta V_{STAT}\left(x_1,x_2\right)\right)=\left\langle x_1,x_2\right e^{-\beta H}\left x_1,x_2\right\rangle ,$	$\exp\left(-\beta V_{STAT}\left(x_1,x_2\right)\right)=\left\langle x_1,x_2\right e^{-\beta H}\left x_1,x_2\right\rangle ,$
cardona-qft-methods_ EQ0236 (2)	113	0.763	$[X]=a_0+a_1\mathbb{L}+\cdots+a_r\mathbb{L}^r,$	$[X]=a_0+a_1\mathbb{L}+\cdots+a_r\mathbb{L}^r,$	$[X]=a_0+a_1\mathbb{L}+\cdots+a_r\mathbb{L}^r,$
cardona-qft-methods_ EQ0883	343	0.769	$H^{n-p}(M)\times H^p(M)\rightarrow \mathbf{k}$	$H^{n-p}(M)\times H^p(M)\rightarrow \mathbf{k}$	$H^{n-p}(M)\times H^p(M)\rightarrow \mathbf{k}$
cardona-qft-methods_ EQ0181 (7)	92	0.772	$\mathrm{t}-\mathrm{Ind}_G\colon K_G\left(T_G^*M\right)\longrightarrow \widehat{R}(G)$	$\mathrm{t}-\mathrm{Ind}_G\colon K_G\left(T_G^*M\right)\longrightarrow \widehat{R}(G)$	$\mathrm{t}-\mathrm{Ind}_G\colon K_G\left(T_G^*M\right)\longrightarrow \widehat{R}(G)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0408	186	■ 0.772	$\delta B \backslash \left(\left(\phi^r \right) \right _{\partial M} \right) + \int_{\partial M} \Theta(\phi, \delta \phi) = 0$.	$\delta B(\phi^r _{\partial M}) + \int_{\partial M} \Theta(\phi, \delta \phi) = 0.$	$\delta B(\phi^r _{\partial M}) + \int_{\partial M} \Theta(\phi, \delta \phi) = 0.$
cardona-qft-methods_ EQ0022	25	■ 0.776	$W \operatorname{Res} (P_1 P_2) = \operatorname{Res}_{s=0} \operatorname{Tr} (P_2 \mathcal{D} ^{-s} P_1) = \operatorname{Res}_{s=0} \operatorname{Tr} (P_2 P_1 \mathcal{D} ^{-s}) = W \operatorname{Res} (P_2 P_1)$	$W \operatorname{Res} (P_1 P_2) = \operatorname{Res}_{s=0} \operatorname{Tr} (P_2 \mathcal{D} ^{-s} P_1) = \operatorname{Res}_{s=0} \operatorname{Tr} (P_2 P_1 \mathcal{D} ^{-s}) = W \operatorname{Res} (P_2 P_1)$	$W \operatorname{Res} (P_1 P_2) = \operatorname{Res}_{s=0} \operatorname{Tr} (P_2 \mathcal{D} ^{-s} P_1) = \operatorname{Res}_{s=0} \operatorname{Tr} (P_2 P_1 \mathcal{D} ^{-s}) = W \operatorname{Res} (P_2 P_1)$
cardona-qft-methods_ EQ0928	353	■ 0.782	$H \, H^* \left(C^*(M) \# G, C^*(M) \# G \right) \cong H_*^{\text{pro}} \left(\mathbb{L}(M \times_G EG)^{-T(M \times_G EG)} \right)$	$HH^*(C^*(M) \# G, C^*(M) \# G) \cong H_*^{\text{pro}} \left(\mathbb{L}(M \times_G EG)^{-T(M \times_G EG)} \right)$	$HH^*(C^*(M) \# G, C^*(M) \# G) \cong H_*^{\text{pro}} \left(\mathbb{L}(M \times_G EG)^{-T(M \times_G EG)} \right).$
cardona-qft-methods_ EQ0435	208	■ 0.790	$\begin{array}{l} \mathbf{2}_L \otimes \mathbf{1}^0 \oplus \mathbf{2}_R \otimes \mathbf{1}^0 \oplus \mathbf{2}_L \otimes \mathbf{3}^0 \oplus \mathbf{2}_R \otimes \mathbf{3}^0 \\ \oplus \mathbf{1} \otimes \mathbf{2}_L^0 \oplus \mathbf{1} \otimes \mathbf{2}_R^0 \oplus \mathbf{3} \otimes \mathbf{2}_L^0 \oplus \mathbf{3} \otimes \mathbf{2}_R^0 \end{array}$	$\begin{array}{l} \mathbf{2}_L \otimes \mathbf{1}^0 \oplus \mathbf{2}_R \otimes \mathbf{1}^0 \oplus \mathbf{2}_L \otimes \mathbf{3}^0 \oplus \mathbf{2}_R \otimes \mathbf{3}^0 \\ \oplus \mathbf{1} \otimes \mathbf{2}_L^0 \oplus \mathbf{1} \otimes \mathbf{2}_R^0 \oplus \mathbf{3} \otimes \mathbf{2}_L^0 \oplus \mathbf{3} \otimes \mathbf{2}_R^0 \end{array}$	$\begin{array}{l} \mathbf{2}_L \otimes \mathbf{1}^0 \oplus \mathbf{2}_R \otimes \mathbf{1}^0 \oplus \mathbf{2}_L \otimes \mathbf{3}^0 \oplus \mathbf{2}_R \otimes \mathbf{3}^0 \\ \oplus \mathbf{1} \otimes \mathbf{2}_L^0 \oplus \mathbf{1} \otimes \mathbf{2}_R^0 \oplus \mathbf{3} \otimes \mathbf{2}_L^0 \oplus \mathbf{3} \otimes \mathbf{2}_R^0 \end{array}$
cardona-qft-methods_ EQ0838 (17)	317	■ 0.790	$F^{(10)} = F + \operatorname{vol}_4 \wedge \lambda(*F),$	$F^{(10)} = F + \operatorname{vol}_4 \wedge \lambda(*F),$	$F^{(10)} = F + \operatorname{vol}_4 \wedge \lambda(*F) \, ,$
cardona-qft-methods_ EQ0959 (11)	361	■ 0.793	$\mathcal{C}: X_\Gamma \backslash \Sigma_n \rightarrow X_{\Gamma^\vee} \backslash \Sigma_n,$	$\mathcal{C}: X_\Gamma \backslash \Sigma_n \rightarrow X_{\Gamma^\vee} \backslash \Sigma_n,$	$\mathcal{C}: X_\Gamma \backslash \Sigma_n \rightarrow X_{\Gamma^\vee} \backslash \Sigma_n \, ,$
cardona-qft-methods_ EQ0359 (30)	171	■ 0.794	$\begin{aligned} & d\,C_3^{Lor} = R^{ab}R_{ab} = \mathbf{P}_4 \\ & d\,C_3^{Tor} = T^aT_a - e^ae^bR_{ab} = \mathbf{N}_4 \end{aligned}$	$\begin{aligned} dC_3^{Lor} &= R^{ab}R_{ab} = \mathbf{P}_4 \\ dC_3^{Tor} &= T^aT_a - e^ae^bR_{ab} = \mathbf{N}_4 \end{aligned}$	$\begin{aligned} dC_3^{Lor} &= R^{ab}R_{ab} = \mathbf{P}_4, \\ dC_3^{Tor} &= T^aT_a - e^ae^bR_{ab} = \mathbf{N}_4. \end{aligned}$
cardona-qft-methods_ EQ0818 (18)	309	■ 0.794	$\begin{aligned} & 6 \rightarrow 3 + \overline{3} \\ & 15 \rightarrow 8 + 3 + \overline{3} + 1 \\ & 10 + 10^* \rightarrow 6 + \overline{6} + 3 + \overline{3} + 1 + 1. \end{aligned}$	$\begin{aligned} & 6 \rightarrow 3 + \overline{3} \\ & 15 \rightarrow 8 + 3 + \overline{3} + 1 \\ & 10 + 10^* \rightarrow 6 + \overline{6} + 3 + \overline{3} + 1 + 1. \end{aligned}$	$\begin{aligned} & \mathbf{6} \rightarrow \mathbf{3} + \overline{\mathbf{3}} \, , \\ & \mathbf{15} \rightarrow \mathbf{8} + \mathbf{3} + \overline{\mathbf{3}} + \mathbf{1} \, , \\ & \mathbf{10} + \mathbf{10}^* \rightarrow \mathbf{6} + \overline{\mathbf{6}} + \mathbf{3} + \overline{\mathbf{3}} + \mathbf{1} + \mathbf{1} \, . \end{aligned}$
cardona-qft-methods_ EQ0906	349	■ 0.798	$\begin{aligned} & G \times P_G X \rightarrow P_G X \\ & ((f,k),g) \mapsto (fg,g^{-1}kg) \, . \end{aligned}$	$\begin{aligned} G \times P_G X &\rightarrow P_G X \\ ((f,k),g) &\mapsto (fg,g^{-1}kg) \, . \end{aligned}$	$\begin{aligned} G \times P_G X &\rightarrow P_G X \\ ((f,k),g) &\mapsto (fg,g^{-1}kg). \end{aligned}$
cardona-qft-methods_ EQ0362 (2)	174	■ 0.805	$\begin{aligned} & \delta \omega_b^a = d\lambda_b^a + \omega_c^a \lambda_b^c - \omega_b^c \lambda_c^a \\ & = D_\omega \lambda_b^a, \end{aligned}$	$\begin{aligned} \delta \omega_b^a &= d\lambda_b^a + \omega_c^a \lambda_b^c - \omega_b^c \lambda_c^a \\ &= D_\omega \lambda_b^a, \end{aligned}$	$\begin{aligned} \delta \omega_b^a &= d\lambda_b^a + \omega_c^a \lambda_b^c - \omega_b^c \lambda_c^a \\ &= D_\omega \lambda_b^a, \end{aligned}$
cardona-qft-methods_ EQ0146	82	■ 0.808	$\sigma_L(\Delta)(\xi_1,\xi_2,\dots,\xi_n) = \xi_1^2 + \xi_2^2 + \dots + \xi_n^2,$	$\sigma_L(\Delta)(\xi_1,\xi_2,\dots,\xi_n) = \xi_1^2 + \xi_2^2 + \dots + \xi_n^2,$	$\sigma_L(\Delta)(\xi_1,\xi_2,\dots,\xi_n) = \xi_1^2 + \xi_2^2 + \dots + \xi_n^2,$
cardona-qft-methods_ EQ0177 (4)	90	■ 0.808	$T_G^*M := \{ \xi \in T^*M \mid \xi \text{ is perpendicular to the orbits of } G \}.$	$T_G^*M := \{ \xi \in T^*M \mid \xi \text{ is perpendicular to the orbits of } G \}.$	$T_G^*M \, := \, \{ \xi \in T^*M \mid \xi \text{ is perpendicular to the orbits of } G \}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0059	39	■ 0.810	$\begin{aligned} k_{\{t\}}(x, y) &= \left\langle x, e^{t\Delta} y \right\rangle \\ &= \sum_{n,m=1}^{\infty} \overline{\psi_n(x)} \psi_n(y) e^{t\lambda_n}. \end{aligned}$	$k_t(x,y) = \langle x, e^{t\Delta} y \rangle = \sum_{n,m=1}^{\infty} \langle x, \psi_m \rangle \langle \psi_m, e^{t\Delta} \psi_n \rangle \langle \psi_n, y \rangle$ $= \sum_{n,m=1}^{\infty} \overline{\psi_n(x)} \psi_n(y) e^{t\lambda_n}.$	$k_t(x,y) = \langle x, e^{t\Delta} y \rangle = \sum_{n,m=1}^{\infty} \langle x, \psi_m \rangle \langle \psi_m, e^{t\Delta} \psi_n \rangle \langle \psi_n, y \rangle$ $= \sum_{n,m=1}^{\infty} \overline{\psi_n(x)} \psi_n(y) e^{t\lambda_n}.$
cardona-qft-methods_ EQ0254	119	■ 0.811	$P_{\{G\}}(\underline{\text{t}}, p) = \sum_C s_C \prod_{e \in C} t_e,$	$P_G(t,p) = \sum_C s_C \prod_{e \in C} t_e,$	$P_G(\underline{t}, p) = \sum_C s_C \prod_{e \in C} t_e \quad ,$
cardona-qft-methods_ EQ0758	286	■ 0.813	$\begin{aligned} &\text{string energy} \longleftrightarrow \text{scaling dimension} \\ &E(\lambda) = \Delta(\lambda). \end{aligned}$	$\text{string energy} \longleftrightarrow \text{scaling dimension}$ $E(\lambda) = \Delta(\lambda).$	$\text{string energy} \longleftrightarrow \text{scaling dimension}$ $E(\lambda) = \Delta(\lambda).$
cardona-qft-methods_ EQ0185 (1)	93	■ 0.816	$c(v)^2 = - v ^2 \text{ Id. }$	$c(v)^2 = - v ^2 \text{ Id.}$	$c(v)^2 = - v ^2 \text{ Id.}$
cardona-qft-methods_ EQ0615 (23)	255	■ 0.817	$\begin{aligned} S_{\{N\}} &= \left\langle \tau_{i,i+1} \mid i \in \{1, 2, \dots, N-1\}, \tau_{i,i+1}^2 = \mathbb{I}, \right. \\ &\quad \left. \tau_{i,i+1} \tau_{i+1,i+2} \tau_{i,i+1} = \tau_{i+1,i+2} \tau_{i,i+1} \tau_{i+1,i+2} \right\rangle \end{aligned}$	$S_N = \langle \tau_{i,i+1} \mid i \in \{1, 2, \dots, N-1\}, \tau_{i,i+1}^2 = \mathbb{I},$ $\tau_{i,i+1} \tau_{i+1,i+2} \tau_{i,i+1} = \tau_{i+1,i+2} \tau_{i,i+1} \tau_{i+1,i+2} \rangle$	$S_N = \langle \tau_{i,i+1} \mid i \in \{1, 2, \dots, N-1\}, \tau_{i,i+1}^2 = \mathbb{I},$ $\tau_{i,i+1} \tau_{i+1,i+2} \tau_{i,i+1} = \tau_{i+1,i+2} \tau_{i,i+1} \tau_{i+1,i+2} \rangle$
cardona-qft-methods_ EQ0320 (6)	160	■ 0.820	$\begin{aligned} \phi_{\parallel}^a(x) &\equiv \phi^a(x+dx) + dx^\mu \omega_{b\mu}^a(x) \phi^b(x) \\ &= \phi^a(x) + dx^\mu [\partial_\mu \phi^a(x) + \omega_{b\mu}^a(x) \phi^b(x)]. \end{aligned}$	$\phi_{\parallel}^a(x) \equiv \phi^a(x+dx) + dx^\mu \omega_{b\mu}^a(x) \phi^b(x)$ $= \phi^a(x) + dx^\mu [\partial_\mu \phi^a(x) + \omega_{b\mu}^a(x) \phi^b(x)].$	$\phi_{\parallel}^a(x) \equiv \phi^a(x+dx) + dx^\mu \omega_{b\mu}^a(x) \phi^b(x)$ $= \phi^a(x) + dx^\mu [\partial_\mu \phi^a(x) + \omega_{b\mu}^a(x) \phi^b(x)].$
cardona-qft-methods_ EQ0650 (57)	260	■ 0.831	$\varphi_{\theta}^* = \exp \frac{i}{2} \partial \wedge P \varphi_0,$	$\varphi_{\theta}^* = \exp \frac{i}{2} \partial \wedge P \varphi_0,$	$\varphi_{\theta}^* = \exp \frac{i}{2} \partial \wedge P \varphi_0,$
cardona-qft-methods_ EQ0029	28	■ 0.834	$\operatorname{Tr}_{\{D \text{ i } x\}}(P) = \frac{1}{d} \operatorname{WRes}(P) = \frac{1}{d} \int_M \int_{S^*M} \sigma_{-d}^P(x, \xi) d\xi dx $	$\operatorname{Tr}_{Dix}(P) = \frac{1}{d} \operatorname{WRes}(P) = \frac{1}{d} \int_M \int_{S^*M} \sigma_{-d}^P(x, \xi) d\xi dx $	$\operatorname{Tr}_{Dix}(P) = \frac{1}{d} \operatorname{WRes}(P) = \frac{1}{d} \int_M \int_{S^*M} \sigma_{-d}^P(x, \xi) d\xi dx $
cardona-qft-methods_ EQ0519 (5)	237	■ 0.835	$f * g(x) = f e^{i/2 \overleftarrow{\partial}_\mu \theta^{\mu\nu} \overrightarrow{\partial}_\nu} g$	$f * g(x) = f e^{i/2 \overleftarrow{\partial}_\mu \theta^{\mu\nu} \overrightarrow{\partial}_\nu} g$	$f * g(x) = f e^{i/2 \overleftarrow{\partial}_\mu \theta^{\mu\nu} \overrightarrow{\partial}_\nu} g.$
cardona-qft-methods_ EQ0613 (21)	254	■ 0.835	$\begin{aligned} &\int \prod_i d\mu(p_i) c_{\Delta p_1}^\dagger c_{\Delta p_2}^\dagger \dots c_{\Delta p_N}^\dagger 0\rangle e_{\Delta p_1} \otimes e_{\Delta p_1} \otimes \dots \otimes e_{\Delta p_N} = \\ &\int \prod_i d\mu(p_i) c_{p_1}^\dagger c_{p_2}^\dagger \dots c_{p_N}^\dagger 0\rangle e_{p_1} \otimes e_{p_1} \otimes \dots \otimes e_{p_N}, \end{aligned}$	$\int \prod_i d\mu(p_i) c_{\Delta p_1}^\dagger c_{\Delta p_2}^\dagger \dots c_{\Delta p_N}^\dagger 0\rangle e_{\Delta p_1} \otimes e_{\Delta p_1} \otimes \dots \otimes e_{\Delta p_N} =$ $\int \prod_i d\mu(p_i) c_{p_1}^\dagger c_{p_2}^\dagger \dots c_{p_N}^\dagger 0\rangle e_{p_1} \otimes e_{p_1} \otimes \dots \otimes e_{p_N}$	$\int \prod_i d\mu(p_i) c_{\Delta p_1}^\dagger c_{\Delta p_2}^\dagger \dots c_{\Delta p_N}^\dagger 0\rangle e_{\Delta p_1} \otimes e_{\Delta p_1} \otimes \dots \otimes e_{\Delta p_N} =$ $\int \prod_i d\mu(p_i) c_{p_1}^\dagger c_{p_2}^\dagger \dots c_{p_N}^\dagger 0\rangle e_{p_1} \otimes e_{p_1} \otimes \dots \otimes e_{p_N},$
cardona-qft-methods_ EQ0512 (5)	230	■ 0.840	$\partial_t x_j = \beta_j^{\text{SM}} + \beta_j^{\text{grav}},$	$\partial_t x_j = \beta_j^{\text{SM}} + \beta_j^{\text{grav}},$	$\partial_t x_j = \beta_j^{\text{SM}} + \beta_j^{\text{grav}},$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0046	35	■ 0.840	$\mathcal{C}\ell_x M = \left\{ \begin{array}{l} M_{2^m}(\mathbb{C}) \text{ when } d=2m \text{ is even,} \\ M_{2^m}(\mathbb{C}) \oplus M_{2^m}(\mathbb{C}) \text{ when } d=2m+1. \end{array} \right.$		$\mathcal{C}\ell_x M = \left\{ \begin{array}{l} M_{2^m}(\mathbb{C}) \text{ when } d=2m \text{ is even,} \\ M_{2^m}(\mathbb{C}) \oplus M_{2^m}(\mathbb{C}) \text{ when } d=2m+1. \end{array} \right.$
cardona-qft-methods_ EQ0749	284	■ 0.842	$\text{Super} \left[AdS_5 \times S^5 \right] := \frac{\widetilde{\text{PSU}}(2,2 4)}{\text{SO}(1,4) \times \text{SO}(5)}$		$\text{Super} \left[AdS_5 \times S^5 \right] := \frac{\widetilde{\text{PSU}}(2,2 4)}{\text{SO}(1,4) \times \text{SO}(5)},$
cardona-qft-methods_ EQ0335 (21)	164	■ 0.848	$D \left(R^a{}_b \phi^b \right) = R^a{}_b \wedge D \phi^b.$		$D(R^a{}_b \phi^b) = R^a{}_b \wedge D \phi^b.$
cardona-qft-methods_ EQ0898	347	■ 0.850	$\begin{gathered} A \stackrel{L}{\otimes}_{A^e} M = B(A) \stackrel{L}{\otimes}_{A^e} M \\ \mathcal{RH} \downarrow_{A^e} (A, M) = \text{Hom}_{A^e}(B(A), M) \end{gathered}$		$A \stackrel{L}{\otimes}_{A^e} M = B(A) \stackrel{L}{\otimes}_{A^e} M$ $\mathcal{RH}om_{A^e}(A, M) = \text{Hom}_{A^e}(B(A), M)$
cardona-qft-methods_ EQ0100	58	■ 0.850	$\text{Tr} \left(e^{-t \mathcal{D}^2} \right) = \frac{1}{t^2} \left(\frac{2}{3} + \frac{2}{3} t + \sum_{k=0}^n a_k t^{k+2} + \mathcal{O}(t^{n+3}) \right),$		$\text{Tr}(e^{-t \mathcal{D}^2}) = \frac{1}{t^2} \left(\frac{2}{3} + \frac{2}{3} t + \sum_{k=0}^n a_k t^{k+2} + \mathcal{O}(t^{n+3}) \right),$
cardona-qft-methods_ EQ0957 (9)	361	■ 0.850	$\mathcal{C} \left(X_\Gamma \cap \left(\mathbb{P}_k^{n-1} \setminus \Sigma_n \right) \right) = X_{\Gamma^\vee} \cap \left(\mathbb{P}_k^{n-1} \setminus \Sigma_n \right)$		$\mathcal{C} \left(X_\Gamma \cap \left(\mathbb{P}_k^{n-1} \setminus \Sigma_n \right) \right) = X_{\Gamma^\vee} \cap \left(\mathbb{P}_k^{n-1} \setminus \Sigma_n \right)$
cardona-qft-methods_ EQ0284	137	■ 0.856	$\Psi_{G_{2e}} = t_e t_{e'} \Psi_{G \setminus e} + (t_e + t_{e'}) \Psi_{G/e}.$		$\Psi_{G_{2e}} = t_e t_{e'} \Psi_{G \setminus e} + (t_e + t_{e'}) \Psi_{G/e}.$
cardona-qft-methods_ EQ0337 (23)	164	■ 0.856	$e^a \equiv e^a_\mu(x) dx^\mu \quad \text{and} \quad \omega^a_b \equiv \omega^a_{b\mu}(x) dx^\mu.$		$e^a \equiv e^a_\mu(x) dx^\mu \quad \text{and} \quad \omega^a_b \equiv \omega^a_{b\mu}(x) dx^\mu.$
cardona-qft-methods_ EQ0037	29	■ 0.857	$\tau_\lambda(T_1+T_2) - \tau_\lambda(T_1) - \tau_\lambda(T_2) \underset{\lambda \rightarrow \infty}{\simeq} \mathcal{O}\left(\frac{\log(\log \lambda)}{\log \lambda}\right), \text{ when } 0 \leq T_1, T_2 \in \mathcal{L}^{1,\infty}.$		$\tau_\lambda(T_1+T_2) - \tau_\lambda(T_1) - \tau_\lambda(T_2) \underset{\lambda \rightarrow \infty}{\simeq} \mathcal{O}\left(\frac{\log(\log \lambda)}{\log \lambda}\right), \text{ when } 0 \leq T_1, T_2 \in \mathcal{L}^{1,\infty}.$
cardona-qft-methods_ EQ0068	42	■ 0.858	$\begin{aligned} \text{WRes}(D^{-p}) &= \frac{2}{\Gamma(p/2)} a_{d-p}(D^2) \\ &= \frac{2}{(4\pi)^{d/2} \Gamma(p/2)} \int_M \text{tr}(k_{(d-p)/2}(x,x)) d\text{vol}_g(x) \end{aligned}$		$\begin{aligned} \text{WRes}(D^{-p}) &= \frac{2}{\Gamma(p/2)} a_{d-p}(D^2) \\ &= \frac{2}{(4\pi)^{d/2} \Gamma(p/2)} \int_M \text{tr}(k_{(d-p)/2}(x,x)) d\text{vol}_g(x). \end{aligned}$
cardona-qft-methods_ EQ0265	125	■ 0.863	$\langle G \rangle = \text{'propagator'} \cdot \langle G_1 \rangle \cdot \langle G_2 \rangle,$		$\langle G \rangle = \text{'propagator'} \cdot \langle G_1 \rangle \cdot \langle G_2 \rangle,$
cardona-qft-methods_ EQ0261	123	■ 0.872	$t_1 t_2 + t_1 t_3 + t_1 t_5 + t_2 t_4 + t_2 t_5 + t_3 t_4 + t_3 t_5 + t_4 t_5,$		$t_1 t_2 + t_1 t_3 + t_1 t_5 + t_2 t_4 + t_2 t_5 + t_3 t_4 + t_3 t_5 + t_4 t_5,$
cardona-qft-methods_ EQ0967 (7)	362	■ 0.873	$[X_{\Gamma^\vee}] = [\Sigma_n] = \frac{(1 + \mathbb{T})^n - 1 - \mathbb{T}^n}{\mathbb{T}} = \sum_{i=1}^{n-1} \binom{n}{i} \mathbb{T}^{n-1-i}.$		$[X_{\Gamma^\vee}] = [\Sigma_n] = \frac{(1 + \mathbb{T})^n - 1 - \mathbb{T}^n}{\mathbb{T}} = \sum_{i=1}^{n-1} \binom{n}{i} \mathbb{T}^{n-1-i}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E08338 (4)	165	■ 0.874	$\begin{aligned} \mathbf{P}_{2k} &=: R^{a_1}_{a_2} R^{a_2}_{a_3} \cdots R^{a_k}_{a_1} \\ v_k &=: e_{a_1} R^{a_1}_{a_2} R^{a_2}_{a_3} \cdots R^{a_k}_b e^b, \text{ odd } k \\ \tau_k &=: T_{a_1} R^{a_1}_{a_2} R^{a_2}_{a_3} \cdots R^{a_k}_b T^b, \text{ even } k \\ \zeta_k &=: e_{a_1} R^{a_1}_{a_2} R^{a_2}_{a_3} \cdots R^{a_k}_b T^b \\ \mathbf{E}_D &=: \epsilon_{a_1 a_2 \cdots a_D} R^{a_1 a_2} R^{a_3 a_4} \cdots R^{a_{D-1} a_D}, \text{ even } D \\ L_p &=: \epsilon_{a_1 a_2 \cdots a_D} R^{a_1 a_2} R^{a_3 a_4} \cdots R^{a_{2p-1} a_{2p}} e^{a_{2p+1}} \cdots e^{a_D}. \end{aligned}$	$\begin{aligned} \mathbf{P}_{2k} &=: R^{a_1}_{a_2} R^{a_2}_{a_3} \cdots R^{a_k}_{a_1} \\ v_k &=: e_{a_1} R^{a_1}_{a_2} R^{a_2}_{a_3} \cdots R^{a_k}_b e^b, \text{ odd } k \\ \tau_k &=: T_{a_1} R^{a_1}_{a_2} R^{a_2}_{a_3} \cdots R^{a_k}_b T^b, \text{ even } k \\ \zeta_k &=: e_{a_1} R^{a_1}_{a_2} R^{a_2}_{a_3} \cdots R^{a_k}_b T^b \\ \mathbf{E}_D &=: \epsilon_{a_1 a_2 \cdots a_D} R^{a_1 a_2} R^{a_3 a_4} \cdots R^{a_{D-1} a_D}, \text{ even } D \\ L_p &=: \epsilon_{a_1 a_2 \cdots a_D} R^{a_1 a_2} R^{a_3 a_4} \cdots R^{a_{2p-1} a_{2p}} e^{a_{2p+1}} \cdots e^{a_D}. \end{aligned}$	$\begin{aligned} \mathbf{P}_{2k} &=: R^{a_1}_{a_2} R^{a_2}_{a_3} \cdots R^{a_k}_{a_1} \\ v_k &=: e_{a_1} R^{a_1}_{a_2} R^{a_2}_{a_3} \cdots R^{a_k}_b e^b, \text{ odd } k \\ \tau_k &=: T_{a_1} R^{a_1}_{a_2} R^{a_2}_{a_3} \cdots R^{a_k}_b T^b, \text{ even } k \\ \zeta_k &=: e_{a_1} R^{a_1}_{a_2} R^{a_2}_{a_3} \cdots R^{a_k}_b T^b \\ \mathbf{E}_D &=: \epsilon_{a_1 a_2 \cdots a_D} R^{a_1 a_2} R^{a_3 a_4} \cdots R^{a_{D-1} a_D}, \text{ even } D \\ L_p &=: \epsilon_{a_1 a_2 \cdots a_D} R^{a_1 a_2} R^{a_3 a_4} \cdots R^{a_{2p-1} a_{2p}} e^{a_{2p+1}} \cdots e^{a_D}. \end{aligned}$
cardona-qft-methods_ E09915	351	■ 0.874	$H \, H_* \left((C^*(X) \# G, C^*(X) \# G) \cong \text{Tor}^*_{\mathbb{Z}G} \left(\mathbb{Z}, C^*(X) \otimes_{C^*(X)^e}^L C^*(X) \# G \right) \right)$	$HH_* \left((C^*(X) \# G, C^*(X) \# G) \cong \text{Tor}^*_{\mathbb{Z}G} \left(\mathbb{Z}, C^*(X) \otimes_{C^*(X)^e}^L C^*(X) \# G \right) \right)$	$HH_*((C^*(X) \# G, C^*(X) \# G) \cong \text{Tor}^*_{\mathbb{Z}G}(\mathbb{Z}, C^*(X) \otimes_{C^*(X)^e}^L C^*(X) \# G))$
cardona-qft-methods_ E08870	337	■ 0.875	$\begin{aligned} \mathcal{G} &= G \\ L &= \{(g_1, g_2, g_3) \mid (g_1, g_2) \in G \times_{(s,t)} G, g_3 = \mu(g_1, g_2)\}. \\ I &= g \mapsto g^{-1}, g \in G. \end{aligned}$	$\begin{aligned} \mathcal{G} &= G \\ L &= \{(g_1, g_2, g_3) \mid (g_1, g_2) \in G \times_{(s,t)} G, g_3 = \mu(g_1, g_2)\}. \\ I &= g \mapsto g^{-1}, g \in G. \end{aligned}$	$\begin{aligned} \mathcal{G} &= G. \\ L &= \{(g_1, g_2, g_3) \mid (g_1, g_2) \in G \times_{(s,t)} G, g_3 = \mu(g_1, g_2)\}. \\ I &= g \mapsto g^{-1}, g \in G. \end{aligned}$
cardona-qft-methods_ E08670	264	■ 0.875	$\begin{aligned} &\left\langle a_{lm} a_{l'm'}^* \right\rangle_{\theta} \\ &= * \pi^2 \int dk \sum_{l''=0, l'': \text{even}}^{\infty} i^{l+l'} (-1)^{l+m} (2l''+1) k^2 \Delta_l(k) \Delta_{l'}'(k) P_{\Phi}(k) i_{l''}(\theta k H) \\ &\times \sqrt{(2l+1)(2l'+1)} \begin{bmatrix} l & l' & l'' \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} l & l' & l'' \\ -m & m' & 0 \end{bmatrix}, \end{aligned}$	$\begin{aligned} &\left\langle a_{lm} a_{l'm'}^* \right\rangle_{\theta} \\ &= * \pi^2 \int dk \sum_{l''=0, l'': \text{even}}^{\infty} i^{l+l'} (-1)^{l+m} (2l''+1) k^2 \Delta_l(k) \Delta_{l'}'(k) P_{\Phi}(k) i_{l''}(\theta k H) \\ &\times \sqrt{(2l+1)(2l'+1)} \begin{bmatrix} l & l' & l'' \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} l & l' & l'' \\ -m & m' & 0 \end{bmatrix} \end{aligned}$	$\begin{aligned} &\left\langle a_{lm} a_{l'm'}^* \right\rangle_{\theta} \\ &= * \pi^2 \int dk \sum_{l''=0, l'': \text{even}}^{\infty} i^{l+l'} (-1)^{l+m} (2l''+1) k^2 \Delta_l(k) \Delta_{l'}'(k) P_{\Phi}(k) i_{l''}(\theta k H) \\ &\times \sqrt{(2l+1)(2l'+1)} \begin{bmatrix} l & l' & l'' \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} l & l' & l'' \\ -m & m' & 0 \end{bmatrix}, \end{aligned}$
cardona-qft-methods_ E08357 (26)	171	■ 0.877	$\begin{aligned} &\left(\mathbf{N}_4 \right)^2 = \left(T^a T_a - e^a e^b R_{ab} \right)^2, \\ \mathbf{N}_4 \mathbf{P}_4 &= \left(T^a T_a - e^a e^b R_{ab} \right) (R_d^c R_c^d), \end{aligned}$	$\begin{aligned} (\mathbf{N}_4)^2 &= (T^a T_a - e^a e^b R_{ab})^2 \\ \mathbf{N}_4 \mathbf{P}_4 &= (T^a T_a - e^a e^b R_{ab}) (R_d^c R_c^d) \end{aligned}$	$\begin{aligned} (\mathbf{N}_4)^2 &= (T^a T_a - e^a e^b R_{ab})^2, \\ \mathbf{N}_4 \mathbf{P}_4 &= (T^a T_a - e^a e^b R_{ab}) (R_d^c R_c^d), \end{aligned}$
cardona-qft-methods_ E08267	127	■ 0.880	$\Psi_{G_1}(t_1, \dots, t_{n_1}) \Psi_{G_2}(u_1, \dots, u_{n_2}) \neq 0,$	$\Psi_{G_1}(t_1, \dots, t_{n_1}) \Psi_{G_2}(u_1, \dots, u_{n_2}) \neq 0,$	$\Psi_{G_1}(t_1, \dots, t_{n_1}) \Psi_{G_2}(u_1, \dots, u_{n_2}) \neq 0 \quad ,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0314 (18)	156	■ 0.882	$\begin{aligned} & \& \left[\mathcal{H}_{\perp}(x), \mathcal{H}_{\perp}(y)\right] = g^{ij}(x)\delta(x,y),_i\mathcal{H}_j(y) - g^{ij}(y)\delta(y,x),_i\mathcal{H}_j(x) \\ & \mathcal{H}_i(x), \mathcal{H}_j(y) = \delta(x,y),_i\mathcal{H}_j(y) - \delta(x,y),_j\mathcal{H}_i(y) \\ & \mathcal{H}_{\perp}(x), \mathcal{H}_i(y) = \delta(x,y),_i\mathcal{H}_{\perp}(y) \end{aligned}$	$\begin{aligned} [\mathcal{H}_{\perp}(x), \mathcal{H}_{\perp}(y)] &= g^{ij}(x)\delta(x,y),_i\mathcal{H}_j(y) - g^{ij}(y)\delta(y,x),_i\mathcal{H}_j(x) \\ [\mathcal{H}_i(x), \mathcal{H}_j(y)] &= \delta(x,y),_i\mathcal{H}_j(y) - \delta(x,y),_j\mathcal{H}_i(y) \\ [\mathcal{H}_{\perp}(x), \mathcal{H}_i(y)] &= \delta(x,y),_i\mathcal{H}_{\perp}(y) \end{aligned}$	$\begin{aligned} [\mathcal{H}_{\perp}(x), \mathcal{H}_{\perp}(y)] &= g^{ij}(x)\delta(x,y),_i\mathcal{H}_j(y) - g^{ij}(y)\delta(y,x),_i\mathcal{H}_j(x) \\ [\mathcal{H}_i(x), \mathcal{H}_j(y)] &= \delta(x,y),_i\mathcal{H}_j(y) - \delta(x,y),_j\mathcal{H}_i(y) \\ [\mathcal{H}_{\perp}(x), \mathcal{H}_i(y)] &= \delta(x,y),_i\mathcal{H}_{\perp}(y) \end{aligned},$
cardona-qft-methods_ EQ0175 (3)	90	■ 0.890	$\operatorname{Ind}_G(\tilde{\mathcal{D}}) := \bigoplus_{V \in \operatorname{Irr}(G)} (\dim V) (\dim \operatorname{Ker}(\mathcal{D}) - \dim \operatorname{Ker}(\mathcal{D})) V.$	$\operatorname{Ind}_G(\tilde{\mathcal{D}}) := \bigoplus_{V \in \operatorname{Irr}(G)} (\dim V) (\dim \operatorname{Ker}(\mathcal{D}) - \dim \operatorname{Ker}(\mathcal{D})) V.$	$\operatorname{Ind}_G(\tilde{\mathcal{D}}) := \bigoplus_{V \in \operatorname{Irr}(G)} (\dim V) (\dim \operatorname{Ker}(\mathcal{D}) - \dim \operatorname{Ker}(\mathcal{D})) V.$
cardona-qft-methods_ EQ0820 (20)	310	■ 0.890	$\begin{aligned} & \Phi_+ = \eta_+^1 \otimes \eta_+^{2\dagger} = -\frac{i}{8} \omega \wedge e^{-iv \wedge v'} , \\ & \Phi_- = \eta_+^1 \otimes \eta_-^{2\dagger} = -\frac{1}{8} e^{-ij} \wedge (v + iv') . \end{aligned}$	$\begin{aligned} \Phi_+ &= \eta_+^1 \otimes \eta_+^{2\dagger} = -\frac{i}{8} \omega \wedge e^{-iv \wedge v'} , \\ \Phi_- &= \eta_+^1 \otimes \eta_-^{2\dagger} = -\frac{1}{8} e^{-ij} \wedge (v + iv') . \end{aligned}$	$\begin{aligned} \Phi_+ &= \eta_+^1 \otimes \eta_+^{2\dagger} = -\frac{i}{8} \omega \wedge e^{-iv \wedge v'} , \\ \Phi_- &= \eta_+^1 \otimes \eta_-^{2\dagger} = -\frac{1}{8} e^{-ij} \wedge (v + iv') . \end{aligned}$
cardona-qft-methods_ EQ0449	211	■ 0.896	$Y_{\downarrow 1} = V_1 Y_{\downarrow 1} V_3^*, \quad Y_{\uparrow 1} = V_2 Y_{\uparrow 1} V_3^*, \quad Y_R = V_2 Y_R \bar{V}_2^*,$	$Y'_{\downarrow 1} = V_1 Y_{\downarrow 1} V_3^*, \quad Y'_{\uparrow 1} = V_2 Y_{\uparrow 1} V_3^*, \quad Y'_R = V_2 Y_R \bar{V}_2^*,$	$Y'_{\downarrow 1} = V_1 Y_{\downarrow 1} V_3^*, \quad Y'_{\uparrow 1} = V_2 Y_{\uparrow 1} V_3^*, \quad Y'_R = V_2 Y_R \bar{V}_2^*,$
cardona-qft-methods_ EQ0427	205	■ 0.896	$J^2 = \varepsilon, \quad JD = \varepsilon' DJ, \quad \text{and} \quad J\gamma = \varepsilon'' \gamma J,$	$J^2 = \varepsilon, \quad JD = \varepsilon' DJ, \quad \text{and} \quad J\gamma = \varepsilon'' \gamma J,$	$J^2 = \varepsilon, \quad JD = \varepsilon' DJ, \quad \text{and} \quad J\gamma = \varepsilon'' \gamma J,$
cardona-qft-methods_ EQ0728	277	■ 0.905	$M_a(u) = \begin{pmatrix} \hat{A}(u) & \hat{B}(u) \\ \hat{C}(u) & \hat{D}(u) \end{pmatrix},$	$M_a(u) = \begin{pmatrix} \hat{A}(u) & \hat{B}(u) \\ \hat{C}(u) & \hat{D}(u) \end{pmatrix},$	$M_a(u) = \begin{pmatrix} \hat{A}(u) & \hat{B}(u) \\ \hat{C}(u) & \hat{D}(u) \end{pmatrix},$
cardona-qft-methods_ EQ0900	347	■ 0.907	$C_* (LM^{-TM}) \stackrel{\sim}{\rightarrow} \mathcal{R} \mathcal{H} om_{C^*(M)^e} (C^*(M), C^*(M))$	$C_* (LM^{-TM}) \stackrel{\sim}{\rightarrow} \mathcal{R} \mathcal{H} om_{C^*(M)^e} (C^*(M), C^*(M))$	$C_* (LM^{-TM}) \stackrel{\sim}{\rightarrow} \mathcal{R} \mathcal{H} om_{C^*(M)^e} (C^*(M), C^*(M))$
cardona-qft-methods_ EQ0979 (19)	364	■ 0.907	$[X_{\Gamma^\vee}] = \frac{(\mathbb{T}+1)^n-1}{\mathbb{T}} - \frac{\mathbb{T}^n-(-1)^n}{\mathbb{T}} - n\mathbb{T}^{n-2}.$	$[X_{\Gamma^\vee}] = \frac{(\mathbb{T}+1)^n-1}{\mathbb{T}} - \frac{\mathbb{T}^n-(-1)^n}{\mathbb{T}} - n\mathbb{T}^{n-2}.$	$[X_{\Gamma^\vee}] = \frac{(\mathbb{T}+1)^n-1}{\mathbb{T}} - \frac{\mathbb{T}^n-(-1)^n}{\mathbb{T}} - n\mathbb{T}^{n-2}.$
cardona-qft-methods_ EQ0549 (18)	244	■ 0.911	$\begin{aligned} & m_r[(S \otimes \operatorname{Id} \otimes \operatorname{Id})(\operatorname{Id} \otimes \Delta)\Delta(\alpha)] = e \otimes \alpha \\ & m'_r[(\operatorname{Id} \otimes S \otimes \operatorname{Id})(\operatorname{Id} \otimes \Delta)\Delta(\alpha)] = \alpha \otimes e, \quad \alpha \in H. \end{aligned}$	$\begin{aligned} m_r[(S \otimes \operatorname{Id} \otimes \operatorname{Id})(\operatorname{Id} \otimes \Delta)\Delta(\alpha)] &= e \otimes \alpha \\ m'_r[(\operatorname{Id} \otimes S \otimes \operatorname{Id})(\operatorname{Id} \otimes \Delta)\Delta(\alpha)] &= \alpha \otimes e, \quad \alpha \in H. \end{aligned}$	$\begin{aligned} m_r[(S \otimes \operatorname{Id} \otimes \operatorname{Id})(\operatorname{Id} \otimes \Delta)\Delta(\alpha)] &= e \otimes \alpha \\ m'_r[(\operatorname{Id} \otimes S \otimes \operatorname{Id})(\operatorname{Id} \otimes \Delta)\Delta(\alpha)] &= \alpha \otimes e, \quad \alpha \in H. \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0238	114	■ 0.922	$c(E)=1+c_{\{1\}}(E)+c_{\{2\}}(E)+\cdots+c_{\{\mathrm{rk}\}}(E)$.	$c(E) = 1 + c_1(E) + c_2(E) + \cdots + c_{\mathrm{rk}} E(E).$	$c(E) = 1 + c_1(E) + c_2(E) + \cdots + c_{\mathrm{rk}\, E}(E)$.
cardona-qft-methods_ EQ0264	125	■ 0.926	$\sum_{\text{graphs } G}\langle G\rangle\langle G\rangle=\exp\left(\sum_{\text{connected graphs } G}\langle G\rangle\right)$.	$\sum_{\text{graphs } G} \langle G \rangle = \exp \left(\sum_{\text{connected graphs } G} \langle G \rangle \right).$	$\sum_{\text{graphs } G} \langle G \rangle = \exp \left(\sum_{\text{connected graphs } G} \langle G \rangle \right)$.
cardona-qft-methods_ EQ0768 (9)	294	■ 0.927	$\left[\alpha_n^P,\alpha_m^Q\right]=\left[\tilde{\alpha}_n^P,\tilde{\alpha}_m^Q\right]=n\delta_{n+m}\eta^{PQ},\quad\left[x^P,p^Q\right]=i\eta^{PQ}.$	$\left[\alpha_n^P,\alpha_m^Q\right]=\left[\tilde{\alpha}_n^P,\tilde{\alpha}_m^Q\right]=n\delta_{n+m}\eta^{PQ},\quad\left[x^P,p^Q\right]=i\eta^{PQ}.$	$\left[\alpha_n^P,\alpha_m^Q\right]=\left[\tilde{\alpha}_n^P,\tilde{\alpha}_m^Q\right]=n\delta_{n+m}\eta^{PQ},\quad\left[x^P,p^Q\right]=i\eta^{PQ}.$
cardona-qft-methods_ EQ0098 (35)	56	■ 0.928	$\begin{aligned} & \mathsf{a}_k\left(\mathcal{D}^2\right)=\frac{1}{2}\Gamma\left(\frac{d-k}{2}\right)f\left \mathcal{D}\right ^{-d+k} \quad \text{for } k=0,\cdots,d-1, \\ & \mathsf{a}_d\left(\mathcal{D}^2\right)=\dim\mathcal{D}+\zeta_{\mathcal{D}^2}(0). \end{aligned}$	$a_k\left(\mathcal{D}^2\right)=\frac{1}{2}\Gamma\left(\frac{d-k}{2}\right)f\left \mathcal{D}\right ^{-d+k} \quad \text{for } k=0,\cdots,d-1,$ $a_d\left(\mathcal{D}^2\right)=\dim\mathcal{D}+\zeta_{\mathcal{D}^2}(0).$	$a_k(\mathcal{D}^2)=\tfrac{1}{2}\Gamma(\tfrac{d-k}{2})\oint \mathcal{D} ^{-d+k} \quad \text{for } k=0,\cdots,d-1,$ $a_d(\mathcal{D}^2)=\dim\mathcal{D}+\zeta_{\mathcal{D}^2}(0).$
cardona-qft-methods_ EQ0734 (1)	279	■ 0.929	$\begin{aligned} S=&\frac{N}{\lambda}\int\frac{\mathrm{d}^4x}{(2\pi)^4}\mathrm{Tr}\left(\frac{1}{4}F^{\mu\nu}F_{\mu\nu}+\frac{1}{2}(D^\mu)^*\varphi_mD_\mu\varphi_m+\bar\psi_\alpha^a(\sigma_\mu)^{\dot\alpha\beta}D^\mu\psi_{a\beta}\right. \\ &-\frac{1}{4}[\varphi^m,\varphi^n][\varphi_m,\varphi_n] \\ &\left.-\frac{i}{2}\psi_{a\alpha}(\Sigma_m)^{ab}\varepsilon^{\alpha\beta}[\varphi^m\psi_{a\beta}]-\frac{i}{2}\bar\psi_\alpha^a(\Sigma_m)^{ab}\varepsilon^{\dot\alpha\dot\beta}\left[\varphi^m\bar\psi_{a\dot\beta}\right]\right) \\ &+ \text{ghosts} + \text{gauge fixing} + \text{topological term} \end{aligned}$	$S=\frac{N}{\lambda}\int\frac{\mathrm{d}^4x}{(2\pi)^4}\mathrm{Tr}\left(\frac{1}{4}F^{\mu\nu}F_{\mu\nu}+\frac{1}{2}(D^\mu)^*\varphi_mD_\mu\varphi_m+\bar\psi_\alpha^a(\sigma_\mu)^{\dot\alpha\beta}D^\mu\psi_{a\beta}\right. \\ \left.-\frac{1}{4}[\varphi^m,\varphi^n][\varphi_m,\varphi_n] \right. \\ \left.-\frac{i}{2}\psi_{a\alpha}(\Sigma_m)^{ab}\varepsilon^{\alpha\beta}[\varphi^m\psi_{a\beta}]-\frac{i}{2}\bar\psi_\alpha^a(\Sigma_m)^{ab}\varepsilon^{\dot\alpha\dot\beta}\left[\varphi^m\bar\psi_{a\dot\beta}\right]\right) \\ + \text{ghosts} + \text{gauge fixing} + \text{topological term}$	$S=\frac{N}{\lambda}\int\frac{\mathrm{d}^4x}{(2\pi)^4}\mathrm{Tr}\left(\frac{1}{4}F^{\mu\nu}F_{\mu\nu}+\frac{1}{2}(D^\mu)^*\varphi_mD_\mu\varphi_m+\bar\psi_\alpha^a(\sigma_\mu)^{\dot\alpha\beta}D^\mu\psi_{a\beta}\right. \\ \left.-\frac{1}{4}[\varphi^m,\varphi^n][\varphi_m,\varphi_n] \right. \\ \left.-\frac{i}{2}\psi_{a\alpha}(\Sigma_m)^{ab}\varepsilon^{\alpha\beta}[\varphi^m\psi_{a\beta}]-\frac{i}{2}\bar\psi_\alpha^a(\Sigma_m)^{ab}\varepsilon^{\dot\alpha\dot\beta}[\varphi^m\bar\psi_{a\dot\beta}]\right) \\ + \text{ghosts} + \text{gauge fixing} + \text{topological term} . \quad (1)$
cardona-qft-methods_ EQ0972 (12)	363	■ 0.929	$\left[X_{\backslash\Gamma}\backslash\Sigma_n\right]=\left[X_{\backslash\Gamma^{\vee}}\backslash\Sigma_n\right]$.	$[X_\Gamma\backslash\Sigma_n]=[X_{\Gamma^\vee}\backslash\Sigma_n].$	$[X_\Gamma\backslash\Sigma_n]=[X_{\Gamma^\vee}\backslash\Sigma_n].$
cardona-qft-methods_ EQ0398	181	■ 0.929	$\begin{aligned} & \mathsf{d}\,\mathsf{L}_{\{3\}}^{\{\text{Exotic}\}}\,\&=\mathsf{R}_{\{b\}}^{\{a\}}\mathsf{R}_{\{a\}}^{\{b\}}\,\mathrm{p}\mathrm{m}\,\frac{2}{\mathrm{l}^{\{2\}}}\,\mathrm{l}\mathrm{e}\mathrm{f}\mathrm{t}\left(\mathsf{T}^{\{a\}}\mathsf{T}_{\{a\}}\cdot\mathsf{e}^{\{a\}}\mathsf{e}^{\{b\}}\mathsf{R}_{\{a\,b\}}\right)\,\&=\mathsf{F}_{\{B\}}^{\{A\}}\mathsf{F}_{\{A\}}^{\{B\}} \end{aligned}$	$dL_3^{\mathrm{Exotic}}=R_b^aR_a^b\pm\frac{2}{l^2}\left(T^aT_a-e^ae^bR_{ab}\right)\\=F_B^AF_A^B$	$dL_3^{Exotic}=R_b^aR_a^b\pm\frac{2}{l^2}\left(T^aT_a-e^ae^bR_{ab}\right)\\=F_B^AF_A^B.$
cardona-qft-methods_ EQ0044	33	■ 0.933	$i[D,f]=i\left[d+d^*,f\right]=i\left(i\sigma_1^d(df)-i\sigma_1^{d^*}(df)\right)=-i(\epsilon+\iota)(df).$	$i[D,f]=i\left[d+d^*,f\right]=i\left(i\sigma_1^d(df)-i\sigma_1^{d^*}(df)\right)=-i(\epsilon+\iota)(df).$	$i[D,f]=i\left[d+d^*,f\right]=i\left(i\sigma_1^d(df)-i\sigma_1^{d^*}(df)\right)=-i(\epsilon+\iota)(df).$
cardona-qft-methods_ EQ0112	73	■ 0.933	$\operatorname{Ind}(A)=-\operatorname{Ind}\left(A^*\right)=\operatorname{dimKer}(A)-\operatorname{dimKer}\left(A^*\right).$	$\operatorname{Ind}(A)=-\operatorname{Ind}(A^*)=\dim\operatorname{Ker}(A)-\dim\operatorname{Ker}(A^*).$	$\operatorname{Ind}(A)=-\operatorname{Ind}(A^*)=\dim\operatorname{Ker}(A)-\dim\operatorname{Ker}(A^*).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0845 (3)	329	■ 0.933	$\delta S = \int_{\Sigma} \frac{\delta \mathcal{L}}{\delta X} \delta X + \frac{\delta \mathcal{L}}{\delta \eta} \delta \eta + \text{boundary terms}$	$\delta S = \int_{\Sigma} \frac{\delta \mathcal{L}}{\delta X} \delta X + \frac{\delta \mathcal{L}}{\delta \eta} \delta \eta + \text{boundary terms}$	$\delta S = \int_{\Sigma} \frac{\delta \mathcal{L}}{\delta X} \delta X + \frac{\delta \mathcal{L}}{\delta \eta} \delta \eta + \text{boundary terms}.$
cardona-qft-methods_ EQ0840	32	■ 0.936	$\begin{gathered} \epsilon(\omega_1) \omega_2 = \omega_1 \wedge \omega_2, \\ \iota(\omega) (\omega_1 \wedge \dots \wedge \omega_m) = \sum_{j=1}^m (-1)^{j-1} g(\omega, \omega_j) \omega_1 \wedge \dots \wedge \widehat{\omega_j} \wedge \dots \wedge \omega_m \end{gathered}$	$\epsilon(\omega_1) \omega_2 = \omega_1 \wedge \omega_2, \\ \iota(\omega) (\omega_1 \wedge \dots \wedge \omega_m) = \sum_{j=1}^m (-1)^{j-1} g(\omega, \omega_j) \omega_1 \wedge \dots \wedge \widehat{\omega_j} \wedge \dots \wedge \omega_m$	$\epsilon(\omega_1) \omega_2 = \omega_1 \wedge \omega_2, \\ \iota(\omega) (\omega_1 \wedge \dots \wedge \omega_m) = \sum_{j=1}^m (-1)^{j-1} g(\omega, \omega_j) \omega_1 \wedge \dots \wedge \widehat{\omega_j} \wedge \dots \wedge \omega_m$
cardona-qft-methods_ EQ0745	283	■ 0.941	$\langle \mathcal{O}(x), \overline{\mathcal{O}}(y) \rangle = \frac{1}{(x-y)^{2\Delta}} \sim \frac{1}{(x-y)^{2\Delta_0}} (1 - g_{\text{YM}}^2 \gamma_1 \log[(x-y)^2 \Lambda^2]),$	$\langle \mathcal{O}(x), \overline{\mathcal{O}}(y) \rangle = \frac{1}{(x-y)^{2\Delta}} \sim \frac{1}{(x-y)^{2\Delta_0}} (1 - g_{\text{YM}}^2 \gamma_1 \log[(x-y)^2 \Lambda^2]),$	$\langle \mathcal{O}(x), \overline{\mathcal{O}}(y) \rangle = \frac{1}{(x-y)^{2\Delta}} \sim \frac{1}{(x-y)^{2\Delta_0}} (1 - g_{\text{YM}}^2 \gamma_1 \log[(x-y)^2 \Lambda^2]),$
cardona-qft-methods_ EQ0661 (11)	262	■ 0.943	$S^{(2)} = \frac{-i^2}{2!} \int d^4x d^4y T(H_I(x), H_I(y)) = \frac{-i^2}{2!} \int d^4x d^4y (\theta(x_0 - y_0) [H_I(x), H_I(y)] + H_I(y) H_I(x))$	$S^{(2)} = \frac{-i^2}{2!} \int d^4x d^4y T(H_I(x), H_I(y)) = \frac{-i^2}{2!} \int d^4x d^4y (\theta(x_0 - y_0) [H_I(x), H_I(y)] + H_I(y) H_I(x))$	$S^{(2)} = \frac{-i^2}{2!} \int d^4x d^4y T(H_I(x), H_I(y)) = \frac{-i^2}{2!} \int d^4x d^4y (\theta(x_0 - y_0) [H_I(x), H_I(y)] + H_I(y) H_I(x)),$
cardona-qft-methods_ EQ0308 (12)	154	■ 0.944	$d(\delta_{\text{gauge}} \mathcal{C}_{2k-1} = 0).$	$d(\delta_{\text{gauge}} \mathcal{C}_{2k-1} = 0).$	$d(\delta_{\text{gauge}} \mathcal{C}_{2k-1} = 0).$
cardona-qft-methods_ EQ0333 (19)	163	■ 0.947	$R_b^a = \bar{R}_b^a + \bar{D} \kappa_b^a + \kappa_c^a \kappa_b^c,$	$R_b^a = \bar{R}_b^a + \bar{D} \kappa_b^a + \kappa_c^a \kappa_b^c,$	$R_b^a = \bar{R}_b^a + \bar{D} \kappa_b^a + \kappa_c^a \kappa_b^c,$
cardona-qft-methods_ EQ0432	208	■ 0.947	$\mathcal{B} = \oplus^{4-\text{times}} M_2(\mathbb{C}) \oplus M_6(\mathbb{C}).$	$\mathcal{B} = \oplus^{4-\text{times}} M_2(\mathbb{C}) \oplus M_6(\mathbb{C}).$	$\mathcal{B} = \oplus^{4-\text{times}} M_2(\mathbb{C}) \oplus M_6(\mathbb{C}).$
cardona-qft-methods_ EQ0278	135	■ 0.947	$(\mathbb{T} - 1) \mathbb{T} (\mathbb{T} + 1)^2 + \mathbb{T} (\mathbb{T} + 1)^2 + (\mathbb{T} + 1) \mathbb{T} (\mathbb{T} + 1) = \mathbb{T} (\mathbb{T} + 1)^3.$	$(\mathbb{T} - 1) \mathbb{T} (\mathbb{T} + 1)^2 + \mathbb{T} (\mathbb{T} + 1)^2 + (\mathbb{T} + 1) \mathbb{T} (\mathbb{T} + 1) = \mathbb{T} (\mathbb{T} + 1)^3.$	$(\mathbb{T} - 1) \mathbb{T} (\mathbb{T} + 1)^2 + \mathbb{T} (\mathbb{T} + 1)^2 + (\mathbb{T} + 1) \mathbb{T} (\mathbb{T} + 1) = \mathbb{T} (\mathbb{T} + 1)^3.$
cardona-qft-methods_ EQ0132	78	■ 0.948	$f = (f_1, \dots, f_N) : \mathbb{R}^n \longrightarrow \mathbb{R}^N, \quad f_j \in C^\infty(\mathbb{R}^n).$	$f = (f_1, \dots, f_N) : \mathbb{R}^n \longrightarrow \mathbb{R}^N, \quad f_j \in C^\infty(\mathbb{R}^n).$	$f = (f_1, \dots, f_N) : \mathbb{R}^n \longrightarrow \mathbb{R}^N, \quad f_j \in C^\infty(\mathbb{R}^n).$
cardona-qft-methods_ EQ0237 (3)	113	■ 0.948	$N_q(X) = a_0 + a_1 q + \dots + a_r q^r,$	$N_q(X) = a_0 + a_1 q + \dots + a_r q^r,$	$N_q(X) = a_0 + a_1 q + \dots + a_r q^r,$
cardona-qft-methods_ EQ0316 (2)	159	■ 0.949	$ds^2 = \eta_{ab} dz^a dz^b = \eta_{ab} e_\mu^a(x) dx^\mu e_\nu^b(x) dx^\nu = g_{\mu\nu}(x) dx^\mu dx^\nu,$	$ds^2 = \eta_{ab} dz^a dz^b = \eta_{ab} e_\mu^a(x) dx^\mu e_\nu^b(x) dx^\nu = g_{\mu\nu}(x) dx^\mu dx^\nu,$	$ds^2 = \eta_{ab} dz^a dz^b = \eta_{ab} e_\mu^a(x) dx^\mu e_\nu^b(x) dx^\nu = g_{\mu\nu}(x) dx^\mu dx^\nu,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0363	174	■ 0.949	$\begin{aligned} \delta e^a &= -\lambda^a{}_c e^c \\ \delta R^a{}_b &= -(\lambda^a{}_c R^{cb} + \lambda^b{}_c R^{ac}), \\ \delta \epsilon_{abc} &= -(\lambda^d{}_a \epsilon_{dbc} + \lambda^d{}_b \epsilon_{adc} + \lambda^d{}_c \epsilon_{abd}) \equiv 0. \end{aligned}$	$\delta e^a = -\lambda^a{}_c e^c$ $\delta R^{ab} = -(\lambda^a{}_c R^{cb} + \lambda^b{}_c R^{ac})$ $\delta \epsilon_{abc} = -(\lambda^d{}_a \epsilon_{dbc} + \lambda^d{}_b \epsilon_{adc} + \lambda^d{}_c \epsilon_{abd}) \equiv 0.$	$\delta e^a = -\lambda^a{}_c e^c$ $\delta R^{ab} = -(\lambda^a{}_c R^{cb} + \lambda^b{}_c R^{ac}),$ $\delta \epsilon_{abc} = -(\lambda^d{}_a \epsilon_{dbc} + \lambda^d{}_b \epsilon_{adc} + \lambda^d{}_c \epsilon_{abd}) \equiv 0.$
cardona-qft-methods_ EQ0837 (16)	317	■ 0.950	$F_{\mathrm{IIA}} = F_0 \pm F_2 + F_4 \pm F_6, \quad F_{\mathrm{IIB}} = F_1 \pm F_3 + F_5.$	$F_{\mathrm{IIA}} \pm = F_0 \pm F_2 + F_4 \pm F_6, \quad F_{\mathrm{IIB}} \pm = F_1 \pm F_3 + F_5.$	$F_{\mathrm{IIA}} \pm = F_0 \pm F_2 + F_4 \pm F_6, \quad F_{\mathrm{IIB}} \pm = F_1 \pm F_3 + F_5.$
cardona-qft-methods_ EQ0186	93	■ 0.950	$c(v) \, c(u) + c(u) \, c(v) = -2 \langle v, u \rangle \mathrm{Id}, \quad \text{for all } u, v \in V.$	$c(v)c(u) + c(u)c(v) = -2\langle v, u \rangle \mathrm{Id}, \quad \text{for all } u, v \in V.$	$c(v) \, c(u) + c(u) \, c(v) = -2 \langle v, u \rangle \mathrm{Id}, \quad \text{for all } u, v \in V.$
cardona-qft-methods_ EQ0931	353	■ 0.950	$E G_{\{n\}} / C(g) \times E G_{\{n\}} / C(h) \stackrel{\Delta}{\longleftarrow} E G_{\{n\}} / (C(g) \cap C(h)) \longrightarrow E G_{\{n\}} / C(gh)$	$EG_n/C(g) \times EG_n/C(h) \stackrel{\Delta}{\longleftarrow} EG_n/(C(g) \cap C(h)) \longrightarrow EG_n/C(gh)$	$EG_n/C(g) \times EG_n/C(h) \stackrel{\Delta}{\longleftarrow} EG_n/(C(g) \cap C(h)) \longrightarrow EG_n/C(gh)$
cardona-qft-methods_ EQ0360 (31)	171	■ 0.951	$C_7^{Lor} = \mathrm{Tr} \left[\omega(d\omega)^3 + \frac{8}{5} \omega^3(d\omega)^2 + \frac{4}{5} \omega^2(d\omega)\omega(d\omega) + 2\omega^5(d\omega) + \frac{4}{7} \omega^7 \right]$	$C_7^{Lor} = \mathrm{Tr} \left[\omega(d\omega)^3 + \frac{8}{5} \omega^3(d\omega)^2 + \frac{4}{5} \omega^2(d\omega)\omega(d\omega) + 2\omega^5(d\omega) + \frac{4}{7} \omega^7 \right]$	$C_7^{Lor} = \mathrm{Tr}[\omega(d\omega)^3 + \frac{8}{5} \omega^3(d\omega)^2 + \frac{4}{5} \omega^2(d\omega)\omega(d\omega) + 2\omega^5(d\omega) + \frac{4}{7} \omega^7],$
cardona-qft-methods_ EQ0315 (1)	159	■ 0.953	$dz^a = e^a_\mu(x) dx^\mu,$	$dz^a = e^a_\mu(x) dx^\mu,$	$dz^a = e^a_\mu(x) dx^\mu,$
cardona-qft-methods_ EQ0765 (6)	293	■ 0.953	$\begin{aligned} X^M(\tau, \sigma) &= X_R^M(\sigma^-) + X_L^M(\sigma^+) \\ X_R^M(\sigma^-) &= \frac{1}{2} x^M + \alpha' p^M \sigma^- + i \left(\frac{\alpha'}{2} \right)^{1/2} \sum_{n \neq 0} \frac{1}{n} \alpha_n^M e^{-2in\sigma^-} \\ X_L^M(\sigma^+) &= \frac{1}{2} x^M + \alpha' p^M \sigma^+ + i \left(\frac{\alpha'}{2} \right)^{1/2} \sum_{n \neq 0} \frac{1}{n} \tilde{\alpha}_n^M e^{-2in\sigma^+}, \end{aligned}$	$X^M(\tau, \sigma) = X_R^M(\sigma^-) + X_L^M(\sigma^+)$ $X_R^M(\sigma^-) = \frac{1}{2} x^M + \alpha' p^M \sigma^- + i \left(\frac{\alpha'}{2} \right)^{1/2} \sum_{n \neq 0} \frac{1}{n} \alpha_n^M e^{-2in\sigma^-}$ $X_L^M(\sigma^+) = \frac{1}{2} x^M + \alpha' p^M \sigma^+ + i \left(\frac{\alpha'}{2} \right)^{1/2} \sum_{n \neq 0} \frac{1}{n} \tilde{\alpha}_n^M e^{-2in\sigma^+},$	$X^M(\tau, \sigma) = X_R^M(\sigma^-) + X_L^M(\sigma^+)$ $X_R^M(\sigma^-) = \frac{1}{2} x^M + \alpha' p^M \sigma^- + i \left(\frac{\alpha'}{2} \right)^{1/2} \sum_{n \neq 0} \frac{1}{n} \alpha_n^M e^{-2in\sigma^-}$ $X_L^M(\sigma^+) = \frac{1}{2} x^M + \alpha' p^M \sigma^+ + i \left(\frac{\alpha'}{2} \right)^{1/2} \sum_{n \neq 0} \frac{1}{n} \tilde{\alpha}_n^M e^{-2in\sigma^+},$
cardona-qft-methods_ EQ0736 (6)	280	■ 0.954	$\begin{aligned} \{ [M_{\mu\nu}, K_\rho] &= i(\eta_{\rho\nu} K_\mu - \eta_{\mu\rho} K_\nu), \\ [P_\mu, K_\nu] &= 2i(\eta_{\mu\nu} \mathcal{D} - M_{\mu\nu}). \end{aligned}$	$[M_{\mu\nu}, K_\rho] = i(\eta_{\rho\nu} K_\mu - \eta_{\mu\rho} K_\nu),$ $[P_\mu, K_\nu] = 2i(\eta_{\mu\nu} \mathcal{D} - M_{\mu\nu}).$	$[M_{\mu\nu}, K_\rho] = i(\eta_{\rho\nu} K_\mu - \eta_{\mu\rho} K_\nu),$ $[P_\mu, K_\nu] = 2i(\eta_{\mu\nu} \mathcal{D} - M_{\mu\nu}).$
cardona-qft-methods_ EQ0323 (9)	161	■ 0.954	$\varphi_{ }^\mu(x) = \varphi^\mu(x) + dx^\lambda \left[\partial_\lambda \varphi^\mu(x) + \Gamma_{\lambda\rho}^\mu(x) \varphi^\rho(x) \right].$	$\varphi_{ }^\mu(x) = \varphi^\mu(x) + dx^\lambda \left[\partial_\lambda \varphi^\mu(x) + \Gamma_{\lambda\rho}^\mu(x) \varphi^\rho(x) \right].$	$\varphi_{ }^\mu(x) = \varphi^\mu(x) + dx^\lambda [\partial_\lambda \varphi^\mu(x) + \Gamma_{\lambda\rho}^\mu(x) \varphi^\rho(x)].$
cardona-qft-methods_ EQ0147	83	■ 0.957	$\mathcal{D} := -i \frac{d}{dt} + \sin t : C^\infty(M) \rightarrow C^\infty(M).$	$\mathcal{D} := -i \frac{d}{dt} + \sin t : C^\infty(M) \rightarrow C^\infty(M).$	$\mathcal{D} := -i \frac{d}{dt} + \sin t : C^\infty(M) \rightarrow C^\infty(M).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0665	40	■ 0.957	$\begin{aligned} \omega_{\nu} &= \frac{1}{2} g_{\nu\mu} (\mathbb{A}^{\mu} + g^{\sigma\varepsilon} \Gamma_{\sigma\varepsilon}^{\mu} \text{id}_V) \\ E &= \mathbb{B} - g^{\nu\mu} (\partial_{\nu} \omega_{\mu} + \omega_{\nu} \omega_{\mu} - \omega_{\sigma} \Gamma_{\nu\mu}^{\sigma}) . \end{aligned}$	$\omega_{\nu} = \frac{1}{2} g_{\nu\mu} (\mathbb{A}^{\mu} + g^{\sigma\varepsilon} \Gamma_{\sigma\varepsilon}^{\mu} \text{id}_V)$ $E = \mathbb{B} - g^{\nu\mu} (\partial_{\nu} \omega_{\mu} + \omega_{\nu} \omega_{\mu} - \omega_{\sigma} \Gamma_{\nu\mu}^{\sigma}) .$	$\omega_{\nu} = \frac{1}{2} g_{\nu\mu} (\mathbb{A}^{\mu} + g^{\sigma\varepsilon} \Gamma_{\sigma\varepsilon}^{\mu} \text{id}_V) ,$ $E = \mathbb{B} - g^{\nu\mu} (\partial_{\nu} \omega_{\mu} + \omega_{\nu} \omega_{\mu} - \omega_{\sigma} \Gamma_{\nu\mu}^{\sigma}) .$
cardona-qft-methods_ EQ0618 (26)	256	■ 0.957	$\left(f_1 f_2\right)(p)=f_1(p) \ f_2(p) \ .$	$(f_1 f_2)(p) = f_1(p) f_2(p).$	$(f_1 f_2)(p) = f_1(p) f_2(p).$
cardona-qft-methods_ EQ0751	284	■ 0.957	$\left.\frac{30 \mid 32}{10 \mid 0 \oplus 10 \mid 0} = 10 \right 32,$	$\frac{30 \mid 32}{10 \mid 0 \oplus 10 \mid 0} = 10 \Big 32,$	$\frac{30 \mid 32}{10 \mid 0 \oplus 10 \mid 0} = 10 \Big 32,$
cardona-qft-methods_ EQ0816 (16)	309	■ 0.958	$J_{mn}=\mp 2\mathrm{i}\eta_{\pm}^{\dagger}\gamma_{mn}\eta_{\pm}\quad \Omega_{mnp}=-2\mathrm{i}\eta_{-}^{\dagger}\gamma_{mnp}\eta_{+}.$	$J_{mn}=\mp 2\mathrm{i}\eta_{\pm}^{\dagger}\gamma_{mn}\eta_{\pm}\quad \Omega_{mnp}=-2\mathrm{i}\eta_{-}^{\dagger}\gamma_{mnp}\eta_{+}.$	$J_{mn}=\mp 2\mathrm{i}\eta_{\pm}^{\dagger}\gamma_{mn}\eta_{\pm}\quad \Omega_{mnp}=-2\mathrm{i}\eta_{-}^{\dagger}\gamma_{mnp}\eta_{+}.$
cardona-qft-methods_ EQ0488	221	■ 0.958	$\begin{aligned} a_0(x,P) &= (4\pi)^{-m/2} \text{Tr}(\text{id}) \\ a_2(x,P) &= (4\pi)^{-m/2} \text{Tr}\left(-\frac{R}{6} \text{id} + E\right) \\ a_4(x,P) &= (4\pi)^{-m/2} \frac{1}{360} \text{Tr}\left(-12 R_{;\mu}{}^{\mu} + 5 R^2 - 2 R_{\mu\nu} R^{\mu\nu} \right. \\ &\quad \left. + 2 R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 60 R E + 180 E^2 + 60 E_{;\mu}{}^{\mu} \right. \\ &\quad \left. + 30 \Omega_{\mu\nu} \Omega^{\mu\nu}\right). \end{aligned}$	$a_0(x,P) = (4\pi)^{-m/2} \text{Tr}(\text{id})$ $a_2(x,P) = (4\pi)^{-m/2} \text{Tr}\left(-\frac{R}{6} \text{id} + E\right)$ $a_4(x,P) = (4\pi)^{-m/2} \frac{1}{360} \text{Tr}\left(-12 R_{;\mu}{}^{\mu} + 5 R^2 - 2 R_{\mu\nu} R^{\mu\nu} \right. \\ \left. + 2 R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 60 R E + 180 E^2 + 60 E_{;\mu}{}^{\mu} \right. \\ \left. + 30 \Omega_{\mu\nu} \Omega^{\mu\nu}\right) .$	$a_0(x,P) = (4\pi)^{-m/2} \text{Tr}(\text{id})$ $a_2(x,P) = (4\pi)^{-m/2} \text{Tr}\left(-\frac{R}{6} \text{id} + E\right)$ $a_4(x,P) = (4\pi)^{-m/2} \frac{1}{360} \text{Tr}\left(-12 R_{;\mu}{}^{\mu} + 5 R^2 - 2 R_{\mu\nu} R^{\mu\nu} \right. \\ \left. + 2 R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 60 R E + 180 E^2 + 60 E_{;\mu}{}^{\mu} \right. \\ \left. + 30 \Omega_{\mu\nu} \Omega^{\mu\nu}\right).$
cardona-qft-methods_ EQ0500	226	■ 0.959	$\frac{2f_2\Lambda_{unif}^2}{f_0} \leq \mathfrak{c}(\Lambda_{unif}) \leq \frac{6f_2\Lambda_{unif}^2}{f_0}$	$\frac{2f_2\Lambda_{unif}^2}{f_0} \leq \mathfrak{c}(\Lambda_{unif}) \leq \frac{6f_2\Lambda_{unif}^2}{f_0}$	$\frac{2f_2\Lambda_{unif}^2}{f_0} \leq \mathfrak{c}(\Lambda_{unif}) \leq \frac{6f_2\Lambda_{unif}^2}{f_0}$
cardona-qft-methods_ EQ0234 (1)	112	■ 0.961	$[\widetilde{X}]-[E]=[X]-[Z]$	$[\widetilde{X}]-[E]=[X]-[Z]$	$[\widetilde{X}]-[E]=[X]-[Z]$
cardona-qft-methods_ EQ0262	123	■ 0.961	$\hat{Y}_{\{G\}} \text{ mixed-Tate } \Longrightarrow \left[\hat{Y}_G\right] \in \mathbb{Z}[\mathbb{L}] \Longrightarrow N_q(G) \text{ is a polynomial in } q.$	$\hat{Y}_G \text{ mixed-Tate } \Longrightarrow \left[\hat{Y}_G\right] \in \mathbb{Z}[\mathbb{L}] \Longrightarrow N_q(G) \text{ is a polynomial in } q.$	$\hat{Y}_G \text{ mixed-Tate } \Longrightarrow \left[\hat{Y}_G\right] \in \mathbb{Z}[\mathbb{L}] \Longrightarrow N_q(G) \text{ is a polynomial in } q.$
cardona-qft-methods_ EQ0332	163	■ 0.963	$d\,e^{\{a\}}+\overline{\omega}_{\{b\}}^a\wedge e^{\{b\}}\equiv 0,\quad \text{and}\quad T^a=\kappa_b^a\wedge e^b\,.$	$de^a+\bar{\omega}_b^a\wedge e^b\equiv 0,\quad \text{and}\quad T^a=\kappa_b^a\wedge e^b.$	$de^a+\bar{\omega}_b^a\wedge e^b\equiv 0,\quad \text{and}\quad T^a=\kappa_b^a\wedge e^b.$
cardona-qft-methods_ EQ0142 (6)	81	■ 0.963	$\sigma_L(\mathcal{D})(x_0,\xi)=\sum_{ \alpha =k}a_\alpha(x_0)\xi_1^{\alpha_1}\cdots\xi_n^{\alpha_n}.$	$\sigma_L(\mathcal{D})(x_0,\xi)=\sum_{ \alpha =k}a_\alpha(x_0)\xi_1^{\alpha_1}\cdots\xi_n^{\alpha_n}.$	$\sigma_L(\mathcal{D})(x_0,\xi)=\sum_{ \alpha =k}a_\alpha(x_0)\xi_1^{\alpha_1}\cdots\xi_n^{\alpha_n}.$
cardona-qft-methods_ EQ0454	214	■ 0.964	$\left(C^{\infty}(M,\mathcal{E}),L^2(M,\mathcal{E}\otimes S),\mathcal{D}_{\mathcal{E}}\right),$	$(C^{\infty}(M,\mathcal{E}),L^2(M,\mathcal{E}\otimes S),\mathcal{D}_{\mathcal{E}}),$	$(C^{\infty}(M,\mathcal{E}),L^2(M,\mathcal{E}\otimes S),\mathcal{D}_{\mathcal{E}}),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E00086	50	■ 0.965	$\begin{gathered} \operatorname{Tad}_{\mathcal{D}+A}(d-k)=- (d-k) \int A \mathcal{D} \mathcal{D} ^{-(d-k)-2}, \quad \forall k \neq d \\ \operatorname{Tad}_{\mathcal{D}+A}(0)=- \int A \mathcal{D}^{-1}. \end{gathered}$		
cardona-qft-methods_ E00106 (1)	72	■ 0.965	$\operatorname{Ind}(A) := \operatorname{dimKer}(A) - \operatorname{dimCoker}(A) \in \mathbb{Z},$	$\operatorname{Ind}(A) := \dim \operatorname{Ker}(A) - \dim \operatorname{Coker}(A) \in \mathbb{Z},$	$\operatorname{Ind}(A) := \dim \operatorname{Ker}(A) - \dim \operatorname{Coker}(A) \in \mathbb{Z},$
cardona-qft-methods_ E00272	131	■ 0.966	$C_{G_1 \amalg G_2}(T) = \chi(Y_{G_1}) \chi(Y_{G_2}) T^2 + \text{h.o.t} :$	$C_{G_1 \amalg G_2}(T) = \chi(Y_{G_1}) \chi(Y_{G_2}) T^2 + \text{h.o.t} :$	$C_{G_1 \amalg G_2}(T) = \chi(Y_{G_1}) \chi(Y_{G_2}) T^2 + \text{h.o.t} :$
cardona-qft-methods_ E00055	38	■ 0.970	$\sigma_d^{\mathcal{P}_{g'}}(x,\xi) = e^{-h(x)/2} U_{g',g}^{-1}(x) \sigma_d^{\mathcal{P}_g}(x,\xi) U_{g',g}(x) e^{-h(x)/2}, \quad \xi \in T_x^* M.$	$\sigma_d^{\mathcal{P}_{g'}}(x,\xi) = e^{-h(x)/2} U_{g',g}^{-1}(x) \sigma_d^{\mathcal{P}_g}(x,\xi) U_{g',g}(x) e^{-h(x)/2}, \quad \xi \in T_x^* M.$	$\sigma_d^{\mathcal{P}_{g'}}(x,\xi) = e^{-h(x)/2} U_{g',g}^{-1}(x) \sigma_d^{\mathcal{P}_g}(x,\xi) U_{g',g}(x) e^{-h(x)/2}, \quad \xi \in T_x^* M.$
cardona-qft-methods_ E00321 (7)	161	■ 0.971	$\begin{aligned} d\,x^{\mu}\left[\partial_{\mu}\phi^a(x)+\omega_{b\mu}^a(x)\phi^b(x)\right]&=dx^{\mu}D_{\mu}\phi^a(x)\\&=D\phi^a(x). \end{aligned}$	$dx^{\mu}\left[\partial_{\mu}\phi^a(x)+\omega_{b\mu}^a(x)\phi^b(x)\right]=dx^{\mu}D_{\mu}\phi^a(x)=D\phi^a(x).$	$dx^{\mu}[\partial_{\mu}\phi^a(x)+\omega_{b\mu}^a(x)\phi^b(x)]=dx^{\mu}D_{\mu}\phi^a(x)=D\phi^a(x).$
cardona-qft-methods_ E00969 (9)	363	■ 0.971	$X_{\{\Gamma\}}=\left\{t\in\mathbb{P}_{\mathbb{P}_k}^{n-1}\mid t_1+t_2+\cdots+t_n=0\right\}=: \mathcal{L}.$	$X_{\Gamma}=\left\{t\in\mathbb{P}_k^{n-1}\mid t_1+t_2+\cdots+t_n=0\right\}=: \mathcal{L}.$	$X_{\Gamma}=\left\{t\in\mathbb{P}_k^{n-1}\mid t_1+t_2+\cdots+t_n=0\right\}=: \mathcal{L}.$
cardona-qft-methods_ E00452	213	■ 0.972	$\begin{gathered} \mathcal{A}=\mathcal{A}_{\{1\}}\otimes\mathcal{A}_{\{2\}}\quad\mathcal{H}=\mathcal{H}_{\{1\}}\otimes\mathcal{H}_{\{2\}}\\ D=D_{\{1\}}\otimes 1+\gamma_1\otimes D_{\{2\}}\\ \gamma=\gamma_1\otimes\gamma_2\quad J=J_1\otimes J_2. \end{gathered}$	$\begin{aligned} \mathcal{A} &= \mathcal{A}_1 \otimes \mathcal{A}_2 \quad \mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2 \\ D &= D_1 \otimes 1 + \gamma_1 \otimes D_2 \\ \gamma &= \gamma_1 \otimes \gamma_2 \quad J = J_1 \otimes J_2 \end{aligned}$	$\begin{aligned} \mathcal{A} &= \mathcal{A}_1 \otimes \mathcal{A}_2 \quad \mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2 \\ D &= D_1 \otimes 1 + \gamma_1 \otimes D_2 \\ \gamma &= \gamma_1 \otimes \gamma_2 \quad J = J_1 \otimes J_2. \end{aligned}$
cardona-qft-methods_ E00971 (11)	363	■ 0.974	$\Psi_{\Gamma^{\vee}}(t)=t_2t_3\cdots t_n+t_1t_3\cdots t_n+\cdots+t_1t_2\cdots\hat{t}_i\cdots t_n+\cdots+t_1t_2\cdots t_{n-1},$	$\Psi_{\Gamma^{\vee}}(t)=t_2t_3\cdots t_n+t_1t_3\cdots t_n+\cdots+t_1t_2\cdots\hat{t}_i\cdots t_n+\cdots+t_1t_2\cdots t_{n-1},$	$\Psi_{\Gamma^{\vee}}(t)=t_2t_3\cdots t_n+t_1t_3\cdots t_n+\cdots+t_1t_2\cdots\hat{t}_i\cdots t_n+\cdots+t_1t_2\cdots t_{n-1},$
cardona-qft-methods_ E00591 (25)	251	■ 0.974	$\begin{aligned} \phi\otimes_{S_{\theta}}\chi&\equiv\left(\frac{1+\tau_0}{2}\right)(\phi\otimes\chi),\\ \phi\otimes_{A_{\theta}}\chi&\equiv\left(\frac{1-\tau_0}{2}\right)(\phi\otimes\chi), \end{aligned}$	$\begin{aligned} \phi\otimes_{S_{\theta}}\chi&\equiv\left(\frac{1+\tau_0}{2}\right)(\phi\otimes\chi),\\ \phi\otimes_{A_{\theta}}\chi&\equiv\left(\frac{1-\tau_0}{2}\right)(\phi\otimes\chi), \end{aligned}$	$\begin{aligned} \phi\otimes_{S_{\theta}}\chi&\equiv\left(\frac{1+\tau_0}{2}\right)(\phi\otimes\chi),\\ \phi\otimes_{A_{\theta}}\chi&\equiv\left(\frac{1-\tau_0}{2}\right)(\phi\otimes\chi), \end{aligned}$
cardona-qft-methods_ E00960 (12)	361	■ 0.974	$\pi_{\{2\}}:\pi_{\{1\}}^{-1}\left(X_{\{\Gamma\}}\backslash\Sigma_n\right)\rightarrow X_{\{\Gamma\}}\backslash\Sigma_n.$	$\pi_2:\pi_1^{-1}\left(X_{\Gamma}\backslash\Sigma_n\right)\rightarrow X_{\Gamma^{\vee}}\backslash\Sigma_n.$	$\pi_2:\pi_1^{-1}\left(X_{\Gamma}\backslash\Sigma_n\right)\rightarrow X_{\Gamma^{\vee}}\backslash\Sigma_n.$
cardona-qft-methods_ E00302 (6)	151	■ 0.975	$\delta I=\int_M\delta\mathcal{C}\wedge\star j=\int_M(d\Omega)\wedge\star j=\int_Md(\Omega\wedge\star j)-\int_M\Omega\wedge d\star j,$	$\delta I=\int_M\delta\mathcal{C}\wedge\star j=\int_M(d\Omega)\wedge\star j=\int_Md(\Omega\wedge\star j)-\int_M\Omega\wedge d\star j,$	$\delta I=\int_M\delta\mathcal{C}\wedge\star j=\int_M(d\Omega)\wedge\star j=\int_Md(\Omega\wedge\star j)-\int_M\Omega\wedge d\star j,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E00358 (28)	171	■ 0.976	$\begin{aligned} C_3^{\text{Lor}} &= \omega_b^a d\omega_a^b + \frac{2}{3} \omega_b^a \omega_c^b \omega_a^c \\ C_3^{\text{Tor}} &= e^a T_a. \end{aligned}$	$C_3^{\text{Lor}} = \omega_b^a d\omega_a^b + \frac{2}{3} \omega_b^a \omega_c^b \omega_a^c$ $C_3^{\text{Tor}} = e^a T_a.$	$C_3^{\text{Lor}} = \omega_b^a d\omega_a^b + \frac{2}{3} \omega_b^a \omega_c^b \omega_a^c$ $C_3^{\text{Tor}} = e^a T_a.$
cardona-qft-methods_ E00027 (8)	27	■ 0.976	$\operatorname{WRes}\left((1+\Delta)^{-d/2}\right)=\operatorname{Vol}\left(\mathbb{S}^{\mathrm{d}-1}\right)=\frac{2\pi^{\mathrm{d}/2}}{\Gamma(\mathrm{d}/2)}.$	$\operatorname{WRes}\left((1+\Delta)^{-d/2}\right)=\operatorname{Vol}\left(\mathbb{S}^{\mathrm{d}-1}\right)=\frac{2\pi^{\mathrm{d}/2}}{\Gamma(\mathrm{d}/2)}.$	$\operatorname{WRes}\left((1+\Delta)^{-d/2}\right)=\operatorname{Vol}\left(\mathbb{S}^{\mathrm{d}-1}\right)=\frac{2\pi^{\mathrm{d}/2}}{\Gamma(\mathrm{d}/2)}.$
cardona-qft-methods_ E00336 (22)	164	■ 0.978	$D\,T^{\{a\}}=R_{\{b\}}^{\{a\}}\wedge e^{\{b\}}\,.$	$DT^a = R_b^a \wedge e^b.$	$DT^a = R_b^a \wedge e^b.$
cardona-qft-methods_ E00925	352	■ 0.978	$\left(P_G M\times_G EG_n\right)^{-e_0^*TM_n}.$	$\left(P_G M\times_G EG_n\right)^{-e_0^*TM_n}.$	$\left(P_G M\times_G EG_n\right)^{-e_0^*TM_n}.$
cardona-qft-methods_ E00303 (7)	152	■ 0.979	$\begin{aligned} \mathbf{A} &= \mathbf{A}_\mu dx^\mu \\ &= A_\mu^a \mathbf{K}_a dx^\mu \end{aligned}$	$\mathbf{A} = \mathbf{A}_\mu dx^\mu$ $= A_\mu^a \mathbf{K}_a dx^\mu$	$\mathbf{A} = \mathbf{A}_\mu dx^\mu$ $= A_\mu^a \mathbf{K}_a dx^\mu,$
cardona-qft-methods_ E00489	221	■ 0.980	$\int \frac{1}{2} D_\mu \mathbf{H} ^2 \sqrt{g} d^4 x;$	$\int \frac{1}{2} D_\mu \mathbf{H} ^2 \sqrt{g} d^4 x;$	$\int \frac{1}{2} D_\mu \mathbf{H} ^2 \sqrt{g} d^4 x;$
cardona-qft-methods_ E00268	127	■ 0.981	$\Psi_{G_1}(t_1,\ldots,t_{n_1})\neq 0 \quad \text{and} \quad \Psi_{G_2}(u_1,\ldots,u_{n_2})\neq 0,$	$\Psi_{G_1}(t_1,\ldots,t_{n_1})\neq 0 \quad \text{and} \quad \Psi_{G_2}(u_1,\ldots,u_{n_2})\neq 0,$	$\Psi_{G_1}(t_1,\ldots,t_{n_1})\neq 0 \quad \text{and} \quad \Psi_{G_2}(u_1,\ldots,u_{n_2})\neq 0 \quad ,$
cardona-qft-methods_ E00095	56	■ 0.983	$g\left(t\,\mathrm{mathcal{D}}^2\right)=\int_0^\infty e^{-st\mathcal{D}^2}\phi(s)ds,\mathrm{Tr}\left(g\left(t\mathcal{D}^2\right)\right)\underset{t\downarrow 0}{\sim}\sum_{\alpha\in\mathrm{Sp}^+}a_\alpha t^\alpha\int_0^\infty s^\alpha g(s)ds$	$g\left(t\mathcal{D}^2\right)=\int_0^\infty e^{-st\mathcal{D}^2}\phi(s)ds,\mathrm{Tr}\left(g\left(t\mathcal{D}^2\right)\right)\underset{t\downarrow 0}{\sim}\sum_{\alpha\in\mathrm{Sp}^+}a_\alpha t^\alpha\int_0^\infty s^\alpha g(s)ds$	$g(t\mathcal{D}^2)=\int_0^\infty e^{-st\mathcal{D}^2}\phi(s)ds,\mathrm{Tr}\left(g(t\mathcal{D}^2)\right)\underset{t\downarrow 0}{\sim}\sum_{\alpha\in\mathrm{Sp}^+}a_\alpha t^\alpha\int_0^\infty s^\alpha g(s)ds.$
cardona-qft-methods_ E00379 (20)	177	■ 0.983	$L_4=\epsilon_{abcd}R^{ab}e^ce^d\,.$	$L_4 = \epsilon_{abcd}R^{ab}e^ce^d.$	$L_4 = \epsilon_{abcd}R^{ab}e^ce^d.$
cardona-qft-methods_ E00490	221	■ 0.983	$\frac{1}{4}G_{\mu\nu}^i\bar{G}^{\mu\nu i}+\frac{1}{4}F_{\mu\nu}^\alpha\bar{F}^{\mu\nu\alpha}+\frac{1}{4}B_{\mu\nu}\bar{B}^{\mu\nu}.$	$\frac{1}{4}G_{\mu\nu}^i\bar{G}^{\mu\nu i}+\frac{1}{4}F_{\mu\nu}^\alpha\bar{F}^{\mu\nu\alpha}+\frac{1}{4}B_{\mu\nu}\bar{B}^{\mu\nu}.$	$\frac{1}{4}G_{\mu\nu}^i\bar{G}^{\mu\nu i}+\frac{1}{4}F_{\mu\nu}^\alpha\bar{F}^{\mu\nu\alpha}+\frac{1}{4}B_{\mu\nu}\bar{B}^{\mu\nu}.$
cardona-qft-methods_ E00698	271	■ 0.983	$\hat{H}=\sum_{l=1}^L\mathcal{H}_{l,l+1}.$	$\hat{H} = \sum_{l=1}^L \mathcal{H}_{l,l+1}.$	$\hat{H} = \sum_{l=1}^L \mathcal{H}_{l,l+1}.$
cardona-qft-methods_ E00368 (7)	175	■ 0.984	$\begin{aligned} \delta\omega^{ab} &= d\lambda^{ab} + \omega_c^a \lambda^{cb} + \omega_c^b \lambda^{ac} \pm [e^a \lambda^b - \lambda^a e^b] l^{-2} \\ \delta e^a &= d\lambda^a + \omega_b^a \lambda^b - \lambda_b^a e^b. \end{aligned}$	$\delta\omega^{ab} = d\lambda^{ab} + \omega_c^a \lambda^{cb} + \omega_c^b \lambda^{ac} \pm [e^a \lambda^b - \lambda^a e^b] l^{-2}$ $\delta e^a = d\lambda^a + \omega_b^a \lambda^b - \lambda_b^a e^b.$	$\delta\omega^{ab} = d\lambda^{ab} + \omega_c^a \lambda^{cb} + \omega_c^b \lambda^{ac} \pm [e^a \lambda^b - \lambda^a e^b] l^{-2}$ $\delta e^a = d\lambda^a + \omega_b^a \lambda^b - \lambda_b^a e^b.$
cardona-qft-methods_ E00730	277	■ 0.984	$\left(\frac{u_k+\frac{i}{2}}{u_k-\frac{i}{2}}\right)^L=\prod_{\substack{j=1\\j\neq k}}^M\frac{u_k-u_j+i}{u_k-u_j-i}.$	$\left(\frac{u_k+\frac{i}{2}}{u_k-\frac{i}{2}}\right)^L=\prod_{\substack{j=1\\j\neq k}}^M\frac{u_k-u_j+i}{u_k-u_j-i}.$	$\left(\frac{u_k+\frac{i}{2}}{u_k-\frac{i}{2}}\right)^L=\prod_{\substack{j=1\\j\neq k}}^M\frac{u_k-u_j+i}{u_k-u_j-i}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0781 (22)	298	■ 0.984	$X^{\mathrm{M}}(\tau,\sigma)=x^{\mathrm{M}}+2\,\alpha^{\mathrm{\prime}}\,p^{\mathrm{M}}\,\tau+\mathrm{i}\,(2\alpha^{\mathrm{\prime}})^{1/2}\sum_{n\neq 0}\frac{1}{n}\alpha_n^{\mathrm{M}}e^{-\mathrm{i}n\tau}\cos(n\sigma)\;.$	$X^M(\tau,\sigma)=x^M+2\alpha'p^M\tau+\mathrm{i}\,(2\alpha')^{1/2}\sum_{n\neq 0}\frac{1}{n}\alpha_n^Me^{-\mathrm{i}n\tau}\cos(n\sigma).$	$X^M(\tau,\sigma)=x^M+2\alpha'p^M\tau+\mathrm{i}\,(2\alpha')^{1/2}\sum_{n\neq 0}\frac{1}{n}\alpha_n^Me^{-\mathrm{i}n\tau}\cos(n\sigma)\;.$
cardona-qft-methods_ EQ0434	208	■ 0.985	$\mathcal{E}=\mathbf{2}_L\otimes\mathbf{1}^0\oplus\mathbf{2}_R\otimes\mathbf{1}^0\oplus\mathbf{2}_L\otimes\mathbf{3}^0\oplus\mathbf{2}_R\otimes\mathbf{3}^0,$	$\mathcal{E}=\mathbf{2}_L\otimes\mathbf{1}^0\oplus\mathbf{2}_R\otimes\mathbf{1}^0\oplus\mathbf{2}_L\otimes\mathbf{3}^0\oplus\mathbf{2}_R\otimes\mathbf{3}^0,$	$\mathcal{E}=\mathbf{2}_L\otimes\mathbf{1}^0\oplus\mathbf{2}_R\otimes\mathbf{1}^0\oplus\mathbf{2}_L\otimes\mathbf{3}^0\oplus\mathbf{2}_R\otimes\mathbf{3}^0,$
cardona-qft-methods_ EQ0485	221	■ 0.985	$\begin{aligned} &\mathfrak{a}=\operatorname{Tr}\left(Y_{\nu}^{\dagger}Y_{\nu}+Y_e^{\dagger}Y_e+3\left(Y_u^{\dagger}Y_u+Y_d^{\dagger}Y_d\right)\right.\\ &\left.\left(Y_{\nu}^{\dagger}Y_{\nu}\right)^2+\left(Y_e^{\dagger}Y_e\right)^2+3\left(Y_u^{\dagger}Y_u\right)^2+3\left(Y_d^{\dagger}Y_d\right)^2\right)\\ &\mathfrak{b}=\operatorname{Tr}\left(MM^{\dagger}\right)\\ &\mathfrak{d}=\operatorname{Tr}\left(MM^{\dagger}\right)^2\\ &\mathfrak{e}=\operatorname{Tr}\left(MM^{\dagger}Y_{\nu}^{\dagger}Y_{\nu}\right). \end{aligned}$	$\begin{aligned} \mathfrak{a} &= \operatorname{Tr}\left(Y_{\nu}^{\dagger}Y_{\nu}+Y_e^{\dagger}Y_e+3\left(Y_u^{\dagger}Y_u+Y_d^{\dagger}Y_d\right)\right)\\ \mathfrak{b} &= \operatorname{Tr}\left(\left(Y_{\nu}^{\dagger}Y_{\nu}\right)^2+\left(Y_e^{\dagger}Y_e\right)^2+3\left(Y_u^{\dagger}Y_u\right)^2+3\left(Y_d^{\dagger}Y_d\right)^2\right)\\ \mathfrak{c} &= \operatorname{Tr}\left(MM^{\dagger}\right)\\ \mathfrak{d} &= \operatorname{Tr}\left(MM^{\dagger}\right)^2\\ \mathfrak{e} &= \operatorname{Tr}\left(MM^{\dagger}Y_{\nu}^{\dagger}Y_{\nu}\right). \end{aligned}$	$\begin{aligned} \mathfrak{a} &= \operatorname{Tr}(Y_{\nu}^{\dagger}Y_{\nu}+Y_e^{\dagger}Y_e+3(Y_u^{\dagger}Y_u+Y_d^{\dagger}Y_d))\\ \mathfrak{b} &= \operatorname{Tr}((Y_{\nu}^{\dagger}Y_{\nu})^2+(Y_e^{\dagger}Y_e)^2+3(Y_u^{\dagger}Y_u)^2+3(Y_d^{\dagger}Y_d)^2)\\ \mathfrak{c} &= \operatorname{Tr}(MM^{\dagger})\\ \mathfrak{d} &= \operatorname{Tr}(MM^{\dagger})^2\\ \mathfrak{e} &= \operatorname{Tr}(MM^{\dagger}Y_{\nu}^{\dagger}Y_{\nu}). \end{aligned}$
cardona-qft-methods_ EQ0085 (30)	49	■ 0.985	$\begin{aligned} &\mathfrak{f}\left D_A\right ^{-d}=f\left D\right ^{-d},\\ &\mathfrak{f}\left D_A\right ^{-(d-1)}=f\left D\right ^{-(d-1)}-\left(\frac{d-1}{2}\right)fX\left D\right ^{-d-1},\\ &\mathfrak{f}\left D_A\right ^{-(d-2)}=f\left D\right ^{-(d-2)}+\frac{d-2}{2}\left(-fX\left D\right ^{-d}+\frac{d}{4}fX^2\left D\right ^{-2-d}\right). \end{aligned}$	$\begin{aligned} f\left D_A\right ^{-d} &= f\left D\right ^{-d},\\ f\left D_A\right ^{-(d-1)} &= f\left D\right ^{-(d-1)}-\left(\frac{d-1}{2}\right)fX\left D\right ^{-d-1},\\ f\left D_A\right ^{-(d-2)} &= f\left D\right ^{-(d-2)}+\frac{d-2}{2}\left(-fX\left D\right ^{-d}+\frac{d}{4}fX^2\left D\right ^{-2-d}\right). \end{aligned}$	$\begin{aligned} \mathfrak{f}\left D_A\right ^{-d} &= \mathfrak{f}\left D\right ^{-d},\\ \mathfrak{f}\left D_A\right ^{-(d-1)} &= \mathfrak{f}\left D\right ^{-(d-1)}-\left(\frac{d-1}{2}\right)\mathfrak{f}X\left D\right ^{-d-1},\\ \mathfrak{f}\left D_A\right ^{-(d-2)} &= \mathfrak{f}\left D\right ^{-(d-2)}+\frac{d-2}{2}\left(-\mathfrak{f}X\left D\right ^{-d}+\frac{d}{4}\mathfrak{f}X^2\left D\right ^{-2-d}\right). \end{aligned}$
cardona-qft-methods_ EQ0250 (1)	118	■ 0.986	$S_{\mathrm{\scriptscriptstyle eff}}(\phi)=\sum_G\langle G\rangle$	$S_{\mathrm{eff}}(\phi)=\sum_G\langle G\rangle$	$S_{\mathrm{eff}}(\phi)=\sum_G\langle G\rangle$
cardona-qft-methods_ EQ0102	61	■ 0.986	$U_{\mathrm{k}}U_{\mathrm{q}}=\mathrm{e}^{-\frac{\mathrm{i}}{2}\mathrm{k}\cdot\Theta\mathrm{q}}U_{\mathrm{k}+\mathrm{q}},\text{ and }U_{\mathrm{k}}U_{\mathrm{q}}=\mathrm{e}^{-\mathrm{i}\mathrm{k}\cdot\Theta\mathrm{q}}U_{\mathrm{q}}U_{\mathrm{k}}$	$U_kU_q=e^{-\frac{\mathrm{i}}{2}k\cdot\Theta q}U_{k+q},\text{ and }U_kU_q=e^{-\mathrm{i}k\cdot\Theta q}U_qU_k$	$U_kU_q=e^{-\frac{\mathrm{i}}{2}k\cdot\Theta q}U_{k+q},\text{ and }U_kU_q=e^{-\mathrm{i}k\cdot\Theta q}U_qU_k$
cardona-qft-methods_ EQ0445	211	■ 0.986	$S_{\mathrm{3}}=\left(\begin{array}{cccc} 0&0&Y_{(\uparrow 3)}^*&0\\ 0&0&0&Y_{(\downarrow 3)}^*\\ Y_{(\uparrow 3)}&0&0&0\\ 0&Y_{(\downarrow 3)}&0&0 \end{array}\right)$	$S_3=\left(\begin{array}{cccc} 0&0&Y_{(\uparrow 3)}^*&0\\ 0&0&0&Y_{(\downarrow 3)}^*\\ Y_{(\uparrow 3)}&0&0&0\\ 0&Y_{(\downarrow 3)}&0&0 \end{array}\right)$	$S_3=\left(\begin{array}{cccc} 0&0&Y_{(\uparrow 3)}^*&0\\ 0&0&0&Y_{(\downarrow 3)}^*\\ Y_{(\uparrow 3)}&0&0&0\\ 0&Y_{(\downarrow 3)}&0&0 \end{array}\right)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0441	210	■ 0.987	$\begin{array}{ccccc} & \uparrow & \otimes \mathbf{1}^0 & \downarrow & \otimes \mathbf{1}^0 & \uparrow & \otimes \mathbf{3}^0 & \downarrow & \otimes \mathbf{3}^0 \\ \mathbf{2}_L & & -1 & & -1 & & \frac{1}{3} & & \frac{1}{3} \\ \mathbf{2}_R & & 0 & & -2 & & \frac{4}{3} & & -\frac{2}{3} \end{array}$		
cardona-qft-methods_ EQ0678 (26)	46	■ 0.987	$\mathrm{cal}_D\{\widetilde{A}\}=\mathrm{cal}_D\{A\},$ $\quad \widetilde{A}=A+\epsilon JAJ^{-1}, \quad A=A^*.$	$\mathcal{D}_{\widetilde{A}}=\mathcal{D}+\widetilde{A}, \quad \widetilde{A}=A+\epsilon JAJ^{-1}, \quad A=A^*.$	$\mathcal{D}_{\widetilde{A}}=\mathcal{D}+\widetilde{A}, \quad \widetilde{A}=A+\epsilon JAJ^{-1}, \quad A=A^*.$
cardona-qft-methods_ EQ0164	88	■ 0.988	$\mathrm{t}\text{-}\mathrm{Ind}_G:K_G(M)\rightarrow K_G(\{pt\})\simeq R(G).$	$\mathrm{t}\text{-}\mathrm{Ind}_G:K_G(M)\rightarrow K_G(\{pt\})\simeq R(G).$	$\mathrm{t}\text{-}\mathrm{Ind}_G:K_G(M)\rightarrow K_G(\{pt\})\simeq R(G).$
cardona-qft-methods_ EQ0956 (8)	361	■ 0.988	$\Psi_{\Gamma}(t)=\left(\prod_{i=1}^nt_i\right)\Psi_{\Gamma^{\vee}}\left(\frac{1}{t}\right),$	$\Psi_{\Gamma}(t)=\left(\prod_{i=1}^nt_i\right)\Psi_{\Gamma^{\vee}}\left(\frac{1}{t}\right),$	$\Psi_{\Gamma}(t)=\left(\prod_{i=1}^nt_i\right)\Psi_{\Gamma^{\vee}}\left(\frac{1}{t}\right),$
cardona-qft-methods_ EQ0655 (5)	262	■ 0.988	$\begin{aligned} \langle \vartheta \rho_{\theta}(x) \eta_{\theta}(y) 0 \rangle &= e^{-\frac{1}{2} \frac{\partial}{\partial x^{\mu}} \theta^{\mu \nu} \frac{\partial}{\partial y^{\nu}}} \langle 0 \rho_0(x) \eta_0(y) 0 \rangle \\ &= e^{-\frac{1}{2} \frac{\partial}{\partial x^{\mu}} \theta^{\mu \nu} \frac{\partial}{\partial y^{\nu}}} F(x-y) = F(x-y) \\ &= \langle 0 \rho_0(x) \eta_0(y) 0 \rangle \end{aligned}$	$\begin{aligned} \langle 0 \rho_{\theta}(x) \eta_{\theta}(y) 0 \rangle &= e^{-\frac{1}{2} \frac{\partial}{\partial x^{\mu}} \theta^{\mu \nu} \frac{\partial}{\partial y^{\nu}}} \langle 0 \rho_0(x) \eta_0(y) 0 \rangle \\ &= e^{-\frac{1}{2} \frac{\partial}{\partial x^{\mu}} \theta^{\mu \nu} \frac{\partial}{\partial y^{\nu}}} F(x-y) = F(x-y) \\ &= \langle 0 \rho_0(x) \eta_0(y) 0 \rangle \end{aligned}$	$\begin{aligned} \langle 0 \rho_{\theta}(x) \eta_{\theta}(y) 0 \rangle &= e^{-\frac{1}{2} \frac{\partial}{\partial x^{\mu}} \theta^{\mu \nu} \frac{\partial}{\partial y^{\nu}}} \langle 0 \rho_0(x) \eta_0(y) 0 \rangle \\ &= e^{-\frac{1}{2} \frac{\partial}{\partial x^{\mu}} \theta^{\mu \nu} \frac{\partial}{\partial y^{\nu}}} F(x-y) = F(x-y) \\ &= \langle 0 \rho_0(x) \eta_0(y) 0 \rangle \end{aligned}$
cardona-qft-methods_ EQ0375 (16)	176	■ 0.989	$\Upsilon_0^{AB}=\left[\begin{array}{cc}:\eta^{ab}&0\\0&0\end{array}\right],$	$\Upsilon_0^{AB}=\left[\begin{array}{cc}:\eta^{ab}&0\\0&0\end{array}\right],$	$\Upsilon_0^{AB}=\left[\begin{array}{cc}:\eta^{ab}&0\\0&0\end{array}\right],$
cardona-qft-methods_ EQ0884	343	■ 0.989	$P\cdot D:H_p(M)\cong\left(H^p(M)\right)^*\cong H^{n-p}(M).$	$P.D:H_p(M)\cong\left(H^p(M)\right)^*\cong H^{n-p}(M).$	$P.D:H_p(M)\cong\left(H^p(M)\right)^*\cong H^{n-p}(M).$
cardona-qft-methods_ EQ0937	354	■ 0.989	$H_{\ast}^{\ast}(\mathrm{L}\,\mathrm{B}\,\mathrm{G}^{-T\,\mathrm{B}\,\mathrm{G}})\cong H^{\ast}(\mathrm{L}\,\mathrm{B}\,\mathrm{G}).$	$H_{\ast}^{pro}\left(LBG^{-TBG}\right)\cong H^{\ast}(LBG).$	$H_{\ast}^{pro}\left(LBG^{-TBG}\right)\cong H^{\ast}(LBG).$
cardona-qft-methods_ EQ0897	347	■ 0.989	$\begin{aligned} & H\,H_{\ast}(A,M)\cong \\ & \operatorname{Tor}_{A\otimes A^{op}}^{\ast}(A,M)\,\,\& \\ & H\,H^{\ast}(A,M)\cong \operatorname{Ext}_{A\otimes A^{op}}^{\ast}(A,M) \end{aligned}$	$\begin{aligned} HH_{\ast}(A,M) &\cong \operatorname{Tor}_{A\otimes A^{op}}^{\ast}(A,M) \\ HH^{\ast}(A,M) &\cong \operatorname{Ext}_{A\otimes A^{op}}^{\ast}(A,M) \end{aligned}$	$\begin{aligned} HH_{\ast}(A,M) &\cong \operatorname{Tor}_{A\otimes A^{op}}^{\ast}(A,M) \\ HH^{\ast}(A,M) &\cong \operatorname{Ext}_{A\otimes A^{op}}^{\ast}(A,M) \end{aligned}$
cardona-qft-methods_ EQ0513 (6)	230	■ 0.990	$\beta_j^{\mathrm{grav}}=\frac{a_j}{8\pi}\frac{\Lambda^2}{M_P^2(\Lambda)}x_j,$	$\beta_j^{\mathrm{grav}}=\frac{a_j}{8\pi}\frac{\Lambda^2}{M_P^2(\Lambda)}x_j,$	$\beta_j^{\mathrm{grav}}=\frac{a_j}{8\pi}\frac{\Lambda^2}{M_P^2(\Lambda)}x_j,$
cardona-qft-methods_ EQ0002	18	■ 0.990	$\left \partial_x^\alpha\partial_y^\gamma\partial_\xi^\beta a(x,y,\xi)\right \leq C_{K\alpha\beta\gamma}(1+ \xi)^{\Re(m)- \beta },\quad\forall x,y\in K,\xi\in\mathbb{R}^d$	$\left \partial_x^\alpha\partial_y^\gamma\partial_\xi^\beta a(x,y,\xi)\right \leq C_{K\alpha\beta\gamma}(1+ \xi)^{\Re(m)- \beta },\quad\forall x,y\in K,\xi\in\mathbb{R}^d$	$ \partial_x^\alpha\partial_y^\gamma\partial_\xi^\beta a(x,y,\xi) \leq C_{K\alpha\beta\gamma}(1+ \xi)^{\Re(m)- \beta },\quad\forall x,y\in K,\xi\in\mathbb{R}^d.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0199 (9)	95	■ 0.990	$\mathcal{D}=\left(\begin{array}{ll} 0 & 1 \\ 1 & 0 \end{array}\right)\frac{1}{i}\frac{\partial}{\partial x_1}+\left(\begin{array}{ll} 0 & -i \\ i & 0 \end{array}\right)\frac{1}{i}\frac{\partial}{\partial x_2}+\left(\begin{array}{ll} 1 & 0 \\ 0 & -1 \end{array}\right)\frac{1}{i}\frac{\partial}{\partial x_3}.$	$\mathcal{D}=\left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right)\frac{1}{i}\frac{\partial}{\partial x_1}+\left(\begin{array}{cc} 0 & -i \\ i & 0 \end{array}\right)\frac{1}{i}\frac{\partial}{\partial x_2}+\left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right)\frac{1}{i}\frac{\partial}{\partial x_3}.$	$\mathcal{D}=\left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right)\frac{1}{i}\frac{\partial}{\partial x_1}+\left(\begin{array}{cc} 0 & -i \\ i & 0 \end{array}\right)\frac{1}{i}\frac{\partial}{\partial x_2}+\left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right)\frac{1}{i}\frac{\partial}{\partial x_3}.$
cardona-qft-methods_ EQ0301 (5)	151	■ 0.990	$I=\int \mathcal{C}\wedge \star j,$	$I=\int \mathcal{C}\wedge \star j,$	$I=\int \mathcal{C}\wedge \star j,$
cardona-qft-methods_ EQ0409 (39)	187	■ 0.991	$B_{2n}=-\kappa n\int_0^1 dt\int_0^t ds\,\epsilon\theta e\left(\widetilde{R}+t^2\theta^2+s^2e^2\right)^{n-1}$	$B_{2n}=-\kappa n\int_0^1 dt\int_0^t ds\epsilon\theta e\left(\widetilde{R}+t^2\theta^2+s^2e^2\right)^{n-1}$	$B_{2n}=-\kappa n\int_0^1 dt\int_0^t ds\,\epsilon\theta e\left(\widetilde{R}+t^2\theta^2+s^2e^2\right)^{n-1},$
cardona-qft-methods_ EQ0787 (28)	299	■ 0.991	$\begin{array}{rc} \mathrm{R}: & \Psi^M(\tau,0)=\Psi^M(\tau,2\pi) \\ \mathrm{NS}: & \Psi^M(\tau,0)=-\Psi^M(\tau,2\pi) \end{array}$	$\begin{array}{l} \mathrm{R}: \quad \Psi^M(\tau,0)=\Psi^M(\tau,2\pi) \\ \mathrm{NS}: \quad \Psi^M(\tau,0)=-\Psi^M(\tau,2\pi) \end{array}$	$\begin{array}{l} \mathrm{R}: \quad \Psi^M(\tau,0)=\Psi^M(\tau,2\pi) \\ \mathrm{NS}: \quad \Psi^M(\tau,0)=-\Psi^M(\tau,2\pi) \end{array}$
cardona-qft-methods_ EQ0995	374	■ 0.991	$\Delta X^{\alpha}=\sum_{\beta\leq\alpha}\binom{\alpha}{\beta}X^{\beta}\otimes X^{\alpha-\beta}$	$\Delta X^{\alpha}=\sum_{\beta\leq\alpha}\binom{\alpha}{\beta}X^{\beta}\otimes X^{\alpha-\beta}$	$\Delta X^{\alpha}=\sum_{\beta\leq\alpha}\binom{\alpha}{\beta}X^{\beta}\otimes X^{\alpha-\beta},$
cardona-qft-methods_ EQ1003	375	■ 0.991	$\begin{aligned} f_{\{a\}b}(x) &= \sum_{\alpha}\frac{x^{\alpha}}{\alpha!}a\,b\left(X^{\alpha}\right)=\sum_{\alpha}\frac{x^{\alpha}}{\alpha!}(a\otimes b)\left(\Delta X^{\alpha}\right) \\ &= \sum_{\alpha}\frac{x^{\alpha}}{\alpha!}(a\otimes b)\left(\sum_{\beta\leq\alpha}\binom{\alpha}{\beta}X^{\beta}\otimes X^{\alpha-\beta}\right) \\ &= \sum_{\beta}\sum_{\gamma}\frac{x^{\beta}x^{\gamma}}{\beta!\gamma!}a\left(X^{\beta}\right)b\left(X^{\gamma}\right)=f_a(x)f_b(x). \end{aligned}$	$\begin{aligned} f_{ab}(x) &= \sum_{\alpha}\frac{x^{\alpha}}{\alpha!}ab\left(X^{\alpha}\right)=\sum_{\alpha}\frac{x^{\alpha}}{\alpha!}(a\otimes b)\left(\Delta X^{\alpha}\right) \\ &= \sum_{\alpha}\frac{x^{\alpha}}{\alpha!}(a\otimes b)\left(\sum_{\beta\leq\alpha}\binom{\alpha}{\beta}X^{\beta}\otimes X^{\alpha-\beta}\right) \\ &= \sum_{\beta}\sum_{\gamma}\frac{x^{\beta}x^{\gamma}}{\beta!\gamma!}a\left(X^{\beta}\right)b\left(X^{\gamma}\right)=f_a(x)f_b(x). \end{aligned}$	$\begin{aligned} f_{ab}(x) &= \sum_{\alpha}\frac{x^{\alpha}}{\alpha!}ab\left(X^{\alpha}\right)=\sum_{\alpha}\frac{x^{\alpha}}{\alpha!}(a\otimes b)\left(\Delta X^{\alpha}\right) \\ &= \sum_{\alpha}\frac{x^{\alpha}}{\alpha!}(a\otimes b)\left(\sum_{\beta\leq\alpha}\binom{\alpha}{\beta}X^{\beta}\otimes X^{\alpha-\beta}\right) \\ &= \sum_{\beta}\sum_{\gamma}\frac{x^{\beta}x^{\gamma}}{\beta!\gamma!}a\left(X^{\beta}\right)b\left(X^{\gamma}\right)=f_a(x)f_b(x). \end{aligned}$
cardona-qft-methods_ EQ0118 (6)	76	■ 0.991	$\left\{\bigoplus_{V\in\operatorname{Irr}(G)}m_VV\mid m_V\in\mathbb{Z},\,\text{only finitely many numbers }m_V\neq 0\right\}.$	$\left\{\bigoplus_{V\in\operatorname{Irr}(G)}m_VV\mid m_V\in\mathbb{Z},\,\text{only finitely many numbers }m_V\neq 0\right\}.$	$\left\{\bigoplus_{V\in\operatorname{Irr}(G)}m_VV\mid m_V\in\mathbb{Z},\,\text{only finitely many numbers }m_V\neq 0\right\}.$
cardona-qft-methods_ EQ0835 (14)	317	■ 0.991	$\begin{aligned} &e^{-2A+\phi}(\mathrm{d}-H\wedge)(e^{2A-\phi}\Phi_+)=\mathrm{d}A\wedge\bar{\Phi}_+-\frac{\mathrm{i}}{16}e^{\phi+A}*F_{\mathrm{IIB}-}\\ &e^{-2A+\phi}(\mathrm{d}-H\wedge)(e^{2A-\phi}\Phi_-)=0 \end{aligned}$	$\begin{aligned} e^{-2A+\phi}(\mathrm{d}-H\wedge)(e^{2A-\phi}\Phi_+) &= \mathrm{d}A\wedge\bar{\Phi}_+-\frac{\mathrm{i}}{16}e^{\phi+A}*F_{\mathrm{IIB}-}\\ e^{-2A+\phi}(\mathrm{d}-H\wedge)(e^{2A-\phi}\Phi_-) &= 0 \end{aligned}$	$\begin{aligned} e^{-2A+\phi}(\mathrm{d}-H\wedge)(e^{2A-\phi}\Phi_+) &= \mathrm{d}A\wedge\bar{\Phi}_+-\frac{\mathrm{i}}{16}e^{\phi+A}*F_{\mathrm{IIB}-}\\ e^{-2A+\phi}(\mathrm{d}-H\wedge)(e^{2A-\phi}\Phi_-) &= 0, \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0088	50	■ 0.992	$\operatorname{Tr}\!\left(P\,e^{\{-t\,\not{D}^{\{2\}}\}}\right)$ $\underset{t\,\downarrow 0^+}{\sim}\sum_{k=0}^{\infty}a_kt^{(k-\operatorname{ord}(P)-d)/2}+\sum_{k=0}^{\infty}(-a'_k\log t+b_k)t^k$	$\operatorname{Tr}\!\left(P\,e^{-t\mathcal{D}^2}\right)_{t\,\downarrow 0^+}\sim\sum_{k=0}^{\infty}a_kt^{(k-\operatorname{ord}(P)-d)/2}+\sum_{k=0}^{\infty}(-a'_k\log t+b_k)t^k,$	
cardona-qft-methods_ EQ0502	226	■ 0.992	$\begin{gathered} \zeta=\exp\left(2\,\pi\,i\,/\,3\right)\,\,\,U_{\{P\,M\,N\,S\}}\!\left(\Lambda_{unif}\right)=\frac{1}{3}\!\left(\begin{array}{ccc} 1&\zeta&\zeta^2\\ \zeta&1&\zeta\\ \zeta^2&\zeta&1 \end{array}\right) \end{gathered}$	$\zeta=\exp(2\pi i/3)$ $U_{PMNS}\left(\Lambda_{unif}\right)=\frac{1}{3}\left(\begin{array}{ccc} 1&\zeta&\zeta^2\\ \zeta&1&\zeta\\ \zeta^2&\zeta&1 \end{array}\right)$	$\zeta=\exp(2\pi i/3)$ $U_{PMNS}(\Lambda_{unif})=\frac{1}{3}\left(\begin{array}{ccc} 1&\zeta&\zeta^2\\ \zeta&1&\zeta\\ \zeta^2&\zeta&1 \end{array}\right)$
cardona-qft-methods_ EQ0141 (5)	81	■ 0.992	$\sigma_L(\mathcal{D})(x_0,\xi)\,e:=\lim_{t\rightarrow\infty}t^{-k}\mathcal{D}\left(e^{itf(x)}s(x)\right)\Big _{x=x_0}.$	$\sigma_L(\mathcal{D})(x_0,\xi)\,e:=\lim_{t\rightarrow\infty}t^{-k}\mathcal{D}\left(e^{itf(x)}s(x)\right)\Big _{x=x_0}.$	$\sigma_L(\mathcal{D})(x_0,\xi)\,e\, :=\, \lim_{t\rightarrow\infty}t^{-k}\mathcal{D}\big(e^{itf(x)}s(x)\big)\Big _{x=x_0}.$
cardona-qft-methods_ EQ0461	216	■ 0.992	$V^{\prime}=-V-\frac{1}{3}\left(\begin{array}{ccc} \Lambda & 0 & 0 \\ 0 & \Lambda & 0 \\ 0 & 0 & \Lambda \end{array}\right)=-V-\frac{1}{3}\Lambda 1_3.$	$V'=-V-\frac{1}{3}\left(\begin{array}{ccc} \Lambda & 0 & 0 \\ 0 & \Lambda & 0 \\ 0 & 0 & \Lambda \end{array}\right)=-V-\frac{1}{3}\Lambda 1_3.$	$V'=-V-\frac{1}{3}\left(\begin{array}{ccc} \Lambda & 0 & 0 \\ 0 & \Lambda & 0 \\ 0 & 0 & \Lambda \end{array}\right)=-V-\frac{1}{3}\Lambda 1_3.$
cardona-qft-methods_ EQ0372 (12)	176	■ 0.993	$\Upsilon^{AB}=\left[\begin{array}{cc} \eta^{ab} & 0 \\ 0 & \mp 1 \end{array}\right],$	$\Upsilon^{AB}=\left[\begin{array}{cc} \eta^{ab} & 0 \\ 0 & \mp 1 \end{array}\right],$	$\Upsilon^{AB}=\left[\begin{array}{cc} \eta^{ab} & 0 \\ 0 & \mp 1 \end{array}\right],$
cardona-qft-methods_ EQ0374 (14)	176	■ 0.993	$\begin{aligned} \delta\omega^{ab}&=d\lambda^{ab}+\omega_c^a\lambda^{cb}+\omega_c^b\lambda^{ac}=D_\omega\lambda^{ab} \\ \delta e^a&=d\lambda^a+\omega_b^a\lambda^b-\lambda_b^ae^b=D_\omega\lambda^a. \end{aligned}$	$\delta\omega^{ab}=d\lambda^{ab}+\omega_c^a\lambda^{cb}+\omega_c^b\lambda^{ac}=D_\omega\lambda^{ab}$ $\delta e^a=d\lambda^a+\omega_b^a\lambda^b-\lambda_b^ae^b=D_\omega\lambda^a$	$\delta\omega^{ab}=d\lambda^{ab}+\omega_c^a\lambda^{cb}+\omega_c^b\lambda^{ac}=D_\omega\lambda^{ab}$ $\delta e^a=d\lambda^a+\omega_b^a\lambda^b-\lambda_b^ae^b=D_\omega\lambda^a.$
cardona-qft-methods_ EQ0605 (14)	253	■ 0.993	$e_{\{p\}}(x)=\exp\,-i\,p\cdot x,\quad\left p_{\{0\}}\right =\left \mathbf{p}^2+m^2\right ^{\frac{1}{2}},\quad d\mu(p)=\frac{d^dp}{2\left p_0\right },$	$e_p(x)=\exp -ip\cdot x,\quad p_0 =(\mathbf{p}^2+m^2)^{\frac{1}{2}},\quad d\mu(p)=\frac{d^dp}{2 p_0 },$	$e_p(x)=\exp -ip\cdot x,\quad p_0 =(\mathbf{p}^2+m^2)^{\frac{1}{2}},\quad d\mu(p)=\frac{d^dp}{2 p_0 },$
cardona-qft-methods_ EQ0740 (8)	282	■ 0.993	$\begin{aligned} Q\cdot\varphi&\sim\psi,\quad\bar{Q}\cdot\varphi\sim\bar{\psi} \\ Q\cdot\psi&\sim F+g_{YM}[\varphi,\varphi],\quad\bar{Q}\cdot\psi\sim\mathcal{D}\varphi \end{aligned}$	$Q\cdot\varphi\sim\psi,\quad\bar{Q}\cdot\varphi\sim\bar{\psi}$ $Q\cdot\psi\sim F+g_{YM}[\varphi,\varphi],\quad\bar{Q}\cdot\psi\sim\mathcal{D}\varphi$	$Q\cdot\varphi\sim\psi\qquad,\qquad\bar{Q}\cdot\varphi\sim\bar{\psi}$ $Q\cdot\psi\sim F+g_{YM}[\varphi,\varphi]\qquad,\qquad\bar{Q}\cdot\psi\sim\mathcal{D}\varphi$
cardona-qft-methods_ EQ0075	45	■ 0.993	$\Psi(\mathcal{A})=\{T\, \,\forall N\in\mathbb{N}\exists P\in\mathcal{D}(\mathcal{A}),R\in OP^{-N},p\in\mathbb{N}\,\text{s.t.}\,T=P \mathcal{D} ^{-p}+R\}.$	$\Psi(\mathcal{A})=\{T\, \,\forall N\in\mathbb{N}\exists P\in\mathcal{D}(\mathcal{A}),R\in OP^{-N},p\in\mathbb{N}\,\text{s.t.}\,T=P \mathcal{D} ^{-p}+R\}.$	$\Psi(\mathcal{A})=\{T\, \,\forall N\in\mathbb{N}\exists P\in\mathcal{D}(\mathcal{A}),R\in OP^{-N},p\in\mathbb{N}\,\text{s.t.}\,T=P \mathcal{D} ^{-p}+R\}.$
cardona-qft-methods_ EQ0927	352	■ 0.993	$H_*^{pro}\left(\mathbb{L}(M\times_G EG)^{-T(M\times_G EG)}\right):=\lim_{\leftarrow n}H_*\left((P_G M\times_G EG_n)^{-e_0^*T(M\times_G EG_n)}\right)$	$H_*^{pro}\left(\mathbb{L}(M\times_G EG)^{-T(M\times_G EG)}\right):=\lim_{\leftarrow n}H_*\left((P_G M\times_G EG_n)^{-e_0^*T(M\times_G EG_n)}\right)$	$H_*^{pro}(\mathbb{L}(M\times_G EG)^{-T(M\times_G EG)}) := \lim_{\leftarrow n} H_*((P_G M\times_G EG_n)^{-e_0^*T(M\times_G EG_n)})$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0899	347	■ 0.993	$\begin{gathered} H\,H_*(A,M)=H^*\left(A\stackrel{L}{\otimes}_{A^e}M\right)\\ \otimes_{A^e}M\right)\\ \mathcal{H}\mathcal{H}^*(A,M)=H_*\left(\mathcal{R}Hom_{A^e}(A,M)\right) \end{gathered}$	$HH_*(A,M)=H^*\left(A\stackrel{L}{\otimes}_{A^e}M\right)$ $HH^*(A,M)=H_*\left(\mathcal{R}Hom_{A^e}(A,M)\right)$	$HH_*(A,M)=H^*(A\stackrel{L}{\otimes}_{A^e}M)$ $HH^*(A,M)=H_*(\mathcal{R}Hom_{A^e}(A,M))$
cardona-qft-methods_ EQ0138	80	■ 0.994	$\phi_i:U_i\stackrel{\sim}{\rightarrow}V_i\subset\mathbb{R}^n\quad(i=1,\ldots,m)$	$\phi_i:U_i\stackrel{\sim}{\rightarrow}V_i\subset\mathbb{R}^n\quad(i=1,\ldots,m)$	$\phi_i:U_i\stackrel{\sim}{\rightarrow}V_i\subset\mathbb{R}^n\quad(i=1,\ldots,m)$
cardona-qft-methods_ EQ0292	142	■ 0.994	$\left[B\,\ell_BX\right]_{SB}-[E]_{SB}=[X]_{SB}-[B]_{SB},$	$[B\ell_BX]_{SB}-[E]_{SB}=[X]_{SB}-[B]_{SB},$	$[B\ell_BX]_{SB}-[E]_{SB}=[X]_{SB}-[B]_{SB}\quad,$
cardona-qft-methods_ EQ0917	351	■ 0.994	$\operatorname{Tor}_{C^*(X)^e}(C^*(X),C^*(X)\#G)\cong H^*(P_GX).$	$\operatorname{Tor}_{C^*(X)^e}(C^*(X),C^*(X)\#G)\cong H^*(P_GX).$	$\operatorname{Tor}_{C^*(X)^e}(C^*(X),C^*(X)\#G)\cong H^*(P_GX).$
cardona-qft-methods_ EQ0209	97	■ 0.994	$E^{+}:=\bigoplus_{j\text{ even}}\Lambda^j(T^*M\otimes\mathcal{C}),\quad E^{-}:=\bigoplus_{j\text{ odd}}\Lambda^j(T^*M\otimes\mathcal{C}).$	$E^{+}:=\bigoplus_{j\text{ even}}\Lambda^j(T^*M\otimes\mathcal{C}),\quad E^{-}:=\bigoplus_{j\text{ odd}}\Lambda^j(T^*M\otimes\mathcal{C}).$	$E^{+}:=\bigoplus_{j\text{ even}}\Lambda^j(T^*M\otimes\mathcal{C}),\quad E^{-}:=\bigoplus_{j\text{ odd}}\Lambda^j(T^*M\otimes\mathcal{C}).$
cardona-qft-methods_ EQ0263	125	■ 0.994	$\langle G\rangle=\prod_i\langle G_i\rangle/\text{extra symmetries}$	$\langle G\rangle=\prod_i\langle G_i\rangle/\text{extra symmetries}$	$\langle G\rangle=\prod_i\langle G_i\rangle/\text{extra symmetries}\quad.$
cardona-qft-methods_ EQ0422 (7)	193	■ 0.994	$\int_M\Theta\operatorname{Tr}[F\wedge F]=\int_{\partial\bar{M}}j\wedge\operatorname{Tr}[A\wedge dA],$	$\int_M\Theta\operatorname{Tr}[F\wedge F]=\int_{\partial\bar{M}}j\wedge\operatorname{Tr}[A\wedge dA],$	$\int_M\Theta Tr[F\wedge F]=\int_{\partial\bar{M}}j\wedge Tr[A\wedge dA],$
cardona-qft-methods_ EQ0888	345	■ 0.994	$\circ:H_p(LM)\otimes H_q(LM)\rightarrow H_{p+q-n}(LM).$	$\circ:H_p(LM)\otimes H_q(LM)\rightarrow H_{p+q-n}(LM).$	$\circ:H_p(LM)\otimes H_q(LM)\rightarrow H_{p+q-n}(LM).$
cardona-qft-methods_ EQ0364 (3)	174	■ 0.994	$\begin{aligned} \delta e^a&=d\lambda^a+\omega_b^a\lambda^b\\ &=D_\omega\lambda^a \end{aligned}$	$\delta e^a=d\lambda^a+\omega_b^a\lambda^b$ $=D_\omega\lambda^a$	$\delta e^a=d\lambda^a+\omega_b^a\lambda^b,$ $=D_\omega\lambda^a$
cardona-qft-methods_ EQ0412 (42)	188	■ 0.995	$\mathcal{T}_{2n+1}(A,\bar{A})=\mathcal{C}_{2n+1}(A)-\mathcal{C}_{2n+1}(\bar{A})+d\mathcal{B}_{2n}(A,\bar{A}).$	$\mathcal{T}_{2n+1}(A,\bar{A})=\mathcal{C}_{2n+1}(A)-\mathcal{C}_{2n+1}(\bar{A})+d\mathcal{B}_{2n}(A,\bar{A}).$	$\mathcal{T}_{2n+1}(A,\bar{A})=\mathcal{C}_{2n+1}(A)-\mathcal{C}_{2n+1}(\bar{A})+d\mathcal{B}_{2n}(A,\bar{A}).$
cardona-qft-methods_ EQ0942 (2)	358	■ 0.995	$[\emptyset]=0.$	$[\emptyset]=0.$	$[\emptyset]=0.$
cardona-qft-methods_ EQ0349 (17)	168	■ 0.995	$\begin{aligned} &\&\mathcal{E}_{a\,b_2\cdots b_D}^{(p)}:=\epsilon_{ab_2\cdots b_D}R^{b_2b_3}\cdots R^{b_{2p}b_{2p+1}}e^{b_{2p+2}}\cdots e^{b_D},\\ &\mathcal{E}_{ab}^{(p)}:=\epsilon_{aba_3\cdots a_D}R^{a_3a_4}\cdots R^{a_{2p-1}a_{2p}}T^{a_{2p+1}}e^{a_{2p+2}}\cdots e^{a_D}. \end{aligned}$	$\mathcal{E}_a^{(p)}:=\epsilon_{ab_2\cdots b_D}R^{b_2b_3}\cdots R^{b_{2p}b_{2p+1}}e^{b_{2p+2}}\cdots e^{b_D},$ $\mathcal{E}_{ab}^{(p)}:=\epsilon_{aba_3\cdots a_D}R^{a_3a_4}\cdots R^{a_{2p-1}a_{2p}}T^{a_{2p+1}}e^{a_{2p+2}}\cdots e^{a_D}.$	$\mathcal{E}_a^{(p)}:=\epsilon_{ab_2\cdots b_D}R^{b_2b_3}\cdots R^{b_{2p}b_{2p+1}}e^{b_{2p+2}}\cdots e^{b_D},$ $\mathcal{E}_{ab}^{(p)}:=\epsilon_{aba_3\cdots a_D}R^{a_3a_4}\cdots R^{a_{2p-1}a_{2p}}T^{a_{2p+1}}e^{a_{2p+2}}\cdots e^{a_D}.$
cardona-qft-methods_ EQ0796 (37)	302	■ 0.995	$N\,S-N\,S:\text{ type IIA and IIB }\mathbf{8}_v\otimes\mathbf{8}_v=\mathbf{1}\oplus\mathbf{28}\oplus\mathbf{35}=\Phi\oplus\mathbf{B}_2\oplus\mathbf{G}_{\mathrm{MN}}$	$NS-NS:\text{ type IIA and IIB }\mathbf{8}_v\otimes\mathbf{8}_v=\mathbf{1}\oplus\mathbf{28}\oplus\mathbf{35}=\Phi\oplus\mathbf{B}_2\oplus\mathbf{G}_{\mathrm{MN}}$	$NS-NS:\text{ type IIA and IIB }\mathbf{8}_v\otimes\mathbf{8}_v=\mathbf{1}\oplus\mathbf{28}\oplus\mathbf{35}=\Phi\oplus\mathbf{B}_2\oplus\mathbf{G}_{\mathrm{MN}}$
cardona-qft-methods_ EQ0471	218	■ 0.995	$\operatorname{Tr}(f(D/\Lambda))\sim\sum_kf_k\Lambda^k\int D ^{-k}+f(0)\zeta_D(0)+o(1),$	$\operatorname{Tr}(f(D/\Lambda))\sim\sum_kf_k\Lambda^k\int D ^{-k}+f(0)\zeta_D(0)+o(1),$	$\operatorname{Tr}(f(D/\Lambda))\sim\sum_kf_k\Lambda^k\int D ^{-k}+f(0)\zeta_D(0)+o(1),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0574 (9)	249	■ 0.995	$\widehat{x}_{\mu} \star \widehat{x}_{\nu} - \widehat{x}_{\nu} \star \widehat{x}_{\mu} = \left[\widehat{x}_{\mu}, \widehat{x}_{\nu} \right]_{\star} = i \theta_{\mu\nu} \quad \mu, \nu = 0, 1, \dots d.$	$\widehat{x}_{\mu} \star \widehat{x}_{\nu} - \widehat{x}_{\nu} \star \widehat{x}_{\mu} = [\widehat{x}_{\mu}, \widehat{x}_{\nu}]_{\star} = i \theta_{\mu\nu} \quad \mu, \nu = 0, 1, \dots d.$	$\widehat{x}_{\mu} \star \widehat{x}_{\nu} - \widehat{x}_{\nu} \star \widehat{x}_{\mu} = [\widehat{x}_{\mu}, \widehat{x}_{\nu}]_{\star} = i \theta_{\mu\nu} \quad \mu, \nu = 0, 1, \dots d.$
cardona-qft-methods_ EQ0111	73	■ 0.996	$\operatorname{Ind}(B \ A) = \operatorname{Ind}(A) + \operatorname{Ind}(B).$	$\operatorname{Ind}(BA) = \operatorname{Ind}(A) + \operatorname{Ind}(B).$	$\operatorname{Ind}(BA) = \operatorname{Ind}(A) + \operatorname{Ind}(B).$
cardona-qft-methods_ EQ0429	206	■ 0.996	$\mathcal{A} = \oplus_{i=1}^N M_{n_i}(\mathbb{K}_i)$	$\mathcal{A} = \oplus_{i=1}^N M_{n_i}(\mathbb{K}_i)$	$\mathcal{A} = \oplus_{i=1}^N M_{n_i}(\mathbb{K}_i)$
cardona-qft-methods_ EQ0494	223	■ 0.996	$\begin{gathered} 16 \ \pi^{\{2\}} \ \beta_{g_i} = b_i \ g_i^3 \\ \quad \text{with } \left(b_{SU(3)}, b_{SU(2)}, b_{U(1)} \right) = \left(-7, -\frac{19}{6}, \frac{41}{10} \right) \\ 16 \pi^2 \ \beta_{Y_u} = Y_u \left(\frac{3}{2} Y_u^{\dagger} Y_u - \frac{3}{2} Y_d^{\dagger} Y_d + \mathfrak{a} - \frac{17}{20} g_1^2 - \frac{9}{4} g_2^2 - 8 g_3^2 \right) \\ 16 \pi^2 \ \beta_{Y_d} = Y_d \left(\frac{3}{2} Y_d^{\dagger} Y_d - \frac{3}{2} Y_u^{\dagger} Y_u + \mathfrak{a} - \frac{1}{4} g_1^2 - \frac{9}{4} g_2^2 - 8 g_3^2 \right) \\ 16 \pi^2 \ \beta_{Y_{\nu}} = Y_{\nu} \left(\frac{3}{2} Y_{\nu}^{\dagger} Y_{\nu} - \frac{3}{2} Y_e^{\dagger} Y_e + \mathfrak{a} - \frac{9}{20} g_1^2 - \frac{9}{4} g_2^2 \right) \\ 16 \pi^2 \ \beta_{Y_e} = Y_e \left(\frac{3}{2} Y_e^{\dagger} Y_e - \frac{3}{2} Y_{\nu}^{\dagger} Y_{\nu} + \mathfrak{a} - \frac{9}{4} g_1^2 - \frac{9}{4} g_2^2 \right) \\ 16 \pi^2 \ \beta_M = Y_{\nu} Y_{\nu}^{\dagger} M + M (Y_{\nu} Y_{\nu}^{\dagger})^T \\ 16 \pi^2 \ \beta_{\lambda} = 6 \lambda^2 - 3 \lambda \left(3 g_2^2 + \frac{3}{5} g_1^2 \right) + 3 g_2^4 + \frac{3}{2} \left(\frac{3}{5} g_1^2 + g_2^2 \right)^2 + 4 \lambda \mathfrak{a} - 8 \mathfrak{b}. \end{gathered}$	$\begin{gathered} 16 \pi^2 \beta_{g_i} = b_i g_i^3 \quad \text{with } (b_{SU(3)}, b_{SU(2)}, b_{U(1)}) = \left(-7, -\frac{19}{6}, \frac{41}{10} \right) \\ 16 \pi^2 \beta_{Y_u} = Y_u \left(\frac{3}{2} Y_u^{\dagger} Y_u - \frac{3}{2} Y_d^{\dagger} Y_d + \mathfrak{a} - \frac{17}{20} g_1^2 - \frac{9}{4} g_2^2 - 8 g_3^2 \right) \\ 16 \pi^2 \beta_{Y_d} = Y_d \left(\frac{3}{2} Y_d^{\dagger} Y_d - \frac{3}{2} Y_u^{\dagger} Y_u + \mathfrak{a} - \frac{1}{4} g_1^2 - \frac{9}{4} g_2^2 - 8 g_3^2 \right) \\ 16 \pi^2 \beta_{Y_{\nu}} = Y_{\nu} \left(\frac{3}{2} Y_{\nu}^{\dagger} Y_{\nu} - \frac{3}{2} Y_e^{\dagger} Y_e + \mathfrak{a} - \frac{9}{20} g_1^2 - \frac{9}{4} g_2^2 \right) \\ 16 \pi^2 \beta_{Y_e} = Y_e \left(\frac{3}{2} Y_e^{\dagger} Y_e - \frac{3}{2} Y_{\nu}^{\dagger} Y_{\nu} + \mathfrak{a} - \frac{9}{4} g_1^2 - \frac{9}{4} g_2^2 \right) \\ 16 \pi^2 \beta_M = Y_{\nu} Y_{\nu}^{\dagger} M + M (Y_{\nu} Y_{\nu}^{\dagger})^T \\ 16 \pi^2 \beta_{\lambda} = 6 \lambda^2 - 3 \lambda \left(3 g_2^2 + \frac{3}{5} g_1^2 \right) + 3 g_2^4 + \frac{3}{2} \left(\frac{3}{5} g_1^2 + g_2^2 \right)^2 + 4 \lambda \mathfrak{a} - 8 \mathfrak{b} \end{gathered}$	$\begin{gathered} 16 \pi^2 \ \beta_{g_i} = b_i \ g_i^3 \quad \text{with } (b_{SU(3)}, b_{SU(2)}, b_{U(1)}) = \left(-7, -\frac{19}{6}, \frac{41}{10} \right) \\ 16 \pi^2 \ \beta_{Y_u} = Y_u \left(\frac{3}{2} Y_u^{\dagger} Y_u - \frac{3}{2} Y_d^{\dagger} Y_d + \mathfrak{a} - \frac{17}{20} g_1^2 - \frac{9}{4} g_2^2 - 8 g_3^2 \right) \\ 16 \pi^2 \ \beta_{Y_d} = Y_d \left(\frac{3}{2} Y_d^{\dagger} Y_d - \frac{3}{2} Y_u^{\dagger} Y_u + \mathfrak{a} - \frac{1}{4} g_1^2 - \frac{9}{4} g_2^2 - 8 g_3^2 \right) \\ 16 \pi^2 \ \beta_{Y_{\nu}} = Y_{\nu} \left(\frac{3}{2} Y_{\nu}^{\dagger} Y_{\nu} - \frac{3}{2} Y_e^{\dagger} Y_e + \mathfrak{a} - \frac{9}{20} g_1^2 - \frac{9}{4} g_2^2 \right) \\ 16 \pi^2 \ \beta_{Y_e} = Y_e \left(\frac{3}{2} Y_e^{\dagger} Y_e - \frac{3}{2} Y_{\nu}^{\dagger} Y_{\nu} + \mathfrak{a} - \frac{9}{4} g_1^2 - \frac{9}{4} g_2^2 \right) \\ 16 \pi^2 \ \beta_M = Y_{\nu} Y_{\nu}^{\dagger} M + M (Y_{\nu} Y_{\nu}^{\dagger})^T \\ 16 \pi^2 \ \beta_{\lambda} = 6 \lambda^2 - 3 \lambda (3 g_2^2 + \frac{3}{5} g_1^2) + 3 g_2^4 + \frac{3}{2} (\frac{3}{5} g_1^2 + g_2^2)^2 + 4 \lambda \mathfrak{a} - 8 \mathfrak{b}. \end{gathered}$
cardona-qft-methods_ EQ0241	115	■ 0.996	$\kappa_*^*: A_* X \rightarrow A_* \{\mathrm{pt}\} = \mathbb{Z},$	$\kappa_* : A_* X \rightarrow A_* \{\mathrm{pt}\} = \mathbb{Z},$	$\kappa_* : A_* X \rightarrow A_* \{\mathrm{pt}\} = \mathbb{Z} \quad ,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0279	136	■ 0.996	$\begin{aligned} \sum_{m \geq 0} \mathbb{U}(G_{me}) \frac{s^m}{m!} &= \frac{e^{\mathbb{T}s} - e^{-s}}{\mathbb{T} + 1} \mathbb{U}(G) + \frac{e^{\mathbb{T}s} + \mathbb{T}e^{-s}}{\mathbb{T} + 1} \mathbb{U}(G \setminus e) \\ &\quad + \left(s e^{\mathbb{T}s} - \frac{e^{\mathbb{T}s} - e^{-s}}{\mathbb{T} + 1} \right) \mathbb{U}(G/e). \end{aligned}$	$\sum_{m \geq 0} \mathbb{U}(G_{me}) \frac{s^m}{m!} = \frac{e^{\mathbb{T}s} - e^{-s}}{\mathbb{T} + 1} \mathbb{U}(G) + \frac{e^{\mathbb{T}s} + \mathbb{T}e^{-s}}{\mathbb{T} + 1} \mathbb{U}(G \setminus e) + \left(s e^{\mathbb{T}s} - \frac{e^{\mathbb{T}s} - e^{-s}}{\mathbb{T} + 1} \right) \mathbb{U}(G/e)$	$\sum_{m \geq 0} \mathbb{U}(G_{me}) \frac{s^m}{m!} = \frac{e^{\mathbb{T}s} - e^{-s}}{\mathbb{T} + 1} \mathbb{U}(G) + \frac{e^{\mathbb{T}s} + \mathbb{T}e^{-s}}{\mathbb{T} + 1} \mathbb{U}(G \setminus e) + \left(s e^{\mathbb{T}s} - \frac{e^{\mathbb{T}s} - e^{-s}}{\mathbb{T} + 1} \right) \mathbb{U}(G/e).$
cardona-qft-methods_ EQ0588 (21)	251	■ 0.996	$\tau_0 \Delta_\theta(\Lambda) \neq \Delta_\theta(\Lambda) \tau_0,$	$\tau_0 \Delta_\theta(\Lambda) \neq \Delta_\theta(\Lambda) \tau_0,$	$\tau_0 \Delta_\theta(\Lambda) \neq \Delta_\theta(\Lambda) \tau_0,$
cardona-qft-methods_ EQ0709	273	■ 0.996	$e^{\mathrm{i} \, p_k L} = \prod_{j \neq k}^M S(p_k, p_j) \quad \text{for} \quad k = 1, \dots, M,$	$e^{ip_k L} = \prod_{j \neq k}^M S(p_k, p_j) \quad \text{for} \quad k = 1, \dots, M,$	$e^{ip_k L} = \prod_{j \neq k}^M S(p_k, p_j) \quad \text{for} \quad k = 1, \dots, M,$
cardona-qft-methods_ EQ0803 (3)	306	■ 0.996	$\emptyset \longrightarrow T^*M \longrightarrow E \xrightarrow{\pi} TM \longrightarrow 0.$	$0 \longrightarrow T^*M \longrightarrow E \xrightarrow{\pi} TM \longrightarrow 0.$	$0 \longrightarrow T^*M \longrightarrow E \xrightarrow{\pi} TM \longrightarrow 0.$
cardona-qft-methods_ EQ0202 (10)	96	■ 0.997	$\mathcal{D} = d + d^*.$	$\mathcal{D} = d + d^*.$	$\mathcal{D} = d + d^*.$
cardona-qft-methods_ EQ0290	139	■ 0.997	$\begin{aligned} \sum_{m \geq 0} C_{G_{(m+1)e}}(t) \frac{s^m}{m!} &= \left(e^{ts} - e^{(t-1)s} \right) C_{G_{2e}}(t) \\ &\quad - \left((t-1)e^{ts} - te^{(t-1)s} \right) C_G(t) + t \left((s-1)e^{ts} + e^{(t-1)s} \right) C_{G/e}(t). \end{aligned}$	$\sum_{m \geq 0} C_{G_{(m+1)e}}(t) \frac{s^m}{m!} = \left(e^{ts} - e^{(t-1)s} \right) C_{G_{2e}}(t) - \left((t-1)e^{ts} - te^{(t-1)s} \right) C_G(t) + t \left((s-1)e^{ts} + e^{(t-1)s} \right) C_{G/e}(t)$	$\sum_{m \geq 0} C_{G_{(m+1)e}}(t) \frac{s^m}{m!} = \left(e^{ts} - e^{(t-1)s} \right) C_{G_{2e}}(t) - \left((t-1)e^{ts} - te^{(t-1)s} \right) C_G(t) + t \left((s-1)e^{ts} + e^{(t-1)s} \right) C_{G/e}(t).$
cardona-qft-methods_ EQ0439	210	■ 0.997	$\begin{gathered} \mathcal{A}_F = \left\{ \left(\lambda, q_L, \lambda, m \right) \mid \lambda \in \mathbb{C}, q_L \in \mathbb{H}, m \in M_3(\mathbb{C}) \right\} \\ \sim \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C}), \end{gathered}$	$\mathcal{A}_F = \{ (\lambda, q_L, \lambda, m) \mid \lambda \in \mathbb{C}, q_L \in \mathbb{H}, m \in M_3(\mathbb{C}) \} \sim \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C}),$	$\mathcal{A}_F = \{ (\lambda, q_L, \lambda, m) \mid \lambda \in \mathbb{C}, q_L \in \mathbb{H}, m \in M_3(\mathbb{C}) \} \sim \mathbb{C} \oplus \mathbb{H} \oplus M_3(\mathbb{C}),$
cardona-qft-methods_ EQ0958 (10)	361	■ 0.997	$X_{\Gamma^\vee} = \pi_2 \left(\pi_1^{-1} (X_\Gamma) \right),$	$X_{\Gamma^\vee} = \pi_2 \left(\pi_1^{-1} (X_\Gamma) \right),$	$X_{\Gamma^\vee} = \pi_2 \left(\pi_1^{-1} (X_\Gamma) \right),$
cardona-qft-methods_ EQ0804 (4)	306	■ 0.997	$x_{(\alpha)} + \xi_{(\alpha)} = a_{(\alpha\beta)} \cdot x_{(\beta)} + \left[a_{(\alpha\beta)}^{-T} \xi_{(\beta)} - \iota_{a_{(\alpha\beta)}x_{(\beta)}} b_{(\alpha\beta)} \right],$	$x_{(\alpha)} + \xi_{(\alpha)} = a_{(\alpha\beta)} \cdot x_{(\beta)} + \left[a_{(\alpha\beta)}^{-T} \xi_{(\beta)} - \iota_{a_{(\alpha\beta)}x_{(\beta)}} b_{(\alpha\beta)} \right],$	$x_{(\alpha)} + \xi_{(\alpha)} = a_{(\alpha\beta)} \cdot x_{(\beta)} + \left[a_{(\alpha\beta)}^{-T} \xi_{(\beta)} - \iota_{a_{(\alpha\beta)}x_{(\beta)}} b_{(\alpha\beta)} \right],$
cardona-qft-methods_ EQ0157	86	■ 0.997	$j: T^*M \hookrightarrow \mathbb{R}^N \oplus \mathbb{R}^N \simeq \mathcal{C}^N.$	$j: T^*M \hookrightarrow \mathbb{R}^N \oplus \mathbb{R}^N \simeq \mathcal{C}^N.$	$j: T^*M \hookrightarrow \mathbb{R}^N \oplus \mathbb{R}^N \simeq \mathcal{C}^N.$
cardona-qft-methods_ EQ0286	138	■ 0.997	$C_{G_{2e}}(T) = (2T-1)C_G(T) - T(T-1)C_{G \setminus e}(T) + C_{G/e}(T).$	$C_{G_{2e}}(T) = (2T-1)C_G(T) - T(T-1)C_{G \setminus e}(T) + C_{G/e}(T).$	$C_{G_{2e}}(T) = (2T-1)C_G(T) - T(T-1)C_{G \setminus e}(T) + C_{G/e}(T).$
cardona-qft-methods_ EQ0543	243	■ 0.997	$\Delta \circ m(a \otimes b) = (m_{13} \otimes m_{24})(\Delta \otimes \Delta)(a \otimes b).$	$\Delta \circ m(a \otimes b) = (m_{13} \otimes m_{24})(\Delta \otimes \Delta)(a \otimes b).$	$\Delta \circ m(a \otimes b) = (m_{13} \otimes m_{24})(\Delta \otimes \Delta)(a \otimes b).$
cardona-qft-methods_ EQ0811 (11)	308	■ 0.997	$\Phi_\pm = \eta_+^1 \otimes \eta_\pm^{2\dagger}.$	$\Phi_\pm = \eta_+^1 \otimes \eta_\pm^{2\dagger}.$	$\Phi_\pm = \eta_+^1 \otimes \eta_\pm^{2\dagger}.$
cardona-qft-methods_ EQ0987	370	■ 0.997	$\mathbb{D}\pi(X)\xi = X\xi = \lim_{t \rightarrow 0} \frac{1}{t} (\exp(tX)\xi - \xi),$	$\mathbb{D}\pi(X)\xi = X\xi = \lim_{t \rightarrow 0} \frac{1}{t} (\exp(tX)\xi - \xi),$	$\mathbb{D}\pi(X)\xi = X\xi = \lim_{t \rightarrow 0} \frac{1}{t} (\exp(tX)\xi - \xi),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0081 (28)	48	■ 0.998	$\mathcal{D}_{\mathcal{A}}=\mathcal{D}+\widetilde{\mathcal{A}} \text{ where } \widetilde{\mathcal{A}}=A+\varepsilon JAJ^{-1}, \quad \mathcal{D}_{\mathcal{A}}=\mathcal{D}+P_{\mathcal{A}}$	$\mathcal{D}_A = \mathcal{D} + \tilde{A} \text{ where } \tilde{A} = A + \varepsilon JAJ^{-1}, \quad D_A = \mathcal{D}_A + P_A$	$\mathcal{D}_A = \mathcal{D} + \tilde{A} \text{ where } \tilde{A} = A + \varepsilon JAJ^{-1}, \quad D_A = \mathcal{D}_A + P_A$
cardona-qft-methods_ EQ0169	89	■ 0.998	$g_{\{1\}} \cdot \left(x, g_{\{2\}}\right) := \left(x, g_{\{1\}} g_{\{2\}}\right), \quad x \in N, g_{\{1\}}, g_{\{2\}} \in G.$	$g_1 \cdot (x, g_2) := (x, g_1 g_2), \quad x \in N, g_1, g_2 \in G.$	$g_1 \cdot (x, g_2) := (x, g_1 g_2), \quad x \in N, g_1, g_2 \in G.$
cardona-qft-methods_ EQ0542	243	■ 0.998	$m\left(\int d\mu(g)\alpha(g)g\otimes\int d\mu(h)\beta(h)h\right)=\int d\mu(g)d\mu(h)\alpha(g)\beta(h)gh.$	$m\left(\int d\mu(g)\alpha(g)g\otimes\int d\mu(h)\beta(h)h\right)=\int d\mu(g)d\mu(h)\alpha(g)\beta(h)gh.$	$m\left(\int d\mu(g)\alpha(g)g\otimes\int d\mu(h)\beta(h)h\right)=\int d\mu(g)d\mu(h)\alpha(g)\beta(h)gh.$
cardona-qft-methods_ EQ0617 (25)	256	■ 0.998	$f_{\{1\}}\,f_{\{2\}}\in C^{\infty}(M),$	$f_1f_2\in C^\infty(M),$	$f_1f_2\in C^\infty(M),$
cardona-qft-methods_ EQ0638 (45)	258	■ 0.998	$\varphi_{\theta}=\varphi_0e^{-\frac{i}{2}\overleftarrow{\partial}\wedge P}.$	$\varphi_{\theta}=\varphi_0e^{-\frac{i}{2}\overleftarrow{\partial}\wedge P}.$	$\varphi_{\theta}=\varphi_0e^{-\frac{i}{2}\overleftarrow{\partial}\wedge P}.$
cardona-qft-methods_ EQ0664 (15)	263	■ 0.998	$\theta\left(x_{\{0\}}-y_{\{0\}}\right)=\theta\left((\Lambda x)_{\{0\}}-(\Lambda y)_{\{0\}}\right),$	$\theta\left(x_0-y_0\right)=\theta\left((\Lambda x)_0-(\Lambda y)_0\right),$	$\theta(x_0-y_0)=\theta((\Lambda x)_0-(\Lambda y)_0),$
cardona-qft-methods_ EQ0983 (3)	367	■ 0.998	$Q=\left\{t_i\mid \text{ such that } t_i \text{ appears in } q(t)\right\}.$	$Q=\left\{t_i\mid \text{ such that } t_i \text{ appears in } q(t)\right\}.$	$Q=\left\{t_i\mid \text{ such that } t_i \text{ appears in } q(t)\right\}.$
cardona-qft-methods_ EQ0052	37	■ 0.998	$\begin{aligned} &\left\langle U_{g',g}\psi',U_{g',g}\psi'\right\rangle_{\mathcal{H}_g}=\int_M U_{g,g'}\psi' _g^2dvol_g=\int_M I_{g',g}\psi' _g^2f_{g',g}dvol_g\\ &=\int_M \psi' _{g'}^2dvol_{g'}=\langle\psi_{g'},\psi_{g'}\rangle_{\mathcal{H}_{g'}} \end{aligned}$	$\begin{aligned} \langle U_{g',g}\psi',U_{g',g}\psi'\rangle_{\mathcal{H}_g} &= \int_M U_{g,g'}\psi' _g^2 dvol_g = \int_M I_{g',g}\psi' _g^2 f_{g',g} dvol_g \\ &= \int_M \psi' _{g'}^2 dvol_{g'} = \langle \psi_{g'}, \psi_{g'} \rangle_{\mathcal{H}_{g'}} \end{aligned}$	$\begin{aligned} \langle U_{g',g}\psi',U_{g',g}\psi'\rangle_{\mathcal{H}_g} &= \int_M U_{g,g'}\psi' _g^2 dvol_g = \int_M I_{g',g}\psi' _g^2 f_{g',g} dvol_g \\ &= \int_M \psi' _{g'}^2 dvol_{g'} = \langle \psi_{g'}, \psi_{g'} \rangle_{\mathcal{H}_{g'}}. \end{aligned}$
cardona-qft-methods_ EQ0006	19	■ 0.998	$\begin{aligned} &\sigma(x,\xi)\in S^m\left(U\times\mathbb{R}^d\right)\longleftrightarrow k_\sigma(x,y)\in C_c^\infty\left(U\times U\times\mathbb{R}^d\right)\\ &\qquad\qquad\qquad\longleftrightarrow A=Op(\sigma)\in\Psi DO^m \end{aligned}$	$\begin{aligned} \sigma(x,\xi)\in S^m\left(U\times\mathbb{R}^d\right) &\longleftrightarrow k_\sigma(x,y)\in C_c^\infty\left(U\times U\times\mathbb{R}^d\right)\\ &\longleftrightarrow A=Op(\sigma)\in\Psi DO^m \end{aligned}$	$\begin{aligned} \sigma(x,\xi)\in S^m(U\times\mathbb{R}^d) &\longleftrightarrow k_\sigma(x,y)\in C_c^\infty(U\times U\times\mathbb{R}^d)\\ &\longleftrightarrow A=Op(\sigma)\in\Psi DO^m \end{aligned}$
cardona-qft-methods_ EQ0010	19	■ 0.998	$\sigma^{\phi_*P}(x,\xi)=\sigma_m^P\big(\phi^{-1}(x),(d\phi)^t\xi\big)+\sum_{ \alpha >0}\frac{(-i)^\alpha}{\alpha!}\phi_\alpha(x,\xi)\partial_\xi^\alpha\sigma^P\big(\phi^{-1}(x),(d\phi)^t\xi\big)$	$\sigma^{\phi_*P}(x,\xi)=\sigma_m^P\big(\phi^{-1}(x),(d\phi)^t\xi\big)+\sum_{ \alpha >0}\frac{(-i)^\alpha}{\alpha!}\phi_\alpha(x,\xi)\partial_\xi^\alpha\sigma^P\big(\phi^{-1}(x),(d\phi)^t\xi\big)$	$\sigma^{\phi_*P}(x,\xi)=\sigma_m^P\big(\phi^{-1}(x),(d\phi)^t\xi\big)+\sum_{ \alpha >0}\frac{(-i)^\alpha}{\alpha!}\phi_\alpha(x,\xi)\partial_\xi^\alpha\sigma^P\big(\phi^{-1}(x),(d\phi)^t\xi\big)$
cardona-qft-methods_ EQ0074	45	■ 0.998	$\nabla\left(OP^\alpha\right)\subset OP^{\alpha+1}.$	$\nabla\left(OP^\alpha\right)\subset OP^{\alpha+1}.$	$\nabla\left(OP^\alpha\right)\subset OP^{\alpha+1}.$
cardona-qft-methods_ EQ0083 (29)	48	■ 0.998	$\left D_{\mathcal{A}}\right ^{-s}\sim\left D\right ^{-s}+\sum_{p=1}^N K_p(Y,s)\left D\right ^{-s}\mod OP^{-N-1-\Re(s)}$	$ D_A ^{-s}\sim D ^{-s}+\sum_{p=1}^N K_p(Y,s) D ^{-s}\mod OP^{-N-1-\Re(s)}$	$ D_A ^{-s}\sim D ^{-s}+\sum_{p=1}^N K_p(Y,s) D ^{-s}\mod OP^{-N-1-\Re(s)}$
cardona-qft-methods_ EQ0395	180	■ 0.998	$L_{2n-1}^{(A)dS}=\kappa\cdot\epsilon_{A_1\cdots A_{2n}}\left[W(dW)^{n-1}+a_3W^3(dW)^{n-2}+\cdots a_{2n-1}W^{2n-1}\right],$	$L_{2n-1}^{(A)dS}=\kappa\cdot\epsilon_{A_1\cdots A_{2n}}\left[W(dW)^{n-1}+a_3W^3(dW)^{n-2}+\cdots a_{2n-1}W^{2n-1}\right],$	$L_{2n-1}^{(A)dS}=\kappa\cdot\epsilon_{A_1\cdots A_{2n}}\left[W(dW)^{n-1}+a_3W^3(dW)^{n-2}+\cdots a_{2n-1}W^{2n-1}\right],$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0413 (43)	188	■ 0.998	$\begin{aligned} & \text{\&A} \rightarrow A^{\prime}=\Lambda^{\{-1\}} \text{\&A} \\ & \Lambda+\Lambda^{\{-1\}} \text{d} \Lambda \quad \& \bar{\text{\&A}} \rightarrow \bar{A}^{\prime}=\bar{\Lambda}^{\{-1\}} \bar{\text{\&A}} \\ & \bar{\Lambda}+\bar{\Lambda}^{\{-1\}} \text{d} \bar{\Lambda} \end{aligned}$	$A \rightarrow A' = \Lambda^{-1} A \Lambda + \Lambda^{-1} d \Lambda$ $\bar{A} \rightarrow \bar{A}' = \bar{\Lambda}^{-1} \bar{A} \bar{\Lambda} + \bar{\Lambda}^{-1} d \bar{\Lambda}$	$A \rightarrow A' = \Lambda^{-1} A \Lambda + \Lambda^{-1} d \Lambda$ $\bar{A} \rightarrow \bar{A}' = \bar{\Lambda}^{-1} \bar{A} \bar{\Lambda} + \bar{\Lambda}^{-1} d \bar{\Lambda}$
cardona-qft-methods_ EQ0437	209	■ 0.998	$J_{\text{\&F}}^{\{2\}}=1, \quad \text{\&J}_{\text{\&F}} \text{\&gamma}_{\text{\&F}}=-\text{\&gamma}_{\text{\&F}} J_{\text{\&F}}.$	$J_F^2 = 1, \quad J_F \gamma_F = -\gamma_F J_F.$	$J_F^2 = 1, \quad J_F \gamma_F = -\gamma_F J_F.$
cardona-qft-methods_ EQ0438	209	■ 0.998	$q(\lambda)=\left(\begin{array}{l} \lambda \\ 0 \end{array}\right) \quad q(\lambda) \uparrow\rangle=\lambda \uparrow\rangle, \quad q(\lambda) \downarrow\rangle=\bar{\lambda} \downarrow\rangle.$	$q(\lambda)=\left(\begin{array}{cc} \lambda & 0 \\ 0 & \bar{\lambda} \end{array}\right) \quad q(\lambda) \uparrow\rangle=\lambda \uparrow\rangle, \quad q(\lambda) \downarrow\rangle=\bar{\lambda} \downarrow\rangle.$	$q(\lambda)=\left(\begin{array}{cc} \lambda & 0 \\ 0 & \bar{\lambda} \end{array}\right) \quad q(\lambda) \uparrow\rangle=\lambda \uparrow\rangle, \quad q(\lambda) \downarrow\rangle=\bar{\lambda} \downarrow\rangle.$
cardona-qft-methods_ EQ0530 (10)	241	■ 0.998	$U^{\{2\}}(g)=U^{\{1\}}(g) \otimes U^{\{1\}}(g),$	$U^{(2)}(g)=U^{(1)}(g) \otimes U^{(1)}(g),$	$U^{(2)}(g)=U^{(1)}(g) \otimes U^{(1)}(g),$
cardona-qft-methods_ EQ0580	250	■ 0.998	$\Lambda: x \rightarrow \Lambda(x) \in \mathbb{R}^N.$	$\Lambda: x \rightarrow \Lambda(x) \in \mathbb{R}^N.$	$\Lambda: x \rightarrow \Lambda(x) \in \mathbb{R}^N.$
cardona-qft-methods_ EQ0874	338	■ 0.998	$\begin{aligned} & \text{\&mathcal{G}} \&=T^{\{*\}}(P \text{\&M}) . \quad \& \text{\&L} \&= \\ & \left\{\left(\left(X_{\{1\}}, \eta_{\{1\}}\right),\left(X_{\{2\}}, \eta_{\{2\}}\right)\right.\right. \\ & \left.\left.\left(X_{\{3\}}, \eta_{\{3\}}\right)\right\} \in \mathcal{C}_{\Pi}^3\right\} \\ & \mid\left(X_1 * X_2, \eta_1 * \eta_2\right) \sim\left(X_3 * \eta_3\right)\} . \\ & \text{\&I} \& \left(X, \eta\right) \mapsto\left(\phi^* X, \phi^* \eta\right)\} . \end{aligned}$	$\mathcal{G}=T^*(PM).$ $L=\left\{\left(X_1, \eta_1\right),\left(X_2, \eta_2\right),\left(X_3, \eta_3\right) \in C_{\Pi}^3 \mid\left(X_1 * X_2, \eta_1 * \eta_2\right) \sim\left(X_3 * \eta_3\right)\right\}.$ $I=\left(X, \eta\right) \mapsto\left(\phi^* X, \phi^* \eta\right)\}.$	$\mathcal{G}=T^*(PM).$ $L=\left\{\left(X_1, \eta_1\right),\left(X_2, \eta_2\right),\left(X_3, \eta_3\right) \in C_{\Pi}^3 \mid\left(X_1 * X_2, \eta_1 * \eta_2\right) \sim\left(X_3 * \eta_3\right)\right\}.$ $I=\left(X, \eta\right) \mapsto\left(\phi^* X, \phi^* \eta\right)\}.$
cardona-qft-methods_ EQ0965 (5)	362	■ 0.998	$X_{\backslash \Gamma} \backslash \Sigma_n=\emptyset=X_{\backslash \Gamma^{\vee}} \backslash \Sigma_n.$	$X_{\Gamma} \backslash \Sigma_n = \emptyset = X_{\Gamma^{\vee}} \backslash \Sigma_n.$	$X_{\Gamma} \backslash \Sigma_n = \emptyset = X_{\Gamma^{\vee}} \backslash \Sigma_n.$
cardona-qft-methods_ EQ0377 (18)	177	■ 0.998	$S_{0(D-1,1)} \hookrightarrow \left\{ \begin{array}{l} S_{0(D,1)}, \\ S_{0(D,2)}, \end{array} \right. \& \Upsilon^{AB} = \text{diag}(\eta^{\text{ab}}, +1) \& \Upsilon^{AB} = \text{diag}(\eta^{\text{ab}}, -1)$	$SO(D-1,1) \hookrightarrow \left\{ \begin{array}{ll} SO(D,1), & \Upsilon^{AB} = \text{diag}(\eta^{\text{ab}}, +1) \\ SO(D-1,2), & \Upsilon^{AB} = \text{diag}(\eta^{\text{ab}}, -1) \end{array} \right.$	$SO(D-1,1) \hookrightarrow \left\{ \begin{array}{ll} SO(D,1), & \Upsilon^{AB} = \text{diag}(\eta^{\text{ab}}, +1) \\ SO(D-1,2), & \Upsilon^{AB} = \text{diag}(\eta^{\text{ab}}, -1) \end{array} \right.$
cardona-qft-methods_ EQ0116	75	■ 0.999	$\operatorname{Ker}(A)=\bigoplus_{V\in\operatorname{Irr}(G)}m_V^+V\quad\operatorname{Coker}(A)=\bigoplus_{V\in\operatorname{Irr}(G)}m_V^-V.$	$\operatorname{Ker}(A)=\bigoplus_{V\in\operatorname{Irr}(G)}m_V^+V;\quad\operatorname{Coker}(A)=\bigoplus_{V\in\operatorname{Irr}(G)}m_V^-V.$	$\operatorname{Ker}(A)=\bigoplus_{V\in\operatorname{Irr}(G)}m_V^+V;\quad\operatorname{Coker}(A)=\bigoplus_{V\in\operatorname{Irr}(G)}m_V^-V.$
cardona-qft-methods_ EQ0411 (41)	188	■ 0.999	$d\,\text{\&mathcal{T}}_{2n-1}(A,\,\bar{A})=\text{\&mathcal{P}}_{2n}(A)-\text{\&mathcal{P}}_{2n}(\bar{A}),$	$d\mathcal{T}_{2n-1}(A,\bar{A})=\mathcal{P}_{2n}(A)-\mathcal{P}_{2n}(\bar{A}),$	$d\mathcal{T}_{2n-1}(A,\bar{A})=\mathcal{P}_{2n}(A)-\mathcal{P}_{2n}(\bar{A}),$
cardona-qft-methods_ EQ0415 (46)	189	■ 0.999	$I[A,\,\bar{A}]=\int_M\text{\&mathcal{C}}(A)+\int_{\bar{M}}\text{\&mathcal{C}}(\bar{A})+\int_{\partial M}\text{\&mathcal{B}}(A,\bar{A}),$	$I[A,\bar{A}]=\int_M\mathcal{C}(A)+\int_{\bar{M}}\mathcal{C}(\bar{A})+\int_{\partial M}\mathcal{B}(A,\bar{A}),$	$I[A,\bar{A}]=\int_M\mathcal{C}(A)+\int_{\bar{M}}\mathcal{C}(\bar{A})+\int_{\partial M}\mathcal{B}(A,\bar{A}),$
cardona-qft-methods_ EQ0448	211	■ 0.999	$\text{\&mathcal{C}}_{\{3\}}=(K\times K)\backslash(G\times G)/K,$	$\mathcal{C}_3=(K\times K)\backslash(G\times G)/K,$	$\mathcal{C}_3=(K\times K)\backslash(G\times G)/K,$
cardona-qft-methods_ EQ0612 (20)	254	■ 0.999	$\left(\Lambda e_p\right)(x)=e_p\left(\Lambda^{-1} x\right)=e_{\Lambda p(x)},$	$\left(\Lambda e_p\right)(x)=e_p\left(\Lambda^{-1} x\right)=e_{\Lambda p(x)},$	$\left(\Lambda e_p\right)(x)=e_p\left(\Lambda^{-1} x\right)=e_{\Lambda p(x)},$
cardona-qft-methods_ EQ0624 (32)	257	■ 0.999	$\Delta_{\theta}(g)=F_{\theta}^{-1}(g\otimes g)F_{\theta};\quad F_{\theta}=\exp^{\frac{i}{2}\hat{P}_{\mu}\otimes\theta_{\mu\nu}\hat{P}_{\nu}},$	$\Delta_{\theta}(g)=F_{\theta}^{-1}(g\otimes g)F_{\theta};\quad F_{\theta}=\exp^{\frac{i}{2}\hat{P}_{\mu}\otimes\theta_{\mu\nu}\hat{P}_{\nu}},$	$\Delta_{\theta}(g)=F_{\theta}^{-1}(g\otimes g)F_{\theta};\quad F_{\theta}=\exp^{\frac{i}{2}\hat{P}_{\mu}\otimes\theta_{\mu\nu}\hat{P}_{\nu}},$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E00744	283	■ 0.999	$[K, \mathcal{O}] = 0 \quad , \quad [S, \mathcal{O}] = 0 \quad , \quad [\bar{S}, \mathcal{O}] = 0.$	$[K, \mathcal{O}] = 0 \quad , \quad [S, \mathcal{O}] = 0 \quad , \quad [\bar{S}, \mathcal{O}] = 0.$	$[K, \mathcal{O}] = 0 \quad , \quad [S, \mathcal{O}] = 0 \quad , \quad [\bar{S}, \mathcal{O}] = 0.$
cardona-qft-methods_ E00767 (8)	293	■ 0.999	$\eta_{MN} \partial_{\sigma^+} X_L^M \partial_{\sigma^+} X_L^N = \eta_{MN} \partial_{\sigma^-} X_R^M \partial_{\sigma^-} X_R^N = 0.$	$\eta_{MN} \partial_{\sigma^+} X_L^M \partial_{\sigma^+} X_L^N = \eta_{MN} \partial_{\sigma^-} X_R^M \partial_{\sigma^-} X_R^N = 0.$	$\eta_{MN} \partial_{\sigma^+} X_L^M \partial_{\sigma^+} X_L^N = \eta_{MN} \partial_{\sigma^-} X_R^M \partial_{\sigma^-} X_R^N = 0.$
cardona-qft-methods_ E00943 (3)	358	■ 0.999	$[X] \ \text{cdot}[Y] = [X \ \text{times} \ Y] \ .$	$[X] \cdot [Y] = [X \times Y].$	$[X] \cdot [Y] = [X \times Y].$
cardona-qft-methods_ E00989	371	■ 0.999	$A \ B = \text{the closure of } \left. A \right _{\mathcal{D}}, \quad \mathcal{D} = B^{-1} \operatorname{dom} A.$	$AB = \text{ the closure of } AB _{\mathcal{D}}, \quad \mathcal{D} = B^{-1} \operatorname{dom} A.$	$AB = \text{the closure of } AB _{\mathcal{D}}, \quad \mathcal{D} = B^{-1} \operatorname{dom} A.$
cardona-qft-methods_ E00289	138	■ 0.999	$\begin{aligned} C_{G_{(m+3)e}}(T) = & \ (3\ T-1)\ C_{G_{(m+2)e}}(T) - (3T^2-2T)\ C_{G_{(m+1)e}}(T) \\ & + (T^3-T^2)\ C_{G_{me}}(T) \end{aligned}$	$C_{G_{(m+3)e}}(T) = (3T-1)C_{G_{(m+2)e}}(T) - (3T^2-2T)\ C_{G_{(m+1)e}}(T) \\ + (T^3-T^2)\ C_{G_{me}}(T)$	$C_{G_{(m+3)e}}(T) = (3T-1)C_{G_{(m+2)e}}(T) - (3T^2-2T)\ C_{G_{(m+1)e}}(T) \\ + (T^3-T^2)\ C_{G_{me}}(T)$
cardona-qft-methods_ E00389 (27)	179	■ 0.999	$\mathbf{E}_{2n} = \epsilon_{A_1 \cdots A_{2n}} F^{A_1 A_2} \ldots F^{A_{2n-1} A_{2n}}.$	$\mathbf{E}_{2n} = \epsilon_{A_1 \cdots A_{2n}} F^{A_1 A_2} \ldots F^{A_{2n-1} A_{2n}}.$	$\mathbf{E}_{2n} = \epsilon_{A_1 \cdots A_{2n}} F^{A_1 A_2} \ldots F^{A_{2n-1} A_{2n}}.$
cardona-qft-methods_ E00392 (30)	180	■ 0.999	$I_{2n-1} = \frac{\kappa}{l} \int_M \int_0^1 dt \epsilon_{a_1 \cdots a_{2n-1}} R_t^{a_1 a_2} \ldots R_t^{a_{2n-3} a_{2n-2}} e^{a_{2n-1}},$	$I_{2n-1} = \frac{\kappa}{l} \int_M \int_0^1 dt \epsilon_{a_1 \cdots a_{2n-1}} R_t^{a_1 a_2} \ldots R_t^{a_{2n-3} a_{2n-2}} e^{a_{2n-1}},$	$I_{2n-1} = \frac{\kappa}{l} \int_M \int_0^1 dt \epsilon_{a_1 \cdots a_{2n-1}} R_t^{a_1 a_2} \ldots R_t^{a_{2n-3} a_{2n-2}} e^{a_{2n-1}},$
cardona-qft-methods_ E00405 (37)	184	■ 0.999	$\begin{aligned} L_{2n}^{BI} = & \epsilon_{a_1 b_1 a_2 b_2 \cdots a_n b_n} \left[R^{a_1 b_1} + \frac{1}{l^2} e^{a_1} e^{b_1} \right] \cdots \left[R^{a_n b_n} + \frac{1}{l^2} e^{a_n} e^{b_n} \right] \\ = & \operatorname{Pfaff} \left[R^{ab} + \frac{1}{l^2} e^a e^b \right]. \end{aligned}$	$L_{2n}^{BI} = \epsilon_{a_1 b_1 a_2 b_2 \cdots a_n b_n} \left[R^{a_1 b_1} + \frac{1}{l^2} e^{a_1} e^{b_1} \right] \cdots \left[R^{a_n b_n} + \frac{1}{l^2} e^{a_n} e^{b_n} \right] \\ = \operatorname{Pfaff} \left[R^{ab} + \frac{1}{l^2} e^a e^b \right].$	$L_{2n}^{BI} = \epsilon_{a_1 b_1 a_2 b_2 \cdots a_n b_n} \left[R^{a_1 b_1} + \frac{1}{l^2} e^{a_1} e^{b_1} \right] \cdots \left[R^{a_n b_n} + \frac{1}{l^2} e^{a_n} e^{b_n} \right] \\ = \operatorname{Pfaff} \left[R^{ab} + \frac{1}{l^2} e^a e^b \right].$
cardona-qft-methods_ E00436	209	■ 0.999	$J_{\mathcal{F}}^2 = 1, \quad \xi b = J_{\mathcal{F}} b^* J_{\mathcal{F}} \xi \quad \xi \in \mathcal{M}_{\mathcal{F}}, b \in \mathcal{A}_{LR}.$	$J_F^2 = 1, \quad \xi b = J_F b^* J_F \xi \quad \xi \in \mathcal{M}_F, b \in \mathcal{A}_{LR}.$	$J_F^2 = 1, \quad \xi b = J_F b^* J_F \xi \quad \xi \in \mathcal{M}_F, b \in \mathcal{A}_{LR}.$
cardona-qft-methods_ E00498	226	■ 0.999	$\lambda(\Lambda_{\text{unif}}) = \frac{\pi^2}{2f_0} \frac{\mathfrak{b}(\Lambda_{\text{unif}})}{\mathfrak{a}(\Lambda_{\text{unif}})^2}$	$\lambda(\Lambda_{\text{unif}}) = \frac{\pi^2}{2f_0} \frac{\mathfrak{b}(\Lambda_{\text{unif}})}{\mathfrak{a}(\Lambda_{\text{unif}})^2}$	$\lambda(\Lambda_{unif}) = \frac{\pi^2}{2f_0} \frac{\mathfrak{b}(\Lambda_{unif})}{\mathfrak{a}(\Lambda_{unif})^2}$
cardona-qft-methods_ E00520 (6)	237	■ 0.999	$\hat{x}_{\mu} \star \hat{x}_{\nu} - \hat{x}_{\nu} \star \hat{x}_{\mu} = [\hat{x}_{\mu}, \hat{x}_{\nu}]_{\star} = i \theta_{\mu\nu}, \quad \mu, \nu = 0, 1, \dots, d.$	$\hat{x}_{\mu} \star \hat{x}_{\nu} - \hat{x}_{\nu} \star \hat{x}_{\mu} = [\hat{x}_{\mu}, \hat{x}_{\nu}]_{\star} = i \theta_{\mu\nu}, \quad \mu, \nu = 0, 1, \dots, d.$	$\hat{x}_{\mu} \star \hat{x}_{\nu} - \hat{x}_{\nu} \star \hat{x}_{\mu} = [\hat{x}_{\mu}, \hat{x}_{\nu}]_{\star} = i \theta_{\mu\nu}, \quad \mu, \nu = 0, 1, \dots, d.$
cardona-qft-methods_ E00592 (1)	252	■ 0.999	$(a, \Lambda) \in \mathcal{P} : (a, \Lambda) x = \Lambda x + a.$	$(a, \Lambda) \in \mathcal{P} : (a, \Lambda) x = \Lambda x + a.$	$(a, \Lambda) \in \mathcal{P} : (a, \Lambda) x = \Lambda x + a.$
cardona-qft-methods_ E00598 (7)	252	■ 0.999	$(\operatorname{Ad} U \otimes V) \Delta_0((a, \Lambda)) \varphi = \varphi$	$(\operatorname{Ad} U \otimes V) \Delta_0((a, \Lambda)) \varphi = \varphi$	$(\operatorname{Ad} U \otimes V) \Delta_0((a, \Lambda)) \varphi = \varphi$
cardona-qft-methods_ E00623 (31)	257	■ 0.999	$\mathcal{P} \ni (a, \Delta) : \alpha \rightarrow (a, \Delta) \alpha, \quad ((a, \Delta) \alpha)(x) = \alpha \left(((a, \Delta)^{-1} x) \right).$	$\mathcal{P} \ni (a, \Delta) : \alpha \rightarrow (a, \Delta) \alpha, \quad ((a, \Delta) \alpha)(x) = \alpha \left(((a, \Delta)^{-1} x) \right).$	$\mathcal{P} \ni (a, \Delta) : \alpha \rightarrow (a, \Delta) \alpha, \quad ((a, \Delta) \alpha)(x) = \alpha \left(((a, \Delta)^{-1} x) \right).$
cardona-qft-methods_ E00678 (3)	269	■ 0.999	$\left[S^{\mathfrak{i}}, S^{\mathfrak{j}}\right]=i \, \epsilon^{\mathfrak{i} \, \mathfrak{j} \, \mathfrak{k}} S^{\mathfrak{k}} \, .$	$[S^i, S^j] = i \epsilon^{ijk} S^k.$	$[S^i, S^j] = i \epsilon^{ijk} S^k.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0697	271	■ 0.999	$\mathrm{H}_{\{l, \, l+1\}}=2\left(\mathrm{H}_{\{l, \, l+1\}}-\mathrm{P}_{\{l, \, l+1\}}\right)=\left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & +2 & -2 & 0 \\ 0 & -2 & +2 & 0 \\ 0 & 0 & 0 & 0 \end{array}\right),$	$\mathcal{H}_{l,l+1} = 2 \left(\mathbb{I}_{l,l+1} - \mathbb{P}_{l,l+1} \right) = \left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & +2 & -2 & 0 \\ 0 & -2 & +2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right),$	$\mathcal{H}_{l,l+1} = 2 \left(\mathbb{I}_{l,l+1} - \mathbb{P}_{l,l+1} \right) = \left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & +2 & -2 & 0 \\ 0 & -2 & +2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right),$
cardona-qft-methods_ EQ0699	271	■ 0.999	$\left[\vec{S}_{\mathrm{tot}}, \hat{H}\right]=0,$	$\left[\vec{S}_{\mathrm{tot}}, \hat{H}\right] = 0,$	$[\vec{S}_{\mathrm{tot}}, \hat{H}] = 0,$
cardona-qft-methods_ EQ0823 (1)	312	■ 0.999	$\mathrm{d} s^2=e^{2 A(y)} \eta_{\mu \nu} \mathrm{d} x^{\mu} \mathrm{d} x^{\nu}+g_{m n} \mathrm{d} y^m \mathrm{d} y^n, \quad \mu=0,1,2,3 \quad m=1, \ldots, 6$	$\mathrm{d} s^2=e^{2 A(y)} \eta_{\mu \nu} \mathrm{d} x^{\mu} \mathrm{d} x^{\nu}+g_{m n} \mathrm{d} y^m \mathrm{d} y^n, \quad \mu=0,1,2,3 \quad m=1, \ldots, 6$	$\mathrm{d} s^2=e^{2 A(y)} \eta_{\mu \nu} \mathrm{d} x^{\mu} \mathrm{d} x^{\nu}+g_{m n} \mathrm{d} y^m \mathrm{d} y^n, \quad \mu=0,1,2,3 \quad m=1, \ldots, 6$
cardona-qft-methods_ EQ0842 (1)	329	■ 0.999	$\Pi^{\wedge\{s \, r\}}(x)\left(\partial_r\right) \Pi^{\wedge\{l \, k\}}(x)+\Pi^{\wedge\{k \, r\}}(x)\left(\partial_r\right) \Pi^{\wedge\{s \, l\}}(x)+\Pi^{\wedge\{l \, r\}}(x)\left(\partial_r\right) \Pi^{\wedge\{k \, s\}}(x)=0,$	$\Pi^{sr}(x)(\partial_r)\Pi^{lk}(x)+\Pi^{kr}(x)(\partial_r)\Pi^{sl}(x)+\Pi^{lr}(x)(\partial_r)\Pi^{ks}(x)=0,$	$\Pi^{sr}(x)(\partial_r)\Pi^{lk}(x)+\Pi^{kr}(x)(\partial_r)\Pi^{sl}(x)+\Pi^{lr}(x)(\partial_r)\Pi^{ks}(x)=0,$
cardona-qft-methods_ EQ0887	344	■ 0.999	$H_{\{p\}}(M) \otimes H_{\{q\}}(M) \longrightarrow H_{\{p+q\}}(M \times M) \xrightarrow{\tau_{\Delta*}} H_{p+q}\left(M^{TM}\right) \xrightarrow{Thom} H_{p+q-n}(M)$	$H_p(M) \otimes H_q(M) \longrightarrow H_{p+q}(M \times M) \xrightarrow{\tau_{\Delta*}} H_{p+q}\left(M^{TM}\right) \xrightarrow{Thom} H_{p+q-n}(M)$	$H_p(M) \otimes H_q(M) \longrightarrow H_{p+q}(M \times M) \xrightarrow{\tau_{\Delta*}} H_{p+q}(M^{TM}) \xrightarrow{Thom} H_{p+q-n}(M)$
cardona-qft-methods_ EQ0988	371	■ 0.999	$A+B=\text{the closure of } A _{\mathcal{D}}+B _{\mathcal{D}}, \quad \mathcal{D}=\text{dom } A\cap\text{dom } B.$	$A+B= \text{ the closure of } A _{\mathcal{D}}+B _{\mathcal{D}}, \quad \mathcal{D}=\text{dom } A\cap\text{dom } B.$	$A+B= \text{ the closure of } A _{\mathcal{D}}+B _{\mathcal{D}}, \quad \mathcal{D}=\text{dom } A\cap\text{dom } B.$
cardona-qft-methods_ EQ0759	287	■ 0.999	$\begin{aligned} &\uparrow=Z=\text{nothing}, \quad \& \\ &\downarrow=M_{A\dot{A}}=8+8 \text{ magnons transforming under } \frac{s}{u}(2 \mid 2) \otimes \overline{\mathfrak{su}}(2 \mid 2), \end{aligned}$	$\begin{aligned} \uparrow &= Z = \text{ nothing } , \\ \downarrow &= M_{A\dot{A}} = 8 + 8 \text{ magnons transforming under } \mathfrak{su}(2 \mid 2) \otimes \overline{\mathfrak{su}}(2 \mid 2), \end{aligned}$	$\begin{aligned} \uparrow &= Z = \text{ nothing } , \\ \downarrow &= M_{A\dot{A}} = 8 + 8 \text{ magnons transforming under } \mathfrak{su}(2 2) \otimes \overline{\mathfrak{su}}(2 2) , \end{aligned}$
cardona-qft-methods_ EQ0939	31	■ 1.000	$\lim _{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=0}^N \mu_n\left((1+\Delta)^{-d / 2}\right)=\frac{\alpha}{\Gamma(d / 2+1)}=\frac{\pi^{d / 2}}{\Gamma(d / 2+1)} .$	$\lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=0}^N \mu_n \left((1 + \Delta)^{-d/2} \right) = \frac{\alpha}{\Gamma(d/2 + 1)} = \frac{\pi^{d/2}}{\Gamma(d/2 + 1)}.$	$\lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=0}^N \mu_n \left((1 + \Delta)^{-d/2} \right) = \frac{\alpha}{\Gamma(d/2+1)} = \frac{\pi^{d/2}}{\Gamma(d/2+1)}.$
cardona-qft-methods_ EQ0847 (13)	35	■ 1.000	$\not{D}=-i\,c\,\circ\nabla^S.$	$\not{D} = -i c \circ \nabla^S.$	$\not{D} = -i\,c\,\circ\,\nabla^S.$
cardona-qft-methods_ EQ0987	50	■ 1.000	$f\,A\,\mathcal{D}^{-1}=0,\forall A\in\Omega^1_{\mathcal{D}}(\mathcal{A})\iff f\,a\,b=f\,a\alpha(b),\forall a,b\in\mathcal{A},$	$fAD^{-1}=0,\forall A\in\Omega^1_{\mathcal{D}}(\mathcal{A})\iff fab=f\alpha(b),\forall a,b\in\mathcal{A},$	$f\,AD^{-1}=0,\forall A\in\Omega^1_{\mathcal{D}}(\mathcal{A})\iff f\,ab=f\,a\alpha(b),\forall a,b\in\mathcal{A},$
cardona-qft-methods_ EQ0125	77	■ 1.000	$D^{\alpha}:=D_1^{\alpha_1}D_2^{\alpha_1}\cdots D_n^{\alpha_n}:C^{\infty}(\mathbb{R}^n)\longrightarrow C^{\infty}(\mathbb{R}^n).$	$D^{\alpha} := D_1^{\alpha_1} D_2^{\alpha_1} \cdots D_n^{\alpha_n} : C^{\infty}(\mathbb{R}^n) \longrightarrow C^{\infty}(\mathbb{R}^n).$	$D^{\alpha} := D_1^{\alpha_1} D_2^{\alpha_1} \cdots D_n^{\alpha_n} : C^{\infty}(\mathbb{R}^n) \longrightarrow C^{\infty}(\mathbb{R}^n).$
cardona-qft-methods_ EQ0135	79	■ 1.000	$\mathcal{D}:H_K^m\left(\mathbb{R}^n,\mathbb{R}^N\right)\longrightarrow L^2\left(\mathbb{R}^n,\mathbb{R}^N\right),\quad\text{for }m>k.$	$\mathcal{D}:H_K^m\left(\mathbb{R}^n,\mathbb{R}^N\right)\longrightarrow L^2\left(\mathbb{R}^n,\mathbb{R}^N\right),\quad\text{for }m>k.$	$\mathcal{D}:H_K^m(\mathbb{R}^n,\mathbb{R}^N)\longrightarrow L^2(\mathbb{R}^n,\mathbb{R}^N),\quad\text{for }m>k.$
cardona-qft-methods_ EQ0162	87	■ 1.000	$\mathcal{D}:C^{\infty}(M,E)\longrightarrow C^{\infty}(M,E),$	$\mathcal{D}:C^{\infty}(M,E)\longrightarrow C^{\infty}(M,E),$	$\mathcal{D}:C^{\infty}(M,E)\longrightarrow C^{\infty}(M,E),$
cardona-qft-methods_ EQ0187 (2)	93	■ 1.000	$\sigma_1=\left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right),\quad \sigma_2=\left(\begin{array}{cc} 0 & -i \\ i & 0 \end{array}\right),\quad \sigma_3=\left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right).$	$\sigma_1 = \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array} \right), \quad \sigma_2 = \left(\begin{array}{cc} 0 & -i \\ i & 0 \end{array} \right), \quad \sigma_3 = \left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array} \right).$	$\sigma_1 = \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array} \right), \quad \sigma_2 = \left(\begin{array}{cc} 0 & -i \\ i & 0 \end{array} \right), \quad \sigma_3 = \left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array} \right).$
cardona-qft-methods_ EQ0283	137	■ 1.000	$\mathbb{L}\cdot\left[\left(Y_{G\backslash e}\cup Y_{G/e}\right)\backslash Y_{G\backslash e}\right],$	$\mathbb{L}\cdot\left[\left(Y_{G\backslash e}\cup Y_{G/e}\right)\backslash Y_{G\backslash e}\right],$	$\mathbb{L}\cdot\left[\left(Y_{G\setminus e}\cup Y_{G/e}\right)\setminus Y_{G\setminus e}\right],$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0287	138	■ 1.000	$T(1+T)^{\{2\}} \quad , \quad \quad \quad (1+T)^{\{2\}} \quad \quad , \quad \quad T(1+T)$	$T(1+T)^2 \quad , \quad (1+T)^2 \quad , \quad T(1+T)$	$T(1+T)^2 \quad , \quad (1+T)^2 \quad , \quad T(1+T)$
cardona-qft-methods_ EQ0296	143	■ 1.000	$\mathbb{U}(G)=\left\{\begin{array}{l} 0 \quad \text{mod}(\mathbb{L}) \quad \text{if } G \text{ has edges that are not looping edges} \\ (-1)^n \quad \text{mod}(\mathbb{L}) \quad \text{if } G \text{ has } n \text{ looping edges and no other edge.} \end{array}\right.$	$\mathbb{U}(G)=\left\{\begin{array}{l} 0 \quad \text{mod}(\mathbb{L}) \quad \text{if } G \text{ has edges that are not looping edges} \\ (-1)^n \quad \text{mod}(\mathbb{L}) \quad \text{if } G \text{ has } n \text{ looping edges and no other edge.} \end{array}\right.$	$\mathbb{U}(G)=\left\{\begin{array}{l} 0 \quad \text{mod}(\mathbb{L}) \quad \text{if } G \text{ has edges that are not looping edges} \\ (-1)^n \quad \text{mod}(\mathbb{L}) \quad \text{if } G \text{ has } n \text{ looping edges and no other edge.} \end{array}\right.$
cardona-qft-methods_ EQ0329 (16)	162	■ 1.000	$\begin{aligned} R_b^a &= d\omega_b^a + \omega_c^a \wedge \omega_b^c \\ &= \frac{1}{2} R_{b\mu\nu}^a dx^\mu \wedge dx^\nu \end{aligned}$	$\begin{aligned} R_b^a &= d\omega_b^a + \omega_c^a \wedge \omega_b^c \\ &= \frac{1}{2} R_{b\mu\nu}^a dx^\mu \wedge dx^\nu \end{aligned}$	$\begin{aligned} R_b^a &= d\omega_b^a + \omega_c^a \wedge \omega_b^c \\ &= \frac{1}{2} R_{b\mu\nu}^a dx^\mu \wedge dx^\nu. \end{aligned}$
cardona-qft-methods_ EQ0334 (20)	163	■ 1.000	$D R_b^a = d R_b^a + \omega_c^a \wedge R_b^c - \omega_b^c \wedge R_c^a \equiv 0.$	$DR_b^a = dR_b^a + \omega_c^a \wedge R_b^c - \omega_b^c \wedge R_c^a \equiv 0.$	$DR_b^a = dR_b^a + \omega_c^a \wedge R_b^c - \omega_b^c \wedge R_c^a \equiv 0.$
cardona-qft-methods_ EQ0343 (11)	167	■ 1.000	$\begin{aligned} I_2 &= \kappa \int_M \epsilon_{ab} [\alpha_1 R^{ab} + \alpha_0 e^a e^b] \\ &= \kappa \int_M \sqrt{ g } (\alpha_1 R + 2\alpha_0) d^2x \\ &= \kappa \alpha_1 \cdot \mathbf{E}_2 + 2\kappa \alpha_0 \cdot V_2 \end{aligned}$	$\begin{aligned} I_2 &= \kappa \int_M \epsilon_{ab} [\alpha_1 R^{ab} + \alpha_0 e^a e^b] \\ &= \kappa \int_M \sqrt{ g } (\alpha_1 R + 2\alpha_0) d^2x \\ &= \kappa \alpha_1 \cdot \mathbf{E}_2 + 2\kappa \alpha_0 \cdot V_2 \end{aligned}$	$\begin{aligned} I_2 &= \kappa \int_M \epsilon_{ab} [\alpha_1 R^{ab} + \alpha_0 e^a e^b] \\ &= \kappa \int_M \sqrt{ g } (\alpha_1 R + 2\alpha_0) d^2x \\ &= \kappa \alpha_1 \cdot \mathbf{E}_2 + 2\kappa \alpha_0 \cdot V_2. \end{aligned}$
cardona-qft-methods_ EQ0351 (20)	168	■ 1.000	$\omega^a{}_{b\mu} = -E_b^\nu (\partial_\mu e_\nu^a - \Gamma_{\mu\nu}^\lambda e_\lambda^a),$	$\omega^a{}_{b\mu} = -E_b^\nu (\partial_\mu e_\nu^a - \Gamma_{\mu\nu}^\lambda e_\lambda^a),$	$\omega^a{}_{b\mu} = -E_b^\nu (\partial_\mu e_\nu^a - \Gamma_{\mu\nu}^\lambda e_\lambda^a),$
cardona-qft-methods_ EQ0354 (22)	170	■ 1.000	$v_1=e^a e^b R_{ab}, \quad \tau_0=T^a T_a, \quad \mathbf{P}_4=R^{ab} R_{ab}.$	$v_1=e^a e^b R_{ab}, \quad \tau_0=T^a T_a, \quad \mathbf{P}_4=R^{ab} R_{ab}.$	$v_1=e^a e^b R_{ab}, \quad \tau_0=T^a T_a, \quad \mathbf{P}_4=R^{ab} R_{ab}.$
cardona-qft-methods_ EQ0355 (23)	170	■ 1.000	$\mathbf{N}_4=T^a T_a-e^a e^b R_{ab},$	$\mathbf{N}_4=T^a T_a-e^a e^b R_{ab},$	$\mathbf{N}_4=T^a T_a-e^a e^b R_{ab},$
cardona-qft-methods_ EQ0378 (19)	177	■ 1.000	$S \, 0(D-1,1) \hookrightarrow I \, S \, 0(D-1,1) \, .$	$SO(D-1,1) \hookrightarrow ISO(D-1,1).$	$SO(D-1,1) \hookrightarrow ISO(D-1,1).$
cardona-qft-methods_ EQ0380 (21)	178	■ 1.000	$\begin{aligned} \delta L_4 &= 2\epsilon_{abcd} R^{ab} e^c \delta e^d \\ &= d \left(2\epsilon_{abcd} R^{ab} e^c \lambda^d \right) - 2\epsilon_{abcd} R^{ab} T^c \lambda^d \end{aligned}$	$\begin{aligned} \delta L_4 &= 2\epsilon_{abcd} R^{ab} e^c \delta e^d \\ &= d \left(2\epsilon_{abcd} R^{ab} e^c \lambda^d \right) - 2\epsilon_{abcd} R^{ab} T^c \lambda^d \end{aligned}$	$\begin{aligned} \delta L_4 &= 2\epsilon_{abcd} R^{ab} e^c \delta e^d \\ &= d(2\epsilon_{abcd} R^{ab} e^c \lambda^d) - 2\epsilon_{abcd} R^{ab} T^c \lambda^d. \end{aligned}$
cardona-qft-methods_ EQ0388	179	■ 1.000	$d\left(\delta L_3^{AdS}\right)=0,$	$d\left(\delta L_3^{AdS}\right)=0,$	$d\left(\delta L_3^{AdS}\right)=0,$
cardona-qft-methods_ EQ0414 (45)	188	■ 1.000	$\bar{\Lambda}(x)-\Lambda(x)=0, \quad \text{for } x \in \partial M.$	$\bar{\Lambda}(x)-\Lambda(x)=0, \quad \text{for } x \in \partial M.$	$\bar{\Lambda}(x)-\Lambda(x)=0, \quad \text{for } x \in \partial M.$
cardona-qft-methods_ EQ0443	211	■ 1.000	$D(Y)=\left(\begin{array}{cc} S & T^* \\ T & \bar{S} \end{array}\right) \quad \text{with} \quad S=S_1 \oplus (S_3 \otimes 1_3)$	$D(Y)=\left(\begin{array}{cc} S & T^* \\ T & \bar{S} \end{array}\right) \quad \text{with} \quad S=S_1 \oplus (S_3 \otimes 1_3)$	$D(Y)=\left(\begin{array}{cc} S & T^* \\ T & \bar{S} \end{array}\right) \quad \text{with} \quad S=S_1 \oplus (S_3 \otimes 1_3),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0444	211	■ 1.000	$S_{\{1\}}=\left(\begin{array}{cccc} 0 & 0 & Y_{(\uparrow 1)}^* & 0 \\ 0 & 0 & 0 & Y_{(\downarrow 1)}^* \\ Y_{(\uparrow 1)} & 0 & 0 & 0 \\ 0 & Y_{(\downarrow 1)} & 0 & 0 \end{array}\right)$	$S_1=\left(\begin{array}{cccc} 0 & 0 & Y_{(\uparrow 1)}^* & 0 \\ 0 & 0 & 0 & Y_{(\downarrow 1)}^* \\ Y_{(\uparrow 1)} & 0 & 0 & 0 \\ 0 & Y_{(\downarrow 1)} & 0 & 0 \end{array}\right)$	$S_1=\left(\begin{array}{cccc} 0 & 0 & Y_{(\uparrow 1)}^* & 0 \\ 0 & 0 & 0 & Y_{(\downarrow 1)}^* \\ Y_{(\uparrow 1)} & 0 & 0 & 0 \\ 0 & Y_{(\downarrow 1)} & 0 & 0 \end{array}\right)$
cardona-qft-methods_ EQ0514 (7)	230	■ 1.000	$M_{\{P\}^{\wedge 2}(\backslash\Lambda\bda)=M_{\{P\}^{\wedge 2}+2\ \rho_{\{0\}}\ \backslash\Lambda\bda^{\wedge 2}},$	$M_P^2(\Lambda)=M_P^2+2\rho_0\Lambda^2,$	$M_P^2(\Lambda)=M_P^2+2\rho_0\Lambda^2,$
cardona-qft-methods_ EQ0547 (17)	244	■ 1.000	$(\mathrm{Id}\,\,\otimes\,\,\epsilon)\,\Delta=\mathrm{Id}=(\epsilon\,\,\otimes\,\,\mathrm{Id})\Delta=(\epsilon\,\,\otimes\,\,\epsilon)\,\Delta.$	$(\mathrm{Id}\,\,\otimes\,\,\epsilon)\Delta=\mathrm{Id}=(\epsilon\,\,\otimes\,\,\mathrm{Id})\Delta.$	$(\mathrm{Id}\,\,\otimes\,\,\epsilon)\Delta=\mathrm{Id}=(\epsilon\,\,\otimes\,\,\mathrm{Id})\Delta.$
cardona-qft-methods_ EQ0568 (2)	248	■ 1.000	$\widetilde{R}(\omega)=\int_{-\infty}^{\infty} dt e^{i\omega t} R(t)=\int_0^{\infty} e^{i\omega t} R(t)$	$\widetilde{R}(\omega)=\int_{-\infty}^{\infty} dt e^{i\omega t} R(t)=\int_0^{\infty} e^{i\omega t} R(t)$	$\widetilde{R}(\omega)=\int_{-\infty}^{\infty} dt e^{i\omega t} R(t)=\int_0^{\infty} e^{i\omega t} R(t)$
cardona-qft-methods_ EQ0573 (8)	249	■ 1.000	$f\ast g(x)=f\,e^{\frac{i}{2}\,\mathrm{vec}\{\partial_{\mu}\}_{\mu}\,\theta^{\wedge}\{\mu\,N\}\,\mathrm{vec}\{\partial_{\mu}\}_{\mu}}\,g$	$f\ast g(x)=fe^{\frac{i}{2}\vec{\partial}_{\mu}\theta^{\mu N}\vec{\partial}_{\nu}}g$	$f\ast g(x)=fe^{\frac{i}{2}\vec{\partial}_{\mu}\theta^{\mu N}\vec{\partial}_{\nu}}g$
cardona-qft-methods_ EQ0621 (29)	256	■ 1.000	$m_{\{\theta\}}(\alpha\otimes\beta)=m_{\{0\}}\,\mathcal{F}_{\{\theta\}}(\alpha\otimes\beta)\quad\alpha,\beta\in\mathcal{A}_{\theta}(M^{d+1}),\quad\mathcal{F}_{\theta}=e^{\frac{i}{2}\theta_{\mu\nu}\partial_{\mu}\otimes\partial_{\nu}},$	$m_{\theta}(\alpha\otimes\beta)=m_0\mathcal{F}_{\theta}(\alpha\otimes\beta)\quad\alpha,\beta\in\mathcal{A}_{\theta}(M^{d+1}),\quad\mathcal{F}_{\theta}=e^{\frac{i}{2}\theta_{\mu\nu}\partial_{\mu}\otimes\partial_{\nu}},$	$m_{\theta}(\alpha\otimes\beta)=m_0\mathcal{F}_{\theta}(\alpha\otimes\beta)\quad\alpha,\beta\in\mathcal{A}_{\theta}(M^{d+1}),\quad\mathcal{F}_{\theta}=e^{\frac{i}{2}\theta_{\mu\nu}\partial_{\mu}\otimes\partial_{\nu}},$
cardona-qft-methods_ EQ0634	258	■ 1.000	$\begin{array}{l} \int \prod_i d\mu(p_i) U_{\theta}(\Lambda) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle \exp^{\frac{i}{2}(\Lambda p_1) \wedge (\Lambda p_2)} \exp^{\frac{i}{2}(p_1) \wedge (p_2)} e_{\Lambda p_1} \otimes e_{\Lambda p_2} = \\ \int \prod_i d\mu(p_i) U_{\theta}(\Lambda) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e_{\Lambda p_1} \otimes e_{\Lambda p_2} = \\ \int \prod_i d\mu(p_i) U_{\theta}(\Lambda) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e_{\Lambda p_1} \otimes e_{\Lambda p_2} \end{array}$	$\int \prod_i d\mu(p_i) U_{\theta}(\Lambda) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle \exp^{\frac{i}{2}(\Lambda p_1) \wedge (\Lambda p_2)} \exp^{\frac{i}{2}(p_1) \wedge (p_2)} e_{\Lambda p_1} \otimes e_{\Lambda p_2} = \\ \int \prod_i d\mu(p_i) U_{\theta}(\Lambda) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e_{\Lambda p_1} \otimes e_{\Lambda p_2} = \\ \int \prod_i d\mu(p_i) U_{\theta}(\Lambda) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e_{\Lambda p_1} \otimes e_{\Lambda p_2}$	$\int \prod_i d\mu(p_i) U_{\theta}(\Lambda) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle \exp^{\frac{i}{2}(\Lambda p_1) \wedge (\Lambda p_2)} \exp^{\frac{i}{2}(p_1) \wedge (p_2)} e_{\Lambda p_1} \otimes e_{\Lambda p_2} = \\ \int \prod_i d\mu(p_i) U_{\theta}(\Lambda) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e_{\Lambda p_1} \otimes e_{\Lambda p_2} = \\ \int \prod_i d\mu(p_i) U_{\theta}(\Lambda) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e_{\Lambda p_1} \otimes e_{\Lambda p_2}$
cardona-qft-methods_ EQ0643 (51)	259	■ 1.000	$\begin{aligned} e_{\{p_1\}}\otimes_{S_{\theta}}e_{\{p_2\}} &= \frac{1}{2}[e_{p_1}\otimes e_{p_2} + \mathcal{F}_{\theta}^{-2}e_{p_2}\otimes e_{p_1}] \\ &= \frac{1}{2}[e_{p_1}\otimes e_{p_2} + e^{ip_2\wedge p_1}e_{p_2}\otimes e_{p_1}] \\ &= e^{ip_2\wedge p_1}e_{p_2}\otimes e_{p_1}. \end{aligned}$	$e_{p_1}\otimes_{S_{\theta}}e_{p_2}=\frac{1}{2}[e_{p_1}\otimes e_{p_2}+\mathcal{F}_{\theta}^{-2}e_{p_2}\otimes e_{p_1}]$ $=\frac{1}{2}[e_{p_1}\otimes e_{p_2}+e^{ip_2\wedge p_1}e_{p_2}\otimes e_{p_1}]$ $=e^{ip_2\wedge p_1}e_{p_2}\otimes e_{p_1}.$	$e_{p_1}\otimes_{S_{\theta}}e_{p_2}=\frac{1}{2}[e_{p_1}\otimes e_{p_2}+\mathcal{F}_{\theta}^{-2}e_{p_2}\otimes e_{p_1}]$ $=\frac{1}{2}[e_{p_1}\otimes e_{p_2}+e^{ip_2\wedge p_1}e_{p_2}\otimes e_{p_1}]$ $=e^{ip_2\wedge p_1}e_{p_2}\otimes e_{p_1}.$
cardona-qft-methods_ EQ0646 (53)	260	■ 1.000	$a_{\{p_1\}}^{\dagger}a_{\{p_2\}}^{\dagger}=e^{ip_1\wedge p_2}a_{p_2}^{\dagger}a_{p_1}^{\dagger}.$	$a_{p_1}^{\dagger}a_{p_2}^{\dagger}=e^{ip_1\wedge p_2}a_{p_2}^{\dagger}a_{p_1}^{\dagger}.$	$a_{p_1}^{\dagger}a_{p_2}^{\dagger}=e^{ip_1\wedge p_2}a_{p_2}^{\dagger}a_{p_1}^{\dagger}.$
cardona-qft-methods_ EQ0660 (10)	262	■ 1.000	$\left[H_I(x),\ H_I(y)\right]=0,$	$[H_I(x),H_I(y)]=0,$	$[H_I(x),H_I(y)]=0,$
cardona-qft-methods_ EQ0684	269	■ 1.000	$S^{\{+\}}=\frac{1}{2}\,\sigma^{\{+\}}=\left(\begin{array}{ll} 0 & 1 \\ 0 & 0 \end{array}\right),\quad\quad S^{\{-\}}=\frac{1}{2}\,\sigma^{\{-\}}=\left(\begin{array}{ll} 0 & 0 \\ 1 & 0 \end{array}\right),$	$S^+=\frac{1}{2}\sigma^+=\left(\begin{array}{cc} 0 & 1 \\ 0 & 0 \end{array}\right),\quad\quad S^-=\frac{1}{2}\sigma^-=\left(\begin{array}{cc} 0 & 0 \\ 1 & 0 \end{array}\right),$	$S^+=\frac{1}{2}\sigma^+=\left(\begin{array}{cc} 0 & 1 \\ 0 & 0 \end{array}\right),\quad\quad S^-=\frac{1}{2}\sigma^-=\left(\begin{array}{cc} 0 & 0 \\ 1 & 0 \end{array}\right),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E0695 (10)	271	■ 1.000	$\begin{aligned} \hat{H} &= \sum_{l=1}^L \left(1 - \vec{\sigma}_l \cdot \vec{\sigma}_{l+1} \right) \\ &= \sum_{l=1}^L \left(1 - \sigma_l^3 \sigma_{l+1}^3 - \frac{1}{2} \sigma_l^+ \sigma_{l+1}^- - \frac{1}{2} \sigma_l^- \sigma_{l+1}^+ \right) \\ &= 2 \sum_{l=1}^L (\mathbb{I}_{l,l+1} - \mathbb{P}_{l,l+1}) . \end{aligned}$	$\hat{H} = \sum_{l=1}^L (1 - \vec{\sigma}_l \cdot \vec{\sigma}_{l+1})$ $= \sum_{l=1}^L \left(1 - \sigma_l^3 \sigma_{l+1}^3 - \frac{1}{2} \sigma_l^+ \sigma_{l+1}^- - \frac{1}{2} \sigma_l^- \sigma_{l+1}^+ \right)$ $= 2 \sum_{l=1}^L (\mathbb{I}_{l,l+1} - \mathbb{P}_{l,l+1}) .$	$\hat{H} = \sum_{l=1}^L (1 - \vec{\sigma}_l \cdot \vec{\sigma}_{l+1})$ $= \sum_{l=1}^L \left(1 - \sigma_l^3 \sigma_{l+1}^3 - \frac{1}{2} \sigma_l^+ \sigma_{l+1}^- - \frac{1}{2} \sigma_l^- \sigma_{l+1}^+ \right)$ $= 2 \sum_{l=1}^L (\mathbb{I}_{l,l+1} - \mathbb{P}_{l,l+1}) .$
cardona-qft-methods_ E0718	275	■ 1.000	$R_{\{a, b\}} \left(u - u' \right) M_{\{a\}}(u) M_{\{b\}} \left(u' \right) = M_{\{b\}} \left(u' \right) M_{\{a\}}(u) R_{\{a, b\}} \left(u - u' \right) .$	$R_{a,b}(u-u') M_a(u) M_b(u') = M_b(u') M_a(u) R_{a,b}(u-u') .$	$R_{a,b}(u-u') M_a(u) M_b(u') = M_b(u') M_a(u) R_{a,b}(u-u') .$
cardona-qft-methods_ E0766 (7)	293	■ 1.000	$T^{\alpha\beta} \equiv -\frac{2\pi}{\sqrt{\gamma}} \frac{\delta S}{\delta \gamma_{\alpha\beta}} = -\frac{1}{\alpha'} \left(\partial^\alpha X^M \partial^\beta X_M - \frac{1}{2} \gamma^{\alpha\beta} \gamma_{\lambda\rho} \partial^\lambda X^M \partial^\rho X_M \right) = 0 .$	$T^{\alpha\beta} \equiv -\frac{2\pi}{\sqrt{\gamma}} \frac{\delta S}{\delta \gamma_{\alpha\beta}} = -\frac{1}{\alpha'} \left(\partial^\alpha X^M \partial^\beta X_M - \frac{1}{2} \gamma^{\alpha\beta} \gamma_{\lambda\rho} \partial^\lambda X^M \partial^\rho X_M \right) = 0 .$	$T^{\alpha\beta} \equiv -\frac{2\pi}{\sqrt{\gamma}} \frac{\delta S}{\delta \gamma_{\alpha\beta}} = -\frac{1}{\alpha'} \left(\partial^\alpha X^M \partial^\beta X_M - \frac{1}{2} \gamma^{\alpha\beta} \gamma_{\lambda\rho} \partial^\lambda X^M \partial^\rho X_M \right) = 0 .$
cardona-qft-methods_ E0775 (16)	295	■ 1.000	$\begin{aligned} Z &= \int \mathcal{D}X \mathcal{D}\gamma e^{-S} \\ &= \int \mathcal{D}X \mathcal{D}\gamma e^{-S_0} \left(1 + \frac{1}{4\pi\alpha'} \int d\sigma d\tau \sqrt{\gamma} \gamma^{\alpha\beta} h_{MN} \partial_\alpha X^M \partial_\beta X^N + \dots \right) , \end{aligned}$	$Z = \int \mathcal{D}X \mathcal{D}\gamma e^{-S}$ $= \int \mathcal{D}X \mathcal{D}\gamma e^{-S_0} \left(1 + \frac{1}{4\pi\alpha'} \int d\sigma d\tau \sqrt{\gamma} \gamma^{\alpha\beta} h_{MN} \partial_\alpha X^M \partial_\beta X^N + \dots \right)$	$Z = \int \mathcal{D}X \mathcal{D}\gamma e^{-S} \quad (16)$ $= \int \mathcal{D}X \mathcal{D}\gamma e^{-S_0} \left(1 + \frac{1}{4\pi\alpha'} \int d\sigma d\tau \sqrt{\gamma} \gamma^{\alpha\beta} h_{MN} \partial_\alpha X^M \partial_\beta X^N + \dots \right) ,$
cardona-qft-methods_ E0789 (21)	297	■ 1.000	$\begin{aligned} \mathrm{S} &= \frac{1}{2} \kappa_0^2 \int d^D X (-G)^{1/2} e^{-2\Phi} \left[R + 4 \nabla_M \Phi \nabla^M \Phi - \frac{1}{12} H_{MNP} H^{MNP} - \frac{2(D-26)}{3\alpha'} + O(\alpha') \right] \end{aligned}$	$S = \frac{1}{2\kappa_0^2} \int d^D X (-G)^{1/2} e^{-2\Phi} \left[R + 4 \nabla_M \Phi \nabla^M \Phi - \frac{1}{12} H_{MNP} H^{MNP} - \frac{2(D-26)}{3\alpha'} + O(\alpha') \right]$	$S = \frac{1}{2\kappa_0^2} \int d^D X (-G)^{1/2} e^{-2\Phi} \left[R + 4 \nabla_M \Phi \nabla^M \Phi - \frac{1}{12} H_{MNP} H^{MNP} - \frac{2(D-26)}{3\alpha'} + O(\alpha') \right] .$
cardona-qft-methods_ E0819 (19)	310	■ 1.000	$\begin{aligned} \Phi_- &= -\frac{i}{8} \Omega \leftrightarrow \mathcal{J}_1 \\ \Phi_+ &= \frac{1}{8} e^{-iJ} \leftrightarrow \mathcal{J}_2 \end{aligned}$	$\Phi_- = -\frac{i}{8} \Omega \leftrightarrow \mathcal{J}_1$ $\Phi_+ = \frac{1}{8} e^{-iJ} \leftrightarrow \mathcal{J}_2$	$\Phi_- = -\frac{i}{8} \Omega \leftrightarrow \mathcal{J}_1$ $\Phi_+ = \frac{1}{8} e^{-iJ} \leftrightarrow \mathcal{J}_2$
cardona-qft-methods_ E0851	330	■ 1.000	$[df, dg] := d\{f, g\}, \forall f, g \in \mathcal{C}^\infty(M),$	$[df, dg] := d\{f, g\}, \forall f, g \in \mathcal{C}^\infty(M),$	$[df, dg] := d\{f, g\}, \forall f, g \in \mathcal{C}^\infty(M),$
cardona-qft-methods_ E0863	334	■ 1.000	$L_3 \circ (L_1 \times L_1) = L_1 .$	$L_3 \circ (L_1 \times L_1) = L_1 .$	$L_3 \circ (L_1 \times L_1) = L_1 .$
cardona-qft-methods_ E0947 (7)	359	■ 1.000	$[X] - [X \cap Y] = [X \backslash (X \cap Y)] = [(X \cup Y) \backslash Y] = [X \cup Y] - [Y],$	$[X] - [X \cap Y] = [X \setminus (X \cap Y)] = [(X \cup Y) \setminus Y] = [X \cup Y] - [Y],$	$[X] - [X \cap Y] = [X \setminus (X \cap Y)] = [(X \cup Y) \setminus Y] = [X \cup Y] - [Y],$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0992	372	■ 1.000	$\backslash\mathrm{cal}\{S\}=\mathrm{text}\{ \text{ the multiplicative set generated by } \}$ $\backslash\mathrm{left}\{ \{ : 1+E^{\dagger} E \mid E \in \mathcal{E} \} \backslash\mathrm{right}\}$	$\mathcal{S} = \text{ the multiplicative set generated by } \{ : 1 + E^{\dagger} E \mid E \in \mathcal{E} : \}$	$\mathcal{S} = \text{ the multiplicative set generated by } \{ : 1 + E^{\dagger} E \mid E \in \mathcal{E} : \}$
cardona-qft-methods_ EQ0001	18	■ 1.000	$\backslash\mathrm{left} \backslash\mathrm{partial}_{\{x\}}^{\{\alpha\}} \backslash\mathrm{partial}_{\{x\}}^{\{\beta\}}\backslash\mathrm{left}(\backslash\mathrm{sigma}-\backslash\mathrm{sum}_{\{j<N\}} \backslash\mathrm{sigma}_{\{m-j\}}\backslash\mathrm{right}\})(x, \{x\})\backslash\mathrm{right} \backslash\mathrm{leq} \backslash\mathrm{C}_{\{N \alpha \beta\}}(x) \{x\} ^{\{\Re(m)-N- \beta \}}$	$\left \partial_x^\alpha \partial_\xi^\beta \left(\sigma - \sum_{j < N} \sigma_{m-j} \right) (x, \xi) \right \leq C_{N\alpha\beta}(x) \xi ^{\Re(m)-N- \beta }$	$ \partial_x^\alpha \partial_\xi^\beta (\sigma - \sum_{j < N} \sigma_{m-j})(x, \xi) \leq C_{N\alpha\beta}(x) \xi ^{\Re(m)-N- \beta }$
cardona-qft-methods_ EQ0003 (1)	18	■ 1.000	$\backslash\mathrm{sigma}(\backslash\mathrm{cdot}, D)(u)(x)=\backslash\mathrm{sigma}(x, D)(u)=\backslash\mathrm{frac}\{1\}\{(2 \pi)^{\{d\}} \backslash\mathrm{int}_{\{\backslash\mathrm{mathbb{R}}^{\{d\}}\}} \backslash\mathrm{sigma}(x, \{x\}) \backslash\mathrm{widehat}\{u\}(\{x\}) e^{\{i x \cdot \mathrm{cdot} \{x\}} d \{x\}}$	$\sigma(\cdot, D)(u)(x) = \sigma(x, D)(u) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \sigma(x, \xi) \widehat{u}(\xi) e^{ix \cdot \xi} d\xi$	$\sigma(\cdot, D)(u)(x) = \sigma(x, D)(u) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \sigma(x, \xi) \widehat{u}(\xi) e^{ix \cdot \xi} d\xi$
cardona-qft-methods_ EQ0004	18	■ 1.000	$k^{\{\backslash\mathrm{sigma}(x, D)\}}(x, y)=\backslash\mathrm{frac}\{1\}\{(2 \pi)^{\{d\}} \backslash\mathrm{int}_{\{\backslash\mathrm{mathbb{R}}^{\{d\}}\}} \backslash\mathrm{sigma}(x, \{x\}) e^{\{i(x-y) \cdot \mathrm{cdot} \{x\}} d \{x\}}=\backslash\mathrm{check}\{\backslash\mathrm{sigma}\}_{\{x\}} \backslash\mathrm{rightarrow} y\}(x, x-y)$	$k^{\sigma(x, D)}(x, y) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \sigma(x, \xi) e^{i(x-y) \cdot \xi} d\xi = \check{\sigma}_{\xi \rightarrow y}(x, x - y)$	$k^{\sigma(x, D)}(x, y) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} \sigma(x, \xi) e^{i(x-y) \cdot \xi} d\xi = \check{\sigma}_{\xi \rightarrow y}(x, x - y)$
cardona-qft-methods_ EQ0005 (2)	18	■ 1.000	$k^{\{a\}}(x, y)=\backslash\mathrm{frac}\{1\}\{(2 \pi)^{\{d\}} \backslash\mathrm{int}_{\{\backslash\mathrm{mathbb{R}}^{\{d\}}\}} a(x, y, \{x\}) e^{\{i(x-y) \cdot \mathrm{cdot} \{x\}} d \{x\}}$	$k^a(x, y) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} a(x, y, \xi) e^{i(x-y) \cdot \xi} d\xi$	$k^a(x, y) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} a(x, y, \xi) e^{i(x-y) \cdot \xi} d\xi.$
cardona-qft-methods_ EQ0007	19	■ 1.000	$\backslash\mathrm{sigma}^{\{A\}} \backslash\mathrm{simeq} \backslash\mathrm{sum}_{\{\alpha\}} \backslash\mathrm{frac}\{(-i)^{\{\alpha\}}\}\{\alpha!\} \backslash\mathrm{partial}_{\{x\}}^{\{\alpha\}} \backslash\mathrm{partial}_{\{y\}}^{\{\alpha\}} k_{\{\backslash\mathrm{sigma}\}^{\{A\}}}(x, y, \{x\})_{\{\mid y=x\}}$	$\sigma^A \simeq \sum_{\alpha} \frac{(-i)^\alpha}{\alpha!} \partial_\xi^\alpha \partial_y^\alpha k_\sigma^A(x, y, \xi) _{y=x}$	$\sigma^A \simeq \sum_{\alpha} \frac{(-i)^\alpha}{\alpha!} \partial_\xi^\alpha \partial_y^\alpha k_\sigma^A(x, y, \xi) _{y=x}$
cardona-qft-methods_ EQ0008	19	■ 1.000	$k_{\{\backslash\mathrm{sigma}\}^{\{A\}}}(x, y)=\backslash\mathrm{frac}\{1\}\{(2 \pi)^{\{d\}} \backslash\mathrm{int}_{\{\backslash\mathrm{mathbb{R}}^{\{d\}}\}} e^{\{i(x-y) \cdot \mathrm{cdot} \{x\}} k^{\{A\}}(x, y, \{x\}) d \{x\}}$	$k_\sigma^A(x, y) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{i(x-y) \cdot \xi} k^A(x, y, \xi) d\xi$	$k_\sigma^A(x, y) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{i(x-y) \cdot \xi} k^A(x, y, \xi) d\xi,$
cardona-qft-methods_ EQ0009	19	■ 1.000	$\backslash\mathrm{sigma}^{\{P_1\} P_2}(x, \{x\}) \backslash\mathrm{simeq} \backslash\mathrm{sum}_{\{\alpha \in \mathbb{N}^d\}} \backslash\mathrm{frac}\{(-i)^{\{\alpha\}}\}\{\alpha!\} \backslash\mathrm{partial}_{\{x\}}^{\{\alpha\}} \backslash\mathrm{sigma}^{\{P_1\}}(x, \{x\}) \backslash\mathrm{partial}_{\{x\}}^{\{\alpha\}} \backslash\mathrm{sigma}^{\{P_2\}}(x, \{x\}) .$	$\sigma^{P_1 P_2}(x, \xi) \simeq \sum_{\alpha \in \mathbb{N}^d} \frac{(-i)^\alpha}{\alpha!} \partial_\xi^\alpha \sigma^{P_1}(x, \xi) \partial_x^\alpha \sigma^{P_2}(x, \xi).$	$\sigma^{P_1 P_2}(x, \xi) \simeq \sum_{\alpha \in \mathbb{N}^d} \frac{(-i)^\alpha}{\alpha!} \partial_\xi^\alpha \sigma^{P_1}(x, \xi) \partial_x^\alpha \sigma^{P_2}(x, \xi).$
cardona-qft-methods_ EQ0011	20	■ 1.000	$\backslash\mathrm{sigma}_{\{m\}}^{\{\phi_{\{*\}} P\}}(x, \{x\})=\backslash\mathrm{sigma}_{\{m\}}^{\{P\}}\backslash\mathrm{left}(\backslash\mathrm{phi}^{\{-1\}}(x), (d \backslash\mathrm{phi})^{\{t\}} \{x\}\backslash\mathrm{right}) .$	$\sigma_m^{\phi_* P}(x, \xi) = \sigma_m^P(\phi^{-1}(x), (d\phi)^t \xi).$	$\sigma_m^{\phi_* P}(x, \xi) = \sigma_m^P(\phi^{-1}(x), (d\phi)^t \xi).$
cardona-qft-methods_ EQ0013 (3)	21	■ 1.000	$\backslash\mathrm{sigma}_{\{-m\}}^{\{P\}}(x, \{x\})=\backslash\mathrm{lim}_{\{t \rightarrow \infty\}} t^{\{-m\}}\backslash\mathrm{left}(e^{\{-i t h\}} \cdot P \cdot e^{\{i t h\}}\backslash\mathrm{right})(x) \backslash\mathrm{text}\{ \text{ for } x \in M, \{x\} \in T_{\{x\}}^{\{*\}} M$	$\sigma_{-m}^P(x, \xi) = \lim_{t \rightarrow \infty} t^{-m} (e^{-ith} \cdot P \cdot e^{ith})(x) \text{ for } x \in M, \xi \in T_x^* M$	$\sigma_{-m}^P(x, \xi) = \lim_{t \rightarrow \infty} t^{-m} (e^{-ith} \cdot P \cdot e^{ith})(x) \text{ for } x \in M, \xi \in T_x^* M$
cardona-qft-methods_ EQ0014	22	■ 1.000	$k^{\{P\}}(x, y)=\backslash\mathrm{sum}_{\{-(m+d) \leq j \leq 0\}} a_{\{j\}}(x, x-y)-c_{\{P\}}(x) \backslash\mathrm{log} x-y +\backslash\mathrm{mathcal}\{0\}(1)$	$k^P(x, y) = \sum_{-(m+d) \leq j \leq 0} a_j(x, x - y) - c_P(x) \log x - y + \mathcal{O}(1)$	$k^P(x, y) = \sum_{-(m+d) \leq j \leq 0} a_j(x, x - y) - c_P(x) \log x - y + \mathcal{O}(1)$
cardona-qft-methods_ EQ0015 (4)	22	■ 1.000	$c_{\{P\}}(x)=\backslash\mathrm{frac}\{1\}\{(2 \pi)^{\{d\}} \backslash\mathrm{int}_{\{\backslash\mathrm{mathbb{S}}^{\{d-1\}}\}} \backslash\mathrm{sigma}_{\{-d\}}^{\{P\}}(x, \{x\}) d \{x\}}$	$c_P(x) = \frac{1}{(2\pi)^d} \int_{\mathbb{S}^{d-1}} \sigma_{-d}^P(x, \xi) d\xi$	$c_P(x) = \frac{1}{(2\pi)^d} \int_{\mathbb{S}^{d-1}} \sigma_{-d}^P(x, \xi) d\xi.$
cardona-qft-methods_ EQ0016	22	■ 1.000	$\backslash\mathrm{operatorname}\{tr\}\backslash\mathrm{left}(k^{\{P\}}(x, y)\backslash\mathrm{right})=\backslash\mathrm{sum}_{\{j=-(m+d)\}^{\{0\}} } a_{\{j\}}(x, x-y)-c_{\{P\}}(x) \backslash\mathrm{log} x-y +\backslash\mathrm{mathcal}\{0\}(1),$	$\mathrm{tr} \left(k^P(x, y) \right) = \sum_{j=-(m+d)}^0 a_j(x, x - y) - c_P(x) \log x - y + \mathcal{O}(1),$	$\mathrm{tr} \left(k^P(x, y) \right) = \sum_{j=-(m+d)}^0 a_j(x, x - y) - c_P(x) \log x - y + \mathcal{O}(1),$
cardona-qft-methods_ EQ0018 (6)	22	■ 1.000	$c_{\{\phi_{\{*\}} P\}}(x)=\phi_{\{*\}}\backslash\mathrm{left}(c_{\{P\}}(x)\backslash\mathrm{right}) .$	$c_{\phi_* P}(x) = \phi_* (c_P(x)).$	$c_{\phi_* P}(x) = \phi_* (c_P(x)).$
cardona-qft-methods_ EQ0020 (8)	24	■ 1.000	$\backslash\mathrm{text}\{ \text{ WRes } P=-\backslash\mathrm{int}_{\{M\}} c_{\{P\}}(x) d x $	$\mathrm{WRes} \, P = - \int_M c_P(x) dx $	$\mathrm{WRes} \, P = - \int_M c_P(x) dx $

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0021	25	■ 1.000	$\begin{aligned} \operatorname{Tr}(\phi P) &= \int_U \phi(x) k^P(x, x) dx \\ &= \int_U \phi(x) L(\sigma(x, \cdot)) dx \end{aligned}$	$\operatorname{Tr}(\phi P) = \int_U \phi(x) k^P(x, x) dx = \frac{1}{(2\pi)^d} \int_U \phi(x) \sigma(x, \xi) d\xi dx $	$\operatorname{Tr}(\phi P) = \int_U \phi(x) k^P(x, x) dx = \frac{1}{(2\pi)^d} \int_U \phi(x) \sigma(x, \xi) d\xi dx $ $= \int_U \phi(x) L(\sigma(x, \cdot)) dx ,$
cardona-qft-methods_ EQ0023 (9)	26	■ 1.000	$\text{if } \phi \in \operatorname{Diff}(M), W \operatorname{Res}(P) = W \operatorname{Res}(\phi_* P)$	$\text{if } \phi \in \operatorname{Diff}(M), W \operatorname{Res}(P) = W \operatorname{Res}(\phi_* P)$	$\text{if } \phi \in \operatorname{Diff}(M), W \operatorname{Res}(P) = W \operatorname{Res}(\phi_* P)$
cardona-qft-methods_ EQ0024	26	■ 1.000	$\left[x^j, \sigma \right] = i \partial_{\xi_j} \sigma, \quad [\xi_j, \sigma] = -i \partial_{x^j} \sigma.$	$[x^j, \sigma] = i \partial_{\xi_j} \sigma, \quad [\xi_j, \sigma] = -i \partial_{x^j} \sigma.$	$[x^j, \sigma] = i \partial_{\xi_j} \sigma, \quad [\xi_j, \sigma] = -i \partial_{x^j} \sigma.$
cardona-qft-methods_ EQ0025	26	■ 1.000	$\sigma_{-d}^P = \sum_{k=1}^d \partial_{\xi_k}^2 \tilde{h} = -i \sum_{k=1}^d [\partial_{\xi_k} \tilde{h}, x^k]$	$\sigma_{-d}^P = \sum_{k=1}^d \partial_{\xi_k}^2 \tilde{h} = -i \sum_{k=1}^d [\partial_{\xi_k} \tilde{h}, x^k]$	$\sigma_{-d}^P = \sum_{k=1}^d \partial_{\xi_k}^2 \tilde{h} = -i \sum_{k=1}^d [\partial_{\xi_k} \tilde{h}, x^k]$
cardona-qft-methods_ EQ0030	28	■ 1.000	$\begin{aligned} \mu_n(T) &= \mu_n(T^*) = \mu_n(T). \\ T \in \mathcal{L}^1(\mathcal{H}) \text{ (meaning } \ T\ _1 = \operatorname{Tr}(T) < \infty) &\iff \sum_{n \in \mathbb{N}} \mu_n(T) < \infty. \\ \mu_n(ATB) &\leq \ A\ \mu_n(T) \ B\ \text{ when } A, B \in \mathcal{B}(\mathcal{H}). \\ \mu_N(UTU^*) &= \mu_N(T) \text{ when } U \text{ is a unitary.} \end{aligned}$	$\mu_n(T) = \mu_n(T^*) = \mu_n(T).$ $T \in \mathcal{L}^1(\mathcal{H}) \text{ (meaning } \ T\ _1 = \operatorname{Tr}(T) < \infty) \iff \sum_{n \in \mathbb{N}} \mu_n(T) < \infty.$ $\mu_n(ATB) \leq \ A\ \mu_n(T) \ B\ \text{ when } A, B \in \mathcal{B}(\mathcal{H}).$ $\mu_N(UTU^*) = \mu_N(T) \text{ when } U \text{ is a unitary.}$	$\mu_n(T) = \mu_n(T^*) = \mu_n(T).$ $T \in \mathcal{L}^1(\mathcal{H}) \text{ (meaning } \ T\ _1 = \operatorname{Tr}(T) < \infty) \iff \sum_{n \in \mathbb{N}} \mu_n(T) < \infty.$ $\mu_n(ATB) \leq \ A\ \mu_n(T) \ B\ \text{ when } A, B \in \mathcal{B}(\mathcal{H}).$ $\mu_N(UTU^*) = \mu_N(T) \text{ when } U \text{ is a unitary.}$
cardona-qft-methods_ EQ0032	29	■ 1.000	$\sigma_N(T) = \inf \{ \ x\ _1 + N \ y\ \mid T = x + y \text{ with } x \in \mathcal{L}^1(\mathcal{H}), y \in \mathcal{K}(\mathcal{H}) \}.$	$\sigma_N(T) = \inf \{ \ x\ _1 + N \ y\ \mid T = x + y \text{ with } x \in \mathcal{L}^1(\mathcal{H}), y \in \mathcal{K}(\mathcal{H}) \}.$	$\sigma_N(T) = \inf \{ \ x\ _1 + N \ y\ \mid T = x + y \text{ with } x \in \mathcal{L}^1(\mathcal{H}), y \in \mathcal{K}(\mathcal{H}) \}.$
cardona-qft-methods_ EQ0033	29	■ 1.000	$\sigma_\lambda(T) = \inf \{ \ x\ _1 + \lambda \ y\ \mid T = x + y \text{ with } x \in \mathcal{L}^1(\mathcal{H}), y \in \mathcal{K}(\mathcal{H}) \}.$	$\sigma_\lambda(T) = \inf \{ \ x\ _1 + \lambda \ y\ \mid T = x + y \text{ with } x \in \mathcal{L}^1(\mathcal{H}), y \in \mathcal{K}(\mathcal{H}) \}.$	$\sigma_\lambda(T) = \inf \{ \ x\ _1 + \lambda \ y\ \mid T = x + y \text{ with } x \in \mathcal{L}^1(\mathcal{H}), y \in \mathcal{K}(\mathcal{H}) \}.$
cardona-qft-methods_ EQ0034	29	■ 1.000	$\mathcal{L}^{1,\infty} = \left\{ T \in \mathcal{K}(\mathcal{H}) \mid \ T\ _{1,\infty} = \sup_{\lambda \geq e} \frac{\sigma_\lambda(T)}{\log \lambda} < \infty \right\}.$	$\mathcal{L}^{1,\infty} = \left\{ T \in \mathcal{K}(\mathcal{H}) \mid \ T\ _{1,\infty} = \sup_{\lambda \geq e} \frac{\sigma_\lambda(T)}{\log \lambda} < \infty \right\}.$	$\mathcal{L}^{1,\infty} = \{ T \in \mathcal{K}(\mathcal{H}) \mid \ T\ _{1,\infty} = \sup_{\lambda \geq e} \frac{\sigma_\lambda(T)}{\log \lambda} < \infty \}.$
cardona-qft-methods_ EQ0036	29	■ 1.000	$\mathcal{L}^{1,+} = \left\{ T \in \mathcal{K}(\mathcal{H}) \mid \ T\ _{1,+} = \sup_{N \geq 2} \frac{\sigma_N(T)}{\log(N)} < \infty \right\}.$	$\mathcal{L}^{1,+} = \left\{ T \in \mathcal{K}(\mathcal{H}) \mid \ T\ _{1,+} = \sup_{N \geq 2} \frac{\sigma_N(T)}{\log(N)} < \infty \right\}.$	$\mathcal{L}^{1,+} = \{ T \in \mathcal{K}(\mathcal{H}) \mid \ T\ _{1,+} = \sup_{N \geq 2} \frac{\sigma_N(T)}{\log(N)} < \infty \}.$
cardona-qft-methods_ EQ0038	29	■ 1.000	$ \tau_\lambda(T_1 + T_2) - \tau_\lambda(T_1) - \tau_\lambda(T_2) \leq \left(\frac{\log 2(2 + \log \log \lambda)}{\log \lambda} \right) \ T_1 + T_2\ _{1,\infty},$	$ \tau_\lambda(T_1 + T_2) - \tau_\lambda(T_1) - \tau_\lambda(T_2) \leq \left(\frac{\log 2(2 + \log \log \lambda)}{\log \lambda} \right) \ T_1 + T_2\ _{1,\infty},$	$ \tau_\lambda(T_1 + T_2) - \tau_\lambda(T_1) - \tau_\lambda(T_2) \leq \left(\frac{\log 2(2 + \log \log \lambda)}{\log \lambda} \right) \ T_1 + T_2\ _{1,\infty},$
cardona-qft-methods_ EQ0042 (12)	32	■ 1.000	$\sigma_1^d(\omega) = i\epsilon(\omega), \quad \sigma_1^{d*}(\omega) = -i\iota(\omega).$	$\sigma_1^d(\omega) = i\epsilon(\omega), \quad \sigma_1^{d*}(\omega) = -i\iota(\omega).$	$\sigma_1^d(\omega) = i\epsilon(\omega), \quad \sigma_1^{d*}(\omega) = -i\iota(\omega).$
cardona-qft-methods_ EQ0045	33	■ 1.000	$D_\nabla = -ic \circ \nabla, \quad \Gamma(M, E) \xrightarrow{\nabla} \Gamma(M, T^*M \otimes E) \xrightarrow{c \otimes 1} \Gamma(M, E),$	$D_\nabla = -ic \circ \nabla, \quad \Gamma(M, E) \xrightarrow{\nabla} \Gamma(M, T^*M \otimes E) \xrightarrow{c \otimes 1} \Gamma(M, E),$	$D_\nabla = -ic \circ \nabla, \quad \Gamma(M, E) \xrightarrow{\nabla} \Gamma(M, T^*M \otimes E) \xrightarrow{c \otimes 1} \Gamma(M, E),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0048	35	■ 1.000	$c\left(d\,x^j\right)\left(x_0\right)=\gamma^j,\quad \sigma_1^D\left(x_0,\xi\right)=c\left(\xi\right)=\gamma^j\xi_j$		
cardona-qft-methods_ EQ0050	36	■ 1.000	$\Delta^S=-\operatorname{Tr}_g\left(\nabla^S\circ\nabla^S\right):\Gamma^\infty(M,S)\rightarrow\Gamma^\infty(M,S).$		
cardona-qft-methods_ EQ0051	37	■ 1.000	$U_{g',g}:=\sqrt{f_{g,g'}}I_{g',g}:\mathcal{H}_{g'}\rightarrow\mathcal{H}_g.$		
cardona-qft-methods_ EQ0053 (15)	37	■ 1.000	$D_{g'}:\mathcal{H}_g\rightarrow\mathcal{H}_g,\quad D_{g'}:=U_{g,g'}^{-1}\mathcal{D}_{g'}U_{g,g'}.$		
cardona-qft-methods_ EQ0054	37	■ 1.000	$\begin{aligned} &\&\not D_{g^{\prime}}I_{g,g^{\prime}}\psi=e^{-h}I_{g,g^{\prime}}\left(\mathcal{D}_g\psi-i\frac{d-1}{2}c_g(\operatorname{grad}h)\psi\right),\\ &\mathcal{D}_{g'}=e^{-\frac{d+1}{2}h}I_{g,g'}\mathcal{D}_gI_{g,g'}^{-1}e^{\frac{d-1}{2}h},\\ &D_{g'}=e^{-h/2}\mathcal{D}_ge^{-h/2}. \end{aligned}$		
cardona-qft-methods_ EQ0056 (16)	38	■ 1.000	$k_t(x,y)=\frac{1}{(4\pi t)^{d/2}}e^{- x-y ^2/4t}\text{ for }x\neq y$		
cardona-qft-methods_ EQ0057	38	■ 1.000	$\partial_t k_t(x,y)=\Delta_x k_t(x,y),\quad \forall t>0,x,y\in\mathbb{R}^d$		
cardona-qft-methods_ EQ0058	39	■ 1.000	$-\Delta\phi=\lambda\psi\text{ in }U,\quad \psi=0\text{ on }\partial U.$		
cardona-qft-methods_ EQ0060	39	■ 1.000	$k_t(x,x)\underset{t\downarrow 0^+}{\sim}\sum_{k=0}^{\infty}a_k(x)t^{(-d+k)/m}$		
cardona-qft-methods_ EQ0061 (17)	39	■ 1.000	$k(t,f,P)\underset{t\downarrow 0^+}{\sim}\sum_{k=0}^{\infty}a_k(f,P)t^{(-d+k)/m}.$		
cardona-qft-methods_ EQ0062 (18)	40	■ 1.000	$P=-\left(g^{\mu\nu}\partial_\mu\partial_\nu+\mathbb{A}^\mu\partial_\mu+\mathbb{B}\right)$		
cardona-qft-methods_ EQ0063	40	■ 1.000	$P=-\left(\operatorname{Tr}_g\nabla^2+E\right),\quad \nabla^2(X,Y):=[\nabla_X,\nabla_Y]-\nabla_{\nabla_X^L C}Y,$		
cardona-qft-methods_ EQ0064	40	■ 1.000	$\operatorname{Tr}_g\nabla^2:=g^{\mu\nu}\left(\nabla_\mu\nabla_\nu-\Gamma_{\mu\nu}^\rho\nabla_\rho\right)$		
cardona-qft-methods_ EQ0067 (20)	41	■ 1.000	$\operatorname{Tr}\left(e^{-tD^2}\right)\underset{t\downarrow 0^+}{\sim}\sum_{j\geq 0}t^{(j-d)/2}a_j\left(D^2\right)$		
cardona-qft-methods_ EQ0071 (23)	43	■ 1.000	$\left[\pi(a),\pi(b)^\circ\right]=0,\quad \left[[\mathcal{D},\pi(a)],\pi(b)^\circ\right]=0,\forall a,b\in\mathcal{A}$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E00072 (24)	44	■ 1.000	$d\left(\phi_1,\phi_2\right)=\sup \left\{\left \phi_1(a)-\phi_2(a)\right \mid \left[\mathcal{D},\pi(a)\right]\leq 1,a\in \mathcal{A}\right\}$	$d(\phi_1,\phi_2)=\sup \{ \phi_1(a)-\phi_2(a) \mid \ [\mathcal{D},\pi(a)]\ \leq 1,a\in \mathcal{A}\}$	$d(\phi_1,\phi_2)=\sup \{ \phi_1(a)-\phi_2(a) \mid \ [\mathcal{D},\pi(a)]\ \leq 1,a\in \mathcal{A}\}$
cardona-qft-methods_ E00073	45	■ 1.000	$0\leq \alpha\leq \beta\implies 0\leq \alpha+\beta\leq \beta,\quad 0\leq \alpha\leq \beta\implies 0\leq \alpha+\beta\leq \beta\text{ if }\alpha\leq \beta,\quad \delta(0\leq \alpha\leq \beta)\subset 0\leq \alpha\leq \beta$	$OP^\alpha OP^\beta\subset OP^{\alpha+\beta},\quad OP^\alpha\subset OP^\beta\text{ if }\alpha\leq \beta,\quad \delta(OP^\alpha)\subset OP^\alpha$	$OP^\alpha OP^\beta\subset OP^{\alpha+\beta},\quad OP^\alpha\subset OP^\beta\text{ if }\alpha\leq \beta,\quad \delta(OP^\alpha)\subset OP^\alpha$
cardona-qft-methods_ E00076 (25)	45	■ 1.000	$\zeta_{\mathcal{D}}^P(s):=\mathrm{Tr}\left(P \mathcal{D} ^{-s}\right)$	$\zeta_{\mathcal{D}}^P:s\in\mathbb{C}\rightarrow\mathrm{Tr}\left(P \mathcal{D} ^{-s}\right)$	$\zeta_{\mathcal{D}}^P:s\in\mathbb{C}\rightarrow\mathrm{Tr}\left(P \mathcal{D} ^{-s}\right)$
cardona-qft-methods_ E00079	47	■ 1.000	$\sigma_z(T)= D ^zT D ^{-z},\quad z\in\mathbb{C},\quad \epsilon(T)=\nabla(T)D^{-2}$	$\sigma_z(T)= D ^zT D ^{-z},\quad z\in\mathbb{C},\quad \epsilon(T)=\nabla(T)D^{-2}$	$\sigma_z(T)= D ^zT D ^{-z},\quad z\in\mathbb{C},\quad \epsilon(T)=\nabla(T)D^{-2},$
cardona-qft-methods_ E00080 (27)	47	■ 1.000	$\sigma_z(T)\sim\sum_{r=0}^Ng(z,r)\varepsilon^r(T)\pmod{OP^{-N-1+q}}$	$\sigma_z(T)\sim\sum_{r=0}^Ng(z,r)\varepsilon^r(T)\pmod{OP^{-N-1+q}}$	$\sigma_z(T)\sim\sum_{r=0}^Ng(z,r)\varepsilon^r(T)\pmod{OP^{-N-1+q}}$
cardona-qft-methods_ E00082	48	■ 1.000	$Y\sim\sum_{p=1}^N\sum_{k_1,\dots,k_p=0}^{N-p}\frac{(-1)^{ k _1+p+1}}{ k _1+p}\nabla^{k_p}(X\nabla^{k_{p-1}}(\dots X\nabla^{k_1}(X)\dots))D^{-2(k _1+p)},$	$Y\sim\sum_{p=1}^N\sum_{k_1,\dots,k_p=0}^{N-p}\frac{(-1)^{ k _1+p+1}}{ k _1+p}\nabla^{k_p}(X\nabla^{k_{p-1}}(\dots X\nabla^{k_1}(X)\dots))D^{-2(k _1+p)},$	$Y\sim\sum_{p=1}^N\sum_{k_1,\dots,k_p=0}^{N-p}\frac{(-1)^{ k _1+p+1}}{ k _1+p}\nabla^{k_p}(X\nabla^{k_{p-1}}(\dots X\nabla^{k_1}(X)\dots))D^{-2(k _1+p)},$
cardona-qft-methods_ E00084	49	■ 1.000	$\sum_{p=1}^k\sum_{r_1,\dots,r_p=0}^{k-p}\mathrm{Res}_{s=d-k}h(s,r,p)\mathrm{Tr}\left(\varepsilon^{r_1}(Y)\dots\varepsilon^{r_p}(Y) D ^{-s}\right),$	$\sum_{p=1}^k\sum_{r_1,\dots,r_p=0}^{k-p}\mathrm{Res}_{s=d-k}h(s,r,p)\mathrm{Tr}\left(\varepsilon^{r_1}(Y)\dots\varepsilon^{r_p}(Y) D ^{-s}\right),$	$\sum_{p=1}^k\sum_{r_1,\dots,r_p=0}^{k-p}\mathrm{Res}_{s=d-k}h(s,r,p)\mathrm{Tr}\left(\varepsilon^{r_1}(Y)\dots\varepsilon^{r_p}(Y) D ^{-s}\right),$
cardona-qft-methods_ E00089	51	■ 1.000	$\zeta_{\mathcal{D}_A}(0)=c_1\int_M\left(5R^2-8R_{\mu\nu}r^{\mu\nu}-7R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}dvol+c_2\int_M\mathrm{tr}(F_{\mu\nu}F^{\mu\nu})dvol\right)$	$\zeta_{\mathcal{D}_A}(0)=c_1\int_M\left(5R^2-8R_{\mu\nu}r^{\mu\nu}-7R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}dvol+c_2\int_M\mathrm{tr}(F_{\mu\nu}F^{\mu\nu})dvol\right)$	$\zeta_{\mathcal{D}_A}(0)=c_1\int_M(5R^2-8R_{\mu\nu}r^{\mu\nu}-7R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}dvol+c_2\int_M\mathrm{tr}(F_{\mu\nu}F^{\mu\nu})dvol,$
cardona-qft-methods_ E00090	52	■ 1.000	$\mathcal{A}:=\mathcal{A}_1\otimes\mathcal{A}_2,\quad \mathcal{D}:=\mathcal{D}_1\otimes 1+\chi_1\otimes\mathcal{D}_2,\quad \mathcal{H}:=\mathcal{H}_1\otimes\mathcal{H}_2.$	$\mathcal{A}:=\mathcal{A}_1\otimes\mathcal{A}_2,\quad \mathcal{D}:=\mathcal{D}_1\otimes 1+\chi_1\otimes\mathcal{D}_2,\quad \mathcal{H}:=\mathcal{H}_1\otimes\mathcal{H}_2.$	$\mathcal{A}:=\mathcal{A}_1\otimes\mathcal{A}_2,\quad \mathcal{D}:=\mathcal{D}_1\otimes 1+\chi_1\otimes\mathcal{D}_2,\quad \mathcal{H}:=\mathcal{H}_1\otimes\mathcal{H}_2.$
cardona-qft-methods_ E00091	52	■ 1.000	$\mathrm{Res}_{s=d_1+d_2}(\zeta_{\mathcal{D}}(s))=\frac{1}{2}\frac{\Gamma(d_1/2)\Gamma(d_2/2)}{\Gamma(d/2)}\mathrm{Res}_{s=d_1}(\zeta_{\mathcal{D}_1}(s))\mathrm{Res}_{s=d_2}(\zeta_{\mathcal{D}_2}(s)).$	$\mathrm{Res}_{s=d_1+d_2}(\zeta_{\mathcal{D}}(s))=\frac{1}{2}\frac{\Gamma(d_1/2)\Gamma(d_2/2)}{\Gamma(d/2)}\mathrm{Res}_{s=d_1}(\zeta_{\mathcal{D}_1}(s))\mathrm{Res}_{s=d_2}(\zeta_{\mathcal{D}_2}(s)).$	$\mathrm{Res}_{s=d_1+d_2}(\zeta_{\mathcal{D}}(s))=\frac{1}{2}\frac{\Gamma(d_1/2)\Gamma(d_2/2)}{\Gamma(d/2)}\mathrm{Res}_{s=d_1}(\zeta_{\mathcal{D}_1}(s))\mathrm{Res}_{s=d_2}(\zeta_{\mathcal{D}_2}(s)).$
cardona-qft-methods_ E00092 (31)	55	■ 1.000	$\mathrm{Tr}\left(e^{-t\mathcal{D}^2}\right)\underset{t\downarrow 0}{\sim}\sum_{\alpha\in Sd}a_\alpha t^\alpha\quad\text{with }a_\alpha\neq 0.$	$\mathrm{Tr}\left(e^{-t\mathcal{D}^2}\right)\underset{t\downarrow 0}{\sim}\sum_{\alpha\in Sd}a_\alpha t^\alpha\quad\text{with }a_\alpha\neq 0.$	$\mathrm{Tr}\left(e^{-t\mathcal{D}^2}\right)\underset{t\downarrow 0}{\sim}\sum_{\alpha\in Sd}a_\alpha t^\alpha\quad\text{with }a_\alpha\neq 0.$
cardona-qft-methods_ E00093 (32)	55	■ 1.000	$a_\alpha=\frac{1}{2}\mathrm{Res}_{s=-2\alpha}(\Gamma(s/2)\zeta_{\mathcal{D}}(s)).$	$a_\alpha=\frac{1}{2}\mathrm{Res}_{s=-2\alpha}(\Gamma(s/2)\zeta_{\mathcal{D}}(s)).$	$a_\alpha=\frac{1}{2}\mathrm{Res}_{s=-2\alpha}(\Gamma(s/2)\zeta_{\mathcal{D}}(s)).$
cardona-qft-methods_ E00094 (33)	55	■ 1.000	$\mathrm{Tr}(f(\mathcal{D}/\Lambda))\underset{\Lambda\rightarrow+\infty}{\sim}\sum_{\beta\in Sd^+}f_\beta\Lambda^\beta\int \mathcal{D} ^\beta+f(0)\zeta_{\mathcal{D}}(0)+\dots$	$\mathrm{Tr}(f(\mathcal{D}/\Lambda))\underset{\Lambda\rightarrow+\infty}{\sim}\sum_{\beta\in Sd^+}f_\beta\Lambda^\beta\int \mathcal{D} ^\beta+f(0)\zeta_{\mathcal{D}}(0)+\dots$	$\mathrm{Tr}(f(\mathcal{D}/\Lambda))\underset{\Lambda\rightarrow+\infty}{\sim}\sum_{\beta\in Sd^+}f_\beta\Lambda^\beta\int \mathcal{D} ^\beta+f(0)\zeta_{\mathcal{D}}(0)+\dots$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0097 (34)	56	■ 1.000	$\operatorname{Tr}\left(e^{-t\,\mathrm{D}^2}\right)$ $\underset{t\downarrow 0}{\sim}\sum_{k\in\{0,\cdots,d\}}t^{(k-d)/2}a_k(\mathcal{D}^2)+\cdots,$	$\operatorname{Tr}\left(e^{-t\mathcal{D}^2}\right)\underset{t\downarrow 0}{\sim}\sum_{k\in\{0,\cdots,d\}}t^{(k-d)/2}a_k(\mathcal{D}^2)+\cdots,$	$\operatorname{Tr}\left(e^{-t\mathcal{D}^2}\right)\underset{t\downarrow 0}{\sim}\sum_{k\in\{0,\cdots,d\}}t^{(k-d)/2}a_k(\mathcal{D}^2)+\cdots,$
cardona-qft-methods_ EQ0101	58	■ 1.000	$a_{\{k\}}:=\frac{(-1)^k4}{3\,k!}\left(\frac{B_{2\,k+2}}{2\,k+2}-\frac{B_{2k+4}}{2\,k+4}\right)$	$a_k:=\frac{(-1)^k4}{3k!}\left(\frac{B_{2k+2}}{2k+2}-\frac{B_{2k+4}}{2k+4}\right)$	$a_k:=\frac{(-1)^k4}{3\,k!}\left(\frac{B_{2k+2}}{2k+2}-\frac{B_{2k+4}}{2k+4}\right)$
cardona-qft-methods_ EQ0103	61	■ 1.000	$\mathrm{D}=-i\,\delta_\mu\,\otimes\,\gamma^\mu,$	$\mathcal{D}=-i\delta_\mu\otimes\gamma^\mu,$	$\mathcal{D}=-i\delta_\mu\otimes\gamma^\mu,$
cardona-qft-methods_ EQ0107 (2)	72	■ 1.000	$\operatorname{Ind}(A)=\operatorname{dim}\,\mathrm{H}_1-$ $\operatorname{dim}\,\mathrm{H}_2$	$\operatorname{Ind}(A)=\dim\mathcal{H}_1-\dim\mathcal{H}_2$	$\operatorname{Ind}(A)=\dim\mathcal{H}_1-\dim\mathcal{H}_2$
cardona-qft-methods_ EQ0108 (3)	73	■ 1.000	$\mathrm{H}=\mathcal{L}_2:=\left\{\left(x_1,x_2,\ldots\right)\in\mathcal{C},\sum_{j=1}^{\infty}\left x_j\right ^2<\infty\right\},$	$\mathcal{H}=l_2:=\left\{\left(x_1,x_2,\ldots\left x_j\in\mathcal{C},\sum_{j=1}^{\infty}\left x_j\right ^2<\infty\right\},\right.$	$\mathcal{H}=l_2:=\left\{\left(x_1,x_2,\ldots\left x_j\in\mathcal{C},\sum_{j=1}^{\infty}\left x_j\right ^2<\infty\right\},\right.$
cardona-qft-methods_ EQ0109 (4)	73	■ 1.000	$T\left(x_1,x_2,x_3,\ldots\right):=\left(x_3,x_4,\ldots\right).$	$T\left(x_1,x_2,x_3,\ldots\right):=\left(x_3,x_4,\ldots\right).$	$T\left(x_1,x_2,x_3,\ldots\right):=\left(x_3,x_4,\ldots\right).$
cardona-qft-methods_ EQ0110	73	■ 1.000	$\operatorname{Ker}(T)=\left\{\left(x_1,x_2,0,0,\ldots\right)\in\mathcal{H}\mid x_1,x_2\in\mathbb{C}\right\}.$	$\operatorname{Ker}(T)=\{(x_1,x_2,0,0,\ldots)\in\mathcal{H}\mid x_1,x_2\in\mathbb{C}\}.$	$\operatorname{Ker}(T)=\left\{\left(x_1,x_2,0,0,\ldots\right)\in\mathcal{H}\mid x_1,x_2\in\mathbb{C}\right\}.$
cardona-qft-methods_ EQ0113 (5)	73	■ 1.000	$(A+K)\,x=0$	$(A+K)x=0$	$(A+K)\,x=0$
cardona-qft-methods_ EQ0114	74	■ 1.000	$U=\bigoplus_{V\in\operatorname{Irr}(G)}m_VV.$	$U=\bigoplus_{V\in\operatorname{Irr}(G)}m_VV.$	$U=\bigoplus_{V\in\operatorname{Irr}(G)}m_VV.$
cardona-qft-methods_ EQ0115	74	■ 1.000	$g\,A\,x=A\,g\,x,\quad\text{for all}\quad x\in\mathcal{H}_1.$	$gAx=Agx,\quad\text{for all}\quad x\in\mathcal{H}_1.$	$gAx=Agx,\quad\text{for all}\quad x\in\mathcal{H}_1.$
cardona-qft-methods_ EQ0117	75	■ 1.000	$V-U\sim A-B\,\Longlefttrightarrow\,V\oplus B\simeq A\oplus U,$	$V-U\sim A-B\iff V\oplus B\simeq A\oplus U,$	$V-U\sim A-B\iff V\oplus B\simeq A\oplus U,$
cardona-qft-methods_ EQ0119 (7)	76	■ 1.000	$\widehat{R}(G)=\left\{\bigoplus_{V\in\operatorname{Irr}(G)}m_VV\mid m_V\in\mathbb{Z}\right\}.$	$\widehat{R}(G)=\left\{\bigoplus_{V\in\operatorname{Irr}(G)}m_VV\mid m_V\in\mathbb{Z}\right\}.$	$\widehat{R}(G)=\left\{\bigoplus_{V\in\operatorname{Irr}(G)}m_VV\mid m_V\in\mathbb{Z}\right\}.$
cardona-qft-methods_ EQ0120 (8)	76	■ 1.000	$\operatorname{Ind}_G(A):=\operatorname{Ker}(A)-\operatorname{Coker}(A)=\bigoplus_{V\in\operatorname{Irr}(G)}\left(m_V^+-m_V^-\right)V\in R(G).$	$\operatorname{Ind}_G(A):=\operatorname{Ker}(A)-\operatorname{Coker}(A)=\bigoplus_{V\in\operatorname{Irr}(G)}\left(m_V^+-m_V^-\right)V\in R(G).$	$\operatorname{Ind}_G(A):=\operatorname{Ker}(A)-\operatorname{Coker}(A)=\bigoplus_{V\in\operatorname{Irr}(G)}\left(m_V^+-m_V^-\right)V\in R(G).$
cardona-qft-methods_ EQ0121	76	■ 1.000	$G:=\mathbb{Z}_2=\{1,-1\}.$	$G:=\mathbb{Z}_2=\{1,-1\}.$	$G:=\mathbb{Z}_2=\{1,-1\}.$
cardona-qft-methods_ EQ0122	76	■ 1.000	$q\left(x_1,x_2,\ldots\right)=\left(x_2,x_1,x_4,x_3,\ldots\right).$	$q\left(x_1,x_2,\ldots\right)=\left(x_2,x_1,x_4,x_3,\ldots\right).$	$q\left(x_1,x_2,\ldots\right)=\left(x_2,x_1,x_4,x_3,\ldots\right).$
cardona-qft-methods_ EQ0123	77	■ 1.000	$D_{\{j\}}f=\frac{1}{i}\frac{\partial f}{\partial x_j},\quad j=1,\ldots,n.$	$D_jf=\frac{1}{i}\frac{\partial f}{\partial x_j},\quad j=1,\ldots,n.$	$D_jf=\frac{1}{i}\frac{\partial f}{\partial x_j},\quad j=1,\ldots,n.$
cardona-qft-methods_ EQ0124	77	■ 1.000	$ \alpha :=\alpha_1+\cdots+\alpha_n$	$ \alpha :=\alpha_1+\cdots+\alpha_n$	$ \alpha :=\alpha_1+\cdots+\alpha_n$
cardona-qft-methods_ EQ0126 (1)	77	■ 1.000	$\mathrm{D}=\sum_{ \alpha \leq k}a_{\alpha}(x)D^{\alpha}:\mathcal{C}^{\infty}(\mathbb{R}^n)\longrightarrow\mathcal{C}^{\infty}(\mathbb{R}^n),$	$\mathcal{D}=\sum_{ \alpha \leq k}a_{\alpha}(x)D^{\alpha}:\mathcal{C}^{\infty}(\mathbb{R}^n)\longrightarrow\mathcal{C}^{\infty}(\mathbb{R}^n),$	$\mathcal{D}=\sum_{ \alpha \leq k}a_{\alpha}(x)D^{\alpha}:\mathcal{C}^{\infty}(\mathbb{R}^n)\longrightarrow\mathcal{C}^{\infty}(\mathbb{R}^n),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0127	77	■ 1.000	$\langle f, g \rangle := \int_{\mathbb{R}^n} f(x) \overline{g(x)} dx$	$\langle f, g \rangle := \int_{\mathbb{R}^n} f(x) \overline{g(x)} dx$	$\langle f, g \rangle := \int_{\mathbb{R}^n} f(x) \overline{g(x)} dx.$
cardona-qft-methods_ EQ0128 (2)	77	■ 1.000	$\langle f, g \rangle_k := \sum_{ \alpha =k} \langle D^\alpha f, D^\alpha g \rangle.$	$\langle f, g \rangle_k := \sum_{ \alpha =k} \langle D^\alpha f, D^\alpha g \rangle.$	$\langle f, g \rangle_k := \sum_{ \alpha =k} \langle D^\alpha f, D^\alpha g \rangle.$
cardona-qft-methods_ EQ0129 (3)	77	■ 1.000	$\ f\ _k := \sqrt{\langle f, f \rangle_k}$	$\ f\ _k := \sqrt{\langle f, f \rangle_k}$	$\ f\ _k := \sqrt{\langle f, f \rangle_k}$
cardona-qft-methods_ EQ0130	78	■ 1.000	$H^k(\mathbb{R}^n) = \{f : \mathbb{R}^n \rightarrow \mathcal{C} \mid D^\alpha f \in L^2(\mathbb{R}^n) \text{ for all } \alpha \in \mathbb{Z}_+^n \text{ with } \alpha \leq k\}.$	$H^k(\mathbb{R}^n) = \{f : \mathbb{R}^n \rightarrow \mathcal{C} \mid D^\alpha f \in L^2(\mathbb{R}^n) \text{ for all } \alpha \in \mathbb{Z}_+^n \text{ with } \alpha \leq k\}.$	$H^k(\mathbb{R}^n) = \{f : \mathbb{R}^n \rightarrow \mathcal{C} \mid D^\alpha f \in L^2(\mathbb{R}^n) \text{ for all } \alpha \in \mathbb{Z}_+^n \text{ with } \alpha \leq k\}.$
cardona-qft-methods_ EQ0131	78	■ 1.000	$H_K^k(\mathbb{R}^n) = \{f \in H^k(\mathbb{R}^n) \mid \text{supp}(f) \subset K\}.$	$H_K^k(\mathbb{R}^n) = \{f \in H^k(\mathbb{R}^n) \mid \text{supp}(f) \subset K\}.$	$H_K^k(\mathbb{R}^n) = \{f \in H^k(\mathbb{R}^n) \mid \text{supp}(f) \subset K\}.$
cardona-qft-methods_ EQ0133	79	■ 1.000	$a_\alpha(x) = \{a_{\alpha,ij}(x)\}_{\substack{1 \leq i \leq N_2 \\ 1 \leq j \leq N_1}} \in \text{Mat}_{N_1 \times N_2}, \quad a_{\alpha,ij}(x) \in C^\infty(\mathbb{R}^n).$	$a_\alpha(x) = \{a_{\alpha,ij}(x)\}_{\substack{1 \leq i \leq N_2 \\ 1 \leq j \leq N_1}} \in \text{Mat}_{N_1 \times N_2}, \quad a_{\alpha,ij}(x) \in C^\infty(\mathbb{R}^n).$	$a_\alpha(x) = \{a_{\alpha,ij}(x)\}_{\substack{1 \leq i \leq N_2 \\ 1 \leq j \leq N_1}} \in \text{Mat}_{N_1 \times N_2}, \quad a_{\alpha,ij}(x) \in C^\infty(\mathbb{R}^n).$
cardona-qft-methods_ EQ0134	79	■ 1.000	$\mathcal{D} : H^k(\mathbb{R}^n, \mathbb{R}^N) \longrightarrow L^2(\mathbb{R}^n, \mathbb{R}^N)$	$\mathcal{D} : H^k(\mathbb{R}^n, \mathbb{R}^N) \longrightarrow L^2(\mathbb{R}^n, \mathbb{R}^N)$	$\mathcal{D} : H^k(\mathbb{R}^n, \mathbb{R}^N) \longrightarrow L^2(\mathbb{R}^n, \mathbb{R}^N)$
cardona-qft-methods_ EQ0136	79	■ 1.000	$\pi(s(x)) = x, \quad \text{for all } x \in M.$	$\pi(s(x)) = x, \quad \text{for all } x \in M.$	$\pi(s(x)) = x, \quad \text{for all } x \in M.$
cardona-qft-methods_ EQ0137 (4)	80	■ 1.000	$\mathcal{D} : C^\infty(M, E) \longrightarrow C^\infty(M, F)$	$\mathcal{D} : C^\infty(M, E) \longrightarrow C^\infty(M, F)$	$\mathcal{D} : C^\infty(M, E) \longrightarrow C^\infty(M, F)$
cardona-qft-methods_ EQ0139	80	■ 1.000	$\mathcal{D}_i : C^\infty(V_i, \mathbb{R}^{N_1}) \longrightarrow C^\infty(V_i, \mathbb{R}^{N_2}).$	$\mathcal{D}_i : C^\infty(V_i, \mathbb{R}^{N_1}) \longrightarrow C^\infty(V_i, \mathbb{R}^{N_2}).$	$\mathcal{D}_i : C^\infty(V_i, \mathbb{R}^{N_1}) \longrightarrow C^\infty(V_i, \mathbb{R}^{N_2}).$
cardona-qft-methods_ EQ0140	81	■ 1.000	$\mathcal{D} : H_K^m(M, E) \longrightarrow L^2(M, F).$	$\mathcal{D} : H_K^m(M, E) \longrightarrow L^2(M, F).$	$\mathcal{D} : H_K^m(M, E) \longrightarrow L^2(M, F).$
cardona-qft-methods_ EQ0143	81	■ 1.000	$\sigma(\mathcal{D})(x_0, \xi) = \sum_{ \alpha \leq k} a_\alpha(x_0) \xi_1^{\alpha_1} \cdots \xi_n^{\alpha_n}.$	$\sigma(\mathcal{D})(x_0, \xi) = \sum_{ \alpha \leq k} a_\alpha(x_0) \xi_1^{\alpha_1} \cdots \xi_n^{\alpha_n}.$	$\sigma(\mathcal{D})(x_0, \xi) = \sum_{ \alpha \leq k} a_\alpha(x_0) \xi_1^{\alpha_1} \cdots \xi_n^{\alpha_n}.$
cardona-qft-methods_ EQ0144	82	■ 1.000	$\sigma_L(\mathcal{D})(x, \xi) : \pi^* E_{(x, \xi)} \rightarrow \pi^* F_{(x, \xi)}.$	$\sigma_L(\mathcal{D})(x, \xi) : \pi^* E_{(x, \xi)} \rightarrow \pi^* F_{(x, \xi)}.$	$\sigma_L(\mathcal{D})(x, \xi) : \pi^* E_{(x, \xi)} \rightarrow \pi^* F_{(x, \xi)}.$
cardona-qft-methods_ EQ0145	82	■ 1.000	$\Delta = -\frac{\partial^2}{\partial x_1^2} - \frac{\partial^2}{\partial x_2^2} - \cdots - \frac{\partial^2}{\partial x_n^2}$	$\Delta = -\frac{\partial^2}{\partial x_1^2} - \frac{\partial^2}{\partial x_2^2} - \cdots - \frac{\partial^2}{\partial x_n^2}$	$\Delta = -\frac{\partial^2}{\partial x_1^2} - \frac{\partial^2}{\partial x_2^2} - \cdots - \frac{\partial^2}{\partial x_n^2}$
cardona-qft-methods_ EQ0150 (2)	85	■ 1.000	$\sigma_1 \oplus \text{Id} : E_1 \oplus \mathcal{C}^{k_1} \rightarrow F_1 \oplus \mathcal{C}^{k_1}, \quad \text{and} \quad \sigma_2 \oplus \text{Id} : E_2 \oplus \mathcal{C}^{k_2} \rightarrow F_2 \oplus \mathcal{C}^{k_2},$	$\sigma_1 \oplus \text{Id} : E_1 \oplus \mathcal{C}^{k_1} \rightarrow F_1 \oplus \mathcal{C}^{k_1}, \quad \text{and} \quad \sigma_2 \oplus \text{Id} : E_2 \oplus \mathcal{C}^{k_2} \rightarrow F_2 \oplus \mathcal{C}^{k_2},$	$\sigma_1 \oplus \text{Id} : E_1 \oplus \mathcal{C}^{k_1} \rightarrow F_1 \oplus \mathcal{C}^{k_1}, \quad \text{and} \quad \sigma_2 \oplus \text{Id} : E_2 \oplus \mathcal{C}^{k_2} \rightarrow F_2 \oplus \mathcal{C}^{k_2},$
cardona-qft-methods_ EQ0151	85	■ 1.000	$K(X) = \{\sigma : E \rightarrow F \mid \sigma(x) \text{ is invertible outside of a compact set}\} / \sim$	$K(X) = \{\sigma : E \rightarrow F \mid \sigma(x) \text{ is invertible outside of a compact set}\} / \sim$	$K(X) = \{\sigma : E \rightarrow F \mid \sigma(x) \text{ is invertible outside of a compact set}\} / \sim$
cardona-qft-methods_ EQ0152	85	■ 1.000	$\dim E - \dim F \in \mathbb{Z}.$	$\dim E - \dim F \in \mathbb{Z}.$	$\dim E - \dim F \in \mathbb{Z}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0153	85	■ 1.000	$K(\backslash\{p\ t\})\ \backslash\mathrm{cong}\ \backslash\mathrm{mathbb{Z}}$	$K(\{pt\}) \cong \mathbb{Z}$	$K(\{pt\}) \cong \mathbb{Z}.$
cardona-qft-methods_ EQ0155	86	■ 1.000	$j_{!}!: K(Y)\ \backslash\mathrm{rightrightarrow}\ K(X),$	$j_{!}!: K(Y) \rightarrow K(X),$	$j_{!}: K(Y) \rightarrow K(X),$
cardona-qft-methods_ EQ0156	86	■ 1.000	$i_{!}!: K(p\ t)\ \backslash\mathrm{longrightrightarrow}\ K\backslash\mathrm{left}(\backslash\mathrm{mathcal{C}}^{\{N\}}\ \backslash\mathrm{right})$	$i_{!}: K(pt) \longrightarrow K\left(\mathcal{C}^N\right)$	$i_{!}: K(pt) \longrightarrow K(\mathcal{C}^N)$
cardona-qft-methods_ EQ0159	87	■ 1.000	$g\ \backslash\mathrm{mapsto}\ \backslash\mathrm{psi}(g): E\ \backslash\mathrm{rightrightarrow}\ E,\ \backslash\mathrm{quad}\ g\ \backslash\mathrm{in}\ G,$	$g \mapsto \psi(g): E \rightarrow E,\quad g \in G,$	$g \mapsto \psi(g): E \rightarrow E,\quad g \in G,$
cardona-qft-methods_ EQ0160	87	■ 1.000	$\backslash\mathrm{pi}\ \backslash\mathrm{circ}\ \backslash\mathrm{psi}(g)=\backslash\mathrm{phi}\ \backslash\mathrm{circ}\ \backslash\mathrm{pi}(g)\ .$	$\pi \circ \psi(g) = \phi \circ \pi(g).$	$\pi \circ \psi(g) = \phi \circ \pi(g).$
cardona-qft-methods_ EQ0161 (5)	87	■ 1.000	$g\ \backslash\mathrm{mapsto}\ l_{\{g\}}: C^{\backslash\mathrm{infty}}(M,\ E)\ \backslash\mathrm{longrightrightarrow}\ C^{\backslash\mathrm{infty}}(M,\ E),\ \backslash\mathrm{quad}\ l_{\{g\}}(f)(x):=g\ \backslash\mathrm{cdot}\ f\backslash\mathrm{left}(g^{\{-1\}}\ \backslash\mathrm{cdot}\ x\backslash\mathrm{right})\ .$	$g \mapsto l_g: C^\infty(M, E) \longrightarrow C^\infty(M, E),\quad l_g(f)(x) := g \cdot f\left(g^{-1} \cdot x\right).$	$g \mapsto l_g: C^\infty(M, E) \longrightarrow C^\infty(M, E),\quad l_g(f)(x) := g \cdot f(g^{-1} \cdot x).$
cardona-qft-methods_ EQ0163 (6)	88	■ 1.000	$K_{\{G\}}(\backslash\{p\ t\})=R(G),$	$K_G(\{pt\}) = R(G),$	$K_G(\{pt\}) = R(G),$
cardona-qft-methods_ EQ0165	88	■ 1.000	$\backslash\mathrm{mathcal{D}}: C^{\backslash\mathrm{infty}}(M,\ E)\ \backslash\mathrm{rightrightarrow}\ C^{\backslash\mathrm{infty}}(M,\ F)$	$\mathcal{D}: C^\infty(M, E) \rightarrow C^\infty(M, F)$	$\mathcal{D}: C^\infty(M, E) \rightarrow C^\infty(M, F)$
cardona-qft-methods_ EQ0166	88	■ 1.000	$\backslash\mathrm{sigma}_{\{L\}}(\backslash\mathrm{mathcal{D}}): \backslash\mathrm{pi}^{\{*\}}\ E\ \backslash\mathrm{longrightrightarrow}\ \backslash\mathrm{pi}^{\{*\}}\ F,$	$\sigma_L(\mathcal{D}): \pi^*E \longrightarrow \pi^*F,$	$\sigma_L(\mathcal{D}): \pi^*E \longrightarrow \pi^*F,$
cardona-qft-methods_ EQ0167	89	■ 1.000	$\backslash\mathrm{operatorname{Ind}}_{\{G\}}(\backslash\mathrm{mathcal{D}})=\backslash\mathrm{mathrm{t}}\text{-}\backslash\mathrm{operatorname{Ind}}_{\{G\}}(\backslash\mathrm{sigma})\ .$	$\mathrm{Ind}_G(\mathcal{D}) = \mathrm{t} - \mathrm{Ind}_G(\sigma).$	$\mathrm{Ind}_G(\mathcal{D}) = \mathrm{t}\text{-}\mathrm{Ind}_G(\sigma).$
cardona-qft-methods_ EQ0168	89	■ 1.000	$M:=N\ \backslash\mathrm{times}\ G$	$M := N \times G$	$M := N \times G$
cardona-qft-methods_ EQ0170	89	■ 1.000	$\backslash\mathrm{tilde}(\backslash\mathrm{mathcal{D}}): C^{\backslash\mathrm{infty}}(M)\ \backslash\mathrm{longrightrightarrow}\ C^{\backslash\mathrm{infty}}(M)$	$\tilde{\mathcal{D}}: C^\infty(M) \longrightarrow C^\infty(M)$	$\tilde{\mathcal{D}}: C^\infty(M) \longrightarrow C^\infty(M)$
cardona-qft-methods_ EQ0171	89	■ 1.000	$\backslash\mathrm{mathcal{D}}=\backslash\mathrm{sum}_{\{ \backslash\mathrm{alpha} \ \backslash\mathrm{leq}\ k\}}\ a_{\{\backslash\mathrm{alpha}\}}(x)\ \backslash\mathrm{frac}\{\backslash\mathrm{partial}^{\{\backslash\mathrm{alpha}_{\{1\}}\}}\{\backslash\mathrm{partial}\ x_{\{1\}}^{\{\backslash\mathrm{alpha}_{\{1\}}\}}\}\ \backslash\mathrm{cdot}\ \backslash\mathrm{frac}\{\backslash\mathrm{partial}^{\{\backslash\mathrm{alpha}_{\{n\}}\}}\{\backslash\mathrm{partial}\ x_{\{n\}}^{\{\backslash\mathrm{alpha}_{\{n\}}\}}\}}\ .$	$\mathcal{D} = \sum_{ \alpha \leq k} a_\alpha(x) \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \cdots \frac{\partial^{\alpha_n}}{\partial x_n^{\alpha_n}}.$	$\mathcal{D} = \sum_{ \alpha \leq k} a_\alpha(x) \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \cdots \frac{\partial^{\alpha_n}}{\partial x_n^{\alpha_n}}.$
cardona-qft-methods_ EQ0172	89	■ 1.000	$\backslash\mathrm{tilde}(\backslash\mathrm{mathcal{D}}):=\backslash\mathrm{sum}_{\{ \backslash\mathrm{alpha} \ \backslash\mathrm{leq}\ k\}}\ a_{\{\backslash\mathrm{alpha}\}}(x)\ \backslash\mathrm{frac}\{\backslash\mathrm{partial}^{\{\backslash\mathrm{alpha}_{\{1\}}\}}\{\backslash\mathrm{partial}\ x_{\{1\}}^{\{\backslash\mathrm{alpha}_{\{1\}}\}}\}\ \backslash\mathrm{cdot}\ \backslash\mathrm{frac}\{\backslash\mathrm{partial}^{\{\backslash\mathrm{alpha}_{\{n\}}\}}\{\backslash\mathrm{partial}\ x_{\{n\}}^{\{\backslash\mathrm{alpha}_{\{n\}}\}}\}}\ .$	$\tilde{\mathcal{D}} := \sum_{ \alpha \leq k} a_\alpha(x) \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \cdots \frac{\partial^{\alpha_n}}{\partial x_n^{\alpha_n}}.$	$\tilde{\mathcal{D}} := \sum_{ \alpha \leq k} a_\alpha(x) \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \cdots \frac{\partial^{\alpha_n}}{\partial x_n^{\alpha_n}}.$
cardona-qft-methods_ EQ0173 (1)	90	■ 1.000	$\backslash\mathrm{operatorname{Ker}}(\backslash\mathrm{tilde}(\backslash\mathrm{mathcal{D}}))=\backslash\mathrm{operatorname{Ker}}(\backslash\mathrm{mathcal{D}})\ \backslash\mathrm{otimes}\ L^{\{2\}}(G),\ \backslash\mathrm{quad}\ \backslash\mathrm{operatorname{Coker}}(\backslash\mathrm{tilde}(\backslash\mathrm{mathcal{D}}))=\backslash\mathrm{operatorname{Coker}}(\backslash\mathrm{mathcal{D}})\ \backslash\mathrm{otimes}\ L^{\{2\}}(G)\ .$	$\mathrm{Ker}(\tilde{\mathcal{D}}) = \mathrm{Ker}(\mathcal{D}) \otimes L^2(G),\quad \mathrm{Coker}(\tilde{\mathcal{D}}) = \mathrm{Coker}(\mathcal{D}) \otimes L^2(G).$	$\mathrm{Ker}(\tilde{\mathcal{D}}) = \mathrm{Ker}(\mathcal{D}) \otimes L^2(G),\quad \mathrm{Coker}(\tilde{\mathcal{D}}) = \mathrm{Coker}(\mathcal{D}) \otimes L^2(G).$
cardona-qft-methods_ EQ0174 (2)	90	■ 1.000	$L^{\{2\}}(G)=\backslash\mathrm{bigoplus}_{\{V\ \backslash\mathrm{in}\ \backslash\mathrm{operatorname{Irr}}(G)\}}(\backslash\mathrm{operatorname{dim}}\ V)\ V\ .$	$L^2(G) = \bigoplus_{V \in \mathrm{Irr}(G)} (\dim V) V.$	$L^2(G) = \bigoplus_{V \in \mathrm{Irr}(G)} (\dim V) V.$
cardona-qft-methods_ EQ0176	90	■ 1.000	$\backslash\mathrm{mathcal{O}}(x):=\backslash\{g\ \backslash\mathrm{cdot}\ x\ \backslash\mathrm{mid}\ g\ \backslash\mathrm{in}\ G\}\ .$	$\mathcal{O}(x) := \{g \cdot x \mid g \in G\}.$	$\mathcal{O}(x) := \left\{g \cdot x \middle g \in G\right\}.$
cardona-qft-methods_ EQ0178	91	■ 1.000	$\backslash\mathrm{mathcal{D}}: C^{\backslash\mathrm{infty}}(M,\ E)\ \backslash\mathrm{longrightrightarrow}\ C^{\backslash\mathrm{infty}}(M,\ F)$	$\mathcal{D}: C^\infty(M, E) \longrightarrow C^\infty(M, F)$	$\mathcal{D}: C^\infty(M, E) \longrightarrow C^\infty(M, F)$
cardona-qft-methods_ EQ0179 (5)	91	■ 1.000	$\backslash\mathrm{operatorname{Ker}}(\backslash\mathrm{mathcal{D}})=\backslash\mathrm{bigoplus}_{\{V\ \backslash\mathrm{in}\ \backslash\mathrm{operatorname{Irr}}(G)\}}\ m_{\{V\}}^{\{+\}}\ V,\ \backslash\mathrm{quad}\ \backslash\mathrm{operatorname{Coker}}(\backslash\mathrm{mathcal{D}})=\backslash\mathrm{bigoplus}_{\{V\ \backslash\mathrm{in}\ \backslash\mathrm{operatorname{Irr}}(G)\}}\ m_{\{V\}}^{\{-}\}\ V\ .$	$\mathrm{Ker}(\mathcal{D}) = \bigoplus_{V \in \mathrm{Irr}(G)} m_V^+ V,\quad \mathrm{Coker}(\mathcal{D}) = \bigoplus_{V \in \mathrm{Irr}(G)} m_V^- V.$	$\mathrm{Ker}(\mathcal{D}) = \bigoplus_{V \in \mathrm{Irr}(G)} m_V^+ V,\quad \mathrm{Coker}(\mathcal{D}) = \bigoplus_{V \in \mathrm{Irr}(G)} m_V^- V.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0180 (6)	91	■ 1.000	$\operatorname{Ind}_{\{G\}}(\mathcal{D}) := \bigoplus_{V \in \operatorname{Irr}(G)} (m_V^+ - m_V^-) V \in \widehat{R}(G),$	$\operatorname{Ind}_G(\mathcal{D}) := \bigoplus_{V \in \operatorname{Irr}(G)} (m_V^+ - m_V^-) V \in \widehat{R}(G),$	$\operatorname{Ind}_G(\mathcal{D}) := \bigoplus_{V \in \operatorname{Irr}(G)} (m_V^+ - m_V^-) V \in \widehat{R}(G),$
cardona-qft-methods_ EQ0182 (8)	92	■ 1.000	$\operatorname{t}\text{-}\operatorname{Ind}_{\{G\}}(\sigma) := \operatorname{Ind}_{\{G\}}(\sigma) \in \widehat{R}(G).$	$\operatorname{t} - \operatorname{Ind}_G(\sigma) := \operatorname{Ind}_G(\mathcal{D}).$	$\operatorname{t}\text{-}\operatorname{Ind}_G(\sigma) := \operatorname{Ind}_G(\mathcal{D}).$
cardona-qft-methods_ EQ0183	92	■ 1.000	$\operatorname{Ind}_{\{G\}}(\mathcal{D}) = \operatorname{t}\text{-}\operatorname{Ind}_{\{G\}}(\sigma) \in \widehat{R}(G).$	$\operatorname{Ind}_G(\mathcal{D}) = \operatorname{t} - \operatorname{Ind}_G(\sigma) \in \widehat{R}(G).$	$\operatorname{Ind}_G(\mathcal{D}) = \operatorname{t}\text{-}\operatorname{Ind}_G(\sigma) \in \widehat{R}(G).$
cardona-qft-methods_ EQ0184	93	■ 1.000	$c: V \rightarrow \operatorname{End}(W)$	$c: V \rightarrow \operatorname{End}(W)$	$c: V \rightarrow \operatorname{End}(W)$
cardona-qft-methods_ EQ0188 (3)	93	■ 1.000	$c: \mathbb{R}^3 \rightarrow \operatorname{End}(\mathcal{C}^2), \quad c(x_1, x_2, x_3) := \frac{1}{i} (x_1 \sigma_1 + x_2 \sigma_2 + x_3 \sigma_3)$	$c: \mathbb{R}^3 \rightarrow \operatorname{End}(\mathcal{C}^2), \quad c(x_1, x_2, x_3) := \frac{1}{i} (x_1 \sigma_1 + x_2 \sigma_2 + x_3 \sigma_3)$	$c: \mathbb{R}^3 \rightarrow \operatorname{End}(\mathcal{C}^2), \quad c(x_1, x_2, x_3) := \frac{1}{i} (x_1 \sigma_1 + x_2 \sigma_2 + x_3 \sigma_3)$
cardona-qft-methods_ EQ0189	93	■ 1.000	$W = \Lambda^*(V^{\mathcal{C}})$	$W = \Lambda^*(V^{\mathcal{C}})$	$W = \Lambda^*(V^{\mathcal{C}})$
cardona-qft-methods_ EQ0190	93	■ 1.000	$\varepsilon(v) \alpha := v \wedge \alpha, \quad \alpha \in W.$	$\varepsilon(v) \alpha := v \wedge \alpha, \quad \alpha \in W.$	$\varepsilon(v) \alpha := v \wedge \alpha, \quad \alpha \in W.$
cardona-qft-methods_ EQ0191 (4)	93	■ 1.000	$c(v) := \varepsilon_v - \iota_v: W \rightarrow W.$	$c(v) := \varepsilon_v - \iota_v: W \rightarrow W.$	$c(v) := \varepsilon_v - \iota_v: W \rightarrow W.$
cardona-qft-methods_ EQ0192 (5)	94	■ 1.000	$c: T^*M \rightarrow \operatorname{End}(E),$	$c: T^*M \rightarrow \operatorname{End}(E),$	$c: T^*M \rightarrow \operatorname{End}(E),$
cardona-qft-methods_ EQ0193	94	■ 1.000	$c: T_x^*M \rightarrow \operatorname{End}(E_x)$	$c: T_x^*M \rightarrow \operatorname{End}(E_x)$	$c: T_x^*M \rightarrow \operatorname{End}(E_x)$
cardona-qft-methods_ EQ0194	94	■ 1.000	$c(v) \omega = v \wedge \omega - \iota_v \omega.$	$c(v) \omega = v \wedge \omega - \iota_v \omega.$	$c(v) \omega = v \wedge \omega - \iota_v \omega.$
cardona-qft-methods_ EQ0195 (6)	94	■ 1.000	$\nabla_u(c(v)s) = c(\nabla_u^{LC} v)s + c(v)\nabla_u s,$	$\nabla_u(c(v)s) = c(\nabla_u^{LC} v)s + c(v)\nabla_u s,$	$\nabla_u(c(v)s) = c(\nabla_u^{LC} v)s + c(v)\nabla_u s,$
cardona-qft-methods_ EQ0196 (7)	95	■ 1.000	$\mathcal{D} := \sum_{j=1}^n c(e_j) \nabla_{e_j}: C^\infty(M, E) \rightarrow C^\infty(M, E),$	$\mathcal{D} := \sum_{j=1}^n c(e_j) \nabla_{e_j}: C^\infty(M, E) \rightarrow C^\infty(M, E),$	$\mathcal{D} := \sum_{j=1}^n c(e_j) \nabla_{e_j}: C^\infty(M, E) \rightarrow C^\infty(M, E),$
cardona-qft-methods_ EQ0197 (8)	95	■ 1.000	$\sigma_L(\mathcal{D})(x, \xi) = i c(\xi).$	$\sigma_L(\mathcal{D})(x, \xi) = i c(\xi).$	$\sigma_L(\mathcal{D})(x, \xi) = i c(\xi).$
cardona-qft-methods_ EQ0198	95	■ 1.000	$\nabla_{\partial/\partial x_i} = \frac{\partial}{\partial x_i}.$	$\nabla_{\partial/\partial x_i} = \frac{\partial}{\partial x_i}.$	$\nabla_{\partial/\partial x_i} = \frac{\partial}{\partial x_i}.$
cardona-qft-methods_ EQ0200	95	■ 1.000	$d^*: \Omega^*(M, E) \rightarrow \Omega^{*-1}(M, E)$	$d^*: \Omega^*(M, E) \rightarrow \Omega^{*-1}(M, E)$	$d^*: \Omega^*(M, E) \rightarrow \Omega^{*-1}(M, E)$
cardona-qft-methods_ EQ0203 (11)	96	■ 1.000	$\operatorname{Ker}(\mathcal{D}) \simeq H^*(M).$	$\operatorname{Ker}(\mathcal{D}) \simeq H^*(M).$	$\operatorname{Ker}(\mathcal{D}) \simeq H^*(M).$
cardona-qft-methods_ EQ0204 (12)	96	■ 1.000	$E = E^+ \oplus E^-,$	$E = E^+ \oplus E^-,$	$E = E^+ \oplus E^-,$
cardona-qft-methods_ EQ0205	96	■ 1.000	$c(v): E^\pm \rightarrow E^\mp.$	$c(v): E^\pm \rightarrow E^\mp.$	$c(v): E^\pm \rightarrow E^\mp.$
cardona-qft-methods_ EQ0206	96	■ 1.000	$\nabla_v: C^\infty(M, E^\pm) \rightarrow C^\infty(M, E^\pm).$	$\nabla_v: C^\infty(M, E^\pm) \rightarrow C^\infty(M, E^\pm).$	$\nabla_v: C^\infty(M, E^\pm) \rightarrow C^\infty(M, E^\pm).$
cardona-qft-methods_ EQ0207	96	■ 1.000	$\mathcal{D}: C^\infty(M, E^\pm) \rightarrow C^\infty(M, E^\mp).$	$\mathcal{D}: C^\infty(M, E^\pm) \rightarrow C^\infty(M, E^\mp).$	$\mathcal{D}: C^\infty(M, E^\pm) \rightarrow C^\infty(M, E^\mp).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0208 (13)	97	■ 1.000	$\backslash\mathrm{cal}{D}=\backslash\mathrm{left}(\backslash\mathrm{begin}{array}{cc} \ 0 \ \& \ \mathrm{cal}{D}^{\{-}\ \backslash \ \backslash \ \mathrm{cal}{D}^{\{+\}} \ \& \ 0 \ \backslash\mathrm{end}{array}\backslash\mathrm{right}) \ .$	$\mathcal{D} = \left(\begin{array}{cc} 0 & \mathcal{D}^- \\ \mathcal{D}^+ & 0 \end{array} \right).$	$\mathcal{D} \ = \ \left(\begin{array}{cc} 0 & \mathcal{D}^- \\ \mathcal{D}^+ & 0 \end{array} \right).$
cardona-qft-methods_ EQ0210	97	■ 1.000	$\backslash\mathrm{operatorname}{Ind}\backslash\mathrm{left}(\backslash\mathrm{cal}{D}^{\{+\}}\backslash\mathrm{right})=\backslash\mathrm{sum}_{\{j=0\}^{\{n\}}(-1)^{\{n\}} \ \backslash\mathrm{operatorname}{dim} \ H^{\{j\}}(M)$	$\mathrm{Ind}\left(\mathcal{D}^+\right)=\sum_{j=0}^n(-1)^n\dim H^j(M)$	$\mathrm{Ind}(\mathcal{D}^+) \ = \ \sum_{j=0}^n(-1)^n\dim H^j(M).$
cardona-qft-methods_ EQ0211	97	■ 1.000	$\backslash\Gamma\omega:=\mathrm{i}^{\{p(p-1)+l\}} \ * \ \omega, \ \backslash\mathrm{quad} \ \omega\in \backslash\Lambda^p(M) \ \backslash\mathrm{otimes} \ \backslash\mathrm{cal}{C} \ .$	$\Gamma\omega := i^{p(p-1)+l} * \omega, \quad \omega \in \Lambda^p(M) \otimes \mathcal{C}.$	$\Gamma\omega \ := \ i^{p(p-1)+l} * \omega, \quad \omega \in \Lambda^p(M) \otimes \mathcal{C}.$
cardona-qft-methods_ EQ0212	98	■ 1.000	$\backslash\mathrm{operatorname}{Ind}_{\{G\}}\backslash\mathrm{left}(\backslash\mathrm{cal}{D}^{\{+\}}\backslash\mathrm{right}) \ \backslash\mathrm{in} \ R(G) \ .$	$\mathrm{Ind}_G\left(\mathcal{D}^+\right) \in R(G).$	$\mathrm{Ind}_G(\mathcal{D}^+) \ \in \ R(G).$
cardona-qft-methods_ EQ0213 (1)	98	■ 1.000	$\backslash\mathrm{mathbf}{v}: M \ \backslash\mathrm{rightarrow} \ \backslash\mathrm{mathfrak}{g}=\backslash\mathrm{operatorname}{Lie} \ G \ .$	$\mathbf{v}: M \rightarrow \mathfrak{g} = \mathrm{Lie}\,G.$	$\mathbf{v}: M \rightarrow \mathfrak{g} = \mathrm{Lie}\,G.$
cardona-qft-methods_ EQ0215	99	■ 1.000	$G=S^{\{1\}}=\{z \ \backslash\mathrm{in} \ \backslash\mathrm{cal}{C}: z =1\}$	$G = S^1 = \{z \in \mathcal{C} : z = 1\}$	$G \ = \ S^1 \ = \ \left\{ z \in \mathcal{C} : z = 1 \right\}$
cardona-qft-methods_ EQ0216 (3)	99	■ 1.000	$v(x) \ \backslash\mathrm{neq} \ 0, \ \backslash\mathrm{quad} \ \backslash\mathrm{text} \ \{ \ \mathrm{for} \ \mathrm{all} \ \} \ \backslash\mathrm{quad} \ x \ \backslash\mathrm{notin} \ K \ .$	$v(x) \neq 0, \quad \text{for all} \quad x \notin K.$	$v(x) \neq 0, \quad \text{for all} \quad x \notin K.$
cardona-qft-methods_ EQ0218	100	■ 1.000	$c(\backslash\mathrm{x}i+v): \ \backslash\mathrm{pi}^{\{*\}} \ E^{\{+\}} \ \backslash\mathrm{rightarrow} \ \backslash\mathrm{pi}^{\{*\}} \ E^{\{-}\} \ .$	$c(\xi + v) : \pi^*E^+ \rightarrow \pi^*E^-.$	$c(\xi + v) : \pi^*E^+ \rightarrow \pi^*E^-.$
cardona-qft-methods_ EQ0219	100	■ 1.000	$\backslash\mathrm{mathrm}{t}-\backslash\mathrm{operatorname}{Ind}_{\{G\}}(c(\backslash\mathrm{x}i+v)) \ \backslash\mathrm{in} \ \backslash\mathrm{widehat}{R}(G) \ .$	$t - \mathrm{Ind}_G(c(\xi + v)) \in \widehat{R}(G).$	$t\text{-}\mathrm{Ind}_G\left(c(\xi + v)\right) \ \in \ \widehat{R}(G).$
cardona-qft-methods_ EQ0220 (5)	100	■ 1.000	$\backslash\mathrm{mathrm}{t}-\backslash\mathrm{operatorname}{Ind}_{\{G\}}(E, \ \backslash\mathrm{mathbf}{v}):=\backslash\mathrm{mathrm}{t}-\backslash\mathrm{operatorname}{Ind}_{\{G\}}(c(\backslash\mathrm{x}i+v)) \ .$	$t - \mathrm{Ind}_G(E, \mathbf{v}) := t - \mathrm{Ind}_G(c(\xi + v)).$	$t\text{-}\mathrm{Ind}_G(E, \mathbf{v}) := t\text{-}\mathrm{Ind}_G\left(c(\xi + v)\right).$
cardona-qft-methods_ EQ0221 (6)	101	■ 1.000	$\backslash\mathrm{cal}{D}_{\{f \ v\}}=\backslash\mathrm{cal}{D}+\mathrm{i} \ c(f \ v),$	$\mathcal{D}_{fv} = \mathcal{D} + ic(fv),$	$\mathcal{D}_{fv} \ = \ \mathcal{D} \ + \ ic(fv),$
cardona-qft-methods_ EQ0222 (7)	101	■ 1.000	$\backslash\mathrm{operatorname}{Ker} \ \backslash\mathrm{cal}{D}_{\{f \ v\}}^{\{+\}}=\backslash\mathrm{sum}_{\{V \ \backslash\mathrm{in} \ \backslash\mathrm{operatorname}{Irr}\}(G)\} \ m_{\{V\}}^{\{+\}} \ \backslash\mathrm{cdot} \ V, \ \backslash\mathrm{quad} \ \backslash\mathrm{operatorname}{Coker} \ \backslash\mathrm{cal}{D}_{\{f \ v\}}^{\{+\}}=\backslash\mathrm{sum}_{\{V \ \backslash\mathrm{in} \ \backslash\mathrm{operatorname}{Irr}\}(G)\} \ m_{\{V\}}^{\{-}\} \ \backslash\mathrm{cdot} \ V,$	$\mathrm{Ker}\,\mathcal{D}_{fv}^+ = \sum_{V \in \mathrm{Irr}(G)} m_V^+ \cdot V, \quad \mathrm{Coker}\,\mathcal{D}_{fv}^+ = \sum_{V \in \mathrm{Irr}(G)} m_V^- \cdot V,$	$\mathrm{Ker}\,\mathcal{D}_{fv}^+ = \sum_{V \in \mathrm{Irr}(G)} m_V^+ \cdot V, \quad \mathrm{Coker}\,\mathcal{D}_{fv}^+ = \sum_{V \in \mathrm{Irr}(G)} m_V^- \cdot V,$
cardona-qft-methods_ EQ0223 (8)	101	■ 1.000	$\backslash\mathrm{operatorname}{Ind}_{\{G\}}\backslash\mathrm{left}(\backslash\mathrm{cal}{D}_{\{f \ v\}}^{\{+\}} \ \backslash\mathrm{right})=\backslash\mathrm{sum}_{\{V \ \backslash\mathrm{in} \ \backslash\mathrm{operatorname}{Irr}\}(G)\} \ \backslash\mathrm{left}(m_{\{V\}}^{\{+\}}-m_{\{V\}}^{\{-}\}\backslash\mathrm{right}) \ \backslash\mathrm{cdot} \ V \ \backslash\mathrm{in} \ \backslash\mathrm{widehat}{R}(G) \ .$	$\mathrm{Ind}_G\left(\mathcal{D}_{fv}^+\right) = \sum_{V \in \mathrm{Irr}(G)} (m_V^+ - m_V^-) \cdot V \in \widehat{R}(G).$	$\mathrm{Ind}_G(\mathcal{D}_{fv}^+) \ = \ \sum_{V \in \mathrm{Irr}(G)} (m_V^+ - m_V^-) \cdot V \in \widehat{R}(G).$
cardona-qft-methods_ EQ0224 (9)	101	■ 1.000	$\backslash\mathrm{operatorname}{Ind}_{\{G\}}(\backslash\mathrm{cal}{D}, \ \backslash\mathrm{mathbf}{v}):=\backslash\mathrm{operatorname}{Ind}_{\{G\}}\backslash\mathrm{left}(\backslash\mathrm{cal}{D}_{\{f \ v\}}^{\{+\}}\backslash\mathrm{right}),$	$\mathrm{Ind}_G(\mathcal{D}, \mathbf{v}) := \mathrm{Ind}_G\left(\mathcal{D}_{fv}^+\right),$	$\mathrm{Ind}_G(\mathcal{D}, \mathbf{v}) \ := \ \mathrm{Ind}_G(\mathcal{D}_{fv}^+),$
cardona-qft-methods_ EQ0225	102	■ 1.000	$E^{\{ \ \backslash\mathrm{pm}\}}=M \ \backslash\mathrm{times} \ \backslash\mathrm{mathbb}{R}$	$E^\pm = M \times \mathbb{R}$	$E^\pm \ = \ M \times \mathbb{R}$
cardona-qft-methods_ EQ0226	102	■ 1.000	$c(\backslash\mathrm{x}i):=\frac{1}{i}\backslash\mathrm{left}(\backslash\mathrm{x}i_{\{1\}} \ \backslash\mathrm{sigma}_{\{1\}}+\backslash\mathrm{x}i_{\{2\}} \ \backslash\mathrm{sigma}_{\{2\}}\backslash\mathrm{right}), \ \backslash\mathrm{quad} \ \backslash\mathrm{x}i=\backslash\mathrm{left}(\backslash\mathrm{x}i_{\{1\}}, \ \backslash\mathrm{x}i_{\{2\}} \ \backslash\mathrm{right}) \ \backslash\mathrm{in} \ T^{\{*\}} M \ \backslash\mathrm{simeq} \ \backslash\mathrm{mathbb}{R}^{\{2\}},$	$c(\xi) := \frac{1}{i} \left(\xi_1 \sigma_1 + \xi_2 \sigma_2 \right), \quad \xi = (\xi_1, \xi_2) \in T^*M \simeq \mathbb{R}^2,$	$c(\xi) \ := \ \frac{1}{i} \left(\xi_1 \sigma_1 + \xi_2 \sigma_2 \right), \quad \xi = (\xi_1, \xi_2) \in T^*M \simeq \mathbb{R}^2,$
cardona-qft-methods_ EQ0228 (10)	102	■ 1.000	$\backslash\mathrm{operatorname}{Ind}(\backslash\mathrm{cal}{D}, \ \backslash\mathrm{mathbf}{v})=\backslash\mathrm{mathrm}{t}-\backslash\mathrm{operatorname}{Ind}(E, \ \backslash\mathrm{mathbf}{v}) \ .$	$\mathrm{Ind}(\mathcal{D}, \mathbf{v}) = t - \mathrm{Ind}(E, \mathbf{v}).$	$\mathrm{Ind}(\mathcal{D}, \mathbf{v}) \ = \ t\text{-}\mathrm{Ind}(E, \mathbf{v}).$
cardona-qft-methods_ EQ0229 (11)	102	■ 1.000	$\backslash\mathrm{operatorname}{Ind}_{\{G\}}(\backslash\mathrm{cal}{D}, \ \backslash\mathrm{mathbf}{v})=\backslash\mathrm{operatorname}{Ind}_{\{G\}}(\backslash\mathrm{cal}{D}) \ .$	$\mathrm{Ind}_G(\mathcal{D}, \mathbf{v}) = \mathrm{Ind}_G(\mathcal{D}).$	$\mathrm{Ind}_G(\mathcal{D}, \mathbf{v}) \ = \ \mathrm{Ind}_G(\mathcal{D}).$
cardona-qft-methods_ EQ0230	103	■ 1.000	$M \ \backslash\mathrm{backslash} \ \backslash\Sigma=M_{\{1\}} \ \backslash\mathrm{sqcup} \ M_{\{2\}} \ .$	$M \backslash \Sigma = M_1 \sqcup M_2.$	$M \backslash \Sigma \ = \ M_1 \sqcup M_2.$
cardona-qft-methods_ EQ0231 (12)	103	■ 1.000	$\backslash\mathrm{operatorname}{Ind}_{\{G\}}(\backslash\mathrm{cal}{D}, \ \backslash\mathrm{mathbf}{v})=\backslash\mathrm{operatorname}{Ind}_{\{G\}}\backslash\mathrm{left}(\backslash\mathrm{cal}{D}_{\{1\}}, \ \backslash\mathrm{mathbf}{v}_{\{1\}}\backslash\mathrm{right})+\backslash\mathrm{operatorname}{Ind}_{\{G\}}\backslash\mathrm{left}(\backslash\mathrm{cal}{D}_{\{2\}}, \ \backslash\mathrm{mathbf}{v}_{\{2\}}\backslash\mathrm{right}) \ .$	$\mathrm{Ind}_G(\mathcal{D}, \mathbf{v}) = \mathrm{Ind}_G\left(\mathcal{D}_1, \mathbf{v}_1\right) + \mathrm{Ind}_G\left(\mathcal{D}_2, \mathbf{v}_2\right).$	$\mathrm{Ind}_G(\mathcal{D}, \mathbf{v}) \ = \ \mathrm{Ind}_G(\mathcal{D}_1, \mathbf{v}_1) \ + \ \mathrm{Ind}_G(\mathcal{D}_2, \mathbf{v}_2).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0232	109	■ 1.000	$\chi(X)=s_{\{0\}}-s_{\{1\}}+s_{\{2\}}-\cdots$	$\chi(X) = s_0 - s_1 + s_2 - \cdots$	$\chi(X) = s_0 - s_1 + s_2 - \cdots$
cardona-qft-methods_ EQ0235	112	■ 1.000	$\left[\mathbb{P}^{n-1}\right]=1+\mathbb{L}+\cdots+\mathbb{L}^n=\frac{\mathbb{L}^n-1}{\mathbb{L}-1}$	$\left[\mathbb{P}^{n-1}\right] = 1 + \mathbb{L} + \cdots + \mathbb{L}^n = \frac{\mathbb{L}^n - 1}{\mathbb{L} - 1}$	$\left[\mathbb{P}^{n-1}\right] = 1 + \mathbb{L} + \cdots + \mathbb{L}^n = \frac{\mathbb{L}^n - 1}{\mathbb{L} - 1}$
cardona-qft-methods_ EQ0239 (4)	114	■ 1.000	$\int c(TX) \cap [X] = \chi(X)$	$\int c(TX) \cap [X] = \chi(X)$	$\int c(TX) \cap [X] = \chi(X) \quad .$
cardona-qft-methods_ EQ0240	115	■ 1.000	$f_{\{*\}}\left(\mathbb{1}_{\{Z\}}\right)(p)=\chi\left(f^{-1}(p) \cap Z\right)$	$f_{*}\left(\mathbb{1}_Z\right)(p)=\chi\left(f^{-1}(p) \cap Z\right)$	$f_{*}\left(\mathbb{1}_Z\right)(p)=\chi\left(f^{-1}(p) \cap Z\right)$
cardona-qft-methods_ EQ0242 (5)	116	■ 1.000	$\int c_{\{\mathrm{SM}\}}(X)=\kappa_{*} c_{\{\mathrm{SM}\}}\left(\mathbb{1}_{\{X\}}\right)=c_{\{\mathrm{SM}\}}\left(\kappa_{*}\mathbb{1}_{\{X\}}\right)=\chi(X) c_{\{\mathrm{SM}\}}\left(\mathbb{1}_{\mathrm{pt}}\right)=\chi(X)$	$\int c_{\mathrm{SM}}(X)=\kappa_{*} c_{\mathrm{SM}}\left(\mathbb{1}_X\right)=c_{\mathrm{SM}}\left(\kappa_{*} \mathbb{1}_X\right)=\chi(X) c_{\mathrm{SM}}\left(\mathbb{1}_{\mathrm{pt}}\right)=\chi(X)$	$\int c_{\mathrm{SM}}(X)=\kappa_{*} c_{\mathrm{SM}}\left(\mathbb{1}_X\right)=c_{\mathrm{SM}}\left(\kappa_{*} \mathbb{1}_X\right)=\chi(X) c_{\mathrm{SM}}\left(\mathbb{1}_{\mathrm{pt}}\right)=\chi(X)$
cardona-qft-methods_ EQ0243	116	■ 1.000	$\begin{aligned} c_{\{\mathrm{SM}\}}(X) &= c_{\{\mathrm{SM}\}}\left(\mathbb{1}_X\right) &= c_{\{\mathrm{SM}\}}\left(\mathbb{1}_Z+\mathbb{1}_{X \backslash Z}\right) \\ &= c_{\{\mathrm{SM}\}}\left(\mathbb{1}_Z\right)+c_{\{\mathrm{SM}\}}\left(\mathbb{1}_{X \backslash Z}\right) &= c_{\{\mathrm{SM}\}}(Z)+c_{\{\mathrm{SM}\}}(X \backslash Z) \end{aligned}$	$\begin{aligned} c_{\mathrm{SM}}(X) &= c_{\mathrm{SM}}\left(\mathbb{1}_X\right) = c_{\mathrm{SM}}\left(\mathbb{1}_Z+\mathbb{1}_{X \backslash Z}\right) \\ &= c_{\mathrm{SM}}\left(\mathbb{1}_Z\right)+c_{\mathrm{SM}}\left(\mathbb{1}_{X \backslash Z}\right) = c_{\mathrm{SM}}(Z)+c_{\mathrm{SM}}(X \backslash Z) \end{aligned}$	$\begin{aligned} c_{\mathrm{SM}}(X) &= c_{\mathrm{SM}}\left(\mathbb{1}_X\right) = c_{\mathrm{SM}}\left(\mathbb{1}_Z+\mathbb{1}_{X \backslash Z}\right) \\ &= c_{\mathrm{SM}}\left(\mathbb{1}_Z\right)+c_{\mathrm{SM}}\left(\mathbb{1}_{X \backslash Z}\right) = c_{\mathrm{SM}}(Z)+c_{\mathrm{SM}}(X \backslash Z) \end{aligned}$
cardona-qft-methods_ EQ0244	116	■ 1.000	$c_{\{\mathrm{SM}\}}\left(\mathbb{P}^1\right)=c\left(T \mathbb{P}^1\right) \cap\left[\mathbb{P}^1\right]=\left(1+c_1\left(T \mathbb{P}^1\right)\right) \cap\left[\mathbb{P}^1\right]=\left[\mathbb{P}^1\right]+\chi\left(\mathbb{P}^1\right)\left[\mathbb{P}^0\right]=\left[\mathbb{P}^1\right]+2\left[\mathbb{P}^0\right] .$	$c_{\mathrm{SM}}\left(\mathbb{P}^1\right)=c\left(T \mathbb{P}^1\right) \cap\left[\mathbb{P}^1\right]=\left(1+c_1\left(T \mathbb{P}^1\right)\right) \cap\left[\mathbb{P}^1\right]=\left[\mathbb{P}^1\right]+\chi\left(\mathbb{P}^1\right)\left[\mathbb{P}^0\right]=\left[\mathbb{P}^1\right]+2\left[\mathbb{P}^0\right] .$	$c_{\mathrm{SM}}\left(\mathbb{P}^1\right)=c\left(T \mathbb{P}^1\right) \cap\left[\mathbb{P}^1\right]=\left(1+c_1\left(T \mathbb{P}^1\right)\right) \cap\left[\mathbb{P}^1\right]=\left[\mathbb{P}^1\right]+\chi\left(\mathbb{P}^1\right)\left[\mathbb{P}^0\right]=\left[\mathbb{P}^1\right]+2\left[\mathbb{P}^0\right] .$
cardona-qft-methods_ EQ0245 (6)	116	■ 1.000	$c_{\{\mathrm{SM}\}}\left(\mathbb{P}^n\right)=\left[\mathbb{P}^n\right]+\binom{n+1}{1}\left[\mathbb{P}^{n-1}\right]+\binom{n+1}{2}\left[\mathbb{P}^{n-2}\right]+\cdots+\binom{n+1}{n}\left[\mathbb{P}^0\right] .$	$c_{\mathrm{SM}}\left(\mathbb{P}^n\right)=\left[\mathbb{P}^n\right]+\binom{n+1}{1}\left[\mathbb{P}^{n-1}\right]+\binom{n+1}{2}\left[\mathbb{P}^{n-2}\right]+\cdots+\binom{n+1}{n}\left[\mathbb{P}^0\right] .$	$c_{\mathrm{SM}}\left(\mathbb{P}^n\right)=\left[\mathbb{P}^n\right]+\binom{n+1}{1}\left[\mathbb{P}^{n-1}\right]+\binom{n+1}{2}\left[\mathbb{P}^{n-2}\right]+\cdots+\binom{n+1}{n}\left[\mathbb{P}^0\right] .$
cardona-qft-methods_ EQ0246	116	■ 1.000	$c_{\{\mathrm{SM}\}}\left(\mathbb{P}^n\right)=\left(1+H\right)^{n+1} \cap\left[\mathbb{P}^n\right]$	$c_{\mathrm{SM}}\left(\mathbb{P}^n\right)=\left(1+H\right)^{n+1} \cap\left[\mathbb{P}^n\right]$	$c_{\mathrm{SM}}\left(\mathbb{P}^n\right)=\left(1+H\right)^{n+1} \cap\left[\mathbb{P}^n\right]$
cardona-qft-methods_ EQ0247	117	■ 1.000	$c_{\{\mathrm{SM}\}}(C)=c_{\{\mathrm{SM}\}}\left(\mathbb{1}_C\right)=c(T C) \cap[C]=\left(1+c_1(T C)\right) \cap[C]=2\left[\mathbb{P}^1\right]+2\left[\mathbb{P}^0\right] .$	$c_{\mathrm{SM}}(C)=c_{\mathrm{SM}}\left(\mathbb{1}_C\right)=c(T C) \cap[C]=\left(1+c_1(T C)\right) \cap[C]=2\left[\mathbb{P}^1\right]+2\left[\mathbb{P}^0\right] .$	$c_{\mathrm{SM}}(C)=c_{\mathrm{SM}}\left(\mathbb{1}_C\right)=c(T C) \cap[C]=\left(1+c_1(T C)\right) \cap[C]=2\left[\mathbb{P}^1\right]+2\left[\mathbb{P}^0\right] .$
cardona-qft-methods_ EQ0248	117	■ 1.000	$c_{\{\mathrm{SM}\}}\left(L_1\right)+c_{\{\mathrm{SM}\}}\left(L_2\right)-c_{\{\mathrm{SM}\}}\left(L_1 \cap L_2\right)=2 c_{\{\mathrm{SM}\}}\left(\mathbb{P}^1\right)-c_{\{\mathrm{SM}\}}\left(\mathbb{P}^0\right)=2\left[\mathbb{P}^1\right]+3\left[\mathbb{P}^0\right] .$	$c_{\mathrm{SM}}\left(L_1\right)+c_{\mathrm{SM}}\left(L_2\right)-c_{\mathrm{SM}}\left(L_1 \cap L_2\right)=2 c_{\mathrm{SM}}\left(\mathbb{P}^1\right)-c_{\mathrm{SM}}\left(\mathbb{P}^0\right)=2\left[\mathbb{P}^1\right]+3\left[\mathbb{P}^0\right] .$	$c_{\mathrm{SM}}\left(L_1\right)+c_{\mathrm{SM}}\left(L_2\right)-c_{\mathrm{SM}}\left(L_1 \cap L_2\right)=2 c_{\mathrm{SM}}\left(\mathbb{P}^1\right)-c_{\mathrm{SM}}\left(\mathbb{P}^0\right)=2\left[\mathbb{P}^1\right]+3\left[\mathbb{P}^0\right] .$
cardona-qft-methods_ EQ0249	118	■ 1.000	$S(\phi)=\int L(\phi) d^D x$	$S(\phi)=\int L(\phi) d^D x$	$S(\phi)=\int L(\phi) d^D x$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0251	118	■ 1.000	$G(\phi)=\frac{1}{N!}\int\limits_{\sum_i p_i=0}\hat{\phi}(p_1)\cdots\hat{\phi}(p_N)U(G(p_1,\dots,p_N))dp_1\cdots dp_N$	$G(\phi)=\frac{1}{N!}\int\limits_{\sum_i p_i=0}\hat{\phi}(p_1)\cdots\hat{\phi}(p_N)U(G(p_1,\dots,p_N))dp_1\cdots dp_N$	$G(\phi)=\frac{1}{N!}\int\limits_{\sum_i p_i=0}\hat{\phi}(p_1)\cdots\hat{\phi}(p_N)U(G(p_1,\dots,p_N))dp_1\cdots dp_N$
cardona-qft-methods_ EQ0252	118	■ 1.000	$U(G(p_1,\dots,p_N))=\int I_G(k_1,\dots,k_\ell,p_1,\dots,p_N)d^Dk_1\cdots d^Dk_\ell$	$U(G(p_1,\dots,p_N))=\int I_G(k_1,\dots,k_\ell,p_1,\dots,p_N)d^Dk_1\cdots d^Dk_\ell$	$U(G(p_1,\dots,p_N))=\int I_G(k_1,\dots,k_\ell,p_1,\dots,p_N)d^Dk_1\cdots d^Dk_\ell$
cardona-qft-methods_ EQ0253 (2)	119	■ 1.000	$U(G)=\frac{\Gamma(n-D\ell/2)}{(4\pi)^{\ell D/2}}\int_{\sigma_n}\frac{P_G(\underline{t},p)^{-n+D\ell/2}\omega_n}{\Psi_G(\underline{t})^{-n+D(\ell+1)/2}}$	$U(G)=\frac{\Gamma(n-D\ell/2)}{(4\pi)^{\ell D/2}}\int_{\sigma_n}\frac{P_G(\underline{t},p)^{-n+D\ell/2}\omega_n}{\Psi_G(\underline{t})^{-n+D(\ell+1)/2}}$	$U(G)=\frac{\Gamma(n-D\ell/2)}{(4\pi)^{\ell D/2}}\int_{\sigma_n}\frac{P_G(\underline{t},p)^{-n+D\ell/2}\omega_n}{\Psi_G(\underline{t})^{-n+D(\ell+1)/2}}$
cardona-qft-methods_ EQ0255	119	■ 1.000	$U(G)=\frac{\Gamma(2-D/2)}{(4\pi)^{D/2}}\int_{\sigma_2}\frac{(p^2t_1t_2)^{-2+D/2}\omega_2}{(t_1+t_2)^{-2+D}}$	$U(G)=\frac{\Gamma(2-D/2)}{(4\pi)^{D/2}}\int_{\sigma_2}\frac{(p^2t_1t_2)^{-2+D/2}\omega_2}{(t_1+t_2)^{-2+D}}$	$U(G)=\frac{\Gamma(2-D/2)}{(4\pi)^{D/2}}\int_{\sigma_2}\frac{(p^2t_1t_2)^{-2+D/2}\omega_2}{(t_1+t_2)^{-2+D}}$
cardona-qft-methods_ EQ0256	120	■ 1.000	$\int_{\text{cycle } \sigma} \frac{\text{diff. form}}{\Psi_G(t)^M}$	$\int_{\text{cycle } \sigma} \frac{\text{diff. form}}{\Psi_G(t)^M}$	$\int_{\text{cycle } \sigma} \frac{\text{diff. form}}{\Psi_G(t)^M}$
cardona-qft-methods_ EQ0257	120	■ 1.000	$\zeta(n_1,\dots,n_r)=\sum_{0\leq k_1\leq\dots\leq k_r}\frac{1}{k_1^{n_1}\cdots k_r^{n_r}}$	$\zeta(n_1,\dots,n_r)=\sum_{0\leq k_1\leq\dots\leq k_r}\frac{1}{k_1^{n_1}\cdots k_r^{n_r}}$	$\zeta(n_1,\dots,n_r)=\sum_{0\leq k_1\leq\dots\leq k_r}\frac{1}{k_1^{n_1}\cdots k_r^{n_r}}$
cardona-qft-methods_ EQ0258	121	■ 1.000	$\Psi_G(\underline{t}_1,\dots,\underline{t}_n):=\sum_T\prod_{e\notin T}t_e$	$\Psi_G(t_1,\dots,t_n):=\sum_T\prod_{e\notin T}t_e$	$\Psi_G(t_1,\dots,t_n):=\sum_T\prod_{e\notin T}t_e$
cardona-qft-methods_ EQ0259	122	■ 1.000	$\Psi_G(\underline{t})=t_2t_3+t_1t_3+t_1t_2=t_1t_2t_3\left(\frac{1}{t_1}+\frac{1}{t_2}+\frac{1}{t_3}\right)$	$\Psi_G(\underline{t})=t_2t_3+t_1t_3+t_1t_2=t_1t_2t_3\left(\frac{1}{t_1}+\frac{1}{t_2}+\frac{1}{t_3}\right)$	$\Psi_G(\underline{t})=t_2t_3+t_1t_3+t_1t_2=t_1t_2t_3\left(\frac{1}{t_1}+\frac{1}{t_2}+\frac{1}{t_3}\right)$
cardona-qft-methods_ EQ0260	122	■ 1.000	$\Psi_G(\underline{t})=t_1\cdots t_n\left(\frac{1}{t_1}+\cdots+\frac{1}{t_n}\right)$	$\Psi_G(\underline{t})=t_1\cdots t_n\left(\frac{1}{t_1}+\cdots+\frac{1}{t_n}\right)$	$\Psi_G(\underline{t})=t_1\cdots t_n\left(\frac{1}{t_1}+\cdots+\frac{1}{t_n}\right)$
cardona-qft-methods_ EQ0266	127	■ 1.000	$\Psi_G(t_1,\dots,t_{n_1},u_1,\dots,u_{n_2})\neq 0$	$\Psi_G(t_1,\dots,t_{n_1},u_1,\dots,u_{n_2})\neq 0$	$\Psi_G(t_1,\dots,t_{n_1},u_1,\dots,u_{n_2})\neq 0$
cardona-qft-methods_ EQ0269	129	■ 1.000	$c_{\mathrm{SM}}(\mathbb{1}_{\hat{X}})=a_0[\mathbb{P}^0]+a_1[\mathbb{P}^1]+\cdots+a_n[\mathbb{P}^n]$	$c_{\mathrm{SM}}(\mathbb{1}_{\hat{X}})=a_0[\mathbb{P}^0]+a_1[\mathbb{P}^1]+\cdots+a_n[\mathbb{P}^n]$	$c_{\mathrm{SM}}(\mathbb{1}_{\hat{X}})=a_0[\mathbb{P}^0]+a_1[\mathbb{P}^1]+\cdots+a_n[\mathbb{P}^n]$
cardona-qft-methods_ EQ0270	130	■ 1.000	$\begin{aligned} c_{\mathrm{SM}}(\mathbb{1}_{\hat{X}}) &= c_{\mathrm{SM}}(\mathbb{1}_{\mathbb{P}^n} - \mathbb{1}_{\mathbb{P}^{n-1}}) \\ &= ((1+H)^{n+1} - (1+H)^n H) \cap [\mathbb{P}^n] = (1+H)^n \cap [\mathbb{P}^n] \\ &= \sum_{i=0}^n \binom{n}{i} [\mathbb{P}^i] \rightsquigarrow F_{\mathbb{A}^n}(T) = \sum_{i=0}^n \binom{n}{i} T^i \end{aligned}$	$c_{\mathrm{SM}}(\mathbb{1}_{\hat{X}}) = c_{\mathrm{SM}}(\mathbb{1}_{\mathbb{P}^n} - \mathbb{1}_{\mathbb{P}^{n-1}}) = ((1+H)^{n+1} - (1+H)^n H) \cap [\mathbb{P}^n] = (1+H)^n \cap [\mathbb{P}^n] = \sum_{i=0}^n \binom{n}{i} [\mathbb{P}^i] \rightsquigarrow F_{\mathbb{A}^n}(T) = \sum_{i=0}^n \binom{n}{i} T^i$	$c_{\mathrm{SM}}(\mathbb{1}_{\hat{X}}) = c_{\mathrm{SM}}(\mathbb{1}_{\mathbb{P}^n} - \mathbb{1}_{\mathbb{P}^{n-1}}) = ((1+H)^{n+1} - (1+H)^n H) \cap [\mathbb{P}^n] = (1+H)^n \cap [\mathbb{P}^n] = \sum_{i=0}^n \binom{n}{i} [\mathbb{P}^i] \rightsquigarrow F_{\mathbb{A}^n}(T) = \sum_{i=0}^n \binom{n}{i} T^i$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0271	130	■ 1.000	$\begin{aligned} C_{\{G\}}(T) &= F_{\mathbb{A}^n \setminus \mathbb{A}^{n-1}}(T) = (1+T)^n - (1+T)^{n-1} \\ &= (1+T-1)(1+T)^{n-1} = T(1+T)^{n-1} \end{aligned}$	$C_G(T) = F_{\mathbb{A}^n \setminus \mathbb{A}^{n-1}}(T) = (1+T)^n - (1+T)^{n-1} \\ = (1+T-1)(1+T)^{n-1} = T(1+T)^{n-1}$	$C_G(T) = F_{\mathbb{A}^n \setminus \mathbb{A}^{n-1}}(T) = (1+T)^n - (1+T)^{n-1} \\ = (1+T-1)(1+T)^{n-1} = T(1+T)^{n-1}$
cardona-qft-methods_ EQ0273	132	■ 1.000	$T_{\{G\}}(x, y) = T_{G/e}(x, y) + T_{G \setminus e}(x, y)$	$T_G(x, y) = T_{G/e}(x, y) + T_{G \setminus e}(x, y)$	$T_G(x, y) = T_{G/e}(x, y) + T_{G \setminus e}(x, y)$
cardona-qft-methods_ EQ0274	133	■ 1.000	$\gamma^{b_0(G)} \alpha^{\#V(G) - b_0(G)} \beta^{b_1(G)} T_G \left(\frac{\gamma x}{\alpha}, \frac{y}{\beta} \right)$	$\gamma^{b_0(G)} \alpha^{\#V(G) - b_0(G)} \beta^{b_1(G)} T_G \left(\frac{\gamma x}{\alpha}, \frac{y}{\beta} \right)$	$\gamma^{b_0(G)} \alpha^{\#V(G) - b_0(G)} \beta^{b_1(G)} T_G \left(\frac{\gamma x}{\alpha}, \frac{y}{\beta} \right)$
cardona-qft-methods_ EQ0275	134	■ 1.000	$T_{\{G_1\} \cup G_2}(x, y) = T_{\{G_1\}}(x, y) \cdot T_{G_2}(x, y)$	$T_{G_1 \cup G_2}(x, y) = T_{G_1}(x, y) \cdot T_{G_2}(x, y)$	$T_{G_1 \cup G_2}(x, y) = T_{G_1}(x, y) \cdot T_{G_2}(x, y)$
cardona-qft-methods_ EQ0276	134	■ 1.000	$\begin{aligned} T_{G_2e}(x, y) &= T_G(x, y) + y T_{G/e}(x, y) \\ &= T_{G \setminus e}(x, y) + (y+1) T_{G/e}(x, y) \end{aligned}$	$T_{G_{2e}}(x, y) = T_G(x, y) + y T_{G/e}(x, y) \\ = T_{G \setminus e}(x, y) + (y+1) T_{G/e}(x, y)$	$T_{G_{2e}}(x, y) = T_G(x, y) + y T_{G/e}(x, y) \\ = T_{G \setminus e}(x, y) + (y+1) T_{G/e}(x, y)$
cardona-qft-methods_ EQ0277	134	■ 1.000	$\sum_{m \geq 0} T_{G_{me}}(x, y) \frac{s^m}{m!} = e^s \left(T_{G \setminus e}(x, y) + \frac{e^{(y-1)s} - 1}{y-1} T_{G/e}(x, y) \right)$	$\sum_{m \geq 0} T_{G_{me}}(x, y) \frac{s^m}{m!} = e^s \left(T_{G \setminus e}(x, y) + \frac{e^{(y-1)s} - 1}{y-1} T_{G/e}(x, y) \right)$	$\sum_{m \geq 0} T_{G_{me}}(x, y) \frac{s^m}{m!} = e^s \left(T_{G \setminus e}(x, y) + \frac{e^{(y-1)s} - 1}{y-1} T_{G/e}(x, y) \right) \quad ,$
cardona-qft-methods_ EQ0280	136	■ 1.000	$\chi \left(Y_{\{G_m\}} \right) = (-1)^{m-1} \left(\chi(Y_G) - \chi(Y_{G/e}) \right),$	$\chi(Y_{G_{me}}) = (-1)^{m-1} \left(\chi(Y_G) - \chi(Y_{G/e}) \right),$	$\chi(Y_{G_{me}}) = (-1)^{m-1} \left(\chi(Y_G) - \chi(Y_{G/e}) \right) \quad ,$
cardona-qft-methods_ EQ0281 (1)	137	■ 1.000	$\Psi_{\{G\}} = t_e \Psi_{G \setminus e} + \Psi_{G/e}$	$\Psi_G = t_e \Psi_{G \setminus e} + \Psi_{G/e}$	$\Psi_G = t_e \Psi_{G \setminus e} + \Psi_{G/e} \quad .$
cardona-qft-methods_ EQ0285	137	■ 1.000	$C_{\{G\}}(T) = C_{G \setminus e, G/e}(T) + (T-1) C_{G \setminus e}(T)$	$C_G(T) = C_{G \setminus e, G/e}(T) + (T-1) C_{G \setminus e}(T)$	$C_G(T) = C_{G \setminus e, G/e}(T) + (T-1) C_{G \setminus e}(T) \quad .$
cardona-qft-methods_ EQ0288	138	■ 1.000	$C_{\{G\}}(T) = (2T-1) T(1+T)^2 - T(T-1)(1+T)^2 + T(1+T) = T(1+T)(1+T+T^2)$	$C_G(T) = (2T-1) T(1+T)^2 - T(T-1)(1+T)^2 + T(1+T) = T(1+T)(1+T+T^2)$	$C_G(T) = (2T-1) T(1+T)^2 - T(T-1)(1+T)^2 + T(1+T) = T(1+T)(1+T+T^2)$
cardona-qft-methods_ EQ0291	139	■ 1.000	$\Psi_{\{G\}} = t_e \Psi_{G \setminus e} + \Psi_{G/e}$	$\Psi_G = t_e \Psi_{G \setminus e} + \Psi_{G/e}$	$\Psi_G = t_e \Psi_{G \setminus e} + \Psi_{G/e}$
cardona-qft-methods_ EQ0293	142	■ 1.000	$\mathbb{U}(G) \equiv -\mathbb{U}(G \setminus e) \pmod{\mathbb{L}}$	$\mathbb{U}(G) \equiv -\mathbb{U}(G \setminus e) \pmod{\mathbb{L}}$	$\mathbb{U}(G) \equiv -\mathbb{U}(G \setminus e) \pmod{\mathbb{L}}$
cardona-qft-methods_ EQ0294	142	■ 1.000	$\mathbb{U}(G) = (\mathbb{L}-1) \mathbb{U}(G \setminus e) \equiv -\mathbb{U}(G \setminus e) \pmod{\mathbb{L}};$	$\mathbb{U}(G) = (\mathbb{L}-1) \mathbb{U}(G \setminus e) \equiv -\mathbb{U}(G \setminus e) \pmod{\mathbb{L}};$	$\mathbb{U}(G) = (\mathbb{L}-1) \mathbb{U}(G \setminus e) \equiv -\mathbb{U}(G \setminus e) \pmod{\mathbb{L}};$
cardona-qft-methods_ EQ0295	142	■ 1.000	$\mathbb{U}(G) = \mathbb{L} \mathbb{U}(G \setminus e) \equiv 0 \pmod{\mathbb{L}}$	$\mathbb{U}(G) = \mathbb{L} \mathbb{U}(G \setminus e) \equiv 0 \pmod{\mathbb{L}}$	$\mathbb{U}(G) = \mathbb{L} \mathbb{U}(G \setminus e) \equiv 0 \pmod{\mathbb{L}}$
cardona-qft-methods_ EQ0297 (1)	150	■ 1.000	$\mathbf{A}_{\mu}(x) \rightarrow \mathbf{A}'_{\mu}(x) = g^{-1}(x) [\mathbf{A}_{\mu}(x) + \partial_{\mu}] g(x),$	$\mathbf{A}_{\mu}(x) \rightarrow \mathbf{A}'_{\mu}(x) = g^{-1}(x) [\mathbf{A}_{\mu}(x) + \partial_{\mu}] g(x),$	$\mathbf{A}_{\mu}(x) \rightarrow \mathbf{A}'_{\mu}(x) = g^{-1}(x) [\mathbf{A}_{\mu}(x) + \partial_{\mu}] g(x),$
cardona-qft-methods_ EQ0298 (2)	150	■ 1.000	$\mathcal{C}(\mathbf{A}') = \mathcal{C}(\mathbf{A}) + d\Omega,$	$\mathcal{C}(\mathbf{A}') = \mathcal{C}(\mathbf{A}) + d\Omega,$	$\mathcal{C}(\mathbf{A}') = \mathcal{C}(\mathbf{A}) + d\Omega,$
cardona-qft-methods_ EQ0299 (3)	150	■ 1.000	$L(q, \dot{q}) dt \rightarrow L(q', \dot{q}') dt' = L(q, \dot{q}) dt + d\Omega(q, t).$	$L(q, \dot{q}) dt \rightarrow L(q', \dot{q}') dt' = L(q, \dot{q}) dt + d\Omega(q, t)$	$L(q, \dot{q}) dt \rightarrow L(q', \dot{q}') dt' = L(q, \dot{q}) dt + d\Omega(q, t).$
cardona-qft-methods_ EQ0300 (4)	150	■ 1.000	$\int \mathbf{A}_{\mu}(x) j^{\mu}(x) d^n x$	$\int \mathbf{A}_{\mu}(x) j^{\mu}(x) d^n x$	$\int \mathbf{A}_{\mu}(x) j^{\mu}(x) d^n x.$
cardona-qft-methods_ EQ0304 (8)	152	■ 1.000	$\mathbf{A}(x) \xrightarrow{g} \mathbf{A}'(x) = g^{-1}(x) \mathbf{A}(x) g(x) + g^{-1}(x) dg(x).$	$\mathbf{A}(x) \xrightarrow{g} \mathbf{A}'(x) = g^{-1}(x) \mathbf{A}(x) g(x) + g^{-1}(x) dg(x).$	$\mathbf{A}(x) \xrightarrow{g} \mathbf{A}'(x) = g^{-1}(x) \mathbf{A}(x) g(x) + g^{-1}(x) dg(x).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0305 (9)	152	■ 1.000	$\mathbf{F} \xrightarrow{\mathbf{g}} \mathbf{F}' = g \mathbf{F} g^{-1}$ $\mathbf{F}^{g^{-1}}$.	$\mathbf{F} \xrightarrow{g} \mathbf{F}' = g \mathbf{F} g^{-1}$.	$\mathbf{F} \xrightarrow{g} \mathbf{F}' = g \mathbf{F} g^{-1}$.
cardona-qft-methods_ EQ0306 (10)	152	■ 1.000	$\operatorname{Tr}(\mathbf{F}^k)$,	$\operatorname{Tr}(\mathbf{F}^k)$,	$\operatorname{Tr}(\mathbf{F}^k)$,
cardona-qft-methods_ EQ0307 (11)	154	■ 1.000	$\delta_{\text{gauge}} P_{2k} = d(\delta_{\text{gauge}} \mathcal{C}_{2k-1})$,	$\delta_{\text{gauge}} P_{2k} = d(\delta_{\text{gauge}} \mathcal{C}_{2k-1})$,	$\delta_{\text{gauge}} P_{2k} = d(\delta_{\text{gauge}} \mathcal{C}_{2k-1})$,
cardona-qft-methods_ EQ0309 (13)	154	■ 1.000	$\delta_{\text{gauge}} \mathcal{C}_{2k-1} = d \Omega$	$\delta_{\text{gauge}} \mathcal{C}_{2k-1} = d \Omega$	$\delta_{\text{gauge}} \mathcal{C}_{2k-1} = d \Omega$. ■
cardona-qft-methods_ EQ0310 (14)	154	■ 1.000	$I[A] = \int_{M^{2n-1}} \delta_{\text{gauge}} \mathcal{C}_{2n-1} = \int_{M^{2n-1}} d \Omega$	$\delta_{\text{gauge}} I[A] = \int_{M^{2n-1}} \delta_{\text{gauge}} \mathcal{C}_{2n-1} = \int_{M^{2n-1}} d \Omega$	$\delta_{\text{gauge}} I[A] = \int_{M^{2n-1}} \delta_{\text{gauge}} \mathcal{C}_{2n-1} = \int_{M^{2n-1}} d \Omega$,
cardona-qft-methods_ EQ0311 (15)	155	■ 1.000	$x^\mu \rightarrow x'^\mu = x^\mu + \xi^\mu(x)$.	$x^\mu \rightarrow x'^\mu = x^\mu + \xi^\mu(x)$.	$x^\mu \rightarrow x'^\mu = x^\mu + \xi^\mu(x)$.
cardona-qft-methods_ EQ0312 (16)	155	■ 1.000	$v^\mu(x) = L_\nu^\mu(x) v^\nu(x)$, where $L_\nu^\mu(x) = \frac{\partial x'^\mu}{\partial x^\nu} = \delta_\nu^\mu + \frac{\partial \xi^\mu}{\partial x^\nu}$.	$v'^\mu(x') = L_\nu^\mu(x) v^\nu(x)$, where $L_\nu^\mu(x) = \frac{\partial x'^\mu}{\partial x^\nu} = \delta_\nu^\mu + \frac{\partial \xi^\mu}{\partial x^\nu}$.	$v'^\mu(x') = L_\nu^\mu(x) v^\nu(x)$, where $L_\nu^\mu(x) = \frac{\partial x'^\mu}{\partial x^\nu} = \delta_\nu^\mu + \frac{\partial \xi^\mu}{\partial x^\nu}$.
cardona-qft-methods_ EQ0313 (17)	155	■ 1.000	$\begin{aligned} v^\mu(x) &= L_\nu^\mu(x) v^\nu(x) \\ v^\mu(x) &= L_\nu^\mu(x) v^\nu(x) + \frac{\partial \xi^\mu}{\partial x^\nu} v^\nu(x) - \xi^\lambda(x) \partial_\lambda v^\mu(x) \end{aligned}$	$\begin{aligned} v'^\mu(x) &= L_\nu^\mu(x) v^\nu(x) - \xi^\lambda(x) \partial_\lambda v^\mu(x) \\ &= v^\mu(x) + \frac{\partial \xi^\mu}{\partial x^\nu} v^\nu(x) - \xi^\lambda(x) \partial_\lambda v^\mu(x) \end{aligned}$	$\begin{aligned} v'^\mu(x) &= L_\nu^\mu(x) v^\nu(x) - \xi^\lambda(x) \partial_\lambda v^\mu(x) \\ &= v^\mu(x) + \frac{\partial \xi^\mu}{\partial x^\nu} v^\nu(x) - \xi^\lambda(x) \partial_\lambda v^\mu(x) \end{aligned}$
cardona-qft-methods_ EQ0317 (3)	159	■ 1.000	$g_{\mu\nu}(x) \equiv \eta_{ab} e_\mu^a(x) e_\nu^b(x)$,	$g_{\mu\nu}(x) \equiv \eta_{ab} e_\mu^a(x) e_\nu^b(x)$,	$g_{\mu\nu}(x) \equiv \eta_{ab} e_\mu^a(x) e_\nu^b(x)$,
cardona-qft-methods_ EQ0318 (4)	160	■ 1.000	$e_\mu^a(x) \longrightarrow e_\mu'^a(x) = \Lambda_b^a(x) e_\mu^b(x)$,	$e_\mu^a(x) \longrightarrow e_\mu'^a(x) = \Lambda_b^a(x) e_\mu^b(x)$,	$e_\mu^a(x) \longrightarrow e_\mu'^a(x) = \Lambda_b^a(x) e_\mu^b(x)$,
cardona-qft-methods_ EQ0319 (5)	160	■ 1.000	$\Lambda_c^a(x) \Lambda_d^b(x) \eta_{ab} = \eta_{cd}$,	$\Lambda_c^a(x) \Lambda_d^b(x) \eta_{ab} = \eta_{cd}$,	$\Lambda_c^a(x) \Lambda_d^b(x) \eta_{ab} = \eta_{cd}$,
cardona-qft-methods_ EQ0322 (8)	161	■ 1.000	$\omega_{b\mu}^a(x) \xrightarrow{\Lambda} \omega_{b\mu}'^a(x) = \Lambda_c^a(x) \Lambda_b^d(x) \omega_{d\mu}^c(x) + \Lambda_c^a(x) \partial_\mu \Lambda_b^c(x)$.	$\omega_{b\mu}^a(x) \xrightarrow{\Lambda} \omega_{b\mu}'^a(x) = \Lambda_c^a(x) \Lambda_b^d(x) \omega_{d\mu}^c(x) + \Lambda_c^a(x) \partial_\mu \Lambda_b^c(x)$.	$\omega_{b\mu}^a(x) \xrightarrow{\Lambda} \omega_{b\mu}'^a(x) = \Lambda_c^a(x) \Lambda_b^d(x) \omega_{d\mu}^c(x) + \Lambda_c^a(x) \partial_\mu \Lambda_b^c(x)$.
cardona-qft-methods_ EQ0327 (14)	162	■ 1.000	$d(\alpha_p) = d^2 \alpha_p = 0$.	$d(d\alpha_p) = d^2 \alpha_p = 0$.	$d(d\alpha_p) = d^2 \alpha_p = 0$.
cardona-qft-methods_ EQ0328 (15)	162	■ 1.000	$\begin{aligned} D^2 \phi^a &= D[d\phi^a + \omega_b^a \phi^b] \\ &= d[d\phi^a + \omega_b^a \phi^b] + \omega_b^a [d\phi^b + \omega_c^b \phi^c] \\ &= [d\omega_b^a + \omega_c^a \wedge \omega_b^c] \phi^b . \end{aligned}$	$\begin{aligned} D^2 \phi^a &= D[d\phi^a + \omega_b^a \phi^b] \\ &= d[d\phi^a + \omega_b^a \phi^b] + \omega_b^a [d\phi^b + \omega_c^b \phi^c] \\ &= [d\omega_b^a + \omega_c^a \wedge \omega_b^c] \phi^b . \end{aligned}$	$\begin{aligned} D^2 \phi^a &= D[d\phi^a + \omega_b^a \phi^b] \\ &= d[d\phi^a + \omega_b^a \phi^b] + \omega_b^a [d\phi^b + \omega_c^b \phi^c] \\ &= [d\omega_b^a + \omega_c^a \wedge \omega_b^c] \phi^b . \end{aligned}$
cardona-qft-methods_ EQ0331 (18)	163	■ 1.000	$T^a = de^a + \omega_b^a \wedge e^b$	$T^a = de^a + \omega_b^a \wedge e^b$	$T^a = de^a + \omega_b^a \wedge e^b$,
cardona-qft-methods_ EQ0340 (8)	166	■ 1.000	$\mathbf{P}_D = \mathbf{P}_{4k_1} \mathbf{P}_{4k_2} \cdots \mathbf{P}_{4k_r}$, $\cdots \mathbf{P}_{4k_r}$,	$\mathbf{P}_D = \mathbf{P}_{4k_1} \mathbf{P}_{4k_2} \cdots \mathbf{P}_{4k_r}$,	$\mathbf{P}_D = \mathbf{P}_{4k_1} \mathbf{P}_{4k_2} \cdots \mathbf{P}_{4k_r}$,

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0341 (9)	166	■ 1.000	$I_{\{D\}}[e, \omega] = \kappa \int_M \sum_{p=0}^{[D/2]} \alpha_p L_p^D$	$I_D[e, \omega] = \kappa \int_M \sum_{p=0}^{[D/2]} \alpha_p L_p^D$	$I_D[e, \omega] = \kappa \int_M \sum_{p=0}^{[D/2]} \alpha_p L_p^D$
cardona-qft-methods_ EQ0342 (10)	166	■ 1.000	$L_{\{p\}}^D = \epsilon_{a_1 \dots a_D} R^{a_1 a_2} \dots R^{a_{2p-1} a_{2p}} e^{a_{2p+1}} \dots e^{a_D}$	$L_p^D = \epsilon_{a_1 \dots a_D} R^{a_1 a_2} \dots R^{a_{2p-1} a_{2p}} e^{a_{2p+1}} \dots e^{a_D}$	$L_p^D = \epsilon_{a_1 \dots a_D} R^{a_1 a_2} \dots R^{a_{2p-1} a_{2p}} e^{a_{2p+1}} \dots e^{a_D}$. ■
cardona-qft-methods_ EQ0344 (12)	167	■ 1.000	$\begin{aligned} I_{\{4\}} &= \kappa \int_M \epsilon_{abcd} \left[\alpha_2 R^{ab} R^{cd} + \alpha_1 R^{ab} e^c e^d + \alpha_0 e^a e^b e^c e^d \right] \\ &= -\kappa \int_M \sqrt{ g } \left[\alpha_2 (R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta} - 4R^{\alpha\beta} R_{\alpha\beta} + R^2) + 2\alpha_1 R + 24\alpha_0 \right] d^4x \\ &= -\kappa \alpha_2 \cdot \mathbf{E}_4 - 2\alpha_1 \int_M \sqrt{ g } R d^4x - 24\kappa \alpha_0 \cdot V_4. \end{aligned}$	$\begin{aligned} I_4 &= \kappa \int_M \epsilon_{abcd} [\alpha_2 R^{ab} R^{cd} + \alpha_1 R^{ab} e^c e^d + \alpha_0 e^a e^b e^c e^d] \\ &= -\kappa \int_M \sqrt{ g } [\alpha_2 (R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta} - 4R^{\alpha\beta} R_{\alpha\beta} + R^2) + 2\alpha_1 R + 24\alpha_0] d^4x \\ &= -\kappa \alpha_2 \cdot \mathbf{E}_4 - 2\alpha_1 \int_M \sqrt{ g } R d^4x - 24\kappa \alpha_0 \cdot V_4 \end{aligned}$	$\begin{aligned} I_4 &= \kappa \int_M \epsilon_{abcd} [\alpha_2 R^{ab} R^{cd} + \alpha_1 R^{ab} e^c e^d + \alpha_0 e^a e^b e^c e^d] \\ &= -\kappa \int_M \sqrt{ g } [\alpha_2 (R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta} - 4R^{\alpha\beta} R_{\alpha\beta} + R^2) + 2\alpha_1 R + 24\alpha_0] d^4x \\ &= -\kappa \alpha_2 \cdot \mathbf{E}_4 - 2\alpha_1 \int_M \sqrt{ g } R d^4x - 24\kappa \alpha_0 \cdot V_4. \end{aligned} \quad (12)$
cardona-qft-methods_ EQ0345 (13)	167	■ 1.000	$\epsilon_{abcd} R^{ab} R^{cd} e^e = \sqrt{ g } [R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta} - 4R^{\alpha\beta} R_{\alpha\beta} + R^2] d^5x$	$\epsilon_{abcde} R^{ab} R^{cd} e^e = \sqrt{ g } [R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta} - 4R^{\alpha\beta} R_{\alpha\beta} + R^2] d^5x$	$\epsilon_{abcde} R^{ab} R^{cd} e^e = \sqrt{ g } [R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta} - 4R^{\alpha\beta} R_{\alpha\beta} + R^2] d^5x$
cardona-qft-methods_ EQ0346 (14)	168	■ 1.000	$\delta I_D = \int [\delta e^a \mathcal{E}_a + \delta \omega^{ab} \mathcal{E}_{ab}] = 0$	$\delta I_D = \int [\delta e^a \mathcal{E}_a + \delta \omega^{ab} \mathcal{E}_{ab}] = 0$	$\delta I_D = \int [\delta e^a \mathcal{E}_a + \delta \omega^{ab} \mathcal{E}_{ab}] = 0,$
cardona-qft-methods_ EQ0347 (15)	168	■ 1.000	$\mathcal{E}_a = \sum_{p=0}^{[\frac{D-1}{2}]} \alpha_p (D-2p) \mathcal{E}_a^{(p)} = 0$	$\mathcal{E}_a = \sum_{p=0}^{[\frac{D-1}{2}]} \alpha_p (D-2p) \mathcal{E}_a^{(p)} = 0$	$\mathcal{E}_a = \sum_{p=0}^{[\frac{D-1}{2}]} \alpha_p (D-2p) \mathcal{E}_a^{(p)} = 0,$
cardona-qft-methods_ EQ0348 (16)	168	■ 1.000	$\mathcal{E}_{ab} = \sum_{p=1}^{[\frac{D-1}{2}]} \alpha_p p (D-2p) \mathcal{E}_{ab}^{(p)} = 0$	$\mathcal{E}_{ab} = \sum_{p=1}^{[\frac{D-1}{2}]} \alpha_p p (D-2p) \mathcal{E}_{ab}^{(p)} = 0$	$\mathcal{E}_{ab} = \sum_{p=1}^{[\frac{D-1}{2}]} \alpha_p p (D-2p) \mathcal{E}_{ab}^{(p)} = 0,$
cardona-qft-methods_ EQ0350 (19)	168	■ 1.000	$T^a = de^a + \omega^a_b e^b = 0,$	$T^a = de^a + \omega^a_b e^b = 0,$	$T^a = de^a + \omega^a_b e^b = 0,$
cardona-qft-methods_ EQ0352	169	■ 1.000	$I_D[g] = \int_M d^D x \sqrt{g} [\alpha'_0 + \alpha'_1 R + \alpha'_2 (R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta} - 4R^{\alpha\beta} R_{\alpha\beta} + R^2) + \dots]$	$I_D[g] = \int_M d^D x \sqrt{g} [\alpha'_0 + \alpha'_1 R + \alpha'_2 (R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta} - 4R^{\alpha\beta} R_{\alpha\beta} + R^2) + \dots]$	$I_D[g] = \int_M d^D x \sqrt{g} [\alpha'_0 + \alpha'_1 R + \alpha'_2 (R^{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta} - 4R^{\alpha\beta} R_{\alpha\beta} + R^2) + \dots]$
cardona-qft-methods_ EQ0353 (21)	170	■ 1.000	$\zeta_0 = e^a T_a,$	$\zeta_0 = e^a T_a,$	$\zeta_0 = e^a T_a,$
cardona-qft-methods_ EQ0361 (1)	173	■ 1.000	$L_{\{3\}} = \epsilon_{abc} R^{ab} e^c$	$L_3 = \epsilon_{abc} R^{ab} e^c$	$L_3 = \epsilon_{abc} R^{ab} e^c$
cardona-qft-methods_ EQ0365 (4)	174	■ 1.000	$\delta \omega^{ab} = 0,$	$\delta \omega^{ab} = 0,$	$\delta \omega^{ab} = 0,$
cardona-qft-methods_ EQ0366 (5)	174	■ 1.000	$\delta L_3 = d[\epsilon_{abc} R^{ab} \lambda^c],$	$\delta L_3 = d[\epsilon_{abc} R^{ab} \lambda^c],$	$\delta L_3 = d[\epsilon_{abc} R^{ab} \lambda^c],$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0367 (6)	175	■ 1.000	$L_{\{3\}^{\{A\ d\ S\}}=\epsilon_{abc}\left(R^{\{a\ b\ c\}}\left(\frac{1}{3}\frac{1}{l^2}\right)e^{\{a\}}e^{\{b\}}e^{\{c\}}\right)$	$L_3^{AdS} = \epsilon_{abc} \left(R^{ab} e^c \pm \frac{1}{3l^2} e^a e^b e^c \right)$	$L_3^{AdS} = \epsilon_{abc} (R^{ab} e^c \pm \frac{1}{3l^2} e^a e^b e^c),$
cardona-qft-methods_ EQ0369 (9)	175	■ 1.000	$\begin{aligned} \delta \left[\begin{array}{cc} \omega^{ab} & e^a l^{-l} \\ -e^b l^{-l} & 0 \end{array} \right] &= d \left[\begin{array}{cc} \lambda^{ab} & \lambda^a l^{-l} \\ -\lambda^b l^{-l} & 0 \end{array} \right] \\ &+ \left[\begin{array}{cc} \omega_c^a & e^a l^{-l} \\ -e_c l^{-l} & 0 \end{array} \right] \left[\begin{array}{cc} \lambda^{cb} & \lambda^c l^{-1} \\ \pm \lambda^b l^{-1} & 0 \end{array} \right] \\ &- \left[\begin{array}{cc} \lambda^{ac} & \lambda^a l^{-1} \\ -\lambda^c l^{-1} & 0 \end{array} \right] \left[\begin{array}{cc} \omega_c^b & e_c l^{-1} \\ \pm e^b l^{-1} & 0 \end{array} \right]. \end{aligned}$	$\delta \left[\begin{array}{cc} \omega^{ab} & e^a l^{-l} \\ -e^b l^{-l} & 0 \end{array} \right] = d \left[\begin{array}{cc} \lambda^{ab} & \lambda^a l^{-l} \\ -\lambda^b l^{-l} & 0 \end{array} \right] \\ + \left[\begin{array}{cc} \omega_c^a & e^a l^{-l} \\ -e_c l^{-l} & 0 \end{array} \right] \left[\begin{array}{cc} \lambda^{cb} & \lambda^c l^{-1} \\ \pm \lambda^b l^{-1} & 0 \end{array} \right] \\ - \left[\begin{array}{cc} \lambda^{ac} & \lambda^a l^{-1} \\ -\lambda^c l^{-1} & 0 \end{array} \right] \left[\begin{array}{cc} \omega_c^b & e_c l^{-1} \\ \pm e^b l^{-1} & 0 \end{array} \right].$	$\delta \left[\begin{array}{cc} \omega^{ab} & e^a l^{-l} \\ -e^b l^{-l} & 0 \end{array} \right] = d \left[\begin{array}{cc} \lambda^{ab} & \lambda^a l^{-l} \\ -\lambda^b l^{-l} & 0 \end{array} \right] \\ + \left[\begin{array}{cc} \omega_c^a & e^a l^{-l} \\ -e_c l^{-l} & 0 \end{array} \right] \left[\begin{array}{cc} \lambda^{cb} & \lambda^c l^{-1} \\ \pm \lambda^b l^{-1} & 0 \end{array} \right] \\ - \left[\begin{array}{cc} \lambda^{ac} & \lambda^a l^{-1} \\ -\lambda^c l^{-1} & 0 \end{array} \right] \left[\begin{array}{cc} \omega_c^b & e_c l^{-1} \\ \pm e^b l^{-1} & 0 \end{array} \right].$
cardona-qft-methods_ EQ0370 (11)	175	■ 1.000	$\delta W^{AB} = d\Lambda^{AB} + W_C^A \Lambda^{CB} - \Lambda^{AC} W_C^B,$	$\delta W^{AB} = d\Lambda^{AB} + W_C^A \Lambda^{CB} - \Lambda^{AC} W_C^B,$	$\delta W^{AB} = d\Lambda^{AB} + W_C^A \Lambda^{CB} - \Lambda^{AC} W_C^B,$
cardona-qft-methods_ EQ0371 (10)	175	■ 1.000	$\begin{aligned} W^{AB} &= \left[\begin{array}{cc} \omega^{ab} & e^a l^{-1} \\ -e^b l^{-1} & 0 \end{array} \right] \\ \Lambda^{AB} &= \left[\begin{array}{cc} \lambda^{ab} & \lambda^a l^{-1} \\ -\lambda^b l^{-1} & 0 \end{array} \right], \end{aligned}$	$W^{AB} = \left[\begin{array}{cc} \omega^{ab} & e^a l^{-1} \\ -e^b l^{-1} & 0 \end{array} \right] \\ \Lambda^{AB} = \left[\begin{array}{cc} \lambda^{ab} & \lambda^a l^{-1} \\ -\lambda^b l^{-1} & 0 \end{array} \right],$	$W^{AB} = \left[\begin{array}{cc} \omega^{ab} & e^a l^{-1} \\ -e^b l^{-1} & 0 \end{array} \right] \\ \Lambda^{AB} = \left[\begin{array}{cc} \lambda^{ab} & \lambda^a l^{-1} \\ -\lambda^b l^{-1} & 0 \end{array} \right],$
cardona-qft-methods_ EQ0373 (13)	176	■ 1.000	$D_W \Upsilon^{AB} = d\Upsilon^{AB} + W_C^A \Upsilon^{CB} + W_C^B \Upsilon^{AC} = 0.$	$D_W \Upsilon^{AB} = d\Upsilon^{AB} + W_C^A \Upsilon^{CB} + W_C^B \Upsilon^{AC} = 0.$	$D_W \Upsilon^{AB} = d\Upsilon^{AB} + W_C^A \Upsilon^{CB} + W_C^B \Upsilon^{AC} = 0.$
cardona-qft-methods_ EQ0376 (17)	177	■ 1.000	$L_{\{3\}^{\{E\ H\}}}=\epsilon_{abc}R^{\{a\ b\ c\}}R^{\{a\ b\}}e^{\{c\}},$	$L_3^{EH} = \epsilon_{abc} R^{ab} e^c,$	$L_3^{EH} = \epsilon_{abc} R^{ab} e^c,$
cardona-qft-methods_ EQ0381 (22)	178	■ 1.000	$L_{\{2\ n+1\}}=\epsilon_{a_1\cdots a_{2\ n+1}}R^{\{a_1\}a_{\{2\}}\cdots R^{\{a_{2\ n-1}\}a_{\{2\ n\}}}e^{\{a_{\{2\ n+1\}}\}},$	$L_{2n+1} = \epsilon_{a_1\cdots a_{2n+1}} R^{a_1 a_2} \dots R^{a_{2n-1} a_{2n}} e^{a_{2n+1}},$	$L_{2n+1} = \epsilon_{a_1\cdots a_{2n+1}} R^{a_1 a_2} \dots R^{a_{2n-1} a_{2n}} e^{a_{2n+1}},$
cardona-qft-methods_ EQ0382	178	■ 1.000	$F^{\{A\ B\}}=d\ W^{\{A\ B\}}+W_{\{C\}}^{\{A\}}\ W^{\{C\ B\}},$	$F^{AB} = dW^{AB} + W_C^A W^{CB},$	$F^{AB} = dW^{AB} + W_C^A W^{CB},$
cardona-qft-methods_ EQ0383 (23)	178	■ 1.000	$F^{\{A\ B\}}=\left[\begin{array}{cc} R^{\{a\ b\}}\left(\frac{1}{3}\frac{1}{l^2}\right)e^{\{a\}}e^{\{b\}} & l^{-1}T^a \\ -l^{-1}T^b & 0 \end{array}\right].$	$F^{AB} = \left[\begin{array}{cc} R^{ab} \pm l^{-2} e^a e^b & l^{-1} T^a \\ -l^{-1} T^b & 0 \end{array} \right].$	$F^{AB} = \left[\begin{array}{cc} R^{ab} \pm l^{-2} e^a e^b & l^{-1} T^a \\ -l^{-1} T^b & 0 \end{array} \right].$
cardona-qft-methods_ EQ0384 (24)	178	■ 1.000	$\mathbf{E}_4=\epsilon_{ABCD}F^{\{A\ B\}}F^{\{C\ D\}}.$	$\mathbf{E}_4 = \epsilon_{ABCD} F^{AB} F^{CD}.$	$\mathbf{E}_4 = \epsilon_{ABCD} F^{AB} F^{CD}.$
cardona-qft-methods_ EQ0385 (25)	179	■ 1.000	$\begin{aligned} \mathbf{E}_4 &= 4\epsilon_{abc} (R^{ab} \pm l^{-2} e^a e^b) l^{-1} T^a \\ &= \frac{4}{l} d \left[\epsilon_{abc} \left(R^{ab} \pm \frac{1}{3l^2} e^a e^b \right) e^c \right] \end{aligned}$	$\mathbf{E}_4 = 4\epsilon_{abc} (R^{ab} \pm l^{-2} e^a e^b) l^{-1} T^a \\ = \frac{4}{l} d \left[\epsilon_{abc} \left(R^{ab} \pm \frac{1}{3l^2} e^a e^b \right) e^c \right]$	$\mathbf{E}_4 = 4\epsilon_{abc} (R^{ab} \pm l^{-2} e^a e^b) l^{-1} T^a \\ = \frac{4}{l} d \left[\epsilon_{abc} \left(R^{ab} \pm \frac{1}{3l^2} e^a e^b \right) e^c \right],$
cardona-qft-methods_ EQ0387	179	■ 1.000	$\delta \left(dL_3^{AdS} \right) = 0.$	$\delta \left(dL_3^{AdS} \right) = 0.$	$\delta \left(dL_3^{AdS} \right) = 0.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0390 (28)	180	■ 1.000	$L_{2n-1}^{\{A\}dS} = \sum_{p=0}^{n-1} \bar{\alpha}_p L_p^{2n-1}$	$L_{2n-1}^{\{A\}dS} = \sum_{p=0}^{n-1} \bar{\alpha}_p L_p^{2n-1}$	$L_{2n-1}^{\{A\}dS} = \sum_{p=0}^{n-1} \bar{\alpha}_p L_p^{2n-1},$
cardona-qft-methods_ EQ0391 (29)	180	■ 1.000	$\bar{\alpha}_p = \kappa \cdot \frac{(\pm 1)^{p+1} l^{2p-D}}{(D-2p)} \binom{n-1}{p}, p = 1, 2, \dots, n-1 = \frac{D-1}{2},$	$\bar{\alpha}_p = \kappa \cdot \frac{(\pm 1)^{p+1} l^{2p-D}}{(D-2p)} \binom{n-1}{p}, p = 1, 2, \dots, n-1 = \frac{D-1}{2},$	$\bar{\alpha}_p = \kappa \cdot \frac{(\pm 1)^{p+1} l^{2p-D}}{(D-2p)} \binom{n-1}{p}, p = 1, 2, \dots, n-1 = \frac{D-1}{2},$
cardona-qft-methods_ EQ0393 (31)	180	■ 1.000	$L_5^{\{A\}dS} = \frac{\kappa}{l} \epsilon_{abcde} \left[e^a R^{bc} R^{de} \pm \frac{2}{3l^2} e^a e^b e^c R^{de} + \frac{1}{5l^4} e^a e^b e^c e^d e^e \right].$	$L_5^{\{A\}dS} = \frac{\kappa}{l} \epsilon_{abcde} \left[e^a R^{bc} R^{de} \pm \frac{2}{3l^2} e^a e^b e^c R^{de} + \frac{1}{5l^4} e^a e^b e^c e^d e^e \right].$	$L_5^{\{A\}dS} = \frac{\kappa}{l} \epsilon_{abcde} \left[e^a R^{bc} R^{de} \pm \frac{2}{3l^2} e^a e^b e^c R^{de} + \frac{1}{5l^4} e^a e^b e^c e^d e^e \right].$
cardona-qft-methods_ EQ0394	180	■ 1.000	$\tilde{e}^a = l^{-1} e^a,$	$\tilde{e}^a = l^{-1} e^a,$	$\tilde{e}^a = l^{-1} e^a,$
cardona-qft-methods_ EQ0396	181	■ 1.000	$L_3^{\{A\}dS} = \epsilon_{abc} \left(R^{ab} \pm \frac{e^a e^b}{3l^2} \right) e^c$	$L_3^{\{A\}dS} = \epsilon_{abc} \left(R^{ab} \pm \frac{e^a e^b}{3l^2} \right) e^c$	$L_3^{\{A\}dS} = \epsilon_{abc} (R^{ab} \pm \frac{e^a e^b}{3l^2}) e^c$
cardona-qft-methods_ EQ0400 (33)	183	■ 1.000	$\kappa \left[\int_{\Omega} \mathbf{E}_{2n} - \int_{\Omega'} \mathbf{E}_{2n} \right] = 2n\pi\hbar.$	$\kappa \left[\int_{\Omega} \mathbf{E}_{2n} - \int_{\Omega'} \mathbf{E}_{2n} \right] = 2n\pi\hbar.$	$\kappa \left[\int_{\Omega} \mathbf{E}_{2n} - \int_{\Omega'} \mathbf{E}_{2n} \right] = 2n\pi\hbar.$
cardona-qft-methods_ EQ0401 (34)	183	■ 1.000	$\kappa \left[\int_{\Omega} \mathbf{E}_{2n} + \int_{\tilde{\Omega}'} \mathbf{E}_{2n} \right] = 2n\pi\hbar.$	$\kappa \left[\int_{\Omega} \mathbf{E}_{2n} + \int_{\tilde{\Omega}'} \mathbf{E}_{2n} \right] = 2n\pi\hbar.$	$\kappa \left[\int_{\Omega} \mathbf{E}_{2n} + \int_{\tilde{\Omega}'} \mathbf{E}_{2n} \right] = 2n\pi\hbar.$
cardona-qft-methods_ EQ0402	183	■ 1.000	$\kappa = n\hbar,$	$\kappa = n\hbar,$	$\kappa = n\hbar,$
cardona-qft-methods_ EQ0403 (35)	183	■ 1.000	$\epsilon_{ab_1 b_2 \dots b_{2n}} \left[R^{b_1 b_2} + \frac{1}{l^2} e^{b_1} e^{b_2} \right] \dots \left[R^{b_{2n-1} b_{2n}} + \frac{1}{l^2} e^{b_{2n-1}} e^{b_{2n}} \right] = 0$	$\epsilon_{ab_1 b_2 \dots b_{2n}} \left[R^{b_1 b_2} + \frac{1}{l^2} e^{b_1} e^{b_2} \right] \dots \left[R^{b_{2n-1} b_{2n}} + \frac{1}{l^2} e^{b_{2n-1}} e^{b_{2n}} \right] = 0$	$\epsilon_{ab_1 b_2 \dots b_{2n}} \left[R^{b_1 b_2} + \frac{1}{l^2} e^{b_1} e^{b_2} \right] \dots \left[R^{b_{2n-1} b_{2n}} + \frac{1}{l^2} e^{b_{2n-1}} e^{b_{2n}} \right] = 0$
cardona-qft-methods_ EQ0404 (36)	184	■ 1.000	$\epsilon_{abb_2 \dots b_{2n}} \left[R^{b_2 b_3} + \frac{1}{l^2} e^{b_2} e^{b_3} \right] \dots \left[R^{b_{2n-2} b_{2n-1}} + \frac{1}{l^2} e^{b_{2n-2}} e^{b_{2n-1}} \right] T^{b_{2n}} = 0$	$\epsilon_{abb_2 \dots b_{2n}} \left[R^{b_2 b_3} + \frac{1}{l^2} e^{b_2} e^{b_3} \right] \dots \left[R^{b_{2n-2} b_{2n-1}} + \frac{1}{l^2} e^{b_{2n-2}} e^{b_{2n-1}} \right] T^{b_{2n}} = 0$	$\epsilon_{abb_2 \dots b_{2n}} \left[R^{b_2 b_3} + \frac{1}{l^2} e^{b_2} e^{b_3} \right] \dots \left[R^{b_{2n-2} b_{2n-1}} + \frac{1}{l^2} e^{b_{2n-2}} e^{b_{2n-1}} \right] T^{b_{2n}} = 0$
cardona-qft-methods_ EQ0406	185	■ 1.000	$\text{Pfaff}[M] = \epsilon_{a_1 b_1 a_2 b_2 \dots a_n b_n} M^{a_1 b_1} \dots M^{a_n b_n} = \sqrt{\det[M]}.$	$\text{Pfaff}[M] = \epsilon_{a_1 b_1 a_2 b_2 \dots a_n b_n} M^{a_1 b_1} \dots M^{a_n b_n} = \sqrt{\det[M]}.$	$\text{Pfaff}[M] = \epsilon_{a_1 b_1 a_2 b_2 \dots a_n b_n} M^{a_1 b_1} \dots M^{a_n b_n} = \sqrt{\det[M]}.$
cardona-qft-methods_ EQ0407 (38)	186	■ 1.000	$\delta I[\phi] = \int_M \delta \phi^r \frac{\delta L}{\delta \phi^r} + \int_{\partial M} \Theta(\phi, \delta \phi).$	$\delta I[\phi] = \int_M \delta \phi^r \frac{\delta L}{\delta \phi^r} + \int_{\partial M} \Theta(\phi, \delta \phi).$	$\delta I[\phi] = \int_M \delta \phi^r \frac{\delta L}{\delta \phi^r} + \int_{\partial M} \Theta(\phi, \delta \phi).$
cardona-qft-methods_ EQ0410 (40)	187	■ 1.000	$I_{2n-1} = \int_M L_{2n-1}^{\{A\}dS} + \int_{\partial M} B_{2n}$	$I_{2n-1} = \int_M L_{2n-1}^{\{A\}dS} + \int_{\partial M} B_{2n}$	$I_{2n-1} = \int_M L_{2n-1}^{\{A\}dS} + \int_{\partial M} B_{2n},$
cardona-qft-methods_ EQ0416 (1)	190	■ 1.000	$I_{\text{Int}} = \int_{\Gamma} A_{\mu} j^{\mu} d^4 x$	$I_{\text{Int}} = \int_{\Gamma} A_{\mu} j^{\mu} d^4 x$	$I_{\text{Int}} = \int_{\Gamma} A_{\mu} j^{\mu} d^4 x,$
cardona-qft-methods_ EQ0417 (2)	191	■ 1.000	$I_{\text{Int}} = q \int_{\Gamma} A_{\mu}(z) dz^{\mu} = q \int_{\Gamma} A$	$I_{\text{Int}} = q \int_{\Gamma} A_{\mu}(z) dz^{\mu} = q \int_{\Gamma} A$	$I_{\text{Int}} = q \int_{\Gamma} A_{\mu}(z) dz^{\mu} = q \int_{\Gamma} A,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0418 (3)	192	■ 1.000	$I[\mathbf{A};\mathbf{j}]=\int\left\langle\mathbf{j}_{2p}\wedge\mathfrak{C}_{2p+1}(\mathbf{A})\right\rangle,$	$I[\mathbf{A};\mathbf{j}]=\int\langle\mathbf{j}_{2p}\wedge\mathfrak{C}_{2p+1}(\mathbf{A})\rangle,$	$I[\mathbf{A};\mathbf{j}]=\int\langle\mathbf{j}_{2p}\wedge\mathfrak{C}_{2p+1}(\mathbf{A})\rangle,$
cardona-qft-methods_ EQ0419 (4)	192	■ 1.000	$\mathbf{j}_{2p}=qj^{a_1a_2\cdots a_s}\mathbf{K}_{a_1}\mathbf{K}_{a_2}\cdots\mathbf{K}_{a_s}\delta(\Gamma)dx^{\alpha_1}\wedge dx^{\alpha_2}\wedge\cdots dx^{\alpha_{D-2p-1}},$	$\mathbf{j}_{2p}=qj^{a_1a_2\cdots a_s}\mathbf{K}_{a_1}\mathbf{K}_{a_2}\cdots\mathbf{K}_{a_s}\delta(\Gamma)dx^{\alpha_1}\wedge dx^{\alpha_2}\wedge\cdots dx^{\alpha_{D-2p-1}},$	$\mathbf{j}_{2p}=qj^{a_1a_2\cdots a_s}\mathbf{K}_{a_1}\mathbf{K}_{a_2}\cdots\mathbf{K}_{a_s}\delta(\Gamma)dx^{\alpha_1}\wedge dx^{\alpha_2}\wedge\cdots dx^{\alpha_{D-2p-1}},$
cardona-qft-methods_ EQ0420 (5)	192	■ 1.000	$D\mathbf{j}_{2p}=d\mathbf{j}_{2p}+[\mathbf{A},\mathbf{j}_{2p}]=0,$	$D\mathbf{j}_{2p}=d\mathbf{j}_{2p}+[\mathbf{A},\mathbf{j}_{2p}]=0,$	$D\mathbf{j}_{2p}=d\mathbf{j}_{2p}+[\mathbf{A},\mathbf{j}_{2p}]=0,$
cardona-qft-methods_ EQ0421 (6)	193	■ 1.000	$I[A]=\frac{1}{2}\int_M(\mathrm{Tr}[F\wedge *F]-\Theta(x)\mathrm{Tr}[F\wedge F]),$	$I[A]=\frac{1}{2}\int_M(\mathrm{Tr}[F\wedge *F]-\Theta(x)\mathrm{Tr}[F\wedge F]),$	$I[A]=\frac{1}{2}\int_M(\mathrm{Tr}[F\wedge *F]-\Theta(x)\mathrm{Tr}[F\wedge F]),$
cardona-qft-methods_ EQ0423	202	■ 1.000	$S_{EH}(g_{\mu\nu})=\frac{1}{16\pi G}\int_M R\sqrt{g}\,d^4x$	$S_{EH}(g_{\mu\nu})=\frac{1}{16\pi G}\int_MR\sqrt{g}d^4x$	$S_{EH}(g_{\mu\nu})=\frac{1}{16\pi G}\int_MR\sqrt{g}\,d^4x,$
cardona-qft-methods_ EQ0425	203	■ 1.000	$\int_M\left(\frac{1}{2} D_\mu\mathbf{H} ^2-\mu_0^2 \mathbf{H} ^2-\xi_0R \mathbf{H} ^2+\lambda_0 \mathbf{H} ^4\right)\sqrt{g}d^4x.$	$\int_M\left(\frac{1}{2} D_\mu\mathbf{H} ^2-\mu_0^2 \mathbf{H} ^2-\xi_0R \mathbf{H} ^2+\lambda_0 \mathbf{H} ^4\right)\sqrt{g}d^4x.$	$\int_M\left(\frac{1}{2} D_\mu\mathbf{H} ^2-\mu_0^2 \mathbf{H} ^2-\xi_0R \mathbf{H} ^2+\lambda_0 \mathbf{H} ^4\right)\sqrt{g}\,d^4x.$
cardona-qft-methods_ EQ0426	204	■ 1.000	$[\gamma,a]=0,\forall a\in\mathcal{A},\quad\text{and}\quad D\gamma=-\gamma D.$	$[\gamma,a]=0,\forall a\in\mathcal{A},\quad\text{and}\quad D\gamma=-\gamma D.$	$[\gamma,a]=0,\,\forall a\in\mathcal{A},\quad\text{and}\quad D\gamma=-\gamma D.$
cardona-qft-methods_ EQ0428	205	■ 1.000	$\left[[D,a],b^0\right]=0\quad\forall a,b\in\mathcal{A}.$	$\left[[D,a],b^0\right]=0\quad\forall a,b\in\mathcal{A}.$	$[[D,a],b^0]=0\quad\quad\forall a,b\in\mathcal{A}.$
cardona-qft-methods_ EQ0430	207	■ 1.000	$\mathcal{A}_{LR}=\mathbb{C}\oplus\mathbb{H}_L\oplus\mathbb{H}_R\oplus M_3(\mathbb{C}).$	$\mathcal{A}_{LR}=\mathbb{C}\oplus\mathbb{H}_L\oplus\mathbb{H}_R\oplus M_3(\mathbb{C}).$	$\mathcal{A}_{LR}=\mathbb{C}\oplus\mathbb{H}_L\oplus\mathbb{H}_R\oplus M_3(\mathbb{C}).$
cardona-qft-methods_ EQ0431	208	■ 1.000	$\mathrm{Ad}(u)\xi=u\xi u^*\quad\forall\xi\in\mathcal{M}.$	$\mathrm{Ad}(u)\xi=u\xi u^*\quad\forall\xi\in\mathcal{M}.$	$\mathrm{Ad}(u)\xi=u\xi u^*\quad\forall\xi\in\mathcal{M}.$
cardona-qft-methods_ EQ0433	208	■ 1.000	$\mathcal{M}^0=\{\bar{\xi};\xi\in\mathcal{M}\},\quad a\bar{\xi}b=\overline{b^*\xi a^*}$	$\mathcal{M}^0=\{\bar{\xi};\xi\in\mathcal{M}\},\quad a\bar{\xi}b=\overline{b^*\xi a^*}$	$\mathcal{M}^0=\{\bar{\xi}\,;\,\xi\in\mathcal{M}\},\quad a\bar{\xi}b=\overline{b^*\xi a^*}$
cardona-qft-methods_ EQ0440	210	■ 1.000	$\mathrm{SU}(\mathcal{A}_F)\sim\mathrm{U}(1)\times\mathrm{SU}(2)\times\mathrm{SU}(3),$	$\mathrm{SU}(\mathcal{A}_F)\sim\mathrm{U}(1)\times\mathrm{SU}(2)\times\mathrm{SU}(3),$	$\mathrm{SU}(\mathcal{A}_F)\sim\mathrm{U}(1)\times\mathrm{SU}(2)\times\mathrm{SU}(3),$
cardona-qft-methods_ EQ0442	211	■ 1.000	$\mathbb{C}_F\subset\mathcal{A}_F,\quad\mathbb{C}_F=\{(\lambda,\lambda,0),\lambda\in\mathbb{C}\}.$	$\mathbb{C}_F\subset\mathcal{A}_F,\quad\mathbb{C}_F=\{(\lambda,\lambda,0),\lambda\in\mathbb{C}\}.$	$\mathbb{C}_F\subset\mathcal{A}_F,\quad\mathbb{C}_F=\{(\lambda,\lambda,0),\lambda\in\mathbb{C}\}.$
cardona-qft-methods_ EQ0446	211	■ 1.000	$T:E_R=\uparrow_R\otimes\mathbf{1}^0\rightarrow J_F E_R.$	$T:E_R=\uparrow_R\otimes\mathbf{1}^0\rightarrow J_F E_R.$	$T:E_R=\uparrow_R\otimes\mathbf{1}^0\rightarrow J_F E_R.$
cardona-qft-methods_ EQ0447	211	■ 1.000	$Y_{(\downarrow 3)}^{\prime}=W_1Y_{(\downarrow 3)}W_3^*,\quad Y_{(\uparrow 3)}^{\prime}=W_2Y_{(\uparrow 3)}W_3^*$	$Y_{(\downarrow 3)}^{\prime}=W_1Y_{(\downarrow 3)}W_3^*,\quad Y_{(\uparrow 3)}^{\prime}=W_2Y_{(\uparrow 3)}W_3^*$	$Y_{(\downarrow 3)}^{\prime}=W_1Y_{(\downarrow 3)}W_3^*,\quad Y_{(\uparrow 3)}^{\prime}=W_2Y_{(\uparrow 3)}W_3^*$
cardona-qft-methods_ EQ0450	212	■ 1.000	$\begin{gathered} Y_{(\uparrow 3)}=\delta_{(\uparrow 3)}\quad Y_{(\downarrow 3)}=U_{CKM}\delta_{(\downarrow 3)}U_{CKM}^*\\ \quad Y_{(\downarrow 3)}=U_{\{C\,K\,M\}}\delta_{(\downarrow 3)}U_{\{C\,K\,M\}}^*\quad Y_{(\uparrow 1)}=U_{\{P\,M\,N\,S\}}\delta_{(\uparrow 1)}U_{\{P\,M\,N\,S\}}^*\\ \quad Y_{(\downarrow 1)}=\delta_{(\downarrow 1)}\end{gathered}$	$Y_{(\uparrow 3)}=\delta_{(\uparrow 3)}\quad Y_{(\downarrow 3)}=U_{CKM}\delta_{(\downarrow 3)}U_{CKM}^*\\ Y_{(\uparrow 1)}=U_{PMNS}^*\delta_{(\uparrow 1)}U_{PMNS}\quad Y_{(\downarrow 1)}=\delta_{(\downarrow 1)}$	$Y_{(\uparrow 3)}=\delta_{(\uparrow 3)}\quad Y_{(\downarrow 3)}=U_{CKM}\delta_{(\downarrow 3)}U_{CKM}^*\\ Y_{(\uparrow 1)}=U_{PMNS}^*\delta_{(\uparrow 1)}U_{PMNS}\quad Y_{(\downarrow 1)}=\delta_{(\downarrow 1)}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0451	212	■ 1.000	$U=\left(\begin{array}{ccc} c_1 & -s_1 & c_3 \\ s_1 & c_1 & c_2 \\ \delta & c_1 & c_2 \end{array}\right)$	$U = \begin{pmatrix} c_1 & -s_1 c_3 & -s_1 s_3 \\ s_1 c_2 & c_1 c_2 c_3 - s_2 s_3 e_\delta & c_1 c_2 s_3 + s_2 c_3 e_\delta \\ s_1 s_2 & c_1 s_2 c_3 + c_2 s_3 e_\delta & c_1 s_2 s_3 - c_2 c_3 e_\delta \end{pmatrix}$	$U = \begin{pmatrix} c_1 & -s_1 c_3 & -s_1 s_3 \\ s_1 c_2 & c_1 c_2 c_3 - s_2 s_3 e_\delta & c_1 c_2 s_3 + s_2 c_3 e_\delta \\ s_1 s_2 & c_1 s_2 c_3 + c_2 s_3 e_\delta & c_1 s_2 s_3 - c_2 c_3 e_\delta \end{pmatrix}$
cardona-qft-methods_ EQ0453	214	■ 1.000	$\begin{gathered} \mathcal{A} = C^\infty(M) \otimes \mathcal{A}_F \\ \mathcal{H} = L^2(M, S) \otimes \mathcal{H}_F \\ D = \not{D}_M \otimes 1 + \gamma_5 \otimes D_F \end{gathered}$	$\mathcal{A} = C^\infty(M) \otimes \mathcal{A}_F = C^\infty(M, \mathcal{A}_F)$ $\mathcal{H} = L^2(M, S) \otimes \mathcal{H}_F = L^2(M, S \otimes \mathcal{H}_F)$ $D = \not{D}_M \otimes 1 + \gamma_5 \otimes D_F$	$\mathcal{A} = C^\infty(M) \otimes \mathcal{A}_F = C^\infty(M, \mathcal{A}_F)$ $\mathcal{H} = L^2(M, S) \otimes \mathcal{H}_F = L^2(M, S \otimes \mathcal{H}_F)$ $D = \not{D}_M \otimes 1 + \gamma_5 \otimes D_F$
cardona-qft-methods_ EQ0455	215	■ 1.000	$\xi b = b^0 \xi, \quad \xi \in \mathcal{H}, \quad b \in \mathcal{A}$	$\xi b = b^0 \xi, \quad \xi \in \mathcal{H}, \quad b \in \mathcal{A}$	$\xi b = b^0 \xi, \quad \xi \in \mathcal{H}, \quad b \in \mathcal{A}$
cardona-qft-methods_ EQ0456	215	■ 1.000	$\xi \in \mathcal{H} \rightarrow \text{Ad}(u)\xi = u\xi u^* \quad \xi \in \mathcal{H}$	$\xi \in \mathcal{H} \rightarrow \text{Ad}(u)\xi = u\xi u^* \quad \xi \in \mathcal{H}$	$\xi \in \mathcal{H} \rightarrow \text{Ad}(u)\xi = u\xi u^* \quad \xi \in \mathcal{H}$
cardona-qft-methods_ EQ0457	215	■ 1.000	$D \rightarrow D_A = D + A + \varepsilon' J A J^{-1}$	$D \rightarrow D_A = D + A + \varepsilon' J A J^{-1}$	$D \rightarrow D_A = D + A + \varepsilon' J A J^{-1}$
cardona-qft-methods_ EQ0458	215	■ 1.000	$A = \sum a_j [D, b_j], \quad a_j, b_j \in \mathcal{A}.$	$A = \sum a_j [D, b_j], \quad a_j, b_j \in \mathcal{A}.$	$A = \sum a_j [D, b_j], \quad a_j, b_j \in \mathcal{A}.$
cardona-qft-methods_ EQ0459	216	■ 1.000	$\text{SU}(\mathcal{A}_F) = \{u \in \text{U}(\mathcal{A}_F) \mid \det(u) = 1\},$	$\text{SU}(\mathcal{A}_F) = \{u \in \text{U}(\mathcal{A}_F) \mid \det(u) = 1\},$	$\text{SU}(\mathcal{A}_F) = \{u \in \text{U}(\mathcal{A}_F) \mid \det(u) = 1\},$
cardona-qft-methods_ EQ0460	216	■ 1.000	$\text{SU}(\mathcal{A}_F) \sim \text{U}(1) \times \text{SU}(2) \times \text{SU}(3),$	$\text{SU}(\mathcal{A}_F) \sim \text{U}(1) \times \text{SU}(2) \times \text{SU}(3),$	$\text{SU}(\mathcal{A}_F) \sim \text{U}(1) \times \text{SU}(2) \times \text{SU}(3),$
cardona-qft-methods_ EQ0462	216	■ 1.000	$\left(\begin{array}{cccc} \frac{4}{3}\Lambda + V & 0 & 0 & 0 \\ 0 & -\frac{2}{3}\Lambda + V & 0 & 0 \\ 0 & 0 & Q_{11} + \frac{1}{3}\Lambda + V & Q_{12} \\ 0 & 0 & Q_{21} & Q_{22} + \frac{1}{3}\Lambda + V \end{array}\right)$	$\left(\begin{array}{cccc} \frac{4}{3}\Lambda + V & 0 & 0 & 0 \\ 0 & -\frac{2}{3}\Lambda + V & 0 & 0 \\ 0 & 0 & Q_{11} + \frac{1}{3}\Lambda + V & Q_{12} \\ 0 & 0 & Q_{21} & Q_{22} + \frac{1}{3}\Lambda + V \end{array}\right)$	$\left(\begin{array}{cccc} \frac{4}{3}\Lambda + V & 0 & 0 & 0 \\ 0 & -\frac{2}{3}\Lambda + V & 0 & 0 \\ 0 & 0 & Q_{11} + \frac{1}{3}\Lambda + V & Q_{12} \\ 0 & 0 & Q_{21} & Q_{22} + \frac{1}{3}\Lambda + V \end{array}\right)$
cardona-qft-methods_ EQ0463	217	■ 1.000	$\left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & -2\Lambda & 0 & 0 \\ 0 & 0 & Q_{11} - \Lambda & Q_{12} \\ 0 & 0 & Q_{21} & Q_{22} - \Lambda \end{array}\right)$	$\left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & -2\Lambda & 0 & 0 \\ 0 & 0 & Q_{11} - \Lambda & Q_{12} \\ 0 & 0 & Q_{21} & Q_{22} - \Lambda \end{array}\right)$	$\left(\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & -2\Lambda & 0 & 0 \\ 0 & 0 & Q_{11} - \Lambda & Q_{12} \\ 0 & 0 & Q_{21} & Q_{22} - \Lambda \end{array}\right)$
cardona-qft-methods_ EQ0464	217	■ 1.000	$\sum_i a_i [\gamma_5 \otimes D_F, a'_i](x) \Big _{\mathcal{H}_f} = \gamma_5 \otimes (A_q^{(0,1)} + A_\ell^{(0,1)})$	$\sum_i a_i [\gamma_5 \otimes D_F, a'_i](x) \Big _{\mathcal{H}_f} = \gamma_5 \otimes (A_q^{(0,1)} + A_\ell^{(0,1)})$	$\sum_i a_i [\gamma_5 \otimes D_F, a'_i](x) \Big _{\mathcal{H}_f} = \gamma_5 \otimes (A_q^{(0,1)} + A_\ell^{(0,1)})$
cardona-qft-methods_ EQ0466	217	■ 1.000	$q = \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}.$	$q = \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}.$	$q = \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0468	217	■ 1.000	$-E=\frac{1}{4}\,s\,\otimes\mathrm{id}+\sum_{\mu<\nu}\gamma^{\mu}\gamma^{\nu}\otimes\mathbb{F}_{\mu\nu}-i\gamma_5\gamma^{\mu}\otimes\mathbb{M}(D_{\mu}\varphi)+1_4\otimes(D^{0,1})^2,$	$-E=\frac{1}{4}s\otimes\mathrm{id}+\sum_{\mu<\nu}\gamma^{\mu}\gamma^{\nu}\otimes\mathbb{F}_{\mu\nu}-i\gamma_5\gamma^{\mu}\otimes\mathbb{M}(D_{\mu}\varphi)+1_4\otimes(D^{0,1})^2$	$-E=\frac{1}{4}\,s\otimes\mathrm{id}+\sum_{\mu<\nu}\gamma^{\mu}\gamma^{\nu}\otimes\mathbb{F}_{\mu\nu}-i\gamma_5\gamma^{\mu}\otimes\mathbb{M}(D_{\mu}\varphi)+1_4\otimes(D^{0,1})^2,$
cardona-qft-methods_ EQ0469	218	■ 1.000	$D_{\mu}\varphi=\partial_{\mu}\varphi+\frac{i}{2}g_2W_{\mu}^{\alpha}\varphi\sigma^{\alpha}-\frac{i}{2}g_1B_{\mu}\varphi$	$D_{\mu}\varphi=\partial_{\mu}\varphi+\frac{i}{2}g_2W_{\mu}^{\alpha}\varphi\sigma^{\alpha}-\frac{i}{2}g_1B_{\mu}\varphi$	$D_{\mu}\varphi=\partial_{\mu}\varphi+\frac{i}{2}g_2W_{\mu}^{\alpha}\varphi\sigma^{\alpha}-\frac{i}{2}g_1B_{\mu}\varphi$
cardona-qft-methods_ EQ0470	218	■ 1.000	$\operatorname{Tr}(f(D/\Lambda)),$	$\operatorname{Tr}(f(D/\Lambda)),$	$\operatorname{Tr}(f(D/\Lambda)),$
cardona-qft-methods_ EQ0472	218	■ 1.000	$\operatorname{Tr}\left(e^{-t\,\Delta}\right)\sim\sum a_{\alpha}\,t^{\alpha}\quad(t\rightarrow0)$	$\operatorname{Tr}\left(e^{-t\Delta}\right)\sim\sum a_{\alpha}t^{\alpha}\quad(t\rightarrow0)$	$\operatorname{Tr}(e^{-t\Delta})\sim\sum a_{\alpha}t^{\alpha}\quad(t\rightarrow0)$
cardona-qft-methods_ EQ0473	218	■ 1.000	$\zeta_D(s)=\operatorname{Tr}\left(\Delta^{-s/2}\right).$	$\zeta_D(s)=\operatorname{Tr}\left(\Delta^{-s/2}\right).$	$\zeta_D(s)=\operatorname{Tr}(\Delta^{-s/2}).$
cardona-qft-methods_ EQ0474	218	■ 1.000	$\operatorname{Res}_{s=-2\alpha}\zeta_D(s)=\frac{2a_{\alpha}}{\Gamma(-\alpha)}.$	$\operatorname{Res}_{s=-2\alpha}\zeta_D(s)=\frac{2a_{\alpha}}{\Gamma(-\alpha)}.$	$\operatorname{Res}_{s=-2\alpha}\zeta_D(s)=\frac{2a_{\alpha}}{\Gamma(-\alpha)}.$
cardona-qft-methods_ EQ0475	218	■ 1.000	$ D ^{-s}=\Delta^{-s/2}=\frac{1}{\Gamma\left(\frac{s}{2}\right)}\int_0^{\infty}e^{-t\Delta}t^{s/2-1}dt$	$ D ^{-s}=\Delta^{-s/2}=\frac{1}{\Gamma\left(\frac{s}{2}\right)}\int_0^{\infty}e^{-t\Delta}t^{s/2-1}dt$	$ D ^{-s}=\Delta^{-s/2}=\frac{1}{\Gamma\left(\frac{s}{2}\right)}\int_0^{\infty}e^{-t\Delta}t^{s/2-1}dt$
cardona-qft-methods_ EQ0476	219	■ 1.000	$\frac{1}{\Gamma\left(\frac{s}{2}\right)}\sim\frac{s}{2}\quad\text{as}\quad s\rightarrow0$	$\frac{1}{\Gamma\left(\frac{s}{2}\right)}\sim\frac{s}{2}\quad\text{as}\quad s\rightarrow0$	$\frac{1}{\Gamma\left(\frac{s}{2}\right)}\sim\frac{s}{2}\quad\text{as}\quad s\rightarrow0.$
cardona-qft-methods_ EQ0477	219	■ 1.000	$\int_0^{\infty}\operatorname{Tr}\left(e^{-t\Delta}\right)t^{s/2-1}dt$	$\int_0^{\infty}\operatorname{Tr}\left(e^{-t\Delta}\right)t^{s/2-1}dt$	$\int_0^{\infty}\operatorname{Tr}(e^{-t\Delta})t^{s/2-1}dt$
cardona-qft-methods_ EQ0479	219	■ 1.000	$\frac{1}{2}\left\langle J\tilde{\xi},D_A\tilde{\xi}\right\rangle$	$\frac{1}{2}\left\langle J\tilde{\xi},D_A\tilde{\xi}\right\rangle$	$\frac{1}{2}\langle J\tilde{\xi},D_A\tilde{\xi}\rangle$
cardona-qft-methods_ EQ0480	219	■ 1.000	$\mathcal{H}_{cl}^+=\left\{\xi\in\mathcal{H}_{cl}\mid\gamma\xi=\xi\right\}$	$\mathcal{H}_{cl}^+=\{\xi\in\mathcal{H}_{cl}\mid\gamma\xi=\xi\}$	$\mathcal{H}_{cl}^+=\{\xi\in\mathcal{H}_{cl}\mid\gamma\xi=\xi\}$
cardona-qft-methods_ EQ0482	220	■ 1.000	$\frac{1}{2}\left\langle J\tilde{\xi},D_A\tilde{\xi}\right\rangle$	$\frac{1}{2}\left\langle J\tilde{\xi},D_A\tilde{\xi}\right\rangle$	$\frac{1}{2}\langle J\tilde{\xi},D_A\tilde{\xi}\rangle$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0483	220	■ 1.000	$\begin{aligned} S &= \frac{1}{\pi^2} \left(48 f_4 \Lambda^4 - f_2 \Lambda^2 \mathfrak{c} + \frac{f_0}{4} \mathfrak{d} \right) \int \sqrt{g} d^4 x \\ &+ \frac{96 f_2 \Lambda^2 - f_0 \mathfrak{c}}{24 \pi^2} \int R \sqrt{g} d^4 x \\ &+ \frac{f_0}{10 \pi^2} \int \left(\frac{11}{6} R^* R^* - 3 C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} \right) \sqrt{g} d^4 x \\ &+ \frac{(-2 \mathfrak{a} f_2 \Lambda^2 + \mathfrak{c} f_0)}{\pi^2} \int \varphi ^2 \sqrt{g} d^4 x \\ &+ \frac{f_0 \mathfrak{a}}{2 \pi^2} \int D_\mu \varphi ^2 \sqrt{g} d^4 x \\ &- \frac{f_0 \mathfrak{a}}{12 \pi^2} \int R \varphi ^2 \sqrt{g} d^4 x \\ &+ \frac{f_0 \mathfrak{b}}{2 \pi^2} \int \varphi ^4 \sqrt{g} d^4 x \\ &+ \frac{f_0}{2 \pi^2} \int \left(g_3^2 G_{\mu\nu}^i G^{\mu\nu i} + g_2^2 F_{\mu\nu}^\alpha F^{\mu\nu\alpha} + \frac{5}{3} g_1^2 B_{\mu\nu} B^{\mu\nu} \right) \sqrt{g} d^4 x, \end{aligned}$	$S = \frac{1}{\pi^2} \left(48 f_4 \Lambda^4 - f_2 \Lambda^2 \mathfrak{c} + \frac{f_0}{4} \mathfrak{d} \right) \int \sqrt{g} d^4 x$ $+ \frac{96 f_2 \Lambda^2 - f_0 \mathfrak{c}}{24 \pi^2} \int R \sqrt{g} d^4 x$ $+ \frac{f_0}{10 \pi^2} \int \left(\frac{11}{6} R^* R^* - 3 C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} \right) \sqrt{g} d^4 x$ $+ \frac{(-2 \mathfrak{a} f_2 \Lambda^2 + \mathfrak{c} f_0)}{\pi^2} \int \varphi ^2 \sqrt{g} d^4 x$ $+ \frac{f_0 \mathfrak{a}}{2 \pi^2} \int D_\mu \varphi ^2 \sqrt{g} d^4 x$ $- \frac{f_0 \mathfrak{a}}{12 \pi^2} \int R \varphi ^2 \sqrt{g} d^4 x$ $+ \frac{f_0 \mathfrak{b}}{2 \pi^2} \int \varphi ^4 \sqrt{g} d^4 x$ $+ \frac{f_0}{2 \pi^2} \int \left(g_3^2 G_{\mu\nu}^i G^{\mu\nu i} + g_2^2 F_{\mu\nu}^\alpha F^{\mu\nu\alpha} + \frac{5}{3} g_1^2 B_{\mu\nu} B^{\mu\nu} \right) \sqrt{g} d^4 x,$	$S = \frac{1}{\pi^2} (48 f_4 \Lambda^4 - f_2 \Lambda^2 \mathfrak{c} + \frac{f_0}{4} \mathfrak{d}) \int \sqrt{g} d^4 x$ $+ \frac{96 f_2 \Lambda^2 - f_0 \mathfrak{c}}{24 \pi^2} \int R \sqrt{g} d^4 x$ $+ \frac{f_0}{10 \pi^2} \int \left(\frac{11}{6} R^* R^* - 3 C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} \right) \sqrt{g} d^4 x$ $+ \frac{(-2 \mathfrak{a} f_2 \Lambda^2 + \mathfrak{c} f_0)}{\pi^2} \int \varphi ^2 \sqrt{g} d^4 x$ $+ \frac{f_0 \mathfrak{a}}{2 \pi^2} \int D_\mu \varphi ^2 \sqrt{g} d^4 x$ $- \frac{f_0 \mathfrak{a}}{12 \pi^2} \int R \varphi ^2 \sqrt{g} d^4 x$ $+ \frac{f_0 \mathfrak{b}}{2 \pi^2} \int \varphi ^4 \sqrt{g} d^4 x$ $+ \frac{f_0}{2 \pi^2} \int (g_3^2 G_{\mu\nu}^i G^{\mu\nu i} + g_2^2 F_{\mu\nu}^\alpha F^{\mu\nu\alpha} + \frac{5}{3} g_1^2 B_{\mu\nu} B^{\mu\nu}) \sqrt{g} d^4 x,$
cardona-qft-methods_ EQ0484	220	■ 1.000	$f_{\{k\}} = \int_0^\infty f(v) v^{k-1} dv$	$f_k = \int_0^\infty f(v) v^{k-1} dv$	$f_k = \int_0^\infty f(v) v^{k-1} dv.$
cardona-qft-methods_ EQ0486	221	■ 1.000	$\operatorname{Tr} e^{-tP} \sim \sum_{n \geq 0} t^{\frac{n-m}{2}} \int_M a_n(x, P) dv(x)$	$\operatorname{Tr} e^{-tP} \sim \sum_{n \geq 0} t^{\frac{n-m}{2}} \int_M a_n(x, P) dv(x)$	$\operatorname{Tr} e^{-tP} \sim \sum_{n \geq 0} t^{\frac{n-m}{2}} \int_M a_n(x, P) dv(x)$
cardona-qft-methods_ EQ0487	221	■ 1.000	$\begin{gathered} \nabla_{\mu} = \partial_{\mu} + \omega'_{\mu}, \quad \omega'_{\mu} = \frac{1}{2} g_{\mu\nu} (A^{\nu} + \Gamma^{\nu} \cdot \text{id}) \\ E = B - g^{\mu\nu} (\partial_{\mu} \omega'_{\nu} + \omega'_{\mu} \omega'_{\nu} - \Gamma_{\mu\nu}^{\rho} \omega'_{\rho}) \\ \Omega_{\mu\nu} = \partial_{\mu} \omega'_{\nu} - \partial_{\nu} \omega'_{\mu} + [\omega'_{\mu}, \omega'_{\nu}] \end{gathered}$	$\nabla_{\mu} = \partial_{\mu} + \omega'_{\mu}, \quad \omega'_{\mu} = \frac{1}{2} g_{\mu\nu} (A^{\nu} + \Gamma^{\nu} \cdot \text{id})$ $E = B - g^{\mu\nu} (\partial_{\mu} \omega'_{\nu} + \omega'_{\mu} \omega'_{\nu} - \Gamma_{\mu\nu}^{\rho} \omega'_{\rho})$ $\Omega_{\mu\nu} = \partial_{\mu} \omega'_{\nu} - \partial_{\nu} \omega'_{\mu} + [\omega'_{\mu}, \omega'_{\nu}]$	$\nabla_{\mu} = \partial_{\mu} + \omega'_{\mu}, \quad \omega'_{\mu} = \frac{1}{2} g_{\mu\nu} (A^{\nu} + \Gamma^{\nu} \cdot \text{id})$ $E = B - g^{\mu\nu} (\partial_{\mu} \omega'_{\nu} + \omega'_{\mu} \omega'_{\nu} - \Gamma_{\mu\nu}^{\rho} \omega'_{\rho})$ $\Omega_{\mu\nu} = \partial_{\mu} \omega'_{\nu} - \partial_{\nu} \omega'_{\mu} + [\omega'_{\mu}, \omega'_{\nu}].$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E00491	222	■ 1.000	$\begin{aligned} S &= \frac{1}{2} \kappa_0^2 \int R \sqrt{g} d^4x + \gamma_0 \int \sqrt{g} d^4x \\ &+ \alpha_0 \int C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} \sqrt{g} d^4x + \tau_0 \int R^* R^* \sqrt{g} d^4x \\ &+ \frac{1}{2} \int DH ^2 \sqrt{g} d^4x - \mu_0^2 \int H ^2 \sqrt{g} d^4x \\ &- \xi_0 \int R H ^2 \sqrt{g} d^4x + \lambda_0 \int H ^4 \sqrt{g} d^4x \\ &+ \frac{1}{4} \int (G_{\mu\nu}^i G^{\mu\nu i} + F_{\mu\nu}^\alpha F^{\mu\nu\alpha} + B_{\mu\nu} B^{\mu\nu}) \sqrt{g} d^4x \end{aligned}$	$S = \frac{1}{2\kappa_0^2} \int R \sqrt{g} d^4x + \gamma_0 \int \sqrt{g} d^4x \\ + \alpha_0 \int C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} \sqrt{g} d^4x + \tau_0 \int R^* R^* \sqrt{g} d^4x \\ + \frac{1}{2} \int DH ^2 \sqrt{g} d^4x - \mu_0^2 \int H ^2 \sqrt{g} d^4x \\ - \xi_0 \int R H ^2 \sqrt{g} d^4x + \lambda_0 \int H ^4 \sqrt{g} d^4x \\ + \frac{1}{4} \int (G_{\mu\nu}^i G^{\mu\nu i} + F_{\mu\nu}^\alpha F^{\mu\nu\alpha} + B_{\mu\nu} B^{\mu\nu}) \sqrt{g} d^4x$	$S = \frac{1}{2\kappa_0^2} \int R \sqrt{g} d^4x + \gamma_0 \int \sqrt{g} d^4x \\ + \alpha_0 \int C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} \sqrt{g} d^4x + \tau_0 \int R^* R^* \sqrt{g} d^4x \\ + \frac{1}{2} \int DH ^2 \sqrt{g} d^4x - \mu_0^2 \int H ^2 \sqrt{g} d^4x \\ - \xi_0 \int R H ^2 \sqrt{g} d^4x + \lambda_0 \int H ^4 \sqrt{g} d^4x \\ + \frac{1}{4} \int (G_{\mu\nu}^i G^{\mu\nu i} + F_{\mu\nu}^\alpha F^{\mu\nu\alpha} + B_{\mu\nu} B^{\mu\nu}) \sqrt{g} d^4x$
cardona-qft-methods_ E00493	222	■ 1.000	$\frac{g^2 f_0}{2\pi^2} = \frac{1}{4}.$	$\frac{g^2 f_0}{2\pi^2} = \frac{1}{4}.$	$\frac{g^2 f_0}{2\pi^2} = \frac{1}{4}.$
cardona-qft-methods_ E00495	225	■ 1.000	$\partial_t x_i(t) = \beta_{x_i}(x(t)),$	$\partial_t x_i(t) = \beta_{x_i}(x(t)),$	$\partial_t x_i(t) = \beta_{x_i}(x(t)),$
cardona-qft-methods_ E00496 (1)	225	■ 1.000	$\beta_1 = \frac{41}{96\pi^2} g_1^3, \quad \beta_2 = -\frac{19}{96\pi^2} g_2^3, \quad \beta_3 = -\frac{7}{16\pi^2} g_3^3.$	$\beta_1 = \frac{41}{96\pi^2} g_1^3, \quad \beta_2 = -\frac{19}{96\pi^2} g_2^3, \quad \beta_3 = -\frac{7}{16\pi^2} g_3^3.$	$\beta_1 = \frac{41}{96\pi^2} g_1^3, \quad \beta_2 = -\frac{19}{96\pi^2} g_2^3, \quad \beta_3 = -\frac{7}{16\pi^2} g_3^3.$
cardona-qft-methods_ E00497 (2)	225	■ 1.000	$g_1(0) = 0.3575, \quad g_2(0) = 0.6514, \quad g_3(0) = 1.221$	$g_1(0) = 0.3575, \quad g_2(0) = 0.6514, \quad g_3(0) = 1.221$	$g_1(0) = 0.3575, \quad g_2(0) = 0.6514, \quad g_3(0) = 1.221$
cardona-qft-methods_ E00499	226	■ 1.000	$\frac{\sqrt{a} f_0}{\pi} = \frac{2M_W}{g}$	$\frac{\sqrt{a} f_0}{\pi} = \frac{2M_W}{g}$	$\frac{\sqrt{a} f_0}{\pi} = \frac{2M_W}{g}$
cardona-qft-methods_ E00503 (4)	227	■ 1.000	$\delta_{(\uparrow 1)} = \frac{1}{246} \begin{pmatrix} 12.2 \times 10^{-9} & 0 & 0 \\ 0 & 170 \times 10^{-6} & 0 \\ 0 & 0 & 15.5 \times 10^{-3} \end{pmatrix}$ $Y_\nu = U_{PMNS}^\dagger \delta_{(\uparrow 1)} U_{PMNS}$	$\delta_{(\uparrow 1)} = \frac{1}{246} \begin{pmatrix} 12.2 \times 10^{-9} & 0 & 0 \\ 0 & 170 \times 10^{-6} & 0 \\ 0 & 0 & 15.5 \times 10^{-3} \end{pmatrix}$ $Y_\nu = U_{PMNS}^\dagger \delta_{(\uparrow 1)} U_{PMNS}$	$\delta_{(\uparrow 1)} = \frac{1}{246} \begin{pmatrix} 12.2 \times 10^{-9} & 0 & 0 \\ 0 & 170 \times 10^{-6} & 0 \\ 0 & 0 & 15.5 \times 10^{-3} \end{pmatrix}$ $Y_\nu = U_{PMNS}^\dagger \delta_{(\uparrow 1)} U_{PMNS}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0504 (3)	227	■ 1.000	$\begin{gathered} \beta_y=\frac{1}{16\pi^2}\left(\frac{9}{2}y^3-8g_3^2y-\frac{9}{4}g_2^2y-\frac{17}{12}g_1^2y\right)\\ \beta_\lambda=\frac{1}{16\pi^2}\left(24\lambda^2+12\lambda y^2-9\lambda\left(g_2^2+\frac{1}{3}g_1^2\right)-6y^4+\frac{9}{8}g_2^4+\frac{3}{8}g_1^4+\frac{3}{4}g_2^2g_1^2\right), \end{gathered}$	$\beta_y=\frac{1}{16\pi^2}\left(\frac{9}{2}y^3-8g_3^2y-\frac{9}{4}g_2^2y-\frac{17}{12}g_1^2y\right).$ $\beta_\lambda=\frac{1}{16\pi^2}\left(24\lambda^2+12\lambda y^2-9\lambda\left(g_2^2+\frac{1}{3}g_1^2\right)-6y^4+\frac{9}{8}g_2^4+\frac{3}{8}g_1^4+\frac{3}{4}g_2^2g_1^2\right),$	$\beta_y=\frac{1}{16\pi^2}\left(\frac{9}{2}y^3-8g_3^2y-\frac{9}{4}g_2^2y-\frac{17}{12}g_1^2y\right). \quad (3)$ $\beta_\lambda=\frac{1}{16\pi^2}\left(24\lambda^2+12\lambda y^2-9\lambda(g_2^2+\frac{1}{3}g_1^2)-6y^4+\frac{9}{8}g_2^4+\frac{3}{8}g_1^4+\frac{3}{4}g_2^2g_1^2\right),$
cardona-qft-methods_ EQ0505	228	■ 1.000	$\frac{f_0}{2\pi^2}\int b \varphi ^4\sqrt{g}d^4x=\frac{\pi^2}{2f_0}\frac{\mathfrak{b}}{\mathfrak{a}^2}\int \mathbf{H} ^4\sqrt{g}d^4x$	$\frac{f_0}{2\pi^2}\int b \varphi ^4\sqrt{g}d^4x=\frac{\pi^2}{2f_0}\frac{\mathfrak{b}}{\mathfrak{a}^2}\int \mathbf{H} ^4\sqrt{g}d^4x$	$\frac{f_0}{2\pi^2}\int b \varphi ^4\sqrt{g}d^4x=\frac{\pi^2}{2f_0}\frac{\mathfrak{b}}{\mathfrak{a}^2}\int \mathbf{H} ^4\sqrt{g}d^4x$
cardona-qft-methods_ EQ0506	228	■ 1.000	$\tilde{\lambda}(\Lambda)=g_3^2\frac{\mathfrak{b}}{\mathfrak{a}^2}.$	$\tilde{\lambda}(\Lambda)=g_3^2\frac{\mathfrak{b}}{\mathfrak{a}^2}.$	$\tilde{\lambda}(\Lambda)=g_3^2\frac{\mathfrak{b}}{\mathfrak{a}^2}.$
cardona-qft-methods_ EQ0507	228	■ 1.000	$\frac{d\lambda}{dt}=\lambda\gamma+\frac{1}{8\pi^2}(12\lambda^2+B)$	$\frac{d\lambda}{dt}=\lambda\gamma+\frac{1}{8\pi^2}(12\lambda^2+B)$	$\frac{d\lambda}{dt}=\lambda\gamma+\frac{1}{8\pi^2}(12\lambda^2+B)$
cardona-qft-methods_ EQ0508	228	■ 1.000	$\gamma=\frac{1}{16\pi^2}(12y_t^2-9g_2^2-3g_1^2) \quad B=\frac{3}{16}(3g_2^4+2g_1^2g_2^2+g_1^4)-3y_t^4$	$\gamma=\frac{1}{16\pi^2}(12y_t^2-9g_2^2-3g_1^2) \quad B=\frac{3}{16}(3g_2^4+2g_1^2g_2^2+g_1^4)-3y_t^4$	$\gamma=\frac{1}{16\pi^2}(12y_t^2-9g_2^2-3g_1^2) \quad B=\frac{3}{16}(3g_2^4+2g_1^2g_2^2+g_1^4)-3y_t^4$
cardona-qft-methods_ EQ0509	228	■ 1.000	$m_H^2=8\lambda\frac{M^2}{g^2}, \quad m_H=\sqrt{2\lambda}\frac{2M}{g}.$	$m_H^2=8\lambda\frac{M^2}{g^2}, \quad m_H=\sqrt{2\lambda}\frac{2M}{g}.$	$m_H^2=8\lambda\frac{M^2}{g^2}, \quad m_H=\sqrt{2\lambda}\frac{2M}{g}.$
cardona-qft-methods_ EQ0510	229	■ 1.000	$\int\left(\frac{1}{2\eta}C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma}-\frac{\omega}{3\eta}R^2+\frac{\theta}{\eta}R^*R^*\right)\sqrt{g}d^4x$	$\int\left(\frac{1}{2\eta}C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma}-\frac{\omega}{3\eta}R^2+\frac{\theta}{\eta}R^*R^*\right)\sqrt{g}d^4x$	$\int\left(\frac{1}{2\eta}C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma}-\frac{\omega}{3\eta}R^2+\frac{\theta}{\eta}R^*R^*\right)\sqrt{g}d^4x$
cardona-qft-methods_ EQ0511	229	■ 1.000	$\begin{aligned} &\& \beta_{\eta}=-\frac{1}{(4\pi)^2}\frac{133}{10}\eta^2 \\ &\& \beta_{\omega}=-\frac{1}{(4\pi)^2}\frac{25+1098\omega+200\omega^2}{60}\eta \\ &\& \beta_{\theta}=\frac{1}{(4\pi)^2}\frac{7(56-171\theta)}{90}\eta. \end{aligned}$	$\begin{aligned} &\beta_{\eta}=-\frac{1}{(4\pi)^2}\frac{133}{10}\eta^2 \\ &\beta_{\omega}=-\frac{1}{(4\pi)^2}\frac{25+1098\omega+200\omega^2}{60}\eta \\ &\beta_{\theta}=\frac{1}{(4\pi)^2}\frac{7(56-171\theta)}{90}\eta. \end{aligned}$	$\begin{aligned} &\beta_{\eta}=-\frac{1}{(4\pi)^2}\frac{133}{10}\eta^2 \\ &\beta_{\omega}=-\frac{1}{(4\pi)^2}\frac{25+1098\omega+200\omega^2}{60}\eta \\ &\beta_{\theta}=\frac{1}{(4\pi)^2}\frac{7(56-171\theta)}{90}\eta. \end{aligned}$
cardona-qft-methods_ EQ0516 (2)	236	■ 1.000	$\Delta x\Delta p\geq\frac{\hbar}{2},$	$\Delta x\Delta p\geq\frac{\hbar}{2},$	$\Delta x\Delta p\geq\frac{\hbar}{2},$
cardona-qft-methods_ EQ0517 (3)	236	■ 1.000	$xy-yx=i\theta,$	$xy-yx=i\theta,$	$xy-yx=i\theta,$
cardona-qft-methods_ EQ0518 (4)	236	■ 1.000	$\Delta x\Delta y\geq\frac{\theta}{2},$	$\Delta x\Delta y\geq\frac{\theta}{2},$	$\Delta x\Delta y\geq\frac{\theta}{2},$
cardona-qft-methods_ EQ0521 (1)	239	■ 1.000	$U(g)=\bigoplus_NU^{(N)}(g),$	$U(g)=\bigoplus_NU^{(N)}(g),$	$U(g)=\bigoplus_NU^{(N)}(g),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0522 (2)	239	■ 1.000	$U^{\{N\}}(g)\left \rho_{\{1\}}\right\rangle\!\!\!\rangle\otimes\left \rho_{\{2\}}\right\rangle\!\!\!\rangle\otimes\cdots\otimes\left \rho_{\{N\}}\right\rangle\!\!\!\rangle=U^{\{1\}}(g)\left \rho_{\{1\}}\right\rangle\!\!\!\rangle\otimes\left \rho_{\{2\}}\right\rangle\!\!\!\rangle\otimes\cdots\otimes\left \rho_{\{N\}}\right\rangle\!\!\!\rangle$	$U^{(N)}(g) \rho_1\rangle\otimes \rho_2\rangle\otimes\cdots\otimes \rho_N\rangle=U^{(1)}(g) \rho_1\rangle\otimes U^{(1)}(g) \rho_2\rangle\otimes\cdots\otimes U^{(1)}(g) \rho_N\rangle.$	$U^{(N)}(g) \rho_1\rangle\otimes \rho_2\rangle\otimes\cdots\otimes \rho_N\rangle=U^{(1)}(g) \rho_1\rangle\otimes U^{(1)}(g) \rho_2\rangle\otimes\cdots\otimes U^{(1)}(g) \rho_N\rangle.$
cardona-qft-methods_ EQ0523 (3)	240	■ 1.000	$\alpha^*=\sum_{g\in G}\alpha(g)g,$	$\alpha^*=\sum_{g\in G}\alpha(g)g,$	$\alpha^*=\sum_{g\in G}\alpha(g)g,$
cardona-qft-methods_ EQ0524 (4)	240	■ 1.000	$\alpha^*\beta^*:=\sum_{g\in G}(\alpha*\beta)(g)g,$	$\alpha^*\beta^*:=\sum_{g\in G}(\alpha*\beta)(g)g,$	$\alpha^*\beta^*:=\sum_{g\in G}(\alpha*\beta)(g)g,$
cardona-qft-methods_ EQ0525 (5)	240	■ 1.000	$(\alpha*\beta)(g)=\sum_{h\in G}\alpha(gh^{-1})\beta(h).$	$(\alpha*\beta)(g)=\sum_{h\in G}\alpha(gh^{-1})\beta(h).$	$(\alpha*\beta)(g)=\sum_{h\in G}\alpha(gh^{-1})\beta(h).$
cardona-qft-methods_ EQ0526 (6)	240	■ 1.000	$\alpha^*=\int d\mu(g)\alpha(g)g.$	$\alpha^*=\int d\mu(g)\alpha(g)g.$	$\alpha^*=\int d\mu(g)\alpha(g)g.$
cardona-qft-methods_ EQ0527 (7)	240	■ 1.000	$\begin{aligned} U^{\{N\}}\left(\alpha^*\right) &= \int d\mu(g)\alpha(g)U^{\{N\}}(g) \\ U\left(\alpha^*\right) &= \int d\mu(g)\alpha(g)U(g) \end{aligned}$	$\begin{aligned} U^{(N)}\left(\alpha^*\right) &= \int d\mu(g)\alpha(g)U^{(N)}(g) \\ U\left(\alpha^*\right) &= \int d\mu(g)\alpha(g)U(g) \end{aligned}$	$\begin{aligned} U^{(N)}(\alpha^*) &= \int d\mu(g)\alpha(g)U^{(N)}(g), \\ U(\alpha^*) &= \int d\mu(g)\alpha(g)U(g). \end{aligned}$
cardona-qft-methods_ EQ0528 (8)	241	■ 1.000	$U^{\{1\}}(g)\left \rho\right\rangle\!\!\!\rangle= \sigma\rangle D_{\sigma\rho}(g),$	$U^{(1)}(g) \rho\rangle= \sigma\rangle D_{\sigma\rho}(g),$	$U^{(1)}(g) \rho\rangle= \sigma\rangle D_{\sigma\rho}(g),$
cardona-qft-methods_ EQ0529 (9)	241	■ 1.000	$\Delta(g)=g\otimes g.$	$\Delta(g)=g\otimes g.$	$\Delta(g)=g\otimes g.$
cardona-qft-methods_ EQ0531	241	■ 1.000	$\begin{aligned} &U^{\{3\}}(g)=\left(U^{\{2\}}\otimes U^{\{1\}}\right)\Delta(g), \\ &U^{\{4\}}(g)=\left(U^{\{3\}}\otimes U^{\{1\}}\right)\Delta(g), \end{aligned}$	$\begin{aligned} U^{(3)}(g) &= \left(U^{(2)}\otimes U^{(1)}\right)\Delta(g), \\ U^{(4)}(g) &= \left(U^{(3)}\otimes U^{(1)}\right)\Delta(g), \end{aligned}$	$\begin{aligned} U^{(3)}(g) &= \left(U^{(2)}\otimes U^{(1)}\right)\Delta(g), \\ U^{(4)}(g) &= \left(U^{(3)}\otimes U^{(1)}\right)\Delta(g), \end{aligned}$
cardona-qft-methods_ EQ0532 (11)	241	■ 1.000	$U^{\{N\}}(g)=U^{\{1\}}(g)\otimes U^{\{1\}}(g)\otimes\cdots\otimes U^{\{1\}}(g).$	$U^{(N)}(g)=U^{(1)}(g)\otimes U^{(1)}(g)\otimes\cdots\otimes U^{(1)}(g).$	$U^{(N)}(g)=U^{(1)}(g)\otimes U^{(1)}(g)\otimes\cdots\otimes U^{(1)}(g).$
cardona-qft-methods_ EQ0533 (12)	241	■ 1.000	$(\Delta\otimes\mathrm{Id})\Delta=(\mathrm{Id}\otimes\Delta)\Delta.$	$(\Delta\otimes\mathrm{Id})\Delta=(\mathrm{Id}\otimes\Delta)\Delta.$	$(\Delta\otimes\mathrm{Id})\Delta=(\mathrm{Id}\otimes\Delta)\Delta.$
cardona-qft-methods_ EQ0534 (13)	241	■ 1.000	$\mathbb{C}G\rightarrow\bigotimes_{N\text{ factors}}\mathbb{C}G.$	$\mathbb{C}G\rightarrow\bigotimes_{N\text{ factors}}\mathbb{C}G.$	$\mathbb{C}G\rightarrow\bigotimes_{N\text{ factors}}\mathbb{C}G.$
cardona-qft-methods_ EQ0535 (14)	241	■ 1.000	$\mathbb{C}G\rightarrow\bigotimes_{(N+1)\text{ factors}}\mathbb{C}G.$	$\mathbb{C}G\rightarrow\bigotimes_{(N+1)\text{ factors}}\mathbb{C}G.$	$\mathbb{C}G\rightarrow\bigotimes_{(N+1)\text{ factors}}\mathbb{C}G.$
cardona-qft-methods_ EQ0536 (15)	242	■ 1.000	$\Delta_\theta(g)=F_\theta^{-1}(g\otimes g)F_\theta,$	$\Delta_\theta(g)=F_\theta^{-1}(g\otimes g)F_\theta,$	$\Delta_\theta(g)=F_\theta^{-1}(g\otimes g)F_\theta,$
cardona-qft-methods_ EQ0537	242	■ 1.000	$F_\theta=e^{\frac{1}{2}P_\mu\theta^{\mu\nu}\otimes P_\nu},$	$F_\theta=e^{\frac{1}{2}P_\mu\theta^{\mu\nu}\otimes P_\nu},$	$F_\theta=e^{\frac{1}{2}P_\mu\theta^{\mu\nu}\otimes P_\nu},$
cardona-qft-methods_ EQ0538	242	■ 1.000	$\epsilon:G\rightarrow\mathbb{C},\quad\epsilon(g)=1.$	$\epsilon:G\rightarrow\mathbb{C},\quad\epsilon(g)=1.$	$\epsilon:G\rightarrow\mathbb{C},\quad\epsilon(g)=1.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E08539	242	■ 1.000	$\backslash\epsilon\mathrm{psilon}:\int\mathrm{d}\mu(g)\backslash\alpha\mathrm{pha}(g)\,g\rightarrow\int\mathrm{d}\mu(g)\backslash\alpha\mathrm{pha}(g)$	$\epsilon:\int d\mu(g)\alpha(g)g\rightarrow\int d\mu(g)\alpha(g)$	$\epsilon:\int d\mu(g)\alpha(g)g\rightarrow\int d\mu(g)\alpha(g)$
cardona-qft-methods_ E08548	243	■ 1.000	$S:\hat{\alpha}=\int\mathrm{d}\mu(g)\backslash\alpha\mathrm{pha}(g)\,g\rightarrow S(\hat{\alpha})=\int\mathrm{d}\mu(g)\backslash\alpha\mathrm{pha}(g)\,S(g)$	$S:\hat{\alpha}=\int d\mu(g)\alpha(g)g\rightarrow S(\hat{\alpha})=\int d\mu(g)\alpha(g)S(g)$	$S:\hat{\alpha}=\int d\mu(g)\alpha(g)g\rightarrow S(\hat{\alpha})=\int d\mu(g)\alpha(g)S(g).$
cardona-qft-methods_ E08541	243	■ 1.000	$m(g\otimes h)=gh,\quad;g,h\in G.$	$m(g\otimes h)=gh,\quad;g,h\in G.$	$m(g\otimes h)=gh,\quad;g,h\in G.$
cardona-qft-methods_ E08544	243	■ 1.000	$\left(m_{13}\otimes m_{24}\right)\left(a_1\otimes a_2\otimes a_3\otimes a_4\right)=a_1a_3\otimes a_2a_4.$	$(m_{13}\otimes m_{24})(a_1\otimes a_2\otimes a_3\otimes a_4)=a_1a_3\otimes a_2a_4.$	$(m_{13}\otimes m_{24})(a_1\otimes a_2\otimes a_3\otimes a_4)=a_1a_3\otimes a_2a_4.$
cardona-qft-methods_ E08545	243	■ 1.000	$\left(\int\mathrm{d}\mu(g)\backslash\alpha\mathrm{pha}(g)\,g\right)^{*}=\int\mathrm{d}\mu(g)\backslash\bar{\alpha}\mathrm{pha}(g)\,g^{-1}$	$\left(\int d\mu(g)\alpha(g)g\right)^{*}=\int d\mu(g)\bar{\alpha}(g)g^{-1}$	$\left(\int d\mu(g)\alpha(g)g\right)^{*}=\int d\mu(g)\bar{\alpha}(g)g^{-1}.$
cardona-qft-methods_ E08546 (16)	244	■ 1.000	$\Delta:H\rightarrow H\otimes H$	$\Delta:H\rightarrow H\otimes H$	$\Delta:H\rightarrow H\otimes H$
cardona-qft-methods_ E08550	244	■ 1.000	$m(\alpha\otimes\beta)=\alpha\beta.$	$m(\alpha\otimes\beta)=\alpha\beta.$	$m(\alpha\otimes\beta)=\alpha\beta.$
cardona-qft-methods_ E08551	244	■ 1.000	$\left(m_{13}\otimes m_{24}\right)\left(\alpha_1\otimes\alpha_2\otimes\alpha_3\otimes\alpha_4\right)=\alpha_1\alpha_3\otimes\alpha_2\alpha_4.$	$(m_{13}\otimes m_{24})(\alpha_1\otimes\alpha_2\otimes\alpha_3\otimes\alpha_4)=\alpha_1\alpha_3\otimes\alpha_2\alpha_4.$	$(m_{13}\otimes m_{24})(\alpha_1\otimes\alpha_2\otimes\alpha_3\otimes\alpha_4)=\alpha_1\alpha_3\otimes\alpha_2\alpha_4.$
cardona-qft-methods_ E08552 (19)	244	■ 1.000	$\Delta m(\alpha\otimes\beta)=\left(m_{13}\otimes m_{24}\right)(\Delta\otimes\Delta)(\alpha\otimes\beta).$	$\Delta m(\alpha\otimes\beta)=(m_{13}\otimes m_{24})(\Delta\otimes\Delta)(\alpha\otimes\beta).$	$\Delta m(\alpha\otimes\beta)=(m_{13}\otimes m_{24})(\Delta\otimes\Delta)(\alpha\otimes\beta).$
cardona-qft-methods_ E08553	245	■ 1.000	$(F\otimes Id)(\Delta\otimes Id)F=(Id\otimes F)(Id\otimes\Delta)F,\quad(\epsilon\otimes Id)F=(Id\otimes\epsilon)F=1.$	$(F\otimes Id)(\Delta\otimes Id)F=(Id\otimes F)(Id\otimes\Delta)F,\quad(\epsilon\otimes Id)F=(Id\otimes\epsilon)F=1.$	$(F\otimes Id)(\Delta\otimes Id)F=(Id\otimes F)(Id\otimes\Delta)F,\quad(\epsilon\otimes Id)F=(Id\otimes\epsilon)F=1.$
cardona-qft-methods_ E08554	245	■ 1.000	$\Delta(g)=g\otimes g,\quad g\in G\subset GCG,$	$\Delta(g)=g\otimes g,\quad g\in G\subset GCG,$	$\Delta(g)=g\otimes g,\quad g\in G\subset GCG,$
cardona-qft-methods_ E08555	245	■ 1.000	$\xi\otimes\eta\rightarrow(\rho\otimes\rho)\Delta(g)\xi\otimes\eta=\rho(g)\xi\otimes\rho(g)\eta.$	$\xi\otimes\eta\rightarrow(\rho\otimes\rho)\Delta(g)\xi\otimes\eta=\rho(g)\xi\otimes\rho(g)\eta.$	$\xi\otimes\eta\rightarrow(\rho\otimes\rho)\Delta(g)\xi\otimes\eta=\rho(g)\xi\otimes\rho(g)\eta.$
cardona-qft-methods_ E08556	245	■ 1.000	$\tau(\xi\otimes\eta)=\eta\otimes\xi.$	$\tau(\xi\otimes\eta)=\eta\otimes\xi.$	$\tau(\xi\otimes\eta)=\eta\otimes\xi.$
cardona-qft-methods_ E08557	246	■ 1.000	$\tau_{i,i+1}=\underbrace{1\otimes\cdots\otimes 1}_{(i-1)\text{ factors}}\otimes\tau\otimes\underbrace{1\otimes\cdots\otimes 1}_{(N-i-1)\text{ factors}}.$	$\tau_{i,i+1}=\underbrace{1\otimes\cdots\otimes 1}_{(i-1)\text{ factors}}\otimes\tau\otimes\underbrace{1\otimes\cdots\otimes 1}_{(N-i-1)\text{ factors}}.$	$\tau_{i,i+1}=\underbrace{1\otimes\cdots\otimes 1}_{(i-1)\text{ factors}}\otimes\tau\otimes\underbrace{1\otimes\cdots\otimes 1}_{(N-i-1)\text{ factors}}.$
cardona-qft-methods_ E08559	246	■ 1.000	$\Delta^{\mathrm{op}}(\eta)=(\eta^{\{2\}}\otimes\eta^{\{2\}})$	$\Delta^{\mathrm{op}}(\eta)=\eta^{(2)}\otimes\eta^{(2)}$	$\Delta^{\mathrm{op}}(\eta)=\eta^{(2)}\otimes\eta^{(2)}.$
cardona-qft-methods_ E08560	247	■ 1.000	$R_{i,i+1}=\underbrace{1\otimes 1\otimes\cdots\otimes 1}_{(i-1)\text{ factors}}\otimes R\otimes\underbrace{1\otimes 1\otimes\cdots\otimes 1}_{(N-i-1)\text{ factors}}$	$R_{i,i+1}=\underbrace{1\otimes 1\otimes\cdots\otimes 1}_{(i-1)\text{ factors}}\otimes R\otimes\underbrace{1\otimes 1\otimes\cdots\otimes 1}_{(N-i-1)\text{ factors}}$	$R_{i,i+1}=\underbrace{1\otimes 1\otimes\cdots\otimes 1}_{(i-1)\text{ factors}}\otimes R\otimes\underbrace{1\otimes 1\otimes\cdots\otimes 1}_{(N-i-1)\text{ factors}}$
cardona-qft-methods_ E08561	247	■ 1.000	$\mathcal{R}_{i,i+1}=\tau_{i,i+1}(\rho\otimes\rho\otimes\cdots\otimes\rho)R_{i,i+1}$	$\mathcal{R}_{i,i+1}=\tau_{i,i+1}(\rho\otimes\rho\otimes\cdots\otimes\rho)R_{i,i+1}$	$\mathcal{R}_{i,i+1}=\tau_{i,i+1}(\rho\otimes\rho\otimes\cdots\otimes\rho)R_{i,i+1}$
cardona-qft-methods_ E08562	247	■ 1.000	$R_{i+1,i+2}R_{i,i+2}R_{i,i+1}=R_{i,i+1}R_{i,i+2}R_{i+1,i+2}.$	$R_{i+1,i+2}R_{i,i+2}R_{i,i+1}=R_{i,i+1}R_{i,i+2}R_{i+1,i+2}.$	$R_{i+1,i+2}R_{i,i+2}R_{i,i+1}=R_{i,i+1}R_{i,i+2}R_{i+1,i+2}.$
cardona-qft-methods_ E08563	247	■ 1.000	$\Delta_{\theta}\{g\}=F_{\theta}^{-1}(g\otimes g)F_{\theta}$	$\Delta_{\theta}(g)=F_{\theta}^{-1}(g\otimes g)F_{\theta}$	$\Delta_{\theta}(g)=F_{\theta}^{-1}(g\otimes g)F_{\theta}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0564	247	■ 1.000	$\tau_{\{\theta\}}=(\rho \otimes \rho) F_{\{\theta\}}^{-1} \tau(\rho \otimes \rho) F_{\{\theta\}}$	$\tau_{\theta}=(\rho \otimes \rho) F_{\theta}^{-1} \tau(\rho \otimes \rho) F_{\theta}$	$\tau_{\theta}=(\rho \otimes \rho) F_{\theta}^{-1} \tau(\rho \otimes \rho) F_{\theta}$
cardona-qft-methods_ EQ0565	247	■ 1.000	$\tau_{\{\theta\}}=\tau(\rho \otimes \rho) F_{\{\theta\}}^2,$	$\tau_{\theta}=\tau(\rho \otimes \rho) F_{\theta}^2,$	$\tau_{\theta}=\tau(\rho \otimes \rho) F_{\theta}^2,$
cardona-qft-methods_ EQ0566	247	■ 1.000	$R_{\{\theta\}}=F_{\{\theta\}}^2$.	$R_{\theta}=F_{\theta}^2.$	$R_{\theta}=F_{\theta}^2.$
cardona-qft-methods_ EQ0567 (1)	248	■ 1.000	$R(t)=0 \quad t<0$	$R(t)=0 \quad t<0$	$R(t)=0 \quad t<0$
cardona-qft-methods_ EQ0569 (3)	248	■ 1.000	$\Im \omega>0$.	$\Im \omega>0.$	$\Im \omega>0.$
cardona-qft-methods_ EQ0570 (4)	248	■ 1.000	$\begin{gathered} {[\rho(x), \eta(y)] = 0 \quad \text{if} \\ \left(x^0 - y^0\right)^2 - (\vec{x} - \vec{y})^2 < 0} \end{gathered}$	$[\rho(x), \eta(y)] = 0 \quad \text{if} \\ \left(x^0 - y^0\right)^2 - (\vec{x} - \vec{y})^2 < 0$	$[\rho(x), \eta(y)] = 0 \quad \text{if} \\ \left(x^0 - y^0\right)^2 - (\vec{x} - \vec{y})^2 < 0$
cardona-qft-methods_ EQ0571 (6)	249	■ 1.000	$[\varphi(x), \chi(y)]_{-} = 0 \quad x \sim y$	$[\varphi(x), \chi(y)]_{-} = 0 \quad x \sim y$	$[\varphi(x), \chi(y)]_{-} = 0 \quad x \sim y$
cardona-qft-methods_ EQ0572 (7)	249	■ 1.000	$\left.\left.\left[\psi_{\alpha}^{(1)}(x), \psi_{\alpha}^{(2)}(y)\right]\right]_{+} = 0 \quad x \sim y\right.$	$\left[\psi_{\alpha}^{(1)}(x), \psi_{\alpha}^{(2)}(y)\right]_{+} = 0 \quad x \sim y$	$[\psi_{\alpha}^{(1)}(x), \psi_{\alpha}^{(2)}(y)]_{+} = 0 \quad x \sim y$
cardona-qft-methods_ EQ0575 (10)	249	■ 1.000	$g \rightarrow [U_1 \otimes U_2](g \otimes g)$	$g \rightarrow [U_1 \otimes U_2](g \otimes g)$	$g \rightarrow [U_1 \otimes U_2](g \otimes g)$
cardona-qft-methods_ EQ0576 (11)	250	■ 1.000	$\begin{aligned} \Delta: G &\rightarrow G \otimes G, \\ g &\rightarrow \Delta(g) := g \otimes g, \end{aligned}$	$\Delta: G \rightarrow G \otimes G,$ $g \rightarrow \Delta(g) := g \otimes g,$	$\Delta: G \rightarrow G \otimes G,$ $g \rightarrow \Delta(g) := g \otimes g,$
cardona-qft-methods_ EQ0577 (12)	250	■ 1.000	$F_{\{\theta\}}=e^{\frac{i}{2}\partial_{\mu}\otimes\theta^{\mu\nu}\partial_{\nu}}=\text{"Twist element"}$	$F_{\theta}=e^{\frac{i}{2}\partial_{\mu}\otimes\theta^{\mu\nu}\partial_{\nu}}=\text{"Twist element"}$	$F_{\theta}=e^{\frac{i}{2}\partial_{\mu}\otimes\theta^{\mu\nu}\partial_{\nu}}=\text{"Twist element"}$
cardona-qft-methods_ EQ0578 (13)	250	■ 1.000	$f \star g=m_{\{0\}}\left[F_{\{\theta\}} f \otimes g\right],$	$f \star g=m_0\left[F_{\theta} f \otimes g\right],$	$f \star g=m_0[F_{\theta} f \otimes g],$
cardona-qft-methods_ EQ0579 (14)	250	■ 1.000	$m_{\{0\}}(\alpha \otimes \beta)(x)=\alpha(x) \beta(x)$	$m_0(\alpha \otimes \beta)(x)=\alpha(x) \beta(x)$	$m_0(\alpha \otimes \beta)(x)=\alpha(x) \beta(x)$
cardona-qft-methods_ EQ0581	250	■ 1.000	$\Lambda: \alpha \rightarrow \Lambda^* \alpha, \quad \Lambda^* \alpha(x)=\alpha\left[\Lambda^{-1}(x)\right] .$	$\Lambda: \alpha \rightarrow \Lambda^* \alpha, \quad \left(\Lambda^* \alpha\right)(x)=\alpha\left[\Lambda^{-1}(x)\right] .$	$\Lambda: \alpha \rightarrow \Lambda^* \alpha, \quad \left(\Lambda^* \alpha\right)(x)=\alpha\left[\Lambda^{-1}(x)\right].$
cardona-qft-methods_ EQ0582 (15)	250	■ 1.000	$\Delta_{\{\theta\}}(\Lambda)=F_{\{\theta\}}^{-1}(\Lambda \otimes \lambda) F_{\{\theta\}} .$	$\Delta_{\theta}(\Lambda)=F_{\theta}^{-1}(\Lambda \otimes \lambda) F_{\theta}.$	$\Delta_{\theta}(\Lambda)=F_{\theta}^{-1}(\Lambda \otimes \lambda) F_{\theta}.$
cardona-qft-methods_ EQ0583 (16)	250	■ 1.000	$\phi \otimes_{S, A} \chi \equiv \frac{1}{2}(\phi \otimes \chi \pm \chi \otimes \phi)$	$\phi \otimes_{S, A} \chi \equiv \frac{1}{2}(\phi \otimes \chi \pm \chi \otimes \phi)$	$\phi \otimes_{S, A} \chi \equiv \frac{1}{2}(\phi \otimes \chi \pm \chi \otimes \phi)$
cardona-qft-methods_ EQ0584 (17)	251	■ 1.000	$\tau_{\{0\}}(\phi \otimes \chi)=\chi \otimes \phi$	$\tau_0(\phi \otimes \chi)=\chi \otimes \phi$	$\tau_0(\phi \otimes \chi)=\chi \otimes \phi$
cardona-qft-methods_ EQ0585 (18)	251	■ 1.000	$\tau_{\{0\}} \Delta_{\{0\}}(\Lambda)=\Delta_{\{0\}}(\Lambda) \tau_{\{0\}}$	$\tau_0 \Delta_0(\Lambda)=\Delta_0(\Lambda) \tau_0$	$\tau_0 \Delta_0(\Lambda)=\Delta_0(\Lambda) \tau_0$
cardona-qft-methods_ EQ0586 (19)	251	■ 1.000	$\left(\frac{1 \pm \tau_0}{2}\right)(\phi \otimes \chi) .$	$\left(\frac{1 \pm \tau_0}{2}\right)(\phi \otimes \chi) .$	$\left(\frac{1 \pm \tau_0}{2}\right)(\phi \otimes \chi) .$
cardona-qft-methods_ EQ0587 (20)	251	■ 1.000	$\tau_{\{0\}} F_{\{\theta\}}=F_{\{\theta\}}^{-1} \tau_{\{0\}},$	$\tau_0 F_{\theta}=F_{\theta}^{-1} \tau_0,$	$\tau_0 F_{\theta}=F_{\theta}^{-1} \tau_0,$
cardona-qft-methods_ EQ0589 (22)	251	■ 1.000	$\tau_{\{\theta\}} \equiv F_{\{\theta\}}^{-1} \tau_0 F_{\{\theta\}}, \quad \tau_{\{\theta\}}^2=1 \otimes 1,$	$\tau_{\theta} \equiv F_{\theta}^{-1} \tau_0 F_{\theta}, \quad \tau_{\theta}^2=1 \otimes 1,$	$\tau_{\theta} \equiv F_{\theta}^{-1} \tau_0 F_{\theta}, \quad \tau_{\theta}^2=1 \otimes 1,$
cardona-qft-methods_ EQ0590 (23)	251	■ 1.000	$\Delta_{\{\theta\}}(\Lambda)=F_{\{\theta\}}^{-1} \Lambda \otimes \Lambda F_{\{\theta\}} .$	$\Delta_{\theta}(\Lambda)=F_{\theta}^{-1} \Lambda \otimes \Lambda F_{\theta}.$	$\Delta_{\theta}(\Lambda)=F_{\theta}^{-1} \Lambda \otimes \Lambda F_{\theta}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E08593 (2)	252	■ 1.000	$U:(a,\Lambda)\rightarrow U(a,\Lambda)$	$U:(a,\Lambda)\rightarrow U(a,\Lambda)$	$U:(a,\Lambda)\rightarrow U(a,\Lambda)$
cardona-qft-methods_ E08594 (3)	252	■ 1.000	$U(a,\Lambda)\varphi(x)U(a,\Lambda)^{-1}=\varphi((a,\Delta)x)$	$U(a,\Lambda)\varphi(x)U(a,\Lambda)^{-1}=\varphi((a,\Delta)x)$	$U(a,\Lambda)\varphi(x)U(a,\Lambda)^{-1}=\varphi((a,\Delta)x)$
cardona-qft-methods_ E08595 (4)	252	■ 1.000	$\operatorname{Ad} U(a,\Lambda)\varphi=U(a,\Lambda)\varphi U(a,\Lambda)^{-1}$	$\operatorname{Ad} U(a,\Lambda)\varphi=U(a,\Lambda)\varphi U(a,\Lambda)^{-1}$	$\operatorname{Ad} U(a,\Lambda)\varphi=U(a,\Lambda)\varphi U(a,\Lambda)^{-1}$
cardona-qft-methods_ E08596 (5)	252	■ 1.000	$(a,\Delta):\alpha\rightarrow(a,\Lambda)\triangleright\alpha,\quad [(a,\Lambda)\alpha](x)\quad\alpha\in S(M^{d+1})$	$(a,\Delta):\alpha\rightarrow(a,\Lambda)\triangleright\alpha,\quad [(a,\Lambda)\alpha](x)\quad\alpha\in S(M^{d+1})$	$(a,\Delta):\alpha\rightarrow(a,\Lambda)\triangleright\alpha,\quad [(a,\Lambda)\alpha](x)\quad\alpha\in S(M^{d+1})$
cardona-qft-methods_ E08597 (6)	252	■ 1.000	$\Delta_0((a,\Lambda))=(a,\Lambda)\otimes(a,\Lambda)$	$\Delta_0((a,\Lambda))=(a,\Lambda)\otimes(a,\Lambda)$	$\Delta_0((a,\Lambda))=(a,\Lambda)\otimes(a,\Lambda)$
cardona-qft-methods_ E08599 (8)	252	■ 1.000	$\varphi\left(x_1\right)\varphi\left(x_2\right)\ldots\varphi\left(x_N\right).$	$\varphi\left(x_1\right)\varphi\left(x_2\right)\ldots\varphi\left(x_N\right).$	$\varphi\left(x_1\right)\varphi\left(x_2\right)\ldots\varphi\left(x_N\right).$
cardona-qft-methods_ E08600 (9)	253	■ 1.000	$(\varphi\otimes\ldots\otimes\varphi)\left(x_1,x_2,\ldots,x_N\right)=\varphi\left(x_1\right)\varphi\left(x_2\right)\ldots\varphi\left(x_N\right)$	$(\varphi\otimes\ldots\otimes\varphi)\left(x_1,x_2,\ldots,x_N\right)=\varphi\left(x_1\right)\varphi\left(x_2\right)\ldots\varphi\left(x_N\right)$	$(\varphi\otimes\ldots\otimes\varphi)\left(x_1,x_2,\ldots,x_N\right)=\varphi\left(x_1\right)\varphi\left(x_2\right)\ldots\varphi\left(x_N\right)$
cardona-qft-methods_ E08601 (10)	253	■ 1.000	$(\underbrace{\mathbb{I}\otimes\mathbb{I}\otimes\ldots\mathbb{I}}_{N-1}\otimes\mathbb{I}\otimes\mathbb{I}\otimes\ldots\mathbb{I}\otimes\Delta_0)\ldots\Delta_0$	$(\underbrace{\mathbb{I}\otimes\mathbb{I}\otimes\mathbb{I}\otimes\ldots\mathbb{I}}_{N-1}\otimes\mathbb{I}\otimes\mathbb{I}\otimes\ldots\mathbb{I}\otimes\Delta_0)\ldots\Delta_0$	$(\underbrace{\mathbb{I}\otimes\mathbb{I}\otimes\mathbb{I}\otimes\ldots\mathbb{I}}_{N-1}\otimes\mathbb{I}\otimes\mathbb{I}\otimes\ldots\mathbb{I}\otimes\Delta_0)\ldots\Delta_0$
cardona-qft-methods_ E08602 (11)	253	■ 1.000	$\Delta_0^N(a,\Lambda)=(a,\Lambda)\otimes(a,\Lambda)\otimes\cdots\otimes(a,\Lambda).$	$\Delta_0^N(a,\Lambda)=(a,\Lambda)\otimes(a,\Lambda)\otimes\cdots\otimes(a,\Lambda).$	$\Delta_0^N(a,\Lambda)=(a,\Lambda)\otimes(a,\Lambda)\otimes\cdots\otimes(a,\Lambda).$
cardona-qft-methods_ E08603 (12)	253	■ 1.000	$U(a,\Lambda)\left(\varphi^{\otimes N}\left((a,\Lambda)^{-1}x_1,\ldots,(a,\Lambda)^{-1}x_N\right)\right)U(a,\Lambda)^{-1}=\varphi^{\otimes N}\left(x_1,x_2,\ldots,x_N\right)$	$U(a,\Lambda)\left(\varphi^{\otimes N}\left((a,\Lambda)^{-1}x_1,\ldots,(a,\Lambda)^{-1}x_N\right)\right)U(a,\Lambda)^{-1}=\varphi^{\otimes N}\left(x_1,x_2,\ldots,x_N\right)$	$U(a,\Lambda)\left(\varphi^{\otimes N}\left((a,\Lambda)^{-1}x_1,\ldots,(a,\Lambda)^{-1}x_N\right)\right)U(a,\Lambda)^{-1}=\varphi^{\otimes N}\left(x_1,x_2,\ldots,x_N\right)$
cardona-qft-methods_ E08604 (13)	253	■ 1.000	$\varphi=\int d\mu(p)\left(c_p^\dagger e_p+c_p e_{-p}\right)=\varphi^{(-)}+\varphi^{(+)}$	$\varphi=\int d\mu(p)\left(c_p^\dagger e_p+c_p e_{-p}\right)=\varphi^{(-)}+\varphi^{(+)}$	$\varphi=\int d\mu(p)\left(c_p^\dagger e_p+c_p e_{-p}\right)=\varphi^{(-)}+\varphi^{(+)}$
cardona-qft-methods_ E08606	253	■ 1.000	$\begin{array}{r} \varphi^{(-)}\left(x_1\right)\varphi^{(-)}\left(x_2\right)\ldots\varphi^{(-)}\left(x_N\right) 0\rangle=\\ \int\prod_i d\mu\left(p_i\right)c_{p_1}^\dagger\left(x_1\right)c_{p_2}^\dagger\left(x_2\right)\ldots c_{p_N}^\dagger\left(x_N\right) 0\rangle e_{p_1}\left(x_1\right)e_{p_2}\left(x_2\right)\ldots e_{p_N}\left(x_N\right) \end{array}$	$\varphi^{(-)}\left(x_1\right)\varphi^{(-)}\left(x_2\right)\ldots\varphi^{(-)}\left(x_N\right) 0\rangle=\\ \int\prod_i d\mu\left(p_i\right)c_{p_1}^\dagger\left(x_1\right)c_{p_2}^\dagger\left(x_2\right)\ldots c_{p_N}^\dagger\left(x_N\right) 0\rangle e_{p_1}\left(x_1\right)e_{p_2}\left(x_2\right)\ldots e_{p_N}\left(x_N\right)$	$\varphi^{(-)}\left(x_1\right)\varphi^{(-)}\left(x_2\right)\ldots\varphi^{(-)}\left(x_N\right) 0\rangle=\\ \int\prod_i d\mu\left(p_i\right)c_{p_1}^\dagger\left(x_1\right)c_{p_2}^\dagger\left(x_2\right)\ldots c_{p_N}^\dagger\left(x_N\right) 0\rangle e_{p_1}\left(x_1\right)e_{p_2}\left(x_2\right)\ldots e_{p_N}\left(x_N\right)$
cardona-qft-methods_ E08607 (15)	253	■ 1.000	$\left[P_\mu,c_p^\dagger\right]=P_\mu c_p^\dagger,\quad P_\mu 0\rangle=0$	$\left[P_\mu,c_p^\dagger\right]=P_\mu c_p^\dagger,\quad P_\mu 0\rangle=0$	$\left[P_\mu,c_p^\dagger\right]=P_\mu c_p^\dagger,\quad P_\mu 0\rangle=0$
cardona-qft-methods_ E08608 (16)	254	■ 1.000	$\mathcal{P}_\mu e_p=-p_\mu e_p$	$\mathcal{P}_\mu e_p=-p_\mu e_p$	$\mathcal{P}_\mu e_p=-p_\mu e_p$
cardona-qft-methods_ E08609 (17)	254	■ 1.000	$\Delta_0\left(\mathcal{P}_\mu\right)=\mathbb{I}\otimes\mathcal{P}_\mu+\mathcal{P}_\mu\otimes\mathbb{I}$	$\Delta_0\left(\mathcal{P}_\mu\right)=\mathbb{I}\otimes\mathcal{P}_\mu+\mathcal{P}_\mu\otimes\mathbb{I}$	$\Delta_0\left(\mathcal{P}_\mu\right)=\mathbb{I}\otimes\mathcal{P}_\mu+\mathcal{P}_\mu\otimes\mathbb{I}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0610 (18)	254	■ 1.000	$\left(\mathbb{I} \otimes \mathbb{I} \otimes \mathbb{I} \dots \mathbb{I} \otimes \Delta_0\right) \dots \Delta_0(\mathcal{P}_\mu) e_{p_1} \otimes e_{p_2} \otimes \dots e_{p_N} = - \sum_i p_{ij} e_{p_1} \otimes e_{p_2} \otimes \dots e_{p_N}.$	$(\mathbb{I} \otimes \mathbb{I} \otimes \mathbb{I} \dots \mathbb{I} \otimes \Delta_0) \dots \Delta_0(\mathcal{P}_\mu) e_{p_1} \otimes e_{p_2} \otimes \dots e_{p_N} = - \sum_i p_{ij} e_{p_1} \otimes e_{p_2} \otimes \dots e_{p_N}.$	$(\mathbb{I} \otimes \mathbb{I} \otimes \mathbb{I} \dots \mathbb{I} \otimes \Delta_0) \dots \Delta_0(\mathcal{P}_\mu) e_{p_1} \otimes e_{p_2} \otimes \dots e_{p_N} = - \sum_i p_{ij} e_{p_1} \otimes e_{p_2} \otimes \dots e_{p_N}.$
cardona-qft-methods_ EQ0614 (22)	254	■ 1.000	$d \, \mu \left(\Lambda^{-1} p_i \right) = d \, \mu \left(p_i \right)$	$d\mu\left(\Lambda^{-1}p_i\right)=d\mu\left(p_i\right)$	$d\mu(\Lambda^{-1}p_i)=d\mu(p_i)$
cardona-qft-methods_ EQ0616 (24)	255	■ 1.000	$\left(\alpha^* f\right)(p)=f(\alpha p)$	$(\alpha^* f)(p)=f(\alpha p)$	$(\alpha^* f)(p)=f(\alpha p)$
cardona-qft-methods_ EQ0619 (27)	256	■ 1.000	$U(a,\Lambda)\varphi^2\left((a,\Lambda)^{-1}x\right)U(a,\Lambda)^{-1}=\varphi^2(x).$	$U(a,\Lambda)\varphi^2\left((a,\Lambda)^{-1}x\right)U(a,\Lambda)^{-1}=\varphi^2(x).$	$U(a,\Lambda)\varphi^2((a,\Lambda)^{-1}x)U(a,\Lambda)^{-1}=\varphi^2(x).$
cardona-qft-methods_ EQ0620 (28)	256	■ 1.000	$U(a,\Delta)\Psi\left((a,\Delta)^{-1}x\right)U(a,\Delta)^{-1}=\Psi$	$U(a,\Delta)\Psi\left((a,\Delta)^{-1}x\right)U(a,\Delta)^{-1}=\Psi$	$U(a,\Delta)\Psi((a,\Delta)^{-1}x)U(a,\Delta)^{-1}=\Psi$
cardona-qft-methods_ EQ0622 (30)	257	■ 1.000	$m_{\theta}(\gamma\otimes\delta)=\gamma(x)\delta(x),\quad \gamma,\delta\in\mathcal{A}_0\left(M^{(d+1)}\right).$	$m_{\theta}(\gamma\otimes\delta)=\gamma(x)\delta(x),\quad \gamma,\delta\in\mathcal{A}_0\left(M^{(d+1)}\right).$	$m_{\theta}(\gamma\otimes\delta)=\gamma(x)\delta(x),\quad \gamma,\delta\in\mathcal{A}_0(M^{(d+1)}).$
cardona-qft-methods_ EQ0625 (33)	257	■ 1.000	$U_{\theta}(a,\Lambda)\varphi_{\theta}\left((a,\Lambda)^{-1}x\right)U_{\theta}(a,\Lambda)^{-1}=\varphi_{\theta}(x)$	$U_{\theta}(a,\Lambda)\varphi_{\theta}\left((a,\Lambda)^{-1}x\right)U_{\theta}(a,\Lambda)^{-1}=\varphi_{\theta}(x)$	$U_{\theta}(a,\Lambda)\varphi_{\theta}((a,\Lambda)^{-1}x)U_{\theta}(a,\Lambda)^{-1}=\varphi_{\theta}(x)$
cardona-qft-methods_ EQ0626 (34)	257	■ 1.000	$\varphi=\int d\mu(p)\left[a_p^{\dagger}e_p+a_pe_{-p}\right]=\varphi_{\theta}^{(-)}+\varphi_{\theta}^{(+)},\quad d\mu(p)=\frac{d^dp}{2 p_0 }.$	$\varphi=\int d\mu(p)\left[a_p^{\dagger}e_p+a_pe_{-p}\right]=\varphi_{\theta}^{(-)}+\varphi_{\theta}^{(+)},\quad d\mu(p)=\frac{d^dp}{2 p_0 }.$	$\varphi=\int d\mu(p)[a_p^{\dagger}e_p+a_pe_{-p}]=\varphi_{\theta}^{(-)}+\varphi_{\theta}^{(+)},\quad d\mu(p)=\frac{d^dp}{2 p_0 }.$
cardona-qft-methods_ EQ0627 (35)	257	■ 1.000	$a_p 0\rangle=0,\forall p.$	$a_p 0\rangle=0,\forall p.$	$a_p 0\rangle=0,\forall p.$
cardona-qft-methods_ EQ0628 (36)	257	■ 1.000	$U_{\theta}(a,\Lambda)a_p^{\dagger}U_{\theta}(a,\Lambda)^{-1}=a_{\Lambda p}^{\dagger},\quad U_{\theta}(a,\Lambda)a_pU_{\theta}(a,\Lambda)^{-1}=a_{\Delta p}$	$U_{\theta}(a,\Lambda)a_p^{\dagger}U_{\theta}(a,\Lambda)^{-1}=a_{\Lambda p}^{\dagger},\quad U_{\theta}(a,\Lambda)a_pU_{\theta}(a,\Lambda)^{-1}=a_{\Delta p}$	$U_{\theta}(a,\Lambda)a_p^{\dagger}U_{\theta}(a,\Lambda)^{-1}=a_{\Lambda p}^{\dagger},\quad U_{\theta}(a,\Lambda)a_pU_{\theta}(a,\Lambda)^{-1}=a_{\Delta p}$
cardona-qft-methods_ EQ0629 (37)	257	■ 1.000	$U_{\theta}(a,\Lambda) 0\rangle=a_{\Lambda p}^{\dagger} 0\rangle.$	$U_{\theta}(a,\Lambda) 0\rangle=a_{\Lambda p}^{\dagger} 0\rangle.$	$U_{\theta}(a,\Lambda) 0\rangle=a_{\Lambda p}^{\dagger} 0\rangle.$
cardona-qft-methods_ EQ0630 (38)	258	■ 1.000	$\int \prod_i d\mu(p_i) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e_{p_1} \otimes e_{p_2}$	$\int \prod_i d\mu(p_i) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e_{p_1} \otimes e_{p_2}$	$\int \prod_i d\mu(p_i) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e_{p_1} \otimes e_{p_2}$
cardona-qft-methods_ EQ0631 (39)	258	■ 1.000	$\Delta_{\theta}(P_{\mu})e_{p_1}\otimes e_{p_2}=(\mathbb{I}\otimes P_{\mu}+P_{\mu}\otimes\mathbb{I})e_{p_1}\otimes e_{p_2}=-\left(\sum_i p_{i\mu}e_{p_1}\otimes e_{p_2}\right).$	$\Delta_{\theta}(P_{\mu})e_{p_1}\otimes e_{p_2}=(\mathbb{I}\otimes P_{\mu}+P_{\mu}\otimes\mathbb{I})e_{p_1}\otimes e_{p_2}=-\left(\sum_i p_{i\mu}e_{p_1}\otimes e_{p_2}\right).$	$\Delta_{\theta}(P_{\mu})e_{p_1}\otimes e_{p_2}=(\mathbb{I}\otimes P_{\mu}+P_{\mu}\otimes\mathbb{I})e_{p_1}\otimes e_{p_2}=-\left(\sum_i p_{i\mu}e_{p_1}\otimes e_{p_2}\right).$
cardona-qft-methods_ EQ0632 (40)	258	■ 1.000	$\left[P_{\mu}^{\theta},a_p^{\dagger}\right]=P_{\mu}a_p^{\dagger}$	$\left[P_{\mu}^{\theta},a_p^{\dagger}\right]=P_{\mu}a_p^{\dagger}$	$\left[P_{\mu}^{\theta},a_p^{\dagger}\right]=P_{\mu}a_p^{\dagger},$
cardona-qft-methods_ EQ0633 (41)	258	■ 1.000	$\Delta_{\theta}(\Lambda)\triangleright e_{p_1}\otimes e_{p_2}=\mathcal{F}_{\theta}^{-1}(\Lambda\otimes\Lambda)\mathcal{F}e_{p_1}\otimes e_{p_2}.$	$\Delta_{\theta}(\Lambda)\triangleright e_{p_1}\otimes e_{p_2}=\mathcal{F}_{\theta}^{-1}(\Lambda\otimes\Lambda)\mathcal{F}e_{p_1}\otimes e_{p_2}.$	$\Delta_{\theta}(\Lambda)\triangleright e_{p_1}\otimes e_{p_2}=\mathcal{F}_{\theta}^{-1}(\Lambda\otimes\Lambda)\mathcal{F}e_{p_1}\otimes e_{p_2}.$
cardona-qft-methods_ EQ0635 (42)	258	■ 1.000	$a_p^{\dagger}=c_p^{\dagger}e^{\frac{i}{2}p\wedge P}$	$a_p^{\dagger}=c_p^{\dagger}e^{\frac{i}{2}p\wedge P}$	$a_p^{\dagger}=c_p^{\dagger}e^{\frac{i}{2}p\wedge P}$
cardona-qft-methods_ EQ0636 (43)	258	■ 1.000	$U_{\theta}(a,\Lambda)=U_0(a,\Lambda)=U(a,\Lambda).$	$U_{\theta}(a,\Lambda)=U_0(a,\Lambda)=U(a,\Lambda).$	$U_{\theta}(a,\Lambda)=U_0(a,\Lambda)=U(a,\Lambda).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0637 (44)	258	■ 1.000	$a_{\{p\}}=e^{\{-\frac{i}{2}p\wedge P\}}c_{\{p\}}=c_{\{p\}}e^{\{-\frac{i}{2}p\wedge P\}}$	$a_p = e^{-\frac{i}{2}p\wedge P}c_p = c_pe^{-\frac{i}{2}p\wedge P}$	$a_p = e^{-\frac{i}{2}p\wedge P}c_p = c_pe^{-\frac{i}{2}p\wedge P},$
cardona-qft-methods_ EQ0639 (46)	259	■ 1.000	$\begin{gathered} U(a, \Lambda) \varphi_{\theta}(x_1) \varphi_{\theta}(x_2) \dots \varphi_{\theta}(x_N) U(a, \Lambda)^{-1} 0\rangle = \\ \varphi_{\theta}((a, \Lambda)(x_1)) \varphi_{\theta}((a, \Lambda)(x_2)) \dots \varphi_{\theta}((a, \Lambda)(x_N)) 0\rangle \end{gathered}$	$U(a, \Lambda) \varphi_{\theta}(x_1) \varphi_{\theta}(x_2) \dots \varphi_{\theta}(x_N) U(a, \Lambda)^{-1} 0\rangle = \\ \varphi_{\theta}((a, \Lambda)(x_1)) \varphi_{\theta}((a, \Lambda)(x_2)) \dots \varphi_{\theta}((a, \Lambda)(x_N)) 0\rangle$	$U(a, \Lambda) \varphi_{\theta}(x_1) \varphi_{\theta}(x_2) \dots \varphi_{\theta}(x_N) U(a, \Lambda)^{-1} 0\rangle = \\ \varphi_{\theta}((a, \Lambda)(x_1)) \varphi_{\theta}((a, \Lambda)(x_2)) \dots \varphi_{\theta}((a, \Lambda)(x_N)) 0\rangle$
cardona-qft-methods_ EQ0640 (47)	259	■ 1.000	$\begin{gathered} \tau_{\theta} = \mathcal{F}_{\theta}^{-1} \tau_0 \mathcal{F}, \\ \tau_0 \alpha \otimes \beta := \beta \otimes \alpha, \\ \tau_0 \alpha \otimes \beta := \beta \otimes \alpha, \end{gathered}$	$\tau_{\theta} = \mathcal{F}_{\theta}^{-1} \tau_0 \mathcal{F}, \\ \tau_0 \alpha \otimes \beta := \beta \otimes \alpha,$	$\tau_{\theta} = \mathcal{F}_{\theta}^{-1} \tau_0 \mathcal{F}, \\ \tau_0 \alpha \otimes \beta := \beta \otimes \alpha,$
cardona-qft-methods_ EQ0641 (49)	259	■ 1.000	$\tau_0^2 = \mathbb{I} \quad \tau_{\theta}^2 = \mathbb{I}.$	$\tau_0^2 = \mathbb{I} \quad \tau_{\theta}^2 = \mathbb{I}.$	$\tau_0^2 = \mathbb{I} \quad \tau_{\theta}^2 = \mathbb{I}.$
cardona-qft-methods_ EQ0642 (50)	259	■ 1.000	$e_{\{p_1\}} \otimes S_{\theta} e_{\{p_2\}} = \frac{\mathbb{I} \pm \tau_{\theta}}{2} e_{\{p_1\}} \otimes e_{\{p_2\}}$	$e_{p_1} \otimes S_{\theta} e_{p_2} = \frac{\mathbb{I} \pm \tau_{\theta}}{2} e_{p_1} \otimes e_{p_2}$	$e_{p_1} \otimes S_{\theta} e_{p_2} = \frac{\mathbb{I} \pm \tau_{\theta}}{2} e_{p_1} \otimes e_{p_2}$
cardona-qft-methods_ EQ0644 (48)	259	■ 1.000	$\begin{gathered} \int \prod_i d\mu(p_i) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e_{p_1} \otimes S_{\theta} e_{\Lambda p_2} \\ = \int \prod_i d\mu(p_i) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e^{ip_2 \wedge p_1} e_{\Lambda p_2} \otimes S_{\theta} e_{\Lambda p_1} \end{gathered}$	$\int \prod_i d\mu(p_i) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e_{p_1} \otimes S_{\theta} e_{\Lambda p_2} \\ = \int \prod_i d\mu(p_i) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e^{ip_2 \wedge p_1} e_{\Lambda p_2} \otimes S_{\theta} e_{\Lambda p_1}$	$\int \prod_i d\mu(p_i) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e_{p_1} \otimes S_{\theta} e_{\Lambda p_2} \\ = \int \prod_i d\mu(p_i) a_{p_1}^{\dagger} a_{p_2}^{\dagger} 0\rangle e^{ip_2 \wedge p_1} e_{\Lambda p_2} \otimes S_{\theta} e_{\Lambda p_1}$
cardona-qft-methods_ EQ0645 (52)	260	■ 1.000	$= \int \prod_i d\mu(p_i) (\exp^{ip_1 \wedge p_2} a_{p_2}^{\dagger} a_{p_1}^{\dagger}) 0\rangle e^{ip_1 \wedge p_2} e_{\Lambda p_1} \otimes S_{\theta} e_{\Lambda p_2}.$	$= \int \prod_i d\mu(p_i) (\exp^{ip_1 \wedge p_2} a_{p_2}^{\dagger} a_{p_1}^{\dagger}) 0\rangle e^{ip_1 \wedge p_2} e_{\Lambda p_1} \otimes S_{\theta} e_{\Lambda p_2}.$	$= \int \prod_i d\mu(p_i) (\exp^{ip_1 \wedge p_2} a_{p_2}^{\dagger} a_{p_1}^{\dagger}) 0\rangle e^{ip_1 \wedge p_2} e_{\Lambda p_1} \otimes S_{\theta} e_{\Lambda p_2}.$
cardona-qft-methods_ EQ0647 (54)	260	■ 1.000	$\tau_{\theta}^{ij} = \mathcal{F}_{\theta}^{-1} \tau_0^{ij} \mathcal{F} = \mathcal{F}_{\theta}^{-2} \tau_0^{ij},$	$\tau_{\theta}^{ij} = \mathcal{F}_{\theta}^{-1} \tau_0^{ij} \mathcal{F} = \mathcal{F}_{\theta}^{-2} \tau_0^{ij},$	$\tau_{\theta}^{ij} = \mathcal{F}_{\theta}^{-1} \tau_0^{ij} \mathcal{F} = \mathcal{F}_{\theta}^{-2} \tau_0^{ij},$
cardona-qft-methods_ EQ0648 (55)	260	■ 1.000	$\left(\tau_0^{ij}\right)^2 = \mathbb{I}$	$\left(\tau_0^{ij}\right)^2 = \mathbb{I}$	$\left(\tau_0^{ij}\right)^2 = \mathbb{I}$
cardona-qft-methods_ EQ0649 (56)	260	■ 1.000	$\tau_{\theta}^{i,i+1} \tau_{\theta}^{i+1,i+2} \tau_{\theta}^{i,i+1} = \tau_{\theta}^{i+1,i+2} \tau_{\theta}^{i,i+1} \tau_{\theta}^{i+1,i+2}$	$\tau_{\theta}^{i,i+1} \tau_{\theta}^{i+1,i+2} \tau_{\theta}^{i,i+1} = \tau_{\theta}^{i+1,i+2} \tau_{\theta}^{i,i+1} \tau_{\theta}^{i+1,i+2}$	$\tau_{\theta}^{i,i+1} \tau_{\theta}^{i+1,i+2} \tau_{\theta}^{i,i+1} = \tau_{\theta}^{i+1,i+2} \tau_{\theta}^{i,i+1} \tau_{\theta}^{i+1,i+2}$
cardona-qft-methods_ EQ0651 (58)	260	■ 1.000	$\varphi_{\theta}^* = \varphi_0^* e - \frac{i}{2} \overleftarrow{\partial} \wedge P$	$\varphi_{\theta}^* = \varphi_0^* e - \frac{i}{2} \overleftarrow{\partial} \wedge P$	$\varphi_{\theta}^* = \varphi_0^* e - \frac{i}{2} \overleftarrow{\partial} \wedge P.$
cardona-qft-methods_ EQ0653 (3)	261	■ 1.000	$\tau_{\theta} x_1, x_2\rangle = - x_1, x_2\rangle.$	$\tau_{\theta} x_1, x_2\rangle = - x_1, x_2\rangle.$	$\tau_{\theta} x_1, x_2\rangle = - x_1, x_2\rangle.$
cardona-qft-methods_ EQ0656 (6)	262	■ 1.000	$\langle [0 \rho_{\theta}(x), \eta_{\theta}(y)] 0 \rangle = 0.$	$\langle [0 \rho_{\theta}(x), \eta_{\theta}(y)] 0 \rangle = 0.$	$\langle [0 \rho_{\theta}(x), \eta_{\theta}(y)] 0 \rangle = 0.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0657 (7)	262	■ 1.000	$\omega_{\beta}(\rho_{\theta}(x)\eta_{\theta}(y))=\frac{\mathrm{Tr}\left[e^{-\beta H}\rho_{\theta}(x)\eta_{\theta}(y)\right]}{\mathrm{Tr}e^{-\beta H}},$	$\omega_{\beta}(\rho_{\theta}(x)\eta_{\theta}(y))=\frac{\mathrm{Tr}\left[e^{-\beta H}\rho_{\theta}(x)\eta_{\theta}(y)\right]}{\mathrm{Tr}e^{-\beta H}},$	
cardona-qft-methods_ EQ0658 (8)	262	■ 1.000	$ds^2=dt^2-a(t)^2\left[dr^2+r^2d\Omega^2\right]$	$ds^2=dt^2-a(t)^2\left[dr^2+r^2d\Omega^2\right]$	
cardona-qft-methods_ EQ0662 (13)	263	■ 1.000	$U(\Lambda)S^{(2)}U^{-1}(\Lambda)=\frac{-i^2}{2!}\int d^4xd^4y\left(\theta(x_0-y_0)[H_I(x),H_I(y)]+H_I(y)H_I(x)\right)$	$U(\Lambda)S^{(2)}U^{-1}(\Lambda)=\frac{-i^2}{2!}\int d^4xd^4y\left(\theta(x_0-y_0)[H_I(x),H_I(y)]+H_I(y)H_I(x)\right)$	
cardona-qft-methods_ EQ0663 (14)	263	■ 1.000	$\theta\left(x_0-y_0\right)\left[H_I(\Lambda x),H_I(\Lambda y)\right]=\theta\left((\Lambda x)_0-(\Lambda y)_0\right)\left[H_I(\Lambda x),H_I(\Lambda y)\right]$	$\theta(x_0-y_0)[H_I(\Lambda x),H_I(\Lambda y)]=\theta((\Lambda x)_0-(\Lambda y)_0)[H_I(\Lambda x),H_I(\Lambda y)]$	
cardona-qft-methods_ EQ0665 (16)	263	■ 1.000	$\left[H_I(\Lambda x),H_I(\Lambda y)\right]=0,\quad x\sim y,$	$[H_I(\Lambda x),H_I(\Lambda y)]=0,\quad x\sim y,$	
cardona-qft-methods_ EQ0666 (17)	263	■ 1.000	$\theta_{0i}\left(\vec{P}_{inc}\right)_i$	$\theta_{0i}\left(\vec{P}_{inc}\right)_i$	
cardona-qft-methods_ EQ0667 (18)	263	■ 1.000	$\theta_{0i}\left(\vec{P}_{inc}\right)_i=0$	$\theta_{0i}\left(\vec{P}_{inc}\right)_i=0$	
cardona-qft-methods_ EQ0668 (19)	264	■ 1.000	$\frac{\Delta T(\widehat{n})}{T}=\sum_{lm}a_{lm}Y_{lm}(\widehat{n}).$	$\frac{\Delta T(\widehat{n})}{T}=\sum_{lm}a_{lm}Y_{lm}(\widehat{n}).$	
cardona-qft-methods_ EQ0669 (20)	264	■ 1.000	$a_{lm}=4\pi(-i)^l\int\frac{d^3k}{(2\pi)^3}\Phi(k)\Delta_l^T(k)Y_{lm}^*(\widehat{k})$	$a_{lm}=4\pi(-i)^l\int\frac{d^3k}{(2\pi)^3}\Phi(k)\Delta_l^T(k)Y_{lm}^*(\widehat{k})$	
cardona-qft-methods_ EQ0671 (21)	264	■ 1.000	$P_{\Phi}=\left.\frac{8\pi GH^2}{9\varepsilon k^3}\right _{aH=k}$	$P_{\Phi}=\frac{8\pi GH^2}{9\varepsilon k^3} _{aH=k}$	
cardona-qft-methods_ EQ0672 (22)	264	■ 1.000	$\varphi(\alpha,x)=\left(\frac{\alpha}{3}\right)^{\frac{3}{2}}\int d^3x'\varphi_{\theta}(x)\exp^{-\alpha(x'-x)^2},$	$\varphi(\alpha,x)=\left(\frac{\alpha}{3}\right)^{\frac{3}{2}}\int d^3x'\varphi_{\theta}(x)\exp^{-\alpha(x'-x)^2},$	
cardona-qft-methods_ EQ0673 (23)	264	■ 1.000	$\Delta\varphi_{\theta}(\alpha,x_1)\Delta\varphi_{\theta}(\alpha,x_2)\geq\langle 0 \varphi_{\theta}(\alpha,x_1),\varphi_{\theta}(\alpha,x_2) 0\rangle$	$\Delta\varphi_{\theta}(\alpha,x_1)\Delta\varphi_{\theta}(\alpha,x_2)\geq\langle 0 \varphi_{\theta}(\alpha,x_1),\varphi_{\theta}(\alpha,x_2) 0\rangle$	
cardona-qft-methods_ EQ0675 (1)	268	■ 1.000	$\vec{S}=\frac{1}{2}\vec{\sigma},$	$\vec{S}=\frac{1}{2}\vec{\sigma},$	
cardona-qft-methods_ EQ0676 (2)	268	■ 1.000	$\sigma^1=\left(\begin{array}{cc}0&+1\\+1&0\end{array}\right),\quad\sigma^2=\left(\begin{array}{cc}0&-i\\+i&0\end{array}\right),\quad\sigma^3=\left(\begin{array}{cc}+1&0\\0&-1\end{array}\right).$	$\sigma^1=\left(\begin{array}{cc}0&+1\\+1&0\end{array}\right),\quad\sigma^2=\left(\begin{array}{cc}0&-i\\+i&0\end{array}\right),\quad\sigma^3=\left(\begin{array}{cc}+1&0\\0&-1\end{array}\right).$	
cardona-qft-methods_ EQ0677	268	■ 1.000	$\left[\sigma^i,\sigma^j\right]=2i\epsilon^{ijk}\sigma^k$	$[\sigma^i,\sigma^j]=2i\epsilon^{ijk}\sigma^k,$	
cardona-qft-methods_ EQ0679 (4)	269	■ 1.000	$s=0,\frac{1}{2},1,\frac{3}{2},\ldots,$	$s=0,\frac{1}{2},1,\frac{3}{2},\ldots,$	

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0680 (5)	269	■ 1.000	$\vec{S} \cdot \text{state} = 0,$	$\vec{S} \cdot \text{state}\rangle = 0,$	$\vec{S} \cdot \text{state}\rangle = 0,$
cardona-qft-methods_ EQ0682 (6)	269	■ 1.000	$S^{\pm} = S^1 \pm i S^2$	$S^{\pm} = S^1 \pm i S^2$	$S^{\pm} = S^1 \pm i S^2.$
cardona-qft-methods_ EQ0683	269	■ 1.000	$\left[S^3, S^{\pm}\right] = \pm S^{\pm}, \quad \left[S^{+}, S^{-}\right] = 2 S^3$	$[S^3, S^{\pm}] = \pm S^{\pm}, \quad [S^{+}, S^{-}] = 2 S^3$	$[S^3, S^{\pm}] = \pm S^{\pm}, \quad [S^{+}, S^{-}] = 2 S^3.$
cardona-qft-methods_ EQ0685	269	■ 1.000	$ \uparrow\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \downarrow\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$	$ \uparrow\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \downarrow\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$	$ \uparrow\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \downarrow\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix}.$
cardona-qft-methods_ EQ0686	269	■ 1.000	$S^3 \uparrow\rangle = +\frac{1}{2} \uparrow\rangle, \quad S^3 \downarrow\rangle = -\frac{1}{2} \downarrow\rangle,$	$S^3 \uparrow\rangle = +\frac{1}{2} \uparrow\rangle, \quad S^3 \downarrow\rangle = -\frac{1}{2} \downarrow\rangle,$	$S^3 \uparrow\rangle = +\frac{1}{2} \uparrow\rangle, \quad S^3 \downarrow\rangle = -\frac{1}{2} \downarrow\rangle,$
cardona-qft-methods_ EQ0688 (7)	270	■ 1.000	$\mathcal{H} = \mathbb{C}^2 \otimes \dots \otimes \mathbb{C}^2 = (\mathbb{C}^2)^{\otimes L}$	$\mathcal{H} = \mathbb{C}^2 \otimes \dots \otimes \mathbb{C}^2 = (\mathbb{C}^2)^{\otimes L}$	$\mathcal{H} = \mathbb{C}^2 \otimes \dots \otimes \mathbb{C}^2 = (\mathbb{C}^2)^{\otimes L}$
cardona-qft-methods_ EQ0689	270	■ 1.000	$ \uparrow\uparrow \dots \downarrow\uparrow\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \dots \otimes \begin{pmatrix} 0 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$	$ \uparrow\uparrow \dots \downarrow\uparrow\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \dots \otimes \begin{pmatrix} 0 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$	$ \uparrow\uparrow \dots \downarrow\uparrow\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \dots \otimes \begin{pmatrix} 0 \\ 1 \end{pmatrix} \otimes \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$
cardona-qft-methods_ EQ0690	270	■ 1.000	$\frac{1}{2} \otimes \frac{1}{2} = 1 \oplus 0.$	$\frac{1}{2} \otimes \frac{1}{2} = 1 \oplus 0.$	$\frac{1}{2} \otimes \frac{1}{2} = 1 \oplus 0.$
cardona-qft-methods_ EQ0691	270	■ 1.000	$\left\{ \uparrow\uparrow\rangle, \frac{1}{\sqrt{2}} \left(\uparrow\downarrow\rangle + \downarrow\uparrow\rangle \right), \frac{1}{\sqrt{2}} \left(\uparrow\downarrow\rangle - \downarrow\uparrow\rangle \right) \right\}.$	$\left\{ \uparrow\uparrow\rangle, \frac{1}{\sqrt{2}} \left(\uparrow\downarrow\rangle + \downarrow\uparrow\rangle \right), \frac{1}{\sqrt{2}} \left(\uparrow\downarrow\rangle - \downarrow\uparrow\rangle \right) \right\}.$	$\{ \uparrow\uparrow\rangle, \frac{1}{\sqrt{2}}(\uparrow\downarrow\rangle + \downarrow\uparrow\rangle), \downarrow\downarrow\rangle, \frac{1}{\sqrt{2}}(\uparrow\downarrow\rangle - \downarrow\uparrow\rangle)\}.$
cardona-qft-methods_ EQ0692 (8)	270	■ 1.000	$\hat{H} = 4 \sum_{l=1}^L \left(\frac{1}{4} - \vec{S}_l \cdot \vec{S}_{l+1} \right),$	$\hat{H} = 4 \sum_{l=1}^L \left(\frac{1}{4} - \vec{S}_l \cdot \vec{S}_{l+1} \right),$	$\hat{H} = 4 \sum_{l=1}^L \left(\frac{1}{4} - \vec{S}_l \cdot \vec{S}_{l+1} \right),$
cardona-qft-methods_ EQ0693	270	■ 1.000	$\vec{S}_{L+1} = \vec{S}_1.$	$\vec{S}_{L+1} = \vec{S}_1.$	$\vec{S}_{L+1} = \vec{S}_1.$
cardona-qft-methods_ EQ0694 (9)	271	■ 1.000	$\hat{H} \psi\rangle = E \psi\rangle, \quad \psi\rangle \in (\mathbb{C}^2)^{\otimes L},$	$\hat{H} \psi\rangle = E \psi\rangle, \quad \psi\rangle \in (\mathbb{C}^2)^{\otimes L},$	$\hat{H} \psi\rangle = E \psi\rangle, \quad \psi\rangle \in (\mathbb{C}^2)^{\otimes L},$
cardona-qft-methods_ EQ0696	271	■ 1.000	$\mathbb{P}_{l,l+1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$	$\mathbb{P}_{l,l+1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$	$\mathbb{P}_{l,l+1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$
cardona-qft-methods_ EQ0700	272	■ 1.000	$\hat{U} = \mathbb{P}_{1,2} \dots \mathbb{P}_{L-1,L}.$	$\hat{U} = \mathbb{P}_{1,2} \dots \mathbb{P}_{L-1,L}.$	$\hat{U} = \mathbb{P}_{1,2} \dots \mathbb{P}_{L-1,L}.$
cardona-qft-methods_ EQ0701 (11)	272	■ 1.000	$\hat{U}^L = \mathbb{I}.$	$\hat{U}^L = \mathbb{I}.$	$\hat{U}^L = \mathbb{I}.$
cardona-qft-methods_ EQ0702 (12)	272	■ 1.000	$U = e^{2\pi i n / L},$	$U = e^{2\pi i n / L},$	$U = e^{2\pi i n / L},$
cardona-qft-methods_ EQ0703 (13)	272	■ 1.000	$\left(\frac{u_k + \frac{i}{2}}{u_k - \frac{i}{2}} \right)^L = \prod_{j \neq k}^M \frac{u_k - u_j + i}{u_k - u_j - i}, \quad k = 1, \dots, M$	$\left(\frac{u_k + \frac{i}{2}}{u_k - \frac{i}{2}} \right)^L = \prod_{j \neq k}^M \frac{u_k - u_j + i}{u_k - u_j - i}, \quad k = 1, \dots, M$	$\left(\frac{u_k + \frac{i}{2}}{u_k - \frac{i}{2}} \right)^L = \prod_{j \neq k}^M \frac{u_k - u_j + i}{u_k - u_j - i}, \quad k = 1, \dots, M.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0704	272	■ 1.000	$E=2 \sum_{k=1}^M \frac{1}{u_k^2+\frac{1}{4}},$ $\quad U=\prod_{k=1}^M \frac{u_k+\frac{i}{2}}{u_k-\frac{i}{2}}\frac{1}{u_k^2+\frac{1}{4}},$	$E=2\sum_{k=1}^M\frac{1}{u_k^2+\frac{1}{4}},\quad U=\prod_{k=1}^M\frac{u_k+\frac{i}{2}}{u_k-\frac{i}{2}}.$	$E=2\sum_{k=1}^M\frac{1}{u_k^2+\frac{1}{4}},\quad U=\prod_{k=1}^M\frac{u_k+\frac{i}{2}}{u_k-\frac{i}{2}}.$
cardona-qft-methods_ EQ0705 (1)	272	■ 1.000	$ \psi\rangle=\sum_{1\leq l_1<\cdots< l_M\leq L} \psi\rangle\langle\psi _{l_1,\ldots,l_M}S_{l_1}^-\cdots S_{l_M}^- 0\rangle,$	$ \psi\rangle=\sum_{1\leq l_1<\cdots< l_M\leq L}\psi(l_1,\ldots,l_M)S_{l_1}^-\cdots S_{l_M}^- 0\rangle,$	$ \psi\rangle=\sum_{1\leq l_1<\cdots< l_M\leq L}\psi(l_1,\ldots,l_M)S_{l_1}^-\cdots S_{l_M}^- 0\rangle,$
cardona-qft-methods_ EQ0706 (2)	273	■ 1.000	$\psi(l_1,\ldots,l_M)=\sum_{\tau\in S_n}A(\tau)e^{ip_{\tau_1}l_1+\cdots+ip_{\tau_M}l_M},$	$\psi(l_1,\ldots,l_M)=\sum_{\tau\in S_n}A(\tau)e^{ip_{\tau_1}l_1+\cdots+ip_{\tau_M}l_M}$	$\psi(l_1,\ldots,l_M)=\sum_{\tau\in S_n}A(\tau)e^{ip_{\tau_1}l_1+\cdots+ip_{\tau_M}l_M},$
cardona-qft-methods_ EQ0707	273	■ 1.000	$A(\tau)=\operatorname{sgn}(\tau)\prod_{j<k}(e^{ip_k+ip_j}-2e^{ip_k}+1),$	$A(\tau)=\operatorname{sgn}(\tau)\prod_{j<k}(e^{ip_k+ip_j}-2e^{ip_k}+1),$	$A(\tau)=\operatorname{sgn}(\tau)\prod_{j<k}(e^{ip_k+ip_j}-2e^{ip_k}+1),$
cardona-qft-methods_ EQ0708	273	■ 1.000	$E=\sum_{k=1}^M8\sin^2\left(\frac{p_k}{2}\right).$	$E=\sum_{k=1}^M8\sin^2\left(\frac{p_k}{2}\right).$	$E=\sum_{k=1}^M8\sin^2\left(\frac{p_k}{2}\right).$
cardona-qft-methods_ EQ0710	273	■ 1.000	$S(p_k,p_j)=-\frac{e^{ip_k+ip_j}-2e^{ip_k}+1}{e^{ip_k+ip_j}-2e^{ip_j}+1}.$	$S(p_k,p_j)=-\frac{e^{ip_k+ip_j}-2e^{ip_k}+1}{e^{ip_k+ip_j}-2e^{ip_j}+1}.$	$S(p_k,p_j)=-\frac{e^{ip_k+ip_j}-2e^{ip_k}+1}{e^{ip_k+ip_j}-2e^{ip_j}+1}.$
cardona-qft-methods_ EQ0711	273	■ 1.000	$e^{ip_k}=\frac{u_k+\frac{i}{2}}{u_k-\frac{i}{2}}\iff\frac{1}{2}\cot\left(\frac{p_k}{2}\right)=u_k,$	$e^{ip_k}=\frac{u_k+\frac{i}{2}}{u_k-\frac{i}{2}}\iff\frac{1}{2}\cot\left(\frac{p_k}{2}\right)=u_k,$	$e^{ip_k}=\frac{u_k+\frac{i}{2}}{u_k-\frac{i}{2}}\iff\frac{1}{2}\cot\left(\frac{p_k}{2}\right)=u_k,$
cardona-qft-methods_ EQ0713	274	■ 1.000	$R_{a,b}(u-u')R_{a,c}(u)R_{b,c}(u')=R_{b,c}(u')R_{a,c}(u)R_{a,b}(u-u')$	$R_{a,b}(u-u')R_{a,c}(u)R_{b,c}(u')=R_{b,c}(u')R_{a,c}(u)R_{a,b}(u-u')$	$R_{a,b}(u-u')R_{a,c}(u)R_{b,c}(u')=R_{b,c}(u')R_{a,c}(u)R_{a,b}(u-u')$
cardona-qft-methods_ EQ0714	274	■ 1.000	$R_{a,b}(u-u')=$	$R_{a,b}(u-u')=$	$R_{a,b}(u-u')=$
cardona-qft-methods_ EQ0715	275	■ 1.000	$R_{a,b}(u-u')L_{a,l}(u)L_{b,l}(u')=L_{b,l}(u')L_{a,l}(u)R_{a,b}(u-u').$	$R_{a,b}(u-u')L_{a,l}(u)L_{b,l}(u')=L_{b,l}(u')L_{a,l}(u)R_{a,b}(u-u').$	$R_{a,b}(u-u')L_{a,l}(u)L_{b,l}(u')=L_{b,l}(u')L_{a,l}(u)R_{a,b}(u-u').$
cardona-qft-methods_ EQ0719	276	■ 1.000	$\hat{T}(u)\hat{T}(u')=\hat{T}(u')\hat{T}(u)$	$\hat{T}(u)\hat{T}(u')=\hat{T}(u')\hat{T}(u)$	$\hat{T}(u)\hat{T}(u')=\hat{T}(u')\hat{T}(u)$
cardona-qft-methods_ EQ0720	276	■ 1.000	$[\hat{T}(u),\hat{T}(u')]=0.$	$[\hat{T}(u),\hat{T}(u')]=0.$	$[\hat{T}(u),\hat{T}(u')]=0.$
cardona-qft-methods_ EQ0721	276	■ 1.000	$\frac{1}{i^L}\hat{T}\left(\frac{i}{2}\right)=\operatorname{tr}_a(P_{a,L}\cdots P_{a,L-1}\cdots P_{a,1})=P_{12}P_{23}\cdots P_{L-1,L}=\hat{U}=e^{i\hat{P}},$	$\frac{1}{i^L}\hat{T}\left(\frac{i}{2}\right)=\operatorname{tr}_a(P_{a,L}\cdots P_{a,L-1}\cdots P_{a,1})=P_{12}P_{23}\cdots P_{L-1,L}=\hat{U}=e^{i\hat{P}}$	$\frac{1}{i^L}\hat{T}\left(\frac{i}{2}\right)=\operatorname{tr}_a(P_{a,L}\cdots P_{a,L-1}\cdots P_{a,1})=P_{12}P_{23}\cdots P_{L-1,L}=\hat{U}=e^{i\hat{P}},$
cardona-qft-methods_ EQ0722	276	■ 1.000	$i\hat{T}(u)^{-1}\frac{\mathrm{d}}{\mathrm{d}u}\hat{T}(u)\Big _{u=\frac{i}{2}}=\sum_{l=1}^LP_{l-1,l}.$	$i\hat{T}(u)^{-1}\frac{\mathrm{d}}{\mathrm{d}u}\hat{T}(u)\Big _{u=\frac{i}{2}}=\sum_{l=1}^LP_{l-1,l}.$	$i\hat{T}(u)^{-1}\frac{\mathrm{d}}{\mathrm{d}u}\hat{T}(u)\Big _{u=\frac{i}{2}}=\sum_{l=1}^LP_{l-1,l}.$
cardona-qft-methods_ EQ0723	276	■ 1.000	$\hat{H}=2\sum_{l=1}^L(1_{l,l+1}-P_{l,l+1})=2L-2i\frac{\mathrm{d}}{\mathrm{d}u}\log\hat{T}(u)\Big _{u=\frac{i}{2}}.$	$\hat{H}=2\sum_{l=1}^L(1_{l,l+1}-P_{l,l+1})=2L-2i\frac{\mathrm{d}}{\mathrm{d}u}\log\hat{T}(u)\Big _{u=\frac{i}{2}}.$	$\hat{H}=2\sum_{l=1}^L(1_{l,l+1}-P_{l,l+1})=2L-2i\frac{\mathrm{d}}{\mathrm{d}u}\log\hat{T}(u)\Big _{u=\frac{i}{2}}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0724	276	■ 1.000	$\log \hat{T}(u)=\hat{\mathcal{Q}}_1+\left(u-\frac{i}{2}\right)\hat{\mathcal{Q}}_2+\left(u-\frac{i}{2}\right)^2\hat{\mathcal{Q}}_3+\left(u-\frac{i}{2}\right)^3\hat{\mathcal{Q}}_4+\ldots,$	$\log \hat{T}(u)=\hat{Q}_1+\left(u-\frac{i}{2}\right)\hat{Q}_2+\left(u-\frac{i}{2}\right)^2\hat{Q}_3+\left(u-\frac{i}{2}\right)^3\hat{Q}_4+\ldots,$	$\log \hat{T}(u)=\hat{Q}_1+(u-\frac{i}{2})\hat{Q}_2+(u-\frac{i}{2})^2\hat{Q}_3+(u-\frac{i}{2})^3\hat{Q}_4+\ldots,$
cardona-qft-methods_ EQ0725	276	■ 1.000	$\hat{\mathcal{Q}}_1\propto\hat{P},\quad\hat{\mathcal{Q}}_2\propto\hat{H}-2L.$	$\hat{Q}_1\propto\hat{P},\quad\hat{Q}_2\propto\hat{H}-2L.$	$\hat{Q}_1\propto\hat{P},\quad\hat{Q}_2\propto\hat{H}-2L.$
cardona-qft-methods_ EQ0726	276	■ 1.000	$\left[\hat{\mathcal{Q}}_n,\hat{\mathcal{Q}}_m\right]=0$	$\left[\hat{Q}_n,\hat{Q}_m\right]=0$	$[\hat{Q}_n,\hat{Q}_m]=0.$
cardona-qft-methods_ EQ0727	277	■ 1.000	$\hat{T}(u) \psi\rangle=T(u) \psi\rangle.$	$\hat{T}(u) \psi\rangle=T(u) \psi\rangle.$	$\hat{T}(u) \psi\rangle=T(u) \psi\rangle.$
cardona-qft-methods_ EQ0729	277	■ 1.000	$ \psi\rangle=\hat{B}(u_1)\cdots\hat{B}(u_M) 0\rangle$	$ \psi\rangle=\hat{B}(u_1)\cdots\hat{B}(u_M) 0\rangle$	$ \psi\rangle=\hat{B}(u_1)\cdots\hat{B}(u_M) 0\rangle$
cardona-qft-methods_ EQ0731	277	■ 1.000	$T(u)=\left(u+\frac{i}{2}\right)^L\prod_{j=1}^M\frac{u-u_j-i}{u-u_j}+\left(u-\frac{i}{2}\right)^L\prod_{j=1}^M\frac{u-u_j+i}{u-u_j}.$	$T(u)=\left(u+\frac{i}{2}\right)^L\prod_{j=1}^M\frac{u-u_j-i}{u-u_j}+\left(u-\frac{i}{2}\right)^L\prod_{j=1}^M\frac{u-u_j+i}{u-u_j}.$	$T(u)=(u+\frac{i}{2})^L\prod_{j=1}^M\frac{u-u_j-i}{u-u_j}+(u-\frac{i}{2})^L\prod_{j=1}^M\frac{u-u_j+i}{u-u_j}.$
cardona-qft-methods_ EQ0732	278	■ 1.000	$Q(u)=\prod_{j=1}^M(u-u_j).$	$Q(u)=\prod_{j=1}^M(u-u_j).$	$Q(u)=\prod_{j=1}^M(u-u_j).$
cardona-qft-methods_ EQ0733	278	■ 1.000	$T(u)Q(u)=\left(u+\frac{i}{2}\right)^LQ(u-i)+\left(u-\frac{i}{2}\right)^LQ(u+i).$	$T(u)Q(u)=\left(u+\frac{i}{2}\right)^LQ(u-i)+\left(u-\frac{i}{2}\right)^LQ(u+i).$	$T(u)Q(u)=(u+\frac{i}{2})^LQ(u-i)+(u-\frac{i}{2})^LQ(u+i).$
cardona-qft-methods_ EQ0737 (7)	281	■ 1.000	$[\mathcal{D},P_\mu]=iP_\mu\quad,\quad[\mathcal{D},K_\mu]=-iK_\mu$	$[\mathcal{D},P_\mu]=iP_\mu\quad,\quad[\mathcal{D},K_\mu]=-iK_\mu$	$[\mathcal{D},P_\mu]=iP_\mu\quad,\quad[\mathcal{D},K_\mu]=-iK_\mu$
cardona-qft-methods_ EQ0738	281	■ 1.000	$L=\begin{array}{ c c }\hline S^3 & S^+ \\ \hline S^- & -S^3 \\ \hline\end{array}.$	$L=\begin{array}{ c c }\hline S^3 & S^+ \\ \hline S^- & -S^3 \\ \hline\end{array}.$	$L=\begin{array}{ c c }\hline S^3 & S^+ \\ \hline S^- & -S^3 \\ \hline\end{array}.$
cardona-qft-methods_ EQ0739 (9)	282	■ 1.000	$\begin{aligned} &\left\{Q_{a\alpha},\bar{Q}_{\dot{\alpha}}^b\right\}=(\sigma^\mu)_{\alpha\dot{\alpha}}P_\mu\delta_a^b\\ &[\mathcal{D},Q_{a\alpha}]=\frac{i}{2}Q_{a\alpha} \end{aligned}$	$\begin{aligned} &\{Q_{a\alpha},\bar{Q}_{\dot{\alpha}}^b\}=(\sigma^\mu)_{\alpha\dot{\alpha}}P_\mu\delta_a^b\\ &[\mathcal{D},Q_{a\alpha}]=\frac{i}{2}Q_{a\alpha} \end{aligned}$	$\begin{aligned} &\{Q_{a\alpha},\bar{Q}_{\dot{\alpha}}^b\}=(\sigma^\mu)_{\alpha\dot{\alpha}}P_\mu\delta_a^b\\ &[\mathcal{D},Q_{a\alpha}]=\frac{i}{2}Q_{a\alpha} \end{aligned}$
cardona-qft-methods_ EQ0741	282	■ 1.000	$\mathcal{O}\equiv\mathcal{O}_{(x)}=\mathrm{Tr}\left(ZX\bar{Z}(DY)\left(D^5Y\right)\psi\left(D^3\psi\right)\bar{\psi}F\left(D^7\bar{F}\right)...\right)_{(x)}.$	$\mathcal{O}\equiv\mathcal{O}_{(x)}=\mathrm{Tr}\left(ZX\bar{Z}(DY)\left(D^5Y\right)\psi\left(D^3\psi\right)\bar{\psi}F\left(D^7\bar{F}\right)...\right)_{(x)}.$	$\mathcal{O}\equiv\mathcal{O}_{(x)}=\mathrm{Tr}\Big(ZX\bar{Z}(DY)(D^5Y)\psi(D^3\psi)\bar{\psi}F(D^7\bar{F})...\Big)_{(x)}.$
cardona-qft-methods_ EQ0742	282	■ 1.000	$[\mathcal{D},\mathcal{O}]=i\Delta\mathcal{O},$	$[\mathcal{D},\mathcal{O}]=i\Delta\mathcal{O},$	$[\mathcal{D},\mathcal{O}]=i\Delta\mathcal{O},$
cardona-qft-methods_ EQ0743	282	■ 1.000	$\gamma=g_{YM}^2\,\underbrace{\gamma_1}_{1-loop}+g_{YM}^4\,\underbrace{\gamma_2}_{2-loop}+\ldots$	$\gamma=g_{YM}^2\,\underbrace{\gamma_1}_{1-loop}+g_{YM}^4\,\underbrace{\gamma_2}_{2-loop}+\ldots$	$\gamma=g_{YM}^2\,\underbrace{\gamma_1}_{1-loop}+g_{YM}^4\,\underbrace{\gamma_2}_{2-loop}+\ldots$
cardona-qft-methods_ EQ0746	283	■ 1.000	$\langle(\bar{Z}^j)(x)(Z^j)(y)\rangle=\frac{1}{(x-y)^{2j}}$	$\langle(\bar{Z}^j)(x)(Z^j)(y)\rangle=\frac{1}{(x-y)^{2j}}$	$\langle(\bar{Z}^j)(x)(Z^j)(y)\rangle=\frac{1}{(x-y)^{2j}}$
cardona-qft-methods_ EQ0750	284	■ 1.000	$\mathfrak{psu}(2,2\mid 4)=\begin{array}{ c c }\hline \mathfrak{u}(2,2) & \text{supercharges} \\ \hline \text{supercharges} & \mathfrak{u}(4) \\ \hline\end{array}$	$\mathfrak{psu}(2,2\mid 4)=\begin{array}{ c c }\hline \mathfrak{u}(2,2) & \text{supercharges} \\ \hline \text{supercharges} & \mathfrak{u}(4) \\ \hline\end{array}$	$\mathfrak{psu}(2,2 4)=\begin{array}{ c c }\hline \mathfrak{u}(2,2) & \text{supercharges} \\ \hline \text{supercharges} & \mathfrak{u}(4) \\ \hline\end{array}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0752 (1)	284	■ 1.000	$S=\frac{R^{\{2\}}{\{\alpha^{\{\prime\}}\}}\int\mathrm{d}\,\tau\,\frac{\mathrm{d}}{\mathrm{d}\,\tau}\left[\frac{1}{2}\,\partial_aZ^M\partial^aZ_M+\frac{1}{2}\partial_aY^N\partial^aY_N\right]+\text{fermions}$	$S=\frac{R^2}{\alpha'}\int d\tau\frac{d\sigma}{2\pi}\left(\frac{1}{2}\partial_aZ^M\partial^aZ_M+\frac{1}{2}\partial_aY^N\partial^aY_N\right)+\text{fermions}$	$S=\frac{R^2}{\alpha'}\int d\tau\frac{d\sigma}{2\pi}\left(\frac{1}{2}\partial_aZ^M\partial^aZ_M+\frac{1}{2}\partial_aY^N\partial^aY_N\right)+\text{fermions},$
cardona-qft-methods_ EQ0753	285	■ 1.000	$\sqrt{\lambda}=\frac{R^2}{\alpha^{\{\prime\}}}\quad,\quad 4\pi g_s=\frac{\lambda}{N}.$	$\sqrt{\lambda}=\frac{R^2}{\alpha'}\quad,\quad 4\pi g_s=\frac{\lambda}{N}.$	$\sqrt{\lambda}=\frac{R^2}{\alpha'}\quad,\quad 4\pi g_s=\frac{\lambda}{N}.$
cardona-qft-methods_ EQ0754	285	■ 1.000	$S_{\{SYM\}}=\int\frac{N}{\lambda}\frac{d^4x}{4\pi^2}\text{Tr}\{\dots\}\longleftrightarrow S_{\sigma-\text{model}}=\sqrt{\lambda}\int d\tau\frac{d\sigma}{2\pi}\{\dots\},$	$S_{SYM}=\int\frac{N}{\lambda}\frac{d^4x}{4\pi^2}\text{Tr}\{\dots\}\longleftrightarrow S_{\sigma-\text{model}}=\sqrt{\lambda}\int d\tau\frac{d\sigma}{2\pi}\{\dots\},$	$S_{SYM}=\int\frac{N}{\lambda}\frac{d^4x}{4\pi^2}Tr\{\dots\}\longleftrightarrow S_{\sigma-model}=\sqrt{\lambda}\int d\tau\frac{d\sigma}{2\pi}\{\dots\},$
cardona-qft-methods_ EQ0755	285	■ 1.000	$Z_{\{4\}}+Z_{\{5\}}=\frac{R}{z},\quad\text{quad}\left(Z_{\{0\}},\,Z_{\{1\}},\,Z_{\{2\}},\,Z_{\{3\}}\right)=\frac{R}{z}x_{\{\mu\}},\quad\text{quad}\,\mathrm{d}\,S_{\{Ad\,S_5\}}^2=\frac{R^2}{z^2}\left(dx^\mu dx_\mu+dz^2\right).$	$Z_4+Z_5=\frac{R}{z},\quad(Z_0,Z_1,Z_2,Z_3)=\frac{R}{z}x_\mu,\quad ds_{AdS_5}^2=\frac{R^2}{z^2}\left(dx^\mu dx_\mu+dz^2\right).$	$Z_4+Z_5=\frac{R}{z},\quad(Z_0,Z_1,Z_2,Z_3)=\frac{R}{z}x_\mu,\quad ds_{AdS_5}^2=\frac{R^2}{z^2}\left(dx^\mu dx_\mu+dz^2\right).$
cardona-qft-methods_ EQ0756	286	■ 1.000	$Y_{\{1\}}+i\,Y_{\{2\}}=r_{\{1\}}\,e^{i\,\phi_{\{1\}}},\quad\text{quad}\,Y_{\{3\}}+i\,Y_{\{4\}}=r_{\{2\}}\,e^{i\,\phi_{\{2\}}},\quad\text{quad}\,Y_{\{5\}}+i\,Y_{\{6\}}=r_{\{3\}}\,e^{i\,\phi_{\{3\}}},$	$Y_1+iY_2=r_1e^{i\phi_1},\quad Y_3+iY_4=r_2e^{i\phi_2},\quad Y_5+iY_6=r_3e^{i\phi_3}$	$Y_1+iY_2=r_1e^{i\phi_1},\quad Y_3+iY_4=r_2e^{i\phi_2},\quad Y_5+iY_6=r_3e^{i\phi_3},$
cardona-qft-methods_ EQ0757	286	■ 1.000	$Z_{\{0\}}-i\,Z_{\{5\}}=\rho_{\{3\}}\,e^{i\,t},\quad\text{quad}\,Z_{\{1\}}+i\,Z_{\{2\}}=\rho_{\{1\}}\,e^{i\,\alpha_{\{1\}}},\quad\text{quad}\,Z_{\{3\}}+i\,Z_{\{4\}}=\rho_{\{2\}}\,e^{i\,\alpha_{\{2\}}},$	$Z_0-iZ_5=\rho_3e^{it},\quad Z_1+iZ_2=\rho_1e^{i\alpha_1},\quad ZY_3+iZ_4=\rho_2e^{i\alpha_2}$	$Z_0-iZ_5=\rho_3e^{it},\quad Z_1+iZ_2=\rho_1e^{i\alpha_1},\quad ZY_3+iZ_4=\rho_2e^{i\alpha_2}.$
cardona-qft-methods_ EQ0760 (1)	292	■ 1.000	$S=-\frac{1}{4\pi\alpha^{\{\prime\}}}\int\mathrm{d}\sigma\,\mathrm{d}\tau\sqrt{\gamma}\gamma^{\alpha\beta}\eta_{MN}\partial_\alpha X^M\partial_\beta X^N$	$S=-\frac{1}{4\pi\alpha'}\int\mathrm{d}\sigma\,\mathrm{d}\tau\sqrt{\gamma}\gamma^{\alpha\beta}\eta_{MN}\partial_\alpha X^M\partial_\beta X^N$	$S=-\frac{1}{4\pi\alpha'}\int\mathrm{d}\sigma\mathrm{d}\tau\sqrt{\gamma}\gamma^{\alpha\beta}\eta_{MN}\partial_\alpha X^M\partial_\beta X^N,$
cardona-qft-methods_ EQ0761 (2)	292	■ 1.000	$T=\frac{1}{2\pi\alpha^{\{\prime\}}}\frac{1}{l_s^{\{2\}}}.$	$T=\frac{1}{2\pi\alpha'}=\frac{1}{2\pi l_s^2}.$	$T=\frac{1}{2\pi\alpha'}=\frac{1}{2\pi l_s^2}.$
cardona-qft-methods_ EQ0762 (3)	293	■ 1.000	$\left(\frac{\partial^2}{\partial\sigma^2}-\frac{\partial^2}{\partial\tau^2}\right)X^M(\tau,\sigma)=0$	$\left(\frac{\partial^2}{\partial\sigma^2}-\frac{\partial^2}{\partial\tau^2}\right)X^M(\tau,\sigma)=0$	$\left(\frac{\partial^2}{\partial\sigma^2}-\frac{\partial^2}{\partial\tau^2}\right)X^M(\tau,\sigma)=0.$
cardona-qft-methods_ EQ0763 (4)	293	■ 1.000	$\frac{\partial}{\partial\sigma^+}\frac{\partial}{\partial\sigma^-}X^M(\tau,\sigma)=0$	$\frac{\partial}{\partial\sigma^+}\frac{\partial}{\partial\sigma^-}X^M(\tau,\sigma)=0$	$\frac{\partial}{\partial\sigma^+}\frac{\partial}{\partial\sigma^-}X^M(\tau,\sigma)=0,$
cardona-qft-methods_ EQ0764 (5)	293	■ 1.000	$X^{\{M\}}(\tau,\sigma)=X_{\{R\}}^{\{M\}}\left(\sigma^-\right)+X_{\{L\}}^{\{M\}}\left(\sigma^+\right).$	$X^M(\tau,\sigma)=X_R^M(\sigma^-)+X_L^M(\sigma^+).$	$X^M(\tau,\sigma)=X_R^M(\sigma^-)+X_L^M(\sigma^+).$
cardona-qft-methods_ EQ0769 (10)	294	■ 1.000	$M^{\{2\}}=\frac{2}{\alpha^{\{\prime\}}}\left(\sum_{n=1}^\infty\alpha_n\cdot\alpha_{-n}+\tilde{\alpha}_n\cdot\tilde{\alpha}_{-n}-2\right)$	$M^2=\frac{2}{\alpha'}\left(\sum_{n=1}^\infty\alpha_n\cdot\alpha_{-n}+\tilde{\alpha}_n\cdot\tilde{\alpha}_{-n}-2\right)$	$M^2=\frac{2}{\alpha'}\left(\sum_{n=1}^\infty\alpha_n\cdot\alpha_{-n}+\tilde{\alpha}_n\cdot\tilde{\alpha}_{-n}-2\right)$
cardona-qft-methods_ EQ0770 (11)	294	■ 1.000	$\hat{L}_{\{0\}}=\frac{1}{2}\left(\sum_{n=1}^\infty\alpha_n\cdot\alpha_{-n}-\tilde{\alpha}_n\cdot\tilde{\alpha}_{-n}\right)$	$\hat{L}_0=\frac{1}{2}\left(\sum_{n=1}^\infty\alpha_n\cdot\alpha_{-n}-\tilde{\alpha}_n\cdot\tilde{\alpha}_{-n}\right)$	$\hat{L}_0=\frac{1}{2}\left(\sum_{n=1}^\infty\alpha_n\cdot\alpha_{-n}-\tilde{\alpha}_n\cdot\tilde{\alpha}_{-n}\right)$
cardona-qft-methods_ EQ0771 (12)	294	■ 1.000	$\left \xi_{\{MN\}}\right\rangle\equiv\xi_M\tilde{\xi}_N\alpha_1^M\tilde{\alpha}_1^N\left 0\right\rangle$	$ \xi_{MN}\rangle\equiv\xi_M\tilde{\xi}_N\alpha_1^M\tilde{\alpha}_1^N 0\rangle$	$ \xi_{MN}\rangle\equiv\xi_M\tilde{\xi}_N\alpha_1^M\tilde{\alpha}_1^N 0\rangle$
cardona-qft-methods_ EQ0772 (13)	294	■ 1.000	$\xi_{\{MN\}}=\xi_{\{MN\}}^s+\xi_{\{t\}}\eta_{\{MN\}}+\xi_{\{MN\}}^a$	$\xi_{MN}=\xi_{MN}^s+\xi_t\eta_{MN}+\xi_{MN}^a$	$\xi_{MN}=\xi_{MN}^s+\xi_t\eta_{MN}+\xi_{MN}^a,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E00774 (15)	295	■ 1.000	$S=-\frac{1}{4}\pi\,\alpha^{\prime}\int\mathrm{d}\sigma\,\mathrm{d}\tau\sqrt{\gamma}\,\gamma^{\alpha\beta}G_{MN}(X)\partial_{\alpha}X^M\partial_{\beta}X^N$	$S=-\frac{1}{4\pi\alpha'}\int\mathrm{d}\sigma\,\mathrm{d}\tau\sqrt{\gamma}\,\gamma^{\alpha\beta}G_{MN}(X)\partial_{\alpha}X^M\partial_{\beta}X^N$	$S=-\frac{1}{4\pi\alpha'}\int\mathrm{d}\sigma\mathrm{d}\tau\sqrt{\gamma}\,\gamma^{\alpha\beta}G_{MN}(X)\partial_{\alpha}X^M\partial_{\beta}X^N\,,$
cardona-qft-methods_ E00776 (17)	295	■ 1.000	$\begin{aligned} S= & -\frac{1}{4}\pi\,\alpha^{\prime}\int\mathrm{d}\sigma\,\mathrm{d}\tau\sqrt{\gamma}\,\gamma^{\alpha\beta}G_{MN}(X)\partial_{\alpha}X^M\partial_{\beta}X^N \\ & +\epsilon^{\alpha\beta}B_{MN}(X)\partial_{\alpha}X^M\partial_{\beta}X^N \\ & +\alpha'\Phi R\end{aligned}$	$S=-\frac{1}{4\pi\alpha'}\int\mathrm{d}\sigma\,\mathrm{d}\tau\sqrt{\gamma}\left[(\gamma^{\alpha\beta}g_{MN}(X)+\mathrm{i}\epsilon^{\alpha\beta}B_{MN}(X))\partial_{\alpha}X^M\partial_{\beta}X^N+\alpha'\Phi R\right]$	$S=-\frac{1}{4\pi\alpha'}\int\mathrm{d}\sigma\mathrm{d}\tau\sqrt{\gamma}\left[(\gamma^{\alpha\beta}g_{MN}(X)+\mathrm{i}\epsilon^{\alpha\beta}B_{MN}(X))\partial_{\alpha}X^M\partial_{\beta}X^N+\alpha'\Phi R\right]\,.$
cardona-qft-methods_ E00777 (18)	296	■ 1.000	$\chi=\frac{1}{4\pi}\int\mathrm{d}^2\sigma(-\gamma)^{1/2}R\,.$	$\chi=\frac{1}{4\pi}\int\mathrm{d}^2\sigma(-\gamma)^{1/2}R.$	$\chi=\frac{1}{4\pi}\int\mathrm{d}^2\sigma(-\gamma)^{1/2}R\,.$
cardona-qft-methods_ E00782 (23)	298	■ 1.000	$M^2=\frac{1}{\alpha'}\left(\sum_{n=1}^{\infty}\alpha_n\cdot\alpha_{-n}-1\right)\,.$	$M^2=\frac{1}{\alpha'}\left(\sum_{n=1}^{\infty}\alpha_n\cdot\alpha_{-n}-1\right)\,.$	$M^2=\frac{1}{\alpha'}\left(\sum_{n=1}^{\infty}\alpha_n\cdot\alpha_{-n}-1\right)\,.$
cardona-qft-methods_ E00783 (24)	298	■ 1.000	$\left \xi_M\right\rangle\equiv\xi_M\alpha_1^M\left 0\right\rangle$	$ \xi_M\rangle\equiv\xi_M\alpha_1^M 0\rangle$	$ \xi_M\rangle\equiv\xi_M\alpha_1^M 0\rangle$
cardona-qft-methods_ E00784 (25)	298	■ 1.000	$\int_{\partial M}d\tau A_M\partial_\tau X^M$	$\int_{\partial M}d\tau A_M\partial_\tau X^M$	$\int_{\partial M}d\tau A_M\partial_\tau X^M$
cardona-qft-methods_ E00785 (26)	298	■ 1.000	$S=-\frac{1}{4}\int\mathrm{d}^D X\,e^{-\Phi}F_{MN}F^{MN}+O(\alpha')$	$S=-\frac{1}{4}\int\mathrm{d}^D X\,e^{-\Phi}F_{MN}F^{MN}+O(\alpha')$	$S=-\frac{1}{4}\int\mathrm{d}^D X\,e^{-\Phi}F_{MN}F^{MN}+O(\alpha')\,,$
cardona-qft-methods_ E00786 (27)	299	■ 1.000	$S=\frac{1}{4\pi}\int_M\mathrm{d}^2\sigma\eta_{MN}\left[\frac{1}{\alpha'}\partial X^M\bar{\partial}X^N+\Psi^M\bar{\partial}\Psi^N+\tilde{\Psi}^M\bar{\partial}\tilde{\Psi}^N\right]$	$S=\frac{1}{4\pi}\int_M\mathrm{d}^2\sigma\eta_{MN}\left[\frac{1}{\alpha'}\partial X^M\bar{\partial}X^N+\Psi^M\bar{\partial}\Psi^N+\tilde{\Psi}^M\bar{\partial}\tilde{\Psi}^N\right]$	$S=\frac{1}{4\pi}\int_M\mathrm{d}^2\sigma\eta_{MN}\left[\frac{1}{\alpha'}\partial X^M\bar{\partial}X^N+\Psi^M\bar{\partial}\Psi^N+\tilde{\Psi}^M\bar{\partial}\tilde{\Psi}^N\right]$
cardona-qft-methods_ E00788 (29)	299	■ 1.000	$\left\{\psi_r^M,\psi_s^N\right\}=\left\{\tilde{\psi}_r^M,\tilde{\psi}_s^N\right\}=\eta^{MN}\delta_{r+s}\,.$	$\left\{\psi_r^M,\psi_s^N\right\}=\left\{\tilde{\psi}_r^M,\tilde{\psi}_s^N\right\}=\eta^{MN}\delta_{r+s}\,.$	$\{\psi_r^M,\psi_s^N\}=\{\tilde{\psi}_r^M,\tilde{\psi}_s^N\}=\eta^{MN}\delta_{r+s}\,.$
cardona-qft-methods_ E00789 (30)	300	■ 1.000	$M^2=\frac{1}{\alpha'}\left(\sum_{n=1}^{\infty}\alpha_n\cdot\alpha_{-n}+\sum_r r\psi_r\cdot\psi_{-r}-a+\text{ same with tilde }\right),$	$M^2=\frac{1}{\alpha'}\left(\sum_{n=1}^{\infty}\alpha_n\cdot\alpha_{-n}+\sum_r r\psi_r\cdot\psi_{-r}-a+\text{ same with tilde }\right),$	$M^2=\frac{1}{\alpha'}\left(\sum_{n=1}^{\infty}\alpha_n\cdot\alpha_{-n}+\sum_r r\psi_r\cdot\psi_{-r}-a+\text{ same with tilde }\right),$
cardona-qft-methods_ E00790 (31)	300	■ 1.000	$\tilde{L}_0=\left(\sum_{n=1}^{\infty}\alpha_n\cdot\alpha_{-n}+\sum_r r\psi_r\cdot\psi_{-r}-a-\text{ same with tilde }\right)\,.$	$\tilde{L}_0=\left(\sum_{n=1}^{\infty}\alpha_n\cdot\alpha_{-n}+\sum_r r\psi_r\cdot\psi_{-r}-a-\text{ same with tilde }\right)\,.$	$\tilde{L}_0=\left(\sum_{n=1}^{\infty}\alpha_n\cdot\alpha_{-n}+\sum_r r\psi_r\cdot\psi_{-r}-a-\text{ same with tilde }\right)\,.$
cardona-qft-methods_ E00792 (33)	300	■ 1.000	$R:\left \psi_0^M\right 0\rangle\rightarrow\left s\right\rangle=\left \pm\frac{1}{2},\pm\frac{1}{2},\pm\frac{1}{2},\pm\frac{1}{2}\right\rangle$	$R:\psi_0^M 0\rangle\rightarrow s\rangle=\left \pm\frac{1}{2},\pm\frac{1}{2},\pm\frac{1}{2},\pm\frac{1}{2}\right\rangle$	$R:\psi_0^M 0\rangle\rightarrow s\rangle=\left \pm\frac{1}{2},\pm\frac{1}{2},\pm\frac{1}{2},\pm\frac{1}{2}\right\rangle$
cardona-qft-methods_ E00793 (34)	301	■ 1.000	$NS:\left \psi_{1/2}^M\right 0\rangle,$	$NS:\psi_{1/2}^M 0\rangle,$	$NS:\psi_{1/2}^M 0\rangle\,,$
cardona-qft-methods_ E00797 (38)	302	■ 1.000	$M^2=\frac{n^2}{R^2}+\frac{m^2R^2}{\alpha'^2}+\frac{2}{\alpha'}(N+\tilde{N}-2)$	$M^2=\frac{n^2}{R^2}+\frac{m^2R^2}{\alpha'^2}+\frac{2}{\alpha'}(N+\tilde{N}-2)$	$M^2=\frac{n^2}{R^2}+\frac{m^2R^2}{\alpha'^2}+\frac{2}{\alpha'}(N+\tilde{N}-2)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E08798 (39)	304	■ 1.000	$S_{\{D\ B\ I\}}=-T_{\{p\}}\int\mathrm{d}^p\mathrm{d}^{p+1}\xi\mathrm{e}^{-\Phi}\sqrt{\mathrm{det}(G_{ab}+B_{ab}+2\pi\alpha'F_{ab})}\sqrt{\mathrm{det}(G_{ab}+B_{ab}+2\pi\alpha'F_{ab})}\mathrm{d}^p\mathrm{d}^{p+1}\xi\mathrm{e}^{-\Phi}\sqrt{\mathrm{det}(G_{ab}+B_{ab}+2\pi\alpha'F_{ab})}$	$S_{DBI}=-T_p\int\mathrm{d}^{p+1}\xi e^{-\Phi}\sqrt{\det(G_{ab}+B_{ab}+2\pi\alpha'F_{ab})}$	$S_{DBI}=-T_p\int\mathrm{d}^{p+1}\xi e^{-\Phi}\sqrt{\det(G_{ab}+B_{ab}+2\pi\alpha'F_{ab})}$
cardona-qft-methods_ E08799 (40)	304	■ 1.000	$S_{\{C\ S\}}=\mathrm{i}T_{\{p\}}\int_{p+1}\mathrm{e}^{2\pi\alpha'F+B}\wedge C$	$S_{CS}=\mathrm{i}T_p\int_{p+1}e^{2\pi\alpha'F+B}\wedge C$	$S_{CS}=\mathrm{i}T_p\int_{p+1}e^{2\pi\alpha'F+B}\wedge C$
cardona-qft-methods_ E08800 (41)	304	■ 1.000	$G_{\{a\ b\}}(\xi)=\frac{\partial X^M}{\partial \xi^a}\frac{\partial X^N}{\partial \xi^b}G_{MN}(X)$	$G_{ab}(\xi)=\frac{\partial X^M}{\partial \xi^a}\frac{\partial X^N}{\partial \xi^b}G_{MN}(X)$	$G_{ab}(\xi)=\frac{\partial X^M}{\partial \xi^a}\frac{\partial X^N}{\partial \xi^b}G_{MN}(X)$
cardona-qft-methods_ E08801 (1)	305	■ 1.000	$\mathcal{G}=\left(\begin{array}{ll}0&1\\1&0\end{array}\right)$	$\mathcal{G}=\left(\begin{array}{cc}0&1\\1&0\end{array}\right)$	$\mathcal{G}=\left(\begin{array}{cc}0&1\\1&0\end{array}\right)$
cardona-qft-methods_ E08802 (2)	306	■ 1.000	$\mathcal{B}=\left(\begin{array}{ll}0&0\\B&0\end{array}\right),$	$\mathcal{B}=\left(\begin{array}{cc}0&0\\B&0\end{array}\right),$	$\mathcal{B}=\left(\begin{array}{cc}0&0\\B&0\end{array}\right),$
cardona-qft-methods_ E08805 (5)	306	■ 1.000	$\Lambda_{\{\alpha\beta\}}+\Lambda_{\{\beta\gamma\}}+\Lambda_{\{\gamma\alpha\}}=g_{(\alpha\beta\gamma)}\mathrm{d}g_{(\alpha\beta\gamma)}$	$\Lambda_{(\alpha\beta)}+\Lambda_{(\beta\gamma)}+\Lambda_{(\gamma\alpha)}=g_{(\alpha\beta\gamma)}\mathrm{d}g_{(\alpha\beta\gamma)}$	$\Lambda_{(\alpha\beta)}+\Lambda_{(\beta\gamma)}+\Lambda_{(\gamma\alpha)}=g_{(\alpha\beta\gamma)}\mathrm{d}g_{(\alpha\beta\gamma)}$
cardona-qft-methods_ E08806 (6)	306	■ 1.000	$\mathcal{J}_1\equiv\left(\begin{array}{cc}I&0\\0&-I^t\end{array}\right)$	$\mathcal{J}_1\equiv\left(\begin{array}{cc}I&0\\0&-I^t\end{array}\right)$	$\mathcal{J}_1\equiv\left(\begin{array}{cc}I&0\\0&-I^t\end{array}\right),$
cardona-qft-methods_ E08807 (7)	307	■ 1.000	$\mathcal{J}_2\equiv\left(\begin{array}{cc}0&-J^{-1}\\J&0\end{array}\right).$	$\mathcal{J}_2\equiv\left(\begin{array}{cc}0&-J^{-1}\\J&0\end{array}\right).$	$\mathcal{J}_2\equiv\left(\begin{array}{cc}0&-J^{-1}\\J&0\end{array}\right).$
cardona-qft-methods_ E08810 (10)	308	■ 1.000	$\mathcal{H}=\mathcal{G}\mathcal{J}_1\mathcal{J}_2.$	$\mathcal{H}=\mathcal{G}\mathcal{J}_1\mathcal{J}_2.$	$\mathcal{H}=\mathcal{G}\mathcal{J}_1\mathcal{J}_2.$
cardona-qft-methods_ E08812 (12)	308	■ 1.000	$\eta_{+}^1\otimes\eta_{\pm}^{2\dagger}=\frac{1}{4}\sum_{k=0}^6\frac{1}{k!}\eta_{\pm}^{2\dagger}\gamma_{i_1\dots i_k}\eta_{+}^1\gamma^{i_k\dots i_1}$	$\eta_{+}^1\otimes\eta_{\pm}^{2\dagger}=\frac{1}{4}\sum_{k=0}^6\frac{1}{k!}\eta_{\pm}^{2\dagger}\gamma_{i_1\dots i_k}\eta_{+}^1\gamma^{i_k\dots i_1}$	$\eta_{+}^1\otimes\eta_{\pm}^{2\dagger}=\frac{1}{4}\sum_{k=0}^6\frac{1}{k!}\eta_{\pm}^{2\dagger}\gamma_{i_1\dots i_k}\eta_{+}^1\gamma^{i_k\dots i_1}$
cardona-qft-methods_ E08813 (13)	308	■ 1.000	$\sum_k\frac{1}{k!}C_{i_1\dots i_k}^{(k)}\gamma_{\alpha\beta}^{i_i\dots i_k}\longleftrightarrow\sum_k\frac{1}{k!}C_{i_1\dots i_k}^{(k)}\mathrm{d}x^{i_i}\wedge\dots\wedge\mathrm{d}x^{i_k}$	$\sum_k\frac{1}{k!}C_{i_1\dots i_k}^{(k)}\gamma_{\alpha\beta}^{i_i\dots i_k}\longleftrightarrow\sum_k\frac{1}{k!}C_{i_1\dots i_k}^{(k)}\mathrm{d}x^{i_i}\wedge\dots\wedge\mathrm{d}x^{i_k}$	$\sum_k\frac{1}{k!}C_{i_1\dots i_k}^{(k)}\gamma_{\alpha\beta}^{i_i\dots i_k}\longleftrightarrow\sum_k\frac{1}{k!}C_{i_1\dots i_k}^{(k)}\mathrm{d}x^{i_i}\wedge\dots\wedge\mathrm{d}x^{i_k}.$
cardona-qft-methods_ E08814 (14)	309	■ 1.000	$\Phi_{\pm}\rightarrow e^{-B}\Phi_{\pm}\equiv\Phi_{\pm}^D$	$\Phi_{\pm}\rightarrow e^{-B}\Phi_{\pm}\equiv\Phi_{\pm}^D$	$\Phi_{\pm}\rightarrow e^{-B}\Phi_{\pm}\equiv\Phi_{\pm}^D$
cardona-qft-methods_ E08815 (15)	309	■ 1.000	$\Phi_{+}=e^{-\mathrm{i}J},\quad\Phi_{-}=-\mathrm{i}\Omega,$	$\Phi_{+}=e^{-\mathrm{i}J},\quad\Phi_{-}=-\mathrm{i}\Omega,$	$\Phi_{+}=e^{-\mathrm{i}J},\quad\Phi_{-}=-\mathrm{i}\Omega,$
cardona-qft-methods_ E08817 (17)	309	■ 1.000	$J\wedge\Omega=0,\quad J\wedge J\wedge J=\frac{3\mathrm{i}}{4}\Omega\wedge\bar{\Omega},$	$J\wedge\Omega=0,\quad J\wedge J\wedge J=\frac{3\mathrm{i}}{4}\Omega\wedge\bar{\Omega},$	$J\wedge\Omega=0,\quad J\wedge J\wedge J=\frac{3\mathrm{i}}{4}\Omega\wedge\bar{\Omega},$
cardona-qft-methods_ E08821 (21)	311	■ 1.000	$[X+\zeta,Y+\eta]_H=[X+\zeta,Y+\eta]_C+\iota_X\iota_YH,$	$[X+\zeta,Y+\eta]_H=[X+\zeta,Y+\eta]_C+\iota_X\iota_YH,$	$[X+\zeta,Y+\eta]_H=[X+\zeta,Y+\eta]_C+\iota_X\iota_YH,$
cardona-qft-methods_ E08822	311	■ 1.000	$(\mathrm{d}-H\wedge)\Phi=0$	$(\mathrm{d}-H\wedge)\Phi=0$	$(\mathrm{d}-H\wedge)\Phi=0$
cardona-qft-methods_ E08825 (4)	313	■ 1.000	$F^{(10)}=\mathrm{d}C-H\wedge C+m\mathrm{e}^B=\hat{F}-H\wedge C$	$F^{(10)}=\mathrm{d}C-H\wedge C+m\mathrm{e}^B=\hat{F}-H\wedge C$	$F^{(10)}=\mathrm{d}C-H\wedge C+m\mathrm{e}^B=\hat{F}-H\wedge C$
cardona-qft-methods_ E08826 (5)	313	■ 1.000	$F_n^{(10)}=(-1)^{\mathrm{Int}[n/2]}\star F_{10-n}^{(10)},$	$F_n^{(10)}=(-1)^{\mathrm{Int}[n/2]}\star F_{10-n}^{(10)},$	$F_n^{(10)}=(-1)^{\mathrm{Int}[n/2]}\star F_{10-n}^{(10)},$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ E08827 (6)	314	■ 1.000	$d\,H=0,\quad d\,F^{\{10\}}-H\,\wedge\,F^{\{10\}}=0\quad.$	$dH=0,\quad dF^{(10)}-H\wedge F^{(10)}=0.$	$dH=0\,,\quad dF^{(10)}-H\wedge F^{(10)}=0\,.$
cardona-qft-methods_ E08828 (7)	314	■ 1.000	$\tilde{\nabla}_{\mu}\epsilon+\frac{e^{-A}}{2}\left(\gamma_{\mu}\gamma_5\otimes\not{\nabla}A\right)\epsilon=0$	$\tilde{\nabla}_{\mu}\epsilon+\frac{e^{-A}}{2}\left(\gamma_{\mu}\gamma_5\otimes\not{\nabla}A\right)\epsilon=0$	$\tilde{\nabla}_{\mu}\epsilon+\frac{e^{-A}}{2}(\gamma_{\mu}\gamma_5\otimes\nabla A)\epsilon=0$
cardona-qft-methods_ E08829 (8)	314	■ 1.000	$\left[\tilde{\nabla}_{\mu},\tilde{\nabla}_{\nu}\right]\epsilon=-\frac{1}{2}\left(\nabla_mA\right)\left(\nabla^mA\right)\gamma_{\mu\nu}\epsilon$	$\left[\tilde{\nabla}_{\mu},\tilde{\nabla}_{\nu}\right]\epsilon=-\frac{1}{2}\left(\nabla_mA\right)\left(\nabla^mA\right)\gamma_{\mu\nu}\epsilon$	$[\tilde{\nabla}_{\mu},\tilde{\nabla}_{\nu}]\epsilon=-\frac{1}{2}(\nabla_mA)(\nabla^mA)\gamma_{\mu\nu}\epsilon$
cardona-qft-methods_ E08830 (9)	314	■ 1.000	$\left[\tilde{\nabla}_{\mu},\tilde{\nabla}_{\nu}\right]\epsilon=\frac{1}{4}\tilde{R}_{\mu\nu\lambda\rho}\gamma^{\lambda\rho}\epsilon=0$	$\left[\tilde{\nabla}_{\mu},\tilde{\nabla}_{\nu}\right]\epsilon=\frac{1}{4}\tilde{R}_{\mu\nu\lambda\rho}\gamma^{\lambda\rho}\epsilon=0$	$[\tilde{\nabla}_{\mu},\tilde{\nabla}_{\nu}]\epsilon=\frac{1}{4}\tilde{R}_{\mu\nu\lambda\rho}\gamma^{\lambda\rho}\epsilon=0$
cardona-qft-methods_ E08832 (11)	314	■ 1.000	$\epsilon_{\mathrm{IIB}}^A=\xi_+^A\otimes\eta_+^A+\xi_-^A\otimes\eta_-^A,\quad\epsilon_{\eta_+^A+\xi_+^A+\xi_-^A}=\epsilon_{\eta_+^A+\xi_-^A},\quad A=1,2\quad.$	$\epsilon_{\mathrm{IIB}}^A=\xi_+^A\otimes\eta_+^A+\xi_-^A\otimes\eta_-^A,\quad A=1,2.$	$\epsilon_{\mathrm{IIB}}^A=\xi_+^A\otimes\eta_+^A+\xi_-^A\otimes\eta_-^A,\quad A=1,2\,.$
cardona-qft-methods_ E08833 (12)	314	■ 1.000	$\nabla_m\eta_{\pm}^A=0\quad.$	$\nabla_m\eta_{\pm}^A=0.$	$\nabla_m\eta_{\pm}^A=0\,.$
cardona-qft-methods_ E08836 (15)	317	■ 1.000	$ \Phi_+ = \Phi_- =e^A$	$ \Phi_+ = \Phi_- =e^A$	$ \Phi_+ = \Phi_- =e^A$
cardona-qft-methods_ E08839 (18)	317	■ 1.000	$\lambda\left(A_n\right)=\left(-1\right)^{\mathrm{Int}\left[n/2\right]}A_n\,.$	$\lambda\left(A_n\right)=\left(-1\right)^{\mathrm{Int}\left[n/2\right]}A_n.$	$\lambda(A_n)=\left(-1\right)^{\mathrm{Int}\left[n/2\right]}A_n\,.$
cardona-qft-methods_ E08840	328	■ 1.000	$\{f,g\}=\Pi(d\,f,d\,g)$	$\{f,g\}=\Pi(df,dg)$	$\{f,g\}=\Pi(df,dg)$
cardona-qft-methods_ E08841	328	■ 1.000	$\{f,g\,h\}=g\{f,h\}+h\{f,g\},\,\forall f,g,h\in\mathcal{C}^{\infty}(M)\,.$	$\{f,gh\}=g\{f,h\}+h\{f,g\},\forall f,g,h\in\mathcal{C}^{\infty}(M).$	$\{f,gh\}=g\{f,h\}+h\{f,g\},\forall f,g,h\in\mathcal{C}^{\infty}(M).$
cardona-qft-methods_ E08843	329	■ 1.000	$S(X,\eta):=\int_{\Sigma}\langle\eta,dX\rangle+\frac{1}{2}\langle\eta,(\Pi^{\#}\circ X)\eta\rangle$	$S(X,\eta):=\int_{\Sigma}\langle\eta,dX\rangle+\frac{1}{2}\langle\eta,(\Pi^{\#}\circ X)\eta\rangle$	$S(X,\eta):=\int_{\Sigma}\langle\eta,dX\rangle+\frac{1}{2}\langle\eta,(\Pi^{\#}\circ X)\eta\rangle,$
cardona-qft-methods_ E08844	329	■ 1.000	$\mathrm{EL}=\{\text{ Solutions of the Euler-Lagrange equations }\}\subset\Phi,$	$\mathrm{EL}=\{\text{ Solutions of the Euler-Lagrange equations }\}\subset\Phi,$	$\mathrm{EL}=\{\text{Solutions of the Euler-Lagrange equations}\}\subset\Phi,$
cardona-qft-methods_ E08847	330	■ 1.000	$\Phi_{\partial}:=\{\text{ vector bundle morphisms between }T(\partial\Sigma)\text{ and }T^*M\}\,.$	$\Phi_{\partial}:=\{\text{ vector bundle morphisms between }T(\partial\Sigma)\text{ and }T^*M\}\,.$	$\Phi_{\partial}:=\{\text{vector bundle morphisms between }T(\partial\Sigma)\text{ and }T^*M\}.$
cardona-qft-methods_ E08848	330	■ 1.000	$L_{\Sigma}:=p(E\,L)\,.$	$L_{\Sigma}:=p(EL).$	$L_{\Sigma}:=p(EL).$
cardona-qft-methods_ E08849	330	■ 1.000	$C_{\Pi}:=\{(X,\eta)\, \,dX=\pi^{\#}(X)\eta,X:\partial\Sigma\rightarrow M,\eta\in\Gamma(T^*I\otimes X^*(T^*M))\}\,.$	$C_{\Pi}:=\{(X,\eta)\, \,dX=\pi^{\#}(X)\eta,X:\partial\Sigma\rightarrow M,\eta\in\Gamma(T^*I\otimes X^*(T^*M))\}\,.$	$C_{\Pi}:=\{(X,\eta) dX=\pi^{\#}(X)\eta,\,X:\partial\Sigma\rightarrow M,\,\eta\in\Gamma(T^*I\otimes X^*(T^*M))\}.$
cardona-qft-methods_ E08850	330	■ 1.000	$[X,f\,Y]_A=f[X,Y]+\rho_*(X)(f)Y,\,\forall X,Y\in\Gamma(A),\,f\in\mathcal{C}^{\infty}(M)\,.$	$[X,fY]_A=f[X,Y]+\rho_*(X)(f)Y,\forall X,Y\in\Gamma(A),f\in\mathcal{C}^{\infty}(M).$	$[X,fY]_A=f[X,Y]+\rho_*(X)(f)Y,\forall X,Y\in\Gamma(A),f\in\mathcal{C}^{\infty}(M).$
cardona-qft-methods_ E08852	331	■ 1.000	$\delta_A f:=\rho^*df,$	$\delta_A f:=\rho^*df,$	$\delta_A f:=\rho^*df,$
cardona-qft-methods_ E08853	331	■ 1.000	$\langle\delta_A\alpha,X\wedge Y\rangle:=-\langle\alpha,[X,Y]_A\rangle+\langle\delta\langle\alpha,X\rangle,Y\rangle-\langle\delta\langle\alpha,Y\rangle,X\rangle,$	$\langle\delta_A\alpha,X\wedge Y\rangle:=-\langle\alpha,[X,Y]_A\rangle+\langle\delta\langle\alpha,X\rangle,Y\rangle-\langle\delta\langle\alpha,Y\rangle,X\rangle,$	$\langle\delta_A\alpha,X\wedge Y\rangle:=-\langle\alpha,[X,Y]_A\rangle+\langle\delta\langle\alpha,X\rangle,Y\rangle-\langle\delta\langle\alpha,Y\rangle,X\rangle,$
cardona-qft-methods_ E08854	331	■ 1.000	$\delta_A\varphi^*=\varphi^*\delta_B\,.$	$\delta_A\varphi^*=\varphi^*\delta_B.$	$\delta_A\varphi^*=\varphi^*\delta_B.$
cardona-qft-methods_ E08856	332	■ 1.000	$G_{r_{\mu}}:=\{(a,b,c)\in G^3\, \,c=\mu(a,b)\}$	$G_{r_{\mu}}:=\{(a,b,c)\in G^3\, \,c=\mu(a,b)\}$	$Gr_{\mu}:=\{(a,b,c)\in G^3 c=\mu(a,b)\}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0878	339	■ 1.000	$\langle a, b \rangle_H := \langle \bar{a}, b \rangle,$	$\langle a, b \rangle_H := \langle \bar{a}, b \rangle,$	$\langle a, b \rangle_H := \langle \bar{a}, b \rangle,$
cardona-qft-methods_ EQ0879	339	■ 1.000	$\langle m(a, b), c \rangle_H = \langle a, m(b, c) \rangle_H$	$\langle m(a, b), c \rangle_H = \langle a, m(b, c) \rangle_H$	$\langle m(a, b), c \rangle_H = \langle a, m(b, c) \rangle_H$
cardona-qft-methods_ EQ0880	339	■ 1.000	$e := \sum_i m(e_i, \overline{e_i})$	$e := \sum_i m(e_i, \overline{e_i})$	$e := \sum_i m(e_i, \overline{e_i})$
cardona-qft-methods_ EQ0881	339	■ 1.000	$\begin{aligned} P: H &\rightarrow H \\ a &\mapsto m(a, e) \end{aligned}$	$\begin{aligned} P: H &\rightarrow H \\ a &\mapsto m(a, e) \end{aligned}$	$\begin{aligned} P: H &\rightarrow H \\ a &\mapsto m(a, e) \end{aligned}$
cardona-qft-methods_ EQ0882	342	■ 1.000	$H_p(LM) \otimes H_q(LM) \rightarrow H_{p+q-n}(LM),$	$H_p(LM) \otimes H_q(LM) \rightarrow H_{p+q-n}(LM),$	$H_p(LM) \otimes H_q(LM) \rightarrow H_{p+q-n}(LM),$
cardona-qft-methods_ EQ0885	344	■ 1.000	$\tau_\Delta: M \times M \rightarrow (M \times M) / \nu_\Delta^c$	$\tau_\Delta: M \times M \rightarrow (M \times M) / \nu_\Delta^c$	$\tau_\Delta: M \times M \rightarrow (M \times M) / \nu_\Delta^c$
cardona-qft-methods_ EQ0886	344	■ 1.000	$\tau_\Delta: M \times M \rightarrow M^{TM}$	$\tau_\Delta: M \times M \rightarrow M^{TM}$	$\tau_\Delta: M \times M \rightarrow M^{TM}$
cardona-qft-methods_ EQ0891	345	■ 1.000	$LM^{-TM} \wedge LM^{-TM} \rightarrow LM^{-TM},$	$LM^{-TM} \wedge LM^{-TM} \rightarrow LM^{-TM},$	$LM^{-TM} \wedge LM^{-TM} \rightarrow LM^{-TM},$
cardona-qft-methods_ EQ0892	346	■ 1.000	$H_p(M^{-TM}) \cong H_{p+n}(M)$	$H_p(M^{-TM}) \cong H_{p+n}(M)$	$H_p(M^{-TM}) \cong H_{p+n}(M)$
cardona-qft-methods_ EQ0893	346	■ 1.000	$M^{-TM} := \Sigma^{-k} M^{\nu(e)}$	$M^{-TM} := \Sigma^{-k} M^{\nu(e)}$	$M^{-TM} := \Sigma^{-k} M^{\nu(e)}$
cardona-qft-methods_ EQ0894	346	■ 1.000	$\Delta^*: M^{-TM} \wedge M^{-TM} = (M \times M)^{-2TM} \rightarrow M^{TM-2TM} = M^{-TM}$	$\Delta^*: M^{-TM} \wedge M^{-TM} = (M \times M)^{-2TM} \rightarrow M^{TM-2TM} = M^{-TM}$	$\Delta^*: M^{-TM} \wedge M^{-TM} = (M \times M)^{-2TM} \rightarrow M^{TM-2TM} = M^{-TM}$
cardona-qft-methods_ EQ0901	347	■ 1.000	$H_{*+n}(LM) \cong HH^*(C^*(M), C^*(M)).$	$H_{*+n}(LM) \cong HH^*(C^*(M), C^*(M)).$	$H_{*+n}(LM) \cong HH^*(C^*(M), C^*(M)).$
cardona-qft-methods_ EQ0902	348	■ 1.000	$G_x = \{g \in G \mid g\bar{x} = \bar{x}\}$	$G_x = \{g \in G \mid g\bar{x} = \bar{x}\}$	$G_x = \{g \in G \mid g\bar{x} = \bar{x}\}$
cardona-qft-methods_ EQ0903	349	■ 1.000	$P_G M \times_G E G \simeq L(M \times_G E G)$	$P_G M \times_G E G \simeq L(M \times_G E G).$	$P_G M \times_G E G \simeq L(M \times_G E G).$
cardona-qft-methods_ EQ0904	349	■ 1.000	$H_*(\mathcal{L}(M \times_G E G); \mathbb{Q}) \cong H_*(P_G M; \mathbb{Q})^G$	$H_*(\mathcal{L}(M \times_G E G); \mathbb{Q}) \cong H_*(P_G M; \mathbb{Q})^G$	$H_*(\mathcal{L}(M \times_G E G); \mathbb{Q}) \cong H_*(P_G M; \mathbb{Q})^G$
cardona-qft-methods_ EQ0905	349	■ 1.000	$P_G X := \bigsqcup_{g \in G} P_g X \times \{g\} \quad \text{with} \quad P_g X := \{f: [0, 1] \rightarrow X \mid f(0)g = f(1)\}$	$P_G X := \bigsqcup_{g \in G} P_g X \times \{g\} \quad \text{with} \quad P_g X := \{f: [0, 1] \rightarrow X \mid f(0)g = f(1)\}$	$P_G X := \bigsqcup_{g \in G} P_g X \times \{g\} \quad \text{with} \quad P_g X := \{f: [0, 1] \rightarrow X \mid f(0)g = f(1)\}$
cardona-qft-methods_ EQ0907	350	■ 1.000	$P_g M := \{(\phi, \psi) \in P_g M \times P_h M \mid \phi(1) = \psi(0)\}$	$P_g M := \{(\phi, \psi) \in P_g M \times P_h M \mid \phi(1) = \psi(0)\}$	$P_g M := \{(\phi, \psi) \in P_g M \times P_h M \mid \phi(1) = \psi(0)\}$
cardona-qft-methods_ EQ0909	350	■ 1.000	$P_G M^{-TM} := \bigsqcup_{g \in G} P_g M^{-TM}$	$P_G M^{-TM} := \bigsqcup_{g \in G} P_g M^{-TM}$	$P_G M^{-TM} := \bigsqcup_{g \in G} P_g M^{-TM}$
cardona-qft-methods_ EQ0910	350	■ 1.000	$\mu: P_G M^{-TM} \wedge P_G M^{-TM} \rightarrow P_G M^{-TM}.$	$\mu: P_G M^{-TM} \wedge P_G M^{-TM} \rightarrow P_G M^{-TM}.$	$\mu: P_G M^{-TM} \wedge P_G M^{-TM} \rightarrow P_G M^{-TM}.$
cardona-qft-methods_ EQ0911	350	■ 1.000	$H_*(P_G M^{-TM}; \mathbb{Q})^G \cong H_{*+n}(L(M \times_G E G); \mathbb{Q})$	$H_*(P_G M^{-TM}; \mathbb{Q})^G \cong H_{*+n}(L(M \times_G E G); \mathbb{Q})$	$H_*(P_G M^{-TM}; \mathbb{Q})^G \cong H_{*+n}(L(M \times_G E G); \mathbb{Q})$
cardona-qft-methods_ EQ0912	351	■ 1.000	$HH^*(C^*(M; \mathbb{Q}) \# G, C^*(M; \mathbb{Q}) \# G) \cong H_*(P_G M^{-TM}; \mathbb{Q})^G$	$HH^*(C^*(M; \mathbb{Q}) \# G, C^*(M; \mathbb{Q}) \# G) \cong H_*(P_G M^{-TM}; \mathbb{Q})^G$	$HH^*(C^*(M; \mathbb{Q}) \# G, C^*(M; \mathbb{Q}) \# G) \cong H_*(P_G M^{-TM}; \mathbb{Q})^G$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0913	351	■ 1.000	$C^{\{*\}}(M) \setminus \# G := C^{\{*\}}(M) \otimes \mathbb{Q}G \quad (\alpha, g) \cdot (\beta, h) := (\alpha \cdot g^* \beta, gh) .$	$C^*(M) \# G := C^*(M) \otimes \mathbb{Q}G \quad (\alpha, g) \cdot (\beta, h) := (\alpha \cdot g^* \beta, gh) .$	$C^*(M) \# G := C^*(M) \otimes \mathbb{Q}G \quad (\alpha, g) \cdot (\beta, h) := (\alpha \cdot g^* \beta, gh) .$
cardona-qft-methods_ EQ0914	351	■ 1.000	$H \, H^{\{*\}} \left(C^{\{*\}}(M ; \mathbb{Q}) \setminus \# G, C^{\{*\}}(M ; \mathbb{Q}) \setminus \# G \right) \cong \operatorname{Ext}_{\mathbb{Q}G}^* \left(\mathbb{Q}, C_*(P_G M^{-TM}) \right) .$	$HH^*(C^*(M; \mathbb{Q}) \# G, C^*(M; \mathbb{Q}) \# G) \cong \operatorname{Ext}_{\mathbb{Q}G}^*(\mathbb{Q}, C_*(P_G M^{-TM}; \mathbb{Q})) .$	$HH^*(C^*(M; \mathbb{Q}) \# G, C^*(M; \mathbb{Q}) \# G) \cong \operatorname{Ext}_{\mathbb{Q}G}^*(\mathbb{Q}, C_*(P_G M^{-TM}; \mathbb{Q})) .$
cardona-qft-methods_ EQ0918	351	■ 1.000	$C^{\{-*\}}(M) \rightarrowtail C_* \left(M^{\{-T M\}} \right)$	$C^{-*}(M) \rightarrow C_*(M^{-TM})$	$C^{-*}(M) \rightarrow C_*(M^{-TM})$
cardona-qft-methods_ EQ0920	352	■ 1.000	$H \, H^{\{*\}} \left(C^{\{*\}}(M ; \mathbb{Q}) \setminus \# G, C^{\{*\}}(M ; \mathbb{Q}) \setminus \# G \right) \cong \operatorname{Ext}_{\mathbb{Q}G}^* \left(\mathbb{Q}, C_*(P_G M^{-TM}) \right) .$	$HH^*(C^*(M; \mathbb{Q}) \# G, C^*(M; \mathbb{Q}) \# G) \cong \operatorname{Ext}_{\mathbb{Q}G}^*(\mathbb{Q}, C_*(P_G M^{-TM}; \mathbb{Q}))$	$HH^*(C^*(M; \mathbb{Q}) \# G, C^*(M; \mathbb{Q}) \# G) \cong \operatorname{Ext}_{\mathbb{Q}G}^*(\mathbb{Q}, C_*(P_G M^{-TM}; \mathbb{Q}))$
cardona-qft-methods_ EQ0921	352	■ 1.000	$H \, H^{\{*\}} \left(C^{\{*\}}(M ; \mathbb{Q}) \setminus \# G, C^{\{*\}}(M ; \mathbb{Q}) \setminus \# G \right) \cong H_* \left(P_G M^{-TM}; \mathbb{Q} \right)^G .$	$HH^*(C^*(M; \mathbb{Q}) \# G, C^*(M; \mathbb{Q}) \# G) \cong H_* \left(P_G M^{-TM}; \mathbb{Q} \right)^G .$	$HH^*(C^*(M; \mathbb{Q}) \# G, C^*(M; \mathbb{Q}) \# G) \cong H_* \left(P_G M^{-TM}; \mathbb{Q} \right)^G .$
cardona-qft-methods_ EQ0922	352	■ 1.000	$H \, H^{\{*\}} \left(C^{\{*\}}(M) \setminus \# G, C^{\{*\}}(M) \setminus \# G \right)$	$HH^*(C^*(M) \# G, C^*(M) \# G)$	$HH^*(C^*(M) \# G, C^*(M) \# G)$
cardona-qft-methods_ EQ0923	352	■ 1.000	$\operatorname{Ext}_{\mathbb{Z}G}^* \left(\mathbb{Z}, C_* \left(P_G M^{-TM} \right) \right) \cong H^* \left(\left(C^*(EG) \otimes C_* \left(P_G M^{-TM} \right) \right)^G \right)$	$\operatorname{Ext}_{\mathbb{Z}G}^*(\mathbb{Z}, C_*(P_G M^{-TM})) \cong H^* \left(\left(C^*(EG) \otimes C_*(P_G M^{-TM}) \right)^G \right)$	$\operatorname{Ext}_{\mathbb{Z}G}^*(\mathbb{Z}, C_*(P_G M^{-TM})) \cong H^* \left(\left(C^*(EG) \otimes C_*(P_G M^{-TM}) \right)^G \right)$
cardona-qft-methods_ EQ0924	352	■ 1.000	$P_G M \times_G EG_n \xrightarrow{e_0} M \times_G EG_n$	$P_G M \times_G EG_n \xrightarrow{e_0} M \times_G EG_n$	$P_G M \times_G EG_n \xrightarrow{e_0} M \times_G EG_n$
cardona-qft-methods_ EQ0926	352	■ 1.000	$\left(P_G M \times_G EG_n \right)^{-e_0^* T(M_n)} \leftarrow \left(P_G M \times_G EG_{n+1} \right)^{-e_0^* T(M_{n+1})} \leftarrow \dots$	$(P_G M \times_G EG_n)^{-e_0^* T(M_n)} \leftarrow (P_G M \times_G EG_{n+1})^{-e_0^* T(M_{n+1})} \leftarrow \dots$	$(P_G M \times_G EG_n)^{-e_0^* T(M_n)} \leftarrow (P_G M \times_G EG_{n+1})^{-e_0^* T(M_{n+1})} \leftarrow \dots$
cardona-qft-methods_ EQ0929	353	■ 1.000	$\left[G^{ad} / G \right] \cong \bigsqcup_{(g)} [* / C(g)]$	$[G^{ad} / G] \cong \bigsqcup_{(g)} [* / C(g)]$	$[G^{ad} / G] \cong \bigsqcup_{(g)} [* / C(g)]$
cardona-qft-methods_ EQ0930 (g)	353	■ 1.000	$\bigsqcup B \, C(g)_n^{\{-T \, B \, C(g)_n\}}$	$\bigsqcup BC(g)_n^{-TBC(g)_n}$	$\bigsqcup BC(g)_n^{-TBC(g)_n}$
cardona-qft-methods_ EQ0932	354	■ 1.000	$H_* \left(EG_n / C(g) \times EG_n / C(h) \right) \rightarrow H_{*-k_n} \left(EG_n / (C(g) \cap C(h)) \right)$	$H_*(EG_n / C(g) \times EG_n / C(h)) \rightarrow H_{*-k_n}(EG_n / (C(g) \cap C(h)))$	$H_*(EG_n / C(g) \times EG_n / C(h)) \rightarrow H_{*-k_n}(EG_n / (C(g) \cap C(h)))$
cardona-qft-methods_ EQ0933	354	■ 1.000	$H_* \left(EG_n / (C(g) \cap C(h)) \right) \rightarrow H_* \left(EG_n / C(gh) \right)$	$H_*(EG_n / (C(g) \cap C(h))) \rightarrow H_*(EG_n / C(gh))$	$H_*(EG_n / (C(g) \cap C(h))) \rightarrow H_*(EG_n / C(gh))$
cardona-qft-methods_ EQ0934	354	■ 1.000	$H^* \left(EG_n / (C(g) \cap C(h)) \right) \rightarrow H^* \left(EG_n / C(gh) \right)$	$H^*(EG_n / (C(g) \cap C(h))) \rightarrow H^*(EG_n / C(gh))$	$H^*(EG_n / (C(g) \cap C(h))) \rightarrow H^*(EG_n / C(gh))$
cardona-qft-methods_ EQ0935	354	■ 1.000	$H^* \left(B(C(g) \cap C(h)) \right) \rightarrow H^* \left(BC(gh) \right) .$	$H^*(B(C(g) \cap C(h))) \rightarrow H^*(BC(gh)) .$	$H^*(B(C(g) \cap C(h))) \rightarrow H^*(BC(gh)) .$
cardona-qft-methods_ EQ0938	355	■ 1.000	$H_*^{pro} \left(LBG^{-TBG} \right) \cong H^* \left(LBG \right)$	$H_*^{pro}(LBG^{-TBG}) \cong H^*(LBG)$	$H_*^{pro}(LBG^{-TBG}) \cong H^*(LBG)$
cardona-qft-methods_ EQ0939	355	■ 1.000	$LBG \leftarrow \operatorname{Map}(\Theta, BG) \rightarrow LBG \times LBG$	$LBG \leftarrow \operatorname{Map}(\Theta, BG) \rightarrow LBG \times LBG$	$LBG \leftarrow \operatorname{Map}(\Theta, BG) \rightarrow LBG \times LBG$
cardona-qft-methods_ EQ0940	355	■ 1.000	$H^* \left(LBG \right) \otimes H^* \left(LBG \right) \rightarrow H^* \left(LBG \right)$	$H^*(LBG) \otimes H^*(LBG) \rightarrow H^*(LBG)$	$H^*(LBG) \otimes H^*(LBG) \rightarrow H^*(LBG)$
cardona-qft-methods_ EQ0941 (1)	358	■ 1.000	$[X] - [Y] = [X \setminus Y],$	$[X] - [Y] = [X \setminus Y],$	$[X] - [Y] = [X \setminus Y],$
cardona-qft-methods_ EQ0944 (4)	358	■ 1.000	$\left[\mathbb{A}_{\mathbb{A}}^n \right] = \mathbb{L}^n .$	$[\mathbb{A}_k^n] = \mathbb{L}^n .$	$[\mathbb{A}_k^n] = \mathbb{L}^n .$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0945 (5)	359	■ 1.000	$\mathbb{T}=\left[\mathbb{A}_{\mathbb{k}}^1\right]-\left[\mathbb{A}_{\mathbb{k}}^0\right]=\mathbb{L}-1,$ $\mathbb{A}_{\mathbb{k}}^{\{0\}}\right]=\mathbb{L}-1,$	$\mathbb{T} = \left[\mathbb{A}_{\mathbb{k}}^1\right] - \left[\mathbb{A}_{\mathbb{k}}^0\right] = \mathbb{L} - 1,$	$\mathbb{T} = \left[\mathbb{A}_{\mathbb{k}}^1\right] - \left[\mathbb{A}_{\mathbb{k}}^0\right] = \mathbb{L} - 1,$
cardona-qft-methods_ EQ0946 (6)	359	■ 1.000	$[X \cup Y]=[X]+[Y]-[X \cap Y]$	$[X \cup Y] = [X] + [Y] - [X \cap Y]$	$[X \cup Y] = [X] + [Y] - [X \cap Y]$
cardona-qft-methods_ EQ0948 (8)	359	■ 1.000	$\left[\mathbb{P}_{\mathbb{k}}^{\mathbb{n}}\right]=\frac{\mathbb{L}^{n+1}-1}{\mathbb{L}-1},$	$[\mathbb{P}_{\mathbb{k}}^n] = \frac{\mathbb{L}^{n+1} - 1}{\mathbb{L} - 1},$	$[\mathbb{P}_{\mathbb{k}}^n] = \frac{\mathbb{L}^{n+1} - 1}{\mathbb{L} - 1},$
cardona-qft-methods_ EQ0949 (9)	359	■ 1.000	$\left[\mathbb{P}_{\mathbb{k}}^{\mathbb{n}}\right]=1+\mathbb{L}+\mathbb{L}^2+\cdots+\mathbb{L}^n,$	$[\mathbb{P}_{\mathbb{k}}^n] = 1 + \mathbb{L} + \mathbb{L}^2 + \cdots + \mathbb{L}^n,$	$[\mathbb{P}_{\mathbb{k}}^n] = 1 + \mathbb{L} + \mathbb{L}^2 + \cdots + \mathbb{L}^n,$
cardona-qft-methods_ EQ0950 (1)	359	■ 1.000	$\Psi_{\Gamma}(\mathfrak{t})=\sum_{\mathbb{T} \subset \Gamma} \prod_{e \notin E(\mathbb{T})} t_e,$	$\Psi_{\Gamma}(t) = \sum_{T \subset \Gamma} \prod_{e \notin E(T)} t_e,$	$\Psi_{\Gamma}(t) = \sum_{T \subset \Gamma} \prod_{e \notin E(T)} t_e,$
cardona-qft-methods_ EQ0951 (2)	360	■ 1.000	$X_{\Gamma}=\left\{t=\left(t_1:t_2:\cdots:t_n\right) \in \mathbb{P}_{\mathbb{k}}^{n-1} \mid \Psi_{\Gamma}(\mathfrak{t})=0\right\}.$	$X_{\Gamma} = \left\{t = (t_1 : t_2 : \cdots : t_n) \in \mathbb{P}_k^{n-1} \mid \Psi_{\Gamma}(t) = 0\right\}.$	$X_{\Gamma} = \left\{t = (t_1 : t_2 : \cdots : t_n) \in \mathbb{P}_k^{n-1} \mid \Psi_{\Gamma}(t) = 0\right\}.$
cardona-qft-methods_ EQ0952 (4)	360	■ 1.000	$\begin{gathered} \mathcal{C}: \mathbb{P}_{\mathbb{k}}^{\mathbb{n}-1} \rightarrow \mathbb{P}_{\mathbb{k}}^{\mathbb{n}-1} \\ \left(t_1:t_2:\cdots:t_n\right) \mapsto \left(\frac{1}{t_1}:\frac{1}{t_2}:\cdots:\frac{1}{t_n}\right) \end{gathered}$	$\mathcal{C}: \mathbb{P}_k^{n-1} \rightarrow \mathbb{P}_k^{n-1}$ $(t_1 : t_2 : \cdots : t_n) \mapsto \left(\frac{1}{t_1} : \frac{1}{t_2} : \cdots : \frac{1}{t_n}\right)$	$\mathcal{C}: \mathbb{P}_k^{n-1} \rightarrow \mathbb{P}_k^{n-1}$ $(t_1 : t_2 : \cdots : t_n) \mapsto \left(\frac{1}{t_1} : \frac{1}{t_2} : \cdots : \frac{1}{t_n}\right)$
cardona-qft-methods_ EQ0953 (5)	360	■ 1.000	$\Sigma_{\mathbb{n}}=\left\{\left(t_1:t_2:\cdots:t_n\right) \in \mathbb{P}_{\mathbb{k}}^{n-1} \mid \prod_{i=1}^n t_i=0\right\}.$	$\Sigma_n = \left\{(t_1 : t_2 : \cdots : t_n) \in \mathbb{P}_k^{n-1} \mid \prod_{i=1}^n t_i = 0\right\}.$	$\Sigma_n = \left\{(t_1 : t_2 : \cdots : t_n) \in \mathbb{P}_k^{n-1} \mid \prod_{i=1}^n t_i = 0\right\}.$
cardona-qft-methods_ EQ0954 (6)	361	■ 1.000	$\mathcal{C} \circ \pi_1=\pi_2.$	$\mathcal{C} \circ \pi_1 = \pi_2.$	$\mathcal{C} \circ \pi_1 = \pi_2.$
cardona-qft-methods_ EQ0955 (7)	361	■ 1.000	$t_1 t_{n+1}=t_2 t_{n+2}=\cdots=t_n t_{2 n},$	$t_1 t_{n+1} = t_2 t_{n+2} = \cdots = t_n t_{2n},$	$t_1 t_{n+1} = t_2 t_{n+2} = \cdots = t_n t_{2n},$
cardona-qft-methods_ EQ0961 (1)	362	■ 1.000	$\Psi_{\Gamma}(\mathfrak{t})=1,$	$\Psi_{\Gamma}(t) = 1,$	$\Psi_{\Gamma}(t) = 1,$
cardona-qft-methods_ EQ0962 (2)	362	■ 1.000	$X_{\Gamma}=\emptyset.$	$X_{\Gamma} = \emptyset.$	$X_{\Gamma} = \emptyset.$
cardona-qft-methods_ EQ0963 (3)	362	■ 1.000	$\Psi_{\Gamma^{\vee}}(\mathfrak{t})=t_1 t_2 \cdots t_n,$	$\Psi_{\Gamma^{\vee}}(t) = t_1 t_2 \cdots t_n,$	$\Psi_{\Gamma^{\vee}}(t) = t_1 t_2 \cdots t_n,$
cardona-qft-methods_ EQ0964 (4)	362	■ 1.000	$X_{\Gamma^{\vee}}=\left\{t \in \mathbb{P}_{\mathbb{k}}^{n-1} \mid \Psi_{\Gamma^{\vee}}(\mathfrak{t})=0\right\}=\Sigma_{\mathbb{n}}.$	$X_{\Gamma^{\vee}} = \left\{t \in \mathbb{P}_k^{n-1} \mid \Psi_{\Gamma^{\vee}}(t) = 0\right\} = \Sigma_n.$	$X_{\Gamma^{\vee}} = \left\{t \in \mathbb{P}_k^{n-1} \mid \Psi_{\Gamma^{\vee}}(t) = 0\right\} = \Sigma_n.$
cardona-qft-methods_ EQ0966 (6)	362	■ 1.000	$\left[X_{\Gamma}\right]=0.$	$[X_{\Gamma}] = 0.$	$[X_{\Gamma}] = 0.$
cardona-qft-methods_ EQ0968 (8)	363	■ 1.000	$\Psi_{\Gamma}(\mathfrak{t})=t_1+t_2+\cdots+t_n,$	$\Psi_{\Gamma}(t) = t_1 + t_2 + \cdots + t_n,$	$\Psi_{\Gamma}(t) = t_1 + t_2 + \cdots + t_n,$
cardona-qft-methods_ EQ0970 (10)	363	■ 1.000	$\left[\mathcal{L}\right]=\left[X_{\Gamma}\right]=\left[\mathbb{P}_{\mathbb{k}}^{n-2}\right]=\frac{\mathbb{L}^{n-1}-1}{\mathbb{L}-1}=\frac{(\mathbb{T}+1)^{n-1}-1}{\mathbb{T}}.$	$[\mathcal{L}] = [X_{\Gamma}] = \left[\mathbb{P}_k^{n-2}\right] = \frac{\mathbb{L}^{n-1} - 1}{\mathbb{L} - 1} = \frac{(\mathbb{T} + 1)^{n-1} - 1}{\mathbb{T}}.$	$[\mathcal{L}] = [X_{\Gamma}] = \left[\mathbb{P}_k^{n-2}\right] = \frac{\mathbb{L}^{n-1} - 1}{\mathbb{L} - 1} = \frac{(\mathbb{T} + 1)^{n-1} - 1}{\mathbb{T}}.$
cardona-qft-methods_ EQ0973 (13)	363	■ 1.000	$\mathcal{H}: g(\mathbb{T}) \mapsto \frac{g(\mathbb{T})-g(-1)}{\mathbb{T}+1}$	$\mathcal{H}: g(\mathbb{T}) \mapsto \frac{g(\mathbb{T}) - g(-1)}{\mathbb{T} + 1}$	$\mathcal{H}: g(\mathbb{T}) \mapsto \frac{g(\mathbb{T}) - g(-1)}{\mathbb{T} + 1}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ0974 (14)	364	■ 1.000	$\frac{g(\mathbb{T}-g(-1))}{\mathbb{T}+1}=\frac{g(\mathbb{T}-1)^{n-1}-1}{\mathbb{T}}=\left[\mathbb{P}^{n-1}\right],$	$\frac{g(\mathbb{T}-g(-1))}{\mathbb{T}+1}=\frac{(\mathbb{T}-1)^{n-1}-1}{\mathbb{T}}=\left[\mathbb{P}^{n-1}\right],$	$\frac{g(\mathbb{T}-g(-1))}{\mathbb{T}+1}=\frac{(\mathbb{T}-1)^{n-1}-1}{\mathbb{T}}=\left[\mathbb{P}^{n-1}\right],$
cardona-qft-methods_ EQ0975 (15)	364	■ 1.000	$\left[\mathcal{L}\cap\Sigma_n\right]=\frac{g(\mathbb{T})-g(-1)}{\mathbb{T}+1}=\frac{(1+\mathbb{T})^{n-1}-1}{\mathbb{T}}-\frac{\mathbb{T}^{n-1}-(-1)^{n-1}}{\mathbb{T}+1},$	$[\mathcal{L}\cap\Sigma_n]=\frac{g(\mathbb{T})-g(-1)}{\mathbb{T}+1}=\frac{(1+\mathbb{T})^{n-1}-1}{\mathbb{T}}-\frac{\mathbb{T}^{n-1}-(-1)^{n-1}}{\mathbb{T}+1},$	$[\mathcal{L}\cap\Sigma_n]=\frac{g(\mathbb{T})-g(-1)}{\mathbb{T}+1}=\frac{(1+\mathbb{T})^{n-1}-1}{\mathbb{T}}-\frac{\mathbb{T}^{n-1}-(-1)^{n-1}}{\mathbb{T}+1},$
cardona-qft-methods_ EQ0976 (16)	364	■ 1.000	$\left[X_{\Gamma^\vee}\backslash\Sigma_n\right]=\left[\mathcal{L}\backslash\Sigma_n\right]=\left[\mathcal{L}\right]-\left[\mathcal{L}\cap\Sigma_n\right]=\frac{\mathbb{T}^{n-1}-(-1)^{n-1}}{\mathbb{T}+1}.$	$[X_{\Gamma^\vee}\backslash\Sigma_n]=[\mathcal{L}\backslash\Sigma_n]=[\mathcal{L}]-[\mathcal{L}\cap\Sigma_n]=\frac{\mathbb{T}^{n-1}-(-1)^{n-1}}{\mathbb{T}+1}.$	$[X_{\Gamma^\vee}\backslash\Sigma_n]=[\mathcal{L}\backslash\Sigma_n]=[\mathcal{L}]-[\mathcal{L}\cap\Sigma_n]=\frac{\mathbb{T}^{n-1}-(-1)^{n-1}}{\mathbb{T}+1}.$
cardona-qft-methods_ EQ0977 (17)	364	■ 1.000	$\mathcal{I}=\left(t_2t_3\cdots t_n,t_1t_3\cdots t_n,\dots,t_1t_2\cdots\hat{t}_i\cdots t_n,\dots,t_1t_2\cdots t_{n-1}\right).$	$\mathcal{I}=\left(t_2t_3\cdots t_n,t_1t_3\cdots t_n,\dots,t_1t_2\cdots\hat{t}_i\cdots t_n,\dots,t_1t_2\cdots t_{n-1}\right).$	$\mathcal{I}=\left(t_2t_3\cdots t_n,t_1t_3\cdots t_n,\dots,t_1t_2\cdots\hat{t}_i\cdots t_n,\dots,t_1t_2\cdots t_{n-1}\right).$
cardona-qft-methods_ EQ0980 (20)	365	■ 1.000	$\left[X_{\Gamma^\vee}\backslash\Sigma\right]+\left[X_{\Gamma^\vee}\cap\Sigma\right]=\left[X_{\Gamma^\vee}\backslash\Sigma\right]+[\mathcal{S}_n],$	$[X_{\Gamma^\vee}]=[X_{\Gamma^\vee}\backslash\Sigma]+[X_{\Gamma^\vee}\cap\Sigma]=[X_{\Gamma^\vee}\backslash\Sigma]+[\mathcal{S}_n],$	$[X_{\Gamma^\vee}]=[X_{\Gamma^\vee}\backslash\Sigma]+[X_{\Gamma^\vee}\cap\Sigma]=[X_{\Gamma^\vee}\backslash\Sigma]+[\mathcal{S}_n],$
cardona-qft-methods_ EQ0981 (1)	366	■ 1.000	$\Psi_\Gamma(t)=p(t)q(t)$	$\Psi_\Gamma(t)=p(t)q(t)$	$\Psi_\Gamma(t)=p(t)q(t)$
cardona-qft-methods_ EQ0982 (2)	367	■ 1.000	$P=\{t_i\mid \text{ such that } t_i \text{ appears in } p(t)\}$	$P=\{t_i\mid \text{ such that } t_i \text{ appears in } p(t)\}$	$P=\{t_i\mid \text{ such that } t_i \text{ appears in } p(t)\}$
cardona-qft-methods_ EQ0984	369	■ 1.000	$XY-YX-[X,Y],\quad X,Y\in\tilde{m}g.$	$XY-YX-[X,Y],\quad X,Y\in\tilde{m}g.$	$XY-YX-[X,Y],\quad X,Y\in\tilde{m}g.$
cardona-qft-methods_ EQ0985	369	■ 1.000	$gf(x)=f(g^{-1}x),\quad f\in C(M).$	$gf(x)=f(g^{-1}x),\quad f\in C(M).$	$gf(x)=f(g^{-1}x),\quad f\in C(M).$
cardona-qft-methods_ EQ0986	369	■ 1.000	$e_\xi:g\in G\mapsto\langle J\xi,g\nu\rangle\in\mathbb{C}.$	$e_\xi:g\in G\mapsto\langle J\xi,g\nu\rangle\in\mathbb{C}.$	$e_\xi:g\in G\mapsto\langle J\xi,g\nu\rangle\in\mathbb{C}.$
cardona-qft-methods_ EQ0990	371	■ 1.000	$\omega(A)=\langle\nu,Av\rangle,\quad A\in\mathcal{A}.$	$\omega(A)=\langle\nu,Av\rangle,\quad A\in\mathcal{A}.$	$\omega(A)=\langle\nu,Av\rangle,\quad A\in\mathcal{A}.$
cardona-qft-methods_ EQ0991	372	■ 1.000	$\langle[A],[B]\rangle=\omega(A^\dagger B),\quad A,B\in\mathcal{A},$	$\langle[A],[B]\rangle=\omega(A^\dagger B),\quad A,B\in\mathcal{A},$	$\langle[A],[B]\rangle=\omega(A^\dagger B),\quad A,B\in\mathcal{A},$
cardona-qft-methods_ EQ0993	373	■ 1.000	$\Delta\mathcal{I}\subseteq\mathcal{I}\otimes\mathcal{T}(\tilde{m}g)+\mathcal{T}(\tilde{m}g)\otimes\mathcal{I}.$	$\Delta\mathcal{I}\subseteq\mathcal{I}\otimes\mathcal{T}(\tilde{m}g)+\mathcal{T}(\tilde{m}g)\otimes\mathcal{I}.$	$\Delta\mathcal{I}\subseteq\mathcal{I}\otimes\mathcal{T}(\tilde{m}g)+\mathcal{T}(\tilde{m}g)\otimes\mathcal{I}.$
cardona-qft-methods_ EQ0996	374	■ 1.000	$ab(E)=a\otimes b(\Delta E),\quad a,b\in\mathcal{E}^\dagger,E\in\mathcal{E}.$	$ab(E)=a\otimes b(\Delta E),\quad a,b\in\mathcal{E}^\dagger,E\in\mathcal{E}.$	$ab(E)=a\otimes b(\Delta E),\quad a,b\in\mathcal{E}^\dagger,E\in\mathcal{E}.$
cardona-qft-methods_ EQ0997 (1)	374	■ 1.000	$Ea(F)=a(\overline{E}^\dagger F),\quad E,F\in\mathcal{E},a\in\mathcal{E}^\dagger.$	$Ea(F)=a(\overline{E}^\dagger F),\quad E,F\in\mathcal{E},a\in\mathcal{E}^\dagger.$	$Ea(F)=a(\overline{E}^\dagger F),\quad E,F\in\mathcal{E},a\in\mathcal{E}^\dagger.$
cardona-qft-methods_ EQ0998	374	■ 1.000	$(\forall a,b\in\mathcal{E}^\dagger)\delta(ab)=\delta(a)b+a\delta(b).$	$(\forall a,b\in\mathcal{E}^\dagger)\delta(ab)=\delta(a)b+a\delta(b).$	$(\forall a,b\in\mathcal{E}^\dagger)\delta(ab)=\delta(a)b+a\delta(b).$
cardona-qft-methods_ EQ0999	374	■ 1.000	$X(ab)=X((a\otimes b)\circ\Delta)=(a\otimes b)\circ\Delta\circ X^\dagger(\cdot).$	$X(ab)=X((a\otimes b)\circ\Delta)=(a\otimes b)\circ\Delta\circ X^\dagger(\cdot).$	$X(ab)=X((a\otimes b)\circ\Delta)=(a\otimes b)\circ\Delta\circ X^\dagger(\cdot).$
cardona-qft-methods_ EQ1002	375	■ 1.000	$a\in\mathcal{E}^\dagger\mapsto f_a\in\mathbb{C}[[x_1,\dots x_n]],\quad f_a(x)=\sum_{\alpha\in\mathbb{N}^n}\frac{x^\alpha}{\alpha!}a(X^\alpha).$	$a\in\mathcal{E}^\dagger\mapsto f_a\in\mathbb{C}[[x_1,\dots x_n]],\quad f_a(x)=\sum_{\alpha\in\mathbb{N}^n}\frac{x^\alpha}{\alpha!}a(X^\alpha).$	$a\in\mathcal{E}^\dagger\mapsto f_a\in\mathbb{C}[[x_1,\dots x_n]],\quad f_a(x)=\sum_{\alpha\in\mathbb{N}^n}\frac{x^\alpha}{\alpha!}a(X^\alpha).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
cardona-qft-methods_ EQ1004	375	■ 1.000	$a\,b(1)=a\,\otimes b(\Delta 1)=a(1)\,b(1)\;.$	$ab(1) = a \otimes b(\Delta 1) = a(1)b(1).$	$ab(1) = a \otimes b(\Delta 1) = a(1)b(1).$
cardona-qft-methods_ EQ1005	375	■ 1.000	$a^{\dagger}(E+f\,F)=\overline{a(E)}+f\,\overline{a(F)},$ $\quad E,\,F\in\mathcal{E}_{\mathbb{R}}\;.$	$a^{\dagger}(E+fF) = \overline{a(E)} + f\overline{a(F)}, \quad E, F \in \mathcal{E}_{\mathbb{R}}.$	$a^{\dagger}(E+\,f\,F) = \overline{a(E)} + \,f\,\overline{a(F)}, \quad E, F \in \mathcal{E}_{\mathbb{R}}.$
cardona-qft-methods_ EQ1006	375	■ 1.000	$\chi_{\{0\}}\left(a^{\dagger}\right)=a^{\dagger}(1)=$ $\overline{a(1)}=\overline{\chi_{\{0\}}(a)}\;.$	$\chi_0\left(a^{\dagger}\right) = a^{\dagger}(1) = \overline{a(1)} = \overline{\chi_0(a)}.$	$\chi_0(a^{\dagger}) = a^{\dagger}(1) = \overline{a(1)} = \overline{\chi_0(a)}.$
cardona-qft-methods_ EQ1007	376	■ 1.000	$\xi\in\mathcal{F}\mapsto e_{\xi}\in\mathcal{E}^{\dagger}, \quad e_{\xi}(E)=\langle J\xi,E\nu\rangle,$ $\quad e_{\xi}(E)=\langle J\xi,E\nu\rangle,$	$\xi \in \mathcal{F} \mapsto e_{\xi} \in \mathcal{E}^{\dagger}, \quad e_{\xi}(E) = \langle J\xi, E\nu\rangle,$	$\xi \in \mathcal{F} \mapsto e_{\xi} \in \mathcal{E}^{\dagger}, \quad e_{\xi}(E) = \langle J\xi, E\nu\rangle,$
cardona-qft-methods_ EQ1008	376	■ 1.000	$M=\left\{:\chi\in\mathcal{A}^{\dagger}\mid\chi(a\,b)=\right.$ $\left.\chi(a)\,\chi(b),\,\chi\left(a^{\dagger}\right)=\overline{\chi(a)}:\right\}.$	$M = \left\{:\chi \in \mathcal{A}^{\dagger} \mid \chi(ab) = \chi(a)\chi(b), \chi\left(a^{\dagger}\right) = \overline{\chi(a)}:\right\}.$	$M = \left\{:\chi \in \mathcal{A}^{\dagger} \mid \chi(ab) = \chi(a)\chi(b), \chi(a^{\dagger}) = \overline{\chi(a)}:\right\}.$
cardona-qft-methods_ EQ1009	376	■ 1.000	$E\,e_{\xi}(F)=\langle J\xi,\bar{E}^{\dagger}F\nu\rangle=\langle\bar{E}J\xi,F\nu\rangle=\langle JE\xi,F\nu\rangle=e_{E\xi}(F).$ $\quad E\,e_{\xi}(F)=\langle J\xi,\bar{E}^{\dagger}F\nu\rangle=\langle\bar{E}J\xi,F\nu\rangle=\langle JE\xi,F\nu\rangle=e_{E\xi}(F).$	$Ee_{\xi}(F) = \langle J\xi, \bar{E}^{\dagger}F\nu\rangle = \langle \bar{E}J\xi, F\nu\rangle = \langle JE\xi, F\nu\rangle = e_{E\xi}(F).$	$Ee_{\xi}(F) = \langle J\xi, \overline{E}^{\dagger}F\nu\rangle = \langle \overline{E}J\xi, F\nu\rangle = \langle JE\xi, F\nu\rangle = e_{E\xi}(F).$
cardona-qft-methods_ EQ1010	376	■ 1.000	$\omega\in\mathcal{E}^{\dagger}, \quad \omega(E)=e_{\nu}(E)=\langle\nu,E\nu\rangle$ $\quad \omega(E)=\langle\nu,E\nu\rangle$	$\omega \in \mathcal{E}^{\dagger}, \quad \omega(E) = e_{\nu}(E) = \langle \nu, E\nu\rangle$	$\omega \in \mathcal{E}^{\dagger}, \quad \omega(E) = e_{\nu}(E) = \langle \nu, E\nu\rangle$
cardona-qft-methods_ EQ1011	377	■ 1.000	$\tilde{m}g=\mathbb{R}X, \quad \mathcal{F}=\mathcal{H}=\mathbb{C}^2,$ $\quad X=\left(\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right), \quad \nu=\left(\begin{array}{c} 1 \\ 0 \end{array}\right).$ $\quad \tilde{m}g=\mathbb{R}X, \quad \mathcal{F}=\mathcal{H}=\mathbb{C}^2, \quad X=\left(\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right), \quad \nu=\left(\begin{array}{c} 1 \\ 0 \end{array}\right).$	$\tilde{m}g = \mathbb{R}X, \quad \mathcal{F} = \mathcal{H} = \mathbb{C}^2, \quad X = \left(\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right), \quad \nu = \left(\begin{array}{c} 1 \\ 0 \end{array}\right).$	$\tilde{m}g = \mathbb{R}X, \quad \mathcal{F} = \mathcal{H} = \mathbb{C}^2, \quad X = \left(\begin{array}{cc} 0 & 1 \\ -1 & 0 \end{array}\right), \quad \nu = \left(\begin{array}{c} 1 \\ 0 \end{array}\right).$