

cardona-qft-methods: table evidence

Tables

Identifier	Page	Conf.	LaTeX source	Rendered	Image																																																																								
cardona-qft-methods_ TAB_001	43	■ 1.000	$\begin{tabular}{t}{ l l l l l l l l l }\hline \backslash(d\backslash) \& 0 \& 1 \& 2 \& 3 \& 4 \& 5 \& 6 \& 7 \backslash\backslash\hline \backslash(\epsilon\backslash) \& 1 \& -1 \& 1 \& 1 \& 1 \& -1 \& 1 \& 1 \backslash\backslash\hline \backslash(\epsilon^{\prime}\backslash) \& 1 \& 1 \& -1 \& -1 \& -1 \& -1 \& 1 \& 1 \backslash\backslash\hline \backslash(\epsilon^{\prime\prime}\backslash) \& 1 \& \& -1 \& \& 1 \& \& -1 \& \backslash\backslash\hline\end{tabular}$	<table><tr><td>d</td><td>0</td><td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td><td>7</td></tr><tr><td>ϵ</td><td>1</td><td>-1</td><td>1</td><td>1</td><td>1</td><td>-1</td><td>1</td><td>1</td></tr><tr><td>ϵ'</td><td>1</td><td>1</td><td>-1</td><td>-1</td><td>-1</td><td>-1</td><td>1</td><td>1</td></tr><tr><td>ϵ''</td><td>1</td><td></td><td>-1</td><td></td><td>1</td><td></td><td>-1</td><td></td></tr></table>	d	0	1	2	3	4	5	6	7	ϵ	1	-1	1	1	1	-1	1	1	ϵ'	1	1	-1	-1	-1	-1	1	1	ϵ''	1		-1		1		-1		<table><tr><td>d</td><td>0</td><td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td><td>7</td></tr><tr><td>ϵ</td><td>1</td><td>-1</td><td>1</td><td>1</td><td>1</td><td>-1</td><td>1</td><td>1</td></tr><tr><td>ϵ'</td><td>1</td><td>1</td><td>-1</td><td>-1</td><td>-1</td><td>-1</td><td>1</td><td>1</td></tr><tr><td>ϵ''</td><td>1</td><td></td><td>-1</td><td></td><td>1</td><td></td><td>-1</td><td></td></tr></table>	d	0	1	2	3	4	5	6	7	ϵ	1	-1	1	1	1	-1	1	1	ϵ'	1	1	-1	-1	-1	-1	1	1	ϵ''	1		-1		1		-1	
d	0	1	2	3	4	5	6	7																																																																					
ϵ	1	-1	1	1	1	-1	1	1																																																																					
ϵ'	1	1	-1	-1	-1	-1	1	1																																																																					
ϵ''	1		-1		1		-1																																																																						
d	0	1	2	3	4	5	6	7																																																																					
ϵ	1	-1	1	1	1	-1	1	1																																																																					
ϵ'	1	1	-1	-1	-1	-1	1	1																																																																					
ϵ''	1		-1		1		-1																																																																						
cardona-qft-methods_ TAB_002	181	■ 0.546	$\begin{tabular}{t}{ l l l }\hline \backslash(D=3\backslash) \text{CS Lagrangians \& Characteristic classes \& Groups} \backslash\backslash\hline \backslash[L_3^{\wedge(A)} d S]=\epsilon_{a b c}\backslashleft(R^{\wedge{a} b} \backslashpm \backslashfrac{e^{\wedge{a}} e^{\wedge{b}}}{3 l^2}\backslashright) e^{\wedge{c}} \backslash] \& \backslash(\mathbf{E}_4=\epsilon_{a b c}\backslashleft(R^{\wedge{a} b} \backslashpm \backslashfrac{e^{\wedge{a}} e^{\wedge{b}}}{l^2}\backslashright) T^{\wedge{c}}\backslash) \& \backslash\begin{tabular}{t}{ l }\hline \backslash(S 0(4)\backslash), \backslash\backslash(S 0(3,1)\backslash) \backslash\backslash\text{or} \backslash(S 0(2,2)\backslash) \backslash\end{tabular} \backslash\backslash\hline \backslash(L_3^{\wedge}\backslash\text{text {Lor}})=\omega^{\wedge{a}}\{ \}_{\wedge{b}} d \omega^{\wedge{b}}\{ \}_{\wedge{a}}+\backslashfrac{2}{3}\omega^{\wedge{a}}\{ \}_{\wedge{b}} \omega^{\wedge{b}}\{ \}_{\wedge{c}} \omega^{\wedge{c}}\{ \}_{\wedge{a}} \backslash) \& \backslash(\mathbf{P}_4^{\wedge}\backslash\text{text {Lor}})=R^{\wedge{a}}\{ \}_{\wedge{b}} R^{\wedge{b}}\{ \}_{\wedge{a}}\backslash) \& \backslash(S 0(2,1)\backslash) \backslash\backslash\hline \backslash(L_3^{\wedge}\backslash\text{T o r})=e^{\wedge{a}} T_{\wedge{a}}\backslash) \& \backslash(\mathbf{N}_4=T^{\wedge{a}} T_{\wedge{a}}-e^{\wedge{a}} e^{\wedge{b}} R_{a b})\backslash) \& \backslash(S 0(2,1)\backslash) \backslash\backslash\hline \backslash(L_3^{\wedge}\backslash\text{U(1)})=A d A\backslash) \& \backslash(\mathbf{P}_4^{\wedge}\backslash\text{U(1)})=F F\backslash) \& \backslash(U(1)\backslash) \backslash\backslash\hline \backslash(L_3^{\wedge}\backslash\text{S U(N)})=\operatorname{Tr}\backslashleft[\mathbf{A} d \mathbf{A}+\backslashfrac{2}{3} \mathbf{A} \mathbf{A} \mathbf{A}\right] \backslash) \& \backslash(\mathbf{P}_4^{\wedge}\backslash\text{S U(4)})=\operatorname{Tr}[\mathbf{F} F]\backslash) \& \backslash(S U(N)\backslash) \backslash\backslash\hline\end{tabular}$	(not rendered)	<table><tr><th>$D = 3$ CS Lagrangians</th><th>Characteristic classes</th><th>Groups</th></tr><tr><td>$L_3^{(A)dS} = \epsilon_{abc}(R^{ab} \pm \frac{e^a e^b}{3l^2})e^c$</td><td>$\mathbf{E}_4 = \epsilon_{abc}(R^{ab} \pm \frac{e^a e^b}{l^2})T^c$</td><td>$SO(4)$, $SO(3, 1)$ or $SO(2, 2)$</td></tr><tr><td>$L_3^{Lor} = \omega^a_b d\omega^b_a + \frac{2}{3}\omega^a_b \omega^b_c \omega^c_a$</td><td>$\mathbf{P}_4^{Lor} = R^a_b R^b_a$</td><td>$SO(2, 1)$</td></tr><tr><td>$L_3^{Tor} = e^a T_a$</td><td>$\mathbf{N}_4 = T^a T_a - e^a e^b R_{ab}$</td><td>$SO(2, 1)$</td></tr><tr><td>$L_3^{U(1)} = AdA$</td><td>$\mathbf{P}_4^{U(1)} = FF$</td><td>$U(1)$</td></tr><tr><td>$L_3^{SU(N)} = Tr[\mathbf{A}d\mathbf{A} + \frac{2}{3}\mathbf{A}\mathbf{A}\mathbf{A}]$</td><td>$\mathbf{P}_4^{SU(4)} = Tr[\mathbf{F}\mathbf{F}]$</td><td>$SU(N)$</td></tr></table>	$D = 3$ CS Lagrangians	Characteristic classes	Groups	$L_3^{(A)dS} = \epsilon_{abc}(R^{ab} \pm \frac{e^a e^b}{3l^2})e^c$	$\mathbf{E}_4 = \epsilon_{abc}(R^{ab} \pm \frac{e^a e^b}{l^2})T^c$	$SO(4)$, $SO(3, 1)$ or $SO(2, 2)$	$L_3^{Lor} = \omega^a_b d\omega^b_a + \frac{2}{3}\omega^a_b \omega^b_c \omega^c_a$	$\mathbf{P}_4^{Lor} = R^a_b R^b_a$	$SO(2, 1)$	$L_3^{Tor} = e^a T_a$	$\mathbf{N}_4 = T^a T_a - e^a e^b R_{ab}$	$SO(2, 1)$	$L_3^{U(1)} = AdA$	$\mathbf{P}_4^{U(1)} = FF$	$U(1)$	$L_3^{SU(N)} = Tr[\mathbf{A}d\mathbf{A} + \frac{2}{3}\mathbf{A}\mathbf{A}\mathbf{A}]$	$\mathbf{P}_4^{SU(4)} = Tr[\mathbf{F}\mathbf{F}]$	$SU(N)$																																																						
$D = 3$ CS Lagrangians	Characteristic classes	Groups																																																																											
$L_3^{(A)dS} = \epsilon_{abc}(R^{ab} \pm \frac{e^a e^b}{3l^2})e^c$	$\mathbf{E}_4 = \epsilon_{abc}(R^{ab} \pm \frac{e^a e^b}{l^2})T^c$	$SO(4)$, $SO(3, 1)$ or $SO(2, 2)$																																																																											
$L_3^{Lor} = \omega^a_b d\omega^b_a + \frac{2}{3}\omega^a_b \omega^b_c \omega^c_a$	$\mathbf{P}_4^{Lor} = R^a_b R^b_a$	$SO(2, 1)$																																																																											
$L_3^{Tor} = e^a T_a$	$\mathbf{N}_4 = T^a T_a - e^a e^b R_{ab}$	$SO(2, 1)$																																																																											
$L_3^{U(1)} = AdA$	$\mathbf{P}_4^{U(1)} = FF$	$U(1)$																																																																											
$L_3^{SU(N)} = Tr[\mathbf{A}d\mathbf{A} + \frac{2}{3}\mathbf{A}\mathbf{A}\mathbf{A}]$	$\mathbf{P}_4^{SU(4)} = Tr[\mathbf{F}\mathbf{F}]$	$SU(N)$																																																																											

Identifier	Page	Conf.	LaTeX source	Rendered	Image								
cardona-qft-methods_ TAB_008	287	■ 0.514	$\begin{tabular}[t]{t l l}\hline \mathfrak{s} \mathfrak{u}(2 \mid 2) & \text{create } (8_{\{B\}} + 8_{\{F\}}) \text{ magnons} \\ \hline \text{destroy } 8 + 8 \text{ magnons} & \overline{\mathfrak{s} \mathfrak{u}(2 \mid 2)} \\ \hline \end{tabular}$	<table><tr><td>$\mathfrak{su}(2 \mid 2)$</td><td>create $8_B + 8_F$ magnons</td></tr><tr><td>destroy $8 + 8$ magnons</td><td>$\mathfrak{su}(2 \mid 2)$</td></tr></table>	$\mathfrak{su}(2 \mid 2)$	create $8_B + 8_F$ magnons	destroy $8 + 8$ magnons	$\mathfrak{su}(2 \mid 2)$	<table><tr><td>$\mathfrak{su}(2 2)$</td><td>create $8_B + 8_F$ magnons</td></tr><tr><td>destroy $8 + 8$ magnons</td><td>$\overline{\mathfrak{su}(2 2)}$</td></tr></table>	$\mathfrak{su}(2 2)$	create $8_B + 8_F$ magnons	destroy $8 + 8$ magnons	$\overline{\mathfrak{su}(2 2)}$
$\mathfrak{su}(2 \mid 2)$	create $8_B + 8_F$ magnons												
destroy $8 + 8$ magnons	$\mathfrak{su}(2 \mid 2)$												
$\mathfrak{su}(2 2)$	create $8_B + 8_F$ magnons												
destroy $8 + 8$ magnons	$\overline{\mathfrak{su}(2 2)}$												