

fong-spivak-seven-sketches: formula evidence

The page, the confidence and the picture of an inline formula are its HOST LINE’s — a formula has none of its own. A line’s confidence is not a formula’s.

Inline formulas (first occurrence)

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_8_F00001	8	■ 0.976	a	a	2.1 Getting from a to b
fong-spivak-seven-sketches_8_F00002	8	■ 0.976	b	b	2.1 Getting from a to b
fong-spivak-seven-sketches_9_F00003	9	■ 0.986	$\mathcal{V}$	$\mathcal{V}$	2.3.4 $\mathcal{V}$ -variations on preorders and metric spaces
fong-spivak-seven-sketches_9_F00004	9	■ 1.000	Σ	Σ	3.4.3 Left and right pushforward functors, Σ and Π
fong-spivak-seven-sketches_9_F00005	9	■ 1.000	Π	Π	3.4.3 Left and right pushforward functors, Σ and Π
fong-spivak-seven-sketches_10_F00006	10	■ 0.991	$\mathbf{Mat}(R)$	$\mathbf{Mat}(R)$	5.4.1 A presentation of $\mathbf{Mat}(R)$
fong-spivak-seven-sketches_11_F00007	11	■ 1.000	Ω	Ω	7.4.1 The subobject classifier Ω in a sheaf topos
fong-spivak-seven-sketches_11_F00008	11	■ 0.545	$\mathbb{I}\mathbb{R}$	$\mathbb{I}\mathbb{R}$	7.5.2 Sheaves on $\mathbb{I}\mathbb{R}$
fong-spivak-seven-sketches_13_F00009	13	■ 0.999	$f: X \rightarrow Y$	$f: X \rightarrow Y$	maps. A map $f: X \rightarrow Y$ is a kind of observation of object X via a specified relationship
fong-spivak-seven-sketches_13_F00010	13	■ 0.999	X	X	maps. A map $f: X \rightarrow Y$ is a kind of observation of object X via a specified relationship
fong-spivak-seven-sketches_13_F00011	13	■ 1.000	Y	Y	it has with another object, Y . For example, think of X as the subject of an experiment
fong-spivak-seven-sketches_14_F00012	14	■ 1.000	f	f	Asking which aspects of X one wants to preserve under the observation f becomes
fong-spivak-seven-sketches_14_F00013	14	■ 1.000	$\mathbb{R}$	$\mathbb{R}$	functions f from $\mathbb{R}$ to $\mathbb{R}$, and we can think of them as observations: rather than view
fong-spivak-seven-sketches_14_F00014	14	■ 1.000	x	x	x “directly”, we only observe $f(x)$. Out of all the functions $f: \mathbb{R} \rightarrow \mathbb{R}$, only some of
fong-spivak-seven-sketches_14_F00015	14	■ 1.000	$f(x)$	$f(x)$	x “directly”, we only observe $f(x)$. Out of all the functions $f: \mathbb{R} \rightarrow \mathbb{R}$, only some of
fong-spivak-seven-sketches_14_F00016	14	■ 1.000	$f: \mathbb{R} \rightarrow \mathbb{R}$	$f: \mathbb{R} \rightarrow \mathbb{R}$	x “directly”, we only observe $f(x)$. Out of all the functions $f: \mathbb{R} \rightarrow \mathbb{R}$, only some of
fong-spivak-seven-sketches_14_F00017	14	■ 0.714	$x \leq y$	$x \leq y$	(a) <i>order-preserving</i> if $x \leq y$ implies $f(x) \leq f(y)$, for all $x, y \in \mathbb{R}$; ¹
fong-spivak-seven-sketches_14_F00018	14	■ 0.714	$f(x) \leq f(y)$	$f(x) \leq f(y)$	(a) <i>order-preserving</i> if $x \leq y$ implies $f(x) \leq f(y)$, for all $x, y \in \mathbb{R}$; ¹
fong-spivak-seven-sketches_14_F00019	14	■ 0.714	$x, y \in \mathbb{R}$	$x, y \in \mathbb{R}$	(a) <i>order-preserving</i> if $x \leq y$ implies $f(x) \leq f(y)$, for all $x, y \in \mathbb{R}$; ¹
fong-spivak-seven-sketches_14_F00020	14	■ 1.000	$ x-y = f(x)-f(y) $	$ x-y = f(x)-f(y) $	(b) <i>metric-preserving</i> if $ x-y = f(x)-f(y) $;
fong-spivak-seven-sketches_14_F00021	14	■ 1.000	$f(x+y)=f(x)+f(y)$	$f(x+y)=f(x)+f(y)$	(c) <i>addition-preserving</i> if $f(x+y)=f(x)+f(y)$.
fong-spivak-seven-sketches_14_F00022	14	—	$\square$	$\square$	—
fong-spivak-seven-sketches_14_F00023	14	■ 0.766	x, y	x, y	¹ We are often taught to view functions $f: \mathbb{R} \rightarrow \mathbb{R}$ as plots on an (x, y) -axis, where x is the domain

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_F00024	14	■ 1.000	y	y	(independent) variable and y is the codomain (dependent) variable. In this book, we do not adhere to
fong-spivak-seven-sketches_F00025	14	■ 1.000	$f(x)=x^2$	$f(x) = x^2$	example consider the function $f(x) = x^2$. Then f being order-preserving would say that for any $x, y \in \mathbb{R}$,
fong-spivak-seven-sketches_F00026	14	■ 1.000	$x^2 \leq y^2$	$x^2 \leq y^2$	if $x \leq y$ then $x^2 \leq y^2$; is that true?
fong-spivak-seven-sketches_F00027	15	■ 1.000	c	c	tions are also transitive, meaning that if a is connected to b , and b is connected to c , then
fong-spivak-seven-sketches_F00028	15	■ 1.000	a, b	a, b	a is connected to c ; that is, all a, b , and c are connected. Friendship is not transitive—my
fong-spivak-seven-sketches_F00029	15	■ 1.000	Φ	Φ	is observing this system. Her observation of interest, which we call Φ , extracts a single
fong-spivak-seven-sketches_F00030	16	■ 1.000	A	A	Joining our simple systems. Joining two systems A and B is performed simply by
fong-spivak-seven-sketches_F00031	16	■ 1.000	B	B	Joining our simple systems. Joining two systems A and B is performed simply by
fong-spivak-seven-sketches_F00032	16	■ 1.000	$A \vee B$	$A \vee B$	it $A \vee B$, has a connection between points x and y if there are some points $z_1, \dots, z_n$
fong-spivak-seven-sketches_F00033	16	■ 1.000	$z_{\{1\}}, \ldots, z_{\{n\}}$	$z_1, \dots, z_n$	it $A \vee B$, has a connection between points x and y if there are some points $z_1, \dots, z_n$
fong-spivak-seven-sketches_F00034	16	■ 0.562	$z_{\{1\}}$	z_1	such that each of the following are true in at least one of A or B : x is connected to z_1 ,
fong-spivak-seven-sketches_F00035	16	■ 1.000	$z_{\{i\}}$	z_i	z_i is connected to z_{i+1} , and z_n is connected to y . In a three-point system, the above
fong-spivak-seven-sketches_F00036	16	■ 1.000	$z_{\{i+1\}}$	z_{i+1}	z_i is connected to z_{i+1} , and z_n is connected to y . In a three-point system, the above
fong-spivak-seven-sketches_F00037	16	■ 1.000	$z_{\{n\}}$	z_n	z_i is connected to z_{i+1} , and z_n is connected to y . In a three-point system, the above
fong-spivak-seven-sketches_F00038	16	■ 0.506	$\Phi(\text{\textcircled{\tiny{O}}}) = \Phi(\text{\textcircled{\tiny{O}}}) =$	$\Phi(\text{\textcircled{\tiny{O}}}) = \Phi(\text{\textcircled{\tiny{O}}}) =$	We see above in Eq. (1.2) that $\Phi(\text{\textcircled{\tiny{O}}}) = \Phi(\text{\textcircled{\tiny{O}}}) = \text{false}$: in both cases $\bullet$ is not connected
fong-spivak-seven-sketches_F00039	16	■ 0.322	$\Phi(\text{\textcircled{\tiny{O}}} \vee \text{\textcircled{\tiny{O}}}) = \Phi(\text{\textcircled{\tiny{O}}} \vee \text{\textcircled{\tiny{O}}}) =$	$\Phi(\text{\textcircled{\tiny{O}}} \vee \text{\textcircled{\tiny{O}}}) = \Phi(\text{\textcircled{\tiny{O}}} \vee \text{\textcircled{\tiny{O}}}) =$	$\Phi(\text{\textcircled{\tiny{O}}} \vee \text{\textcircled{\tiny{O}}}) = \Phi(\text{\textcircled{\tiny{O}}} \vee \text{\textcircled{\tiny{O}}}) = \text{true}$: in the joined system, $\bullet$ is connected to $*$. The question that
fong-spivak-seven-sketches_F00040	16	■ 0.322	$\bullet$	$\bullet$	$\Phi(\text{\textcircled{\tiny{O}}} \vee \text{\textcircled{\tiny{O}}}) = \Phi(\text{\textcircled{\tiny{O}}} \vee \text{\textcircled{\tiny{O}}}) = \text{true}$: in the joined system, $\bullet$ is connected to $*$. The question that
fong-spivak-seven-sketches_F00041	17	■ 1.000	$A \leq B$	$A \leq B$	systems A and B , we say that $A \leq B$ if, whenever x is connected to y in A , then x is
fong-spivak-seven-sketches_F00042	17	■ 1.000	$\leq$	$\leq$	This notion of $\leq$ leads to the following diagram:
fong-spivak-seven-sketches_F00043	17	■ 1.000	$A \leq (A \vee B)$	$A \leq (A \vee B)$	smallest system that is bigger than both A and B . That is, $A \leq (A \vee B)$ and $B \leq (A \vee B)$,
fong-spivak-seven-sketches_F00044	17	■ 1.000	$B \leq (A \vee B)$	$B \leq (A \vee B)$	smallest system that is bigger than both A and B . That is, $A \leq (A \vee B)$ and $B \leq (A \vee B)$,
fong-spivak-seven-sketches_F00045	17	■ 0.999	C	C	and for any C , if $A \leq C$ and $B \leq C$ then $(A \vee B) \leq C$. Let’s walk through this with an
fong-spivak-seven-sketches_F00046	17	■ 0.999	$A \leq C$	$A \leq C$	and for any C , if $A \leq C$ and $B \leq C$ then $(A \vee B) \leq C$. Let’s walk through this with an
fong-spivak-seven-sketches_F00047	17	■ 0.999	$B \leq C$	$B \leq C$	and for any C , if $A \leq C$ and $B \leq C$ then $(A \vee B) \leq C$. Let’s walk through this with an
fong-spivak-seven-sketches_F00048	17	■ 0.999	$(A \vee B) \leq C$	$(A \vee B) \leq C$	and for any C , if $A \leq C$ and $B \leq C$ then $(A \vee B) \leq C$. Let’s walk through this with an
fong-spivak-seven-sketches_F00049	17	■ 0.993	$\{\bullet, *\}$	$\{\bullet, *\}$	1. Write down all the partitions of a two element set $\{\bullet, *\}$, order them as above,
fong-spivak-seven-sketches_F00050	18	■ 1.000	$\mathbb{B} = \{$	$\mathbb{B} = \{$	The set $\mathbb{B} = \{\text{true}, \text{false}\}$ of booleans also has an order, $\text{false} \leq \text{true}$:
fong-spivak-seven-sketches_F00051	18	■ 1.000	$\}$	$\}$	The set $\mathbb{B} = \{\text{true}, \text{false}\}$ of booleans also has an order, $\text{false} \leq \text{true}$:

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F00052	18	■ 1.000	<code>\not \leq</code>	$\not\leq$	Thus $\text{false} \leq \text{false}$, $\text{false} \leq \text{true}$, and $\text{true} \leq \text{true}$, but $\text{true} \not\leq \text{false}$. In other
fong-spivak-seven-sketches-F00053	18	■ 0.637	<code>B.^{2}</code>	$B.^2$	words, $A \leq B$ if A implies B . ²
fong-spivak-seven-sketches-F00054	18	■ 1.000	<code>A, B</code>	A, B	For any A, B in $\mathbb{B}$, we can again write $A \vee B$ to mean the least element that is greater
fong-spivak-seven-sketches-F00055	18	■ 1.000	<code>\mathbb{B}</code>	$\mathbb{B}$	For any A, B in $\mathbb{B}$, we can again write $A \vee B$ to mean the least element that is greater
fong-spivak-seven-sketches-F00056	18	■ 0.760	<code>\vee</code>	$\vee$	2. $\text{false} \vee \text{true}$?
fong-spivak-seven-sketches-F00057	18	■ 1.000	<code>\Phi(A)=\Phi(B)=</code>	$\Phi(A) = \Phi(B) =$	with $\Phi(A) = \Phi(B) = \text{false}$, so $\Phi(A) \vee \Phi(B) = \text{false}$, but where $\Phi(A \vee B) = \text{true}$. In
fong-spivak-seven-sketches-F00058	18	■ 1.000	<code>\Phi(A) \vee \Phi(B)=</code>	$\Phi(A) \vee \Phi(B) =$	with $\Phi(A) = \Phi(B) = \text{false}$, so $\Phi(A) \vee \Phi(B) = \text{false}$, but where $\Phi(A \vee B) = \text{true}$. In
fong-spivak-seven-sketches-F00059	18	■ 1.000	<code>\Phi(A \vee B)=</code>	$\Phi(A \vee B) =$	with $\Phi(A) = \Phi(B) = \text{false}$, so $\Phi(A) \vee \Phi(B) = \text{false}$, but where $\Phi(A \vee B) = \text{true}$. In
fong-spivak-seven-sketches-F00060	18	■ 1.000	<code>P</code>	P	² In mathematical logic, false implies true but true does not imply false . That is “ P implies Q ”
fong-spivak-seven-sketches-F00061	18	■ 1.000	<code>Q</code>	Q	² In mathematical logic, false implies true but true does not imply false . That is “ P implies Q ”
fong-spivak-seven-sketches-F00062	19	■ 1.000	<code>a, b, c</code>	a, b, c	certain tree, or the names of your favorite fruits, or simply some symbols a, b, c . For
fong-spivak-seven-sketches-F00063	19	■ 1.000	<code>A=\{h, 1\}</code>	$A = \{h, 1\}$	example, we write $A = \{h, 1\}$ to denote the set, called A , that contains exactly two
fong-spivak-seven-sketches-F00064	19	■ 0.911	<code>h</code>	h	elements, one called h and one called 1. The set $\{h, h, 1, h, 1\}$ is exactly the same as A
fong-spivak-seven-sketches-F00065	19	■ 0.911	<code>\{h, h, 1, h, 1\}</code>	$\{h, h, 1, h, 1\}$	elements, one called h and one called 1. The set $\{h, h, 1, h, 1\}$ is exactly the same as A
fong-spivak-seven-sketches-F00066	19	■ 1.000	<code>x \in X</code>	$x \in X$	than once in the notation doesn’t change the set. ³ For an arbitrary set X , we write $x \in X$
fong-spivak-seven-sketches-F00067	19	■ 1.000	<code>h \in A</code>	$h \in A$	if x is element of X ; so we have $h \in A$ and $1 \in A$, but $0 \notin A$.
fong-spivak-seven-sketches-F00068	19	■ 1.000	<code>1 \in A</code>	$1 \in A$	if x is element of X ; so we have $h \in A$ and $1 \in A$, but $0 \notin A$.
fong-spivak-seven-sketches-F00069	19	■ 1.000	<code>0 \notin A</code>	$0 \notin A$	if x is element of X ; so we have $h \in A$ and $1 \in A$, but $0 \notin A$.
fong-spivak-seven-sketches-F00070	19	■ 1.000	<code>\varnothing</code>	$\emptyset$	• $\emptyset$ denotes the empty set; it has no elements.
fong-spivak-seven-sketches-F00071	19	■ 0.999	<code>\mathbb{N}</code>	$\mathbb{N}$	• $\mathbb{N}$ denotes the set of <i>natural numbers</i> ; it has elements $0, 1, 2, 3, \dots, 90^{717}, \dots$
fong-spivak-seven-sketches-F00072	19	■ 0.999	<code>0,1,2,3, \ldots, 90^{717}, \ldots</code>	$0, 1, 2, 3, \dots, 90^{717}, \dots$	• $\mathbb{N}$ denotes the set of <i>natural numbers</i> ; it has elements $0, 1, 2, 3, \dots, 90^{717}, \dots$.
fong-spivak-seven-sketches-F00073	19	■ 0.992	<code>\underline{n}</code>	$\underline{n}$	• $\underline{n}$, for any $n \in \mathbb{N}$, denotes the n^{th} ordinal; it has n elements $1, 2, \dots, n$. For example,
fong-spivak-seven-sketches-F00074	19	■ 0.992	<code>n \in \mathbb{N}</code>	$n \in \mathbb{N}$	• $\underline{n}$, for any $n \in \mathbb{N}$, denotes the n^{th} ordinal; it has n elements $1, 2, \dots, n$. For example,
fong-spivak-seven-sketches-F00075	19	■ 0.992	<code>n^{\text{\text{th}}}</code>	n^{th}	• $\underline{n}$, for any $n \in \mathbb{N}$, denotes the n^{th} ordinal; it has n elements $1, 2, \dots, n$. For example,
fong-spivak-seven-sketches-F00076	19	■ 0.992	<code>n</code>	n	• $\underline{n}$, for any $n \in \mathbb{N}$, denotes the n^{th} ordinal; it has n elements $1, 2, \dots, n$. For example,
fong-spivak-seven-sketches-F00077	19	■ 0.992	<code>1,2, \ldots, n</code>	$1, 2, \dots, n$	• $\underline{n}$, for any $n \in \mathbb{N}$, denotes the n^{th} ordinal; it has n elements $1, 2, \dots, n$. For example,
fong-spivak-seven-sketches-F00078	19	■ 1.000	<code>\underline{0}=\varnothing, \underline{1}=\{1\}</code>	$\underline{0} = \emptyset, \underline{1} = \{1\}$	$\underline{0} = \emptyset, \underline{1} = \{1\}$, and $\underline{5} = \{1, 2, 3, 4, 5\}$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_19 F00079	19	■ 1.000	<code>\underline{5}=\{1,2,3,4,5\}</code>	$\underline{5} = \{1, 2, 3, 4, 5\}$	$\underline{0} = \emptyset, \underline{1} = \{1\}$, and $\underline{5} = \{1, 2, 3, 4, 5\}$.
fong-spivak-seven-sketches_19 F00080	19	■ 0.152	<code>\mathbb{Z}</code>	$\mathbb{Z}$	• $\mathbb{Z}$, the set of <i>integers</i> ; it has elements $\dots, -2, -1, 0, 1, 2, \dots, 90^{717}, \dots$
fong-spivak-seven-sketches_19 F00081	19	■ 0.152	<code>90^{717}, \ldots</code>	$90^{717}, \dots$	• $\mathbb{Z}$, the set of <i>integers</i> ; it has elements $\dots, -2, -1, 0, 1, 2, \dots, 90^{717}, \dots$
fong-spivak-seven-sketches_19 F00082	19	■ 1.000	<code>\pi, 3.14, 5 * \sqrt{2}, e, e^2, -1457, 90^{717}</code>	$\pi, 3.14, 5 * \sqrt{2}, e, e^2, -1457, 90^{717}$	• $\mathbb{R}$, the set of <i>real numbers</i> ; it has elements like $\pi, 3.14, 5 * \sqrt{2}, e, e^2, -1457, 90^{717}$, etc.
fong-spivak-seven-sketches_19 F00083	19	■ 1.000	<code>X \subseteq Y</code>	$X \subseteq Y$	Given sets X and Y , we say that X is a <i>subset</i> of Y , and write $X \subseteq Y$, if every element
fong-spivak-seven-sketches_19 F00084	19	■ 1.000	<code>\{h\} \subseteq A</code>	$\{h\} \subseteq A$	in X is also in Y . For example $\{h\} \subseteq A$. Note that the empty set $\emptyset := \{\}$ is a subset
fong-spivak-seven-sketches_19 F00085	19	■ 1.000	<code>\varnothing:=\{ \}</code>	$\emptyset := \{\}$	in X is also in Y . For example $\{h\} \subseteq A$. Note that the empty set $\emptyset := \{\}$ is a subset
fong-spivak-seven-sketches_19 F00086	19	■ 0.808	<code>\{y \in Y \mid P(y)\}</code>	$\{y \in Y \mid P(y)\}$	element of Y , we write $y \in Y \mid P(y)$ to mean the subset of those y 's that satisfy P .
fong-spivak-seven-sketches_19 F00087	19	■ 0.808	<code>y^{\prime}</code>	y'	element of Y , we write $\{y \in Y \mid P(y)\}$ to mean the subset of those y 's that satisfy P .
fong-spivak-seven-sketches_19 F00088	19	■ 1.000	<code>\mathbb{N}=\{n \in \mathbb{Z} \mid n \geq 0\}</code>	$\mathbb{N} = \{n \in \mathbb{Z} \mid n \geq 0\}$	1. Is it true that $\mathbb{N} = \{n \in \mathbb{Z} \mid n \geq 0\}$?
fong-spivak-seven-sketches_19 F00089	19	■ 1.000	<code>\mathbb{N}=\{n \in \mathbb{Z} \mid n \geq 1\}</code>	$\mathbb{N} = \{n \in \mathbb{Z} \mid n \geq 1\}$	2. Is it true that $\mathbb{N} = \{n \in \mathbb{Z} \mid n \geq 1\}$?
fong-spivak-seven-sketches_19 F00090	19	■ 1.000	<code>\varnothing=\{n \in \mathbb{Z} \mid 1 < n < 2\}</code>	$\emptyset = \{n \in \mathbb{Z} \mid 1 < n < 2\}$	3. Is it true that $\emptyset = \{n \in \mathbb{Z} \mid 1 < n < 2\}$?
fong-spivak-seven-sketches_19 F00091	19	■ 1.000	<code>X_{1}</code>	X_1	If both X_1 and X_2 are subsets of Y , their <i>union</i> , denoted $X_1 \cup X_2$, is also a subset of
fong-spivak-seven-sketches_19 F00092	19	■ 1.000	<code>X_{2}</code>	X_2	If both X_1 and X_2 are subsets of Y , their <i>union</i> , denoted $X_1 \cup X_2$, is also a subset of
fong-spivak-seven-sketches_19 F00093	19	■ 1.000	<code>X_{1} \cup X_{2}</code>	$X_1 \cup X_2$	If both X_1 and X_2 are subsets of Y , their <i>union</i> , denoted $X_1 \cup X_2$, is also a subset of
fong-spivak-seven-sketches_19 F00094	19	■ 0.988	<code>h, 1</code>	$h, 1$	³ If you want a notion where “ $h, 1$ ” is different than “ $h, h, 1, h, 1$ ”, you can use something called <i>bags</i> ,
fong-spivak-seven-sketches_19 F00095	19	■ 0.988	<code>h, h, 1, h, 1</code>	$h, h, 1, h, 1$	³ If you want a notion where “ $h, 1$ ” is different than “ $\overline{h, h, 1, h, 1}$ ”, you can use something called <i>bags</i> ,
fong-spivak-seven-sketches_19 F00096	19	■ 0.829	<code>Z:=</code>	$Z :=$	⁴ When we write $Z := \text{foo}$, it means “assign the meaning <i>foo</i> to variable Z ”, whereas $Z = \text{foo}$ means
fong-spivak-seven-sketches_19 F00097	19	■ 1.000	<code>Z</code>	Z	simply that Z is equal to <i>foo</i> , perhaps as discovered via some calculation. In particular, $Z := \text{foo}$ implies
fong-spivak-seven-sketches_19 F00098	19	■ 1.000	<code>Z=</code>	$Z =$	$Z = \text{foo}$ but not vice versa; indeed it <i>would not</i> be proper to write $3 + 2 := 5$ or $\{\} := \emptyset$.
fong-spivak-seven-sketches_19 F00099	19	■ 1.000	<code>3+2:=5</code>	$3 + 2 := 5$	$Z = \text{foo}$ but not vice versa; indeed it <i>would not</i> be proper to write $3 + 2 := 5$ or $\{\} := \emptyset$.
fong-spivak-seven-sketches_19 F00100	19	■ 1.000	<code>\}:=\varnothing</code>	$\} := \emptyset$	$Z = \text{foo}$ but not vice versa; indeed it <i>would not</i> be proper to write $3 + 2 := 5$ or $\{\} := \emptyset$.
fong-spivak-seven-sketches_20 F00101	20	■ 1.000	<code>Y=\{1,2,3,4\}</code>	$Y = \{1, 2, 3, 4\}$	For example if $Y = \{1, 2, 3, 4\}$ and $X_1 = \{1, 2\}$ and $X_2 = \{2, 4\}$, then $X_1 \cup X_2 = \{1, 2, 4\}$.
fong-spivak-seven-sketches_20 F00102	20	■ 1.000	<code>X_{1}=\{1,2\}</code>	$X_1 = \{1, 2\}$	For example if $Y = \{1, 2, 3, 4\}$ and $X_1 = \{1, 2\}$ and $X_2 = \{2, 4\}$, then $X_1 \cup X_2 = \{1, 2, 4\}$.
fong-spivak-seven-sketches_20 F00103	20	■ 1.000	<code>X_{2}=\{2,4\}</code>	$X_2 = \{2, 4\}$	For example if $Y = \{1, 2, 3, 4\}$ and $X_1 = \{1, 2\}$ and $X_2 = \{2, 4\}$, then $X_1 \cup X_2 = \{1, 2, 4\}$.
fong-spivak-seven-sketches_20 F00104	20	■ 1.000	<code>X_{1} \cup X_{2}=\{1,2,4\}</code>	$X_1 \cup X_2 = \{1, 2, 4\}$	For example if $Y = \{1, 2, 3, 4\}$ and $X_1 = \{1, 2\}$ and $X_2 = \{2, 4\}$, then $X_1 \cup X_2 = \{1, 2, 4\}$.
fong-spivak-seven-sketches_20 F00105	20	■ 1.000	<code>\varnothing \cup X=X</code>	$\emptyset \cup X = X$	Note that $\emptyset \cup X = X$ for any $X \subseteq Y$.
fong-spivak-seven-sketches_20 F00106	20	■ 0.997	<code>X_{1} \cap X_{2}</code>	$X_1 \cap X_2$	Similarly, if both X_1 and X_2 are subsets of Y , then their <i>intersection</i> , denoted $X_1 \cap X_2$,

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_20 F00107	20	■ 1.000	$\{1,2,3\} \cap \{2,5\} = \{2\}$	$\{1,2,3\} \cap \{2,5\} = \{2\}$	X_1 and in X_2 , and no others. So $\{1,2,3\} \cap \{2,5\} = \{2\}$.
fong-spivak-seven-sketches_20 F00108	20	■ 1.000	$X_{\{0\}} = \varnothing, X_{\{1\}} = \{1\}, X_{\{2\}} = \{1,2\}$	$X_0 = \varnothing, X_1 = \{1\}, X_2 = \{1,2\}$	sets $X_0 = \varnothing, X_1 = \{1\}, X_2 = \{1,2\}$, etc. as subsets of $\mathbb{N}$, and we want to know what the
fong-spivak-seven-sketches_20 F00109	20	■ 0.984	$\bigcup_{n \in \mathbb{N}} X_n$	$\bigcup_{n \in \mathbb{N}} X_n$	union of all of them is. This union is written $\bigcup_{n \in \mathbb{N}} X_n$, and it is the subset of $\mathbb{N}$ that
fong-spivak-seven-sketches_20 F00110	20	■ 1.000	$X_{\{n\}}$	X_n	contains every element of every X_n , but no others. Namely, $\bigcup_{n \in \mathbb{N}} X_n = \{n \in \mathbb{N} \mid n \geq 1\}$.
fong-spivak-seven-sketches_20 F00111	20	■ 1.000	$\bigcup_{n \in \mathbb{N}} X_n = \{n \in \mathbb{N} \mid n \geq 1\}$	$\bigcup_{n \in \mathbb{N}} X_n = \{n \in \mathbb{N} \mid n \geq 1\}$	contains every element of every X_n , but no others. Namely, $\bigcup_{n \in \mathbb{N}} X_n = \{n \in \mathbb{N} \mid n \geq 1\}$.
fong-spivak-seven-sketches_20 F00112	20	■ 1.000	$\bigcap_{n \in \mathbb{N}} X_n$	$\bigcap_{n \in \mathbb{N}} X_n$	Similarly one can write $\bigcap_{n \in \mathbb{N}} X_n$ for the intersection of all of them, which will be empty
fong-spivak-seven-sketches_20 F00113	20	■ 0.715	$X \times Y$	$X \times Y$	Given two sets X and Y , the <i>product</i> $X \times Y$ of X and Y is the set of pairs (x, y) , where
fong-spivak-seven-sketches_20 F00114	20	■ 1.000	$y \in Y$	$y \in Y$	$x \in X$ and $y \in Y$.
fong-spivak-seven-sketches_20 F00115	20	■ 1.000	$X \sqcup Y$	$X \sqcup Y$	in common. Given two sets X and Y , their <i>disjoint union</i> $X \sqcup Y$ is the set of pairs of the
fong-spivak-seven-sketches_20 F00116	20	■ 0.994	$(x, 1)$	$(x, 1)$	form $(x, 1)$ or $(y, 2)$, where $x \in X$ and $y \in Y$.
fong-spivak-seven-sketches_20 F00117	20	■ 0.994	$(y, 2)$	$(y, 2)$	form $(x, 1)$ or $(y, 2)$, where $x \in X$ and $y \in Y$.
fong-spivak-seven-sketches_20 F00118	20	■ 1.000	$A := \{h, 1\}$	$A := \{h, 1\}$	<i>Exercise 1.11.</i> Let $A := \{h, 1\}$ and $B := \{1, 2, 3\}$.
fong-spivak-seven-sketches_20 F00119	20	■ 1.000	$B := \{1, 2, 3\}$	$B := \{1, 2, 3\}$	<i>Exercise 1.11.</i> Let $A := \{h, 1\}$ and $B := \{1, 2, 3\}$.
fong-spivak-seven-sketches_20 F00120	20	■ 1.000	$A \times B$	$A \times B$	3. There are six elements in $A \times B$; write them out.
fong-spivak-seven-sketches_20 F00121	20	■ 1.000	$A \sqcup B$	$A \sqcup B$	4. There are five elements of $A \sqcup B$; write them out.
fong-spivak-seven-sketches_20 F00122	20	■ 1.000	$A \cup B$	$A \cup B$	$A \cup B$; write them out.
fong-spivak-seven-sketches_20 F00123	20	■ 0.999	$R \subseteq X \times Y$	$R \subseteq X \times Y$	Definition 1.12. Let X and Y be sets. A <i>relation between X and Y</i> is a subset $R \subseteq X \times Y$.
fong-spivak-seven-sketches_20 F00124	20	■ 0.973	$R \subseteq X \times X$	$R \subseteq X \times X$	A <i>binary relation on X</i> is a relation between X and X , i.e. a subset $R \subseteq X \times X$.
fong-spivak-seven-sketches_20 F00125	20	■ 1.000	$R \subseteq A \times A$	$R \subseteq A \times A$	It is convenient to use something called <i>infix notation</i> for binary relations $R \subseteq A \times A$.
fong-spivak-seven-sketches_20 F00126	20	■ 0.651	$a \star b$	$a \star b$	This means one picks a symbol, say $\star$, and writes $a \star b$ to mean $(a, b) \in R$.
fong-spivak-seven-sketches_20 F00127	20	■ 0.651	$(a, b) \in R$	$(a, b) \in R$	This means one picks a symbol, say $\star$, and writes $a \star b$ to mean $(a, b) \in R$.
fong-spivak-seven-sketches_20 F00128	20	■ 1.000	$(5, 6) \in R$	$(5, 6) \in R$	$(5, 6) \in R$, we write $5 \leq 6$.
fong-spivak-seven-sketches_20 F00129	20	■ 1.000	$5 \leq 6$	$5 \leq 6$	$(5, 6) \in R$, we write $5 \leq 6$.
fong-spivak-seven-sketches_20 F00130	20	■ 0.975	$=, \approx, <, >$	$=, \approx, <, >$	Other examples of infix notation for relations are $=, \approx, <, >$. In number theory, they
fong-spivak-seven-sketches_21 F00131	21	■ 1.000	$p \in P$	$p \in P$	Definition 1.14. If A is a set, a <i>partition</i> of A consists of a set P and, for each $p \in P$, a
fong-spivak-seven-sketches_21 F00132	21	■ 1.000	$A_p \subseteq A$	$A_p \subseteq A$	nonempty subset $A_p \subseteq A$, such that
fong-spivak-seven-sketches_21 F00133	21	■ 1.000	$\{A_p\}_{p \in P}$	$\{A_p\}_{p \in P}$	We may denote the partition by $\{A_p\}_{p \in P}$. We refer to P as the set of <i>part labels</i> and if

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_1 F00161	22	■ 1.000	$A_{\{p\}} \cap A_{\{q\}} = \varnothing$	$A_p \cap A_q = \emptyset$	2. Show that if $p \neq q$, i.e. if A_p and A_q are not exactly the same set, then $A_p \cap A_q = \emptyset$.
fong-spivak-seven-sketches_2 F00162	22	■ 0.562	$A = \bigcup_{p \in P} A_{\{p\}}$	$A = \bigcup_{p \in P} A_p$	3. Show that $A = \bigcup_{p \in P} A_p$.
fong-spivak-seven-sketches_3 F00163	22	■ 1.000	$A / \sim$	$A / \sim$	<i>quotient</i> $A / \sim$ of A under $\sim$ is the set of parts of the corresponding partition.
fong-spivak-seven-sketches_4 F00164	22	■ 1.000	S	S	Definition 1.22. Let S and T be sets. A <i>function from S to T</i> is a subset $F \subseteq S \times T$ such
fong-spivak-seven-sketches_5 F00165	22	■ 1.000	T	T	Definition 1.22. Let S and T be sets. A <i>function from S to T</i> is a subset $F \subseteq S \times T$ such
fong-spivak-seven-sketches_6 F00166	22	■ 1.000	$F \subseteq S \times T$	$F \subseteq S \times T$	Definition 1.22. Let S and T be sets. A <i>function from S to T</i> is a subset $F \subseteq S \times T$ such
fong-spivak-seven-sketches_7 F00167	22	■ 1.000	$s \in S$	$s \in S$	that for all $s \in S$ there exists a unique $t \in T$ with $(s, t) \in F$.
fong-spivak-seven-sketches_8 F00168	22	■ 1.000	$t \in T$	$t \in T$	that for all $s \in S$ there exists a unique $t \in T$ with $(s, t) \in F$.
fong-spivak-seven-sketches_9 F00169	22	■ 1.000	$(s, t) \in F$	$(s, t) \in F$	that for all $s \in S$ there exists a unique $t \in T$ with $(s, t) \in F$.
fong-spivak-seven-sketches_10 F00170	22	■ 1.000	F	F	The function F is often denoted $F: S \rightarrow T$. From now on, we write $F(s) = t$, or
fong-spivak-seven-sketches_11 F00171	22	■ 1.000	$F: S \rightarrow T$	$F: S \rightarrow T$	The function F is often denoted $F: S \rightarrow T$. From now on, we write $F(s) = t$, or
fong-spivak-seven-sketches_12 F00172	22	■ 1.000	$F(s)=t$	$F(s) = t$	The function F is often denoted $F: S \rightarrow T$. From now on, we write $F(s) = t$, or
fong-spivak-seven-sketches_13 F00173	22	■ 1.000	$s \mapsto t$	$s \mapsto t$	sometimes $s \mapsto t$, to mean $(s, t) \in F$. For any $t \in T$, the <i>preimage of t along F</i> is the
fong-spivak-seven-sketches_14 F00174	22	■ 1.000	t	t	sometimes $s \mapsto t$, to mean $(s, t) \in F$. For any $t \in T$, the <i>preimage of t along F</i> is the
fong-spivak-seven-sketches_15 F00175	22	■ 1.000	$\{s \in S \mid F(s)=t\}$	$\{s \in S \mid F(s) = t\}$	subset $\{s \in S \mid F(s) = t\}$.
fong-spivak-seven-sketches_16 F00176	22	■ 1.000	$s_{\{1\}}, s_{\{2\}} \in S$	$s_1, s_2 \in S$	$F(s) = t$. A function is called <i>injective</i> , or an <i>injection</i> , if for all $t \in T$ and $s_1, s_2 \in S$
fong-spivak-seven-sketches_17 F00177	22	■ 1.000	$F(s_{\{1\}})=t$	$F(s_1) = t$	with $F(s_1) = t$ and $F(s_2) = t$, we have $s_1 = s_2$. A function is called <i>bijective</i> if it is both
fong-spivak-seven-sketches_18 F00178	22	■ 1.000	$F(s_{\{2\}})=t$	$F(s_2) = t$	with $F(s_1) = t$ and $F(s_2) = t$, we have $s_1 = s_2$. A function is called <i>bijective</i> if it is both
fong-spivak-seven-sketches_19 F00179	22	■ 1.000	$s_{\{1\}}=s_{\{2\}}$	$s_1 = s_2$	with $F(s_1) = t$ and $F(s_2) = t$, we have $s_1 = s_2$. A function is called <i>bijective</i> if it is both
fong-spivak-seven-sketches_20 F00180	23	■ 0.413	$\rightarrow, \rightarrowtail, \twoheadrightarrow, \stackrel{\cong}{\rightarrow}$	$\rightarrow, \rightarrowtail, \twoheadrightarrow, \stackrel{\cong}{\rightarrow}$	We use various decorations on arrows, $\rightarrow, \rightarrowtail, \twoheadrightarrow, \stackrel{\cong}{\rightarrow}$ to denote these special sorts of
fong-spivak-seven-sketches_21 F00181	23	■ 0.832	id_X	id_X	a set X , denoted id_X . It is the bijective function $\mathrm{id}_X(x) = x$.
fong-spivak-seven-sketches_22 F00182	23	■ 0.832	$\operatorname{id}_X(x)=x$	$\mathrm{id}_X(x) = x$	a set X , denoted id_X . It is the bijective function $\mathrm{id}_X(x) = x$.
fong-spivak-seven-sketches_23 F00183	23	■ 1.000	$f: A \rightarrow B$	$f: A \rightarrow B$	1. Find two sets A and B and a function $f: A \rightarrow B$ that is injective but not surjective.
fong-spivak-seven-sketches_24 F00184	23	■ 1.000	$f: A \rightarrow \varnothing$	$f: A \rightarrow \emptyset$	<i>Exercise 1.25.</i> Suppose that A is a set and $f: A \rightarrow \emptyset$ is a function to the empty set.
fong-spivak-seven-sketches_25 F00185	23	■ 1.000	$f: A \rightarrow P$	$f: A \rightarrow P$	functions out of A . Given a surjective function $f: A \twoheadrightarrow P$, where P is any other set, the
fong-spivak-seven-sketches_26 F00186	23	■ 1.000	$f^{-1}(p) \subseteq A$	$f^{-1}(p) \subseteq A$	preimages $f^{-1}(p) \subseteq A$, one for each element $p \in P$, form a partition of A . Here is an
fong-spivak-seven-sketches_27 F00187	24	■ 1.000	$S:=\{11,12,13,21,22,23\}$	$S := \{11, 12, 13, 21, 22, 23\}$	Consider the partition of $S := \{11, 12, 13, 21, 22, 23\}$ shown below:

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches- F00188	24	■ 1.000	$P=\{a, b, c, d\}$	$P = \{a, b, c, d\}$	It has been partitioned into four parts, so let $P = \{a, b, c, d\}$ and let $f: S \twoheadrightarrow P$ be given
fong-spivak-seven-sketches- F00189	24	■ 1.000	$f: S \rightarrowtail P$	$f: S \rightarrow P$	It has been partitioned into four parts, so let $P = \{a, b, c, d\}$ and let $f: S \twoheadrightarrow P$ be given
fong-spivak-seven-sketches- F00190	24	■ 1.000	$F: X \rightarrowtail Y$	$F: X \rightarrow Y$	Definition 1.28. If $F: X \rightarrow Y$ is a function and $G: Y \rightarrow Z$ is a function, their <i>composite</i>
fong-spivak-seven-sketches- F00191	24	■ 1.000	$G: Y \rightarrowtail Z$	$G: Y \rightarrow Z$	Definition 1.28. If $F: X \rightarrow Y$ is a function and $G: Y \rightarrow Z$ is a function, their <i>composite</i>
fong-spivak-seven-sketches- F00192	24	■ 1.000	$X \rightarrowtail Z$	$X \rightarrow Z$	is the function $X \rightarrow Z$ defined to be $G(F(x))$ for any $x \in X$. It is often denoted $G \circ F$, but
fong-spivak-seven-sketches- F00193	24	■ 1.000	$G(F(x))$	$G(F(x))$	is the function $X \rightarrow Z$ defined to be $G(F(x))$ for any $x \in X$. It is often denoted $G \circ F$, but
fong-spivak-seven-sketches- F00194	24	■ 1.000	$G \circ F$	$G \circ F$	is the function $X \rightarrow Z$ defined to be $G(F(x))$ for any $x \in X$. It is often denoted $G \circ F$, but
fong-spivak-seven-sketches- F00195	24	■ 1.000	$F; G$	$F; G$	we prefer to denote it $F \circ G$. It takes any element $x \in X$, evaluates F to get an element
fong-spivak-seven-sketches- F00196	24	■ 1.000	$F(x) \in Y$	$F(x) \in Y$	$F(x) \in Y$ and then evaluates G to get an element $G(F(x))$.
fong-spivak-seven-sketches- F00197	24	■ 1.000	G	G	$F(x) \in Y$ and then evaluates G to get an element $G(F(x))$.
fong-spivak-seven-sketches- F00198	24	■ 1.000	$\{1\} \rightarrowtail X$	$\{1\} \rightarrow X$	$\{1\} \rightarrow X$, namely the function sending 1 to x . For example, the three functions
fong-spivak-seven-sketches- F00199	24	■ 0.974	$\{1\} \rightarrowtail \{1,2,3\}$	$\{1\} \rightarrow \{1,2,3\}$	$\{1\} \rightarrow \{1,2,3\}$ shown below correspond to the three elements of $\{1,2,3\}$:
fong-spivak-seven-sketches- F00200	24	■ 0.974	$\{1,2,3\}$	$\{1,2,3\}$	$\{1\} \rightarrow \{1,2,3\}$ shown below correspond to the three elements of $\{1,2,3\}$:
fong-spivak-seven-sketches- F00201	24	■ 1.000	$x:\{1\} \rightarrowtail X$	$x: \{1\} \rightarrow X$	$x: \{1\} \rightarrow X$. Then evaluating F at x is given by the composite, $F(x) = x \circ F$.
fong-spivak-seven-sketches- F00202	24	■ 1.000	$F(x)=x; F$	$F(x) = x; F$	$x: \{1\} \rightarrow X$. Then evaluating F at x is given by the composite, $F(x) = x \circ F$.
fong-spivak-seven-sketches- F00203	24	■ 1.000	$x \leq x$	$x \leq x$	(a) $x \leq x$; and
fong-spivak-seven-sketches- F00204	24	■ 1.000	$y \leq z$	$y \leq z$	(b) if $x \leq y$ and $y \leq z$, then $x \leq z$.
fong-spivak-seven-sketches- F00205	24	■ 1.000	$x \leq z$	$x \leq z$	(b) if $x \leq y$ and $y \leq z$, then $x \leq z$.
fong-spivak-seven-sketches- F00206	25	■ 0.608	$y \leq x$	$y \leq x$	$y \leq x$, we write $x \cong y$ and say x and y are <i>equivalent</i> . We call a pair $(X, \leq)$ consisting
fong-spivak-seven-sketches- F00207	25	■ 0.608	$x \cong y$	$x \cong y$	$y \leq x$, we write $x \cong y$ and say x and y are <i>equivalent</i> . We call a pair $(X, \leq)$ consisting
fong-spivak-seven-sketches- F00208	25	■ 0.608	$(X, \leq)$	$(X, \leq)$	$y \leq x$, we write $x \cong y$ and say x and y are <i>equivalent</i> . We call a pair $(X, \leq)$ consisting
fong-spivak-seven-sketches- F00209	25	■ 0.999	$(X, =)$	$(X, =)$	$(X, =)$. This means that the only order relations on X are of the form $x \leq x$; if $x \neq y$
fong-spivak-seven-sketches- F00210	25	■ 0.999	$x \neq y$	$x \neq y$	$(X, =)$. This means that the only order relations on X are of the form $x \leq x$; if $x \neq y$
fong-spivak-seven-sketches- F00211	25	■ 0.924	$X, \leq$	$X, \leq$	preorder $(X, \leq)$ by equipping it with the total binary relation $X \times X \subseteq X \times X$. This is a
fong-spivak-seven-sketches- F00212	25	■ 0.924	$X \times X \subseteq X \times X$	$X \times X \subseteq X \times X$	preorder $(X, \leq)$ by equipping it with the total binary relation $X \times X \subseteq X \times X$. This is a
fong-spivak-seven-sketches- F00213	25	■ 1.000	$x=y$	$x = y$	(c) $x \cong y$ implies $x = y$.
fong-spivak-seven-sketches- F00214	26	■ 1.000	$G=(V, A, s, t)$	$G = (V, A, s, t)$	Definition 1.36. A graph $G = (V, A, s, t)$ consists of a set V whose elements are called

Identifier	Page	Conf.	LaTeX source	Rendered	Image																		
fong-spivak-seven-sketches-F00215	26	1.000	V	V	Definition 1.36. A graph $G = (V, A, s, t)$ consists of a set V whose elements are called																		
fong-spivak-seven-sketches-F00216	26	0.999	$s, t: A \rightarrow V$	$s, t: A \rightarrow V$	<i>vertices</i> , a set A whose elements are called <i>arrows</i> , and two functions $s, t: A \rightarrow V$ known																		
fong-spivak-seven-sketches-F00217	26	1.000	$s(a)=v$	$s(a) = v$	as the <i>source</i> and <i>target</i> functions respectively. Given $a \in A$ with $s(a) = v$ and $t(a) = w$,																		
fong-spivak-seven-sketches-F00218	26	1.000	$t(a)=w$	$t(a) = w$	as the <i>source</i> and <i>target</i> functions respectively. Given $a \in A$ with $s(a) = v$ and $t(a) = w$,																		
fong-spivak-seven-sketches-F00219	26	1.000	v	v	we say that a is an arrow from v to w .																		
fong-spivak-seven-sketches-F00220	26	1.000	w	w	we say that a is an arrow from v to w .																		
fong-spivak-seven-sketches-F00221	26	1.000	$V=\{1,2,3,4\}$	$V = \{1, 2, 3, 4\}$	It has $V = \{1, 2, 3, 4\}$ and $A = \{a, b, c, d, e\}$. The source and target functions, $s, t: A \rightarrow$																		
fong-spivak-seven-sketches-F00222	26	1.000	$A=\{a, b, c, d, e\}$	$A = \{a, b, c, d, e\}$	It has $V = \{1, 2, 3, 4\}$ and $A = \{a, b, c, d, e\}$. The source and target functions, $s, t: A \rightarrow$																		
fong-spivak-seven-sketches-F00223	26	1.000	$s, t: A \rightarrow$	$s, t: A \rightarrow$	It has $V = \{1, 2, 3, 4\}$ and $A = \{a, b, c, d, e\}$. The source and target functions, $s, t: A \rightarrow$																		
fong-spivak-seven-sketches-F00224	26	0.630	$s(a) \in V$	$s(a) \in V$	<table><tr><th>arrow a</th><th>source $s(a) \in V$</th><th>target $t(a) \in V$</th></tr><tr><td>a</td><td>1</td><td>?</td></tr><tr><td>b</td><td>1</td><td>3</td></tr><tr><td>c</td><td>?</td><td>?</td></tr><tr><td>d</td><td>?</td><td>?</td></tr><tr><td>e</td><td>?</td><td>?</td></tr></table>	arrow a	source $s(a) \in V$	target $t(a) \in V$	a	1	?	b	1	3	c	?	?	d	?	?	e	?	?
arrow a	source $s(a) \in V$	target $t(a) \in V$																					
a	1	?																					
b	1	3																					
c	?	?																					
d	?	?																					
e	?	?																					
fong-spivak-seven-sketches-F00225	26	0.630	$t(a) \in V$	$t(a) \in V$	<table><tr><th>arrow a</th><th>source $s(a) \in V$</th><th>target $t(a) \in V$</th></tr><tr><td>a</td><td>1</td><td>?</td></tr><tr><td>b</td><td>1</td><td>3</td></tr><tr><td>c</td><td>?</td><td>?</td></tr><tr><td>d</td><td>?</td><td>?</td></tr><tr><td>e</td><td>?</td><td>?</td></tr></table>	arrow a	source $s(a) \in V$	target $t(a) \in V$	a	1	?	b	1	3	c	?	?	d	?	?	e	?	?
arrow a	source $s(a) \in V$	target $t(a) \in V$																					
a	1	?																					
b	1	3																					
c	?	?																					
d	?	?																					
e	?	?																					
fong-spivak-seven-sketches-F00226	26	0.630	d	d	<table><tr><th>arrow a</th><th>source $s(a) \in V$</th><th>target $t(a) \in V$</th></tr><tr><td>a</td><td>1</td><td>?</td></tr><tr><td>b</td><td>1</td><td>3</td></tr><tr><td>c</td><td>?</td><td>?</td></tr><tr><td>d</td><td>?</td><td>?</td></tr><tr><td>e</td><td>?</td><td>?</td></tr></table>	arrow a	source $s(a) \in V$	target $t(a) \in V$	a	1	?	b	1	3	c	?	?	d	?	?	e	?	?
arrow a	source $s(a) \in V$	target $t(a) \in V$																					
a	1	?																					
b	1	3																					
c	?	?																					
d	?	?																					
e	?	?																					
fong-spivak-seven-sketches-F00227	26	0.630	e	e	<table><tr><th>arrow a</th><th>source $s(a) \in V$</th><th>target $t(a) \in V$</th></tr><tr><td>a</td><td>1</td><td>?</td></tr><tr><td>b</td><td>1</td><td>3</td></tr><tr><td>c</td><td>?</td><td>?</td></tr><tr><td>d</td><td>?</td><td>?</td></tr><tr><td>e</td><td>?</td><td>?</td></tr></table>	arrow a	source $s(a) \in V$	target $t(a) \in V$	a	1	?	b	1	3	c	?	?	d	?	?	e	?	?
arrow a	source $s(a) \in V$	target $t(a) \in V$																					
a	1	?																					
b	1	3																					
c	?	?																					
d	?	?																					
e	?	?																					
fong-spivak-seven-sketches-F00228	26	1.000	$1 \rightarrow 2$	$1 \rightarrow 2$	are infinitely many paths $1 \rightarrow 2$ because one can loop and loop and loop through d as																		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_26 F00229	26	0.833	$(P, \leq)$	$(P, \leq)$	graph $G = (V, A, s, t)$ that gives a <i>presentation</i> of a preorder $(P, \leq)$. The elements of P
fong-spivak-seven-sketches_26 F00230	26	0.842	$v \leq w$	$v \leq w$	are the vertices V in G , and the order $\leq$ is given by $v \leq w$ iff ⁶ there is a path $v \rightarrow w$.
fong-spivak-seven-sketches_26 F00231	26	0.842	$^{\{6\}}$	6	are the vertices V in G , and the order $\leq$ is given by $v \leq w$ iff ⁶ there is a path $v \rightarrow w$.
fong-spivak-seven-sketches_26 F00232	26	0.842	$v \rightarrowtail w$	$v \rightarrow w$	are the vertices V in G , and the order $\leq$ is given by $v \leq w$ iff ⁶ there is a path $v \rightarrow w$.
fong-spivak-seven-sketches_26 F00233	26	1.000	$v \rightarrowtail v$	$v \rightarrow v$	For any vertex v , there is always a path $v \rightarrow v$, and this translates into the reflexivity
fong-spivak-seven-sketches_26 F00234	26	1.000	$u \rightarrowtail v$	$u \rightarrow v$	law from Definition 1.30. The fact that paths $u \rightarrow v$ and $v \rightarrow w$ can be concatenated to
fong-spivak-seven-sketches_26 F00235	26	1.000	$u \rightarrowtail w$	$u \rightarrow w$	a path  translates into the transitivity law.
fong-spivak-seven-sketches_27 F00236	27	0.959	$\left(p_{\{1\}}, p_{\{2\}}\right)$	(p_1, p_2)	That is, what are the elements of P and write down every pair (p_1, p_2) for which $p_1 \leq p_2$.
fong-spivak-seven-sketches_27 F00237	27	0.959	$p_{\{1\}} \leq p_{\{2\}}$	$p_1 \leq p_2$	That is, what are the elements of P and write down every pair (p_1, p_2) for which $p_1 \leq p_2$.
fong-spivak-seven-sketches_27 F00238	27	0.999	$\{\bullet, \circ, *\}$	$\{\bullet, \circ, *\}$	<i>Exercise 1.42.</i> Let X be the set of partitions of $\{\bullet, \circ, *\}$; it has five elements and an
fong-spivak-seven-sketches_27 F00239	27	0.929	(x, y)	(x, y)	(x, y) of elements in X such that $x \leq y$. There should be 12.
fong-spivak-seven-sketches_27 F00240	27	1.000	$\mathbb{N} := \{0, 1, 2, 3, \ldots\}$	$\mathbb{N} := \{0, 1, 2, 3, \ldots\}$	<i>Example 1.45</i> (Natural numbers). The natural numbers  are a pre-
fong-spivak-seven-sketches_27 F00241	27	1.000	$0 \leq 1$	$0 \leq 1$	order with the order given by the usual size ordering, e.g. $0 \leq 1$ and $5 \leq 100$. This is
fong-spivak-seven-sketches_27 F00242	27	1.000	$5 \leq 100$	$5 \leq 100$	order with the order given by the usual size ordering, e.g. $0 \leq 1$ and $5 \leq 100$. This is
fong-spivak-seven-sketches_27 F00243	27	1.000	$m \leq n$	$m \leq n$	a total order: either $m \leq n$ or $n \leq m$ for all m, n . One can see that its Hasse diagram
fong-spivak-seven-sketches_27 F00244	27	1.000	$n \leq m$	$n \leq m$	a total order: either $m \leq n$ or $n \leq m$ for all m, n . One can see that its Hasse diagram
fong-spivak-seven-sketches_27 F00245	27	1.000	m, n	m, n	a total order: either $m \leq n$ or $n \leq m$ for all m, n . One can see that its Hasse diagram
fong-spivak-seven-sketches_27 F00246	27	1.000	$a \leq b$	$a \leq b$	and b —namely all those in the middle row—which satisfy neither $a \leq b$ nor $b \leq a$.
fong-spivak-seven-sketches_27 F00247	27	1.000	$b \leq a$	$b \leq a$	and b —namely all those in the middle row—which satisfy neither $a \leq b$ nor $b \leq a$.
fong-spivak-seven-sketches_27 F00248	27	1.000	$n=m$	$n = m$	$n = m$. Another ordering is the reverse ordering, like $5 \leq 3$ and $3 \leq 2$, like how golf is
fong-spivak-seven-sketches_27 F00249	27	1.000	$5 \leq 3$	$5 \leq 3$	$n = m$. Another ordering is the reverse ordering, like $5 \leq 3$ and $3 \leq 2$, like how golf is
fong-spivak-seven-sketches_27 F00250	27	1.000	$3 \leq 2$	$3 \leq 2$	$n = m$. Another ordering is the reverse ordering, like $5 \leq 3$ and $3 \leq 2$, like how golf is
fong-spivak-seven-sketches_27 F00251	27	1.000	m	m	m without remainder. In this ordering $2 \leq 4$, for example, but $2 \not\leq 3$, since there is a
fong-spivak-seven-sketches_27 F00252	27	1.000	$2 \leq 4$	$2 \leq 4$	m without remainder. In this ordering $2 \leq 4$, for example, but $2 \not\leq 3$, since there is a
fong-spivak-seven-sketches_27 F00253	27	1.000	$2 \not\leq 3$	$2 \not\leq 3$	m without remainder. In this ordering $2 \leq 4$, for example, but $2 \not\leq 3$, since there is a
fong-spivak-seven-sketches_27 F00254	27	0.999	$1, 2, \ldots, 10$	$1, 2, \ldots, 10$	<i>Exercise 1.46.</i> Write down the numbers  and draw an arrow $a \rightarrow b$ if a
fong-spivak-seven-sketches_27 F00255	27	0.999	$a \rightarrowtail b$	$a \rightarrow b$	<i>Exercise 1.46.</i> Write down the numbers $1, 2, \ldots, 10$ and draw an arrow $a \rightarrow b$ if a

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F00256	28	■ 0.967	$-500 \leq -499 \leq 0 \leq \sqrt{2} \leq 100 / 3$	$-500 \leq -499 \leq 0 \leq \sqrt{2} \leq 100/3$	ordering”, e.g. $-500 \leq -499 \leq 0 \leq \sqrt{2} \leq 100/3$.
fong-spivak-seven-sketches-F00257	28	■ 1.000	$\mathrm{P}(X)$	$P(X)$	set of X ; we denote it $P(X)$. The power set can naturally be given an order by inclusion
fong-spivak-seven-sketches-F00258	28	■ 1.000	$X=\{0,1,2\}$	$X = \{0, 1, 2\}$	For example, taking $X = \{0, 1, 2\}$, we depict $P(X)$ as
fong-spivak-seven-sketches-F00259	28	■ 0.975	$\mathrm{P}\backslash\{1,2, \ldots, n\},\{ \}^{\mathrm{a}}$	$P\{1,2,\dots,n\},^a$	See the cube? The Hasse diagram for the power set of a finite set, say $P\{1,2,\dots,n\}$, ^a
fong-spivak-seven-sketches-F00260	28	■ 1.000	$\mathrm{P} \ X$	PX	^a Note that we omit the parentheses here, writing PX instead of $P(X)$; throughout this book we will
fong-spivak-seven-sketches-F00261	28	■ 0.963	$\mathrm{P}(\varnothing), \mathrm{P}\backslash\{1\}$	$P(\emptyset), P\{1\}$	<i>Exercise 1.51.</i> Draw the Hasse diagrams for $P(\emptyset)$, $P\{1\}$, and $P\{1,2\}$.
fong-spivak-seven-sketches-F00262	28	■ 0.963	$\mathrm{P}\backslash\{1,2\}$	$P\{1,2\}$	<i>Exercise 1.51.</i> Draw the Hasse diagrams for $P(\emptyset)$, $P\{1\}$, and $P\{1,2\}$.
fong-spivak-seven-sketches-F00263	29	■ 1.000	$\operatorname{Prt}(A)$	$\operatorname{Prt}(A)$	think about how we might order the set $\operatorname{Prt}(A)$ of <i>all partitions</i> of A , for some set A . In
fong-spivak-seven-sketches-F00264	29	■ 1.000	$q \in Q$	$q \in Q$	a partition P is <i>finer</i> than a partition Q if, for every part $p \in P$ there is a part $q \in Q$ such
fong-spivak-seven-sketches-F00265	29	■ 1.000	$A_{\{p\}} \subseteq A_{\{q\}}$	$A_p \subseteq A_q$	that $A_p \subseteq A_q$. We could also say that Q is <i>coarser</i> than P .
fong-spivak-seven-sketches-F00266	29	■ 1.000	$g: A \rightarrow Q$	$g: A \rightarrow Q$	tions out of A . Then $f: A \twoheadrightarrow P$ is finer than $g: A \twoheadrightarrow Q$ if there is a function $h: P \rightarrow Q$
fong-spivak-seven-sketches-F00267	29	■ 1.000	$h: P \rightarrow Q$	$h: P \rightarrow Q$	tions out of A . Then $f: A \twoheadrightarrow P$ is finer than $g: A \twoheadrightarrow Q$ if there is a function $h: P \rightarrow Q$
fong-spivak-seven-sketches-F00268	29	■ 1.000	$f ; h=g$	$f;h = g$	such that $f \circ h = g$.
fong-spivak-seven-sketches-F00269	29	■ 0.795	$P, \leq$	$P, \leq$	<i>Example 1.54</i> (Upper sets). Given a preorder $(P, \leq)$, an <i>upper set</i> in P is a subset U of P
fong-spivak-seven-sketches-F00270	29	■ 0.795	U	U	<i>Example 1.54</i> (Upper sets). Given a preorder $(P, \leq)$, an <i>upper set</i> in P is a subset U of P
fong-spivak-seven-sketches-F00271	29	■ 1.000	$p \in U$	$p \in U$	satisfying the condition that if $p \in U$ and $p \leq q$, then $q \in U$. “If p is an element then so
fong-spivak-seven-sketches-F00272	29	■ 1.000	$p \leq q$	$p \leq q$	satisfying the condition that if $p \in U$ and $p \leq q$, then $q \in U$. “If p is an element then so
fong-spivak-seven-sketches-F00273	29	■ 1.000	$q \in U$	$q \in U$	satisfying the condition that if $p \in U$ and $p \leq q$, then $q \in U$. “If p is an element then so
fong-spivak-seven-sketches-F00274	29	■ 1.000	p	p	satisfying the condition that if $p \in U$ and $p \leq q$, then $q \in U$. “If p is an element then so
fong-spivak-seven-sketches-F00275	29	■ 0.912	$\mathrm{U}(P)$	$U(P)$	is anything bigger.” Write $U(P)$ for the set of upper sets in P . We can give the set U an
fong-spivak-seven-sketches-F00276	29	■ 1.000	$U \leq V$	$U \leq V$	order by letting $U \leq V$ if U is contained in V .
fong-spivak-seven-sketches-F00277	29	■ 0.998	$(\mathbb{B}, \leq)$	$(\mathbb{B}, \leq)$	For example, if $(\mathbb{B}, \leq)$ is the booleans (Example 1.34), then its preorder of uppersets
fong-spivak-seven-sketches-F00278	29	■ 0.457	$\mathrm{U}(\mathbb{B})$	$U(\mathbb{B})$	$U(\mathbb{B})$ is
fong-spivak-seven-sketches-F00279	29	■ 0.749	$\subseteq \mathbb{B}$	$\subseteq \mathbb{B}$	The subset $\{\text{false}\} \subseteq \mathbb{B}$ is not an upper set, because $\text{false} \leq \text{true}$ and $\text{true} \notin \{\text{false}\}$.
fong-spivak-seven-sketches-F00280	29	■ 0.749	$\notin \{$	$\notin \{$	The subset $\{\text{false}\} \subseteq \mathbb{B}$ is not an upper set, because $\text{false} \leq \text{true}$ and $\text{true} \notin \{\text{false}\}$.
fong-spivak-seven-sketches-F00281	29	■ 0.998	$(Q, \leq)$	$(Q, \leq)$	<i>Example 1.56</i> (Product preorder). Given preorders $(P, \leq)$ and $(Q, \leq)$, we may define a
fong-spivak-seven-sketches-F00282	29	■ 1.000	$P \times Q$	$P \times Q$	preorder structure on the product set $P \times Q$ by setting $(p, q) \leq (p', q')$ if and only if

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F00283	29	■ 1.000	$(p, q) \leq \left(p^{\{\text{prime}\}}, q^{\{\text{prime}\}}\right)$	$(p, q) \leq (p', q')$	preorder structure on the product set $P \times Q$ by setting $\boxed{p, q} \leq \boxed{p', q'}$ if and only if
fong-spivak-seven-sketches-F00284	29	■ 1.000	$p \leq p^{\{\text{prime}\}}$	$p \leq p'$	$p \leq p'$ and $q \leq q'$. We call this the <i>product preorder</i> . This is a basic example of a more
fong-spivak-seven-sketches-F00285	29	■ 1.000	$q \leq q^{\{\text{prime}\}}$	$q \leq q'$	$p \leq p'$ and $q \leq q'$. We call this the <i>product preorder</i> . This is a basic example of a more
fong-spivak-seven-sketches-F00286	30	■ 0.889	$(P, \leq \mathrm{op})$	$(P, \leq \mathrm{op})$	preorder $(P, \leq^{\mathrm{op}})$ to have the same set of elements, but with $p \leq^{\mathrm{op}} q$ if and only if $q \leq p$.
fong-spivak-seven-sketches-F00287	30	■ 0.889	$p \leq \{ \}^{\{\mathrm{op}\}} q$	$p \leq^{\mathrm{op}} q$	preorder $(P, \leq^{\mathrm{op}})$ to have the same set of elements, but with $p \leq^{\mathrm{op}} q$ if and only if $q \leq p$.
fong-spivak-seven-sketches-F00288	30	■ 0.889	$q \leq p$	$q \leq p$	preorder $(P, \leq^{\mathrm{op}})$ to have the same set of elements, but with $p \leq^{\mathrm{op}} q$ if and only if $q \leq p$.
fong-spivak-seven-sketches-F00289	30	■ 0.988	$A, \leq_A$	$A, \leq_A$	Definition 1.59. A <i>monotone map</i> between preorders $(A, \leq_A)$ and $(B, \leq_B)$ is a function
fong-spivak-seven-sketches-F00290	30	■ 0.988	$B, \leq_B$	$B, \leq_B$	Definition 1.59. A <i>monotone map</i> between preorders $(A, \leq_A)$ and $(B, \leq_B)$ is a function
fong-spivak-seven-sketches-F00291	30	■ 0.872	$x, y \in A$	$x, y \in A$	$f: A \rightarrow B$ such that, for all elements $x, y \in A$, if $x \leq_A y$ then $f(x) \leq_B f(y)$.
fong-spivak-seven-sketches-F00292	30	■ 0.872	$x \leq_A y$	$x \leq_A y$	$f: A \rightarrow B$ such that, for all elements $x, y \in A$, if $x \leq_A y$ then $f(x) \leq_B f(y)$.
fong-spivak-seven-sketches-F00293	30	■ 0.872	$f(x) \leq_B f(y)$	$f(x) \leq_B f(y)$	$f: A \rightarrow B$ such that, for all elements $x, y \in A$, if $x \leq_A y$ then $f(x) \leq_B f(y)$.
fong-spivak-seven-sketches-F00294	30	■ 1.000	$A \rightarrow B$	$A \rightarrow B$	A monotone map $A \rightarrow B$ between two preorders associates to each element of
fong-spivak-seven-sketches-F00295	30	■ 1.000	$x \in A$	$x \in A$	from each element $x \in A$ to its image $f(x) \in B$. Note that the order must be preserved
fong-spivak-seven-sketches-F00296	30	■ 1.000	$f(x) \in B$	$f(x) \in B$	from each element $x \in A$ to its image $f(x) \in B$. Note that the order must be preserved
fong-spivak-seven-sketches-F00297	30	■ 1.000	$f(y)$	$f(y)$	lefthand preorder A , then the image $f(x)$ will be above the image $f(y)$ in the righthand
fong-spivak-seven-sketches-F00298	31	■ 1.000	$\mathbb{B} \rightarrow \mathbb{N}$	$\mathbb{B} \rightarrow \mathbb{N}$	preorder of natural numbers from Example 1.45. The map $\mathbb{B} \rightarrow \mathbb{N}$ sending <code>false</code> to 17
fong-spivak-seven-sketches-F00299	31	■ 1.000	$x \rightarrow y$	$x \rightarrow y$	because it preserves order: whenever there is a path $x \rightarrow y$ upstairs, there is a path
fong-spivak-seven-sketches-F00300	31	■ 1.000	$F(x) \rightarrow F(y)$	$F(x) \rightarrow F(y)$	$\boxed{F(x) \rightarrow F(y)}$ downstairs.
fong-spivak-seven-sketches-F00301	31	■ 0.988	$ \cdot : \mathrm{P}(X) \rightarrow \mathbb{N}$	$ \cdot : \mathrm{P}(X) \rightarrow \mathbb{N}$	from Example 1.50. The map $ \cdot : \mathrm{P}(X) \rightarrow \mathbb{N}$ sending each subset S to its number of
fong-spivak-seven-sketches-F00302	31	■ 1.000	$ S $	$ S $	elements $ S $, also called its <i>cardinality</i> , is a monotone map.
fong-spivak-seven-sketches-F00303	31	■ 1.000	$0 \leq 1 \leq 2 \leq 3$	$0 \leq 1 \leq 2 \leq 3$	2. Draw the Hasse diagram for the preorder $0 \leq 1 \leq 2 \leq 3$.
fong-spivak-seven-sketches-F00304	32	■ 0.997	$ \cdot $	$ \cdot $	3. Draw the cardinality map $ \cdot $ from Example 1.62 as dashed lines between them.
fong-spivak-seven-sketches-F00305	32	■ 0.995	$\mathrm{U}(P) \rightarrow \mathrm{P}(P)$	$\mathrm{U}(P) \rightarrow \mathrm{P}(P)$	$(P, \leq)$, the map $\mathrm{U}(P) \rightarrow \mathrm{P}(P)$ sending each upper set of $(P, \leq)$ to itself—considered as
fong-spivak-seven-sketches-F00306	32	■ 0.939	$\mathrm{P}(\mathbb{B})$	$\mathrm{P}(\mathbb{B})$	Example 1.54, and you drew the Hasse diagram for $\mathrm{P}(\mathbb{B})$ in Exercise 1.51. Now draw
fong-spivak-seven-sketches-F00307	32	■ 1.000	$\uparrow p := \{p' \in P \mid p \leq p'\}$	$\uparrow p := \{p' \in P \mid p \leq p'\}$	1. Show that the set $\boxed{\uparrow p := \{p' \in P \mid p \leq p'\}}$ is an upper set, for any $p \in P$.
fong-spivak-seven-sketches-F00308	32	■ 0.671	$\uparrow: P^{\mathrm{op}} \rightarrow \mathrm{U}(P)$	$\uparrow: P^{\mathrm{op}} \rightarrow \mathrm{U}(P)$	2. Show that this construction defines a monotone map $\uparrow: P^{\mathrm{op}} \rightarrow \mathrm{U}(P)$.
fong-spivak-seven-sketches-F00309	32	■ 1.000	$\uparrow(p') \subseteq \uparrow(p)$	$\uparrow(p') \subseteq \uparrow(p)$	3. Show that if $p \leq p'$ in P if and only if $\uparrow(p') \subseteq \uparrow(p)$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F00310	32	■ 1.000	$b \geq a \leq c$	$b \geq a \leq c$	4. Draw a picture of the map $\uparrow$ in the case where P is the preorder ($b \geq a \leq c$) from
fong-spivak-seven-sketches F00311	32	■ 1.000	$f: \left(P, \leq_P\right) \rightarrow \left(Q, \leq_Q\right)$	$f: (P, \leq_P) \rightarrow (Q, \leq_Q)$	Exercise 1.67. As you yourself well know, a monotone map $f: (P, \leq_P) \rightarrow (Q, \leq_Q)$
fong-spivak-seven-sketches F00312	32	■ 1.000	$f: P \rightarrow Q$	$f: P \rightarrow Q$	consists of a function $f: P \rightarrow Q$ that satisfies a “monotonicity” property. Show that
fong-spivak-seven-sketches F00313	32	■ 1.000	$\left(P, \leq_P\right)$	$(P, \leq_P)$	when $(P, \leq_P)$ is a discrete preorder, then <i>every</i> function $P \rightarrow Q$ satisfies the monotonicity
fong-spivak-seven-sketches F00314	32	■ 1.000	$P \rightarrow Q$	$P \rightarrow Q$	when $(P, \leq_P)$ is a discrete preorder, then <i>every</i> function $P \rightarrow Q$ satisfies the monotonicity
fong-spivak-seven-sketches F00315	32	■ 1.000	$\leq_Q$	$\leq_Q$	property, regardless of the order $\leq_Q$.
fong-spivak-seven-sketches F00316	32	■ 1.000	$\operatorname{Prt}(X)$	$\operatorname{Prt}(X)$	Example 1.68. Recall from Example 1.52 that given a set X we define $\operatorname{Prt}(X)$ to be the
fong-spivak-seven-sketches F00317	32	■ 1.000	$s: X \rightarrow P$	$s: X \rightarrow P$	$s: X \twoheadrightarrow P$ for some set P .
fong-spivak-seven-sketches F00318	32	■ 1.000	$f^*: \operatorname{Prt}(Y) \rightarrow \operatorname{Prt}(X)$	$f^*: \operatorname{Prt}(Y) \rightarrow \operatorname{Prt}(X)$	Any surjective function $f: X \twoheadrightarrow Y$ induces a monotone map $f^*: \operatorname{Prt}(Y) \rightarrow \operatorname{Prt}(X)$,
fong-spivak-seven-sketches F00319	32	■ 0.994	$s: Y \rightarrow P$	$s: Y \rightarrow P$	going “backwards.” It is defined by sending a partition $s: Y \twoheadrightarrow P$ to the composite
fong-spivak-seven-sketches F00320	32	■ 0.905	s	s	$f \circ s: X \twoheadrightarrow P$.
fong-spivak-seven-sketches F00321	32	■ 0.905	$X \rightarrow P$	$X \rightarrow P$	$f \circ s: X \twoheadrightarrow P$
fong-spivak-seven-sketches F00322	32	■ 1.000	$f^*(P)$	$f^*(P)$	partitions P and Q of Y , and then find $f^*(P)$ and $f^*(Q)$.
fong-spivak-seven-sketches F00323	32	■ 1.000	$f^*(Q)$	$f^*(Q)$	partitions P and Q of Y , and then find $f^*(P)$ and $f^*(Q)$.
fong-spivak-seven-sketches F00324	33	■ 0.979	$\left(Q, \leq_Q\right)$	$(Q, \leq_Q)$	If $(Q, \leq_Q)$ and $(R, \leq_R)$ are preorders and $f: P \rightarrow Q$ and $g: Q \rightarrow R$ are monotone,
fong-spivak-seven-sketches F00325	33	■ 0.979	$\left(R, \leq_R\right)$	$(R, \leq_R)$	If $(Q, \leq_Q)$ and $(R, \leq_R)$ are preorders and $f: P \rightarrow Q$ and $g: Q \rightarrow R$ are monotone,
fong-spivak-seven-sketches F00326	33	■ 0.979	$g: Q \rightarrow R$	$g: Q \rightarrow R$	If $(Q, \leq_Q)$ and $(R, \leq_R)$ are preorders and $f: P \rightarrow Q$ and $g: Q \rightarrow R$ are monotone,
fong-spivak-seven-sketches F00327	33	■ 0.960	$(f; g): P \rightarrow R$	$(f; g): P \rightarrow R$	then $(f \circ g): P \rightarrow R$ is also monotone.
fong-spivak-seven-sketches F00328	33	■ 0.629	$\operatorname{id}_P: P \rightarrow P$	$\operatorname{id}_P: P \rightarrow P$	identity function $\operatorname{id}_P: P \rightarrow P$ is a monotone map $(P, \leq) \rightarrow (P, \leq^{\operatorname{op}})$ if and only if for
fong-spivak-seven-sketches F00329	33	■ 0.629	$(P, \leq) \rightarrow (P, \leq^{\operatorname{op}})$	$(P, \leq) \rightarrow (P, \leq^{\operatorname{op}})$	identity function $\operatorname{id}_P: P \rightarrow P$ is a monotone map $(P, \leq) \rightarrow (P, \leq^{\operatorname{op}})$ if and only if for
fong-spivak-seven-sketches F00330	33	■ 1.000	$p, q \in P$	$p, q \in P$	all $p, q \in P$ we have $q \leq p$ whenever $p \leq q$. For historical reasons connected to linear
fong-spivak-seven-sketches F00331	33	■ 0.997	$p_1 = p_2$	$p_1 = p_2$	made into a discrete preorder ($p_1 \leq p_2$ iff $p_1 = p_2$), and the round-trips return us where
fong-spivak-seven-sketches F00332	33	■ 1.000	$g: Q \rightarrow P$	$g: Q \rightarrow P$	is called an <i>isomorphism</i> if there exists a monotone function $g: Q \rightarrow P$ such that
fong-spivak-seven-sketches F00333	33	■ 0.555	$f; g = \operatorname{id}_P$	$f; g = \operatorname{id}_P$	$f \circ g = \operatorname{id}_P$ and $g \circ f = \operatorname{id}_Q$. This means that for any $p \in P$ and $q \in Q$, we have
fong-spivak-seven-sketches F00334	33	■ 0.555	$g \circ f = \operatorname{id}_Q$	$g \circ f = \operatorname{id}_Q$	$f \circ g = \operatorname{id}_P$ and $g \circ f = \operatorname{id}_Q$. This means that for any $p \in P$ and $q \in Q$, we have
fong-spivak-seven-sketches F00335	33	■ 1.000	g	g	We refer to g as the <i>inverse</i> of f , and vice versa: f is the inverse of g .
fong-spivak-seven-sketches F00336	34	■ 1.000	P, Q	P, Q	Example 1.76. Here are the Hasse diagrams for three preorders P, Q , and R , all of which

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-34 F00337	34	1.000	R	R	<i>Example 1.76.</i> Here are the Hasse diagrams for three preorders P , Q , and R , all of which
fong-spivak-seven-sketches-34 F00338	34	1.000	$f(a)=v, f(b)=w, f(c)=x, f(d)=y$	$f(a) = v, f(b) = w, f(c) = x, f(d) = y$	The map $f: P \rightarrow Q$ given by $f(a) = v, f(b) = w, f(c) = x, f(d) = y$, and $f(e) = z$ has
fong-spivak-seven-sketches-34 F00339	34	1.000	$f(e)=z$	$f(e) = z$	The map $f: P \rightarrow Q$ given by $f(a) = v, f(b) = w, f(c) = x, f(d) = y$, and $f(e) = z$ has
fong-spivak-seven-sketches-34 F00340	34	1.000	$x \rightarrowtail z$	$x \rightarrow z$	is an arrow $x \rightarrow z$ in the Hasse diagram for R and not one in Q , but in fact this arrow
fong-spivak-seven-sketches-34 F00341	34	1.000	$v \rightarrowtail y$	$v \rightarrow y$	drawn an arrow $v \rightarrow y$ in either Q or R and it would not have changed the preorder.
fong-spivak-seven-sketches-34 F00342	34	0.546	$\operatorname{Prt}(\{*, \bullet, \circ\}) \rightarrowtail \mathbb{B}$	$\operatorname{Prt}(\{*, \bullet, \circ\}) \rightarrow \mathbb{B}$	‘Is $\bullet$ connected to $*$?’ is a monotone map $\operatorname{Prt}(\{*, \bullet, \circ\}) \rightarrow \mathbb{B}$; see also Eq. (1.5).
fong-spivak-seven-sketches-34 F00343	34	1.000	$P \rightarrowtail \mathbb{B}$	$P \rightarrow \mathbb{B}$	Proposition 1.78. Let P be a preorder. Monotone maps $P \rightarrow \mathbb{B}$ are in one-to-one
fong-spivak-seven-sketches-34 F00344	34	0.656	$f: P \rightarrowtail \mathbb{B}$	$f: P \rightarrow \mathbb{B}$	<i>Proof.</i> Let $f: P \rightarrow \mathbb{B}$ be a monotone map. We will show that the subset $f^{-1}(\mathbf{true}) \subseteq P$
fong-spivak-seven-sketches-34 F00345	34	0.656	$f^{-1}(\mathbf{true})$	$f^{-1}(\mathbf{true})$	<i>Proof.</i> Let $f: P \rightarrow \mathbb{B}$ be a monotone map. We will show that the subset $f^{-1}(\mathbf{true}) \subseteq P$
fong-spivak-seven-sketches-34 F00346	34	0.656	$f^{-1}(\mathbf{true}) \subseteq P$	$f^{-1}(\mathbf{true}) \subseteq P$	<i>Proof.</i> Let $f: P \rightarrow \mathbb{B}$ be a monotone map. We will show that the subset $f^{-1}(\mathbf{true}) \subseteq P$
fong-spivak-seven-sketches-34 F00347	34	1.000	$p \in f^{-1}(\mathbf{true})$	$p \in f^{-1}(\mathbf{true})$	is an upper set. Suppose $p \in f^{-1}(\mathbf{true})$ and $p \leq q$; then $\mathbf{true} = f(p) \leq f(q)$. But in
fong-spivak-seven-sketches-34 F00348	34	1.000	$f(p) \leq f(q)$	$f(p) \leq f(q)$	is an upper set. Suppose $p \in f^{-1}(\mathbf{true})$ and $p \leq q$; then $\mathbf{true} = f(p) \leq f(q)$. But in
fong-spivak-seven-sketches-34 F00349	34	0.842	$f(p) \leq f(q)$	$f(p) \leq f(q)$	$\mathbb{B}$, if $\mathbf{true} \leq f(q)$ then $\mathbf{true} = f(q)$. This implies $q \in f^{-1}(\mathbf{true})$ and thus shows that
fong-spivak-seven-sketches-34 F00350	34	0.842	$f(p) \leq f(q)$	$f(p) \leq f(q)$	$\mathbb{B}$, if $\mathbf{true} \leq f(q)$ then $\mathbf{true} = f(q)$. This implies $q \in f^{-1}(\mathbf{true})$ and thus shows that
fong-spivak-seven-sketches-34 F00351	34	0.842	$q \in f^{-1}(\mathbf{true})$	$q \in f^{-1}(\mathbf{true})$	$\mathbb{B}$, if $\mathbf{true} \leq f(q)$ then $\mathbf{true} = f(q)$. This implies $q \in f^{-1}(\mathbf{true})$ and thus shows that
fong-spivak-seven-sketches-34 F00352	34	1.000	$f^{-1}(\mathbf{true})$	$f^{-1}(\mathbf{true})$	$f^{-1}(\mathbf{true})$ is an upper set.
fong-spivak-seven-sketches-34 F00353	34	1.000	$f_U: P \rightarrowtail \mathbb{B}$	$f_U: P \rightarrow \mathbb{B}$	Conversely, if U is an upper set in P , define $f_U: P \rightarrow \mathbb{B}$ such that $f_U(p) = \mathbf{true}$ when
fong-spivak-seven-sketches-34 F00354	34	1.000	$f_U(p) =$	$f_U(p) =$	Conversely, if U is an upper set in P , define $f_U: P \rightarrow \mathbb{B}$ such that $f_U(p) = \mathbf{true}$ when
fong-spivak-seven-sketches-34 F00355	34	1.000	$p \notin U$	$p \notin U$	$p \in U$, and $f_U(p) = \mathbf{false}$ when $p \notin U$. This is a monotone map, because if $p \leq q$, then
fong-spivak-seven-sketches-34 F00356	34	1.000	$f(p) =$	$f(p) =$	either $p \in U$, so $q \in U$ and $f(p) = \mathbf{true} = f(q)$, or $p \notin U$, so $f(p) = \mathbf{false} \leq f(q)$.
fong-spivak-seven-sketches-34 F00357	34	0.996	$f^*: \mathcal{U}(Q) \rightarrowtail \mathcal{U}(P)$	$f^*: \mathcal{U}(Q) \rightarrow \mathcal{U}(P)$	map. Then we can define a monotone map $f^*: \mathcal{U}(Q) \rightarrow \mathcal{U}(P)$ sending an upper set
fong-spivak-seven-sketches-34 F00358	34	1.000	$U \subseteq Q$	$U \subseteq Q$	$U \subseteq Q$ to the upper set $f^{-1}(U) \subseteq P$. We call this the <i>pullback along f</i> .
fong-spivak-seven-sketches-34 F00359	34	1.000	$f^{-1}(U) \subseteq P$	$f^{-1}(U) \subseteq P$	$U \subseteq Q$ to the upper set $f^{-1}(U) \subseteq P$. We call this the <i>pullback along f</i> .
fong-spivak-seven-sketches-34 F00360	34	1.000	f^*	f^*	can be understood in terms of composition. Indeed, show that the f^* is defined by
fong-spivak-seven-sketches-34 F00361	34	1.000	$u: Q \rightarrowtail \mathbb{B}$	$u: Q \rightarrow \mathbb{B}$	taking $u: Q \rightarrow \mathbb{B}$ to $(f \circ u): P \rightarrow \mathbb{B}$.
fong-spivak-seven-sketches-34 F00362	34	1.000	$(f \circ u): P \rightarrowtail \mathbb{B}$	$(f \circ u): P \rightarrow \mathbb{B}$	taking $u: Q \rightarrow \mathbb{B}$ to $(f \circ u): P \rightarrow \mathbb{B}$.
fong-spivak-seven-sketches-35 F00363	35	0.637	$(\mathbb{R}, \leq)$	$(\mathbb{R}, \leq)$	Consider the preorder $(\mathbb{R}, \leq)$ of real numbers ordered in the usual way. The subset

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F00364	35	■ 1.000	$\mathbb{N} \subseteq \mathbb{R}$	$\mathbb{N} \subseteq \mathbb{R}$	$\mathbb{N} \subseteq \mathbb{R}$ has many lower bounds, namely -1.5 is a lower bound: every element of $\mathbb{N}$ is
fong-spivak-seven-sketches F00365	35	■ 1.000	$\left\{ \left\{ \frac{1}{n+1} \right\} \mid n \in \mathbb{N} \right\} \subseteq \mathbb{R}$	$\left\{ \frac{1}{n+1} \mid n \in \mathbb{N} \right\} \subseteq \mathbb{R}$	1. Why is 0 a lower bound for $\left\{ \frac{1}{n+1} \mid n \in \mathbb{N} \right\} \subseteq \mathbb{R}$?
fong-spivak-seven-sketches F00366	35	■ 0.719	$A \subseteq P$	$A \subseteq P$	Definition 1.81. Let $(P, \leq)$ be a preorder, and let $A \subseteq P$ be a subset. We say that an
fong-spivak-seven-sketches F00367	35	■ 1.000	$p \leq a$	$p \leq a$	(a) for all $a \in A$, we have $p \leq a$, and
fong-spivak-seven-sketches F00368	35	■ 1.000	q	q	(b) for all q such that $q \leq a$ for all $a \in A$, we have that $q \leq p$.
fong-spivak-seven-sketches F00369	35	■ 1.000	$q \leq a$	$q \leq a$	(b) for all q such that $q \leq a$ for all $a \in A$, we have that $q \leq p$.
fong-spivak-seven-sketches F00370	35	■ 1.000	$p = \bigwedge A, p = \bigwedge_{a \in A} a$	$p = \bigwedge A, p = \bigwedge_{a \in A} a$	We write $p = \bigwedge A, p = \bigwedge_{a \in A} a$, or, if the dummy variable a is clear from context, just
fong-spivak-seven-sketches F00371	35	■ 0.999	$p = \bigwedge_{A} a$	$p = \bigwedge_A a$	$p = \bigwedge_A a$. If A just consists of two elements, say $A = \{a, b\}$, we can denote $\bigwedge A$ simply
fong-spivak-seven-sketches F00372	35	■ 0.999	$A = \{a, b\}$	$A = \{a, b\}$	$p = \bigwedge_A a$. If A just consists of two elements, say $A = \{a, b\}$, we can denote $\bigwedge A$ simply
fong-spivak-seven-sketches F00373	35	■ 0.999	$\bigwedge A$	$\bigwedge A$	$p = \bigwedge_A a$. If A just consists of two elements, say $A = \{a, b\}$, we can denote $\bigwedge A$ simply
fong-spivak-seven-sketches F00374	35	■ 1.000	$a \wedge b$	$a \wedge b$	by $a \wedge b$.
fong-spivak-seven-sketches F00375	35	■ 1.000	$a \leq p$	$a \leq p$	(a) for all $a \in A$ we have $a \leq p$, and
fong-spivak-seven-sketches F00376	35	■ 1.000	$a \leq q$	$a \leq q$	(b) for all q such that $a \leq q$ for all $a \in A$, we have that $p \leq q$.
fong-spivak-seven-sketches F00377	35	■ 0.990	$p = \bigvee A$	$p = \bigvee A$	We write $p = \bigvee A$ or $p = \bigvee_{a \in A} a$, or when $A = \{a, b\}$ we may simply write $p = a \vee b$.
fong-spivak-seven-sketches F00378	35	■ 0.990	$p = \bigvee_{a \in A} a$	$p = \bigvee_{a \in A} a$	We write $p = \bigvee A$ or $p = \bigvee_{a \in A} a$, or when $A = \{a, b\}$ we may simply write $p = a \vee b$.
fong-spivak-seven-sketches F00379	35	■ 0.990	$p = a \vee b$	$p = a \vee b$	We write $p = \bigvee A$ or $p = \bigvee_{a \in A} a$, or when $A = \{a, b\}$ we may simply write $p = a \vee b$.
fong-spivak-seven-sketches F00380	36	■ 0.980	$p = \bigwedge A$	$p = \bigwedge A$	$p = \bigwedge A$ and $q = \bigwedge A$ with $p \neq q$, which does not make sense if $p \neq q$!
fong-spivak-seven-sketches F00381	36	■ 0.980	$q = \bigwedge A$	$q = \bigwedge A$	$p = \bigwedge A$ and $q = \bigwedge A$ with $p \neq q$, which does not make sense if $p \neq q$!
fong-spivak-seven-sketches F00382	36	■ 0.980	$p \neq q$	$p \neq q$	$p = \bigwedge A$ and $q = \bigwedge A$ with $p \neq q$, which does not make sense if $p \neq q$!
fong-spivak-seven-sketches F00383	36	■ 1.000	p, q	p, q	meets p, q are automatically isomorphic: the very definition of meet forces both $p \leq q$
fong-spivak-seven-sketches F00384	36	■ 1.000	$p \cong q$	$p \cong q$	and $q \leq p$, and thus we have $p \cong q$. So for any $x \in P$, we have $p \leq x$ iff $q \leq x$ and
fong-spivak-seven-sketches F00385	36	■ 1.000	$x \in P$	$x \in P$	and $q \leq p$, and thus we have $p \cong q$. So for any $x \in P$, we have $p \leq x$ iff $q \leq x$ and
fong-spivak-seven-sketches F00386	36	■ 1.000	$p \leq x$	$p \leq x$	and $q \leq p$, and thus we have $p \cong q$. So for any $x \in P$, we have $p \leq x$ iff $q \leq x$ and
fong-spivak-seven-sketches F00387	36	■ 1.000	$q \leq x$	$q \leq x$	and $q \leq p$, and thus we have $p \cong q$. So for any $x \in P$, we have $p \leq x$ iff $q \leq x$ and
fong-spivak-seven-sketches F00388	36	■ 1.000	$x \leq p$	$x \leq p$	$x \leq p$ iff $x \leq q$. Thus as long as we are only interested in elements of P based on their
fong-spivak-seven-sketches F00389	36	■ 1.000	$x \leq q$	$x \leq q$	$x \leq p$ iff $x \leq q$. Thus as long as we are only interested in elements of P based on their

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F00390	36	■ 1.000	$P=\{p, q, r\}$	$P = \{p, q, r\}$	a subset A need not have a meet or a join. Consider the three element set $P = \{p, q, r\}$
fong-spivak-seven-sketches F00391	36	■ 1.000	$A=\{p, q\}$	$A = \{p, q\}$	with the discrete ordering. The set $A = \{p, q\}$ does not have a join in P because if x
fong-spivak-seven-sketches F00392	36	■ 0.965	$\{a, b\}$	$\{a, b\}$	Let A be the subset $\{a, b\}$ in the preorder specified by this Hasse diagram. Then both
fong-spivak-seven-sketches F00393	36	■ 0.998	$c \leq d$	$c \leq d$	less than d . Note that, as in Remark 1.82, $c \leq d$ and $d \leq c$, so $c \cong d$. Such will always
fong-spivak-seven-sketches F00394	36	■ 0.998	$d \leq c$	$d \leq c$	less than d . Note that, as in Remark 1.82, $c \leq d$ and $d \leq c$, so $c \cong d$. Such will always
fong-spivak-seven-sketches F00395	36	■ 0.998	$c \cong d$	$c \cong d$	less than d . Note that, as in Remark 1.82, $c \leq d$ and $d \leq c$, so $c \cong d$. Such will always
fong-spivak-seven-sketches F00396	36	■ 0.519	$A=\{p\}$	$A = \{p\}$	<i>Exercise 1.85.</i> Let $(P, \leq)$ be a preorder and $p \in P$ an element. Consider the set $A = \{p\}$
fong-spivak-seven-sketches F00397	36	■ 1.000	$\wedge A \cong p$	$\wedge A \cong p$	1. Show that $\wedge A \cong p$.
fong-spivak-seven-sketches F00398	36	■ 1.000	$\wedge A = p$	$\wedge A = p$	2. Show that if P is in fact a partial order, then $\wedge A = p$.
fong-spivak-seven-sketches F00399	36	■ 0.999	$\wedge$	$\wedge$	3. Are the analogous facts true when $\wedge$ is replaced by $\vee$?
fong-spivak-seven-sketches F00400	37	■ 1.000	$p \vee p = p \wedge p = p$	$p \vee p = p \wedge p = p$	<i>Example 1.86.</i> In any partial order P , we have $p \vee p = p \wedge p = p$. The reason is that our
fong-spivak-seven-sketches F00401	37	■ 0.796	$p \vee p$	$p \vee p$	notation says $p \vee p$ means $\vee\{p, p\}$. But $\{p, p\} = \{p\}$ (see Section 1.2.1), so $p \vee p = p$ by
fong-spivak-seven-sketches F00402	37	■ 0.796	$\vee\{p, p\}$	$\vee\{p, p\}$	notation says $p \vee p$ means $\vee\{p, p\}$. But $\{p, p\} = \{p\}$ (see Section 1.2.1), so $p \vee p = p$ by
fong-spivak-seven-sketches F00403	37	■ 0.796	$\{p, p\} = \{p\}$	$\{p, p\} = \{p\}$	notation says $p \vee p$ means $\vee\{p, p\}$. But $\{p, p\} = \{p\}$ (see Section 1.2.1), so $p \vee p = p$ by
fong-spivak-seven-sketches F00404	37	■ 0.796	$p \vee p = p$	$p \vee p = p$	notation says $p \vee p$ means $\vee\{p, p\}$. But $\{p, p\} = \{p\}$ (see Section 1.2.1), so $p \vee p = p$ by
fong-spivak-seven-sketches F00405	37	■ 0.999	$A, B \subseteq X$	$A, B \subseteq X$	<i>Example 1.87.</i> In a power set $P(X)$, the meet of a collection of subsets, say $A, B \subseteq X$ is
fong-spivak-seven-sketches F00406	37	■ 1.000	$A \wedge B = A \cap B$	$A \wedge B = A \cap B$	their intersection $A \wedge B = A \cap B$, while the join is their union, $A \vee B = A \cup B$.
fong-spivak-seven-sketches F00407	37	■ 1.000	$A \vee B = A \cup B$	$A \vee B = A \cup B$	their intersection $A \wedge B = A \cap B$, while the join is their union, $A \vee B = A \cup B$.
fong-spivak-seven-sketches F00408	37	■ 1.000	$n \mid m$	$n \mid m$	<i>Exercise 1.90.</i> Recall the division ordering on $\mathbb{N}$ from Example 1.45: we write $n \mid m$
fong-spivak-seven-sketches F00409	37	■ 0.706	$A \subseteq B \subseteq P$	$A \subseteq B \subseteq P$	Proposition 1.91. Suppose $(P, \leq)$ is a preorder and $A \subseteq B \subseteq P$ are subsets that have
fong-spivak-seven-sketches F00410	37	■ 1.000	$\wedge B \leq \wedge A$	$\wedge B \leq \wedge A$	meets. Then $\wedge B \leq \wedge A$.
fong-spivak-seven-sketches F00411	37	■ 0.998	$\bigvee A \leq \bigvee B$	$\bigvee A \leq \bigvee B$	Similarly, if A and B have joins, then $\bigvee A \leq \bigvee B$.
fong-spivak-seven-sketches F00412	37	■ 1.000	$m = \wedge A$	$m = \wedge A$	<i>Proof.</i> Let $m = \wedge A$ and $n = \wedge B$. Then for any $a \in A$ we also have $a \in B$, so $n \leq a$
fong-spivak-seven-sketches F00413	37	■ 1.000	$n = \wedge B$	$n = \wedge B$	<i>Proof.</i> Let $m = \wedge A$ and $n = \wedge B$. Then for any $a \in A$ we also have $a \in B$, so $n \leq a$
fong-spivak-seven-sketches F00414	37	■ 1.000	$a \in B$	$a \in B$	<i>Proof.</i> Let $m = \wedge A$ and $n = \wedge B$. Then for any $a \in A$ we also have $a \in B$, so $n \leq a$
fong-spivak-seven-sketches F00415	37	■ 1.000	$n \leq a$	$n \leq a$	<i>Proof.</i> Let $m = \wedge A$ and $n = \wedge B$. Then for any $a \in A$ we also have $a \in B$, so $n \leq a$
fong-spivak-seven-sketches F00416	37	■ 0.843	$A^{\{\prime\}}$	A'	because m is A 's greatest lower bound. The second claim is proved similarly.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_38_F00417	38	1.000	$\Phi: P \rightarrow Q$	$\Phi: P \rightarrow Q$	map $\Phi: P \rightarrow Q$ is a phenomenon (we might say “feature”) of P as observed by Q . He
fong-spivak-seven-sketches_38_F00418	38	1.000	$f(a \wedge b) \cong$	$f(a \wedge b) \cong$	Definition 1.92. We say that a monotone map $f: P \rightarrow Q$ <i>preserves meets</i> if $f(a \wedge b) \cong$
fong-spivak-seven-sketches_38_F00419	38	0.997	$f(a) \wedge f(b)$	$f(a) \wedge f(b)$	$f(a) \wedge f(b)$ for all $a, b \in P$. We similarly say f <i>preserves joins</i> if $f(a \vee b) \cong f(a) \vee f(b)$
fong-spivak-seven-sketches_38_F00420	38	0.997	$a, b \in P$	$a, b \in P$	$f(a) \wedge f(b)$ for all $a, b \in P$. We similarly say f <i>preserves joins</i> if $f(a \vee b) \cong f(a) \vee f(b)$
fong-spivak-seven-sketches_38_F00421	38	0.997	$f(a \vee b) \cong f(a) \vee f(b)$	$f(a \vee b) \cong f(a) \vee f(b)$	$f(a) \wedge f(b)$ for all $a, b \in P$. We similarly say f <i>preserves joins</i> if $f(a \vee b) \cong f(a) \vee f(b)$
fong-spivak-seven-sketches_38_F00422	38	1.000	$f(a) \vee f(b)$	$f(a) \vee f(b)$	a and b , then the left hand side $f(a) \vee f(b)$ may be interpreted as the combination of
fong-spivak-seven-sketches_38_F00423	38	1.000	$f(a \vee b)$	$f(a \vee b)$	$f(a \vee b)$ is the observation of the combined system $a \vee b$. The inequality implies that
fong-spivak-seven-sketches_38_F00424	38	1.000	$a \vee b$	$a \vee b$	$f(a \vee b)$ is the observation of the combined system $a \vee b$. The inequality implies that
fong-spivak-seven-sketches_38_F00425	38	1.000	$f(a \vee b) \neq f(a) \vee f(b)$	$f(a \vee b) \neq f(a) \vee f(b)$	$f(a \vee b) \neq f(a) \vee f(b)$, but in the subsequent text we seemed to imply there would be
fong-spivak-seven-sketches_38_F00426	38	1.000	$f(a), f(b) \in Q$	$f(a), f(b) \in Q$	Prove that for any monotone map $f: P \rightarrow Q$, if $a, b \in P$ have a join and $f(a), f(b) \in Q$
fong-spivak-seven-sketches_38_F00427	38	1.000	$f(a) \vee f(b) \leq f(a \vee b)$	$f(a) \vee f(b) \leq f(a \vee b)$	have a join, then indeed $f(a) \vee f(b) \leq f(a \vee b)$.
fong-spivak-seven-sketches_39_F00428	39	0.889	$(3 \times -): \mathbb{Z} \rightarrow \mathbb{R}$	$(3 \times -): \mathbb{Z} \rightarrow \mathbb{R}$	<i>Example 1.97.</i> Consider the map $(3 \times -): \mathbb{Z} \rightarrow \mathbb{R}$ which sends $x \in \mathbb{Z}$ to $3x$, which we can
fong-spivak-seven-sketches_39_F00429	39	0.889	$x \in \mathbb{Z}$	$x \in \mathbb{Z}$	<i>Example 1.97.</i> Consider the map $(3 \times -): \mathbb{Z} \rightarrow \mathbb{R}$ which sends $x \in \mathbb{Z}$ to $3x$, which we can
fong-spivak-seven-sketches_39_F00430	39	0.889	$3x$	$3x$	<i>Example 1.97.</i> Consider the map $(3 \times -): \mathbb{Z} \rightarrow \mathbb{R}$ which sends $x \in \mathbb{Z}$ to $3x$, which we can
fong-spivak-seven-sketches_39_F00431	39	0.711	$3x \in \mathbb{Z} \subseteq \mathbb{R}$	$3x \in \mathbb{Z} \subseteq \mathbb{R}$	consider as a real number $3x \in \mathbb{Z} \subseteq \mathbb{R}$. Let’s find a left adjoint for the map $(3 \times -)$.
fong-spivak-seven-sketches_39_F00432	39	0.711	$3 \times -$	$3 \times -$	consider as a real number $3x \in \mathbb{Z} \subseteq \mathbb{R}$. Let’s find a left adjoint for the map $(3 \times -)$.
fong-spivak-seven-sketches_39_F00433	39	1.000	$\lceil z \rceil$	$\lceil z \rceil$	Write $\lceil z \rceil$ for the smallest natural number above $z \in \mathbb{R}$, and write $\lfloor z \rfloor$ for the largest
fong-spivak-seven-sketches_39_F00434	39	1.000	$z \in \mathbb{R}$	$z \in \mathbb{R}$	Write $\lceil z \rceil$ for the smallest natural number above $z \in \mathbb{R}$, and write $\lfloor z \rfloor$ for the largest
fong-spivak-seven-sketches_39_F00435	39	1.000	$\lfloor z \rfloor$	$\lfloor z \rfloor$	Write $\lceil z \rceil$ for the smallest natural number above $z \in \mathbb{R}$, and write $\lfloor z \rfloor$ for the largest
fong-spivak-seven-sketches_39_F00436	39	0.582	$\lceil 3.14 \rceil = 4$	$\lceil 3.14 \rceil = 4$	integer below $z \in \mathbb{R}$, e.g. $\lceil 3.14 \rceil = 4$ and $\lfloor 3.14 \rfloor = 3$. ^a As the left adjoint $\mathbb{R} \rightarrow \mathbb{Z}$, let’s see
fong-spivak-seven-sketches_39_F00437	39	0.582	$\lfloor 3.14 \rfloor = 3$	$\lfloor 3.14 \rfloor = 3$. ^a	integer below $z \in \mathbb{R}$, e.g. $\lceil 3.14 \rceil = 4$ and $\lfloor 3.14 \rfloor = 3$. ^a As the left adjoint $\mathbb{R} \rightarrow \mathbb{Z}$, let’s see
fong-spivak-seven-sketches_39_F00438	39	0.582	$\mathbb{R} \rightarrow \mathbb{Z}$	$\mathbb{R} \rightarrow \mathbb{Z}$	integer below $z \in \mathbb{R}$, e.g. $\lceil 3.14 \rceil = 4$ and $\lfloor 3.14 \rfloor = 3$. ^a As the left adjoint $\mathbb{R} \rightarrow \mathbb{Z}$, let’s see
fong-spivak-seven-sketches_39_F00439	39	1.000	$\lceil -/3 \rceil$	$\lceil -/3 \rceil$	if $\lceil -/3 \rceil$ works.
fong-spivak-seven-sketches_39_F00440	39	0.788	$(3 \times -)$	$(3 \times -)$	Success! Thus we have a Galois connection between $\lceil -/3 \rceil$ and $(3 \times -)$.
fong-spivak-seven-sketches_39_F00441	39	1.000	$\lfloor 3 \rfloor = 3 = \lceil 3 \rceil$	$\lfloor 3 \rfloor = 3 = \lceil 3 \rceil$	monthful. Anyway, $\lfloor 3 \rfloor = 3 = \lceil 3 \rceil$.
fong-spivak-seven-sketches_39_F00442	39	0.989	$3 \times$	$3 \times$	<i>Exercise 1.98.</i> In Example 1.97 we found a left adjoint for the monotone map $(3 \times$
fong-spivak-seven-sketches_39_F00443	39	0.945	$-): \mathbb{Z} \rightarrow \mathbb{R}$	$-): \mathbb{Z} \rightarrow \mathbb{R}$	$-): \mathbb{Z} \rightarrow \mathbb{R}$. Now find a right adjoint for the same map, and show it is correct.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches- F00444	39	■ 1.000	$P=Q=\underline{3}$	$P = Q = \underline{3}$	<i>Exercise 1.99.</i> Consider the preorder $P = Q = \underline{3}$.
fong-spivak-seven-sketches- F00445	39	■ 1.000	f, g	f, g	1. Let f, g be the monotone maps shown below:
fong-spivak-seven-sketches- F00446	39	■ 0.999	$1 \leq p, q \leq 3$	$1 \leq p, q \leq 3$	Is it the case that f is left adjoint to g ? Check that for each $1 \leq p, q \leq 3$, one has
fong-spivak-seven-sketches- F00447	39	■ 1.000	$f(p) \leq q$	$f(p) \leq q$	$f(p) \leq q$ iff $p \leq g(q)$.
fong-spivak-seven-sketches- F00448	39	■ 1.000	$p \leq g(q)$	$p \leq g(q)$	$f(p) \leq q$ iff $p \leq g(q)$.
fong-spivak-seven-sketches- F00449	40	■ 1.000	$L: \mathbb{Z} \rightarrow \mathbb{R}$	$L: \mathbb{Z} \rightarrow \mathbb{R}$	1. Does $\lceil -/3 \rceil$ have a left adjoint $L: \mathbb{Z} \rightarrow \mathbb{R}$?
fong-spivak-seven-sketches- F00450	40	■ 1.000	$\operatorname{Prt}(S)$	$\operatorname{Prt}(S)$	Recall from Example 1.52 that we can understand the set $\operatorname{Prt}(S)$ of partitions on a set S
fong-spivak-seven-sketches- F00451	40	■ 1.000	$g: S \rightarrow T$	$g: S \rightarrow T$	Suppose we are given any function $g: S \rightarrow T$. We will show that this function g
fong-spivak-seven-sketches- F00452	40	■ 0.933	$g_!: \operatorname{Prt}(S) \leftrightsquigarrow \operatorname{Prt}(T): g^*$	$g_!: \operatorname{Prt}(S) \leftrightsquigarrow \operatorname{Prt}(T): g^*$	induces a Galois connection $g_!: \operatorname{Prt}(S) \leftrightsquigarrow \operatorname{Prt}(T): g^*$, between preorder of S -partitions
fong-spivak-seven-sketches- F00453	40	■ 0.991	$g_!: \operatorname{Prt}(S) \rightarrow \operatorname{Prt}(T)$	$g_!: \operatorname{Prt}(S) \rightarrow \operatorname{Prt}(T)$	Start with $g: S \rightarrow T$; we first want to understand $g_!: \operatorname{Prt}(S) \rightarrow \operatorname{Prt}(T)$. So start with
fong-spivak-seven-sketches- F00454	40	■ 1.000	$\sim_S$	$\sim_S$	a partition $\sim_S$ of S . To begin the process of obtaining a partition $\sim_T$ on T , say that two
fong-spivak-seven-sketches- F00455	40	■ 1.000	$\sim_T$	$\sim_T$	a partition $\sim_S$ of S . To begin the process of obtaining a partition $\sim_T$ on T , say that two
fong-spivak-seven-sketches- F00456	40	■ 0.997	$t_1, t_2 \in T$	$t_1, t_2 \in T$	elements $t_1, t_2 \in T$ are in the same part, $t_1 \sim_T t_2$, if there exist $s_1, s_2 \in S$ with such
fong-spivak-seven-sketches- F00457	40	■ 0.997	$t_1 \sim_T t_2$	$t_1 \sim_T t_2$	elements $t_1, t_2 \in T$ are in the same part, $t_1 \sim_T t_2$, if there exist $s_1, s_2 \in S$ with such
fong-spivak-seven-sketches- F00458	40	■ 0.990	$s_1 \sim_S s_2$	$s_1 \sim_S s_2$	that $s_1 \sim_S s_2$ and $g(s_1) = t_1$ and $g(s_2) = t_2$. However, the result of doing so will not
fong-spivak-seven-sketches- F00459	40	■ 0.990	$g(s_1) = t_1$	$g(s_1) = t_1$	that $s_1 \sim_S s_2$ and $g(s_1) = t_1$ and $g(s_2) = t_2$. However, the result of doing so will not
fong-spivak-seven-sketches- F00460	40	■ 0.990	$g(s_2) = t_2$	$g(s_2) = t_2$	that $s_1 \sim_S s_2$ and $g(s_1) = t_1$ and $g(s_2) = t_2$. However, the result of doing so will not
fong-spivak-seven-sketches- F00461	41	■ 1.000	$t_2 \sim_T t_3$	$t_2 \sim_T t_3$	necessarily be transitive—you may get $t_1 \sim_T t_2$ and $t_2 \sim_T t_3$ without $t_1 \sim_T^? t_3$ —and
fong-spivak-seven-sketches- F00462	41	■ 1.000	$t_1 \sim_T^? t_3$	$t_1 \sim_T^? t_3$	necessarily be transitive—you may get $t_1 \sim_T t_2$ and $t_2 \sim_T t_3$ without $t_1 \sim_T^? t_3$ —and
fong-spivak-seven-sketches- F00463	41	■ 0.522	$g_!(\sim_S) := \sim_T$	$g_!(\sim_S) := \sim_T$	pieces (take the transitive closure). The result is $g_!(\sim_S) := \sim_T$.
fong-spivak-seven-sketches- F00464	41	■ 0.813	$g^*: \operatorname{Prt}(T) \rightarrow \operatorname{Prt}(S)$	$g^*: \operatorname{Prt}(T) \rightarrow \operatorname{Prt}(S)$	Again starting with g , we want to get the right adjoint $g^*: \operatorname{Prt}(T) \rightarrow \operatorname{Prt}(S)$. So start
fong-spivak-seven-sketches- F00465	41	■ 1.000	$g(s_1) \sim_T g(s_2)$	$g(s_1) \sim_T g(s_2)$	with a partition $\sim_T$ of T . Get a partition $\sim_S$ on S by saying that $s_1 \sim_S s_2$ iff $g(s_1) \sim_T g(s_2)$.
fong-spivak-seven-sketches- F00466	41	■ 1.000	$g^*(\sim_T) := \sim_S$	$g^*(\sim_T) := \sim_S$	The result is $g^*(\sim_T) := \sim_S$.
fong-spivak-seven-sketches- F00467	41	■ 1.000	$S = \{1, 2, 3, 4\}, T = \{12, 3, 4\}$	$S = \{1, 2, 3, 4\}, T = \{12, 3, 4\}$	<i>Example 1.102.</i> Let $S = \{1, 2, 3, 4\}, T = \{12, 3, 4\}$, and $g: S \rightarrow T$ by $g(1) := g(2) := 12$,
fong-spivak-seven-sketches- F00468	41	■ 1.000	$g(1) := g(2) := 12$	$g(1) := g(2) := 12$	<i>Example 1.102.</i> Let $S = \{1, 2, 3, 4\}, T = \{12, 3, 4\}$, and $g: S \rightarrow T$ by $g(1) := g(2) := 12$,
fong-spivak-seven-sketches- F00469	41	■ 0.813	$g(3) := 3$	$g(3) := 3$	$g(3) := 3$, and $g(4) := 4$. The partition shown left below is translated by $g_!$ to the
fong-spivak-seven-sketches- F00470	41	■ 0.813	$g(4) := 4$	$g(4) := 4$	$g(3) := 3$, and $g(4) := 4$. The partition shown left below is translated by $g_!$ to the

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F00471	41	0.813	$g_{\{!\}}$	$g_{!}$	$g(3) := 3$, and $g(4) := 4$. The partition shown left below is translated by $g_{!}$ to the
fong-spivak-seven-sketches-F00472	41	0.999	$c: S \rightarrowtail P$	$c: S \rightarrow P$	different ones and for each one, call it $c: S \rightarrowtail P$, find $g_{!}(c)$, where S, T , and $g: S \rightarrow T$
fong-spivak-seven-sketches-F00473	41	0.999	$g_{\{!\}}(c)$	$g_{!}(c)$	different ones and for each one, call it $c: S \rightarrowtail P$, find $g_{!}(c)$, where S, T , and $g: S \rightarrow T$
fong-spivak-seven-sketches-F00474	41	0.999	S, T	S, T	different ones and for each one, call it $c: S \rightarrowtail P$, find $g_{!}(c)$, where S, T , and $g: S \rightarrow T$
fong-spivak-seven-sketches-F00475	41	1.000	$g^{\{*\}}$	g^*	Here is a picture of how g^* takes a partition on T and “pulls it back” to a partition on S :
fong-spivak-seven-sketches-F00476	41	1.000	$T = \{12, 3, 4\}$	$T = \{12, 3, 4\}$	$T = \{12, 3, 4\}$. Using the same S and $g: S \rightarrow T$ as in Example 1.102, determine the
fong-spivak-seven-sketches-F00477	41	1.000	$g^{\{*\}}(c)$	$g^*(c)$	partition $g^*(c)$ on S for each of the five partitions $c: T \rightarrowtail P$.
fong-spivak-seven-sketches-F00478	41	1.000	$c: T \rightarrowtail P$	$c: T \rightarrow P$	partition $g^*(c)$ on S for each of the five partitions $c: T \rightarrowtail P$.
fong-spivak-seven-sketches-F00479	42	1.000	$d: T \rightarrowtail P^{\{\prime\}}$	$d: T \rightarrow P'$	2. Choose any coarser partition $d: T \rightarrowtail P'$, i.e. where $g_{!}(c) \leq d$.
fong-spivak-seven-sketches-F00480	42	1.000	$g_{\{!\}}(c) \leq d$	$g_{!}(c) \leq d$	2. Choose any coarser partition $d: T \rightarrowtail P'$, i.e. where $g_{!}(c) \leq d$.
fong-spivak-seven-sketches-F00481	42	1.000	$e: T \rightarrowtail Q$	$e: T \rightarrow Q$	3. Choose any non-coarser partition $e: T \rightarrowtail Q$, i.e. where $g_{!}(c) \not\leq e$. (If you can’t do
fong-spivak-seven-sketches-F00482	42	1.000	$g_{\{!\}}(c) \not\leq e$	$g_{!}(c) \not\leq e$	3. Choose any non-coarser partition $e: T \rightarrowtail Q$, i.e. where $g_{!}(c) \not\leq e$. (If you can’t do
fong-spivak-seven-sketches-F00483	42	1.000	$g^{\{*\}}(d)$	$g^*(d)$	4. Find $g^*(d)$ and $g^*(e)$.
fong-spivak-seven-sketches-F00484	42	1.000	$g^{\{*\}}(e)$	$g^*(e)$	4. Find $g^*(d)$ and $g^*(e)$.
fong-spivak-seven-sketches-F00485	42	0.999	$c \leq g^{\{*\}}(d)$	$c \leq g^*(d)$	$g_{!}(c) \not\leq e$, we should have $c \leq g^*(d)$ and $c \not\leq g^*(e)$. Show that this is true.
fong-spivak-seven-sketches-F00486	42	0.999	$c \not\leq g^{\{*\}}(e)$	$c \not\leq g^*(e)$	$g_{!}(c) \not\leq e$, we should have $c \leq g^*(d)$ and $c \not\leq g^*(e)$. Show that this is true.
fong-spivak-seven-sketches-F00487	42	1.000	$q := f(p)$	$q := f(p)$	<i>Proof.</i> Suppose f is left adjoint to g . Take any $p \in P$, and let $q := f(p)$. By reflexivity,
fong-spivak-seven-sketches-F00488	42	1.000	$p \leq g(f(p))$	$p \leq g(f(p))$	this means $p \leq g(f(p))$. The proof that $f(g(q)) \leq q$ is similar.
fong-spivak-seven-sketches-F00489	42	1.000	$f(g(q)) \leq q$	$f(g(q)) \leq q$	this means $p \leq g(f(p))$. The proof that $f(g(q)) \leq q$ is similar.
fong-spivak-seven-sketches-F00490	42	1.000	$g(f(p)) \leq g(q)$	$g(f(p)) \leq g(q)$	$f(p) \leq q$ iff $p \leq g(q)$. Suppose $f(p) \leq q$; then since g is monotonic, $g(f(p)) \leq g(q)$, but
fong-spivak-seven-sketches-F00491	42	1.000	$g^{\{\prime\}}$	g'	That means, for any other right adjoint g' , we have $g(q) \cong g'(q)$ for all $q \in Q$.
fong-spivak-seven-sketches-F00492	42	1.000	$g(q) \cong g^{\{\prime\}}(q)$	$g(q) \cong g'(q)$	That means, for any other right adjoint g' , we have $g(q) \cong g'(q)$ for all $q \in Q$.
fong-spivak-seven-sketches-F00493	43	1.000	$A \subseteq Q$	$A \subseteq Q$	$g: Q \rightarrow P$. Suppose $A \subseteq Q$ any subset, and let $g(A) := \{g(a) \mid a \in A\}$ be its image.
fong-spivak-seven-sketches-F00494	43	1.000	$g(A) := \{g(a) \mid a \in A\}$	$g(A) := \{g(a) \mid a \in A\}$	$g: Q \rightarrow P$. Suppose $A \subseteq Q$ any subset, and let $g(A) := \{g(a) \mid a \in A\}$ be its image.
fong-spivak-seven-sketches-F00495	43	1.000	$\bigwedge A \in Q$	$\bigwedge A \in Q$	Then if A has a meet $\bigwedge A \in Q$ then $g(A)$ has a meet $\bigwedge g(A)$ in P , and we have
fong-spivak-seven-sketches-F00496	43	1.000	$g(A)$	$g(A)$	Then if A has a meet $\bigwedge A \in Q$ then $g(A)$ has a meet $\bigwedge g(A)$ in P , and we have

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_1 F00497	43	■ 1.000	$\wedge g(A)$	$\bigwedge g(A)$	Then if A has a meet $\bigwedge A \in Q$ then $g(A)$ has a meet $\bigwedge g(A)$ in P , and we have
fong-spivak-seven-sketches_1 F00498	43	■ 1.000	$\bigvee A \in P$	$\bigvee A \in P$	any subset that has a join $\bigvee A \in P$, then $f(A)$ has a join $\bigvee f(A)$ in Q , and we have
fong-spivak-seven-sketches_1 F00499	43	■ 1.000	$f(A)$	$f(A)$	any subset that has a join $\bigvee A \in P$, then $f(A)$ has a join $\bigvee f(A)$ in Q , and we have
fong-spivak-seven-sketches_1 F00500	43	■ 1.000	$\bigvee f(A)$	$\bigvee f(A)$	any subset that has a join $\bigvee A \in P$, then $f(A)$ has a join $\bigvee f(A)$ in Q , and we have
fong-spivak-seven-sketches_1 F00501	43	■ 1.000	$m := \wedge A$	$m := \bigwedge A$	f . Let $A \subseteq Q$ be any subset and let $m := \bigwedge A$ be its meet. Then since g is monotone
fong-spivak-seven-sketches_1 F00502	43	■ 1.000	$g(m) \leq g(a)$	$g(m) \leq g(a)$	$g(m) \leq g(a)$ for all $a \in A$, so $g(m)$ is a lower bound for the set $g(A)$. We will be done if
fong-spivak-seven-sketches_1 F00503	43	■ 1.000	$g(m)$	$g(m)$	$g(m) \leq g(a)$ for all $a \in A$, so $g(m)$ is a lower bound for the set $g(A)$. We will be done if
fong-spivak-seven-sketches_1 F00504	43	■ 1.000	$b \leq g(a)$	$b \leq g(a)$	$b \leq g(a)$ and we want to show $b \leq g(m)$. Then by definition of g being a right adjoint
fong-spivak-seven-sketches_1 F00505	43	■ 1.000	$b \leq g(m)$	$b \leq g(m)$	$b \leq g(a)$ and we want to show $b \leq g(m)$. Then by definition of g being a right adjoint
fong-spivak-seven-sketches_1 F00506	43	■ 1.000	$f(b) \leq a$	$f(b) \leq a$	(Definition 1.95), we also have $f(b) \leq a$. This means that $f(b)$ is a lower bound for A in
fong-spivak-seven-sketches_1 F00507	43	■ 1.000	$f(b)$	$f(b)$	(Definition 1.95), we also have $f(b) \leq a$. This means that $f(b)$ is a lower bound for A in
fong-spivak-seven-sketches_1 F00508	43	■ 1.000	$f(b) \leq m$	$f(b) \leq m$	Q . Since the meet m is the greatest lower bound, we have $f(b) \leq m$. Once again using
fong-spivak-seven-sketches_1 F00509	43	■ 1.000	$f(3.9) := 4$	$f(3.9) := 4$	as possible but with $f(3.9) := 4$. Both are f and g monotonic, and one can check that g
fong-spivak-seven-sketches_1 F00510	44	■ 1.000	$1 \vee 2 = 4$	$1 \vee 2 = 4$	is right adjoint to f (see Exercise 1.114). But g does not preserve joins because $1 \vee 2 = 4$
fong-spivak-seven-sketches_1 F00511	44	■ 1.000	$g(1) \vee g(2) = 1 \vee 2 = 3.9 \neq 4 = g(4)$	$g(1) \vee g(2) = 1 \vee 2 = 3.9 \neq 4 = g(4)$	holds in Q , whereas $g(1) \vee g(2) = 1 \vee 2 = 3.9 \neq 4 = g(4)$ in P .
fong-spivak-seven-sketches_1 F00512	44	■ 1.000	$\{q' \in Q \mid p' \leq g(q')\} \subseteq \{q \in Q \mid p \leq g(q)\}$	$\{q' \in Q \mid p' \leq g(q')\} \subseteq \{q \in Q \mid p \leq g(q)\}$	need to show it is monotone. So suppose that $p \leq p'$. Then $\{q' \in Q \mid p' \leq g(q')\} \subseteq \{q \in Q \mid p \leq g(q)\}$. By Proposition 1.91, this implies $f(p) \leq f(p')$. Thus f is monotone.
fong-spivak-seven-sketches_1 F00513	44	■ 1.000	$Q \mid p \leq g(q)\}$	$Q \mid p \leq g(q)\}$	$Q \mid p \leq g(q)\}$. By Proposition 1.91, this implies $f(p) \leq f(p')$. Thus f is monotone.
fong-spivak-seven-sketches_1 F00514	44	■ 1.000	$f(p) \leq f(p')$	$f(p) \leq f(p')$	$Q \mid p \leq g(q)\}$. By Proposition 1.91, this implies $f(p) \leq f(p')$. Thus f is monotone.
fong-spivak-seven-sketches_1 F00515	44	■ 1.000	$p_0 \leq g(f(p_0))$	$p_0 \leq g(f(p_0))$	By Proposition 1.111, it suffices to show that $p_0 \leq g(f(p_0))$ and that $f(g(q_0)) \leq q_0$
fong-spivak-seven-sketches_1 F00516	44	■ 1.000	$f(g(q_0)) \leq q_0$	$f(g(q_0)) \leq q_0$	By Proposition 1.111, it suffices to show that $p_0 \leq g(f(p_0))$ and that $f(g(q_0)) \leq q_0$
fong-spivak-seven-sketches_1 F00517	44	■ 1.000	$p_0 \in P$	$p_0 \in P$	for all $p_0 \in P$ and $q_0 \in Q$. For the first, we have
fong-spivak-seven-sketches_1 F00518	44	■ 1.000	$q_0 \in Q$	$q_0 \in Q$	for all $p_0 \in P$ and $q_0 \in Q$. For the first, we have
fong-spivak-seven-sketches_1 F00519	44	■ 1.000	p_0	p_0	where the first inequality follows from the fact that if p_0 is below every element of a
fong-spivak-seven-sketches_1 F00520	44	■ 1.000	$\{q_0\} \subseteq \{q \in Q \mid g(q_0) \leq q\}$	$\{q_0\} \subseteq \{q \in Q \mid g(q_0) \leq q\}$	where the first inequality follows from Proposition 1.91 since $\{q_0\} \subseteq \{q \in Q \mid g(q_0) \leq q\}$
fong-spivak-seven-sketches_1 F00521	44	■ 0.999	$g(q)\}$	$g(q)\}$	$g(q)\}$, and the fact that $\bigwedge \{q_0\} = q_0$.
fong-spivak-seven-sketches_1 F00522	44	■ 0.999	$\bigwedge \{q_0\} = q_0$	$\bigwedge \{q_0\} = q_0$	$g(q)\}$, and the fact that $\bigwedge \{q_0\} = q_0$.
fong-spivak-seven-sketches_1 F00523	44	■ 1.000	$f^*: \mathbf{P}(Y) \rightarrow \mathbf{P}(X)$	$f^*: \mathbf{P}(Y) \rightarrow \mathbf{P}(X)$	Then we have the monotone map $f^*: \mathbf{P}(Y) \rightarrow \mathbf{P}(X)$ that category theorists call

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F00551	46	1.000	$j(x)$	$j(x)$	means that if an expression x can be rewritten into expression y , then the reduction $j(x)$
fong-spivak-seven-sketches-F00552	46	1.000	$j(y)$	$j(y)$	can be rewritten into $j(y)$. Moreover, the requirement $x \leq j(x)$ implies that j can only
fong-spivak-seven-sketches-F00553	46	1.000	$x \leq j(x)$	$x \leq j(x)$	can be rewritten into $j(y)$. Moreover, the requirement $x \leq j(x)$ implies that j can only
fong-spivak-seven-sketches-F00554	46	1.000	j	j	can be rewritten into $j(y)$. Moreover, the requirement $x \leq j(x)$ implies that j can only
fong-spivak-seven-sketches-F00555	46	1.000	$j(j(x))=j(x)$	$j(j(x)) = j(x)$	$j(j(x)) = j(x)$ implies if you try to reduce an expression that has already been reduced,
fong-spivak-seven-sketches-F00556	46	0.771	$\{ \}_{j}$	j	fix_j to have elements the fixed points of j ; that is,
fong-spivak-seven-sketches-F00557	46	1.000	$j(p)$	$j(p)$	Note that $j(p)$ is a fixed point for all $p \in P$, since $j(j(p)) \cong j(p)$.
fong-spivak-seven-sketches-F00558	46	0.502	$j: P \rightarrow$	$j: P \rightarrow$	We define an adjunction with left adjoint $j: P \rightarrow \text{fix}_j$ sending p to $j(p)$, and right
fong-spivak-seven-sketches-F00559	46	0.502	$\{ \}_{j}$	j	We define an adjunction with left adjoint $j: P \rightarrow \text{fix}_j$ sending p to $j(p)$, and right
fong-spivak-seven-sketches-F00560	46	0.910	$\{ \}_{j} \rightarrow P$	$j \rightarrow P$	adjoint $g: \text{fix}_j \rightarrow P$ simply the inclusion of the sub-preorder. To see it's really an
fong-spivak-seven-sketches-F00561	46	0.990	$q \in \text{fix}_j$	$q \in \text{fix}_j$	adjunction, we need to see that for any $p \in P$ and $q \in \text{fix}_j$, we have $j(p) \leq q$ if and
fong-spivak-seven-sketches-F00562	46	0.990	$j(p) \leq q$	$j(p) \leq q$	adjunction, we need to see that for any $p \in P$ and $q \in \text{fix}_j$, we have $j(p) \leq q$ if and
fong-spivak-seven-sketches-F00563	46	1.000	$j(p) \leq j(q) \cong q$	$j(p) \leq j(q) \cong q$	transitivity. Conversely, since q is a fixed point, $p \leq q$ implies $j(p) \leq j(q) \cong q$.
fong-spivak-seven-sketches-F00564	47	0.999	$p \Rightarrow q$	$p \Rightarrow q$	$p \Rightarrow q$. Since $p \Rightarrow p$ and since whenever $p \Rightarrow q$ and $q \Rightarrow r$, we also have $p \Rightarrow r$, this is
fong-spivak-seven-sketches-F00565	47	0.999	$p \Rightarrow p$	$p \Rightarrow p$	$p \Rightarrow q$. Since $p \Rightarrow p$ and since whenever $p \Rightarrow q$ and $q \Rightarrow r$, we also have $p \Rightarrow r$, this is
fong-spivak-seven-sketches-F00566	47	0.999	$q \Rightarrow r$	$q \Rightarrow r$	$p \Rightarrow q$. Since $p \Rightarrow p$ and since whenever $p \Rightarrow q$ and $q \Rightarrow r$, we also have $p \Rightarrow r$, this is
fong-spivak-seven-sketches-F00567	47	0.999	$p \Rightarrow r$	$p \Rightarrow r$	$p \Rightarrow q$. Since $p \Rightarrow p$ and since whenever $p \Rightarrow q$ and $q \Rightarrow r$, we also have $p \Rightarrow r$, this is
fong-spivak-seven-sketches-F00568	47	1.000	$p \Rightarrow j(p)$	$p \Rightarrow j(p)$	propositions to propositions, for which $p \Rightarrow j(p)$ and $j(j(p)) = j(p)$. An example of a
fong-spivak-seven-sketches-F00569	47	1.000	$j(j(p))=j(p)$	$j(j(p)) = j(p)$	propositions to propositions, for which $p \Rightarrow j(p)$ and $j(j(p)) = j(p)$. An example of a
fong-spivak-seven-sketches-F00570	47	0.966	$p^{\prime\prime}$	p''	Bob is in San Diego, p'' means $B \Rightarrow p$. Let's see why $B \Rightarrow -$ is a closure operator.
fong-spivak-seven-sketches-F00571	47	0.966	$B \Rightarrow p$	$B \Rightarrow p$	Bob is in San Diego, p'' means $B \Rightarrow p$. Let's see why $B \Rightarrow -$ is a closure operator.
fong-spivak-seven-sketches-F00572	47	0.966	$B \Rightarrow -$	$B \Rightarrow -$	Bob is in San Diego, p'' means $B \Rightarrow p$. Let's see why $B \Rightarrow -$ is a closure operator.
fong-spivak-seven-sketches-F00573	47	1.000	$p^{\prime}$	p'	“assuming Bob is in San Diego it is the case that, assuming Bob is in San Diego, p' is
fong-spivak-seven-sketches-F00574	47	0.622	$\mathbf{Rel}(S)$	$\mathbf{Rel}(S)$	Given any set S , there is a set $\mathbf{Rel}(S)$ of binary relations on S . An element $R \in \mathbf{Rel}(S)$
fong-spivak-seven-sketches-F00575	47	0.622	$R \in \mathbf{Rel}(S)$	$R \in \mathbf{Rel}(S)$	Given any set S , there is a set $\mathbf{Rel}(S)$ of binary relations on S . An element $R \in \mathbf{Rel}(S)$
fong-spivak-seven-sketches-F00576	47	1.000	$R \subseteq S \times S$	$R \subseteq S \times S$	is formally a subset $R \subseteq S \times S$. The set $\mathbf{Rel}(S)$ can be given an order via the subset
fong-spivak-seven-sketches-F00577	47	1.000	$R \subseteq R^{\prime}$	$R \subseteq R'$	relation, $R \subseteq R'$, i.e. if whenever $R(s_1, s_2)$ holds then so does $R'(s_1, s_2)$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-47 F00578	47	■ 1.000	$R\left(s_1, s_2\right)$	$R(s_1, s_2)$	relation, $R \subseteq R'$, i.e. if whenever $R(s_1, s_2)$ holds then so does $R'(s_1, s_2)$.
fong-spivak-seven-sketches-47 F00579	47	■ 1.000	$R^{\prime}\left(s_1, s_2\right)$	$R'(s_1, s_2)$	relation, $R \subseteq R'$, i.e. if whenever $R(s_1, s_2)$ holds then so does $R'(s_1, s_2)$.
fong-spivak-seven-sketches-47 F00580	47	■ 0.999	$\operatorname{\mathbf{Rel}}(\{1\})$	$\mathbf{Rel}(\{1\})$	For example, the Hasse diagram for $\mathbf{Rel}(\{1\})$ is:
fong-spivak-seven-sketches-47 F00581	47	■ 0.998	$\operatorname{\mathbf{Rel}}(\{1,2\})$	$\mathbf{Rel}(\{1,2\})$	<i>Exercise 1.124.</i> Draw the Hasse diagram for the preorder $\mathbf{Rel}(\{1,2\})$ of all binary
fong-spivak-seven-sketches-47 F00582	47	■ 0.755	$\{1,2\}$	$\{1,2\}$	relations on the set $\{1,2\}$.
fong-spivak-seven-sketches-47 F00583	47	■ 1.000	$\operatorname{\mathbf{Pos}}(S)$	$\mathbf{Pos}(S)$	For any set S , there is also a set $\mathbf{Pos}(S)$, consisting of all the preorder relations on S .
fong-spivak-seven-sketches-47 F00584	47	■ 0.961	$\sqsubseteq$	$\sqsubseteq$	In fact there is a preorder structure $\sqsubseteq$ on $\mathbf{Pos}(S)$, again given by inclusion: $\leq$ is below
fong-spivak-seven-sketches-47 F00585	47	■ 0.982	$\leq^{\prime}$	$\leq'$	$\leq'$ (we'll write $\leq \sqsubseteq \leq'$) if $a \leq b$ implies $a \leq' b$ for every $a, b \in S$. A preorder of preorder
fong-spivak-seven-sketches-47 F00586	47	■ 0.982	$\leq \leq \leq^{\prime}$	$\leq \leq \leq'$	$\leq'$ (we'll write $\leq \sqsubseteq \leq'$) if $a \leq b$ implies $a \leq' b$ for every $a, b \in S$. A preorder of preorder
fong-spivak-seven-sketches-47 F00587	47	■ 0.982	$a \leq^{\prime} b$	$a \leq' b$	$\leq'$ (we'll write $\leq \sqsubseteq \leq'$) if $a \leq b$ implies $a \leq' b$ for every $a, b \in S$. A preorder of preorder
fong-spivak-seven-sketches-47 F00588	47	■ 0.982	$a, b \in S$	$a, b \in S$	$\leq'$ (we'll write $\leq \sqsubseteq \leq'$) if $a \leq b$ implies $a \leq' b$ for every $a, b \in S$. A preorder of preorder
fong-spivak-seven-sketches-47 F00589	47	■ 1.000	$\operatorname{\mathbf{Pos}}(S) \rightarrow \operatorname{\mathbf{Rel}}(S)$	$\mathbf{Pos}(S) \rightarrow \mathbf{Rel}(S)$	$\mathbf{Pos}(S) \rightarrow \mathbf{Rel}(S)$. This is the right adjoint of a Galois connection. Its left adjoint
fong-spivak-seven-sketches-47 F00590	47	■ 0.977	$\operatorname{\mathbf{Cl}}: \operatorname{\mathbf{Rel}}(S) \rightarrow \operatorname{\mathbf{Pos}}(S)$	$\mathbf{Cl}: \mathbf{Rel}(S) \rightarrow \mathbf{Pos}(S)$	is a monotone map $\mathbf{Cl}: \mathbf{Rel}(S) \rightarrow \mathbf{Pos}(S)$ given by taking any relation R , writing it in
fong-spivak-seven-sketches-47 F00591	47	■ 1.000	$s \leq s$	$s \leq s$	infix notation using $\leq$, and taking the reflexive and transitive closure, i.e. adding $s \leq s$
fong-spivak-seven-sketches-47 F00592	47	■ 1.000	$s \leq u$	$s \leq u$	for every s and adding $s \leq u$ whenever $s \leq t$ and $t \leq u$.
fong-spivak-seven-sketches-47 F00593	47	■ 1.000	$s \leq t$	$s \leq t$	for every s and adding $s \leq u$ whenever $s \leq t$ and $t \leq u$.
fong-spivak-seven-sketches-47 F00594	47	■ 1.000	$t \leq u$	$t \leq u$	for every s and adding $s \leq u$ whenever $s \leq t$ and $t \leq u$.
fong-spivak-seven-sketches-47 F00595	47	■ 1.000	$S = \{1,2,3\}$	$S = \{1,2,3\}$	<i>Exercise 1.125.</i> Let $S = \{1,2,3\}$. Let's try to understand the adjunction discussed
fong-spivak-seven-sketches-47 F00596	47	■ 1.000	$U(\leq)$	$U(\leq)$	1. Come up with any preorder relation $\leq$ on S , and define $U(\leq)$ to be the subset
fong-spivak-seven-sketches-47 F00597	47	■ 1.000	$U(\leq) := \left\{ \left(s_1, s_2 \right) \mid s_1 \leq s_2 \right\} \subseteq S \times S$	$U(\leq) := \{(s_1, s_2) \mid s_1 \leq s_2\} \subseteq S \times S$	i.e. $U(\leq)$ is the image of $\leq$ under the inclusion
fong-spivak-seven-sketches-48 F00598	48	■ 1.000	$Q \subseteq S \times S$	$Q \subseteq S \times S$	2. Come up with any two binary relations $Q \subseteq S \times S$ and $Q' \subseteq S \times S$ such that
fong-spivak-seven-sketches-48 F00599	48	■ 1.000	$Q^{\prime} \subseteq S \times S$	$Q' \subseteq S \times S$	2. Come up with any two binary relations $Q \subseteq S \times S$ and $Q' \subseteq S \times S$ such that
fong-spivak-seven-sketches-48 F00600	48	■ 1.000	$Q \subseteq U(\leq)$	$Q \subseteq U(\leq)$	$Q \subseteq U(\leq)$ but $Q' \not\subseteq U(\leq)$. Note that your choice of Q, Q' do not have to come
fong-spivak-seven-sketches-48 F00601	48	■ 1.000	$Q^{\prime} \not\subseteq U(\leq)$	$Q' \not\subseteq U(\leq)$	$Q \subseteq U(\leq)$ but $Q' \not\subseteq U(\leq)$. Note that your choice of Q, Q' do not have to come
fong-spivak-seven-sketches-48 F00602	48	■ 1.000	$Q, Q^{\prime}$	Q, Q'	$Q \subseteq U(\leq)$ but $Q' \not\subseteq U(\leq)$. Note that your choice of Q, Q' do not have to come
fong-spivak-seven-sketches-48 F00603	48	■ 0.973	$\operatorname{\mathbf{Cl}}(Q) \sqsubseteq \leq$	$\mathbf{Cl}(Q) \sqsubseteq \leq$	show that $\mathbf{Cl}(Q) \sqsubseteq \leq$, where $\sqsubseteq$ is the order on $\mathbf{Pos}(S)$, defined immediately above
fong-spivak-seven-sketches-48 F00604	48	■ 0.973	$\operatorname{\mathbf{Pos}}(S)$	$\mathbf{Pos}(S)$	show that $\mathbf{Cl}(Q) \sqsubseteq \leq$, where $\sqsubseteq$ is the order on $\mathbf{Pos}(S)$, defined immediately above

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_1 F00605	48	■ 0.986	$\mathrm{Cl}\left(Q^{\prime}\right) \backslash \subseteq \leq$	$\mathrm{Cl}\left(Q^{\prime}\right) \not\subseteq \leq$	4. Concretely show that $\mathrm{Cl}\left(Q^{\prime}\right) \not\subseteq \leq$.
fong-spivak-seven-sketches_1 F00606	48	■ 1.000	$a \otimes b$	$a \otimes b$	idea of a monoidal structure on preorders. This allows us to construct an element $a \otimes b$
fong-spivak-seven-sketches_1 F00607	51	■ 0.953	$\boldsymbol{a}$	$\boldsymbol{a}$	2.1 Getting from a to b
fong-spivak-seven-sketches_1 F00608	51	■ 0.953	$\boldsymbol{b}$	$\boldsymbol{b}$	2.1 Getting from a to b
fong-spivak-seven-sketches_1 F00609	53	■ 1.000	$d \in[0, \infty]$	$d \in[0, \infty]$	the question of getting from a to b has an answer $d \in[0, \infty]$, a cost.
fong-spivak-seven-sketches_1 F00610	53	■ 1.000	$I \in X$	$I \in X$	(i) an element $I \in X$, called the <i>monoidal unit</i> , and
fong-spivak-seven-sketches_1 F00611	53	■ 0.983	$\otimes: X \times X \rightarrow X$	$\otimes: X \times X \rightarrow X$	(ii) a function $\otimes: X \times X \rightarrow X$, called the <i>monoidal product</i> .
fong-spivak-seven-sketches_1 F00612	53	■ 0.997	$\otimes\left(x_1, x_2\right)=$	$\otimes\left(x_1, x_2\right)=$	These constituents must satisfy the following properties, where we write $\otimes\left(x_1, x_2\right)=$
fong-spivak-seven-sketches_1 F00613	53	■ 1.000	$x_1 \otimes x_2$	$x_1 \otimes x_2$	$x_1 \otimes x_2$:
fong-spivak-seven-sketches_1 F00614	54	■ 1.000	$x_1, x_2, y_1, y_2 \in X$	$x_1, x_2, y_1, y_2 \in X$	(a) for all $x_1, x_2, y_1, y_2 \in X$, if $x_1 \leq y_1$ and $x_2 \leq y_2$, then $x_1 \otimes x_2 \leq y_1 \otimes y_2$,
fong-spivak-seven-sketches_1 F00615	54	■ 1.000	$x_1 \leq y_1$	$x_1 \leq y_1$	(a) for all $x_1, x_2, y_1, y_2 \in X$, if $x_1 \leq y_1$ and $x_2 \leq y_2$, then $x_1 \otimes x_2 \leq y_1 \otimes y_2$,
fong-spivak-seven-sketches_1 F00616	54	■ 1.000	$x_2 \leq y_2$	$x_2 \leq y_2$	(a) for all $x_1, x_2, y_1, y_2 \in X$, if $x_1 \leq y_1$ and $x_2 \leq y_2$, then $x_1 \otimes x_2 \leq y_1 \otimes y_2$,
fong-spivak-seven-sketches_1 F00617	54	■ 1.000	$x_1 \otimes x_2 \leq y_1 \otimes y_2$	$x_1 \otimes x_2 \leq y_1 \otimes y_2$	(a) for all $x_1, x_2, y_1, y_2 \in X$, if $x_1 \leq y_1$ and $x_2 \leq y_2$, then $x_1 \otimes x_2 \leq y_1 \otimes y_2$,
fong-spivak-seven-sketches_1 F00618	54	■ 1.000	$I \otimes x=x$	$I \otimes x=x$	(b) for all $x \in X$, the equations $I \otimes x=x$ and $x \otimes I=x$ hold,
fong-spivak-seven-sketches_1 F00619	54	■ 1.000	$x \otimes I=x$	$x \otimes I=x$	(b) for all $x \in X$, the equations $I \otimes x=x$ and $x \otimes I=x$ hold,
fong-spivak-seven-sketches_1 F00620	54	■ 1.000	$x, y, z \in X$	$x, y, z \in X$	(c) for all $x, y, z \in X$, the equation $(x \otimes y) \otimes z=x \otimes(y \otimes z)$ holds, and
fong-spivak-seven-sketches_1 F00621	54	■ 1.000	$(x \otimes y) \otimes z=x \otimes(y \otimes z)$	$(x \otimes y) \otimes z=x \otimes(y \otimes z)$	(c) for all $x, y, z \in X$, the equation $(x \otimes y) \otimes z=x \otimes(y \otimes z)$ holds, and
fong-spivak-seven-sketches_1 F00622	54	■ 1.000	$x \otimes y=y \otimes x$	$x \otimes y=y \otimes x$	(d) for all $x, y \in X$, the equation $x \otimes y=y \otimes x$ holds.
fong-spivak-seven-sketches_1 F00623	54	■ 0.849	$(X, \leq, I, \otimes)$	$(X, \leq, I, \otimes)$	A preorder equipped with a symmetric monoidal structure, $(X, \leq, I, \otimes)$, is called a
fong-spivak-seven-sketches_1 F00624	54	■ 1.000	I	I	Anyone can propose a set X , an order $\leq$ on X , an element I in X , and a binary
fong-spivak-seven-sketches_1 F00625	54	■ 0.933	$\cong$	$\cong$	<i>Remark 2.3.</i> It is often useful to replace $=$ with $\cong$ throughout Definition 2.2. The result is
fong-spivak-seven-sketches_1 F00626	54	■ 0.585	$\{*\}$	$\{*\}$	units we have seen include I (as in the definition), 0, 1, true, false, $\{*\}$, and more.
fong-spivak-seven-sketches_1 F00627	54	■ 0.999	$-5 \leq \sqrt{2}$	$-5 \leq \sqrt{2}$	numbers; e.g. $-5 \leq \sqrt{2}$. We propose 0 as a monoidal unit and $+: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ as a
fong-spivak-seven-sketches_1 F00628	54	■ 0.999	$+: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$	$+: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$	numbers; e.g. $-5 \leq \sqrt{2}$. We propose 0 as a monoidal unit and $+: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ as a
fong-spivak-seven-sketches_1 F00629	54	■ 0.902	$(\mathbb{R}, \leq, 0,+)$	$(\mathbb{R}, \leq, 0,+)$	monoidal product. Does $(\mathbb{R}, \leq, 0,+)$ satisfy the conditions of Definition 2.2?
fong-spivak-seven-sketches_1 F00630	54	■ 1.000	$x_1+x_2 \leq y_1+y_2$	$x_1+x_2 \leq y_1+y_2$	If $x_1 \leq y_1$ and $x_2 \leq y_2$, it is true that $x_1+x_2 \leq y_1+y_2$. It is also true that $0+x=x$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_1 F00631	54	■ 1.000	$0+x=x$	$0 + x = x$	If $x_1 \leq y_1$ and $x_2 \leq y_2$, it is true that $x_1 + x_2 \leq y_1 + y_2$. It is also true that $0 + x = x$
fong-spivak-seven-sketches_1 F00632	54	■ 0.989	$x+0=x$	$x + 0 = x$	and $x + 0 = x$, that $(x + y) + z = x + (y + z)$, and that $x + y = y + x$. Thus $(\mathbb{R}, \leq, 0, +)$
fong-spivak-seven-sketches_1 F00633	54	■ 0.989	$(x+y)+z=x+(y+z)$	$(x + y) + z = x + (y + z)$	and $x + 0 = x$, that $(x + y) + z = x + (y + z)$, and that $x + y = y + x$. Thus $(\mathbb{R}, \leq, 0, +)$
fong-spivak-seven-sketches_1 F00634	54	■ 0.989	$x+y=y+x$	$x + y = y + x$	and $x + 0 = x$, that $(x + y) + z = x + (y + z)$, and that $x + y = y + x$. Thus $(\mathbb{R}, \leq, 0, +)$
fong-spivak-seven-sketches_1 F00635	54	■ 0.554	M	M	Example 2.6. A monoid consists of a set M , a function $*$: $M \times M \rightarrow M$ called the <i>monoid</i>
fong-spivak-seven-sketches_1 F00636	54	■ 0.554	$M \times M \rightarrow M$	$M \times M \rightarrow M$	Example 2.6. A monoid consists of a set M , a function $*$: $M \times M \rightarrow M$ called the <i>monoid</i>
fong-spivak-seven-sketches_1 F00637	54	■ 1.000	$e \in M$	$e \in M$	<i>multiplication</i> , and an element $e \in M$ called the <i>monoid unit</i> , such that, when you write
fong-spivak-seven-sketches_1 F00638	54	■ 0.944	$*(m, n)$	$*(m, n)$	$*(m, n)$ as $m * n$, i.e. using infix notation, the equations
fong-spivak-seven-sketches_1 F00639	54	■ 0.944	$m * n$	$m * n$	$*(m, n)$ as $m * n$, i.e. using infix notation, the equations
fong-spivak-seven-sketches_1 F00640	54	■ 1.000	$m, n, p \in M$	$m, n, p \in M$	hold for all $m, n, p \in M$. It is called <i>commutative</i> if also $m * n = n * m$.
fong-spivak-seven-sketches_1 F00641	54	■ 1.000	$m * n = n * m$	$m * n = n * m$	hold for all $m, n, p \in M$. It is called <i>commutative</i> if also $m * n = n * m$.
fong-spivak-seven-sketches_1 F00642	54	■ 0.912	$\mathbf{Disc}_S$	$\mathbf{Disc}_S$	Every set S determines a discrete preorder $\mathbf{Disc}_S$ (where $m \leq n$ iff $m = n$; see
fong-spivak-seven-sketches_1 F00643	54	■ 0.912	$m=n$	$m = n$	Every set S determines a discrete preorder $\mathbf{Disc}_S$ (where $m \leq n$ iff $m = n$; see
fong-spivak-seven-sketches_1 F00644	54	■ 1.000	$(M, e, *)$	$(M, e, *)$	Example 1.32), and it is easy to check that if $(M, e, *)$ is a commutative monoid then
fong-spivak-seven-sketches_1 F00645	54	■ 0.770	$(\mathbf{Disc}_M, =, e, *)$	$(\mathbf{Disc}_M, =, e, *)$	$(\mathbf{Disc}_M, =, e, *)$ is a symmetric monoidal preorder.
fong-spivak-seven-sketches_1 F00646	54	■ 0.770	$(\mathbf{Disc}_M, =, e, *)$	$(\mathbf{Disc}_M, =, e, *)$	$(\mathbf{Disc}_M, =, e, *)$ is a symmetric monoidal preorder.
fong-spivak-seven-sketches_1 F00647	55	■ 0.896	$(M, *, e)$	$(M, *, e)$	Exercise 2.8. We said it was easy to check that if $(M, *, e)$ is a commutative monoid
fong-spivak-seven-sketches_1 F00648	55	■ 0.447	$(\mathbf{Disc}_M, =, *, e)$	$(\mathbf{Disc}_M, =, *, e)$	then $(\mathbf{Disc}_M, =, *, e)$ is a symmetric monoidal preorder. Are we telling the truth? $\diamond$
fong-spivak-seven-sketches_1 F00649	55	■ 1.000	H	H	Let H be the set of all poker hands, where a hand means a choice of five cards from the
fong-spivak-seven-sketches_1 F00650	55	■ 1.000	$h \leq h'$	$h \leq h'$	standard 52-card deck. As an order, put $h \leq h'$ if h' beats or equals h in poker.
fong-spivak-seven-sketches_1 F00651	55	■ 1.000	h'	h'	standard 52-card deck. As an order, put $h \leq h'$ if h' beats or equals h in poker.
fong-spivak-seven-sketches_1 F00652	55	■ 1.000	$\otimes: H \times H \rightarrow H$	$\otimes: H \times H \rightarrow H$	One could propose a monoidal product $\otimes: H \times H \rightarrow H$ by assigning $h_1 \otimes h_2$ to be
fong-spivak-seven-sketches_1 F00653	55	■ 1.000	$h_1 \otimes h_2$	$h_1 \otimes h_2$	One could propose a monoidal product $\otimes: H \times H \rightarrow H$ by assigning $h_1 \otimes h_2$ to be
fong-spivak-seven-sketches_1 F00654	55	■ 1.000	h_1	h_1	“the best hand one can form out of the ten cards in h_1 and h_2 .” If some cards are in
fong-spivak-seven-sketches_1 F00655	55	■ 1.000	h_2	h_2	“the best hand one can form out of the ten cards in h_1 and h_2 .” If some cards are in
fong-spivak-seven-sketches_1 F00656	55	■ 0.355	$\{2 \odot, 5 \diamond, 6 \diamond, 6 \star, 7 \star\} = \{2 \diamond, 3 \diamond, 4 \diamond, 5 \diamond, 6 \circlearrowright\}$	$\{2 \odot, 5 \diamond, 6 \diamond, 6 \star, 7 \star\} = \{2 \diamond, 3 \diamond, 4 \diamond, 5 \diamond, 6 \circlearrowright\}$	$\{2 \heartsuit, 5 \heartsuit, 6 \heartsuit, 6 \clubsuit, 7 \spadesuit\} = \{2 \heartsuit, 3 \heartsuit, 4 \heartsuit, 5 \heartsuit, 6 \heartsuit\}$, because the latter is the best hand you
fong-spivak-seven-sketches_1 F00657	55	■ 1.000	$h_1 \leq i_1$	$h_1 \leq i_1$	could be the case that $h_1 \leq i_1$ and $h_2 \leq i_2$, and yet <i>not</i> be the case that $h_1 \otimes h_2 \leq i_1 \otimes i_2$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F00658	55	■ 1.000	$h_{\{2\}} \leq i_{\{2\}}$	$h_2 \leq i_2$	could be the case that $h_1 \leq i_1$ and $h_2 \leq i_2$, and yet <i>not</i> be the case that $h_1 \otimes h_2 \leq i_1 \otimes i_2$.
fong-spivak-seven-sketches F00659	55	■ 1.000	$h_{\{1\}} \otimes h_{\{2\}} \leq i_{\{1\}} \otimes i_{\{2\}}$	$h_1 \otimes h_2 \leq i_1 \otimes i_2$	could be the case that $h_1 \leq i_1$ and $h_2 \leq i_2$, and yet <i>not</i> be the case that $h_1 \otimes h_2 \leq i_1 \otimes i_2$.
fong-spivak-seven-sketches F00660	55	■ 0.925	$h_{\{1\}} \otimes h_{\{2\}} = \{10\star, J\star, Q\star, K\star, A\star, \mathbf{0} \star, \mathbf{K} \star, \mathbf{A} \star\}$	$h_1 \otimes h_2 = \{10\star, J\star, Q\star, K\star, A\star\}$	Here, $h_1 \leq i_1$ and $h_2 \leq i_2$, but $h_1 \otimes h_2 = \{10\clubsuit, 1\clubsuit, 0\clubsuit, K\clubsuit, A\clubsuit\}$ is the best possible
fong-spivak-seven-sketches F00661	55	■ 0.999	$i_{\{1\}} \otimes i_{\{2\}} = \{5\bullet, 5\diamond, 6\diamond, 6\heartsuit, Q\heartsuit\}$	$i_1 \otimes i_2 = \{5\clubsuit, 5\heartsuit, 6\heartsuit, 6\heartsuit, Q\heartsuit\}$	hand and beats $h_1 \otimes h_2 = \{5\clubsuit, 5\heartsuit, 6\heartsuit, 6\heartsuit, Q\heartsuit\}$.
fong-spivak-seven-sketches F00662	55	■ 0.973	$0, 1, 2$	$0, 1, 2$	We can chain some number of these $\leq$ -relationships—say 0, 1, 2, or 3 of them—together
fong-spivak-seven-sketches F00663	56	■ 0.974	$(x \in X)$	$(x \in X)$	$(x \in X)$, relationship ($\leq$), and combination (using transitivity and $\otimes$), and so do wiring
fong-spivak-seven-sketches F00664	56	■ 0.974	$(\leq)$	$(\leq)$	$(x \in X)$, relationship ($\leq$), and combination (using transitivity and $\otimes$), and so do wiring
fong-spivak-seven-sketches F00665	56	■ 1.000	$x \otimes y$	$x \otimes y$	draw one wire labeled $x \otimes y$.
fong-spivak-seven-sketches F00666	57	■ 0.997	$x_{\{1\}} \otimes x_{\{2\}} \leq y_{\{1\}} \otimes y_{\{2\}} \otimes y_{\{3\}}$	$x_1 \otimes x_2 \leq y_1 \otimes y_2 \otimes y_3$	The validity of the second box corresponds to the inequality $x_1 \otimes x_2 \leq y_1 \otimes y_2 \otimes y_3$.
fong-spivak-seven-sketches F00667	57	■ 0.990	$4 \leq 7$	$4 \leq 7$	And here we express a couple facts about $(\mathbb{R}, \leq, 0, +)$ in this language: $4 \leq 7$ and
fong-spivak-seven-sketches F00668	57	■ 1.000	$2+5 \leq -1+5+3$	$2 + 5 \leq -1 + 5 + 3$	$2 + 5 \leq -1 + 5 + 3$.
fong-spivak-seven-sketches F00669	58	■ 1.000	$x_{\{1\}} \otimes x_{\{2\}} \leq$	$x_1 \otimes x_2 \leq$	structure (Definition 2.2). Property (a) says that if $x_1 \leq y_1$ and $x_2 \leq y_2$ then $x_1 \otimes x_2 \leq$
fong-spivak-seven-sketches F00670	58	■ 1.000	$y_{\{1\}} \otimes y_{\{2\}}$	$y_1 \otimes y_2$	$y_1 \otimes y_2$. This corresponds to the idea that stacking any two valid boxes in parallel is
fong-spivak-seven-sketches F00671	59	■ 0.985	$\mathcal{X} = (X, \leq, I, \otimes)$	$\mathcal{X} = (X, \leq, I, \otimes)$	Wiring diagrams as graphical proofs Given a monoidal preorder $\mathcal{X} = (X, \leq, I, \otimes)$, a
fong-spivak-seven-sketches F00672	59	■ 0.381	$\mathcal{X}$	$\mathcal{X}$	wiring diagram is a graphical proof of something about $\mathcal{X}$. Each box in the diagram
fong-spivak-seven-sketches F00673	59	■ 1.000	$v+w+u$	$v + w + u$	technically there is no such thing as $v + w + u$, for example. Instead, there is $(v + w) + u$
fong-spivak-seven-sketches F00674	59	■ 1.000	$(v+w)+u$	$(v + w) + u$	technically there is no such thing as $v + w + u$, for example. Instead, there is $(v + w) + u$
fong-spivak-seven-sketches F00675	59	■ 1.000	$v+(w+u)$	$v + (w + u)$	and $v + (w + u)$, and so on.
fong-spivak-seven-sketches F00676	61	■ 1.000	$2 \mathbf{H}_{\{2\}} + 2 \mathbf{Na}$	$2\text{H}_2\text{O} + 2\text{Na}$	The collection on the left, $2\text{H}_2\text{O} + 2\text{Na}$ is called the <i>reactant</i> , and the collection on the
fong-spivak-seven-sketches F00677	61	■ 1.000	$2 \mathbf{NaOH} + \mathbf{H}_{\{2\}}$	$2\text{NaOH} + \text{H}_2$	right, $2\text{NaOH} + \text{H}_2$ is called the <i>product</i> .
fong-spivak-seven-sketches F00678	61	■ 0.997	$\in$	$\in$	of atoms and molecules, sometimes called <i>materials</i> . So we have $\text{NaCl} \in \text{Mat}$ and
fong-spivak-seven-sketches F00679	61	■ 0.543	$4 \mathbf{H}_{\{2\}} + 6 \mathbf{Ne} \in$	$4\text{H}_2\text{O} + 6\text{Ne} \in$	$4\text{H}_2\text{O} + 6\text{Ne} \in \text{Mat}$.
fong-spivak-seven-sketches F00680	61	■ 1.000	k	k	Here k is the catalyst because it is found both in the reactant and the product of the
fong-spivak-seven-sketches F00681	61	■ 1.000	$x+y \rightarrow z$	$x + y \rightarrow z$	reaction. It is said to catalyze the reaction $x + y \rightarrow z$. The idea is that the reaction
fong-spivak-seven-sketches F00682	61	■ 1.000	$x+y+k \rightarrow z+k$	$x + y + k \rightarrow z + k$	corresponds to the composite reaction $x + y + k \rightarrow z + k$.
fong-spivak-seven-sketches F00683	62	■ 1.000	$x \leq I$	$x \leq I$	(e) $x \leq I$ for all $x \in X$.
fong-spivak-seven-sketches F00684	62	■ 1.000	$x \rightarrow 0$	$x \rightarrow 0$	chemistry section, we would write instead $x \rightarrow 0$: any x yields nothing. But this is


Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches- F00685	63	■ 1.000	10^{23}	10^{23}	manipulate—each item produced. In chemistry, when you have 10^{23} of substance A
fong-spivak-seven-sketches- F00686	64	■ 1.000	$x \leq x+x$	$x \leq x + x$	(f) $x \leq x + x$ for all $x \in X$.
fong-spivak-seven-sketches- F00687	64	■ 0.904	$=0$	$= 0$	false = 0 and true = 1, then $\wedge$ corresponds to the usual multiplication operation $*$.
fong-spivak-seven-sketches- F00688	64	■ 0.904	$=1$	$= 1$	false = 0 and true = 1, then $\wedge$ corresponds to the usual multiplication operation $*$.
fong-spivak-seven-sketches- F00689	64	■ 0.299	$= (\mathbb{B}, \leq$	$= (\mathbb{B}, \leq$	preorder which we denote Bool := $(\mathbb{B}, \leq, \text{true}, \wedge)$.
fong-spivak-seven-sketches- F00690	64	■ 0.299	$\Lambda)$	$\Lambda)$	preorder which we denote Bool := $(\mathbb{B}, \leq, \text{true}, \wedge)$.
fong-spivak-seven-sketches- F00691	64	■ 0.986	$\mathcal{V} = (V, \leq, I, \otimes)$	$\mathcal{V} = (V, \leq, I, \otimes)$	monoidal preorder $\mathcal{V} = (V, \leq, I, \otimes)$ means “letting $\mathcal{V}$ structure the question of getting
fong-spivak-seven-sketches- F00692	64	■ 0.928	$\otimes$	$\otimes$	relation $\leq$, the monoidal unit I , and the monoidal product $\otimes$ —play a role in how
fong-spivak-seven-sketches- F00693	65	■ 0.182	$(\mathbb{B}, \leq$	$(\mathbb{B}, \leq$	For example, let’s look at the case of Bool = $(\mathbb{B}, \leq, \text{true}, \wedge)$. The fact that its
fong-spivak-seven-sketches- F00694	65	■ 0.182	$\wedge)$	$\wedge)$	For example, let’s look at the case of Bool = $(\mathbb{B}, \leq, \text{true}, \wedge)$. The fact that its
fong-spivak-seven-sketches- F00695	65	■ 0.666	$\mathbb{B}, \leq$	$\mathbb{B}, \leq$	<i>Exercise 2.29.</i> Let $(\mathbb{B}, \leq)$ be as above, but now consider the monoidal product to be $\vee$
fong-spivak-seven-sketches- F00696	65	■ 0.991	$(\mathbb{N}, \leq)$	$(\mathbb{N}, \leq)$	preorder $(\mathbb{N}, \leq)$ of natural numbers, where $\leq$ is the usual ordering ($0 \leq 1$ and $5 \leq 16$).
fong-spivak-seven-sketches- F00697	65	■ 0.991	$5 \leq 16$	$5 \leq 16$	preorder $(\mathbb{N}, \leq)$ of natural numbers, where $\leq$ is the usual ordering ($0 \leq 1$ and $5 \leq 16$).
fong-spivak-seven-sketches- F00698	65	■ 0.994	$6+4=10$	$6 + 4 = 10$	where the monoidal unit is 0 and the monoidal product is $+$, i.e. $6 + 4 = 10$. It is easy
fong-spivak-seven-sketches- F00699	65	■ 0.992	$*$	$*$	product is $*$, i.e. $6 * 4 = 24$. What should the monoidal unit be?
fong-spivak-seven-sketches- F00700	65	■ 0.992	$6 * 4=24$	$6 * 4 = 24$	product is $*$, i.e. $6 * 4 = 24$. What should the monoidal unit be?
fong-spivak-seven-sketches- F00701	65	■ 1.000	$m \mid n$	$m \mid n$	a “divisibility” order on $\mathbb{N}$: we write $m n$ to mean that m divides into n without
fong-spivak-seven-sketches- F00702	65	■ 1.000	$1 \mid m$	$1 \mid m$	remainder. So $1 m$ for all m and $4 12$.
fong-spivak-seven-sketches- F00703	65	■ 1.000	$4 \mid 12$	$4 \mid 12$	remainder. So $1 m$ for all m and $4 12$.
fong-spivak-seven-sketches- F00704	65	■ 0.995	$(\mathbb{N},)$	$(\mathbb{N},)$	There is a monoidal structure on $(\mathbb{N},)$, where the monoidal unit is 1 and the
fong-spivak-seven-sketches- F00705	65	■ 0.912	$x_{\{1\}} \mid y_{\{1\}}$	$x_1 \mid y_1$	monoidal product is $*$, i.e. $6 * 4 = 24$. Then if $x_1 y_1$ and $x_2 y_2$, then $(x_1 * x_2) (y_1 * y_2)$.
fong-spivak-seven-sketches- F00706	65	■ 0.912	$x_{\{2\}} \mid y_{\{2\}}$	$x_2 \mid y_2$	monoidal product is $*$, i.e. $6 * 4 = 24$. Then if $x_1 y_1$ and $x_2 y_2$, then $(x_1 * x_2) (y_1 * y_2)$.
fong-spivak-seven-sketches- F00707	65	■ 0.912	$\left(x_{\{1\}} * x_{\{2\}}\right) \mid \left(y_{\{1\}} * y_{\{2\}}\right)$	$(x_1 * x_2) \mid (y_1 * y_2)$	monoidal product is $*$, i.e. $6 * 4 = 24$. Then if $x_1 y_1$ and $x_2 y_2$, then $(x_1 * x_2) (y_1 * y_2)$.
fong-spivak-seven-sketches- F00708	65	■ 1.000	$p_{\{1\}}, p_{\{2\}} \in \mathbb{N}$	$p_1, p_2 \in \mathbb{N}$	Indeed, if there is some $p_1, p_2 \in \mathbb{N}$ such that $x_1 * p_1 = y_1$ and $x_2 * p_2 = y_2$, then
fong-spivak-seven-sketches- F00709	65	■ 1.000	$x_{\{1\}} * p_{\{1\}}=y_{\{1\}}$	$x_1 * p_1 = y_1$	Indeed, if there is some $p_1, p_2 \in \mathbb{N}$ such that $x_1 * p_1 = y_1$ and $x_2 * p_2 = y_2$, then
fong-spivak-seven-sketches- F00710	65	■ 1.000	$x_{\{2\}} * p_{\{2\}}=y_{\{2\}}$	$x_2 * p_2 = y_2$	Indeed, if there is some $p_1, p_2 \in \mathbb{N}$ such that $x_1 * p_1 = y_1$ and $x_2 * p_2 = y_2$, then
fong-spivak-seven-sketches- F00711	65	■ 1.000	$\left(p_{\{1\}} * p_{\{2\}}\right) * \left(x_{\{1\}} * x_{\{2\}}\right) = y_{\{1\}} * y_{\{2\}}$	$(p_1 * p_2) * (x_1 * x_2) = y_1 * y_2$	$p_1 * p_2) * (x_1 * x_2) = y_1 * y_2$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_66_F00712	66	■ 1.000	$\mathrm{P}(S)$	$P(S)$	<i>Exercise 2.35.</i> Let S be a set and let $P(S)$ be its power set, the set of all subsets of
fong-spivak-seven-sketches_66_F00713	66	■ 1.000	$\varnothing \subseteq S$	$\emptyset \subseteq S$	S , including the empty subset, $\emptyset \subseteq S$, and the “everything” subset, $S \subseteq S$. We can
fong-spivak-seven-sketches_66_F00714	66	■ 1.000	$S \subseteq S$	$S \subseteq S$	S , including the empty subset, $\emptyset \subseteq S$, and the “everything” subset, $S \subseteq S$. We can
fong-spivak-seven-sketches_66_F00715	66	■ 1.000	$A \subseteq B$	$A \subseteq B$	give $P(S)$ an order: $A \leq B$ is given by the subset relation $A \subseteq B$, as discussed in
fong-spivak-seven-sketches_66_F00716	66	■ 1.000	$A \cap B$	$A \cap B$	unit S and monoidal product given by intersection $A \cap B$.
fong-spivak-seven-sketches_66_F00717	66	■ 0.902	$\{ \ }^{\mathbb{N}}$	$\mathbb{N}$	<i>Exercise 2.36.</i> Let $\mathrm{Prop}^{\mathbb{N}}$ denote the set of all mathematical statements one can make
fong-spivak-seven-sketches_66_F00718	66	■ 0.901	$\operatorname{Prop}^{\mathbb{N}}$	$\mathrm{Prop}^{\mathbb{N}}$	if and only if the other is true. For example “ n is prime” is an element of $\mathrm{Prop}^{\mathbb{N}}$, and so
fong-spivak-seven-sketches_66_F00719	66	■ 0.993	$n=2$	$n = 2$	are “ $n = 2$ ” and “ $n \geq 11$.” The statements “ $n + 2 = 5$ ” and “ n is the least odd prime”
fong-spivak-seven-sketches_66_F00720	66	■ 0.993	$n \geq 11$	$n \geq 11$	are “ $n = 2$ ” and “ $n \geq 11$.” The statements “ $n + 2 = 5$ ” and “ n is the least odd prime”
fong-spivak-seven-sketches_66_F00721	66	■ 0.993	$n+2=5$	$n + 2 = 5$	are “ $n = 2$ ” and “ $n \geq 11$.” The statements “ $n + 2 = 5$ ” and “ n is the least odd prime”
fong-spivak-seven-sketches_66_F00722	66	■ 0.992	$P, Q \in \operatorname{Prop}^{\mathbb{N}}$	$P, Q \in \mathrm{Prop}^{\mathbb{N}}$	are considered the same. Given $P, Q \in \mathrm{Prop}^{\mathbb{N}}$, we say $P \leq Q$ if for all $n \in \mathbb{N}$, whenever
fong-spivak-seven-sketches_66_F00723	66	■ 0.992	$P \leq Q$	$P \leq Q$	are considered the same. Given $P, Q \in \mathrm{Prop}^{\mathbb{N}}$, we say $P \leq Q$ if for all $n \in \mathbb{N}$, whenever
fong-spivak-seven-sketches_66_F00724	66	■ 1.000	$P(n)$	$P(n)$	$P(n)$ is true, so is $Q(n)$.
fong-spivak-seven-sketches_66_F00725	66	■ 1.000	$Q(n)$	$Q(n)$	$P(n)$ is true, so is $Q(n)$.
fong-spivak-seven-sketches_66_F00726	66	■ 1.000	$[0, \infty]$	$[0, \infty]$	<i>Example 2.37</i> (Lawvere’s monoidal preorder, Cost). Let $[0, \infty]$ denote the set of non-
fong-spivak-seven-sketches_66_F00727	66	■ 0.999	$0, 1, 15.33 \overline{3}$	$0, 1, 15.33\overline{3}$	negative real numbers—such as $0, 1, 15.33\overline{3}$, and 2π —together with ∞ . Consider the
fong-spivak-seven-sketches_66_F00728	66	■ 0.999	2π	2π	negative real numbers—such as $0, 1, 15.33\overline{3}$, and 2π —together with ∞ . Consider the
fong-spivak-seven-sketches_66_F00729	66	■ 0.999	∞	∞	negative real numbers—such as $0, 1, 15.33\overline{3}$, and 2π —together with ∞ . Consider the
fong-spivak-seven-sketches_66_F00730	66	■ 0.725	$([0, \infty], \geq)$	$([0, \infty], \geq)$	preorder $([0, \infty], \geq)$, with the usual notion of $\geq$, where of course $\infty \geq x$ for all $x \in [0, \infty]$.
fong-spivak-seven-sketches_66_F00731	66	■ 0.725	$\infty \geq x$	$\infty \geq x$	preorder $([0, \infty], \geq)$, with the usual notion of $\geq$, where of course $\infty \geq x$ for all $x \in [0, \infty]$.
fong-spivak-seven-sketches_66_F00732	66	■ 0.725	$x \in [0, \infty]$	$x \in [0, \infty]$	preorder $([0, \infty], \geq)$, with the usual notion of $\geq$, where of course $\infty \geq x$ for all $x \in [0, \infty]$.
fong-spivak-seven-sketches_66_F00733	66	■ 0.983	$x + \infty = \infty$	$x + \infty = \infty$	the monoidal product is $+$. In particular, $x + \infty = \infty$ for any $x \in [0, \infty]$. Let’s call this
fong-spivak-seven-sketches_67_F00734	67	■ 0.999	$(X, \leq)^{\mathrm{op}} := (X, \geq)$	$(X, \leq)^{\mathrm{op}} := (X, \geq)$	flip the order: $(X, \leq)^{\mathrm{op}} := (X, \geq)$; see Example 1.58. Proposition 2.38 says that if the
fong-spivak-seven-sketches_67_F00735	67	■ 0.955	$\mathcal{X} = (X, \leq)$	$\mathcal{X} = (X, \leq)$	Proposition 2.38. Suppose $\mathcal{X} = (X, \leq)$ is a preorder and $\mathcal{X}^{\mathrm{op}} = (X, \geq)$ is its opposite. If
fong-spivak-seven-sketches_67_F00736	67	■ 0.955	$\mathcal{X}^{\mathrm{op}} = (X, \geq)$	$\mathcal{X}^{\mathrm{op}} = (X, \geq)$	Proposition 2.38. Suppose $\mathcal{X} = (X, \leq)$ is a preorder and $\mathcal{X}^{\mathrm{op}} = (X, \geq)$ is its opposite. If
fong-spivak-seven-sketches_67_F00737	67	■ 0.930	$(X, \leq I, \otimes)$	$(X, \leq I, \otimes)$	$(X, \leq, I, \otimes)$ is a symmetric monoidal preorder then so is its opposite, $(X, \geq, I, \otimes)$.
fong-spivak-seven-sketches_67_F00738	67	■ 0.930	$(X, \geq I, \otimes)$	$(X, \geq, I, \otimes)$	$(X, \leq, I, \otimes)$ is a symmetric monoidal preorder then so is its opposite, $(X, \geq, I, \otimes)$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F00739	67	0.838	$x_{\{1\}} \geq y_{\{1\}}$	$x_1 \geq y_1$	<i>Proof.</i> Let's first check monotonicity. Suppose $x_1 \geq y_1$ and $x_2 \geq y_2$ in $\mathcal{X}^{\text{op}}$; we need to
fong-spivak-seven-sketches F00740	67	0.838	$x_{\{2\}} \geq y_{\{2\}}$	$x_2 \geq y_2$	<i>Proof.</i> Let's first check monotonicity. Suppose $x_1 \geq y_1$ and $x_2 \geq y_2$ in $\mathcal{X}^{\text{op}}$; we need to
fong-spivak-seven-sketches F00741	67	0.838	$X^{\{\text{op}\}}$	X^{op}	<i>Proof.</i> Let's first check monotonicity. Suppose $x_1 \geq y_1$ and $x_2 \geq y_2$ in $\mathcal{X}^{\text{op}}$; we need to
fong-spivak-seven-sketches F00742	67	0.999	$x_{\{1\}} \otimes x_{\{2\}} \geq y_{\{1\}} \otimes y_{\{2\}}$	$x_1 \otimes x_2 \geq y_1 \otimes y_2$	show that $x_1 \otimes x_2 \geq y_1 \otimes y_2$. But by definition of opposite order, we have $y_1 \leq x_1$ and
fong-spivak-seven-sketches F00743	67	0.999	$y_{\{1\}} \leq x_{\{1\}}$	$y_1 \leq x_1$	show that $x_1 \otimes x_2 \geq y_1 \otimes y_2$. But by definition of opposite order, we have $y_1 \leq x_1$ and
fong-spivak-seven-sketches F00744	67	0.558	$y_{\{2\}} \leq x_{\{2\}}$	$y_2 \leq x_2$	$y_2 \leq x_2$ in $\mathcal{X}$, and thus $y_1 \otimes y_2 \leq x_1 \otimes x_2$ in $\mathcal{X}$. Thus indeed $x_1 \otimes x_2 \geq y_1 \otimes y_2$ in $\mathcal{X}^{\text{op}}$.
fong-spivak-seven-sketches F00745	67	0.558	$y_{\{1\}} \otimes y_{\{2\}} \leq x_{\{1\}} \otimes x_{\{2\}}$	$y_1 \otimes y_2 \leq x_1 \otimes x_2$	$y_2 \leq x_2$ in $\mathcal{X}$, and thus $y_1 \otimes y_2 \leq x_1 \otimes x_2$ in $\mathcal{X}$. Thus indeed $x_1 \otimes x_2 \geq y_1 \otimes y_2$ in $\mathcal{X}^{\text{op}}$.
fong-spivak-seven-sketches F00746	67	0.996	$\{\}^{\{\mathrm{op}\}}$	op	Cost ^{op} is too.
fong-spivak-seven-sketches F00747	67	1.000	$x \cong x^{\{\prime\}}$	$x \cong x'$	relation $\cong$ on X , where $x \cong x'$ iff both $x \leq x'$ and $x' \leq x$.
fong-spivak-seven-sketches F00748	67	1.000	$x \leq x^{\{\prime\}}$	$x \leq x'$	relation $\cong$ on X , where $x \cong x'$ iff both $x \leq x'$ and $x' \leq x$.
fong-spivak-seven-sketches F00749	67	1.000	$x^{\{\prime\}} \leq x$	$x' \leq x$	relation $\cong$ on X , where $x \cong x'$ iff both $x \leq x'$ and $x' \leq x$.
fong-spivak-seven-sketches F00750	67	1.000	$\mathcal{P} = \left(P, \leq_P, I_P, \otimes_P \right)$	$\mathcal{P} = (P, \leq_P, I_P, \otimes_P)$	Definition 2.41. Let $\mathcal{P} = (P, \leq_P, I_P, \otimes_P)$ and $\mathcal{Q} = (Q, \leq_Q, I_Q, \otimes_Q)$ be monoidal preorders.
fong-spivak-seven-sketches F00751	67	1.000	$\mathcal{Q} = \left(Q, \leq_Q, I_Q, \otimes_Q \right)$	$\mathcal{Q} = (Q, \leq_Q, I_Q, \otimes_Q)$	Definition 2.41. Let $\mathcal{P} = (P, \leq_P, I_P, \otimes_P)$ and $\mathcal{Q} = (Q, \leq_Q, I_Q, \otimes_Q)$ be monoidal preorders.
fong-spivak-seven-sketches F00752	67	0.995	$\mathcal{P}$	$\mathcal{P}$	A monoidal monotone from $\mathcal{P}$ to $\mathcal{Q}$ is a monotone map $f: (P, \leq_P) \rightarrow (Q, \leq_Q)$, satisfying
fong-spivak-seven-sketches F00753	67	1.000	$I_{\{0\}} \leq_{\{0\}} f(I_{\{P\}})$	$I_Q \leq_Q f(I_P)$	(a) $I_Q \leq_Q f(I_P)$, and
fong-spivak-seven-sketches F00754	67	1.000	$f(p_{\{1\}}) \otimes_{\{0\}} f(p_{\{2\}}) \leq_{\{0\}} f(p_{\{1\}} \otimes_{\{P\}} p_{\{2\}})$	$f(p_1) \otimes_Q f(p_2) \leq_Q f(p_1 \otimes_P p_2)$	(b) $f(p_1) \otimes_Q f(p_2) \leq_Q f(p_1 \otimes_P p_2)$
fong-spivak-seven-sketches F00755	67	1.000	$p_{\{1\}}, p_{\{2\}} \in P$	$p_1, p_2 \in P$	for all $p_1, p_2 \in P$.
fong-spivak-seven-sketches F00756	68	1.000	$I_{\{0\}} \cong f(I_{\{P\}})$	$I_Q \cong f(I_P)$	(a') $I_Q \cong f(I_P)$, and
fong-spivak-seven-sketches F00757	68	1.000	$f(p_{\{1\}}) \otimes_{\{0\}} f(p_{\{2\}}) \cong f(p_{\{1\}} \otimes_{\{P\}} p_{\{2\}})$	$f(p_1) \otimes_Q f(p_2) \cong f(p_1 \otimes_P p_2)$	(b') $f(p_1) \otimes_Q f(p_2) \cong f(p_1 \otimes_P p_2)$;
fong-spivak-seven-sketches F00758	68	0.895	$I_{\{0\}} = f(I_{\{P\}})$	$I_Q = f(I_P)$	(a'') $I_Q = f(I_P)$, and
fong-spivak-seven-sketches F00759	68	0.995	$f(p_{\{1\}}) \otimes_{\{0\}} f(p_{\{2\}}) = f(p_{\{1\}} \otimes_{\{P\}} p_{\{2\}})$	$f(p_1) \otimes_Q f(p_2) = f(p_1 \otimes_P p_2)$	(b'') $f(p_1) \otimes_Q f(p_2) = f(p_1 \otimes_P p_2)$.
fong-spivak-seven-sketches F00760	68	1.000	$i: (\mathbb{N}, \leq, 0, +) \rightarrow (\mathbb{R}, \leq, 0, +)$	$i: (\mathbb{N}, \leq, 0, +) \rightarrow (\mathbb{R}, \leq, 0, +)$	<i>Example 2.42.</i> There is a monoidal monotone $i: (\mathbb{N}, \leq, 0, +) \rightarrow (\mathbb{R}, \leq, 0, +)$, where $i(n) =$
fong-spivak-seven-sketches F00761	68	1.000	$i(n) =$	$i(n) =$	<i>Example 2.42.</i> There is a monoidal monotone $i: (\mathbb{N}, \leq, 0, +) \rightarrow (\mathbb{R}, \leq, 0, +)$, where $i(n) =$
fong-spivak-seven-sketches F00762	68	1.000	$i(m) \leq i(n)$	$i(m) \leq i(n)$	n for all $n \in \mathbb{N}$. It is clearly monotonic, $m \leq n$ implies $i(m) \leq i(n)$. It is even strict
fong-spivak-seven-sketches F00763	68	1.000	$i(0) = 0$	$i(0) = 0$	monoidal because $i(0) = 0$ and $i(m + n) = i(m) + i(n)$.
fong-spivak-seven-sketches F00764	68	1.000	$i(m+n) = i(m) + i(n)$	$i(m + n) = i(m) + i(n)$	monoidal because $i(0) = 0$ and $i(m + n) = i(m) + i(n)$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F00765	68	0.998	$f: (\mathbb{R}, \leq, 0, +) \rightarrow (\mathbb{N}, \leq, 0, +)$	$f: (\mathbb{R}, \leq, 0, +) \rightarrow (\mathbb{N}, \leq, 0, +)$	There is also a monoidal monotone $f: (\mathbb{R}, \leq, 0, +) \rightarrow (\mathbb{N}, \leq, 0, +)$ going the other
fong-spivak-seven-sketches F00766	68	1.000	$f(x) := \lfloor x \rfloor$	$f(x) := \lfloor x \rfloor$	way. Here $f(x) := \lfloor x \rfloor$ is the floor function, e.g. $f(3.14) = 3$. It is monotonic because
fong-spivak-seven-sketches F00767	68	1.000	$f(3.14) = 3$	$f(3.14) = 3$	way. Here $f(x) := \lfloor x \rfloor$ is the floor function, e.g. $f(3.14) = 3$. It is monotonic because
fong-spivak-seven-sketches F00768	68	1.000	$f(0) = 0$	$f(0) = 0$	$x \leq y$ implies $f(x) \leq f(y)$. Also $f(0) = 0$ and $f(x) + f(y) \leq f(x + y)$, so it is a monoidal
fong-spivak-seven-sketches F00769	68	1.000	$f(x) + f(y) \leq f(x + y)$	$f(x) + f(y) \leq f(x + y)$	$x \leq y$ implies $f(x) \leq f(y)$. Also $f(0) = 0$ and $f(x) + f(y) \leq f(x + y)$, so it is a monoidal
fong-spivak-seven-sketches F00770	68	1.000	$f(0.5) + f(0.5) \neq f(0.5 + 0.5)$	$f(0.5) + f(0.5) \neq f(0.5 + 0.5)$	monotone. But it is not strict or even strong because $f(0.5) + f(0.5) \neq f(0.5 + 0.5)$.
fong-spivak-seven-sketches F00771	68	0.754	$= ([0, \infty], \geq, 0, +)$	$= ([0, \infty], \geq, 0, +)$	Recall Bool = $(\mathbb{B}, \leq, \text{true}, \wedge)$ from Example 2.27 and Cost = $([0, \infty], \geq, 0, +)$ from
fong-spivak-seven-sketches F00772	68	0.905	$:= \infty$	$:= \infty$	Example 2.37. There is a monoidal monotone $g: \mathbf{Bool} \rightarrow \mathbf{Cost}$, given by $g(\text{false}) := \infty$
fong-spivak-seven-sketches F00773	68	0.998	$:= 0$	$:= 0$	and $g(\text{true}) := 0$.
fong-spivak-seven-sketches F00774	68	0.683	$g: (\mathbb{B}, \leq$	$g: (\mathbb{B}, \leq$	1. Check that the map $g: (\mathbb{B}, \leq, \text{true}, \wedge) \rightarrow ([0, \infty], \geq, 0, +)$ presented above indeed
fong-spivak-seven-sketches F00775	68	0.683	$\wedge) \rightarrow ([0, \infty], \geq, 0, +)$	$\wedge) \rightarrow ([0, \infty], \geq, 0, +)$	1. Check that the map $g: (\mathbb{B}, \leq, \text{true}, \wedge) \rightarrow ([0, \infty], \geq, 0, +)$ presented above indeed
fong-spivak-seven-sketches F00776	68	1.000	$d, u: [0, \infty] \rightarrow \mathbb{B}$	$d, u: [0, \infty] \rightarrow \mathbb{B}$	functions $d, u: [0, \infty] \rightarrow \mathbb{B}$ defined as follows:
fong-spivak-seven-sketches F00777	68	1.000	u	u	4. Is u monotonic?
fong-spivak-seven-sketches F00778	69	0.929	$(\mathbb{N}, \leq, 1, *)$	$(\mathbb{N}, \leq, 1, *)$	1. Is $(\mathbb{N}, \leq, 1, *)$ a monoidal preorder, where $*$ is the usual multiplication of natural
fong-spivak-seven-sketches F00779	69	0.998	$(\mathbb{N}, \leq, 0, +) \rightarrow (\mathbb{N}, \leq$	$(\mathbb{N}, \leq, 0, +) \rightarrow (\mathbb{N}, \leq$	2. If not, why not? If so, does there exist a monoidal monotone $(\mathbb{N}, \leq, 0, +) \rightarrow (\mathbb{N}, \leq$
fong-spivak-seven-sketches F00780	69	0.833	$, 1, *)$	$, 1, *)$	$, 1, *)$? If not, why not? If so, find it.
fong-spivak-seven-sketches F00781	69	0.935	$(\mathbb{Z}, \leq, *, 1)$	$(\mathbb{Z}, \leq, *, 1)$	3. Is $(\mathbb{Z}, \leq, *, 1)$ a monoidal preorder?
fong-spivak-seven-sketches F00782	69	0.949	$\operatorname{Ob}(\mathcal{X})$	$\operatorname{Ob}(\mathcal{X})$	(i) one specifies a set $\operatorname{Ob}(\mathcal{X})$, elements of which are called <i>objects</i> ;
fong-spivak-seven-sketches F00783	69	1.000	$\mathcal{X}(x, y) \in V$	$\mathcal{X}(x, y) \in V$	(ii) for every two objects x, y , one specifies an element $\mathcal{X}(x, y) \in V$, called the <i>hom-</i>
fong-spivak-seven-sketches F00784	69	0.631	$x \in \operatorname{Ob}(X)$	$x \in \operatorname{Ob}(X)$	(a) for every object $x \in \operatorname{Ob}(\mathcal{X})$ we have $I \leq \mathcal{X}(x, x)$, and
fong-spivak-seven-sketches F00785	69	0.631	$I \leq \mathcal{X}(x, x)$	$I \leq \mathcal{X}(x, x)$	(a) for every object $x \in \operatorname{Ob}(\mathcal{X})$ we have $I \leq \mathcal{X}(x, x)$, and
fong-spivak-seven-sketches F00786	69	0.416	$x, y, z \in \operatorname{Ob}(X)$	$x, y, z \in \operatorname{Ob}(X)$	(b) for every three objects $x, y, z \in \operatorname{Ob}(\mathcal{X})$ we have $\mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \leq \mathcal{X}(x, z)$.
fong-spivak-seven-sketches F00787	69	0.416	$\mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \leq \mathcal{X}(x, z)$	$\mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \leq \mathcal{X}(x, z)$	(b) for every three objects $x, y, z \in \operatorname{Ob}(\mathcal{X})$, we have $\mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \leq \mathcal{X}(x, z)$.
fong-spivak-seven-sketches F00788	69	0.999	$\mathcal{X}(x, y)$	$\mathcal{X}(x, y)$	name for $\mathcal{X}(x, y)$ might be <i>mapping object</i> , but we use “hom” mainly because it is an important jargon word
fong-spivak-seven-sketches F00789	70	0.986	$\operatorname{Ob}(\mathcal{X}) = \{p, q, r, s, t\}$	$\operatorname{Ob}(\mathcal{X}) = \{p, q, r, s, t\}$	the elements of the preorder, i.e. $\operatorname{Ob}(\mathcal{X}) = \{p, q, r, s, t\}$. Next, for every pair of objects
fong-spivak-seven-sketches F00790	70	1.000	$t \not\leq s$	$t \not\leq s$	if otherwise. So for example, since $s \leq t$ and $t \not\leq s$, we have $\mathcal{X}(s, t) = \text{true}$ and
fong-spivak-seven-sketches F00791	70	1.000	$\mathcal{X}(s, t) =$	$\mathcal{X}(s, t) =$	if otherwise. So for example, since $s \leq t$ and $t \not\leq s$, we have $\mathcal{X}(s, t) = \text{true}$ and

Identifier	Page	Conf.	LaTeX source	Rendered	Image																																				
fong-spivak-seven-sketches_70 F00792	70	■ 0.989	$X(t, s) =$	$X(t, s) =$	$\mathcal{X}(t, s) = \text{false}$. Recalling from Example 2.27 that the monoidal unit I of Bool is true ,																																				
fong-spivak-seven-sketches_70 F00793	70	■ 0.598	$x(x, y)$	$x(x, y)$	(x, y) th entry is simply the hom-object $\mathcal{X}(x, y)$. So, for example, the above preorder in																																				
fong-spivak-seven-sketches_70 F00794	70	■ 0.598	r	r	<table border="1"> <thead> <tr> <th>$\mathcal{X}$</th><th>p</th><th>q</th><th>r</th><th>s</th><th>t</th></tr> </thead> <tbody> <tr> <th>p</th><td>true</td><td>true</td><td>true</td><td>true</td><td>true</td></tr> <tr> <th>q</th><td>false</td><td>true</td><td>false</td><td>true</td><td>true</td></tr> <tr> <th>r</th><td>false</td><td>false</td><td>true</td><td>true</td><td>true</td></tr> <tr> <th>s</th><td>false</td><td>false</td><td>false</td><td>true</td><td>true</td></tr> <tr> <th>t</th><td>false</td><td>false</td><td>false</td><td>false</td><td>true</td></tr> </tbody> </table>	$\mathcal{X}$	p	q	r	s	t	p	true	true	true	true	true	q	false	true	false	true	true	r	false	false	true	true	true	s	false	false	false	true	true	t	false	false	false	false	true
$\mathcal{X}$	p	q	r	s	t																																				
p	true	true	true	true	true																																				
q	false	true	false	true	true																																				
r	false	false	true	true	true																																				
s	false	false	false	true	true																																				
t	false	false	false	false	true																																				
fong-spivak-seven-sketches_71 F00795	71	■ 0.503	$x, y \in \mathrm{Ob}(X)$	$x, y \in \mathrm{Ob}(X)$	(ii) for every $x, y \in \mathrm{Ob}(\mathcal{X})$ an element $\mathcal{X}(x, y) \in \mathbb{B}$, i.e. either $\mathcal{X}(x, y) = \text{true}$ or																																				
fong-spivak-seven-sketches_71 F00796	71	■ 0.503	$\mathcal{X}(x, y) \in \mathbb{B}$	$\mathcal{X}(x, y) \in \mathbb{B}$	(ii) for every $x, y \in \mathrm{Ob}(\mathcal{X})$ an element $\mathcal{X}(x, y) \in \mathbb{B}$, i.e. either $\mathcal{X}(x, y) = \text{true}$ or																																				
fong-spivak-seven-sketches_71 F00797	71	■ 0.503	$\mathcal{X}(x, y) =$	$\mathcal{X}(x, y) =$	(ii) for every $x, y \in \mathrm{Ob}(\mathcal{X})$ an element $\mathcal{X}(x, y) \in \mathbb{B}$, i.e. either $\mathcal{X}(x, y) = \text{true}$ or																																				
fong-spivak-seven-sketches_71 F00798	71	■ 0.958	$X(x, y) =$	$X(x, y) =$	$\mathcal{X}(x, y) = \text{false}$.																																				
fong-spivak-seven-sketches_71 F00799	71	■ 0.507	$X := \operatorname{Ob}(X)$	$X := \mathrm{Ob}(X)$	objects of $\mathcal{X}$. So let's call the preorder $(X, \leq)$, and let $X := \mathrm{Ob}(\mathcal{X})$. For the $\leq$ relation,																																				
fong-spivak-seven-sketches_71 F00800	71	■ 0.952	$\leq \mathcal{X}(x, x)$	$\leq \mathcal{X}(x, x)$	(a) for every element $x \in X$ we have $\text{true} \leq \mathcal{X}(x, x)$,																																				
fong-spivak-seven-sketches_71 F00801	71	■ 0.798	$\mathcal{X}(x, y) \wedge \mathcal{X}(y, z) \leq \mathcal{X}(x, z)$	$\mathcal{X}(x, y) \wedge \mathcal{X}(y, z) \leq \mathcal{X}(x, z)$	(b) for every three elements $x, y, z \in X$ we have $\mathcal{X}(x, y) \wedge \mathcal{X}(y, z) \leq \mathcal{X}(x, z)$.																																				
fong-spivak-seven-sketches_71 F00802	71	■ 0.998	$b \in$	$b \in$	For $b \in \mathbf{Bool}$, if $\text{true} \leq b$ then $b = \text{true}$, so the first statement says $\mathcal{X}(x, x) = \text{true}$,																																				
fong-spivak-seven-sketches_71 F00803	71	■ 0.998	$\leq b$	$\leq b$	For $b \in \mathbf{Bool}$, if $\text{true} \leq b$ then $b = \text{true}$, so the first statement says $\mathcal{X}(x, x) = \text{true}$,																																				
fong-spivak-seven-sketches_71 F00804	71	■ 0.998	$b =$	$b =$	For $b \in \mathbf{Bool}$, if $\text{true} \leq b$ then $b = \text{true}$, so the first statement says $\mathcal{X}(x, x) = \text{true}$,																																				
fong-spivak-seven-sketches_71 F00805	71	■ 0.998	$\mathcal{X}(x, x) =$	$\mathcal{X}(x, x) =$	For $b \in \mathbf{Bool}$, if $\text{true} \leq b$ then $b = \text{true}$, so the first statement says $\mathcal{X}(x, x) = \text{true}$,																																				
fong-spivak-seven-sketches_71 F00806	71	■ 0.691	$b \in \mathbb{B}$	$b \in \mathbb{B}$	$\text{false} \leq b$ for all $b \in \mathbb{B}$, the only way statement (b) has any force is if $\mathcal{X}(x, y) = \text{true}$																																				
fong-spivak-seven-sketches_71 F00807	71	■ 0.943	$\mathcal{X}(y, z) =$	$\mathcal{X}(y, z) =$	and $\mathcal{X}(y, z) = \text{true}$, in which case it forces $\mathcal{X}(x, z) = \text{true}$. This condition exactly																																				
fong-spivak-seven-sketches_71 F00808	71	■ 0.943	$\mathcal{X}(x, z) =$	$\mathcal{X}(x, z) =$	and $\mathcal{X}(y, z) = \text{true}$, in which case it forces $\mathcal{X}(x, z) = \text{true}$. This condition exactly																																				
fong-spivak-seven-sketches_71 F00809	71	■ 0.766	(X, d)	(X, d)	Definition 2.51. A metric space (X, d) consists of:																																				
fong-spivak-seven-sketches_71 F00810	71	■ 1.000	$d: X \times X \rightarrow \mathbb{R}_{\geq 0}$	$d: X \times X \rightarrow \mathbb{R}_{\geq 0}$	(ii) a function $d: X \times X \rightarrow \mathbb{R}_{\geq 0}$, where $d(x, y)$ is called the <i>distance between x and y</i> .																																				
fong-spivak-seven-sketches_71 F00811	71	■ 1.000	$d(x, y)$	$d(x, y)$	(ii) a function $d: X \times X \rightarrow \mathbb{R}_{\geq 0}$, where $d(x, y)$ is called the <i>distance between x and y</i> .																																				
fong-spivak-seven-sketches_72 F00812	72	■ 1.000	$d(x, x) = 0$	$d(x, x) = 0$	(a) for every $x \in X$, we have $d(x, x) = 0$,																																				
fong-spivak-seven-sketches_72 F00813	72	■ 1.000	$d(x, y) = 0$	$d(x, y) = 0$	(b) for every $x, y \in X$, if $d(x, y) = 0$ then $x = y$,																																				
fong-spivak-seven-sketches_72 F00814	72	■ 1.000	$d(x, y) = d(y, x)$	$d(x, y) = d(y, x)$	(c) for every $x, y \in X$, we have $d(x, y) = d(y, x)$, and																																				
fong-spivak-seven-sketches_72 F00815	72	■ 1.000	$d(x, y) + d(y, z) \geq d(x, z)$	$d(x, y) + d(y, z) \geq d(x, z)$	(d) for every $x, y, z \in X$, we have $d(x, y) + d(y, z) \geq d(x, z)$.																																				

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_72 F00816	72	0.772	$d: X \times X \rightarrow [0, \infty] = \mathbb{R}_{\geq 0} \cup \{\infty\}$	$d: X \times X \rightarrow [0, \infty] = \mathbb{R}_{\geq 0} \cup \{\infty\}$	If we ask instead in (ii) for a function $d: X \times X \rightarrow [0, \infty] = \mathbb{R}_{\geq 0} \cup \{\infty\}$, we call (X, d)
fong-spivak-seven-sketches_72 F00817	72	0.772	X, d	X, d	If we ask instead in (ii) for a function $d: X \times X \rightarrow [0, \infty] = \mathbb{R}_{\geq 0} \cup \{\infty\}$, we call (X, d)
fong-spivak-seven-sketches_72 F00818	72	1.000	z	z	The triangle inequality says that when plotting a route from x to z , the distance is
fong-spivak-seven-sketches_72 F00819	72	1.000	$x \rightarrow y \rightarrow z$	$x \rightarrow y \rightarrow z$	always at most what you get by choosing an intermediate point y and going  .
fong-spivak-seven-sketches_72 F00820	72	1.000	$3+5 \geq 7.2$	$3 + 5 \geq 7.2$	It can be invoked three different ways in the above picture: $3 + 5 \geq 7.2$, but also
fong-spivak-seven-sketches_72 F00821	72	1.000	$5+7.2 \geq 3$	$5 + 7.2 \geq 3$	$5 + 7.2 \geq 3$ and $3 + 7.2 \geq 5$. Oh yeah, and $5 + 3 \geq 7.2$, $7.2 + 5 \geq 3$ and $7.2 + 3 \geq 5$.
fong-spivak-seven-sketches_72 F00822	72	1.000	$3+7.2 \geq 5$	$3 + 7.2 \geq 5$	$5 + 7.2 \geq 3$ and $3 + 7.2 \geq 5$. Oh yeah, and $5 + 3 \geq 7.2$, $7.2 + 5 \geq 3$ and $7.2 + 3 \geq 5$.
fong-spivak-seven-sketches_72 F00823	72	1.000	$5+3 \geq 7.2, 7.2+5 \geq 3$	$5 + 3 \geq 7.2, 7.2 + 5 \geq 3$	$5 + 7.2 \geq 3$ and $3 + 7.2 \geq 5$. Oh yeah, and $5 + 3 \geq 7.2$, $7.2 + 5 \geq 3$ and $7.2 + 3 \geq 5$.
fong-spivak-seven-sketches_72 F00824	72	1.000	$7.2+3 \geq 5$	$7.2 + 3 \geq 5$	$5 + 7.2 \geq 3$ and $3 + 7.2 \geq 5$. Oh yeah, and $5 + 3 \geq 7.2$, $7.2 + 5 \geq 3$ and $7.2 + 3 \geq 5$.
fong-spivak-seven-sketches_72 F00825	72	0.764	$d(x, y) = ? d(y, x)$	$d(x, y) = ? d(y, x)$	are any hills, the symmetry axiom, $d(x, y) = ? d(y, x)$, fails: it's easier to get from x
fong-spivak-seven-sketches_72 F00826	72	0.689	$d(\mathrm{US})$	$d(\mathrm{US}$	distance in the worst-case scenario?" So $d(\mathrm{US}, \mathrm{Spain})$ is the distance from somewhere
fong-spivak-seven-sketches_72 F00827	72	0.689	)	)	distance in the worst-case scenario?" So $d(\mathrm{US}, \mathrm{Spain})$ is the distance from somewhere
fong-spivak-seven-sketches_72 F00828	72	0.998	$d($	$d($	It also violates another condition, because $d(\mathrm{Boston}, \mathrm{US}) = 0$. No matter where you
fong-spivak-seven-sketches_72 F00829	72	0.998	) = 0	) = 0	It also violates another condition, because $d(\mathrm{Boston}, \mathrm{US}) = 0$. No matter where you
fong-spivak-seven-sketches_72 F00830	72	0.852	$U \subseteq X$	$U \subseteq X$	³ The Hausdorff distance gives a metric on the set of all subsets $U \subseteq X$ of a given metric space (X, d) .
fong-spivak-seven-sketches_73 F00831	73	0.547	) $\neq 0$	) $\neq 0$	$d(\mathrm{US}, \mathrm{Boston}) \neq 0$.
fong-spivak-seven-sketches_73 F00832	73	0.867	$\{r \in \mathbb{R} \mid r < 0\}$	$\{r \in \mathbb{R} \mid r < 0\}$	between two regions is often infinite, e.g. the distance between  and $\{0\}$
fong-spivak-seven-sketches_73 F00833	73	0.996	$(\mathbb{R}, d)$	$(\mathbb{R}, d)$	as subsets of $(\mathbb{R}, d)$ is infinite.
fong-spivak-seven-sketches_73 F00834	73	0.899	$x, y \in \mathrm{Ob}(\mathcal{X})$	$x, y \in \mathrm{Ob}(\mathcal{X})$	(ii) for every $x, y \in \mathrm{Ob}(\mathcal{X})$ an element $\mathcal{X}(x, y) \in [0, \infty]$.
fong-spivak-seven-sketches_73 F00835	73	0.899	$\mathcal{X}(x, y) \in [0, \infty]$	$\mathcal{X}(x, y) \in [0, \infty]$	(ii) for every $x, y \in \mathrm{Ob}(\mathcal{X})$ an element $\mathcal{X}(x, y) \in [0, \infty]$.
fong-spivak-seven-sketches_73 F00836	73	1.000	$0 \geq d(x, x)$	$0 \geq d(x, x)$	(a) $0 \geq d(x, x)$ for all $x \in X$, and
fong-spivak-seven-sketches_73 F00837	73	1.000	$d(x, x) \in [0, \infty]$	$d(x, x) \in [0, \infty]$	Since $d(x, x) \in [0, \infty]$, if $0 \geq d(x, x)$ then $d(x, x) = 0$. So the first condition is equivalent
fong-spivak-seven-sketches_73 F00838	73	1.000	$d(x, y) := y - x $	$d(x, y) := y - x $	hence a Lawvere metric space structure. Namely  , the absolute value
fong-spivak-seven-sketches_73 F00839	73	1.000	$d(3, 7) = 4$	$d(3, 7) = 4$	of the difference. So $d(3, 7) = 4$.
fong-spivak-seven-sketches_73 F00840	73	0.214	$\mathbb{R}_{\geq 0, \geq, 0, +}$	$\mathbb{R}_{\geq 0, \geq, 0, +}$	Exercise 2.55. Consider the symmetric monoidal preorder $(\mathbb{R}_{\geq 0}, \geq, 0, +)$, which is
fong-spivak-seven-sketches_73 F00841	73	0.824	$\left(\mathbb{R}_{\geq 0, \geq, 0, +}\right)$	$(\mathbb{R}_{\geq 0}, \geq, 0, +)$	the difference between a Lawvere metric space and a $(\mathbb{R}_{\geq 0}, \geq, 0, +)$ -category in the sense
fong-spivak-seven-sketches_73 F00842	73	1.000	$d(U, V) := \max(d_L(U, V), d_L(V, U))$	$d(U, V) := \max(d_L(U, V), d_L(V, U))$	formula and symmetrize it: $d(U, V) := \max(d_L(U, V), d_L(V, U))$. We happen to see the unsymmetrized

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_74 F00843	74	■ 1.000	$w \geq 0$	$w \geq 0$	are labeled with numbers $w \geq 0$ —into a Lawvere metric space. In fact, we shall
fong-spivak-seven-sketches_74 F00844	74	■ 0.985	$d_{\{X\}}$	d_X	Given a weighted graph, one forms a metric d_X on its set X of vertices by setting $d(p, q)$
fong-spivak-seven-sketches_74 F00845	74	■ 0.985	$d(p, q)$	$d(p, q)$	Given a weighted graph, one forms a metric d_X on its set X of vertices by setting $d(p, q)$
fong-spivak-seven-sketches_74 F00846	74	■ 0.669	$d\left(\nearrow^{\prime}\right)$	$d(\nearrow')$	$\begin{array}{c ccc} d(\nearrow) & x & y & z \\ \hline x & 0 & 4 & 3 \\ y & 3 & 0 & 6 \\ z & 7 & 4 & 0 \end{array}$
fong-spivak-seven-sketches_74 F00847	74	■ 0.999	$M_{\{G\}}$	M_G	taking G and getting a square matrix M_G , whose rows and columns are indexed by the
fong-spivak-seven-sketches_75 F00848	75	■ 0.577	$M_{\{Y\}} :=$	$M_Y :=$	$M_Y := \begin{array}{c ccc} \nearrow & x & y & z \\ \hline x & 0 & 4 & 3 \\ y & 3 & 0 & \infty \\ z & \infty & 4 & 0 \end{array}$
fong-spivak-seven-sketches_75 F00849	75	■ 0.998	$d_{\{Y\}}$	d_Y	that the more difficult “distance matrices” d_Y , such as Eq. (2.57), can be obtained from
fong-spivak-seven-sketches_75 F00850	75	■ 0.992	$M_{\{Y\}}$	M_Y	the easy graph matrices M_Y , such as Eq. (2.59), by repeating a certain sort of “matrix
fong-spivak-seven-sketches_75 F00851	75	■ 0.699	$M_{\{X\}}$	M_X	<i>Exercise 2.60.</i> Fill out the matrix M_X associated to the graph X in Eq. (2.56):
fong-spivak-seven-sketches_75 F00852	75	■ 0.608	D	D	$M_X = \begin{array}{c cccc} \nearrow & A & B & C & D \\ \hline A & 0 & ? & ? & ? \\ B & 2 & 0 & \infty & ? \\ C & ? & ? & ? & ? \\ D & ? & ? & ? & ? \end{array}$
fong-spivak-seven-sketches_75 F00853	75	■ 0.608	$M_{\{X\}} =$	$M_X =$	$M_X = \begin{array}{c cccc} \nearrow & A & B & C & D \\ \hline A & 0 & ? & ? & ? \\ B & 2 & 0 & \infty & ? \\ C & ? & ? & ? & ? \\ D & ? & ? & ? & ? \end{array}$
fong-spivak-seven-sketches_75 F00854	75	■ 0.993	$\mathcal{M} := (\mathrm{P}(M), \subseteq, M, \cap)$	$\mathcal{M} := (P(M), \subseteq, M, \cap)$	<i>Exercise 2.62.</i> Let M be a set and let $\mathcal{M} := (P(M), \subseteq, M, \cap)$ be the monoidal preorder
fong-spivak-seven-sketches_75 F00855	75	■ 0.870	$\mathcal{M}$	$\mathcal{M}$	modes of transportation (e.g. car, boat, foot). Then an $\mathcal{M}$ -category $\mathcal{X}$ tells you all the
fong-spivak-seven-sketches_75 F00856	75	■ 0.966	$a, b \in \mathrm{Ob}(\mathcal{X})$	$a, b \in \mathrm{Ob}(\mathcal{X})$	modes that will get you from a all the way to b , for any two points $a, b \in \mathrm{Ob}(\mathcal{X})$.”
fong-spivak-seven-sketches_75 F00857	75	■ 0.595	$M = \{$	$M = \{$	of $M = \{\text{car}, \text{boat}, \text{foot}\}$.
fong-spivak-seven-sketches_76 F00858	76	■ 1.000	$\mathcal{W} := (\mathbb{N} \cup \{\infty\}, \leq, \infty, \min)$	$\mathcal{W} := (\mathbb{N} \cup \{\infty\}, \leq, \infty, \min)$	<i>Exercise 2.63.</i> Consider the monoidal preorder $\mathcal{W} := (\mathbb{N} \cup \{\infty\}, \leq, \infty, \min)$.
fong-spivak-seven-sketches_76 F00859	76	■ 1.000	$\mathbb{N} \cup \{\infty\}$	$\mathbb{N} \cup \{\infty\}$	1. Draw a small graph labeled by elements of $\mathbb{N} \cup \{\infty\}$.
fong-spivak-seven-sketches_76 F00860	76	■ 1.000	$\mathcal{W}$	$\mathcal{W}$	3. Prove that this matrix is the matrix of hom-objects for a $\mathcal{W}$ -category. This will

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_76 F00861	76	■ 1.000	$f: \mathcal{V} \rightarrow \mathcal{W}$	$f: \mathcal{V} \rightarrow \mathcal{W}$	base. This allows us to use a monoidal monotone $f: \mathcal{V} \rightarrow \mathcal{W}$ to construct $\mathcal{W}$ -categories
fong-spivak-seven-sketches_76 F00862	76	■ 1.000	$\mathcal{V} \rightarrow \mathcal{W}$	$\mathcal{V} \rightarrow \mathcal{W}$	Any monoidal monotone $\mathcal{V} \rightarrow \mathcal{W}$ between symmetric monoidal preorders lets us
fong-spivak-seven-sketches_76 F00863	76	■ 0.999	$\mathcal{C}$	$\mathcal{C}$	Construction 2.64. Let $f: \mathcal{V} \rightarrow \mathcal{W}$ be a monoidal monotone. Given a $\mathcal{V}$ -category $\mathcal{C}$, one
fong-spivak-seven-sketches_76 F00864	76	■ 1.000	$\mathcal{C}_f$	$\mathcal{C}_f$	forms the associated $\mathcal{W}$ -category, say $\mathcal{C}_f$ as follows.
fong-spivak-seven-sketches_76 F00865	76	■ 0.825	$\operatorname{Ob}(\mathcal{C}_f) := \operatorname{Ob}(\mathcal{C})$	$\operatorname{Ob}(\mathcal{C}_f) := \operatorname{Ob}(\mathcal{C})$	(i) We take the same objects: $\operatorname{Ob}(\mathcal{C}_f) := \operatorname{Ob}(\mathcal{C})$.
fong-spivak-seven-sketches_76 F00866	76	■ 0.963	$c, d \in \operatorname{Ob}(\mathcal{C})$	$c, d \in \operatorname{Ob}(\mathcal{C})$	(ii) For any $c, d \in \operatorname{Ob}(\mathcal{C})$, put $\mathcal{C}_f(c, d) := f(\mathcal{C}(c, d))$.
fong-spivak-seven-sketches_76 F00867	76	■ 0.963	$\mathcal{C}_f(c, d) := f(\mathcal{C}(c, d))$	$\mathcal{C}_f(c, d) := f(\mathcal{C}(c, d))$	(ii) For any $c, d \in \operatorname{Ob}(\mathcal{C})$, put $\mathcal{C}_f(c, d) := f(\mathcal{C}(c, d))$.
fong-spivak-seven-sketches_76 F00868	76	■ 1.000	$c \in \mathcal{C}$	$c \in \mathcal{C}$	(a) for every $c \in \mathcal{C}$, we have
fong-spivak-seven-sketches_77 F00869	77	■ 0.964	$c, d, e \in \operatorname{Ob}(\mathcal{C})$	$c, d, e \in \operatorname{Ob}(\mathcal{C})$	(b) for every $c, d, e \in \operatorname{Ob}(\mathcal{C})$ we have
fong-spivak-seven-sketches_77 F00870	77	■ 1.000	$f: [0, \infty] \rightarrow \{\text{true}, \text{false}\}$	$f: [0, \infty] \rightarrow \{\text{true}, \text{false}\}$	<i>Example 2.65.</i> As an example, consider the function $f: [0, \infty] \rightarrow \{\text{true}, \text{false}\}$ given
fong-spivak-seven-sketches_77 F00871	77	■ 0.651	$x_f \neq x_g$	$x_f \neq x_g$	$\mathcal{X}_f \neq \mathcal{X}_g$.
fong-spivak-seven-sketches_77 F00872	77	■ 0.606	$F: \mathcal{X} \rightarrow \mathcal{Y}$	$F: \mathcal{X} \rightarrow \mathcal{Y}$	Definition 2.69. Let $\mathcal{X}$ and $\mathcal{Y}$ be $\mathcal{V}$ -categories. A $\mathcal{V}$ -functor from $\mathcal{X}$ to $\mathcal{Y}$, denoted $F: \mathcal{X} \rightarrow \mathcal{Y}$,
fong-spivak-seven-sketches_77 F00873	77	■ 0.961	$F: \operatorname{Ob}(\mathcal{X}) \rightarrow \operatorname{Ob}(\mathcal{Y})$	$F: \operatorname{Ob}(\mathcal{X}) \rightarrow \operatorname{Ob}(\mathcal{Y})$	(i) a function $F: \operatorname{Ob}(\mathcal{X}) \rightarrow \operatorname{Ob}(\mathcal{Y})$
fong-spivak-seven-sketches_77 F00874	77	■ 0.365	$x_1, x_2 \in \operatorname{Ob}(\mathcal{X})$	$x_1, x_2 \in \operatorname{Ob}(\mathcal{X})$	(a) for all $x_1, x_2 \in \operatorname{Ob}(\mathcal{X})$, one has $\mathcal{X}(x_1, x_2) \leq \mathcal{Y}(F(x_1), F(x_2))$.
fong-spivak-seven-sketches_77 F00875	77	■ 0.365	$\mathcal{X}(x_1, x_2) \leq \mathcal{Y}(F(x_1), F(x_2))$	$\mathcal{X}(x_1, x_2) \leq \mathcal{Y}(F(x_1), F(x_2))$	(a) for all $x_1, x_2 \in \operatorname{Ob}(\mathcal{X})$, one has $\mathcal{X}(x_1, x_2) \leq \mathcal{Y}(F(x_1), F(x_2))$.
fong-spivak-seven-sketches_78 F00876	78	■ 0.829	$X(x_1, x_2) =$	$X(x_1, x_2) =$	orders are Bool -categories, where $\mathcal{X}(x_1, x_2) = \text{true}$ is denoted $x_1 \leq x_2$. One would hope
fong-spivak-seven-sketches_78 F00877	78	■ 0.829	$x_1 \leq x_2$	$x_1 \leq x_2$	orders are Bool -categories, where $\mathcal{X}(x_1, x_2) = \text{true}$ is denoted $x_1 \leq x_2$. One would hope
fong-spivak-seven-sketches_78 F00878	78	■ 1.000	$(X, \leq_X) \rightarrow (Y, \leq_Y)$	$(X, \leq_X) \rightarrow (Y, \leq_Y)$	and that's true. A monotone map $(X, \leq_X) \rightarrow (Y, \leq_Y)$ is a function $F: X \rightarrow Y$ such that
fong-spivak-seven-sketches_78 F00879	78	■ 1.000	$x_1, x_2 \in X$	$x_1, x_2 \in X$	for every $x_1, x_2 \in X$, if $x_1 \leq_X x_2$ then $F(x_1) \leq_Y F(x_2)$. In other words, we have
fong-spivak-seven-sketches_78 F00880	78	■ 1.000	$x_1 \leq_X x_2$	$x_1 \leq_X x_2$	for every $x_1, x_2 \in X$, if $x_1 \leq_X x_2$ then $F(x_1) \leq_Y F(x_2)$. In other words, we have
fong-spivak-seven-sketches_78 F00881	78	■ 1.000	$F(x_1) \leq_Y F(x_2)$	$F(x_1) \leq_Y F(x_2)$	for every $x_1, x_2 \in X$, if $x_1 \leq_X x_2$ then $F(x_1) \leq_Y F(x_2)$. In other words, we have
fong-spivak-seven-sketches_78 F00882	78	■ 0.996	$\mathcal{V} =$	$\mathcal{V} =$	where the above $\leq$ takes place in the enriching category $\mathcal{V} = \mathbf{Bool}$; this is exactly the
fong-spivak-seven-sketches_78 F00883	78	■ 0.990	(X, d_X)	(X, d_X)	does not increase. That is, given Lawvere metric spaces (X, d_X) and (Y, d_Y) , a Cost -
fong-spivak-seven-sketches_78 F00884	78	■ 0.990	(Y, d_Y)	(Y, d_Y)	does not increase. That is, given Lawvere metric spaces (X, d_X) and (Y, d_Y) , a Cost -
fong-spivak-seven-sketches_78 F00885	78	■ 0.982	$d_X(x_1, x_2) \geq d_Y(F(x_1), F(x_2))$	$d_X(x_1, x_2) \geq d_Y(F(x_1), F(x_2))$	$d_X(x_1, x_2) \geq d_Y(F(x_1), F(x_2))$.
fong-spivak-seven-sketches_78 F00886	78	■ 0.971	$\mathcal{X}^{\operatorname{op}}$	$\mathcal{X}^{\operatorname{op}}$	$\mathcal{V}$ -category $\mathcal{X}$ is denoted $\mathcal{X}^{\operatorname{op}}$ and is defined by

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_78 F00887	78	0.784	$\operatorname{Ob}\{\left(\operatorname{cal}{X}^{\text{op}}\right)\}:=\operatorname{Ob}\{\operatorname{cal}{X}\}$	$\operatorname{Ob}(\mathcal{X}^{\text{op}}) := \operatorname{Ob}(\mathcal{X})$	(i) $\operatorname{Ob}(\mathcal{X}^{\text{op}}) := \operatorname{Ob}(\mathcal{X})$, and
fong-spivak-seven-sketches_78 F00888	78	0.959	$x, y \in \operatorname{cal}{X}$	$x, y \in \mathcal{X}$	(ii) for all $x, y \in \mathcal{X}$, we have $\mathcal{X}^{\text{op}}(x, y) := \mathcal{X}(y, x)$.
fong-spivak-seven-sketches_78 F00889	78	0.959	$\operatorname{cal}{X}^{\operatorname{op}}(x, y) := \operatorname{cal}{X}(y, x)$	$\mathcal{X}^{\text{op}}(x, y) := \mathcal{X}(y, x)$	(ii) for all $x, y \in \mathcal{X}$, we have $\mathcal{X}^{\text{op}}(x, y) := \mathcal{X}(y, x)$.
fong-spivak-seven-sketches_78 F00890	78	0.334	$\operatorname{t} : \operatorname{frac}{X} \rightarrow X^{\operatorname{op}}$	$t : \mathfrak{X} \rightarrow X^{\text{op}}$	A $\mathcal{V}$ -category $\mathcal{X}$ is a <i>dagger</i> $\mathcal{V}$ -category if the identity function is a $\mathcal{V}$ -functor $\dagger : \mathcal{X} \rightarrow \mathcal{X}^{\text{op}}$.
fong-spivak-seven-sketches_78 F00891	78	0.470	$I \leq X(x, y)$	$I \leq X(x, y)$	And a <i>skeletal</i> $\mathcal{V}$ -category is one in which if $I \leq \mathcal{X}(x, y)$ and $I \leq \mathcal{X}(y, x)$, then $x = y$.
fong-spivak-seven-sketches_78 F00892	78	0.470	$I \leq X(y, x)$	$I \leq X(y, x)$	And a <i>skeletal</i> $\mathcal{V}$ -category is one in which if $I \leq \mathcal{X}(x, y)$ and $I \leq \mathcal{X}(y, x)$, then $x = y$.
fong-spivak-seven-sketches_78 F00893	78	0.868	$\operatorname{cal}{Y}$	$\mathcal{Y}$	If $\mathcal{V} = (V, \leq, I, \otimes)$ is a symmetric monoidal preorder and $\mathcal{X}$ and $\mathcal{Y}$ are $\mathcal{V}$ -categories, then
fong-spivak-seven-sketches_79 F00894	79	0.966	$\operatorname{cal}{X} \times \operatorname{cal}{Y}$	$\mathcal{X} \times \mathcal{Y}$	to be the $\mathcal{V}$ -category $\mathcal{X} \times \mathcal{Y}$ with
fong-spivak-seven-sketches_79 F00895	79	0.382	$\operatorname{Ob}\{\operatorname{cal}{X} \times \operatorname{cal}{Y}\} := \operatorname{Ob}\{\operatorname{cal}{X}\} \times \operatorname{Ob}\{\operatorname{cal}{Y}\}$	$\operatorname{Ob}(\mathcal{X} \times \mathcal{Y}) := \operatorname{Ob}(\mathcal{X}) \times \operatorname{Ob}(\mathcal{Y})$	(i) $\operatorname{Ob}(\mathcal{X} \times \mathcal{Y}) := \operatorname{Ob}(\mathcal{X}) \times \operatorname{Ob}(\mathcal{Y})$,
fong-spivak-seven-sketches_79 F00896	79	0.711	$(\operatorname{cal}{X} \times \operatorname{cal}{Y})\left(\left(x, y\right),\left(x^{\prime}, y^{\prime}\right)\right) := \operatorname{cal}{X}\left(x, x^{\prime}\right) \otimes \operatorname{cal}{Y}\left(y, y^{\prime}\right)$	$(\mathcal{X} \times \mathcal{Y})((x, y), (x', y')) := \mathcal{X}(x, x') \otimes \mathcal{Y}(y, y')$	(ii) $(\mathcal{X} \times \mathcal{Y})((x, y), (x', y')) := \mathcal{X}(x, x') \otimes \mathcal{Y}(y, y')$,
fong-spivak-seven-sketches_79 F00897	79	0.334	$\left(x^{\prime}, y^{\prime}\right)$	(x', y')	for two objects (x, y) and (x', y') in $\operatorname{Ob}(\mathcal{X} \times \mathcal{Y})$.
fong-spivak-seven-sketches_79 F00898	79	0.334	$\operatorname{Ob}\{\operatorname{cal}{X} \times y\}$	$\operatorname{Ob}(\mathcal{X} \times y)$	for two objects (x, y) and (x', y') in $\operatorname{Ob}(\mathcal{X} \times \mathcal{Y})$.
fong-spivak-seven-sketches_79 F00899	79	0.603	$(x, y) \in \operatorname{Ob}(x \times y)$	$(x, y) \in \operatorname{Ob}(x \times y)$	1. Check that for every object $(x, y) \in \operatorname{Ob}(\mathcal{X} \times \mathcal{Y})$ we have $I \leq (\mathcal{X} \times \mathcal{Y})((x, y), (x, y))$.
fong-spivak-seven-sketches_79 F00900	79	0.603	$I \leq(x \times y)\left((x, y),(x, y)\right)$	$I \leq (x \times y)((x, y), (x, y))$	1. Check that for every object $(x, y) \in \operatorname{Ob}(\mathcal{X} \times \mathcal{Y})$ we have $I \leq (\mathcal{X} \times \mathcal{Y})((x, y), (x, y))$.
fong-spivak-seven-sketches_79 F00901	79	1.000	$\left(x_1, y_1\right),\left(x_2, y_2\right)$	$(x_1, y_1), (x_2, y_2)$	2. Check that for every three objects $(x_1, y_1), (x_2, y_2)$, and (x_3, y_3) , we have
fong-spivak-seven-sketches_79 F00902	79	1.000	$\left(x_3, y_3\right)$	(x_3, y_3)	2. Check that for every three objects $(x_1, y_1), (x_2, y_2)$, and (x_3, y_3) , we have
fong-spivak-seven-sketches_79 F00903	79	1.000	$\left(P, \leq_P\right) \times\left(Q, \leq_Q\right)$	$(P, \leq_P) \times (Q, \leq_Q)$	When taking the product of two preorders $(P, \leq_P) \times (Q, \leq_Q)$, as first described in
fong-spivak-seven-sketches_79 F00904	79	0.960	$\left(p_1, q_1\right) \leq\left(p_2, q_2\right)$	$(p_1, q_1) \leq (p_2, q_2)$	Example 1.56, we say that $(p_1, q_1) \leq (p_2, q_2)$ iff both $p_1 \leq p_2$ AND $q_1 \leq q_2$; the AND is
fong-spivak-seven-sketches_79 F00905	79	0.960	$q_1 \leq q_2$	$q_1 \leq q_2$	Example 1.56, we say that $(p_1, q_1) \leq (p_2, q_2)$ iff both $p_1 \leq p_2$ AND $q_1 \leq q_2$; the AND is
fong-spivak-seven-sketches_79 F00906	79	0.640	$x \times y$	$x \times y$	objects in $\mathcal{X} \times \mathcal{Y}$. And the distance $d_{\mathcal{X} \times \mathcal{Y}}((x, y), (x', y'))$ between any two points is given
fong-spivak-seven-sketches_79 F00907	79	0.640	$d_{X \times Y}\left(\left(x, y\right),\left(x^{\prime}, y^{\prime}\right)\right)$	$d_{\mathcal{X} \times \mathcal{Y}}((x, y), (x', y'))$	objects in $\mathcal{X} \times \mathcal{Y}$. And the distance $d_{\mathcal{X} \times \mathcal{Y}}((x, y), (x', y'))$ between any two points is given
fong-spivak-seven-sketches_79 F00908	79	1.000	$d_X\left(x, x^{\prime}\right)+d_Y\left(y, y^{\prime}\right)$	$d_X(x, x') + d_Y(y, y')$	by the sum $d_X(x, x') + d_Y(y, y')$.
fong-spivak-seven-sketches_80 F00909	80	1.000	$\mathbb{R} \times \mathbb{R}$	$\mathbb{R} \times \mathbb{R}$	Example 2.54). Form the Cost -product $\mathbb{R} \times \mathbb{R}$. What is the distance from (5,6) to
fong-spivak-seven-sketches_80 F00910	80	1.000	(5,6)	(5,6)	Example 2.54). Form the Cost -product $\mathbb{R} \times \mathbb{R}$. What is the distance from (5,6) to
fong-spivak-seven-sketches_80 F00911	80	0.876	(-1,4)	(-1,4)	(-1,4)? Hint: apply Definition 2.74; the answer is not $\sqrt{40}$.
fong-spivak-seven-sketches_80 F00912	80	0.876	$\sqrt{40}$	$\sqrt{40}$	(-1,4)? Hint: apply Definition 2.74; the answer is not $\sqrt{40}$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image																																																	
fong-spivak-seven-sketches-F00913	80	0.764	A, p	A, p	<table><tr><th>$\mathcal{X} \times \mathcal{Y}$</th><th>$(A, p)$</th><th>$(B, p)$</th><th>$(C, p)$</th><th>$(A, q)$</th><th>$(B, q)$</th><th>$(C, q)$</th></tr><tr><td>$(A, p)$</td><td>0</td><td>2</td><td>5</td><td>5</td><td>7</td><td>10</td></tr><tr><td>(B, p)</td><td>∞</td><td>0</td><td>3</td><td>∞</td><td>5</td><td>8</td></tr><tr><td>(C, p)</td><td>∞</td><td>∞</td><td>0</td><td>∞</td><td>∞</td><td>5</td></tr><tr><td>(A, q)</td><td>8</td><td>10</td><td>13</td><td>0</td><td>2</td><td>5</td></tr><tr><td>(B, q)</td><td>∞</td><td>8</td><td>11</td><td>∞</td><td>0</td><td>3</td></tr><tr><td>(C, q)</td><td>∞</td><td>∞</td><td>8</td><td>∞</td><td>∞</td><td>0</td></tr></table>	$\mathcal{X} \times \mathcal{Y}$	(A, p)	(B, p)	(C, p)	(A, q)	(B, q)	(C, q)	(A, p)	0	2	5	5	7	10	(B, p)	∞	0	3	∞	5	8	(C, p)	∞	∞	0	∞	∞	5	(A, q)	8	10	13	0	2	5	(B, q)	∞	8	11	∞	0	3	(C, q)	∞	∞	8	∞	∞	0
$\mathcal{X} \times \mathcal{Y}$	(A, p)	(B, p)	(C, p)	(A, q)	(B, q)	(C, q)																																																
(A, p)	0	2	5	5	7	10																																																
(B, p)	∞	0	3	∞	5	8																																																
(C, p)	∞	∞	0	∞	∞	5																																																
(A, q)	8	10	13	0	2	5																																																
(B, q)	∞	8	11	∞	0	3																																																
(C, q)	∞	∞	8	∞	∞	0																																																
fong-spivak-seven-sketches-F00914	80	0.764	(B, p)	(B, p)	<table><tr><th>$\mathcal{X} \times \mathcal{Y}$</th><th>$(A, p)$</th><th>$(B, p)$</th><th>$(C, p)$</th><th>$(A, q)$</th><th>$(B, q)$</th><th>$(C, q)$</th></tr><tr><td>$(A, p)$</td><td>0</td><td>2</td><td>5</td><td>5</td><td>7</td><td>10</td></tr><tr><td>(B, p)</td><td>∞</td><td>0</td><td>3</td><td>∞</td><td>5</td><td>8</td></tr><tr><td>(C, p)</td><td>∞</td><td>∞</td><td>0</td><td>∞</td><td>∞</td><td>5</td></tr><tr><td>(A, q)</td><td>8</td><td>10</td><td>13</td><td>0</td><td>2</td><td>5</td></tr><tr><td>(B, q)</td><td>∞</td><td>8</td><td>11</td><td>∞</td><td>0</td><td>3</td></tr><tr><td>(C, q)</td><td>∞</td><td>∞</td><td>8</td><td>∞</td><td>∞</td><td>0</td></tr></table>	$\mathcal{X} \times \mathcal{Y}$	(A, p)	(B, p)	(C, p)	(A, q)	(B, q)	(C, q)	(A, p)	0	2	5	5	7	10	(B, p)	∞	0	3	∞	5	8	(C, p)	∞	∞	0	∞	∞	5	(A, q)	8	10	13	0	2	5	(B, q)	∞	8	11	∞	0	3	(C, q)	∞	∞	8	∞	∞	0
$\mathcal{X} \times \mathcal{Y}$	(A, p)	(B, p)	(C, p)	(A, q)	(B, q)	(C, q)																																																
(A, p)	0	2	5	5	7	10																																																
(B, p)	∞	0	3	∞	5	8																																																
(C, p)	∞	∞	0	∞	∞	5																																																
(A, q)	8	10	13	0	2	5																																																
(B, q)	∞	8	11	∞	0	3																																																
(C, q)	∞	∞	8	∞	∞	0																																																
fong-spivak-seven-sketches-F00915	80	0.764	(C, p)	(C, p)	<table><tr><th>$\mathcal{X} \times \mathcal{Y}$</th><th>$(A, p)$</th><th>$(B, p)$</th><th>$(C, p)$</th><th>$(A, q)$</th><th>$(B, q)$</th><th>$(C, q)$</th></tr><tr><td>$(A, p)$</td><td>0</td><td>2</td><td>5</td><td>5</td><td>7</td><td>10</td></tr><tr><td>(B, p)</td><td>∞</td><td>0</td><td>3</td><td>∞</td><td>5</td><td>8</td></tr><tr><td>(C, p)</td><td>∞</td><td>∞</td><td>0</td><td>∞</td><td>∞</td><td>5</td></tr><tr><td>(A, q)</td><td>8</td><td>10</td><td>13</td><td>0</td><td>2</td><td>5</td></tr><tr><td>(B, q)</td><td>∞</td><td>8</td><td>11</td><td>∞</td><td>0</td><td>3</td></tr><tr><td>(C, q)</td><td>∞</td><td>∞</td><td>8</td><td>∞</td><td>∞</td><td>0</td></tr></table>	$\mathcal{X} \times \mathcal{Y}$	(A, p)	(B, p)	(C, p)	(A, q)	(B, q)	(C, q)	(A, p)	0	2	5	5	7	10	(B, p)	∞	0	3	∞	5	8	(C, p)	∞	∞	0	∞	∞	5	(A, q)	8	10	13	0	2	5	(B, q)	∞	8	11	∞	0	3	(C, q)	∞	∞	8	∞	∞	0
$\mathcal{X} \times \mathcal{Y}$	(A, p)	(B, p)	(C, p)	(A, q)	(B, q)	(C, q)																																																
(A, p)	0	2	5	5	7	10																																																
(B, p)	∞	0	3	∞	5	8																																																
(C, p)	∞	∞	0	∞	∞	5																																																
(A, q)	8	10	13	0	2	5																																																
(B, q)	∞	8	11	∞	0	3																																																
(C, q)	∞	∞	8	∞	∞	0																																																
fong-spivak-seven-sketches-F00916	80	0.764	(A, q)	(A, q)	<table><tr><th>$\mathcal{X} \times \mathcal{Y}$</th><th>$(A, p)$</th><th>$(B, p)$</th><th>$(C, p)$</th><th>$(A, q)$</th><th>$(B, q)$</th><th>$(C, q)$</th></tr><tr><td>$(A, p)$</td><td>0</td><td>2</td><td>5</td><td>5</td><td>7</td><td>10</td></tr><tr><td>(B, p)</td><td>∞</td><td>0</td><td>3</td><td>∞</td><td>5</td><td>8</td></tr><tr><td>(C, p)</td><td>∞</td><td>∞</td><td>0</td><td>∞</td><td>∞</td><td>5</td></tr><tr><td>(A, q)</td><td>8</td><td>10</td><td>13</td><td>0</td><td>2</td><td>5</td></tr><tr><td>(B, q)</td><td>∞</td><td>8</td><td>11</td><td>∞</td><td>0</td><td>3</td></tr><tr><td>(C, q)</td><td>∞</td><td>∞</td><td>8</td><td>∞</td><td>∞</td><td>0</td></tr></table>	$\mathcal{X} \times \mathcal{Y}$	(A, p)	(B, p)	(C, p)	(A, q)	(B, q)	(C, q)	(A, p)	0	2	5	5	7	10	(B, p)	∞	0	3	∞	5	8	(C, p)	∞	∞	0	∞	∞	5	(A, q)	8	10	13	0	2	5	(B, q)	∞	8	11	∞	0	3	(C, q)	∞	∞	8	∞	∞	0
$\mathcal{X} \times \mathcal{Y}$	(A, p)	(B, p)	(C, p)	(A, q)	(B, q)	(C, q)																																																
(A, p)	0	2	5	5	7	10																																																
(B, p)	∞	0	3	∞	5	8																																																
(C, p)	∞	∞	0	∞	∞	5																																																
(A, q)	8	10	13	0	2	5																																																
(B, q)	∞	8	11	∞	0	3																																																
(C, q)	∞	∞	8	∞	∞	0																																																
fong-spivak-seven-sketches-F00917	80	0.764	(B, q)	(B, q)	<table><tr><th>$\mathcal{X} \times \mathcal{Y}$</th><th>$(A, p)$</th><th>$(B, p)$</th><th>$(C, p)$</th><th>$(A, q)$</th><th>$(B, q)$</th><th>$(C, q)$</th></tr><tr><td>$(A, p)$</td><td>0</td><td>2</td><td>5</td><td>5</td><td>7</td><td>10</td></tr><tr><td>(B, p)</td><td>∞</td><td>0</td><td>3</td><td>∞</td><td>5</td><td>8</td></tr><tr><td>(C, p)</td><td>∞</td><td>∞</td><td>0</td><td>∞</td><td>∞</td><td>5</td></tr><tr><td>(A, q)</td><td>8</td><td>10</td><td>13</td><td>0</td><td>2</td><td>5</td></tr><tr><td>(B, q)</td><td>∞</td><td>8</td><td>11</td><td>∞</td><td>0</td><td>3</td></tr><tr><td>(C, q)</td><td>∞</td><td>∞</td><td>8</td><td>∞</td><td>∞</td><td>0</td></tr></table>	$\mathcal{X} \times \mathcal{Y}$	(A, p)	(B, p)	(C, p)	(A, q)	(B, q)	(C, q)	(A, p)	0	2	5	5	7	10	(B, p)	∞	0	3	∞	5	8	(C, p)	∞	∞	0	∞	∞	5	(A, q)	8	10	13	0	2	5	(B, q)	∞	8	11	∞	0	3	(C, q)	∞	∞	8	∞	∞	0
$\mathcal{X} \times \mathcal{Y}$	(A, p)	(B, p)	(C, p)	(A, q)	(B, q)	(C, q)																																																
(A, p)	0	2	5	5	7	10																																																
(B, p)	∞	0	3	∞	5	8																																																
(C, p)	∞	∞	0	∞	∞	5																																																
(A, q)	8	10	13	0	2	5																																																
(B, q)	∞	8	11	∞	0	3																																																
(C, q)	∞	∞	8	∞	∞	0																																																

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F00918	80	0.764	(C, q)	(C, q)	$\begin{array}{c ccc ccc} \mathcal{X} \times \mathcal{Y} & (A, p) & (B, p) & (C, p) & (A, q) & (B, q) & (C, q) \\ \hline (A, p) & 0 & 2 & 5 & 5 & 7 & 10 \\ (B, p) & \infty & 0 & 3 & \infty & 5 & 8 \\ (C, p) & \infty & \infty & 0 & \infty & \infty & 5 \\ \hline (A, q) & 8 & 10 & 13 & 0 & 2 & 5 \\ (B, q) & \infty & 8 & 11 & \infty & 0 & 3 \\ (C, q) & \infty & \infty & 8 & \infty & \infty & 0 \end{array}$
fong-spivak-seven-sketches F00919	80	0.764	(A, p)	(A, p)	$\begin{array}{c ccc ccc} \mathcal{X} \times \mathcal{Y} & (A, p) & (B, p) & (C, p) & (A, q) & (B, q) & (C, q) \\ \hline (A, p) & 0 & 2 & 5 & 5 & 7 & 10 \\ (B, p) & \infty & 0 & 3 & \infty & 5 & 8 \\ (C, p) & \infty & \infty & 0 & \infty & \infty & 5 \\ \hline (A, q) & 8 & 10 & 13 & 0 & 2 & 5 \\ (B, q) & \infty & 8 & 11 & \infty & 0 & 3 \\ (C, q) & \infty & \infty & 8 & \infty & \infty & 0 \end{array}$
fong-spivak-seven-sketches F00920	80	0.482	$\mathfrak{y}$	η	like $\mathfrak{X}$ —for every entry of $\mathcal{Y}$. Make sure you can see each block as the $\mathfrak{X}$ -matrix shifted
fong-spivak-seven-sketches F00921	81	1.000	$v, w \in V$	$v, w \in V$	monoidal closed (or just closed) if, for every two elements $v, w \in V$, there is an element
fong-spivak-seven-sketches F00922	81	1.000	$a, v, w \in V$	$a, v, w \in V$	for all $a, v, w \in V$.
fong-spivak-seven-sketches F00923	81	1.000	$v \in V$	$v \in V$	preorder is monoidal closed iff, given any $v \in V$, the map $(- \otimes v): V \rightarrow V$ given by
fong-spivak-seven-sketches F00924	81	1.000	$(- \otimes v): V \rightarrow V$	$(- \otimes v): V \rightarrow V$	preorder is monoidal closed iff, given any $v \in V$, the map $(- \otimes v): V \rightarrow V$ given by
fong-spivak-seven-sketches F00925	81	0.747	$(v \circ -): V \rightarrow V$	$(v \circ -): V \rightarrow V$	multiplying with v has a right adjoint. We write this right adjoint $(v \multimap -): V \rightarrow V$.
fong-spivak-seven-sketches F00926	81	1.000	$(- \otimes v)$	$(- \otimes v)$	1. Using Definition 2.2, show that $(- \otimes v)$ is monotone.
fong-spivak-seven-sketches F00927	81	0.959	$((v \circ w) \otimes v) \leq w$	$((v \circ w) \otimes v) \leq w$	2. Supposing that $\mathcal{V}$ is closed, show that for all $v, w \in V$ we have $((v \multimap w) \otimes v) \leq w$.
fong-spivak-seven-sketches F00928	81	0.702	$(v \circ -)$	$(v \circ -)$	3. Using 2., show that $(v \multimap -)$ is monotone.
fong-spivak-seven-sketches F00929	81	0.950	$x, y \in [0, \infty]$	$x, y \in [0, \infty]$	deed, for any $x, y \in [0, \infty]$, define $x \multimap y := \max(0, y - x)$. Then, for any $a, x, y \in [0, \infty]$,
fong-spivak-seven-sketches F00930	81	0.950	$x \multimap y := \max(0, y - x)$	$x \multimap y := \max(0, y - x)$	deed, for any $x, y \in [0, \infty]$, define $x \multimap y := \max(0, y - x)$. Then, for any $a, x, y \in [0, \infty]$,
fong-spivak-seven-sketches F00931	81	0.950	$a, x, y \in [0, \infty]$	$a, x, y \in [0, \infty]$	deed, for any $x, y \in [0, \infty]$, define $x \multimap y := \max(0, y - x)$. Then, for any $a, x, y \in [0, \infty]$,
fong-spivak-seven-sketches F00932	82	0.746	$\leq p \multimap q$	$\leq p \multimap q$	as in Definition 2.79. Note that $\text{false} \leq p \multimap q$, for any p, q no matter what $\multimap$ is,
fong-spivak-seven-sketches F00933	82	0.746	$\rightarrow$	$\rightarrow$	as in Definition 2.79. Note that $\text{false} \leq p \multimap q$, for any p, q no matter what $\multimap$ is,
fong-spivak-seven-sketches F00934	82	0.987	$a =$	$a =$	because false is less than everything. But using $a = \text{false}$, $p = \text{true}$, and $q = \text{false}$,
fong-spivak-seven-sketches F00935	82	0.987	$p =$	$p =$	because false is less than everything. But using $a = \text{false}$, $p = \text{true}$, and $q = \text{false}$,
fong-spivak-seven-sketches F00936	82	0.987	$q =$	$q =$	because false is less than everything. But using $a = \text{false}$, $p = \text{true}$, and $q = \text{false}$,

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F00937	82	■ 0.999	$(a \vee p) \not\leq q$	$(a \vee p) \not\leq q$	we then get a contradiction: $(a \vee p) \not\leq q$ and yet $a \leq (p \multimap q)$.
fong-spivak-seven-sketches F00938	82	■ 0.999	$a \leq (p \multimap q)$	$a \leq (p \multimap q)$	we then get a contradiction: $(a \vee p) \not\leq q$ and yet $a \leq (p \multimap q)$.
fong-spivak-seven-sketches F00939	82	■ 1.000	c, d	c, d	say that for every two material collections c, d one can form a material collection $c \multimap d$
fong-spivak-seven-sketches F00940	82	■ 1.000	$c \rightarrowtail d$	$c \rightarrow d$	say that for every two material collections c, d one can form a material collection $c \multimap d$
fong-spivak-seven-sketches F00941	82	■ 1.000	$2 \mathrm{H}_{\{2\}} + 2 \mathrm{Na} \rightarrowtail 2 \mathrm{NaOH} + \mathrm{H}_2$	$2\mathrm{H}_2\mathrm{O} + 2\mathrm{Na} \rightarrow 2\mathrm{NaOH} + \mathrm{H}_2$	Or more down to earth, since we have the reaction $2\mathrm{H}_2\mathrm{O} + 2\mathrm{Na} \rightarrow 2\mathrm{NaOH} + \mathrm{H}_2$, we
fong-spivak-seven-sketches F00942	82	■ 0.426	$2 \mathrm{Na} \rightarrowtail (2 \mathrm{NaOH} + \mathrm{H}_2)$	$2\mathrm{Na} \rightarrow (2\mathrm{NaOH} + \mathrm{H}_2)$	But it is not so far-fetched: this hypothetical new substance $(2\mathrm{Na} \multimap (2\mathrm{NaOH} + \mathrm{H}_2))$
fong-spivak-seven-sketches F00943	82	■ 0.935	$\mathcal{V} = (V, \leq, I, \otimes, \multimap)$	$\mathcal{V} = (V, \leq, I, \otimes, \multimap)$	Proposition 2.87. Suppose $\mathcal{V} = (V, \leq, I, \otimes, \multimap)$ is a symmetric monoidal preorder that
fong-spivak-seven-sketches F00944	82	■ 1.000	$\multimap v : (V, \leq) \rightarrowtail (V, \leq)$	$\multimap v : (V, \leq) \rightarrow (V, \leq)$	(a) For every $v \in V$, the monotone map $\multimap v : (V, \leq) \rightarrow (V, \leq)$ is left adjoint to
fong-spivak-seven-sketches F00945	82	■ 0.913	$v \rightarrowtail - : (V, \leq) \rightarrowtail (V, \leq)$	$v \rightarrow - : (V, \leq) \rightarrow (V, \leq)$	$v \multimap - : (V, \leq) \rightarrow (V, \leq)$.
fong-spivak-seven-sketches F00946	82	■ 1.000	$A \subseteq V$	$A \subseteq V$	(b) For any element $v \in V$ and set of elements $A \subseteq V$, if the join $\bigvee_{a \in A} a$ exists then
fong-spivak-seven-sketches F00947	82	■ 1.000	$\bigvee_{a \in A} a$	$\bigvee_{a \in A} a$	(b) For any element $v \in V$ and set of elements $A \subseteq V$, if the join $\bigvee_{a \in A} a$ exists then
fong-spivak-seven-sketches F00948	82	■ 1.000	$\bigvee_{a \in A} v \otimes a$	$\bigvee_{a \in A} v \otimes a$	so does $\bigvee_{a \in A} v \otimes a$ and we have
fong-spivak-seven-sketches F00949	82	■ 1.000	$v \otimes (v \multimap w) \leq w$	$v \otimes (v \multimap w) \leq w$	(c) For any $v, w \in V$, we have $v \otimes (v \multimap w) \leq w$.
fong-spivak-seven-sketches F00950	82	■ 1.000	$v \cong (I \multimap v)$	$v \cong (I \multimap v)$	(d) For any $v \in V$, we have $v \cong (I \multimap v)$.
fong-spivak-seven-sketches F00951	82	■ 1.000	$u, v, w \in V$	$u, v, w \in V$	(e) For any $u, v, w \in V$, we have $(u \multimap v) \otimes (v \multimap w) \leq (u \multimap w)$.
fong-spivak-seven-sketches F00952	82	■ 1.000	$(u \multimap v) \otimes (v \multimap w) \leq (u \multimap w)$	$(u \multimap v) \otimes (v \multimap w) \leq (u \multimap w)$	(e) For any $u, v, w \in V$, we have $(u \multimap v) \otimes (v \multimap w) \leq (u \multimap w)$.
fong-spivak-seven-sketches F00953	83	■ 0.748	$(v \multimap -)$	$(v \multimap -)$	(a) The definition of $(\multimap v)$ being left adjoint to $(v \multimap -)$ is exactly the condition
fong-spivak-seven-sketches F00954	83	■ 0.999	$(v \multimap w) \leq (v \multimap w)$	$(v \multimap w) \leq (v \multimap w)$	in Proposition 1.107. More concretely, from reflexivity $v \multimap w \leq (v \multimap w)$, we
fong-spivak-seven-sketches F00955	83	■ 0.971	$(v \multimap w) \otimes v \leq w$	$(v \multimap w) \otimes v \leq w$	obtain $v \multimap w \otimes v \leq w$ Eq. (2.80), and we are done by symmetry, which says
fong-spivak-seven-sketches F00956	83	■ 0.988	$v \otimes (v \multimap w) = (v \multimap w) \otimes v$	$v \otimes (v \multimap w) = (v \multimap w) \otimes v$	$v \otimes (v \multimap w) = (v \multimap w) \otimes v$.
fong-spivak-seven-sketches F00957	83	■ 1.000	$v \otimes I = v \leq v$	$v \otimes I = v \leq v$	(d) Since $v \otimes I = v \leq v$, Eq. (2.80) says $v \leq (I \multimap v)$. For the other direction, we have
fong-spivak-seven-sketches F00958	83	■ 1.000	$v \leq (I \multimap v)$	$v \leq (I \multimap v)$	(d) Since $v \otimes I = v \leq v$, Eq. (2.80) says $v \leq (I \multimap v)$. For the other direction, we have
fong-spivak-seven-sketches F00959	83	■ 0.909	$(I \multimap v) = I \otimes (I \multimap v) \leq v$	$(I \multimap v) = I \otimes (I \multimap v) \leq v$	$(I \multimap v) = I \otimes (I \multimap v) \leq v$ by (c).
fong-spivak-seven-sketches F00960	83	■ 0.966	$u \otimes (u \multimap v) \otimes (v \multimap w) \leq w$	$u \otimes (u \multimap v) \otimes (v \multimap w) \leq w$	(e) To obtain this inequality, we just need $u \otimes (u \multimap v) \otimes (v \multimap w) \leq w$. But this
fong-spivak-seven-sketches F00961	83	■ 0.994	$v, w \in \mathrm{Ob}(\mathcal{V})$	$v, w \in \mathrm{Ob}(\mathcal{V})$	<i>Remark 2.89.</i> We can consider $\mathcal{V}$ to be enriched in itself. That is, for every $v, w \in \mathrm{Ob}(\mathcal{V})$,
fong-spivak-seven-sketches F00962	83	■ 0.899	$\mathcal{V}(v, w) := (v \multimap w) \in \mathcal{V}$	$\mathcal{V}(v, w) := (v \multimap w) \in \mathcal{V}$	we can define $\mathcal{V}(v, w) := (v \multimap w) \in \mathcal{V}$. For this to really be an enrichment, we just need
fong-spivak-seven-sketches F00963	83	■ 0.858	$I \leq \mathcal{X}(x, x) = (x \multimap x)$	$I \leq \mathcal{X}(x, x) = (x \multimap x)$	to check the two conditions of Definition 2.46. The first condition $I \leq \mathcal{X}(x, x) = (x \multimap x)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_1 F00964	83	■ 1.000	$I \otimes x \leq x$	$I \otimes x \leq x$	is satisfied because $I \otimes x \leq x$. The second condition is satisfied by Proposition 2.87(e).
fong-spivak-seven-sketches_1 F00965	83	■ 0.844	$\mathcal{V} = (V, \leq, I, \otimes, \multimap)$	$\mathcal{V} = (V, \leq, I, \otimes, \multimap)$	$\mathcal{V} = (V, \leq, I, \otimes, \multimap)$ that has all joins: $\bigvee A$ exists for every $A \subseteq V$. In particular, we often
fong-spivak-seven-sketches_1 F00966	83	■ 0.844	$\bigvee A$	$\bigvee A$	$\mathcal{V} = (V, \leq, I, \otimes, \multimap)$ that has all joins: $\bigvee A$ exists for every $A \subseteq V$. In particular, we often
fong-spivak-seven-sketches_1 F00967	83	■ 1.000	$0 := \vee \varnothing$	$0 := \bigvee \emptyset$	denote the empty join by $0 := \bigvee \emptyset$.
fong-spivak-seven-sketches_1 F00968	83	■ 1.000	$A \subseteq [0, \infty]$	$A \subseteq [0, \infty]$	Cost is a quantale, we take an arbitrary set of elements $A \subseteq [0, \infty]$ and ask if it has a
fong-spivak-seven-sketches_1 F00969	83	■ 0.564	$\vee A$	$\bigvee A$	join $\bigvee A$. To be a join, it needs to satisfy two properties:
fong-spivak-seven-sketches_1 F00970	83	■ 0.959	$a \geq \bigvee A$	$a \geq \bigvee A$	a. $a \geq \bigvee A$ for all $a \in A$, and
fong-spivak-seven-sketches_1 F00971	83	■ 0.969	$b \in [0, \infty]$	$b \in [0, \infty]$	b. if $b \in [0, \infty]$ is any element such that $a \geq b$ for all $a \in A$, then $\bigvee A \geq b$.
fong-spivak-seven-sketches_1 F00972	83	■ 0.969	$a \geq b$	$a \geq b$	b. if $b \in [0, \infty]$ is any element such that $a \geq b$ for all $a \in A$, then $\bigvee A \geq b$.
fong-spivak-seven-sketches_1 F00973	83	■ 0.969	$\bigvee A \geq b$	$\bigvee A \geq b$	b. if $b \in [0, \infty]$ is any element such that $a \geq b$ for all $a \in A$, then $\bigvee A \geq b$.
fong-spivak-seven-sketches_1 F00974	84	■ 0.728	$A \cdot \{ \}^5$	A^5	bound, of A^5 For example, if $A = \{2, 3\}$ then $\bigvee A = 2$. We have joins for infinite sets
fong-spivak-seven-sketches_1 F00975	84	■ 0.728	$A = \{2, 3\}$	$A = \{2, 3\}$	bound, of A^5 For example, if $A = \{2, 3\}$ then $\bigvee A = 2$. We have joins for infinite sets
fong-spivak-seven-sketches_1 F00976	84	■ 0.728	$\bigvee A = 2$	$\bigvee A = 2$	bound, of A^5 For example, if $A = \{2, 3\}$ then $\bigvee A = 2$. We have joins for infinite sets
fong-spivak-seven-sketches_1 F00977	84	■ 0.573	$B = \{2.5, 2.05, 2.005, \dots\}$	$B = \{2.5, 2.05, 2.005, \dots\}$	too: if $B = \{2.5, 2.05, 2.005, \dots\}$, its infimum is 2. Finally, in order to say that $([0, \infty], \geq)$
fong-spivak-seven-sketches_1 F00978	84	■ 1.000	$A = \varnothing$	$A = \emptyset$	has all joins, we need a join to exist for the empty set $A = \emptyset$ too. The first condition
fong-spivak-seven-sketches_1 F00979	84	■ 0.994	$\bigvee \varnothing \geq b$	$\bigvee \emptyset \geq b$	$b \in [0, \infty]$ we have $\bigvee \emptyset \geq b$; this means $\bigvee \emptyset = \infty$.
fong-spivak-seven-sketches_1 F00980	84	■ 0.994	$\bigvee \varnothing = \infty$	$\bigvee \emptyset = \infty$	$b \in [0, \infty]$ we have $\bigvee \emptyset \geq b$; this means $\bigvee \emptyset = \infty$.
fong-spivak-seven-sketches_1 F00981	84	■ 0.998	$\vee \varnothing$	$\bigvee \emptyset$	1. What is $\bigvee \emptyset$, which we generally denote 0, in the case
fong-spivak-seven-sketches_1 F00982	84	■ 0.408	$\mathcal{V} = \mathbf{Bool} = (\mathbb{B}, \leq$	$\mathcal{V} = \mathbf{Bool} = (\mathbb{B}, \leq$	a. $\mathcal{V} = \mathbf{Bool} = (\mathbb{B}, \leq, \text{true}, \wedge)$?
fong-spivak-seven-sketches_1 F00983	84	■ 0.959	$\mathcal{V} = \mathbf{Cost} = ([0, \infty], \geq, 0, +)$	$\mathcal{V} = \mathbf{Cost} = ([0, \infty], \geq, 0, +)$	b. $\mathcal{V} = \mathbf{Cost} = ([0, \infty], \geq, 0, +)$?
fong-spivak-seven-sketches_1 F00984	84	■ 1.000	$x \vee y$	$x \vee y$	2. What is the join $x \vee y$ in the case
fong-spivak-seven-sketches_1 F00985	84	■ 0.724	$\mathcal{V} = \mathbf{Bool}$	$\mathcal{V} = \mathbf{Bool}$	a. $\mathcal{V} = \mathbf{Bool}$, and $x, y \in \mathbb{B}$ are booleans?
fong-spivak-seven-sketches_1 F00986	84	■ 0.724	$x, y \in \mathbb{B}$	$x, y \in \mathbb{B}$	a. $\mathcal{V} = \mathbf{Bool}$, and $x, y \in \mathbb{B}$ are booleans?
fong-spivak-seven-sketches_1 F00987	84	■ 0.939	$\mathbf{Bool} = (\mathbb{B}, \leq$	$\mathbf{Bool} = (\mathbb{B}, \leq$	<i>Exercise 2.93.</i> Show that $\mathbf{Bool} = (\mathbb{B}, \leq, \text{true}, \wedge)$ is a quantale.
fong-spivak-seven-sketches_1 F00988	84	■ 0.884	$\mathcal{P}(S), \subseteq, S, \cap$	$\mathcal{P}(S), \subseteq, S, \cap$	<i>Exercise 2.94.</i> Let S be a set and recall the power set monoidal preorder $(\mathcal{P}(S), \subseteq, S, \cap)$
fong-spivak-seven-sketches_1 F00989	84	■ 1.000	$\mathcal{P} = (P, \leq)$	$\mathcal{P} = (P, \leq)$	Proposition 2.96. Let $\mathcal{P} = (P, \leq)$ be a preorder. It has all joins iff it has all meets.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-1 F00990	84	■ 1.000	$A \subseteq \mathcal{P}$	$A \subseteq \mathcal{P}$	Suppose $\mathcal{P}$ has all joins and suppose that $A \subseteq \mathcal{P}$ is a subset for which we want
fong-spivak-seven-sketches-1 F00991	84	■ 1.000	$M_A := \{p \in P \mid p \leq a$	$M_A := \{p \in P \mid p \leq a$	the meet. Consider the set $M_A := \{p \in P \mid p \leq a \text{ for all } a \in A\}$ of elements below
fong-spivak-seven-sketches-1 F00992	84	■ 1.000	$a \in A\}$	$a \in A\}$	the meet. Consider the set $M_A := \{p \in P \mid p \leq a \text{ for all } a \in A\}$ of elements below
fong-spivak-seven-sketches-1 F00993	84	■ 1.000	$m_A := \bigvee_{p \in M_A} p$	$m_A := \bigvee_{p \in M_A} p$	everything in A . Let $m_A := \bigvee_{p \in M_A} p$ be their join. We claim that m_A is a meet for A .
fong-spivak-seven-sketches-1 F00994	84	■ 1.000	m_A	m_A	everything in A . Let $m_A := \bigvee_{p \in M_A} p$ be their join. We claim that m_A is a meet for A .
fong-spivak-seven-sketches-1 F00995	84	■ 1.000	$m_A \leq a$	$m_A \leq a$	We first need to know that for any $a \in A$ we have $m_A \leq a$, but this is by definition of
fong-spivak-seven-sketches-1 F00996	84	■ 1.000	$p \in M_A$	$p \in M_A$	join: since all $p \in M_A$ satisfy $p \leq a$, so does their join $m_A \leq a$. We second need to know
fong-spivak-seven-sketches-1 F00997	84	■ 1.000	$m' \in P$	$m' \in P$	that for any $m' \in P$ with $m' \leq a$ for all $a \in A$, we have $m' \leq m$. But every such m' is
fong-spivak-seven-sketches-1 F00998	84	■ 1.000	$m' \leq a$	$m' \leq a$	that for any $m' \in P$ with $m' \leq a$ for all $a \in A$, we have $m' \leq m$. But every such m' is
fong-spivak-seven-sketches-1 F00999	84	■ 1.000	$m' \leq m$	$m' \leq m$	that for any $m' \in P$ with $m' \leq a$ for all $a \in A$, we have $m' \leq m$. But every such m' is
fong-spivak-seven-sketches-1 F01000	84	■ 1.000	m'	m'	that for any $m' \in P$ with $m' \leq a$ for all $a \in A$, we have $m' \leq m$. But every such m' is
fong-spivak-seven-sketches-1 F01001	84	■ 1.000	M_A	M_A	actually an element of M_A and m is their join, so $m' \leq m$. This completes the proof.
fong-spivak-seven-sketches-1 F01002	84	■ 0.666	$\{3.1, 3.01, 3.001, \dots\}$	$\{3.1, 3.01, 3.001, \dots\}$	largest number that is $\leq$ everything in A . For example, the infimum of $\{3.1, 3.01, 3.001, \dots\}$ is 3. But note
fong-spivak-seven-sketches-1 F01003	85	■ 1.000	$V \subseteq X$	$V \subseteq X$	and $V \subseteq X$, we can generalize the usual Hausdorff distance, on the left below, to the
fong-spivak-seven-sketches-1 F01004	85	■ 0.302	$u \in U$	$u \in U$	answers the question “can I get into V from every $u \in U$,” i.e. $\forall u \in U. \exists v \in V. u \leq v$. Or
fong-spivak-seven-sketches-1 F01005	85	■ 0.302	$\forall u \in U. \exists v \in V. u \leq v$	$\forall u \in U. \exists v \in V. u \leq v$	answers the question “can I get into V from every $u \in U$,” i.e. $\forall u \in U. \exists v \in V. u \leq v$. Or
fong-spivak-seven-sketches-1 F01006	85	■ 0.999	$\mathcal{V} = \mathbf{P}(M)$	$\mathcal{V} = \mathbf{P}(M)$	for another example, use $\mathcal{V} = \mathbf{P}(M)$ with its interpretation as modes of transportation,
fong-spivak-seven-sketches-1 F01007	85	■ 1.000	$d(U, V) \in \mathbf{P}(M)$	$d(U, V) \in \mathbf{P}(M)$	as in Exercise 2.62. Then the Hausdorff distance $d(U, V) \in \mathbf{P}(M)$ tells us those modes
fong-spivak-seven-sketches-1 F01008	85	■ 1.000	$- \otimes v : V \rightarrow V$	$- \otimes v : V \rightarrow V$	Eq. (2.88) holds. We need to show that Eq. (2.88) holds then $- \otimes v : V \rightarrow V$ has a right
fong-spivak-seven-sketches-1 F01009	85	■ 0.995	$v \multimap -$	$v \multimap -$	adjoint $v \multimap -$. This is just the adjoint functor theorem, Theorem 1.115. It says we can
fong-spivak-seven-sketches-1 F01010	85	■ 1.000	$v \multimap w$	$v \multimap w$	define $v \multimap w$ to be
fong-spivak-seven-sketches-1 F01011	85	■ 0.703	$\sum$	$\sum$	We will get a formula where joins stand in for the sum operation $\sum$, and $\otimes$ stands in
fong-spivak-seven-sketches-1 F01012	85	■ 1.000	$\mathcal{V} = (V, \leq, \otimes, I)$	$\mathcal{V} = (V, \leq, \otimes, I)$	Definition 2.100. Let $\mathcal{V} = (V, \leq, \otimes, I)$ be a quantale. Given sets X and Y , a <i>matrix with</i>
fong-spivak-seven-sketches-1 F01013	85	■ 1.000	$M : X \times Y \rightarrow V$	$M : X \times Y \rightarrow V$	<i>entries in $\mathcal{V}$</i> , or simply a <i>$\mathcal{V}$-matrix</i> , is a function $M : X \times Y \rightarrow V$. For any $x \in X$ and
fong-spivak-seven-sketches-1 F01014	85	■ 0.999	$M(x, y)$	$M(x, y)$	$y \in Y$, we call $M(x, y)$ the <i>(x, y)-entry</i> .
fong-spivak-seven-sketches-1 F01015	86	■ 1.000	$N : Y \times Z \rightarrow V$	$N : Y \times Z \rightarrow V$	Here is how you multiply $\mathcal{V}$ -matrices $M : X \times Y \rightarrow V$ and $N : Y \times Z \rightarrow V$. Their
fong-spivak-seven-sketches-1 F01016	86	■ 0.997	$(M * N) : X \times Z \rightarrow V$	$(M * N) : X \times Z \rightarrow V$	product is defined to be the matrix $(M * N) : X \times Z \rightarrow V$, whose entries are given by
fong-spivak-seven-sketches-1 F01017	86	■ 0.595	$\mathcal{V} = \mathbf{Bool}$	$\mathcal{V} = \mathbf{Bool}$	<i>Example 2.102.</i> Let $\mathcal{V} = \mathbf{Bool}$. Here is an example of matrix multiplication $M * N$. Here

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_1 F01018	86	0.595	$M * N$	$M * N$	<i>Example 2.102.</i> Let $\mathcal{V} = \mathbf{Bool}$. Here is an example of matrix multiplication $M * N$. Here
fong-spivak-seven-sketches_1 F01019	86	0.996	$X = \{1, 2, 3\}, Y = \{1, 2\}$	$X = \{1, 2, 3\}, Y = \{1, 2\}$	$X = \{1, 2, 3\}, Y = \{1, 2\}$, and $Z = \{1, 2, 3\}$, matrices $M: X \times Y \rightarrow \mathbb{B}$ and $N: Y \times Z \rightarrow \mathbb{B}$
fong-spivak-seven-sketches_1 F01020	86	0.996	$Z = \{1, 2, 3\}$	$Z = \{1, 2, 3\}$	$X = \{1, 2, 3\}, Y = \{1, 2\}$, and $Z = \{1, 2, 3\}$, matrices $M: X \times Y \rightarrow \mathbb{B}$ and $N: Y \times Z \rightarrow \mathbb{B}$
fong-spivak-seven-sketches_1 F01021	86	0.996	$M: X \times Y \rightarrow \mathbb{B}$	$M: X \times Y \rightarrow \mathbb{B}$	$X = \{1, 2, 3\}, Y = \{1, 2\}$, and $Z = \{1, 2, 3\}$, matrices $M: X \times Y \rightarrow \mathbb{B}$ and $N: Y \times Z \rightarrow \mathbb{B}$
fong-spivak-seven-sketches_1 F01022	86	0.996	$N: Y \times Z \rightarrow \mathbb{B}$	$N: Y \times Z \rightarrow \mathbb{B}$	$X = \{1, 2, 3\}, Y = \{1, 2\}$, and $Z = \{1, 2, 3\}$, matrices $M: X \times Y \rightarrow \mathbb{B}$ and $N: Y \times Z \rightarrow \mathbb{B}$
fong-spivak-seven-sketches_1 F01023	86	1.000	$I_X: X \times X \rightarrow V$	$I_X: X \times X \rightarrow V$	The identity V -matrix on a set X is $I_X: X \times X \rightarrow V$ given by
fong-spivak-seven-sketches_1 F01024	86	0.964	2×2	2×2	<i>Exercise 2.103.</i> Write down the 2×2 -identity matrix for each of the quantales $(\mathbb{N}, \leq, 1, *)$,
fong-spivak-seven-sketches_1 F01025	86	0.171	$([0, \infty], \geq, 0, +)$	$([0, \infty], \geq, 0, +)$	$\mathbf{Bool} = (\mathbb{B}, \leq, \mathbf{true}, \wedge)$, and $\mathbf{Cost} = ([0, \infty], \geq, 0, +)$.
fong-spivak-seven-sketches_1 F01026	86	1.000	$I_X * M = M$	$I_X * M = M$	has $I_X * M = M$.
fong-spivak-seven-sketches_1 F01027	86	1.000	$M: W \times X \rightarrow V, N: X \times Y \rightarrow V$	$M: W \times X \rightarrow V, N: X \times Y \rightarrow V$	2. Prove the <i>associative law</i> : for any matrices $M: W \times X \rightarrow V, N: X \times Y \rightarrow V$, and
fong-spivak-seven-sketches_1 F01028	86	1.000	$P: Y \times Z \rightarrow V$	$P: Y \times Z \rightarrow V$	$P: Y \times Z \rightarrow V$, one has $(M * N) * P = M * (N * P)$.
fong-spivak-seven-sketches_1 F01029	86	1.000	$(M * N) * P = M * (N * P)$	$(M * N) * P = M * (N * P)$	$P: Y \times Z \rightarrow V$, one has $(M * N) * P = M * (N * P)$.
fong-spivak-seven-sketches_1 F01030	87	0.999	M_Y^2	M_Y^2	When we multiply M_Y by itself using the formula Eq. (2.101), the result M_Y^2 tells us
fong-spivak-seven-sketches_1 F01031	87	1.000	M_Y^3	M_Y^3	the length of the shortest path traversing 2 edges or fewer. Similarly M_Y^3 tells us about
fong-spivak-seven-sketches_1 F01032	87	1.000	$M_Y^2 = M_Y^3$	$M_Y^2 = M_Y^3$	One sees that the powers stabilize: $M_Y^2 = M_Y^3$; as soon as that happens one has the
fong-spivak-seven-sketches_1 F01033	87	1.000	M_Y^n	M_Y^n	matrix of distances, d_Y . Indeed M_Y^n records the lengths of the shortest path traverse n
fong-spivak-seven-sketches_1 F01034	87	0.911	M_X^2, M_X^3	M_X^2, M_X^3	Calculate M_X^2, M_X^3 , and M_X^4 . Check that M_X^4 is what you got for the distance matrix in
fong-spivak-seven-sketches_1 F01035	87	0.911	M_X^4	M_X^4	Calculate M_X^2, M_X^3 , and M_X^4 . Check that M_X^4 is what you got for the distance matrix in
fong-spivak-seven-sketches_1 F01036	89	1.000	$\$ 2.5$	$\$2.5$	technology) budgets, and that the market for data integration software was \$2.5 billion
fong-spivak-seven-sketches_1 F01037	89	0.798	8%	8%	in 2007 and increasing at a rate of more than 8% per year. In other words, it is a major
fong-spivak-seven-sketches_1 F01038	91	0.984	$\mathcal{C} \rightarrow$	$\mathcal{C} \rightarrow$	by a ‘set-valued’ functor $\mathcal{C} \rightarrow \mathbf{Set}$, and that databases can be mapped to each other via
fong-spivak-seven-sketches_1 F01039	91	1.000	$\mathcal{C} \rightarrow \mathcal{D}$	$\mathcal{C} \rightarrow \mathcal{D}$	functors $\mathcal{C} \rightarrow \mathcal{D}$. In other words, there is a relatively large overlap between database
fong-spivak-seven-sketches_1 F01040	93	1.000	$\{ \}_{2} \mathrm{Ob}(\mathcal{C})$	${}^2\mathrm{Ob}(\mathcal{C})$	(i) one specifies a collection ² $\mathrm{Ob}(\mathcal{C})$, elements of which are called <i>objects</i> .
fong-spivak-seven-sketches_1 F01041	93	0.963	$\mathcal{C}(c, d), \{ \}_{3}$	$\mathcal{C}(c, d), {}^3$	(ii) for every two objects c, d , one specifies a set $\mathcal{C}(c, d)$, ³ elements of which are called
fong-spivak-seven-sketches_1 F01042	93	0.964	$c \in \mathrm{Ob}(\mathcal{C})$	$c \in \mathrm{Ob}(\mathcal{C})$	(iii) for every object $c \in \mathrm{Ob}(\mathcal{C})$, one specifies a morphism $\mathrm{id}_c \in \mathcal{C}(c, c)$, called the
fong-spivak-seven-sketches_1 F01043	93	0.964	$\mathrm{id}_c \in \mathcal{C}(c, c)$	$\mathrm{id}_c \in \mathcal{C}(c, c)$	(iii) for every object $c \in \mathrm{Ob}(\mathcal{C})$, one specifies a morphism $\mathrm{id}_c \in \mathcal{C}(c, c)$, called the
fong-spivak-seven-sketches_1 F01044	93	0.939	$f \in \mathcal{C}(c, d)$	$f \in \mathcal{C}(c, d)$	(iv) for every three objects $c, d, e \in \mathrm{Ob}(\mathcal{C})$ and morphisms $f \in \mathcal{C}(c, d)$ and $g \in \mathcal{C}(d, e)$,

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_1 F01045	93	■ 0.939	$g \in \mathcal{C}(d, e)$	$g \in \mathcal{C}(d, e)$	(iv) for every three objects $c, d, e \in \text{Ob}(\mathcal{C})$ and morphisms $f \in \mathcal{C}(c, d)$ and $g \in \mathcal{C}(d, e)$,
fong-spivak-seven-sketches_1 F01046	93	■ 0.998	$f ; g \in \mathcal{C}(c, e)$	$f ; g \in \mathcal{C}(c, e)$	one specifies a morphism $f \circ g \in \mathcal{C}(c, e)$, called <i>the composite of f and g</i> .
fong-spivak-seven-sketches_1 F01047	93	■ 0.948	$f: c \rightarrow d$	$f: c \rightarrow d$	to denote elements $f \in \mathcal{C}(c, d)$ as $f: c \rightarrow d$. Here, c is called the <i>domain</i> of f , and d is
fong-spivak-seven-sketches_1 F01048	93	■ 0.620	$\text{id}_c; f = f$	$\text{id}_c; f = f$	does nothing: $\text{id}_c \circ f = f$ and $f \circ \text{id}_d = f$.
fong-spivak-seven-sketches_1 F01049	93	■ 0.620	$f ; \text{id}_d = f$	$f; \text{id}_d = f$	does nothing: $\text{id}_c \circ f = f$ and $f \circ \text{id}_d = f$.
fong-spivak-seven-sketches_1 F01050	93	■ 1.000	$f: c_0 \rightarrow c_1, g: c_1 \rightarrow c_2$	$f: c_0 \rightarrow c_1, g: c_1 \rightarrow c_2$	(b) <i>associativity</i> : for any three morphisms $f: c_0 \rightarrow c_1, g: c_1 \rightarrow c_2$, and $h: c_2 \rightarrow c_3$,
fong-spivak-seven-sketches_1 F01051	93	■ 1.000	$h: c_2 \rightarrow c_3$	$h: c_2 \rightarrow c_3$	(b) <i>associativity</i> : for any three morphisms $f: c_0 \rightarrow c_1, g: c_1 \rightarrow c_2$, and $h: c_2 \rightarrow c_3$,
fong-spivak-seven-sketches_1 F01052	93	■ 1.000	$(f ; g) ; h = f ; (g ; h)$	$(f;g);h = f;(g;h)$	the following are equal: $(f \circ g) \circ h = f \circ (g \circ h)$. We write this composite simply
fong-spivak-seven-sketches_1 F01053	93	■ 0.947	$f ; g ; h$	$f;g;h$	as $f \circ g \circ h$.
fong-spivak-seven-sketches_1 F01054	93	■ 0.998	$\mathcal{C}(c, d)$	$\mathcal{C}(c, d)$	³ This set $\mathcal{C}(c, d)$ is often denoted $\text{Hom}_{\mathcal{C}}(c, d)$, and called the “hom-set from c to d .” The word “hom”
fong-spivak-seven-sketches_1 F01055	93	■ 0.998	$\text{Hom}_{\mathcal{C}}(c, d)$	$\text{Hom}_{\mathcal{C}}(c, d)$	³ This set $\mathcal{C}(c, d)$ is often denoted $\text{Hom}_{\mathcal{C}}(c, d)$, and called the “hom-set from c to d .” The word “hom”
fong-spivak-seven-sketches_1 F01056	94	■ 1.000	$p: v \rightarrow w$	$p: v \rightarrow w$	write <u>$p: v \rightarrow w$</u> . To every vertex v , there is a trivial path, containing no arrows, starting
fong-spivak-seven-sketches_1 F01057	94	■ 0.999	id_v	id_v	and ending at v ; we often denote it by id_v or simply by v . We may also concatenate
fong-spivak-seven-sketches_1 F01058	94	■ 1.000	$q: w \rightarrow x$	$q: w \rightarrow x$	paths: given $p: v \rightarrow w$ and <u>$q: w \rightarrow x$</u> , their concatenation is denoted $p \circ q$, and it goes
fong-spivak-seven-sketches_1 F01059	94	■ 1.000	$p ; q$	$p; q$	paths: given $p: v \rightarrow w$ and $q: w \rightarrow x$, their concatenation is denoted $p \circ q$, and it goes
fong-spivak-seven-sketches_1 F01060	94	■ 1.000	$v \rightarrow x$	$v \rightarrow x$	<u>$v \rightarrow x$</u> .
fong-spivak-seven-sketches_1 F01061	94	■ 0.980	$(V, \leq)$	$(V, \leq)$	In Chapter 1, we used graphs to depict preorders $(V, \leq)$: the vertices form the
fong-spivak-seven-sketches_1 F01062	94	■ 0.999	$\text{Free}(G)$	$\text{Free}(G)$	Definition 3.7. For any graph $G = (V, A, s, t)$, we can define a category Free (G), called
fong-spivak-seven-sketches_1 F01063	94	■ 0.754	v_1	v_1	It has two objects v_1 and v_2 , and three morphisms: $\text{id}_{v_1}: v_1 \rightarrow v_1, f_1: v_1 \rightarrow v_2$, and
fong-spivak-seven-sketches_1 F01064	94	■ 0.754	v_2	v_2	It has two objects v_1 and v_2 , and three morphisms: $\text{id}_{v_1}: v_1 \rightarrow v_1, f_1: v_1 \rightarrow v_2$, and
fong-spivak-seven-sketches_1 F01065	94	■ 0.754	$\text{id}_{v_1}: v_1 \rightarrow v_1, f_1: v_1 \rightarrow v_2, \text{id}_{v_2}: v_2 \rightarrow v_2$	$\text{id}_{v_1}: v_1 \rightarrow v_1, f_1: v_1 \rightarrow v_2$	It has two objects v_1 and v_2 , and three morphisms: $\text{id}_{v_1}: v_1 \rightarrow v_1, f_1: v_1 \rightarrow v_2$, and
fong-spivak-seven-sketches_1 F01066	94	■ 0.607	$\text{id}_{v_2}: v_2 \rightarrow v_2$	$\text{id}_{v_2}: v_2 \rightarrow v_2$	$\text{id}_{v_2}: v_2 \rightarrow v_2$. Here id_{v_1} is the path of length 0 starting and ending at v_1, f_1 is the path
fong-spivak-seven-sketches_1 F01067	94	■ 0.607	id_{v_1}	id_{v_1}	$\text{id}_{v_2}: v_2 \rightarrow v_2$. Here id_{v_1} is the path of length 0 starting and ending at v_1, f_1 is the path
fong-spivak-seven-sketches_1 F01068	94	■ 0.607	v_1, f_1	v_1, f_1	$\text{id}_{v_2}: v_2 \rightarrow v_2$. Here id_{v_1} is the path of length 0 starting and ending at v_1, f_1 is the path
fong-spivak-seven-sketches_1 F01069	94	■ 0.997	f_1	f_1	of length 1 consisting of just the arrow f_1 , and id_{v_2} is the length 0 path at v_2 . As our
fong-spivak-seven-sketches_1 F01070	94	■ 0.997	id_{v_2}	id_{v_2}	of length 1 consisting of just the arrow f_1 , and id_{v_2} is the length 0 path at v_2 . As our
fong-spivak-seven-sketches_1 F01071	94	■ 0.562	$f_1 = f_1$	$f_1 = f_1$	for v_2 . As composition is given by concatenation, we have, for example $\text{id}_{v_1} \circ f_1 = f_1$,

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_94 F01072	94	0.524	$\mathrm{id}_{\{v_2\}}\{ \}_9 \mathrm{id}_{\{v_2\}} = \mathrm{id}_{\{v_2\}}$	$\mathrm{id}_{v_2} \mathrm{id}_{v_2} = \mathrm{id}_{v_2}$	$\mathrm{id}_{v_2} \circ \mathrm{id}_{v_2} = \mathrm{id}_{v_2}$, and so on.
fong-spivak-seven-sketches_95 F01073	95	0.621	$\mathbf{0}$	$\mathbf{0}$	2. What is the category $\mathbf{0}$?
fong-spivak-seven-sketches_95 F01074	95	1.000	$\mathbf{n}$	$\mathbf{n}$	3. What is the formula for the number of morphisms in $\mathbf{n}$ for arbitrary $n \in \mathbb{N}$?
fong-spivak-seven-sketches_95 F01075	95	0.968	$\{z, s, (s ; s), (s ; s ; s), \ldots\}$	$\{z, s, (s; s), (s; s; s), \dots\}$	$\{z, s, (s \circ s), (s \circ s \circ s), \dots\}$, where we write z for the length 0 path on z ; it represents
fong-spivak-seven-sketches_95 F01076	95	0.967	$\mathrm{id}_{\{z\}}$	id_z	the morphism id_z . There is a one-to-one correspondence between Path and the natural
fong-spivak-seven-sketches_95 F01077	95	1.000	$\mathbb{N} = \{0, 1, 2, 3, \ldots\}$	$\mathbb{N} = \{0, 1, 2, 3, \dots\}$	numbers, $\mathbb{N} = \{0, 1, 2, 3, \dots\}$.
fong-spivak-seven-sketches_95 F01078	95	0.966	$*: M \times M \rightarrow M$	$*: M \times M \rightarrow M$	a tuple $(M, *, e)$ where $*: M \times M \rightarrow M$ is a function and $e \in M$ is an element, and
fong-spivak-seven-sketches_95 F01079	95	1.000	$m * 1 = m = 1 * m$	$m * 1 = m = 1 * m$	$m * 1 = m = 1 * m$ and $(m * n) * p = m * (n * p)$.
fong-spivak-seven-sketches_95 F01080	95	1.000	$(m * n) * p = m * (n * p)$	$(m * n) * p = m * (n * p)$	$m * 1 = m = 1 * m$ and $(m * n) * p = m * (n * p)$.
fong-spivak-seven-sketches_95 F01081	95	0.851	$M := \mathcal{C}(\bullet, \bullet)$	$M := \mathcal{C}(\bullet, \bullet)$	connection. Given a category $\mathcal{C}$ with one object, say $\bullet$, let $M := \mathcal{C}(\bullet, \bullet)$, let $e = \mathrm{id}_\bullet$, and
fong-spivak-seven-sketches_95 F01082	95	0.851	$e = \mathrm{id}_{\{\bullet\}}$	$e = \mathrm{id}_\bullet$	connection. Given a category $\mathcal{C}$ with one object, say $\bullet$, let $M := \mathcal{C}(\bullet, \bullet)$, let $e = \mathrm{id}_\bullet$, and
fong-spivak-seven-sketches_95 F01083	95	0.922	$*: \mathcal{C}(\bullet, \bullet) \times \mathcal{C}(\bullet, \bullet) \rightarrow \mathcal{C}(\bullet, \bullet)$	$*: \mathcal{C}(\bullet, \bullet) \times \mathcal{C}(\bullet, \bullet) \rightarrow \mathcal{C}(\bullet, \bullet)$	let $*: \mathcal{C}(\bullet, \bullet) \times \mathcal{C}(\bullet, \bullet) \rightarrow \mathcal{C}(\bullet, \bullet)$ be the composition operation $* = \circ$. The associativity
fong-spivak-seven-sketches_95 F01084	95	0.922	$* = \circ$	$* = \circ$	let $*: \mathcal{C}(\bullet, \bullet) \times \mathcal{C}(\bullet, \bullet) \rightarrow \mathcal{C}(\bullet, \bullet)$ be the composition operation $* = \circ$. The associativity
fong-spivak-seven-sketches_95 F01085	95	1.000	$m, n \in \mathbb{N}$	$m, n \in \mathbb{N}$	numbers $n \in \mathbb{N}$. Paths can be concatenated. Given numbers $m, n \in \mathbb{N}$, what number
fong-spivak-seven-sketches_96 F01086	96	0.920	$f \circ h$	$f \circ h$	because $f \circ h$ and $g \circ i$ are made equal. This category is called the “commutative square.”
fong-spivak-seven-sketches_96 F01087	96	0.920	$g \circ i$	$g \circ i$	because $f \circ h$ and $g \circ i$ are made equal. This category is called the “commutative square.”
fong-spivak-seven-sketches_96 F01088	96	0.547	$g \circ i$	$g \circ i$	One might say “the missing one is $g \circ i$,” but that is not quite right: $g \circ i$ is there too,
fong-spivak-seven-sketches_96 F01089	96	0.795	$f ; h$	$f; h$	because it is equal to $f \circ h$. As usual, A denotes id_A , etc.
fong-spivak-seven-sketches_96 F01090	96	0.795	id_A	id_A	because it is equal to $f \circ h$. As usual, A denotes id_A , etc.
fong-spivak-seven-sketches_97 F01091	97	0.499	$s ; s = z$	$s; s = z$	In $\mathcal{C}$ we have the equation $s \circ s = z$. But this implies $s \circ s \circ s = z \circ s = s$! And similarly
fong-spivak-seven-sketches_97 F01092	97	0.499	$s ; s ; s = z ; s = s$	$s; s; s = z; s = s$	In $\mathcal{C}$ we have the equation $s \circ s = z$. But this implies $s \circ s \circ s = z \circ s = s$! And similarly
fong-spivak-seven-sketches_97 F01093	97	0.493	$; s = z$	$; s = z$	we have $s \circ s \circ s \circ s = z \circ z = z$. The set of morphisms in $\mathcal{C}$ is in fact merely $\{z, s\}$, with
fong-spivak-seven-sketches_97 F01094	97	0.493	$= z$	$= z$	we have $s \circ s \circ s \circ s = z \circ z = z$. The set of morphisms in $\mathcal{C}$ is in fact merely $\{z, s\}$, with
fong-spivak-seven-sketches_97 F01095	97	0.493	$\{z, s\}$	$\{z, s\}$	we have $s \circ s \circ s \circ s = z \circ z = z$. The set of morphisms in $\mathcal{C}$ is in fact merely $\{z, s\}$, with
fong-spivak-seven-sketches_97 F01096	97	0.556	$z = z$	$z = z$	composition described by $s \circ s = z \circ z = z$, and $z \circ s = s \circ z = s$. In group theory, one
fong-spivak-seven-sketches_97 F01097	97	0.556	$z ; s = s ; z = s$	$z; s = s; z = s$	composition described by $s \circ s = z \circ z = z$, and $z \circ s = s \circ z = s$. In group theory, one
fong-spivak-seven-sketches_97 F01098	97	1.000	$\mathbb{Z} / 2 \mathbb{Z}$	$\mathbb{Z} / 2\mathbb{Z}$	would speak of a group called $\mathbb{Z} / 2\mathbb{Z}$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F011099	97	■ 1.000	$\mathcal{D}$	$\mathcal{D}$	<i>Exercise 3.19.</i> Write down all the morphisms in the category $\mathcal{D}$ from Example 3.18.
fong-spivak-seven-sketches-F011100	97	■ 1.000	$\operatorname{Ob}(\mathcal{P}) = P$	$\operatorname{Ob}(\mathcal{P}) = P$	follows. The objects of $\mathcal{P}$ are precisely the elements of P ; that is, $\operatorname{Ob}(\mathcal{P}) = P$. As for
fong-spivak-seven-sketches-F011101	97	■ 1.000	$p \rightarrow q$	$p \rightarrow q$	morphisms, $\mathcal{P}$ has exactly one morphism $p \rightarrow q$ if $p \leq q$ and no morphisms $p \rightarrow q$ if
fong-spivak-seven-sketches-F011102	97	■ 1.000	$p \not\leq q$	$p \not\leq q$	$p \not\leq q$. The fact that $\leq$ is reflexive ensures that every object has an identity, and the fact
fong-spivak-seven-sketches-F011103	98	■ 1.000	$(C, \leq)$	$(C, \leq)$	$(C, \leq)$ from it by destroying the distinction between any two parallel morphisms. That
fong-spivak-seven-sketches-F011104	98	■ 0.985	$C := \operatorname{Ob}(\mathcal{C})$	$C := \operatorname{Ob}(\mathcal{C})$	is, let $C := \operatorname{Ob}(\mathcal{C})$, and put $c_1 \leq c_2$ iff $\mathcal{C}(c_1, c_2) \neq \emptyset$. If there is one, or two, or fifty,
fong-spivak-seven-sketches-F011105	98	■ 0.985	$c_1 \leq c_2$	$c_1 \leq c_2$	is, let $C := \operatorname{Ob}(\mathcal{C})$, and put $c_1 \leq c_2$ iff $\mathcal{C}(c_1, c_2) \neq \emptyset$. If there is one, or two, or fifty,
fong-spivak-seven-sketches-F011106	98	■ 0.985	$\mathcal{C}(c_1, c_2) \neq \varnothing$	$\mathcal{C}(c_1, c_2) \neq \emptyset$	is, let $C := \operatorname{Ob}(\mathcal{C})$, and put $c_1 \leq c_2$ iff $\mathcal{C}(c_1, c_2) \neq \emptyset$. If there is one, or two, or fifty,
fong-spivak-seven-sketches-F011107	98	■ 1.000	$c_1 \rightarrow c_2$	$c_1 \rightarrow c_2$	or infinitely many morphisms $c_1 \rightarrow c_2$ in $\mathcal{C}$, the preorder reflection does not see the
fong-spivak-seven-sketches-F011108	99	■ 0.810	$\operatorname{Ob}(\mathcal{C})$	$\operatorname{Ob}(\mathcal{C})$	(i) $\operatorname{Ob}(\mathbf{Set})$ is the collection of all sets.
fong-spivak-seven-sketches-F011109	99	■ 0.998	$\mathbf{Set}(S, T) = \{f : S \rightarrow T \mid f \text{ is a function}\}$	$\mathbf{Set}(S, T) = \{f : S \rightarrow T \mid f \text{ is a function}\}$	(ii) If S and T are sets, then $\mathbf{Set}(S, T) = \{f : S \rightarrow T \mid f \text{ is a function}\}$.
fong-spivak-seven-sketches-F011110	99	■ 0.958	$\operatorname{id}_S : S \rightarrow S$	$\operatorname{id}_S : S \rightarrow S$	(iii) For each set S , the identity morphism is the function $\operatorname{id}_S : S \rightarrow S$ given by
fong-spivak-seven-sketches-F011111	99	■ 0.908	$\operatorname{id}_S(s) := s$	$\operatorname{id}_S(s) := s$	$\operatorname{id}_S(s) := s$ for each $s \in S$.
fong-spivak-seven-sketches-F011112	99	■ 1.000	$f : S \rightarrow T$	$f : S \rightarrow T$	(iv) Given $f : S \rightarrow T$ and $g : T \rightarrow U$, their composite is the function $f \circ g : S \rightarrow U$
fong-spivak-seven-sketches-F011113	99	■ 1.000	$g : T \rightarrow U$	$g : T \rightarrow U$	(iv) Given $f : S \rightarrow T$ and $g : T \rightarrow U$, their composite is the function $f \circ g : S \rightarrow U$
fong-spivak-seven-sketches-F011114	99	■ 1.000	$f \circ g : S \rightarrow U$	$f \circ g : S \rightarrow U$	(iv) Given $f : S \rightarrow T$ and $g : T \rightarrow U$, their composite is the function $f \circ g : S \rightarrow U$
fong-spivak-seven-sketches-F011115	99	■ 1.000	$(f \circ g)(s) := g(f(s))$	$(f \circ g)(s) := g(f(s))$	given by $(f \circ g)(s) := g(f(s))$.
fong-spivak-seven-sketches-F011116	99	■ 1.000	$\underline{2} = \{1, 2\}$	$\underline{2} = \{1, 2\}$	<i>Exercise 3.25.</i> Let $\underline{2} = \{1, 2\}$ and $\underline{3} = \{1, 2, 3\}$. These are objects in the category $\mathbf{Set}$
fong-spivak-seven-sketches-F011117	99	■ 1.000	$\underline{3} = \{1, 2, 3\}$	$\underline{3} = \{1, 2, 3\}$	<i>Exercise 3.25.</i> Let $\underline{2} = \{1, 2\}$ and $\underline{3} = \{1, 2, 3\}$. These are objects in the category $\mathbf{Set}$
fong-spivak-seven-sketches-F011118	99	■ 0.952	$\mathbf{Set}(\underline{2}, \underline{3})$	$\mathbf{Set}(\underline{2}, \underline{3})$	discussed in Definition 3.24. Write down all the elements of the set $\mathbf{Set}(\underline{2}, \underline{3})$; there
fong-spivak-seven-sketches-F011119	100	■ 0.984	$\mathcal{C}^{\operatorname{op}}$	$\mathcal{C}^{\operatorname{op}}$	<i>Example 3.27.</i> Let $\mathcal{C}$ be a category. Its <i>opposite</i> , denoted $\mathcal{C}^{\operatorname{op}}$, is the category with
fong-spivak-seven-sketches-F011120	100	■ 0.268	$\operatorname{Ob}(\mathcal{C}^{\operatorname{op}}) := \operatorname{Ob}(\mathcal{C})$	$\operatorname{Ob}(\mathcal{C}^{\operatorname{op}}) := \operatorname{Ob}(\mathcal{C})$	the same objects, $\operatorname{Ob}(\mathcal{C}^{\operatorname{op}}) := \operatorname{Ob}(\mathcal{C})$, and for any two objects $c, d \in \operatorname{Ob}(\mathcal{C})$, one has
fong-spivak-seven-sketches-F011121	100	■ 0.984	$\mathcal{C}^{\operatorname{op}}(c, d) := \mathcal{C}(d, c)$	$\mathcal{C}^{\operatorname{op}}(c, d) := \mathcal{C}(d, c)$	$\mathcal{C}^{\operatorname{op}}(c, d) := \mathcal{C}(d, c)$. Identities and composition are as in $\mathcal{C}$.
fong-spivak-seven-sketches-F011122	100	■ 0.980	$\{\mathbf{m}, \square\}$	$\{\mathbf{m}, \square\}$	example, in the category $\mathbf{Set}$, one can exchange the set $\{\mathbf{m}, \square\}$ for the set $\{0, 1\}$ and
fong-spivak-seven-sketches-F011123	100	■ 0.980	$\{0, 1\}$	$\{0, 1\}$	example, in the category $\mathbf{Set}$, one can exchange the set $\{\mathbf{m}, \square\}$ for the set $\{0, 1\}$ and
fong-spivak-seven-sketches-F011124	100	■ 0.999	$a \cong b$	$a \cong b$	has a preorder with elements a, b , such that $a \leq b$ and $b \leq a$, i.e. $a \cong b$, then a and b are
fong-spivak-seven-sketches-F011125	100	■ 1.000	$c \leq a$	$c \leq a$	know that $c \leq a$ iff $c \leq b$, and $c \geq a$ iff $c \geq b$. The notion of isomorphism formalizes

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_100 F01126	100	■ 1.000	$c \leq b$	$c \leq b$	know that $c \leq a$ iff $c \leq b$, and $c \geq a$ iff $c \geq b$. The notion of isomorphism formalizes
fong-spivak-seven-sketches_100 F01127	100	■ 1.000	$c \geq a$	$c \geq a$	know that $c \leq a$ iff $c \leq b$, and $c \geq a$ iff $c \geq b$. The notion of isomorphism formalizes
fong-spivak-seven-sketches_100 F01128	100	■ 1.000	$c \geq b$	$c \geq b$	know that $c \leq a$ iff $c \leq b$, and $c \geq a$ iff $c \geq b$. The notion of isomorphism formalizes
fong-spivak-seven-sketches_100 F01129	100	■ 0.931	$g: B \rightarrow A$	$g: B \rightarrow A$	morphism $g: B \rightarrow A$ satisfying $f \circ g = \text{id}_A$ and $g \circ f = \text{id}_B$. In this case we call f and g
fong-spivak-seven-sketches_100 F01130	100	■ 0.931	$f; g = \text{id}_A$	$f; g = \text{id}_A$	morphism $g: B \rightarrow A$ satisfying $f \circ g = \text{id}_A$ and $g \circ f = \text{id}_B$. In this case we call f and g
fong-spivak-seven-sketches_100 F01131	100	■ 0.931	$g; f = \text{id}_B$	$g; f = \text{id}_B$	morphism $g: B \rightarrow A$ satisfying $f \circ g = \text{id}_A$ and $g \circ f = \text{id}_B$. In this case we call f and g
fong-spivak-seven-sketches_100 F01132	100	■ 1.000	$g = f^{-1}$	$g = f^{-1}$	<i>inverses</i> , and we often write $g = f^{-1}$, or equivalently $f = g^{-1}$. We also say that A and B
fong-spivak-seven-sketches_100 F01133	100	■ 1.000	$f = g^{-1}$	$f = g^{-1}$	<i>inverses</i> , and we often write $g = f^{-1}$, or equivalently $f = g^{-1}$. We also say that A and B
fong-spivak-seven-sketches_100 F01134	100	■ 1.000	$A := \{a, b, c\}$	$A := \{a, b, c\}$	<i>Example 3.29.</i> The set $A := \{a, b, c\}$ and the set $\underline{3} = \{1, 2, 3\}$ are isomorphic; that is,
fong-spivak-seven-sketches_100 F01135	100	■ 1.000	$f: A \rightarrow \underline{3}$	$f: A \rightarrow \underline{3}$	there exists an isomorphism $f: A \rightarrow \underline{3}$ given by $f(a) = 2, f(b) = 1, f(c) = 3$. The
fong-spivak-seven-sketches_100 F01136	100	■ 1.000	$f(a)=2, f(b)=1, f(c)=3$	$f(a) = 2, f(b) = 1, f(c) = 3$	there exists an isomorphism $f: A \rightarrow \underline{3}$ given by $f(a) = 2, f(b) = 1, f(c) = 3$. The
fong-spivak-seven-sketches_100 F01137	100	■ 0.954	$A \in$	$A \in$	understood in terms of isomorphisms in FinSet . Namely, for any finite set $A \in \mathbf{FinSet}$,
fong-spivak-seven-sketches_100 F01138	100	■ 1.000	$A \cong \underline{n}$	$A \cong \underline{n}$	its cardinality is the number $n \in \mathbb{N}$ such that there exists an isomorphism $A \cong \underline{n}$. Georg
fong-spivak-seven-sketches_100 F01139	100	■ 1.000	$f^{-1}: \underline{3} \rightarrow A$	$f^{-1}: \underline{3} \rightarrow A$	1. What is the inverse $f^{-1}: \underline{3} \rightarrow A$ of the function f given in Example 3.29?
fong-spivak-seven-sketches_100 F01140	100	■ 0.998	$A \rightarrow \underline{3}$	$A \rightarrow \underline{3}$	2. How many distinct isomorphisms are there $A \rightarrow \underline{3}$?
fong-spivak-seven-sketches_100 F01141	100	■ 1.000	id_c	id_c	identity id_c is an isomorphism.
fong-spivak-seven-sketches_101 F01142	101	■ 0.880	$\mathbf{Free}(G)$	$\mathbf{Free}(G)$	<i>Exercise 3.33.</i> Let G be a graph, and let $\mathbf{Free}(G)$ be the corresponding free category.
fong-spivak-seven-sketches_101 F01143	101	■ 1.000	$f: \underline{2} \rightarrow \underline{3}$	$f: \underline{2} \rightarrow \underline{3}$	Consider the functions $f: \underline{2} \rightarrow \underline{3}$ and $g: \underline{3} \rightarrow \underline{2}$ drawn below:
fong-spivak-seven-sketches_101 F01144	101	■ 1.000	$g: \underline{3} \rightarrow \underline{2}$	$g: \underline{3} \rightarrow \underline{2}$	Consider the functions $f: \underline{2} \rightarrow \underline{3}$ and $g: \underline{3} \rightarrow \underline{2}$ drawn below:
fong-spivak-seven-sketches_101 F01145	101	■ 0.330	$f; g = \text{id}_{\underline{2}}$	$f; g = \text{id}_{\underline{2}}$	Then the reader should be able to instantly check that $f \circ g = \text{id}_{\underline{2}}$ but $g \circ f \neq \text{id}_{\underline{3}}$. Thus f
fong-spivak-seven-sketches_101 F01146	101	■ 0.330	$g \circ f \neq \text{id}_{\underline{3}}$	$g \circ f \neq \text{id}_{\underline{3}}$	Then the reader should be able to instantly check that $f \circ g = \text{id}_{\underline{2}}$ but $g \circ f \neq \text{id}_{\underline{3}}$. Thus f
fong-spivak-seven-sketches_103 F01147	103	■ 0.998	$F: \mathcal{C} \rightarrow \mathcal{D}$	$F: \mathcal{C} \rightarrow \mathcal{D}$	$F: \mathcal{C} \rightarrow \mathcal{D}$,
fong-spivak-seven-sketches_103 F01148	103	■ 0.989	$F(c) \in \text{Ob}(\mathcal{D})$	$F(c) \in \text{Ob}(\mathcal{D})$	(i) for every object $c \in \text{Ob}(\mathcal{C})$, one specifies an object $F(c) \in \text{Ob}(\mathcal{D})$;
fong-spivak-seven-sketches_103 F01149	103	■ 0.203	$f: c_1 \rightarrow c_2$	$f: c_1 \rightarrow c_2$	(ii) for every morphism $f: c_1 \rightarrow c_2$ in $\mathcal{C}$, one specifies a morphism $F(f): F(c_1) \rightarrow$
fong-spivak-seven-sketches_103 F01150	103	■ 0.203	$\mathcal{C}_{\text{one}}$	$\mathcal{C}_{\text{one}}$	(ii) for every morphism $f: c_1 \rightarrow c_2$ in $\mathcal{C}$, one specifies a morphism $F(f): F(c_1) \rightarrow$
fong-spivak-seven-sketches_103 F01151	103	■ 0.203	$F(f): F(c_1) \rightarrow$	$F(f): F(c_1) \rightarrow$	(ii) for every morphism $f: c_1 \rightarrow c_2$ in $\mathcal{C}$, one specifies a morphism $F(f): F(c_1) \rightarrow$
fong-spivak-seven-sketches_103 F01152	103	■ 1.000	$F(c_2)$	$F(c_2)$	$F(c_2)$ in $\mathcal{D}$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_103 F01153	103	0.628	$F(\mathrm{id}_c)=\mathrm{id}_{F(c)}$	$F(\mathrm{id}_c) = \mathrm{id}_{F(c)}$	(a) for every object $c \in \mathrm{Ob}(\mathcal{C})$, we have $F(\mathrm{id}_c) = \mathrm{id}_{F(c)}$.
fong-spivak-seven-sketches_103 F01154	103	0.870	$c_1, c_2, c_3 \in \operatorname{Ob}(\mathcal{C})$	$c_1, c_2, c_3 \in \mathrm{Ob}(\mathcal{C})$	(b) for every three objects $c_1, c_2, c_3 \in \mathrm{Ob}(\mathcal{C})$ and two morphisms $f \in \mathcal{C}(c_1, c_2)$, $g \in$
fong-spivak-seven-sketches_103 F01155	103	0.870	$f \in \mathcal{C}(c_1, c_2), g \in$	$f \in \mathcal{C}(c_1, c_2), g \in$	(b) for every three objects $c_1, c_2, c_3 \in \mathrm{Ob}(\mathcal{C})$ and two morphisms $f \in \mathcal{C}(c_1, c_2)$, $g \in$
fong-spivak-seven-sketches_103 F01156	103	1.000	$\mathcal{C}(c_2, c_3)$	$\mathcal{C}(c_2, c_3)$	$\mathcal{C}(c_2, c_3)$, the equation $F(f \circ g) = F(f) \circ F(g)$ holds in $\mathcal{D}$.
fong-spivak-seven-sketches_103 F01157	103	1.000	$F(f \circ g)=F(f) \circ F(g)$	$F(f \circ g) = F(f) \circ F(g)$	$\mathcal{C}(c_2, c_3)$, the equation $F(f \circ g) = F(f) \circ F(g)$ holds in $\mathcal{D}$.
fong-spivak-seven-sketches_103 F01158	103	0.975	$F: 2 \rightarrow 3$	$F: 2 \rightarrow 3$	Example 3.36. For example, here we draw three functors $F: 2 \rightarrow 3$:
fong-spivak-seven-sketches_103 F01159	103	1.000	n_0	n_0	path to the trivial path, i.e. the identity on n_0 . In the middle diagram $F(m_0) = n_0$,
fong-spivak-seven-sketches_103 F01160	103	1.000	$F(m_0)=n_0$	$F(m_0) = n_0$	path to the trivial path, i.e. the identity on n_0 . In the middle diagram $F(m_0) = n_0$,
fong-spivak-seven-sketches_103 F01161	103	1.000	$F(f_1)=g_1$	$F(f_1) = g_1$	$F(f_1) = g_1$, and $F(m_1) = n_1$. In the right-hand diagram, $F(m_0) = n_0$, $F(m_1) = n_2$, and
fong-spivak-seven-sketches_103 F01162	103	1.000	$F(m_1)=n_1$	$F(m_1) = n_1$	$F(f_1) = g_1$, and $F(m_1) = n_1$. In the right-hand diagram, $F(m_0) = n_0$, $F(m_1) = n_2$, and
fong-spivak-seven-sketches_103 F01163	103	1.000	$F(m_0)=n_0, F(m_1)=n_2$	$F(m_0) = n_0, F(m_1) = n_2$	$F(f_1) = g_1$, and $F(m_1) = n_1$. In the right-hand diagram, $F(m_0) = n_0$, $F(m_1) = n_2$, and
fong-spivak-seven-sketches_103 F01164	103	0.996	$F(f_1)=g_1 \circ g_2$	$F(f_1) = g_1 \circ g_2$	$F(f_1) = g_1 \circ g_2$.
fong-spivak-seven-sketches_103 F01165	103	0.772	$2 \rightarrow 3$	$2 \rightarrow 3$	the remaining functors $2 \rightarrow 3$.
fong-spivak-seven-sketches_104 F01166	104	1.000	$\mathcal{F}$	$\mathcal{F}$	There are lots of functors from the free square category (let's call it $\mathcal{F}$) to the commutative
fong-spivak-seven-sketches_104 F01167	104	1.000	$F: \mathcal{F} \rightarrow \mathcal{C}$	$F: \mathcal{F} \rightarrow \mathcal{C}$	However, there is exactly one functor $F: \mathcal{F} \rightarrow \mathcal{C}$ that sends A' to A , B' to B , C' to C ,
fong-spivak-seven-sketches_104 F01168	104	1.000	$A, B^{\prime}$	A, B'	However, there is exactly one functor $F: \mathcal{F} \rightarrow \mathcal{C}$ that sends A' to A , B' to B , C' to C ,
fong-spivak-seven-sketches_104 F01169	104	1.000	$B, C^{\prime}$	B, C'	However, there is exactly one functor $F: \mathcal{F} \rightarrow \mathcal{C}$ that sends A' to A , B' to B , C' to C ,
fong-spivak-seven-sketches_104 F01170	104	1.000	$D^{\prime}$	D'	and D' to D . That is, once we have made this decision about how F acts on objects,
fong-spivak-seven-sketches_104 F01171	104	0.994	$\mathcal{C}=\bullet \rightarrow \bullet$	$\mathcal{C} = \bullet \rightarrow \bullet$	Exercise 3.40. Consider the free categories $\mathcal{C} = \boxed{\bullet \rightarrow \bullet}$ and $\mathcal{D} = \boxed{\bullet \rightrightarrows \bullet}$. Give two
fong-spivak-seven-sketches_104 F01172	104	0.994	$\mathcal{D}=\bullet \rightrightarrows \bullet$	$\mathcal{D} = \bullet \rightrightarrows \bullet$	Exercise 3.40. Consider the free categories $\mathcal{C} = \boxed{\bullet \rightarrow \bullet}$ and $\mathcal{D} = \boxed{\bullet \rightrightarrows \bullet}$. Give two
fong-spivak-seven-sketches_104 F01173	104	1.000	$F, G: \mathcal{C} \rightarrow \mathcal{D}$	$F, G: \mathcal{C} \rightarrow \mathcal{D}$	functors $F, G: \mathcal{C} \rightarrow \mathcal{D}$ that act the same on objects but differently on morphisms.
fong-spivak-seven-sketches_104 F01174	104	0.979	$A^{\prime}, B$	A', B	to the free square category $\mathcal{F}$, but <i>none</i> that sends A to A' , B to B' , C to C' , and D to D' .
fong-spivak-seven-sketches_104 F01175	104	0.979	$B^{\prime}, C$	B', C	to the free square category $\mathcal{F}$, but <i>none</i> that sends A to A' , B to B' , C to C' , and D to D' .
fong-spivak-seven-sketches_104 F01176	104	0.979	$C^{\prime}$	C'	to the free square category $\mathcal{F}$, but <i>none</i> that sends A to A' , B to B' , C to C' , and D to D' .
fong-spivak-seven-sketches_104 F01177	104	0.352	$f^{\circ}; h=g^{\circ} i$	$f^{\circ}; h = g^{\circ} i$	The reason is that if F were such a functor, then since $f \circ g = g \circ i$ in $\mathcal{C}$, we would have
fong-spivak-seven-sketches_104 F01178	104	1.000	$F(f \circ h)=F(g \circ i)$	$F(f \circ h) = F(g \circ i)$	$F(f \circ h) = F(g \circ i)$, but then the rules of functors would let us reason as follows:
fong-spivak-seven-sketches_104 F01179	104	0.932	$f^{\prime}; h^{\prime}=g^{\prime} i^{\prime}$	$f'; h' = g' i'$	The resulting equation, $f' \circ h' = g' \circ i'$ does not hold in $\mathcal{F}$ because it is a free category

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_104 F01180	104	■ 0.999	$\mathcal{N}$	$\mathcal{N}$	For example, consider the preorder $(\mathbb{N}, \leq)$ considered as a category $\mathcal{N}$ with objects
fong-spivak-seven-sketches_104 F01181	104	■ 1.000	$\operatorname{Ob}(\mathcal{N})=\mathbb{N}$	$\operatorname{Ob}(\mathcal{N}) = \mathbb{N}$	$\operatorname{Ob}(\mathcal{N}) = \mathbb{N}$ and a unique morphism $m \rightarrow n$ iff $m \leq n$. A functor $F: \mathcal{N} \rightarrow \mathcal{N}$ sends
fong-spivak-seven-sketches_104 F01182	104	■ 1.000	$m \rightarrowtail n$	$m \rightarrow n$	$\operatorname{Ob}(\mathcal{N}) = \mathbb{N}$ and a unique morphism $m \rightarrow n$ iff $m \leq n$. A functor $F: \mathcal{N} \rightarrow \mathcal{N}$ sends
fong-spivak-seven-sketches_104 F01183	104	■ 1.000	$F: \mathcal{N} \rightarrow \mathcal{N}$	$F: \mathcal{N} \rightarrow \mathcal{N}$	$\operatorname{Ob}(\mathcal{N}) = \mathbb{N}$ and a unique morphism $m \rightarrow n$ iff $m \leq n$. A functor $F: \mathcal{N} \rightarrow \mathcal{N}$ sends
fong-spivak-seven-sketches_104 F01184	104	■ 1.000	$F(n) \in \mathbb{N}$	$F(n) \in \mathbb{N}$	each object $n \in \mathbb{N}$ to an object $F(n) \in \mathbb{N}$. It must send morphisms in $\mathcal{N}$ to morphisms
fong-spivak-seven-sketches_104 F01185	104	■ 1.000	$F(m) \rightarrowtail F(n)$	$F(m) \rightarrow F(n)$	$F(m) \rightarrow F(n)$. In other words, if $m \leq n$, then we had better have $F(m) \leq F(n)$. But as
fong-spivak-seven-sketches_104 F01186	104	■ 1.000	$F(m) \leq F(n)$	$F(m) \leq F(n)$	$F(m) \rightarrow F(n)$. In other words, if $m \leq n$, then we had better have $F(m) \leq F(n)$. But as
fong-spivak-seven-sketches_105 F01187	105	■ 1.000	$\mathbb{N} \rightarrowtail \mathbb{N}$	$\mathbb{N} \rightarrow \mathbb{N}$	Thus a functor $F: \mathcal{N} \rightarrow \mathcal{N}$ and a monotone map $\mathbb{N} \rightarrow \mathbb{N}$ are the same thing.
fong-spivak-seven-sketches_105 F01188	105	■ 0.873	$\operatorname{id}_{\mathcal{C}}: \mathcal{C} \rightarrowtail \mathcal{C}$	$\operatorname{id}_{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{C}$	1. Given any category $\mathcal{C}$, show that there exists a functor $\operatorname{id}_{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{C}$, known as the
fong-spivak-seven-sketches_105 F01189	105	■ 1.000	$\operatorname{Ob}(\mathcal{C})$	$\operatorname{Ob}(\mathcal{C})$	Note that a functor $\mathcal{C} \rightarrow \mathcal{D}$ consists of a function from $\operatorname{Ob}(\mathcal{C})$ to $\operatorname{Ob}(\mathcal{D})$ and for each
fong-spivak-seven-sketches_105 F01190	105	■ 1.000	$\operatorname{Ob}(\mathcal{D})$	$\operatorname{Ob}(\mathcal{D})$	Note that a functor $\mathcal{C} \rightarrow \mathcal{D}$ consists of a function from $\operatorname{Ob}(\mathcal{C})$ to $\operatorname{Ob}(\mathcal{D})$ and for each
fong-spivak-seven-sketches_105 F01191	105	■ 0.987	$c_1, c_2 \in \mathcal{C}$	$c_1, c_2 \in \mathcal{C}$	pair of objects $c_1, c_2 \in \mathcal{C}$ a function from $\mathcal{C}(c_1, c_2)$ to $\mathcal{D}(F(c_1), F(c_2))$.
fong-spivak-seven-sketches_105 F01192	105	■ 0.987	$\mathcal{C}(c_1, c_2)$	$\mathcal{C}(c_1, c_2)$	pair of objects $c_1, c_2 \in \mathcal{C}$ a function from $\mathcal{C}(c_1, c_2)$ to $\mathcal{D}(F(c_1), F(c_2))$.
fong-spivak-seven-sketches_105 F01193	105	■ 0.987	$\mathcal{D}(F(c_1), F(c_2))$	$\mathcal{D}(F(c_1), F(c_2))$	pair of objects $c_1, c_2 \in \mathcal{C}$ a function from $\mathcal{C}(c_1, c_2)$ to $\mathcal{D}(F(c_1), F(c_2))$.
fong-spivak-seven-sketches_105 F01194	105	■ 0.982	$G: \mathcal{D} \rightarrowtail \mathcal{E}$	$G: \mathcal{D} \rightarrow \mathcal{E}$	2. Show that given $F: \mathcal{C} \rightarrow \mathcal{D}$ and $G: \mathcal{D} \rightarrow \mathcal{E}$, we can define a new functor $(F \circ$
fong-spivak-seven-sketches_105 F01195	105	■ 1.000	$G): \mathcal{C} \rightarrowtail \mathcal{E}$	$G): \mathcal{C} \rightarrow \mathcal{E}$	$G): \mathcal{C} \rightarrow \mathcal{E}$ just by composing functions.
fong-spivak-seven-sketches_105 F01196	105	■ 0.991	$F: \mathcal{C} \rightarrowtail$	$F: \mathcal{C} \rightarrow$	Let $\mathcal{C}$ be a category, and recall the category Set from Definition 3.24. A functor $F: \mathcal{C} \rightarrow$
fong-spivak-seven-sketches_105 F01197	105	■ 0.508	$I: \mathcal{C} \rightarrowtail$	$I: \mathcal{C} \rightarrow$	of the database—is given by a set-valued functor $I: \mathcal{C} \rightarrow \mathbf{Set}$. The only additional
fong-spivak-seven-sketches_105 F01198	105	■ 1.000	$c = \stackrel{\text{string}}{\circ}$	$c = \stackrel{\text{string}}{\circ}$	detail is that for any white node, such as $c = \stackrel{\text{string}}{\circ}$, we want to force I to map to the set
fong-spivak-seven-sketches_105 F01199	105	■ 0.954	5	5	functor $I: \mathcal{C} \rightarrow \mathbf{Set}$. ⁵
fong-spivak-seven-sketches_105 F01200	105	■ 0.976	id_1	id_1	phism id_1 , and no other morphisms. For any functor $F: \mathbf{1} \rightarrow \mathbf{Set}$ one can extract a set
fong-spivak-seven-sketches_105 F01201	105	■ 0.976	$F: \mathbf{1} \rightarrowtail$	$F: \mathbf{1} \rightarrow$	phism id_1 , and no other morphisms. For any functor $F: \mathbf{1} \rightarrow \mathbf{Set}$ one can extract a set
fong-spivak-seven-sketches_105 F01202	105	■ 1.000	$F(1)$	$F(1)$	$F(1)$. Show that for any set S , there is a functor $F_S: \mathbf{1} \rightarrow \mathbf{Set}$ such that $F_S(1) = S$.
fong-spivak-seven-sketches_105 F01203	105	■ 1.000	$F_S: \mathbf{1} \rightarrowtail$	$F_S: \mathbf{1} \rightarrow$	$F(1)$. Show that for any set S , there is a functor $F_S: \mathbf{1} \rightarrow \mathbf{Set}$ such that $F_S(1) = S$.
fong-spivak-seven-sketches_105 F01204	105	■ 1.000	$F_S(1)=S$	$F_S(1) = S$	$F(1)$. Show that for any set S , there is a functor $F_S: \mathbf{1} \rightarrow \mathbf{Set}$ such that $F_S(1) = S$.
fong-spivak-seven-sketches_105 F01205	105	■ 1.000	$r \in I(c)$	$r \in I(c)$	we call a row $r \in I(c)$.
fong-spivak-seven-sketches_106 F01206	106	■ 0.999	$Z:=F(z)$	$Z := F(z)$	What is a set-valued functor $F: \mathcal{C} \rightarrow \mathbf{Set}$? It will consist of a set $Z := F(z)$ and a function

Identifier	Page	Conf.	LaTeX source	Rendered	Image																		
fong-spivak-seven-sketches-F01207	106	1.000	$S:=F(s): Z \rightarrowtail Z$	$S := F(s) : Z \rightarrow Z$	$S := F(s): Z \rightarrow Z$, subject to the requirement that $S \circ S = S$. Here are some examples																		
fong-spivak-seven-sketches-F01208	106	1.000	$S \circ S=S$	$S; S = S$	$S := F(s): Z \rightarrow Z$, subject to the requirement that $S \circ S = S$. Here are some examples																		
fong-spivak-seven-sketches-F01209	106	0.837	$Z=\mathbb{N}$	$Z = \mathbb{N}$	$Z = \mathbb{N}$ is the set of natural numbers and S sends each number to 0. In particular,																		
fong-spivak-seven-sketches-F01210	106	0.917	$13+(2 * 4)$	$13 + (2 * 4)$	Z is the set of all well-formed arithmetic expressions, such as $13+(2*4)$ or -5, that																		
fong-spivak-seven-sketches-F01211	106	0.990	$Z=\mathbb{N}_{\geq 2}$	$Z = \mathbb{N}_{\geq 2}$	$Z = \mathbb{N}_{\geq 2}$, and S sends n to its smallest prime factor. The smallest prime factor of																		
fong-spivak-seven-sketches-F01212	106	1.000	$\mathbb{N}_{\geq 2}$	$\mathbb{N}_{\geq 2}$	<table><tr><th>$\mathbb{N}_{\geq 2}$</th><th>smallest prime factor</th></tr><tr><td>2</td><td>2</td></tr><tr><td>3</td><td>3</td></tr><tr><td>4</td><td>2</td></tr><tr><td>$\vdots$</td><td>$\vdots$</td></tr><tr><td>49</td><td>7</td></tr><tr><td>50</td><td>2</td></tr><tr><td>51</td><td>3</td></tr><tr><td>$\vdots$</td><td>$\vdots$</td></tr></table>	$\mathbb{N}_{\geq 2}$	smallest prime factor	2	2	3	3	4	2	$\vdots$	$\vdots$	49	7	50	2	51	3	$\vdots$	$\vdots$
$\mathbb{N}_{\geq 2}$	smallest prime factor																						
2	2																						
3	3																						
4	2																						
$\vdots$	$\vdots$																						
49	7																						
50	2																						
51	3																						
$\vdots$	$\vdots$																						
fong-spivak-seven-sketches-F01213	107	1.000	$\alpha: F \rightarrowtail G$	$\alpha : F \Rightarrow G$	a natural transformation $\alpha: F \Rightarrow G$,																		
fong-spivak-seven-sketches-F01214	107	0.884	$\alpha_c: F(c) \rightarrowtail G(c)$	$\alpha_c : F(c) \rightarrow G(c)$	(i) for each object $c \in \mathcal{C}$, one specifies a morphism $\alpha_c: F(c) \rightarrow G(c)$ in $\mathcal{D}$, called the																		
fong-spivak-seven-sketches-F01215	107	1.000	α	α	c -component of α .																		
fong-spivak-seven-sketches-F01216	107	1.000	$\alpha: F \rightarrow G$	$\alpha : F \rightarrow G$	A natural transformation $\alpha: F \rightarrow G$ is called a natural isomorphism if each component																		
fong-spivak-seven-sketches-F01217	107	0.996	α_c	α_c	α_c is an isomorphism in $\mathcal{D}$.																		
fong-spivak-seven-sketches-F01218	107	0.988	$D: \mathcal{J} \rightarrowtail \mathcal{C}$	$D : \mathcal{J} \rightarrow \mathcal{C}$	Definition 3.51. A diagram D in $\mathcal{C}$ is a functor $D: \mathcal{J} \rightarrow \mathcal{C}$ from any category $\mathcal{J}$, called																		
fong-spivak-seven-sketches-F01219	107	0.988	$\mathcal{J}$	$\mathcal{J}$	Definition 3.51. A diagram D in $\mathcal{C}$ is a functor $D: \mathcal{J} \rightarrow \mathcal{C}$ from any category $\mathcal{J}$, called																		
fong-spivak-seven-sketches-F01220	107	1.000	$D(f)=D(\text{f}^{\prime})$	$D(f) = D(f')$	the indexing category of the diagram D . We say that D commutes if $D(f) = D(f')$ holds																		
fong-spivak-seven-sketches-F01221	107	0.738	$f, \text{f}^{\prime}: a \rightarrowtail b$	$f, f' : a \rightarrow b$	for every parallel pair of morphisms $f, f': a \rightarrow b$ in $\mathcal{J}$. ⁷																		
fong-spivak-seven-sketches-F01222	107	0.738	$\mathcal{J}^7$	$\mathcal{J}^7$	for every parallel pair of morphisms $f, f': a \rightarrow b$ in $\mathcal{J}$. ⁷																		
fong-spivak-seven-sketches-F01223	107	1.000	$F(c) \rightarrowtailtail G(d)$	$F(c) \rightrightarrows G(d)$	In terms of Eq. (3.50), the only case of two parallel morphisms is that of $F(c) \rightrightarrows G(d)$,																		
fong-spivak-seven-sketches-F01224	107	0.528	$F(f) \circ \alpha_d = \alpha_c \circ G(f)$	$F(f) \circ \alpha_d = \alpha_c \circ G(f)$	so to say that the diagram commutes is to say that $F(f) \circ \alpha_d = \alpha_c \circ G(f)$. This is exactly																		
fong-spivak-seven-sketches-F01225	108	1.000	$\alpha_1: v \rightarrowtail x$	$\alpha_1 : v \rightarrow x$	transformation $\alpha: F \Rightarrow G$ is given by choosing components $\alpha_1: v \rightarrow x$ and $\alpha_2: w \rightarrow y$.																		
fong-spivak-seven-sketches-F01226	108	1.000	$\alpha_2: w \rightarrowtail y$	$\alpha_2 : w \rightarrow y$	transformation $\alpha: F \Rightarrow G$ is given by choosing components $\alpha_1: v \rightarrow x$ and $\alpha_2: w \rightarrow y$.																		
fong-spivak-seven-sketches-F01227	108	1.000	$\alpha_1=c$	$\alpha_1 = c$	We have highlighted the only choice for each in green; namely, $\alpha_1 = c$ and $\alpha_2 = g$.																		
fong-spivak-seven-sketches-F01228	108	1.000	$\alpha_2=g$	$\alpha_2 = g$	We have highlighted the only choice for each in green; namely, $\alpha_1 = c$ and $\alpha_2 = g$.																		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_1_F01229	108	0.150	$\mathcal{C} \xrightarrow[\underset{G}{\leftarrow}]{\underset{F}{\rightarrow}} D$	$\mathcal{C} \xrightarrow[\underset{G}{\leftarrow}]{\underset{F}{\rightarrow}} D$	the notation $\mathcal{C} \begin{array}{c} \xrightarrow{F} \\ \Downarrow \alpha \\ \xrightarrow{G} \end{array} \mathcal{D}?$
fong-spivak-seven-sketches_1_F01230	108	0.959	$A, B: \mathbf{1} \rightarrow$	$A, B: \mathbf{1} \rightarrow$	a set. So suppose A and B are sets considered as functors $A, B: \mathbf{1} \rightarrow \mathbf{Set}$. A natural
fong-spivak-seven-sketches_1_F01231	108	1.000	$\mathcal{D}^{\mathcal{C}}$	$\mathcal{D}^{\mathcal{C}}$	Definition 3.54. Let $\mathcal{C}$ and $\mathcal{D}$ be categories. We denote by $\mathcal{D}^{\mathcal{C}}$ the category whose objects
fong-spivak-seven-sketches_1_F01232	108	0.678	$\mathcal{D}^{\mathcal{P}}(F, G)$	$\mathcal{D}^{\mathcal{P}}(F, G)$	are functors $F: \mathcal{C} \rightarrow \mathcal{D}$ and whose morphisms $\mathcal{D}^{\mathcal{C}}(F, G)$ are the natural transformations
fong-spivak-seven-sketches_1_F01233	108	0.962	$\mathcal{D}^{\mathrm{e}}$	$\mathcal{D}^{\mathrm{e}}$	Exercise 3.55. Let's look more deeply at how $\mathcal{D}^{\mathcal{C}}$ is a category.
fong-spivak-seven-sketches_1_F01234	108	0.875	$F \in \mathcal{D}^{\mathcal{C}}$	$F \in \mathcal{D}^{\mathcal{C}}$	2. Propose an identity natural transformation on any object $F \in \mathcal{D}^{\mathcal{C}}$, and check that
fong-spivak-seven-sketches_1_F01235	108	0.774	$\mathbf{Set}^{\mathbf{1}}$	$\mathbf{Set}^{\mathbf{1}}$	Example 3.56. In our new language, Example 3.53 says that $\mathbf{Set}^{\mathbf{1}}$ is equivalent to $\mathbf{Set}$.
fong-spivak-seven-sketches_1_F01236	108	0.923	$(F_0, F_1, F_2, \dots)$	$(F_0, F_1, F_2, \dots)$	sequence $(F_0, F_1, F_2, \dots)$ of natural numbers, i.e. $F_0 \leq F_1 \leq F_2 \leq \dots$. If G is another
fong-spivak-seven-sketches_1_F01237	108	0.923	$F_0 \leq F_1 \leq F_2 \leq \dots$	$F_0 \leq F_1 \leq F_2 \leq \dots$	sequence $(F_0, F_1, F_2, \dots)$ of natural numbers, i.e. $F_0 \leq F_1 \leq F_2 \leq \dots$. If G is another
fong-spivak-seven-sketches_1_F01238	109	1.000	$\alpha_n: F_n \rightarrow G_n$	$\alpha_n: F_n \rightarrow G_n$	ponent $\alpha_n: F_n \rightarrow G_n$ has no data, it just tells us a fact: that $F_n \leq G_n$. And the naturality
fong-spivak-seven-sketches_1_F01239	109	1.000	$F_n \leq G_n$	$F_n \leq G_n$	ponent $\alpha_n: F_n \rightarrow G_n$ has no data, it just tells us a fact: that $F_n \leq G_n$. And the naturality
fong-spivak-seven-sketches_1_F01240	109	0.987	$F \Rightarrow G$	$F \Rightarrow G$	$F \Rightarrow G$ are the same. In other words, the category $\mathcal{N}^{\mathcal{N}}$ is itself a preorder; namely the
fong-spivak-seven-sketches_1_F01241	109	0.987	$\mathcal{N}^{\mathcal{N}}$	$\mathcal{N}^{\mathcal{N}}$	$F \Rightarrow G$ are the same. In other words, the category $\mathcal{N}^{\mathcal{N}}$ is itself a preorder; namely the
fong-spivak-seven-sketches_1_F01242	109	1.000	$F, G: \mathcal{C} \rightarrow \mathcal{P}$	$F, G: \mathcal{C} \rightarrow \mathcal{P}$	1. For any two functors $F, G: \mathcal{C} \rightarrow \mathcal{P}$, there is at most one natural transformation
fong-spivak-seven-sketches_1_F01243	109	1.000	$F \rightarrow G$	$F \rightarrow G$	$F \rightarrow G$.
fong-spivak-seven-sketches_1_F01244	109	0.999	$F, G: \mathcal{P} \rightarrow \mathcal{C}$	$F, G: \mathcal{P} \rightarrow \mathcal{C}$	2. For any two functors $F, G: \mathcal{P} \rightarrow \mathcal{C}$, there is at most one natural transformation
fong-spivak-seven-sketches_1_F01245	109	0.865	$F; G \cong \mathrm{id}_{\mathbf{Pro}}$	$F; G \cong \mathrm{id}_{\mathbf{Pro}}$	such that there exist natural isomorphisms $F; G \cong \mathrm{id}_{\mathbf{Pro}}$ and $G; F \cong \mathrm{id}_{\mathbf{Bool-Cat}}$ in the
fong-spivak-seven-sketches_1_F01246	109	0.865	$G; F \cong \mathrm{id}_{\mathbf{Bool-Cat}}$	$G; F \cong \mathrm{id}_{\mathbf{Bool-Cat}}$	such that there exist natural isomorphisms $F; G \cong \mathrm{id}_{\mathbf{Pro}}$ and $G; F \cong \mathrm{id}_{\mathbf{Bool-Cat}}$ in the
fong-spivak-seven-sketches_1_F01247	109	1.000	$I, J: \mathcal{C} \rightarrow$	$I, J: \mathcal{C} \rightarrow$	Definition 3.60. Suppose that $\mathcal{C}$ is a database schema and $I, J: \mathcal{C} \rightarrow \mathbf{Set}$ are database
fong-spivak-seven-sketches_1_F01248	109	1.000	$\alpha: I \rightarrow$	$\alpha: I \rightarrow$	instances. An <i>instance homomorphism</i> between them is a natural transformation $\alpha: I \rightarrow$
fong-spivak-seven-sketches_1_F01249	109	0.490	$:=$	$:=$	J . Write $\mathcal{C}\text{-Inst} := \mathbf{Set}^{\mathcal{C}}$ to denote the functor category as defined in Definition 3.54.
fong-spivak-seven-sketches_1_F01250	109	0.490	$^{\mathcal{C}}$	$^{\mathcal{C}}$	J . Write $\mathcal{C}\text{-Inst} := \mathbf{Set}^{\mathcal{C}}$ to denote the functor category as defined in Definition 3.54.
fong-spivak-seven-sketches_1_F01251	110	0.986	$I($	$I($	Here $I(\text{Arrow}) = \{a, b, c, d, e\}$, and $I(\text{Vertex}) = \{1, 2, 3, 4\}$. One can draw the instance
fong-spivak-seven-sketches_1_F01252	110	0.986	$) = \{a, b, c, d, e\}$	$) = \{a, b, c, d, e\}$	Here $I(\text{Arrow}) = \{a, b, c, d, e\}$, and $I(\text{Vertex}) = \{1, 2, 3, 4\}$. One can draw the instance
fong-spivak-seven-sketches_1_F01253	110	0.986	$) = \{1, 2, 3, 4\}$	$) = \{1, 2, 3, 4\}$	Here $I(\text{Arrow}) = \{a, b, c, d, e\}$, and $I(\text{Vertex}) = \{1, 2, 3, 4\}$. One can draw the instance
fong-spivak-seven-sketches_1_F01254	110	0.983	$G, H: \mathbf{Gr} \rightarrow$	$G, H: \mathbf{Gr} \rightarrow$	are called <i>graph homomorphisms</i> . Let's unwind this. Suppose that $G, H: \mathbf{Gr} \rightarrow \mathbf{Set}$
fong-spivak-seven-sketches_1_F01255	110	0.989	$G, H \in \mathbf{Gr}$	$G, H \in \mathbf{Gr}$	are functors (i.e. $\mathbf{Gr}$ -instances); that is, they are objects $G, H \in \mathbf{Gr}\text{-Inst}$. A morphism

Identifier	Page	Conf.	LaTeX source	Rendered	Image																																								
fong-spivak-seven-sketches_110 F01256	110	0.999	$G \rightarrowtail H$	$G \rightarrow H$	$G \rightarrow H$ is a natural transformation $\alpha: G \rightarrow H$ between them; what does that entail?																																								
fong-spivak-seven-sketches_110 F01257	110	0.999	$\alpha: G \rightarrowtail H$	$\alpha: G \rightarrow H$	$G \rightarrow H$ is a natural transformation $\alpha: G \rightarrow H$ between them; what does that entail?																																								
fong-spivak-seven-sketches_110 F01258	110	1.000	$\alpha_{\text{\texttt{Vertex}}}$	α_{Vertex}	α_{Vertex} and α_{Arrow} to respect source and targets in G and H . The left diagram says “start																																								
fong-spivak-seven-sketches_110 F01259	110	1.000	$\alpha_{\text{\texttt{Arrow}}}$	α_{Arrow}	α_{Vertex} and α_{Arrow} to respect source and targets in G and H . The left diagram says “start																																								
fong-spivak-seven-sketches_111 F01260	111	0.980	$G :=$	$G :=$	<table><tr><td></td><td>Arrow</td><td>source</td><td>target</td><td>Vertex</td></tr><tr><td>$G :=$</td><td>a</td><td>1</td><td>2</td><td>1</td></tr><tr><td></td><td>b</td><td>2</td><td>3</td><td>2</td></tr><tr><td></td><td></td><td></td><td></td><td>3</td></tr></table> <table><tr><td></td><td>Arrow</td><td>source</td><td>target</td><td>Vertex</td></tr><tr><td>$H :=$</td><td>c</td><td>4</td><td>5</td><td>4</td></tr><tr><td></td><td>d</td><td>4</td><td>5</td><td>5</td></tr><tr><td></td><td>e</td><td>5</td><td>5</td><td></td></tr></table>		Arrow	source	target	Vertex	$G :=$	a	1	2	1		b	2	3	2					3		Arrow	source	target	Vertex	$H :=$	c	4	5	4		d	4	5	5		e	5	5	
	Arrow	source	target	Vertex																																									
$G :=$	a	1	2	1																																									
	b	2	3	2																																									
				3																																									
	Arrow	source	target	Vertex																																									
$H :=$	c	4	5	4																																									
	d	4	5	5																																									
	e	5	5																																										
fong-spivak-seven-sketches_111 F01261	111	0.980	$H :=$	$H :=$	<table><tr><td></td><td>Arrow</td><td>source</td><td>target</td><td>Vertex</td></tr><tr><td>$G :=$</td><td>a</td><td>1</td><td>2</td><td>1</td></tr><tr><td></td><td>b</td><td>2</td><td>3</td><td>2</td></tr><tr><td></td><td></td><td></td><td></td><td>3</td></tr></table> <table><tr><td></td><td>Arrow</td><td>source</td><td>target</td><td>Vertex</td></tr><tr><td>$H :=$</td><td>c</td><td>4</td><td>5</td><td>4</td></tr><tr><td></td><td>d</td><td>4</td><td>5</td><td>5</td></tr><tr><td></td><td>e</td><td>5</td><td>5</td><td></td></tr></table>		Arrow	source	target	Vertex	$G :=$	a	1	2	1		b	2	3	2					3		Arrow	source	target	Vertex	$H :=$	c	4	5	4		d	4	5	5		e	5	5	
	Arrow	source	target	Vertex																																									
$G :=$	a	1	2	1																																									
	b	2	3	2																																									
				3																																									
	Arrow	source	target	Vertex																																									
$H :=$	c	4	5	4																																									
	d	4	5	5																																									
	e	5	5																																										
fong-spivak-seven-sketches_111 F01262	111	1.000	G, H	G, H	<i>Exercise 3.64.</i> We claim that—with G, H as in Example 3.63—there is exactly one																																								
fong-spivak-seven-sketches_111 F01263	111	0.995	$\alpha_{\text{\texttt{Arrow}}}(a) = d$	$\alpha_{\text{Arrow}}(a) = d$	graph homomorphism $\alpha: G \rightarrow H$ such that $\alpha_{\text{Arrow}}(a) = d$.																																								
fong-spivak-seven-sketches_112 F01264	112	0.598	$F: \mathbf{Gr} \rightarrow \mathbf{DDS}$	$F: \mathbf{Gr} \rightarrow \mathbf{DDS}$	$F: \mathbf{Gr} \rightarrow \mathbf{DDS}$ to migrate data in a way that makes sense to us, the authors. Here we																																								
fong-spivak-seven-sketches_113 F01265	113	0.384	$F_{\circlearrowleft}: \mathbf{Gr} \rightarrow$	$F_{\circlearrowleft}: \mathbf{Gr} \rightarrow$	$F_{\circlearrowleft}: \mathbf{Gr} \rightarrow \mathbf{Set}$. Both F and I respect identities and composition, so $F_{\circlearrowleft}$ does too. Thus																																								
fong-spivak-seven-sketches_113 F01266	113	0.384	$F_{\circlearrowleft} I$	$F_{\circlearrowleft} I$	$F_{\circlearrowleft}: \mathbf{Gr} \rightarrow \mathbf{Set}$. Both F and I respect identities and composition, so $F_{\circlearrowleft}$ does too. Thus																																								
fong-spivak-seven-sketches_114 F01267	114	0.731	$I: \mathcal{D} \rightarrow$	$I: \mathcal{D} \rightarrow$	set-valued functor $I: \mathcal{D} \rightarrow \mathbf{Set}$, we refer to the composite functor $F_{\circlearrowleft} I: \mathcal{C} \rightarrow \mathbf{Set}$ as the																																								
fong-spivak-seven-sketches_114 F01268	114	0.731	$F: I: \mathcal{C} \rightarrow$	$F: I: \mathcal{C} \rightarrow$	set-valued functor $I: \mathcal{D} \rightarrow \mathbf{Set}$, we refer to the composite functor $F_{\circlearrowleft} I: \mathcal{C} \rightarrow \mathbf{Set}$ as the																																								
fong-spivak-seven-sketches_114 F01269	114	1.000	$\alpha: I \Rightarrow J$	$\alpha: I \Rightarrow J$	Given a natural transformation $\alpha: I \Rightarrow J$, there is a natural transformation $\alpha_F: F_{\circlearrowleft} I \Rightarrow$																																								
fong-spivak-seven-sketches_114 F01270	114	1.000	$\alpha_F: F_{\circlearrowleft} I \Rightarrow$	$\alpha_F: F_{\circlearrowleft} I \Rightarrow$	Given a natural transformation $\alpha: I \Rightarrow J$, there is a natural transformation $\alpha_F: F_{\circlearrowleft} I \Rightarrow$																																								
fong-spivak-seven-sketches_114 F01271	114	0.314	$F_{\circlearrowleft} J$	$F_{\circlearrowleft} J$	$F_{\circlearrowleft} J$, whose component $(F_{\circlearrowleft} I)(c) \rightarrow (F_{\circlearrowleft} J)(c)$ for any $c \in \text{Ob}(\mathcal{C})$ is given by $(\alpha_F)_c := \alpha_{Fc}$.																																								
fong-spivak-seven-sketches_114 F01272	114	0.314	$\left(F_{\circlearrowleft} I\right)(c) \rightarrow \left(F_{\circlearrowleft} J\right)(c)$	$(F_{\circlearrowleft} I)(c) \rightarrow (F_{\circlearrowleft} J)(c)$	$F_{\circlearrowleft} J$, whose component $(F_{\circlearrowleft} I)(c) \rightarrow (F_{\circlearrowleft} J)(c)$ for any $c \in \text{Ob}(\mathcal{C})$ is given by $(\alpha_F)_c := \alpha_{Fc}$.																																								
fong-spivak-seven-sketches_114 F01273	114	0.314	$c \in \text{Ob}(\mathcal{C})$	$c \in \text{Ob}(\mathcal{C})$	$F_{\circlearrowleft} J$, whose component $(F_{\circlearrowleft} I)(c) \rightarrow (F_{\circlearrowleft} J)(c)$ for any $c \in \text{Ob}(\mathcal{C})$ is given by $(\alpha_F)_c := \alpha_{Fc}$.																																								
fong-spivak-seven-sketches_114 F01274	114	0.314	$(\alpha_F)_c := \alpha_{Fc}$	$(\alpha_F)_c := \alpha_{Fc}$	$F_{\circlearrowleft} J$, whose component $(F_{\circlearrowleft} I)(c) \rightarrow (F_{\circlearrowleft} J)(c)$ for any $c \in \text{Ob}(\mathcal{C})$ is given by $(\alpha_F)_c := \alpha_{Fc}$.																																								
fong-spivak-seven-sketches_114 F01275	114	0.338	$\Delta_F: \mathcal{D}$	$\Delta_F: \mathcal{D}$	This uses the data of F to define a functor $\Delta_F: \mathcal{D}\text{-Inst} \rightarrow \mathcal{C}\text{-Inst}$.																																								

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_114 F01276	114	■ 1.000	Δ	Δ	in general. The relevance to databases is that the data migration functor Δ from Defi-
fong-spivak-seven-sketches_114 F01277	114	■ 0.998	$P(a, b)$	$P(a, b)$	Recall from Section 3.2.3 that in a preorder P , a hom-set $P(a, b)$ has one element when
fong-spivak-seven-sketches_114 F01278	114	■ 0.997	$Q(f(p), q) \cong P(p, g(q))$	$Q(f(p), q) \cong P(p, g(q))$	of sets $Q(f(p), q) \cong P(p, g(q))$: either both are one-element sets or both are 0-element
fong-spivak-seven-sketches_114 F01279	114	■ 1.000	$L: \mathcal{C} \rightarrow \mathcal{D}$	$L: \mathcal{C} \rightarrow \mathcal{D}$	Definition 3.70. Let $\mathcal{C}$ and $\mathcal{D}$ be categories, and $L: \mathcal{C} \rightarrow \mathcal{D}$ and $R: \mathcal{D} \rightarrow \mathcal{C}$ be functors.
fong-spivak-seven-sketches_114 F01280	114	■ 1.000	$R: \mathcal{D} \rightarrow \mathcal{C}$	$R: \mathcal{D} \rightarrow \mathcal{C}$	Definition 3.70. Let $\mathcal{C}$ and $\mathcal{D}$ be categories, and $L: \mathcal{C} \rightarrow \mathcal{D}$ and $R: \mathcal{D} \rightarrow \mathcal{C}$ be functors.
fong-spivak-seven-sketches_114 F01281	114	■ 1.000	L	L	We say that L is left adjoint to R (and that R is right adjoint to L) if, for any $c \in \mathcal{C}$ and
fong-spivak-seven-sketches_114 F01282	114	■ 0.844	$d \in \mathcal{D}$	$d \in \mathcal{D}$	$d \in \mathcal{D}$, there is an isomorphism of hom-sets
fong-spivak-seven-sketches_114 F01283	114	■ 0.615	$d.^8$	$d.^8$	that is natural in c and d . ⁸
fong-spivak-seven-sketches_114 F01284	114	■ 0.998	$f: c \rightarrow R(d)$	$f: c \rightarrow R(d)$	Given a morphism $f: c \rightarrow R(d)$ in $\mathcal{C}$, its image $g := \alpha_{c,d}(f)$ is called its <i>mate</i> .
fong-spivak-seven-sketches_114 F01285	114	■ 0.998	$g := \alpha_{c,d}(f)$	$g := \alpha_{c,d}(f)$	Given a morphism $f: c \rightarrow R(d)$ in $\mathcal{C}$, its image $g := \alpha_{c,d}(f)$ is called its <i>mate</i> .
fong-spivak-seven-sketches_114 F01286	114	■ 1.000	$g: L(c) \rightarrow d$	$g: L(c) \rightarrow d$	Similarly, the mate of $g: L(c) \rightarrow d$ is f .
fong-spivak-seven-sketches_115 F01287	115	■ 1.000	$L \dashv R$	$L \dashv R$	To denote an adjunction we write $L \dashv R$, or in diagrams,
fong-spivak-seven-sketches_115 F01288	115	■ 0.801	$B \in \mathrm{Ob}(\mathbf{Set})$	$B \in \mathrm{Ob}(\mathbf{Set})$	<i>Example 3.72.</i> Let $B \in \mathrm{Ob}(\mathbf{Set})$ be any set. There is an adjunction called ‘currying B ,’
fong-spivak-seven-sketches_115 F01289	115	■ 1.000	A, C	A, C	Abstractly we write it as on the left, but what this means is that for any sets A, C , there
fong-spivak-seven-sketches_115 F01290	115	■ 0.912	$B \rightarrow C$	$B \rightarrow C$	and C are sets. Then the set of functions $B \rightarrow C$ is also a set; let’s denote it C^B . It’s
fong-spivak-seven-sketches_115 F01291	115	■ 0.912	C^B	C^B	and C are sets. Then the set of functions $B \rightarrow C$ is also a set; let’s denote it C^B . It’s
fong-spivak-seven-sketches_115 F01292	115	■ 1.000	10^3	10^3	written this way because if C has 10 elements and B has 3 elements then C^B has 10^3
fong-spivak-seven-sketches_115 F01293	115	■ 1.000	$ \mathrm{left} C^B\mathrm{right} = C ^{ B }$	$ C^B = C ^{ B }$	elements, and more generally for any two finite sets $ C^B = C ^{ B }$.
fong-spivak-seven-sketches_115 F01294	115	■ 1.000	$(A \times B) \rightarrow C$	$(A \times B) \rightarrow C$	dence between functions $(A \times B) \rightarrow C$ and functions $A \rightarrow C^B$. Intuitively, if I have a
fong-spivak-seven-sketches_115 F01295	115	■ 1.000	$A \rightarrow C^B$	$A \rightarrow C^B$	dence between functions $(A \times B) \rightarrow C$ and functions $A \rightarrow C^B$. Intuitively, if I have a
fong-spivak-seven-sketches_115 F01296	115	■ 0.996	$b \multimap c$	$b \multimap c$	objects and what we called hom-elements $b \multimap c$ in Definition 2.79.
fong-spivak-seven-sketches_115 F01297	115	■ 0.983	$- \times B$	$- \times B$	<i>Exercise 3.73.</i> In Example 3.72, we discussed an adjunction between functors $- \times B$
fong-spivak-seven-sketches_115 F01298	115	■ 1.000	$(-)^B$	$(-)^B$	and $(-)^B$. But we only said how these functors worked on objects: for an arbitrary set
fong-spivak-seven-sketches_115 F01299	115	■ 1.000	$X \times B$	$X \times B$	X , they return sets $X \times B$ and X^B respectively.
fong-spivak-seven-sketches_115 F01300	115	■ 1.000	X^B	X^B	X , they return sets $X \times B$ and X^B respectively.
fong-spivak-seven-sketches_115 F01301	115	■ 0.953	$\mathcal{C}^{\mathrm{op}} \times \mathcal{D} \rightarrow$	$\mathcal{C}^{\mathrm{op}} \times \mathcal{D} \rightarrow$	⁸ This naturality is between functors $\mathcal{C}^{\mathrm{op}} \times \mathcal{D} \rightarrow \mathbf{Set}$. It says that for any morphisms $f: c' \rightarrow c$ in $\mathcal{C}$
fong-spivak-seven-sketches_115 F01302	115	■ 0.953	$f: c' \rightarrow c$	$f: c' \rightarrow c$	⁸ This naturality is between functors $\mathcal{C}^{\mathrm{op}} \times \mathcal{D} \rightarrow \mathbf{Set}$. It says that for any morphisms $f: c' \rightarrow c$ in $\mathcal{C}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_1 F01303	115	■ 1.000	$\text{g: } d \rightarrow d^{\{\prime\}}$	$g: d \rightarrow d'$	and $g: d \rightarrow d'$ in $\mathcal{D}$, the following diagram commutes:
fong-spivak-seven-sketches_1 F01304	116	■ 1.000	$-\times B: X \times B \rightarrow Y \times B$	$-\times B: X \times B \rightarrow Y \times B$	1. Given a morphism $f: X \rightarrow Y$, what morphism should $-\times B: X \times B \rightarrow Y \times B$
fong-spivak-seven-sketches_1 F01305	116	■ 1.000	$(-)^B: X^B \rightarrow Y^B$	$(-)^B: X^B \rightarrow Y^B$	2. Given a morphism $f: X \rightarrow Y$, what morphism should $(-)^B: X^B \rightarrow Y^B$ return?
fong-spivak-seven-sketches_1 F01306	116	■ 1.000	$+: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$	$+: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$	3. Consider the function $+: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$, which sends $(a, b) \mapsto a + b$. Currying $+$,
fong-spivak-seven-sketches_1 F01307	116	■ 1.000	$(a, b) \mapsto a + b$	$(a, b) \mapsto a + b$	3. Consider the function $+: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$, which sends $(a, b) \mapsto a + b$. Currying $+$,
fong-spivak-seven-sketches_1 F01308	116	■ 1.000	$p: \mathbb{N} \rightarrow \mathbb{N}^{\mathbb{N}}$	$p: \mathbb{N} \rightarrow \mathbb{N}^{\mathbb{N}}$	we get a certain function $p: \mathbb{N} \rightarrow \mathbb{N}^{\mathbb{N}}$. What is $p(3)$?
fong-spivak-seven-sketches_1 F01309	116	■ 1.000	$p(3)$	$p(3)$	we get a certain function $p: \mathbb{N} \rightarrow \mathbb{N}^{\mathbb{N}}$. What is $p(3)$?
fong-spivak-seven-sketches_1 F01310	116	■ 1.000	Δ_F	Δ_F	Given $F: \mathcal{C} \rightarrow \mathcal{D}$, the data migration functor Δ_F turns $\mathcal{D}$ -instances into $\mathcal{C}$ -instances.
fong-spivak-seven-sketches_1 F01311	117	■ 1.000	Σ_F	Σ_F	and get a graph, the migration functors Σ_F and Π_F can be used to turn any graph into
fong-spivak-seven-sketches_1 F01312	117	■ 1.000	Π_F	Π_F	and get a graph, the migration functors Σ_F and Π_F can be used to turn any graph into
fong-spivak-seven-sketches_1 F01313	117	■ 0.910	$J: \mathbf{Gr} \rightarrow$	$J: \mathbf{Gr} \rightarrow$	a discrete dynamical system. That is, given an instance $J: \mathbf{Gr} \rightarrow \mathbf{Set}$, we can get
fong-spivak-seven-sketches_1 F01314	117	■ 1.000	$\Sigma_F(J)$	$\Sigma_F(J)$	instances $\Sigma_F(J)$ and $\Pi_F(J)$ on DDS . This, however, is quite technical, and we leave it
fong-spivak-seven-sketches_1 F01315	117	■ 1.000	$\Pi_F(J)$	$\Pi_F(J)$	instances $\Sigma_F(J)$ and $\Pi_F(J)$ on DDS . This, however, is quite technical, and we leave it
fong-spivak-seven-sketches_1 F01316	117	■ 1.000	(e, f)	(e, f)	The functor Π_F puts into Airline Seat the set of those pairs (e, f) where e is an
fong-spivak-seven-sketches_1 F01317	118	■ 1.000	$\Pi_F(I)$	$\Pi_F(I)$	instance I , in which case $\Pi_F(I)(\text{Airline Seat})$ would be empty. But in other uses of these
fong-spivak-seven-sketches_1 F01318	118	■ 1.000	Δ_F, Σ_F	Δ_F, Σ_F	F between schemas and using the “induced” functors Δ_F , Σ_F , and Π_F . Indeed, in
fong-spivak-seven-sketches_1 F01319	118	■ 0.931	$\mathbf{1}$	$\mathbf{1}$	namely where the target category $\mathcal{D}$ is equal to $\mathbf{1}$; see Exercise 3.12. In this case, there
fong-spivak-seven-sketches_1 F01320	118	■ 0.911	$\mathcal{C} \rightarrow \mathbf{1}$	$\mathcal{C} \rightarrow \mathbf{1}$	is exactly one functor $\mathcal{C} \rightarrow \mathbf{1}$ for any $\mathcal{C}$; let’s denote it
fong-spivak-seven-sketches_1 F01321	118	■ 0.915	$\mathcal{C} \rightarrow \mathbf{1}$	$\mathcal{C} \rightarrow \mathbf{1}$	<i>Exercise 3.76.</i> Describe the functor $!: \mathcal{C} \rightarrow \mathbf{1}$ from Eq. (3.75). Where does it send each
fong-spivak-seven-sketches_1 F01322	118	■ 0.133	$\Sigma_!$	$\Sigma_!$	We want to consider the data migration functors $\Sigma_!: \mathcal{C}\text{-Inst} \rightarrow \mathbf{1}\text{-Inst}$ and $\Pi_!: \mathcal{C}\text{-Inst} \rightarrow$
fong-spivak-seven-sketches_1 F01323	118	■ 0.133	$\Pi_!: \mathcal{C}\text{-Inst} \rightarrow$	$\Pi_!: \mathcal{C}\text{-Inst} \rightarrow$	We want to consider the data migration functors $\Sigma_!: \mathcal{C}\text{-Inst} \rightarrow \mathbf{1}\text{-Inst}$ and $\Pi_!: \mathcal{C}\text{-Inst} \rightarrow$
fong-spivak-seven-sketches_1 F01324	118	■ 0.988	$\Sigma_!(I)$	$\Sigma_!(I)$	Given any schema $\mathcal{C}$ and instance $I: \mathcal{C} \rightarrow \mathbf{Set}$, we will get sets $\Sigma_!(I)$ and $\Pi_!(I)$. Thinking
fong-spivak-seven-sketches_1 F01325	118	■ 0.988	$\Pi_!(I)$	$\Pi_!(I)$	Given any schema $\mathcal{C}$ and instance $I: \mathcal{C} \rightarrow \mathbf{Set}$, we will get sets $\Sigma_!(I)$ and $\Pi_!(I)$. Thinking
fong-spivak-seven-sketches_1 F01326	118	■ 0.985	$I: \mathcal{G} \rightarrow$	$I: \mathcal{G} \rightarrow$	Here’s a sample instance $I: \mathcal{G} \rightarrow \mathbf{Set}$:
fong-spivak-seven-sketches_1 F01327	119	■ 1.000	$\mathcal{G}$	$\mathcal{G}$	<i>Exercise 3.78.</i> Note that $\mathcal{G}$ from Eq. (3.77) is isomorphic to the schema $\mathbf{Gr}$. In Sec-
fong-spivak-seven-sketches_1 F01328	119	■ 0.769	$!: \mathcal{G} \rightarrow \mathbf{1}$	$!: \mathcal{G} \rightarrow \mathbf{1}$	Now we have a unique functor $!: \mathcal{G} \rightarrow \mathbf{1}$, and we want to say what $\Sigma_!(I)$ and $\Pi_!(I)$ give
fong-spivak-seven-sketches_1 F01329	119	■ 1.000	$\Pi_!$	$\Pi_!$	Where do these facts—that $\Pi_!$ and $\Sigma_!$ act the way we said—come from? Everything

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_119 F01330	119	0.788	$\mathbf{1}, \Sigma$	$\mathbf{1}, \Sigma$	reader should keep in mind that, in general and not just for functors to $\mathbf{1}$, Σ -operations
fong-spivak-seven-sketches_120 F01331	120	0.999	$C \rightarrow Z$	$C \rightarrow Z$	each object C of $\mathcal{C}$, there exists a unique morphism $! : C \rightarrow Z$.
fong-spivak-seven-sketches_120 F01332	120	0.539	$\{\bullet\}$	$\{\bullet\}$	Consider some such set $\{\bullet\}$. Then for any other set C we need to check that there is
fong-spivak-seven-sketches_120 F01333	120	0.821	$C \rightarrow \{\bullet\}$	$C \rightarrow \{\bullet\}$	exactly one function $! : C \rightarrow \{\bullet\}$. This unique function is the one that does the only
fong-spivak-seven-sketches_120 F01334	120	1.000	$c \in C$	$c \in C$	thing that can be done: it maps each element $c \in C$ to the element $\bullet \in \{\bullet\}$.
fong-spivak-seven-sketches_120 F01335	120	1.000	$\bullet \in \{\bullet\}$	$\bullet \in \{\bullet\}$	thing that can be done: it maps each element $c \in C$ to the element $\bullet \in \{\bullet\}$.
fong-spivak-seven-sketches_120 F01336	120	0.918	$z \in P$	$z \in P$	<i>Exercise 3.81.</i> Let $(P, \leq)$ be a preorder, let $z \in P$ be an element, and let $\mathcal{P}$ be the
fong-spivak-seven-sketches_120 F01337	120	1.000	$c \in P$	$c \in P$	only if it is a <i>top element</i> in P : that is, if and only if for all $c \in P$ we have $c \leq z$.
fong-spivak-seven-sketches_120 F01338	120	1.000	$c \leq z$	$c \leq z$	only if it is a <i>top element</i> in P : that is, if and only if for all $c \in P$ we have $c \leq z$.
fong-spivak-seven-sketches_120 F01339	120	0.924	$Z^{\prime}$	Z'	<i>Proof.</i> This is a simple, but powerful standard argument. Suppose Z and Z' are both
fong-spivak-seven-sketches_120 F01340	120	1.000	$a : Z \rightarrow Z^{\prime}$	$a : Z \rightarrow Z'$	terminal objects in some category $\mathcal{C}$. Then there exist (unique) maps $a : Z \rightarrow Z'$ and
fong-spivak-seven-sketches_120 F01341	120	1.000	$b : Z^{\prime} \rightarrow Z$	$b : Z' \rightarrow Z$	$b : Z' \rightarrow Z$. Composing these, we get a map $a \circ b : Z \rightarrow Z$. Now since Z is terminal, this
fong-spivak-seven-sketches_120 F01342	120	1.000	$a ; b : Z \rightarrow Z$	$a; b : Z \rightarrow Z$	$b : Z' \rightarrow Z$. Composing these, we get a map $a \circ b : Z \rightarrow Z$. Now since Z is terminal, this
fong-spivak-seven-sketches_120 F01343	120	0.707	$Z \rightarrow Z$	$Z \rightarrow Z$	map $Z \rightarrow Z$ must be unique. But id_Z is also such a map. So we must have $a \circ b = \text{id}_Z$.
fong-spivak-seven-sketches_120 F01344	120	0.707	id_Z	id_Z	map $Z \rightarrow Z$ must be unique. But id_Z is also such a map. So we must have $a \circ b = \text{id}_Z$.
fong-spivak-seven-sketches_120 F01345	120	0.707	$a ; b = \text{id}_Z$	$a; b = \text{id}_Z$	map $Z \rightarrow Z$ must be unique. But id_Z is also such a map. So we must have $a \circ b = \text{id}_Z$.
fong-spivak-seven-sketches_120 F01346	120	0.900	$b ; a = \text{id}_{Z^{\prime}}$	$b; a = \text{id}_{Z'}$	Similarly, we find that $b \circ a = \text{id}_{Z'}$. Thus a is an isomorphism, with inverse b .
fong-spivak-seven-sketches_120 F01347	120	1.000	X, Y	X, Y	Definition 3.86. Let $\mathcal{C}$ be a category, and let X, Y be objects in $\mathcal{C}$. A <i>product</i> of X and Y is
fong-spivak-seven-sketches_120 F01348	120	1.000	$p_X : X \times Y \rightarrow X$	$p_X : X \times Y \rightarrow X$	an object, denoted $X \times Y$, together with morphisms $p_X : X \times Y \rightarrow X$ and $p_Y : X \times Y \rightarrow Y$
fong-spivak-seven-sketches_120 F01349	120	1.000	$p_Y : X \times Y \rightarrow Y$	$p_Y : X \times Y \rightarrow Y$	an object, denoted $X \times Y$, together with morphisms $p_X : X \times Y \rightarrow X$ and $p_Y : X \times Y \rightarrow Y$
fong-spivak-seven-sketches_120 F01350	120	1.000	$f : C \rightarrow X$	$f : C \rightarrow X$	such that for all objects C together with morphisms $f : C \rightarrow X$ and $g : C \rightarrow Y$, there
fong-spivak-seven-sketches_120 F01351	120	1.000	$g : C \rightarrow Y$	$g : C \rightarrow Y$	such that for all objects C together with morphisms $f : C \rightarrow X$ and $g : C \rightarrow Y$, there
fong-spivak-seven-sketches_121 F01352	121	1.000	$C \rightarrow X \times Y$	$C \rightarrow X \times Y$	exists a unique morphism $C \rightarrow X \times Y$, denoted $\langle f, g \rangle$, for which the following diagram
fong-spivak-seven-sketches_121 F01353	121	1.000	$\langle f, g \rangle$	$\langle f, g \rangle$	exists a unique morphism $C \rightarrow X \times Y$, denoted $\langle f, g \rangle$, for which the following diagram
fong-spivak-seven-sketches_121 F01354	121	1.000	$p_X(x, y) := x$	$p_X(x, y) := x$	$p_X(x, y) := x$ and $p_Y(x, y) := y$.
fong-spivak-seven-sketches_121 F01355	121	1.000	$p_Y(x, y) := y$	$p_Y(x, y) := y$	$p_X(x, y) := x$ and $p_Y(x, y) := y$.
fong-spivak-seven-sketches_121 F01356	121	1.000	$\langle f, g \rangle(c) := (f(c), g(c))$	$\langle f, g \rangle(c) := (f(c), g(c))$	to $X \times Y$ such that the required diagram commutes is given by $\langle f, g \rangle(c) := (f(c), g(c))$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F01357	121	1.000	$\underline{6} \times \underline{4}$	$\underline{6} \times \underline{4}$	Here is a picture of the product $\underline{6} \times \underline{4}$ of sets $\underline{6}$ and $\underline{4}$.
fong-spivak-seven-sketches-F01358	121	1.000	$\underline{6}$	$\underline{6}$	Here is a picture of the product $\underline{6} \times \underline{4}$ of sets $\underline{6}$ and $\underline{4}$.
fong-spivak-seven-sketches-F01359	121	1.000	$\underline{4}$	$\underline{4}$	Here is a picture of the product $\underline{6} \times \underline{4}$ of sets $\underline{6}$ and $\underline{4}$.
fong-spivak-seven-sketches-F01360	121	0.943	$x, y \in P$	$x, y \in P$	Exercise 3.88. Let $(P, \leq)$ be a preorder, let $x, y \in P$ be elements, and let $\mathcal{P}$ be the
fong-spivak-seven-sketches-F01361	121	1.000	$x \wedge y$	$x \wedge y$	corresponding category. Show that the product $x \times y$ in $\mathcal{P}$ agrees with their meet $x \wedge y$
fong-spivak-seven-sketches-F01362	122	0.997	$\mathcal{C} \times \mathcal{D}$	$\mathcal{C} \times \mathcal{D}$	Example 3.89. Given two categories $\mathcal{C}$ and $\mathcal{D}$, their product $\mathcal{C} \times \mathcal{D}$ may be given as
fong-spivak-seven-sketches-F01363	122	1.000	(c, d)	(c, d)	follows. The objects of this category are pairs (c, d) , where c is an object of $\mathcal{C}$ and d is an
fong-spivak-seven-sketches-F01364	122	0.887	$(c, d) \rightarrow (c', d')$	$(c, d) \rightarrow (c', d')$	object of $\mathcal{D}$. Similarly, morphisms $(c, d) \rightarrow (c', d')$ are pairs (f, g) where $f: c \rightarrow c'$ is a
fong-spivak-seven-sketches-F01365	122	0.887	(f, g)	(f, g)	object of $\mathcal{D}$. Similarly, morphisms $(c, d) \rightarrow (c', d')$ are pairs (f, g) where $f: c \rightarrow c'$ is a
fong-spivak-seven-sketches-F01366	122	0.887	$f: c \rightarrow c'$	$f: c \rightarrow c'$	object of $\mathcal{D}$. Similarly, morphisms $(c, d) \rightarrow (c', d')$ are pairs (f, g) where $f: c \rightarrow c'$ is a
fong-spivak-seven-sketches-F01367	122	0.944	$(f, g), (f', g') = (f; f', g; g')$	$(f, g), (f', g') = (f; f', g; g')$	given by composing each entry in the pair separately, so $(f, g) \circ (f', g') = (f \circ f', g \circ g')$.
fong-spivak-seven-sketches-F01368	122	1.000	$\mathbf{1} \times \mathbf{2}$	$\mathbf{1} \times \mathbf{2}$	3. What is the product category $\mathbf{1} \times \mathbf{2}$?
fong-spivak-seven-sketches-F01369	122	0.941	$\mathcal{P} \times \mathcal{Q}$	$\mathcal{P} \times \mathcal{Q}$	4. What is the product category $\mathcal{P} \times \mathcal{Q}$ when P and Q are preorders and $\mathcal{P}$ and $\mathcal{Q}$ are
fong-spivak-seven-sketches-F01370	122	1.000	$X \xleftarrow{f} C \xrightarrow{g} Y$	$X \xleftarrow{f} C \xrightarrow{g} Y$	pair of maps $X \xleftarrow{f} C \xrightarrow{g} Y$, there exists a unique map $C \rightarrow X \times Y$ such that certain
fong-spivak-seven-sketches-F01371	122	0.623	(X, Y)	(X, Y)	we'll call $\mathbf{Cone}(X, Y)$, the category of cones over X and Y in $\mathcal{C}$. We will see in Exercise 3.91
fong-spivak-seven-sketches-F01372	122	0.437	$X \xleftarrow{p_X} X \times Y \xrightarrow{p_Y} Y$	$X \xleftarrow{p_X} X \times Y \xrightarrow{p_Y} Y$	that $X \xleftarrow{p_X} X \times Y \xrightarrow{p_Y} Y$ is a terminal object in $\mathbf{Cone}(X, Y)$.
fong-spivak-seven-sketches-F01373	122	0.437	$\mathbf{Cone}(X, Y)$	$\mathbf{Cone}(X, Y)$	that $X \xleftarrow{p_X} X \times Y \xrightarrow{p_Y} Y$ is a terminal object in $\mathbf{Cone}(X, Y)$.
fong-spivak-seven-sketches-F01374	122	0.857	$X \xleftarrow{f'} C' \xrightarrow{g'} Y$	$X \xleftarrow{f'} C' \xrightarrow{g'} Y$	$X \xleftarrow{f} C \xrightarrow{g} Y$ to $X \xleftarrow{f'} C' \xrightarrow{g'} Y$ in $\mathbf{Cone}(X, Y)$ is a morphism $a: C \rightarrow C'$ in $\mathcal{C}$ such that
fong-spivak-seven-sketches-F01375	122	0.857	$a: C \rightarrow C'$	$a: C \rightarrow C'$	$X \xleftarrow{f} C \xrightarrow{g} Y$ to $X \xleftarrow{f'} C' \xrightarrow{g'} Y$ in $\mathbf{Cone}(X, Y)$ is a morphism $a: C \rightarrow C'$ in $\mathcal{C}$ such that
fong-spivak-seven-sketches-F01376	122	0.714	$X \xleftrightarrow{p_X} X \times Y \xrightarrow{p_Y} Y$	$X \xleftrightarrow{p_X} X \times Y \xrightarrow{p_Y} Y$	Exercise 3.91. Check that a product $X \xleftrightarrow{p_X} X \times Y \xrightarrow{p_Y} Y$ is exactly the same as a terminal
fong-spivak-seven-sketches-F01377	123	0.659	$\mathcal{J}$	$\mathcal{J}$	$D: \mathcal{J} \rightarrow \mathcal{C}$. Here $\mathcal{J}$ is called the indexing category of the diagram D .
fong-spivak-seven-sketches-F01378	123	0.996	(C, c_*)	(C, c_*)	Definition 3.92. Let $D: \mathcal{J} \rightarrow \mathcal{C}$ be a diagram. A cone (C, c_*) over D consists of
fong-spivak-seven-sketches-F01379	123	1.000	$C \in \mathcal{C}$	$C \in \mathcal{C}$	(i) an object $C \in \mathcal{C}$;
fong-spivak-seven-sketches-F01380	123	0.915	$j \in \mathcal{J}$	$j \in \mathcal{J}$	(ii) for each object $j \in \mathcal{J}$, a morphism $c_j: C \rightarrow D(j)$.
fong-spivak-seven-sketches-F01381	123	0.915	$c_j: C \rightarrow D(j)$	$c_j: C \rightarrow D(j)$	(ii) for each object $j \in \mathcal{J}$, a morphism $c_j: C \rightarrow D(j)$.
fong-spivak-seven-sketches-F01382	123	0.672	$f: j \rightarrow k$	$f: j \rightarrow k$	(a) for each $f: j \rightarrow k$ in $\mathcal{J}$, we have $c_k = c_j \circ D(f)$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F01383	123	0.672	$c_{\{k\}}=c_{\{j\}} \ ; \ D(f)$	$c_k = c_j; D(f)$	(a) for each $f: j \rightarrow k$ in $\mathcal{J}$, we have $c_k = c_j \circ D(f)$.
fong-spivak-seven-sketches F01384	123	0.997	$\left(C, c_{\{*\}}\right) \rightarrow \left(C^{\{\prime\}}, c_{\{*\}}^{\{\prime\}}\right)$	$(C, c_*) \rightarrow (C', c'_*)$	A morphism of cones $[C, c_*] \rightarrow [C', c'_*]$ is a morphism $a: C \rightarrow C'$ in $\mathcal{C}$ such that for all
fong-spivak-seven-sketches F01385	123	0.271	$c_{\{j\}}=a$	$c_j = a$	$j \in \mathcal{J}$ we have $c_j = a \circ c'_j$. Cones over D , and their morphisms, form a category $\mathbf{Cone}(D)$.
fong-spivak-seven-sketches F01386	123	0.271	$c_{\{j\}}^{\{\prime\}}$	c'_j	$j \in \mathcal{J}$ we have $c_j = a \circ c'_j$. Cones over D , and their morphisms, form a category $\mathbf{Cone}(D)$.
fong-spivak-seven-sketches F01387	123	0.271	$\operatorname{Cone}(D)$	$\mathbf{Cone}(D)$	$j \in \mathcal{J}$ we have $c_j = a \circ c'_j$. Cones over D , and their morphisms, form a category $\mathbf{Cone}(D)$.
fong-spivak-seven-sketches F01388	123	0.742	$\lim (D)$	$\lim(D)$	The limit of D , denoted $\lim(D)$, is the terminal object in the category $\mathbf{Cone}(D)$. Say
fong-spivak-seven-sketches F01389	123	0.997	$\lim (D)=\left(C, c_{\{*\}}\right)$	$\lim(D) = (C, c_*)$	it is the cone $\lim(D) = (C, c_*)$; we refer to C as the <i>limit object</i> and the map c_j for any
fong-spivak-seven-sketches F01390	123	0.997	$c_{\{j\}}$	c_j	it is the cone $\lim(D) = (C, c_*)$; we refer to C as the <i>limit object</i> and the map c_j for any
fong-spivak-seven-sketches F01391	123	0.995	$j^{\{\text{th}\}}$	j^{th}	$j \in \mathcal{J}$ as the j^{th} projection map.
fong-spivak-seven-sketches F01392	123	0.998	$\mathcal{J}=\varnothing$	$\mathcal{J} = \emptyset$	Example 3.93. Terminal objects are limits where the indexing category is empty, $\mathcal{J} = \emptyset$.
fong-spivak-seven-sketches F01393	123	0.198	v, w	v, w	v, w and no arrows, $\mathcal{J} = \begin{array}{ c c } \hline v & w \\ \hline \bullet & \bullet \\ \hline \end{array}$.
fong-spivak-seven-sketches F01394	123	0.198	$\mathcal{J}=$	$\mathcal{J} =$	v, w and no arrows, $\mathcal{J} = \begin{array}{ c c } \hline v & w \\ \hline \bullet & \bullet \\ \hline \end{array}$.
fong-spivak-seven-sketches F01395	123	0.198	$\cdot$	$\cdot$	v, w and no arrows, $\mathcal{J} = \begin{array}{ c c } \hline v & w \\ \hline \bullet & \bullet \\ \hline \end{array}$.
fong-spivak-seven-sketches F01396	124	0.879	V, A, s, t	V, A, s, t	Theorem 3.95. Let $\mathcal{J}$ be a category presented by the finite graph (V, A, s, t) together with
fong-spivak-seven-sketches F01397	124	0.994	$D: \mathcal{J} \rightarrow \mathbf{Set}$	$D: \mathcal{J} \rightarrow$	some equations, and let $D: \mathcal{J} \rightarrow \mathbf{Set}$ be a set-valued functor. Write $V = \{v_1, \dots, v_n\}$.
fong-spivak-seven-sketches F01398	124	0.994	$V=\left\{v_{\{1\}}, \ldots, v_{\{n\}}\right\}$	$V = \{v_1, \dots, v_n\}$	some equations, and let $D: \mathcal{J} \rightarrow \mathbf{Set}$ be a set-valued functor. Write $V = \{v_1, \dots, v_n\}$.
fong-spivak-seven-sketches F01399	124	1.000	$p_{\{i\}}: \left(\lim _{\mathcal{J}} D\right) \rightarrow D\left(v_{\{i\}}\right)$	$p_i: \left(\lim _{\mathcal{J}} D\right) \rightarrow D\left(v_i\right)$	together with the projection maps $p_i: (\lim_{\mathcal{J}} D) \rightarrow D(v_i)$ given by $p_i(d_1, \dots, d_n) := d_i$,
fong-spivak-seven-sketches F01400	124	1.000	$p_{\{i\}}\left(d_{\{1\}}, \ldots, d_{\{n\}}\right):=d_{\{i\}}$	$p_i\left(d_1, \ldots, d_n\right):=d_i$	together with the projection maps $p_i: (\lim_{\mathcal{J}} D) \rightarrow D(v_i)$ given by $p_i(d_1, \dots, d_n) := d_i$,
fong-spivak-seven-sketches F01401	124	0.999	$\mathcal{J}$	J	Example 3.96. If J is the empty graph $\square$, then $n = 0$: there are no vertices. There is ex-
fong-spivak-seven-sketches F01402	124	0.999	$n=0$	$n = 0$	Example 3.96. If J is the empty graph $\square$, then $n = 0$: there are no vertices. There is ex-
fong-spivak-seven-sketches F01403	124	1.000	$D: \mathbf{1} \rightarrow \mathbf{Set}$	$D: \mathbf{1} \rightarrow$	Exercise 3.98. If $D: \mathbf{1} \rightarrow \mathbf{Set}$ is a functor, what is the limit of D ? Compute it using
fong-spivak-seven-sketches F01404	124	0.784	$\left(d_{\{1\}}, \ldots, d_{\{n\}}\right)$	$(d_1, \dots, d_n)$	$d_1, \dots, d_n$ such that $D(a)(d_i) = d_j$ for each morphism $a: i \rightarrow j$ in $\mathcal{J}$ allows us to use
fong-spivak-seven-sketches F01405	124	0.784	$D(a)\left(d_{\{i\}}\right)=d_{\{j\}}$	$D(a)\left(d_i\right)=d_j$	$(d_1, \dots, d_n)$ such that $D(a)(d_i) = d_j$ for each morphism $a: i \rightarrow j$ in $\mathcal{J}$ allows us to use
fong-spivak-seven-sketches F01406	124	0.784	$a: i \rightarrow j$	$a: i \rightarrow j$	$(d_1, \dots, d_n)$ such that $D(a)(d_i) = d_j$ for each morphism $a: i \rightarrow j$ in $\mathcal{J}$ allows us to use
fong-spivak-seven-sketches F01407	124	0.986	$\stackrel{x}{\bullet} \rightarrow \stackrel{a}{\bullet} \stackrel{g}{\leftarrow} \stackrel{y}{\bullet}$	$\stackrel{x}{\bullet} \xrightarrow{f} \stackrel{a}{\bullet} \xleftarrow{g} \stackrel{y}{\bullet}$	Example 3.99. If J is presented by the <i>cospan</i> graph $\begin{array}{ c c c } \hline x & \xrightarrow{f} & a & \xleftarrow{g} & y \\ \hline \bullet & & \bullet & & \bullet \\ \hline \end{array}$, then its limit is

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F01408	124	■ 0.999	$X \xrightarrow{f} A \stackrel{g}{\leftarrow} Y$	$X \xrightarrow{f} A \xleftarrow{g} Y$	known as a <i>pullback</i> . Given the diagram $X \xrightarrow{f} A \xleftarrow{g} Y$, the pullback is the cone shown
fong-spivak-seven-sketches-F01409	125	■ 1.000	$c_{\{a\}}$	c_a	The fact that the diagram commutes means that the diagonal arrow c_a is in some
fong-spivak-seven-sketches-F01410	125	■ 0.579	$X \times_A Y$	$X \times_A Y$	naming the cone point $X \times_A Y$, and adding the $\lrcorner$ symbol, as shown to the right above.
fong-spivak-seven-sketches-F01411	125	■ 1.000	$X = \underline{6}, Y = \underline{4}$	$X = \underline{6}, Y = \underline{4}$	Here is a picture to help us unpack the definition in Set . We take $X = \underline{6}, Y = \underline{4}$, and
fong-spivak-seven-sketches-F01412	125	■ 1.000	$f: \underline{6} \rightarrow A$	$f: \underline{6} \rightarrow A$	The functions $f: \underline{6} \rightarrow A$ and $g: \underline{4} \rightarrow A$ are expressed in the coloring of the dots: for
fong-spivak-seven-sketches-F01413	125	■ 1.000	$g: \underline{4} \rightarrow A$	$g: \underline{4} \rightarrow A$	The functions $f: \underline{6} \rightarrow A$ and $g: \underline{4} \rightarrow A$ are expressed in the coloring of the dots: for
fong-spivak-seven-sketches-F01414	125	■ 1.000	$g(2)=g(4)=$	$g(2) = g(4) =$	example, $\overline{g(2) = g(4)} =$ red, while $f(5) =$ black. The pullback selects pairs $(i, j) \in \underline{6} \times \underline{4}$
fong-spivak-seven-sketches-F01415	125	■ 1.000	$f(5)=$	$f(5) =$	example, $g(2) = g(4) =$ red, while $f(5) =$ black. The pullback selects pairs $(i, j) \in \underline{6} \times \underline{4}$
fong-spivak-seven-sketches-F01416	125	■ 1.000	$(i, j) \in \underline{6} \times \underline{4}$	$(i, j) \in \underline{6} \times \underline{4}$	example, $g(2) = g(4) =$ red, while $f(5) =$ black. The pullback selects pairs $\overline{(i, j) \in \underline{6} \times \underline{4}}$
fong-spivak-seven-sketches-F01417	125	■ 1.000	$f(i)$	$f(i)$	such that $f(i)$ and $g(j)$ have the same color.
fong-spivak-seven-sketches-F01418	125	■ 1.000	$g(j)$	$g(j)$	such that $f(i)$ and $g(j)$ have the same color.
fong-spivak-seven-sketches-F01419	125	■ 0.516	$F^{\text{op}}: \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}^{\text{op}}$	$F^{\text{op}}: \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}^{\text{op}}$	Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor. How should we define its opposite, $F^{\text{op}}: \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}^{\text{op}}$? That
fong-spivak-seven-sketches-F01420	126	■ 0.997	F^{op}	F^{op}	is, how should F^{op} act on objects, and how should it act on morphisms?
fong-spivak-seven-sketches-F01421	126	■ 0.981	$D^{\text{op}}: \mathcal{J}^{\text{op}} \rightarrow \mathcal{C}^{\text{op}}$	$D^{\text{op}}: \mathcal{J}^{\text{op}} \rightarrow \mathcal{C}^{\text{op}}$	Given a diagram $D: \mathcal{J} \rightarrow \mathcal{C}$, we may take the limit of the functor $D^{\text{op}}: \mathcal{J}^{\text{op}} \rightarrow \mathcal{C}^{\text{op}}$.
fong-spivak-seven-sketches-F01422	126	■ 1.000	Δ, Σ, Π	Δ, Σ, Π	have all three data migration functors Δ, Σ, Π ; see [Spi12; SW15b]. The version of this
fong-spivak-seven-sketches-F01423	130	■ 1.000	$\$10 \leq \$20, 5W \leq 6W$	$\$10 \leq \$20, 5W \leq 6W$	$\overline{\$10 \leq \$20, 5W \leq 6W}$ and so on.
fong-spivak-seven-sketches-F01424	131	■ 1.000	$(p, r) \in P \times R$	$(p, r) \in P \times R$	every pair $(p, r) \in P \times R$, where P is the preorder of resources to be produced and R
fong-spivak-seven-sketches-F01425	131	■ 1.000	$\Phi: P \times R \rightarrow$	$\Phi: P \times R \rightarrow$	Feasibility relations hence define a function $\Phi: P \times R \rightarrow \mathbf{Bool}$. For a function
fong-spivak-seven-sketches-F01426	131	■ 1.000	$\Phi(p, r) =$	$\Phi(p, r) =$	(a) If $\Phi(p, r) = \mathbf{true}$ and $p' \leq p$, then $\Phi(p', r) = \mathbf{true}$.
fong-spivak-seven-sketches-F01427	131	■ 1.000	$p^{\text{prime}} \leq p$	$p' \leq p$	(a) If $\Phi(p, r) = \mathbf{true}$ and $p' \leq p$, then $\Phi(p', r) = \mathbf{true}$.
fong-spivak-seven-sketches-F01428	131	■ 1.000	$\Phi(p^{\text{prime}}, r) =$	$\Phi(p', r) =$	(a) If $\Phi(p, r) = \mathbf{true}$ and $p' \leq p$, then $\Phi(p', r) = \mathbf{true}$.
fong-spivak-seven-sketches-F01429	131	■ 1.000	$r \leq r^{\text{prime}}$	$r \leq r'$	(b) If $\Phi(p, r) = \mathbf{true}$ and $r \leq r'$ then $\Phi(p, r') = \mathbf{true}$.
fong-spivak-seven-sketches-F01430	131	■ 1.000	$\Phi(p, r^{\text{prime}}) =$	$\Phi(p, r') =$	(b) If $\Phi(p, r) = \mathbf{true}$ and $r \leq r'$ then $\Phi(p, r') = \mathbf{true}$.
fong-spivak-seven-sketches-F01431	131	■ 1.000	$r^{\text{prime}} \geq r$	$r' \geq r$	(b) also produce p given more resources $r' \geq r$. We will see that these two conditions
fong-spivak-seven-sketches-F01432	131	■ 0.685	$P^{\text{op}} \times R \rightarrow$	$P^{\text{op}} \times R \rightarrow$	are formalized by requiring Φ to be a monotone map $P^{\text{op}} \times R \rightarrow \mathbf{Bool}$.
fong-spivak-seven-sketches-F01433	131	■ 0.516	$5W \leq 10W$	$5W \leq 10W$	wattage, if $5W \leq 10W$, it means that whenever we are provided $10W$, we implicitly also

Identifier	Page	Conf.	LaTeX source	Rendered	Image															
fong-spivak-seven-sketches-F01434	132	0.500	$X=(X, \leq)$	$X = (X, \leq)$	profunctors as feasibility relations. Recall that if $\mathcal{X} = (X, \leq)$ is a preorder, then its															
fong-spivak-seven-sketches-F01435	132	0.982	$X^{\mathrm{\mathrm{op}}}=(X, \geq)$	$X^{\mathrm{op}} = (X, \geq)$	opposite $\mathcal{X}^{\mathrm{op}} = (X, \geq)$ has $x \geq y$ iff $y \leq x$.															
fong-spivak-seven-sketches-F01436	132	0.982	$x \geq y$	$x \geq y$	opposite $\mathcal{X}^{\mathrm{op}} = (X, \geq)$ has $x \geq y$ iff $y \leq x$.															
fong-spivak-seven-sketches-F01437	132	0.521	$X=\mathrm{\left(X, \leq_{\mathcal{X}}\right)}$	$X = (X, \leq_X)$	Definition 4.2. Let $\mathcal{X} = (X, \leq_X)$ and $\mathcal{Y} = (Y, \leq_Y)$ be preorders. A <i>feasibility relation</i> for $\mathcal{X}$															
fong-spivak-seven-sketches-F01438	132	0.521	$y=\mathrm{\left(Y, \leq_{\mathcal{Y}}\right)}$	$y = (Y, \leq_Y)$	Definition 4.2. Let $\mathcal{X} = (X, \leq_X)$ and $\mathcal{Y} = (Y, \leq_Y)$ be preorders. A <i>feasibility relation</i> for $\mathcal{X}$															
fong-spivak-seven-sketches-F01439	132	0.521	$\mathfrak{X}$	$\mathfrak{X}$	Definition 4.2. Let $\mathcal{X} = (X, \leq_X)$ and $\mathcal{Y} = (Y, \leq_Y)$ be preorders. A <i>feasibility relation</i> for $\mathcal{X}$															
fong-spivak-seven-sketches-F01440	132	0.775	$\Phi: \mathcal{X} \rightarrowtail y$	$\Phi: \mathcal{X} \rightarrowtail y$	We denote this by $\Phi: \mathcal{X} \rightarrowtail \mathcal{Y}$.															
fong-spivak-seven-sketches-F01441	132	1.000	$\Phi(x, y)=$	$\Phi(x, y) =$	Given $x \in X$ and $y \in Y$, if $\Phi(x, y) = \text{true}$ we say <i>x can be obtained given y</i> .															
fong-spivak-seven-sketches-F01442	132	1.000	$x^{\mathrm{\prime}} \leq_{\mathcal{X}} x$	$x' \leq_X x$	if $x' \leq_X x$ and $y \leq_Y y'$ then $\Phi(x, y) \leq_{\mathbf{Bool}} \Phi(x', y')$. In other words, if x can be															
fong-spivak-seven-sketches-F01443	132	1.000	$y \leq_{\mathcal{Y}} y^{\mathrm{\prime}}$	$y \leq_Y y'$	if $x' \leq_X x$ and $y \leq_Y y'$ then $\Phi(x, y) \leq_{\mathbf{Bool}} \Phi(x', y')$. In other words, if x can be															
fong-spivak-seven-sketches-F01444	132	1.000	$\Phi(x, y) \leq_{\text{Bool}} \Phi(\mathrm{\left(x^{\mathrm{\prime}}, y^{\mathrm{\prime}}\right)})$	$\Phi(x, y) \leq_{\mathbf{Bool}} \Phi(x', y')$	if $x' \leq_X x$ and $y \leq_Y y'$ then $\Phi(x, y) \leq_{\mathbf{Bool}} \Phi(x', y')$. In other words, if x can be															
fong-spivak-seven-sketches-F01445	132	1.000	$x^{\mathrm{\prime}}$	x'	obtained given y , and if x' is available given x , then x' can be obtained given y . And if															
fong-spivak-seven-sketches-F01446	132	0.605	$\mathcal{X}^{\mathrm{\mathrm{op}}} \times \mathcal{Y}$	$\mathcal{X}^{\mathrm{op}} \times \mathcal{Y}$	1. Draw the Hasse diagram for the preorder $\mathcal{X}^{\mathrm{op}} \times \mathcal{Y}$.															
fong-spivak-seven-sketches-F01447	132	0.994	$\Lambda: \mathcal{X} \rightarrowtail y$	$\Lambda: \mathcal{X} \rightarrowtail y$	2. Write down a profunctor $\Lambda: \mathcal{X} \rightarrowtail \mathcal{Y}$ and, reading $\Lambda(x, y) = \text{true}$ as “my aunt can															
fong-spivak-seven-sketches-F01448	132	0.994	$\Lambda(x, y)=$	$\Lambda(x, y) =$	2. Write down a profunctor $\Lambda: \mathcal{X} \rightarrowtail \mathcal{Y}$ and, reading $\Lambda(x, y) = \text{true}$ as “my aunt can															
fong-spivak-seven-sketches-F01449	132	0.372	$X^{\mathrm{\mathrm{op}}} \times y$	$X^{\mathrm{op}} \times y$	forms an upper set in $\mathcal{X}^{\mathrm{op}} \times \mathcal{Y}$.															
fong-spivak-seven-sketches-F01450	132	0.637	$\Rightarrow: \mathbb{B} \times \mathbb{B} \rightarrowtail \mathbb{B}$	$\Rightarrow: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$	joins $\vee$, and a closure operation, which we’ll write $\Rightarrow: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$. By definition, this															
fong-spivak-seven-sketches-F01451	132	1.000	$b, c, d \in \mathbb{B}$	$b, c, d \in \mathbb{B}$	operation satisfies the property that for all $b, c, d \in \mathbb{B}$ one has															
fong-spivak-seven-sketches-F01452	133	0.986	$c \rightarrowtail d$	$c \Rightarrow d$	<table><tr><th>c</th><th>d</th><th>$c \Rightarrow d$</th></tr><tr><td>true</td><td>true</td><td>true</td></tr><tr><td>true</td><td>false</td><td>false</td></tr><tr><td>false</td><td>true</td><td>true</td></tr><tr><td>false</td><td>false</td><td>true</td></tr></table>	c	d	$c \Rightarrow d$	true	true	true	true	false	false	false	true	true	false	false	true
c	d	$c \Rightarrow d$																		
true	true	true																		
true	false	false																		
false	true	true																		
false	false	true																		
fong-spivak-seven-sketches-F01453	133	0.671	$\Phi: \mathrm{Ob}(\mathcal{X}) \times \mathrm{Ob}(\mathcal{Y}) \rightarrowtail V$	$\Phi: \mathrm{Ob}(\mathcal{X}) \times \mathrm{Ob}(\mathcal{Y}) \rightarrowtail V$	$\Phi: \mathrm{Ob}(\mathcal{X}) \times \mathrm{Ob}(\mathcal{Y}) \rightarrowtail V$ such that for any $x, x' \in \mathcal{X}$ and $y, y' \in \mathcal{Y}$ the following															
fong-spivak-seven-sketches-F01454	133	0.671	$x, x^{\mathrm{\prime}} \in \mathcal{X}$	$x, x' \in \mathcal{X}$	$\Phi: \mathrm{Ob}(\mathcal{X}) \times \mathrm{Ob}(\mathcal{Y}) \rightarrowtail V$ such that for any $x, x' \in \mathcal{X}$ and $y, y' \in \mathcal{Y}$ the following															
fong-spivak-seven-sketches-F01455	133	0.671	$y, y^{\mathrm{\prime}} \in \mathcal{Y}$	$y, y' \in \mathcal{Y}$	$\Phi: \mathrm{Ob}(\mathcal{X}) \times \mathrm{Ob}(\mathcal{Y}) \rightarrowtail V$ such that for any $x, x' \in \mathcal{X}$ and $y, y' \in \mathcal{Y}$ the following															
fong-spivak-seven-sketches-F01456	134	0.821	$\Phi: X \rightarrowtail Y$	$\Phi: X \rightarrowtail Y$	inition 4.8 in the case $\mathcal{V} = \mathbf{Bool}$. One way to imagine a Bool -profunctor $\Phi: X \rightarrowtail Y$ is in															
fong-spivak-seven-sketches-F01457	134	1.000	$W \leq N$	$W \leq N$	Both X and Y are preorders, e.g. with $W \leq N$ and $b \leq a$. With bridges coming from															

Identifier	Page	Conf.	LaTeX source	Rendered	Image																				
fong-spivak-seven-sketches-F01458	134	■ 1.000	N	N	get from points in city X to points in city Y . For example, since there is a path from N																				
fong-spivak-seven-sketches-F01459	134	■ 1.000	E	E	to e and E to a , we have $\Phi(N, e) = \text{true}$ and $\Phi(E, a) = \text{true}$. On the other hand, since																				
fong-spivak-seven-sketches-F01460	134	■ 1.000	$\Phi(N, e) =$	$\Phi(N, e) =$	to e and E to a , we have $\Phi(N, e) = \text{true}$ and $\Phi(E, a) = \text{true}$. On the other hand, since																				
fong-spivak-seven-sketches-F01461	134	■ 1.000	$\Phi(E, a) =$	$\Phi(E, a) =$	to e and E to a , we have $\Phi(N, e) = \text{true}$ and $\Phi(E, a) = \text{true}$. On the other hand, since																				
fong-spivak-seven-sketches-F01462	134	■ 1.000	W	W	there is no path from W to d , we have $\Phi(W, d) = \text{false}$.																				
fong-spivak-seven-sketches-F01463	134	■ 1.000	$\Phi(W, d) =$	$\Phi(W, d) =$	there is no path from W to d , we have $\Phi(W, d) = \text{false}$.																				
fong-spivak-seven-sketches-F01464	134	■ 1.000	$W \leq N \leq c \leq a$	$W \leq N \leq c \leq a$	$W < N < c < a$, etc. This is called the <i>collage</i> of Φ ; we'll explore this in more detail																				
fong-spivak-seven-sketches-F01465	134	■ 1.000	$\Phi(m, n) \in \mathbb{B}$	$\Phi(m, n) \in \mathbb{B}$	$\Phi(m, n) \in \mathbb{B}$. Fill out the Bool -matrix:																				
fong-spivak-seven-sketches-F01466	135	■ 0.637	M_{Φ}	M_{Φ}	<table><tr><th>M_{Φ}</th><th>x</th><th>y</th><th>z</th></tr><tr><td>A</td><td>∞</td><td>∞</td><td>∞</td></tr><tr><td>B</td><td>11</td><td>∞</td><td>∞</td></tr><tr><td>C</td><td>∞</td><td>∞</td><td>∞</td></tr><tr><td>D</td><td>∞</td><td>9</td><td>∞</td></tr></table>	M_{Φ}	x	y	z	A	∞	∞	∞	B	11	∞	∞	C	∞	∞	∞	D	∞	9	∞
M_{Φ}	x	y	z																						
A	∞	∞	∞																						
B	11	∞	∞																						
C	∞	∞	∞																						
D	∞	9	∞																						
fong-spivak-seven-sketches-F01467	135	■ 1.000	$d_X * M_{\Phi} * d_Y$	$d_X * M_{\Phi} * d_Y$	The matrix of distances for the profunctor Φ will be equal to $d_X * M_{\Phi} * d_Y$. In fact, since																				
fong-spivak-seven-sketches-F01468	135	■ 0.835	$\Phi = M_X^3 * M_{\Phi} * M_Y^2$	$\Phi = M_X^3 * M_{\Phi} * M_Y^2$	X has four elements and Y has three, we also know that $\Phi = M_X^3 * M_{\Phi} * M_Y^2$.																				
fong-spivak-seven-sketches-F01469	135	■ 1.000	$M_X^3 * M_{\Phi} * M_Y^2$	$M_X^3 * M_{\Phi} * M_Y^2$	<i>Exercise 4.17.</i> Calculate $M_Y^3 * M_{\Phi} * M_X^3$, remembering to do matrix multiplication																				
fong-spivak-seven-sketches-F01470	136	■ 0.873	$([0, \infty], \leq)$	$([0, \infty], \leq)$	load might be the preorder $([0, \infty], \leq)$, where an element $x \in [0, \infty]$ represents the																				
fong-spivak-seven-sketches-F01471	136	—	100, more_than_\	(not rendered)	—																				
fong-spivak-seven-sketches-F01472	136	■ 0.965	$\Phi: (T \times E) \mapsto \$$	$\Phi: (T \times E) \mapsto \$$	$\Phi: (T \times E) \twoheadrightarrow \$$. We represent this by the box																				
fong-spivak-seven-sketches-F01473	136	■ 0.695	T, E	T, E	where T, E , and $\$$ are the preorders drawn below:																				
fong-spivak-seven-sketches-F01474	136	■ 0.695	$\$$	$\$$	where T, E , and $\$$ are the preorders drawn below:																				
fong-spivak-seven-sketches-F01475	137	■ 0.940	$\$ 500 \text{~}\mathrm{K}$	$\$500 \text{~}\mathrm{K}$	This says, for example, that a good-natured but boring movie costs \$500K to produce																				
fong-spivak-seven-sketches-F01476	137	—	500K given \	(not rendered)	—																				
fong-spivak-seven-sketches-F01477	137	■ 0.928	$\$ 500$	$\$500$	given \$500K”, and “I can provide a mean and boring movie given a good-natured																				
fong-spivak-seven-sketches-F01478	137	—	IM indicates also that "I can provide a mean and boring movie given \	(not rendered)	—																				
fong-spivak-seven-sketches-F01479	137	■ 0.952	$\mathcal{P}, \mathcal{Q}$	$\mathcal{P}, \mathcal{Q}$	two of them in series. Imagine you have cities $\mathcal{P}, \mathcal{Q}$, and $\mathcal{R}$ and you have bridges—and																				
fong-spivak-seven-sketches-F01480	137	■ 0.952	$\mathcal{R}$	$\mathcal{R}$	two of them in series. Imagine you have cities $\mathcal{P}, \mathcal{Q}$, and $\mathcal{R}$ and you have bridges—and																				
fong-spivak-seven-sketches-F01481	138	■ 0.497	$\Phi: \mathcal{P} \mapsto \mathcal{Q}$	$\Phi: \mathcal{P} \mapsto \mathcal{Q}$	hence feasibility matrices—connecting these cities, say $\Phi: \mathcal{P} \twoheadrightarrow \mathcal{Q}$ and $\Psi: \mathcal{Q} \twoheadrightarrow \mathcal{R}$.																				
fong-spivak-seven-sketches-F01482	138	■ 0.497	$\Psi: \mathcal{Q} \rightarrow \mathcal{R}$	$\Psi: \mathcal{Q} \rightarrow \mathcal{R}$	hence feasibility matrices—connecting these cities, say $\Phi: \mathcal{P} \twoheadrightarrow \mathcal{Q}$ and $\Psi: \mathcal{Q} \twoheadrightarrow \mathcal{R}$.																				

Identifier	Page	Conf.	LaTeX source	Rendered	Image																									
fong-spivak-seven-sketches_138 F01483	138	1.000	Ψ	Ψ	The feasibility matrices for Φ (in blue) and Ψ (in red) are:																									
fong-spivak-seven-sketches_138 F01484	138	0.223	$\Phi_{\mathfrak{Y}}^{\mathfrak{Q}} \Psi$	$\Phi_{\mathfrak{Y}}^{\mathfrak{Q}} \Psi$	trying to give a feasibility matrix $\Phi \circ \Psi$ for getting from $\mathcal{P}$ to $\mathcal{R}$. How should this be																									
fong-spivak-seven-sketches_138 F01485	138	0.984	$p \in \mathcal{P}$	$p \in \mathcal{P}$	done? Basically, for every pair $p \in \mathcal{P}$ and $r \in \mathcal{R}$, the navigator searches through $\mathcal{Q}$ for a																									
fong-spivak-seven-sketches_138 F01486	138	0.984	$r \in \mathcal{R}$	$r \in \mathcal{R}$	done? Basically, for every pair $p \in \mathcal{P}$ and $r \in \mathcal{R}$, the navigator searches through $\mathcal{Q}$ for a																									
fong-spivak-seven-sketches_138 F01487	138	0.984	$\mathcal{Q}$	$\mathcal{Q}$	done? Basically, for every pair $p \in \mathcal{P}$ and $r \in \mathcal{R}$, the navigator searches through $\mathcal{Q}$ for a																									
fong-spivak-seven-sketches_138 F01488	138	0.795	$\Phi \circ \Psi$	$\Phi \circ \Psi$	<table><tr><th>$\Phi \circ \Psi$</th><th>x</th><th>y</th></tr><tr><td>N</td><td>false</td><td>true</td></tr><tr><td>E</td><td>false</td><td>true</td></tr><tr><td>W</td><td>true</td><td>true</td></tr><tr><td>S</td><td>true</td><td>true</td></tr></table>	$\Phi \circ \Psi$	x	y	N	false	true	E	false	true	W	true	true	S	true	true										
$\Phi \circ \Psi$	x	y																												
N	false	true																												
E	false	true																												
W	true	true																												
S	true	true																												
fong-spivak-seven-sketches_139 F01489	139	0.604	$\mathcal{X}, \mathcal{Y}$	$\mathcal{X}, \mathcal{Y}$	Definition 4.21. Let $\mathcal{V}$ be a quantale, let $\mathcal{X}, \mathcal{Y}$, and $\mathcal{Z}$ be $\mathcal{V}$ -categories, and let $\Phi: \mathcal{X} \rightarrow \mathcal{Y}$																									
fong-spivak-seven-sketches_139 F01490	139	0.604	$\mathcal{Z}$	$\mathcal{Z}$	Definition 4.21. Let $\mathcal{V}$ be a quantale, let $\mathcal{X}, \mathcal{Y}$, and $\mathcal{Z}$ be $\mathcal{V}$ -categories, and let $\Phi: \mathcal{X} \rightarrow \mathcal{Y}$																									
fong-spivak-seven-sketches_139 F01491	139	0.604	$\Phi: \mathcal{X} \rightarrow \mathcal{Y}$	$\Phi: \mathcal{X} \rightarrow \mathcal{Y}$	Definition 4.21. Let $\mathcal{V}$ be a quantale, let $\mathcal{X}, \mathcal{Y}$, and $\mathcal{Z}$ be $\mathcal{V}$ -categories, and let $\Phi: \mathcal{X} \rightarrow \mathcal{Y}$																									
fong-spivak-seven-sketches_139 F01492	139	0.666	$\Psi: \mathcal{Y} \rightarrow \mathcal{Z}$	$\Psi: \mathcal{Y} \rightarrow \mathcal{Z}$	and $\Psi: \mathcal{Y} \rightarrow \mathcal{Z}$ be $\mathcal{V}$ -profunctors. We define their <i>composite</i> , denoted $\Phi \circ \Psi: \mathcal{X} \rightarrow \mathcal{Z}$ by																									
fong-spivak-seven-sketches_139 F01493	139	0.666	$\Phi; \Psi: \mathcal{X} \rightarrow \mathcal{Z}$	$\Phi; \Psi: \mathcal{X} \rightarrow \mathcal{Z}$	and $\Psi: \mathcal{Y} \rightarrow \mathcal{Z}$ be $\mathcal{V}$ -profunctors. We define their <i>composite</i> , denoted $\Phi \circ \Psi: \mathcal{X} \rightarrow \mathcal{Z}$ by																									
fong-spivak-seven-sketches_139 F01494	139	0.405	$\Phi: \mathcal{X} \rightarrow \mathcal{Y}$	$\Phi: \mathcal{X} \rightarrow \mathcal{Y}$	<i>Exercise 4.22.</i> Consider the Cost -profunctors $\Phi: \mathcal{X} \rightarrow \mathcal{Y}$ and $\Psi: \mathcal{Y} \rightarrow \mathcal{Z}$ shown below:																									
fong-spivak-seven-sketches_139 F01495	139	0.600	$\Phi_{\mathfrak{Y}}^{\mathfrak{Q}} \Psi$	$\Phi_{\mathfrak{Y}}^{\mathfrak{Q}} \Psi$	<table><tr><th>$\Phi \circ \Psi$</th><th>p</th><th>q</th><th>r</th><th>s</th></tr><tr><td>A</td><td>?</td><td>24</td><td>?</td><td>?</td></tr><tr><td>B</td><td>?</td><td>?</td><td>?</td><td>?</td></tr><tr><td>C</td><td>?</td><td>?</td><td>?</td><td>?</td></tr><tr><td>D</td><td>?</td><td>?</td><td>9</td><td>?</td></tr></table>	$\Phi \circ \Psi$	p	q	r	s	A	?	24	?	?	B	?	?	?	?	C	?	?	?	?	D	?	?	9	?
$\Phi \circ \Psi$	p	q	r	s																										
A	?	24	?	?																										
B	?	?	?	?																										
C	?	?	?	?																										
D	?	?	9	?																										
fong-spivak-seven-sketches_139 F01496	139	0.564	$\mathcal{V}$	$\mathcal{V}$	Theorem 4.23. For any skeletal quantale $\mathcal{V}$, there is a category Prof $_{\mathcal{V}}$ whose objects are																									
fong-spivak-seven-sketches_139 F01497	139	0.426	$\mathcal{X} \rightarrow \mathcal{Y}$	$\mathcal{X} \rightarrow \mathcal{Y}$	$\mathcal{V}$ -categories $\mathcal{X}$, whose morphisms are $\mathcal{V}$ -profunctors $\mathcal{X} \rightarrow \mathcal{Y}$, and with composition																									
fong-spivak-seven-sketches_139 F01498	139	0.505	Bool	Bool	Definition 4.24. We define Feas $:=$ Prof Bool .																									
fong-spivak-seven-sketches_140 F01499	140	0.744	$\mathrm{U}_x: \mathcal{X} \rightarrow \mathcal{X}$	$\mathrm{U}_x: \mathcal{X} \rightarrow \mathcal{X}$	Define the <i>unit profunctor</i> $\mathrm{U}_{\mathcal{X}}: \mathcal{X} \rightarrow \mathcal{X}$ on a $\mathcal{V}$ -category $\mathcal{X}$ by the formula																									
fong-spivak-seven-sketches_140 F01500	140	0.988	$\mathcal{X}(x, y) \in \mathcal{V}$	$\mathcal{X}(x, y) \in \mathcal{V}$	pair of elements $x, y \in \mathcal{X}$ an hom-object $\mathcal{X}(x, y) \in \mathcal{V}$. The unit profunctor $\mathrm{U}_{\mathcal{X}}$ just																									
fong-spivak-seven-sketches_140 F01501	140	0.988	U_x	U_x	pair of elements $x, y \in \mathcal{X}$ an hom-object $\mathcal{X}(x, y) \in \mathcal{V}$. The unit profunctor $\mathrm{U}_{\mathcal{X}}$ just																									
fong-spivak-seven-sketches_140 F01502	140	0.666	$\mathrm{U}_{\mathcal{X}}: \mathcal{X} \rightarrow \mathcal{X}$	$\mathrm{U}_{\mathcal{X}}: \mathcal{X} \rightarrow \mathcal{X}$	for the unit profunctor $\mathrm{U}_{\mathcal{X}}: \mathcal{X} \rightarrow \mathcal{X}$.																									
fong-spivak-seven-sketches_140 F01503	140	0.982	$\Phi: \mathcal{P} \rightarrow \mathcal{Q}$	$\Phi: \mathcal{P} \rightarrow \mathcal{Q}$	Lemma 4.27. Composing any profunctor $\Phi: \mathcal{P} \rightarrow \mathcal{Q}$ with either unit profunctor, $\mathrm{U}_{\mathcal{P}}$ or																									
fong-spivak-seven-sketches_140 F01504	140	0.982	$\mathrm{U}_{\mathcal{P}}$	$\mathrm{U}_{\mathcal{P}}$	Lemma 4.27. Composing any profunctor $\Phi: \mathcal{P} \rightarrow \mathcal{Q}$ with either unit profunctor, $\mathrm{U}_{\mathcal{P}}$ or																									

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_140 F01505	140	0.637	U_{Q}	U_Q	U_Q , returns Φ :
fong-spivak-seven-sketches_140 F01506	140	0.296	$\mathrm{U}_{\mathrm{P}}; \Phi = \Phi$	$U_{\mathcal{P}}; \Phi = \Phi$	<i>Proof.</i> We show that $U_{\mathcal{P}} \circ \Phi = \Phi$ holds; proving $\Phi = \Phi \circ U_Q$ is similar. Fix $p \in P$ and
fong-spivak-seven-sketches_140 F01507	140	0.296	$\Phi = \Phi$	$\Phi = \Phi$	<i>Proof.</i> We show that $U_{\mathcal{P}} \circ \Phi = \Phi$ holds; proving $\Phi = \Phi \circ U_Q$ is similar. Fix $p \in P$ and
fong-spivak-seven-sketches_140 F01508	140	0.673	$\Phi \leq \mathrm{U}_{\mathrm{P}}; \Phi$	$\Phi \leq U_{\mathcal{P}}; \Phi$	$q \in Q$. Since $\mathcal{V}$ is skeletal, to prove the equality it's enough to show $\Phi \leq U_{\mathcal{P}} \circ \Phi$ and
fong-spivak-seven-sketches_140 F01509	140	0.996	$\mathrm{U}_{\mathrm{P}}; \Phi \leq \Phi$	$U_{\mathcal{P}}; \Phi \leq \Phi$	$U_{\mathcal{P}} \circ \Phi \leq \Phi$. We have one direction:
fong-spivak-seven-sketches_140 F01510	140	1.000	$\bigvee_{p_1 \in P} \left(\mathcal{P}(p, p_1) \otimes \Phi(p_1, q) \right) \leq \Phi(p, q)$	$\bigvee_{p_1 \in P} (\mathcal{P}(p, p_1) \otimes \Phi(p_1, q)) \leq \Phi(p, q)$	For the other direction, we must show $\bigvee_{p_1 \in P} (\mathcal{P}(p, p_1) \otimes \Phi(p_1, q)) \leq \Phi(p, q)$. But by
fong-spivak-seven-sketches_140 F01511	140	1.000	$\mathcal{P}(p, p_1) \otimes \Phi(p_1, q) \leq \Phi(p, q)$	$\mathcal{P}(p, p_1) \otimes \Phi(p_1, q) \leq \Phi(p, q)$	definition of join, this holds iff $\mathcal{P}(p, p_1) \otimes \Phi(p_1, q) \leq \Phi(p, q)$ is true for each $p_1 \in P$. This
fong-spivak-seven-sketches_140 F01512	140	1.000	$p_1 \in P$	$p_1 \in \mathcal{P}$	definition of join, this holds iff $\mathcal{P}(p, p_1) \otimes \Phi(p_1, q) \leq \Phi(p, q)$ is true for each $p_1 \in P$. This
fong-spivak-seven-sketches_140 F01513	140	0.696	$(=, \leq, \leq, =)$	$(=, \leq, \leq, =)$	1. Justify each of the four steps $(=, \leq, \leq, =)$ in Eq. (4.28).
fong-spivak-seven-sketches_141 F01514	141	0.964	$\leq, \leq$	$\leq, \leq$	3. Justify each of the three steps $(=, \leq, \leq)$ in Eq. (4.29).
fong-spivak-seven-sketches_141 F01515	141	0.567	$\Phi: \mathcal{P} \rightarrow Q, \Psi: Q \rightarrow \mathcal{R}$	$\Phi: \mathcal{P} \rightarrow Q, \Psi: Q \rightarrow \mathcal{R}$	tors $\Phi: \mathcal{P} \rightarrow Q, \Psi: Q \rightarrow \mathcal{R}$, and $\Upsilon: \mathcal{R} \rightarrow S$, we have
fong-spivak-seven-sketches_141 F01516	141	0.567	$\Upsilon: \mathcal{R} \rightarrow S$	$\Upsilon: \mathcal{R} \rightarrow S$	tors $\Phi: \mathcal{P} \rightarrow Q, \Psi: Q \rightarrow \mathcal{R}$, and $\Upsilon: \mathcal{R} \rightarrow S$, we have
fong-spivak-seven-sketches_142 F01517	142	0.426	$\{ \}_{\mathrm{C}}$	C	the category $\mathbf{Cospan}_C$ in Definition 6.45. Second, we might relax what we mean by
fong-spivak-seven-sketches_142 F01518	142	0.977	$F: \mathcal{P} \rightarrow Q$	$F: \mathcal{P} \rightarrow Q$	Definition 4.34. Let $F: \mathcal{P} \rightarrow Q$ be a $\mathcal{V}$ -functor. The <i>companion</i> of F , denoted $\widehat{F}: \mathcal{P} \rightarrowtail Q$
fong-spivak-seven-sketches_142 F01519	142	0.977	$\widehat{F}: \mathcal{P} \rightarrow Q$	$\widehat{F}: \mathcal{P} \rightarrow Q$	Definition 4.34. Let $F: \mathcal{P} \rightarrow Q$ be a $\mathcal{V}$ -functor. The <i>companion</i> of F , denoted $\widehat{F}: \mathcal{P} \rightarrowtail Q$
fong-spivak-seven-sketches_142 F01520	142	0.488	$\check{F}: Q \mapsto \mathcal{P}$	$\check{F}: Q \mapsto \mathcal{P}$	and the <i>conjoint</i> of F , denoted $\check{F}: Q \rightarrowtail \mathcal{P}$ are defined to be the following $\mathcal{V}$ -profunctors:
fong-spivak-seven-sketches_142 F01521	142	0.995	$F(p) \in Q$	$F(p) \in Q$	$F(p) \in Q$.
fong-spivak-seven-sketches_142 F01522	142	1.000	$\widehat{F}$	$\widehat{F}$	picture in that light, they are seeing the companion $\widehat{F}$. But now mentally reverse every
fong-spivak-seven-sketches_142 F01523	142	0.979	$Q \rightarrow \mathcal{P}$	$Q \rightarrow \mathcal{P}$	dotted arrow, and the result would be bridges Q to $\mathcal{P}$. This is a profunctor $Q \rightarrowtail \mathcal{P}$! We
fong-spivak-seven-sketches_142 F01524	142	1.000	$\check{F}$	$\check{F}$	call it $\check{F}$.
fong-spivak-seven-sketches_142 F01525	142	0.710	$\mathrm{id}: \mathcal{P} \rightarrow \mathcal{P}$	$\mathrm{id}: \mathcal{P} \rightarrow \mathcal{P}$	<i>Example 4.35.</i> For any preorder $\mathcal{P}$, there is an identity functor $\mathrm{id}: \mathcal{P} \rightarrow \mathcal{P}$. Its compan-
fong-spivak-seven-sketches_142 F01526	142	0.699	$\widehat{\mathrm{id}} = \check{\mathrm{id}}: \mathcal{P} \rightarrowtail \mathcal{P}$	$\widehat{\mathrm{id}} = \check{\mathrm{id}}: \mathcal{P} \mapsto \mathcal{P}$	ion and conjoint agree $\widehat{\mathrm{id}} = \check{\mathrm{id}}: \mathcal{P} \rightarrowtail \mathcal{P}$. The resulting profunctor is in fact the unit
fong-spivak-seven-sketches_142 F01527	142	1.000	$\mathrm{U}_{\mathcal{P}}$	$U_{\mathcal{P}}$	profunctor, $U_{\mathcal{P}}$, as defined in Eq. (4.25).
fong-spivak-seven-sketches_143 F01528	143	0.991	$\widehat{\mathrm{id}}$	$\widehat{\mathrm{id}}$	<i>Exercise 4.36.</i> Check that the companion $\widehat{\mathrm{id}}$ of $\mathrm{id}: \mathcal{P} \rightarrow \mathcal{P}$ really has the unit profunctor
fong-spivak-seven-sketches_143 F01529	143	0.991	$\mathcal{P} \rightarrow \mathcal{P}$	$\mathcal{P} \rightarrow \mathcal{P}$	<i>Exercise 4.36.</i> Check that the companion $\widehat{\mathrm{id}}$ of $\mathrm{id}: \mathcal{P} \rightarrow \mathcal{P}$ really has the unit profunctor
fong-spivak-seven-sketches_143 F01530	143	0.637	$+: \mathbb{R} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$	$+: \mathbb{R} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$	<i>Example 4.37.</i> Consider the function $+: \mathbb{R} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, sending a triple (a, b, c) of real

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_F01531	143	■ 0.979	$a+b+c \in \mathbb{R}$	$a + b + c \in \mathbb{R}$	numbers to $a + b + c \in \mathbb{R}$. This function is monotonic, because if $(a, b, c) \leq (a', b', c')$ —
fong-spivak-seven-sketches_F01532	143	■ 0.979	$(a, b, c) \leq (a^{\prime}, b^{\prime}, c^{\prime})$	$(a, b, c) \leq (a', b', c')$	numbers to $a + b + c \in \mathbb{R}$. This function is monotonic, because if $(a, b, c) \leq (a', b', c')$ —
fong-spivak-seven-sketches_F01533	143	■ 1.000	$a \leq a^{\prime}$	$a \leq a'$	i.e. if $a \leq a'$ and $b \leq b'$, and $c \leq c'$ —then obviously $a + b + c \leq a' + b' + c'$. Thus it has
fong-spivak-seven-sketches_F01534	143	■ 1.000	$b \leq b^{\prime}$	$b \leq b'$	i.e. if $a \leq a'$ and $b \leq b'$, and $c \leq c'$ —then obviously $a + b + c \leq a' + b' + c'$. Thus it has
fong-spivak-seven-sketches_F01535	143	■ 1.000	$c \leq c^{\prime}$	$c \leq c'$	i.e. if $a \leq a'$ and $b \leq b'$, and $c \leq c'$ —then obviously $a + b + c \leq a' + b' + c'$. Thus it has
fong-spivak-seven-sketches_F01536	143	■ 1.000	$a+b+c \leq a^{\prime}+b^{\prime}+c^{\prime}$	$a + b + c \leq a' + b' + c'$	i.e. if $a \leq a'$ and $b \leq b'$, and $c \leq c'$ —then obviously $a + b + c \leq a' + b' + c'$. Thus it has
fong-spivak-seven-sketches_F01537	143	■ 0.955	$\widehat{+} : (\mathbb{R} \times \mathbb{R} \times \mathbb{R}) \rightarrow \mathbb{R}$	$\widehat{+} : (\mathbb{R} \times \mathbb{R} \times \mathbb{R}) \rightarrow \mathbb{R}$	Its companion $\widehat{+} : (\mathbb{R} \times \mathbb{R} \times \mathbb{R}) \rightarrow \mathbb{R}$ is the function that sends (a, b, c, d) to true if
fong-spivak-seven-sketches_F01538	143	■ 0.955	(a, b, c, d)	(a, b, c, d)	Its companion $\widehat{+} : (\mathbb{R} \times \mathbb{R} \times \mathbb{R}) \rightarrow \mathbb{R}$ is the function that sends (a, b, c, d) to true if
fong-spivak-seven-sketches_F01539	143	■ 1.000	$a+b+c \leq d$	$a + b + c \leq d$	$a + b + c \leq d$ and to false otherwise.
fong-spivak-seven-sketches_F01540	143	■ 1.000	$\check{+}$	$\check{+}$	<i>Exercise 4.38.</i> Let $+ : \mathbb{R} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ be as in Example 4.37. What is its conjoint $\check{+}$?
fong-spivak-seven-sketches_F01541	143	■ 0.994	$G : \mathcal{Q} \rightarrow \mathcal{P}$	$G : \mathcal{Q} \rightarrow \mathcal{P}$	of $\mathcal{V}$ -functors $F : \mathcal{P} \rightarrow \mathcal{Q}$ and $G : \mathcal{Q} \rightarrow \mathcal{P}$ such that the following holds for all $p \in P$ and
fong-spivak-seven-sketches_F01542	143	■ 0.990	$\widehat{F} = \widetilde{G}$	$\widehat{F} = \widetilde{G}$	of the former equals the conjoint of the latter: $\widehat{F} = \widetilde{G}$.
fong-spivak-seven-sketches_F01543	143	■ 1.000	$\widehat{\mathrm{id}} = \widetilde{\mathrm{id}}$	$\widehat{\mathrm{id}} = \widetilde{\mathrm{id}}$	2. Use this to prove that $\widehat{\mathrm{id}} = \widetilde{\mathrm{id}}$, as was stated in Example 4.35.
fong-spivak-seven-sketches_F01544	143	■ 0.998	$\mathbf{Col}(\Phi)$	$\mathbf{Col}(\Phi)$	a $\mathcal{V}$ -profunctor. The <i>collage of Φ</i> , denoted $\mathbf{Col}(\Phi)$ is the $\mathcal{V}$ -category defined as follows:
fong-spivak-seven-sketches_F01545	143	■ 0.976	$\mathrm{Ob}(\mathbf{Col}(\Phi)) := \mathrm{Ob}(\mathcal{X}) \sqcup \mathrm{Ob}(\mathcal{Y})$	$\mathrm{Ob}(\mathbf{Col}(\Phi)) := \mathrm{Ob}(\mathcal{X}) \sqcup \mathrm{Ob}(\mathcal{Y})$	(i) $\mathrm{Ob}(\mathbf{Col}(\Phi)) := \mathrm{Ob}(\mathcal{X}) \sqcup \mathrm{Ob}(\mathcal{Y})$;
fong-spivak-seven-sketches_F01546	143	■ 0.567	$a, b \in \mathrm{Ob}(\mathbf{Col}(\Phi))$	$a, b \in \mathrm{Ob}(\mathbf{Col}(\Phi))$	(ii) For any $a, b \in \mathrm{Ob}(\mathbf{Col}(\Phi))$, define $\mathbf{Col}(\Phi)(a, b) \in \mathcal{V}$ to be
fong-spivak-seven-sketches_F01547	143	■ 0.567	$\mathbf{Col}(\Phi)(a, b) \in \mathcal{V}$	$\mathbf{Col}(\Phi)(a, b) \in \mathcal{V}$	(ii) For any $a, b \in \mathrm{Ob}(\mathbf{Col}(\Phi))$, define $\mathbf{Col}(\Phi)(a, b) \in \mathcal{V}$ to be
fong-spivak-seven-sketches_F01548	143	■ 0.994	$i_x : \mathcal{X} \rightarrow \mathbf{Col}(\Phi)$	$i_x : \mathcal{X} \rightarrow \mathbf{Col}(\Phi)$	There are obvious functors $i_x : \mathcal{X} \rightarrow \mathbf{Col}(\Phi)$ and $i_y : \mathcal{Y} \rightarrow \mathbf{Col}(\Phi)$, sending each object
fong-spivak-seven-sketches_F01549	143	■ 0.994	$i_y : \mathcal{Y} \rightarrow \mathbf{Col}(\Phi)$	$i_y : \mathcal{Y} \rightarrow \mathbf{Col}(\Phi)$	There are obvious functors $i_x : \mathcal{X} \rightarrow \mathbf{Col}(\Phi)$ and $i_y : \mathcal{Y} \rightarrow \mathbf{Col}(\Phi)$, sending each object
fong-spivak-seven-sketches_F01550	144	■ 0.600	$\Phi : x \rightarrow y$	$\Phi : x \rightarrow y$	<i>Example 4.43.</i> Consider the following picture of a Cost -profunctor $\Phi : \mathcal{X} \rightarrow \mathcal{Y}$:
fong-spivak-seven-sketches_F01551	145	■ 1.000	$5+3=8$	$5 + 3 = 8$	that $5 + 3 = 8$. One way to categorify $\mathbb{N}$ is to use the category FinSet of finite sets
fong-spivak-seven-sketches_F01552	145	■ 0.709	$\overline{5} = \{$	$\overline{5} = \{$	sets of that many elements, say $\overline{5} = \{\text{apple, banana, cherry, dragonfruit, elephant}\}$,
fong-spivak-seven-sketches_F01553	145	■ 0.456	$\overline{3} = \{$	$\overline{3} = \{$	$\overline{3} = \{\text{apple, tomato, cantaloupe}\}$, and $\overline{8} = \{\text{Ali, Bob, Carl, Deb, Eli, Fritz, Gem, Helen}\}$
fong-spivak-seven-sketches_F01554	145	■ 0.456	$\overline{8} = \{$	$\overline{8} = \{$	$\overline{3} = \{\text{apple, tomato, cantaloupe}\}$, and $\overline{8} = \{\text{Ali, Bob, Carl, Deb, Eli, Fritz, Gem, Helen}\}$
fong-spivak-seven-sketches_F01555	145	■ 0.895	$\sqcup$	$\sqcup$	respectively. We also replace $+$ with disjoint union of sets $\sqcup$, and the brute property of
fong-spivak-seven-sketches_F01556	145	■ 0.998	$\overline{5} \sqcup \overline{3} \cong \overline{8}$	$\overline{5} \sqcup \overline{3} \cong \overline{8}$	is that, having made these replacements, the analogue of $5 + 3 = 8$ is still true: $\overline{5} \sqcup \overline{3} \cong \overline{8}$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_145 F01557	145	■ 1.000	$\overline{5} \sqcup \overline{3}$	$\overline{5} \sqcup \overline{3}$	Thus it's not the case that $\overline{5} \sqcup \overline{3}$ is <i>equal</i> to our chosen eight-element set $\overline{8}$, but more
fong-spivak-seven-sketches_145 F01558	145	■ 1.000	$\overline{8}$	$\overline{8}$	Thus it's not the case that $\overline{5} \sqcup \overline{3}$ is <i>equal</i> to our chosen eight-element set $\overline{8}$, but more
fong-spivak-seven-sketches_146 F01559	146	■ 1.000	$\mathcal{P}(a, b) =$	$\mathcal{P}(a, b) =$	exists a morphism $a \rightarrow b$," denoted $a \leq b$ or $\mathcal{P}(a, b) = \text{true}$, we instead say "here is a set
fong-spivak-seven-sketches_146 F01560	146	■ 1.000	$x_{\{0\}} \leq x_{\{1\}}$	$x_0 \leq x_1$	whenever $x_0 \leq x_1$. Chaining these together, we could prove facts in our preorder. For
fong-spivak-seven-sketches_146 F01561	146	■ 1.000	$x_{\{0\}} \leq x_{\{3\}}$	$x_0 \leq x_3$	provides a proof that $x_0 \leq x_3$ (the exterior box) using three facts (the interior boxes),
fong-spivak-seven-sketches_146 F01562	146	■ 1.000	$x_{\{0\}} \leq x_{\{1\}}, x_{\{1\}} \leq x_{\{2\}}$	$x_0 \leq x_1, x_1 \leq x_2$	$x_0 \leq x_1, x_1 \leq x_2$, and $x_2 \leq x_3$.
fong-spivak-seven-sketches_146 F01563	146	■ 1.000	$x_{\{2\}} \leq x_{\{3\}}$	$x_2 \leq x_3$	$x_0 \leq x_1, x_1 \leq x_2$, and $x_2 \leq x_3$.
fong-spivak-seven-sketches_146 F01564	146	■ 1.000	$g: B \rightarrow C$	$g: B \rightarrow C$	Suppose given additional morphisms $g: B \rightarrow C$, and $h: C \rightarrow D$. Representing these
fong-spivak-seven-sketches_146 F01565	146	■ 1.000	$h: C \rightarrow D$	$h: C \rightarrow D$	Suppose given additional morphisms $g: B \rightarrow C$, and $h: C \rightarrow D$. Representing these
fong-spivak-seven-sketches_146 F01566	146	■ 0.918	$f ; (g ; h)$	$f ; (g ; h)$	say! We don't know which morphism it is. Is it $f ; (g ; h)$? Or $(f ; g) ; h$? It's good that
fong-spivak-seven-sketches_146 F01567	146	■ 0.918	$(f ; g) ; h$	$(f ; g) ; h$	say! We don't know which morphism it is. Is it $f ; (g ; h)$? Or $(f ; g) ; h$? It's good that
fong-spivak-seven-sketches_146 F01568	146	■ 0.743	$f ; (g ; h) = (f ; g) ; h$	$f ; (g ; h) = (f ; g) ; h$	Associativity says $f ; (g ; h) = (f ; g) ; h$, so it doesn't matter which way we chose to try
fong-spivak-seven-sketches_147 F01569	147	■ 0.647	id_x	id_x	we will see that it is not harmful to draw id_x in any of the following three ways:
fong-spivak-seven-sketches_147 F01570	147	■ 0.926	$f = f = f$	$f = f = f$	property and the associativity property. The unitality says that $\mathrm{id}_x ; f = f = f ; \mathrm{id}_y$ for
fong-spivak-seven-sketches_147 F01571	147	■ 0.926	id_y	id_y	property and the associativity property. The unitality says that $\mathrm{id}_x ; f = f = f ; \mathrm{id}_y$ for
fong-spivak-seven-sketches_147 F01572	147	■ 1.000	$f: x \rightarrow y$	$f: x \rightarrow y$	any $f: x \rightarrow y$. In terms of diagrams this would say
fong-spivak-seven-sketches_148 F01573	148	■ 0.943	$\{1\}, \times$	$\{1\}, \times$	We will soon meet the monoidal category $(\mathbf{Set}, \{1\}, \times)$. The monoidal product will
fong-spivak-seven-sketches_148 F01574	148	■ 1.000	$S \times T = \{(s, t) \mid s \in S, t \in T\}$	$S \times T = \{(s, t) \mid s \in S, t \in T\}$	take two sets, S and T , and return the set $S \times T = \{(s, t) \mid s \in S, t \in T\}$. But whereas
fong-spivak-seven-sketches_148 F01575	148	■ 0.999	$(p \otimes q) \otimes r = p \otimes (q \otimes r)$	$(p \otimes q) \otimes r = p \otimes (q \otimes r)$	for monoidal preorders we had the brute associative property $(p \otimes q) \otimes r = p \otimes (q \otimes r)$,
fong-spivak-seven-sketches_148 F01576	148	■ 1.000	$T \times U$	$T \times U$	and an element in $T \times U$, while the second contains pairs consisting of an element in
fong-spivak-seven-sketches_148 F01577	148	■ 1.000	$S \times T$	$S \times T$	$S \times T$ and an element in U . The sets are not equal, but they are clearly isomorphic,
fong-spivak-seven-sketches_148 F01578	148	■ 1.000	$\alpha_{s, t, u}$	$\alpha_{s, t, u}$	example, above we replaced the associativity equation with an isomorphism $\alpha_{s, t, u}$, but
fong-spivak-seven-sketches_148 F01579	148	■ 0.731	$I \in \mathrm{Ob}(\mathcal{C})$	$I \in \mathrm{Ob}(\mathcal{C})$	(i) an object $I \in \mathrm{Ob}(\mathcal{C})$ called the <i>monoidal unit</i> , and
fong-spivak-seven-sketches_149 F01580	149	■ 0.990	$\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$	$\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$	(ii) a functor $\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$, called the <i>monoidal product</i>
fong-spivak-seven-sketches_149 F01581	149	■ 0.967	$\lambda_c: I \otimes c \cong c$	$\lambda_c: I \otimes c \cong c$	(a) $\lambda_c: I \otimes c \cong c$ for every $c \in \mathrm{Ob}(\mathcal{C})$,
fong-spivak-seven-sketches_149 F01582	149	■ 0.835	$\rho_c: c \otimes I \cong c$	$\rho_c: c \otimes I \cong c$	(b) $\rho_c: c \otimes I \cong c$ for every $c \in \mathrm{Ob}(\mathcal{C})$,
fong-spivak-seven-sketches_149 F01583	149	■ 0.981	$\alpha_{c, d, e}: (c \otimes d) \otimes e \cong c \otimes (d \otimes e)$	$\alpha_{c, d, e}: (c \otimes d) \otimes e \cong c \otimes (d \otimes e)$	(c) $\alpha_{c, d, e}: (c \otimes d) \otimes e \cong c \otimes (d \otimes e)$ for every $c, d, e \in \mathrm{Ob}(\mathcal{C})$, and

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_149 F01584	149	0.964	$\sigma_{\mathcal{C}, d} : c \otimes d \cong d \otimes c$	$\sigma_{\mathcal{C}, d} : c \otimes d \cong d \otimes c$	(d) $\sigma_{c,d} : c \otimes d \cong d \otimes c$ for every $c, d \in \text{Ob}(\mathcal{C})$, called the <i>swap map</i> , such that $\sigma \circ \sigma = \text{id}$.
fong-spivak-seven-sketches_149 F01585	149	0.964	$c, d \in \text{Ob}(\mathcal{C})$	$c, d \in \text{Ob}(\mathcal{C})$	(d) $\sigma_{c,d} : c \otimes d \cong d \otimes c$ for every $c, d \in \text{Ob}(\mathcal{C})$, called the <i>swap map</i> , such that $\sigma \circ \sigma = \text{id}$.
fong-spivak-seven-sketches_149 F01586	149	0.964	$\sigma \circ \sigma = \text{id}$	$\sigma \circ \sigma = \text{id}$	(d) $\sigma_{c,d} : c \otimes d \cong d \otimes c$ for every $c, d \in \text{Ob}(\mathcal{C})$, called the <i>swap map</i> , such that $\sigma \circ \sigma = \text{id}$.
fong-spivak-seven-sketches_149 F01587	149	0.999	$(\mathcal{P}, I, \otimes)$	$(\mathcal{P}, I, \otimes)$	a monoidal preorder is a monoidal category $(\mathcal{P}, I, \otimes)$ where, for every $p, q \in \mathcal{P}$, the set
fong-spivak-seven-sketches_149 F01588	149	0.999	$p, q \in \mathcal{P}$	$p, q \in \mathcal{P}$	a monoidal preorder is a monoidal category $(\mathcal{P}, I, \otimes)$ where, for every $p, q \in \mathcal{P}$, the set
fong-spivak-seven-sketches_149 F01589	149	1.000	$\mathcal{P}(p, q)$	$\mathcal{P}(p, q)$	$\mathcal{P}(p, q)$ has at most one element.
fong-spivak-seven-sketches_149 F01590	149	1.000	$I := \{1\}$	$I := \{1\}$	monoidal unit is some choice of singleton set, say $I := \{1\}$, and the monoidal product
fong-spivak-seven-sketches_149 F01591	149	0.513	$\otimes := \times$	$\otimes := \times$	is $\otimes := \times$. What it means that $\times$ is a functor is that:
fong-spivak-seven-sketches_149 F01592	149	0.513	$\times$	$\times$	is $\otimes := \times$. What it means that $\times$ is a functor is that:
fong-spivak-seven-sketches_149 F01593	149	0.537	$(S, T) \in \text{Ob}(\text{Set} \times \text{Set})$	$(S, T) \in \text{Ob}(\text{Set} \times \text{Set})$	• For any pair of objects, i.e. sets, $(S, T) \in \text{Ob}(\text{Set} \times \text{Set})$, one obtains a set $(S \times T) \in \text{Ob}(\text{Set})$.
fong-spivak-seven-sketches_149 F01594	149	0.537	$(S \times T) \in \text{Ob}(\text{Set})$	$(S \times T) \in \text{Ob}(\text{Set})$	• For any pair of objects, i.e. sets, $(S, T) \in \text{Ob}(\text{Set} \times \text{Set})$, one obtains a set $(S \times T) \in \text{Ob}(\text{Set})$.
fong-spivak-seven-sketches_149 F01595	149	0.988	$\{(s, t) \mid s \in S, t \in T\}$	$\{(s, t) \mid s \in S, t \in T\}$	$\text{Ob}(\text{Set})$. We know what it is: the set of pairs $\{(s, t) \mid s \in S, t \in T\}$.
fong-spivak-seven-sketches_149 F01596	149	1.000	$f : S \rightarrow S'$	$f : S \rightarrow S'$	• For any pair of morphisms, i.e. functions, $f : S \rightarrow S'$ and $g : T \rightarrow T'$, one obtains a
fong-spivak-seven-sketches_149 F01597	149	1.000	$g : T \rightarrow T'$	$g : T \rightarrow T'$	• For any pair of morphisms, i.e. functions, $f : S \rightarrow S'$ and $g : T \rightarrow T'$, one obtains a
fong-spivak-seven-sketches_149 F01598	149	1.000	$(f \times g) : (S \times T) \rightarrow (S' \times T')$	$(f \times g) : (S \times T) \rightarrow (S' \times T')$	function $(f \times g) : (S \times T) \rightarrow (S' \times T')$. It works pointwise: $(f \times g)(s, t) := (f(s), g(t))$.
fong-spivak-seven-sketches_149 F01599	149	1.000	$(f \times g)(s, t) := (f(s), g(t))$	$(f \times g)(s, t) := (f(s), g(t))$	function $(f \times g) : (S \times T) \rightarrow (S' \times T')$. It works pointwise: $(f \times g)(s, t) := (f(s), g(t))$.
fong-spivak-seven-sketches_149 F01600	149	0.441	$\text{id}_S \times \text{id}_T = \text{id}_{S \times T}$	$\text{id}_S \times \text{id}_T = \text{id}_{S \times T}$	• These should preserve identities: $\text{id}_S \times \text{id}_T = \text{id}_{S \times T}$ for any sets S, T .
fong-spivak-seven-sketches_149 F01601	149	1.000	$S \xrightarrow{f} S' \xrightarrow{f'} S''$	$S \xrightarrow{f} S' \xrightarrow{f'} S''$	• These should preserve composition: for any functions $S \xrightarrow{f} S' \xrightarrow{f'} S''$ and $T \xrightarrow{g} T' \xrightarrow{g'} T''$ and $h : S' \rightarrow T'$, one has
fong-spivak-seven-sketches_149 F01602	149	1.000	$T \xrightarrow{g} T' \xrightarrow{g'} T''$	$T \xrightarrow{g} T' \xrightarrow{g'} T''$	• These should preserve composition: for any functions $S \xrightarrow{f} S' \xrightarrow{f'} S''$ and $T \xrightarrow{g} T' \xrightarrow{g'} T''$ and $h : S' \rightarrow T'$, one has
fong-spivak-seven-sketches_149 F01603	149	1.000	$T' \xrightarrow{g'} T''$	$T' \xrightarrow{g'} T''$	$T' \xrightarrow{g'} T''$, one has
fong-spivak-seven-sketches_150 F01604	150	1.000	$\{1\} \times S \cong S$	$\{1\} \times S \cong S$	The four conditions, (a), (b), (c), and (d) give isomorphisms $\{1\} \times S \cong S$, etc. These
fong-spivak-seven-sketches_150 F01605	150	0.742	$\{(1, s) \mid s \in S\} \rightarrow S$	$\{(1, s) \mid s \in S\} \rightarrow S$	maps are obvious in the case of Set , e.g. the function $\{(1, s) \mid s \in S\} \rightarrow S$ sending $(1, s)$
fong-spivak-seven-sketches_150 F01606	150	0.742	$1, s$	$1, s$	maps are obvious in the case of Set , e.g. the function $\{(1, s) \mid s \in S\} \rightarrow S$ sending $(1, s)$
fong-spivak-seven-sketches_150 F01607	150	0.984	$1, \times$	$1, \times$	<i>Exercise 4.50.</i> Consider the monoidal category $(\text{Set}, 1, \times)$, together with the diagram
fong-spivak-seven-sketches_150 F01608	150	0.755	$A = B = C = D = F = G = \mathbb{Z}$	$A = B = C = D = F = G = \mathbb{Z}$	Suppose that $A = B = C = D = F = G = \mathbb{Z}$ and $E = \mathbb{B} = \{\text{true}, \text{false}\}$, and
fong-spivak-seven-sketches_150 F01609	150	0.755	$E = \mathbb{B} = \{\text{true}, \text{false}\}$	$E = \mathbb{B} = \{\text{true}, \text{false}\}$	Suppose that $A = B = C = D = F = G = \mathbb{Z}$ and $E = \mathbb{B} = \{\text{true}, \text{false}\}$, and
fong-spivak-seven-sketches_150 F01610	150	0.976	$f_C(a) = a , f_D(a) = a * 5, g_E(d, b) = d - b$	$f_C(a) = a , f_D(a) = a * 5, g_E(d, b) = d - b$	suppose that $f_C(a) = a , f_D(a) = a * 5, g_E(d, b) = d - b$, and

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_F01611	150	0.976	$d \leq b$	$d \leq b$	suppose that $f_C(a) = a $, $f_D(a) = a * 5$, $g_E(d, b) = “d \leq b,”$ $g_F(d, b) = d - b$, and
fong-spivak-seven-sketches_F01612	150	0.976	$g_{\{F\}}(d, b)=d-b$	$g_F(d, b) = d - b$	suppose that $f_C(a) = a $, $f_D(a) = a * 5$, $g_E(d, b) = “d \leq b,”$ $g_F(d, b) = d - b$, and
fong-spivak-seven-sketches_F01613	150	1.000	$h(c, e)=$	$h(c, e) =$	$h(c, e) \equiv$ if e then c else $1 - c$.
fong-spivak-seven-sketches_F01614	150	1.000	$1-c$	$1 - c$	$h(c, e) =$ if e then c else $1 - c$.
fong-spivak-seven-sketches_F01615	150	1.000	$g_{\{E\}}(5, 3)$	$g_E(5, 3)$	1. What are $g_E(5, 3)$ and $g_F(5, 3)$?
fong-spivak-seven-sketches_F01616	150	1.000	$g_{\{F\}}(5, 3)$	$g_F(5, 3)$	1. What are $g_E(5, 3)$ and $g_F(5, 3)$?
fong-spivak-seven-sketches_F01617	150	1.000	$g_{\{E\}}(3, 5)$	$g_E(3, 5)$	2. What are $g_E(3, 5)$ and $g_F(3, 5)$?
fong-spivak-seven-sketches_F01618	150	1.000	$g_{\{F\}}(3, 5)$	$g_F(3, 5)$	2. What are $g_E(3, 5)$ and $g_F(3, 5)$?
fong-spivak-seven-sketches_F01619	150	0.992	$h(5$	$h(5$	3. What is $h(5, \text{true})$?
fong-spivak-seven-sketches_F01620	150	0.758	$h(-5$	$h(-5$	4. What is $h(-5, \text{true})$?
fong-spivak-seven-sketches_F01621	150	1.000	$A \times B \rightarrowtail G \times F$	$A \times B \rightarrow G \times F$	The whole diagram now defines a function $A \times B \rightarrow G \times F$; call it q .
fong-spivak-seven-sketches_F01622	150	1.000	$q_{\{G\}}(-2, 3)$	$q_G(-2, 3)$	6. What are $q_G(-2, 3)$ and $q_F(-2, 3)$?
fong-spivak-seven-sketches_F01623	150	1.000	$q_{\{F\}}(-2, 3)$	$q_F(-2, 3)$	6. What are $q_G(-2, 3)$ and $q_F(-2, 3)$?
fong-spivak-seven-sketches_F01624	150	1.000	$q_{\{G\}}(2, 3)$	$q_G(2, 3)$	7. What are $q_G(2, 3)$ and $q_F(2, 3)$?
fong-spivak-seven-sketches_F01625	150	1.000	$q_{\{F\}}(2, 3)$	$q_F(2, 3)$	7. What are $q_G(2, 3)$ and $q_F(2, 3)$?
fong-spivak-seven-sketches_F01626	150	0.542	$X(x, y) \in \mathcal{V}$	$X(x, y) \in \mathcal{V}$	(ii) for every pair $x, y \in \text{Ob}(\mathcal{X})$, one specifies an object $\mathcal{X}(x, y) \in \mathcal{V}$, called the <i>hom-</i>
fong-spivak-seven-sketches_F01627	150	0.356	$x \in \mathrm{Ob}(X)$	$x \in \text{Ob}(X)$	(iii) for every $x \in \text{Ob}(\mathcal{X})$, one specifies a morphism $\text{id}_x: I \rightarrow \mathcal{X}(x, x)$ in $\mathcal{V}$, called the
fong-spivak-seven-sketches_F01628	150	0.356	$\mathrm{id}_x: I \rightarrowtail X(x, x)$	$\text{id}_x: I \rightarrow X(x, x)$	(iii) for every $x \in \text{Ob}(\mathcal{X})$, one specifies a morphism $\text{id}_x: I \rightarrow \mathcal{X}(x, x)$ in $\mathcal{V}$, called the
fong-spivak-seven-sketches_F01629	151	0.608	$x, y, z \in \mathrm{Ob}(X)$	$x, y, z \in \text{Ob}(X)$	(iv) for each $x, y, z \in \text{Ob}(\mathcal{X})$, one specifies a morphism $\mathbin{\mathcal{X}}: \mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \rightarrow \mathcal{X}(x, z)$,
fong-spivak-seven-sketches_F01630	151	0.608	$9: \mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \rightarrowtail \mathcal{X}(x, z)$	$9: \mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \rightarrow \mathfrak{X}(x, z)$	(iv) for each $x, y, z \in \text{Ob}(\mathcal{X})$, one specifies a morphism $\mathbin{\mathcal{X}}: \mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \rightarrow \mathcal{X}(x, z)$,
fong-spivak-seven-sketches_F01631	151	1.000	$\mathcal{V}=($	$\mathcal{V} = ($	<i>Exercise 4.52.</i> Recall from Example 4.49 that $\mathcal{V} = (\mathbf{Set}, \{1\}, \times)$ is a symmetric monoidal
fong-spivak-seven-sketches_F01632	151	1.000	$, \{1\}, \times)$	$, \{1\}, \times)$	<i>Exercise 4.52.</i> Recall from Example 4.49 that $\mathcal{V} = (\mathbf{Set}, \{1\}, \times)$ is a symmetric monoidal
fong-spivak-seven-sketches_F01633	151	0.961	$\mathrm{id}_x: I \rightarrowtail \mathcal{X}(x, x)$	$\text{id}_x: I \rightarrow \mathcal{X}(x, x)$	$\text{id}_x: I \rightarrow \mathcal{X}(x, x)$ for the object x simply becomes the requirement (a) that $I \leq \mathcal{X}(x, x)$ is
fong-spivak-seven-sketches_F01634	151	0.961	$I \leq \mathcal{X}(x, x)$	$I \leq \mathcal{X}(x, x)$	$\text{id}_x: I \rightarrow \mathcal{X}(x, x)$ for the object x simply becomes the requirement (a) that $I \leq \mathcal{X}(x, x)$ is
fong-spivak-seven-sketches_F01635	153	0.644	$(\mathcal{C}, I, \otimes)$	$(\mathcal{C}, I, \otimes)$	Definition 4.58. Let $(\mathcal{C}, I, \otimes)$ be a symmetric monoidal category, and $c \in \text{Ob}(\mathcal{C})$ an
fong-spivak-seven-sketches_F01636	153	0.999	$c^* \in \mathrm{Ob}(\mathcal{C})$	$c^* \in \text{Ob}(\mathcal{C})$	(i) an object $c^* \in \text{Ob}(\mathcal{C})$, called the <i>dual of c</i> ,
fong-spivak-seven-sketches_F01637	153	1.000	$\eta_c: I \rightarrowtail c^* \otimes c$	$\eta_c: I \rightarrow c^* \otimes c$	(ii) a morphism $\eta_c: I \rightarrow c^* \otimes c$, called the <i>unit for c</i> ,

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F01638	153	■ 1.000	$\backslash\epsilon_{\text{c}}: c \otimes c^* \rightarrow I$	$\epsilon_c : c \otimes c^* \rightarrow I$	(iii) a morphism $\epsilon_c : c \otimes c^* \rightarrow I$, called the <i>counit</i> for c .
fong-spivak-seven-sketches F01639	153	■ 0.516	$c^{\ast}$	c^*	If for every object $c \in \text{Ob}(\mathcal{C})$ there exists a dual c^* for c , then we say that $(\mathcal{C}, I, \otimes)$ is
fong-spivak-seven-sketches F01640	153	■ 0.997	$\backslash\text{xrightarrow}{c}$	$\xrightarrow{c}$	wire labeled c^* , i.e. $\xrightarrow{c}$ is the same as $\xleftarrow{c^*}$. The cup and cap discussed above are in fact
fong-spivak-seven-sketches F01641	153	■ 0.997	$\backslash\text{stackrel}{c^{\ast}}{\leftarrow}$	$\xleftarrow{c^*}$	wire labeled c^* , i.e. $\xrightarrow{c}$ is the same as $\xleftarrow{c^*}$. The cup and cap discussed above are in fact
fong-spivak-seven-sketches F01642	154	■ 0.877	$\backslash\epsilon_{\text{\text{sound}}}$	ϵ_{sound}	Note that the pictures in Eq. (4.57) correspond to ϵ_{sound} and η_{sound^*} .
fong-spivak-seven-sketches F01643	154	■ 0.877	$\backslash\eta_{\text{\text{sound}^*}}$	η_{sound^*}	Note that the pictures in Eq. (4.57) correspond to ϵ_{sound} and η_{sound^*} .
fong-spivak-seven-sketches F01644	154	■ 0.879	$\backslash\text{mathcal}{C}(b \otimes c, d) \cong \backslash\text{mathcal}{C}(b, c \multimap d)$	$\mathcal{C}(b \otimes c, d) \cong \mathcal{C}(b, c \multimap d)$	object $c \multimap d$ and an isomorphism $\mathcal{C}(b \otimes c, d) \cong \mathcal{C}(b, c \multimap d)$, natural in b . While we will
fong-spivak-seven-sketches F01645	154	■ 1.000	$c^{\prime}$	c'	2. if c^* and c' are both duals to c then there is an isomorphism $c^* \cong c'$; and
fong-spivak-seven-sketches F01646	154	■ 1.000	$c^{\ast} \cong c^{\prime}$	$c^* \cong c'$	2. if c^* and c' are both duals to c then there is an isomorphism $c^* \cong c'$; and
fong-spivak-seven-sketches F01647	154	■ 1.000	$c \cong c^{**}$	$c \cong c^{**}$	3. there is an isomorphism between c and its double-dual, $c \cong c^{**}$.
fong-spivak-seven-sketches F01648	154	■ 0.947	$c^{\ast} \otimes d$	$c^* \otimes d$	To prove 1., the key idea is that for any c and d , the object $c \multimap d$ is given by $c^* \otimes d$,
fong-spivak-seven-sketches F01649	154	■ 0.999	$\backslash\text{mathcal}{C}(b \otimes c, d) \cong \backslash\text{mathcal}{C}(b, c \rightarrow d)$	$\mathcal{C}(b \otimes c, d) \cong \mathcal{C}(b, c \rightarrow d)$	and the natural isomorphism $\mathcal{C}(b \otimes c, d) \cong \mathcal{C}(b, c \multimap d)$ is given by precomposing with
fong-spivak-seven-sketches F01650	154	■ 0.899	$\backslash\text{mathrm{id}}_b \otimes \eta_c$	$\text{id}_b \otimes \eta_c$	$\text{id}_b \otimes \eta_c$.
fong-spivak-seven-sketches F01651	154	■ 1.000	$\backslash\alpha: A \rightarrow B$	$\alpha : A \rightarrow B$	look at than to write down formally. ² If in addition to the corelation $\alpha: A \rightarrow B$ above
fong-spivak-seven-sketches F01652	154	■ 1.000	$\backslash\beta: B \rightarrow C$	$\beta : B \rightarrow C$	we have another corelation $\beta: B \rightarrow C$
fong-spivak-seven-sketches F01653	155	■ 0.999	$\backslash\beta \circ \alpha$	$\beta \circ \alpha$	Then the composite $\beta \circ \alpha$ of our two corelations is given by
fong-spivak-seven-sketches F01654	155	■ 1.000	$\backslash\beta$	β	one to the other staying within equivalence classes of either α or β .
fong-spivak-seven-sketches F01655	155	■ 0.873	$(\varnothing, \sqcup)$	$(\emptyset, \sqcup)$	$(\emptyset, \sqcup)$. This monoidal category is compact closed, with every finite set its own
fong-spivak-seven-sketches F01656	155	■ 1.000	$A \sqcup A := \{(a, 1), (a, 2) \mid a \in A\}$	$A \sqcup A := \{(a, 1), (a, 2) \mid a \in A\}$	$A \sqcup A := \{(a, 1), (a, 2) \mid a \in A\}$ where each part simply consists of the two elements $(a, 1)$
fong-spivak-seven-sketches F01657	155	■ 1.000	$(a, 1)$	$(a, 1)$	$A \sqcup A := \{(a, 1), (a, 2) \mid a \in A\}$ where each part simply consists of the two elements $(a, 1)$
fong-spivak-seven-sketches F01658	155	■ 1.000	$(a, 2)$	$(a, 2)$	and $(a, 2)$ for each $a \in A$. The unit on a finite set A is the corelation $\eta_A: \emptyset \rightarrow A \sqcup A$
fong-spivak-seven-sketches F01659	155	■ 1.000	$\backslash\eta_{\text{\text{A}}}: \varnothing \rightarrow A \sqcup A$	$\eta_A : \emptyset \rightarrow A \sqcup A$	and $(a, 2)$ for each $a \in A$. The unit on a finite set A is the corelation $\eta_A: \emptyset \rightarrow A \sqcup A$
fong-spivak-seven-sketches F01660	155	■ 1.000	$\backslash\epsilon_{\text{\text{A}}}: A \sqcup A \rightarrow \varnothing$	$\epsilon_A : A \sqcup A \rightarrow \emptyset$	$\epsilon_A: A \sqcup A \rightarrow \emptyset$ specied by this same equivalence relation.
fong-spivak-seven-sketches F01661	155	■ 0.997	$\backslash\varnothing \rightarrow \underline{3} \sqcup \underline{3}$	$\emptyset \rightarrow \underline{3} \sqcup \underline{3}$	1. Draw a picture of the unit corelation $\emptyset \rightarrow \underline{3} \sqcup \underline{3}$.
fong-spivak-seven-sketches F01662	155	■ 1.000	$\backslash\underline{3} \sqcup \underline{3} \rightarrow \varnothing$	$\underline{3} \sqcup \underline{3} \rightarrow \emptyset$	2. Draw a picture of the counit corelation $\underline{3} \sqcup \underline{3} \rightarrow \emptyset$.
fong-spivak-seven-sketches F01663	155	■ 0.993	$(\backslash\text{mathcal}{V}, I, \otimes)$	$(\mathcal{V}, I, \otimes)$	extra work to prove that for any skeletal (unital, commutative) quantale $(\mathcal{V}, I, \otimes)$ the

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_155_F01664	155	0.562	$\operatorname{Prof}_{\mathcal{V}}$	$\operatorname{Prof}_{\mathcal{V}}$	profunctor category $\operatorname{Prof}_{\mathcal{V}}$ of Theorem 4.23 is compact closed, so we'll discuss this
fong-spivak-seven-sketches_155_F01665	155	0.863	$\alpha ; \beta$	$\alpha ; \beta$	$\alpha \circ \beta$ on $A \sqcup C$. To do so requires three steps: (i) consider α and β as relations on $A \sqcup B \sqcup C$, (ii) take the
fong-spivak-seven-sketches_155_F01666	155	0.863	$A \sqcup C$	$A \sqcup C$	$\alpha \circ \beta$ on $A \sqcup C$. To do so requires three steps: (i) consider α and β as relations on $A \sqcup B \sqcup C$, (ii) take the
fong-spivak-seven-sketches_155_F01667	155	0.863	$A \sqcup B \sqcup C$	$A \sqcup B \sqcup C$	$\alpha \circ \beta$ on $A \sqcup C$. To do so requires three steps: (i) consider α and β as relations on $A \sqcup B \sqcup C$, (ii) take the
fong-spivak-seven-sketches_155_F01668	155	1.000	$\alpha \subseteq (A \sqcup B) \times (A \sqcup B)$	$\alpha \subseteq (A \sqcup B) \times (A \sqcup B)$	formal description. Note that as binary relations, we have $\alpha \subseteq (A \sqcup B) \times (A \sqcup B)$, and $\beta \subseteq (B \sqcup C) \times (B \sqcup C)$.
fong-spivak-seven-sketches_155_F01669	155	1.000	$\beta \subseteq (B \sqcup C) \times (B \sqcup C)$	$\beta \subseteq (B \sqcup C) \times (B \sqcup C)$	formal description. Note that as binary relations, we have $\alpha \subseteq (A \sqcup B) \times (A \sqcup B)$, and $\beta \subseteq (B \sqcup C) \times (B \sqcup C)$.
fong-spivak-seven-sketches_155_F01670	155	1.000	$\iota_{A \sqcup B} : A \sqcup B \rightarrow A \sqcup B \sqcup C, \iota_{B \sqcup C} : B \sqcup C \rightarrow A \sqcup B \sqcup C$	$\iota_{A \sqcup B} : A \sqcup B \rightarrow A \sqcup B \sqcup C, \iota_{B \sqcup C} : B \sqcup C \rightarrow A \sqcup B \sqcup C$	We also have three inclusions: $\iota_{A \sqcup B} : A \sqcup B \rightarrow A \sqcup B \sqcup C, \iota_{B \sqcup C} : B \sqcup C \rightarrow A \sqcup B \sqcup C$, and $\iota_{A \sqcup C} : A \sqcup C \rightarrow$
fong-spivak-seven-sketches_155_F01671	155	1.000	$\iota_{A \sqcup C} : A \sqcup C \rightarrow$	$\iota_{A \sqcup C} : A \sqcup C \rightarrow$	We also have three inclusions: $\iota_{A \sqcup B} : A \sqcup B \rightarrow A \sqcup B \sqcup C, \iota_{B \sqcup C} : B \sqcup C \rightarrow A \sqcup B \sqcup C$, and $\iota_{A \sqcup C} : A \sqcup C \rightarrow$
fong-spivak-seven-sketches_156_F01672	156	1.000	$\mathcal{V}$	$\mathcal{V}$	Monoidal products in $\operatorname{Prof}_{\mathcal{V}}$ are just product categories. In terms of wiring diagrams,
fong-spivak-seven-sketches_156_F01673	156	0.383	$\Phi : X_1 \mapsto X_2$	$\Phi : X_1 \mapsto X_2$	$\operatorname{Prof}_{\mathcal{V}}$ are $\mathcal{V}$ -profunctors. So given $\mathcal{V}$ -profunctors $\Phi : X_1 \mapsto X_2$ and $\Psi : Y_1 \mapsto Y_2$, one
fong-spivak-seven-sketches_156_F01674	156	0.383	$\Psi : y_1 \rightarrow y_2$	$\Psi : y_1 \rightarrow y_2$	$\operatorname{Prof}_{\mathcal{V}}$ are $\mathcal{V}$ -profunctors. So given $\mathcal{V}$ -profunctors $\Phi : X_1 \mapsto X_2$ and $\Psi : Y_1 \mapsto Y_2$, one
fong-spivak-seven-sketches_156_F01675	156	0.262	$(\Phi \times \Psi) : \mathcal{X}_1 \times y_1 \rightarrow x_2 \times y_2$	$(\Phi \times \Psi) : \mathcal{X}_1 \times y_1 \rightarrow x_2 \times y_2$	defines a $\mathcal{V}$ -profunctor $(\Phi \times \Psi) : \mathcal{X}_1 \times y_1 \rightarrow x_2 \times y_2$ by
fong-spivak-seven-sketches_156_F01676	156	0.857	$\mathbf{Prof}_{\mathbf{Bool}}$	$\mathbf{Prof}_{\mathbf{Bool}}$	<i>Exercise 4.64.</i> Interpret the monoidal products in $\mathbf{Prof}_{\mathbf{Bool}}$ in terms of feasibility.
fong-spivak-seven-sketches_156_F01677	156	1.000	$\Phi \times \Psi$	$\Phi \times \Psi$	sense as the monoidal product of resource preorders $\mathcal{X}$ and $\mathcal{Y}$ and why $\Phi \times \Psi$ makes
fong-spivak-seven-sketches_156_F01678	156	0.593	$(1, 1) = I$	$(1, 1) = I$	$\mathcal{V}$ -category $\mathbf{1}$; it has one object, say 1 , and $(1, 1) = I$ is the monoidal unit of $\mathcal{V}$. We take
fong-spivak-seven-sketches_156_F01679	156	0.467	$X \times \mathbf{1} \rightarrow X$	$X \times \mathbf{1} \rightarrow X$	phisms $X \times \mathbf{1} \mapsto X$ and $\mathbf{1} \times X \mapsto X$ in $\mathbf{Prof}_{\mathcal{V}}$, for any $\mathcal{V}$ -category X . What are they?
fong-spivak-seven-sketches_156_F01680	156	0.467	$\mathbf{1} \times X \rightarrow X$	$\mathbf{1} \times X \rightarrow X$	phisms $X \times \mathbf{1} \mapsto X$ and $\mathbf{1} \times X \mapsto X$ in $\mathbf{Prof}_{\mathcal{V}}$, for any $\mathcal{V}$ -category X . What are they?
fong-spivak-seven-sketches_157_F01681	157	0.894	$\mathcal{X}^{\operatorname{op}}$	$\mathcal{X}^{\operatorname{op}}$	Duals are easy: for every $\mathcal{V}$ -category X , its dual is its opposite category X^{op} (see
fong-spivak-seven-sketches_157_F01682	157	0.225	$\eta_x : \mathbf{1} \rightarrow \mathcal{X}^{\operatorname{op}} \times \mathcal{X}$	$\eta_x : \mathbf{1} \rightarrow \mathcal{X}^{\operatorname{op}} \times \mathcal{X}$	$\mathcal{V}$ -profunctor $\eta_X : \mathbf{1} \mapsto X^{\operatorname{op}} \times X$. By definition, this is a $\mathcal{V}$ -functor
fong-spivak-seven-sketches_157_F01683	157	0.981	$\eta_x(1, x, x') := \mathcal{X}(x, x')$	$\eta_x(1, x, x') := \mathcal{X}(x, x')$	we define it by $\eta_X(1, x, x') := X(x, x')$. Similarly, the counit is the profunctor $\epsilon_X : (X \times$
fong-spivak-seven-sketches_157_F01684	157	0.981	$\epsilon_x : (\mathcal{X} \times$	$\epsilon_x : (\mathcal{X} \times$	we define it by $\eta_X(1, x, x') := X(x, x')$. Similarly, the counit is the profunctor $\epsilon_X : (X \times$
fong-spivak-seven-sketches_157_F01685	157	0.710	$\mathcal{X}^{\operatorname{op}}) \rightarrow \mathbf{1}$	$\mathcal{X}^{\operatorname{op}}) \rightarrow \mathbf{1}$	$X^{\operatorname{op}}) \mapsto \mathbf{1}$, defined by $\epsilon_X(x, x', 1) := X(x, x')$.
fong-spivak-seven-sketches_157_F01686	157	0.710	$\epsilon_X(x, x', 1) := \mathcal{X}(x, x')$	$\epsilon_X(x, x', 1) := \mathcal{X}(x, x')$	$X^{\operatorname{op}}) \mapsto \mathbf{1}$, defined by $\epsilon_X(x, x', 1) := X(x, x')$.
fong-spivak-seven-sketches_160_F01687	160	1.000	$u = 15x$	$u = 15x$	produces two output signals: $u = 15x$ (upper right) and $v = 3x + 21y$ (lower right).
fong-spivak-seven-sketches_160_F01688	160	1.000	$v = 3x + 21y$	$v = 3x + 21y$	produces two output signals: $u = 15x$ (upper right) and $v = 3x + 21y$ (lower right).
fong-spivak-seven-sketches_160_F01689	160	1.000	$x + 7y$	$x + 7y$	adds the second copy of x to the lower signal to get $x + 7y$, and so on.
fong-spivak-seven-sketches_161_F01690	161	0.608	$(\mathcal{C}, 0, +)$	$(\mathcal{C}, 0, +)$	Definition 5.2. A <i>prop</i> is a symmetric strict monoidal category $(\mathcal{C}, 0, +)$ for which




Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_161 F01691	161	■ 0.950	$\operatorname{Ob}\{\mathcal{C}\}=\mathbb{N}$	$\operatorname{Ob}(\mathcal{C}) = \mathbb{N}$	$\operatorname{Ob}(\mathcal{C}) = \mathbb{N}$, the monoidal unit is $0 \in \mathbb{N}$, and the monoidal product on objects is
fong-spivak-seven-sketches_161 F01692	161	■ 0.950	$0 \in \mathbb{N}$	$0 \in \mathbb{N}$	$\operatorname{Ob}(\mathcal{C}) = \mathbb{N}$, the monoidal unit is $0 \in \mathbb{N}$, and the monoidal product on objects is
fong-spivak-seven-sketches_161 F01693	161	■ 0.994	$\mathcal{C}(m, n)$	$\mathcal{C}(m, n)$	(i) a set $\mathcal{C}(m, n)$ of morphisms $m \rightarrow n$, for $m, n \in \mathbb{N}$.
fong-spivak-seven-sketches_161 F01694	161	■ 0.655	$\operatorname{id}_n: n \rightarrow n$	$\operatorname{id}_n: n \rightarrow n$	(ii) for all $n \in \mathbb{N}$, an identity map $\operatorname{id}_n: n \rightarrow n$.
fong-spivak-seven-sketches_161 F01695	161	■ 0.999	$\sigma_{m, n}: m+n \rightarrow n+m$	$\sigma_{m, n}: m + n \rightarrow n + m$	(iii) for all $m, n \in \mathbb{N}$, a symmetry map $\sigma_{m, n}: m + n \rightarrow n + m$.
fong-spivak-seven-sketches_161 F01696	161	■ 1.000	$f: m \rightarrow n$	$f: m \rightarrow n$	(iv) a composition rule: given $f: m \rightarrow n$ and $g: n \rightarrow p$, a map $(f \circ g): m \rightarrow p$.
fong-spivak-seven-sketches_161 F01697	161	■ 1.000	$g: n \rightarrow p$	$g: n \rightarrow p$	(iv) a composition rule: given $f: m \rightarrow n$ and $g: n \rightarrow p$, a map $(f \circ g): m \rightarrow p$.
fong-spivak-seven-sketches_161 F01698	161	■ 1.000	$(f; g): m \rightarrow p$	$(f; g): m \rightarrow p$	(iv) a composition rule: given $f: m \rightarrow n$ and $g: n \rightarrow p$, a map $(f \circ g): m \rightarrow p$.
fong-spivak-seven-sketches_161 F01699	161	■ 1.000	$f: m \rightarrow m^{\prime}$	$f: m \rightarrow m'$	(v) a monoidal product on morphisms: given $f: m \rightarrow m'$ and $g: n \rightarrow n'$, a map
fong-spivak-seven-sketches_161 F01700	161	■ 1.000	$g: n \rightarrow n^{\prime}$	$g: n \rightarrow n'$	(v) a monoidal product on morphisms: given $f: m \rightarrow m'$ and $g: n \rightarrow n'$, a map
fong-spivak-seven-sketches_161 F01701	161	■ 0.963	$(f+g): m+n \rightarrow m^{\prime}+n^{\prime}$	$(f + g): m + n \rightarrow m' + n'$	$(f + g): m + n \rightarrow m' + n'$.
fong-spivak-seven-sketches_162 F01702	162	■ 1.000	$\underline{m}=\{1, \ldots m\}$	$\underline{m} = \{1, \dots m\}$	from $\underline{m} = \{1, \dots m\}$ to $\underline{n} = \{1, \dots, n\}$. (The identities, symmetries, and composition
fong-spivak-seven-sketches_162 F01703	162	■ 1.000	$\underline{n}=\{1, \ldots, n\}$	$\underline{n} = \{1, \dots, n\}$	from $\underline{m} = \{1, \dots m\}$ to $\underline{n} = \{1, \dots, n\}$. (The identities, symmetries, and composition
fong-spivak-seven-sketches_162 F01704	162	■ 1.000	$f+g: m+n \rightarrow m^{\prime}+n^{\prime}$	$f + g: m + n \longrightarrow m' + n'$	functions: that is, given $f: m \rightarrow m'$ and $g: n \rightarrow n'$, we define $(f + g): m + n \longrightarrow m' + n'$
fong-spivak-seven-sketches_162 F01705	162	■ 1.000	$f: 3 \rightarrow 2$	$f: 3 \rightarrow 2$	1. Draw a morphism $f: 3 \rightarrow 2$ and a morphism $g: 2 \rightarrow 4$ in FinSet .
fong-spivak-seven-sketches_162 F01706	162	■ 1.000	$g: 2 \rightarrow 4$	$g: 2 \rightarrow 4$	1. Draw a morphism $f: 3 \rightarrow 2$ and a morphism $g: 2 \rightarrow 4$ in FinSet .
fong-spivak-seven-sketches_162 F01707	162	■ 1.000	$f+g$	$f + g$	2. Draw $f + g$.
fong-spivak-seven-sketches_162 F01708	162	■ 0.998	$\sigma_{m, n}$	$\sigma_{m, n}$	5. Choose $m, n \in \mathbb{N}$, and draw the symmetry map $\sigma_{m, n}$ in FinSet ?
fong-spivak-seven-sketches_162 F01709	162	■ 1.000	$\underline{m} \rightarrow \underline{n}$	$\underline{m} \rightarrow \underline{n}$	bijections $\underline{m} \rightarrow \underline{n}$. Note that in this case morphisms $m \rightarrow n$ only exist when $m = n$;
fong-spivak-seven-sketches_162 F01710	162	■ 0.987	$m \neq n$	$m \neq n$	when $m \neq n$ the homset $\mathbf{Bij}(m, n)$ is empty. Since Bij is a subcategory of FinSet , we
fong-spivak-seven-sketches_162 F01711	162	■ 0.987	$\mathbf{Bij}(m, n)$	$\mathbf{Bij}(m, n)$	when $m \neq n$ the homset $\mathbf{Bij}(m, n)$ is empty. Since Bij is a subcategory of FinSet , we
fong-spivak-seven-sketches_162 F01712	162	■ 0.987	$\mathbf{Bij}$	Bij	when $m \neq n$ the homset $\mathbf{Bij}(m, n)$ is empty. Since Bij is a subcategory of FinSet , we
fong-spivak-seven-sketches_162 F01713	162	■ 1.000	$\underline{m} \sqcup \underline{n}$	$\underline{m} \sqcup \underline{n}$	are partitions on $\underline{m} \sqcup \underline{n}$ (see Example 4.61), is a prop.
fong-spivak-seven-sketches_162 F01714	162	■ 0.952	$R \subseteq \underline{m} \times \underline{n}$	$R \subseteq \underline{m} \times \underline{n}$	Example 5.8. There is a prop Rel for which morphisms $m \rightarrow n$ are relations, $R \subseteq \underline{m} \times \underline{n}$.
fong-spivak-seven-sketches_162 F01715	162	■ 1.000	$S \subseteq \underline{n} \times \underline{p}$	$S \subseteq \underline{n} \times \underline{p}$	The composition of R with $S \subseteq \underline{n} \times \underline{p}$ is
fong-spivak-seven-sketches_163 F01716	163	■ 0.934	$F(n)=n$	$F(n) = n$	(a) F is identity-on-objects, i.e. $F(n) = n$ for all $n \in \operatorname{Ob}(\mathcal{C}) = \operatorname{Ob}(\mathcal{D}) = \mathbb{N}$, and

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F01717	163	0.934	$n \in \operatorname{Ob}(\mathcal{C}) = \operatorname{Ob}(\mathcal{D}) = \mathbb{N}$	$n \in \operatorname{Ob}(\mathcal{C}) = \operatorname{Ob}(\mathcal{D}) = \mathbb{N}$	(a) F is identity-on-objects, i.e. $F(n) = n$ for all $n \in \operatorname{Ob}(\mathcal{C}) = \operatorname{Ob}(\mathcal{D}) = \mathbb{N}$, and
fong-spivak-seven-sketches-F01718	163	0.977	$f: m_1 \rightarrow m_2$	$f: m_1 \rightarrow m_2$	(b) for all $f: m_1 \rightarrow m_2$ and $g: n_1 \rightarrow n_2$ in $\mathcal{C}$, we have $F(f) + F(g) = F(f + g)$ in $\mathcal{D}$.
fong-spivak-seven-sketches-F01719	163	0.977	$g: n_1 \rightarrow n_2$	$g: n_1 \rightarrow n_2$	(b) for all $f: m_1 \rightarrow m_2$ and $g: n_1 \rightarrow n_2$ in $\mathcal{C}$, we have $F(f) + F(g) = F(f + g)$ in $\mathcal{D}$.
fong-spivak-seven-sketches-F01720	163	0.977	$F(f)+F(g)=F(f+g)$	$F(f) + F(g) = F(f + g)$	(b) for all $f: m_1 \rightarrow m_2$ and $g: n_1 \rightarrow n_2$ in $\mathcal{C}$, we have $F(f) + F(g) = F(f + g)$ in $\mathcal{D}$.
fong-spivak-seven-sketches-F01721	163	0.966	i	i	<i>Example 5.12.</i> The inclusion $i: \mathbf{Bij} \rightarrow \mathbf{FinSet}$ is a prop functor. Perhaps more interest-
fong-spivak-seven-sketches-F01722	163	0.503	$F:$	$F:$	ingly, there is a prop functor $F: \mathbf{FinSet} \rightarrow \mathbf{Rel}_{\mathbf{Fin}}$. It sends a function $f: \underline{m} \rightarrow \underline{n}$ to the
fong-spivak-seven-sketches-F01723	163	0.503	$\mathbf{Rel}_{\mathbf{Fin}}$	$\mathbf{Rel}_{\mathbf{Fin}}$	ingly, there is a prop functor $F: \mathbf{FinSet} \rightarrow \mathbf{Rel}_{\mathbf{Fin}}$. It sends a function $f: \underline{m} \rightarrow \underline{n}$ to the
fong-spivak-seven-sketches-F01724	163	0.503	$f: \underline{m} \rightarrow \underline{n}$	$f: \underline{m} \rightarrow \underline{n}$	ingly, there is a prop functor $F: \mathbf{FinSet} \rightarrow \mathbf{Rel}_{\mathbf{Fin}}$. It sends a function $f: \underline{m} \rightarrow \underline{n}$ to the
fong-spivak-seven-sketches-F01725	163	1.000	$F(f) := \{(i, j) \mid f(i) = j\} \subseteq \underline{m} \times \underline{n}$	$F(f) := \{(i, j) \mid f(i) = j\} \subseteq \underline{m} \times \underline{n}$	relation $F(f) := \{(i, j) \mid f(i) = j\} \subseteq \underline{m} \times \underline{n}$.
fong-spivak-seven-sketches-F01726	163	0.616	ι	ι	Definition 5.13. For $m, n \in \mathbb{N}$, an (m, n) -port graph (V, in, out, ι) is specified by
fong-spivak-seven-sketches-F01727	163	0.835	$V \rightarrow \mathbb{N}$	$V \rightarrow \mathbb{N}$	(ii) functions $in, out: V \rightarrow \mathbb{N}$, where $in(v)$ and $out(v)$ are called the <i>in degree</i> and <i>out</i>
fong-spivak-seven-sketches-F01728	163	0.835	$\operatorname{in}(v)$	$in(v)$	(ii) functions $in, out: V \rightarrow \mathbb{N}$, where $in(v)$ and $out(v)$ are called the <i>in degree</i> and <i>out</i>
fong-spivak-seven-sketches-F01729	163	0.835	$\operatorname{out}(v)$	$out(v)$	(ii) functions $in, out: V \rightarrow \mathbb{N}$, where $in(v)$ and $out(v)$ are called the <i>in degree</i> and <i>out</i>
fong-spivak-seven-sketches-F01730	163	0.793	$\iota: \underline{m} \sqcup O \xrightarrow{\cong} I \sqcup \underline{n}$	$\iota: \underline{m} \sqcup O \xrightarrow{\cong} I \sqcup \underline{n}$	(iii) a bijection $\iota: \underline{m} \sqcup O \xrightarrow{\cong} I \sqcup \underline{n}$, where $I = \{(v, i) \mid v \in V, 1 \leq i \leq in(v)\}$ is the set of
fong-spivak-seven-sketches-F01731	163	0.793	$I = \{(v, i) \mid v \in V, 1 \leq i \leq \operatorname{in}(v)\}$	$I = \{(v, i) \mid v \in V, 1 \leq i \leq in(v)\}$	(iii) a bijection $\iota: \underline{m} \sqcup O \xrightarrow{\cong} I \sqcup \underline{n}$, where $I = \{(v, i) \mid v \in V, 1 \leq i \leq in(v)\}$ is the set of
fong-spivak-seven-sketches-F01732	163	1.000	$O = \{(v, i) \mid v \in V, 1 \leq i \leq \operatorname{out}(v)\}$	$O = \{(v, i) \mid v \in V, 1 \leq i \leq out(v)\}$	<i>vertex inputs</i> , and $O = \{(v, i) \mid v \in V, 1 \leq i \leq out(v)\}$ is the set of <i>vertex outputs</i> .
fong-spivak-seven-sketches-F01733	163	0.933	$e_{v,j}^{u,i}: u \rightarrow v$	$e_{v,j}^{u,i}: u \rightarrow v$	construct the graph with vertices V and with an arrow $e_{v,j}^{u,i}: u \rightarrow v$ for every $i, j \in \mathbb{N}$
fong-spivak-seven-sketches-F01734	163	0.933	$i, j \in \mathbb{N}$	$i, j \in \mathbb{N}$	construct the graph with vertices V and with an arrow $e_{v,j}^{u,i}: u \rightarrow v$ for every $i, j \in \mathbb{N}$
fong-spivak-seven-sketches-F01735	163	1.000	$\iota(u, i) = (v, j)$	$\iota(u, i) = (v, j)$	such that $\iota(u, i) = (v, j)$; call it the <i>internal flow graph</i> . If the internal flow graph is
fong-spivak-seven-sketches-F01736	163	1.000	$R_1 + R_2$	$R_1 + R_2$	³ The monoidal product $R_1 + R_2$ of relations $R_1 \subseteq \underline{m}_1 \times \underline{n}_1$ and $R_2 \subseteq \underline{m}_2 \times \underline{n}_2$ is given by $R_1 \sqcup R_2 \subseteq$
fong-spivak-seven-sketches-F01737	163	1.000	$R_1 \subseteq \underline{m}_1 \times \underline{n}_1$	$R_1 \subseteq \underline{m}_1 \times \underline{n}_1$	³ The monoidal product $R_1 + R_2$ of relations $R_1 \subseteq \underline{m}_1 \times \underline{n}_1$ and $R_2 \subseteq \underline{m}_2 \times \underline{n}_2$ is given by $R_1 \sqcup R_2 \subseteq$
fong-spivak-seven-sketches-F01738	163	1.000	$R_2 \subseteq \underline{m}_2 \times \underline{n}_2$	$R_2 \subseteq \underline{m}_2 \times \underline{n}_2$	³ The monoidal product $R_1 + R_2$ of relations $R_1 \subseteq \underline{m}_1 \times \underline{n}_1$ and $R_2 \subseteq \underline{m}_2 \times \underline{n}_2$ is given by $R_1 \sqcup R_2 \subseteq$
fong-spivak-seven-sketches-F01739	163	1.000	$R_1 \sqcup R_2 \subseteq$	$R_1 \sqcup R_2 \subseteq$	³ The monoidal product $R_1 + R_2$ of relations $R_1 \subseteq \underline{m}_1 \times \underline{n}_1$ and $R_2 \subseteq \underline{m}_2 \times \underline{n}_2$ is given by $R_1 \sqcup R_2 \subseteq$
fong-spivak-seven-sketches-F01740	163	0.420	$\left(\underline{m}_1 \times \underline{n}_1 \right) \sqcup \left(\underline{m}_2 \times \underline{n}_2 \right) \subseteq \left(\underline{m}_1 \sqcup \underline{m}_2 \right) \times \left(\underline{n}_1 \sqcup \underline{n}_2 \right)$	$\left(\underline{m}_1 \times \underline{n}_1 \right) \sqcup \left(\underline{m}_2 \times \underline{n}_2 \right) \subseteq \left(\underline{m}_1 \sqcup \underline{m}_2 \right) \times \left(\underline{n}_1 \sqcup \underline{n}_2 \right)$	$\left(\underline{m}_1 \times \underline{n}_1 \right) \sqcup \left(\underline{m}_2 \times \underline{n}_2 \right) \subseteq \left(\underline{m}_1 \sqcup \underline{m}_2 \right) \times \left(\underline{n}_1 \sqcup \underline{n}_2 \right)$.
fong-spivak-seven-sketches-F01741	164	0.999	$m=2$	$m = 2$	<i>Example 5.14.</i> Here is an example of a $(2, 3)$ -port graph, i.e. with $m = 2$ and $n = 3$:

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_F01742	164	■ 0.999	n=3	$n = 3$	<i>Example 5.14.</i> Here is an example of a $(2, 3)$ -port graph, i.e. with $m = 2$ and $n = 3$:
fong-spivak-seven-sketches_F01743	164	■ 1.000	$V=\{a, b, c\}$	$V = \{a, b, c\}$	outer box, and three on the right. The vertex set is $V = \{a, b, c\}$ and, for example
fong-spivak-seven-sketches_F01744	164	■ 0.994	$\operatorname{in}(a)=1$	$\operatorname{in}(a) = 1$	$\operatorname{in}(a) = 1$ and $\operatorname{out}(a) = 3$, so we draw one port on the left-hand side and three ports on
fong-spivak-seven-sketches_F01745	164	■ 0.994	$\operatorname{out}(a)=3$	$\operatorname{out}(a) = 3$	$\operatorname{in}(a) = 1$ and $\operatorname{out}(a) = 3$, so we draw one port on the left-hand side and three ports on
fong-spivak-seven-sketches_F01746	164	■ 0.996	(m, n)	(m, n)	A category $\mathbf{PG}$ whose morphisms are port graphs. Given an (m, n) -port graph
fong-spivak-seven-sketches_F01747	164	■ 0.179	$(V$	$(V$	$(V, \operatorname{in}, \operatorname{out}, \iota)$ and an (n, p) -port graph $(V', \operatorname{in}', \operatorname{out}', \iota')$, we may compose them to produce
fong-spivak-seven-sketches_F01748	164	■ 0.179	$\iota)$	$\iota)$	$(V, \operatorname{in}, \operatorname{out}, \iota)$ and an (n, p) -port graph $(V', \operatorname{in}', \operatorname{out}', \iota')$, we may compose them to produce
fong-spivak-seven-sketches_F01749	164	■ 0.179	(n, p)	(n, p)	$(V, \operatorname{in}, \operatorname{out}, \iota)$ and an (n, p) -port graph $(V', \operatorname{in}', \operatorname{out}', \iota')$, we may compose them to produce
fong-spivak-seven-sketches_F01750	164	■ 0.179	$V^{\{\prime\}}$	V'	$(V, \operatorname{in}, \operatorname{out}, \iota)$ and an (n, p) -port graph $(V', \operatorname{in}', \operatorname{out}', \iota')$, we may compose them to produce
fong-spivak-seven-sketches_F01751	164	■ 0.179	$\iota^{\{\prime\}}$	ι'	$(V, \operatorname{in}, \operatorname{out}, \iota)$ and an (n, p) -port graph $(V', \operatorname{in}', \operatorname{out}', \iota')$, we may compose them to produce
fong-spivak-seven-sketches_F01752	164	■ 0.147	(m, p)	(m, p)	an (m, p) -port graph $(V \sqcup V', [\operatorname{in}, \operatorname{in}'], [\operatorname{out}, \operatorname{out}'], \iota'')$. Here $[\operatorname{in}, \operatorname{in}']$ denotes the function
fong-spivak-seven-sketches_F01753	164	■ 0.147	$\left(V \sqcup V^{\{\prime\}}, [\right.$	$(V \sqcup V', [$	an (m, p) -port graph $(V \sqcup V', [\operatorname{in}, \operatorname{in}'], [\operatorname{out}, \operatorname{out}'], \iota'')$. Here $[\operatorname{in}, \operatorname{in}']$ denotes the function
fong-spivak-seven-sketches_F01754	164	■ 0.147	$], [$	$], [$	an (m, p) -port graph $(V \sqcup V', [\operatorname{in}, \operatorname{in}'], [\operatorname{out}, \operatorname{out}'], \iota'')$. Here $[\operatorname{in}, \operatorname{in}']$ denotes the function
fong-spivak-seven-sketches_F01755	164	■ 0.147	$\left.], \iota^{\{\prime\prime\}}\right)$	$], \iota'')$	an (m, p) -port graph $(V \sqcup V', [\operatorname{in}, \operatorname{in}'], [\operatorname{out}, \operatorname{out}'], \iota'')$. Here $[\operatorname{in}, \operatorname{in}']$ denotes the function
fong-spivak-seven-sketches_F01756	164	■ 0.777	$V \sqcup V^{\{\prime\}} \rightarrow \mathbb{N}$	$V \sqcup V' \rightarrow \mathbb{N}$	$V \sqcup V' \rightarrow \mathbb{N}$ which maps elements of V according to in , and elements of V' according
fong-spivak-seven-sketches_F01757	164	■ 0.191	$\operatorname{in}^{\{\prime\}}$	in'	to in' , and similarly for $[\operatorname{out}, \operatorname{out}']$. The bijection $\iota'': \underline{m} \sqcup O \sqcup O' \rightarrow I \sqcup I' \sqcup \underline{p}$ is defined
fong-spivak-seven-sketches_F01758	164	■ 0.191	$\iota^{\{\prime\prime\}}: \underline{m} \sqcup O \sqcup O' \rightarrow I \sqcup I' \sqcup \underline{p}$	$\iota'': \underline{m} \sqcup O \sqcup O' \rightarrow I \sqcup I' \sqcup \underline{p}$	to in' , and similarly for $[\operatorname{out}, \operatorname{out}']$. The bijection $\iota'': \underline{m} \sqcup O \sqcup O' \rightarrow I \sqcup I' \sqcup \underline{p}$ is defined
fong-spivak-seven-sketches_F01759	165	■ 0.911	$\operatorname{Ob}(\mathbf{PG})=\mathbb{N}$	$\operatorname{Ob}(\mathbf{PG}) = \mathbb{N}$	We thus have a category $\mathbf{PG}$, whose objects are natural numbers $\operatorname{Ob}(\mathbf{PG}) = \mathbb{N}$,
fong-spivak-seven-sketches_F01760	165	■ 0.965	$\mathbf{PG}(m, n)=\{(V, \operatorname{in}, \operatorname{out}, \iota) \mid$	$\mathbf{PG}(m, n) = \{(V, \operatorname{in}, \operatorname{out}, \iota) \mid$	whose morphisms are port graphs $\mathbf{PG}(m, n) = \{(V, \operatorname{in}, \operatorname{out}, \iota) \mid$ as in Definition 5.13}.
fong-spivak-seven-sketches_F01761	165	■ 0.996	(n, n)	(n, n)	Composition of port graphs is as above, and the identity port graph on n is the (n, n) -
fong-spivak-seven-sketches_F01762	165	■ 0.928	$\left(\varnothing, !, !, \operatorname{id}_{\underline{n}}\right)$	$(\varnothing, !, !, \operatorname{id}_{\underline{n}})$	port graph $(\varnothing, !, !, \operatorname{id}_{\underline{n}})$, where $!: \varnothing \rightarrow \mathbb{N}$ is the unique function. The identity on an
fong-spivak-seven-sketches_F01763	165	■ 0.928	$\varnothing \rightarrow \mathbb{N}$	$\varnothing \rightarrow \mathbb{N}$	port graph $(\varnothing, !, !, \operatorname{id}_{\underline{n}})$, where $!: \varnothing \rightarrow \mathbb{N}$ is the unique function. The identity on an
fong-spivak-seven-sketches_F01764	165	■ 0.423	$G:=(V$	$G := (V$	monoidal product of two port graphs $G := (V, \operatorname{in}, \operatorname{out}, \iota)$ and $G' := (V', \operatorname{in}', \operatorname{out}', \iota')$ is
fong-spivak-seven-sketches_F01765	165	■ 0.423	$G^{\{\prime\}}:=\left(V^{\{\prime\}}, \operatorname{in}^{\{\prime\}}, \operatorname{out}^{\{\prime\}}, \iota^{\{\prime\}}\right)$	$G' := (V', \operatorname{in}', \operatorname{out}', \iota')$	monoidal product of two port graphs $G := (V, \operatorname{in}, \operatorname{out}, \iota)$ and $G' := (V', \operatorname{in}', \operatorname{out}', \iota')$ is
fong-spivak-seven-sketches_F01766	165	■ 0.863	$(\varnothing, !, !, !)$	$(\varnothing, !, !, !)$	The monoidal unit is $(\varnothing, !, !, !)$.
fong-spivak-seven-sketches_F01767	165	■ 1.000	$R \subseteq P \times P$	$R \subseteq P \times P$	of taking the reflexive, transitive closure of a relation. That is, given a relation $R \subseteq P \times P$,
fong-spivak-seven-sketches_F01768	165	■ 0.798	(V, A, s, t)	(V, A, s, t)	we consider a graph (V, A, s, t) as defining a relation $\{(s(a), t(a)) \mid a \in A\} \subseteq V \times V$, and

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_F01769	165	0.798	$\{(s(a), t(a)) \mid a \in A\} \subseteq V \times V$	$\{(s(a), t(a)) \mid a \in A\} \subseteq V \times V$	we consider a graph (V, A, s, t) as defining a relation $\{(s(a), t(a)) \mid a \in A\} \subseteq V \times V$ and
fong-spivak-seven-sketches_F01770	166	0.821	$P, \leq_P$	$P, \leq_P$	Exercise 5.20. Let P be a set, let $R \subseteq P \times P$ a relation, let $(P, \leq_P)$ be the preorder
fong-spivak-seven-sketches_F01771	166	1.000	$R(x, y)$	$R(x, y)$	1. Suppose that for every $x, y \in P$, if $R(x, y)$ then $f(x) \leq f(y)$. Show that f defines
fong-spivak-seven-sketches_F01772	166	1.000	$f(x) \leq_Q f(y)$	$f(x) \leq_Q f(y)$	every $x, y \in P$, if $R(x, y)$ then $f(x) \leq_Q f(y)$.
fong-spivak-seven-sketches_F01773	166	0.989	P, Q, R	P, Q, R	Exercise 5.21. Let P, Q, R , etc. be as in Exercise 5.20. We want to see that the universal
fong-spivak-seven-sketches_F01774	166	1.000	$a, b \in Q$	$a, b \in Q$	1. Suppose that for every $a, b \in Q$, if $a \leq b$ then $(g(a), g(b)) \in R$. Is it automatically
fong-spivak-seven-sketches_F01775	166	1.000	$(g(a), g(b)) \in R$	$(g(a), g(b)) \in R$	1. Suppose that for every $a, b \in Q$, if $a \leq b$ then $(g(a), g(b)) \in R$. Is it automatically
fong-spivak-seven-sketches_F01776	166	1.000	$g: (Q, \leq_Q) \rightarrow (P, \leq_P)$	$g: (Q, \leq_Q) \rightarrow (P, \leq_P)$	true that g defines a monotone map $g: (Q, \leq_Q) \rightarrow (P, \leq_P)$?
fong-spivak-seven-sketches_F01777	167	0.998	$\operatorname{Mor}(\mathcal{C})$	$\operatorname{Mor}(\mathcal{C})$	$\mathcal{C}$ be another category whose set of morphisms is denoted $\operatorname{Mor}(\mathcal{C})$.
fong-spivak-seven-sketches_F01778	167	0.996	$\operatorname{Mor}(\mathcal{C}) \rightarrow$	$\operatorname{Mor}(\mathcal{C}) \rightarrow$	1. Someone tells you that there are “domain and codomain” functions $\operatorname{dom}, \operatorname{cod}: \operatorname{Mor}(\mathcal{C}) \rightarrow$
fong-spivak-seven-sketches_F01779	167	0.615	$\operatorname{Ob}(\mathcal{C})$	$\operatorname{Ob}(\mathcal{C})$	$\operatorname{Ob}(\mathcal{C})$; interpret this statement.
fong-spivak-seven-sketches_F01780	167	1.000	$\mathcal{G} \rightarrow \mathcal{C}$	$\mathcal{G} \rightarrow \mathcal{C}$	2. Show that the set of functors $\mathcal{G} \rightarrow \mathcal{C}$ are in one-to-one correspondence with the
fong-spivak-seven-sketches_F01781	167	0.999	$f: V \rightarrow \operatorname{Ob}(\mathcal{C})$	$f: V \rightarrow \operatorname{Ob}(\mathcal{C})$	set of pairs of functions (f, g) , where $f: V \rightarrow \operatorname{Ob}(\mathcal{C})$ and $g: A \rightarrow \operatorname{Mor}(\mathcal{C})$ for
fong-spivak-seven-sketches_F01782	167	0.999	$g: A \rightarrow \operatorname{Mor}(\mathcal{C})$	$g: A \rightarrow \operatorname{Mor}(\mathcal{C})$	set of pairs of functions (f, g) , where $f: V \rightarrow \operatorname{Ob}(\mathcal{C})$ and $g: A \rightarrow \operatorname{Mor}(\mathcal{C})$ for
fong-spivak-seven-sketches_F01783	167	1.000	$\operatorname{dom}(g(a)) = f(s(a))$	$\operatorname{dom}(g(a)) = f(s(a))$	which $\operatorname{dom}(g(a)) = f(s(a))$ and $\operatorname{cod}(g(a)) = f(t(a))$ for all a .
fong-spivak-seven-sketches_F01784	167	1.000	$\operatorname{cod}(g(a)) = f(t(a))$	$\operatorname{cod}(g(a)) = f(t(a))$	which $\operatorname{dom}(g(a)) = f(s(a))$ and $\operatorname{cod}(g(a)) = f(t(a))$ for all a .
fong-spivak-seven-sketches_F01785	167	1.000	$\operatorname{Loop}(A)$	$\operatorname{Loop}(A)$	one-object categories. For any set A , there is a graph $\operatorname{Loop}(A)$ with one vertex and with
fong-spivak-seven-sketches_F01786	167	1.000	$A = \{a\}$	$A = \{a\}$	1. What are the elements of the free monoid on the set $A = \{a\}$?
fong-spivak-seven-sketches_F01787	167	1.000	$\{a\}$	$\{a\}$	2. Can you find a well-known monoid that is isomorphic to the free monoid on $\{a\}$?
fong-spivak-seven-sketches_F01788	167	0.999	G, s, t	G, s, t	Definition 5.25. A prop signature is a tuple (G, s, t) , where G is a set and $s, t: G \rightarrow \mathbb{N}$
fong-spivak-seven-sketches_F01789	167	0.999	$s, t: G \rightarrow \mathbb{N}$	$s, t: G \rightarrow \mathbb{N}$	Definition 5.25. A prop signature is a tuple (G, s, t) , where G is a set and $s, t: G \rightarrow \mathbb{N}$
fong-spivak-seven-sketches_F01790	167	1.000	$g \in G$	$g \in G$	are functions; each element $g \in G$ is called a generator and $s(g), t(g) \in \mathbb{N}$ are called its
fong-spivak-seven-sketches_F01791	167	1.000	$s(g), t(g) \in \mathbb{N}$	$s(g), t(g) \in \mathbb{N}$	are functions; each element $g \in G$ is called a generator and $s(g), t(g) \in \mathbb{N}$ are called its
fong-spivak-seven-sketches_F01792	167	0.860	s, t	s, t	in-arity and out-arity. We often denote (G, s, t) simply by G , taking s, t to be implicit.
fong-spivak-seven-sketches_F01793	167	0.577	$\Gamma = (V$	$\Gamma = (V$	A G -labeling of a port graph $\Gamma = (V, in, out, \iota)$ is a function $\ell: V \rightarrow G$ such that the
fong-spivak-seven-sketches_F01794	167	0.577	$\ell: V \rightarrow G$	$\ell: V \rightarrow G$	A G -labeling of a port graph $\Gamma = (V, in, out, \iota)$ is a function $\ell: V \rightarrow G$ such that the
fong-spivak-seven-sketches_F01795	167	0.999	$s(\ell(v)) = \operatorname{in}(v)$	$s(\ell(v)) = \operatorname{in}(v)$	arities agree: $s(\ell(v)) = \operatorname{in}(v)$ and $t(\ell(v)) = \operatorname{out}(v)$ for each $v \in V$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_167 F01796	167	■ 0.999	$t(\ell(v))=\operatorname{out}\{v\}$	$t(\ell(v)) = \operatorname{out}(v)$	arities agree: $s(\ell(v)) = \operatorname{in}(v)$ and $t(\ell(v)) = \operatorname{out}(v)$ for each $v \in V$.
fong-spivak-seven-sketches_167 F01797	167	■ 1.000	$(m, n) \in \mathbb{N} \times \mathbb{N}$	$(m, n) \in \mathbb{N} \times \mathbb{N}$	⁴ The arity of a prop morphism is a pair $(m, n) \in \mathbb{N} \times \mathbb{N}$, where m is the number of inputs and n is the
fong-spivak-seven-sketches_168 F01798	168	■ 0.852	$\ell^{\prime}$	ℓ'	port graphs $\mathbf{PG}$ (see Eq. (5.17)); the labelings (the ℓ' 's) are just carried along.
fong-spivak-seven-sketches_168 F01799	168	■ 1.000	ℓ	ℓ	the right. In wiring diagrams, we depict the labeling function $\ell: V \rightarrow G$ by using ℓ to
fong-spivak-seven-sketches_168 F01800	168	■ 1.000	$G=\{f: 1 \rightarrow 1, g: 2 \rightarrow 2, h: 2 \rightarrow 1\}$	$G = \{f: 1 \rightarrow 1, g: 2 \rightarrow 2, h: 2 \rightarrow 1\}$	with the same generator. For example, if $G = \{f: 1 \rightarrow 1, g: 2 \rightarrow 2, h: 2 \rightarrow 1\}$, then the
fong-spivak-seven-sketches_168 F01801	168	■ 0.987	$3 \rightarrow 2$	$3 \rightarrow 2$	following is a morphism $3 \rightarrow 2$ in $\mathbf{Free}(G)$:
fong-spivak-seven-sketches_168 F01802	168	■ 1.000	$V \rightarrow \varnothing$	$V \rightarrow \emptyset$	must have a labelling function of the form $V \rightarrow \emptyset$, and hence we must have $V = \emptyset$;
fong-spivak-seven-sketches_168 F01803	168	■ 1.000	$V=\varnothing$	$V = \emptyset$	must have a labelling function of the form $V \rightarrow \emptyset$, and hence we must have $V = \emptyset$;
fong-spivak-seven-sketches_168 F01804	168	■ 1.000	(n, m)	(n, m)	see Exercise 1.25. Thus the only morphisms (n, m) are those given by port graphs
fong-spivak-seven-sketches_168 F01805	168	■ 0.997	$(\varnothing, !, !, \sigma)$	$(\emptyset, !, !, \sigma)$	$(\emptyset, !, !, \sigma)$, where $\sigma: n \rightarrow m$ is a bijection.
fong-spivak-seven-sketches_168 F01806	168	■ 0.997	$\sigma: n \rightarrow m$	$\sigma: n \rightarrow m$	$(\emptyset, !, !, \sigma)$, where $n \rightarrow m$ is a bijection.
fong-spivak-seven-sketches_168 F01807	168	■ 0.985	$(m, n) \in \mathbb{N}^2$	$(m, n) \in \mathbb{N}^2$	i.e. having one generating morphism for each $(m, n) \in \mathbb{N}^2$. Show that $\mathbf{Free}(G)$ is the
fong-spivak-seven-sketches_168 F01808	168	■ 0.886	$\operatorname{prop} \mathcal{C}$	$\operatorname{prop} \mathcal{C}$	for any prop $\mathcal{C}$, the prop functors $\mathbf{Free}(G) \rightarrow \mathcal{C}$ are in one-to-one correspondence with
fong-spivak-seven-sketches_168 F01809	168	■ 0.886	$\operatorname{Free}(G) \rightarrow \mathcal{C}$	$\mathbf{Free}(G) \rightarrow \mathcal{C}$	for any prop $\mathcal{C}$, the prop functors $\mathbf{Free}(G) \rightarrow \mathcal{C}$ are in one-to-one correspondence with
fong-spivak-seven-sketches_168 F01810	168	■ 1.000	$G \rightarrow \mathcal{C}$	$G \rightarrow \mathcal{C}$	functions $G \rightarrow \mathcal{C}$ that send each $g \in G$ to a morphism $s(g) \rightarrow t(g)$ in $\mathcal{C}$.
fong-spivak-seven-sketches_168 F01811	168	■ 1.000	$s(g) \rightarrow t(g)$	$s(g) \rightarrow t(g)$	functions $G \rightarrow \mathcal{C}$ that send each $g \in G$ to a morphism $s(g) \rightarrow t(g)$ in $\mathcal{C}$.
fong-spivak-seven-sketches_169 F01812	169	■ 0.995	$e: m \rightarrow n$	$e: m \rightarrow n$	G -generated prop expression, or simply expression $e: m \rightarrow n$, where $m, n \in \mathbb{N}$, inductively
fong-spivak-seven-sketches_169 F01813	169	■ 0.976	$\operatorname{id}_0: 0 \rightarrow 0$	$\operatorname{id}_0: 0 \rightarrow 0$	• The empty morphism $\operatorname{id}_0: 0 \rightarrow 0$, the identity morphism $\operatorname{id}_1: 1 \rightarrow 1$, and the
fong-spivak-seven-sketches_169 F01814	169	■ 0.976	$\operatorname{id}_1: 1 \rightarrow 1$	$\operatorname{id}_1: 1 \rightarrow 1$	• The empty morphism $\operatorname{id}_0: 0 \rightarrow 0$, the identity morphism $\operatorname{id}_1: 1 \rightarrow 1$, and the
fong-spivak-seven-sketches_169 F01815	169	■ 1.000	$\sigma: 2 \rightarrow 2$	$\sigma: 2 \rightarrow 2$	symmetry $\sigma: 2 \rightarrow 2$ are expressions. ⁵
fong-spivak-seven-sketches_169 F01816	169	■ 1.000	$g: s(g) \rightarrow t(g)$	$g: s(g) \rightarrow t(g)$	• the generators $g \in G$ are expressions $g: s(g) \rightarrow t(g)$.
fong-spivak-seven-sketches_169 F01817	169	■ 1.000	$\alpha: m \rightarrow n$	$\alpha: m \rightarrow n$	• if $\alpha: m \rightarrow n$ and $\beta: p \rightarrow q$ are expressions, then $\alpha + \beta: m + p \rightarrow n + q$ is an
fong-spivak-seven-sketches_169 F01818	169	■ 1.000	$\beta: p \rightarrow q$	$\beta: p \rightarrow q$	• if $\alpha: m \rightarrow n$ and $\beta: p \rightarrow q$ are expressions, then $\alpha + \beta: m + p \rightarrow n + q$ is an
fong-spivak-seven-sketches_169 F01819	169	■ 1.000	$\alpha + \beta: m + p \rightarrow n + q$	$\alpha + \beta: m + p \rightarrow n + q$	• if $\alpha: m \rightarrow n$ and $\beta: p \rightarrow q$ are expressions, then $\alpha + \beta: m + p \rightarrow n + q$ is an
fong-spivak-seven-sketches_169 F01820	169	■ 1.000	$\beta: n \rightarrow p$	$\beta: n \rightarrow p$	• if $\alpha: m \rightarrow n$ and $\beta: n \rightarrow p$ are expressions, then $\alpha \circ \beta: m \rightarrow p$ is an expression.
fong-spivak-seven-sketches_169 F01821	169	■ 1.000	$\alpha; \beta: m \rightarrow p$	$\alpha; \beta: m \rightarrow p$	• if $\alpha: m \rightarrow n$ and $\beta: n \rightarrow p$ are expressions, then $\alpha \circ \beta: m \rightarrow p$ is an expression.
fong-spivak-seven-sketches_169 F01822	169	■ 1.000	$\operatorname{Expr}(G)$	$\operatorname{Expr}(G)$	We write $\operatorname{Expr}(G)$ for the set of expressions in G . If $e: m \rightarrow n$ is an expression, we refer

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_F01823	169	1.000	$f: 1 \rightarrow 1$	$f: 1 \rightarrow 1$	• 
fong-spivak-seven-sketches_F01824	169	0.963	$f: \mathrm{id}_1: 1 \rightarrow 1$	$f: \mathrm{id}_1: 1 \rightarrow 1$	• 
fong-spivak-seven-sketches_F01825	169	0.997	$h+\mathrm{id}_1: 3 \rightarrow 2$	$h + \mathrm{id}_1: 3 \rightarrow 2$	• 
fong-spivak-seven-sketches_F01826	169	0.551	$\left(h+\mathrm{id}_1\right) ; \sigma ; g ; \sigma: 3 \rightarrow 2$	$(h + \mathrm{id}_1); \sigma; g; \sigma: 3 \rightarrow 2$	• 
fong-spivak-seven-sketches_F01827	169	0.981	$f ; \mathrm{id}_1$	$f; \mathrm{id}_1$	phism. For example, we want to consider $f \circ \mathrm{id}_1$ and f to be the same morphism,
fong-spivak-seven-sketches_F01828	169	0.708	$f ; \mathrm{id}_1=f$	$f; \mathrm{id}_1 = f$	since the unitality axiom for categories says $f \circ \mathrm{id}_1 = f$. Nonetheless, we only consider
fong-spivak-seven-sketches_F01829	169	0.923	σ	σ	⁵ One can think of σ as the “swap” icon $\bowtie: 2 \rightarrow 2$
fong-spivak-seven-sketches_F01830	169	0.923	$\mathrm{X}: 2 \rightarrow 2$	$X: 2 \rightarrow 2$	⁵ One can think of σ as the “swap” icon $\bowtie: 2 \rightarrow 2$
fong-spivak-seven-sketches_F01831	170	0.975	$\left(g+\mathrm{id}_1\right) \stackrel{\circ}{\circ} \left(\mathrm{id}_1+\sigma\right) \stackrel{\circ}{\circ} \left(\mathrm{id}_1+g\right) \stackrel{\circ}{\circ} \left(h+\mathrm{id}_1\right) \stackrel{\circ}{\circ} \left(\sigma+\mathrm{id}_1\right) \stackrel{\circ}{\circ} \left(\mathrm{id}_1+h\right) \stackrel{\circ}{\circ} \sigma \stackrel{\circ}{\circ} g$	$(g + \mathrm{id}_1) \stackrel{\circ}{\circ} (\mathrm{id}_1 + \sigma) \stackrel{\circ}{\circ} (\mathrm{id}_1 + g) \stackrel{\circ}{\circ} (h + \mathrm{id}_1)$	Here the result is 
fong-spivak-seven-sketches_F01832	170	1.000	$G=\{f: 1 \rightarrow 1, g: 2 \rightarrow$	$G = \{f: 1 \rightarrow 1, g: 2 \rightarrow$	<i>Exercise 5.32.</i> Consider again the free prop on generators $G = \{f: 1 \rightarrow 1, g: 2 \rightarrow$
fong-spivak-seven-sketches_F01833	170	0.285	$2, h: 2 \rightarrow 1\}$	$2, h: 2 \rightarrow 1\}$	$2, h: 2 \rightarrow 1\}$. Draw a picture of $(f + \mathrm{id}_1 + \mathrm{id}_1) \circ (\sigma + \mathrm{id}_1) \circ (\mathrm{id}_1 + h) \circ \sigma \circ g$, where $\sigma: 2 \rightarrow 2$
fong-spivak-seven-sketches_F01834	170	0.285	$\left(f+\mathrm{id}_1+\mathrm{id}_1\right) \stackrel{\circ}{\circ} \left(\sigma+\mathrm{id}_1\right) \stackrel{\circ}{\circ} \left(\mathrm{id}_1+h\right) \stackrel{\circ}{\circ} \sigma \stackrel{\circ}{\circ} g$	$(f + \mathrm{id}_1 + \mathrm{id}_1) \stackrel{\circ}{\circ} (\sigma + \mathrm{id}_1) \stackrel{\circ}{\circ} (\mathrm{id}_1 + h) \stackrel{\circ}{\circ} \sigma \stackrel{\circ}{\circ} g$	$2, h: 2 \rightarrow 1\}$. Draw a picture of 
fong-spivak-seven-sketches_F01835	170	0.817	G, s, t, E	G, s, t, E	Rough Definition 5.33. A <i>presentation</i> (G, s, t, E) for a prop is a set G , functions
fong-spivak-seven-sketches_F01836	170	1.000	$E \subseteq \operatorname{Expr}(G) \times \operatorname{Expr}(G)$	$E \subseteq \operatorname{Expr}(G) \times \operatorname{Expr}(G)$	$s, t: G \rightarrow \mathbb{N}$, and a set $E \subseteq \operatorname{Expr}(G) \times \operatorname{Expr}(G)$ of pairs of G -generated prop expressions,
fong-spivak-seven-sketches_F01837	170	1.000	e_1	e_1	such that e_1 and e_2 have the same arity for each $(e_1, e_2) \in E$. We refer to G as the set of
fong-spivak-seven-sketches_F01838	170	1.000	e_2	e_2	such that e_1 and e_2 have the same arity for each $(e_1, e_2) \in E$. We refer to G as the set of
fong-spivak-seven-sketches_F01839	170	1.000	$\left(e_1, e_2\right) \in E$	$(e_1, e_2) \in E$	such that e_1 and e_2 have the same arity for each $(e_1, e_2) \in E$. We refer to G as the set of
fong-spivak-seven-sketches_F01840	170	0.936	(G, s, t, E)	(G, s, t, E)	The prop $\mathfrak{G}$ <i>presented</i> by the presentation (G, s, t, E) is the prop whose morphisms
fong-spivak-seven-sketches_F01841	170	1.000	$e_1=e_2$	$e_1 = e_2$	are elements in $\operatorname{Expr}(G)$, quotiented by both the equations $e_1 = e_2$ where $(e_1, e_2) \in E$,
fong-spivak-seven-sketches_F01842	171	1.000	$g \in G, f(g)$	$g \in G, f(g)$	• for all $g \in G$, $f(g)$ is a morphism $s(g) \rightarrow t(g)$, and
fong-spivak-seven-sketches_F01843	171	1.000	$f\left(e_1\right)=f\left(e_2\right)$	$f\left(e_1\right)=f\left(e_2\right)$	• for all $(e_1, e_2) \in E$, we have that $f\left(e_1\right)=f\left(e_2\right)$ in $\mathcal{C}$, where $f(e)$ denotes the
fong-spivak-seven-sketches_F01844	171	1.000	$f(e)$	$f(e)$	• for all $(e_1, e_2) \in E$, we have that $f\left(e_1\right)=f\left(e_2\right)$ in $\mathcal{C}$, where $f(e)$ denotes the
fong-spivak-seven-sketches_F01845	171	0.686	(G, s, t)	(G, s, t)	<i>Exercise 5.35.</i> Is it the case that the free prop on generators (G, s, t) , defined in
fong-spivak-seven-sketches_F01846	171	0.793	$G, s, t, \varnothing$	$G, s, t, \varnothing$	Definition 5.25, is the same thing as the prop presented by $(G, s, t, \varnothing)$, having no
fong-spivak-seven-sketches_F01847	171	0.997	$R, 0,+, 1, *$	$R, 0, +, 1, *$	Definition 5.36. A <i>rig</i> is a tuple $(R, 0, +, 1, *)$, where R is a set, $0, 1 \in R$ are elements,

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_171 F01848	171	■ 0.997	$0, 1 \in R$	$0, 1 \in R$	Definition 5.36. A rig is a tuple $(R, 0, +, 1, *)$, where R is a set, $0, 1 \in R$ are elements,
fong-spivak-seven-sketches_171 F01849	171	■ 0.775	$R \times R \rightarrow R$	$R \times R \rightarrow R$	and $+, *: R \times R \rightarrow R$ are functions, such that
fong-spivak-seven-sketches_171 F01850	171	■ 0.924	$R, +, 0$	$R, +, 0$	(a) $(R, +, 0)$ is a commutative monoid,
fong-spivak-seven-sketches_171 F01851	171	■ 0.911	$R, *, 1$	$R, *, 1$	(b) $(R, *, 1)$ is a monoid, ⁸ and
fong-spivak-seven-sketches_171 F01852	171	■ 1.000	$a * (b+c) = a * b + a * c$	$a * (b + c) = a * b + a * c$	(c) $a * (b + c) = a * b + a * c$ and $(a + b) * c = a * c + b * c$ for all $a, b, c \in R$.
fong-spivak-seven-sketches_171 F01853	171	■ 1.000	$(a+b) * c = a * c + b * c$	$(a + b) * c = a * c + b * c$	(c) $a * (b + c) = a * b + a * c$ and $(a + b) * c = a * c + b * c$ for all $a, b, c \in R$.
fong-spivak-seven-sketches_171 F01854	171	■ 1.000	$a, b, c \in R$	$a, b, c \in R$	(c) $a * (b + c) = a * b + a * c$ and $(a + b) * c = a * c + b * c$ for all $a, b, c \in R$.
fong-spivak-seven-sketches_171 F01855	171	■ 1.000	$a * 0 = 0 = 0 * a$	$a * 0 = 0 = 0 * a$	(d) $a * 0 = 0 = 0 * a$ for all $a \in R$.
fong-spivak-seven-sketches_171 F01856	171	■ 1.000	$a \in R$	$a \in R$	(d) $a * 0 = 0 = 0 * a$ for all $a \in R$.
fong-spivak-seven-sketches_171 F01857	171	■ 0.800	$(\mathbb{N}, 0, +, 1, *)$	$(\mathbb{N}, 0, +, 1, *)$	Example 5.37. The natural numbers form a rig $(\mathbb{N}, 0, +, 1, *)$.
fong-spivak-seven-sketches_171 F01858	171	■ 0.952	$(R, *, 1)$	$(R, *, 1)$	⁸ Note that we did not demand that $(R, *, 1)$ be commutative; we will see a naturally-arising example
fong-spivak-seven-sketches_172 F01859	172	■ 0.737	$V, 0, \vee, I, \otimes$	$V, 0, \vee, I, \otimes$	Example 5.39. Any quantale $\mathcal{V} = (V, \leq, I, \otimes)$ determines a rig $(V, 0, \vee, I, \otimes)$, where $0 =$
fong-spivak-seven-sketches_172 F01860	172	■ 0.737	$0 =$	$0 =$	Example 5.39. Any quantale $\mathcal{V} = (V, \leq, I, \otimes)$ determines a rig $(V, 0, \vee, I, \otimes)$, where $0 =$
fong-spivak-seven-sketches_172 F01861	172	■ 0.971	$\bigvee \varnothing$	$\bigvee \varnothing$	$\bigvee \varnothing$ is the empty join. See Definition 2.79.
fong-spivak-seven-sketches_172 F01862	172	■ 1.000	$\operatorname{Mat}_n(R)$	$\operatorname{Mat}_n(R)$	Example 5.40. If R is a rig and $n \in \mathbb{N}$ is any natural number, then the set $\operatorname{Mat}_n(R)$ of
fong-spivak-seven-sketches_172 F01863	172	■ 1.000	$(n \times n)$	$(n \times n)$	$(n \times n)$ -matrices in R forms a rig. A matrix $M \in \operatorname{Mat}_n(R)$ is a function $M: \underline{n} \times \underline{n} \rightarrow$
fong-spivak-seven-sketches_172 F01864	172	■ 1.000	$M \in \operatorname{Mat}_n(R)$	$M \in \operatorname{Mat}_n(R)$	$(n \times n)$ -matrices in R forms a rig. A matrix $M \in \operatorname{Mat}_n(R)$ is a function $M: \underline{n} \times \underline{n} \rightarrow$
fong-spivak-seven-sketches_172 F01865	172	■ 1.000	$M: \underline{n} \times \underline{n} \rightarrow$	$M: \underline{n} \times \underline{n} \rightarrow$	$(n \times n)$ -matrices in R forms a rig. A matrix $M \in \operatorname{Mat}_n(R)$ is a function $M: \underline{n} \times \underline{n} \rightarrow$
fong-spivak-seven-sketches_172 F01866	172	■ 1.000	$M+N$	$M + N$	R . Addition $M + N$ of matrices is given by $(M + N)(i, j) := M(i, j) + N(i, j)$ and
fong-spivak-seven-sketches_172 F01867	172	■ 1.000	$(M+N)(i, j) := M(i, j) + N(i, j)$	$(M + N)(i, j) := M(i, j) + N(i, j)$	R . Addition $M + N$ of matrices is given by $(M + N)(i, j) := M(i, j) + N(i, j)$ and
fong-spivak-seven-sketches_172 F01868	172	■ 0.977	$(M * N)(i, j) := \sum_{k \in \underline{n}} M(i, k) * N(k, j)$	$(M * N)(i, j) := \sum_{k \in \underline{n}} M(i, k) * N(k, j)$	multiplication $M * N$ is given by $(M * N)(i, j) := \sum_{k \in \underline{n}} M(i, k) * N(k, j)$. The 0-matrix is
fong-spivak-seven-sketches_172 F01869	172	■ 1.000	$0(i, j) := 0$	$0(i, j) := 0$	$0(i, j) := 0$ for all $i, j \in \underline{n}$. Note that $\operatorname{Mat}_n(R)$ is generally not commutative.
fong-spivak-seven-sketches_172 F01870	172	■ 1.000	$i, j \in \underline{n}$	$i, j \in \underline{n}$	$0(i, j) := 0$ for all $i, j \in \underline{n}$. Note that $\operatorname{Mat}_n(R)$ is generally not commutative.
fong-spivak-seven-sketches_172 F01871	172	■ 1.000	$1 \in \operatorname{Mat}_n(R)$	$1 \in \operatorname{Mat}_n(R)$	its multiplicative identity $1 \in \operatorname{Mat}_n(R)$?
fong-spivak-seven-sketches_172 F01872	172	■ 1.000	$\operatorname{Mat}_n(\mathbb{N})$	$\operatorname{Mat}_n(\mathbb{N})$	that $\operatorname{Mat}_n(\mathbb{N})$ is not commutative, where $\mathbb{N}$ is as in Example 5.37.
fong-spivak-seven-sketches_172 F01873	172	■ 0.993	$(\mathbb{R}, 0, +, 1, *)$	$\mathbb{R}, 0, +, 1, *$	Example 5.42. Any ring forms a rig. In particular, the real numbers $(\mathbb{R}, 0, +, 1, *)$ are a
fong-spivak-seven-sketches_172 F01874	172	■ 1.000	$M=R$	$M = R$	R -module M . We will not discuss this here, keeping to the simpler requirement that $M = R$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_173 F01875	173	■ 1.000	$R=\mathbb{N}$	$R = \mathbb{N}$	For example, consider the rig $R = \mathbb{N}$ from Example 5.37, where the scalars are the
fong-spivak-seven-sketches_173 F01876	173	■ 1.000	$a=15 \ x$	$a = 15x$	$a = 15x$ and $b = 3x + 21y$.
fong-spivak-seven-sketches_173 F01877	173	■ 1.000	$b=3 \ x+21 \ y$	$b = 3x + 21y$	$a = 15x$ and $b = 3x + 21y$.
fong-spivak-seven-sketches_174 F01878	174	■ 0.999	$15 \ x$	$15x$	finally amplified by 3. Thus, if x is the first input signal, then this contributes $15x$ to the
fong-spivak-seven-sketches_174 F01879	174	■ 1.000	$3 \ x=x+2 \ x$	$3x = x + 2x$	there are two paths to it from each input. The top input thus contributes $3x = x + 2x$,
fong-spivak-seven-sketches_174 F01880	174	■ 1.000	$21 \ y$	$21y$	$21y$.
fong-spivak-seven-sketches_174 F01881	174	■ 1.000	$(x, \ y, \ z)$	(x, y, z)	the movement of a particle. For example, consider a particle whose position (x, y, z) in
fong-spivak-seven-sketches_175 F01882	175	■ 1.000	$s, \ t: G_{\{R\}} \rightarrow \mathbb{N}$	$s, t : G_R \rightarrow \mathbb{N}$	and let $s, t: G_R \rightarrow \mathbb{N}$ be given by the number of dangling wires on the left and right of
fong-spivak-seven-sketches_175 F01883	175	■ 0.686	$\mathbf{Free}(G_{\{R\}})$	$\mathbf{Free}(G_R)$	prop $\mathbf{Free}(G_R)$ on this set G_R of generators. We define $\mathbf{SFG}_R := \mathbf{Free}(G_R)$.
fong-spivak-seven-sketches_175 F01884	175	■ 0.686	$G_{\{R\}}$	G_R	prop $\mathbf{Free}(G_R)$ on this set G_R of generators. We define $\mathbf{SFG}_R := \mathbf{Free}(G_R)$.
fong-spivak-seven-sketches_175 F01885	175	■ 0.686	$\mathbf{SFG}_{\{R\}} := \mathbf{Free}(G_{\{R\}})$	$\mathbf{SFG}_R := \mathbf{Free}(G_R)$	prop $\mathbf{Free}(G_R)$ on this set G_R of generators. We define $\mathbf{SFG}_R := \mathbf{Free}(G_R)$.
fong-spivak-seven-sketches_175 F01886	175	■ 0.896	$\mathbf{Free}(G_{\{R\}})$	$\mathbf{Free}(G_R)$	port graphs, morphisms in $\mathbf{Free}(G_R)$ should be drawn slightly differently. For example,
fong-spivak-seven-sketches_176 F01887	176	■ 0.978	$(m \times n)$	$(m \times n)$	An $(m \times n)$ -matrix M with values in R is a function $M: (\underline{m} \times \underline{n}) \rightarrow R$. Given an
fong-spivak-seven-sketches_176 F01888	176	■ 0.978	$M: (\underline{m} \times \underline{n}) \rightarrow R$	$M : (\underline{m} \times \underline{n}) \rightarrow R$	An $(m \times n)$ -matrix M with values in R is a function $M: (\underline{m} \times \underline{n}) \rightarrow R$. Given an
fong-spivak-seven-sketches_176 F01889	176	■ 0.815	$(n \times p)$	$(n \times p)$	$(m \times n)$ -matrix M and an $(n \times p)$ -matrix N , their <i>composite</i> is the $(m \times p)$ -matrix $M \circ N$
fong-spivak-seven-sketches_176 F01890	176	■ 0.815	$(m \times p)$	$(m \times p)$	$(m \times n)$ -matrix M and an $(n \times p)$ -matrix N , their <i>composite</i> is the $(m \times p)$ -matrix $M \circ N$
fong-spivak-seven-sketches_176 F01891	176	■ 0.815	$M \circ N$	$M \circ N$	$(m \times n)$ -matrix M and an $(n \times p)$ -matrix N , their <i>composite</i> is the $(m \times p)$ -matrix $M \circ N$
fong-spivak-seven-sketches_176 F01892	176	■ 0.958	$a \in \underline{m}$	$a \in \underline{m}$	defined as follows for any $a \in \underline{m}$ and $c \in \underline{p}$:
fong-spivak-seven-sketches_176 F01893	176	■ 0.958	$c \in \underline{p}$	$c \in \underline{p}$	defined as follows for any $a \in \underline{m}$ and $c \in \underline{p}$:
fong-spivak-seven-sketches_176 F01894	176	■ 0.871	$\sum_{b \in \underline{n}}$	$\sum_{b \in \underline{n}}$	Here the $\sum_{b \in \underline{n}}$ just means repeated addition (using the rig R 's $+$ operation), as usual.
fong-spivak-seven-sketches_176 F01895	176	■ 1.000	$A \ v$	Av	by multiplication in the order Av , where v is a column vector. In keeping with our
fong-spivak-seven-sketches_176 F01896	176	■ 1.000	$v \circ A$	$v \circ A$	composition convention, we use the opposite order, $v \circ A$, where v is a row vector. See
fong-spivak-seven-sketches_176 F01897	176	■ 1.000	$A: m \rightarrow n$	$A : m \rightarrow n$	monoidal product is given by the direct sum of matrices: given matrices $A: m \rightarrow n$
fong-spivak-seven-sketches_176 F01898	176	■ 1.000	$b: p \rightarrow q$	$b : p \rightarrow q$	and $b: p \rightarrow q$, we define $A + B: m + p \rightarrow n + q$ to be the block matrix
fong-spivak-seven-sketches_176 F01899	176	■ 1.000	$A+B: m+p \rightarrow n+q$	$A + B : m + p \rightarrow n + q$	and $b: p \rightarrow q$, we define $A + B: m + p \rightarrow n + q$ to be the block matrix
fong-spivak-seven-sketches_176 F01900	176	■ 1.000	$m \times q$	$m \times q$	where each 0 represents a matrix of zeros of the appropriate dimension $(m \times q$ and
fong-spivak-seven-sketches_176 F01901	176	■ 1.000	$n \times p$	$n \times p$	$n \times p)$. We refer to any combination of multiplication and direct sum as a <i>interconnection</i>

Identifier	Page	Conf.	LaTeX source	Rendered	Image																								
fong-spivak-seven-sketches-F01902	176	1.000	A+B	$A + B$	What is the direct sum matrix $A + B$?																								
fong-spivak-seven-sketches-F01903	177	1.000	1×1	1×1	matrix. For example, we interpret amplification by $a \in R$ as the 1×1 matrix $(a): 1 \rightarrow 1$:																								
fong-spivak-seven-sketches-F01904	177	1.000	$(a): 1 \rightarrow 1$	$(a): 1 \rightarrow 1$	matrix. For example, we interpret amplification by $a \in R$ as the 1×1 matrix $(a): 1 \rightarrow 1$:																								
fong-spivak-seven-sketches-F01905	177	1.000	$x \in R$	$x \in R$	it is an operation that takes an input $x \in R$ and returns $a * x$. Similarly, we can interpret																								
fong-spivak-seven-sketches-F01906	177	1.000	$a * x$	$a * x$	it is an operation that takes an input $x \in R$ and returns $a * x$. Similarly, we can interpret																								
fong-spivak-seven-sketches-F01907	177	0.897	$\supset$	$\supset$	$\supset$ as the 2×1 matrix $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$: it is an operation that takes a row vector consisting of two																								
fong-spivak-seven-sketches-F01908	177	0.897	2×1	2×1	$\supset$ as the 2×1 matrix $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$: it is an operation that takes a row vector consisting of two																								
fong-spivak-seven-sketches-F01909	177	0.897	$\binom{1}{1}$	$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	$\supset$ as the 2×1 matrix $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$: it is an operation that takes a row vector consisting of two																								
fong-spivak-seven-sketches-F01910	177	1.000	$x+y$	$x + y$	inputs, x and y , and returns $x + y$. Here is a table showing the interpretation of each																								
fong-spivak-seven-sketches-F01911	177	0.492	$1 \rightarrow 1$	$1 \rightarrow 1$	<table><thead><tr><th>generator</th><th>icon</th><th>matrix</th><th>arity</th></tr></thead><tbody><tr><td>amplify by $a \in R$</td><td></td><td>(a)</td><td>$1 \rightarrow 1$</td></tr><tr><td>add</td><td></td><td>$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$</td><td>$2 \rightarrow 1$</td></tr><tr><td>zero</td><td></td><td>$()$</td><td>$0 \rightarrow 1$</td></tr><tr><td>copy</td><td></td><td>$(1 \ 1)$</td><td>$1 \rightarrow 2$</td></tr><tr><td>discard</td><td></td><td>$()$</td><td>$1 \rightarrow 0$</td></tr></tbody></table>	generator	icon	matrix	arity	amplify by $a \in R$		(a)	$1 \rightarrow 1$	add		$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	$2 \rightarrow 1$	zero		$()$	$0 \rightarrow 1$	copy		$(1 \ 1)$	$1 \rightarrow 2$	discard		$()$	$1 \rightarrow 0$
generator	icon	matrix	arity																										
amplify by $a \in R$		(a)	$1 \rightarrow 1$																										
add		$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	$2 \rightarrow 1$																										
zero		$()$	$0 \rightarrow 1$																										
copy		$(1 \ 1)$	$1 \rightarrow 2$																										
discard		$()$	$1 \rightarrow 0$																										
fong-spivak-seven-sketches-F01912	177	0.492	$2 \rightarrow 1$	$2 \rightarrow 1$	<table><thead><tr><th>generator</th><th>icon</th><th>matrix</th><th>arity</th></tr></thead><tbody><tr><td>amplify by $a \in R$</td><td></td><td>(a)</td><td>$1 \rightarrow 1$</td></tr><tr><td>add</td><td></td><td>$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$</td><td>$2 \rightarrow 1$</td></tr><tr><td>zero</td><td></td><td>$()$</td><td>$0 \rightarrow 1$</td></tr><tr><td>copy</td><td></td><td>$(1 \ 1)$</td><td>$1 \rightarrow 2$</td></tr><tr><td>discard</td><td></td><td>$()$</td><td>$1 \rightarrow 0$</td></tr></tbody></table>	generator	icon	matrix	arity	amplify by $a \in R$		(a)	$1 \rightarrow 1$	add		$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	$2 \rightarrow 1$	zero		$()$	$0 \rightarrow 1$	copy		$(1 \ 1)$	$1 \rightarrow 2$	discard		$()$	$1 \rightarrow 0$
generator	icon	matrix	arity																										
amplify by $a \in R$		(a)	$1 \rightarrow 1$																										
add		$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	$2 \rightarrow 1$																										
zero		$()$	$0 \rightarrow 1$																										
copy		$(1 \ 1)$	$1 \rightarrow 2$																										
discard		$()$	$1 \rightarrow 0$																										
fong-spivak-seven-sketches-F01913	177	0.492	$0 \rightarrow 1$	$0 \rightarrow 1$	<table><thead><tr><th>generator</th><th>icon</th><th>matrix</th><th>arity</th></tr></thead><tbody><tr><td>amplify by $a \in R$</td><td></td><td>(a)</td><td>$1 \rightarrow 1$</td></tr><tr><td>add</td><td></td><td>$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$</td><td>$2 \rightarrow 1$</td></tr><tr><td>zero</td><td></td><td>$()$</td><td>$0 \rightarrow 1$</td></tr><tr><td>copy</td><td></td><td>$(1 \ 1)$</td><td>$1 \rightarrow 2$</td></tr><tr><td>discard</td><td></td><td>$()$</td><td>$1 \rightarrow 0$</td></tr></tbody></table>	generator	icon	matrix	arity	amplify by $a \in R$		(a)	$1 \rightarrow 1$	add		$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	$2 \rightarrow 1$	zero		$()$	$0 \rightarrow 1$	copy		$(1 \ 1)$	$1 \rightarrow 2$	discard		$()$	$1 \rightarrow 0$
generator	icon	matrix	arity																										
amplify by $a \in R$		(a)	$1 \rightarrow 1$																										
add		$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	$2 \rightarrow 1$																										
zero		$()$	$0 \rightarrow 1$																										
copy		$(1 \ 1)$	$1 \rightarrow 2$																										
discard		$()$	$1 \rightarrow 0$																										

Identifier	Page	Conf.	LaTeX source	Rendered	Image																								
					<table><tr><th>generator</th><th>icon</th><th>matrix</th><th>arity</th></tr><tr><td>amplify by $a \in R$</td><td></td><td>(a)</td><td>$1 \rightarrow 1$</td></tr><tr><td>add</td><td></td><td>$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$</td><td>$2 \rightarrow 1$</td></tr><tr><td>zero</td><td></td><td>$()$</td><td>$0 \rightarrow 1$</td></tr><tr><td>copy</td><td></td><td>$(1 \ 1)$</td><td>$1 \rightarrow 2$</td></tr><tr><td>discard</td><td></td><td>$()$</td><td>$1 \rightarrow 0$</td></tr></table>	generator	icon	matrix	arity	amplify by $a \in R$		(a)	$1 \rightarrow 1$	add		$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	$2 \rightarrow 1$	zero		$()$	$0 \rightarrow 1$	copy		$(1 \ 1)$	$1 \rightarrow 2$	discard		$()$	$1 \rightarrow 0$
generator	icon	matrix	arity																										
amplify by $a \in R$		(a)	$1 \rightarrow 1$																										
add		$\begin{pmatrix} 1 \\ 1 \end{pmatrix}$	$2 \rightarrow 1$																										
zero		$()$	$0 \rightarrow 1$																										
copy		$(1 \ 1)$	$1 \rightarrow 2$																										
discard		$()$	$1 \rightarrow 0$																										
fong-spivak-seven-sketches_177 F01914	177	0.492	$1 \rightarrow 0$	$1 \rightarrow 0$																									
fong-spivak-seven-sketches_177 F01915	177	1.000	$0 \times n$	$0 \times n$	dimensions. In linear algebra it is unusual to consider matrices of the form $0 \times n$ or $n \times 0$																								
fong-spivak-seven-sketches_177 F01916	177	1.000	$n \times 0$	$n \times 0$	dimensions. In linear algebra it is unusual to consider matrices of the form $0 \times n$ or $n \times 0$																								
fong-spivak-seven-sketches_177 F01917	177	0.749	0×3	0×3	you can multiply a 0×3 matrix by a $3 \times n$ matrix for any n , but you can not multiply it																								
fong-spivak-seven-sketches_177 F01918	177	0.749	$3 \times n$	$3 \times n$	you can multiply a 0×3 matrix by a $3 \times n$ matrix for any n , but you can not multiply it																								
fong-spivak-seven-sketches_177 F01919	177	1.000	$2 \times n$	$2 \times n$	by a $2 \times n$ matrix.																								
fong-spivak-seven-sketches_178 F01920	178	0.999	$S: \mathbf{SFG}_R \rightarrow \mathbf{Mat}(R)$	$S: \mathbf{SFG}_R \rightarrow \mathbf{Mat}(R)$	Theorem 5.53. There is a prop functor $S: \mathbf{SFG}_R \rightarrow \mathbf{Mat}(R)$ that sends the generators																								
fong-spivak-seven-sketches_178 F01921	178	1.000	$S(g)$	$S(g)$	We have now constructed a matrix $S(g)$ from any signal flow graph g . But how can																								
fong-spivak-seven-sketches_178 F01922	178	1.000	$m \times n$	$m \times n$	and for the generators in Definition 5.45, the associated matrix has dimension $m \times n$,																								
fong-spivak-seven-sketches_178 F01923	178	1.000	(i, j)	(i, j)	where m is the number of inputs and n the number of outputs, with (i, j) th entry																								
fong-spivak-seven-sketches_178 F01924	178	1.000	$\mathrm{id}_0: 0 \rightarrow 0, \mathrm{id}_1: 1 \rightarrow 1, \sigma: 2 \rightarrow 2$	$\mathrm{id}_0: 0 \rightarrow 0, \mathrm{id}_1: 1 \rightarrow 1, \sigma: 2 \rightarrow 2$	built from the morphisms $\mathrm{id}_0: 0 \rightarrow 0, \mathrm{id}_1: 1 \rightarrow 1, \sigma: 2 \rightarrow 2$, and the generators in G_R ,																								
fong-spivak-seven-sketches_178 F01925	178	1.000	$(\alpha + \beta): (m + p) \rightarrow (n + q)$	$(\alpha + \beta): (m + p) \rightarrow (n + q)$	• if $\alpha: m \rightarrow n$ and $\beta: p \rightarrow q$ are expressions, then $(\alpha + \beta): (m + p) \rightarrow (n + q)$ is an																								
fong-spivak-seven-sketches_178 F01926	178	0.999	$\mathrm{id}_0, \mathrm{id}_1$	$\mathrm{id}_0, \mathrm{id}_1$	is true when g is equal to id_0 , id_1 , and σ .																								
fong-spivak-seven-sketches_178 F01927	178	0.682	$0 \rightarrow 0$	$0 \rightarrow 0$	$(): 0 \rightarrow 0$. The morphisms id_1 , σ , and $a \in R$ map to the identity matrix, the swap																								
fong-spivak-seven-sketches_178 F01928	178	0.682	id_1, σ	id_1, σ	$(): 0 \rightarrow 0$. The morphisms id_1 , σ , and $a \in R$ map to the identity matrix, the swap																								
fong-spivak-seven-sketches_178 F01929	178	0.712	(a)	(a)	matrix, and the scalar matrix (a) respectively:																								
fong-spivak-seven-sketches_178 F01930	178	0.848	$n = p$	$n = p$	holds for (i) $\alpha \circ \beta$ (when $n = p$) and for (ii) $\alpha + \beta$ (in general).																								
fong-spivak-seven-sketches_178 F01931	178	0.848	$\alpha + \beta$	$\alpha + \beta$	holds for (i) $\alpha \circ \beta$ (when $n = p$) and for (ii) $\alpha + \beta$ (in general).																								
fong-spivak-seven-sketches_179 F01932	179	1.000	$\beta: n \rightarrow q$	$\beta: n \rightarrow q$	Here $\alpha: m \rightarrow n$ and $\beta: n \rightarrow q$ are signal flow graphs, assumed to obey the proposition.																								
fong-spivak-seven-sketches_179 F01933	179	0.994	$j \in \underline{n}$	$j \in \underline{n}$	k —of the amplification along that path. So let’s also fix some $j \in \underline{n}$, and consider paths																								
fong-spivak-seven-sketches_179 F01934	179	0.779	$\alpha /$	$\alpha /$	some j th output of α /input of β , the amplification that i contributes to k is																								
fong-spivak-seven-sketches_179 F01935	179	0.606	$S(\alpha) ; S(\beta)$	$S(\alpha); S(\beta)$	This is exactly the formula for matrix multiplication, which is composition $S(\alpha) \circ S(\beta)$																								

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_F01936	179	0.358	$\operatorname{Mat}(R)$	$\operatorname{Mat}(R)$	in the prop $\mathbf{Mat}(R)$; see Definition 5.50. So $\alpha \circ \beta$ obeys the proposition when α and β
fong-spivak-seven-sketches_F01937	179	0.684	$m+i$	$m+i$	of $\alpha + \beta$, the i th input of β is the $(m+i)$ th output of $\alpha + \beta$, and the j th output of β is the
fong-spivak-seven-sketches_F01938	179	0.999	$(n+j)$	$(n+j)$	$(n+j)$ th output of $\alpha + \beta$. This means that the matrix with (i,j) th entry describing the
fong-spivak-seven-sketches_F01939	179	1.000	$S(\alpha)+S(\beta)=S(\alpha+\beta)$	$S(\alpha) + S(\beta) = S(\alpha + \beta)$	amplification of the i th input that contributes to the j th output is $S(\alpha) + S(\beta) = S(\alpha + \beta)$,
fong-spivak-seven-sketches_F01940	181	0.989	$M \in \operatorname{Mat}(R)$	$M \in \operatorname{Mat}(R)$	Proposition 5.56. Given any matrix $M \in \mathbf{Mat}(R)$, there exists a signal flow graph
fong-spivak-seven-sketches_F01941	181	1.000	$g \in \mathbf{SFG}_R$	$g \in \mathbf{SFG}_R$	$g \in \mathbf{SFG}_R$ such that such that $S(g) = M$.
fong-spivak-seven-sketches_F01942	181	1.000	$S(g)=M$	$S(g) = M$	$g \in \mathbf{SFG}_R$ such that such that $S(g) = M$.
fong-spivak-seven-sketches_F01943	181	0.890	$M \in \mathbf{Mat}(R)$	$M \in \mathbf{Mat}(R)$	<i>Proof sketch.</i> Let $M \in \mathbf{Mat}(R)$ be an $(m \times n)$ -matrix. We want a signal flow graph g
fong-spivak-seven-sketches_F01944	181	1.000	$S(g)(i, j)$	$S(g)(i, j)$	such that $S(g) = M$. In particular, to compute $S(g)(i, j)$, we know that we can simply
fong-spivak-seven-sketches_F01945	181	1.000	$M(i, j)$	$M(i, j)$	and that this path has exactly one scalar multiplication icon, namely $M(i, j)$.
fong-spivak-seven-sketches_F01946	181	0.939	$\left(\begin{array}{cc} a & b \\ c & d \end{array}\right)$	$\left(\begin{array}{cc} a & b \\ c & d \end{array}\right)$	from just considering the case of 2×2 -matrices. Suppose M is the 2×2 -matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$.
fong-spivak-seven-sketches_F01947	181	1.000	$\left(\begin{array}{l} 0 \\ 1 \\ 2 \end{array}\right)$	$\left(\begin{array}{l} 0 \\ 1 \\ 2 \end{array}\right)$	1. $\begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix}$
fong-spivak-seven-sketches_F01948	181	1.000	$\left(\begin{array}{ll} 0 & 0 \\ 0 & 0 \end{array}\right)$	$\left(\begin{array}{cc} 0 & 0 \\ 0 & 0 \end{array}\right)$	2. $\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$
fong-spivak-seven-sketches_F01949	181	1.000	$\left(\begin{array}{lll} 1 & 2 & 3 \\ 4 & 5 & 6 \end{array}\right)$	$\left(\begin{array}{ccc} 1 & 2 & 3 \\ 4 & 5 & 6 \end{array}\right)$	3. $\boxed{\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}}$
fong-spivak-seven-sketches_F01950	182	1.000	$\mathbf{SFG}_R$	$\mathbf{SFG}_R$	the same as the set of generators for $\mathbf{SFG}_R$; see Definition 5.45.
fong-spivak-seven-sketches_F01951	182	1.000	$a, b \in R$	$a, b \in R$	We have the following equations for any $a, b \in R$:
fong-spivak-seven-sketches_F01952	183	0.997	$\binom{0}{6}$	$\binom{0}{6}$	<i>Example 5.61.</i> Both of the signal flow graphs below represent the same matrix, $\binom{0}{6}$:
fong-spivak-seven-sketches_F01953	183	0.549	$R=(\mathbb{N}, 0, +, 1, *)$	$R = (\mathbb{N}, 0, +, 1, *)$	1. Let $R = (\mathbb{N}, 0, +, 1, *)$. By examining the presentation of $\mathbf{Mat}(R)$ in Theorem 5.60,
fong-spivak-seven-sketches_F01954	183	1.000	$R=\mathbb{N} / 3 \mathbb{N}$	$R = \mathbb{N}/3\mathbb{N}$	2. Now suppose the rig is $R = \mathbb{N}/3\mathbb{N}$; if you do not know what this means, just
fong-spivak-seven-sketches_F01955	184	0.931	M, μ, η	M, μ, η	Definition 5.65. A monoid object (M, μ, η) in a symmetric monoidal category $(\mathcal{C}, I, \otimes)$ is
fong-spivak-seven-sketches_F01956	184	0.931	$\mathcal{C}, I, \otimes$	$\mathcal{C}, I, \otimes$	Definition 5.65. A monoid object (M, μ, η) in a symmetric monoidal category $(\mathcal{C}, I, \otimes)$ is
fong-spivak-seven-sketches_F01957	184	1.000	$\mu: M \otimes M \rightarrow M$	$\mu: M \otimes M \rightarrow M$	an object M of $\mathcal{C}$ together with morphisms $\mu: M \otimes M \rightarrow M$ and $\eta: I \rightarrow M$ such that
fong-spivak-seven-sketches_F01958	184	1.000	$\eta: I \rightarrow M$	$\eta: I \rightarrow M$	an object M of $\mathcal{C}$ together with morphisms $\mu: M \otimes M \rightarrow M$ and $\eta: I \rightarrow M$ such that

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F01959	184	0.939	$(\backslash\mathrm{id} \otimes \backslash\mathrm{id}) ; \backslash\mu=(\backslash\mathrm{id} \otimes \backslash\mu) ; \backslash\mu$	$(\mu \otimes \mathrm{id}); \mu = (\mathrm{id} \otimes \mu); \mu$	(a) $\mu \otimes \mathrm{id} \circ \mu = (\mathrm{id} \otimes \mu) \circ \mu$ and
fong-spivak-seven-sketches-F01960	184	1.000	$(\backslash\eta \otimes \mathrm{id}) ; \backslash\mu=\backslash\mathrm{id}=(\backslash\mathrm{id} \otimes \backslash\eta) \otimes \backslash\mu$	$(\eta \otimes \mathrm{id}); \mu = \mathrm{id} = (\mathrm{id} \otimes \eta); \mu$	(b) $\eta \otimes \mathrm{id} \circ \mu = \mathrm{id} = (\mathrm{id} \otimes \eta) \circ \mu$.
fong-spivak-seven-sketches-F01961	184	0.959	$\backslash\sigma_{M,M} ; \backslash\mu=\backslash\mu$	$\sigma_{M,M}; \mu = \mu$	(c) $\sigma_{M,M} \circ \mu = \mu$.
fong-spivak-seven-sketches-F01962	184	1.000	$\backslash\sigma_{M,M}$	$\sigma_{M,M}$	where $\sigma_{M,M}$ is the swap map on M in $\mathcal{C}$. We often denote it simply by σ .
fong-spivak-seven-sketches-F01963	184	1.000	$\backslash\mu: M \times M \rightarrowtail M$	$\mu: M \times M \rightarrow M$	Example 2.6; see also Example 3.13. That is, it is a set M , a function $\mu: M \times M \rightarrow M$,
fong-spivak-seven-sketches-F01964	184	0.935	$\backslash\eta(1) \in M$	$\eta(1) \in M$	which we denote by infix notation $*$, and an element $\eta(1) \in M$, which we denote by e ,
fong-spivak-seven-sketches-F01965	184	1.000	$(a * b) * c=a * (b * c)$	$(a * b) * c = a * (b * c)$	satisfying $a * b * c = a * (b * c)$ and $a * e = a = e * a$.
fong-spivak-seven-sketches-F01966	184	1.000	$a * e=a=e * a$	$a * e = a = e * a$	satisfying $a * b * c = a * (b * c)$ and $a * e = a = e * a$.
fong-spivak-seven-sketches-F01967	184	0.784	$\backslash\mu: \mathbb{R} \times \mathbb{R} \rightarrowtail \mathbb{R}$	$\mu: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$	1. Show that if $\mu: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is defined by $\mu(a, b) = a * b$ and if $\eta \in \mathbb{R}$ is defined to
fong-spivak-seven-sketches-F01968	184	0.784	$\backslash\mu(a, b)=a * b$	$\mu(a, b) = a * b$	1. Show that if $\mu: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is defined by $\mu(a, b) = a * b$ and if $\eta \in \mathbb{R}$ is defined to
fong-spivak-seven-sketches-F01969	184	0.784	$\backslash\eta \in \mathbb{R}$	$\eta \in \mathbb{R}$	1. Show that if $\mu: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is defined by $\mu(a, b) = a * b$ and if $\eta \in \mathbb{R}$ is defined to
fong-spivak-seven-sketches-F01970	184	1.000	$\backslash\eta=1$	$\eta = 1$	be $\eta = 1$, then $(\mathbb{R}, *, 1)$ satisfies all three conditions of Definition 5.65.
fong-spivak-seven-sketches-F01971	184	1.000	$(\mathbb{R}, *, 1)$	$(\mathbb{R}, *, 1)$	be $\eta = 1$, then $(\mathbb{R}, *, 1)$ satisfies all three conditions of Definition 5.65.
fong-spivak-seven-sketches-F01972	184	0.995	$\backslash\mu(a, b)=a+b$	$\mu(a, b) = a + b$	2. Show that if $\mu: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is defined by $\mu(a, b) = a + b$ and if $\eta \in \mathbb{R}$ is defined to
fong-spivak-seven-sketches-F01973	184	0.985	$\backslash\eta=0$	$\eta = 0$	be $\eta = 0$, then $(\mathbb{R}, +, 0)$ satisfies all three conditions of Definition 5.65.
fong-spivak-seven-sketches-F01974	184	0.985	$(\mathbb{R}, +, 0)$	$(\mathbb{R}, +, 0)$	be $\eta = 0$, then $(\mathbb{R}, +, 0)$ satisfies all three conditions of Definition 5.65.
fong-spivak-seven-sketches-F01975	184	0.457	$\backslash\mu=\succ-$	$\mu = \succ -$	Example 5.68. Graphically, we can depict $\mu = \succ-$ and $\eta = -$. Then axioms (a), (b), and
fong-spivak-seven-sketches-F01976	184	0.457	$\backslash\eta=\backslash\sigma$	$\eta = \sigma$	Example 5.68. Graphically, we can depict $\mu = \succ-$ and $\eta = -$. Then axioms (a), (b), and
fong-spivak-seven-sketches-F01977	184	0.315	(R)	(R)	following: the triple $(1, \succ-, -)$ is a commutative monoid object in the prop $\mathbf{Mat}(R)$.
fong-spivak-seven-sketches-F01978	185	0.665	$U: \operatorname{Mat}(R) \rightarrowtail$	$U: \operatorname{Mat}(R) \rightarrow$	Exercise 5.69. For any rig R , there is a functor $U: \mathbf{Mat}(R) \rightarrow \mathbf{Set}$, sending the object
fong-spivak-seven-sketches-F01979	185	0.999	$R^{\{n\}}$	R^n	$n \in \mathbb{N}$ to the set R^n , and sending a morphism (matrix) $M: m \rightarrow n$ to the function
fong-spivak-seven-sketches-F01980	185	0.999	$M: m \rightarrowtail n$	$M: m \rightarrow n$	$n \in \mathbb{N}$ to the set R^n , and sending a morphism (matrix) $M: m \rightarrow n$ to the function
fong-spivak-seven-sketches-F01981	185	1.000	$R^{\{m\}} \rightarrowtail R^{\{n\}}$	$R^m \rightarrow R^n$	$R^m \rightarrow R^n$ given by vector-matrix multiplication.
fong-spivak-seven-sketches-F01982	185	0.881	$\{1\}$	$\{1\}$	is $\{1\}$, a set with one element, and the monoidal product is $\times$ (see Example 4.49).
fong-spivak-seven-sketches-F01983	185	0.828	$(M, \backslash\mu, \backslash\eta)$	(M, μ, η)	2. Show that if (M, μ, η) is a monoid object in $\mathbf{Mat}(R)$ then $(U(M), U(\mu), U(\eta))$ is
fong-spivak-seven-sketches-F01984	185	0.828	$(U(M), U(\backslash\mu), U(\backslash\eta))$	$(U(M), U(\mu), U(\eta))$	2. Show that if (M, μ, η) is a monoid object in $\mathbf{Mat}(R)$ then $U(M), U(\mu), U(\eta)$ is

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F01985	185	■ 0.960	$\operatorname{prop} \operatorname{Mat}(R)$	$\operatorname{prop} \operatorname{Mat}(R)$	in the prop $\mathbf{Mat}(R)$. If $R = \mathbb{R}$ is the rig of real numbers, this means that we have
fong-spivak-seven-sketches-F01986	185	■ 0.960	$R = \mathbb{R}$	$R = \mathbb{R}$	in the prop $\mathbf{Mat}(R)$. If $R = \mathbb{R}$ is the rig of real numbers, this means that we have
fong-spivak-seven-sketches-F01987	185	■ 0.212	$\mathbf{Mat}(R)$	$\mathbf{Mat}(R)$	<i>Example 5.70.</i> The triple $(1, \multimap, \multimap)$ in $\mathbf{Mat}(R)$ forms a commutative monoid object in
fong-spivak-seven-sketches-F01988	185	■ 0.157	$\operatorname{Mat}(R)^{\operatorname{op}}$	$\operatorname{Mat}(R)^{\operatorname{op}}$	$\mathbf{Mat}(R)^{\operatorname{op}}$. We hence also say that $(1, \multimap, \multimap)$ forms a <i>co-commutative comonoid object</i> in
fong-spivak-seven-sketches-F01989	185	■ 0.245	$\varnothing, \mathbb{N}$	$\varnothing, \mathbb{N}$	(CMon, $\otimes, \mathbb{N}$) of commutative monoids with tensor product.
fong-spivak-seven-sketches-F01990	185	■ 1.000	$\mu: 2 \rightarrow 1$	$\mu: 2 \rightarrow 1$	<i>Remark 5.74.</i> If we present a prop $\mathcal{M}$ using two generators $\mu: 2 \rightarrow 1$ and $\eta: 0 \rightarrow 1$,
fong-spivak-seven-sketches-F01991	185	■ 1.000	$\eta: 0 \rightarrow 1$	$\eta: 0 \rightarrow 1$	<i>Remark 5.74.</i> If we present a prop $\mathcal{M}$ using two generators $\mu: 2 \rightarrow 1$ and $\eta: 0 \rightarrow 1$,
fong-spivak-seven-sketches-F01992	185	■ 1.000	$\mathcal{M} \rightarrow \mathcal{C}$	$\mathcal{M} \rightarrow \mathcal{C}$	in $\mathcal{C}$ correspond to strict monoidal functors $\mathcal{M} \rightarrow \mathcal{C}$. This sort of idea leads to the study
fong-spivak-seven-sketches-F01993	186	■ 0.998	$g: m \rightarrow n$	$g: m \rightarrow n$	The behavioral approach. A signal flow graph $g: m \rightarrow n$ takes an input $x \in R^m$ and
fong-spivak-seven-sketches-F01994	186	■ 0.998	$x \in R^m$	$x \in R^m$	The behavioral approach. A signal flow graph $g: m \rightarrow n$ takes an input $x \in R^m$ and
fong-spivak-seven-sketches-F01995	186	■ 1.000	$y \in R^n$	$y \in R^n$	gives an output $y \in R^n$. In fact, since this is all we care about, we might just think
fong-spivak-seven-sketches-F01996	186	■ 1.000	$\operatorname{B}(g) \subseteq R^m \times R^n$	$\operatorname{B}(g) \subseteq R^m \times R^n$	(x, y) . We'll call this set the <i>behavior</i> of g and denote it $\operatorname{B}(g) \subseteq R^m \times R^n$. For example,
fong-spivak-seven-sketches-F01997	186	■ 1.000	$(x, (x, x))$	$(x, (x, x))$	copy-behavior. Similarly, $(x, (x, x))$ is copy behavior for every $x \in R$, thus we have
fong-spivak-seven-sketches-F01998	186	■ 0.954	g^{op}	g^{op}	g^{op} , we must have $g^{\operatorname{op}}: n \rightarrow m$. Thus if $\operatorname{B}(g) \subseteq R^m \times R^n$ then we need $\operatorname{B}(g^{\operatorname{op}}) \subseteq R^n \times R^m$.
fong-spivak-seven-sketches-F01999	186	■ 0.954	$g^{\operatorname{op}}: n \rightarrow m$	$g^{\operatorname{op}}: n \rightarrow m$	g^{op} , we must have $g^{\operatorname{op}}: n \rightarrow m$. Thus if $\operatorname{B}(g) \subseteq R^m \times R^n$ then we need $\operatorname{B}(g^{\operatorname{op}}) \subseteq R^n \times R^m$.
fong-spivak-seven-sketches-F02000	186	■ 0.954	$\operatorname{B}(g^{\operatorname{op}}) \subseteq R^n \times R^m$	$\operatorname{B}(g^{\operatorname{op}}) \subseteq R^n \times R^m$	g^{op} , we must have $g^{\operatorname{op}}: n \rightarrow m$. Thus if $\operatorname{B}(g) \subseteq R^m \times R^n$ then we need $\operatorname{B}(g^{\operatorname{op}}) \subseteq R^n \times R^m$.
fong-spivak-seven-sketches-F02001	186	■ 0.996	(b, a)	(b, a)	One simple way to do this is to replace each (a, b) with (b, a) , so we would have
fong-spivak-seven-sketches-F02002	186	■ 0.854	$\operatorname{B}(\multimap)$	$\operatorname{B}(\multimap)$	1. What is the behavior $\operatorname{B}(\multimap)$ of the reversed addition icon $\multimap: 1 \rightarrow 2$?
fong-spivak-seven-sketches-F02003	186	■ 0.854	$\multimap: 1 \rightarrow 2$	$\multimap: 1 \rightarrow 2$	1. What is the behavior $\operatorname{B}(\multimap)$ of the reversed addition icon $\multimap: 1 \rightarrow 2$?
fong-spivak-seven-sketches-F02004	186	■ 0.833	$\operatorname{B}(\multimap)$	$\operatorname{B}(\multimap)$	2. What is the behavior $\operatorname{B}(\multimap)$ of the reversed copy icon, $\multimap: 2 \rightarrow 1$?
fong-spivak-seven-sketches-F02005	186	■ 0.833	$\multimap: 2 \rightarrow 1$	$\multimap: 2 \rightarrow 1$	2. What is the behavior $\operatorname{B}(\multimap)$ of the reversed copy icon, $\multimap: 2 \rightarrow 1$?
fong-spivak-seven-sketches-F02006	187	■ 1.000	$B \subseteq R^m \times R^n$	$B \subseteq R^m \times R^n$	of a signal flow graph $m \rightarrow n$ is a subset $B \subseteq R^m \times R^n$, i.e. a relation. Why not try to
fong-spivak-seven-sketches-F02007	187	■ 1.000	$N: n \rightarrow p$	$N: n \rightarrow p$	flow graphs with matrices $M: m \rightarrow n$ and $N: n \rightarrow p$, we see that their behaviors are
fong-spivak-seven-sketches-F02008	187	■ 1.000	$B_1 := \{(x, Mx) \mid x \in R^m\}$	$B_1 := \{(x, Mx) \mid x \in R^m\}$	the relations $B_1 := \{(x, Mx) \mid x \in R^m\}$ and $B_2 := \{(y, Ny) \mid y \in R^n\}$, while the behavior
fong-spivak-seven-sketches-F02009	187	■ 1.000	$B_2 := \{(y, Ny) \mid y \in R^n\}$	$B_2 := \{(y, Ny) \mid y \in R^n\}$	the relations $B_1 := \{(x, Mx) \mid x \in R^m\}$ and $B_2 := \{(y, Ny) \mid y \in R^n\}$, while the behavior
fong-spivak-seven-sketches-F02010	187	■ 0.999	$\{(x, x; M; N) \mid x \in R^m\}$	$\{(x, x; M; N) \mid x \in R^m\}$	of $M \circ N$ is the relation $\{(x, x \circ M \circ N) \mid x \in R^m\}$. This is a case of relation composition.
fong-spivak-seven-sketches-F02011	187	■ 0.881	$B_1 \subseteq R^m \times R^n$	$B_1 \subseteq R^m \times R^n$	Given relations $B_1 \subseteq R^m \times R^n$ and $B_2 \subseteq R^n \times R^p$, their composite $B_1 \circ B_2 \subseteq R^m \times R^p$ is

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02012	187	0.881	$B_{\{2\}} \subseteq R^{\{n\}} \times R^{\{p\}}$	$B_2 \subseteq R^n \times R^p$	Given relations $B_1 \subseteq R^m \times R^n$ and $B_2 \subseteq R^n \times R^p$, their composite $B_1 \circ B_2 \subseteq R^m \times R^p$ is
fong-spivak-seven-sketches-F02013	187	0.881	$B_{\{1\}} ; B_{\{2\}} \subseteq R^{\{m\}} \times R^{\{p\}}$	$B_1; B_2 \subseteq R^m \times R^p$	Given relations $B_1 \subseteq R^m \times R^n$ and $B_2 \subseteq R^n \times R^p$, their composite $B_1 \circ B_2 \subseteq R^m \times R^p$ is
fong-spivak-seven-sketches-F02014	187	0.953	$\mathbf{Rel}_R$	$\mathbf{Rel}_R$	Definition 5.79. Let R be a rig. We define the prop $\mathbf{Rel}_R$ of R -relations to have subsets
fong-spivak-seven-sketches-F02015	187	1.000	$C \subseteq R^{\{p\}} \times R^{\{q\}}$	$C \subseteq R^p \times R^q$	prop $\mathbf{Rel}_R$. If $B \subseteq R^m \times R^n$ and $C \subseteq R^p \times R^q$ are morphisms in $\mathbf{Rel}_R$, write down $B + C$
fong-spivak-seven-sketches-F02016	187	1.000	$B + C$	$B + C$	prop $\mathbf{Rel}_R$. If $B \subseteq R^m \times R^n$ and $C \subseteq R^p \times R^q$ are morphisms in $\mathbf{Rel}_R$, write down $B + C$
fong-spivak-seven-sketches-F02017	187	0.755	$g \in G_R$	$g \in G_R$	g^{op} for the mirror image of g , for each $g \in G_R$. So let's write $G_R^{\text{op}} := \{g^{\text{op}} \mid g \in G_R\}$ for
fong-spivak-seven-sketches-F02018	187	0.755	$G_R^{\text{op}} := \{g^{\text{op}} \mid g \in G_R\}$	$G_R^{\text{op}} := \{g^{\text{op}} \mid g \in G_R\}$	g^{op} for the mirror image of g , for each $g \in G_R$. So let's write $G_R^{\text{op}} := \{g^{\text{op}} \mid g \in G_R\}$ for
fong-spivak-seven-sketches-F02019	187	0.998	$\mathbf{SFG}_R^+$	$\mathbf{SFG}_R^+$	We call a morphism in the prop $\mathbf{SFG}_R^+$ a (<i>non-simplified</i>) <i>signal flow graph</i> : these extend
fong-spivak-seven-sketches-F02020	188	1.000	$g: m \rightarrow n, h: \ell \rightarrow n$	$g: m \rightarrow n, h: \ell \rightarrow n$	<i>Exercise 5.82.</i> Let $g: m \rightarrow n, h: \ell \rightarrow n$ be signal flow graphs. Note that $h^{\text{op}}: n \rightarrow \ell$ is
fong-spivak-seven-sketches-F02021	188	1.000	$h^{\text{op}}: n \rightarrow \ell$	$h^{\text{op}}: n \rightarrow \ell$	<i>Exercise 5.82.</i> Let $g: m \rightarrow n, h: \ell \rightarrow n$ be signal flow graphs. Note that $h^{\text{op}}: n \rightarrow \ell$ is
fong-spivak-seven-sketches-F02022	188	0.467	$g_9^{\circ}(h^{\text{op}})$	$g_9^{\circ}(h^{\text{op}})$	a signal flow graph, and we can form the composite $g \circ (h^{\text{op}})$:
fong-spivak-seven-sketches-F02023	188	0.274	$g^{\circ}(h^{\text{op}}) \subseteq R^m \times R^{\ell}$	$g^{\circ}(h^{\text{op}}) \subseteq R^m \times R^{\ell}$	Show that the behavior of $g \circ (h^{\text{op}}) \subseteq R^m \times R^{\ell}$ is equal to
fong-spivak-seven-sketches-F02024	188	0.807	$g: m \rightarrow n, h: m \rightarrow p$	$g: m \rightarrow n, h: m \rightarrow p$	<i>Exercise 5.83.</i> Let $g: m \rightarrow n, h: m \rightarrow p$ be signal flow graphs. Note that $(g^{\text{op}}): n \rightarrow m$
fong-spivak-seven-sketches-F02025	188	0.807	$(g^{\text{op}}): n \rightarrow m$	$(g^{\text{op}}): n \rightarrow m$	<i>Exercise 5.83.</i> Let $g: m \rightarrow n, h: m \rightarrow p$ be signal flow graphs. Note that $(g^{\text{op}}): n \rightarrow m$
fong-spivak-seven-sketches-F02026	188	0.856	$g^{\text{op}}; h$	$g^{\text{op}}; h$	is a signal flow graph, and we can form the composite $g^{\text{op}} \circ h$
fong-spivak-seven-sketches-F02027	188	0.963	$S(g) \in \mathbf{Mat}(R)$	$S(g) \in \mathbf{Mat}(R)$	$S(g) \in \mathbf{Mat}(R)$ be the associated $(m \times n)$ -matrix (see Theorem 5.53).
fong-spivak-seven-sketches-F02028	188	0.994	$b_1, b_2 \in \mathbf{B}(g)$	$b_1, b_2 \in \mathbf{B}(g)$	subspace. That is, if $b_1, b_2 \in \mathbf{B}(g)$ then so are $b_1 + b_2$ and $r * b_1$, for any $r \in R$. $\diamond$
fong-spivak-seven-sketches-F02029	188	0.994	$b_1 + b_2$	$b_1 + b_2$	subspace. That is, if $b_1, b_2 \in \mathbf{B}(g)$ then so are $b_1 + b_2$ and $r * b_1$, for any $r \in R$. $\diamond$
fong-spivak-seven-sketches-F02030	188	0.994	$r * b_1$	$r * b_1$	subspace. That is, if $b_1, b_2 \in \mathbf{B}(g)$ then so are $b_1 + b_2$ and $r * b_1$, for any $r \in R$. $\diamond$
fong-spivak-seven-sketches-F02031	188	0.994	$r \in R$	$r \in R$	subspace. That is, if $b_1, b_2 \in \mathbf{B}(g)$ then so are $b_1 + b_2$ and $r * b_1$, for any $r \in R$. $\diamond$
fong-spivak-seven-sketches-F02032	189	0.848	$\mathbf{LinRel}_R$	$\mathbf{LinRel}_R$	$\mathbf{LinRel}_R$. That is, one might want to show that there is a sub-prop of the prop $\mathbf{Rel}_R$
fong-spivak-seven-sketches-F02033	189	0.989	$(x, y) \in B$	$(x, y) \in B$	that B is linear. In other words, where for any $(x, y) \in B$ and $r \in R$, the element
fong-spivak-seven-sketches-F02034	189	0.730	$(r * x, r * y) \in R^m \times R^n$	$(r * x, r * y) \in R^m \times R^n$	$(r * x, r * y) \in R^m \times R^n$ is in B , and for any $(x', y') \in B$, the element $(x + x', y + y')$ is in B .
fong-spivak-seven-sketches-F02035	189	0.730	$(x', y') \in B$	$(x', y') \in B$	$(r * x, r * y) \in R^m \times R^n$ is in B , and for any $(x', y') \in B$, the element $(x + x', y + y')$ is in B .
fong-spivak-seven-sketches-F02036	189	0.730	$(x + x', y + y')$	$(x + x', y + y')$	$(r * x, r * y) \in R^m \times R^n$ is in B , and for any $(x', y') \in B$, the element $(x + x', y + y')$ is in B .
fong-spivak-seven-sketches-F02037	189	0.512	$G_R \sqcup G_R^{\text{op}}$	$G_R \sqcup G_R^{\text{op}}$	set is $G_R \sqcup G_R^{\text{op}}$ as in Eq. (5.81) and whose equations include those from Theorem 5.60,
fong-spivak-seven-sketches-F02038	189	1.000	η_1	η_1	Eq. (5.86) serve as the η_1 and ϵ_1 from Definition 4.58. Using monoidal products of these

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_189 F02039	189	1.000	$\backslash\epsilon_{1}$	ϵ_1	Eq. (5.86) serve as the η_1 and e_1 from Definition 4.58. Using monoidal products of these
fong-spivak-seven-sketches_190 F02040	190	1.000	$n=n^{*}$	$n = n^*$	is dual to itself, $n = n^*$.
fong-spivak-seven-sketches_190 F02041	190	1.000	$\mathbb{R}\left[s, s^{-1}\right]$	$\mathbb{R}\left[s, s^{-1}\right]$	$\mathbb{R}[s, s^{-1}]$ of polynomials in s and s^{-1} with coefficients in $\mathbb{R}$ and where $ss^{-1} = s^{-1}s = 1$.
fong-spivak-seven-sketches_190 F02042	190	1.000	s^{-1}	s^{-1}	$\mathbb{R}[s, s^{-1}]$ of polynomials in s and s^{-1} with coefficients in $\mathbb{R}$ and where $ss^{-1} = s^{-1}s = 1$.
fong-spivak-seven-sketches_190 F02043	190	1.000	$s\ s^{-1}=s^{-1}\ s=1$	$ss^{-1} = s^{-1}s = 1$	$\mathbb{R}[s, s^{-1}]$ of polynomials in s and s^{-1} with coefficients in $\mathbb{R}$ and where $ss^{-1} = s^{-1}s = 1$.
fong-spivak-seven-sketches_190 F02044	190	1.000	$a * u + b * v$	$a * u + b * v$	will take some linear combination $a * u + b * v$ of the desired speed and actual speed,
fong-spivak-seven-sketches_190 F02045	190	1.000	$\{(F, u, v) \mid u = v\}$	$\{(F, u, v) \mid u = v\}$	this signal flow graph close to the relation $\{(F, u, v) \mid u = v\}$? By phrasing problems in
fong-spivak-seven-sketches_195 F02046	195	0.872	$\mathcal{C}^{\prime}$	$\mathcal{C}'$	nomenon of the category $\mathcal{C}$. Our goal is to obtain $\mathcal{C}$'s colimits more directly, as a kind
fong-spivak-seven-sketches_195 F02047	195	0.444	$\operatorname{Cospan}_{\mathcal{C}}$	$\operatorname{Cospan}_{\mathcal{C}}$	that of <i>cospans in $\mathcal{C}$</i> , denoted $\operatorname{Cospan}_{\mathcal{C}}$ —that can conveniently package $\mathcal{C}$'s colimits in
fong-spivak-seven-sketches_195 F02048	195	1.000	$s_{\{m, n\}}: m \rightarrow n$	$s_{m,n} : m \rightarrow n$	(m, n) , an icon $s_{m,n} : m \rightarrow n$. These icons, known as <i>spiders</i> , ³ are drawn as follows:
fong-spivak-seven-sketches_197 F02049	197	1.000	$\varnothing \in \mathcal{C}$	$\varnothing \in \mathcal{C}$	Definition 6.1. Let $\mathcal{C}$ be a category. An <i>initial object</i> in $\mathcal{C}$ is an object $\varnothing \in \mathcal{C}$ such that for
fong-spivak-seven-sketches_197 F02050	197	1.000	$!_T: \varnothing \rightarrow T$	$!_T : \varnothing \rightarrow T$	each object T in $\mathcal{C}$ there exists a unique morphism $!_T : \varnothing \rightarrow T$.
fong-spivak-seven-sketches_197 F02051	197	0.844	$A, \leq$	$A, \leq$	1. $(A, \leq)$ has no initial object.
fong-spivak-seven-sketches_197 F02052	197	0.994	$(A, \leq)$	$(A, \leq)$	2. $(A, \leq)$ has exactly one initial object.
fong-spivak-seven-sketches_197 F02053	197	1.000	$\varnothing \rightarrow T$	$\varnothing \rightarrow T$	is a unique function $\varnothing \rightarrow T$, since $\varnothing$ has no elements.
fong-spivak-seven-sketches_197 F02054	197	0.615	$T \in \operatorname{Ob}(\mathcal{C})$	$T \in \operatorname{Ob}(\mathcal{C})$	show that for each object $T \in \operatorname{Ob}(\mathcal{C})$ (i.e. for both $T = A$ and $T = B$) there is a unique
fong-spivak-seven-sketches_197 F02055	197	0.615	$T=A$	$T = A$	show that for each object $T \in \operatorname{Ob}(\mathcal{C})$ (i.e. for both $T = A$ and $T = B$) there is a unique
fong-spivak-seven-sketches_197 F02056	197	0.615	$T=B$	$T = B$	show that for each object $T \in \operatorname{Ob}(\mathcal{C})$ (i.e. for both $T = A$ and $T = B$) there is a unique
fong-spivak-seven-sketches_197 F02057	197	0.999	$\varnothing = ? A$	$\varnothing = ? A$	morphism $\varnothing \rightarrow T$. Trying the case $\varnothing = ? A$ this condition fails when $T = B$: there are
fong-spivak-seven-sketches_197 F02058	197	1.000	$\varnothing = ? B$	$\varnothing = ? B$	two morphisms $A \rightarrow B$, not one. And trying the case $\varnothing = ? B$ this condition fails when
fong-spivak-seven-sketches_197 F02059	197	1.000	$B \rightarrow A$	$B \rightarrow A$	$T = A$: there are zero morphisms $B \rightarrow A$, not one.
fong-spivak-seven-sketches_197 F02060	197	0.374	$\boxed{\bullet}$	$\boxed{\bullet}$	1. $\boxed{\bullet}$
fong-spivak-seven-sketches_197 F02061	197	0.816	$\stackrel{a}{\bullet} \rightarrow \stackrel{b}{\bullet} \rightarrow \stackrel{c}{\bullet}$ $\rightarrow \stackrel{c}{\bullet}$	$\stackrel{a}{\bullet} \rightarrow \stackrel{b}{\bullet} \rightarrow \stackrel{c}{\bullet}$	2. $\left \begin{array}{c} \stackrel{a}{\bullet} \rightarrow \stackrel{b}{\bullet} \rightarrow \stackrel{c}{\bullet} \end{array} \right.$
fong-spivak-seven-sketches_197 F02062	197	0.981	$\stackrel{a}{\bullet} \stackrel{b}{\bullet}$	$\stackrel{a}{\bullet} \stackrel{b}{\bullet}$	3. $\left \begin{array}{cc} \stackrel{a}{\bullet} & \stackrel{b}{\bullet} \end{array} \right.$
fong-spivak-seven-sketches_197 F02063	197	0.985	$\left(R, 0_R, +_R, 1_R, *_R\right)$	$(R, 0_R, +_R, 1_R, *_R)$	$(R, 0_R, +_R, 1_R, *_R)$ to $(S, 0_S, +_S, 1_S, *_S)$ is a function $f: R \rightarrow S$ such that $f(0_R) = 0_S$,
fong-spivak-seven-sketches_197 F02064	197	0.985	$\left(S, 0_S, +_S, 1_S, *_S\right)$	$(S, 0_S, +_S, 1_S, *_S)$	$(R, 0_R, +_R, 1_R, *_R)$ to $(S, 0_S, +_S, 1_S, *_S)$ is a function $f: R \rightarrow S$ such that $f(0_R) = 0_S$,

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_197 F02065	197	■ 0.985	$f: R \rightarrow S$	$f: R \rightarrow S$	$(R, 0_R, +_R, 1_R, *_R)$ to $(S, 0_S, +_S, 1_S, *_S)$ is a function $f: R \rightarrow S$ such that $f(0_R) = 0_S$,
fong-spivak-seven-sketches_197 F02066	197	■ 0.985	$f(\text{left}(0_R)\text{right})=0_S$	$f(0_R) = 0_S$	$(R, 0_R, +_R, 1_R, *_R)$ to $(S, 0_S, +_S, 1_S, *_S)$ is a function $f: R \rightarrow S$ such that $f(0_R) = 0_S$,
fong-spivak-seven-sketches_197 F02067	197	■ 0.979	$f(\text{left}(r_1)+\{ \}_R \text{right})=f(\text{left}(r_1)\text{right})+s \text{ f}(\text{left}(r_2)\text{right})$	$f(r_1 +_R r_2) = f(r_1) +_S f(r_2)$	$f(r_1 +_R r_2) = f(r_1) +_S f(r_2)$, etc.
fong-spivak-seven-sketches_198 F02068	198	■ 1.000	$c_{\{1\}}$	c_1	Exercise 6.10. Let $\mathcal{C}$ be a category, and suppose that c_1 and c_2 are initial objects. Find
fong-spivak-seven-sketches_198 F02069	198	■ 1.000	$c_{\{2\}}$	c_2	Exercise 6.10. Let $\mathcal{C}$ be a category, and suppose that c_1 and c_2 are initial objects. Find
fong-spivak-seven-sketches_198 F02070	198	■ 0.978	$\iota_A: A \rightarrow$	$\iota_A: A \rightarrow$	is an object, which we denote $A + B$, together with a pair of morphisms $(\iota_A: A \rightarrow$
fong-spivak-seven-sketches_198 F02071	198	■ 0.977	$A+B, \iota_B: B \rightarrow A+B$	$A + B, \iota_B: B \rightarrow A + B$	$A + B, \iota_B: B \rightarrow A + B)$ such that for all objects T and pairs of morphisms $(f: A \rightarrow$
fong-spivak-seven-sketches_198 F02072	198	■ 0.977	$(f: A \rightarrow$	$(f: A \rightarrow$	$A + B, \iota_B: B \rightarrow A + B)$ such that for all objects T and pairs of morphisms $(f: A \rightarrow$
fong-spivak-seven-sketches_198 F02073	198	■ 0.994	$T, g: B \rightarrow T$	$T, g: B \rightarrow T$	$T, g: B \rightarrow T)$, there exists a unique morphism $[f, g]: A + B \rightarrow T$ such that the following
fong-spivak-seven-sketches_198 F02074	198	■ 0.994	$[f, g]: A+B \rightarrow T$	$[f, g]: A + B \rightarrow T$	$T, g: B \rightarrow T)$, there exists a unique morphism $[f, g]: A + B \rightarrow T$ such that the following
fong-spivak-seven-sketches_198 F02075	198	■ 0.999	$[f, g]$	$[f, g]$	We call $[f, g]$ the <i>copairing</i> of f and g .
fong-spivak-seven-sketches_198 F02076	198	■ 1.000	$\iota_A: A \rightarrow A \sqcup B$	$\iota_A: A \rightarrow A \sqcup B$	the disjoint union $A \sqcup B$ together with the inclusion functions $\iota_A: A \rightarrow A \sqcup B$ and
fong-spivak-seven-sketches_199 F02077	199	■ 1.000	$\iota_B: B \rightarrow A \sqcup B$	$\iota_B: B \rightarrow A \sqcup B$	$\iota_B: B \rightarrow A \sqcup B$
fong-spivak-seven-sketches_199 F02078	199	■ 1.000	$f: A \rightarrow T$	$f: A \rightarrow T$	Suppose we have functions $f: A \rightarrow T$ and $g: B \rightarrow T$ for some other set T ,
fong-spivak-seven-sketches_199 F02079	199	■ 1.000	$g: B \rightarrow T$	$g: B \rightarrow T$	Suppose we have functions $f: A \rightarrow T$ and $g: B \rightarrow T$ for some other set T ,
fong-spivak-seven-sketches_199 F02080	199	■ 0.591	$[f, g]: A \sqcup B \rightarrow T$	$[f, g]: A \sqcup B \rightarrow T$	$[f, g]: A \sqcup B \rightarrow T$ such that $\iota_A \circ [f, g] = f$ and $\iota_B \circ [f, g] = g$. What is it? Any element
fong-spivak-seven-sketches_199 F02081	199	■ 0.591	ι_A	ι_A	$[f, g]: A \sqcup B \rightarrow T$ such that $\iota_A \circ [f, g] = f$ and $\iota_B \circ [f, g] = g$. What is it? Any element
fong-spivak-seven-sketches_199 F02082	199	■ 0.591	$[f, g]=f$	$[f, g] = f$	$[f, g]: A \sqcup B \rightarrow T$ such that $\iota_A \circ [f, g] = f$ and $\iota_B \circ [f, g] = g$. What is it? Any element
fong-spivak-seven-sketches_199 F02083	199	■ 0.591	ι_B	ι_B	$[f, g]: A \sqcup B \rightarrow T$ such that $\iota_A \circ [f, g] = f$ and $\iota_B \circ [f, g] = g$. What is it? Any element
fong-spivak-seven-sketches_199 F02084	199	■ 0.591	$[f, g]=g$	$[f, g] = g$	$[f, g]: A \sqcup B \rightarrow T$ such that $\iota_A \circ [f, g] = f$ and $\iota_B \circ [f, g] = g$. What is it? Any element
fong-spivak-seven-sketches_199 F02085	199	■ 0.976	$x \in A \sqcup B$	$x \in A \sqcup B$	$x \in A \sqcup B$ is either ‘from A ’ or ‘from B ’, i.e. either there is some $a \in A$ with $x = \iota_A(a)$ or
fong-spivak-seven-sketches_199 F02086	199	■ 0.976	$x=\iota_A(a)$	$x = \iota_A(a)$	$x \in A \sqcup B$ is either ‘from A ’ or ‘from B ’, i.e. either there is some $a \in A$ with $x = \iota_A(a)$ or
fong-spivak-seven-sketches_199 F02087	199	■ 1.000	$b \in B$	$b \in B$	there is some $b \in B$ with $x = \iota_B(b)$. By Eq. (6.12), we must have:
fong-spivak-seven-sketches_199 F02088	199	■ 1.000	$x=\iota_B(b)$	$x = \iota_B(b)$	there is some $b \in B$ with $x = \iota_B(b)$. By Eq. (6.12), we must have:
fong-spivak-seven-sketches_199 F02089	199	■ 1.000	$T=\{a, b, c, \dots, z\}$	$T = \{a, b, c, \dots, z\}$	Exercise 6.16. Suppose $T = \{a, b, c, \dots, z\}$ is the set of letters in the alphabet, and let
fong-spivak-seven-sketches_199 F02090	199	■ 1.000	$f($	$f($	element of A to the first letter of its label, e.g. $f(\text{apple}) = a$. Let $g: B \rightarrow T$ be the
fong-spivak-seven-sketches_199 F02091	199	■ 1.000	$)=a$	$) = a$	element of A to the first letter of its label, e.g. $f(\text{apple}) = a$. Let $g: B \rightarrow T$ be the

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02092	199	0.999	$g($	$g($	function sending each element of B to the last letter of its label, e.g. $g(\text{apple}) = e$. Write
fong-spivak-seven-sketches-F02093	199	0.999	$)=e$	$) = e$	function sending each element of B to the last letter of its label, e.g. $g(\text{apple}) = e$. Write
fong-spivak-seven-sketches-F02094	199	1.000	$[f, g](x)$	$[f, g](x)$	down the function $[f, g](x)$ for all eight elements of $A \sqcup B$.
fong-spivak-seven-sketches-F02095	199	1.000	$f: A \rightarrow C, g: B \rightarrow C$	$f: A \rightarrow C, g: B \rightarrow C$	Exercise 6.17. Let $f: A \rightarrow C, g: B \rightarrow C$, and $h: C \rightarrow D$ be morphisms in a category $\mathcal{C}$
fong-spivak-seven-sketches-F02096	199	0.538	$\iota_A \sim: [f, g] = f$	$\iota_A: [f, g] = f$	1. $\boxed{\iota_A \circ [f, g] = f}$.
fong-spivak-seven-sketches-F02097	199	0.982	$\iota_B: [f, g] = g$	$\iota_B: [f, g] = g$	2. $\boxed{\iota_B \circ [f, g] = g}$.
fong-spivak-seven-sketches-F02098	199	1.000	$[f, g]; h = [f; h, g; h]$	$[f, g]; h = [f; h, g; h]$	3. $\boxed{[f, g] \circ h = [f \circ h, g \circ h]}$.
fong-spivak-seven-sketches-F02099	199	1.000	$\left[\iota_A, \iota_B\right] = \mathrm{id}_{A+B}$	$[\iota_A, \iota_B] = \mathrm{id}_{A+B}$	4. $\boxed{[\iota_A, \iota_B] = \mathrm{id}_{A+B}}$.
fong-spivak-seven-sketches-F02100	199	0.914	$(\mathcal{C}, +, \emptyset)$	$(\mathcal{C}, +, \emptyset)$	denoted $\emptyset$. Then $(\mathcal{C}, +, \emptyset)$ is a symmetric monoidal category (recall Definition 4.45). In
fong-spivak-seven-sketches-F02101	199	0.827	$\mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$	$\mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$	1. Show that $+$ extends to a functor $\mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$. In particular, how does it act on
fong-spivak-seven-sketches-F02102	199	0.771	$\mathcal{C} \times \mathcal{C}$	$\mathcal{C} \times \mathcal{C}$	morphisms in $\mathcal{C} \times \mathcal{C}$?
fong-spivak-seven-sketches-F02103	199	1.000	$A + \emptyset \rightarrow A$	$A + \emptyset \rightarrow A$	are isomorphisms $A + \emptyset \rightarrow A$ and $\emptyset + A \rightarrow A$.
fong-spivak-seven-sketches-F02104	199	1.000	$\emptyset + A \rightarrow A$	$\emptyset + A \rightarrow A$	are isomorphisms $A + \emptyset \rightarrow A$ and $\emptyset + A \rightarrow A$.
fong-spivak-seven-sketches-F02105	199	1.000	$(A+B)+C \rightarrow A+(B+C)$	$(A+B)+C \rightarrow A+(B+C)$	a) $\boxed{A+B+C \rightarrow A+(B+C)}$.
fong-spivak-seven-sketches-F02106	199	1.000	$A+B \rightarrow B+A$	$A+B \rightarrow B+A$	b) $A+B \rightarrow B+A$.
fong-spivak-seven-sketches-F02107	200	1.000	$f: A \rightarrow X$	$f: A \rightarrow X$	Definition 6.19. Let $\mathcal{C}$ be a category and let $f: A \rightarrow X$ and $g: A \rightarrow Y$ be morphisms
fong-spivak-seven-sketches-F02108	200	1.000	$g: A \rightarrow Y$	$g: A \rightarrow Y$	Definition 6.19. Let $\mathcal{C}$ be a category and let $f: A \rightarrow X$ and $g: A \rightarrow Y$ be morphisms
fong-spivak-seven-sketches-F02109	200	0.786	$X +_{\{ \}_A} Y$	$X +_A Y$	in $\mathcal{C}$ that have a common domain. The <i>pushout</i> $X +_A Y$ is the colimit of the diagram
fong-spivak-seven-sketches-F02110	200	0.857	$X +_{\{ \}_A} Y$	$X +_A Y$	In more detail, a pushout consists of (i) an object $X +_A Y$ and (ii) morphisms $\iota_X: X \rightarrow$
fong-spivak-seven-sketches-F02111	200	0.857	$\iota_X: X \rightarrow$	$\iota_X: X \rightarrow$	In more detail, a pushout consists of (i) an object $X +_A Y$ and (ii) morphisms $\iota_X: X \rightarrow$
fong-spivak-seven-sketches-F02112	200	0.905	$\iota_Y: Y \rightarrow X +_{\{ \}_A} Y$	$\iota_Y: Y \rightarrow X +_A Y$	$X +_A Y$ and $\iota_Y: Y \rightarrow X +_A Y$ satisfying (a) and (b) below.
fong-spivak-seven-sketches-F02113	200	0.371	$\ulcorner$	$\ulcorner$	commutes. (We will explain the “ $\ulcorner$ ” symbol below.)
fong-spivak-seven-sketches-F02114	200	1.000	$x: X \rightarrow T, y: Y \rightarrow T$	$x: X \rightarrow T, y: Y \rightarrow T$	(b) For all objects T and morphisms $\boxed{x: X \rightarrow T, y: Y \rightarrow T}$, if the diagram
fong-spivak-seven-sketches-F02115	200	0.933	$t: X +_{\{ \}_A} Y \rightarrow T$	$t: X +_A Y \rightarrow T$	commutes, then there exists a unique morphism $t: X +_A Y \rightarrow T$ such that
fong-spivak-seven-sketches-F02116	201	0.989	ι_X	ι_X	We further call ι_X the <i>pushout of g along f</i> , and similarly ι_Y the <i>pushout of f along g</i> .
fong-spivak-seven-sketches-F02117	201	0.989	ι_Y	ι_Y	We further call ι_X the <i>pushout of g along f</i> , and similarly ι_Y the <i>pushout of f along g</i> .
fong-spivak-seven-sketches-F02118	201	1.000	$B \leftarrow A \rightarrow C$	$B \leftarrow A \rightarrow C$	pushout of a diagram $B \leftarrow A \rightarrow C$ is equal to the coproduct $B \sqcup C$: namely, both are

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_20119 F02119	201	■ 1.000	$B \sqcup C$	$B \sqcup C$	pushout of a diagram $B \leftarrow A \rightarrow C$ is equal to the coproduct $B \sqcup C$: namely, both are
fong-spivak-seven-sketches_20120 F02120	201	■ 1.000	$B \vee C$	$B \vee C$	equal to the join $B \vee C$.
fong-spivak-seven-sketches_20121 F02121	201	■ 0.867	$i: A \rightarrowtail A'$	$i: A \rightarrow A'$	$i: A \rightarrow A'$ and $j: X \rightarrow X'$, we can take X' to be the pushout $X +_A A'$, i.e. the following
fong-spivak-seven-sketches_20122 F02122	201	■ 0.867	$j: X \rightarrowtail X'$	$j: X \rightarrow X'$	$i: A \rightarrow A'$ and $j: X \rightarrow X'$, we can take X' to be the pushout $X +_A A'$, i.e. the following
fong-spivak-seven-sketches_20123 F02123	201	■ 0.867	$X'^{\prime}$	X'	$i: A \rightarrow A'$ and $j: X \rightarrow X'$, we can take X' to be the pushout $X +_A A'$, i.e. the following
fong-spivak-seven-sketches_20124 F02124	201	■ 0.867	$X + \{ \}_A A'^{\prime}$	$X + {}_A A'$	$i: A \rightarrow A'$ and $j: X \rightarrow X'$, we can take X' to be the pushout $X +_A A'$, i.e. the following
fong-spivak-seven-sketches_20125 F02125	201	■ 0.847	$f'^{\prime} := i^{-1}: f: j$	$f' := i^{-1}: f: j$	where $f' := i^{-1} \circ f \circ j$. To see this, observe that if there is any object T such that the
fong-spivak-seven-sketches_20126 F02126	201	■ 0.997	$f; x = i; a$	$f; x = i; a$	then $f \circ x = i \circ a$, and so we are forced to take $x': X \rightarrow T$ to be $x' := j^{-1} \circ x$. This makes
fong-spivak-seven-sketches_20127 F02127	201	■ 0.997	$x'^{\prime}: X \rightarrowtail T$	$x': X \rightarrow T$	then $f \circ x = i \circ a$, and so we are forced to take $x': X \rightarrow T$ to be $x' := j^{-1} \circ x$. This makes
fong-spivak-seven-sketches_20128 F02128	201	■ 0.997	$x'^{\prime} := j^{-1}; x$	$x' := j^{-1}; x$	then $f \circ x = i \circ a$, and so we are forced to take $x': X \rightarrow T$ to be $x' := j^{-1} \circ x$. This makes
fong-spivak-seven-sketches_20129 F02129	201	■ 0.560	$f'^{\prime}; x'^{\prime} = i^{-1}; f; j; j^{-1}; x = i^{-1}; i; a = a$	$f'; x' = i^{-1}; f; j; j^{-1}; x = i^{-1}; i; a = a$	because $f' \circ x' = i^{-1} \circ f \circ j \circ j^{-1} \circ x = i^{-1} \circ i \circ a = a$.
fong-spivak-seven-sketches_20130 F02130	201	■ 0.998	$_S$	S	Exercise 6.24. For any set S , we have the discrete category $\mathbf{Disc}_S$, with S as objects
fong-spivak-seven-sketches_202131 F02131	202	■ 1.000	$\{f(a) \sim g(a) \mid a \in A\}$	$\{f(a) \sim g(a) \mid a \in A\}$	relation $\{f(a) \sim g(a) \mid a \in A\}$.
fong-spivak-seven-sketches_202132 F02132	202	■ 1.000	$f(a)$	$f(a)$	a connection between $f(a)$ in X and $g(a)$ in Y . The pushout is the set of connected
fong-spivak-seven-sketches_202133 F02133	202	■ 1.000	$g(a)$	$g(a)$	a connection between $f(a)$ in X and $g(a)$ in Y . The pushout is the set of connected
fong-spivak-seven-sketches_202134 F02134	202	■ 1.000	$f: \underline{4} \rightarrowtail \underline{5}$	$f: \underline{4} \rightarrow \underline{5}$	Exercise 6.26. What is the pushout of the functions $f: \underline{4} \rightarrow \underline{5}$ and $g: \underline{4} \rightarrow \underline{3}$ pictured
fong-spivak-seven-sketches_202135 F02135	202	■ 1.000	$g: \underline{4} \rightarrowtail \underline{3}$	$g: \underline{4} \rightarrow \underline{3}$	Exercise 6.26. What is the pushout of the functions $f: \underline{4} \rightarrow \underline{5}$ and $g: \underline{4} \rightarrow \underline{3}$ pictured
fong-spivak-seven-sketches_202136 F02136	202	■ 1.000	$X, Y \in$	$X, Y \in$	Example 6.27. Suppose a category $\mathcal{C}$ has an initial object $\emptyset$. For any two objects $X, Y \in$
fong-spivak-seven-sketches_202137 F02137	202	■ 0.785	$f: \varnothing \rightarrowtail X$	$f: \emptyset \rightarrow X$	$\text{Ob } \mathcal{C}$, there is a unique morphism $f: \emptyset \rightarrow X$ and a unique morphism $g: \emptyset \rightarrow Y$; this
fong-spivak-seven-sketches_202138 F02138	202	■ 0.785	$g: \varnothing \rightarrowtail Y$	$g: \emptyset \rightarrow Y$	$\text{Ob } \mathcal{C}$, there is a unique morphism $f: \emptyset \rightarrow X$ and a unique morphism $g: \emptyset \rightarrow Y$; this
fong-spivak-seven-sketches_202139 F02139	202	■ 0.884	$X \stackrel{f}{\leftarrow} \varnothing \stackrel{g}{\rightarrow} Y$	$X \xleftarrow{f} \emptyset \xrightarrow{g} Y$	The diagram $X \xleftarrow{f} \emptyset \xrightarrow{g} Y$ has a pushout in $\mathcal{C}$ iff X and Y have a coproduct in $\mathcal{C}$,
fong-spivak-seven-sketches_202140 F02140	202	■ 1.000	$X + Y$	$X + Y$	coproduct $X + Y$; then the diagram to the left
fong-spivak-seven-sketches_202141 F02141	202	■ 0.897	$X + Y \rightarrowtail T$	$X + Y \rightarrow T$	is a unique map $X + Y \rightarrow T$ making the diagram as in Eq. (6.21) commute (why? ²).
fong-spivak-seven-sketches_202142 F02142	202	■ 0.615	$X + \varnothing Y \cong X + Y$	$X + \emptyset Y \cong X + Y$	This shows that $X + Y$ is a pushout, $X +_{\emptyset} Y \cong X + Y$.
fong-spivak-seven-sketches_202143 F02143	202	■ 0.986	$X + \varnothing Y$	$X + \emptyset Y$	Similarly, if a pushout $X +_{\emptyset} Y$ exists, then it satisfies the universal property of the
fong-spivak-seven-sketches_202144 F02144	202	■ 0.991	$A = X = Y = \mathbb{N}$	$A = X = Y = \mathbb{N}$	Example 6.29. Let $A = X = Y = \mathbb{N}$. Consider the functions $f: A \rightarrow X$ and $g: A \rightarrow Y$
fong-spivak-seven-sketches_203145 F02145	203	■ 1.000	$f(a) := \lfloor a / 2 \rfloor$	$f(a) := \lfloor a / 2 \rfloor$	given by the ‘floor’ functions, $f(a) := \lfloor a / 2 \rfloor$ and $g(a) := \lfloor (a + 1) / 2 \rfloor$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02146	203	■ 1.000	$g(a) := \lfloor (a+1)/2 \rfloor$	$g(a) := \lfloor (a+1)/2 \rfloor$	given by the ‘floor’ functions, $f(a) := \lfloor a/2 \rfloor$ and $g(a) := \lfloor (a+1)/2 \rfloor$.
fong-spivak-seven-sketches-F02147	203	■ 1.000	$x: X \rightarrow T$	$x: X \rightarrow T$	If T is any other set and we have maps $x: X \rightarrow T$ and $y: Y \rightarrow T$ that commute with
fong-spivak-seven-sketches-F02148	203	■ 1.000	$y: Y \rightarrow T$	$y: Y \rightarrow T$	If T is any other set and we have maps $x: X \rightarrow T$ and $y: Y \rightarrow T$ that commute with
fong-spivak-seven-sketches-F02149	203	■ 1.000	$f \cong x = g \cong y$	$f \cong x = g \cong y$	f and g , i.e. $f \cong x = g \cong y$, then this commutativity implies that
fong-spivak-seven-sketches-F02150	203	■ 0.781	$f(1)=0$	$f(1) = 0$	In other words, Y ’s 0 and X ’s 0 go to the same place in T , say t . But since $f(1) = 0$
fong-spivak-seven-sketches-F02151	203	■ 0.686	$g(1)=1$	$g(1) = 1$	and $g(1) = 1$, we also have that $t = x(0) = x(f(1)) = y(g(1)) = y(1)$. This means Y ’s 1
fong-spivak-seven-sketches-F02152	203	■ 0.686	$t=x(0)=x(f(1))=y(g(1))=y(1)$	$t = x(0) = x(f(1)) = y(g(1)) = y(1)$	and $g(1) = 1$, we also have that $t = x(0) = x(f(1)) = y(g(1)) = y(1)$. This means Y ’s 1
fong-spivak-seven-sketches-F02153	203	■ 1.000	$g(2)=1$	$g(2) = 1$	goes to t also. But since $g(2) = 1$ and $f(2) = 1$, we also have that $t = g(1) = y(g(2)) =$
fong-spivak-seven-sketches-F02154	203	■ 1.000	$f(2)=1$	$f(2) = 1$	goes to t also. But since $g(2) = 1$ and $f(2) = 1$, we also have that $t = g(1) = y(g(2)) =$
fong-spivak-seven-sketches-F02155	203	■ 1.000	$t=g(1)=y(g(2))=$	$t = g(1) = y(g(2)) =$	goes to t also. But since $g(2) = 1$ and $f(2) = 1$, we also have that $t = g(1) = y(g(2)) =$
fong-spivak-seven-sketches-F02156	203	■ 0.900	$x(f(2))=x(1)$	$x(f(2)) = x(1)$	$x(f(2)) = x(1)$, which means that X ’s 1 also goes to t . One can keep repeating this
fong-spivak-seven-sketches-F02157	203	■ 0.996	$X \sqcup_A Y \cong \{1\}$	$X \sqcup_A Y \cong \{1\}$	induction, one can prove that the pushout is in fact a 1-element set, $X \sqcup_A Y \cong \{1\}$.
fong-spivak-seven-sketches-F02158	203	■ 0.832	$j \in \operatorname{Ob} \mathcal{J}$	$j \in \operatorname{Ob} \mathcal{J}$	must have finitely many objects too, because each object $j \in \operatorname{Ob} \mathcal{J}$ has its own identity morphism id_j .
fong-spivak-seven-sketches-F02159	203	■ 0.832	id_j	id_j	must have finitely many objects too, because each object $j \in \operatorname{Ob} \mathcal{J}$ has its own identity morphism id_j .
fong-spivak-seven-sketches-F02160	204	■ 0.996	$\operatorname{colim}_{\mathcal{J}} D$	$\operatorname{colim}_{\mathcal{J}} D$	Definition 6.30. We say that a category $\mathcal{C}$ has finite colimits if a colimit, $\operatorname{colim}_{\mathcal{J}} D$, exists
fong-spivak-seven-sketches-F02161	204	■ 0.890	$\mathbf{0} \rightarrow \mathcal{C}$	$\mathbf{0} \rightarrow \mathcal{C}$	!: $\mathbf{0} \rightarrow \mathcal{C}$, where $\mathbf{0}$ is the category with no objects and no morphisms, and ! is the unique
fong-spivak-seven-sketches-F02162	205	■ 1.000	$(v, d) \sim (w, e)$	$(v, d) \sim (w, e)$	generated by putting $(v, d) \sim (w, e)$ if there is an arrow $a: v \rightarrow w$ in J such that
fong-spivak-seven-sketches-F02163	205	■ 1.000	$a: v \rightarrow w$	$a: v \rightarrow w$	generated by putting $(v, d) \sim (w, e)$ if there is an arrow $a: v \rightarrow w$ in J such that
fong-spivak-seven-sketches-F02164	205	■ 0.996	$D(a)(d)=e$	$D(a)(d) = e$	$D(a)(d) = e$. Then this set, together with the functions $\iota_v: D(v) \rightarrow \operatorname{colim}_{\mathcal{J}} D$ given by
fong-spivak-seven-sketches-F02165	205	■ 0.996	$\iota_v: D(v) \rightarrow \operatorname{colim}_{\mathcal{J}} D$	$\iota_v: D(v) \rightarrow \operatorname{colim}_{\mathcal{J}} D$	$D(a)(d) = e$. Then this set, together with the functions $\iota_v: D(v) \rightarrow \operatorname{colim}_{\mathcal{J}} D$ given by
fong-spivak-seven-sketches-F02166	205	■ 1.000	$d \in D(v)$	$d \in D(v)$	sending $d \in D(v)$ to its equivalence class, constitutes a colimit of D .
fong-spivak-seven-sketches-F02167	205	■ 0.654	$\mathfrak{J} =$	$\mathfrak{J} =$	<i>Example 6.39.</i> A coproduct is a colimit on the graph $\mathcal{J} = \begin{array}{cc} v_1 & v_2 \\ \bullet & \bullet \end{array}$. A functor $D: \mathcal{J} \rightarrow \mathbf{Set}$
fong-spivak-seven-sketches-F02168	205	■ 0.654	$D: \mathfrak{J} \rightarrow$	$D: \mathfrak{J} \rightarrow$	<i>Example 6.39.</i> A coproduct is a colimit on the graph $\mathcal{J} = \begin{array}{cc} v_1 & v_2 \\ \bullet & \bullet \end{array}$. A functor $D: \mathcal{J} \rightarrow \mathbf{Set}$
fong-spivak-seven-sketches-F02169	205	■ 1.000	$X := D(v_1)$	$X := D(v_1)$	can be identified with a choice of two sets, $X := D(v_1)$ and $Y := D(v_2)$. Since there are
fong-spivak-seven-sketches-F02170	205	■ 1.000	$Y := D(v_2)$	$Y := D(v_2)$	can be identified with a choice of two sets, $X := D(v_1)$ and $Y := D(v_2)$. Since there are
fong-spivak-seven-sketches-F02171	206	■ 0.716	$\mathbf{1} = \boxed{\bullet}$	$\mathbf{1} = \boxed{\bullet}$	<i>Example 6.40.</i> If $\mathcal{J}$ is the category $\mathbf{1} = \boxed{\bullet}$, the formula in Theorem 6.37 yields the set
fong-spivak-seven-sketches-F02172	206	■ 1.000	$D(v)$	$D(v)$	This is isomorphic to the set $D(v)$. In other words, if X is a set considered as a diagram

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_206 F02173	206	0.824	$X: \mathbf{1} \rightarrow$	$X : \mathbf{1} \rightarrow$	$X: \mathbf{1} \rightarrow \mathbf{Set}$, then its colimit (like its limit) is just X again.
fong-spivak-seven-sketches_206 F02174	206	1.000	$X \stackrel{f}{\leftarrow} N \xrightarrow{g} Y$	$X \xleftarrow{f} N \xrightarrow{g} Y$	diagram $X \xleftarrow{f} N \xrightarrow{g} Y$ —agree with the description we gave in Example 6.25.
fong-spivak-seven-sketches_206 F02175	206	0.994	$X \xrightarrow[g]{f} Y$	$X \xrightarrow[g]{f} Y$	Consider some diagram $X \xrightarrow[g]{f} Y$ on this graph in $\mathbf{Set}$. The coequalizer of this
fong-spivak-seven-sketches_206 F02176	206	1.000	$y \sim y^{\prime}$	$y \sim y'$	declaring $y \sim y'$ whenever there exists x in X such that $f(x) = y$ and $g(x) = y'$.
fong-spivak-seven-sketches_206 F02177	206	1.000	$f(x)=y$	$f(x) = y$	declaring $y \sim y'$ whenever there exists x in X such that $f(x) = y$ and $g(x) = y'$.
fong-spivak-seven-sketches_206 F02178	206	1.000	$g(x)=y^{\prime}$	$g(x) = y'$	declaring $y \sim y'$ whenever there exists x in X such that $f(x) = y$ and $g(x) = y'$.
fong-spivak-seven-sketches_206 F02179	206	0.777	$f, g: X \rightarrow Y$	$f, g: X \rightarrow Y$	circles $\bullet$, the blue and red arrows respectively define functions $f, g: X \rightarrow Y$. Let's
fong-spivak-seven-sketches_206 F02180	206	1.000	$A \rightarrow N \leftarrow B$	$A \rightarrow N \leftarrow B$	common object $A \rightarrow N \leftarrow B$. The common object N is called the <i>apex</i> of the cospan
fong-spivak-seven-sketches_207 F02181	207	0.749	$N \stackrel{g}{\leftarrow} B \xrightarrow{h} P$	$N \xleftarrow{g} B \xrightarrow{h} P$	Then, if a pushout of $N \xleftarrow{g} B \xrightarrow{h} P$ exists in $\mathcal{C}$, as shown on the left, we can extract a
fong-spivak-seven-sketches_207 F02182	207	1.000	$C \in \mathrm{Ob} \, \mathcal{C}$	$C \in \mathrm{Ob} \mathcal{C}$	need an identity on every object $C \in \mathrm{Ob} \, \mathcal{C}$; but that's not hard: use $C \rightarrow C \leftarrow C$ where
fong-spivak-seven-sketches_207 F02183	207	1.000	$C \rightarrow C \leftarrow C$	$C \rightarrow C \leftarrow C$	need an identity on every object $C \in \mathrm{Ob} \, \mathcal{C}$; but that's not hard: use $C \rightarrow C \leftarrow C$ where
fong-spivak-seven-sketches_207 F02184	207	1.000	$A \rightarrow B \rightarrow C$	$A \rightarrow B \rightarrow C$	so we don't know that every two sequential morphisms $A \rightarrow B \rightarrow C$ can be composed.
fong-spivak-seven-sketches_207 F02185	207	1.000	$A \rightarrow P \leftarrow B$	$A \rightarrow P \leftarrow B$	between their respective apexes. That is, the cospan $A \rightarrow P \leftarrow B$ and $A \rightarrow P' \leftarrow B$
fong-spivak-seven-sketches_207 F02186	207	1.000	$A \rightarrow P^{\prime} \leftarrow B$	$A \rightarrow P' \leftarrow B$	between their respective apexes. That is, the cospan $A \rightarrow P \leftarrow B$ and $A \rightarrow P' \leftarrow B$
fong-spivak-seven-sketches_207 F02187	207	1.000	$P \cong P^{\prime}$	$P \cong P'$	in the diagram shown left below are equivalent iff there is an isomorphism $P \cong P'$
fong-spivak-seven-sketches_207 F02188	207	0.761	$\{ \}_{{\text{\mathcal{C}}}}$	$\mathcal{C}$	business: $\mathbf{Cospan}_{\mathcal{C}}$ will form a category. But in fact, we are very close to getting more.
fong-spivak-seven-sketches_207 F02189	207	0.743	$\operatorname{Cospan}_{\mathcal{C}}$	$\operatorname{Cospan}_{\mathcal{C}}$	If we also demand that $\mathcal{C}$ has an initial object $\emptyset$ as well, then we can upgrade $\mathbf{Cospan}_{\mathcal{C}}$
fong-spivak-seven-sketches_208 F02190	208	0.839	$_{\mathcal{C}}$	c	$\mathbf{Cospan}_{\mathcal{C}}$ with the same objects as $\mathcal{C}$, i.e. $\operatorname{Ob}(\mathbf{Cospan}_{\mathcal{C}}) = \operatorname{Ob}(\mathcal{C})$, where the morphisms
fong-spivak-seven-sketches_208 F02191	208	0.839	$\operatorname{Ob}(\left(\right.$	$\operatorname{Ob} ($	$\mathbf{Cospan}_{\mathcal{C}}$ with the same objects as $\mathcal{C}$, i.e. $\operatorname{Ob}(\mathbf{Cospan}_{\mathcal{C}}) = \operatorname{Ob}(\mathcal{C})$, where the morphisms
fong-spivak-seven-sketches_208 F02192	208	0.839	$\left. _{\mathcal{C}} \right) = \operatorname{Ob}(\operatorname{Ob}(\mathcal{C}))$	$c) = \operatorname{Ob}(\mathcal{C})$	$\mathbf{Cospan}_{\mathcal{C}}$ with the same objects as $\mathcal{C}$, i.e. $\operatorname{Ob}(\mathbf{Cospan}_{\mathcal{C}}) = \operatorname{Ob}(\mathcal{C})$, where the morphisms
fong-spivak-seven-sketches_208 F02193	208	0.370	$_{\mathcal{C}}, \varnothing, +$	$\mathcal{C}, \emptyset, +$	There is a symmetric monoidal structure on this category, denoted $(\mathbf{Cospan}_{\mathcal{C}}, \emptyset, +)$.
fong-spivak-seven-sketches_208 F02194	208	1.000	$A + \varnothing \cong A$	$A + \emptyset \cong A$	coproduct. The coherence isomorphisms, e.g. $A + \emptyset \cong A$, can be defined in a similar
fong-spivak-seven-sketches_208 F02195	208	0.699	$\{ \}_{\mathcal{C}}, \varnothing, +$	$\mathcal{C}, \emptyset, +$	It is a straightforward but time-consuming exercise to verify that $(\mathbf{Cospan}_{\mathcal{C}}, \emptyset, +)$
fong-spivak-seven-sketches_208 F02196	208	0.955	$_{\text{\texttt{FinSet}}}$	$\mathbf{FinSet}$	symmetric monoidal category $\mathbf{Cospan}_{\mathbf{FinSet}}$. What does it look like? It looks a lot like
fong-spivak-seven-sketches_208 F02197	208	1.000	$\{ \}_{\text{\texttt{FinSet}}}$	$\mathbf{FinSet}$	The objects of $\mathbf{Cospan}_{\mathbf{FinSet}}$ are finite sets; here let's draw them as collections of $\bullet$'s.
fong-spivak-seven-sketches_208 F02198	208	0.958	$A \xrightarrow{f} N \stackrel{g}{\leftarrow} B$	$A \xrightarrow{f} N \xleftarrow{g} B$	six element set. Below are two depictions of a cospan $A \xrightarrow{f} N \xleftarrow{g} B$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_208 F02199	208	■ 1.000	$g(b)$	$g(b)$	arrows from each $a \in A$ to $f(a)$ and each $b \in B$ to $g(b)$. In the depiction on the right, we
fong-spivak-seven-sketches_209 F02200	209	■ 0.904	$A+B \rightarrowtail B+C$	$A+B \rightarrow B+C$	Draw their monoidal product as a morphism $A+B \rightarrow B+C$ in $\mathbf{Cospan}_{\mathbf{FinSet}}$.
fong-spivak-seven-sketches_209 F02201	209	■ 0.904	$\operatorname{Cospan}_{\text{FinSet}}$	$\operatorname{Cospan}_{\mathbf{FinSet}}$	Draw their monoidal product as a morphism $A+B \rightarrow B+C$ in $\mathbf{Cospan}_{\mathbf{FinSet}}$.
fong-spivak-seven-sketches_210 F02202	210	■ 0.992	(X, μ, η)	(X, μ, η)	To make this precise, recall from Definition 5.65 that a commutative monoid (X, μ, η)
fong-spivak-seven-sketches_210 F02203	210	■ 1.000	$\mu: X \otimes X \rightarrowtail X$	$\mu: X \otimes X \rightarrow X$	$\mu: X \otimes X \rightarrow X$
fong-spivak-seven-sketches_210 F02204	210	■ 1.000	$\eta: I \rightarrowtail X$	$\eta: I \rightarrow X$	$\eta: I \rightarrow X$
fong-spivak-seven-sketches_210 F02205	210	■ 0.963	$X \otimes X$	$X \otimes X$	where $\bowtie$ is the symmetry on $X \otimes X$. A cocommutative comonoid (X, δ, ϵ) is an
fong-spivak-seven-sketches_210 F02206	210	■ 0.963	(X, δ, ϵ)	(X, δ, ϵ)	where $\bowtie$ is the symmetry on $X \otimes X$. A cocommutative comonoid (X, δ, ϵ) is an
fong-spivak-seven-sketches_210 F02207	210	■ 1.000	$\delta: X \rightarrowtail X \otimes X, \epsilon: X \rightarrowtail I$	$\delta: X \rightarrow X \otimes X, \epsilon: X \rightarrow I$	object X with maps $\delta: X \rightarrow X \otimes X, \epsilon: X \rightarrow I$, obeying the mirror images of the laws in
fong-spivak-seven-sketches_210 F02208	210	■ 1.000	μ, η, δ	μ, η, δ	comonoid, and consider a wiring diagram built only from the icons μ, η, δ , and ϵ ,
fong-spivak-seven-sketches_210 F02209	210	■ 1.000	ϵ	ϵ	comonoid, and consider a wiring diagram built only from the icons μ, η, δ , and ϵ ,
fong-spivak-seven-sketches_210 F02210	210	■ 1.000	μ	μ	how the morphisms μ and δ , the merger and the splitter, interact.
fong-spivak-seven-sketches_210 F02211	210	■ 1.000	δ	δ	how the morphisms μ and δ , the merger and the splitter, interact.
fong-spivak-seven-sketches_210 F02212	210	■ 0.953	$(\mathcal{C}, \otimes, I)$	$(\mathcal{C}, \otimes, I)$	Definition 6.52. Let X be an object in a symmetric monoidal category $(\mathcal{C}, \otimes, I)$. A Frobenius
fong-spivak-seven-sketches_210 F02213	210	■ 0.574	$(\mu, \eta, \delta, \epsilon)$	$(\mu, \eta, \delta, \epsilon)$	structure on X consists of a 4-tuple $(\mu, \eta, \delta, \epsilon)$ such that (X, μ, η) is a commutative
fong-spivak-seven-sketches_211 F02214	211	■ 1.000	$X^{\otimes m} \rightarrowtail X^{\otimes n}$	$X^{\otimes m} \rightarrow X^{\otimes n}$	With these two equations, it turns out that two morphisms $X^{\otimes m} \rightarrow X^{\otimes n}$ —defined
fong-spivak-seven-sketches_211 F02215	211	■ 0.961	μ, δ	μ, δ	by composing and tensoring identities on X and the noteworthy morphisms μ, δ , etc.—
fong-spivak-seven-sketches_211 F02216	211	■ 0.508	$X, \mu, \eta, \delta, \epsilon$	$X, \mu, \eta, \delta, \epsilon$	Definition 6.54. Let $(X, \mu, \eta, \delta, \epsilon)$ be a Frobenius monoid in a monoidal category
fong-spivak-seven-sketches_211 F02217	211	■ 0.999	$s_{m,n}: X^{\otimes m} \rightarrowtail X^{\otimes n}$	$s_{m,n}: X^{\otimes m} \rightarrow X^{\otimes n}$	$(\mathcal{C}, I, \otimes)$. Let $m, n \in \mathbb{N}$. Define $s_{m,n}: X^{\otimes m} \rightarrow X^{\otimes n}$ to be the following morphism
fong-spivak-seven-sketches_211 F02218	211	■ 0.799	$(m-1) \mu$	$(m-1)\mu$	It can be written formally as $(m-1) \mu$'s followed by $(n-1) \delta$'s, with special cases when
fong-spivak-seven-sketches_211 F02219	211	■ 0.799	$(n-1) \delta$	$(n-1)\delta$	It can be written formally as $(m-1) \mu$'s followed by $(n-1) \delta$'s, with special cases when
fong-spivak-seven-sketches_211 F02220	211	■ 1.000	$m=0$	$m=0$	$m=0$ or $n=0$.
fong-spivak-seven-sketches_211 F02221	211	■ 0.892	$s_{m,n}$	$s_{m,n}$	We call $s_{m,n}$ the spider of type (m, n) , and can draw it more simply as the icon
fong-spivak-seven-sketches_211 F02222	211	■ 1.000	$\mu, \eta, \delta, \epsilon, \sigma$	$\mu, \eta, \delta, \epsilon, \sigma$	talking, if a wire is spiderable, it means that we have a bunch of morphisms $\mu, \eta, \delta, \epsilon, \sigma$
fong-spivak-seven-sketches_211 F02223	211	■ 0.781	$(X, \mu, \eta, \delta, \epsilon)$	$(X, \mu, \eta, \delta, \epsilon)$	Theorem 6.55. Let $(X, \mu, \eta, \delta, \epsilon)$ be a Frobenius monoid in a monoidal category $(\mathcal{C}, I, \otimes)$.
fong-spivak-seven-sketches_211 F02224	211	■ 1.000	$f: X^{\otimes m} \rightarrowtail X^{\otimes n}$	$f: X^{\otimes m} \rightarrow X^{\otimes n}$	Suppose that we have a map $f: X^{\otimes m} \rightarrow X^{\otimes n}$ each constructed from spiders and the
fong-spivak-seven-sketches_211 F02225	211	■ 1.000	$\sigma: X^{\otimes 2} \rightarrowtail X^{\otimes 2}$	$\sigma: X^{\otimes 2} \rightarrow X^{\otimes 2}$	symmetry map $\sigma: X^{\otimes 2} \rightarrow X^{\otimes 2}$ using composition and the monoidal product, and such

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02226	211	1.000	$f=s_{\{m, n\}}$	$f = s_{m,n}$	$f = s_{m,n}$.
fong-spivak-seven-sketches-F02227	211	1.000	$X \otimes X \rightarrow X \otimes X \otimes X$	$X \otimes X \rightarrow X \otimes X \otimes X$	morphisms $X \otimes X \rightarrow X \otimes X \otimes X$ in the following list are necessarily equal?
fong-spivak-seven-sketches-F02228	212	0.937	$G:=\{\mu, \eta, \delta, \epsilon\}$	$G := \{\mu, \eta, \delta, \epsilon\}$	Theorem 6.58. Consider the four-element set $G := \{\mu, \eta, \delta, \epsilon\}$ and define $in, out: G \rightarrow$
fong-spivak-seven-sketches-F02229	212	0.937	$G \rightarrow$	$G \rightarrow$	Theorem 6.58. Consider the four-element set $G := \{\mu, \eta, \delta, \epsilon\}$ and define $in, out: G \rightarrow$
fong-spivak-seven-sketches-F02230	212	0.976	(G, E)	(G, E)	Then the free prop on (G, E) is equivalent, as a symmetric monoidal category, ^a to
fong-spivak-seven-sketches-F02231	213	0.985	$X, \mu_X, \delta_X, \eta_X, \epsilon_X$	$X, \mu_X, \delta_X, \eta_X, \epsilon_X$	which each object X is equipped with a Frobenius structure $(X, \mu_X, \delta_X, \eta_X, \epsilon_X)$ such
fong-spivak-seven-sketches-F02232	214	0.991	$\eta_I=\mathrm{id}_I=\epsilon_I$	$\eta_I = \mathrm{id}_I = \epsilon_I$	for all objects X, Y , and such that $\eta_I = \mathrm{id}_I = \epsilon_I$.
fong-spivak-seven-sketches-F02233	214	0.978	$\mu_X, \delta_X, \eta_X, \epsilon_X$	$\mu_X, \delta_X, \eta_X, \epsilon_X$	Frobenius morphisms $\mu_X, \delta_X, \eta_X, \epsilon_X$ for each object X are constructed using the uni-
fong-spivak-seven-sketches-F02234	214	0.794	$\underline{1}$	$\underline{1}$	ject $\underline{1}$ in $\mathbf{Cosp}_{\mathbf{FinSet}}$ using both the function and wiring depictions as in Example 6.46.
fong-spivak-seven-sketches-F02235	214	0.792	$\varnothing, \sqcup$	$\varnothing, \sqcup$	<i>Example 6.64.</i> Recall the monoidal category $(\mathbf{Corel}, \varnothing, \sqcup)$ from Example 4.61; its objects
fong-spivak-seven-sketches-F02236	214	0.578	$\mu_X: X \sqcup X \rightarrow X$	$\mu_X : X \sqcup X \rightarrow X$	corelation $\mu_X: X \sqcup X \rightarrow X$ such that two elements of $X \sqcup X \sqcup X$ are equivalent if and
fong-spivak-seven-sketches-F02237	214	0.578	$X \sqcup X \sqcup X$	$X \sqcup X \sqcup X$	corelation $\mu_X: X \sqcup X \rightarrow X$ such that two elements of $X \sqcup X \sqcup X$ are equivalent if and
fong-spivak-seven-sketches-F02238	214	1.000	$\delta_X: X \rightarrow X \sqcup X$	$\delta_X : X \rightarrow X \sqcup X$	only if they come from the same underlying element of X . Define $\delta_X: X \rightarrow X \sqcup X$ in
fong-spivak-seven-sketches-F02239	214	1.000	$\eta_X: \varnothing \rightarrow X$	$\eta_X : \varnothing \rightarrow X$	the same way, and define $\eta_X: \varnothing \rightarrow X$ and $\epsilon_X: X \rightarrow \varnothing$ such that no two elements of
fong-spivak-seven-sketches-F02240	214	1.000	$\epsilon_X: X \rightarrow \varnothing$	$\epsilon_X : X \rightarrow \varnothing$	the same way, and define $\eta_X: \varnothing \rightarrow X$ and $\epsilon_X: X \rightarrow \varnothing$ such that no two elements of
fong-spivak-seven-sketches-F02241	214	1.000	$X=\varnothing \sqcup X=X \sqcup \varnothing$	$X = \varnothing \sqcup X = X \sqcup \varnothing$	$X = \varnothing \sqcup X = X \sqcup \varnothing$ are equivalent.
fong-spivak-seven-sketches-F02242	214	1.000	$(X, \mu_X, \eta_X, \delta_X, \epsilon_X)$	$(X, \mu_X, \eta_X, \delta_X, \epsilon_X)$	These maps define a special commutative Frobenius monoid $(X, \mu_X, \eta_X, \delta_X, \epsilon_X)$,
fong-spivak-seven-sketches-F02243	216	0.384	$(\mathcal{C}, I_{\mathcal{C}}, \otimes_{\mathcal{C}})$	$(\mathcal{C}, I_{\mathcal{C}}, \otimes_{\mathcal{C}})$	Rough Definition 6.68. Let $(\mathcal{C}, I_{\mathcal{C}}, \otimes_{\mathcal{C}})$ and $(\mathcal{D}, I_{\mathcal{D}}, \otimes_{\mathcal{D}})$ be symmetric monoidal cate-
fong-spivak-seven-sketches-F02244	216	0.384	$(\mathcal{D}, I_{\mathcal{D}}, \otimes_{\mathcal{D}})$	$(\mathcal{D}, I_{\mathcal{D}}, \otimes_{\mathcal{D}})$	Rough Definition 6.68. Let $(\mathcal{C}, I_{\mathcal{C}}, \otimes_{\mathcal{C}})$ and $(\mathcal{D}, I_{\mathcal{D}}, \otimes_{\mathcal{D}})$ be symmetric monoidal cate-
fong-spivak-seven-sketches-F02245	216	0.994	(F, φ)	(F, φ)	gories. To specify a <i>symmetric monoidal functor</i> (F, φ) between them,
fong-spivak-seven-sketches-F02246	216	0.984	$\varphi_I: I_{\mathcal{D}} \rightarrow F(I_{\mathcal{C}})$	$\varphi_I : I_{\mathcal{D}} \rightarrow F(I_{\mathcal{C}})$	(ii) one specifies a morphism $\varphi_I: I_{\mathcal{D}} \rightarrow F(I_{\mathcal{C}})$.
fong-spivak-seven-sketches-F02247	216	0.899	$c_1, c_2 \in \mathrm{Ob}(\mathcal{C})$	$c_1, c_2 \in \mathrm{Ob}(\mathcal{C})$	(iii) for each $c_1, c_2 \in \mathrm{Ob}(\mathcal{C})$, one specifies a morphism
fong-spivak-seven-sketches-F02248	216	1.000	φ	φ	We call the various maps φ <i>coherence maps</i> . We require the coherence maps to obey
fong-spivak-seven-sketches-F02249	216	1.000	φ_I	φ_I	monoidal structures on $\mathcal{C}$ and $\mathcal{D}$. If φ_I and φ_{c_1, c_2} are isomorphisms for all c_1, c_2 , we say
fong-spivak-seven-sketches-F02250	216	1.000	φ_{c_1, c_2}	φ_{c_1, c_2}	monoidal structures on $\mathcal{C}$ and $\mathcal{D}$. If φ_I and φ_{c_1, c_2} are isomorphisms for all c_1, c_2 , we say
fong-spivak-seven-sketches-F02251	216	1.000	c_1, c_2	c_1, c_2	monoidal structures on $\mathcal{C}$ and $\mathcal{D}$. If φ_I and φ_{c_1, c_2} are isomorphisms for all c_1, c_2 , we say
fong-spivak-seven-sketches-F02252	216	1.000	$S \in$	$S \in$	sending a set $S \in \mathbf{Set}$ to its set of subsets $P(S) := \{R \subseteq S\}$. It acts on morphisms by

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02253	216	■ 1.000	$\mathrm{P}(S) := \{R \subseteq S\}$	$P(S) := \{R \subseteq S\}$	sending a set $S \in \mathbf{Set}$ to its set of subsets $P(S) := \{R \subseteq S\}$. It acts on morphisms by
fong-spivak-seven-sketches-F02254	216	■ 1.000	$\operatorname{im}_f: \mathrm{P}(S) \rightarrow \mathrm{P}(T)$	$\operatorname{im}_f: P(S) \rightarrow P(T)$	sending a function $f: S \rightarrow T$ to the image map $\operatorname{im}_f: P(S) \rightarrow P(T)$, which maps $R \subseteq S$
fong-spivak-seven-sketches-F02255	216	■ 1.000	$R \subseteq S$	$R \subseteq S$	sending a function $f: S \rightarrow T$ to the image map $\operatorname{im}_f: P(S) \rightarrow P(T)$, which maps $R \subseteq S$
fong-spivak-seven-sketches-F02256	216	■ 1.000	$\{f(r) \mid r \in R\} \subseteq T$	$\{f(r) \mid r \in R\} \subseteq T$	to $\{f(r) \mid r \in R\} \subseteq T$.
fong-spivak-seven-sketches-F02257	216	■ 1.000	$(\{1\}, \times)$	$(\{1\}, \times)$	Now consider the symmetric monoidal structure $(\{1\}, \times)$ on $\mathbf{Set}$ from Example 4.49.
fong-spivak-seven-sketches-F02258	216	■ 1.000	$\varphi_I: \{1\} \rightarrow$	$\varphi_I: \{1\} \rightarrow$	To make P a symmetric monoidal functor, we need to specify a function $\varphi_I: \{1\} \rightarrow$
fong-spivak-seven-sketches-F02259	216	■ 0.997	$\mathrm{P}(\{1\})$	$P(\{1\})$	$P(\{1\})$ and for all sets S and T , a functor $\varphi_{S,T}: P(S) \times P(T) \rightarrow P(S \times T)$. One possibility is
fong-spivak-seven-sketches-F02260	216	■ 0.997	$\varphi_{S,T}: \mathrm{P}(S) \times \mathrm{P}(T) \rightarrow \mathrm{P}(S \times T)$	$\varphi_{S,T}: P(S) \times P(T) \rightarrow P(S \times T)$	$P(\{1\})$ and for all sets S and T , a functor $\varphi_{S,T}: P(S) \times P(T) \rightarrow P(S \times T)$. One possibility is
fong-spivak-seven-sketches-F02261	216	■ 1.000	$\varphi_I(1)$	$\varphi_I(1)$	to define $\varphi_I(1)$ to be the maximal subset $\{1\} \subseteq \{1\}$, and given subsets $A \subseteq S$ and $B \subseteq T$,
fong-spivak-seven-sketches-F02262	216	■ 1.000	$\{1\} \subseteq \{1\}$	$\{1\} \subseteq \{1\}$	to define $\varphi_I(1)$ to be the maximal subset $\{1\} \subseteq \{1\}$, and given subsets $A \subseteq S$ and $B \subseteq T$,
fong-spivak-seven-sketches-F02263	216	■ 1.000	$A \subseteq S$	$A \subseteq S$	to define $\varphi_I(1)$ to be the maximal subset $\{1\} \subseteq \{1\}$, and given subsets $A \subseteq S$ and $B \subseteq T$,
fong-spivak-seven-sketches-F02264	216	■ 1.000	$B \subseteq T$	$B \subseteq T$	to define $\varphi_I(1)$ to be the maximal subset $\{1\} \subseteq \{1\}$, and given subsets $A \subseteq S$ and $B \subseteq T$,
fong-spivak-seven-sketches-F02265	216	■ 1.000	$\varphi_{S,T}(A, B)$	$\varphi_{S,T}(A, B)$	to define $\varphi_{S,T}(A, B)$ to be the product subset $A \times B \subseteq S \times T$. With these definitions,
fong-spivak-seven-sketches-F02266	216	■ 1.000	$A \times B \subseteq S \times T$	$A \times B \subseteq S \times T$	to define $\varphi_{S,T}(A, B)$ to be the product subset $A \times B \subseteq S \times T$. With these definitions,
fong-spivak-seven-sketches-F02267	216	■ 0.957	(P, φ)	(P, φ)	(P, φ) is a symmetric monoidal functor.
fong-spivak-seven-sketches-F02268	216	■ 1.000	$\varphi_{S,T}$	$\varphi_{S,T}$	Exercise 6.70. Check that the maps $\varphi_{S,T}$ defined in Example 6.69 are natural in S and
fong-spivak-seven-sketches-F02269	217	■ 1.000	2Ω	2Ω	as a resistor of resistance 2Ω). For this circuit to be a morphism in some category, i.e.
fong-spivak-seven-sketches-F02270	217	■ 1.000	$A \rightarrow N$	$A \rightarrow N$	of the functions $A \rightarrow N$ and $B \rightarrow N$ are indicated by the grey arrows. This forms a
fong-spivak-seven-sketches-F02271	217	■ 1.000	$B \rightarrow N$	$B \rightarrow N$	of the functions $A \rightarrow N$ and $B \rightarrow N$ are indicated by the grey arrows. This forms a
fong-spivak-seven-sketches-F02272	218	■ 1.000	$(F, \varphi): (\mathcal{C}, +) \rightarrow$	$(F, \varphi): (\mathcal{C}, +) \rightarrow$	Definition 6.75. Let $\mathcal{C}$ be a category with finite colimits, and $(F, \varphi): (\mathcal{C}, +) \rightarrow (\mathbf{Set}, \times)$
fong-spivak-seven-sketches-F02273	218	■ 1.000	$\times)$	$\times)$	Definition 6.75. Let $\mathcal{C}$ be a category with finite colimits, and $(F, \varphi): (\mathcal{C}, +) \rightarrow (\mathbf{Set}, \times)$
fong-spivak-seven-sketches-F02274	218	■ 0.672	$A \xrightarrow{i} N \stackrel{o}{\leftarrow} B$	$A \xrightarrow{i} N \stackrel{o}{\leftarrow} B$	$A \xrightarrow{i} N \stackrel{o}{\leftarrow} B$ in $\mathcal{C}$ together with an element $s \in F(N)$. ⁵ We call (F, φ) the <i>decoration</i>
fong-spivak-seven-sketches-F02275	218	■ 0.672	$s \in F(N)$.	$s \in F(N)$. ⁵	$A \xrightarrow{i} N \stackrel{o}{\leftarrow} B$ in $\mathcal{C}$ together with an element $s \in F(N)$. ⁵ We call (F, φ) the <i>decoration</i>
fong-spivak-seven-sketches-F02276	218	■ 0.596	$\mathcal{C} =$	$\mathcal{C} =$	The intuition here is to use $\mathcal{C} = \mathbf{FinSet}$, and, for each object $N \in \mathbf{FinSet}$, the functor
fong-spivak-seven-sketches-F02277	218	■ 0.596	$N \in$	$N \in$	The intuition here is to use $\mathcal{C} = \mathbf{FinSet}$, and, for each object $N \in \mathbf{FinSet}$, the functor
fong-spivak-seven-sketches-F02278	218	■ 1.000	$F(N)$	$F(N)$	available decorations on N nodes, taken from the set $F(N)$.
fong-spivak-seven-sketches-F02279	218	■ 0.980	$(A \xrightarrow{f} N \stackrel{g}{\leftarrow} B, s)$	$(A \xrightarrow{f} N \stackrel{g}{\leftarrow} B, s)$	Let $(A \xrightarrow{f} N \stackrel{g}{\leftarrow} B, s)$ and $(B \xrightarrow{h} P \stackrel{k}{\leftarrow} C, t)$ represent decorated cospans. Their

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02280	218	0.980	$(B \xrightarrow{h} P \stackrel{k}{\leftarrow} C, t)$	$(B \xrightarrow{h} P \xleftarrow{k} C, t)$	Let $(A \xrightarrow{f} N \xleftarrow{g} B, s)$ and $(B \xrightarrow{h} P \xleftarrow{k} C, t)$ represent decorated cospans. Their
fong-spivak-seven-sketches-F02281	218	1.000	$B \xrightarrow{h} P \stackrel{k}{\leftarrow} C$	$B \xrightarrow{h} P \xleftarrow{k} C$	composite is represented by the composite of the cospans $A \xrightarrow{f} N \xleftarrow{g} B$ and $B \xrightarrow{h} P \xleftarrow{k} C$,
fong-spivak-seven-sketches-F02282	219	0.998	$F \left(N +_{\{B\}} P \right)$	$F(N +_B P)$	paired with the following element of $F(N +_B P)$:
fong-spivak-seven-sketches-F02283	219	0.875	$_{\{F\}}$	F	tor $(F, \varphi): (\mathcal{C}, +) \longrightarrow (\mathbf{Set}, \times)$, there is a hypergraph category $\mathbf{Cospan}_F$ whose objects
fong-spivak-seven-sketches-F02284	219	0.581	$_{\{\text{C}\}}$	C	$\mathbf{Cospan}_C$.
fong-spivak-seven-sketches-F02285	219	0.876	$_{\{\mathrm{e}\}}$	e	<i>Exercise 6.78.</i> Suppose you’re worried that the notation $\mathbf{Cospan}_e$ looks like the
fong-spivak-seven-sketches-F02286	219	0.579	$\operatorname{Cospan}_{\{F\}}$	$\mathbf{Cospan}_F$	notation $\mathbf{Cospan}_F$, even though they’re very different. An expert tells you “they’re not
fong-spivak-seven-sketches-F02287	219	1.000	$F(c) := \{*\}$	$F(c) := \{*\}$	so different; one is a special case of the other. Just use the constant functor $F(c) := \{*\}$.”
fong-spivak-seven-sketches-F02288	219	1.000	$\mathbb{R}^{+}$	$\mathbb{R}^+$	$\mathbb{R}^+$. But we write C this way to remind us that it consists of circuit components. If we
fong-spivak-seven-sketches-F02289	219	1.000	$\ell: A \rightarrow C$	$\ell: A \rightarrow C$	source and target functions, together with a function $\ell: A \rightarrow C$ labeling each edge with
fong-spivak-seven-sketches-F02290	219	1.000	$V = \{1, 2\}, A = \{e\}, s(e) = 1$	$V = \{1, 2\}, A = \{e\}, s(e) = 1$	For example, we might have the simple case of $V = \{1, 2\}, A = \{e\}, s(e) = 1$,
fong-spivak-seven-sketches-F02291	219	0.562	$t(e) = 2$	$t(e) = 2$	$t(e) = 2$ —so e is an edge from 1 to 2—and $\ell(e) = 3\Omega$. This represents a resistor with
fong-spivak-seven-sketches-F02292	219	0.562	$\ell(e) = 3 \Omega$	$\ell(e) = 3\Omega$	$t(e) = 2$ —so e is an edge from 1 to 2—and $\ell(e) = 3\Omega$. This represents a resistor with
fong-spivak-seven-sketches-F02293	219	1.000	3Ω	3Ω	resistance 3Ω :
fong-spivak-seven-sketches-F02294	220	1.000	$V = \{1, 2\}, A = \{e\}$	$V = \{1, 2\}, A = \{e\}$	resistor could also be represented by the ‘reversed graph’, with data $V = \{1, 2\}, A = \{e\}$,
fong-spivak-seven-sketches-F02295	220	1.000	$s(e) = 2, t(e) = 1$	$s(e) = 2, t(e) = 1$	$s(e) = 2, t(e) = 1$, and $\ell(e) = 3F$.
fong-spivak-seven-sketches-F02296	220	1.000	$\ell(e) = 3 F$	$\ell(e) = 3F$	$s(e) = 2, t(e) = 1$, and $\ell(e) = 3F$.
fong-spivak-seven-sketches-F02297	220	0.983	V, A, s, t, ℓ	V, A, s, t, ℓ	<i>Exercise 6.79.</i> Write a tuple (V, A, s, t, ℓ) that represents the circuit in Eq. (6.71).
fong-spivak-seven-sketches-F02298	220	0.951	ψ	ψ	$(\mathbf{Circ}, \psi)$. We start by defining the functor part
fong-spivak-seven-sketches-F02299	220	1.000	$f: V \rightarrow V^{\prime}$	$f: V \rightarrow V'$	On morphisms, $\mathbf{Circ}$ sends a function $f: V \rightarrow V'$ to the function
fong-spivak-seven-sketches-F02300	220	0.693	$c \in \operatorname{Circ}(\underline{4})$	$c \in \mathbf{Circ}(\underline{4})$	<i>Exercise 6.80.</i> To understand this functor better, let $c \in \mathbf{Circ}(\underline{4})$ be the circuit
fong-spivak-seven-sketches-F02301	220	1.000	$f: \underline{4} \rightarrow \underline{3}$	$f: \underline{4} \rightarrow \underline{3}$	and let $f: \underline{4} \rightarrow \underline{3}$ be the function
fong-spivak-seven-sketches-F02302	220	1.000	$\operatorname{Circ}(f)(c)$	$\mathbf{Circ}(f)(c)$	Draw a picture of the circuit $\mathbf{Circ}(f)(c)$.
fong-spivak-seven-sketches-F02303	220	1.000	$\psi_{V, V^{\prime}}$	$\psi_{V, V'}$	coherence maps $\psi_{V, V'}$ for finite sets V, V' , we define
fong-spivak-seven-sketches-F02304	220	1.000	$V, V^{\prime}$	V, V'	coherence maps $\psi_{V, V'}$ for finite sets V, V' , we define
fong-spivak-seven-sketches-F02305	220	1.000	$V + V^{\prime}$	$V + V'$	considers them together as a circuit on the disjoint union of vertices $V + V'$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02306	220	■ 1.000	$\psi_{\{\underline{2}\}, \underline{2}}(b, s) \in \operatorname{Circ}(\underline{2} + \underline{2}) = \operatorname{Circ}(\underline{4})$	$\psi_{\underline{2}, \underline{2}}(b, s) \in \operatorname{Circ}(\underline{2} + \underline{2}) = \operatorname{Circ}(\underline{4})$	$\psi_{\underline{2}, \underline{2}}(b, s) \in \operatorname{Circ}(\underline{2} + \underline{2}) = \operatorname{Circ}(\underline{4})$ should be, and draw a picture.
fong-spivak-seven-sketches-F02307	221	■ 0.339	(	(	$(\operatorname{Circ}, \psi): (\mathbf{FinSet}, +) \rightarrow (\mathbf{Set}, \times)$, we can construct our promised hypergraph category
fong-spivak-seven-sketches-F02308	221	■ 0.339	$\psi): ($	$\psi): ($	$(\operatorname{Circ}, \psi): (\mathbf{FinSet}, +) \rightarrow (\mathbf{Set}, \times)$, we can construct our promised hypergraph category
fong-spivak-seven-sketches-F02309	221	■ 0.339	) \rightarrow (	) \rightarrow (	$(\operatorname{Circ}, \psi): (\mathbf{FinSet}, +) \rightarrow (\mathbf{Set}, \times)$, we can construct our promised hypergraph category
fong-spivak-seven-sketches-F02310	221	■ 0.945	$\{ \}_{\text{Circ}}$	Circ	Our notation for this category is $\mathbf{Cospan}_{\operatorname{Circ}}$. Following Theorem 6.77, the objects
fong-spivak-seven-sketches-F02311	221	■ 0.147	$\underline{2}$	$\underline{2}$	collections of white circles $\circ$. For example, we'll represent the object $\underline{2}$ of $\mathbf{Cospan}_{\operatorname{Circ}}$ as
fong-spivak-seven-sketches-F02312	221	■ 0.147	$\{ \}_{\text{Circ}}$	Circ	collections of white circles $\circ$. For example, we'll represent the object $\underline{2}$ of $\mathbf{Cospan}_{\operatorname{Circ}}$ as
fong-spivak-seven-sketches-F02313	221	■ 0.999	$\underline{m} \rightarrow \underline{p} \leftarrow \underline{n}$	$\underline{m} \rightarrow \underline{p} \leftarrow \underline{n}$	that is, $\operatorname{cospan} \underline{m} \rightarrow \underline{p} \leftarrow \underline{n}$ together with an element c of $\operatorname{Circ}(\underline{p})$. As an example,
fong-spivak-seven-sketches-F02314	221	■ 0.999	$\operatorname{Circ}(\underline{p})$	$\operatorname{Circ}(\underline{p})$	that is, $\operatorname{cospan} \underline{m} \rightarrow \underline{p} \leftarrow \underline{n}$ together with an element c of $\operatorname{Circ}(\underline{p})$. As an example,
fong-spivak-seven-sketches-F02315	221	■ 1.000	$\underline{1} \xrightarrow{i_1} \underline{2} \xleftarrow{i_2} \underline{1}$	$\underline{1} \xrightarrow{i_1} \underline{2} \xleftarrow{i_2} \underline{1}$	consider the $\operatorname{cospan} \underline{1} \xrightarrow{i_1} \underline{2} \xleftarrow{i_2} \underline{1}$ where $i_1(1) = 1$ and $i_2(1) = 2$, equipped with the
fong-spivak-seven-sketches-F02316	221	■ 1.000	$i_1(1)=1$	$i_1(1) = 1$	consider the $\operatorname{cospan} \underline{1} \xrightarrow{i_1} \underline{2} \xleftarrow{i_2} \underline{1}$ where $i_1(1) = 1$ and $i_2(1) = 2$, equipped with the
fong-spivak-seven-sketches-F02317	221	■ 1.000	$i_2(1)=2$	$i_2(1) = 2$	consider the $\operatorname{cospan} \underline{1} \xrightarrow{i_1} \underline{2} \xleftarrow{i_2} \underline{1}$ where $i_1(1) = 1$ and $i_2(1) = 2$, equipped with the
fong-spivak-seven-sketches-F02318	221	■ 0.760	$(V, A, s, t, \ell) \in \operatorname{Circ}(\underline{2})$	$(V, A, s, t, \ell) \in \operatorname{Circ}(\underline{2})$	some C-circuit $(V, A, s, t, \ell) \in \operatorname{Circ}(\underline{2})$. What is it?
fong-spivak-seven-sketches-F02319	221	■ 0.578	$\underline{1} \rightarrow \underline{1}$	$\underline{1} \rightarrow \underline{1}$	Since this and the circuit in (6.83) are both morphisms $\underline{1} \rightarrow \underline{1}$, we may compose them
fong-spivak-seven-sketches-F02320	221	■ 1.000	$\operatorname{Circ}([\iota_N, \iota_P])(\psi_{N,P}(s, t))$	$\operatorname{Circ}([\iota_N, \iota_P])(\psi_{N,P}(s, t))$	new decoration, we use the formula $\operatorname{Circ}([\iota_N, \iota_P])(\psi_{N,P}(s, t))$ from (6.76). Again, this is
fong-spivak-seven-sketches-F02321	222	■ 0.962	$\underline{1} = \{\circ\}$	$\underline{1} = \{\circ\}$	(We simply ignore the decorations for now.) If we pushout over the common set $\underline{1} = \{\circ\}$,
fong-spivak-seven-sketches-F02322	222	■ 0.307	$\psi_{\{2, \underline{2}\}}$	$\psi_{\underline{2}, \underline{2}}$	new decoration. You told us what the map $\psi_{\underline{2}, \underline{2}}$ does in Exercise 6.82. It takes the two
fong-spivak-seven-sketches-F02323	222	■ 0.834	$\operatorname{Circ}(\underline{4})$	$\operatorname{Circ}(\underline{4})$	in $\operatorname{Circ}(\underline{4})$. So this is what the $\psi_{N,P}(s, t)$ part means. What does the $[\iota_N, \iota_P]$ mean? Recall
fong-spivak-seven-sketches-F02324	222	■ 0.834	$\psi_{N,P}(s, t)$	$\psi_{N,P}(s, t)$	in $\operatorname{Circ}(\underline{4})$. So this is what the $\psi_{N,P}(s, t)$ part means. What does the $[\iota_N, \iota_P]$ mean? Recall
fong-spivak-seven-sketches-F02325	222	■ 0.834	$[\iota_N, \iota_P]$	$[\iota_N, \iota_P]$	in $\operatorname{Circ}(\underline{4})$. So this is what the $\psi_{N,P}(s, t)$ part means. What does the $[\iota_N, \iota_P]$ mean? Recall
fong-spivak-seven-sketches-F02326	222	■ 0.963	$\operatorname{Cospan}_{\operatorname{Circ}}$	$\operatorname{Cospan}_{\operatorname{Circ}}$	diagram as morphisms in $\mathbf{Cospan}_{\operatorname{Circ}}$, and compute their composite. Does it match the
fong-spivak-seven-sketches-F02327	222	■ 0.930	$A \rightarrow M \leftarrow B$	$A \rightarrow M \leftarrow B$	we again just work in $\mathbf{Cospan}_{\mathbf{FinSet}}$: given two cospans $A \rightarrow M \leftarrow B$ and $C \rightarrow N \leftarrow D$,
fong-spivak-seven-sketches-F02328	222	■ 0.930	$C \rightarrow N \leftarrow D$	$C \rightarrow N \leftarrow D$	we again just work in $\mathbf{Cospan}_{\mathbf{FinSet}}$: given two cospans $A \rightarrow M \leftarrow B$ and $C \rightarrow N \leftarrow D$,
fong-spivak-seven-sketches-F02329	222	■ 1.000	$A + C \rightarrow M + N \leftarrow B + D$	$A + C \rightarrow M + N \leftarrow B + D$	we take their coproduct $\operatorname{cospan} \underline{A + C \rightarrow M + N \leftarrow B + D}$. And for decorations, we
fong-spivak-seven-sketches-F02330	222	■ 0.997	$\psi_{M,N}: \operatorname{Circ}(M) \times \operatorname{Circ}(N) \rightarrow \operatorname{Circ}(M+N)$	$\psi_{M,N}: \operatorname{Circ}(M) \times \operatorname{Circ}(N) \rightarrow \operatorname{Circ}(M+N)$	use the map $\psi_{M,N}: \operatorname{Circ}(M) \times \operatorname{Circ}(N) \rightarrow \operatorname{Circ}(M+N)$. So, for example, suppose we
fong-spivak-seven-sketches-F02331	223	■ 1.000	$\eta: 0 \rightarrow 2$	$\eta: 0 \rightarrow 2$	Exercise 6.88. Write x for the open circuit in (6.87). Also define cospans $\eta: 0 \rightarrow 2$ and

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches- F02332	223	■ 0.915	$\eta: 2 \rightarrow 0$	$\eta: 2 \rightarrow 0$	$\eta: 2 \rightarrow 0$ as follows:
fong-spivak-seven-sketches- F02333	223	■ 0.967	$(\underline{1}, \varnothing, !, !, !) \in \operatorname{Circ}(\underline{1})^6$	$(\underline{1}, \varnothing, !, !, !) \in \operatorname{Circ}(\underline{1})^6$	where each of these are decorated by the empty circuit $(\underline{1}, \varnothing, !, !, !) \in \operatorname{Circ}(\underline{1})^6$
fong-spivak-seven-sketches- F02334	223	■ 0.782	$\eta; x; \epsilon$	$\eta; x; \epsilon$	Compute the composite $\eta \circ x \circ \epsilon$ in $\mathbf{Cospan}_{\operatorname{Circ}}$. This is a morphism $\underline{0} \rightarrow \underline{0}$; we call
fong-spivak-seven-sketches- F02335	223	■ 0.782	$\underline{0} \rightarrow \underline{0}$	$\underline{0} \rightarrow \underline{0}$	Compute the composite $\eta \circ x \circ \epsilon$ in $\mathbf{Cospan}_{\operatorname{Circ}}$. This is a morphism $\underline{0} \rightarrow \underline{0}$; we call
fong-spivak-seven-sketches- F02336	223	■ 0.574	$\mathcal{O}$	$\mathcal{O}$	structures from operad functors $\mathcal{O} \rightarrow \mathbf{Set}$.
fong-spivak-seven-sketches- F02337	225	■ 1.000	f, g, h, k	f, g, h, k	The cost is that the ‘boxes’ f, g, h, k in Eq. (6.90) no longer have a left/right dis-
fong-spivak-seven-sketches- F02338	225	■ 0.997	$\mathcal{M} \rightarrow$	$\mathcal{M} \rightarrow$	called the theory of monoids as a prop $\mathcal{M}$, and define monoids using functors $\mathcal{M} \rightarrow \mathbf{Set}$;
fong-spivak-seven-sketches- F02339	225	■ 0.992	$\text{FinSet} \rightarrow$	$\text{FinSet} \rightarrow$	$\mathbf{Cospan}_{\text{FinSet}} \rightarrow \mathbf{Set}$.
fong-spivak-seven-sketches- F02340	225	■ 0.725	$t_1, \dots, t_n, t$	$t_1, \dots, t_n, t$	(ii) for each tuple $(t_1, \dots, t_n, t)$ of types, one specifies a set $\mathcal{O}(t_1, \dots, t_n; t)$, whose
fong-spivak-seven-sketches- F02341	225	■ 0.725	$\mathcal{O}(t_1, \dots, t_n; t)$	$\mathcal{O}(t_1, \dots, t_n; t)$	(ii) for each tuple $(t_1, \dots, t_n, t)$ of types, one specifies a set $\mathcal{O}(t_1, \dots, t_n; t)$, whose
fong-spivak-seven-sketches- F02342	225	■ 0.623	$(t_1, \dots, t_n; t)$	$(t_1, \dots, t_n; t)$	elements are called <i>operations of arity</i> $(t_1, \dots, t_n; t)$;
fong-spivak-seven-sketches- F02343	225	■ 0.988	$(s_1, \dots, s_m, t_i)$	$(s_1, \dots, s_m, t_i)$	(iii) for each pair of tuples $(s_1, \dots, s_m, t_i)$ and $(t_1, \dots, t_n, t)$, one specifies a function
fong-spivak-seven-sketches- F02344	225	■ 0.988	$(t_1, \dots, t_n, t)$	$(t_1, \dots, t_n, t)$	(iii) for each pair of tuples $(s_1, \dots, s_m, t_i)$ and $(t_1, \dots, t_n, t)$, one specifies a function
fong-spivak-seven-sketches- F02345	225	■ 0.739	$\operatorname{id}_t \in \mathcal{O}(t; t)$	$\operatorname{id}_t \in \mathcal{O}(t; t)$	(iv) for each type t , one specifies an operation $\operatorname{id}_t \in \mathcal{O}(t; t)$ called the <i>identity operation</i> .
fong-spivak-seven-sketches- F02346	225	■ 1.000	$m+n-1$	$m+n-1$	operation of arity $m+n-1$: we have m arguments to fill in m , and the remaining $n-1$
fong-spivak-seven-sketches- F02347	225	■ 1.000	$n-1$	$n-1$	operation of arity $m+n-1$: we have m arguments to fill in m , and the remaining $n-1$
fong-spivak-seven-sketches- F02348	226	■ 0.943	$\circ_i$	$\circ_i$	the substitution function $\circ_i$, which says we obtain the operation $f \circ_i g \in \mathcal{O}(m+n-1)$.
fong-spivak-seven-sketches- F02349	226	■ 0.943	$f \circ_i g \in \mathcal{O}(m+n-1)$	$f \circ_i g \in \mathcal{O}(m+n-1)$	the substitution function $\circ_i$, which says we obtain the operation $f \circ_i g \in \mathcal{O}(m+n-1)$.
fong-spivak-seven-sketches- F02350	226	■ 0.835	$X_1 \times \dots \times X_n \rightarrow Y$	$X_1 \times \dots \times X_n \rightarrow Y$	(ii) Functions $X_1 \times \dots \times X_n \rightarrow Y$ as operations of arity $(X_1, \dots, X_n; Y)$.
fong-spivak-seven-sketches- F02351	226	■ 0.835	$X_1, \dots, X_n; Y$	$X_1, \dots, X_n; Y$	(ii) Functions $X_1 \times \dots \times X_n \rightarrow Y$ as operations of arity $(X_1, \dots, X_n; Y)$.
fong-spivak-seven-sketches- F02352	227	■ 0.985	$f \in \operatorname{Set}(W_1, \dots, W_m; X_i), g \in \operatorname{Set}(X_1, \dots, X_n; Y)$	$f \in \operatorname{Set}(W_1, \dots, W_m; X_i), g \in \operatorname{Set}(X_1, \dots, X_n; Y)$	where $f \in \operatorname{Set}(W_1, \dots, W_m; X_i), g \in \operatorname{Set}(X_1, \dots, X_n; Y)$, and hence $g \circ_i f$ is a
fong-spivak-seven-sketches- F02353	227	■ 0.985	$g \circ_i f$	$g \circ_i f$	where $f \in \operatorname{Set}(W_1, \dots, W_m; X_i), g \in \operatorname{Set}(X_1, \dots, X_n; Y)$, and hence $g \circ_i f$ is a
fong-spivak-seven-sketches- F02354	227	■ 0.994	$\operatorname{id}_X \in \operatorname{Set}(X; X)$	$\operatorname{id}_X \in \operatorname{Set}(X; X)$	(iv) Identities $\operatorname{id}_X \in \operatorname{Set}(X; X)$ are given by the identity function $\operatorname{id}_X: X \rightarrow X$.
fong-spivak-seven-sketches- F02355	227	■ 0.994	$\operatorname{id}_X: X \rightarrow X$	$\operatorname{id}_X: X \rightarrow X$	(iv) Identities $\operatorname{id}_X \in \operatorname{Set}(X; X)$ are given by the identity function $\operatorname{id}_X: X \rightarrow X$.
fong-spivak-seven-sketches- F02356	227	■ 1.000	$a \in \mathbb{N}$	$a \in \mathbb{N}$	(i) Natural numbers $a \in \mathbb{N}$ as types.
fong-spivak-seven-sketches- F02357	227	■ 0.998	$\underline{a_1} + \dots + \underline{a_n} \rightarrow \underline{p} \leftarrow \underline{b}$	$\underline{a_1} + \dots + \underline{a_n} \rightarrow \underline{p} \leftarrow \underline{b}$	(ii) Cospans $\underline{a_1} + \dots + \underline{a_n} \rightarrow \underline{p} \leftarrow \underline{b}$ of finite sets as operations of arity $(a_1, \dots, a_n; b)$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches- F02358	227	0.998	$\backslash\mathrm{left}(a_{\{1\}}, \backslash\mathrm{ldots}, a_{\{n\}} ; b\backslash\mathrm{right})$	$(a_1, \dots, a_n; b)$	(ii) Cospans $\underline{a_1} + \dots + \underline{a_n} \rightarrow \underline{p} \leftarrow \underline{b}$ of finite sets as operations of arity $(a_1, \dots, a_n; b)$.
fong-spivak-seven-sketches- F02359	227	0.180	$\backslash\mathrm{operatorname{id}}_{\{a\}} \backslash\mathrm{in} \backslash\mathrm{operatorname{Set}}(a ; a)$	$\mathrm{id}_a \in \mathrm{Set}(a; a)$	(iv) Identities $\mathrm{id}_a \in \mathbf{Set}(a; a)$ just given by the identity $\mathrm{cospan} \underline{a} \xrightarrow{\mathrm{id}_a} \underline{a} \xleftarrow{\mathrm{id}_a} \underline{a}$.
fong-spivak-seven-sketches- F02360	227	0.180	$\backslash\mathrm{operatorname{cospan}} \backslash\mathrm{underline}{a} \backslash\mathrm{xrightarrow}{\backslash\mathrm{mathrm{id}}_{\backslash\mathrm{underline}{a}}} \backslash\mathrm{underline}{a} \backslash\mathrm{stackrel}{\backslash\mathrm{mathrm{id}}_{\backslash\mathrm{underline}{a}}}{\backslash\mathrm{text} \{ a \}} \backslash\mathrm{underline}{a}$	$\mathrm{cospan} \underline{a} \xrightarrow{\mathrm{id}_a} \underline{a} \xleftarrow{\mathrm{id}_a} \underline{a}$	(iv) Identities $\mathrm{id}_a \in \mathbf{Set}(a; a)$ just given by the identity $\mathrm{cospan} \underline{a} \xrightarrow{\mathrm{id}_a} \underline{a} \xleftarrow{\mathrm{id}_a} \underline{a}$.
fong-spivak-seven-sketches- F02361	227	0.894	$\backslash\mathrm{text} \{ \mathrm{FinSet} \}, 0, +$	$\mathrm{FinSet}, 0, +$	This is the operadic analogue of the monoidal category $(\mathbf{Cospan}_{\mathrm{FinSet}}, 0, +)$.
fong-spivak-seven-sketches- F02362	227	0.844	$\backslash\mathrm{underline}{3}, \backslash\mathrm{underline}{3}, \backslash\mathrm{underline}{4}, \backslash\mathrm{underline}{2} ; \backslash\mathrm{underline}{3}$	$\underline{3}, \underline{3}, \underline{4}, \underline{2}; \underline{3}$	This is an operation of arity $(\underline{3}, \underline{3}, \underline{4}, \underline{2}; \underline{3})$. Why? The circles marked f and g have 3
fong-spivak-seven-sketches- F02363	227	0.809	$3, 3, 4, 2$	$3, 3, 4, 2$	ports, h has 4 ports, k has 2 ports, and the outer circle has 3 ports: $3, 3, 4, 2; 3$.
fong-spivak-seven-sketches- F02364	227	1.000	$\backslash\mathrm{underline}{3} + \backslash\mathrm{underline}{3} + \backslash\mathrm{underline}{4} + \backslash\mathrm{underline}{2} \backslash\mathrm{xrightarrow}{a} \backslash\mathrm{underline}{p} \backslash\mathrm{stackrel}{b}{\backslash\mathrm{leftarrow}} \backslash\mathrm{underline}{3}$	$\underline{3} + \underline{3} + \underline{4} + \underline{2} \xrightarrow{a} \underline{p} \xleftarrow{b} \underline{3}$	arity is, by (ii), a $\mathrm{cospan} \underline{3} + \underline{3} + \underline{4} + \underline{2} \xrightarrow{a} \underline{p} \xleftarrow{b} \underline{3}$. In the diagram above, the apex $\underline{p}$ is the
fong-spivak-seven-sketches- F02365	227	1.000	$\backslash\mathrm{underline}{p}$	$\underline{p}$	arity is, by (ii), a $\mathrm{cospan} \underline{3} + \underline{3} + \underline{4} + \underline{2} \xrightarrow{a} \underline{p} \xleftarrow{b} \underline{3}$. In the diagram above, the apex $\underline{p}$ is the
fong-spivak-seven-sketches- F02366	227	0.590	$\backslash\mathrm{underline}{7}$	$\underline{7}$	set $\underline{Z}$, because there are 7 nodes $\bullet$ in the diagram. The function a sends each port on
fong-spivak-seven-sketches- F02367	228	0.986	$f \backslash\mathrm{in} \backslash\mathrm{operatorname{Cospan}}(2, 2 ; 2)$	$f \in \mathrm{Cospan}(2, 2; 2)$	1. Consider the following cospan $f \in \mathbf{Cospan}(2, 2; 2)$:
fong-spivak-seven-sketches- F02368	228	1.000	$g \backslash\mathrm{in} \backslash\mathrm{operatorname{Cospan}}(2, 2, 2 ; 0)$	$g \in \mathrm{Cospan}(2, 2, 2; 0)$	2. Draw the wiring diagram corresponding to the following cospan $g \in \mathbf{Cospan}(2, 2, 2; 0)$:
fong-spivak-seven-sketches- F02369	228	1.000	$g \backslash\mathrm{circ}_{\{1\}} f$	$g \circ_1 f$	3. Compute the cospan $g \circ_1 f$. What is its arity?
fong-spivak-seven-sketches- F02370	228	0.461	$\backslash\mathrm{mathcal}{0}_{\backslash\mathrm{mathcal}{C}}$	$\mathcal{O}_{\mathcal{C}}$	Definition 6.97. For any symmetric monoidal category $(\mathcal{C}, I, \otimes)$, there is an operad $\mathcal{O}_{\mathcal{C}}$,
fong-spivak-seven-sketches- F02371	228	0.535	$C_{\{1\}} \backslash\mathrm{otimes} \backslash\mathrm{cdots} \backslash\mathrm{otimes} C_{\{n\}} \backslash\mathrm{rightarrow} D$	$C_1 \otimes \dots \otimes C_n \rightarrow D$	(ii) morphisms $C_1 \otimes \dots \otimes C_n \rightarrow D$ in $\mathcal{C}$ as the operations of arity $(C_1, \dots, C_n; D)$.
fong-spivak-seven-sketches- F02372	228	0.535	$C_{\{1\}}, \backslash\mathrm{ldots}, C_{\{n\}} ; D$	$C_1, \dots, C_n; D$	(ii) morphisms $C_1 \otimes \dots \otimes C_n \rightarrow D$ in $\mathcal{C}$ as the operations of arity $(C_1, \dots, C_n; D)$.
fong-spivak-seven-sketches- F02373	228	0.639	$\backslash\mathrm{operatorname{id}}_{\{a\}} \backslash\mathrm{in} \backslash\mathrm{mathcal}{0}_{\backslash\mathrm{mathcal}{C}}(a ; a)$	$\mathrm{id}_a \in \mathcal{O}_{\mathcal{C}}(a; a)$	(iv) identities $\mathrm{id}_a \in \mathcal{O}_{\mathcal{C}}(a; a)$ defined by id_a .
fong-spivak-seven-sketches- F02374	228	0.639	$\backslash\mathrm{operatorname{id}}_{\{a\}}$	id_a	(iv) identities $\mathrm{id}_a \in \mathcal{O}_{\mathcal{C}}(a; a)$ defined by id_a .
fong-spivak-seven-sketches- F02375	229	1.000	$F: \backslash\mathrm{mathcal}{0} \backslash\mathrm{rightarrow} \backslash\mathrm{mathcal}{P}$	$F: \mathcal{O} \rightarrow \mathcal{P}$	and U respectively. To specify an operad functor $F: \mathcal{O} \rightarrow \mathcal{P}$,
fong-spivak-seven-sketches- F02376	229	1.000	$f: T \backslash\mathrm{rightarrow} U$	$f: T \rightarrow U$	(i) one specifies a function $f: T \rightarrow U$.
fong-spivak-seven-sketches- F02377	229	1.000	$F: \backslash\mathrm{mathcal}{0} \backslash\mathrm{rightarrow}$	$F: \mathcal{O} \rightarrow$	Definition 6.99. An <i>algebra</i> for an operad $\mathcal{O}$ is an operad functor $F: \mathcal{O} \rightarrow \mathbf{Set}$.
fong-spivak-seven-sketches- F02378	229	0.998	$F(t)$	$F(t)$	operad $\mathcal{O}$ and an algebra $F: \mathcal{O} \rightarrow \mathbf{Set}$ will take a type t and return a set $F(t)$ of fillers
fong-spivak-seven-sketches- F02379	229	1.000	$f \backslash\mathrm{in} \backslash\mathrm{mathcal}{0} \backslash\mathrm{left}(t_{\{1\}}, \backslash\mathrm{ldots}, t_{\{n\}} ; t \backslash\mathrm{right})$	$f \in \mathcal{O}(t_1, \dots, t_n; t)$	for box t . Moreover, given an operation (i.e., a wiring diagram) $f \in \mathcal{O}(t_1, \dots, t_n; t)$, we
fong-spivak-seven-sketches- F02380	229	1.000	$F(f)$	$F(f)$	get a function $F(f)$ that takes an element of each set $F(t_i)$, and returns an element of
fong-spivak-seven-sketches- F02381	229	1.000	$F \backslash\mathrm{left}(t_{\{i\}} \backslash\mathrm{right})$	$F(t_i)$	get a function $F(f)$ that takes an element of each set $F(t_i)$, and returns an element of
fong-spivak-seven-sketches- F02382	229	1.000	$t_{\{1\}}, \backslash\mathrm{ldots}, t_{\{n\}}$	$t_1, \dots, t_n$	$F(t)$. For example, it takes n circuits with interface $\boxed{t_1, \dots, t_n}$ respectively, and returns

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02383	229	0.854	$T=\operatorname{Ob}()$	$T = \operatorname{Ob}()$	<i>Example 6.100.</i> For electric circuits, the types are again finite sets, $T = \operatorname{Ob}(\mathbf{FinSet})$,
fong-spivak-seven-sketches-F02384	229	1.000	$\operatorname{Circ}(t)$	$\operatorname{Circ}(t)$	set $\operatorname{Circ}(t)$ of fillers, namely the set of electric circuits with that t -marked terminals. As
fong-spivak-seven-sketches-F02385	229	0.967	$\operatorname{Circ}(\varphi) \in \operatorname{Set}(\operatorname{Circ}(2), \operatorname{Circ}(2), \operatorname{Circ}(2); \operatorname{Circ}(0))$	$\operatorname{Circ}(\varphi) \in \operatorname{Set}(\operatorname{Circ}(2), \operatorname{Circ}(2), \operatorname{Circ}(2); \operatorname{Circ}(0))$	drawn case, we would get a morphism $\operatorname{Circ}(\varphi) \in \operatorname{Set}(\operatorname{Circ}(2), \operatorname{Circ}(2), \operatorname{Circ}(2); \operatorname{Circ}(0))$,
fong-spivak-seven-sketches-F02386	233	1.000	$\in \mathbb{R}$	$\in \mathbb{R}$	some $\mathsf{safe} \in \mathbb{R}$ from other objects. To do so, several components must interact: a
fong-spivak-seven-sketches-F02387	233	1.000	$S^{\prime}$	S'	sensor that approximates the real distance by an internal variable S' , a controller that
fong-spivak-seven-sketches-F02388	234	1.000	$S \geq$	$S \geq$	In order for the system to ensure $S \geq \mathsf{safe}$, we need each of the components to
fong-spivak-seven-sketches-F02389	234	1.000	$S <$	$S <$	the action signal A , e.g. that if at any time $S < \mathsf{safe}$, then within one second it will
fong-spivak-seven-sketches-F02390	235	1.000	$A =$	$A =$	emit the signal $A = \mathsf{go}$. The motor maintains a relationship between A and S , e.g. that
fong-spivak-seven-sketches-F02391	236	1.000	$[a, b]$	$[a, b]$	• S (wire): The behavior of S over a time-interval $[a, b]$ is that of all continuous
fong-spivak-seven-sketches-F02392	236	1.000	$[a, b] \rightarrow \mathbb{R}$	$[a, b] \rightarrow \mathbb{R}$	real-valued functions $[a, b] \rightarrow \mathbb{R}$.
fong-spivak-seven-sketches-F02393	236	0.809	$\left(\left(S^{\prime}, A\right) \mid E q\right)$	$\{(S', A) \mid Eq\}$	• controller (box): the relation $\{(S', A) \mid Eq. (7.2)\}$, i.e. all behavioral pairs (S', A)
fong-spivak-seven-sketches-F02394	236	0.809	$\left(S^{\prime}, A\right)$	(S', A)	• controller (box): the relation $\{(S', A) \mid Eq. (7.2)\}$, i.e. all behavioral pairs (S', A)
fong-spivak-seven-sketches-F02395	236	0.982	$\in P$	$\in P$	to the set $\mathbb{B} = \{\mathsf{false}, \mathsf{true}\}$ of truth values, where $\mathsf{Brendan} \in P$ maps to true because
fong-spivak-seven-sketches-F02396	237	0.994	$\mathcal{E}$	$\mathcal{E}$	Set in many ways. Here are some facts that are true of any topos $\mathcal{E}$:
fong-spivak-seven-sketches-F02397	237	1.000	$1 \rightarrow \Omega$	$1 \xrightarrow{\mathsf{true}} \Omega$	5. $\mathcal{E}$ has a subobject classifier $1 \xrightarrow{\mathsf{true}} \Omega$.
fong-spivak-seven-sketches-F02398	237	0.736	$\mathcal{E} =$	$\mathcal{E} =$	In particular, since Set is a topos, all of the above facts are true for $\mathcal{E} = \mathbf{Set}$. Our first
fong-spivak-seven-sketches-F02399	237	0.999	(B, C, E, F)	(B, C, E, F)	(B, C, E, F) is a pullback. But in the right-hand square, does the corner symbol mean
fong-spivak-seven-sketches-F02400	237	0.939	(A, B, D, E)	(A, B, D, E)	that (A, B, D, E) is a pullback or that (A, C, D, F) is a pullback? It's ambiguous, but as
fong-spivak-seven-sketches-F02401	237	0.939	(A, C, D, F)	(A, C, D, F)	that (A, B, D, E) is a pullback or that (A, C, D, F) is a pullback? It's ambiguous, but as
fong-spivak-seven-sketches-F02402	237	0.980	$\left(B, C, B^{\prime}, C^{\prime}\right)$	(B, C, B', C')	Proposition 7.3. In the commutative diagram below, suppose that the (B, C, B', C')
fong-spivak-seven-sketches-F02403	237	0.574	$\left(A, B, A^{\prime}, B^{\prime}\right)$	(A, B, A', B')	Then the (A, B, A', B') square is a pullback iff the (A, C, A', C') rectangle is a pullback.
fong-spivak-seven-sketches-F02404	237	0.574	$\left(A, C, A^{\prime}, C^{\prime}\right)$	(A, C, A', C')	Then the (A, B, A', B') square is a pullback iff the (A, C, A', C') rectangle is a pullback.
fong-spivak-seven-sketches-F02405	238	1.000	$i: B^{\prime} \rightarrow B$	$i: B' \rightarrow B$	phism. That is, suppose that $i: B' \rightarrow B$ is an isomorphism and $f: A \rightarrow B$ is any
fong-spivak-seven-sketches-F02406	238	1.000	$i^{\prime}$	i'	morphism. Show that i' is an isomorphism, in the following diagram:
fong-spivak-seven-sketches-F02407	238	1.000	$f^{\prime}$	f'	—
fong-spivak-seven-sketches-F02408	239	1.000	$f: C \rightarrow D$	$f: C \rightarrow D$	epi-mono factorizations are. We say that a morphism $f: C \rightarrow D$ in $\mathcal{E}$ has an epi-mono
fong-spivak-seven-sketches-F02409	239	1.000	$\operatorname{im}(f)$	$\operatorname{im}(f)$	factorization if it has an ‘image’; that is, there is an object $\operatorname{im}(f)$, an epimorphism

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_239_F02410	239	0.865	$C \rightarrowtail \operatorname{im}(f)$	$C \rightarrow \operatorname{im}(f)$	$C \twoheadrightarrow \operatorname{im}(f)$, and a monomorphism $\operatorname{im}(f) \hookrightarrow D$, whose composite is f .
fong-spivak-seven-sketches_239_F02411	239	0.865	$\operatorname{im}(f) \succcurlyeq D$	$\operatorname{im}(f) \succ D$	$C \twoheadrightarrow \operatorname{im}(f)$, and a monomorphism $\operatorname{im}(f) \hookrightarrow D$, whose composite is f .
fong-spivak-seven-sketches_239_F02412	239	1.000	$\operatorname{im}(f) =$	$\operatorname{im}(f) =$	function $f: C \rightarrow D$ may be factored as a surjective function onto its image $\operatorname{im}(f) =$
fong-spivak-seven-sketches_239_F02413	239	1.000	$\{f(c) \mid c \in C\}$	$\{f(c) \mid c \in C\}$	$\{f(c) \mid c \in C\}$, followed by the inclusion of this image into the codomain D . Moreover,
fong-spivak-seven-sketches_239_F02414	239	1.000	$f: \underline{3} \rightarrow \underline{3}$	$f: \underline{3} \rightarrow \underline{3}$	<i>Exercise 7.9.</i> Factor the following function $f: \underline{3} \rightarrow \underline{3}$ as an epimorphism followed by
fong-spivak-seven-sketches_239_F02415	239	1.000	$f = (e ; m)$	$f = (e; m)$	epimorphism e and a monomorphism m such that $f = (e \circ m)$ is their composite.
fong-spivak-seven-sketches_239_F02416	239	1.000	$C, D \in \mathcal{C}$	$C, D \in \mathcal{C}$	closure means that for any two objects $C, D \in \mathcal{C}$, there is a ‘hom-object’ $D^C \in \mathcal{C}$ and a
fong-spivak-seven-sketches_239_F02417	239	1.000	$D^C \in \mathcal{C}$	$D^C \in \mathcal{C}$	closure means that for any two objects $C, D \in \mathcal{C}$, there is a ‘hom-object’ $D^C \in \mathcal{C}$ and a
fong-spivak-seven-sketches_239_F02418	239	0.998	$A \in \mathcal{C}$	$A \in \mathcal{C}$	natural isomorphism for any $A \in \mathcal{C}$:
fong-spivak-seven-sketches_239_F02419	239	1.000	$f(-, -): A \times C \rightarrow D$	$f(-, -): A \times C \rightarrow D$	some game $f(-, -): A \times C \rightarrow D$: for whatever action $a \in A$ that you take and action
fong-spivak-seven-sketches_239_F02420	239	1.000	$f(a, c)$	$f(a, c)$	$c \in C$ that I take, $f(a, c)$ is some value in D . Since you’re self-centered but loving, you
fong-spivak-seven-sketches_239_F02421	239	0.993	$f(a, -): C \rightarrow D$	$f(a, -): C \rightarrow D$	do a , you make a game $f(a, -): C \rightarrow D$ for me alone. In the formalism, D^C represents
fong-spivak-seven-sketches_239_F02422	239	0.993	D^C	D^C	do a , you make a game $f(a, -): C \rightarrow D$ for me alone. In the formalism, D^C represents
fong-spivak-seven-sketches_239_F02423	239	0.962	$v, w, x \in V$	$v, w, x \in V$	tion 2.90—and suppose it satisfies the following for all $v, w, x \in V$:
fong-spivak-seven-sketches_239_F02424	239	1.000	$v \leq I$	$v \leq I$	• $v \leq I$,
fong-spivak-seven-sketches_239_F02425	239	1.000	$v \otimes w \leq v$	$v \otimes w \leq v$	• $v \otimes w \leq v$ and $v \otimes w \leq w$, and
fong-spivak-seven-sketches_239_F02426	239	1.000	$v \otimes w \leq w$	$v \otimes w \leq w$	• $v \otimes w \leq v$ and $v \otimes w \leq w$, and
fong-spivak-seven-sketches_239_F02427	239	1.000	$x \leq v$	$x \leq v$	• if $x \leq v$ and $x \leq w$ then $x \leq v \otimes w$.
fong-spivak-seven-sketches_239_F02428	239	1.000	$x \leq w$	$x \leq w$	• if $x \leq v$ and $x \leq w$ then $x \leq v \otimes w$.
fong-spivak-seven-sketches_239_F02429	239	1.000	$x \leq v \otimes w$	$x \leq v \otimes w$	• if $x \leq v$ and $x \leq w$ then $x \leq v \otimes w$.
fong-spivak-seven-sketches_240_F02430	240	0.924	$X \mapsto Y$	$X \mapsto Y$	are injections, and any injection $X \hookrightarrow Y$ is isomorphic to a subset of Y . This gives
fong-spivak-seven-sketches_240_F02431	240	0.925	$\Omega \in \mathcal{E}$	$\Omega \in \mathcal{E}$	minimal object 1. A <i>subobject classifier</i> in $\mathcal{E}$ consists of an object $\Omega \in \mathcal{E}$, together with a
fong-spivak-seven-sketches_240_F02432	240	1.000	$1 \rightarrow \Omega$	$1 \rightarrow \Omega$	monomorphism $\text{true}: 1 \rightarrow \Omega$, satisfying the following property: for any objects X and
fong-spivak-seven-sketches_240_F02433	240	0.899	$m: X \succcurlyeq Y$	$m: X \succ Y$	Y and monomorphism $m: X \hookrightarrow Y$ in $\mathcal{E}$, there is a unique morphism $\ulcorner m \urcorner: Y \rightarrow \Omega$ such
fong-spivak-seven-sketches_240_F02434	240	0.899	$\ulcorner m \urcorner: Y \rightarrow \Omega$	$\ulcorner m \urcorner: Y \rightarrow \Omega$	Y and monomorphism $m: X \hookrightarrow Y$ in $\mathcal{E}$, there is a unique morphism $\ulcorner m \urcorner: Y \rightarrow \Omega$ such
fong-spivak-seven-sketches_240_F02435	240	1.000	$\ulcorner m \urcorner$	$\ulcorner m \urcorner$	We refer to $\ulcorner m \urcorner$ as the <i>characteristic map</i> of m , or we say that $\ulcorner m \urcorner$ <i>classifies</i> m . Conversely,
fong-spivak-seven-sketches_240_F02436	240	1.000	$p: Y \rightarrow \Omega$	$p: Y \rightarrow \Omega$	given any map $p: Y \rightarrow \Omega$, we denote the pullback of true as on the right of Eq. (7.13).

Identifier	Page	Conf.	LaTeX source	Rendered	Image															
fong-spivak-seven-sketches_1 F02437	240	■ 1.000	$Y \rightarrow \Omega$	$Y \rightarrow \Omega$	<i>A predicate on Y is a morphism $Y \rightarrow \Omega$.</i>															
fong-spivak-seven-sketches_1 F02438	241	■ 0.999	$1 \rightarrow$	$1 \rightarrow$	generically called true : $1 \rightarrow \Omega$; in the case of Set it is played by the function $1 \rightarrow$															
fong-spivak-seven-sketches_1 F02439	241	■ 0.980	$m: X \hookrightarrow Y$	$m: X \hookrightarrow Y$	tion 7.12 says that for any injective function $m: X \hookrightarrow Y$ between sets, we are supposed															
fong-spivak-seven-sketches_1 F02440	241	■ 0.998	$\ulcorner m \urcorner: Y \rightarrow \{$	$\ulcorner m \urcorner: Y \rightarrow \{$	to be able to find a characteristic function $\ulcorner m \urcorner: Y \rightarrow \{\text{true}, \text{false}\}$ with some sort of															
fong-spivak-seven-sketches_1 F02441	241	■ 1.000	$\ulcorner m \urcorner(y)$	$\ulcorner m \urcorner(y)$	In other words, if we think of X as a subobject of Y , then we make $\ulcorner m \urcorner(y)$ equal to															
fong-spivak-seven-sketches_1 F02442	241	■ 1.000	$y \in X$	$y \in X$	true iff $y \in X$.															
fong-spivak-seven-sketches_1 F02443	241	■ 1.000	$p: Y \rightarrow \mathbb{B}$	$p: Y \rightarrow \mathbb{B}$	$p: Y \rightarrow \mathbb{B}$, and vice versa. How it works is encoded in Definition 7.12, but the basic															
fong-spivak-seven-sketches_1 F02444	241	■ 1.000	$\{Y \mid p\}$	$\{Y \mid p\}$	This might help explain our abstract notation $\{Y \mid p\}$ in Eq. (7.13).															
fong-spivak-seven-sketches_1 F02445	241	■ 0.998	$X=\mathbb{N}=\{0,1,2, \ldots\}$	$X = \mathbb{N} = \{0, 1, 2, \dots\}$	<i>Exercise 7.16.</i> Let $X = \mathbb{N} = \{0, 1, 2, \dots\}$ and $Y = \mathbb{Z} = \{\dots, -1, 0, 1, 2, \dots\}$; we have															
fong-spivak-seven-sketches_1 F02446	241	■ 0.998	$Y=\mathbb{Z}=\{\ldots,-1,0,1,2, \ldots\}$	$Y = \mathbb{Z} = \{\dots, -1, 0, 1, 2, \dots\}$	<i>Exercise 7.16.</i> Let $X = \mathbb{N} = \{0, 1, 2, \dots\}$ and $Y = \mathbb{Z} = \{\dots, -1, 0, 1, 2, \dots\}$; we have															
fong-spivak-seven-sketches_1 F02447	241	■ 0.985	$\ulcorner m \urcorner: Y \rightarrow \mathbb{B}$	$\ulcorner m \urcorner: Y \rightarrow \mathbb{B}$	$\ulcorner m \urcorner: Y \rightarrow \mathbb{B}$, as in Definition 7.12.															
fong-spivak-seven-sketches_1 F02448	241	■ 1.000	$\ulcorner m \urcorner(-5) \in \mathbb{B}$	$\ulcorner m \urcorner(-5) \in \mathbb{B}$	1. What is $\ulcorner m \urcorner(-5) \in \mathbb{B}$?															
fong-spivak-seven-sketches_1 F02449	241	■ 1.000	$\ulcorner m \urcorner(0) \in \mathbb{B}$	$\ulcorner m \urcorner(0) \in \mathbb{B}$	2. What is $\ulcorner m \urcorner(0) \in \mathbb{B}$?															
fong-spivak-seven-sketches_1 F02450	241	■ 0.915	$\operatorname{id}_{\mathbb{N}}: \mathbb{N} \rightarrow \mathbb{N}$	$\operatorname{id}_{\mathbb{N}}: \mathbb{N} \rightarrow \mathbb{N}$	1. Consider the identity function $\operatorname{id}_{\mathbb{N}}: \mathbb{N} \rightarrow \mathbb{N}$. It is an injection, so it has a charac-															
fong-spivak-seven-sketches_1 F02451	241	■ 0.746	$\ulcorner \operatorname{id}_{\mathbb{N}} \urcorner: \mathbb{N} \rightarrow \mathbb{B}$	$\ulcorner \operatorname{id}_{\mathbb{N}} \urcorner: \mathbb{N} \rightarrow \mathbb{B}$	teristic function $\ulcorner \operatorname{id}_{\mathbb{N}} \urcorner: \mathbb{N} \rightarrow \mathbb{B}$. Give a concrete description of $\ulcorner \operatorname{id}_{\mathbb{N}} \urcorner$, i.e. its exact															
fong-spivak-seven-sketches_1 F02452	241	■ 0.746	$\ulcorner \operatorname{id}_{\mathbb{N}} \urcorner$	$\ulcorner \operatorname{id}_{\mathbb{N}} \urcorner$	teristic function $\ulcorner \operatorname{id}_{\mathbb{N}} \urcorner: \mathbb{N} \rightarrow \mathbb{B}$. Give a concrete description of $\ulcorner \operatorname{id}_{\mathbb{N}} \urcorner$, i.e. its exact															
fong-spivak-seven-sketches_1 F02453	242	■ 0.990	$!_{\mathbb{N}}: \varnothing \rightarrow \mathbb{N}$	$!_{\mathbb{N}}: \varnothing \rightarrow \mathbb{N}$	2. Consider the unique function $!_{\mathbb{N}}: \varnothing \rightarrow \mathbb{N}$ from the empty set. Give a concrete															
fong-spivak-seven-sketches_1 F02454	242	■ 0.470	$\ulcorner !_{\mathbb{N}} \urcorner: \mathbb{N} \rightarrow \mathbb{B}$	$\ulcorner !_{\mathbb{N}} \urcorner: \mathbb{N} \rightarrow \mathbb{B}$	description of $\ulcorner !_{\mathbb{N}} \urcorner: \mathbb{N} \rightarrow \mathbb{B}$.															
fong-spivak-seven-sketches_1 F02455	242	■ 1.000	$1 \rightarrow \mathbb{B} \times \mathbb{B}$	$1 \rightarrow \mathbb{B} \times \mathbb{B}$	Obtaining the AND operation. Consider the function $1 \rightarrow \mathbb{B} \times \mathbb{B}$ picking out the															
fong-spivak-seven-sketches_1 F02456	242	■ 0.257	$\ulcorner \wedge \urcorner$	$\ulcorner \wedge \urcorner$	$\ulcorner (\text{true}, \text{true}) \urcorner: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$. What function is it? By Eq. (7.15) the only element of $\mathbb{B} \times \mathbb{B}$															
fong-spivak-seven-sketches_1 F02457	242	■ 0.257	$\mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$	$\mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$	$\ulcorner (\text{true}, \text{true}) \urcorner: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$. What function is it? By Eq. (7.15) the only element of $\mathbb{B} \times \mathbb{B}$															
fong-spivak-seven-sketches_1 F02458	242	■ 0.257	$\mathbb{B} \times \mathbb{B}$	$\mathbb{B} \times \mathbb{B}$	$\ulcorner (\text{true}, \text{true}) \urcorner: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$. What function is it? By Eq. (7.15) the only element of $\mathbb{B} \times \mathbb{B}$															
fong-spivak-seven-sketches_1 F02459	242	■ 0.990	$\ulcorner ($	$\ulcorner ($	that can be sent to true is $(\text{true}, \text{true})$. Thus $\ulcorner (\text{true}, \text{true}) \urcorner(P, Q) \in \mathbb{B}$ must be given															
fong-spivak-seven-sketches_1 F02460	242	■ 0.990	$\urcorner(P, Q) \in \mathbb{B}$	$\urcorner(P, Q) \in \mathbb{B}$	that can be sent to true is $(\text{true}, \text{true})$. Thus $\ulcorner (\text{true}, \text{true}) \urcorner(P, Q) \in \mathbb{B}$ must be given															
fong-spivak-seven-sketches_1 F02461	242	■ 0.752	$\urcorner(P, Q)$	$\urcorner(P, Q)$	<table><tr><th>P</th><th>Q</th><th>$\ulcorner (\text{true}, \text{true}) \urcorner(P, Q)$</th></tr><tr><td>true</td><td>true</td><td>true</td></tr><tr><td>true</td><td>false</td><td>false</td></tr><tr><td>false</td><td>true</td><td>false</td></tr><tr><td>false</td><td>false</td><td>false</td></tr></table>	P	Q	$\ulcorner (\text{true}, \text{true}) \urcorner(P, Q)$	true	true	true	true	false	false	false	true	false	false	false	false
P	Q	$\ulcorner (\text{true}, \text{true}) \urcorner(P, Q)$																		
true	true	true																		
true	false	false																		
false	true	false																		
false	false	false																		

Identifier	Page	Conf.	LaTeX source	Rendered	Image															
fong-spivak-seven-sketches-F02462	242	■ 1.000	P \wedge Q	$P \wedge Q$	This is exactly the truth table for the AND of P and Q , i.e. for $P \wedge Q$. In other words,															
fong-spivak-seven-sketches-F02463	242	■ 0.965	)\urcorner=\wedge	$)^\urcorner = \wedge$	$\urcorner(\text{true}, \text{true})^\urcorner = \wedge$. Note that this defines $\wedge$ as a function $\wedge: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$, and we use															
fong-spivak-seven-sketches-F02464	242	■ 0.965	\wedge: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}	$\wedge: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$	$\urcorner(\text{true}, \text{true})^\urcorner = \wedge$. Note that this defines $\wedge$ as a function $\wedge: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$, and we use															
fong-spivak-seven-sketches-F02465	242	■ 1.000	x \wedge y:=\wedge(x, y)	$x \wedge y := \wedge(x, y)$	the usual infix notation $x \wedge y := \wedge(x, y)$.															
fong-spivak-seven-sketches-F02466	242	■ 1.000	\vee: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}	$\vee: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$	OR of P and Q , i.e. that of the function $\vee: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$ defining OR, is:															
fong-spivak-seven-sketches-F02467	242	■ 0.785	P \vee Q	$P \vee Q$	<table><tr><td>P</td><td>Q</td><td>$P \vee Q$</td></tr><tr><td>true</td><td>true</td><td>true</td></tr><tr><td>true</td><td>false</td><td>true</td></tr><tr><td>false</td><td>true</td><td>true</td></tr><tr><td>false</td><td>false</td><td>false</td></tr></table>	P	Q	$P \vee Q$	true	true	true	true	false	true	false	true	true	false	false	false
P	Q	$P \vee Q$																		
true	true	true																		
true	false	true																		
false	true	true																		
false	false	false																		
fong-spivak-seven-sketches-F02468	242	■ 1.000	m: X \subseteq \mathbb{B} \times \mathbb{B}	$m: X \subseteq \mathbb{B} \times \mathbb{B}$	$m: X \subseteq \mathbb{B} \times \mathbb{B}$, what subset would X be? By Eq. (7.15), X should be the set of $y \in Y$															
fong-spivak-seven-sketches-F02469	242	■ 0.801	\mathbb{B} \times \{	$\mathbb{B} \times \{$	think of X as the union of $\{\text{true}\} \times \mathbb{B}$ and $\mathbb{B} \times \{\text{true}\}$ —a colimit of limits involving the															
fong-spivak-seven-sketches-F02470	242	■ 1.000	\Omega \times \Omega	$\Omega \times \Omega$	subobject of $\Omega \times \Omega$, and hence classify a map $\Omega \times \Omega \rightarrow \Omega$, in any topos $\mathcal{E}$.															
fong-spivak-seven-sketches-F02471	242	■ 1.000	\Omega \times \Omega \rightarrow \Omega	$\Omega \times \Omega \rightarrow \Omega$	subobject of $\Omega \times \Omega$, and hence classify a map $\Omega \times \Omega \rightarrow \Omega$, in any topos $\mathcal{E}$.															
fong-spivak-seven-sketches-F02472	243	■ 1.000	\neg: \mathbb{B} \rightarrow \mathbb{B}	$\neg: \mathbb{B} \rightarrow \mathbb{B}$	function $\neg: \mathbb{B} \rightarrow \mathbb{B}$ is the characteristic function of some thing, (*?*)).															
fong-spivak-seven-sketches-F02473	243	■ 1.000	P \Rightarrow Q	$P \Rightarrow Q$	<i>Exercise 7.20.</i> Given two booleans P, Q , define $P \Rightarrow Q$ to mean $P = (P \wedge Q)$.															
fong-spivak-seven-sketches-F02474	243	■ 1.000	P=(P \wedge Q)	$P = (P \wedge Q)$	<i>Exercise 7.20.</i> Given two booleans P, Q , define $P \Rightarrow Q$ to mean $P = (P \wedge Q)$.															
fong-spivak-seven-sketches-F02475	243	■ 1.000	m: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}	$m: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$	3. What is the characteristic function $m: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$ for $P \Rightarrow Q$?															
fong-spivak-seven-sketches-F02476	243	■ 1.000	E:={n \in \mathbb{N} \mid n	$E := \{n \in \mathbb{N} \mid n$	<i>Exercise 7.21.</i> Consider the sets $E := \{n \in \mathbb{N} \mid n \text{ is even}\}$, $P := \{n \in \mathbb{N} \mid n \text{ is prime}\}$,															
fong-spivak-seven-sketches-F02477	243	■ 1.000	\}, P:={n \in \mathbb{N} \mid n	$\}, P := \{n \in \mathbb{N} \mid n$	<i>Exercise 7.21.</i> Consider the sets $E := \{n \in \mathbb{N} \mid n \text{ is even}\}$, $P := \{n \in \mathbb{N} \mid n \text{ is prime}\}$,															
fong-spivak-seven-sketches-F02478	243	■ 1.000	T:={n \in \mathbb{N} \mid n \geq 10\}	$T := \{n \in \mathbb{N} \mid n \geq 10\}$	and $T := \{n \in \mathbb{N} \mid n \geq 10\}$. Each is a subset of $\mathbb{N}$, so defines a function $\mathbb{N} \rightarrow \mathbb{B}$.															
fong-spivak-seven-sketches-F02479	243	■ 1.000	\mathbb{N} \rightarrow \mathbb{B}	$\mathbb{N} \rightarrow \mathbb{B}$	and $T := \{n \in \mathbb{N} \mid n \geq 10\}$. Each is a subset of $\mathbb{N}$, so defines a function $\mathbb{N} \rightarrow \mathbb{B}$.															
fong-spivak-seven-sketches-F02480	243	■ 1.000	\ulcorner E \urcorner(17)	$\urcorner E^\urcorner(17)$	1. What is $\urcorner E^\urcorner(17)$?															
fong-spivak-seven-sketches-F02481	243	■ 1.000	\ulcorner P \urcorner(17)	$\urcorner P^\urcorner(17)$	2. What is $\urcorner P^\urcorner(17)$?															
fong-spivak-seven-sketches-F02482	243	■ 1.000	\ulcorner T \urcorner(17)	$\urcorner T^\urcorner(17)$	3. What is $\urcorner T^\urcorner(17)$?															
fong-spivak-seven-sketches-F02483	243	■ 0.995	(\ulcorner E \urcorner \wedge \ulcorner P \urcorner) \vee \ulcorner T \urcorner	$(\urcorner E^\urcorner \wedge \urcorner P^\urcorner) \vee \urcorner T^\urcorner$	4. Name the smallest three elements in the set classified by $(\urcorner E^\urcorner \wedge \urcorner P^\urcorner) \vee \urcorner T^\urcorner$. $\diamond$															
fong-spivak-seven-sketches-F02484	243	■ 1.000	\wedge, \vee, \Rightarrow: \Omega \times \Omega \rightarrow \Omega	$\wedge, \vee, \Rightarrow: \Omega \times \Omega \rightarrow \Omega$	how to understand logic in terms of Ω : there are various maps $\wedge, \vee, \Rightarrow: \Omega \times \Omega \rightarrow \Omega$															
fong-spivak-seven-sketches-F02485	243	■ 1.000	\neg: \Omega \rightarrow \Omega	$\neg: \Omega \rightarrow \Omega$	and $\neg: \Omega \rightarrow \Omega$ etc., which serve as logical connectives. These are operations on truth															

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_11 F02486	244	■ 1.000	$\mathbb{R}^2$	$\mathbb{R}^2$	of the plane $\mathbb{R}^2$ as a space, we might examine only points and regions in it. But if we
fong-spivak-seven-sketches_11 F02487	244	■ 0.729	$\mathcal{C}^{\mathrm{op}} \rightarrow$	$\mathcal{C}^{\mathrm{op}} \rightarrow$	functors $\mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Set}$. ⁵ Such functors are called <i>presheaves on $\mathcal{C}$</i> .
fong-spivak-seven-sketches_11 F02488	244	■ 0.988	$P(c)$	$P(c)$	To each object $c \in \mathcal{C}$, we refer to the set $P(c)$ as <i>the set of sections of P over c</i> . To each
fong-spivak-seven-sketches_11 F02489	244	■ 1.000	$P(f): P(c) \rightarrow P(c')$	$P(f): P(c) \rightarrow P(c')$	morphism $f: c' \rightarrow c$, we refer to the function $P(f): P(c) \rightarrow P(c')$ as the <i>restriction map</i>
fong-spivak-seven-sketches_11 F02490	244	■ 1.000	$s \in P(c)$	$s \in P(c)$	<i>along f</i> . For any section $s \in P(c)$, we may denote $P(f)(s) \in P(c')$, i.e. its restriction
fong-spivak-seven-sketches_11 F02491	244	■ 1.000	$P(f)(s) \in P(c')$	$P(f)(s) \in P(c')$	<i>along f</i> . For any section $s \in P(c)$, we may denote $P(f)(s) \in P(c')$, i.e. its restriction
fong-spivak-seven-sketches_11 F02492	244	■ 1.000	$s _f$	$s _f$	<i>along f</i> , by $s _f$.
fong-spivak-seven-sketches_11 F02493	245	■ 1.000	$\alpha: P \rightarrow Q$	$\alpha: P \rightarrow Q$	If P and Q are presheaves, a <i>morphism</i> $\alpha: P \rightarrow Q$ between them is a natural trans-
fong-spivak-seven-sketches_11 F02494	245	■ 0.852	$\mathbf{ArShp}^{\mathrm{op}} \rightarrow$	$\mathbf{ArShp}^{\mathrm{op}} \rightarrow$	A presheaf on $\mathbf{ArShp}$ is a functor $I: \mathbf{ArShp}^{\mathrm{op}} \rightarrow \mathbf{Set}$, which is a database instance on
fong-spivak-seven-sketches_11 F02495	245	■ 0.532	$\mathbf{ArShp}^{\mathrm{op}}$	$\mathbf{ArShp}^{\mathrm{op}}$	$\mathbf{ArShp}^{\mathrm{op}}$. Note that $\mathbf{ArShp}^{\mathrm{op}}$ is what we called Gr in Section 3.3.5; there we showed
fong-spivak-seven-sketches_11 F02496	245	■ 0.532	$\mathbf{ArShp}^{\mathrm{op}}$	$\mathbf{ArShp}^{\mathrm{op}}$	$\mathbf{ArShp}^{\mathrm{op}}$. Note that $\mathbf{ArShp}^{\mathrm{op}}$ is what we called Gr in Section 3.3.5; there we showed
fong-spivak-seven-sketches_11 F02497	246	■ 1.000	$P(X) = \{U \subseteq X\}$	$P(X) = \{U \subseteq X\}$	Definition 7.25. Let X be a set, and let $P(X) = \{U \subseteq X\}$ denote its set of subsets. A
fong-spivak-seven-sketches_11 F02498	246	■ 0.672	$\mathbf{Op} \subseteq P(X)$	$\mathbf{Op} \subseteq P(X)$	<i>topology</i> on X is a subset $\mathbf{Op} \subseteq P(X)$, elements of which we call <i>open sets</i> , ⁶ satisfying the
fong-spivak-seven-sketches_11 F02499	246	■ 1.000	$X \in \mathbf{Op}$	$X \subseteq X$	(a) Whole set: the subset $X \subseteq X$ is open, i.e. $X \in \mathbf{Op}$.
fong-spivak-seven-sketches_11 F02500	246	■ 1.000	$X \in \mathbf{Op}$	$X \in \mathbf{Op}$	(a) Whole set: the subset $X \subseteq X$ is open, i.e. $X \in \mathbf{Op}$.
fong-spivak-seven-sketches_11 F02501	246	■ 0.871	$U, V \in \mathbf{Op}$	$U, V \in \mathbf{Op}$	(b) Binary intersections: if $U, V \in \mathbf{Op}$ then $(U \cap V) \in \mathbf{Op}$.
fong-spivak-seven-sketches_11 F02502	246	■ 0.871	$(U \cap V) \in \mathbf{Op}$	$(U \cap V) \in \mathbf{Op}$	(b) Binary intersections: if $U, V \in \mathbf{Op}$ then $(U \cap V) \in \mathbf{Op}$.
fong-spivak-seven-sketches_11 F02503	246	■ 0.743	$U_i \in \mathbf{Op}$	$U_i \in \mathbf{Op}$	(c) Arbitrary unions: if I is a set and if we are given an open set $U_i \in \mathbf{Op}$ for each i ,
fong-spivak-seven-sketches_11 F02504	246	■ 0.999	$\left(\bigcup_{i \in I} U_i\right) \in \mathbf{Op}$	$\left(\bigcup_{i \in I} U_i\right) \in \mathbf{Op}$	then their union is also open, $(\bigcup_{i \in I} U_i) \in \mathbf{Op}$. We interpret the particular case
fong-spivak-seven-sketches_11 F02505	246	■ 1.000	$I = \emptyset$	$I = \emptyset$	where $I = \emptyset$ to mean that the empty set is open: $\emptyset \in \mathbf{Op}$.
fong-spivak-seven-sketches_11 F02506	246	■ 1.000	$\emptyset \in \mathbf{Op}$	$\emptyset \in \mathbf{Op}$	where $I = \emptyset$ to mean that the empty set is open: $\emptyset \in \mathbf{Op}$.
fong-spivak-seven-sketches_11 F02507	246	■ 1.000	$U = \bigcup_{i \in I} U_i$	$U = \bigcup_{i \in I} U_i$	If $U = \bigcup_{i \in I} U_i$, we say that $(U_i)_{i \in I}$ <i>covers</i> U .
fong-spivak-seven-sketches_11 F02508	246	■ 1.000	$(U_i)_{i \in I}$	$(U_i)_{i \in I}$	If $U = \bigcup_{i \in I} U_i$, we say that $(U_i)_{i \in I}$ <i>covers</i> U .
fong-spivak-seven-sketches_11 F02509	246	■ 0.630	$(X, \mathbf{Op})$	$(X, \mathbf{Op})$	A pair $(X, \mathbf{Op})$, where X is a set and $\mathbf{Op}$ is a topology on X , is called a <i>topological</i>
fong-spivak-seven-sketches_11 F02510	246	■ 0.630	$\mathbf{Op}$	$\mathbf{Op}$	A pair $(X, \mathbf{Op})$, where X is a set and $\mathbf{Op}$ is a topology on X , is called a <i>topological</i>
fong-spivak-seven-sketches_11 F02511	246	■ 0.914	$X, \mathbf{Op}_X$	$X, \mathbf{Op}_X$	A <i>continuous function</i> between topological spaces $(X, \mathbf{Op}_X)$ and $(Y, \mathbf{Op}_Y)$ is a function

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches- F02512	246	■ 0.914	$Y, \mathbf{0} \mathbf{p}_Y$	$Y, \mathbf{Op}_Y$	A continuous function between topological spaces $(X, \mathbf{Op}_X)$ and $(Y, \mathbf{Op}_Y)$ is a function
fong-spivak-seven-sketches- F02513	246	■ 0.650	$U \in \mathbf{0p}_Y$	$U \in \mathbf{Op}_Y$	$f: X \rightarrow Y$ such that for every $U \in \mathbf{Op}_Y$, the preimage $f^{-1}(U)$ is in $\mathbf{Op}_X$.
fong-spivak-seven-sketches- F02514	246	■ 0.650	$f^{-1}(U)$	$f^{-1}(U)$	$f: X \rightarrow Y$ such that for every $U \in \mathbf{Op}_Y$, the preimage $f^{-1}(U)$ is in $\mathbf{Op}_X$.
fong-spivak-seven-sketches- F02515	246	■ 0.650	$\mathbf{0p}_X$	$\mathbf{Op}_X$	$f: X \rightarrow Y$ such that for every $U \in \mathbf{Op}_Y$, the preimage $f^{-1}(U)$ is in $\mathbf{Op}_X$.
fong-spivak-seven-sketches- F02516	246	■ 1.000	$\epsilon \in \mathbb{R}$	$\epsilon \in \mathbb{R}$	Example 7.26. The usual topology $\mathbf{Op}$ on $\mathbb{R}^2$ is based on ‘ ϵ -balls.’ For any $\epsilon \in \mathbb{R}$ with
fong-spivak-seven-sketches- F02517	246	■ 1.000	$\epsilon > 0$	$\epsilon > 0$	$\epsilon > 0$, and any point $p = (x, y) \in \mathbb{R}^2$, define the ϵ -ball centered at p to be:
fong-spivak-seven-sketches- F02518	246	■ 1.000	$p = (x, y) \in \mathbb{R}^2$	$p = (x, y) \in \mathbb{R}^2$	$\epsilon > 0$, and any point $p = (x, y) \in \mathbb{R}^2$, define the ϵ -ball centered at p to be:
fong-spivak-seven-sketches- F02519	246	■ 1.000	$U \in \mathbf{0p}$	$U \in \mathbf{Op}$	⁶ In other words, we refer to a subset $U \subseteq X$ as open if $U \in \mathbf{Op}$.
fong-spivak-seven-sketches- F02520	247	■ 0.999	$B(x, y; \epsilon)$	$B(x, y; \epsilon)$	In other words, $B(x, y; \epsilon)$ is the set of all points within ϵ of (x, y) .
fong-spivak-seven-sketches- F02521	247	■ 0.974	$U \subseteq \mathbb{R}^2$	$U \subseteq \mathbb{R}^2$	For an arbitrary subset $U \subseteq \mathbb{R}^2$, we call it open and put it in $\mathbf{Op}$ if, for every $(x, y) \in U$
fong-spivak-seven-sketches- F02522	247	■ 0.974	$\in U$	$\in U$	For an arbitrary subset $U \subseteq \mathbb{R}^2$, we call it open and put it in $\mathbf{Op}$ if, for every $(x, y) \in U$
fong-spivak-seven-sketches- F02523	247	■ 1.000	$B(x, y; \epsilon) \subseteq U$	$B(x, y; \epsilon) \subseteq U$	there exists a (small enough) $\epsilon > 0$ such that $B(x, y; \epsilon) \subseteq U$.
fong-spivak-seven-sketches- F02524	247	■ 1.000	$p = (x, y) \in U$	$p = (x, y) \in U$	an open set $U \subseteq \mathbb{R}^2$, a point $p = (x, y) \in U$,
fong-spivak-seven-sketches- F02525	247	■ 1.000	$d(x_1, x_2) := x_1 - x_2 $	$d(x_1, x_2) := x_1 - x_2 $	Exercise 7.27. Consider the set $\mathbb{R}$. It is a metric space with $d(x_1, x_2) := x_1 - x_2 $.
fong-spivak-seven-sketches- F02526	247	■ 1.000	$x \in \mathbb{R}$	$x \in \mathbb{R}$	for each $x \in \mathbb{R}$, define $B(x, \epsilon)$.
fong-spivak-seven-sketches- F02527	247	■ 1.000	$B(x, \epsilon)$	$B(x, \epsilon)$	for each $x \in \mathbb{R}$, define $B(x, \epsilon)$.
fong-spivak-seven-sketches- F02528	247	■ 1.000	$U \subseteq \mathbb{R}$	$U \subseteq \mathbb{R}$	2. When is an arbitrary subset $U \subseteq \mathbb{R}$ called open, in analogy with Example 7.26?
fong-spivak-seven-sketches- F02529	247	■ 1.000	U_1, U_2	U_1, U_2	3. Find three open sets U_1, U_2 , and U in $\mathbb{R}$, such that $(U_i)_{i \in \{1,2\}}$ covers U .
fong-spivak-seven-sketches- F02530	247	■ 1.000	$(U_i)_{i \in \{1,2\}}$	$(U_i)_{i \in \{1,2\}}$	3. Find three open sets U_1, U_2 , and U in $\mathbb{R}$, such that $(U_i)_{i \in \{1,2\}}$ covers U .
fong-spivak-seven-sketches- F02531	247	■ 0.803	$\mathbf{0p}_{\text{crse}} = (\emptyset, X)$	$\mathbf{Op}_{\text{crse}} = (\emptyset, X)$	possible: $\mathbf{Op}_{\text{crse}} = (\emptyset, X)$. There is also a ‘finest’ topology, having as many open sets as
fong-spivak-seven-sketches- F02532	247	■ 0.592	$\mathbf{0p}_{\text{fine}} = \mathcal{P}(X)$	$\mathbf{Op}_{\text{fine}} = \mathcal{P}(X)$	possible: $\mathbf{Op}_{\text{fine}} = \mathcal{P}(X)$. The latter, $(X, \mathcal{P}(X))$ is called the discrete space on the set X .
fong-spivak-seven-sketches- F02533	247	■ 0.592	$(X, \mathcal{P}(X))$	$(X, \mathcal{P}(X))$	possible: $\mathbf{Op}_{\text{fine}} = \mathcal{P}(X)$. The latter, $(X, \mathcal{P}(X))$ is called the discrete space on the set X .
fong-spivak-seven-sketches- F02534	247	■ 0.995	$\mathbf{0p}_{\text{crse}}$	$\mathbf{Op}_{\text{crse}}$	1. Verify that for any set X , what we called $\mathbf{Op}_{\text{crse}}$ in Example 7.28 really is a topology,
fong-spivak-seven-sketches- F02535	247	■ 0.998	$\mathbf{0p}_{\text{fine}}$	$\mathbf{Op}_{\text{fine}}$	2. Verify also that $\mathbf{Op}_{\text{fine}}$ really is a topology.
fong-spivak-seven-sketches- F02536	247	■ 0.762	$(Y, \mathbf{0p}_Y)$	$(Y, \mathbf{Op}_Y)$	3. Show that if $(X, \mathcal{P}(X))$ is discrete and $(Y, \mathbf{Op}_Y)$ is any topological space, then every
fong-spivak-seven-sketches- F02537	247	■ 1.000	$X \rightarrow Y$	$X \rightarrow Y$	function $X \rightarrow Y$ is continuous.
fong-spivak-seven-sketches- F02538	247	■ 1.000	$X = \{1, 2\}$	$X = \{1, 2\}$	Example 7.30. There are four topologies possible on $X = \{1, 2\}$. Two are $\mathbf{Op}_{\text{crse}}$ and

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_1 F02539	247	■ 1.000	$d\left((x, y), \left(x^{\prime}, y^{\prime}\right)\right):=\sqrt{\left(x-x^{\prime}\right)^2+\left(y-y^{\prime}\right)^2}$	$d((x, y), (x', y')) := \sqrt{(x - x')^2 + (y - y')^2}$	⁷ Here, $d((x, y), (x', y')) := \sqrt{(x - x')^2 + (y - y')^2}$ is the usual ‘Euclidean distance’ between two points.
fong-spivak-seven-sketches_1 F02540	248	■ 0.817	$\mathbf{0} \ \mathbf{p}_{\text{fine}}$	$\mathbf{Op}_{\text{fine}}$	$\mathbf{Op}_{\text{fine}}$ from Example 7.28. The other two are:
fong-spivak-seven-sketches_1 F02541	248	■ 0.943	$\{1, 2\}, \ \mathbf{0} \ \mathbf{p}_1$	$\{1, 2\}, \mathbf{Op}_1$	The two topological spaces $(\{1, 2\}, \mathbf{Op}_1)$ and $(\{1, 2\}, \mathbf{Op}_2)$ are isomorphic; either one
fong-spivak-seven-sketches_1 F02542	248	■ 0.943	$\{1, 2\}, \ \mathbf{0} \ \mathbf{p}_2$	$\{1, 2\}, \mathbf{Op}_2$	The two topological spaces $(\{1, 2\}, \mathbf{Op}_1)$ and $(\{1, 2\}, \mathbf{Op}_2)$ are isomorphic; either one
fong-spivak-seven-sketches_1 F02543	248	■ 0.931	$(\mathbf{0} \ \mathbf{p}, \ \subseteq)$	$(\mathbf{Op}, \subseteq)$	$(X, \mathbf{Op})$, the set $\mathbf{Op}$ has the structure of a preorder using the subset relation, $(\mathbf{Op}, \subseteq)$.
fong-spivak-seven-sketches_1 F02544	248	■ 1.000	$U \ \subseteq U$	$U \subseteq U$	It is reflexive because $U \subseteq U$ for any $U \in \mathbf{Op}$, and it is transitive because if $U \subseteq V$ and
fong-spivak-seven-sketches_1 F02545	248	■ 1.000	$U \ \subseteq V$	$U \subseteq V$	It is reflexive because $U \subseteq U$ for any $U \in \mathbf{Op}$, and it is transitive because if $U \subseteq V$ and
fong-spivak-seven-sketches_1 F02546	248	■ 1.000	$V \ \subseteq W$	$V \subseteq W$	$V \subseteq W$ then $U \subseteq W$.
fong-spivak-seven-sketches_1 F02547	248	■ 1.000	$U \ \subseteq W$	$U \subseteq W$	$V \subseteq W$ then $U \subseteq W$.
fong-spivak-seven-sketches_1 F02548	248	■ 1.000	$U, \ V$	U, V	category: its objects are the open sets U and for any U, V the set of morphisms $\mathbf{Op}(U, V)$
fong-spivak-seven-sketches_1 F02549	248	■ 1.000	$\mathbf{0} \ \mathbf{p}(U, \ V)$	$\mathbf{Op}(U, V)$	category: its objects are the open sets U and for any U, V the set of morphisms $\mathbf{Op}(U, V)$
fong-spivak-seven-sketches_1 F02550	248	■ 1.000	$U \ \nsubseteq V$	$U \not\subseteq V$	is empty if $U \not\subseteq V$ and it has one element if $U \subseteq V$.
fong-spivak-seven-sketches_1 F02551	248	■ 0.761	$X, \ \mathbf{0} \ \mathbf{p}_1$	$X, \mathbf{Op}_1$	<i>Exercise 7.31.</i> Recall the Sierpinski space, say $(X, \mathbf{Op}_1)$ from Example 7.30.
fong-spivak-seven-sketches_1 F02552	248	■ 0.686	$X, \ \mathbf{0} \ \mathbf{p}$	$X, \mathbf{Op}$	<i>Exercise 7.32.</i> Given any topological space $(X, \mathbf{Op})$, any subset $Y \subseteq X$ can be given the
fong-spivak-seven-sketches_1 F02553	248	■ 0.686	$Y \ \subseteq X$	$Y \subseteq X$	<i>Exercise 7.32.</i> Given any topological space $(X, \mathbf{Op})$, any subset $Y \subseteq X$ can be given the
fong-spivak-seven-sketches_1 F02554	248	■ 0.955	$\mathbf{0} \ \mathbf{p}_{\{? \ \cap \ \gamma\}}$	$\mathbf{Op}_{? \cap \gamma}$	<i>subspace topology</i> , call it $\mathbf{Op}_{? \cap Y}$. This topology defines any $A \subseteq Y$ to be open, $A \in \mathbf{Op}_{? \cap Y}$,
fong-spivak-seven-sketches_1 F02555	248	■ 0.955	$A \ \subseteq Y$	$A \subseteq Y$	<i>subspace topology</i> , call it $\mathbf{Op}_{? \cap Y}$. This topology defines any $A \subseteq Y$ to be open, $A \in \mathbf{Op}_{? \cap Y}$,
fong-spivak-seven-sketches_1 F02556	248	■ 0.955	$A \ \text{in} \ \mathbf{0} \ \mathbf{p}_{\{? \ \cap \ Y\}}$	$A \in \mathbf{Op}_{? \cap Y}$	<i>subspace topology</i> , call it $\mathbf{Op}_{? \cap Y}$. This topology defines any $A \subseteq Y$ to be open, $A \in \mathbf{Op}_{? \cap Y}$,
fong-spivak-seven-sketches_1 F02557	248	■ 1.000	$B \ \text{in} \ \mathbf{0} \ \mathbf{p}$	$B \in \mathbf{Op}$	if there is an open set $B \in \mathbf{Op}$ such that $A = B \cap Y$.
fong-spivak-seven-sketches_1 F02558	248	■ 1.000	$A=B \ \cap \ Y$	$A = B \cap Y$	if there is an open set $B \in \mathbf{Op}$ such that $A = B \cap Y$.
fong-spivak-seven-sketches_1 F02559	248	■ 1.000	$Y \ \text{in} \ \mathbf{0} \ \mathbf{p}_{\{? \ \cap \ Y\}}$	$Y \in \mathbf{Op}_{? \cap Y}$	1. Find a $B \in \mathbf{Op}$ that shows that the whole set Y is open, i.e. $Y \in \mathbf{Op}_{? \cap Y}$.
fong-spivak-seven-sketches_1 F02560	248	■ 1.000	$\mathbf{0} \ \mathbf{p}_{\{? \ \cap \ Y\}}$	$\mathbf{Op}_{? \cap Y}$	2. Show that $\mathbf{Op}_{? \cap Y}$ is a topology in the sense of Definition 7.25. ⁸
fong-spivak-seven-sketches_1 F02561	248	■ 1.000	$Y \ \hookrightarrow X$	$Y \hookrightarrow X$	3. Show that the inclusion function $Y \hookrightarrow X$ is a continuous function.
fong-spivak-seven-sketches_1 F02562	248	■ 0.555	$\mathbf{0} \ \mathbf{p}, \ \subseteq$	$\mathbf{Op}, \subseteq$	<i>Remark 7.33.</i> Suppose $(X, \mathbf{Op})$ is a topological space, and consider the preorder $(\mathbf{Op}, \subseteq)$
fong-spivak-seven-sketches_1 F02563	248	■ 0.983	$(\mathbf{0} \ \mathbf{p}, \ \subseteq, \ X, \ \cap)$	$(\mathbf{Op}, \subseteq, X, \cap)$	of open sets. It turns out that $(\mathbf{Op}, \subseteq, X, \cap)$ is always a quantale in the sense of
fong-spivak-seven-sketches_1 F02564	248	■ 0.638	$(X, \ \mathrm{Op})$	(X, Op)	If $(X, \mathbf{Op})$ is a topological space and $\mathcal{V}$ the corresponding quantale as in Remark 7.33,

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-1 F02565	248	■ 0.997	$U_{\{i\}} \subseteq X$	$U_i \subseteq X$	⁸ Hint 1: for any set I , collection of sets $(U_i)_{i \in I}$ with $U_i \subseteq X$, and set $V \subseteq X$, one has $(\bigcup_{i \in I} U_i) \cap V =$
fong-spivak-seven-sketches-1 F02566	248	■ 0.997	$\left(\bigcup_{i \in I} U_{\{i\}}\right) \cap V =$	$\left(\bigcup_{i \in I} U_i\right) \cap V =$	⁸ Hint 1: for any set I , collection of sets $(U_i)_{i \in I}$ with $U_i \subseteq X$, and set $V \subseteq X$, one has $(\bigcup_{i \in I} U_i) \cap V =$
fong-spivak-seven-sketches-1 F02567	248	■ 0.798	$\bigcup_{i \in I} \left(U_{\{i\}} \cap V\right)$	$\bigcup_{i \in I} (U_i \cap V)$	$\bigcup_{i \in I} (U_i \cap V)$. Hint 2: for any $U, V, W \subseteq X$, one has $(U \cap W) \cap (V \cap W) = (U \cap V) \cap W$.
fong-spivak-seven-sketches-1 F02568	248	■ 0.798	$U, V, W \subseteq X$	$U, V, W \subseteq X$	$\bigcup_{i \in I} (U_i \cap V)$. Hint 2: for any $U, V, W \subseteq X$, one has $(U \cap W) \cap (V \cap W) = (U \cap V) \cap W$.
fong-spivak-seven-sketches-1 F02569	248	■ 0.798	$(U \cap W) \cap (V \cap W) = (U \cap V) \cap W$	$(U \cap W) \cap (V \cap W) = (U \cap V) \cap W$	$\bigcup_{i \in I} (U_i \cap V)$. Hint 2: for any $U, V, W \subseteq X$, one has $(U \cap W) \cap (V \cap W) = (U \cap V) \cap W$.
fong-spivak-seven-sketches-1 F02570	249	■ 0.770	$P: \mathbf{0}^{\text{op}} \rightarrow$	$P: \mathbf{Op}^{\text{op}} \rightarrow$	$P: \mathbf{Op}^{\text{op}} \rightarrow \mathbf{Set}$. Thus to every open set $U \in \mathbf{Op}$ we have a set $P(U)$, called the set
fong-spivak-seven-sketches-1 F02571	249	■ 0.770	$P(U)$	$P(U)$	$P: \mathbf{Op}^{\text{op}} \rightarrow \mathbf{Set}$. Thus to every open set $U \in \mathbf{Op}$ we have a set $P(U)$, called the set
fong-spivak-seven-sketches-1 F02572	249	■ 1.000	$V \subseteq U$	$V \subseteq U$	of sections over U , and to every inclusion of open sets $V \subseteq U$ we have a function
fong-spivak-seven-sketches-1 F02573	249	■ 1.000	$P(U) \rightarrow P(V)$	$P(U) \rightarrow P(V)$	$P(U) \rightarrow P(V)$ called the <i>restriction</i> . If $s \in P(U)$ is a section over U , we may denote its
fong-spivak-seven-sketches-1 F02574	249	■ 1.000	$s \in P(U)$	$s \in P(U)$	$P(U) \rightarrow P(V)$ called the <i>restriction</i> . If $s \in P(U)$ is a section over U , we may denote its
fong-spivak-seven-sketches-1 F02575	249	■ 1.000	$s _V$	$s _V$	restriction to V by $s _V$. Recall that we say a collection of open sets $(U_i)_{i \in I}$ covers an open
fong-spivak-seven-sketches-1 F02576	249	■ 0.613	$P: \mathbf{0}^{\text{op}} \rightarrow$	$P: \mathbf{Op}^{\text{op}} \rightarrow$	Definition 7.35. Let $(X, \mathbf{Op})$ be a topological space, and let $P: \mathbf{Op}^{\text{op}} \rightarrow \mathbf{Set}$ be a
fong-spivak-seven-sketches-1 F02577	249	■ 1.000	$U_{\{i\}} \in \mathbf{0}^{\text{op}}$	$U_i \in \mathbf{Op}$	Let $(U_i)_{i \in I}$ be a collection of open sets $U_i \in \mathbf{Op}$ covering U . A <i>matching family</i> $(s_i)_{i \in I}$
fong-spivak-seven-sketches-1 F02578	249	■ 1.000	$(s_{\{i\}})_{i \in I}$	$(s_i)_{i \in I}$	Let $(U_i)_{i \in I}$ be a collection of open sets $U_i \in \mathbf{Op}$ covering U . A <i>matching family</i> $(s_i)_{i \in I}$
fong-spivak-seven-sketches-1 F02579	249	■ 1.000	$s_{\{i\}} \in P(U_{\{i\}})$	$s_i \in P(U_i)$	of P -sections over $(U_i)_{i \in I}$ consists of a section $s_i \in P(U_i)$ for each $i \in I$, such that for every
fong-spivak-seven-sketches-1 F02580	249	■ 1.000	$i \in I$	$i \in I$	of P -sections over $(U_i)_{i \in I}$ consists of a section $s_i \in P(U_i)$ for each $i \in I$, such that for every
fong-spivak-seven-sketches-1 F02581	249	■ 1.000	$i, j \in I$	$i, j \in I$	$i, j \in I$ we have
fong-spivak-seven-sketches-1 F02582	249	■ 1.000	$s _{U_{\{i\}}} = s_{\{i\}}$	$s _{U_i} = s_i$	<i>gluing, or glued section</i> , of the matching family if $s _{U_i} = s_i$ holds for all $i \in I$.
fong-spivak-seven-sketches-1 F02583	249	■ 0.992	$\mathbf{Shv}(X, \mathbf{0}^{\text{op}})$	$\mathbf{Shv}(X, \mathbf{Op})$	We denote by $\mathbf{Shv}(X, \mathbf{Op})$ the category of sheaves on X .
fong-spivak-seven-sketches-1 F02584	249	■ 1.000	$U = \emptyset \subseteq X$	$U = \emptyset \subseteq X$	we take $U = \emptyset \subseteq X$ and $I = \emptyset$, then the empty collection of subsets (one for each $i \in I$,
fong-spivak-seven-sketches-1 F02585	249	■ 0.999	$P(\emptyset) \cong \{()\}$	$P(\emptyset) \cong \{()\}$	rarely sufficient) condition is that $P(\emptyset) \cong \{()\}$, i.e. $P(\emptyset)$ must be a set with one element.
fong-spivak-seven-sketches-1 F02586	249	■ 0.999	$P(\emptyset)$	$P(\emptyset)$	rarely sufficient) condition is that $P(\emptyset) \cong \{()\}$, i.e. $P(\emptyset)$ must be a set with one element.
fong-spivak-seven-sketches-1 F02587	250	■ 0.990	$f(a_1) = f(a_2) = a, f(b_1) = b$	$f(a_1) = f(a_2) = a, f(b_1) = b$	the element of Y immediately below it. For example, $f(a_1) = f(a_2) = a, f(b_1) = b$, and
fong-spivak-seven-sketches-1 F02588	250	■ 0.999	$f^{-1}(y) \subseteq X$	$f^{-1}(y) \subseteq X$	For each point $y \in Y$, the preimage set $f^{-1}(y) \subseteq X$ above it is often called the <i>fiber over</i>
fong-spivak-seven-sketches-1 F02589	250	■ 1.000	$f': X \rightarrow Y$	$f': X \rightarrow Y$	4. Gave an example of a function $f': X \rightarrow Y$ for which every fiber has either one or
fong-spivak-seven-sketches-1 F02590	250	■ 0.886	$\mathbf{Sec}_f: \mathbf{0}^{\text{op}}(Y)^{\text{op}} \rightarrow$	$\mathbf{Sec}_f: \mathbf{Op}(Y)^{\text{op}} \rightarrow$	we begin by building a presheaf—i.e. a functor $\mathbf{Sec}_f: \mathbf{Op}(Y)^{\text{op}} \rightarrow \mathbf{Set}$ —and then we'll
fong-spivak-seven-sketches-1 F02591	250	■ 1.000	$\mathbf{Sec}_f$	$\mathbf{Sec}_f$	Define the presheaf $\mathbf{Sec}_f$ on an arbitrary subset $U \subseteq Y$ by:

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02592	250	■ 1.000	$U \subseteq Y$	$U \subseteq Y$	Define the presheaf Sec_f on an arbitrary subset $U \subseteq Y$ by:
fong-spivak-seven-sketches-F02593	250	■ 1.000	$\operatorname{Sec}_f(U)$	$\mathrm{Sec}_f(U)$	One might describe $\mathrm{Sec}_f(U)$ as the set of all ways to pick a ‘cross-section’ of the f
fong-spivak-seven-sketches-F02594	250	■ 1.000	$s \in \operatorname{Sec}_f(U)$	$s \in \mathrm{Sec}_f(U)$	arrangement over U . That is, an element $s \in \mathrm{Sec}_f(U)$ is a choice of one element per
fong-spivak-seven-sketches-F02595	250	■ 0.999	$U = \{a, b\}$	$U = \{a, b\}$	As an example, let’s say $U = \{a, b\}$. How many such s ’s are there in $\mathrm{Sec}_f(U)$? To
fong-spivak-seven-sketches-F02596	251	■ 0.996	$V_1 = \{a, b, c\}$	$V_1 = \{a, b, c\}$	1. Let $V_1 = \{a, b, c\}$. Draw all the sections over it, i.e. all elements of $\mathrm{Sec}_f(V_1)$, as
fong-spivak-seven-sketches-F02597	251	■ 0.996	$\operatorname{Sec}_f(V_1)$	$\mathrm{Sec}_f(V_1)$	1. Let $V_1 = \{a, b, c\}$. Draw all the sections over it, i.e. all elements of $\mathrm{Sec}_f(V_1)$, as
fong-spivak-seven-sketches-F02598	251	■ 1.000	$V_2 = \{a, b, c, d\}$	$V_2 = \{a, b, c, d\}$	2. Let $V_2 = \{a, b, c, d\}$. Again draw all the sections, $\mathrm{Sec}_f(V_2)$.
fong-spivak-seven-sketches-F02599	251	■ 1.000	$\operatorname{Sec}_f(V_2)$	$\mathrm{Sec}_f(V_2)$	2. Let $V_2 = \{a, b, c, d\}$. Again draw all the sections, $\mathrm{Sec}_f(V_2)$.
fong-spivak-seven-sketches-F02600	251	■ 0.999	$V_3 = \{a, b, d, e\}$	$V_3 = \{a, b, d, e\}$	3. Let $V_3 = \{a, b, d, e\}$. How many sections (elements of $\mathrm{Sec}_f(V_3)$) are there?
fong-spivak-seven-sketches-F02601	251	■ 0.999	$\operatorname{Sec}_f(V_3)$	$\mathrm{Sec}_f(V_3)$	3. Let $V_3 = \{a, b, d, e\}$. How many sections (elements of $\mathrm{Sec}_f(V_3)$) are there?
fong-spivak-seven-sketches-F02602	251	■ 0.974	Sec_f	Sec_f	Sec_f on objects, so you are half way to understanding Sec_f as a presheaf. That is, as
fong-spivak-seven-sketches-F02603	251	■ 1.000	$V = \{a\}$	$V = \{a\}$	restriction maps are easy: if $V \subseteq U$, say $V = \{a\}$ and $U = \{a, b\}$, then given a section s
fong-spivak-seven-sketches-F02604	251	■ 1.000	$\operatorname{Sec}_f(\{a, b, c\})$	$\mathrm{Sec}_f(\{a, b, c\})$	1. Write out the sets of sections $\mathrm{Sec}_f(\{a, b, c\})$ and $\mathrm{Sec}_f(\{a, c\})$.
fong-spivak-seven-sketches-F02605	251	■ 1.000	$\operatorname{Sec}_f(\{a, c\})$	$\mathrm{Sec}_f(\{a, c\})$	1. Write out the sets of sections $\mathrm{Sec}_f(\{a, b, c\})$ and $\mathrm{Sec}_f(\{a, c\})$.
fong-spivak-seven-sketches-F02606	251	■ 1.000	$U_1 = \{a, b\}$	$U_1 = \{a, b\}$	sheaf condition, consider the set $U_1 = \{a, b\}$ and $U_2 = \{b, e\}$. These cover the set
fong-spivak-seven-sketches-F02607	251	■ 1.000	$U_2 = \{b, e\}$	$U_2 = \{b, e\}$	sheaf condition, consider the set $U_1 = \{a, b\}$ and $U_2 = \{b, e\}$. These cover the set
fong-spivak-seven-sketches-F02608	251	■ 1.000	$U = \{a, b, e\} = U_1 \cup U_2$	$U = \{a, b, e\} = U_1 \cup U_2$	$U = \{a, b, e\} = U_1 \cup U_2$. By Definition 7.35, a matching family for this cover consists of
fong-spivak-seven-sketches-F02609	251	■ 1.000	U_1	U_1	a section over U_1 and a section over U_2 that agree on the overlap set, $U_1 \cap U_2 = \{b\}$.
fong-spivak-seven-sketches-F02610	251	■ 1.000	U_2	U_2	a section over U_1 and a section over U_2 that agree on the overlap set, $U_1 \cap U_2 = \{b\}$.
fong-spivak-seven-sketches-F02611	251	■ 1.000	$U_1 \cap U_2 = \{b\}$	$U_1 \cap U_2 = \{b\}$	a section over U_1 and a section over U_2 that agree on the overlap set, $U_1 \cap U_2 = \{b\}$.
fong-spivak-seven-sketches-F02612	252	■ 1.000	$s_1 \in \operatorname{Sec}_f(U_1)$	$s_1 \in \mathrm{Sec}_f(U_1)$	So consider $s_1 \in \mathrm{Sec}_f(U_1)$ and $s_2 \in \mathrm{Sec}_f(U_2)$ shown below.
fong-spivak-seven-sketches-F02613	252	■ 1.000	$s_2 \in \operatorname{Sec}_f(U_2)$	$s_2 \in \mathrm{Sec}_f(U_2)$	So consider $s_1 \in \mathrm{Sec}_f(U_1)$ and $s_2 \in \mathrm{Sec}_f(U_2)$ shown below.
fong-spivak-seven-sketches-F02614	252	■ 1.000	g_1	g_1	Since sections g_1 and g_2 agree on the overlap—they both send b to b_2 —the two sections
fong-spivak-seven-sketches-F02615	252	■ 1.000	g_2	g_2	Since sections g_1 and g_2 agree on the overlap—they both send b to b_2 —the two sections
fong-spivak-seven-sketches-F02616	252	■ 1.000	b_2	b_2	Since sections g_1 and g_2 agree on the overlap—they both send b to b_2 —the two sections
fong-spivak-seven-sketches-F02617	252	■ 1.000	$U = \{a, b, e\}$	$U = \{a, b, e\}$	shown in Eq. (7.43) can be glued to form a single section over $U = \{a, b, e\}$:
fong-spivak-seven-sketches-F02618	252	■ 0.968	(s_1, s_2)	(s_1, s_2)	2. For your answer (s_1, s_2) in part 1, can you find a section $s \in \mathrm{Sec}_f(U_1 \cup U_2)$ such

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02619	252	■ 0.968	$s \in \operatorname{Sec}_f(U_1 \cup U_2)$	$s \in \operatorname{Sec}_f(U_1 \cup U_2)$	2. For your answer (s_1, s_2) in part 1, can you find a section $s \in \operatorname{Sec}_f(U_1 \cup U_2)$ such
fong-spivak-seven-sketches-F02620	252	■ 1.000	$\left.s\right _{U_1}=s_1$	$s _{U_1} = s_1$	that $s _{U_1} = s_1$ and $s _{U_2} = s_2$?
fong-spivak-seven-sketches-F02621	252	■ 1.000	$\left.s\right _{U_2}=s_2$	$s _{U_2} = s_2$	that $s _{U_1} = s_1$ and $s _{U_2} = s_2$?
fong-spivak-seven-sketches-F02622	252	■ 0.683	$h_1 \in \operatorname{Sec}_f(U_1)$	$h_1 \in \operatorname{Sec}_f(U_1)$	3. Find a section $h_1 \in \operatorname{Sec}_f(U_1)$ and a section $h_2 \in \operatorname{Sec}_f(U_2)$ that <i>do</i> agree on the
fong-spivak-seven-sketches-F02623	252	■ 0.683	$h_2 \in \operatorname{Sec}_f(U_2)$	$h_2 \in \operatorname{Sec}_f(U_2)$	3. Find a section $h_1 \in \operatorname{Sec}_f(U_1)$ and a section $h_2 \in \operatorname{Sec}_f(U_2)$ that <i>do</i> agree on the
fong-spivak-seven-sketches-F02624	252	■ 0.683	do	<i>do</i>	3. Find a section $h_1 \in \operatorname{Sec}_f(U_1)$ and a section $h_2 \in \operatorname{Sec}_f(U_2)$ that <i>do</i> agree on the
fong-spivak-seven-sketches-F02625	252	■ 0.975	$h \in \operatorname{Sec}_f(U_1 \cup U_2)$	$h \in \operatorname{Sec}_f(U_1 \cup U_2)$	4. Can you find a section $h \in \operatorname{Sec}_f(U_1 \cup U_2)$ such that $h _{U_1} = h_1$ and $h _{U_2} = h_2$? $\diamond$
fong-spivak-seven-sketches-F02626	252	■ 0.975	$\left.h\right _{U_1}=h_1$	$h _{U_1} = h_1$	4. Can you find a section $h \in \operatorname{Sec}_f(U_1 \cup U_2)$ such that $h _{U_1} = h_1$ and $h _{U_2} = h_2$? $\diamond$
fong-spivak-seven-sketches-F02627	252	■ 0.975	$\left.h\right _{U_2}=h_2$	$h _{U_2} = h_2$	4. Can you find a section $h \in \operatorname{Sec}_f(U_1 \cup U_2)$ such that $h _{U_1} = h_1$ and $h _{U_2} = h_2$? $\diamond$
fong-spivak-seven-sketches-F02628	252	■ 0.984	$f: \left(X, \operatorname{\mathbf{Op}}_X\right) \rightarrow \left(Y, \operatorname{\mathbf{Op}}_Y\right)$	$f: (X, \mathbf{Op}_X) \rightarrow (Y, \mathbf{Op}_Y)$	<i>Example 7.45.</i> Let $f: (X, \mathbf{Op}_X) \rightarrow (Y, \mathbf{Op}_Y)$ be a continuous function. Consider the
fong-spivak-seven-sketches-F02629	252	■ 0.901	$\operatorname{Sec}_f: \operatorname{\mathbf{Op}}_Y^{\operatorname{\text{op}}} \rightarrow$	$\operatorname{Sec}_f: \mathbf{Op}_Y^{\operatorname{\text{op}}} \rightarrow$	functor $\operatorname{Sec}_f: \mathbf{Op}_Y^{\operatorname{\text{op}}} \rightarrow$ Set given by
fong-spivak-seven-sketches-F02630	253	■ 0.623	$\operatorname{\mathbf{Op}}_Y$	$\mathbf{Op}_Y$	The morphisms of $\mathbf{Op}_Y$ are inclusions $V \subseteq U$. Given $g: U \rightarrow X$ and $V \subseteq U$, what we
fong-spivak-seven-sketches-F02631	253	■ 0.623	$g: U \rightarrow X$	$g: U \rightarrow X$	The morphisms of $\mathbf{Op}_Y$ are inclusions $V \subseteq U$. Given $g: U \rightarrow X$ and $V \subseteq U$, what we
fong-spivak-seven-sketches-F02632	253	■ 0.996	TM	TM	space TM whose points are pairs (m, v) , where $m \in M$ is a point in the manifold and
fong-spivak-seven-sketches-F02633	253	■ 0.996	(m, v)	(m, v)	space TM whose points are pairs (m, v) , where $m \in M$ is a point in the manifold and
fong-spivak-seven-sketches-F02634	253	■ 0.996	$m \in M$	$m \in M$	space TM whose points are pairs (m, v) , where $m \in M$ is a point in the manifold and
fong-spivak-seven-sketches-F02635	253	■ 1.000	$\pi: TM \rightarrow$	$\pi: TM \rightarrow$	The tangent bundle TM on a manifold M comes with a continuous map $\pi: TM \rightarrow$
fong-spivak-seven-sketches-F02636	253	■ 0.984	$(m, v) \mapsto m$	$(m, v) \mapsto m$	M back down to the manifold, sending $(m, v) \mapsto m$. One might say that π “forgets the
fong-spivak-seven-sketches-F02637	253	■ 0.984	π	π	M back down to the manifold, sending $(m, v) \mapsto m$. One might say that π “forgets the
fong-spivak-seven-sketches-F02638	253	■ 0.960	Sec_π	Sec_π	defines a sheaf Sec_π . It could be called the sheaf of ‘tangent vector sections on M' ’, but
fong-spivak-seven-sketches-F02639	253	■ 0.960	M'	M'	defines a sheaf Sec_π . It could be called the sheaf of ‘tangent vector sections on M' ’, but
fong-spivak-seven-sketches-F02640	253	■ 1.000	$U \subseteq M$	$U \subseteq M$	we spoke of the sheaf of wind velocities on Earth, above. Given an open subset $U \subseteq M$,
fong-spivak-seven-sketches-F02641	253	■ 1.000	$v \in \operatorname{Sec}_\pi(U)$	$v \in \operatorname{Sec}_\pi(U)$	an element $v \in \operatorname{Sec}_\pi(U)$ is called a vector field over U because it continuously assigns
fong-spivak-seven-sketches-F02642	253	■ 1.000	$v(u)$	$v(u)$	a tangent vector $v(u)$ to each point $u \in U$. The tangent vector at u tells us the velocity
fong-spivak-seven-sketches-F02643	253	■ 1.000	$U \neq M$	$U \neq M$	possible to prove that for every open set U , as long as $U \neq M$, there is a vector field
fong-spivak-seven-sketches-F02644	254	■ 1.000	$U \neq X$	$U \neq X$	namely that hair can be well combed whenever $U \neq X$ but cannot be well combed when
fong-spivak-seven-sketches-F02645	254	■ 1.000	$U=X$	$U = X$	$U = X$ —can be thought of as a generative effect, and can be measured by cohomology

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02646	254	■ 0.988	$\boldsymbol{\operatorname{Shv}}(M)$	$\mathbf{Shv}(M)$	that we can consider the category $\mathbf{Shv}(M)$ of sheaves on M ; in fact, such categories are
fong-spivak-seven-sketches-F02647	254	■ 0.485	X, Op	$X, \mathbf{Op}$	<i>Example 7.48.</i> For every topological space $(X, \mathbf{Op})$, we have the topos of sheaves on it.
fong-spivak-seven-sketches-F02648	254	■ 0.928	$\{1, 2\}, \mathbf{Op}_1$	$\{1, 2\}, \mathbf{Op}_1$	<i>Exercise 7.49.</i> Consider the Sierpinski space $(\{1, 2\}, \mathbf{Op}_1)$ from Example 7.30.
fong-spivak-seven-sketches-F02649	254	■ 0.896	$X = \mathbf{1}$	$X = \mathbf{1}$	one-point space $X = \mathbf{1}$ with its unique topology, we find that the category $\mathbf{Set}$ is a topos,
fong-spivak-seven-sketches-F02650	255	■ 1.000	$\wedge, \vee, \neg, \rightarrow, \exists, \forall$	$\wedge, \vee, \neg, \Rightarrow, \exists, \forall$	topological space. There, the same logical symbols $\wedge, \vee, \neg, \Rightarrow, \exists, \forall$ become operations
fong-spivak-seven-sketches-F02651	255	■ 0.975	$\Omega = \mathbb{B}$	$\Omega = \mathbb{B}$	of booleans $\Omega = \mathbb{B}$. In a sheaf topos $\mathcal{E} = \mathbf{Shv}(X, \mathbf{Op})$, the object $\Omega \in \mathcal{E}$ is a sheaf, not
fong-spivak-seven-sketches-F02652	255	■ 0.975	$\mathcal{E} = \mathbf{Shv}(X, \mathbf{Op})$	$\mathcal{E} = \mathbf{Shv}(X, \mathbf{Op})$	of booleans $\Omega = \mathbb{B}$. In a sheaf topos $\mathcal{E} = \mathbf{Shv}(X, \mathbf{Op})$, the object $\Omega \in \mathcal{E}$ is a sheaf, not
fong-spivak-seven-sketches-F02653	255	■ 0.854	$\Omega : \mathbf{Op}^{\operatorname{op}} \rightarrow \mathbf{Set}$	$\Omega : \mathbf{Op}^{\operatorname{op}} \rightarrow$	$\Omega : \mathbf{Op}^{\operatorname{op}} \rightarrow \mathbf{Set}$; it assigns a set $\Omega(U)$ to each open $U \subseteq X$ and comes with a restriction
fong-spivak-seven-sketches-F02654	255	■ 0.854	$\Omega(U)$	$\Omega(U)$	$\Omega : \mathbf{Op}^{\operatorname{op}} \rightarrow \mathbf{Set}$; it assigns a set $\Omega(U)$ to each open $U \subseteq X$ and comes with a restriction
fong-spivak-seven-sketches-F02655	255	■ 1.000	$\Omega(U) \rightarrow \Omega(V)$	$\Omega(U) \rightarrow \Omega(V)$	map $\Omega(U) \rightarrow \Omega(V)$ whenever $V \subseteq U$. So in our quest to understand Ω , we first ask the
fong-spivak-seven-sketches-F02656	256	■ 1.000	$X = \{1\}$	$X = \{1\}$	<i>Exercise 7.52.</i> Let $X = \{1\}$ be the one point space. We said above that its subobject
fong-spivak-seven-sketches-F02657	256	■ 1.000	$W \subseteq V \subseteq U$	$W \subseteq V \subseteq U$	that whenever $W \subseteq V \subseteq U$, the restriction map $\Omega(U) \rightarrow \Omega(V)$ followed by the
fong-spivak-seven-sketches-F02658	256	■ 1.000	$\Omega(V) \rightarrow \Omega(W)$	$\Omega(V) \rightarrow \Omega(W)$	restriction map $\Omega(V) \rightarrow \Omega(W)$ is the same as the restriction map $\Omega(U) \rightarrow \Omega(W)$.
fong-spivak-seven-sketches-F02659	256	■ 1.000	$\Omega(U) \rightarrow \Omega(W)$	$\Omega(U) \rightarrow \Omega(W)$	restriction map $\Omega(V) \rightarrow \Omega(W)$ is the same as the restriction map $\Omega(U) \rightarrow \Omega(W)$.
fong-spivak-seven-sketches-F02660	256	■ 1.000	$V_i \in \Omega(U_i)$	$V_i \in \Omega(U_i)$	suppose that we have a cover $U = \bigcup_{i \in I} U_i$, and suppose given an element $V_i \in \Omega(U_i)$,
fong-spivak-seven-sketches-F02661	256	■ 1.000	$V_i \subseteq U_i$	$V_i \subseteq U_i$	i.e. an open set $V_i \subseteq U_i$, for each $i \in I$. Suppose further that for all $i, j \in I$, it is the
fong-spivak-seven-sketches-F02662	256	■ 1.000	$V_i \cap U_j = V_j \cap U_i$	$V_i \cap U_j = V_j \cap U_i$	case that $V_i \cap U_j = V_j \cap U_i$, i.e. that the elements form a matching family. Define
fong-spivak-seven-sketches-F02663	256	■ 0.995	$V := \bigcup_{i \in I} V_i$	$V := \bigcup_{i \in I} V_i$	$V := \bigcup_{i \in I} V_i$; it is an open subset of U , so we can consider V as an element of $\Omega(U)$.
fong-spivak-seven-sketches-F02664	256	■ 1.000	$(V_i)_{i \in I}$	$(V_i)_{i \in I}$	The following verifies that V is indeed a gluing for the $(V_i)_{i \in I}$:
fong-spivak-seven-sketches-F02665	256	■ 1.000	$V \cap U_j = V_j$	$V \cap U_j = V_j$	In other words $V \cap U_j = V_j$ for any $j \in I$. So our Ω has been upgraded from presheaf
fong-spivak-seven-sketches-F02666	256	■ 1.000	$j \in I$	$j \in I$	In other words $V \cap U_j = V_j$ for any $j \in I$. So our Ω has been upgraded from presheaf
fong-spivak-seven-sketches-F02667	256	■ 1.000	$\{1\} \rightarrow \Omega$	$\{1\} \rightarrow \Omega$	in good standing, we also need to give a sheaf morphism $\text{true}: \{1\} \rightarrow \Omega$. Here
fong-spivak-seven-sketches-F02668	256	■ 0.623	$\{1\} : \mathbf{Op}^{\operatorname{op}} \rightarrow \mathbf{Set}$	$\{1\} : \mathbf{Op}^{\operatorname{op}} \rightarrow$	$\{1\} : \mathbf{Op}^{\operatorname{op}} \rightarrow \mathbf{Set}$ is the terminal sheaf; it maps every open set to the terminal, one
fong-spivak-seven-sketches-F02669	256	■ 1.000	$\{1\} = \{1\}(U) \rightarrow \Omega(U)$	$\{1\} = \{1\}(U) \rightarrow \Omega(U)$	the sheaf morphism that assigns, for every $U \in \mathbf{Op}$ the function $\{1\} = \{1\}(U) \rightarrow \Omega(U)$
fong-spivak-seven-sketches-F02670	256	■ 0.997	$1 \mapsto U$	$1 \mapsto U$	sending $1 \mapsto U$, the largest open set $U \subseteq U$. From now on we denote $\{1\}$ simply as 1.
fong-spivak-seven-sketches-F02671	256	■ 1.000	$U = \emptyset$	$U = \emptyset$	If this U is everything, $U = X$, then P is really true; if U is nothing, $U = \emptyset$, then P is
fong-spivak-seven-sketches-F02672	257	■ 1.000	$(V, V; A)$	$(V, V; A)$	$\text{true}: 1 \rightarrow \Omega$ sends that loop to $(V, V; A)$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02673	257	1.000	$i: H \subseteq G$	$i: H \subseteq G$	Given any graph G and subgraph $i: H \subseteq G$, we need to construct a graph homo-
fong-spivak-seven-sketches-F02674	257	0.931	$\ulcorner H \urcorner$	$\ulcorner H \urcorner$	morphism $\ulcorner H \urcorner: G \rightarrow \Omega$ classifying H . The idea is that for each part of G , we decide
fong-spivak-seven-sketches-F02675	257	0.931	$G \rightarrow \Omega$	$G \rightarrow \Omega$	morphism $\ulcorner H \urcorner: G \rightarrow \Omega$ classifying H . The idea is that for each part of G , we decide
fong-spivak-seven-sketches-F02676	257	1.000	$G^{\prime \prime}$	G''	is its source in H ? is its target in G'' ? both? neither? Based on the answers to these
fong-spivak-seven-sketches-F02677	257	0.589	$(V, \emptyset; \emptyset), (\emptyset, V; \emptyset), (V, V; \emptyset)$	$(V, 0; 0), (0, V; 0), (V, V; 0)$	questions we send a to $(V, 0; 0), (0, V; 0), (V, V; 0)$, or $(0, 0; 0)$, respectively.
fong-spivak-seven-sketches-F02678	257	0.589	$\emptyset; \emptyset$	$0; 0$	questions we send a to $(V, 0; 0), (0, V; 0), (V, V; 0)$, or $(0, 0; 0)$, respectively.
fong-spivak-seven-sketches-F02679	257	1.000	$H \subseteq G$	$H \subseteq G$	<i>Exercise 7.55.</i> Consider the subgraph $H \subseteq G$ shown here:
fong-spivak-seven-sketches-F02680	257	0.907	$\ulcorner H \urcorner: G \rightarrow \Omega$	$\ulcorner H \urcorner: G \rightarrow \Omega$	Find the graph homomorphism $\ulcorner H \urcorner: G \rightarrow \Omega$ classifying it. See Example 7.54.
fong-spivak-seven-sketches-F02681	257	0.615	$U, V \in \Omega(X)$	$U, V \in \Omega(X)$	values $U, V \in \Omega(X)$, then their conjunction ‘ U AND V ’ is their intersection, and their
fong-spivak-seven-sketches-F02682	257	1.000	$\cap$	$\cap$	These formulas are easy to remember, because $\wedge$ looks like $\cap$ and $\vee$ looks like $\cup$. The
fong-spivak-seven-sketches-F02683	257	1.000	$\cup$	$\cup$	These formulas are easy to remember, because $\wedge$ looks like $\cap$ and $\vee$ looks like $\cup$. The
fong-spivak-seven-sketches-F02684	257	1.000	$U \Rightarrow V$	$U \Rightarrow V$	implication $U \Rightarrow V$ is the largest open set R such that $R \cap U \subseteq V$, i.e.
fong-spivak-seven-sketches-F02685	257	1.000	$R \cap U \subseteq V$	$R \cap U \subseteq V$	implication $U \Rightarrow V$ is the largest open set R such that $R \cap U \subseteq V$, i.e.
fong-spivak-seven-sketches-F02686	257	0.668	$\neg U :=$	$\neg U :=$	Finally, the negation of U is given by $\neg U := (U \Rightarrow \text{false})$, and this turns out
fong-spivak-seven-sketches-F02687	257	0.668	$U \Rightarrow$	$U \Rightarrow$	Finally, the negation of U is given by $\neg U := (U \Rightarrow \text{false})$, and this turns out
fong-spivak-seven-sketches-F02688	257	1.000	$R \cap U = \varnothing$	$R \cap U = \emptyset$	$R \cap U = \emptyset$, i.e. the union of all open sets in the complement of U . If you know topology,
fong-spivak-seven-sketches-F02689	257	1.000	$\neg U$	$\neg U$	you might recognize that $\neg U$ is the ‘interior of the complement of U .’
fong-spivak-seven-sketches-F02690	258	1.000	$X = \mathbb{R}$	$X = \mathbb{R}$	<i>Example 7.58.</i> Consider the real line $X = \mathbb{R}$ as a topological space (see Exercise 7.27).
fong-spivak-seven-sketches-F02691	258	1.000	$U = \{x \in \mathbb{R} \mid x < 3\}$	$U = \{x \in \mathbb{R} \mid x < 3\}$	Let $U, V \in \Omega(X)$ be the open sets $U = \{x \in \mathbb{R} \mid x < 3\}$ and $V = \{x \in \mathbb{R} \mid -4 < x < 4\}$.
fong-spivak-seven-sketches-F02692	258	1.000	$V = \{x \in \mathbb{R} \mid -4 < x < 4\}$	$V = \{x \in \mathbb{R} \mid -4 < x < 4\}$	Let $U, V \in \Omega(X)$ be the open sets $U = \{x \in \mathbb{R} \mid x < 3\}$ and $V = \{x \in \mathbb{R} \mid -4 < x < 4\}$.
fong-spivak-seven-sketches-F02693	258	1.000	$U = (-\infty, 3)$	$U = (-\infty, 3)$	Using interval notation, $U = (-\infty, 3)$ and $V = (-4, 4)$. Then
fong-spivak-seven-sketches-F02694	258	1.000	$V = (-4, 4)$	$V = (-4, 4)$	Using interval notation, $U = (-\infty, 3)$ and $V = (-4, 4)$. Then
fong-spivak-seven-sketches-F02695	258	1.000	$U \wedge V = (-4, 3)$	$U \wedge V = (-4, 3)$	• $U \wedge V = (-4, 3)$.
fong-spivak-seven-sketches-F02696	258	1.000	$U \vee V = (-\infty, 4)$	$U \vee V = (-\infty, 4)$	• $U \vee V = (-\infty, 4)$.
fong-spivak-seven-sketches-F02697	258	1.000	$\neg U = (3, \infty)$	$\neg U = (3, \infty)$	• $\neg U = (3, \infty)$.
fong-spivak-seven-sketches-F02698	258	1.000	$\neg V = (-\infty, -4) \cup (4, \infty)$	$\neg V = (-\infty, -4) \cup (4, \infty)$	• $\neg V = (-\infty, -4) \cup (4, \infty)$.
fong-spivak-seven-sketches-F02699	258	0.999	$(U \Rightarrow V) = (-4, \infty)$	$(U \Rightarrow V) = (-4, \infty)$	• $(U \Rightarrow V) = (-4, \infty)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02700	258	■ 1.000	$(V \Rightarrow U) = U$	$(V \Rightarrow U) = U$	• $(V \Rightarrow U) = U$
fong-spivak-seven-sketches-F02701	258	■ 1.000	$U = \mathbb{R} - \{0\}$	$U = \mathbb{R} - \{0\}$	subset $U = \mathbb{R} - \{0\}$.
fong-spivak-seven-sketches-F02702	258	■ 1.000	$\neg \neg U$	$\neg \neg U$	2. What open subset is $\neg \neg U$?
fong-spivak-seven-sketches-F02703	258	■ 1.000	$U \subseteq \neg \neg U$	$U \subseteq \neg \neg U$	3. Is it true that $U \subseteq \neg \neg U$?
fong-spivak-seven-sketches-F02704	258	■ 1.000	$\neg \neg U \subseteq U$	$\neg \neg U \subseteq U$	4. Is it true that $\neg \neg U \subseteq U$?
fong-spivak-seven-sketches-F02705	258	■ 0.942	$\wedge U = U$	$\wedge U = U$	$U \in \mathbf{Op}$, we have $(\text{true} \wedge U) = U$. Which open set is it?
fong-spivak-seven-sketches-F02706	258	■ 0.877	$\vee U =$	$\vee U =$	2. Other things we should expect from true include $(\text{true} \vee U) = \text{true}$ and $(U \Rightarrow$
fong-spivak-seven-sketches-F02707	258	■ 0.877	$(U \Rightarrow$	$(U \Rightarrow$	2. Other things we should expect from true include $(\text{true} \vee U) = \text{true}$ and $(U \Rightarrow$
fong-spivak-seven-sketches-F02708	258	■ 0.559	$) =$	$) =$	$\text{true}) = \text{true}$ and $(\text{true} \Rightarrow U) = U$. Do these hold for your answer to 1?
fong-spivak-seven-sketches-F02709	258	■ 0.559	$\Rightarrow U) = U$	$\Rightarrow U) = U$	$\text{true}) = \text{true}$ and $(\text{true} \Rightarrow U) = U$. Do these hold for your answer to 1?
fong-spivak-seven-sketches-F02710	258	■ 0.974	$\vee U = U$	$\vee U = U$	$U \in \mathbf{Op}$, we have $(\text{false} \vee U) = U$. Which open set is it?
fong-spivak-seven-sketches-F02711	258	■ 1.000	$\wedge U$	$\wedge U$	4. Other things we should expect from false include $(\text{false} \wedge U) = \text{false}$ and
fong-spivak-seven-sketches-F02712	258	■ 1.000	$\pi: E \rightarrow X$	$\pi: E \rightarrow X$	<i>Example 7.61.</i> For a vector bundle $\pi: E \rightarrow X$ over a space X , the corresponding sheaf
fong-spivak-seven-sketches-F02713	258	■ 0.978	Sec_π	Sec_π	is Sec_π corresponding to its sections: to each open set $i_U: U \subseteq X$, we associate the set
fong-spivak-seven-sketches-F02714	258	■ 0.978	$i_U: U \subseteq X$	$i_U: U \subseteq X$	is Sec_π corresponding to its sections: to each open set $i_U: U \subseteq X$, we associate the set
fong-spivak-seven-sketches-F02715	258	■ 0.999	$s: U \rightarrow E$	$s: U \rightarrow E$	of functions $s: U \rightarrow E$ for which $s \circ \pi = i_U$. For example, in the case of the tangent
fong-spivak-seven-sketches-F02716	258	■ 0.999	$s; \pi = i_U$	$s; \pi = i_U$	of functions $s: U \rightarrow E$ for which $s \circ \pi = i_U$. For example, in the case of the tangent
fong-spivak-seven-sketches-F02717	258	■ 1.000	$\pi: TM \rightarrow M$	$\pi: TM \rightarrow M$	bundle $\pi: TM \rightarrow M$ (see Example 7.46), the corresponding sheaf, call it $\mathbf{VF}$, associates
fong-spivak-seven-sketches-F02718	258	■ 0.991	$\mathbf{VF}(U)$	$\mathbf{VF}(U)$	to each U the set $\mathbf{VF}(U)$ of vector fields on U .
fong-spivak-seven-sketches-F02719	258	■ 0.996	$\mathbf{VF} \rightarrow \Omega$	$\mathbf{VF} \rightarrow \Omega$	fields. For example, one could have a predicate $\text{Grad}: \mathbf{VF} \rightarrow \Omega$ that asks for the largest
fong-spivak-seven-sketches-F02720	258	■ 0.991	$\text{Grad}(v)$	$\text{Grad}(v)$	subspace $\text{Grad}(v)$ on which a given vector field v comes from the gradient of some
fong-spivak-seven-sketches-F02721	259	■ 0.585	$p: S \rightarrow \Omega$	$p: S \rightarrow \Omega$	In a topos $\mathcal{E} = \mathbf{Shv}(X, \mathbf{Op})$, a predicate is a sheaf morphism $p: S \rightarrow \Omega$ where $S \in \mathcal{E}$ is
fong-spivak-seven-sketches-F02722	259	■ 0.585	$S \in \mathcal{E}$	$S \in \mathcal{E}$	In a topos $\mathcal{E} = \mathbf{Shv}(X, \mathbf{Op})$, a predicate is a sheaf morphism $p: S \rightarrow \Omega$ where $S \in \mathcal{E}$ is
fong-spivak-seven-sketches-F02723	259	■ 1.000	$p(U): S(U) \rightarrow \Omega(U)$	$p(U): S(U) \rightarrow \Omega(U)$	we get a function $p(U): S(U) \rightarrow \Omega(U)$ for any open set $U \subseteq X$. In the above example—
fong-spivak-seven-sketches-F02724	259	■ 0.972	$\text{Bob} \in S(U)$	$\text{Bob} \in S(U)$	come and go over time), and $\text{Bob} \in S(U)$ is a person existing over a time U , and p is the
fong-spivak-seven-sketches-F02725	259	■ 0.987	$p(\text{Bob})$	$p(\text{Bob})$	predicate “likes the weather,” then $p(\text{Bob})$ is the set of times during which Bob likes
fong-spivak-seven-sketches-F02726	259	■ 0.979	$s \in S(U)$	$s \in S(U)$	weather,” acts on sections $s \in S(U)$, say $s = \text{Bob}$. But by Definition 7.12, any predicate

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02727	259	■ 0.979	$S =$	$s =$	weather,” acts on sections $s \in S(U)$, say $s = \text{Bob}$. But by Definition 7.12, any predicate
fong-spivak-seven-sketches-F02728	259	■ 1.000	$\{S \mid p\} \subseteq S$	$\{S \mid p\} \subseteq S$	$p: S \rightarrow \Omega$ also defines a subobject of $\{S \mid p\} \subseteq S$. Describe the sections of this subsheaf.
fong-spivak-seven-sketches-F02729	259	■ 1.000	$ \Omega^E = \mathcal{E}(S, \Omega)$	$ \Omega^E = \mathcal{E}(S, \Omega)$	set of S -predicates $ \Omega^E = \mathcal{E}(S, \Omega)$ is naturally given the structure of a poset, which we
fong-spivak-seven-sketches-F02730	259	■ 0.917	$p, q: S \rightarrow \Omega$	$p, q: S \rightarrow \Omega$	Given two predicates $p, q: S \rightarrow \Omega$, we say that $p \leq^S q$ if the first implies the second.
fong-spivak-seven-sketches-F02731	259	■ 0.917	$p \leq^S q$	$p \leq^S q$	Given two predicates $p, q: S \rightarrow \Omega$, we say that $p \leq^S q$ if the first implies the second.
fong-spivak-seven-sketches-F02732	259	■ 0.997	$p(s) \subseteq U$	$p(s) \subseteq U$	$p(s) \subseteq U$ and $q(s) \subseteq U$. We say that $p \leq^S q$ if $p(s) \subseteq q(s)$ for all $U \in \mathbf{Op}$ and $s \in S(U)$.
fong-spivak-seven-sketches-F02733	259	■ 0.997	$q(s) \subseteq U$	$q(s) \subseteq U$	$p(s) \subseteq U$ and $q(s) \subseteq U$. We say that $p \leq^S q$ if $p(s) \subseteq q(s)$ for all $U \in \mathbf{Op}$ and $s \in S(U)$.
fong-spivak-seven-sketches-F02734	259	■ 0.997	$p(s) \subseteq q(s)$	$p(s) \subseteq q(s)$	$p(s) \subseteq U$ and $q(s) \subseteq U$. We say that $p \leq^S q$ if $p(s) \subseteq q(s)$ for all $U \in \mathbf{Op}$ and $s \in S(U)$.
fong-spivak-seven-sketches-F02735	259	■ 0.999	$\leq^S$	$\leq^S$	We often drop the superscript from $\leq^S$ and simply write $\leq$. In formal logic notation,
fong-spivak-seven-sketches-F02736	260	■ 1.000	$S = 1$	$S = 1$	In particular, if $S = 1$ is the terminal object, we denote $ \Omega^S $ by $ \Omega $, and refer to elements
fong-spivak-seven-sketches-F02737	260	■ 1.000	$ \Omega^S $	$ \Omega^S $	In particular, if $S = 1$ is the terminal object, we denote $ \Omega^S $ by $ \Omega $, and refer to elements
fong-spivak-seven-sketches-F02738	260	■ 1.000	$ \Omega $	$ \Omega $	In particular, if $S = 1$ is the terminal object, we denote $ \Omega^S $ by $ \Omega $, and refer to elements
fong-spivak-seven-sketches-F02739	260	■ 1.000	$p \in \Omega $	$p \in \Omega $	$p \in \Omega $ as <i>propositions</i> . They are just morphisms $p: 1 \rightarrow \Omega$.
fong-spivak-seven-sketches-F02740	260	■ 1.000	$p: 1 \rightarrow \Omega$	$p: 1 \rightarrow \Omega$	$p \in \Omega $ as <i>propositions</i> . They are just morphisms $p: 1 \rightarrow \Omega$.
fong-spivak-seven-sketches-F02741	260	■ 1.000	$p = q$	$p = q$	$p = q$. The reason is that for any subsets $U, V \subseteq X$, if $U \subseteq V$ and $V \subseteq U$ then $U = V$.
fong-spivak-seven-sketches-F02742	260	■ 1.000	$U, V \subseteq X$	$U, V \subseteq X$	$p = q$. The reason is that for any subsets $U, V \subseteq X$, if $U \subseteq V$ and $V \subseteq U$ then $U = V$.
fong-spivak-seven-sketches-F02743	260	■ 1.000	$U = V$	$U = V$	$p = q$. The reason is that for any subsets $U, V \subseteq X$, if $U \subseteq V$ and $V \subseteq U$ then $U = V$.
fong-spivak-seven-sketches-F02744	260	■ 0.976	$S \in \mathbf{Shv}(X)$	$S \in \mathbf{Shv}(X)$	<i>Exercise 7.64.</i> Give an example of a space X , a sheaf $S \in \mathbf{Shv}(X)$, and two predicates
fong-spivak-seven-sketches-F02745	260	■ 1.000	$p(s) \vdash_{s:S} q(s)$	$p(s) \vdash_{s:S} q(s)$	$p, q: S \rightarrow \Omega$ for which $p(s) \vdash_{s:S} q(s)$ holds. You do not have to be formal.
fong-spivak-seven-sketches-F02746	260	■ 1.000	$\wedge, \vee, \Rightarrow, \neg$	$\wedge, \vee, \Rightarrow, \neg$	All of the logical symbols (true, false, $\wedge, \vee, \Rightarrow, \neg$) from Section 7.4.2 make sense
fong-spivak-seven-sketches-F02747	260	■ 1.000	$(p \wedge q): S \rightarrow \Omega$	$(p \wedge q): S \rightarrow \Omega$	in any such poset $ \Omega^S $. For any two predicates $p, q: S \rightarrow \Omega$, we define $(p \wedge q): S \rightarrow \Omega$
fong-spivak-seven-sketches-F02748	260	■ 1.000	$(p \wedge q)(s) := p(s) \wedge q(s)$	$(p \wedge q)(s) := p(s) \wedge q(s)$	by $(p \wedge q)(s) := p(s) \wedge q(s)$, and similarly for $\vee$. Thus one says that these operations
fong-spivak-seven-sketches-F02749	260	■ 1.000	$n+1$	$n + 1$	exists.’ Each takes in a predicate of $n+1$ variables and returns a predicate of n variables.
fong-spivak-seven-sketches-F02750	260	■ 1.000	$S, T \in \mathbf{Shv}(X, \mathbf{Op})$	$S, T \in \mathbf{Shv}(X, \mathbf{Op})$	<i>Example 7.65.</i> Suppose we have two sheaves $S, T \in \mathbf{Shv}(X, \mathbf{Op})$ and a predicate $p: S \times$
fong-spivak-seven-sketches-F02751	260	■ 1.000	$p: S \times$	$p: S \times$	<i>Example 7.65.</i> Suppose we have two sheaves $S, T \in \mathbf{Shv}(X, \mathbf{Op})$ and a predicate $p: S \times$
fong-spivak-seven-sketches-F02752	260	■ 1.000	$T \rightarrow \Omega$	$T \rightarrow \Omega$	$T \rightarrow \Omega$. Let’s say T represents what’s considered newsworthy and S is again the set
fong-spivak-seven-sketches-F02753	260	■ 1.000	$t \in T(U)$	$t \in T(U)$	of people. So for a subset of time U , a section $t \in T(U)$ is something that’s considered

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches- F02780	261	■ 1.000	$p(\text{left}(\text{left.s} \text{right} _{V_i}, t_i) = V_i)$	$p(s _{V_i}, t_i) = V_i$	which there exists some $t_i \in T(V_i)$ satisfying $p(s _{V_i}, t_i) = V_i$. If the result is U itself, you
fong-spivak-seven-sketches- F02781	261	■ 1.000	$p(t)$	$p(t)$	might be tempted to think “ah, so there exists some $t \in T(U)$ satisfying $p(t)$,” but that
fong-spivak-seven-sketches- F02782	261	■ 0.516	$U = \bigcup U_i$	$U = \bigcup U_i$	is not necessarily so. There is just a cover of $U = \bigcup U_i$ and local sections $t_i \in T(U_i)$,
fong-spivak-seven-sketches- F02783	261	■ 0.516	$t_i \in T(U_i)$	$t_i \in T(U_i)$	is not necessarily so. There is just a cover of $U = \bigcup U_i$ and local sections $t_i \in T(U_i)$,
fong-spivak-seven-sketches- F02784	262	■ 0.951	$t^{\prime}$	$t^{\prime}$	<i>Exercise 7.68.</i> Apply the above definition to the “person s is worried about news t ”
fong-spivak-seven-sketches- F02785	262	■ 1.000	$\{(s, t) \in S \times T \mid p(s, t)\} \subseteq S \times T$	$\{(s, t) \in S \times T \mid p(s, t)\} \subseteq S \times T$	with the subobject classified by p , namely $\{(s, t) \in S \times T \mid p(s, t)\} \subseteq S \times T$, compose
fong-spivak-seven-sketches- F02786	262	■ 0.982	$\pi_S: S \times T \rightarrow S$	$\pi_S: S \times T \rightarrow S$	with the projection $\pi_S: S \times T \rightarrow S$ as on the upper right; then take the epi-mono
fong-spivak-seven-sketches- F02787	262	■ 0.983	$(\Omega^S , \leq^S)$	$(\Omega^S , \leq^S)$	poset $(\Omega^S , \leq^S)$ of predicates on S , where $ \Omega^S = \mathcal{E}(S, \Omega)$ is just the set of morphisms
fong-spivak-seven-sketches- F02788	262	■ 0.983	$ \Omega^S = \mathcal{E}(S, \Omega)$	$ \Omega^S = \mathcal{E}(S, \Omega)$	poset $(\Omega^S , \leq^S)$ of predicates on S , where $ \Omega^S = \mathcal{E}(S, \Omega)$ is just the set of morphisms
fong-spivak-seven-sketches- F02789	262	■ 0.999	$\mathbf{Shv}(X)$	$\mathbf{Shv}(X)$	Definition 7.69. A modality in $\mathbf{Shv}(X)$ is a sheaf morphism $j: \Omega \rightarrow \Omega$ satisfying three
fong-spivak-seven-sketches- F02790	262	■ 0.999	$j: \Omega \rightarrow \Omega$	$j: \Omega \rightarrow \Omega$	Definition 7.69. A modality in $\mathbf{Shv}(X)$ is a sheaf morphism $j: \Omega \rightarrow \Omega$ satisfying three
fong-spivak-seven-sketches- F02791	262	■ 1.000	$p, q \in \Omega(U)$	$p, q \in \Omega(U)$	properties for all $U \subseteq X$ and $p, q \in \Omega(U)$:
fong-spivak-seven-sketches- F02792	262	■ 1.000	$(j; j)(p) \leq j(p)$	$(j; j)(p) \leq j(p)$	(b) $(j; j)(p) \leq j(p)$; and
fong-spivak-seven-sketches- F02793	262	■ 1.000	$j(p \wedge q) = j(p) \wedge j(q)$	$j(p \wedge q) = j(p) \wedge j(q)$	(c) $j(p \wedge q) = j(p) \wedge j(q)$.
fong-spivak-seven-sketches- F02794	262	■ 1.000	$p \in \Omega(U)$	$p \in \Omega(U)$	holds for all $U \subseteq X$ and $p \in \Omega(U)$. Show that for all $q \in \Omega(U)$ we have $j(j(q)) \leq j(q)$ iff
fong-spivak-seven-sketches- F02795	262	■ 1.000	$q \in \Omega(U)$	$q \in \Omega(U)$	holds for all $U \subseteq X$ and $p \in \Omega(U)$. Show that for all $q \in \Omega(U)$ we have $j(j(q)) \leq j(q)$ iff
fong-spivak-seven-sketches- F02796	262	■ 1.000	$j(j(q)) \leq j(q)$	$j(j(q)) \leq j(q)$	holds for all $U \subseteq X$ and $p \in \Omega(U)$. Show that for all $q \in \Omega(U)$ we have $j(j(q)) \leq j(q)$ iff
fong-spivak-seven-sketches- F02797	262	■ 1.000	$j(j(q)) = j(q)$	$j(j(q)) = j(q)$	$j(j(q)) = j(q)$.
fong-spivak-seven-sketches- F02798	262	■ 1.000	$\Omega \rightarrow \Omega$	$\Omega \rightarrow \Omega$	(a) the sheaf morphism $\Omega \rightarrow \Omega$ given by sending q to $p \Rightarrow q$ is a modality.
fong-spivak-seven-sketches- F02799	262	■ 1.000	$p \vee q$	$p \vee q$	(b) the sheaf morphism $\Omega \rightarrow \Omega$ given by sending q to $p \vee q$ is a modality.
fong-spivak-seven-sketches- F02800	262	■ 0.997	$(q \Rightarrow p) \Rightarrow p$	$(q \Rightarrow p) \Rightarrow p$	(c) the sheaf morphism $\Omega \rightarrow \Omega$ given by sending q to $(q \Rightarrow p) \Rightarrow p$ is a modality.
fong-spivak-seven-sketches- F02801	263	■ 1.000	$p(s)$	$p(s)$	$p(s)$, and what does it mean?
fong-spivak-seven-sketches- F02802	263	■ 1.000	$j(p(s))$	$j(p(s))$	3. What sort of thing is $j(p(s))$ and what does it mean?
fong-spivak-seven-sketches- F02803	263	■ 1.000	$p(s) \leq j(p(s))$	$p(s) \leq j(p(s))$	4. Is it true that $p(s) \leq j(p(s))$? Explain briefly.
fong-spivak-seven-sketches- F02804	263	■ 1.000	$j(j(p(s))) = j(p(s))$	$j(j(p(s))) = j(p(s))$	5. Is it true that $j(j(p(s))) = j(p(s))$? Explain briefly.
fong-spivak-seven-sketches- F02805	263	■ 1.000	$q: S \rightarrow \Omega$	$q: S \rightarrow \Omega$	6. Choose another predicate $q: S \rightarrow \Omega$. Is it true that $j(p \wedge q) = j(p) \wedge j(q)$? Explain

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02806	264	0.500	$s \in S(c)$	$s \in S(c)$	Is it true that for every table $c \in \text{Ob}(\mathcal{C})$ and every row $s \in S(c)$ there exists
fong-spivak-seven-sketches-F02807	264	1.000	$t \in T(c)$	$t \in T(c)$	a row $t \in T(c)$ such that $f(s) = t$?
fong-spivak-seven-sketches-F02808	264	1.000	$f(s)=t$	$f(s) = t$	a row $t \in T(c)$ such that $f(s) = t$?
fong-spivak-seven-sketches-F02809	264	1.000	$\left(U_{\{i\}} \subseteq U\right)_{i \in I}$	$(U_i \subseteq U)_{i \in I}$	bundle S , there exists an open covering of $(U_i \subseteq U)_{i \in I}$ of U and a section
fong-spivak-seven-sketches-F02810	264	1.000	$f\left(t_{\{i\}}\right)=\left.s\right _{U_{\{i\}}}$	$f\left(t_i\right)=s _{U_i}$	$t_i \in T(U_i)$ of the bundle T for each $i \in I$, such that $f\left(t_i\right)=s _{U_i}$ is the
fong-spivak-seven-sketches-F02811	264	1.000	$U_{\{i\}}$	U_i	restriction of s to U_i ?
fong-spivak-seven-sketches-F02812	264	0.912	$\boldsymbol{\operatorname{Shv}}\left(\mathbb{R}\right)$	$\mathbf{Shv}(\mathbb{R})$	intervals of time. In other words, we will be interested in the category $\mathbf{Shv}(\mathbb{R})$ of
fong-spivak-seven-sketches-F02813	264	0.773	$\mathbb{I}$	$\mathbb{I}\mathbb{R}$	and $\mathbf{Op}$ is a topology on X ; see Definition 7.25. The set of points for $\mathbb{I}\mathbb{R}$ is that of all
fong-spivak-seven-sketches-F02814	264	1.000	$a<b$	$a < b$	For $a < b$ in $\mathbb{R}$, let $o_{[a,b]}$ denote the set $o_{[a,b]} := \{[d, u] \in \mathbb{I}\mathbb{R} \mid a < d \leq u < b\}$; these are
fong-spivak-seven-sketches-F02815	264	1.000	$o_{\{[a, b]\}}$	$o_{[a,b]}$	For $a < b$ in $\mathbb{R}$, let $o_{[a,b]}$ denote the set $o_{[a,b]} := \{[d, u] \in \mathbb{I}\mathbb{R} \mid a < d \leq u < b\}$; these are
fong-spivak-seven-sketches-F02816	264	1.000	$o_{\{[a, b]\}}:=\{[d, u] \in \mathbb{R} \mid a<d \leq u<b\}$	$o_{[a,b]} := \{[d, u] \in \mathbb{R} \mid a < d \leq u < b\}$	For $a < b$ in $\mathbb{R}$, let $o_{[a,b]}$ denote the set $o_{[a,b]} := \{[d, u] \in \mathbb{I}\mathbb{R} \mid a < d \leq u < b\}$; these are
fong-spivak-seven-sketches-F02817	264	1.000	$o_{\{[0,5]\}}$	$o_{[0,5]}$	Thus for example, $o_{[0,5]}$ is an open set: it contains every $[d, u]$ contained in the
fong-spivak-seven-sketches-F02818	264	1.000	$[d, u]$	$[d, u]$	Thus for example, $o_{[0,5]}$ is an open set: it contains every $[d, u]$ contained in the
fong-spivak-seven-sketches-F02819	264	1.000	$\{x \in \mathbb{R} \mid 0 < x < 5\}$	$\{x \in \mathbb{R} \mid 0 < x < 5\}$	open interval $\{x \in \mathbb{R} \mid 0 < x < 5\}$. Similarly $o_{[4,8]}$ is an open set, but note that
fong-spivak-seven-sketches-F02820	264	1.000	$o_{\{[4,8]\}}$	$o_{[4,8]}$	open interval $\{x \in \mathbb{R} \mid 0 < x < 5\}$. Similarly $o_{[4,8]}$ is an open set, but note that
fong-spivak-seven-sketches-F02821	264	0.965	$o_{\{[0,5]\}} \cup o_{\{[4,8]\}} \neq o_{\{[0,8]\}}$	$o_{[0,5]} \cup o_{[4,8]} \neq o_{[0,8]}$	$o_{[0,5]} \cup o_{[4,8]} \neq o_{[0,8]}$. Indeed, the interval $[2, 6]$ is in the right-hand side but not the left.
fong-spivak-seven-sketches-F02822	265	1.000	$[2,6] \in o_{\{[0,8]\}}$	$[2,6] \in o_{[0,8]}$	1. Explain why $[2,6] \in o_{[0,8]}$.
fong-spivak-seven-sketches-F02823	265	0.996	$[2,6] \notin o_{\{[0,5]\}} \cup o_{\{[4,8]\}}$	$[2,6] \notin o_{[0,5]} \cup o_{[4,8]}$	2. Explain why $[2,6] \notin o_{[0,5]} \cup o_{[4,8]}$.
fong-spivak-seven-sketches-F02824	265	0.964	$\mathbb{R} \subseteq \mathbb{I}$	$\mathbb{R} \subseteq \mathbb{I}\mathbb{R}$	Exercise 7.77. Show that a subset $U \subseteq \mathbb{R}$ is open in the subspace topology of $\mathbb{R} \subseteq \mathbb{I}\mathbb{R}$
fong-spivak-seven-sketches-F02825	265	1.000	$U \cap \mathbb{R}$	$U \cap \mathbb{R}$	iff $U \cap \mathbb{R}$ is open in the usual topology on $\mathbb{R}$ defined in Example 7.26.
fong-spivak-seven-sketches-F02826	265	0.280	$\mathbf{BT}=\mathbf{Shv}\left(\mathbb{I}, \mathbf{Op}\right)$	$\mathbf{BT} = \mathbf{Shv}(\mathbb{I}\mathbb{R}, \mathbf{Op})$	We cannot go into much depth about the sheaf topos $\mathbf{BT} = \mathbf{Shv}(\mathbb{I}\mathbb{R}, \mathbf{Op})$, for reasons
fong-spivak-seven-sketches-F02827	265	0.591	$\mathbb{I}, \mathbf{Op}$	$\mathbb{I}\mathbb{R}, \mathbf{Op}$	What is a sheaf on $\mathbb{I}\mathbb{R}$? A sheaf S on the interval domain $(\mathbb{I}\mathbb{R}, \mathbf{Op})$ is a functor
fong-spivak-seven-sketches-F02828	265	0.864	$S: \mathbf{Op}^{\text{op}} \rightarrow \mathbf{Set}$	$S: \mathbf{Op}^{\text{op}} \rightarrow$	$S: \mathbf{Op}^{\text{op}} \rightarrow \mathbf{Set}$: it assigns to each open set U a set $S(U)$; how should we interpret this?
fong-spivak-seven-sketches-F02829	265	0.864	$S(U)$	$S(U)$	$S: \mathbf{Op}^{\text{op}} \rightarrow \mathbf{Set}$: it assigns to each open set U a set $S(U)$; how should we interpret this?
fong-spivak-seven-sketches-F02830	265	0.997	$\left(s_{\{i\}}\right)$	(s_i)	V . If $U = \bigcup_{i \in I} U_i$ and we can find matching U_i -events (s_i) for each $i \in I$, then the sheaf
fong-spivak-seven-sketches-F02831	265	0.835	$U \subseteq \mathbb{I}$	$U \subseteq \mathbb{I}\mathbb{R}$	We said in Section 7.5.1 that every open set $U \subseteq \mathbb{I}\mathbb{R}$ can be written as the union of
fong-spivak-seven-sketches-F02832	265	1.000	$S\left(o_{\{[a, b]\}}\right)$	$S\left(o_{[a,b]}\right)$	basic open sets $o_{[a,b]}$. This implies that any sheaf S is determined by its values $S(o_{[a,b]})$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches- F02833	265	0.753	$\mathrm{A} \in \mathbf{Shv}(\mathbb{R})$	$A \in \mathbf{Shv}(\mathbb{R})$	<i>Example 7.78.</i> For any set A there is a sheaf $A \in \mathbf{Shv}(\mathbb{R})$ that assigns to each open set U
fong-spivak-seven-sketches- F02834	265	1.000	$\mathrm{A}(U) := A$	$A(U) := A$	the set $A(U) := A$. This allows us to refer to integers, or real numbers, or letters of an
fong-spivak-seven-sketches- F02835	265	1.000	$7 \in \mathbb{N}$	$7 \in \mathbb{N}$	alphabet, as though they were behaviors. What sort of behavior is $7 \in \mathbb{N}$? It is the sort
fong-spivak-seven-sketches- F02836	266	0.881	$\left(X, \mathbf{0}_p\right)$	$(X, \mathbf{Op}_X)$	<i>Example 7.79.</i> Fix any topological space $(X, \mathbf{Op}_X)$. Then there is a sheaf F_X of local
fong-spivak-seven-sketches- F02837	266	0.881	F_X	F_X	<i>Example 7.79.</i> Fix any topological space $(X, \mathbf{Op}_X)$. Then there is a sheaf F_X of local
fong-spivak-seven-sketches- F02838	266	0.614	$U \in \mathbf{0}_p(\mathbb{R})$	$U \in \mathbf{Op}_{\mathbb{R}}$	functions from $\mathbb{R}$ to X . That is, for any open set $U \in \mathbf{Op}_{\mathbb{R}}$, we assign the set $F_X(U) :=$
fong-spivak-seven-sketches- F02839	266	0.614	$F_X(U) :=$	$F_X(U) :=$	functions from $\mathbb{R}$ to X . That is, for any open set $U \in \mathbf{Op}_{\mathbb{R}}$, we assign the set $F_X(U) :=$
fong-spivak-seven-sketches- F02840	266	0.875	$\{f: U \rightarrow X \mid f$	$\{f: U \rightarrow X \mid f$	$\{f: U \rightarrow X \mid f$ is continuous}. There is also the sheaf G_X of local functions on the
fong-spivak-seven-sketches- F02841	266	0.875	G_X	G_X	$\{f: U \rightarrow X \mid f$ is continuous}. There is also the sheaf G_X of local functions on the
fong-spivak-seven-sketches- F02842	266	0.960	$\mathbb{R} \subseteq \mathbb{R}$	$\mathbb{R} \subseteq \mathbb{R}$	subspace $\mathbb{R} \subseteq \mathbb{R}$. That is, for any open set $U \in \mathbf{Op}_{\mathbb{R}}$, we assign the set $G_X(U) :=$
fong-spivak-seven-sketches- F02843	266	0.960	$G_X(U) :=$	$G_X(U) :=$	subspace $\mathbb{R} \subseteq \mathbb{R}$. That is, for any open set $U \in \mathbf{Op}_{\mathbb{R}}$, we assign the set $G_X(U) :=$
fong-spivak-seven-sketches- F02844	266	0.994	$\{f: U \cap \mathbb{R} \rightarrow X \mid f$	$\{f: U \cap \mathbb{R} \rightarrow X \mid f$	$\{f: U \cap \mathbb{R} \rightarrow X \mid f$ is continuous}.
fong-spivak-seven-sketches- F02845	266	0.995	$R \subseteq \mathbb{R}$	$R \subseteq \mathbb{R}$	$(X, \mathbf{Op}_X)$ and any subset $R \subseteq \mathbb{R}$ of the interval domain. Define $H_X(U) := \{f: U \cap R \rightarrow$
fong-spivak-seven-sketches- F02846	266	0.995	$H_X(U) := \{f: U \cap R \rightarrow$	$H_X(U) := \{f: U \cap R \rightarrow$	$(X, \mathbf{Op}_X)$ and any subset $R \subseteq \mathbb{R}$ of the interval domain. Define $H_X(U) := \{f: U \cap R \rightarrow$
fong-spivak-seven-sketches- F02847	266	1.000	$X \mid f$	$X \mid f$	$X \mid f$ is continuous}.
fong-spivak-seven-sketches- F02848	266	1.000	H_X	H_X	1. Is H_X a presheaf? If not, why not; if so, what are the restriction maps?
fong-spivak-seven-sketches- F02849	266	0.715	$(d, d+3)$	$(d, d+3)$	idea is straightforward: for any interval of time, say a three-day interval $(d, d+3)$, let
fong-spivak-seven-sketches- F02850	266	1.000	$L(d, d+3)$	$L(d, d+3)$	$L(d, d+3)$ denote the set of all possible behaviors a person who is in love could possibly
fong-spivak-seven-sketches- F02851	266	0.585	$(\mathbb{I}, \mathbf{Op})$	$(\mathbb{I}, \mathbf{Op})$	BT is the topos of sheaves on the space $(\mathbb{I}, \mathbf{Op})$, as defined in Section 7.5.1. As
fong-spivak-seven-sketches- F02852	267	1.000	$(0, 5)$	$(0, 5)$	Suppose Bob likes the weather throughout the interval $(0, 5)$ and throughout the
fong-spivak-seven-sketches- F02853	267	0.975	$(0, 8)$	$(0, 8)$	not imply that it's been true on $(0, 8)$: there may have been a change of shift, where
fong-spivak-seven-sketches- F02854	267	0.833	$(4, 8)$	$(4, 8)$	This might be true on $(0, 5)$ and true on $(4, 8)$ but not on $(0, 8)$. In order to capture the
fong-spivak-seven-sketches- F02855	267	1.000	$\theta \in \mathbb{R}_{\geq 0}$	$\theta \in \mathbb{R}_{\geq 0}$	are behavior types: the altimeter is a variable whose value $\theta \in \mathbb{R}_{\geq 0}$ is changing
fong-spivak-seven-sketches- F02856	267	0.925	$\rightarrow \Omega$	$\rightarrow \Omega$	$p: \text{dials} \times \text{thrusters} \rightarrow \Omega$. While we have not written out a formal language for p ,
fong-spivak-seven-sketches- F02857	268	1.000	$p(D, T)$	$p(D, T)$	one could imagine the predicate $p(D, T)$ for $D: \text{dials}$ and $T: \text{thrusters}$ as
fong-spivak-seven-sketches- F02858	268	0.997	$@_t$	$@_t$	Here $@_t$ is a modality, as we discussed in Definition 7.69; in fact it turns out to be one
fong-spivak-seven-sketches- F02859	268	0.988	$@_t(q)$	$@_t(q)$	statement $@_t(q)$ says that q is true in some small enough neighborhood around t . So

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F02860	268	0.998	$D \in$	$D \in$	Given an actual playing-out-of-events over a time period U , i.e. actual section $D \in$
fong-spivak-seven-sketches-F02861	268	0.962	(U)	(U)	$\text{dials}(U)$ and section $T \in \text{thrusters}(U)$, the predicate Eq. (7.82) will hold on certain
fong-spivak-seven-sketches-F02862	268	0.962	$T \in$	$T \in$	$\text{dials}(U)$ and section $T \in \text{thrusters}(U)$, the predicate Eq. (7.82) will hold on certain
fong-spivak-seven-sketches-F02863	271	1.000	$f(x)=x+5$	$f(x) = x + 5$	order-preserving: Take $f(x) = x + 5$; if $x \leq y$ then $x + 5 \leq y + 5$, so f is order-preserving. Take
fong-spivak-seven-sketches-F02864	271	1.000	$x+5 \leq y+5$	$x + 5 \leq y + 5$	order-preserving: Take $f(x) = x + 5$; if $x \leq y$ then $x + 5 \leq y + 5$, so f is order-preserving. Take
fong-spivak-seven-sketches-F02865	271	0.416	$g(x) := -x$	$g(x) := -x$	$g(x) := -x$; even though $1 \leq 2$, the required inequality $-1 \leq -2$ does not hold, so g is not
fong-spivak-seven-sketches-F02866	271	0.416	$1 \leq 2$	$1 \leq 2$	$g(x) := -x$; even though $1 \leq 2$, the required inequality $-1 \leq -2$ does not hold, so g is not
fong-spivak-seven-sketches-F02867	271	0.416	$-1 \leq -2$	$-1 \leq -2$	$g(x) := -x$; even though $1 \leq 2$, the required inequality $-1 \leq -2$ does not hold, so g is not
fong-spivak-seven-sketches-F02868	271	0.986	$f(x) := x + 5$	$f(x) := x + 5$	metric-preserving: Take $f(x) := x + 5$; for any x, y we have $ x - y = (x + 5) - (y + 5) $ by the rules of
fong-spivak-seven-sketches-F02869	271	0.986	$ x - y = (x + 5) - (y + 5) $	$ x - y = (x + 5) - (y + 5) $	metric-preserving: Take $f(x) := x + 5$; for any x, y we have $ x - y = (x + 5) - (y + 5) $ by the rules of
fong-spivak-seven-sketches-F02870	271	1.000	$g(x) := 2 * x$	$g(x) := 2 * x$	arithmetic, so $ x - y = f(x) - f(y) $, meaning f preserves metric. Take $g(x) := 2 * x$; then with
fong-spivak-seven-sketches-F02871	271	1.000	$x=1$	$x = 1$	$x = 1$ and $y = 2$ we have $ x - y = 1$ but $ 2x - 2y = 2$, so g does not preserve the metric.
fong-spivak-seven-sketches-F02872	271	1.000	$y=2$	$y = 2$	$x = 1$ and $y = 2$ we have $ x - y = 1$ but $ 2x - 2y = 2$, so g does not preserve the metric.
fong-spivak-seven-sketches-F02873	271	1.000	$ x - y = 1$	$ x - y = 1$	$x = 1$ and $y = 2$ we have $ x - y = 1$ but $ 2x - 2y = 2$, so g does not preserve the metric.
fong-spivak-seven-sketches-F02874	271	1.000	$ 2x - 2y = 2$	$ 2x - 2y = 2$	$x = 1$ and $y = 2$ we have $ x - y = 1$ but $ 2x - 2y = 2$, so g does not preserve the metric.
fong-spivak-seven-sketches-F02875	271	1.000	$f(x) := 3 * x$	$f(x) := 3 * x$	addition-preserving: Take $f(x) := 3 * x$; for any x, y we have $3 * (x + y) = (3 * x) + (3 * y)$, so f preserves
fong-spivak-seven-sketches-F02876	271	1.000	$3 * (x + y) = (3 * x) + (3 * y)$	$3 * (x + y) = (3 * x) + (3 * y)$	addition-preserving: Take $f(x) := 3 * x$; for any x, y we have $3 * (x + y) = (3 * x) + (3 * y)$, so f preserves
fong-spivak-seven-sketches-F02877	271	1.000	$g(x) := x + 1$	$g(x) := x + 1$	addition. Take $g(x) := x + 1$; then with $x = 0$ and $y = 0$, we have $g(x + y) = 1$, but $g(x) + g(y) = 2$,
fong-spivak-seven-sketches-F02878	271	1.000	$x=0$	$x = 0$	addition. Take $g(x) := x + 1$; then with $x = 0$ and $y = 0$, we have $g(x + y) = 1$, but $g(x) + g(y) = 2$,
fong-spivak-seven-sketches-F02879	271	1.000	$y=0$	$y = 0$	addition. Take $g(x) := x + 1$; then with $x = 0$ and $y = 0$, we have $g(x + y) = 1$, but $g(x) + g(y) = 2$,
fong-spivak-seven-sketches-F02880	271	1.000	$g(x + y) = 1$	$g(x + y) = 1$	addition. Take $g(x) := x + 1$; then with $x = 0$ and $y = 0$, we have $g(x + y) = 1$, but $g(x) + g(y) = 2$,
fong-spivak-seven-sketches-F02881	271	1.000	$g(x) + g(y) = 2$	$g(x) + g(y) = 2$	addition. Take $g(x) := x + 1$; then with $x = 0$ and $y = 0$, we have $g(x + y) = 1$, but $g(x) + g(y) = 2$,
fong-spivak-seven-sketches-F02882	272	1.000	$A = (12)(3)(4)$	$A = (12)(3)(4)$	For the remaining parts, we choose $A = (12)(3)(4)$ and $B = (13)(2)(4)$.
fong-spivak-seven-sketches-F02883	272	1.000	$B = (13)(2)(4)$	$B = (13)(2)(4)$	For the remaining parts, we choose $A = (12)(3)(4)$ and $B = (13)(2)(4)$.
fong-spivak-seven-sketches-F02884	272	1.000	$A \vee B = (123)(4)$	$A \vee B = (123)(4)$	3. $A \vee B = (123)(4)$
fong-spivak-seven-sketches-F02885	272	0.622	$\{1, 2, 3\} \cup \{1\} = \{1, 2, 3\}$	$\{1, 2, 3\} \cup \{1\} = \{1, 2, 3\}$	2. The union of $\{1, 2, 3\}$ and $\{1\}$ is $\{1, 2, 3\} \cup \{1\} = \{1, 2, 3\}$.
fong-spivak-seven-sketches-F02886	272	0.992	$\{h, 1\} \times \{1, 2, 3\}$	$\{h, 1\} \times \{1, 2, 3\}$	3. The six elements of $\{h, 1\} \times \{1, 2, 3\}$ are
fong-spivak-seven-sketches-F02887	272	1.000	$\{h, 1\} \sqcup \{1, 2, 3\}$	$\{h, 1\} \sqcup \{1, 2, 3\}$	4. The five elements of $\{h, 1\} \sqcup \{1, 2, 3\}$ are

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches- F02888	272	■ 1.000	$\{h, 1\} \cup \{1, 2, 3\}$	$\{h, 1\} \cup \{1, 2, 3\}$	5. The four elements of $\{h, 1\} \cup \{1, 2, 3\}$ are
fong-spivak-seven-sketches- F02889	272	■ 1.000	$0 \notin \{n \in \mathbb{Z} \mid n \geq 1\}$	$0 \notin \{n \in \mathbb{Z} \mid n \geq 1\}$	2. This is false: $0 \in \mathbb{N}$ but $0 \notin \{n \in \mathbb{Z} \mid n \geq 1\}$.
fong-spivak-seven-sketches- F02890	273	■ 0.994	$p_1^{\wedge \prime}, p_2^{\wedge \prime} \in P^{\wedge \prime}$	$p_1', p_2' \in P'$	1. Given $p \in P$, suppose we had $p_1', p_2' \in P'$ such that $A_p = A_{p_1}'$ and $A_p = A_{p_2}'$. Well then $A_{p_1}' = A_{p_2}'$.
fong-spivak-seven-sketches- F02891	273	■ 0.994	$A_{\{p\}} = A_{\{p_1^{\wedge \prime}\}}^{\wedge \prime}$	$A_p = A_{p_1}'$	1. Given $p \in P$, suppose we had $p_1', p_2' \in P'$ such that $A_p = A_{p_1}'$ and $A_p = A_{p_2}'$. Well then $A_{p_1}' = A_{p_2}'$.
fong-spivak-seven-sketches- F02892	273	■ 0.994	$A_{\{p\}} = A_{\{p_2^{\wedge \prime}\}}^{\wedge \prime}$	$A_p = A_{p_2}'$	1. Given $p \in P$, suppose we had $p_1', p_2' \in P'$ such that $A_p = A_{p_1}'$ and $A_p = A_{p_2}'$. Well then $A_{p_1}' = A_{p_2}'$.
fong-spivak-seven-sketches- F02893	273	■ 0.994	$A_{\{p_1^{\wedge \prime}\}} = A_{\{p_2^{\wedge \prime}\}}$	$A_{p_1}' = A_{p_2}'$	1. Given $p \in P$, suppose we had $p_1', p_2' \in P'$ such that $A_p = A_{p_1}'$ and $A_p = A_{p_2}'$. Well then $A_{p_1}' = A_{p_2}'$.
fong-spivak-seven-sketches- F02894	273	■ 1.000	$A_{\{p_1^{\wedge \prime}\}} \cap A_{\{p_2^{\wedge \prime}\}} = A_{\{p_1^{\wedge \prime}\}}$	$A_{p_1}' \cap A_{p_2}' = A_{p_1}'$	so in particular $A_{p_1}' \cap A_{p_2}' = A_{p_1}'$. By the definition of partition (1.14), $A_{p_1}' \neq \emptyset$, and yet if $p_1 \neq p_2$
fong-spivak-seven-sketches- F02895	273	■ 1.000	$A_{\{p_1^{\wedge \prime}\}} \neq \varnothing$	$A_{p_1}' \neq \emptyset$	so in particular $A_{p_1}' \cap A_{p_2}' = A_{p_1}'$. By the definition of partition (1.14), $A_{p_1}' \neq \emptyset$, and yet if $p_1 \neq p_2$
fong-spivak-seven-sketches- F02896	273	■ 1.000	$p_1 \neq p_2$	$p_1 \neq p_2$	so in particular $A_{p_1}' \cap A_{p_2}' = A_{p_1}'$. By the definition of partition (1.14), $A_{p_1}' \neq \emptyset$, and yet if $p_1 \neq p_2$
fong-spivak-seven-sketches- F02897	273	■ 0.998	$A_{\{p_1^{\wedge \prime}\}} \cap A_{\{p_2^{\wedge \prime}\}} = \varnothing$	$A_{p_1}' \cap A_{p_2}' = \emptyset$	then $A_{p_1}' \cap A_{p_2}' = \emptyset$. This can't be, so we must have $p_1' = p_2'$, as desired.
fong-spivak-seven-sketches- F02898	273	■ 0.998	$p_1^{\wedge \prime} = p_2^{\wedge \prime}$	$p_1' = p_2'$	then $A_{p_1}' \cap A_{p_2}' = \emptyset$. This can't be, so we must have $p_1' = p_2'$, as desired.
fong-spivak-seven-sketches- F02899	273	■ 1.000	$A_{\{p^{\wedge \prime}\}}^{\wedge \prime} \neq \varnothing$	$A_{p'}' \neq \emptyset$	2. Suppose given $p' \in P'$; we want to show that there is a $p \in P$ such that $A_p = A_{p'}'$. Since $A_{p'}' \neq \emptyset$
fong-spivak-seven-sketches- F02900	273	■ 1.000	$a \in A_{\{p^{\wedge \prime}\}}^{\wedge \prime}$	$a \in A_{p'}'$	is nonempty by definition, we can pick some $a \in A_{p'}'$; since $A_{p'}' \subseteq A$, we have $a \in A$. Finally,
fong-spivak-seven-sketches- F02901	273	■ 1.000	$A_{\{p^{\wedge \prime}\}}^{\wedge \prime} \subseteq A$	$A_{p'}' \subseteq A$	is nonempty by definition, we can pick some $a \in A_{p'}'$; since $A_{p'}' \subseteq A$, we have $a \in A$. Finally,
fong-spivak-seven-sketches- F02902	273	■ 1.000	$p^{\wedge \prime \prime} \in P^{\wedge \prime}$	$p'' \in P'$	$A_p = A_{p'}'$. By assumption there is some $p'' \in P'$ with $A_p = A_{p''}'$, so now $a \in A_{p''}'$, and $a \in A_{p'}'$.
fong-spivak-seven-sketches- F02903	273	■ 1.000	$A_{\{p\}} = A_{\{p^{\wedge \prime \prime}\}}^{\wedge \prime}$	$A_p = A_{p''}'$	$A_p = A_{p'}'$. By assumption there is some $p'' \in P'$ with $A_p = A_{p''}'$, so now $a \in A_{p''}'$, and $a \in A_{p'}'$.
fong-spivak-seven-sketches- F02904	273	■ 1.000	$a \in A_{\{p^{\wedge \prime \prime}\}}^{\wedge \prime}$	$a \in A_{p''}'$	$A_p = A_{p'}'$. By assumption there is some $p'' \in P'$ with $A_p = A_{p''}'$, so now $a \in A_{p''}'$, and $a \in A_{p'}'$.
fong-spivak-seven-sketches- F02905	273	■ 1.000	$a \in A_{\{p^{\wedge \prime \prime}\}}^{\wedge \prime} \cap A_{\{p^{\wedge \prime \prime}\}}^{\wedge \prime}$	$a \in A_{p'}' \cap A_{p''}'$	so $a \in A_{p'}' \cap A_{p''}'$. Again by definition, having a nonempty intersection means $p' = p''$. So we
fong-spivak-seven-sketches- F02906	273	■ 1.000	$p^{\wedge \prime} = p^{\wedge \prime \prime}$	$p' = p''$	so $a \in A_{p'}' \cap A_{p''}'$. Again by definition, having a nonempty intersection means $p' = p''$. So we
fong-spivak-seven-sketches- F02907	273	■ 1.000	$A_{\{p\}} = A_{\{p^{\wedge \prime}\}}$	$A_p = A_{p'}$	conclude that $A_p = A_{p'}$.
fong-spivak-seven-sketches- F02908	273	■ 0.987	$a \in A_{\{p\}} \cap A_{\{q\}}$	$a \in A_p \cap A_q$	otherwise and derive a contradiction. So suppose there exists $p \in A_p \cap A_q$; we will show that
fong-spivak-seven-sketches- F02909	273	■ 1.000	$A_{\{p\}} = A_{\{q\}}$	$A_p = A_q$	$A_p = A_q$, which contradicts an earlier hypothesis. To show that these two subsets are equal, it
fong-spivak-seven-sketches- F02910	273	■ 1.000	$a^{\wedge \prime} \in A_{\{p\}}$	$a' \in A_p$	suffices to show that $a' \in A_p$ iff $a' \in A_q$ for all $a' \in A$. Suppose $a' \in A_p$; then because A_p is
fong-spivak-seven-sketches- F02911	273	■ 1.000	$a^{\wedge \prime} \in A_{\{q\}}$	$a' \in A_q$	suffices to show that $a' \in A_p$ iff $a' \in A_q$ for all $a' \in A$. Suppose $a' \in A_p$; then because A_p is
fong-spivak-seven-sketches- F02912	273	■ 1.000	$a^{\wedge \prime} \in A$	$a' \in A$	suffices to show that $a' \in A_p$ iff $a' \in A_q$ for all $a' \in A$. Suppose $a' \in A_p$; then because A_p is
fong-spivak-seven-sketches- F02913	273	■ 1.000	$a \sim a^{\wedge \prime}$	$a \sim a'$	connected, we have $a \sim a'$. And because A_q is closed, $a' \in A_q$. In just the way, if $a' \in A_q$ then

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_273_F02914	273	■ 1.000	$X := \left\{ a^{\prime} \in A \mid a^{\prime} \sim a \right\}$	$X := \{a' \in A \mid a' \sim a\}$	there is some closed and connected subset containing a . Let $X := \{a' \in A \mid a' \sim a\}$; we claim it is
fong-spivak-seven-sketches_273_F02915	273	■ 0.996	$a^{\prime} \in X$	$a' \in X$	closed and connected and contains a . To see X is closed, suppose $a' \in X$ and $b \sim a'$; then $b \sim a$
fong-spivak-seven-sketches_273_F02916	273	■ 0.996	$b \sim a^{\prime}$	$b \sim a'$	closed and connected and contains a . To see X is closed, suppose $a' \in X$ and $b \sim a'$; then $b \sim a$
fong-spivak-seven-sketches_273_F02917	273	■ 0.996	$b \sim a$	$b \sim a$	closed and connected and contains a . To see X is closed, suppose $a' \in X$ and $b \sim a'$; then $b \sim a$
fong-spivak-seven-sketches_273_F02918	273	■ 0.983	$b \in X$	$b \in X$	by transitivity and symmetry of $\sim$, so $b \in X$. To see that X is connected, suppose $b, c \in X$; then
fong-spivak-seven-sketches_273_F02919	273	■ 0.983	$b, c \in X$	$b, c \in X$	by transitivity and symmetry of $\sim$, so $b \in X$. To see that X is connected, suppose $b, c \in X$; then
fong-spivak-seven-sketches_273_F02920	273	■ 0.988	$c \sim a$	$c \sim a$	$b \sim a$ and $c \sim a$ so $b \sim c$ by the transitivity and symmetry of $\sim$. Finally, $a \in X$ by the reflexivity
fong-spivak-seven-sketches_273_F02921	273	■ 0.988	$a \in X$	$a \in X$	$b \sim a$ and $c \sim a$ so $b \sim c$ by the transitivity and symmetry of $\sim$. Finally, $a \in X$ by the reflexivity
fong-spivak-seven-sketches_273_F02922	273	■ 1.000	$\varnothing \rightarrow \{1\}$	$\emptyset \rightarrow \{1\}$	1. The unique function $\emptyset \rightarrow \{1\}$ is injective but not surjective.
fong-spivak-seven-sketches_273_F02923	273	■ 1.000	$\{a, b\} \rightarrow \{1\}$	$\{a, b\} \rightarrow \{1\}$	2. The unique function $\{a, b\} \rightarrow \{1\}$ is surjective but not injective.
fong-spivak-seven-sketches_273_F02924	273	■ 1.000	$F \subseteq A \times \varnothing$	$F \subseteq A \times \emptyset$	By Definition 1.22, a function $f: A \rightarrow \emptyset$ is a subset $F \subseteq A \times \emptyset$ such that for all $a \in A$, there exists a
fong-spivak-seven-sketches_273_F02925	273	■ 0.995	$b \in \varnothing$	$b \in \emptyset$	unique $b \in \emptyset$ with $(a, b) \in F$. But there are no elements $b \in \emptyset$, so if F is to have the above property,
fong-spivak-seven-sketches_273_F02926	273	■ 0.995	$(a, b) \in F$	$(a, b) \in F$	unique $b \in \emptyset$ with $(a, b) \in F$. But there are no elements $b \in \emptyset$, so if F is to have the above property,
fong-spivak-seven-sketches_273_F02927	273	■ 0.995	$b \in^? \varnothing$	$b \in^? \emptyset$	unique $b \in \emptyset$ with $(a, b) \in F$. But there are no elements $b \in \emptyset$, so if F is to have the above property,
fong-spivak-seven-sketches_274_F02928	274	■ 0.986	$\{\bullet, *, \circ\}$	$\{\bullet, *, \circ\}$	Below each partition, we draw a corresponding surjection out of $\{\bullet, *, \circ\}$:
fong-spivak-seven-sketches_274_F02929	274	■ 0.963	$P := V = \{1, 2, 3, 4\}$	$P := V = \{1, 2, 3, 4\}$	working. The preorder $(P, \leq)$ is given by taking $P := V = \{1, 2, 3, 4\}$ and writing $p \leq q$ whenever there
fong-spivak-seven-sketches_275_F02930	275	■ 1.000	$4 \not\leq 6$	$4 \not\leq 6$	No, it is not a total order; for example $4 \not\leq 6$ and $6 \not\leq 4$.
fong-spivak-seven-sketches_275_F02931	275	■ 1.000	$6 \not\leq 4$	$6 \not\leq 4$	No, it is not a total order; for example $4 \not\leq 6$ and $6 \not\leq 4$.
fong-spivak-seven-sketches_275_F02932	275	■ 1.000	$a, b \in \mathbb{R}$	$a, b \in \mathbb{R}$	Yes, the usual $\leq$ ordering is a total order on $\mathbb{R}$: for every $a, b \in \mathbb{R}$ either $a \leq b$ or $b \leq a$.
fong-spivak-seven-sketches_275_F02933	275	■ 0.996	$!: S \rightarrow \{1\}$	$!: S \rightarrow \{1\}$	The coarsest partition on S corresponds to the unique function $!: S \rightarrow \{1\}$. The finest partition on S
fong-spivak-seven-sketches_275_F02934	275	■ 0.967	$\operatorname{id}_S: S \rightarrow S$	$\operatorname{id}_S: S \rightarrow S$	corresponds to the identity function $\operatorname{id}_S: S \rightarrow S$.
fong-spivak-seven-sketches_275_F02935	275	■ 0.505	$\operatorname{U}(X)$	$\operatorname{U}(X)$	such that $p \leq q$ is p itself, so q is definitely in U ! This means that $\operatorname{U}(X)$ contains all subsets of X , so it's
fong-spivak-seven-sketches_275_F02936	275	■ 1.000	$\operatorname{U}(X) = \operatorname{P}(X)$	$\operatorname{U}(X) = \operatorname{P}(X)$	exactly the power set, $\operatorname{U}(X) = \operatorname{P}(X)$.
fong-spivak-seven-sketches_276_F02937	276	■ 1.000	$0 \leq \cdots \leq 3$	$0 \leq \cdots \leq 3$	With $X = \{0, 1, 2\}$, the Hasse diagram for $\operatorname{P}(X)$, the preorder $0 \leq \cdots \leq 3$, and the cardinality map
fong-spivak-seven-sketches_276_F02938	276	■ 1.000	$q \in \uparrow p$	$q \in \uparrow p$	1. Let $q \in \uparrow p$, and suppose $q \leq q'$. Since $q \in \uparrow p$, we have $p \leq q$. Thus by transitivity $p \leq q'$, so
fong-spivak-seven-sketches_276_F02939	276	■ 1.000	$p \leq q^{\prime}$	$p \leq q'$	1. Let $q \in \uparrow p$, and suppose $q \leq q'$. Since $q \in \uparrow p$, we have $p \leq q$. Thus by transitivity $p \leq q'$, so
fong-spivak-seven-sketches_276_F02940	276	■ 0.991	$q^{\prime} \in \uparrow p$	$q' \in \uparrow p$	$q' \in \uparrow p$. Thus $\uparrow p$ is an upper set.
fong-spivak-seven-sketches_276_F02941	276	■ 0.589	$q \leq^{\operatorname{op}} p$	$q \leq^{\operatorname{op}} p$	2. Suppose $p \leq q$ in P ; this means that $q \leq^{\operatorname{op}} p$ in P^{op} . We must show that $\uparrow q \subseteq \uparrow p$. Take any
fong-spivak-seven-sketches_276_F02942	276	■ 0.589	$\uparrow q \subseteq \uparrow p$	$\uparrow q \subseteq \uparrow p$	2. Suppose $p \leq q$ in P ; this means that $q \leq^{\operatorname{op}} p$ in P^{op} . We must show that $\uparrow q \subseteq \uparrow p$. Take any

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_277 F02972	277	■ 0.998	$\Phi(Q)=$	$\Phi(Q) =$	$\mathbb{B}$. So suppose P, Q are partitions with $P \leq Q$; we need to show that if $\Phi(P) = \text{true}$ then $\Phi(Q) = \text{true}$.
fong-spivak-seven-sketches_277 F02973	277	■ 0.997	$x \sim_P y$	$x \sim_P y$	stuff together. Technically, $x \sim_P y$ implies $x \sim_Q y$ for all $x, y \in \{\bullet, *, \circ\}$. Applying this to $\bullet, *$ gives the
fong-spivak-seven-sketches_277 F02974	277	■ 0.997	$x \sim_Q y$	$x \sim_Q y$	stuff together. Technically, $x \sim_P y$ implies $x \sim_Q y$ for all $x, y \in \{\bullet, *, \circ\}$. Applying this to $\bullet, *$ gives the
fong-spivak-seven-sketches_277 F02975	277	■ 0.997	$x, y \in \{\bullet, *, \circ\}$	$x, y \in \{\bullet, *, \circ\}$	stuff together. Technically, $x \sim_P y$ implies $x \sim_Q y$ for all $x, y \in \{\bullet, *, \circ\}$. Applying this to $\bullet, *$ gives the
fong-spivak-seven-sketches_277 F02976	277	■ 0.997	$\bullet, *$	$\bullet, *$	stuff together. Technically, $x \sim_P y$ implies $x \sim_Q y$ for all $x, y \in \{\bullet, *, \circ\}$. Applying this to $\bullet, *$ gives the
fong-spivak-seven-sketches_277 F02977	277	■ 0.642	$U \mapsto f^{-1}(U)$	$U \mapsto f^{-1}(U)$	Given a function $f: P \rightarrow Q$, we have $f^*: \mathcal{U}(Q) \rightarrow \mathcal{U}(P)$ given by $U \mapsto f^{-1}(U)$. But upper sets in Q are
fong-spivak-seven-sketches_277 F02978	277	■ 1.000	$f^*(U)$	$f^*(U)$	classified by monotone maps $u: Q \rightarrow \mathbb{B}$, and similarly for P ; our job is to show that $f^*(U)$ is given by
fong-spivak-seven-sketches_277 F02979	277	■ 1.000	$q \mapsto$	$q \mapsto$	Given an upper set $U \subseteq Q$, let $u: Q \rightarrow \mathbb{B}$ be the corresponding monotone map, which sends $q \mapsto \text{true}$
fong-spivak-seven-sketches_277 F02980	277	■ 0.890	$p \mapsto$	$p \mapsto$	iff $q \in U$. Then $(f \circ u): P \rightarrow \mathbb{B}$ sends $p \mapsto \text{true}$ iff $f(p) \in U$; it corresponds to the upper set
fong-spivak-seven-sketches_277 F02981	277	■ 0.890	$f(p) \in U$	$f(p) \in U$	iff $q \in U$. Then $(f \circ u): P \rightarrow \mathbb{B}$ sends $p \mapsto \text{true}$ iff $f(p) \in U$; it corresponds to the upper set
fong-spivak-seven-sketches_277 F02982	277	■ 1.000	$\{p \in P \mid f(p) \in U\}$	$\{p \in P \mid f(p) \in U\}$	$\{p \in P \mid f(p) \in U\}$ which is exactly $f^{-1}(U)$.
fong-spivak-seven-sketches_277 F02983	277	■ 1.000	$S = \left\{ \frac{1}{n+1} \mid n \in \mathbb{N} \right\}$	$S = \left\{ \frac{1}{n+1} \mid n \in \mathbb{N} \right\}$	1. 0 is a lower bound for $S = \{ \frac{1}{n+1} \mid n \in \mathbb{N} \}$ because $0 \leq \frac{1}{n+1}$ for any $n \in \mathbb{N}$.
fong-spivak-seven-sketches_277 F02984	277	■ 1.000	$0 \leq \frac{1}{n+1}$	$0 \leq \frac{1}{n+1}$	1. 0 is a lower bound for $S = \{ \frac{1}{n+1} \mid n \in \mathbb{N} \}$ because $0 \leq \frac{1}{n+1}$ for any $n \in \mathbb{N}$.
fong-spivak-seven-sketches_277 F02985	277	■ 1.000	$b \leq 0$	$b \leq 0$	2. Suppose that b is a lower bound for S ; we want to see that $b \leq 0$. If one believes to the contrary
fong-spivak-seven-sketches_277 F02986	277	■ 1.000	$0 < b$	$0 < b$	that $0 < b$, then consider $1/b$; it is a real number so we can find a natural number n that's bigger
fong-spivak-seven-sketches_277 F02987	277	■ 1.000	$1 / b$	$1/b$	that $0 < b$, then consider $1/b$; it is a real number so we can find a natural number n that's bigger
fong-spivak-seven-sketches_277 F02988	277	■ 1.000	$1 / b < n < n+1$	$1/b < n < n+1$	$1/b < n < n+1$. This implies $1 < b(n+1)$ and hence $\frac{1}{n+1} < b$, but that is a contradiction of b
fong-spivak-seven-sketches_277 F02989	277	■ 1.000	$1 < b(n+1)$	$1 < b(n+1)$	$1/b < n < n+1$. This implies $1 < b(n+1)$ and hence $\frac{1}{n+1} < b$, but that is a contradiction of b
fong-spivak-seven-sketches_277 F02990	277	■ 1.000	$\frac{1}{n+1} < b$	$\frac{1}{n+1} < b$	$1/b < n < n+1$. This implies $1 < b(n+1)$ and hence $\frac{1}{n+1} < b$, but that is a contradiction of b
fong-spivak-seven-sketches_277 F02991	277	■ 1.000	$a = p$	$a = p$	$q \leq p$. But the only $a \in A$ is $a = p$, so both are obvious.
fong-spivak-seven-sketches_277 F02992	277	■ 1.000	$4 \wedge 6 = 2$	$4 \wedge 6 = 2$	both are 1 and 2, and 2 is higher, so $4 \wedge 6 = 2$. Similar reasoning shows that $4 \vee 6 = 12$. The meet is the
fong-spivak-seven-sketches_277 F02993	277	■ 1.000	$4 \vee 6 = 12$	$4 \vee 6 = 12$	both are 1 and 2, and 2 is higher, so $4 \wedge 6 = 2$. Similar reasoning shows that $4 \vee 6 = 12$. The meet is the
fong-spivak-seven-sketches_278 F02994	278	■ 1.000	$a \leq a \vee b$	$a \leq a \vee b$	Since f is monotone, the facts that $a \leq a \vee b$ and $b \leq a \vee b$ imply that $f(a) \leq f(a \vee b)$ and $f(b) \leq f(a \vee b)$.
fong-spivak-seven-sketches_278 F02995	278	■ 1.000	$b \leq a \vee b$	$b \leq a \vee b$	Since f is monotone, the facts that $a \leq a \vee b$ and $b \leq a \vee b$ imply that $f(a) \leq f(a \vee b)$ and $f(b) \leq f(a \vee b)$.
fong-spivak-seven-sketches_278 F02996	278	■ 1.000	$f(a) \leq f(a \vee b)$	$f(a) \leq f(a \vee b)$	Since f is monotone, the facts that $a \leq a \vee b$ and $b \leq a \vee b$ imply that $f(a) \leq f(a \vee b)$ and $f(b) \leq f(a \vee b)$.
fong-spivak-seven-sketches_278 F02997	278	■ 1.000	$f(b) \leq f(a \vee b)$	$f(b) \leq f(a \vee b)$	Since f is monotone, the facts that $a \leq a \vee b$ and $b \leq a \vee b$ imply that $f(a) \leq f(a \vee b)$ and $f(b) \leq f(a \vee b)$.
fong-spivak-seven-sketches_278 F02998	278	■ 0.771	$\lfloor -/3 \rfloor$	$\lfloor -/3 \rfloor$	By analogy with Example 1.97, the right adjoint for $(3 \times -)$ should be $\lfloor -/3 \rfloor$. But to prove this is correct,
fong-spivak-seven-sketches_278 F02999	278	■ 1.000	$r \in \mathbb{R}$	$r \in \mathbb{R}$	we must show that for any any $r \in \mathbb{R}$ and $z \in \mathbb{Z}$ we have $z \leq \lfloor r/3 \rfloor$ iff $3 * z \leq r$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_278 F03029	278	■ 1.000	$r=.01$	$r = .01$	by definition of adjunction. So take $z = 1$ and $r = .01$; then $\lceil r/3 \rceil = 1$ so $z \leq C(r)$, and hence
fong-spivak-seven-sketches_278 F03030	278	■ 1.000	$\lceil r / 3 \rceil = 1$	$\lceil r/3 \rceil = 1$	by definition of adjunction. So take $z = 1$ and $r = .01$; then $\lceil r/3 \rceil = 1$ so $z \leq C(r)$, and hence
fong-spivak-seven-sketches_278 F03031	278	■ 1.000	$L(1) \leq 0.01$	$L(1) \leq 0.01$	$L(z) \leq r$, i.e. $L(1) \leq 0.01$. In the same way $L(1) \leq r$ for all $r > 0$, so $L(1) \leq 0$. By definition of
fong-spivak-seven-sketches_278 F03032	278	■ 1.000	$L(1) \leq r$	$L(1) \leq r$	$L(z) \leq r$, i.e. $L(1) \leq 0.01$. In the same way $L(1) \leq r$ for all $r > 0$, so $L(1) \leq 0$. By definition of
fong-spivak-seven-sketches_278 F03033	278	■ 1.000	$r > 0$	$r > 0$	$L(z) \leq r$, i.e. $L(1) \leq 0.01$. In the same way $L(1) \leq r$ for all $r > 0$, so $L(1) \leq 0$. By definition of
fong-spivak-seven-sketches_278 F03034	278	■ 1.000	$L(1) \leq 0$	$L(1) \leq 0$	$L(z) \leq r$, i.e. $L(1) \leq 0.01$. In the same way $L(1) \leq r$ for all $r > 0$, so $L(1) \leq 0$. By definition of
fong-spivak-seven-sketches_278 F03035	278	■ 1.000	$1 \leq C(0) = \lceil 0 / 3 \rceil = 0$	$1 \leq C(0) = \lceil 0/3 \rceil = 0$	adjunction $1 \leq C(0) = \lceil 0/3 \rceil = 0$, a contradiction.
fong-spivak-seven-sketches_278 F03036	278	■ 1.000	$c_{\{1\}}, c_{\{2\}}, c_{\{3\}}, c_{\{4\}}$	c_1, c_2, c_3, c_4	Example 1.102. Take c_1, c_2, c_3, c_4 to be the following partitions:
fong-spivak-seven-sketches_279 F03037	279	■ 0.995	$g_{\{1\}}(c_{\{1\}}), g_{\{1\}}(c_{\{2\}}), g_{\{1\}}(c_{\{3\}})$	$g_1(c_1), g_1(c_2), g_1(c_3)$	Then the induced partitions $g_1(c_1), g_1(c_2), g_1(c_3)$, and $g_1(c_4)$ on T are:
fong-spivak-seven-sketches_279 F03038	279	■ 0.995	$g_{\{1\}}(c_{\{4\}})$	$g_1(c_4)$	Then the induced partitions $g_1(c_1), g_1(c_2), g_1(c_3)$, and $g_1(c_4)$ on T are:
fong-spivak-seven-sketches_279 F03039	279	■ 1.000	$S = \{1, 2, 3, 4\}$	$S = \{1, 2, 3, 4\}$	Here are the partitions on $S = \{1, 2, 3, 4\}$ induced via g^* by the five partitions on $T = \{12, 3, 4\}$:
fong-spivak-seven-sketches_279 F03040	279	■ 1.000	$f: P \rightarrowtail Q: g$	$f: P \rightarrowtail Q: g$	Suppose P and Q are preorders, and that $f: P \rightarrowtail Q: g$ are monotone maps.
fong-spivak-seven-sketches_279 F03041	279	■ 1.000	$g(q) \leq g(q)$	$g(q) \leq g(q)$	$q \in Q$. Then starting with the reflexivity fact $g(q) \leq g(q)$, the definition with $p := g(q)$ gives
fong-spivak-seven-sketches_279 F03042	279	■ 1.000	$p := g(q)$	$p := g(q)$	$q \in Q$. Then starting with the reflexivity fact $g(q) \leq g(q)$, the definition with $p := g(q)$ gives
fong-spivak-seven-sketches_279 F03043	279	■ 1.000	$f(p) \leq f(g(q))$	$f(p) \leq f(g(q))$	sides, we have $f(p) \leq f(g(q))$, and then by transitivity $f(g(q)) \leq q$ implies $f(p) \leq q$, as desired.
fong-spivak-seven-sketches_280 F03044	280	■ 1.000	$g, g^{\prime}: Q \rightarrowtail P$	$g, g': Q \rightarrowtail P$	1. Suppose that $f: P \rightarrowtail Q$ has two right adjoints, $g, g': Q \rightarrowtail P$. We want to show that $g(q) \cong g'(q)$
fong-spivak-seven-sketches_280 F03045	280	■ 1.000	$g(q) \leq g^{\prime}(q)$	$g(q) \leq g'(q)$	for all $q \in Q$. We will prove $g(q) \leq g'(q)$; the inequality $g'(q) \leq g(q)$ is similar. To do this, we use
fong-spivak-seven-sketches_280 F03046	280	■ 1.000	$g^{\prime}(q) \leq g(q)$	$g'(q) \leq g(q)$	for all $q \in Q$. We will prove $g(q) \leq g'(q)$; the inequality $g'(q) \leq g(q)$ is similar. To do this, we use
fong-spivak-seven-sketches_280 F03047	280	■ 1.000	$p \leq g^{\prime}(f(p))$	$p \leq g'(f(p))$	the fact that $p \leq g'(f(p))$ and $f(g(q)) \leq q$ for all p, q by Eq. (1.108). Then the trick is to reason as
fong-spivak-seven-sketches_280 F03048	280	■ 0.999	$j := \vee A$	$j := \vee A$	Suppose $f: P \rightarrowtail Q$ is left adjoint to $g: Q \rightarrowtail P$. Let $A \subseteq P$ be any subset and let $j := \vee A$ be its join.
fong-spivak-seven-sketches_280 F03049	280	■ 1.000	$f(a) \leq f(j)$	$f(a) \leq f(j)$	Then since f is monotone $f(a) \leq f(j)$ for all $a \in A$, so $f(j)$ is an upper bound for the set $f(A)$. We
fong-spivak-seven-sketches_280 F03050	280	■ 1.000	$f(j)$	$f(j)$	Then since f is monotone $f(a) \leq f(j)$ for all $a \in A$, so $f(j)$ is an upper bound for the set $f(A)$. We
fong-spivak-seven-sketches_280 F03051	280	■ 1.000	$f(a) \leq b$	$f(a) \leq b$	$f(a) \leq b$ for all $a \in A$. Then by definition of adjunction, we also have $a \leq g(b)$ for all $a \in A$. By
fong-spivak-seven-sketches_280 F03052	280	■ 1.000	$a \leq g(b)$	$a \leq g(b)$	$f(a) \leq b$ for all $a \in A$. Then by definition of adjunction, we also have $a \leq g(b)$ for all $a \in A$. By
fong-spivak-seven-sketches_280 F03053	280	■ 1.000	$j \leq g(b)$	$j \leq g(b)$	definition of join, we have $j \leq g(b)$. Again by definition of adjunction $f(j) \leq b$, as desired.
fong-spivak-seven-sketches_280 F03054	280	■ 1.000	$f(j) \leq b$	$f(j) \leq b$	definition of join, we have $j \leq g(b)$. Again by definition of adjunction $f(j) \leq b$, as desired.

Identifier	Page	Conf.	LaTeX source	Rendered	Image																																																																																											
fong-spivak-seven-sketches F03055	280	■ 0.941	$f(p)$	$f(p)$	<table><tr><th>p</th><th>q</th><th>$f(p)$</th><th>$g(q)$</th><th>$f(p) \leq^? q$</th><th>$p \leq^? g(q)$</th><th>same?</th></tr><tr><td>1</td><td>1</td><td>1</td><td>1</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>1</td><td>2</td><td>1</td><td>2</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>1</td><td>4</td><td>1</td><td>4</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>2</td><td>1</td><td>2</td><td>1</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>2</td><td>2</td><td>2</td><td>2</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>2</td><td>4</td><td>2</td><td>4</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>3.9</td><td>1</td><td>4</td><td>1</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>3.9</td><td>2</td><td>4</td><td>2</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>3.9</td><td>4</td><td>4</td><td>4</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>4</td><td>1</td><td>4</td><td>1</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>4</td><td>2</td><td>4</td><td>2</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>4</td><td>4</td><td>4</td><td>4</td><td>yes</td><td>yes</td><td>yes!</td></tr></table>	p	q	$f(p)$	$g(q)$	$f(p) \leq^? q$	$p \leq^? g(q)$	same?	1	1	1	1	yes	yes	yes!	1	2	1	2	no	no	yes!	1	4	1	4	yes	yes	yes!	2	1	2	1	no	no	yes!	2	2	2	2	yes	yes	yes!	2	4	2	4	yes	yes	yes!	3.9	1	4	1	no	no	yes!	3.9	2	4	2	no	no	yes!	3.9	4	4	4	yes	yes	yes!	4	1	4	1	no	no	yes!	4	2	4	2	no	no	yes!	4	4	4	4	yes	yes	yes!
p	q	$f(p)$	$g(q)$	$f(p) \leq^? q$	$p \leq^? g(q)$	same?																																																																																										
1	1	1	1	yes	yes	yes!																																																																																										
1	2	1	2	no	no	yes!																																																																																										
1	4	1	4	yes	yes	yes!																																																																																										
2	1	2	1	no	no	yes!																																																																																										
2	2	2	2	yes	yes	yes!																																																																																										
2	4	2	4	yes	yes	yes!																																																																																										
3.9	1	4	1	no	no	yes!																																																																																										
3.9	2	4	2	no	no	yes!																																																																																										
3.9	4	4	4	yes	yes	yes!																																																																																										
4	1	4	1	no	no	yes!																																																																																										
4	2	4	2	no	no	yes!																																																																																										
4	4	4	4	yes	yes	yes!																																																																																										
fong-spivak-seven-sketches F03056	280	■ 0.941	$g(q)$	$g(q)$	<table><tr><th>p</th><th>q</th><th>$f(p)$</th><th>$g(q)$</th><th>$f(p) \leq^? q$</th><th>$p \leq^? g(q)$</th><th>same?</th></tr><tr><td>1</td><td>1</td><td>1</td><td>1</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>1</td><td>2</td><td>1</td><td>2</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>1</td><td>4</td><td>1</td><td>4</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>2</td><td>1</td><td>2</td><td>1</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>2</td><td>2</td><td>2</td><td>2</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>2</td><td>4</td><td>2</td><td>4</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>3.9</td><td>1</td><td>4</td><td>1</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>3.9</td><td>2</td><td>4</td><td>2</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>3.9</td><td>4</td><td>4</td><td>4</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>4</td><td>1</td><td>4</td><td>1</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>4</td><td>2</td><td>4</td><td>2</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>4</td><td>4</td><td>4</td><td>4</td><td>yes</td><td>yes</td><td>yes!</td></tr></table>	p	q	$f(p)$	$g(q)$	$f(p) \leq^? q$	$p \leq^? g(q)$	same?	1	1	1	1	yes	yes	yes!	1	2	1	2	no	no	yes!	1	4	1	4	yes	yes	yes!	2	1	2	1	no	no	yes!	2	2	2	2	yes	yes	yes!	2	4	2	4	yes	yes	yes!	3.9	1	4	1	no	no	yes!	3.9	2	4	2	no	no	yes!	3.9	4	4	4	yes	yes	yes!	4	1	4	1	no	no	yes!	4	2	4	2	no	no	yes!	4	4	4	4	yes	yes	yes!
p	q	$f(p)$	$g(q)$	$f(p) \leq^? q$	$p \leq^? g(q)$	same?																																																																																										
1	1	1	1	yes	yes	yes!																																																																																										
1	2	1	2	no	no	yes!																																																																																										
1	4	1	4	yes	yes	yes!																																																																																										
2	1	2	1	no	no	yes!																																																																																										
2	2	2	2	yes	yes	yes!																																																																																										
2	4	2	4	yes	yes	yes!																																																																																										
3.9	1	4	1	no	no	yes!																																																																																										
3.9	2	4	2	no	no	yes!																																																																																										
3.9	4	4	4	yes	yes	yes!																																																																																										
4	1	4	1	no	no	yes!																																																																																										
4	2	4	2	no	no	yes!																																																																																										
4	4	4	4	yes	yes	yes!																																																																																										
fong-spivak-seven-sketches F03057	280	■ 0.941	$p \leq^? g(q)$	$p \leq^? g(q)$	<table><tr><th>p</th><th>q</th><th>$f(p)$</th><th>$g(q)$</th><th>$f(p) \leq^? q$</th><th>$p \leq^? g(q)$</th><th>same?</th></tr><tr><td>1</td><td>1</td><td>1</td><td>1</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>1</td><td>2</td><td>1</td><td>2</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>1</td><td>4</td><td>1</td><td>4</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>2</td><td>1</td><td>2</td><td>1</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>2</td><td>2</td><td>2</td><td>2</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>2</td><td>4</td><td>2</td><td>4</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>3.9</td><td>1</td><td>4</td><td>1</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>3.9</td><td>2</td><td>4</td><td>2</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>3.9</td><td>4</td><td>4</td><td>4</td><td>yes</td><td>yes</td><td>yes!</td></tr><tr><td>4</td><td>1</td><td>4</td><td>1</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>4</td><td>2</td><td>4</td><td>2</td><td>no</td><td>no</td><td>yes!</td></tr><tr><td>4</td><td>4</td><td>4</td><td>4</td><td>yes</td><td>yes</td><td>yes!</td></tr></table>	p	q	$f(p)$	$g(q)$	$f(p) \leq^? q$	$p \leq^? g(q)$	same?	1	1	1	1	yes	yes	yes!	1	2	1	2	no	no	yes!	1	4	1	4	yes	yes	yes!	2	1	2	1	no	no	yes!	2	2	2	2	yes	yes	yes!	2	4	2	4	yes	yes	yes!	3.9	1	4	1	no	no	yes!	3.9	2	4	2	no	no	yes!	3.9	4	4	4	yes	yes	yes!	4	1	4	1	no	no	yes!	4	2	4	2	no	no	yes!	4	4	4	4	yes	yes	yes!
p	q	$f(p)$	$g(q)$	$f(p) \leq^? q$	$p \leq^? g(q)$	same?																																																																																										
1	1	1	1	yes	yes	yes!																																																																																										
1	2	1	2	no	no	yes!																																																																																										
1	4	1	4	yes	yes	yes!																																																																																										
2	1	2	1	no	no	yes!																																																																																										
2	2	2	2	yes	yes	yes!																																																																																										
2	4	2	4	yes	yes	yes!																																																																																										
3.9	1	4	1	no	no	yes!																																																																																										
3.9	2	4	2	no	no	yes!																																																																																										
3.9	4	4	4	yes	yes	yes!																																																																																										
4	1	4	1	no	no	yes!																																																																																										
4	2	4	2	no	no	yes!																																																																																										
4	4	4	4	yes	yes	yes!																																																																																										
fong-spivak-seven-sketches F03058	281	■ 0.999	$B_{\{1\}} := \{a, b\}$	$B_1 := \{a, b\}$	1. Let $B_1 := \{a, b\}$ and $B_2 := \{c\}$. Then $f^*(B_1) = \{a_1\}$ and $f^*(B_2) = \{c_1, c_2\}$.																																																																																											
fong-spivak-seven-sketches F03059	281	■ 0.999	$B_{\{2\}} := \{c\}$	$B_2 := \{c\}$	1. Let $B_1 := \{a, b\}$ and $B_2 := \{c\}$. Then $f^*(B_1) = \{a_1\}$ and $f^*(B_2) = \{c_1, c_2\}$.																																																																																											
fong-spivak-seven-sketches F03060	281	■ 0.999	$f^*\left(B_{\{1\}}\right) = \left\{a_{\{1\}}\right\}$	$f^*(B_1) = \{a_1\}$	1. Let $B_1 := \{a, b\}$ and $B_2 := \{c\}$. Then $f^*(B_1) = \{a_1\}$ and $f^*(B_2) = \{c_1, c_2\}$.																																																																																											
fong-spivak-seven-sketches F03061	281	■ 0.999	$f^*\left(B_{\{2\}}\right) = \left\{c_{\{1\}}, c_{\{2\}}\right\}$	$f^*(B_2) = \{c_1, c_2\}$	1. Let $B_1 := \{a, b\}$ and $B_2 := \{c\}$. Then $f^*(B_1) = \{a_1\}$ and $f^*(B_2) = \{c_1, c_2\}$.																																																																																											
fong-spivak-seven-sketches F03062	281	■ 1.000	$A_{\{1\}} := \varnothing$	$A_1 := \varnothing$	2. Let $A_1 := \varnothing$ and $A_2 := \{a_1, c_1\}$. Then $f_i(A_1) = \varnothing$ and $f_i(A_2) = \{a, c\}$.																																																																																											
fong-spivak-seven-sketches F03063	281	■ 1.000	$A_{\{2\}} := \left\{a_{\{1\}}, c_{\{1\}}\right\}$	$A_2 := \{a_1, c_1\}$	2. Let $A_1 := \varnothing$ and $A_2 := \{a_1, c_1\}$. Then $f_i(A_1) = \varnothing$ and $f_i(A_2) = \{a, c\}$.																																																																																											

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03064	281	■ 1.000	$f_{!}\left(A_{1}\right)=\varnothing$	$f_{!}\left(A_{1}\right)=\varnothing$	2. Let $A_1 := \varnothing$ and $A_2 := \{a_1, c_1\}$. Then $f_!(A_1) = \varnothing$ and $f_!(A_2) = \{a, c\}$.
fong-spivak-seven-sketches F03065	281	■ 1.000	$f_{!}\left(A_{2}\right)=\{a, c\}$	$f_{!}\left(A_{2}\right)=\{a, c\}$	2. Let $A_1 := \varnothing$ and $A_2 := \{a_1, c_1\}$. Then $f_!(A_1) = \varnothing$ and $f_!(A_2) = \{a, c\}$.
fong-spivak-seven-sketches F03066	281	■ 1.000	A_{1}	A_1	3. With the same A_1 and A_2 , we compute $f_*(A_1) = \{b\}$ and $f_*(A_2) = \{a, b\}$.
fong-spivak-seven-sketches F03067	281	■ 1.000	A_{2}	A_2	3. With the same A_1 and A_2 , we compute $f_*(A_1) = \{b\}$ and $f_*(A_2) = \{a, b\}$.
fong-spivak-seven-sketches F03068	281	■ 1.000	$f_{*}\left(A_{1}\right)=\{b\}$	$f_{*}\left(A_{1}\right)=\{b\}$	3. With the same A_1 and A_2 , we compute $f_*(A_1) = \{b\}$ and $f_*(A_2) = \{a, b\}$.
fong-spivak-seven-sketches F03069	281	■ 1.000	$f_{*}\left(A_{2}\right)=\{a, b\}$	$f_{*}\left(A_{2}\right)=\{a, b\}$	3. With the same A_1 and A_2 , we compute $f_*(A_1) = \{b\}$ and $f_*(A_2) = \{a, b\}$.
fong-spivak-seven-sketches F03070	281	■ 1.000	$g(f(p))$	$g(f(p))$	1. It is part of the definition of adjunction (Proposition 1.107) that $p \leq g(f(p))$, and of course $g(f(p))$
fong-spivak-seven-sketches F03071	281	■ 1.000	$(f ; g)(p)$	$(f;g)(p)$	and $(f \circ g)(p)$ mean the same thing.
fong-spivak-seven-sketches F03072	281	■ 1.000	$g(f(g(f(p)))) \leq g(f(p))$	$g(f(g(f(p)))) \leq g(f(p))$	2. We want to show that $g(f(g(f(p)))) \leq g(f(p))$ and $g(f(p)) \leq g(f(g(f(p))))$ for all p . The latter
fong-spivak-seven-sketches F03073	281	■ 1.000	$g(f(p)) \leq g(f(g(f(p))))$	$g(f(p)) \leq g(f(g(f(p))))$	2. We want to show that $g(f(g(f(p)))) \leq g(f(p))$ and $g(f(p)) \leq g(f(g(f(p))))$ for all p . The latter
fong-spivak-seven-sketches F03074	281	■ 1.000	$p^{\prime} \leq g\left(f\left(p^{\prime}\right)\right)$	$p' \leq g\left(f\left(p^{\prime}\right)\right)$	is just the fact that $p' \leq g(f(p'))$ for any p' , applied with $g(f(p))$ in place of p' . The former uses
fong-spivak-seven-sketches F03075	281	■ 1.000	$f(g(f(p))) \leq f(p)$	$f(g(f(p))) \leq f(p)$	that $f(g(q)) \leq q$, with $f(p)$ substituted for q : this gives $f(g(f(p))) \leq f(p)$, and then we apply g
fong-spivak-seven-sketches F03076	281	■ 0.911	$a \ b$	ab	We denote tuples (a, b) by ab for space reasons. So the relation $\{(1, 1), (1, 2), (2, 1)\}$ will be denoted
fong-spivak-seven-sketches F03077	281	■ 0.911	$\{(1, 1),(1, 2),(2, 1)\}$	$\{(1, 1), (1, 2), (2, 1)\}$	We denote tuples (a, b) by ab for space reasons. So the relation $\{(1, 1), (1, 2), (2, 1)\}$ will be denoted
fong-spivak-seven-sketches F03078	281	■ 1.000	$S:=\{1,2,3\}$	$S := \{1, 2, 3\}$	Let $S := \{1, 2, 3\}$.
fong-spivak-seven-sketches F03079	281	■ 1.000	$1 \leq 1, 2 \leq 2$	$1 \leq 1, 2 \leq 2$	1. Let $\leq$ be the preorder with $1 \leq 2$, and of course $1 \leq 1, 2 \leq 2$, and $3 \leq 3$. Then $U(\leq) =$
fong-spivak-seven-sketches F03080	281	■ 1.000	$3 \leq 3$	$3 \leq 3$	1. Let $\leq$ be the preorder with $1 \leq 2$, and of course $1 \leq 1, 2 \leq 2$, and $3 \leq 3$. Then $U(\leq) =$
fong-spivak-seven-sketches F03081	281	■ 1.000	$U(\leq)=$	$U(\leq) =$	1. Let $\leq$ be the preorder with $1 \leq 2$, and of course $1 \leq 1, 2 \leq 2$, and $3 \leq 3$. Then $U(\leq) =$
fong-spivak-seven-sketches F03082	281	■ 0.999	$\{(1, 1),(1, 2),(2, 2),(3, 3)\}$	$\{(1, 1), (1, 2), (2, 2), (3, 3)\}$	$\{(1, 1), (1, 2), (2, 2), (3, 3)\}$
fong-spivak-seven-sketches F03083	281	■ 1.000	$Q:=\{(1,1)\}$	$Q := \{(1, 1)\}$	2. Let $Q := \{(1, 1)\}$ and $Q' := \{(2, 1)\}$.
fong-spivak-seven-sketches F03084	281	■ 1.000	$Q^{\prime}:=\{(2,1)\}$	$Q' := \{(2, 1)\}$	2. Let $Q := \{(1, 1)\}$ and $Q' := \{(2, 1)\}$.
fong-spivak-seven-sketches F03085	281	■ 0.825	$\mathrm{Cl}(Q)$	$\mathrm{Cl}(Q)$	3. The closure $\mathrm{Cl}(Q)$ of Q is the smallest preorder containing $(1, 1)$, which is $\mathrm{Cl}(Q) = \{(1, 1), (2, 2), (3, 3)\}$.
fong-spivak-seven-sketches F03086	281	■ 0.825	$(1,1)$	$(1, 1)$	3. The closure $\mathrm{Cl}(Q)$ of Q is the smallest preorder containing $(1, 1)$, which is $\mathrm{Cl}(Q) = \{(1, 1), (2, 2), (3, 3)\}$.
fong-spivak-seven-sketches F03087	281	■ 0.825	$\mathrm{Cl}(Q)=\{(1,1),(2,2),(3,3)\}$	$\mathrm{Cl}(Q) = \{(1, 1), (2, 2), (3, 3)\}$	3. The closure $\mathrm{Cl}(Q)$ of Q is the smallest preorder containing $(1, 1)$, which is $\mathrm{Cl}(Q) = \{(1, 1), (2, 2), (3, 3)\}$.
fong-spivak-seven-sketches F03088	281	■ 0.793	$\mathrm{Cl}\left(Q^{\prime}\right)=\{(1,1),(2,1),(2,2),(3,3)\}$	$\mathrm{Cl}\left(Q^{\prime}\right)=\{(1, 1), (2, 1), (2, 2), (3, 3)\}$	Similarly, $\mathrm{Cl}(Q') = \{(1, 1), (2, 1), (2, 2), (3, 3)\}$. It is easy to see that $\mathrm{Cl}(Q) \sqsubseteq \leq$ because every or-
fong-spivak-seven-sketches F03089	281	■ 0.999	$(2,1)$	$(2, 1)$	4. It is easy to see that $\mathrm{Cl}(Q') \not\sqsubseteq \leq$ because the ordered pair $(2, 1)$ is in $\mathrm{Cl}(Q')$ but is not in $\leq$.
fong-spivak-seven-sketches F03090	281	■ 0.999	$\mathrm{Cl}\left(Q^{\prime}\right)$	$\mathrm{Cl}\left(Q^{\prime}\right)$	4. It is easy to see that $\mathrm{Cl}(Q') \not\sqsubseteq \leq$ because the ordered pair $(2, 1)$ is in $\mathrm{Cl}(Q')$ but is not in $\leq$.
fong-spivak-seven-sketches F03091	282	■ 1.000	$x_{1}=-1, \ x_{2}=0, \ y_{1}=-1$	$x_1 = -1, x_2 = 0, y_1 = -1$	The expert is right! The proposal violates property (a) when $x_1 = -1$, $x_2 = 0$, $y_1 = -1$, and $y_2 = 1$.
fong-spivak-seven-sketches F03092	282	■ 1.000	$y_{2}=1$	$y_2 = 1$	The expert is right! The proposal violates property (a) when $x_1 = -1$, $x_2 = 0$, $y_1 = -1$, and $y_2 = 1$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F03093	282	0.901	$-1 \leq -1$	$-1 \leq -1$	Indeed $-1 \leq -1$ and $0 \leq 1$, but $-1 * 0 = 0 \not\leq -1 = -1 * 1$.
fong-spivak-seven-sketches-F03094	282	0.901	$-1 * 0 = 0 \not\leq -1 = -1 * 1$	$-1 * 0 = 0 \not\leq -1 = -1 * 1$	Indeed $-1 \not\leq -1$ and $0 \leq 1$, but $-1 * 0 = 0 \not\leq -1 = -1 * 1$.
fong-spivak-seven-sketches-F03095	282	0.940	$(\operatorname{Disc}(M), =, *, e)$	$(\operatorname{Disc}(M), =, *, e)$	To check that $(\operatorname{Disc}(M), =, *, e)$ is a symmetric monoidal preorder, we need to check our proposed data
fong-spivak-seven-sketches-F03096	282	1.000	$x_1 \otimes x_2 = x_1 \otimes x_2$	$x_1 \otimes x_2 = x_1 \otimes x_2$	obeys conditions (a)-(d) of Definition 2.2. Condition (a) just states the tautology that $x_1 \otimes x_2 = x_1 \otimes x_2$,
fong-spivak-seven-sketches-F03097	282	1.000	$u \leq u, v \leq v$	$u \leq u, v \leq v$	2. We use reflexivity when we assert that $u \leq u, v \leq v$ and $z \leq z$, and use transitivity to assert that
fong-spivak-seven-sketches-F03098	282	1.000	$z \leq z$	$z \leq z$	2. We use reflexivity when we assert that $u \leq u, v \leq v$ and $z \leq z$, and use transitivity to assert that
fong-spivak-seven-sketches-F03099	282	1.000	$t + u \leq y + z$	$t + u \leq y + z$	the above sequence of inequalities implies the single inequality $t + u \leq y + z$.
fong-spivak-seven-sketches-F03100	282	1.000	$z \rightarrow w$	$z \rightarrow w$	Condition (a), monotonicity, says that if $x \rightarrow y$ and $z \rightarrow w$ are reactions, then $x + z \rightarrow y + w$ is a
fong-spivak-seven-sketches-F03101	282	1.000	$x + z \rightarrow y + w$	$x + z \rightarrow y + w$	Condition (a), monotonicity, says that if $x \rightarrow y$ and $z \rightarrow w$ are reactions, then $x + z \rightarrow y + w$ is a
fong-spivak-seven-sketches-F03102	282	0.991	$1 \mid 1$	$1 \mid 1$	This proposal is not monotonic: we have $1 \mid 1$ and $1 \mid 2$, but $(1 + 1) \nmid (1 + 2)$.
fong-spivak-seven-sketches-F03103	282	0.991	$1 \mid 2$	$1 \mid 2$	This proposal is not monotonic: we have $1 \mid 1$ and $1 \mid 2$, but $(1 + 1) \nmid (1 + 2)$.
fong-spivak-seven-sketches-F03104	282	0.991	$(1+1) \nmid (1+2)$	$(1 + 1) \nmid (1 + 2)$	This proposal is not monotonic: we have $1 \mid 1$ and $1 \mid 2$, but $(1 + 1) \nmid (1 + 2)$.
fong-spivak-seven-sketches-F03105	283	0.936	$z \leq w$	$z \leq w$	2. We need to show that (a) if $x \leq y$ and $z \leq w$, then $\min(x, z) \leq \min(y, w)$, (b) $\min(x, \text{yes}) = x =$
fong-spivak-seven-sketches-F03106	283	0.936	$\min(x, z) \leq \min(y, w)$	$\min(x, z) \leq \min(y, w)$	2. We need to show that (a) if $x \leq y$ and $z \leq w$, then $\min(x, z) \leq \min(y, w)$, (b) $\min(x, \text{yes}) = x =$
fong-spivak-seven-sketches-F03107	283	0.936	$\min(x$	$\min(x$	2. We need to show that (a) if $x \leq y$ and $z \leq w$, then $\min(x, z) \leq \min(y, w)$, (b) $\min(x, \text{yes}) = x =$
fong-spivak-seven-sketches-F03108	283	0.936	$=x=$	$= x =$	2. We need to show that (a) if $x \leq y$ and $z \leq w$, then $\min(x, z) \leq \min(y, w)$, (b) $\min(x, \text{yes}) = x =$
fong-spivak-seven-sketches-F03109	283	0.948	$\min ($	$\min($	$\min(\text{yes}, x)$, (c) $\min(\min(x, y), z) = \min(x, \min(y, z))$, and (d) $\min(x, y) = \min(y, z)$. The most
fong-spivak-seven-sketches-F03110	283	0.948	$x)$	$x)$	$\min(\text{yes}, x)$, (c) $\min(\min(x, y), z) = \min(x, \min(y, z))$, and (d) $\min(x, y) = \min(y, z)$. The most
fong-spivak-seven-sketches-F03111	283	0.948	$\min(\min(x, y), z) = \min(x, \min(y, z))$	$\min(\min(x, y), z) = \min(x, \min(y, z))$	$\min(\text{yes}, x)$, (c) $\min(\min(x, y), z) = \min(x, \min(y, z))$, and (d) $\min(x, y) = \min(y, z)$. The most
fong-spivak-seven-sketches-F03112	283	0.948	$\min(x, y) = \min(y, z)$	$\min(x, y) = \min(y, z)$	$\min(\text{yes}, x)$, (c) $\min(\min(x, y), z) = \min(x, \min(y, z))$, and (d) $\min(x, y) = \min(y, z)$. The most
fong-spivak-seven-sketches-F03113	283	0.991	$(\operatorname{P}(S), \leq, S, \cap)$	$(P(S), \leq, S, \cap)$	Yes, $(P(S), \leq, S, \cap)$ is a symmetric monoidal preorder.
fong-spivak-seven-sketches-F03114	283	1.000	$(P \wedge Q)(n)$	$(P \wedge Q)(n)$	makes the statement $P \wedge Q$, where $(P \wedge Q)(n)$ is true if $P(n)$ and $Q(n)$ are true, and false otherwise.
fong-spivak-seven-sketches-F03115	283	0.833	$\{ \}^{\mathbb{N}}, \leq$	$\mathbb{N}, \leq$	Then $(\operatorname{Prop}^{\mathbb{N}}, \leq, \text{true}, \wedge)$ forms a symmetric monoidal preorder.
fong-spivak-seven-sketches-F03116	283	0.957	$n + 10 \leq 1$	$n + 10 \leq 1$	as “ $n + 10 \leq 1$ ” or “ n is made of cheese.” We can also define $\vee$ such that $(P \vee Q)(n)$ is true if and only if
fong-spivak-seven-sketches-F03117	283	0.957	$(P \vee Q)(n)$	$(P \vee Q)(n)$	as “ $n + 10 \leq 1$ ” or “ n is made of cheese.” We can also define $\vee$ such that $(P \vee Q)(n)$ is true if and only if
fong-spivak-seven-sketches-F03118	283	0.561	$\left(\operatorname{Prop}^{\mathbb{N}}, \leq \right)$	$(\operatorname{Prop}^{\mathbb{N}}, \leq$	at least one of $P(n)$ and $Q(n)$ is true. Then $(\operatorname{Prop}^{\mathbb{N}}, \leq, \text{false}, \vee)$ forms a symmetric monoidal preorder.
fong-spivak-seven-sketches-F03119	283	0.561	$\left. \vee \right)$	$\vee)$	at least one of $P(n)$ and $Q(n)$ is true. Then $(\operatorname{Prop}^{\mathbb{N}}, \leq, \text{false}, \vee)$ forms a symmetric monoidal preorder.
fong-spivak-seven-sketches-F03120	283	0.676	X^{op}	X^{op}	Unitality and associativity have nothing to do with the order in $\mathcal{X}^{\operatorname{op}}$: they simply state that $I \otimes x =$
fong-spivak-seven-sketches-F03121	283	0.676	$I \otimes x =$	$I \otimes x =$	Unitality and associativity have nothing to do with the order in $\mathcal{X}^{\operatorname{op}}$: they simply state that $I \otimes x =$

Identifier	Page	Conf.	LaTeX source	Rendered	Image																									
fong-spivak-seven-sketches-283-F03122	283	0.964	$x = x \otimes I$	$x = x \otimes I$	$x = x \otimes I$, and $(x \otimes y) \otimes z = x \otimes (y \otimes z)$. Since these are true in $\mathcal{X}$, they are true in $\mathcal{X}^{\text{op}}$. Symmetry is																									
fong-spivak-seven-sketches-283-F03123	283	0.506	$y \otimes x$	$y \otimes x$	slightly trickier, since in only asks that $x \otimes y$ is equivalent to $y \otimes x$. Nonetheless, this is still true in $\mathcal{X}^{\text{op}}$																									
fong-spivak-seven-sketches-283-F03124	283	0.506	$x^{\{\mathrm{op}\}}$	x^{op}	slightly trickier, since in only asks that $x \otimes y$ is equivalent to $y \otimes x$. Nonetheless, this is still true in $\mathcal{X}^{\text{op}}$																									
fong-spivak-seven-sketches-283-F03125	283	0.978	$(x \otimes y) \cong (y \otimes x)$	$(x \otimes y) \cong (y \otimes x)$	because it is true in $\mathcal{X}$. Indeed the fact that $(x \otimes y) \cong (y \otimes x)$ in $\mathcal{X}$ means that $(x \otimes y) \leq (y \otimes x)$ and																									
fong-spivak-seven-sketches-283-F03126	283	0.978	$(x \otimes y) \leq (y \otimes x)$	$(x \otimes y) \leq (y \otimes x)$	because it is true in $\mathcal{X}$. Indeed the fact that $(x \otimes y) \cong (y \otimes x)$ in $\mathcal{X}$ means that $(x \otimes y) \leq (y \otimes x)$ and																									
fong-spivak-seven-sketches-283-F03127	283	0.927	$(y \otimes x) \leq (x \otimes y)$	$(y \otimes x) \leq (x \otimes y)$	$(y \otimes x) \leq (x \otimes y)$ in $\mathcal{X}$, which respectively imply that $(y \otimes x) \leq (x \otimes y)$ and $(x \otimes y) \leq (y \otimes x)$ in $\mathcal{X}^{\text{op}}$, and																									
fong-spivak-seven-sketches-283-F03128	283	0.768	$\operatorname{Cost}^{\{\mathrm{op}\}}$	Cost^{op}	1. The preorder Cost^{op} has underlying set $[0, \infty]$, and the usual increasing order on real numbers																									
fong-spivak-seven-sketches-283-F03129	283	0.999	$= \infty \geq 0 = g$	$= \infty \geq 0 = g$	1. The map g is monotonic as $g(\text{false}) = \infty \geq 0 = g(\text{true})$, satisfies condition (a) since $0 \geq 0 =$																									
fong-spivak-seven-sketches-283-F03130	283	0.999	$0 \geq 0 =$	$0 \geq 0 =$	1. The map g is monotonic as $g(\text{false}) = \infty \geq 0 = g(\text{true})$, satisfies condition (a) since $0 \geq 0 =$																									
fong-spivak-seven-sketches-284-F03131	284	0.966	$f(n)=1$	$f(n) = 1$	2. The map $f(n) = 1$ for all $n \in \mathbb{N}$ defines a monoidal monotone $f: (\mathbb{N}, \leq, 0, +) \rightarrow (\mathbb{N}, \leq, 1, *)$. (In																									
fong-spivak-seven-sketches-284-F03132	284	0.966	$f: (\mathbb{N}, \leq, 0, +) \rightarrow (\mathbb{N}, \leq, 1, *)$	$f: (\mathbb{N}, \leq, 0, +) \rightarrow (\mathbb{N}, \leq, 1, *)$	2. The map $f(n) = 1$ for all $n \in \mathbb{N}$ defines a monoidal monotone $f: (\mathbb{N}, \leq, 0, +) \rightarrow (\mathbb{N}, \leq, 1, *)$. (In																									
fong-spivak-seven-sketches-284-F03133	284	0.523	$\mathbb{Z}, \leq, *, 1$	$\mathbb{Z}, \leq, *, 1$	3. $(\mathbb{Z}, \leq, *, 1)$ is not a monoidal preorder because $*$ is not monotone. Indeed $-1 \leq 0$ but $(-1 * -1) \not\leq$																									
fong-spivak-seven-sketches-284-F03134	284	0.523	$-1 \leq 0$	$-1 \leq 0$	3. $(\mathbb{Z}, \leq, *, 1)$ is not a monoidal preorder because $*$ is not monotone. Indeed $-1 \leq 0$ but $(-1 * -1) \not\leq$																									
fong-spivak-seven-sketches-284-F03135	284	0.523	$-1 * -1$	$-1 * -1$	3. $(\mathbb{Z}, \leq, *, 1)$ is not a monoidal preorder because $*$ is not monotone. Indeed $-1 \leq 0$ but $(-1 * -1) \not\leq$																									
fong-spivak-seven-sketches-284-F03136	284	0.523	$\nsubseteq$	$\not\subseteq$	3. $(\mathbb{Z}, \leq, *, 1)$ is not a monoidal preorder because $*$ is not monotone. Indeed $-1 \leq 0$ but $(-1 * -1) \not\leq$																									
fong-spivak-seven-sketches-284-F03137	284	0.998	$(0 * 0)$	$(0 * 0)$	$(0 * 0)$.																									
fong-spivak-seven-sketches-284-F03138	284	0.654	X_P	X_P	a Bool -category $\mathcal{X}_P$ with P as its set of objects, and with $\mathcal{X}_P(p, q) := \text{true}$ if $p \leq q$, and $\mathcal{X}_P(p, q) :=$																									
fong-spivak-seven-sketches-284-F03139	284	0.654	$X_P(p, q) :=$	$X_P(p, q) :=$	a Bool -category $\mathcal{X}_P$ with P as its set of objects, and with $\mathcal{X}_P(p, q) := \text{true}$ if $p \leq q$, and $\mathcal{X}_P(p, q) :=$																									
fong-spivak-seven-sketches-284-F03140	284	0.880	$\operatorname{Ob}(X_P) = P$	$\text{Ob}(X_P) = P$	we construct a preorder with underlying set $\text{Ob}(\mathcal{X}_P) = P$, and with $p \leq q$ if and only if $\mathcal{X}_P(p, q) =$																									
fong-spivak-seven-sketches-284-F03141	284	0.880	$X_P(p, q) =$	$X_P(p, q) =$	we construct a preorder with underlying set $\text{Ob}(\mathcal{X}_P) = P$, and with $p \leq q$ if and only if $\mathcal{X}_P(p, q) =$																									
fong-spivak-seven-sketches-284-F03142	284	0.335	$(\operatorname{Ob}(\mathscr{X}), \leq)$	$(\text{Ob}(\mathcal{X}), \leq)$	2. Let $\mathcal{X}$ be a Bool -category. By the proof of Theorem 2.49, we construct a preorder $(\text{Ob}(\mathcal{X}), \leq)$,																									
fong-spivak-seven-sketches-284-F03143	284	0.948	$\operatorname{Ob}(X)$	$\text{Ob}(X)$	1., we construct a Bool -category $\mathcal{X}'$ whose set of objects is $\text{Ob}(\mathcal{X})$, and such that $\mathcal{X}'(x, y) = \text{true}$																									
fong-spivak-seven-sketches-284-F03144	284	0.948	$X^{\{\mathrm{prime}\}}(x, y) =$	$X'(x, y) =$	1., we construct a Bool -category $\mathcal{X}'$ whose set of objects is $\text{Ob}(\mathcal{X})$, and such that $\mathcal{X}'(x, y) = \text{true}$																									
fong-spivak-seven-sketches-284-F03145	284	0.600	$(\operatorname{Ob}(X), \leq)$	$(\text{Ob}(X), \leq)$	if and only if $x \leq y$ in $(\text{Ob}(\mathcal{X}), \leq)$. But by construction, this means $\mathcal{X}'(x, y) = \mathcal{X}(x, y)$. So we get																									
fong-spivak-seven-sketches-284-F03146	284	0.600	$X^{\{\mathrm{prime}\}}(x, y) = \mathcal{X}(x, y)$	$X'(x, y) = \mathcal{X}(x, y)$	if and only if $x \leq y$ in $(\text{Ob}(\mathcal{X}), \leq)$. But by construction, this means $\mathcal{X}'(x, y) = \mathcal{X}(x, y)$. So we get																									
fong-spivak-seven-sketches-284-F03147	284	0.563	$\left(\mathbb{R}_{\geq 0}, \geq, 0, +\right)$	$(\mathbb{R}_{\geq 0}, \geq, 0, +)$	a category enriched over $(\mathbb{R}_{\geq 0}, \geq, 0, +)$ is that in the latter, infinite distances are not allowed between																									
fong-spivak-seven-sketches-284-F03148	284	0.718	$d(/)$	$d(/)$	<table><tr><th>$d(\nearrow)$</th><th>A</th><th>B</th><th>C</th><th>D</th></tr><tr><td>A</td><td>0</td><td>6</td><td>3</td><td>11</td></tr><tr><td>B</td><td>2</td><td>0</td><td>5</td><td>5</td></tr><tr><td>C</td><td>5</td><td>3</td><td>0</td><td>8</td></tr><tr><td>D</td><td>11</td><td>9</td><td>6</td><td>0</td></tr></table>	$d(\nearrow)$	A	B	C	D	A	0	6	3	11	B	2	0	5	5	C	5	3	0	8	D	11	9	6	0
$d(\nearrow)$	A	B	C	D																										
A	0	6	3	11																										
B	2	0	5	5																										
C	5	3	0	8																										
D	11	9	6	0																										

Identifier	Page	Conf.	LaTeX source	Rendered	Image																
fong-spivak-seven-sketches-F03149	284	0.890	$\min(\mathcal{X}(x, y), \mathcal{X}(y, z)) \leq \mathcal{X}(x, z)$	$\min(\mathcal{X}(x, y), \mathcal{X}(y, z)) \leq \mathcal{X}(x, z)$	to no, maybe, or yes. Moreover, we must have $\mathfrak{X}(x, x) = \text{yes}$ and $\min(\mathfrak{X}(x, y), \mathfrak{X}(y, z)) \leq \mathfrak{X}(x, z)$ for all																
fong-spivak-seven-sketches-F03150	284	1.000	x, y, z	x, y, z	x, y, z . So a NMY-category can be thought of as set of points together with an statement—no, maybe,																
fong-spivak-seven-sketches-F03151	285	0.978	$X(C, D)$	$X(C, D)$	For example, to compute the hom-object $\mathfrak{X}(C, D)$, we notice that there are two paths: $C \rightarrow A \rightarrow$																
fong-spivak-seven-sketches-F03152	285	0.978	$C \rightarrowtail A \rightarrowtail$	$C \rightarrow A \rightarrow$	For example, to compute the hom-object $\mathfrak{X}(C, D)$, we notice that there are two paths: $C \rightarrow A \rightarrow$																
fong-spivak-seven-sketches-F03153	285	1.000	$B \rightarrowtail D$	$B \rightarrow D$	$B \rightarrow D$ and $C \rightarrow D$. For the first path, the intersection is the set $\{\text{boat}\}$. For the second path, the																
fong-spivak-seven-sketches-F03154	285	1.000	$C \rightarrowtail D$	$C \rightarrow D$	$B \rightarrow D$ and $C \rightarrow D$. For the first path, the intersection is the set $\{\text{boat}\}$. For the second path, the																
fong-spivak-seven-sketches-F03155	285	0.946	$\mathcal{X}(x, y) \cap \mathcal{X}(y, z) \subseteq \mathcal{X}(x, z)$	$\mathcal{X}(x, y) \cap \mathcal{X}(y, z) \subseteq \mathcal{X}(x, z)$	we ensure that $\mathfrak{X}(x, y) \cap \mathfrak{X}(y, z) \subseteq \mathfrak{X}(x, z)$ for all x, y, z .																
fong-spivak-seven-sketches-F03156	285	1.000	$(x, y)^{\text{th}}$	$(x, y)^{\text{th}}$	2. The matrix M with $(x, y)^{\text{th}}$ entry equal to the maximum, taken over all paths p from x to y , of																
fong-spivak-seven-sketches-F03157	285	0.571	$M(/)$	$M(/)$	<table><tr><td>$M(/)$</td><td>A</td><td>B</td><td>C</td></tr><tr><td>A</td><td>∞</td><td>6</td><td>10</td></tr><tr><td>B</td><td>10</td><td>∞</td><td>10</td></tr><tr><td>C</td><td>10</td><td>6</td><td>∞</td></tr></table>	$M(/)$	A	B	C	A	∞	6	10	B	10	∞	10	C	10	6	∞
$M(/)$	A	B	C																		
A	∞	6	10																		
B	10	∞	10																		
C	10	6	∞																		
fong-spivak-seven-sketches-F03158	285	1.000	$\min(M(x, y), M(y, z)) \leq M(x, y)$	$\min(M(x, y), M(y, z)) \leq M(x, y)$	monoidal unit ∞ , and because $\min(M(x, y), M(y, z)) \leq M(x, y)$ for all $x, y, z \in \{A, B, C\}$.																
fong-spivak-seven-sketches-F03159	285	1.000	$x, y, z \in \{A, B, C\}$	$x, y, z \in \{A, B, C\}$	monoidal unit ∞ , and because $\min(M(x, y), M(y, z)) \leq M(x, y)$ for all $x, y, z \in \{A, B, C\}$.																
fong-spivak-seven-sketches-F03160	286	0.900	$d(A, B) = d(B, A) = 5$	$d(A, B) = d(B, A) = 5$	2. Let $\mathfrak{X}$ be the Lawvere metric space with two objects A and B , such that $d(A, B) = d(B, A) = 5$.																
fong-spivak-seven-sketches-F03161	286	0.210	$x_f =$	$x_f =$	Then we have $\mathfrak{X}_f = \begin{array}{ c c } \hline A & B \\ \hline \bullet & \bullet \\ \hline \end{array}$ while $\mathfrak{X}_g = \begin{array}{ c } \hline A \longrightarrow B \\ \hline \bullet \longrightarrow \bullet \\ \hline \end{array}$.																
fong-spivak-seven-sketches-F03162	286	0.210	$x_g = \stackrel{A}{\bullet} \longrightarrow$	$x_g = \stackrel{A}{\bullet} \longrightarrow$	Then we have $\mathfrak{X}_f = \begin{array}{ c c } \hline A & B \\ \hline \bullet & \bullet \\ \hline \end{array}$ while $\mathfrak{X}_g = \begin{array}{ c } \hline A \longrightarrow B \\ \hline \bullet \longrightarrow \bullet \\ \hline \end{array}$.																
fong-spivak-seven-sketches-F03163	286	1.000	$d(x, y) \leq d(y, x)$	$d(x, y) \leq d(y, x)$	so for all x, y we must have $d(x, y) \leq d(y, x)$. But this means also that $d(y, x) \leq d(x, y)$, and so																
fong-spivak-seven-sketches-F03164	286	1.000	$d(y, x) \leq d(x, y)$	$d(y, x) \leq d(x, y)$	so for all x, y we must have $d(x, y) \leq d(y, x)$. But this means also that $d(y, x) \leq d(x, y)$, and so																
fong-spivak-seven-sketches-F03165	286	1.000	$d(y, x) = 0$	$d(y, x) = 0$	Now let's consider the skeletality condition. This says that if $d(x, y) = 0$ and $d(y, x) = 0$, then																
fong-spivak-seven-sketches-F03166	286	0.893	$(x, y) \in \mathcal{X} \times \mathcal{Y}$	$(x, y) \in \mathcal{X} \times \mathcal{Y}$	1. Let $(x, y) \in \mathfrak{X} \times \mathfrak{Y}$. Since $\mathfrak{X}$ and $\mathfrak{Y}$ are $\mathcal{V}$ -categories, we have $I \leq \mathfrak{X}(x, x)$ and $I \leq \mathfrak{Y}(y, y)$. Thus																
fong-spivak-seven-sketches-F03167	286	0.893	$I \leq \mathcal{X}(x, x)$	$I \leq \mathfrak{X}(x, x)$	1. Let $(x, y) \in \mathfrak{X} \times \mathfrak{Y}$. Since $\mathfrak{X}$ and $\mathfrak{Y}$ are $\mathcal{V}$ -categories, we have $I \leq \mathfrak{X}(x, x)$ and $I \leq \mathfrak{Y}(y, y)$. Thus																
fong-spivak-seven-sketches-F03168	286	0.893	$I \leq y(y, y)$	$I \leq y(y, y)$	1. Let $(x, y) \in \mathfrak{X} \times \mathfrak{Y}$. Since $\mathfrak{X}$ and $\mathfrak{Y}$ are $\mathcal{V}$ -categories, we have $I \leq \mathfrak{X}(x, x)$ and $I \leq \mathfrak{Y}(y, y)$. Thus																
fong-spivak-seven-sketches-F03169	286	0.999	$I = I \otimes I \leq X(x, x) \otimes y(y, y) = (\mathcal{X} \times \mathcal{Y})((x, y), (x, y))$	$I = I \otimes I \leq X(x, x) \otimes y(y, y) = (\mathcal{X} \times \mathcal{Y})((x, y), (x, y))$	$I = I \otimes I \leq \mathfrak{X}(x, x) \otimes \mathfrak{Y}(y, y) = (\mathfrak{X} \times \mathfrak{Y})((x, y), (x, y))$.																
fong-spivak-seven-sketches-F03170	286	0.955	$y(y_1, y_2) \otimes \mathcal{X}(x_2, x_3) = \mathcal{X}(x_2, x_3) \otimes y(y_1, y_2)$	$y(y_1, y_2) \otimes \mathcal{X}(x_2, x_3) = \mathcal{X}(x_2, x_3) \otimes y(y_1, y_2)$	3. In particular, we use the symmetry, to conclude that $\mathfrak{Y}(y_1, y_2) \otimes \mathfrak{X}(x_2, x_3) = \mathfrak{X}(x_2, x_3) \otimes \mathfrak{Y}(y_1, y_2)$.																
fong-spivak-seven-sketches-F03171	286	0.990	$(\mathbb{R} \times \mathbb{R})((5, 6), (-1, 4)) = \mathbb{R}(5, -1) + \mathbb{R}(6, 4) = 6 + 2 = 8$	$(\mathbb{R} \times \mathbb{R})((5, 6), (-1, 4)) = \mathbb{R}(5, -1) + \mathbb{R}(6, 4) = 6 + 2 = 8$	We just apply Definition 2.74(ii): $(\mathbb{R} \times \mathbb{R})((5, 6), (-1, 4)) = \mathbb{R}(5, -1) + \mathbb{R}(6, 4) = 6 + 2 = 8$.																
fong-spivak-seven-sketches-F03172	286	1.000	$u \leq u^{\prime}$	$u \leq u^{\prime}$	1. The function $-\otimes v: V \rightarrow V$ is monotone, because if $u \leq u^{\prime}$ then $u \otimes v \leq u^{\prime} \otimes v$ by the monotonicity																
fong-spivak-seven-sketches-F03173	286	1.000	$u \otimes v \leq u^{\prime} \otimes v$	$u \otimes v \leq u^{\prime} \otimes v$	1. The function $-\otimes v: V \rightarrow V$ is monotone, because if $u \leq u^{\prime}$ then $u \otimes v \leq u^{\prime} \otimes v$ by the monotonicity																
fong-spivak-seven-sketches-F03174	286	0.996	$a := (v \multimap w)$	$a := (v \multimap w)$	2. Let $a := (v \multimap w)$ in Eq. (2.80). The right hand side is thus $(v \multimap w) \leq (v \multimap w)$, which is true by																

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_286_F03175	286	■ 0.995	$(v \multimap u) \otimes v \leq u \leq u'$	$(v \multimap u) \otimes v \leq u \leq u'$	3. Let $u \leq u'$. Then, using 2., $(v \multimap u) \otimes v \leq u \leq u'$. Applying Eq. (2.80), we thus have
fong-spivak-seven-sketches_286_F03176	286	■ 1.000	$(v \multimap u) \leq (v \multimap u^{\prime})$	$(v \multimap u) \leq (v \multimap u')$	$(v \multimap u) \leq (v \multimap u')$. This shows that the map $(v \multimap -): V \rightarrow V$ is monotone.
fong-spivak-seven-sketches_286_F03177	286	■ 1.000	$(v \multimap -): V \rightarrow V$	$(v \multimap -): V \rightarrow V$	$(v \multimap u) \leq (v \multimap u')$. This shows that the map $(v \multimap -): V \rightarrow V$ is monotone.
fong-spivak-seven-sketches_286_F03178	286	■ 0.859	$(v \multimap -)$	$(v \multimap -)$	not know $(- \otimes v)$ and $(v \multimap -)$ are monotone maps. We proved this, however, in items 1 and 3
fong-spivak-seven-sketches_287_F03179	287	■ 1.000	$x \Rightarrow y$	$x \Rightarrow y$	We need to find the hom-element. This is given by implication. That is, define the function $x \Rightarrow y$ by
fong-spivak-seven-sketches_287_F03180	287	■ 0.989	$(a \wedge v) \leq w$	$(a \wedge v) \leq w$	Then $(a \wedge v) \leq w$ if and only if $a \leq (v \Rightarrow w)$. Indeed, if $v = \text{false}$ then $a \wedge \text{false} = \text{false}$, and so the
fong-spivak-seven-sketches_287_F03181	287	■ 0.989	$a \leq (v \Rightarrow w)$	$a \leq (v \Rightarrow w)$	Then $(a \wedge v) \leq w$ if and only if $(a \wedge v) \leq w$. Indeed, if $v = \text{false}$ then $a \wedge \text{false} = \text{false}$, and so the
fong-spivak-seven-sketches_287_F03182	287	■ 0.989	$v =$	$v =$	Then $(a \wedge v) \leq w$ if and only if $a \leq (v \Rightarrow w)$. Indeed, if $v = \text{false}$ then $a \wedge \text{false} = \text{false}$, and so the
fong-spivak-seven-sketches_287_F03183	287	■ 0.989	$a \wedge$	$a \wedge$	Then $(a \wedge v) \leq w$ if and only if $a \leq (v \Rightarrow w)$. Indeed, if $v = \text{false}$ then $a \wedge \text{false} = \text{false}$, and so the
fong-spivak-seven-sketches_287_F03184	287	■ 0.879	$=a$	$= a$	$v = \text{true}$, then $a \wedge \text{true} = a$ so the left hand side says $a \leq w$. But $(\text{true} \Rightarrow w) = w$, so the right hand
fong-spivak-seven-sketches_287_F03185	287	■ 0.879	$a \leq w$	$a \leq w$	$v = \text{true}$, then $a \wedge \text{true} = a$ so the left hand side says $a \leq w$. But $(\text{true} \Rightarrow w) = w$, so the right hand
fong-spivak-seven-sketches_287_F03186	287	■ 0.879	$\Rightarrow w) = w$	$\Rightarrow w) = w$	$v = \text{true}$, then $a \wedge \text{true} = a$ so the left hand side says $a \leq w$. But $(\text{true} \Rightarrow w) = w$, so the right hand
fong-spivak-seven-sketches_287_F03187	287	■ 0.980	$(\mathrm{P}(S), \subseteq, S, \cap)$	$(\mathrm{P}(S), \subseteq, S, \cap)$	Yes, the powerset monoidal preorder $(\mathrm{P}(S), \subseteq, S, \cap)$ is a quantale. The hom-object $B \multimap C$ is given by
fong-spivak-seven-sketches_287_F03188	287	■ 0.980	$B \multimap C$	$B \multimap C$	Yes, the powerset monoidal preorder $(\mathrm{P}(S), \subseteq, S, \cap)$ is a quantale. The hom-object $B \multimap C$ is given by
fong-spivak-seven-sketches_287_F03189	287	■ 1.000	$\bar{B} \cup C$	$\bar{B} \cup C$	$\bar{B} \cup C$, where $\bar{B}$ is the complement of B : it contains all elements of S not contained in B . To see that this
fong-spivak-seven-sketches_287_F03190	287	■ 1.000	$\bar{B}$	$\bar{B}$	$\bar{B} \cup C$, where $\bar{B}$ is the complement of B : it contains all elements of S not contained in B . To see that this
fong-spivak-seven-sketches_287_F03191	287	■ 1.000	$(A \cap B) \subseteq C$	$(A \cap B) \subseteq C$	satisfies Eq. (2.80), note that if $(A \cap B) \subseteq C$, then
fong-spivak-seven-sketches_287_F03192	287	■ 1.000	$A \subseteq (\bar{B} \cup C)$	$A \subseteq (\bar{B} \cup C)$	On the other hand, if $A \subseteq (\bar{B} \cup C)$, then
fong-spivak-seven-sketches_287_F03193	287	■ 0.558	$(\bigvee \varnothing) =$	$(\bigvee \varnothing) =$	1a. In Bool , $(\bigvee \varnothing) = \text{false}$, the least element.
fong-spivak-seven-sketches_287_F03194	287	■ 0.946	$(\bigvee \varnothing) = \infty$	$(\bigvee \varnothing) = \infty$	1b. In Cost , $(\bigvee \varnothing) = \infty$. This is because we use the opposite order $\geq$ on $[0, \infty]$, so $\bigvee \varnothing$ is the greatest
fong-spivak-seven-sketches_287_F03195	287	■ 0.974	$(\bigvee \varnothing) = 0$	$(\bigvee \varnothing) = 0$	$(\bigvee \varnothing) = 0$, is a bit confusing! Beware!
fong-spivak-seven-sketches_287_F03196	287	■ 1.000	$\min(x, y)$	$\min(x, y)$	2b. In Cost , $x \vee y$ is the minimum $\min(x, y)$. Again because we use the opposite order on $[0, \infty]$, the
fong-spivak-seven-sketches_287_F03197	287	■ 1.000	$v \in V, 0 \otimes v = 0$	$v \in V, 0 \otimes v = 0$	1. We first use Proposition 2.87 (2) and symmetry to show that for all $v \in V$, $0 \otimes v = 0$.
fong-spivak-seven-sketches_289_F03198	289	■ 0.999	$\left(v, a_1, \dots, a_n\right)$	$(v, a_1, \dots, a_n)$	path in G to be a tuple of the form $(v, a_1, \dots, a_n)$ where $v \in V$ is a vertex, $s(a_1) = v$, and $t(a_i) = s(a_{i+1})$
fong-spivak-seven-sketches_289_F03199	289	■ 0.999	$s\left(a_1\right)=v$	$s\left(a_1\right)=v$	path in G to be a tuple of the form $(v, a_1, \dots, a_n)$ where $v \in V$ is a vertex, $s(a_1) = v$, and $t(a_i) = s(a_{i+1})$
fong-spivak-seven-sketches_289_F03200	289	■ 0.999	$t\left(a_i\right)=s\left(a_{i+1}\right)$	$t\left(a_i\right)=s\left(a_{i+1}\right)$	path in G to be a tuple of the form $(v, a_1, \dots, a_n)$ where $v \in V$ is a vertex, $s(a_1) = v$, and $t(a_i) = s(a_{i+1})$
fong-spivak-seven-sketches_289_F03201	289	■ 1.000	$i \in\{1, \ldots, n-1\}$	$i \in\{1, \ldots, n-1\}$	for all $i \in\{1, \ldots, n-1\}$: the length of this path is n , and this definition makes sense for any $n \in \mathbb{N}$. We
fong-spivak-seven-sketches_289_F03202	289	■ 1.000	$s(p):=v$	$s(p):=v$	say that the source of p is $s(p):=v$ and the target of p is defined to be
fong-spivak-seven-sketches_289_F03203	289	■ 1.000	$p=\left(v, a_1, \ldots, a_m\right)$	$p=\left(v, a_1, \ldots, a_m\right)$	Two paths $p=\left(v, a_1, \ldots, a_m\right)$ and $q=\left(w, b_1, \ldots, b_n\right)$ can be concatenated if $t(p)=s(q)$, in which case

Identifier	Page	Conf.	LaTeX source	Rendered	Image																																																	
fong-spivak-seven-sketches-F03204	289	1.000	$q=\left(w, \text{ } b_{\{1\}}, \text{ } \ldots, \text{ } b_{\{n\}}\right)$	$q = (w, b_1, \ldots, b_n)$	Two paths $p = (v, a_1, \ldots, a_m)$ and $q = (w, b_1, \ldots, b_n)$ can be concatenated if $t(p) = s(q)$, in which case																																																	
fong-spivak-seven-sketches-F03205	289	1.000	$t(p)=s(q)$	$t(p) = s(q)$	Two paths $p = (v, a_1, \ldots, a_m)$ and $q = (w, b_1, \ldots, b_n)$ can be concatenated if $t(p) = s(q)$, in which case																																																	
fong-spivak-seven-sketches-F03206	289	0.870	$p=(v)$	$p = (v)$	We are now ready to check unitality and associativity. A path p is an identity in $\mathbf{Free}(G)$ iff $p = (v)$																																																	
fong-spivak-seven-sketches-F03207	289	0.551	(v)	(v)	for some $v \in V$. It is easy to see from the above that (v) and $(w, b_1, \ldots, b_n)$ can be concatenated iff																																																	
fong-spivak-seven-sketches-F03208	289	0.551	$\left(w, \text{ } b_{\{1\}}, \text{ } \ldots, \text{ } b_{\{n\}}\right)$	$(w, b_1, \ldots, b_n)$	for some $v \in V$. It is easy to see from the above that (v) and $(w, b_1, \ldots, b_n)$ can be concatenated iff																																																	
fong-spivak-seven-sketches-F03209	289	1.000	$v=w$	$v = w$	$v = w$, in which case the result is $(w, b_1, \ldots, b_n)$. Similarly $(v, a_1, \ldots, a_m)$ and (w) can be concatenated																																																	
fong-spivak-seven-sketches-F03210	289	1.000	$\left(v, \text{ } a_{\{1\}}, \text{ } \ldots, \text{ } a_{\{m\}}\right)$	$(v, a_1, \ldots, a_m)$	$v = w$, in which case the result is $(w, b_1, \ldots, b_n)$. Similarly $(v, a_1, \ldots, a_m)$ and (w) can be concatenated																																																	
fong-spivak-seven-sketches-F03211	289	1.000	(w)	(w)	$v = w$, in which case the result is $(w, b_1, \ldots, b_n)$. Similarly $(v, a_1, \ldots, a_m)$ and (w) can be concatenated																																																	
fong-spivak-seven-sketches-F03212	289	1.000	$w=t\left(a_{\{m\}}\right)$	$w = t(a_m)$	iff $w = t(a_m)$, in which case the result is $(v, a_1, \ldots, a_m)$. Finally, for associativity with p and q as above																																																	
fong-spivak-seven-sketches-F03213	289	0.996	$r=\left(x, \text{ } c_{\{1\}}, \text{ } \ldots, \text{ } c_{\{o\}}\right)$	$r = (x, c_1, \ldots, c_o)$	and $r = (x, c_1, \ldots, c_o)$, the formula readily reads that whichever way they are concatenated, $(p \circ q) \circ r$																																																	
fong-spivak-seven-sketches-F03214	289	0.996	$(p \text{ ; } q) \text{ ; } r$	$(p;q);r$	and $r = (x, c_1, \ldots, c_o)$, the formula readily reads that whichever way they are concatenated, $(p \circ q) \circ r$																																																	
fong-spivak-seven-sketches-F03215	289	0.952	$p \text{ ;}(q \text{ ; } r)$	$p;(q;r)$	or $p \circ (q \circ r)$, the result is																																																	
fong-spivak-seven-sketches-F03216	289	0.543	$\boxtimes$	$\boxtimes$	id_{v_2} . We write $\boxtimes$ when the composite does not make sense (i.e. when the target of the first morphism																																																	
fong-spivak-seven-sketches-F03217	289	0.306	$f_{\{1\}} \text{ ; } f_{\{2\}}$	$f_1;f_2$	<table><tr><td>$\nearrow$</td><td>v_1</td><td>f_1</td><td>$f_1 \circ f_2$</td><td>v_2</td><td>f_2</td><td>v_3</td></tr><tr><td>v_1</td><td>v_1</td><td>f_1</td><td>$f_1 \circ f_2$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td></tr><tr><td>f_1</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>f_1</td><td>$f_1 \circ f_2$</td><td>$\boxtimes$</td></tr><tr><td>$f_1 \circ f_2$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$f_1 \circ f_2$</td></tr><tr><td>v_2</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>v_2</td><td>f_2</td><td>$\boxtimes$</td></tr><tr><td>f_2</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>f_2</td></tr><tr><td>v_3</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>v_3</td></tr></table>	$\nearrow$	v_1	f_1	$f_1 \circ f_2$	v_2	f_2	v_3	v_1	v_1	f_1	$f_1 \circ f_2$	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_1	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_1	$f_1 \circ f_2$	$\boxtimes$	$f_1 \circ f_2$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$f_1 \circ f_2$	v_2	$\boxtimes$	$\boxtimes$	$\boxtimes$	v_2	f_2	$\boxtimes$	f_2	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_2	v_3	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	v_3
$\nearrow$	v_1	f_1	$f_1 \circ f_2$	v_2	f_2	v_3																																																
v_1	v_1	f_1	$f_1 \circ f_2$	$\boxtimes$	$\boxtimes$	$\boxtimes$																																																
f_1	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_1	$f_1 \circ f_2$	$\boxtimes$																																																
$f_1 \circ f_2$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$f_1 \circ f_2$																																																
v_2	$\boxtimes$	$\boxtimes$	$\boxtimes$	v_2	f_2	$\boxtimes$																																																
f_2	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_2																																																
v_3	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	v_3																																																
fong-spivak-seven-sketches-F03218	289	0.306	$f_{\{2\}}$	f_2	<table><tr><td>$\nearrow$</td><td>v_1</td><td>f_1</td><td>$f_1 \circ f_2$</td><td>v_2</td><td>f_2</td><td>v_3</td></tr><tr><td>v_1</td><td>v_1</td><td>f_1</td><td>$f_1 \circ f_2$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td></tr><tr><td>f_1</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>f_1</td><td>$f_1 \circ f_2$</td><td>$\boxtimes$</td></tr><tr><td>$f_1 \circ f_2$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$f_1 \circ f_2$</td></tr><tr><td>v_2</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>v_2</td><td>f_2</td><td>$\boxtimes$</td></tr><tr><td>f_2</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>f_2</td></tr><tr><td>v_3</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>v_3</td></tr></table>	$\nearrow$	v_1	f_1	$f_1 \circ f_2$	v_2	f_2	v_3	v_1	v_1	f_1	$f_1 \circ f_2$	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_1	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_1	$f_1 \circ f_2$	$\boxtimes$	$f_1 \circ f_2$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$f_1 \circ f_2$	v_2	$\boxtimes$	$\boxtimes$	$\boxtimes$	v_2	f_2	$\boxtimes$	f_2	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_2	v_3	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	v_3
$\nearrow$	v_1	f_1	$f_1 \circ f_2$	v_2	f_2	v_3																																																
v_1	v_1	f_1	$f_1 \circ f_2$	$\boxtimes$	$\boxtimes$	$\boxtimes$																																																
f_1	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_1	$f_1 \circ f_2$	$\boxtimes$																																																
$f_1 \circ f_2$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$f_1 \circ f_2$																																																
v_2	$\boxtimes$	$\boxtimes$	$\boxtimes$	v_2	f_2	$\boxtimes$																																																
f_2	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_2																																																
v_3	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	v_3																																																
fong-spivak-seven-sketches-F03219	289	0.306	$v_{\{3\}}$	v_3	<table><tr><td>$\nearrow$</td><td>v_1</td><td>f_1</td><td>$f_1 \circ f_2$</td><td>v_2</td><td>f_2</td><td>v_3</td></tr><tr><td>v_1</td><td>v_1</td><td>f_1</td><td>$f_1 \circ f_2$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td></tr><tr><td>f_1</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>f_1</td><td>$f_1 \circ f_2$</td><td>$\boxtimes$</td></tr><tr><td>$f_1 \circ f_2$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$f_1 \circ f_2$</td></tr><tr><td>v_2</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>v_2</td><td>f_2</td><td>$\boxtimes$</td></tr><tr><td>f_2</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>f_2</td></tr><tr><td>v_3</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>$\boxtimes$</td><td>v_3</td></tr></table>	$\nearrow$	v_1	f_1	$f_1 \circ f_2$	v_2	f_2	v_3	v_1	v_1	f_1	$f_1 \circ f_2$	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_1	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_1	$f_1 \circ f_2$	$\boxtimes$	$f_1 \circ f_2$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$f_1 \circ f_2$	v_2	$\boxtimes$	$\boxtimes$	$\boxtimes$	v_2	f_2	$\boxtimes$	f_2	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_2	v_3	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	v_3
$\nearrow$	v_1	f_1	$f_1 \circ f_2$	v_2	f_2	v_3																																																
v_1	v_1	f_1	$f_1 \circ f_2$	$\boxtimes$	$\boxtimes$	$\boxtimes$																																																
f_1	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_1	$f_1 \circ f_2$	$\boxtimes$																																																
$f_1 \circ f_2$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$f_1 \circ f_2$																																																
v_2	$\boxtimes$	$\boxtimes$	$\boxtimes$	v_2	f_2	$\boxtimes$																																																
f_2	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	f_2																																																
v_3	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	$\boxtimes$	v_3																																																

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03220	289	0.306	$f_{\{1\}} \stackrel{\circ}{\circ} f_{\{2\}}$	$f_1 \stackrel{\circ}{\circ} f_2$	
fong-spivak-seven-sketches F03221	289	1.000	$1+2+\cdots+n$	$1 + 2 + \cdots + n$	is $1 + 2 + \cdots + n$. This makes sense because (and the proof strategy would be to verify that) the
fong-spivak-seven-sketches F03222	289	0.818	$n-i$	$n - i$	above graph has n paths of length 0, it has $n - 1$ paths of length 1, and so on: it has $n - i$ paths
fong-spivak-seven-sketches F03223	289	1.000	$0 \leq i \leq n$	$0 \leq i \leq n$	of length i for every $0 \leq i \leq n$.
fong-spivak-seven-sketches F03224	290	0.999	$m+n$	$m + n$	a path of length n results in a path of length $m + n$.
fong-spivak-seven-sketches F03225	290	0.663	$h=j=A$	$h = j = A$	Note that $A \circ f \circ h = j = A \circ g \circ i$.
fong-spivak-seven-sketches F03226	290	0.485	z, s, s	z, s, s	There are four morphisms in $\mathcal{D}$, shown below, namely $z, s, s \circ s$, and $s \circ s \circ s$:
fong-spivak-seven-sketches F03227	290	0.485	$s ; s$	$s; s$	There are four morphisms in $\mathcal{D}$, shown below, namely $z, s, s \circ s$, and $s \circ s \circ s$:
fong-spivak-seven-sketches F03228	291	1.000	$(f(1), f(2))$	$(f(1), f(2))$	A function $f: \underline{2} \rightarrow \underline{3}$ can be described as an ordered pair $(f(1), f(2))$. The nine such functions are
fong-spivak-seven-sketches F03229	291	0.985	$n!$	$n!$	number of isomorphisms between them is n -factorial, often denoted $n!$. So for example there are
fong-spivak-seven-sketches F03230	291	0.999	$5 * 4 * 3 * 2 * 1=120$	$5 * 4 * 3 * 2 * 1 = 120$	$5 * 4 * 3 * 2 * 1 = 120$ isomorphisms between $\{1, 2, 3, 4, 5\}$ and $\{a, b, c, d, e\}$.
fong-spivak-seven-sketches F03231	291	0.999	$\{1, 2, 3, 4, 5\}$	$\{1, 2, 3, 4, 5\}$	$5 * 4 * 3 * 2 * 1 = 120$ isomorphisms between $\{1, 2, 3, 4, 5\}$ and $\{a, b, c, d, e\}$.
fong-spivak-seven-sketches F03232	291	0.999	$\{a, b, c, d, e\}$	$\{a, b, c, d, e\}$	$5 * 4 * 3 * 2 * 1 = 120$ isomorphisms between $\{1, 2, 3, 4, 5\}$ and $\{a, b, c, d, e\}$.
fong-spivak-seven-sketches F03233	291	0.566	$f: c \rightarrow c$	$f: c \rightarrow c$	We have to show that for any object $c \in \mathcal{C}$, the identity id_c has an inverse, i.e. a morphism $f: c \rightarrow c$
fong-spivak-seven-sketches F03234	291	0.761	$\mathrm{id}_c = \mathrm{id}_c$	$\mathrm{id}_c = \mathrm{id}_c$	such that $f \circ \mathrm{id}_c = \mathrm{id}_c$ and $\mathrm{id}_c \circ f = \mathrm{id}_c$. Take $f = \mathrm{id}_c$; this works.
fong-spivak-seven-sketches F03235	291	0.761	$f = \mathrm{id}_c$	$f = \mathrm{id}_c$	such that $f \circ \mathrm{id}_c = \mathrm{id}_c$ and $\mathrm{id}_c \circ f = \mathrm{id}_c$. Take $f = \mathrm{id}_c$; this works.
fong-spivak-seven-sketches F03236	291	1.000	s^n	s^n	morphism is of the form s^n for some $n \in \mathbb{N}$ and composing it with s gives s^{n+1} , which is never
fong-spivak-seven-sketches F03237	291	1.000	s^{n+1}	s^{n+1}	morphism is of the form s^n for some $n \in \mathbb{N}$ and composing it with s gives s^{n+1} , which is never
fong-spivak-seven-sketches F03238	291	1.000	$s^{\{0\}} . p$	$s^0.p$	
fong-spivak-seven-sketches F03239	291	0.803	$p ; q = \mathrm{id}$	$p; q = \mathrm{id}$	is 0. In order for p to be an isomorphism, there must be some q such that $p \circ q = \mathrm{id}$ and $q \circ p = \mathrm{id}$, in
fong-spivak-seven-sketches F03240	291	0.803	$q ; p = \mathrm{id}$	$q; p = \mathrm{id}$	is 0. In order for p to be an isomorphism, there must be some q such that $p \circ q = \mathrm{id}$ and $q \circ p = \mathrm{id}$, in
fong-spivak-seven-sketches F03241	291	0.961	$\mathbf{2} \rightarrow \mathbf{3}$	$\mathbf{2} \rightarrow \mathbf{3}$	The other three functors $\mathbf{2} \rightarrow \mathbf{3}$ are shown here:
fong-spivak-seven-sketches F03242	292	1.000	$g^{\{\prime\}} ; i^{\{\prime\}} \mapsto f ; h$	$g'; i' \mapsto f; h$	If one of these seems different from the rest, it's probably $g' \circ i' \mapsto f \circ h$. But note that in fact also
fong-spivak-seven-sketches F03243	292	0.546	$g^{\{\prime\}} ; i^{\{\prime\}} \mapsto g ; i$	$g'; i' \mapsto g; i$	$g' \circ i' \mapsto g \circ i$ because $g \circ i = f \circ h$, so it's not an outlier after all.
fong-spivak-seven-sketches F03244	292	0.546	$g ; i = f ; h$	$g; i = f; h$	$g' \circ i' \mapsto g \circ i$ because $g \circ i = f \circ h$, so it's not an outlier after all.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches_03245	292	■ 0.449	F, G	F, G	We need to give two functors F, G from $\bullet \xrightarrow{f} \bullet$ to $\bullet \xrightarrow[f_2]{f_1} \bullet$ whose on-objects parts are the same and
fong-spivak-seven-sketches_03246	292	■ 0.449	$\stackrel{\text{a}}{\rightarrow} \stackrel{\text{b}}{\rightarrow}$	$\xrightarrow{a} \xrightarrow{b}$	We need to give two functors F, G from $\bullet \xrightarrow{f} \bullet$ to $\bullet \xrightarrow[f_2]{f_1} \bullet$ whose on-objects parts are the same and
fong-spivak-seven-sketches_03247	292	■ 0.449	$\stackrel{\text{a}^{\prime}}{\underset{\text{f}_2}{\rightarrow}} \stackrel{\text{b}^{\prime}}{\bullet}$	$\xrightarrow[f_2]{a'} \bullet$	We need to give two functors F, G from $\bullet \xrightarrow{f} \bullet$ to $\bullet \xrightarrow[f_2]{f_1} \bullet$ whose on-objects parts are the same and
fong-spivak-seven-sketches_03248	292	■ 0.958	$\operatorname{id}_{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{C}$	$\mathrm{id}_{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{C}$	1. Let $\mathcal{C}$ be a category. Then defining $\mathrm{id}_{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{C}$ by $\mathrm{id}_{\mathcal{C}}(x) = x$ for every object and morphism in
fong-spivak-seven-sketches_03249	292	■ 0.958	$\operatorname{id}_{\mathcal{C}}(x)=x$	$\mathrm{id}_{\mathcal{C}}(x) = x$	1. Let $\mathcal{C}$ be a category. Then defining $\mathrm{id}_{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{C}$ by $\mathrm{id}_{\mathcal{C}}(x) = x$ for every object and morphism in
fong-spivak-seven-sketches_03250 [conf 0.098]	292	■ 0.098	$\operatorname{id}_{\mathcal{C}}(\operatorname{id}_{\mathcal{C}}(c))=\operatorname{id}_{\mathcal{C}}(c)$	identities $\mathrm{id}_{\mathcal{C}}(\mathrm{id}_{\mathcal{C}}) = \mathrm{id}_{\mathcal{C}} = \mathrm{id}_{\mathrm{id}_{\mathcal{C}}(c)}$	$\mathcal{C}$ is a functor because it preserves identities $\mathrm{id}_{\mathcal{C}}(\mathrm{id}_{\mathcal{C}}) = \mathrm{id}_{\mathcal{C}} = \mathrm{id}_{\mathrm{id}_{\mathcal{C}}(c)}$ for each object $c \in \mathrm{Ob}(\mathcal{C})$,
fong-spivak-seven-sketches_03251	292	■ 0.995	$\operatorname{id}_{\mathcal{C}}(f;g)=f;g$	$\mathrm{id}_{\mathcal{C}}(f;g) = f;g = \mathrm{id}_{\mathcal{C}}(f); \mathrm{id}_{\mathcal{C}}(g)$	and it preserves composition $\mathrm{id}_{\mathcal{C}}(f \circ g) = f \circ g = \mathrm{id}_{\mathcal{C}}(f) \circ \mathrm{id}_{\mathcal{C}}(g)$ for each pair of composable
fong-spivak-seven-sketches_03252	292	■ 0.995	$(F;G)(\operatorname{id}_c)=$	$(F;G)(\mathrm{id}_c) =$	it preserves identities and compositions. If $c \in \mathcal{C}$ is an object then $(F \circ G)(\mathrm{id}_c) =$
fong-spivak-seven-sketches_03253	292	■ 0.515	$G(\operatorname{id}_{F(c)})=\operatorname{id}_{G(F(c))}$	$G(\mathrm{id}_{F(c)}) = G(\mathrm{id}_{G(F(c))}) = \mathrm{id}_{G(G(F(c)))}$	$G(\mathrm{id}_c) = G(\mathrm{id}_{F(c)}) = \mathrm{id}_{G(G(F(c)))}$ because F and G preserve identities. If f, g are composable
fong-spivak-seven-sketches_03254	292	■ 0.874	$\operatorname{id}_{\mathcal{C}}$	$\mathrm{id}_{\mathcal{C}}$	and we pre-compose it as above with $\mathrm{id}_{\mathcal{C}}$; it is easy to see that the result will again be F , and
fong-spivak-seven-sketches_03255	292	■ 0.989	$\operatorname{id}_{\mathcal{D}}$	$\mathrm{id}_{\mathcal{D}}$	similarly if we post-compose F with $\mathrm{id}_{\mathcal{D}}$. This gives unitality, and associativity is just as easy,
fong-spivak-seven-sketches_03256	292	■ 1.000	$H: \mathcal{E} \rightarrow \mathcal{F}$	$H: \mathcal{E} \rightarrow \mathcal{F}$	though more wordy. Given F as above and $G: \mathcal{D} \rightarrow \mathcal{E}$ and $H: \mathcal{E} \rightarrow \mathcal{F}$, we need to show that
fong-spivak-seven-sketches_03257	292	■ 0.991	$(F;G);H=F;(G;H)$	$(F;G);H = F;(G;H)$	$(F \circ G) \circ H = F \circ (G \circ H)$. It's a simple application of the definition: for any $x \in \mathcal{C}$, be it an object
fong-spivak-seven-sketches_03258	292	■ 0.991	$x \in \mathcal{C}$	$x \in \mathcal{C}$	$(F \circ G) \circ H = F \circ (G \circ H)$. It's a simple application of the definition: for any $x \in \mathcal{C}$, be it an object
fong-spivak-seven-sketches_03259	292	■ 0.996	$F_S(\operatorname{id}_1)=\operatorname{id}_S$	$F_S(\mathrm{id}_1) = \mathrm{id}_S$	Let $S \in \mathbf{Set}$ be a set. Define $F_S: \mathbf{1} \rightarrow \mathbf{Set}$ by $F_S(1) = S$ and $F_S(\mathrm{id}_1) = \mathrm{id}_S$. With this definition, F_S
fong-spivak-seven-sketches_03260	292	■ 0.996	F_S	F_S	Let $S \in \mathbf{Set}$ be a set. Define $F_S: \mathbf{1} \rightarrow \mathbf{Set}$ by $F_S(1) = S$ and $F_S(\mathrm{id}_1) = \mathrm{id}_S$. With this definition, F_S
fong-spivak-seven-sketches_03261	292	■ 1.000	$D(z)$	$D(z)$	1. Data on this schema, i.e. a set-valued functor, assigns a set $D(z)$ and a function $D(s): D(z) \rightarrow D(z)$,
fong-spivak-seven-sketches_03262	292	■ 1.000	$D(s): D(z) \rightarrow D(z)$	$D(s): D(z) \rightarrow D(z)$	1. Data on this schema, i.e. a set-valued functor, assigns a set $D(z)$ and a function $D(s): D(z) \rightarrow D(z)$,
fong-spivak-seven-sketches_03263	292	■ 0.999	D_z	D_z	of the set D_z :
fong-spivak-seven-sketches_03264	293	■ 1.000	$D(c)$	$D(c)$	2. We could make $D(c)$ the set of people at a “secret Santa” Christmas party, where everyone gives
fong-spivak-seven-sketches_03265	293	■ 0.995	$D(b)$	$D(b)$	a gift to someone, possibly themselves. Take $D(b)$ to be the set of gifts, g the giver function (each
fong-spivak-seven-sketches_03266	293	■ 0.997	$D(a)$	$D(a)$	gift is given by a person), and h the receiver function (each gift is received by a person), $D(a)$ is
fong-spivak-seven-sketches_03267	293	■ 0.999	$d(f): D(a) \rightarrow D(b)$	$d(f): D(a) \rightarrow D(b)$	the set of people who give a gift to themselves, and $d(f): D(a) \rightarrow D(b)$ is the inclusion.
fong-spivak-seven-sketches_03268	293	■ 0.568	$F, G, H \in \mathcal{D}^{\mathcal{C}}$	$F, G, H \in \mathcal{D}^{\mathcal{C}}$	1. The expert packs so much information in so little space! Suppose given three objects $F, G, H \in \mathcal{D}^{\mathcal{C}}$;
fong-spivak-seven-sketches_03269	293	■ 1.000	$F, G, H: \mathcal{C} \rightarrow \mathcal{D}$	$F, G, H: \mathcal{C} \rightarrow \mathcal{D}$	these are functors $F, G, H: \mathcal{C} \rightarrow \mathcal{D}$. Morphisms $\alpha: F \rightarrow G$ and $\beta: G \rightarrow H$ are natural transfor-
fong-spivak-seven-sketches_03270	293	■ 1.000	$\beta: G \rightarrow H$	$\beta: G \rightarrow H$	these are functors $F, G, H: \mathcal{C} \rightarrow \mathcal{D}$. Morphisms $\alpha: F \rightarrow G$ and $\beta: G \rightarrow H$ are natural transfor-
fong-spivak-seven-sketches_03271	293	■ 0.877	$c \in \mathcal{C}, \alpha$	$c \in \mathcal{C}, \alpha$	So for each $c \in \mathcal{C}$, α has a component $\alpha_c: F(c) \rightarrow G(c)$ and β has a component $\beta_c: G(c) \rightarrow H(c)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches- F03272	293	0.877	$\backslash\mathrm{beta}_{\{c\}}: G(c) \rightarrowtail H(c)$	$\beta_c : G(c) \rightarrow H(c)$	So for each $c \in \mathcal{C}$, α has a component $\alpha_c : F(c) \rightarrow G(c)$ and β has a component $\beta_c : G(c) \rightarrowtail H(c)$
fong-spivak-seven-sketches- F03273	293	0.872	$(\alpha ; \backslash\mathrm{beta})_{\{\mathrm{mathcal{C}}\}} := \left(\alpha_{\{\mathrm{mathcal{C}}\}} \backslash\mathrm{beta}_{\{\mathrm{mathcal{C}}\}} \right)$	$(\alpha ; \beta)_c := (\alpha_c ; \beta_c)$	in $\mathcal{D}$. The expert has told us to define $(\alpha \circ \beta)_c := (\alpha_c \circ \beta_c)$, and indeed that is a morphism
fong-spivak-seven-sketches- F03274	293	1.000	$F(c) \rightarrowtail H(c)$	$F(c) \rightarrow H(c)$	$F(c) \rightarrowtail H(c)$.
fong-spivak-seven-sketches- F03275	293	0.995	$\backslash\mathrm{operatorname{id}}_{\{F\}}$	id_F	2. We propose that the identity natural transformation id_F on a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ has as its
fong-spivak-seven-sketches- F03276	293	0.660	$\left(\backslash\mathrm{operatorname{id}}_{\{F\}} \right)_{\{c\}} := \mathrm{operatorname{id}}_{\{F(c)\}}$	$(\mathrm{id}_F)_c := \mathrm{id}_{F(c)}$	c -component the morphism $(\mathrm{id}_F)_c := \mathrm{id}_{F(c)}$ in $\mathcal{D}$, for any c . The naturality square
fong-spivak-seven-sketches- F03277	293	1.000	$\backslash\mathrm{beta} : F \rightarrowtail G$	$\beta : F \rightarrow G$	obviously commutes for any $f : c \rightarrow c'$. And it is unital: post-composing id_F with any $\beta : F \rightarrow G$
fong-spivak-seven-sketches- F03278	293	0.992	$\backslash\mathrm{alpha} : E \rightarrowtail F$	$\alpha : E \rightarrow F$	(and similarly for precomposing with any $\alpha : E \rightarrow F$) results in a natural transformation $\mathrm{id}_F \circ \beta$
fong-spivak-seven-sketches- F03279	293	0.992	$\backslash\mathrm{operatorname{id}}_{\{F\}} ; \backslash\mathrm{beta}$	$\mathrm{id}_F ; \beta$	(and similarly for precomposing with any $\alpha : E \rightarrow F$) results in a natural transformation $\mathrm{id}_F \circ \beta$
fong-spivak-seven-sketches- F03280	293	0.896	$\left(\backslash\mathrm{operatorname{id}}_{\{F\}} \right)_{\{c\}} ; \backslash\mathrm{beta}_{\{c\}} = \left(\backslash\mathrm{operatorname{id}}_{\{F(c)\}} ; \backslash\mathrm{beta}_{\{c\}} \right) = \backslash\mathrm{beta}_{\{c\}}$	$(\mathrm{id}_F)_c ; \beta_c = (\mathrm{id}_{F(c)} ; \beta_c) = \beta_c$	with components $(\mathrm{id}_F)_c \circ \beta_c = (\mathrm{id}_{F(c)} \circ \beta_c) = \beta_c$, and this is just β as desired.
fong-spivak-seven-sketches- F03281	293	1.000	$\backslash\mathrm{alpha}, \backslash\mathrm{beta} : F \rightarrowtail G$	$\alpha, \beta : F \rightarrow G$	1. Suppose that $F, G : \mathcal{C} \rightarrow \mathcal{P}$ are functors and $\alpha, \beta : F \rightarrow G$ are natural transformations; we need to
fong-spivak-seven-sketches- F03282	293	1.000	$\backslash\mathrm{alpha} = \backslash\mathrm{beta}$	$\alpha = \beta$	show that $\alpha = \beta$. It suffices to check that $\alpha_c = \beta_c$ for each object $c \in \mathrm{Ob}(\mathcal{C})$. But α_c and β_c are
fong-spivak-seven-sketches- F03283	293	1.000	$\backslash\mathrm{alpha}_{\{c\}} = \backslash\mathrm{beta}_{\{c\}}$	$\alpha_c = \beta_c$	show that $\alpha = \beta$. It suffices to check that $\alpha_c = \beta_c$ for each object $c \in \mathrm{Ob}(\mathcal{C})$. But α_c and β_c are
fong-spivak-seven-sketches- F03284	293	1.000	$\backslash\mathrm{beta}_{\{c\}}$	β_c	show that $\alpha = \beta$. It suffices to check that $\alpha_c = \beta_c$ for each object $c \in \mathrm{Ob}(\mathcal{C})$. But α_c and β_c are
fong-spivak-seven-sketches- F03285	293	0.998	$F(c) \rightarrowtail G(c)$	$F(c) \rightarrow G(c)$	morphisms $F(c) \rightarrowtail G(c)$ in $\mathcal{P}$, which is a preorder, and the definition of a preorder—considered
fong-spivak-seven-sketches- F03286	293	0.421	$\backslash\mathrm{mathcal{P}} := \backslash\mathrm{mathbf{1}}$	$\mathcal{P} := \mathbf{1}$	2. This is false. Let $\mathcal{P} := \mathbf{1}$, let $\mathcal{C} := \left \begin{array}{c} a \\ \xrightarrow{f_1} b \\ \xrightarrow{f_2} \end{array} \right $, let $F(1) := a$, let $G(1) := b$, let $\alpha_1 := f_1$, and let $\beta_1 := g_2$.
fong-spivak-seven-sketches- F03287	293	0.421	$\backslash\mathrm{mathcal{C}} := \backslash\mathrm{vec{\bullet}} \rightarrowtail [f_2][f_1] \backslash\mathrm{stackrel{b}{\bullet}}$	$\mathcal{C} := \begin{array}{c} \bullet \\ \xrightarrow{f_1} \bullet \\ \xrightarrow{f_2} \end{array}$	2. This is false. Let $\mathcal{P} := \mathbf{1}$, let $\mathcal{C} := \left \begin{array}{c} a \\ \xrightarrow{f_1} b \\ \xrightarrow{f_2} \end{array} \right $, let $F(1) := a$, let $G(1) := b$, let $\alpha_1 := f_1$, and let $\beta_1 := g_2$.
fong-spivak-seven-sketches- F03288	293	0.421	$F(1) := a$	$F(1) := a$	2. This is false. Let $\mathcal{P} := \mathbf{1}$, let $\mathcal{C} := \left \begin{array}{c} a \\ \xrightarrow{f_1} b \\ \xrightarrow{f_2} \end{array} \right $, let $F(1) := a$, let $G(1) := b$, let $\alpha_1 := f_1$, and let $\beta_1 := g_2$.
fong-spivak-seven-sketches- F03289	293	0.421	$G(1) := b$	$G(1) := b$	2. This is false. Let $\mathcal{P} := \mathbf{1}$, let $\mathcal{C} := \left \begin{array}{c} a \\ \xrightarrow{f_1} b \\ \xrightarrow{f_2} \end{array} \right $, let $F(1) := a$, let $G(1) := b$, let $\alpha_1 := f_1$, and let $\beta_1 := g_2$.
fong-spivak-seven-sketches- F03290	293	0.421	$\backslash\mathrm{alpha}_{\{1\}} := f_1$	$\alpha_1 := f_1$	2. This is false. Let $\mathcal{P} := \mathbf{1}$, let $\mathcal{C} := \left \begin{array}{c} a \\ \xrightarrow{f_1} b \\ \xrightarrow{f_2} \end{array} \right $, let $F(1) := a$, let $G(1) := b$, let $\alpha_1 := f_1$, and let $\beta_1 := g_2$.
fong-spivak-seven-sketches- F03291	293	0.421	$\backslash\mathrm{beta}_{\{1\}} := g_2$	$\beta_1 := g_2$	2. This is false. Let $\mathcal{P} := \mathbf{1}$, let $\mathcal{C} := \left \begin{array}{c} a \\ \xrightarrow{f_1} b \\ \xrightarrow{f_2} \end{array} \right $, let $F(1) := a$, let $G(1) := b$, let $\alpha_1 := f_1$, and let $\beta_1 := g_2$.
fong-spivak-seven-sketches- F03292	294	1.000	$\backslash\mathrm{alpha}_{\{\text{Arrow}\}}(b) = e$	$\alpha_{\mathrm{Arrow}}(b) = e$	1. We have $\alpha_{\mathrm{Arrow}}(b) = e$ and $\alpha_{\mathrm{Vertex}}(1) = 4$, $\alpha_{\mathrm{Vertex}}(2) = 5$, and $\alpha_{\mathrm{Vertex}}(3) = 5$.
fong-spivak-seven-sketches- F03293	294	1.000	$\backslash\mathrm{alpha}_{\{\text{Vertex}\}}(1) = 4, \backslash\mathrm{alpha}_{\{\text{Vertex}\}}(2) = 5$	$\alpha_{\mathrm{Vertex}}(1) = 4, \alpha_{\mathrm{Vertex}}(2) = 5$	1. We have $\alpha_{\mathrm{Arrow}}(b) = e$ and $\alpha_{\mathrm{Vertex}}(1) = 4, \alpha_{\mathrm{Vertex}}(2) = 5$, and $\alpha_{\mathrm{Vertex}}(3) = 5$.
fong-spivak-seven-sketches- F03294	294	1.000	$\backslash\mathrm{alpha}_{\{\text{Vertex}\}}(3) = 5$	$\alpha_{\mathrm{Vertex}}(3) = 5$	1. We have $\alpha_{\mathrm{Arrow}}(b) = e$ and $\alpha_{\mathrm{Vertex}}(1) = 4$, $\alpha_{\mathrm{Vertex}}(2) = 5$, and $\alpha_{\mathrm{Vertex}}(3) = 5$.
fong-spivak-seven-sketches- F03295	295	1.000	$X \times B \rightarrowtail Y \times B$	$X \times B \rightarrow Y \times B$	1. Given an arbitrary function $f : X \rightarrow Y$, we need a function $X \times B \rightarrow Y \times B$. We suggest the
fong-spivak-seven-sketches- F03296	295	1.000	$f \times B$	$f \times B$	function which might be denoted $f \times B$; it sends (x, b) to $(f(x), b)$. This assignment is functorial:

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches- F03297	295	■ 1.000	(x, b)	(x, b)	function which might be denoted $f \times B$; it sends (x, b) to $(f(x), b)$. This assignment is functorial:
fong-spivak-seven-sketches- F03298	295	■ 1.000	$(f(x), b)$	$(f(x), b)$	function which might be denoted $f \times B$; it sends (x, b) to $(f(x), b)$. This assignment is functorial:
fong-spivak-seven-sketches- F03299	295	■ 0.988	$\mathrm{id}_{X \times B}$	$\mathrm{id}_{X \times B}$	applied to id_X it returns $\mathrm{id}_{X \times B}$ and it preserves composition.
fong-spivak-seven-sketches- F03300	295	■ 1.000	$X^B \rightarrow Y^B$	$X^B \rightarrow Y^B$	2. Given a function $f: X \rightarrow Y$, we need a function $X^B \rightarrow Y^B$. The canonical function would
fong-spivak-seven-sketches- F03301	295	■ 0.988	f^B	f^B	be denoted f^B ; it sends a function $g: B \rightarrow X$ to the composite $(g \circ f): B \rightarrow X \rightarrow Y$. This is
fong-spivak-seven-sketches- F03302	295	■ 0.988	$g: B \rightarrow X$	$g: B \rightarrow X$	be denoted f^B ; it sends a function $g: B \rightarrow X$ to the composite $(g \circ f): B \rightarrow X \rightarrow Y$. This is
fong-spivak-seven-sketches- F03303	295	■ 0.988	$(g \circ f): B \rightarrow X \rightarrow Y$	$(g \circ f): B \rightarrow X \rightarrow Y$	be denoted f^B ; it sends a function $g: B \rightarrow X$ to the composite $(g \circ f): B \rightarrow X \rightarrow Y$. This is
fong-spivak-seven-sketches- F03304	295	■ 0.986	$f^B(\mathrm{id}_X) = \mathrm{id}_{X^B}$	$f^B(\mathrm{id}_X) = \mathrm{id}_{X^B}$	functorial: applied to id_X it sends g to g , i.e. $f^B(\mathrm{id}_X) = \mathrm{id}_{X^B}$, and applied to the composite
fong-spivak-seven-sketches- F03305	295	■ 1.000	$\left(f_1; f_2\right): X \rightarrow Y \rightarrow Z$	$(f_1; f_2): X \rightarrow Y \rightarrow Z$	$(f_1; f_2): X \rightarrow Y \rightarrow Z$, we have
fong-spivak-seven-sketches- F03306	295	■ 1.000	$g \in X^B$	$g \in X^B$	for any $g \in X^B$.
fong-spivak-seven-sketches- F03307	295	■ 0.999	$\mathbb{N}^{\mathbb{N}}$	$\mathbb{N}^{\mathbb{N}}$	3. If $p: \mathbb{N} \rightarrow \mathbb{N}^{\mathbb{N}}$ is the result of currying $+: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$, then $p(3)$ is an element of $\mathbb{N}^{\mathbb{N}}$, i.e. we have
fong-spivak-seven-sketches- F03308	295	■ 1.000	$p(3): \mathbb{N} \rightarrow \mathbb{N}$	$p(3): \mathbb{N} \rightarrow \mathbb{N}$	$p(3): \mathbb{N} \rightarrow \mathbb{N}$; what function is it? It is the function that adds three. That is $p(3)(n) := n + 3$.
fong-spivak-seven-sketches- F03309	295	■ 1.000	$p(3)(n) := n + 3$	$p(3)(n) := n + 3$	$p(3): \mathbb{N} \rightarrow \mathbb{N}$; what function is it? It is the function that adds three. That is $p(3)(n) := n + 3$.
fong-spivak-seven-sketches- F03310	295	■ 0.533	$!: \mathcal{C} \rightarrow \mathbf{1}$	$!: \mathcal{C} \rightarrow \mathbf{1}$	The functor $!: \mathcal{C} \rightarrow \mathbf{1}$ from Eq. (3.75) sends each object $c \in \mathcal{C}$ to the unique object $1 \in \mathbf{1}$ and sends each
fong-spivak-seven-sketches- F03311	295	■ 0.533	$1 \in \mathbf{1}$	$1 \in \mathbf{1}$	The functor $!: \mathcal{C} \rightarrow \mathbf{1}$ from Eq. (3.75) sends each object $c \in \mathcal{C}$ to the unique object $1 \in \mathbf{1}$ and sends each
fong-spivak-seven-sketches- F03312	295	■ 0.590	$\mathrm{B} = \mathrm{Bob}, 3 = \mathrm{Em}_3$	$B = \mathrm{Bob}, 3 = \mathrm{Em}_3$	Here it is, with names and emails shortened (e.g. $B = \mathrm{Bob}, 3 = \mathrm{Em}_3$):
fong-spivak-seven-sketches- F03313	295	■ 0.999	$c \rightarrow z$	$c \rightarrow z$	An object z is terminal in some category $\mathcal{C}$ if, for every $c \in \mathcal{C}$ there exists a unique morphism $c \rightarrow z$.
fong-spivak-seven-sketches- F03314	296	■ 0.940	$\mathcal{C} \in$	$\mathcal{C} \in$	any object (category) $\mathcal{C} \in \mathbf{Cat}$.
fong-spivak-seven-sketches- F03315	296	■ 0.877	$2V := \bullet \bullet$	$2V := \bullet \bullet$	Consider the graph $2V := \boxed{\bullet \bullet}$ with two vertices and no arrows, and let $\mathcal{C} = \mathbf{Free}(2V)$; it has two
fong-spivak-seven-sketches- F03316	296	■ 0.877	$\mathcal{C} = \mathbf{Free}(2V)$	$\mathcal{C} = \mathbf{Free}(2V)$	Consider the graph $2V := \boxed{\bullet \bullet}$ with two vertices and no arrows, and let $\mathcal{C} = \mathbf{Free}(2V)$; it has two
fong-spivak-seven-sketches- F03317	296	■ 1.000	$z \in \mathcal{P}$	$z \in \mathcal{P}$	A product of x and y in $\mathcal{P}$ is an object $z \in \mathcal{P}$ equipped with maps $z \rightarrow x$ and $z \rightarrow y$ such that for any
fong-spivak-seven-sketches- F03318	296	■ 1.000	$z \rightarrow x$	$z \rightarrow x$	A product of x and y in $\mathcal{P}$ is an object $z \in \mathcal{P}$ equipped with maps $z \rightarrow x$ and $z \rightarrow y$ such that for any
fong-spivak-seven-sketches- F03319	296	■ 1.000	$z \rightarrow y$	$z \rightarrow y$	A product of x and y in $\mathcal{P}$ is an object $z \in \mathcal{P}$ equipped with maps $z \rightarrow x$ and $z \rightarrow y$ such that for any
fong-spivak-seven-sketches- F03320	296	■ 1.000	z'	z'	other object z' and maps $z' \rightarrow x$ and $z' \rightarrow y$, there is a unique morphism $z' \rightarrow z$ making the evident
fong-spivak-seven-sketches- F03321	296	■ 1.000	$z' \rightarrow x$	$z' \rightarrow x$	other object z' and maps $z' \rightarrow x$ and $z' \rightarrow y$, there is a unique morphism $z' \rightarrow z$ making the evident
fong-spivak-seven-sketches- F03322	296	■ 1.000	$z' \rightarrow y$	$z' \rightarrow y$	other object z' and maps $z' \rightarrow x$ and $z' \rightarrow y$, there is a unique morphism $z' \rightarrow z$ making the evident
fong-spivak-seven-sketches- F03323	296	■ 1.000	$z' \rightarrow z$	$z' \rightarrow z$	other object z' and maps $z' \rightarrow x$ and $z' \rightarrow y$, there is a unique morphism $z' \rightarrow z$ making the evident
fong-spivak-seven-sketches- F03324	296	■ 1.000	$z \leq x$	$z \leq x$	diagrams commute. Thus the above becomes: a product of x and y in $\mathcal{P}$ is an object z with $z \leq x$ and
fong-spivak-seven-sketches- F03325	296	■ 1.000	$z \leq y$	$z \leq y$	$z \leq y$ such that for any other z' , if $z' \leq x$ and $z' \leq y$ then $z' \leq z$. This is exactly the definition of meet,

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches- F03352	296	0.521	$\mathcal{D}(\mathrm{v}_1)=A$	$D(v_1) = A$	Suppose $\mathcal{J}$ is the graph  and $D: \mathcal{J} \rightarrow \mathbf{Set}$ is given by two sets, $D(v_1) = A$ and $D(v_2) = B$ for sets
fong-spivak-seven-sketches- F03353	296	0.521	$\mathcal{D}(\mathrm{v}_2)=B$	$D(v_2) = B$	Suppose $\mathcal{J}$ is the graph  and $D: \mathcal{J} \rightarrow \mathbf{Set}$ is given by two sets, $D(v_1) = A$ and $D(v_2) = B$ for sets
fong-spivak-seven-sketches- F03354	297	0.274	$F^{\mathrm{op}}: \mathcal{C}^{\mathrm{op}} \rightarrow \mathcal{D}^{\mathrm{op}}$	$F^{\mathrm{op}}: \mathcal{C}^{\mathrm{op}} \rightarrow \mathcal{D}^{\mathrm{op}}$	Given a functor $F: \mathcal{C} \rightarrow \mathcal{D}$, we define its opposite $F^{\mathrm{op}}: \mathcal{C}^{\mathrm{op}} \rightarrow \mathcal{D}^{\mathrm{op}}$ as follows. For each object $c \in$
fong-spivak-seven-sketches- F03355	297	0.274	$c \in$	$c \in$	Given a functor $F: \mathcal{C} \rightarrow \mathcal{D}$, we define its opposite $F^{\mathrm{op}}: \mathcal{C}^{\mathrm{op}} \rightarrow \mathcal{D}^{\mathrm{op}}$ as follows. For each object $c \in$
fong-spivak-seven-sketches- F03356	297	0.904	$\mathrm{Ob}(\mathcal{C}^{\mathrm{op}})=\mathrm{Ob}(\mathcal{C})$	$\mathrm{Ob}(\mathcal{C}^{\mathrm{op}}) = \mathrm{Ob}(\mathcal{C})$	$\mathrm{Ob}(\mathcal{C}^{\mathrm{op}}) = \mathrm{Ob}(\mathcal{C})$, put $F^{\mathrm{op}}(c) := F(c)$. For each morphism $f: c_1 \rightarrow c_2$ in $\mathcal{C}^{\mathrm{op}}$, we have a corresponding
fong-spivak-seven-sketches- F03357	297	0.904	$F^{\mathrm{op}}(c):=F(c)$	$F^{\mathrm{op}}(c) := F(c)$	$\mathrm{Ob}(\mathcal{C}^{\mathrm{op}}) = \mathrm{Ob}(\mathcal{C})$, put $F^{\mathrm{op}}(c) := F(c)$. For each morphism $f: c_1 \rightarrow c_2$ in $\mathcal{C}^{\mathrm{op}}$, we have a corresponding
fong-spivak-seven-sketches- F03358	297	1.000	$f^{\mathrm{prime}}: c_2 \rightarrow c_1$	$f': c_2 \rightarrow c_1$	morphism $f': c_2 \rightarrow c_1$ in $\mathcal{C}$ and thus a morphism $F(f'): F(c_2) \rightarrow F(c_1)$ in $\mathcal{D}$, and thus a morphism
fong-spivak-seven-sketches- F03359	297	1.000	$F(f^{\mathrm{prime}}): F(c_2) \rightarrow F(c_1)$	$F(f'): F(c_2) \rightarrow F(c_1)$	morphism $f': c_2 \rightarrow c_1$ in $\mathcal{C}$ and thus a morphism $F(f'): F(c_2) \rightarrow F(c_1)$ in $\mathcal{D}$, and thus a morphism
fong-spivak-seven-sketches- F03360	297	0.951	$F(f^{\mathrm{prime}})^{\mathrm{prime}}: F^{\mathrm{op}}(c_1) \rightarrow F^{\mathrm{op}}(c_2)$	$F(f')': F^{\mathrm{op}}(c_1) \rightarrow F^{\mathrm{op}}(c_2)$	$F(f')': F^{\mathrm{op}}(c_1) \rightarrow F^{\mathrm{op}}(c_2)$. Hence we can define $F^{\mathrm{op}}(f) := F(f')'$. Note that the primes ($-'$) are pretty
fong-spivak-seven-sketches- F03361	297	0.951	$F^{\mathrm{op}}(f):=F(f^{\mathrm{prime}})^{\mathrm{prime}}$	$F^{\mathrm{op}}(f) := F(f')'$	$F(f')': F^{\mathrm{op}}(c_1) \rightarrow F^{\mathrm{op}}(c_2)$. Hence we can define $F^{\mathrm{op}}(f) := F(f')'$. Note that the primes ($-'$) are pretty
fong-spivak-seven-sketches- F03362	297	0.951	$(-)^{\mathrm{prime}}$	$(-)^{\mathrm{'}}$	$F(f')': F^{\mathrm{op}}(c_1) \rightarrow F^{\mathrm{op}}(c_2)$. Hence we can define $F^{\mathrm{op}}(f) := F(f')'$. Note that the primes ($-'$) are pretty
fong-spivak-seven-sketches- F03363	297	0.815	F^{op}	F^{op}	It is easy to check that our definition of F^{op} is functorial: it sends identities to identities and composites
fong-spivak-seven-sketches- F03364	298	0.666	$\mathcal{X}^{\mathrm{op}} \times \dagger$	$\mathcal{X}^{\mathrm{op}} \times \dagger$	1. The Hasse diagram for $\mathcal{X}^{\mathrm{op}} \times \mathcal{Y}$ is shown here (ignore the colors):
fong-spivak-seven-sketches- F03365	298	0.299	$\Lambda: \mathcal{X} \mapsto \mathcal{Y}$	$\Lambda: \mathcal{X} \mapsto \mathcal{Y}$	2. There is a profunctor $\Lambda: \mathcal{X} \nrightarrow \mathcal{Y}$, i.e. a functor $\mathcal{X}^{\mathrm{op}} \times \mathcal{Y} \rightarrow \mathbb{B}$, such that, in the picture above, blue
fong-spivak-seven-sketches- F03366	298	0.299	$\mathcal{X}^{\mathrm{op}} \times \mathcal{Y} \rightarrow \mathbb{B}$	$\mathcal{X}^{\mathrm{op}} \times \mathcal{Y} \rightarrow \mathbb{B}$	2. There is a profunctor $\Lambda: \mathcal{X} \nrightarrow \mathcal{Y}$, i.e. a functor $\mathcal{X}^{\mathrm{op}} \times \mathcal{Y} \rightarrow \mathbb{B}$, such that, in the picture above, blue
fong-spivak-seven-sketches- F03367	298	0.301	$x^{\mathrm{op}} \times y$	$x^{\mathrm{op}} \times y$	The preorder $\mathcal{X}^{\mathrm{op}} \times \mathcal{Y}$ describes tasks in decreasing difficulty. For example, (we hope) it is easier for
fong-spivak-seven-sketches- F03368	298	1.000	Λ	Λ	profunctor Λ describes possible states of knowledge for my aunt: she can describe monoids without
fong-spivak-seven-sketches- F03369	298	0.348	$\Phi: \mathcal{X}^{\mathrm{op}} \times \mathcal{Y} \rightarrow \mathcal{V}$	$\Phi: \mathcal{X}^{\mathrm{op}} \times \mathcal{Y} \rightarrow \mathcal{V}$	Recall from Definition 2.41 that a $\mathcal{V}$ -functor $\Phi: \mathcal{X}^{\mathrm{op}} \times \mathcal{Y} \rightarrow \mathcal{V}$ is a function $\Phi: \mathrm{Ob}(\mathcal{X}^{\mathrm{op}} \times \mathcal{Y}) \rightarrow \mathrm{Ob}(\mathcal{V})$
fong-spivak-seven-sketches- F03370	298	0.348	$\Phi: \mathrm{Ob}(\mathcal{X}^{\mathrm{op}} \times \mathcal{Y}) \rightarrow \mathrm{Ob}(\mathcal{V})$	$\Phi: \mathrm{Ob}(\mathcal{X}^{\mathrm{op}} \times \mathcal{Y}) \rightarrow \mathrm{Ob}(\mathcal{V})$	Recall from Definition 2.41 that a $\mathcal{V}$ -functor $\Phi: \mathcal{X}^{\mathrm{op}} \times \mathcal{Y} \rightarrow \mathcal{V}$ is a function $\Phi: \mathrm{Ob}(\mathcal{X}^{\mathrm{op}} \times \mathcal{Y}) \rightarrow \mathrm{Ob}(\mathcal{V})$
fong-spivak-seven-sketches- F03371	298	0.932	$x^{\mathrm{op}} \times y$	$x^{\mathrm{op}} \times y$	such that for all (x, y) and (x', y') in $\mathcal{X}^{\mathrm{op}} \times \mathcal{Y}$ we have
fong-spivak-seven-sketches- F03372	299	0.997	$\Phi: (T \times E) \rightarrow \$$	$\Phi: (T \times E) \rightarrow \$$	Yes, this is valid: it just means that the profunctor $\Phi: (T \times E) \nrightarrow \$$ does not relate (good-natured,
fong-spivak-seven-sketches- F03373	299	0.895	$\Phi(($	$\Phi(($	funny) to any element of $\$$. More formally, it means that $\Phi(\text{good-natured, funny}, p) = \text{false}$ for
fong-spivak-seven-sketches- F03374	299	0.895	$), p) =$	$), p) =$	funny) to any element of $\$$. More formally, it means that $\Phi(\text{good-natured, funny}, p) = \text{false}$ for
fong-spivak-seven-sketches- F03375	299	1.000	$p \in \$ = \{\$100 \text{ K}, \$500 \text{ K}, \$1 \text{ M}\}$	$p \in \$ = \{\$100 \text{ K}, \$500 \text{ K}, \$1 \text{ M}\}$	all $p \in \$ = \{\$100\text{K}, \$500\text{K}, \$1\text{M}\}$. We can interpret this to mean that it is not feasible to produce a
fong-spivak-seven-sketches- F03376	299	0.712	$(\Phi; \Psi)(-, -) = \mathcal{X}(-, D) + 9 + z(r, -)$	$(\Phi; \Psi)(-, -) = \mathcal{X}(-, D) + 9 + z(r, -)$	go through the bridges from D to y and y to r , so in this case $(\Phi \circ \Psi)(-, -) = \mathcal{X}(-, D) + 9 + \mathcal{Z}(r, -)$. This
fong-spivak-seven-sketches- F03377	300	0.432	U_X	U_X	We choose the Cost -category $\mathcal{X}$ from Eq. (2.56). The unit profunctor U_X on $\mathcal{X}$ is described by the bridge
fong-spivak-seven-sketches- F03378	300	0.705	$I \leq \mathcal{P}(p, p)$	$I \leq \mathcal{P}(p, p)$	of $\otimes$ (Definition 2.2(a)) applied to the inequalities $I \leq \mathcal{P}(p, p)$ (the identity law for $\mathcal{P}$ at p ,
fong-spivak-seven-sketches- F03379	300	0.993	$\Phi(p, q) \leq \Phi(p, q)$	$\Phi(p, q) \leq \Phi(p, q)$	Definition 2.46(a)) and $\Phi(p, q) \leq \Phi(p, q)$ (reflexivity of preorder $\mathcal{V}$, Definition 1.30(a)). The third

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03380	300	■ 1.000	$x \leq x \vee y$	$x \leq x \vee y$	step uses the definition of join: for all x and y , since any $x \leq x$, we have $x \leq x \vee y$. The final
fong-spivak-seven-sketches F03381	300	■ 0.998	$I =$	$I =$	2. Note that in Bool , $I = \text{true}$. Since the identity law at p says $\text{true} \leq \mathcal{P}(p, p)$, and true is the
fong-spivak-seven-sketches F03382	300	■ 0.998	$\leq \mathcal{P}(p, p)$	$\leq \mathcal{P}(p, p)$	2. Note that in Bool , $I = \text{true}$. Since the identity law at p says $\text{true} \leq \mathcal{P}(p, p)$, and true is the
fong-spivak-seven-sketches F03383	300	■ 0.972	$\mathcal{P}(p, p) =$	$\mathcal{P}(p, p) =$	largest element of the preorder Bool , we thus have $\mathcal{P}(p, p) = \text{true}$ for all p . This shows that the
fong-spivak-seven-sketches F03384	300	■ 1.000	$\Phi(p, q)$	$\Phi(p, q)$	side of the inequality is equal to $\Phi(p, q)$. We split into two cases. Suppose $\Phi(p, q) = \text{true}$. Then,
fong-spivak-seven-sketches F03385	300	■ 1.000	$\Phi(p, q) =$	$\Phi(p, q) =$	side of the inequality is equal to $\Phi(p, q)$. We split into two cases. Suppose $\Phi(p, q) = \text{true}$. Then,
fong-spivak-seven-sketches F03386	300	■ 0.825	$\mathcal{P}^{\text{op}} \times \mathcal{Q} \rightarrow$	$\mathcal{P}^{\text{op}} \times \mathcal{Q} \rightarrow$	Next, suppose $\Phi(p, q) = \text{false}$. Note that since Φ is a monotone map $\mathcal{P}^{\text{op}} \times \mathcal{Q} \rightarrow \mathbf{Bool}$, if $p \leq p_1$
fong-spivak-seven-sketches F03387	300	■ 0.825	$p \leq p_1$	$p \leq p_1$	Next, suppose $\Phi(p, q) = \text{false}$. Note that since Φ is a monotone map $\mathcal{P}^{\text{op}} \times \mathcal{Q} \rightarrow \mathbf{Bool}$, if $p \leq p_1$
fong-spivak-seven-sketches F03388	300	■ 1.000	$\Phi(p_1, q) \leq \Phi(p, q)$	$\Phi(p_1, q) \leq \Phi(p, q)$	in $\mathcal{P}$, then $\Phi(p_1, q) \leq \Phi(p, q)$ in Bool . Thus if $\mathcal{P}(p, p_1) = \text{true}$ then $\Phi(p_1, q) = \Phi(p, q) = \text{false}$.
fong-spivak-seven-sketches F03389	300	■ 1.000	$\mathcal{P}(p, p_1) =$	$\mathcal{P}(p, p_1) =$	in $\mathcal{P}$, then $\Phi(p_1, q) \leq \Phi(p, q)$ in Bool . Thus if $\mathcal{P}(p, p_1) = \text{true}$ then $\Phi(p_1, q) = \Phi(p, q) = \text{false}$.
fong-spivak-seven-sketches F03390	300	■ 1.000	$\Phi(p_1, q) = \Phi(p, q) =$	$\Phi(p_1, q) = \Phi(p, q) =$	in $\mathcal{P}$, then $\Phi(p_1, q) \leq \Phi(p, q)$ in Bool . Thus if $\mathcal{P}(p, p_1) = \text{true}$ then $\Phi(p_1, q) = \Phi(p, q) = \text{false}$.
fong-spivak-seven-sketches F03391	300	■ 1.000	$\Phi(p_1, q) =$	$\Phi(p_1, q) =$	This implies that for all $p_1 \in \mathcal{P}$, we have either $\mathcal{P}(p, p_1) = \text{false}$ or $\Phi(p_1, q) = \text{false}$, and hence
fong-spivak-seven-sketches F03392	300	■ 0.772	$\bigvee_{p_1 \in \mathcal{P}} \Phi(p_1, q) = \bigvee_{p_1 \in \mathcal{P}} \mathcal{P}(p, p_1) \wedge \Phi(p_1, q)$	$\bigvee_{p_1 \in \mathcal{P}} \mathcal{P}(p, p_1) \wedge \Phi(p_1, q) = \bigvee_{p_1 \in \mathcal{P}} \text{false} = \text{false}$	$\bigvee_{p_1 \in \mathcal{P}} \mathcal{P}(p, p_1) \wedge \Phi(p_1, q) = \bigvee_{p_1 \in \mathcal{P}} \text{false} = \text{false}$.
fong-spivak-seven-sketches F03393	300	■ 0.644	$\Phi(p, q) = \bigvee_{p_1 \in \mathcal{P}} \mathcal{P}(p, p_1) \wedge \Phi(p_1, q)$	$\Phi(p, q) = \bigvee_{p_1 \in \mathcal{P}} \mathcal{P}(p, p_1) \wedge \Phi(p_1, q)$	Thus, in either case, we see that $\Phi(p, q) = \bigvee_{p_1 \in \mathcal{P}} \mathcal{P}(p, p_1) \wedge \Phi(p_1, q)$, as required.
fong-spivak-seven-sketches F03394	301	■ 1.000	$v \otimes I = v$	$v \otimes I = v$	3. The first equation is unitality in monoidal categories, $v \otimes I = v$ for any $v \in V$. The second is that
fong-spivak-seven-sketches F03395	301	■ 0.999	$I \leq Q(q, q)$	$I \leq Q(q, q)$	$I \leq Q(q, q)$ by unitality of enriched categories, see Definition 2.46, together with monotonicity of
fong-spivak-seven-sketches F03396	301	■ 1.000	$v_1 \leq v_2$	$v_1 \leq v_2$	monoidal product: $v_1 \leq v_2$ implies $v \otimes v_1 \leq v \otimes v_2$. The third was shown in Exercise 4.9.
fong-spivak-seven-sketches F03397	301	■ 1.000	$v \otimes v_1 \leq v \otimes v_2$	$v \otimes v_1 \leq v \otimes v_2$	monoidal product: $v_1 \leq v_2$ implies $v \otimes v_1 \leq v \otimes v_2$. The third was shown in Exercise 4.9.
fong-spivak-seven-sketches F03398	301	■ 0.830	$\widehat{\text{id}}: \mathcal{P} \mapsto \mathcal{P}$	$\widehat{\text{id}}: \mathcal{P} \mapsto \mathcal{P}$	This is very straightforward. We wish to check $\widehat{\text{id}}: \mathcal{P} \rightarrow \mathcal{P}$ has the formula $\widehat{\text{id}}(p, q) = \mathcal{P}(p, q)$. By
fong-spivak-seven-sketches F03399	301	■ 0.830	$\widehat{\text{id}}(p, q) = \mathcal{P}(p, q)$	$\widehat{\text{id}}(p, q) = \mathcal{P}(p, q)$	This is very straightforward. We wish to check $\widehat{\text{id}}: \mathcal{P} \rightarrow \mathcal{P}$ has the formula $\widehat{\text{id}}(p, q) = \mathcal{P}(p, q)$. By
fong-spivak-seven-sketches F03400	301	■ 0.963	$\widehat{\text{id}}(p, q) := \mathcal{P}(\text{id}(p), q) = \mathcal{P}(p, q)$	$\widehat{\text{id}}(p, q) := \mathcal{P}(\text{id}(p), q) = \mathcal{P}(p, q)$	Definition 4.34, $\widehat{\text{id}}(p, q) := \mathcal{P}(\text{id}(p), q) = \mathcal{P}(p, q)$. So they're the same.
fong-spivak-seven-sketches F03401	301	■ 0.771	$\check{+}: \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R} \times \mathbb{R}$	$\check{+}: \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R} \times \mathbb{R}$	The conjoint $\check{+}: \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R} \times \mathbb{R}$ sends (a, b, c, d) to $\mathbb{R}(a, b + c + d)$, which is true if $a \leq b + c + d$, and
fong-spivak-seven-sketches F03402	301	■ 0.771	$\mathbb{R}(a, b + c + d)$	$\mathbb{R}(a, b + c + d)$	The conjoint $\check{+}: \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R} \times \mathbb{R}$ sends (a, b, c, d) to $\mathbb{R}(a, b + c + d)$, which is true if $a \leq b + c + d$, and
fong-spivak-seven-sketches F03403	301	■ 0.771	$a \leq b + c + d$	$a \leq b + c + d$	The conjoint $\check{+}: \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R} \times \mathbb{R}$ sends (a, b, c, d) to $\mathbb{R}(a, b + c + d)$, which is true if $a \leq b + c + d$, and
fong-spivak-seven-sketches F03404	301	■ 0.976	$\widehat{F}(p, q) = Q(F(p), q)$	$\widehat{F}(p, q) = Q(F(p), q)$	1. By Definition 4.34, $\widehat{F}(p, q) = Q(F(p), q)$ and $\check{G}(p, q) = Q(p, G(q))$. Since $\mathcal{V}$ is skeletal, F and G are
fong-spivak-seven-sketches F03405	301	■ 0.976	$\check{G}(p, q) = Q(p, G(q))$	$\check{G}(p, q) = Q(p, G(q))$	1. By Definition 4.34, $\widehat{F}(p, q) = Q(F(p), q)$ and $\check{G}(p, q) = Q(p, G(q))$. Since $\mathcal{V}$ is skeletal, F and G are
fong-spivak-seven-sketches F03406	301	■ 0.992	$Q(F(p), q) = Q(p, G(q))$	$Q(F(p), q) = Q(p, G(q))$	$\mathcal{V}$ adjoints if and only if $Q(F(p), q) = Q(p, G(q))$. Thus F and G are adjoints if and only if $\widehat{F} = \check{G}$.
fong-spivak-seven-sketches F03407	301	■ 0.992	$\widehat{F} = \check{G}$	$\widehat{F} = \check{G}$	$\mathcal{V}$ adjoints if and only if $Q(F(p), q) = Q(p, G(q))$. Thus F and G are adjoints if and only if $\widehat{F} = \check{G}$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03408	301	0.898	$\widehat{\mathrm{id}}=\widehat{\mathrm{id}}$	$\mathrm{id} = \mathrm{id}$	
fong-spivak-seven-sketches F03409	301	1.000	$\mathrm{otimes} \colon \mathcal{P} \times \mathcal{P} \rightarrow \mathcal{P}$	$\otimes : \mathcal{P} \times \mathcal{P} \rightarrow \mathcal{P}$	for merely a function $\otimes \colon \mathcal{P} \times \mathcal{P} \rightarrow \mathcal{P}$, while Definition 4.45 calls for a <i>functor</i> .
fong-spivak-seven-sketches F03410	302	1.000	$\left(x_1, x_2\right) \leq \left(y_1, y_2\right)$	$(x_1, x_2) \leq (y_1, y_2)$	says that if $(x_1, x_2) \leq (y_1, y_2)$ in $\mathcal{P} \otimes \mathcal{P}$, then we must have $x_1 \otimes x_2 \leq y_1 \otimes y_2$ in $\mathcal{P}$. So it is also the case
fong-spivak-seven-sketches F03411	302	1.000	$\mathcal{P} \mathrm{otimes} \mathcal{P}$	$\mathcal{P} \otimes \mathcal{P}$	says that if $(x_1, x_2) \leq (y_1, y_2)$ in $\mathcal{P} \otimes \mathcal{P}$, then we must have $x_1 \otimes x_2 \leq y_1 \otimes y_2$ in $\mathcal{P}$. So it is also the case
fong-spivak-seven-sketches F03412	302	1.000	$g_{\{E\}}(5,3)=$	$g_E(5,3) =$	1. $g_E(5,3) = \text{false}$, $g_F(5,3) = 2$.
fong-spivak-seven-sketches F03413	302	1.000	$g_{\{F\}}(5,3)=2$	$g_F(5,3) = 2$	1. $g_E(5,3) = \text{false}$, $g_F(5,3) = 2$.
fong-spivak-seven-sketches F03414	302	1.000	$g_{\{E\}}(3,5)=$	$g_E(3,5) =$	2. $g_E(3,5) = \text{true}$, $g_F(3,5) = -2$.
fong-spivak-seven-sketches F03415	302	1.000	$g_{\{F\}}(3,5)=-2$	$g_F(3,5) = -2$	2. $g_E(3,5) = \text{true}$, $g_F(3,5) = -2$.
fong-spivak-seven-sketches F03416	302	0.988	$)=5$	$) = 5$	3. $h(5, \text{true}) = 5$.
fong-spivak-seven-sketches F03417	302	0.990	$)=-5$	$) = -5$	4. $h(-5, \text{true}) = -5$.
fong-spivak-seven-sketches F03418	302	0.999	$)=6$	$) = 6$	5. $h(-5, \text{false}) = 6$.
fong-spivak-seven-sketches F03419	302	1.000	$q_{\{G\}}(-2,3)=2, q_{\{F\}}(-2,3)=-13$	$q_G(-2,3) = 2, q_F(-2,3) = -13$	6.
fong-spivak-seven-sketches F03420	302	1.000	$q_{\{G\}}(2,3)=-1, q_{\{F\}}(2,3)=7$	$q_G(2,3) = -1, q_F(2,3) = 7$	7.
fong-spivak-seven-sketches F03421	302	0.963	$\mathcal{C}(x, y)$	$\mathcal{C}(x, y)$	a collection of objects and (ii) asks for, for all pairs of objects x, y , a <i>set</i> $\mathcal{C}(x, y)$ of morphisms. Moreover,
fong-spivak-seven-sketches F03422	302	0.430	$\mathrm{id}_x \colon I \rightarrow \mathcal{C}(x, x)$	$\mathrm{id}_x : I \rightarrow \mathcal{C}(x, x)$	recall that the monoidal unit I is the one element set $\{1\}$. This means a morphism $\mathrm{id}_x \colon I \rightarrow \mathcal{C}(x, x)$ is a
fong-spivak-seven-sketches F03423	302	0.650	$\mathrm{id}_x \colon \{1\} \rightarrow \mathcal{C}(x, x)$	$\mathrm{id}_x : \{1\} \rightarrow \mathcal{C}(x, x)$	function This is the same data as simply an element $\mathrm{id}_x = \mathrm{id}_x(1) \in \mathcal{C}(x, x)$; we call
fong-spivak-seven-sketches F03424	302	0.650	$\mathrm{id}_x=\mathrm{id}_x(1) \in \mathcal{C}(x, x)$	$\mathrm{id}_x = \mathrm{id}_x(1) \in \mathcal{C}(x, x)$	function $\mathrm{id}_x \colon \{1\} \rightarrow \mathcal{C}(x, x)$. This is the same data as simply an element $\mathrm{id}_x = \mathrm{id}_x(1) \in \mathcal{C}(x, x)$; we call
fong-spivak-seven-sketches F03425	302	0.975	$\{ \}_9 \colon \mathcal{C}(x, y) \mathrm{otimes} \mathcal{C}(y, z) \rightarrow \mathcal{C}(x, z)$	$_9 \colon \mathcal{C}(x, y) \otimes \mathcal{C}(y, z) \rightarrow \mathcal{C}(x, z)$	this data the identity morphism on x . Finally, a morphism is a function
fong-spivak-seven-sketches F03426	302	0.300	$\mathcal{C}(x, y) \mathrm{times} \mathcal{C}(y, z) \rightarrow \mathcal{C}(x, z)$	$\mathcal{C}(x, y) \times \mathcal{C}(y, z) \rightarrow \mathcal{C}(x, z)$	$\natural \colon \mathcal{C}(x, y) \times \mathcal{C}(y, z) \rightarrow \mathcal{C}(x, z)$; this is exactly the composite required in Definition 3.6 (iv).
fong-spivak-seven-sketches F03427	302	0.701	$I \rightarrow \mathcal{X}(x, x)$	$I \rightarrow \mathcal{X}(x, x)$	An identity element in a Cost -category $\mathcal{X}$ is a morphism $I \rightarrow \mathcal{X}(x, x)$ in $\mathbf{Cost} = ([0, \infty], \geq, 0, +)$, and
fong-spivak-seven-sketches F03428	302	0.937	$0 \geq \mathcal{X}(x, x)$	$0 \geq \mathcal{X}(x, x)$	hence the condition that $0 \geq \mathcal{X}(x, x)$. This implies that $\mathcal{X}(x, x) = 0$. In terms of distances, we interpret
fong-spivak-seven-sketches F03429	302	0.937	$\mathcal{X}(x, x)=0$	$\mathcal{X}(x, x) = 0$	hence the condition that $0 \geq \mathcal{X}(x, x)$. This implies that $\mathcal{X}(x, x) = 0$. In terms of distances, we interpret
fong-spivak-seven-sketches F03430	302	0.949	$\underline{3} \sqcup \underline{3}$	$\underline{3} \sqcup \underline{3}$	
fong-spivak-seven-sketches F03431	303	0.891	$x \in \mathcal{X}$	$x \in \mathcal{X}$	$x \in \mathcal{X}$ and $y \in \mathcal{Y}$, with $(x, y) \leq (x', y')$ iff $x \leq x'$ and $y \leq y'$. That is, if x is available given x' and y is
fong-spivak-seven-sketches F03432	303	0.891	$y \in \mathcal{Y}$	$y \in \mathcal{Y}$	$x \in \mathcal{X}$ and $y \in \mathcal{Y}$, with $(x, y) \leq (x', y')$ iff $x \leq x'$ and $y \leq y'$. That is, if x is available given x' and y is
fong-spivak-seven-sketches F03433	303	0.891	$(x, y) \leq (x^{\prime}, y^{\prime})$	$(x, y) \leq (x', y')$	$x \in \mathcal{X}$ and $y \in \mathcal{Y}$, with $(x, y) \leq (x', y')$ iff $x \leq x'$ and $y \leq y'$. That is, if x is available given x' and y is
fong-spivak-seven-sketches F03434	303	0.891	$y \leq y^{\prime}$	$y \leq y'$	$x \in \mathcal{X}$ and $y \in \mathcal{Y}$, with $(x, y) \leq (x', y')$ iff $x \leq x'$ and $y \leq y'$. That is, if x is available given x' and y is
fong-spivak-seven-sketches F03435	303	0.917	$\Phi \colon x_1 \rightarrow x_2$	$\Phi : x_1 \rightarrow x_2$	Given two profunctors $\Phi \colon \mathcal{X}_1 \nrightarrow \mathcal{X}_2$ and $\Psi \colon \mathcal{Y}_1 \nrightarrow \mathcal{Y}_2$, the profunctor $\Phi \times \Psi$ represents their conjunction,
fong-spivak-seven-sketches F03436	303	0.998	y_1	y_1	i.e. AND. In other words, if y_1 can be obtained given x_1 AND y_2 can be obtained given x_2 , then (y_1, y_2)

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03437	303	0.998	$x_{\{1\}}$	x_1	i.e. AND. In other words, if y_1 can be obtained given x_1 AND y_2 can be obtained given x_2 , then (y_1, y_2)
fong-spivak-seven-sketches F03438	303	0.998	$y_{\{2\}}$	y_2	i.e. AND. In other words, if y_1 can be obtained given x_1 AND y_2 can be obtained given x_2 , then (y_1, y_2)
fong-spivak-seven-sketches F03439	303	0.998	$x_{\{2\}}$	x_2	i.e. AND. In other words, if y_1 can be obtained given x_1 AND y_2 can be obtained given x_2 , then (y_1, y_2)
fong-spivak-seven-sketches F03440	303	0.998	$\left(y_{\{1\}}, y_{\{2\}}\right)$	(y_1, y_2)	i.e. AND. In other words, if y_1 can be obtained given x_1 AND y_2 can be obtained given x_2 , then (y_1, y_2)
fong-spivak-seven-sketches F03441	303	1.000	$\left(x_{\{1\}}, x_{\{2\}}\right)$	(x_1, x_2)	can be obtained given (x_1, x_2) .
fong-spivak-seven-sketches F03442	303	0.670	$\mathcal{X} \times \mathbf{1} \rightarrow \mathcal{X}$	$\mathcal{X} \times \mathbf{1} \rightarrow \mathcal{X}$	The profunctor $\mathcal{X} \times \mathbf{1} \rightarrow \mathcal{X}$ defined by the functor $\alpha: (\mathcal{X} \times \mathbf{1})^{\text{op}} \times \mathcal{X} \rightarrow \mathcal{V}$ that maps $\alpha((x, 1), y) := \mathcal{X}(x, y)$
fong-spivak-seven-sketches F03443	303	0.670	$\alpha: (\mathcal{X} \times \mathbf{1})^{\text{op}} \times \mathcal{X} \rightarrow \mathcal{V}$	$\alpha: (\mathcal{X} \times \mathbf{1})^{\text{op}} \times \mathcal{X} \rightarrow \mathcal{V}$	The profunctor $\mathcal{X} \times \mathbf{1} \rightarrow \mathcal{X}$ defined by the functor $\alpha: (\mathcal{X} \times \mathbf{1})^{\text{op}} \times \mathcal{X} \rightarrow \mathcal{V}$ that maps $\alpha((x, 1), y) := \mathcal{X}(x, y)$
fong-spivak-seven-sketches F03444	303	0.670	$\alpha((x, 1), y) := \mathcal{X}(x, y)$	$\alpha((x, 1), y) := \mathcal{X}(x, y)$	The profunctor $\mathcal{X} \times \mathbf{1} \rightarrow \mathcal{X}$ defined by the functor $\alpha: (\mathcal{X} \times \mathbf{1})^{\text{op}} \times \mathcal{X} \rightarrow \mathcal{V}$ that maps $\alpha((x, 1), y) := \mathcal{X}(x, y)$
fong-spivak-seven-sketches F03445	303	0.948	$\alpha^{-1}: x \mapsto x \times \mathbf{1}$	$\alpha^{-1}: x \mapsto x \times \mathbf{1}$	is an isomorphism. It has inverse $\alpha^{-1}: \mathcal{X} \rightarrow \mathcal{X} \times \mathbf{1}$ defined by $\alpha^{-1}(x, (y, 1)) := \mathcal{X}(x, y)$. To see that
fong-spivak-seven-sketches F03446	303	0.948	$\alpha^{-1}(x, (y, 1)) := x(x, y)$	$\alpha^{-1}(x, (y, 1)) := x(x, y)$	is an isomorphism. It has inverse $\alpha^{-1}: \mathcal{X} \rightarrow \mathcal{X} \times \mathbf{1}$ defined by $\alpha^{-1}(x, (y, 1)) := \mathcal{X}(x, y)$. To see that
fong-spivak-seven-sketches F03447	303	0.943	$\alpha^{-1}; \alpha = \text{U}_{\mathcal{X}}$	$\alpha^{-1}; \alpha = \text{U}_{\mathcal{X}}$	$\alpha^{-1} \circ \alpha = \text{U}_{\mathcal{X}}$, note first that the unit law for $\mathcal{X}$ at z and the definition of join imply
fong-spivak-seven-sketches F03448	303	0.552	$\mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \leq \mathcal{X}(x, z)$	$\mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \leq \mathcal{X}(x, z)$	while composition says $\mathcal{X}(x, y) \otimes \mathcal{X}(y, z) \leq \mathcal{X}(x, z)$ and hence
fong-spivak-seven-sketches F03449	303	0.473	$\alpha^{-1} = \text{U}_{\mathcal{X} \times \mathbf{1}}$	$\alpha^{-1} = \text{U}_{\mathcal{X} \times \mathbf{1}}$	Similarly we can show $\alpha \circ \alpha^{-1} = \text{U}_{\mathcal{X} \times \mathbf{1}}$, and hence that α is an isomorphism $\mathcal{X} \times \mathbf{1} \rightarrow \mathcal{X}$.
fong-spivak-seven-sketches F03450	303	0.556	$\beta((1, x), y) := \mathcal{X}(x, y)$	$\beta((1, x), y) := \mathcal{X}(x, y)$	Moreover, we can similarly show that $\beta((1, x), y) := \mathcal{X}(x, y)$ defines an isomorphism $\beta: \mathbf{1} \times \mathcal{X} \rightarrow \mathcal{X}$.
fong-spivak-seven-sketches F03451	303	0.556	$\beta: \mathbf{1} \times \mathcal{X} \mapsto \mathcal{X}$	$\beta: \mathbf{1} \times \mathcal{X} \mapsto \mathcal{X}$	Moreover, we can similarly show that $\beta((1, x), y) := \mathcal{X}(x, y)$ defines an isomorphism $\beta: \mathbf{1} \times \mathcal{X} \rightarrow \mathcal{X}$.
fong-spivak-seven-sketches F03452	304	0.979	α^{-1}	α^{-1}	is itself the identity (ie. the unit profunctor on $\mathcal{X}$), where α and α^{-1} are the isomorphisms defined in
fong-spivak-seven-sketches F03453	304	0.571	$\Phi(x, y)$	$\Phi(x, y)$	Freely using the distributivity of $\otimes$ over $\vee$, the value $\Phi(x, y)$ of this composite at $(x, y) \in \mathcal{X}^{\text{op}} \times \mathcal{X}$ is
fong-spivak-seven-sketches F03454	304	0.571	$(x, y) \in \mathcal{X}^{\text{op}} \times \mathcal{X}$	$(x, y) \in \mathcal{X}^{\text{op}} \times \mathcal{X}$	Freely using the distributivity of $\otimes$ over $\vee$, the value $\Phi(x, y)$ of this composite at $(x, y) \in \mathcal{X}^{\text{op}} \times \mathcal{X}$ is
fong-spivak-seven-sketches F03455	304	0.857	$\text{U}_{\mathcal{X}}(a, b) = \mathcal{X}(a, b)$	$\text{U}_{\mathcal{X}}(a, b) = \mathcal{X}(a, b)$	with the unit profunctor $\text{U}_{\mathcal{X}}(a, b) = \mathcal{X}(a, b)$ is the identity.
fong-spivak-seven-sketches F03456	305	1.000	$(f; g)(i) = g(f(i))$	$(f; g)(i) = g(f(i))$	given by $(f \circ g)(i) = g(f(i))$ for all $1 \leq i \leq m$.
fong-spivak-seven-sketches F03457	305	1.000	$1 \leq i \leq m$	$1 \leq i \leq m$	given by $(f \circ g)(i) = g(f(i))$ for all $1 \leq i \leq m$.
fong-spivak-seven-sketches F03458	305	0.562	$\text{id}_m: m \rightarrow m$	$\text{id}_m: m \rightarrow m$	4. The identity $\text{id}_m: m \rightarrow m$ is given by $\text{id}_m(i) = i$ for all $1 \leq i \leq m$. Here is a picture of id_2 and id_8 :
fong-spivak-seven-sketches F03459	305	0.562	$\text{id}_m(i) = i$	$\text{id}_m(i) = i$	4. The identity $\text{id}_m: m \rightarrow m$ is given by $\text{id}_m(i) = i$ for all $1 \leq i \leq m$. Here is a picture of id_2 and id_8 :
fong-spivak-seven-sketches F03460	305	0.562	id_2	id_2	4. The identity $\text{id}_m: m \rightarrow m$ is given by $\text{id}_m(i) = i$ for all $1 \leq i \leq m$. Here is a picture of id_2 and id_8 :
fong-spivak-seven-sketches F03461	305	0.562	id_8	id_8	4. The identity $\text{id}_m: m \rightarrow m$ is given by $\text{id}_m(i) = i$ for all $1 \leq i \leq m$. Here is a picture of id_2 and id_8 :
fong-spivak-seven-sketches F03462	305	1.000	$\sigma_{3,5}: 8 \rightarrow 8$	$\sigma_{3,5}: 8 \rightarrow 8$	5. Here is a picture of the symmetry $\sigma_{3,5}: 8 \rightarrow 8$:
fong-spivak-seven-sketches F03463	305	1.000	$m_1 \leq n_1$	$m_1 \leq n_1$	denoted $m \leq n$, and with the property that whenever $m_1 \leq n_1$ and $m_2 \leq n_2$ hold then $m_1 + m_2 \leq n_1 + n_2$
fong-spivak-seven-sketches F03464	305	1.000	$m_2 \leq n_2$	$m_2 \leq n_2$	denoted $m \leq n$, and with the property that whenever $m_1 \leq n_1$ and $m_2 \leq n_2$ hold then $m_1 + m_2 \leq n_1 + n_2$
fong-spivak-seven-sketches F03465	305	1.000	$m_1 + m_2 \leq n_1 + n_2$	$m_1 + m_2 \leq n_1 + n_2$	denoted $m \leq n$, and with the property that whenever $m_1 \leq n_1$ and $m_2 \leq n_2$ hold then $m_1 + m_2 \leq n_1 + n_2$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03466	305	■ 0.970	$d \in \mathbb{N}$	$d \in \mathbb{N}$	2. Take $\leq$ to be the usual order, $m \leq n$ iff there exists $d \in \mathbb{N}$ with $d + m = n$.
fong-spivak-seven-sketches F03467	305	■ 0.970	$d+m=n$	$d + m = n$	2. Take $\leq$ to be the usual order, $m \leq n$ iff there exists $d \in \mathbb{N}$ with $d + m = n$.
fong-spivak-seven-sketches F03468	305	■ 0.996	$m=n+d$	$m = n + d$	3. Take $\leq$ to be the reverse of the usual order, $m \leq n$ iff there exists $d \in \mathbb{N}$ with $m = n + d$.
fong-spivak-seven-sketches F03469	305	■ 0.999	$q \in \mathbb{N}$	$q \in \mathbb{N}$	5. (Non-example.) Take $\leq$ to be the division order, $m \leq n$ iff there exists $q \in \mathbb{N}$ with $m * q = d$. This
fong-spivak-seven-sketches F03470	305	■ 0.999	$m * q=d$	$m * q = d$	5. (Non-example.) Take $\leq$ to be the division order, $m \leq n$ iff there exists $q \in \mathbb{N}$ with $m * q = d$. This
fong-spivak-seven-sketches F03471	305	■ 0.932	$3 \preceq 3$	$3 \preceq 3$	$3 \leq 3$ but not $5 \preceq^? 7$.
fong-spivak-seven-sketches F03472	305	■ 0.932	$5 \leq ? 7$	$5 \leq ? 7$	$3 \leq 3$ but not $5 \leq^? 7$.
fong-spivak-seven-sketches F03473	305	■ 1.000	$\mathbf{Bij}(m, n) := \{f: \underline{m} \rightarrow \underline{n} \mid f$	$\mathbf{Bij}(m, n) := \{f: \underline{m} \rightarrow \underline{n} \mid f$	1. $\mathbf{Bij}(m, n) := \{f: \underline{m} \rightarrow \underline{n} \mid f$ is a bijection}. Note that $\mathbf{Bij}(m, n) = \emptyset$ if $m \neq n$ and it has $n!$
fong-spivak-seven-sketches F03474	305	■ 1.000	$\mathbf{Bij}(m, n) = \varnothing$	$\mathbf{Bij}(m, n) = \emptyset$	1. $\mathbf{Bij}(m, n) := \{f: \underline{m} \rightarrow \underline{n} \mid f$ is a bijection}. Note that $\mathbf{Bij}(m, n) = \emptyset$ if $m \neq n$ and it has $n!$
fong-spivak-seven-sketches F03475	305	■ 1.000	$n \rightarrow n$	$n \rightarrow n$	2. The identity map $n \rightarrow n$ is the bijection $\underline{n} \rightarrow \underline{n}$ sending $i \mapsto i$.
fong-spivak-seven-sketches F03476	305	■ 1.000	$\underline{n} \rightarrow \underline{n}$	$\underline{n} \rightarrow \underline{n}$	2. The identity map $n \rightarrow n$ is the bijection $\underline{n} \rightarrow \underline{n}$ sending $i \mapsto i$.
fong-spivak-seven-sketches F03477	305	■ 1.000	$i \mapsto i$	$i \mapsto i$	2. The identity map $n \rightarrow n$ is the bijection $\underline{n} \rightarrow \underline{n}$ sending $i \mapsto i$.
fong-spivak-seven-sketches F03478	306	■ 1.000	$m+n \rightarrow n+m$	$m + n \rightarrow n + m$	3. The symmetry map $m + n \rightarrow n + m$ is the bijection $\sigma_{m,n}: \underline{m + n} \rightarrow \underline{n + m}$ given by
fong-spivak-seven-sketches F03479	306	■ 1.000	$\sigma_{m,n}: \underline{m+n} \rightarrow \underline{n+m}$	$\sigma_{m,n}: \underline{m + n} \rightarrow \underline{n + m}$	3. The symmetry map $m + n \rightarrow n + m$ is the bijection $\sigma_{m,n}: \underline{m + n} \rightarrow \underline{n + m}$ given by
fong-spivak-seven-sketches F03480	306	■ 1.000	$n \rightarrow p$	$n \rightarrow p$	4. Composition of bijections $m \rightarrow n$ and $n \rightarrow p$ is just their composition as functions, which
fong-spivak-seven-sketches F03481	306	■ 1.000	$(f+g):(m+n) \rightarrow$	$(f + g): (m + n) \rightarrow$	5. Given bijections $f: m \rightarrow m'$ and $g: n \rightarrow n'$, their monoidal product $(f + g): (m + n) \rightarrow$
fong-spivak-seven-sketches F03482	306	■ 1.000	$\left(m^{\prime}+n^{\prime}\right)$	$(m' + n')$	$(m' + n')$ is given by
fong-spivak-seven-sketches F03483	306	■ 0.817	$\operatorname{Corel}(m, n)$	$\operatorname{Corel}(m, n)$	1. $\operatorname{Corel}(m, n)$ is the set of equivalence relations on $\underline{m + n}$.
fong-spivak-seven-sketches F03484	306	■ 0.817	$\underline{m+n}$	$\underline{m + n}$	1. $\operatorname{Corel}(m, n)$ is the set of equivalence relations on $\underline{m + n}$.
fong-spivak-seven-sketches F03485	306	■ 1.000	$i \sim j$	$i \sim j$	relation, i.e. where $i \sim j$ iff $i = j$.
fong-spivak-seven-sketches F03486	306	■ 1.000	$i=j$	$i = j$	relation, i.e. where $i \sim j$ iff $i = j$.
fong-spivak-seven-sketches F03487	306	■ 0.981	$m+n+n+m$	$m + n + n + m$	3. The symmetry map $\sigma_{m,n}$, as an equivalence relation on $\underline{m + n + n + m}$ is “the obvious
fong-spivak-seven-sketches F03488	306	■ 0.999	$ i-j =m+n+n$	$ i - j = m + n + n$	• $ i - j = m + n + n$, or
fong-spivak-seven-sketches F03489	306	■ 0.969	$m+1 \leq i \leq m+n+n$	$m + 1 \leq i \leq m + n + n$	• $m + 1 \leq i \leq m + n + n$ and $m + 1 \leq j \leq m + n + n$ and $ i - j = n$,
fong-spivak-seven-sketches F03490	306	■ 0.969	$m+1 \leq j \leq m+n+n$	$m + 1 \leq j \leq m + n + n$	• $m + 1 \leq i \leq m + n + n$ and $m + 1 \leq j \leq m + n + n$ and $ i - j = n$,
fong-spivak-seven-sketches F03491	306	■ 0.969	$ i-j =n$	$ i - j = n$	• $m + 1 \leq i \leq m + n + n$ and $m + 1 \leq j \leq m + n + n$ and $ i - j = n$.
fong-spivak-seven-sketches F03492	306	■ 1.000	$\dot{\sim}$	$\sim$	4. Composition of an equivalence relation $\sim$ on $\underline{m + n}$ and an equivalence relation $\dot{\sim}$ on $n + p$
fong-spivak-seven-sketches F03493	306	■ 1.000	$n+p$	$n + p$	4. Composition of an equivalence relation $\sim$ on $\underline{m + n}$ and an equivalence relation $\dot{\sim}$ on $n + p$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03494	306	0.624	$\simeq$	$\simeq$	is the equivalence relation $\simeq$ on $m + p$ given by $i \simeq k$ iff there exists $j \in \underline{n}$ with $i \sim j$ and $\overline{j \sim k}$.
fong-spivak-seven-sketches F03495	306	0.624	$m+p$	$m + p$	is the equivalence relation $\simeq$ on $m + p$ given by $i \simeq k$ iff there exists $j \in \underline{n}$ with $i \sim j$ and $\overline{j \sim k}$.
fong-spivak-seven-sketches F03496	306	0.624	$i \simeq k$	$i \simeq k$	is the equivalence relation $\simeq$ on $m + p$ given by $i \simeq k$ iff there exists $j \in \underline{n}$ with $i \sim j$ and $\overline{j \sim k}$.
fong-spivak-seven-sketches F03497	306	0.624	$\bar{j} \dot{\sim} k$	$\bar{j} \dot{\sim} k$	is the equivalence relation $\simeq$ on $m + p$ given by $i \simeq k$ iff there exists $j \in \underline{n}$ with $i \sim j$ and $\overline{j \dot{\sim} k}$.
fong-spivak-seven-sketches F03498	306	0.990	$\overline{m+n}$	$\overline{m + n}$	5. Given equivalence relations $\sim$ on $\overline{m + n}$ and $\sim'$ on $\overline{m' + n'}$, we need an equivalence relation
fong-spivak-seven-sketches F03499	306	0.990	$\sim^{\prime}$	$\sim'$	5. Given equivalence relations $\sim$ on $\overline{m + n}$ and $\sim'$ on $\overline{m' + n'}$, we need an equivalence relation
fong-spivak-seven-sketches F03500	306	0.990	$m^{\prime}+n^{\prime}$	$m' + n'$	5. Given equivalence relations $\sim$ on $\overline{m + n}$ and $\sim'$ on $\overline{m' + n'}$, we need an equivalence relation
fong-spivak-seven-sketches F03501	306	0.931	$\left(\sim+\sim^{\prime}\right)$	$(\sim + \sim')$	$(\sim + \sim')$ on $\overline{m + n + m' + n'}$. We take it to be “the obvious thing,” namely “using $\sim$ on the
fong-spivak-seven-sketches F03502	306	0.931	$m+n+m^{\prime}+n^{\prime}$	$m + n + m' + n'$	$(\sim + \sim')$ on $\overline{m + n + m' + n'}$. We take it to be “the obvious thing,” namely “using $\sim$ on the
fong-spivak-seven-sketches F03503	306	1.000	$i \leq m+n$	$i \leq m + n$	• $i \leq m + n$ and $j \leq m + n$ and $i \sim j$, or
fong-spivak-seven-sketches F03504	306	1.000	$j \leq m+n$	$j \leq m + n$	• $i \leq m + n$ and $j \leq m + n$ and $i \sim j$, or
fong-spivak-seven-sketches F03505	306	1.000	$m+n+1 \leq i$	$m + n + 1 \leq i$	• $m + n + 1 \leq i$ and $m + n + 1 \leq j$ and $i \sim' j$.
fong-spivak-seven-sketches F03506	306	1.000	$m+n+1 \leq j$	$m + n + 1 \leq j$	• $m + n + 1 \leq i$ and $m + n + 1 \leq j$ and $i \sim' j$.
fong-spivak-seven-sketches F03507	306	1.000	$i \sim^{\prime} j$	$i \sim' j$	• $m + n + 1 \leq i$ and $m + n + 1 \leq j$ and $i \sim' j$.
fong-spivak-seven-sketches F03508	306	1.000	$\boldsymbol{\operatorname{Rel}}(m, n)$	$\mathbf{Rel}(m, n)$	1. $\mathbf{Rel}(m, n)$ is the set of relations on the set $\underline{m} \times \underline{n}$, i.e. the set of subsets of $\underline{m} \times \underline{n}$, i.e. its
fong-spivak-seven-sketches F03509	306	1.000	$\underline{m} \times \underline{n}$	$\underline{m} \times \underline{n}$	1. $\mathbf{Rel}(m, n)$ is the set of relations on the set $\underline{m} \times \underline{n}$, i.e. the set of subsets of $\underline{m} \times \underline{n}$, i.e. its
fong-spivak-seven-sketches F03510	306	1.000	$\{(i, j) \in \underline{n} \times \underline{n} \mid i=j\}$	$\{(i, j) \in \underline{n} \times \underline{n} \mid i = j\}$	2. The identity map $\underline{n} \rightarrow \underline{n}$ is the subset $\{(i, j) \in \underline{n} \times \underline{n} \mid i = j\}$.
fong-spivak-seven-sketches F03511	306	0.999	$(i, j) \in (\underline{m+n}) \times (\underline{n+m})$	$(i, j) \in (\underline{m + n}) \times (\underline{n + m})$	3. The symmetry map $m + n \rightarrow n + m$ is the subset of pairs $(i, j) \in (\underline{m + n}) \times (\underline{n + m})$ such that
fong-spivak-seven-sketches F03512	306	0.999	$i \leq m$	$i \leq m$	• $i \leq m$ and $m + 1 \leq j$ and $i + m = j$, or
fong-spivak-seven-sketches F03513	306	0.999	$m+1 \leq j$	$m + 1 \leq j$	• $i \leq m$ and $m + 1 \leq j$ and $i + m = j$, or
fong-spivak-seven-sketches F03514	306	0.999	$i+m=j$	$i + m = j$	• $i \leq m$ and $m + 1 \leq j$ and $i + m = j$, or
fong-spivak-seven-sketches F03515	306	1.000	$m+1 \leq i$	$m + 1 \leq i$	• $m + 1 \leq i$ and $j \leq m$ and $j + m = i$.
fong-spivak-seven-sketches F03516	306	1.000	$j \leq m$	$j \leq m$	• $m + 1 \leq i$ and $j \leq m$ and $j + m = i$.
fong-spivak-seven-sketches F03517	306	1.000	$j+m=i$	$j + m = i$	• $m + 1 \leq i$ and $j \leq m$ and $j + m = i$.
fong-spivak-seven-sketches F03518	306	0.986	$R^{\prime} \subseteq \underline{m}^{\prime} \times \underline{n}^{\prime}$	$R' \subseteq \underline{m'} \times \underline{n'}$	5. Given a relation $R \subseteq \underline{m} \times \underline{n}$ and a relation $R' \subseteq \underline{m'} \times \underline{n'}$, we need a relation $(R + R') \subseteq$
fong-spivak-seven-sketches F03519	306	0.986	$\left(R+R^{\prime}\right) \subseteq$	$(R + R') \subseteq$	5. Given a relation $R \subseteq \underline{m} \times \underline{n}$ and a relation $R' \subseteq \underline{m'} \times \underline{n'}$, we need a relation $(R + R') \subseteq$
fong-spivak-seven-sketches F03520	306	0.999	$\underline{m+m^{\prime}} \times \underline{n+n^{\prime}}$	$\underline{m + m'} \times \underline{n + n'}$	$\underline{m + m'} \times \underline{n + n'}$. As stated in the example (footnote), this can be given by a universal

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03521	306	■ 0.324	$R_{\{1\}} \sqcup R_{\{2\}} \subseteq \left(\underline{m_{\{1\}}} \times \underline{n_{\{1\}}} \right) \sqcup \left(\underline{m_{\{2\}}} \times \underline{n_{\{2\}}} \right) \subseteq \left(\underline{m_{\{1\}} \sqcup m_{\{2\}}} \right) \times \left(\underline{n_{\{1\}} \sqcup n_{\{2\}}} \right)$	$R_1 \sqcup R_2 \subseteq (\underline{m_1} \times \underline{n_1}) \sqcup (\underline{m_2} \times \underline{n_2}) \subseteq (\underline{m_1 \sqcup m_2}) \times (\underline{n_1 \sqcup n_2})$	Given by $R_1 \sqcup R_2 \subseteq (\underline{m_1} \times \underline{n_1}) \sqcup (\underline{m_2} \times \underline{n_2}) \subseteq (\underline{m_1 \sqcup m_2}) \times (\underline{n_1 \sqcup n_2})$.
fong-spivak-seven-sketches F03522	307	■ 1.000	$x \leq_P y$	$\leq_P$	We have a relation $R \subseteq P \times P$ which generates a preorder $\leq_P$ on P , we have an arbitrary preorder
fong-spivak-seven-sketches F03523	307	■ 1.000	$x \leq_P y$	$x \leq_P y$	i.e. that for every $x \leq_P y$ we have $f(x) \leq_Q f(y)$. By definition of P being the reflexive, transitive
fong-spivak-seven-sketches F03524	307	■ 1.000	$x_{\{0\}}, \dots, x_{\{n\}}$	$x_0, \dots, x_n$	closure of R , we have $x \leq_P y$ iff there exists $n \in \mathbb{N}$ and $x_0, \dots, x_n$ in P with $x_0 = x$ and $x_n = y$ and
fong-spivak-seven-sketches F03525	307	■ 1.000	$x_{\{0\}} = x$	$x_0 = x$	closure of R , we have $x \leq_P y$ iff there exists $n \in \mathbb{N}$ and $x_0, \dots, x_n$ in P with $x_0 = x$ and $x_n = y$ and
fong-spivak-seven-sketches F03526	307	■ 1.000	$x_{\{n\}} = y$	$x_n = y$	closure of R , we have $x \leq_P y$ iff there exists $n \in \mathbb{N}$ and $x_0, \dots, x_n$ in P with $x_0 = x$ and $x_n = y$ and
fong-spivak-seven-sketches F03527	307	■ 1.000	$R(x_i, x_{i+1})$	$R(x_i, x_{i+1})$	$R(x_i, x_{i+1})$ for each $0 \leq i \leq n-1$. (The case $n=0$ handles reflexivity.) But then by assumption,
fong-spivak-seven-sketches F03528	307	■ 1.000	$0 \leq i \leq n-1$	$0 \leq i \leq n-1$	$R(x_i, x_{i+1})$ for each $0 \leq i \leq n-1$. (The case $n=0$ handles reflexivity.) But then by assumption,
fong-spivak-seven-sketches F03529	307	■ 1.000	$f(x_i) \leq_Q f(x_{i+1})$	$f(x_i) \leq_Q f(x_{i+1})$	$R(x_i, x_{i+1})$ implies $f(x_i) \leq_Q f(x_{i+1})$ for each i . By induction on i we show that $f(x_0) \leq_Q f(x_i)$
fong-spivak-seven-sketches F03530	307	■ 1.000	$f(x_0) \leq_Q f(x_i)$	$f(x_0) \leq_Q f(x_i)$	$R(x_i, x_{i+1})$ implies $f(x_i) \leq_Q f(x_{i+1})$ for each i . By induction on i we show that $f(x_0) \leq_Q f(x_i)$
fong-spivak-seven-sketches F03531	307	■ 0.994	$n=1, x_{\{0\}} = x$	$n=1, x_0 = x$	above description is with $n=1$, $x_0 = x$, and $x_n = y$, which we said implies $x \leq_P y$.) Since f is
fong-spivak-seven-sketches F03532	307	■ 1.000	$R(g(a), g(b))$	$R(g(a), g(b))$	1. If $R(g(a), g(b))$ holds for all $a \leq_Q b$ then g is monotone, because $R(x, y)$ implies $x \leq_P y$.
fong-spivak-seven-sketches F03533	307	■ 1.000	$a \leq_Q b$	$a \leq_Q b$	1. If $R(g(a), g(b))$ holds for all $a \leq_Q b$ then g is monotone, because $R(x, y)$ implies $x \leq_P y$.
fong-spivak-seven-sketches F03534	307	■ 1.000	$(g(a), g(b)) \notin R$	$(g(a), g(b)) \notin R$	and $(g(a), g(b)) \notin R$. Indeed, take $Q := \{1\}$ to be the free preorder on one element, and take
fong-spivak-seven-sketches F03535	307	■ 1.000	$Q := \{1\}$	$Q := \{1\}$	and $(g(a), g(b)) \notin R$. Indeed, take $Q := \{1\}$ to be the free preorder on one element, and take
fong-spivak-seven-sketches F03536	307	■ 1.000	$P := \{1\}$	$P := \{1\}$	$P := \{1\}$ with $R = \emptyset$. Then the unique function $g: Q \rightarrow P$ is monotone (because $\leq_P$ is reflexive
fong-spivak-seven-sketches F03537	307	■ 1.000	$R = \varnothing$	$R = \emptyset$	$P := \{1\}$ with $R = \emptyset$. Then the unique function $g: Q \rightarrow P$ is monotone (because $\leq_P$ is reflexive
fong-spivak-seven-sketches F03538	307	■ 1.000	$(g(1), g(1)) \notin R$	$(g(1), g(1)) \notin R$	even though R is empty), and yet $(g(1), g(1)) \notin R$.
fong-spivak-seven-sketches F03539	307	■ 1.000	$\operatorname{Mor}(\mathcal{C}) \rightarrow \operatorname{Ob}(\mathcal{C})$	$\operatorname{Mor}(\mathcal{C}) \rightarrow \operatorname{Ob}(\mathcal{C})$	1. To give a function $\operatorname{Mor}(\mathcal{C}) \rightarrow \operatorname{Ob}(\mathcal{C})$ means that for every element $\operatorname{Mor}(\mathcal{C})$ we need to give exactly
fong-spivak-seven-sketches F03540	307	■ 0.992	$q \in \operatorname{Mor}(\mathcal{C})$	$q \in \operatorname{Mor}(\mathcal{C})$	one element of $\operatorname{Ob}(\mathcal{C})$. So for dom we take any $q \in \operatorname{Mor}(\mathcal{C})$, view it as a morphism $q: y \rightarrow z$, and
fong-spivak-seven-sketches F03541	307	■ 0.992	$q: y \rightarrow z$	$q: y \rightarrow z$	one element of $\operatorname{Ob}(\mathcal{C})$. So for dom we take any $q \in \operatorname{Mor}(\mathcal{C})$, view it as a morphism $q: y \rightarrow z$, and
fong-spivak-seven-sketches F03542	307	■ 0.991	$\operatorname{cod}(q) := z$	$\operatorname{cod}(q) := z$	send it to its domain y . Similarly for cod: we put $\operatorname{cod}(q) := z$.
fong-spivak-seven-sketches F03543	307	■ 1.000	$F: \mathcal{G} \rightarrow \mathcal{C}$	$F: \mathcal{G} \rightarrow \mathcal{C}$	2. Suppose first that we are given a functor $F: \mathcal{G} \rightarrow \mathcal{C}$. On objects we have a function $\operatorname{Ob}(\mathcal{G}) \rightarrow \operatorname{Ob}(\mathcal{C})$,
fong-spivak-seven-sketches F03544	307	■ 1.000	$\operatorname{Ob}(\mathcal{G}) \rightarrow \operatorname{Ob}(\mathcal{C})$	$\operatorname{Ob}(\mathcal{G}) \rightarrow \operatorname{Ob}(\mathcal{C})$	2. Suppose first that we are given a functor $F: \mathcal{G} \rightarrow \mathcal{C}$. On objects we have a function $\operatorname{Ob}(\mathcal{G}) \rightarrow \operatorname{Ob}(\mathcal{C})$,
fong-spivak-seven-sketches F03545	307	■ 0.985	$\operatorname{Ob}(\mathcal{G}) = V$	$\operatorname{Ob}(\mathcal{G}) = V$	and this defines f since $\operatorname{Ob}(\mathcal{G}) = V$. On morphisms, first note that the arrows of graph G are ex-
fong-spivak-seven-sketches F03546	307	■ 0.970	$\operatorname{Mor}(\mathcal{G})$	$\operatorname{Mor}(\mathcal{G})$	actly the length=1 paths in G , whereas $\operatorname{Mor}(\mathcal{G})$ is the set of all paths in G , so we have an inclusion
fong-spivak-seven-sketches F03547	307	■ 1.000	$A \subseteq \operatorname{Mor}(\mathcal{G})$	$A \subseteq \operatorname{Mor}(\mathcal{G})$	$A \subseteq \operatorname{Mor}(\mathcal{G})$. The functor F provides a function $\operatorname{Mor}(\mathcal{G}) \rightarrow \operatorname{Mor}(\mathcal{C})$, which we can restrict to A

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03548	307	1.000	$\operatorname{Mor}(\mathcal{G}) \rightarrow \operatorname{Mor}(\mathcal{C})$	$\operatorname{Mor}(\mathcal{G}) \rightarrow \operatorname{Mor}(\mathcal{C})$	$A \subseteq \operatorname{Mor}(\mathcal{G})$. The functor F provides a function $\operatorname{Mor}(\mathcal{G}) \rightarrow \operatorname{Mor}(\mathcal{C})$, which we can restrict to A
fong-spivak-seven-sketches F03549	307	1.000	$\operatorname{dom}(F(r)) = F(\operatorname{dom}(r))$	$\operatorname{dom}(F(r)) = F(\operatorname{dom}(r))$	to obtain $g: A \rightarrow \operatorname{Mor}(\mathcal{C})$. All functors satisfy $\operatorname{dom}(F(r)) = F(\operatorname{dom}(r))$ and $\operatorname{cod}(F(r)) = F(\operatorname{cod}(r))$
fong-spivak-seven-sketches F03550	307	1.000	$\operatorname{cod}(F(r)) = F(\operatorname{cod}(r))$	$\operatorname{cod}(F(r)) = F(\operatorname{cod}(r))$	to obtain $g: A \rightarrow \operatorname{Mor}(\mathcal{C})$. All functors satisfy $\operatorname{dom}(F(r)) = F(\operatorname{dom}(r))$ and $\operatorname{cod}(F(r)) = F(\operatorname{cod}(r))$
fong-spivak-seven-sketches F03551	307	1.000	$r: w \rightarrow x$	$r: w \rightarrow x$	for any $r: w \rightarrow x$. In particular when $r \in A$ is an arrow we have $\operatorname{dom}(r) = s(r)$ and $\operatorname{cod}(r) = t(r)$.
fong-spivak-seven-sketches F03552	307	1.000	$r \in A$	$r \in A$	for any $r: w \rightarrow x$. In particular when $r \in A$ is an arrow we have $\operatorname{dom}(r) = s(r)$ and $\operatorname{cod}(r) = t(r)$.
fong-spivak-seven-sketches F03553	307	1.000	$\operatorname{dom}(r) = s(r)$	$\operatorname{dom}(r) = s(r)$	for any $r: w \rightarrow x$. In particular when $r \in A$ is an arrow we have $\operatorname{dom}(r) = s(r)$ and $\operatorname{cod}(r) = t(r)$.
fong-spivak-seven-sketches F03554	307	1.000	$\operatorname{cod}(r) = t(r)$	$\operatorname{cod}(r) = t(r)$	for any $r: w \rightarrow x$. In particular when $r \in A$ is an arrow we have $\operatorname{dom}(r) = s(r)$ and $\operatorname{cod}(r) = t(r)$.
fong-spivak-seven-sketches F03555	308	0.561	$f: V \rightarrow \operatorname{Ob}(\mathcal{C})$	$f: V \rightarrow \operatorname{Ob}(\mathcal{C})$	Suppose second that we are given a pair of functions (f, g) where $f: V \rightarrow \operatorname{Ob}(\mathcal{C})$ and $g: A \rightarrow$
fong-spivak-seven-sketches F03556	308	0.561	$g: A \rightarrow$	$g: A \rightarrow$	Suppose second that we are given a pair of functions (f, g) where $f: V \rightarrow \operatorname{Ob}(\mathcal{C})$ and $g: A \rightarrow$
fong-spivak-seven-sketches F03557	308	1.000	$p := (v_0, a_1, a_2, \dots, a_n)$	$p := (v_0, a_1, a_2, \dots, a_n)$	objects by f . An arbitrary morphism in $\mathcal{G}$ is a path $p := (v_0, a_1, a_2, \dots, a_n)$ in G , where $v_0 \in V$,
fong-spivak-seven-sketches F03558	308	1.000	$v_0 \in V$	$v_0 \in V$	objects by f . An arbitrary morphism in $\mathcal{G}$ is a path $p := (v_0, a_1, a_2, \dots, a_n)$ in G , where $v_0 \in V$,
fong-spivak-seven-sketches F03559	308	1.000	$a_i \in A, v_0 = s(a_1)$	$a_i \in A, v_0 = s(a_1)$	$a_i \in A, v_0 = s(a_1)$, and $t(a_i) = s(a_{i+1})$ for all $1 \leq i \leq n - 1$. Then $g(a_i)$ is a morphism in $\mathcal{C}$ whose
fong-spivak-seven-sketches F03560	308	1.000	$1 \leq i \leq n - 1$	$1 \leq i \leq n - 1$	$a_i \in A, v_0 = s(a_1)$, and $t(a_i) = s(a_{i+1})$ for all $1 \leq i \leq n - 1$. Then $g(a_i)$ is a morphism in $\mathcal{C}$ whose
fong-spivak-seven-sketches F03561	308	1.000	$g(a_i)$	$g(a_i)$	$a_i \in A, v_0 = s(a_1)$, and $t(a_i) = s(a_{i+1})$ for all $1 \leq i \leq n - 1$. Then $g(a_i)$ is a morphism in $\mathcal{C}$ whose
fong-spivak-seven-sketches F03562	308	1.000	$f(v_0)$	$f(v_0)$	domain is $f(v_0)$ and the morphisms $g(a_i)$ and $g(a_{i+1})$ are composable for every $1 \leq i \leq n - 1$.
fong-spivak-seven-sketches F03563	308	1.000	$g(a_{i+1})$	$g(a_{i+1})$	domain is $f(v_0)$ and the morphisms $g(a_i)$ and $g(a_{i+1})$ are composable for every $1 \leq i \leq n - 1$.
fong-spivak-seven-sketches F03564	308	0.605	$F(p) := \operatorname{id}_{f(v_0)} \circ g(a_1) \circ \dots \circ g(a_n)$	$F(p) := \operatorname{id}_{f(v_0)} \circ g(a_1) \circ \dots \circ g(a_n)$	We then take $F(p) := \operatorname{id}_{f(v_0)} \circ g(a_1) \circ \dots \circ g(a_n)$ to be the composite. It is easy to check that this is
fong-spivak-seven-sketches F03565	308	1.000	(f', g')	(f', g')	them into a functor $F: \mathcal{G} \rightarrow \mathcal{C}$ and then extract the new pair of functions (f', g') , then $f = f'$ and
fong-spivak-seven-sketches F03566	308	1.000	$f = f'$	$f = f'$	them into a functor $F: \mathcal{G} \rightarrow \mathcal{C}$ and then extract the new pair of functions (f', g') , then $f = f'$ and
fong-spivak-seven-sketches F03567	308	0.992	$g = g'$	$g = g'$	$g = g'$. Finally, given a functor $F: \mathcal{G} \rightarrow \mathcal{C}$, we extract the pair of functions (f, g) as above and
fong-spivak-seven-sketches F03568	308	1.000	$F': \mathcal{G} \rightarrow \mathcal{C}$	$F': \mathcal{G} \rightarrow \mathcal{C}$	then turn them into a new functor $F': \mathcal{G} \rightarrow \mathcal{C}$. It is clear that F and F' act the same on objects,
fong-spivak-seven-sketches F03569	308	1.000	F'	F'	then turn them into a new functor $F': \mathcal{G} \rightarrow \mathcal{C}$. It is clear that F and F' act the same on objects,
fong-spivak-seven-sketches F03570	308	0.499	$U(\mathcal{C}) \in$	$U(\mathcal{C}) \in$	3. $(\operatorname{Mor}(\mathcal{C}), \operatorname{Ob}(\mathcal{C}), \operatorname{dom}, \operatorname{cod})$ is a graph; let's denote it $U(\mathcal{C}) \in \mathbf{Grph}$. We have functors Free : $\mathbf{Grph} \rightleftarrows$
fong-spivak-seven-sketches F03571	308	1.000	$a^i * a^j = a^{i+j}$	$a^i * a^j = a^{i+j}$	$a^i * a^j = a^{i+j}$
fong-spivak-seven-sketches F03572	308	1.000	$a^i \mapsto i$	$a^i \mapsto i$	2. This is isomorphic to $\mathbb{N}$, by sending $a^i \mapsto i$.
fong-spivak-seven-sketches F03573	308	0.560	$\operatorname{prop} \operatorname{Free}(G, s, t)$	$\operatorname{prop} \operatorname{Free}(G, s, t)$	We have two props: the prop of port graphs and the free prop Free (G, s, t) where
fong-spivak-seven-sketches F03574	308	0.963	(Γ, ℓ)	(Γ, ℓ)	By Definition 5.25, a morphism $m \rightarrow n$ in Free (G) is a G -labeled port graph, i.e. a pair (Γ, ℓ) , where
fong-spivak-seven-sketches F03575	308	0.997	$\ell(v) \in G$	$\ell(v) \in G$	right-hand ports—an arity—and $\ell(v) \in G$ is supposed to have the correct arity; precisely, $s(\ell(v)) = \operatorname{in}(v)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03576	308	■ 0.997	$s(\ell(v))=in(v)$	$s(\ell(v)) = in(v)$	right-hand ports—an arity—and $\ell(v) \in G$ is supposed to have the correct arity; precisely, $s(\ell(v)) = in(v)$
fong-spivak-seven-sketches F03577	308	■ 1.000	Γ	Γ	graph Γ , as desired.
fong-spivak-seven-sketches F03578	309	■ 0.491	$\left(f+\mathrm{id}_{\{1\}}+\mathrm{id}_{\{1\}}\right) \stackrel{\circ}{\circ} \left(\sigma+\mathrm{id}_{\{1\}}\right) \stackrel{\circ}{\circ} \left(\mathrm{id}_{\{1\}}+h\right) \stackrel{\circ}{\circ} \sigma \stackrel{\circ}{\circ} g$	$(f + \mathrm{id}_1 + \mathrm{id}_1) \stackrel{\circ}{\circ} (\sigma + \mathrm{id}_1) \stackrel{\circ}{\circ} (\mathrm{id}_1 + h) \stackrel{\circ}{\circ} \sigma \stackrel{\circ}{\circ} g$	Here is a picture of $(f + \mathrm{id}_1 + \mathrm{id}_1) \stackrel{\circ}{\circ} (\sigma + \mathrm{id}_1) \stackrel{\circ}{\circ} (\mathrm{id}_1 + h) \stackrel{\circ}{\circ} \sigma \stackrel{\circ}{\circ} g$, in the free prop on generators $G = \{f: 1 \rightarrow$
fong-spivak-seven-sketches F03579	309	■ 0.491	$G=\{f: 1 \rightarrow$	$G = \{f: 1 \rightarrow$	Here is a picture of $(f + \mathrm{id}_1 + \mathrm{id}_1) \stackrel{\circ}{\circ} (\sigma + \mathrm{id}_1) \stackrel{\circ}{\circ} (\mathrm{id}_1 + h) \stackrel{\circ}{\circ} \sigma \stackrel{\circ}{\circ} g$, in the free prop on generators $G = \{f: 1 \rightarrow$
fong-spivak-seven-sketches F03580	309	■ 0.867	$1, g: 2 \rightarrow 2, h: 2 \rightarrow 1\}$:	$1, g: 2 \rightarrow 2, h: 2 \rightarrow 1\} :$	$1, g: 2 \rightarrow 2, h: 2 \rightarrow 1\}$:
fong-spivak-seven-sketches F03581	309	■ 0.996	$\set{1}, \set{2}, \set{3}$	$\{\{1\}, \{2\}, \{3\}\}$	example the quotient of $S = \{1, 2, 3\}$ by the trivial equivalence relation is the set $\{\{1\}, \{2\}, \{3\}\}$. It is
fong-spivak-seven-sketches F03582	309	■ 0.992	$(R, 0, +, 1, *)$	$(R, 0, +, 1, *)$	1. If $(R, 0, +, 1, *)$ is a rig, then the multiplicative identity $1 \in \mathrm{Mat}_n(R)$ is the usual n -by- n identity
fong-spivak-seven-sketches F03583	309	■ 1.000	$n=4$	$n = 4$	of R). So for $n=4$ it is:
fong-spivak-seven-sketches F03584	309	■ 1.000	$A, B \in \operatorname{Mat}_2(\mathbb{N})$	$A, B \in \mathrm{Mat}_2(\mathbb{N})$	2. We choose $n = 2$ and hence need to find two elements $A, B \in \mathrm{Mat}_2(\mathbb{N})$ such that $A * B \neq B * A$.
fong-spivak-seven-sketches F03585	309	■ 1.000	$A * B \neq B * A$	$A * B \neq B * A$	2. We choose $n = 2$ and hence need to find two elements $A, B \in \mathrm{Mat}_2(\mathbb{N})$ such that $A * B \neq B * A$.
fong-spivak-seven-sketches F03586	309	■ 0.737	$(A * B)(1, 1) =$	$(A * B)(1, 1) =$	One can calculate from the multiplication formula (recalled in Example 5.40) says $(A * B)(1, 1) =$
fong-spivak-seven-sketches F03587	309	■ 0.999	$0 * 0 + 1 * 1 = 1$	$0 * 0 + 1 * 1 = 1$	$0 * 0 + 1 * 1 = 1$ and $(B * A)(1, 1) = 0 * 0 + 0 * 0 = 0$, which are not equal.
fong-spivak-seven-sketches F03588	309	■ 0.999	$(B * A)(1, 1) = 0 * 0 + 0 * 0 = 0$	$(B * A)(1, 1) = 0 * 0 + 0 * 0 = 0$	$0 * 0 + 1 * 1 = 1$ and $(B * A)(1, 1) = 0 * 0 + 0 * 0 = 0$, which are not equal.
fong-spivak-seven-sketches F03589	309	■ 0.991	$(16 \ x + 4 \ y, \ x + 4 \ y)$	$(16x + 4y, x + 4y)$	the resulting output signal is $(16x + 4y, x + 4y)$.
fong-spivak-seven-sketches F03590	309	■ 1.000	$A=\left(\begin{array}{lll}3 & 3 & 1 \\ 2 & 0 & 4\end{array}\right)$	$A = \left(\begin{array}{ccc}3 & 3 & 1 \\ 2 & 0 & 4\end{array}\right)$	The monoidal product of $A = \left(\begin{array}{ccc}3 & 3 & 1 \\ 2 & 0 & 4\end{array}\right)$ and $B = \left(\begin{array}{ccc}2 & 5 & 6 \\ 1 & 1 & 1\end{array}\right)$ is
fong-spivak-seven-sketches F03591	309	■ 1.000	$B=\left(\begin{array}{llll}2 & 5 & 6 & 1\end{array}\right)$	$B = \left(\begin{array}{cccc}2 & 5 & 6 & 1\end{array}\right)$	The monoidal product of $A = \left(\begin{array}{ccc}3 & 3 & 1 \\ 2 & 0 & 4\end{array}\right)$ and $B = \left(\begin{array}{ccc}2 & 5 & 6 \\ 1 & 1 & 1\end{array}\right)$ is
fong-spivak-seven-sketches F03592	310	■ 1.000	$c_{\{n\}}$	c_n	• For the first layer g_1 , take the monoidal product of m copies of c_n ,
fong-spivak-seven-sketches F03593	310	■ 0.999	$s_{\{a\}}: 1 \rightarrow 1$	$s_a: 1 \rightarrow 1$	where $s_a: 1 \rightarrow 1$ is the signal flow graph generator “scalar multiplication by a .” This layer
fong-spivak-seven-sketches F03594	310	■ 0.993	$g_{\{3\}}: m \times n \rightarrow m \times n$	$g_3: m \times n \rightarrow m \times n$	signal flow graph $g_3: m \times n \rightarrow m \times n$, that is the composite and monoidal product of swap and
fong-spivak-seven-sketches F03595	310	■ 1.000	$(i-1) \ m+j$	$(i-1)m + j$	identity maps, such that the $(i-1)m + j$ th input is sent to the $(j-1)n + i$ th output, for all $1 \leq i \leq n$
fong-spivak-seven-sketches F03596	310	■ 1.000	$(j-1) \ n+i$	$(j-1)n + i$	identity maps, such that the $(i-1)m + j$ th input is sent to the $(j-1)n + i$ th output, for all $1 \leq i \leq n$
fong-spivak-seven-sketches F03597	310	■ 1.000	$1 \leq i \leq n$	$1 \leq i \leq n$	identity maps, such that the $(i-1)m + j$ th input is sent to the $(j-1)n + i$ th output, for all $1 \leq i \leq n$
fong-spivak-seven-sketches F03598	310	■ 1.000	$1 \leq j \leq m$	$1 \leq j \leq m$	and $1 \leq j \leq m$.
fong-spivak-seven-sketches F03599	311	■ 0.999	$a_{\{m\}}$	a_m	where a_m is the signal flow graph that adds m inputs to produce a single output:
fong-spivak-seven-sketches F03600 [conf 0.087]	311	■ 0.087	$g=g_{\{1\}} \ g_{\{2\}} \ g_{\{3\}} \ g_{\{4\}}: m \rightarrow$	$g = g_1 g_2 g_3 g_4: m \rightarrow$	Using Proposition 5.54, it is a straightforward but tedious calculation to show that $g = g_1 \circ g_2 \circ g_3 \circ g_4: m \rightarrow$
fong-spivak-seven-sketches F03601	312	■ 0.729	$\sigma_{\{M, M\}}: \mu = \mu$	$\sigma_{M, M}: \mu = \mu$	(c) $\sigma_{M, M} \circ \mu = \mu$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03602	312	1.000	$R^{\{0\}} = \{ \{ () \} \}$	$R^0 = \{ \{ () \} \}$	particular, $R^0 = \{ \{ () \} \}$ consists of a single vector of dimension 0.
fong-spivak-seven-sketches F03603	312	0.977	$\operatorname{prop}(\operatorname{Mat}(R))$	$\operatorname{prop}(\operatorname{Mat}(R))$	1. U preserves the monoidal unit because 0 is the monoidal unit of any prop ($\operatorname{Mat}(R)$ is a prop),
fong-spivak-seven-sketches F03604	312	0.874	$R^{\{0\}}$	R^0	$\{1\}$ is the monoidal unit of Set , and R^0 is canonically isomorphic to $\{1\}$. U also preserves the
fong-spivak-seven-sketches F03605	312	0.874	$\{1\} \cdot U$	$\{1\}.U$	$\{1\}$ is the monoidal unit of Set , and R^0 is canonically isomorphic to $\{1\}$. U also preserves the
fong-spivak-seven-sketches F03606	312	1.000	$R^{\{m\}} \times R^{\{n\}} \cong R^{\{m+n\}}$	$R^m \times R^n \cong R^{m+n}$	monoidal product because there is a canonical isomorphism $R^m \times R^n \cong R^{m+n}$.
fong-spivak-seven-sketches F03607	312	0.889	(m, μ, η)	(m, μ, η)	2. A monoid object in $\operatorname{Mat}(R)$ is a tuple (m, μ, η) where $m \in \mathbb{N}$, $\mu: m + m \rightarrow m$, and $\eta: 0 \rightarrow m$
fong-spivak-seven-sketches F03608	312	0.889	$m \in \mathbb{N}, \mu: m+m \rightarrow m$	$m \in \mathbb{N}, \mu: m + m \rightarrow m$	2. A monoid object in $\operatorname{Mat}(R)$ is a tuple (m, μ, η) where $m \in \mathbb{N}$, $\mu: m + m \rightarrow m$, and $\eta: 0 \rightarrow m$
fong-spivak-seven-sketches F03609	312	0.889	$\eta: 0 \rightarrow m$	$\eta: 0 \rightarrow m$	2. A monoid object in $\operatorname{Mat}(R)$ is a tuple (m, μ, η) where $m \in \mathbb{N}$, $\mu: m + m \rightarrow m$, and $\eta: 0 \rightarrow m$
fong-spivak-seven-sketches F03610	312	1.000	$\mu(\eta, x) = x = \mu(x, \eta)$	$\mu(\eta, x) = x = \mu(x, \eta)$	satisfy the properties $\mu(\eta, x) = x = \mu(x, \eta)$ and $\mu(x, \mu(y, z)) = \mu(\mu(x, y), z)$. Note that there is
fong-spivak-seven-sketches F03611	312	1.000	$\mu(x, \mu(y, z)) = \mu(\mu(x, y), z)$	$\mu(x, \mu(y, z)) = \mu(\mu(x, y), z)$	satisfy the properties $\mu(\eta, x) = x = \mu(x, \eta)$ and $\mu(x, \mu(y, z)) = \mu(\mu(x, y), z)$. Note that there is
fong-spivak-seven-sketches F03612	312	0.970	$0 \rightarrow m$	$0 \rightarrow m$	only one morphism $0 \rightarrow m$ in $\operatorname{Mat}(R)$ for any m . It is not hard to show that for any $m \in \mathbb{N}$ there
fong-spivak-seven-sketches F03613	312	0.970	$m \in \mathbb{N}$	$m \in \mathbb{N}$	only one morphism $0 \rightarrow m$ in $\operatorname{Mat}(R)$ for any m . It is not hard to show that for any $m \in \mathbb{N}$ there
fong-spivak-seven-sketches F03614	312	1.000	$m=2, \mu$	$m = 2, \mu$	is only one monoid structure. For example, when $m = 2$, μ must be the following matrix
fong-spivak-seven-sketches F03615	313	0.999	$U(\eta): R^{\{0\}} \rightarrow R^{\{m\}}$	$U(\eta): R^0 \rightarrow R^m$	Anyway, for any monoid (m, μ, η) , the morphism $U(\eta): R^0 \rightarrow R^m$ is given by $U(\eta)(1) :=$
fong-spivak-seven-sketches F03616	313	0.999	$U(\eta)(1) :=$	$U(\eta)(1) :=$	Anyway, for any monoid (m, μ, η) , the morphism $U(\eta): R^0 \rightarrow R^m$ is given by $U(\eta)(1) :=$
fong-spivak-seven-sketches F03617	313	0.985	$(0, \dots, 0)$	$(0, \dots, 0)$	$(0, \dots, 0)$, and the morphism $U(\mu): R^m \times R^m \rightarrow R^m$ is given by
fong-spivak-seven-sketches F03618	313	0.985	$U(\mu): R^{\{m\}} \times R^{\{m\}} \rightarrow R^{\{m\}}$	$U(\mu): R^m \times R^m \rightarrow R^m$	$(0, \dots, 0)$, and the morphism $U(\mu): R^m \times R^m \rightarrow R^m$ is given by
fong-spivak-seven-sketches F03619	313	1.000	$R^{\{m\}}$	R^m	These give R^m the structure of a monoid.
fong-spivak-seven-sketches F03620	313	0.860	$(1, \supset, \sim)$	$(1, \supset, \sim)$	3. The triple $(1, \supset, \sim)$ corresponds to the additive monoid structure on $\mathbb{R}$, e.g. with $(5, 3) \mapsto 8$.
fong-spivak-seven-sketches F03621	313	0.860	$(5, 3) \mapsto 8$	$(5, 3) \mapsto 8$	3. The triple $(1, \supset, \sim)$ corresponds to the additive monoid structure on $\mathbb{R}$, e.g. with $(5, 3) \mapsto 8$.
fong-spivak-seven-sketches F03622	313	0.989	$\left\{ (x, y, z) \in R^{\{3\}} \mid x = \right.$	$\{ (x, y, z) \in R^3 \mid x =$	1. The behavior $B(\neg\lhd)$ of the reversed addition icon $\neg\lhd: 1 \rightarrow 2$ is the relation $\{ (x, y, z) \in R^3 \mid x =$
fong-spivak-seven-sketches F03623	313	1.000	$y+z\}$	$y + z\}$	$y + z\}$.
fong-spivak-seven-sketches F03624	313	0.395	$\left\{ (x, y, z) \in R^{\{3\}} \mid x=y=\right.$	$\{ (x, y, z) \in R^3 \mid x = y =$	2. The behavior $B(\rhd)$ of the reversed copy icon, $\rhd: 2 \rightarrow 1$ is the relation $\{ (x, y, z) \in R^3 \mid x = y =$
fong-spivak-seven-sketches F03625	313	1.000	$z\}$	$z\}$	—
fong-spivak-seven-sketches F03626	313	0.926	$B+C \subseteq R^{\{m+p\}} \times R^{\{n+q\}}$	$B + C \subseteq R^{m+p} \times R^{n+q}$	If $B \subseteq R^m \times R^n$ and $C \subseteq R^p \times R^q$ are morphisms in $\mathbf{Rel}_R$, then take $B + C \subseteq R^{m+p} \times R^{n+q}$ to be the set
fong-spivak-seven-sketches F03627	313	0.624	$\operatorname{B}(\operatorname{g}^{\circ}(\operatorname{h}^{\operatorname{op}})) = \operatorname{B}(g); \operatorname{B}(\operatorname{h}^{\operatorname{op}})$	$B(g^{\circ}(h^{\operatorname{op}})) = B(g); B(h^{\operatorname{op}})$	and by Eq. (5.78), the composite $B(g \circ (h^{\operatorname{op}})) = B(g) \circ B(h^{\operatorname{op}})$ is:
fong-spivak-seven-sketches F03628	313	1.000	$S(h)$	$S(h)$	Since $S(g)$ and $S(h)$ are functions, the above immediately reduces to the desired formula:
fong-spivak-seven-sketches F03629	313	0.829	$\operatorname{g}^{\operatorname{op}}: n \rightarrow m$	$g^{\operatorname{op}}: n \rightarrow m$	The behavior of $g^{\operatorname{op}}: n \rightarrow m$ and $h: m \rightarrow p$ are respectively

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03630	313	■ 0.829	$h: m \rightarrow p$	$h: m \rightarrow p$	The behavior of $g^{\text{op}}: n \rightarrow m$ and $h: m \rightarrow p$ are respectively
fong-spivak-seven-sketches F03631	313	■ 0.882	$\mathrm{B}(\left(\left(g^{\mathrm{op}}\right) ; h\right) ; \mathrm{B}(h)=\mathrm{B}(\left(g^{\mathrm{op}}\right) ; \mathrm{B}(h)$	$B((g^{\text{op}}); h) = B(g^{\text{op}}); B(h)$	and by Eq. (5.78), the composite $\mathrm{B}(g^{\text{op}}) \circ B(h)$ is:
fong-spivak-seven-sketches F03632	313	■ 0.597	$\{y \in R \mid y=0\}$	$\{y \in R \mid y=0\}$	1. The behavior of the 0-reverse $\dashv$ is the subset $\{y \in R \mid y=0\}$, and its n -fold tensor is similarly
fong-spivak-seven-sketches F03633	313	■ 1.000	$\left\{y \in R^{n} \mid y=0\right\}$	$\{y \in R^n \mid y=0\}$	$\{y \in R^n \mid y=0\}$. Composing this relation with $S(g) \subseteq R^m \times R^n$ gives $\{x \in R^m \mid S(g)=0\}$,
fong-spivak-seven-sketches F03634	313	■ 1.000	$S(g) \subseteq R^m \times R^n$	$S(g) \subseteq R^m \times R^n$	$\{y \in R^n \mid y=0\}$. Composing this relation with $S(g) \subseteq R^m \times R^n$ gives $\{x \in R^m \mid S(g)=0\}$,
fong-spivak-seven-sketches F03635	313	■ 1.000	$\left\{x \in R^m \mid S(g)=0\right\}$	$\{x \in R^m \mid S(g)=0\}$	$\{y \in R^n \mid y=0\}$. Composing this relation with $S(g) \subseteq R^m \times R^n$ gives $\{x \in R^m \mid S(g)=0\}$,
fong-spivak-seven-sketches F03636	313	■ 0.494	$\{x \in R\}$	$\{x \in R\}$	2. The behavior of the discard-inverse $\dashv$ is the subset $\{x \in R\}$, i.e. the largest subset of R , and
fong-spivak-seven-sketches F03637	313	■ 1.000	$R^n \subseteq R^n$	$R^n \subseteq R^n$	similarly its m -fold tensor is $R^n \subseteq R^n$. Composing this relation with $S(g) \subseteq R^m \times R^n$ gives
fong-spivak-seven-sketches F03638	313	■ 0.999	$\left\{y \in R^n \mid \text{there exists } x \in R^m \text{ such that } S(g)(x)=y\right\}$	$\{y \in R^n \mid$	$\{y \in R^n \mid$ there exists $x \in R^m$ such that $S(g)(x) = y\}$, which is exactly the image of $S(g)$.
fong-spivak-seven-sketches F03639	313	■ 0.999	$S(g)(x)=y$	$S(g)(x)=y\}$	$\{y \in R^n \mid$ there exists $x \in R^m$ such that $S(g)(x) = y\}$, which is exactly the image of $S(g)$.
fong-spivak-seven-sketches F03640	314	■ 0.994	$\mathrm{B}(g)=\{(x, y) \mid S(g)(x)=y\}$	$B(g)=\{(x, y) \mid S(g)(x)=y\}$	3. For any $g: m \rightarrow n$, we first claim that the behavior $\mathrm{B}(g)=\{(x, y) \mid S(g)(x)=y\}$ is linear, i.e. it
fong-spivak-seven-sketches F03641	314	■ 1.000	$S(g)(x)=y$	$S(g)(x)=y$	so if $S(g)(x)=y$ then $S(g)(rx)=ry$ and $S(g)(x_1+x_2)=S(g)(x_1)+S(g)(x_2)$. Thus we con-
fong-spivak-seven-sketches F03642	314	■ 1.000	$S(g)(rx)=ry$	$S(g)(rx)=ry$	so if $S(g)(x)=y$ then $S(g)(rx)=ry$ and $S(g)(x_1+x_2)=S(g)(x_1)+S(g)(x_2)$. Thus we con-
fong-spivak-seven-sketches F03643	314	■ 1.000	$S(g)\left(x_1+x_2\right)=S(g)\left(x_1\right)+S(g)\left(x_2\right)$	$S(g)\left(x_1+x_2\right)=S(g)\left(x_1\right)+S(g)\left(x_2\right)$	so if $S(g)(x)=y$ then $S(g)(rx)=ry$ and $S(g)(x_1+x_2)=S(g)(x_1)+S(g)(x_2)$. Thus we con-
fong-spivak-seven-sketches F03644	314	■ 1.000	$(x, y) \in \mathrm{B}(g)$	$(x, y) \in B(g)$	clude that $(x, y) \in B(g)$ implies $(rx, ry) \in B(g)$, so it's closed under scalar multiplication, and
fong-spivak-seven-sketches F03645	314	■ 1.000	$(rx, ry) \in \mathrm{B}(g)$	$(rx, ry) \in B(g)$	clude that $(x, y) \in B(g)$ implies $(rx, ry) \in B(g)$, so it's closed under scalar multiplication, and
fong-spivak-seven-sketches F03646	314	■ 0.999	$\left(x_1, y_1\right),\left(x_2, y_2\right) \in \mathrm{B}(g)$	$\left(x_1, y_1\right),\left(x_2, y_2\right) \in B(g)$	$\left(x_1, y_1\right),\left(x_2, y_2\right) \in B(g)$ implies $\left(x_1+x_2, y_1+y_2\right) \in B(g)$ so it's closed under addition. Similarly,
fong-spivak-seven-sketches F03647	314	■ 0.999	$\left(x_1+x_2, y_1+y_2\right) \in \mathrm{B}(g)$	$\left(x_1+x_2, y_1+y_2\right) \in B(g)$	$\left(x_1, y_1\right),\left(x_2, y_2\right) \in B(g)$ implies $\left(x_1+x_2, y_1+y_2\right) \in B(g)$ so it's closed under addition. Similarly,
fong-spivak-seven-sketches F03648	314	■ 1.000	$\mathrm{B}\left(g^{\mathrm{op}}\right)$	$B\left(g^{\text{op}}\right)$	the behavior $\mathrm{B}\left(g^{\text{op}}\right)$ is also linear; the proof is similar.
fong-spivak-seven-sketches F03649	314	■ 1.000	$C \subseteq R^n \times R^p$	$C \subseteq R^n \times R^p$	$B \subseteq R^m \times R^n$ and $C \subseteq R^n \times R^p$ are linear. Take $\left(x_1, z_1\right),\left(x_2, z_2\right) \in B \circ C$ and take $r \in R$.
fong-spivak-seven-sketches F03650	314	■ 1.000	$\left(x_1, z_1\right),\left(x_2, z_2\right) \in B ; C$	$\left(x_1, z_1\right),\left(x_2, z_2\right) \in B ; C$	$B \subseteq R^m \times R^n$ and $C \subseteq R^n \times R^p$ are linear. Take $\left(x_1, z_1\right),\left(x_2, z_2\right) \in B \circ C$ and take $r \in R$.
fong-spivak-seven-sketches F03651	314	■ 1.000	$y_1, y_2 \in R^n$	$y_1, y_2 \in R^n$	By definition, there exist $y_1, y_2 \in R^n$ such that $\left(x_1, y_1\right),\left(x_2, y_2\right) \in B$ and $\left(y_1, z_1\right),\left(y_2, z_2\right) \in C$.
fong-spivak-seven-sketches F03652	314	■ 1.000	$\left(x_1, y_1\right),\left(x_2, y_2\right) \in B$	$\left(x_1, y_1\right),\left(x_2, y_2\right) \in B$	By definition, there exist $y_1, y_2 \in R^n$ such that $\left(x_1, y_1\right),\left(x_2, y_2\right) \in B$ and $\left(y_1, z_1\right),\left(y_2, z_2\right) \in C$.
fong-spivak-seven-sketches F03653	314	■ 1.000	$\left(y_1, z_1\right),\left(y_2, z_2\right) \in C$	$\left(y_1, z_1\right),\left(y_2, z_2\right) \in C$	By definition, there exist $y_1, y_2 \in R^n$ such that $\left(x_1, y_1\right),\left(x_2, y_2\right) \in B$ and $\left(y_1, z_1\right),\left(y_2, z_2\right) \in C$.
fong-spivak-seven-sketches F03654	314	■ 1.000	$\left(r x_1, r y_1\right) \in B$	$\left(r x_1, r y_1\right) \in B$	Since B and C are linear, $\left(r x_1, r y_1\right) \in B$ and $\left(r y_1, r z_1\right) \in C$, and also $\left(x_1+x_2, y_1+y_2\right) \in B$ and
fong-spivak-seven-sketches F03655	314	■ 1.000	$\left(r y_1, r z_1\right) \in C$	$\left(r y_1, r z_1\right) \in C$	Since B and C are linear, $\left(r x_1, r y_1\right) \in B$ and $\left(r y_1, r z_1\right) \in C$, and also $\left(x_1+x_2, y_1+y_2\right) \in B$ and
fong-spivak-seven-sketches F03656	314	■ 1.000	$\left(x_1+x_2, y_1+y_2\right) \in B$	$\left(x_1+x_2, y_1+y_2\right) \in B$	Since B and C are linear, $\left(r x_1, r y_1\right) \in B$ and $\left(r y_1, r z_1\right) \in C$, and also $\left(x_1+x_2, y_1+y_2\right) \in B$ and
fong-spivak-seven-sketches F03657	314	■ 0.996	$\left(y_1+y_2, z_1+z_2\right) \in C$	$\left(y_1+y_2, z_1+z_2\right) \in C$	$\left(y_1+y_2, z_1+z_2\right) \in C$. Hence $\left(r x_1, r z_1\right) \in\left(B \circ C\right)$ and $\left(x_1+x_2, z_1+z_2\right) \in\left(B \circ C\right)$, as desired.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03658	314	0.996	$\left(r_{x_1}, r_{z_1}\right) \in (B ; C)$	$(rx_1, rz_1) \in (B; C)$	$(y_1 + y_2, z_1 + z_2) \in C$. Hence $(rx_1, rz_1) \in (B \S C)$ and $(x_1 + x_2, z_1 + z_2) \in (B \S C)$, as desired.
fong-spivak-seven-sketches F03659	314	0.996	$\left(x_1+x_2, z_1+z_2\right) \in (B ; C)$	$(x_1 + x_2, z_1 + z_2) \in (B; C)$	$(y_1 + y_2, z_1 + z_2) \in C$. Hence $(rx_1, rz_1) \in (B \S C)$ and $(x_1 + x_2, z_1 + z_2) \in (B \S C)$, as desired.
fong-spivak-seven-sketches F03660	314	0.996	$(B ; C) \subseteq R^m \times R^p$	$(B; C) \subseteq R^m \times R^p$	Suppose that $B \subseteq R^m \times R^n$ and $C \subseteq R^n \times R^p$ are linear. Their composite is the relation $(B \S C) \subseteq R^m \times R^p$
fong-spivak-seven-sketches F03661	314	0.999	(x, z)	(x, z)	consisting of all (x, z) for which there exists $y \in R^n$ with $(x, y) \in B$ and $(y, z) \in C$. We want to show
fong-spivak-seven-sketches F03662	314	0.999	$(y, z) \in C$	$(y, z) \in C$	consisting of all (x, z) for which there exists $y \in R^n$ with $(x, y) \in B$ and $(y, z) \in C$. We want to show
fong-spivak-seven-sketches F03663	314	1.000	$(B ; C)$	$(B; C)$	that the set $(B \S C)$ is linear, i.e. closed under scalar multiplication and addition.
fong-spivak-seven-sketches F03664	314	0.976	$(x, z) \in (B ; C)$	$(x, z) \in (B; C)$	For scalar multiplication, take an $(x, z) \in (B \S C)$ and any $r \in R$. Since B is linear, we have $(r * x, r * y) \in B$
fong-spivak-seven-sketches F03665	314	0.976	$(r * x, r * y) \in B$	$(r * x, r * y) \in B$	For scalar multiplication, take an $(x, z) \in (B \S C)$ and any $r \in R$. Since B is linear, we have $(r * x, r * y) \in B$
fong-spivak-seven-sketches F03666	314	0.997	$(r * y, r * z) \in C$	$(r * y, r * z) \in C$	and since C is linear we have $(r * y, r * z) \in C$, so then $(r * x, r * z) \in (B \S C)$. For addition, if we also
fong-spivak-seven-sketches F03667	314	0.997	$(r * x, r * z) \in (B ; C)$	$(r * x, r * z) \in (B; C)$	and since C is linear we have $(r * y, r * z) \in C$, so then $(r * x, r * z) \in (B \S C)$. For addition, if we also
fong-spivak-seven-sketches F03668	314	1.000	$\left(x^{\prime}, z^{\prime}\right) \in (B ; C)$	$(x', z') \in (B; C)$	have $(x', z') \in (B \S C)$ then there is some $y' \in R^n$ with $(x', y') \in B$ and $(y', z') \in C$, so since B and C are
fong-spivak-seven-sketches F03669	314	1.000	$y^{\prime} \in R^n$	$y' \in R^n$	have $(x', z') \in (B \S C)$ then there is some $y' \in R^n$ with $(x', y') \in B$ and $(y', z') \in C$, so since B and C are
fong-spivak-seven-sketches F03670	314	1.000	$\left(y^{\prime}, z^{\prime}\right) \in C$	$(y', z') \in C$	have $(x', z') \in (B \S C)$ then there is some $y' \in R^n$ with $(x', y') \in B$ and $(y', z') \in C$, so since B and C are
fong-spivak-seven-sketches F03671	314	0.998	$\left(x+x^{\prime}, y+y^{\prime}\right) \in B$	$(x + x', y + y') \in B$	linear we have $(x + x', y + y') \in B$ and $(y + y', z + z') \in C$, hence $(x + x', z + z') \in (B \S C)$.
fong-spivak-seven-sketches F03672	314	0.998	$\left(y+y^{\prime}, z+z^{\prime}\right) \in C$	$(y + y', z + z') \in C$	linear we have $(x + x', y + y') \in B$ and $(y + y', z + z') \in C$, hence $(x + x', z + z') \in (B \S C)$.
fong-spivak-seven-sketches F03673	314	0.998	$\left(x+x^{\prime}, z+z^{\prime}\right) \in (B ; C)$	$(x + x', z + z') \in (B; C)$	linear we have $(x + x', y + y') \in B$ and $(y + y', z + z') \in C$, hence $(x + x', z + z') \in (B \S C)$.
fong-spivak-seven-sketches F03674	315	0.361	$\stackrel{\bullet}{a} \bullet \stackrel{\bullet}{b}, \quad \stackrel{\bullet}{a} \rightarrow \stackrel{\bullet}{b}, \quad \stackrel{\bullet}{a} \rightrightarrows \stackrel{\bullet}{b}$ $\stackrel{\bullet}{a} \bullet \stackrel{\bullet}{b} \rightarrow \stackrel{\bullet}{a} \bullet \stackrel{\bullet}{b}, \quad \stackrel{\bullet}{a} \bullet \stackrel{\bullet}{b} \leftrightharpoons \stackrel{\bullet}{a} \bullet \stackrel{\bullet}{b}$	$\begin{array}{ccc} a & & a \\ \bullet \bullet & \bullet \rightarrow \bullet & \bullet \rightrightarrows \bullet \end{array}$	Let $A = \{a, b\}$, and consider the preorders shown here: $\boxed{\bullet \stackrel{a}{\bullet} \bullet}, \boxed{\bullet \stackrel{a}{\bullet} \bullet \rightarrow \bullet \stackrel{a}{\bullet}}, \boxed{\bullet \stackrel{a}{\bullet} \bullet \leftrightharpoons \bullet \stackrel{a}{\bullet} \bullet}$.
fong-spivak-seven-sketches F03675	315	0.998	$a \not \leq b$	$a \not\leq b$	1. The left-most (the discrete preorder on A) has no initial object, because $\overline{a \not\leq b}$ and $b \not\leq a$.
fong-spivak-seven-sketches F03676	315	0.998	$b \not \leq a$	$b \not\leq a$	1. The left-most (the discrete preorder on A) has no initial object, because $\overline{a \not\leq b}$ and $b \not\leq a$.
fong-spivak-seven-sketches F03677	315	1.000	$f\left(1_R\right)=1_S$	$f\left(1_R\right)=1_S$	1. The remaining conditions are that $f(1_R) = 1_S$, and that $f(r_1 * r_2) = f(r_1) * f(r_2)$.
fong-spivak-seven-sketches F03678	315	1.000	$f\left(r_1 *_R r_2\right)=f\left(r_1\right) *_S f\left(r_2\right)$	$f\left(r_1 *_R r_2\right)=f\left(r_1\right) *_S f\left(r_2\right)$	1. The remaining conditions are that $f(1_R) = 1_S$, and that $f(r_1 * r_2) = f(r_1) * f(r_2)$.
fong-spivak-seven-sketches F03679	315	0.966	$R=\left(R, 0_R,+_R, 1_R,{ }^*\right) R$	$R = (R, 0_R, +_R, 1_R, *_R)$	initial means that for any other rig $R = (R, 0_R, +_R, 1_R, *_R)$, there is a unique rig homomorphism
fong-spivak-seven-sketches F03680	315	1.000	$f: \mathbb{N} \rightarrow R$	$f: \mathbb{N} \rightarrow R$	$f: \mathbb{N} \rightarrow R$.
fong-spivak-seven-sketches F03681	315	1.000	$0_R, 1$	$0_R, 1$	What is this homomorphism? Well, to be a rig homomorphism, f must send 0 to 0_R , 1 to 1_R .
fong-spivak-seven-sketches F03682	315	1.000	1_R	1_R	What is this homomorphism? Well, to be a rig homomorphism, f must send 0 to 0_R , 1 to 1_R .
fong-spivak-seven-sketches F03683	315	1.000	$f(m+n)=f(m)+_R f(n)$	$f(m + n) = f(m) +_R f(n)$	Furthermore, we must also have $f(m + n) = f(m) +_R f(n)$, and hence
fong-spivak-seven-sketches F03684	315	0.610	$f(m * n)=f(m) *_R f(n)$	$f(m * n) = f(m) *_R f(n)$	It remains to check $f(m * n) = f(m) *_R f(n)$, and this will follow from distributivity. Noting that

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches-F03685	315	1.000	$f(m * n)$	$f(m * n)$	$f(m * n)$ is equal to the sum of mn copies of 1_R , we have
fong-spivak-seven-sketches-F03686	315	1.000	$m \ n$	mn	$f(m * n)$ is equal to the sum of mn copies of 1_R , we have
fong-spivak-seven-sketches-F03687	315	1.000	$\varnothing \rightarrow c$	$\emptyset \rightarrow c$	unique map $\emptyset \rightarrow c$ coming from the initial object.
fong-spivak-seven-sketches-F03688	315	1.000	$c_{\{1\}} \rightarrow c$	$c_1 \rightarrow c$	If c_1 is initial then by the universal property, for any c there is a unique morphism $c_1 \rightarrow c$; in particular,
fong-spivak-seven-sketches-F03689	315	1.000	$c_{\{2\}} \rightarrow c_{\{1\}}$	$c_2 \rightarrow c_1$	$c_2 \rightarrow c_1$, call it g . But how do we know that f and g are mutually inverse? Well since c_1 is initial
fong-spivak-seven-sketches-F03690	316	0.705	$c_{\{1\}} \rightarrow c_{\{1\}}$	$c_1 \rightarrow c_1$	there is a unique morphism $c_1 \rightarrow c_1$. But we can think of two: id_{c_1} and $f \circ g$. Thus they must be
fong-spivak-seven-sketches-F03691	316	0.705	$\text{id}_{c_{\{1\}}}$	id_{c_1}	there is a unique morphism $c_1 \rightarrow c_1$. But we can think of two: id_{c_1} and $f \circ g$. Thus they must be
fong-spivak-seven-sketches-F03692	316	0.705	$f \ ; \ g$	$f;g$	there is a unique morphism $c_1 \rightarrow c_1$. But we can think of two: id_{c_1} and $f \circ g$. Thus they must be
fong-spivak-seven-sketches-F03693	316	0.735	$f \ ; \ g = \text{id}_{c_{\{1\}}}$	$f;g = \text{id}_{c_1}$	equal. Similarly for c_2 , so we have $f \circ g = \text{id}_{c_1}$ and $g \circ f = \text{id}_{c_2}$, which is the definition of f and g being
fong-spivak-seven-sketches-F03694	316	0.735	$g \ ; \ f = \text{id}_{c_{\{2\}}}$	$g;f = \text{id}_{c_2}$	equal. Similarly for c_2 , so we have $f \circ g = \text{id}_{c_1}$ and $g \circ f = \text{id}_{c_2}$, which is the definition of f and g being
fong-spivak-seven-sketches-F03695	316	1.000	$p+q$	$p + q$	Translating Definition 6.11 to this case, a coproduct $p + q$ is P is an element of P such that $p \leq p + q$ and
fong-spivak-seven-sketches-F03696	316	1.000	$p \leq p+q$	$p \leq p + q$	Translating Definition 6.11 to this case, a coproduct $p + q$ is P is an element of P such that $p \leq p + q$ and
fong-spivak-seven-sketches-F03697	316	1.000	$q \leq p+q$	$q \leq p + q$	$q \leq p + q$, and such that for all elements $x \in P$ with maps $p \leq x$ and $q \leq x$, we have $p + q \leq x$. But this
fong-spivak-seven-sketches-F03698	316	1.000	$p+q \leq x$	$p + q \leq x$	$q \leq p + q$, and such that for all elements $x \in P$ with maps $p \leq x$ and $q \leq x$, we have $p + q \leq x$. But this
fong-spivak-seven-sketches-F03699	316	1.000	$[f, g]: A \sqcup B \rightarrow T$	$[f, g]: A \sqcup B \longrightarrow T$	$[f, g]: A \sqcup B \longrightarrow T$
fong-spivak-seven-sketches-F03700	316	0.729	o	o	$\text{orange}1 \mapsto o$
fong-spivak-seven-sketches-F03701	316	0.768	$\iota_A; [f, g] = f$	$\iota_A; [f, g] = f$	1. The equation $\iota_A \circ [f, g] = f$ is the commutativity of the left hand triangle in the commutative
fong-spivak-seven-sketches-F03702	316	0.586	$\iota_B \circ [f, g] = g$	$\iota_B \circ [f, g] = g$	2. The equation $\iota_B \circ [f, g] = g$ is the commutativity of the right hand triangle in the commutative
fong-spivak-seven-sketches-F03703	316	0.546	$[f, g]: h = [f; h, g; h]$	$[f, g]: h = [f; h, g; h]$	3. The equation $[f, g] \circ h = [f \circ h, g \circ h]$ follows from the universal property of the coproduct. Indeed,
fong-spivak-seven-sketches-F03704	316	1.000	$[f; h, g; h]: A+B \rightarrow D$	$[f; h, g; h]: A + B \rightarrow D$	commutes, and the universal property says there is a unique map $[f \circ h, g \circ h]: A + B \rightarrow D$ for
fong-spivak-seven-sketches-F03705	317	1.000	$+: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$	$+: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$	1. To define a functor $+: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ we must define its action on objects and morphisms. In both
fong-spivak-seven-sketches-F03706	317	0.984	(A, B)	(A, B)	cases, we just take the coproduct. If (A, B) is an object of $\mathcal{C} \times \mathcal{C}$, its image $A + B$ is, as usual, the
fong-spivak-seven-sketches-F03707	317	1.000	$(f, g): (A, B) \rightarrow (C, D)$	$(f, g): (A, B) \rightarrow (C, D)$	coproduct of the two objects of $\mathcal{C}$. If $(f, g): (A, B) \rightarrow (C, D)$ is a morphism, then we can form a
fong-spivak-seven-sketches-F03708	317	0.510	$f+g = [f \text{ right.}]$	$f + g = [f$	morphism $f + g = [f \circ \iota_C, g \circ \iota_D]: A + B \rightarrow C + D$, where $\iota_C: C \rightarrow C + D$ and $\iota_D: D \rightarrow C + D$
fong-spivak-seven-sketches-F03709	317	0.510	ι_C, g	ι_C, g	morphism $f + g = [f \circ \iota_C, g \circ \iota_D]: A + B \rightarrow C + D$, where $\iota_C: C \rightarrow C + D$ and $\iota_D: D \rightarrow C + D$
fong-spivak-seven-sketches-F03710	317	0.510	$[\iota_D \text{ right}]: A+B \rightarrow C+D$	$\iota_D]: A + B \rightarrow C + D$	morphism $f + g = [f \circ \iota_C, g \circ \iota_D]: A + B \rightarrow C + D$, where $\iota_C: C \rightarrow C + D$ and $\iota_D: D \rightarrow C + D$
fong-spivak-seven-sketches-F03711	317	0.510	$\iota_C: C \rightarrow C+D$	$\iota_C: C \rightarrow C + D$	morphism $f + g = [f \circ \iota_C, g \circ \iota_D]: A + B \rightarrow C + D$, where $\iota_C: C \rightarrow C + D$ and $\iota_D: D \rightarrow C + D$
fong-spivak-seven-sketches-F03712	317	0.510	$\iota_D: D \rightarrow C+D$	$\iota_D: D \rightarrow C + D$	morphism $f + g = [f \circ \iota_C, g \circ \iota_D]: A + B \rightarrow C + D$, where $\iota_C: C \rightarrow C + D$ and $\iota_D: D \rightarrow C + D$
fong-spivak-seven-sketches-F03713	317	0.865	$(h, k): (C, D) \rightarrow (E, F)$	$(h, k): (C, D) \rightarrow (E, F)$	a morphism $(h, k): (C, D) \rightarrow (E, F)$ in $\mathcal{C} \times \mathcal{C}$. We need to show that $(f + g) \circ (h + k) = (f \circ h) + (g \circ k)$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03714	317	0.865	$(f+g) ; (h+k) = (f ; h) + (g ; k)$	$(f + g); (h + k) = (f; h) + (g; k)$	a morphism $(h, k): (C, D) \rightarrow (E, F)$ in $\mathcal{C} \times \mathcal{C}$. We need to show that $(f + g) \circ (h + k) = (f \circ h) + (g \circ k)$.
fong-spivak-seven-sketches F03715	317	1.000	$(f \circ h) + (g \circ k) =$	$(f \circ h) + (g \circ k) =$	Indeed, we again use the uniqueness of the copairing in (6.12), this time to show that $(f \circ h) + (g \circ k) =$
fong-spivak-seven-sketches F03716 [conf 0.083]	317	0.083	$\left[f ; h ; \iota_E, g \circ k ; \iota_F \right] = (f+g) \circ (h+k)$ [U+27C28]	(not rendered)	$[f \circ h \circ \iota_E, g \circ k \circ \iota_F] = (f + g) \circ (h + k)$, as required.
fong-spivak-seven-sketches F03717	317	1.000	$!_A: \varnothing \rightarrow A$	$!_A: \emptyset \rightarrow A$	2. Recall the universal property of the initial object gives a unique map $!_A: \emptyset \rightarrow A$. Then the
fong-spivak-seven-sketches F03718	317	0.821	$\left[\text{id}_A, !_A \right]$	$[\text{id}_A, !_A]$	copairing $[\text{id}_A, !_A]$ is a map $A + \emptyset \rightarrow A$. Moreover, it is an isomorphism with inverse $\iota_A: A \rightarrow$
fong-spivak-seven-sketches F03719	317	1.000	$A + \varnothing$	$A + \emptyset$	$A + \emptyset$. Indeed, using the properties in Exercise 6.17 and the universal property of the initial
fong-spivak-seven-sketches F03720	317	0.554	$\iota_A \sim: \sim \left[\text{id}_A, !_A \right] = \text{id}_A$	$\iota_A: [\text{id}_A, !_A] = \text{id}_A$	object, we have $\iota_A \circ [\text{id}_A, !_A] = \text{id}_A$, and
fong-spivak-seven-sketches F03721	317	1.000	$\left[!_A, \text{id}_A \right]: \varnothing + A \rightarrow A$	$[_A, \text{id}_A]: \emptyset + A \rightarrow A$	An analogous argument shows $[_A, \text{id}_A]: \emptyset + A \rightarrow A$ is an isomorphism.
fong-spivak-seven-sketches F03722	317	0.991	$\left[\iota_A, \iota_B + \text{id}_C \right]: A + (B + C) \rightarrow (A + B) + C$	$[\iota_A, \iota_B + \text{id}_C]: A + (B + C) \rightarrow (A + B) + C$	isomorphism, with inverse $[\iota_A, \iota_B + \text{id}_C]: A + (B + C) \rightarrow (A + B) + C$.
fong-spivak-seven-sketches F03723	317	0.999	$\left[\iota_A, \iota_B \right]: A + B \rightarrow B + A$	$[\iota_A, \iota_B]: A + B \rightarrow B + A$	b) The map $[\iota_A, \iota_B]: A + B \rightarrow B + A$ is an isomorphism. Note our notation here is slightly
fong-spivak-seven-sketches F03724	317	0.990	$\iota_A: A \rightarrow A + B$	$\iota_A: A \rightarrow A + B$	confusing: there are two maps named ι_A , (i) $\iota_A: A \rightarrow A + B$, and (ii) $\iota_A: A \rightarrow B + A$, and
fong-spivak-seven-sketches F03725	317	0.990	$\iota_A: A \rightarrow B + A$	$\iota_A: A \rightarrow B + A$	confusing: there are two maps named ι_A , (i) $\iota_A: A \rightarrow A + B$, and (ii) $\iota_A: A \rightarrow B + A$, and
fong-spivak-seven-sketches F03726	317	0.991	$\left[\iota_A, \iota_B \right]: B + A \rightarrow A + B$	$[\iota_A, \iota_B]: B + A \rightarrow A + B$	similarly for ι_B . In the above we mean the map (ii). It has inverse $[\iota_A, \iota_B]: B + A \rightarrow A + B$,
fong-spivak-seven-sketches F03727	317	0.924	Disc_S	Disc_S	has a pushout. The only morphisms in Disc_S are identities, so in particular $A = B = C$, and the
fong-spivak-seven-sketches F03728	317	0.924	$A = B = C$	$A = B = C$	has a pushout. The only morphisms in Disc_S are identities, so in particular $A = B = C$, and the
fong-spivak-seven-sketches F03729	317	1.000	$s^{\prime}$	s'	than one object, because if s' is another object then there is a morphism $s \rightarrow s'$, but the only
fong-spivak-seven-sketches F03730	317	1.000	$s \rightarrow s^{\prime}$	$s \rightarrow s'$	than one object, because if s' is another object then there is a morphism $s \rightarrow s'$, but the only
fong-spivak-seven-sketches F03731	317	1.000	$s = s^{\prime}$	$s = s'$	morphisms in S are identities so $s = s'$. Hence the set S must consist of exactly one element.
fong-spivak-seven-sketches F03732	318	0.995	$\underline{5} \sqcup \underline{3}$	$\underline{5} \sqcup \underline{3}$	equivalence classes in $\underline{5} \sqcup \underline{3}$ generated by the relation $\{f(a) \sim g(a) \mid a \in \underline{4}\}$. If we denote elements of $\underline{5}$
fong-spivak-seven-sketches F03733	318	0.995	$\{f(a) \sim g(a) \mid a \in \underline{4}\}$	$\{f(a) \sim g(a) \mid a \in \underline{4}\}$	equivalence classes in $\underline{5} \sqcup \underline{3}$ generated by the relation $\{f(a) \sim g(a) \mid a \in \underline{4}\}$. If we denote elements of $\underline{5}$
fong-spivak-seven-sketches F03734	318	0.995	$\underline{5}$	$\underline{5}$	equivalence classes in $\underline{5} \sqcup \underline{3}$ generated by the relation $\{f(a) \sim g(a) \mid a \in \underline{4}\}$. If we denote elements of $\underline{5}$
fong-spivak-seven-sketches F03735	318	1.000	$\{1, \dots, 5\}$	$\{1, \dots, 5\}$	as $\underline{1}, \dots, \underline{5}$ and those of $\underline{3}$ as $\{1', 2', 3'\}$, we can redraw the functions f, g :
fong-spivak-seven-sketches F03736	318	1.000	$\underline{3}$	$\underline{3}$	as $\{1, \dots, 5\}$ and those of $\underline{3}$ as $\{1', 2', 3'\}$, we can redraw the functions f, g :
fong-spivak-seven-sketches F03737	318	1.000	$\{1^{\prime}, 2^{\prime}, 3^{\prime}\}$	$\{1', 2', 3'\}$	as $\{1, \dots, 5\}$ and those of $\underline{3}$ as $\underline{1}, \underline{2}, \underline{3}$, we can redraw the functions f, g :
fong-spivak-seven-sketches F03738	318	1.000	$\{1, 1^{\prime}, 3\}, \{2\}, \{4\}$	$\{1, 1', 3\}, \{2\}, \{4\}$	The equivalence classes are $\{1, 1', 3\}, \{2\}, \{4\}$, and $\{5, 2', 3'\}$. These four are exactly the four elements
fong-spivak-seven-sketches F03739	318	1.000	$\{5, 2^{\prime}, 3^{\prime}\}$	$\{5, 2', 3'\}$	The equivalence classes are $\{1, 1', 3\}, \{2\}, \{4\}$, and $\{5, 2', 3'\}$. These four are exactly the four elements
fong-spivak-seven-sketches F03740	318	0.962	$\varnothing \rightarrow X + \Upsilon$	$\emptyset \rightarrow X + \Upsilon$	1. The diagram to the left commutes because $\emptyset$ is initial, and so has a unique map $\emptyset \rightarrow X + Y$. This
fong-spivak-seven-sketches F03741	318	0.947	$f ; \iota_X = g \circ \iota_Y$	$f; \iota_X = g \circ \iota_Y$	implies we must have $f \circ \iota_X = g \circ \iota_Y$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03742	318	■ 0.454	$t: X + \varnothing Y \rightarrow T$	$t: X + \varnothing Y \rightarrow T$	map $t: X +_{\varnothing} Y \rightarrow T$ such that $\iota_X \circ t = x$ and $\iota_Y \circ t = y$. Thus $X +_{\varnothing} Y$ is the coproduct $X + Y$.
fong-spivak-seven-sketches F03743	318	■ 0.454	$\iota_X: X \rightarrow X + Y$	$\iota_X: t = x$	map $t: X +_{\varnothing} Y \rightarrow T$ such that $\iota_X \circ t = x$ and $\iota_Y \circ t = y$. Thus $X +_{\varnothing} Y$ is the coproduct $X + Y$.
fong-spivak-seven-sketches F03744	318	■ 0.454	$\iota_Y: Y \rightarrow X + Y$	$\iota_Y: t = y$	map $t: X +_{\varnothing} Y \rightarrow T$ such that $\iota_X \circ t = x$ and $\iota_Y \circ t = y$. Thus $X +_{\varnothing} Y$ is the coproduct $X + Y$.
fong-spivak-seven-sketches F03745	319	■ 1.000	A, B, X, Y	A, B, X, Y	That is, we need to show that S , together with the maps from A, B, X, Y , and Z , has the required
fong-spivak-seven-sketches F03746	319	■ 1.000	$S \rightarrow T$	$S \rightarrow T$	We need to show there is a unique map $S \rightarrow T$ making everything commute. Since Q is a pushout of
fong-spivak-seven-sketches F03747	319	■ 1.000	$X \leftarrow A \rightarrow Y$	$X \leftarrow A \rightarrow Y$	$X \leftarrow A \rightarrow Y$, there is a unique map $Q \rightarrow T$ making a commutative triangle with Y , and since R is the
fong-spivak-seven-sketches F03748	319	■ 1.000	$Q \rightarrow T$	$Q \rightarrow T$	$X \leftarrow A \rightarrow Y$, there is a unique map $Q \rightarrow T$ making a commutative triangle with Y , and since R is the
fong-spivak-seven-sketches F03749	319	■ 1.000	$Y \leftarrow B \rightarrow Z$	$Y \leftarrow B \rightarrow Z$	pushout of $Y \leftarrow B \rightarrow Z$, there is a unique map $R \rightarrow T$ making a commutative triangle with Y . This
fong-spivak-seven-sketches F03750	319	■ 1.000	$R \rightarrow T$	$R \rightarrow T$	pushout of $Y \leftarrow B \rightarrow Z$, there is a unique map $R \rightarrow T$ making a commutative triangle with Y . This
fong-spivak-seven-sketches F03751	319	■ 0.964	(Y, Q, R, T)	(Y, Q, R, T)	implies that there is a commuting (Y, Q, R, T) square, and hence a unique map $S \rightarrow T$ from its pushout
fong-spivak-seven-sketches F03752	319	■ 0.998	$X + \{ \}_N Y$	$X +_N Y$	The formula in Theorem 6.37 says that the pushout $X +_N Y$ is given by the set of equivalence classes of
fong-spivak-seven-sketches F03753	319	■ 1.000	$X \sqcup N \sqcup Y$	$X \sqcup N \sqcup Y$	$X \sqcup N \sqcup Y$ under the equivalence relation generated by $x \sim n$ if $x = f(n)$, and $y \sim n$ if $y = g(n)$, where
fong-spivak-seven-sketches F03754	319	■ 1.000	$x \sim n$	$x \sim n$	$X \sqcup N \sqcup Y$ under the equivalence relation generated by $x \sim n$ if $x = f(n)$, and $y \sim n$ if $y = g(n)$, where
fong-spivak-seven-sketches F03755	319	■ 1.000	$x = f(n)$	$x = f(n)$	$X \sqcup N \sqcup Y$ under the equivalence relation generated by $x \sim n$ if $x = f(n)$, and $y \sim n$ if $y = g(n)$, where
fong-spivak-seven-sketches F03756	319	■ 1.000	$y \sim n$	$y \sim n$	$X \sqcup N \sqcup Y$ under the equivalence relation generated by $x \sim n$ if $x = f(n)$, and $y \sim n$ if $y = g(n)$, where
fong-spivak-seven-sketches F03757	319	■ 1.000	$y = g(n)$	$y = g(n)$	$X \sqcup N \sqcup Y$ under the equivalence relation generated by $x \sim n$ if $x = f(n)$, and $y \sim n$ if $y = g(n)$, where
fong-spivak-seven-sketches F03758	319	■ 1.000	$x \in X, y \in Y, n \in N$	$x \in X, y \in Y, n \in N$	$x \in X, y \in Y, n \in N$. Since for every $n \in N$ there exists an $x \in X$ such that $x = f(n)$, this set is equal to
fong-spivak-seven-sketches F03759	319	■ 1.000	$n \in N$	$n \in N$	$x \in X, y \in Y, n \in N$. Since for every $n \in N$ there exists an $x \in X$ such that $x = f(n)$, this set is equal to
fong-spivak-seven-sketches F03760	320	■ 1.000	$B \rightarrow D \otimes D$	$B \rightarrow D \otimes D$	morphism $B \rightarrow D \otimes D$.
fong-spivak-seven-sketches F03761	320	■ 1.000	$g: X \rightarrow Y$	$g: X \rightarrow Y$	$f: X \rightarrow Y$ and $g: X \rightarrow Y$ such that
fong-spivak-seven-sketches F03762	320	■ 1.000	$\iota_1: X \rightarrow X + X$	$\iota_1: X \rightarrow X + X$	Write $\iota_1: X \rightarrow X + X$ for the map into the first copy of X in $X + X$, given by the definition of coproduct.
fong-spivak-seven-sketches F03763	320	■ 1.000	$X + X$	$X + X$	Write $\iota_1: X \rightarrow X + X$ for the map into the first copy of X in $X + X$, given by the definition of coproduct.
fong-spivak-seven-sketches F03764	320	■ 0.748	ι_1	ι_1	Then, using the fact that $\iota_1 \circ [\text{id}, \text{id}] = \text{id}$ from Exercise 6.17 1, and the commutativity of the above
fong-spivak-seven-sketches F03765	320	■ 0.748	$[\text{id}, \text{id}] = \text{id}$	$[\text{id}, \text{id}] = \text{id}$	Then, using the fact that $\iota_1 \circ [\text{id}, \text{id}] = \text{id}$ from Exercise 6.17 1, and the commutativity of the above
fong-spivak-seven-sketches F03766	320	■ 0.732	$f = \iota_1; [\text{id}, \text{id}]; f = \iota_1; [\text{id}, \text{id}]; g = g$	$f = \iota_1; [\text{id}, \text{id}]; f = \iota_1; [\text{id}, \text{id}]; g = g$	square, we have $f = \iota_1 \circ [\text{id}, \text{id}] \circ f = \iota_1 \circ [\text{id}, \text{id}] \circ g = g$. This means that $f: X \rightarrow T$ is the unique map
fong-spivak-seven-sketches F03767	320	■ 0.732	$f: X \rightarrow T$	$f: X \rightarrow T$	square, we have $f = \iota_1 \circ [\text{id}, \text{id}] \circ f = \iota_1 \circ [\text{id}, \text{id}] \circ g = g$. This means that $f: X \rightarrow T$ is the unique map
fong-spivak-seven-sketches F03768	321	■ 0.384	$\mathbb{C}$	$\mathbb{C}$	They mean that every category $\mathbf{Cospan}_{\mathbb{C}}$ is equal to a category $\mathbf{Cospan}_F$, for some well-chosen F . They
fong-spivak-seven-sketches F03769	321	■ 0.809	$X \leftarrow N \rightarrow Y$	$X \leftarrow N \rightarrow Y$	$\mathbb{C}$. Next, observe that a morphism in $\mathbf{Cospan}_F$ is a cospan $X \leftarrow N \rightarrow Y$ in $\mathbb{C}$ together with an element
fong-spivak-seven-sketches F03770	321	■ 0.974	$FN = \{*\}$	$FN = \{*\}$	of $FN = \{*\}$. But FN also has a unique element, $*$! So there's no choice here, and we can consider

Identifier	Page	Conf.	LaTeX source	Rendered	Image																								
fong-spivak-seven-sketches-F03771	321	■ 0.974	F N	FN	of $FN = \{*\}$. But FN also has a unique element, $*$! So there's no choice here, and we can consider																								
fong-spivak-seven-sketches-F03772	321	■ 0.758	$\operatorname{Cospan}_{\{F\}=\operatorname{Cospan}_{\{\mathcal{C}\}}}$	$\operatorname{Cospan}_F = \operatorname{Cospan}_{\mathcal{C}}$	simply the usual composition of cospans via pushout, so $\mathbf{Cospan}_F = \mathbf{Cospan}_{\mathcal{C}}$.																								
fong-spivak-seven-sketches-F03773	321	■ 0.484	$\operatorname{Cospan}_{\{\mathcal{C}\}} \rightarrow \operatorname{Cospan}_{\{F\}}$	$\operatorname{Cospan}_{\mathcal{C}} \rightarrow \operatorname{Cospan}_F$	is the identity-on-objects functor $\mathbf{Cospan}_{\mathcal{C}} \rightarrow \mathbf{Cospan}_F$ that simply decorates each cospan with $*$, and																								
fong-spivak-seven-sketches-F03774	321	■ 0.983	$V=\{\mathrm{ul}, \mathrm{ur}, \mathrm{dl}, \mathrm{dr}\}, A=$	$V = \{\mathrm{ul}, \mathrm{ur}, \mathrm{dl}, \mathrm{dr}\}, A =$	We can represent the circuit in Eq. (6.71) by the tuple (V, A, s, t, ℓ) where $V = \{\mathrm{ul}, \mathrm{ur}, \mathrm{dl}, \mathrm{dr}\}$, $A =$																								
fong-spivak-seven-sketches-F03775	321	■ 0.936	$s(-)$	$s(-)$	<table> <tr><td></td><td>r1</td><td>r2</td><td>r3</td><td>c1</td><td>i1</td></tr> <tr><td>$s(-)$</td><td>dl</td><td>ul</td><td>ur</td><td>ul</td><td>dl</td></tr> <tr><td>$t(-)$</td><td>ul</td><td>ur</td><td>dr</td><td>ur</td><td>dr</td></tr> <tr><td>$\ell(-)$</td><td>1Ω</td><td>2Ω</td><td>1Ω</td><td>3F</td><td>1H</td></tr> </table>		r1	r2	r3	c1	i1	$s(-)$	dl	ul	ur	ul	dl	$t(-)$	ul	ur	dr	ur	dr	$\ell(-)$	1Ω	2Ω	1Ω	3F	1H
	r1	r2	r3	c1	i1																								
$s(-)$	dl	ul	ur	ul	dl																								
$t(-)$	ul	ur	dr	ur	dr																								
$\ell(-)$	1Ω	2Ω	1Ω	3F	1H																								
fong-spivak-seven-sketches-F03776	321	■ 0.936	$t(-)$	$t(-)$	<table> <tr><td></td><td>r1</td><td>r2</td><td>r3</td><td>c1</td><td>i1</td></tr> <tr><td>$s(-)$</td><td>dl</td><td>ul</td><td>ur</td><td>ul</td><td>dl</td></tr> <tr><td>$t(-)$</td><td>ul</td><td>ur</td><td>dr</td><td>ur</td><td>dr</td></tr> <tr><td>$\ell(-)$</td><td>1Ω</td><td>2Ω</td><td>1Ω</td><td>3F</td><td>1H</td></tr> </table>		r1	r2	r3	c1	i1	$s(-)$	dl	ul	ur	ul	dl	$t(-)$	ul	ur	dr	ur	dr	$\ell(-)$	1Ω	2Ω	1Ω	3F	1H
	r1	r2	r3	c1	i1																								
$s(-)$	dl	ul	ur	ul	dl																								
$t(-)$	ul	ur	dr	ur	dr																								
$\ell(-)$	1Ω	2Ω	1Ω	3F	1H																								
fong-spivak-seven-sketches-F03777	321	■ 0.936	$\ell(-)$	$\ell(-)$	<table> <tr><td></td><td>r1</td><td>r2</td><td>r3</td><td>c1</td><td>i1</td></tr> <tr><td>$s(-)$</td><td>dl</td><td>ul</td><td>ur</td><td>ul</td><td>dl</td></tr> <tr><td>$t(-)$</td><td>ul</td><td>ur</td><td>dr</td><td>ur</td><td>dr</td></tr> <tr><td>$\ell(-)$</td><td>1Ω</td><td>2Ω</td><td>1Ω</td><td>3F</td><td>1H</td></tr> </table>		r1	r2	r3	c1	i1	$s(-)$	dl	ul	ur	ul	dl	$t(-)$	ul	ur	dr	ur	dr	$\ell(-)$	1Ω	2Ω	1Ω	3F	1H
	r1	r2	r3	c1	i1																								
$s(-)$	dl	ul	ur	ul	dl																								
$t(-)$	ul	ur	dr	ur	dr																								
$\ell(-)$	1Ω	2Ω	1Ω	3F	1H																								
fong-spivak-seven-sketches-F03778	321	■ 0.936	$1 \ \Omega$	1Ω	<table> <tr><td></td><td>r1</td><td>r2</td><td>r3</td><td>c1</td><td>i1</td></tr> <tr><td>$s(-)$</td><td>dl</td><td>ul</td><td>ur</td><td>ul</td><td>dl</td></tr> <tr><td>$t(-)$</td><td>ul</td><td>ur</td><td>dr</td><td>ur</td><td>dr</td></tr> <tr><td>$\ell(-)$</td><td>1Ω</td><td>2Ω</td><td>1Ω</td><td>3F</td><td>1H</td></tr> </table>		r1	r2	r3	c1	i1	$s(-)$	dl	ul	ur	ul	dl	$t(-)$	ul	ur	dr	ur	dr	$\ell(-)$	1Ω	2Ω	1Ω	3F	1H
	r1	r2	r3	c1	i1																								
$s(-)$	dl	ul	ur	ul	dl																								
$t(-)$	ul	ur	dr	ur	dr																								
$\ell(-)$	1Ω	2Ω	1Ω	3F	1H																								
fong-spivak-seven-sketches-F03779	321	■ 0.936	$3 \ F$	$3F$	<table> <tr><td></td><td>r1</td><td>r2</td><td>r3</td><td>c1</td><td>i1</td></tr> <tr><td>$s(-)$</td><td>dl</td><td>ul</td><td>ur</td><td>ul</td><td>dl</td></tr> <tr><td>$t(-)$</td><td>ul</td><td>ur</td><td>dr</td><td>ur</td><td>dr</td></tr> <tr><td>$\ell(-)$</td><td>1Ω</td><td>2Ω</td><td>1Ω</td><td>3F</td><td>1H</td></tr> </table>		r1	r2	r3	c1	i1	$s(-)$	dl	ul	ur	ul	dl	$t(-)$	ul	ur	dr	ur	dr	$\ell(-)$	1Ω	2Ω	1Ω	3F	1H
	r1	r2	r3	c1	i1																								
$s(-)$	dl	ul	ur	ul	dl																								
$t(-)$	ul	ur	dr	ur	dr																								
$\ell(-)$	1Ω	2Ω	1Ω	3F	1H																								
fong-spivak-seven-sketches-F03780	321	■ 0.936	$1 \ H$	$1H$	<table> <tr><td></td><td>r1</td><td>r2</td><td>r3</td><td>c1</td><td>i1</td></tr> <tr><td>$s(-)$</td><td>dl</td><td>ul</td><td>ur</td><td>ul</td><td>dl</td></tr> <tr><td>$t(-)$</td><td>ul</td><td>ur</td><td>dr</td><td>ur</td><td>dr</td></tr> <tr><td>$\ell(-)$</td><td>1Ω</td><td>2Ω</td><td>1Ω</td><td>3F</td><td>1H</td></tr> </table>		r1	r2	r3	c1	i1	$s(-)$	dl	ul	ur	ul	dl	$t(-)$	ul	ur	dr	ur	dr	$\ell(-)$	1Ω	2Ω	1Ω	3F	1H
	r1	r2	r3	c1	i1																								
$s(-)$	dl	ul	ur	ul	dl																								
$t(-)$	ul	ur	dr	ur	dr																								
$\ell(-)$	1Ω	2Ω	1Ω	3F	1H																								
fong-spivak-seven-sketches-F03781	321	■ 1.000	$\psi_{\underline{2}, \underline{2}}(b, s)$	$\psi_{\underline{2}, \underline{2}}(b, s)$	The circuit $\psi_{\underline{2}, \underline{2}}(b, s)$ is the disjoint union of the two labelled graphs b and s :																								
fong-spivak-seven-sketches-F03782	321	■ 0.220	$1 \xrightarrow{f} \underline{2} \stackrel{g}{\leftarrow} \underline{1}$	$1 \xrightarrow{f} \underline{2} \stackrel{g}{\leftarrow} \underline{1}$	The cospan is the cospan $\underline{1} \xrightarrow{f} \underline{2} \stackrel{g}{\leftarrow} \underline{1}$, where $f(1) = 1$ and $g(1) = 2$. The decoration is the C-circuit																								
fong-spivak-seven-sketches-F03783	321	■ 0.220	$f(1)=1$	$f(1) = 1$	The cospan is the cospan $\underline{1} \xrightarrow{f} \underline{2} \stackrel{g}{\leftarrow} \underline{1}$, where $f(1) = 1$ and $g(1) = 2$. The decoration is the C-circuit																								
fong-spivak-seven-sketches-F03784	321	■ 0.220	$g(1)=2$	$g(1) = 2$	The cospan is the cospan $\underline{1} \xrightarrow{f} \underline{2} \stackrel{g}{\leftarrow} \underline{1}$, where $f(1) = 1$ and $g(1) = 2$. The decoration is the C-circuit																								
fong-spivak-seven-sketches-F03785	321	■ 0.954	$(\underline{2}, \{a\}, s, t, \ell)$	$(\underline{2}, \{a\}, s, t, \ell)$	$(\underline{2}, \{a\}, s, t, \ell)$, where $s(a) = 1, t(a) = 2$ and $\ell(a) = \text{battery}$.																								
fong-spivak-seven-sketches-F03786	321	■ 0.954	$s(a)=1, t(a)=2$	$s(a) = 1, t(a) = 2$	$(\underline{2}, \{a\}, s, t, \ell)$, where $s(a) = 1, t(a) = 2$ and $\ell(a) = \text{battery}$.																								
fong-spivak-seven-sketches-F03787	321	■ 0.954	$\ell(a)=$	$\ell(a) =$	$(\underline{2}, \{a\}, s, t, \ell)$, where $s(a) = 1, t(a) = 2$ and $\ell(a) = \text{battery}$.																								
fong-spivak-seven-sketches-F03788	322	■ 1.000	$C:=(V, A, s, t, \ell)$	$C := (V, A, s, t, \ell)$	Recall the circuit $C := (V, A, s, t, \ell)$ from the solution to Exercise 6.79. Then the first decorated cospan																								

Identifier	Page	Conf.	LaTeX source	Rendered	Image																																
fong-spivak-seven-sketches F03789	322	0.462	$\underline{1} \xrightarrow{\text{f}} V \stackrel{\text{g}}{\leftarrow} \underline{2}, f(1)=\mathrm{ul}, g(1)=\mathrm{ur}$	$\underline{1} \xrightarrow{f} V \xleftarrow{g} \underline{2}, f(1) = \text{ul}, g(1) = \text{ur}$	is given by the cospan $\underline{1} \xrightarrow{f} V \xleftarrow{g} \underline{2}, f(1) = \text{ul}, g(1) = \text{ur}$, and $g(2) = \text{ur}$, decorated by circuit C. The																																
fong-spivak-seven-sketches F03790	322	0.462	$g(2)=$	$g(2) =$	is given by the cospan $\underline{1} \xrightarrow{f} V \xleftarrow{g} \underline{2}, f(1) = \text{ul}, g(1) = \text{ur}$, and $g(2) = \text{ur}$, decorated by circuit C. The																																
fong-spivak-seven-sketches F03791	322	0.953	$\underline{2} \xrightarrow{\text{f}^{\prime}} V^{\prime} \stackrel{\text{g}^{\prime}}{\leftarrow} \underline{2}$	$\underline{2} \xrightarrow{f'} V' \xleftarrow{g'} \underline{2}$	second decorated cospan is given by the cospan $\underline{2} \xrightarrow{f'} V' \xleftarrow{g'} \underline{2}$ and the circuit $C' := (V', A', s', t', \ell')$,																																
fong-spivak-seven-sketches F03792	322	0.953	$C^{\prime} := \left(V^{\prime}, A^{\prime}, s^{\prime}, t^{\prime}, \ell^{\prime} \right)$	$C' := (V', A', s', t', \ell')$	second decorated cospan is given by the cospan $\underline{2} \xrightarrow{f'} V' \xleftarrow{g'} \underline{2}$ and the circuit $C' := (V', A', s', t', \ell')$,																																
fong-spivak-seven-sketches F03793	322	1.000	$V^{\prime} = \{l, r, d\}, A^{\prime} = \{\mathrm{r}^1, \mathrm{r}^2\}$	$V' = \{l, r, d\}, A' = \{\text{r1}', \text{r2}'\}$	where $V' = \{l, r, d\}, A' = \{\text{r1}', \text{r2}'\}$, and the functions are given by the tables																																
fong-spivak-seven-sketches F03794	322	0.975	$\text{f}^{\prime}(-)$	$f'(-)$	<table><tr><td></td><td>1</td><td>2</td></tr><tr><td>$f'(-)$</td><td>1</td><td>d</td></tr><tr><td>$g'(-)$</td><td>r</td><td>r</td></tr></table>		1	2	$f'(-)$	1	d	$g'(-)$	r	r																							
	1	2																																			
$f'(-)$	1	d																																			
$g'(-)$	r	r																																			
fong-spivak-seven-sketches F03795	322	0.975	$\text{g}^{\prime}(-)$	$g'(-)$	<table><tr><td></td><td>1</td><td>2</td></tr><tr><td>$f'(-)$</td><td>1</td><td>d</td></tr><tr><td>$g'(-)$</td><td>r</td><td>r</td></tr></table>		1	2	$f'(-)$	1	d	$g'(-)$	r	r																							
	1	2																																			
$f'(-)$	1	d																																			
$g'(-)$	r	r																																			
fong-spivak-seven-sketches F03796	322	0.786	$5 \ \Omega$	5Ω	<table><tr><td></td><td>r1'</td><td>r2'</td></tr><tr><td>$s(-)$</td><td>1</td><td>r</td></tr><tr><td>$t(-)$</td><td>r</td><td>d</td></tr><tr><td>$\ell(-)$</td><td>5Ω</td><td>8Ω</td></tr></table>		r1'	r2'	$s(-)$	1	r	$t(-)$	r	d	$\ell(-)$	5Ω	8Ω																				
	r1'	r2'																																			
$s(-)$	1	r																																			
$t(-)$	r	d																																			
$\ell(-)$	5Ω	8Ω																																			
fong-spivak-seven-sketches F03797	322	0.786	$8 \ \Omega$	8Ω	<table><tr><td></td><td>r1'</td><td>r2'</td></tr><tr><td>$s(-)$</td><td>1</td><td>r</td></tr><tr><td>$t(-)$</td><td>r</td><td>d</td></tr><tr><td>$\ell(-)$</td><td>5Ω</td><td>8Ω</td></tr></table>		r1'	r2'	$s(-)$	1	r	$t(-)$	r	d	$\ell(-)$	5Ω	8Ω																				
	r1'	r2'																																			
$s(-)$	1	r																																			
$t(-)$	r	d																																			
$\ell(-)$	5Ω	8Ω																																			
fong-spivak-seven-sketches F03798	322	0.971	$V \stackrel{\text{g}}{\leftarrow} \underline{2} \xrightarrow{\text{f}^{\prime}} V^{\prime}$	$V \xleftarrow{g} \underline{2} \xrightarrow{f'} V'$	To compose these, we first take the pushout of $V \xleftarrow{g} \underline{2} \xrightarrow{f'} V'$. This gives the a new apex $V'' =$																																
fong-spivak-seven-sketches F03799	322	0.971	$V^{\prime \prime} =$	$V'' =$	To compose these, we first take the pushout of $V \xleftarrow{g} \underline{2} \xrightarrow{f'} V'$. This gives the a new apex $V'' =$																																
fong-spivak-seven-sketches F03800	322	0.823	$\underline{1} \xrightarrow{\text{h}} V^{\prime \prime} \stackrel{\text{k}}{\leftarrow} \underline{2}$	$\underline{1} \xrightarrow{h} V'' \xleftarrow{k} \underline{2}$	$\{\text{ul}, \text{dl}, \text{dr}, \text{m}, \text{r}\}$ with five elements, and composite cospan $\underline{1} \xrightarrow{h} V'' \xleftarrow{k} \underline{2}$ given by $h(1) = \text{ul}, k(1) = \text{r}$																																
fong-spivak-seven-sketches F03801	322	0.823	$h(1)=\mathrm{ul}, k(1)=\mathrm{r}$	$h(1) = \text{ul}, k(1) = \text{r}$	$\{\text{ul}, \text{dl}, \text{dr}, \text{m}, \text{r}\}$ with five elements, and composite cospan $\underline{1} \xrightarrow{h} V'' \xleftarrow{k} \underline{2}$ given by $h(1) = \text{ul}, k(1) = \text{r}$																																
fong-spivak-seven-sketches F03802	322	0.536	$k(2)=\mathrm{m}$	$k(2) = \text{m}$	and $k(2) = \text{m}$. The new circuit is given by $(V,{}'' A + A', s,{}'' t,{}'' \ell'')$ where the functions are given by																																
fong-spivak-seven-sketches F03803	322	0.536	$\left(V^{\prime \prime} A + A^{\prime}, s,{}^{\prime \prime} t,{}^{\prime \prime} \ell^{\prime \prime} \right)$	$(V'' A + A', s,{}'' t,{}'' \ell'')$	and $k(2) = \text{m}$. The new circuit is given by $(V,{}'' A + A', s,{}'' t,{}'' \ell'')$ where the functions are given by																																
fong-spivak-seven-sketches F03804	322	1.000	$s^{\prime \prime}(-)$	$s''(-)$	<table><tr><td></td><td>r1</td><td>r2</td><td>r3</td><td>c1</td><td>i1</td><td>r1'</td><td>r2'</td></tr><tr><td>$s''(-)$</td><td>dl</td><td>ul</td><td>m</td><td>ul</td><td>dl</td><td>m</td><td>r</td></tr><tr><td>$t''(-)$</td><td>ul</td><td>m</td><td>dr</td><td>m</td><td>dr</td><td>r</td><td>m</td></tr><tr><td>$\ell''(-)$</td><td>1Ω</td><td>2Ω</td><td>1Ω</td><td>3F</td><td>1H</td><td>5Ω</td><td>8Ω</td></tr></table>		r1	r2	r3	c1	i1	r1'	r2'	$s''(-)$	dl	ul	m	ul	dl	m	r	$t''(-)$	ul	m	dr	m	dr	r	m	$\ell''(-)$	1Ω	2Ω	1Ω	3F	1H	5Ω	8Ω
	r1	r2	r3	c1	i1	r1'	r2'																														
$s''(-)$	dl	ul	m	ul	dl	m	r																														
$t''(-)$	ul	m	dr	m	dr	r	m																														
$\ell''(-)$	1Ω	2Ω	1Ω	3F	1H	5Ω	8Ω																														
fong-spivak-seven-sketches F03805	322	1.000	$t^{\prime \prime}(-)$	$t''(-)$	<table><tr><td></td><td>r1</td><td>r2</td><td>r3</td><td>c1</td><td>i1</td><td>r1'</td><td>r2'</td></tr><tr><td>$s''(-)$</td><td>dl</td><td>ul</td><td>m</td><td>ul</td><td>dl</td><td>m</td><td>r</td></tr><tr><td>$t''(-)$</td><td>ul</td><td>m</td><td>dr</td><td>m</td><td>dr</td><td>r</td><td>m</td></tr><tr><td>$\ell''(-)$</td><td>1Ω</td><td>2Ω</td><td>1Ω</td><td>3F</td><td>1H</td><td>5Ω</td><td>8Ω</td></tr></table>		r1	r2	r3	c1	i1	r1'	r2'	$s''(-)$	dl	ul	m	ul	dl	m	r	$t''(-)$	ul	m	dr	m	dr	r	m	$\ell''(-)$	1Ω	2Ω	1Ω	3F	1H	5Ω	8Ω
	r1	r2	r3	c1	i1	r1'	r2'																														
$s''(-)$	dl	ul	m	ul	dl	m	r																														
$t''(-)$	ul	m	dr	m	dr	r	m																														
$\ell''(-)$	1Ω	2Ω	1Ω	3F	1H	5Ω	8Ω																														

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03806	322	1.000	$\ell^{\prime}(-)$	$\ell''(-)$	
fong-spivak-seven-sketches F03807	322	1.000	η	η	Composing η and x we have
fong-spivak-seven-sketches F03808	324	0.846	(X, p, q)	(X, p, q)	Suppose first that (A, B, A', B') is a pullback, and take any (X, p, q) as in the following diagram:
fong-spivak-seven-sketches F03809	324	0.444	$q: f^{\prime}; g^{\prime}=p; h_3$	$q: f'; g' = p; h_3$	where $q \circ f' \circ g' = p \circ h_3$. Then by the universal property of the (B, C, B', C') pullback, we get a unique
fong-spivak-seven-sketches F03810	324	0.180	$r; h_2=g$	$r; h_2 = g$	In other words $r \circ h_2 = g \circ f'$ and $r \circ g = p$. Then by the universal property of the (A, B, A', B') pullback,
fong-spivak-seven-sketches F03811	324	0.180	$r; g=p$	$r; g = p$	In other words $r \circ h_2 = g \circ f'$ and $r \circ g = p$. Then by the universal property of the (A, B, A', B') pullback,
fong-spivak-seven-sketches F03812	324	0.991	$r^{\prime}: X \rightarrow A$	$r': X \rightarrow A$	we get a unique dotted arrow $r': X \rightarrow A$ making the right-hand diagram commute, i.e. $r' \circ f = r$ and
fong-spivak-seven-sketches F03813	324	0.991	$r^{\prime}; f=r$	$r'; f = r$	we get a unique dotted arrow $r': X \rightarrow A$ making the right-hand diagram commute, i.e. $r' \circ f = r$ and
fong-spivak-seven-sketches F03814	324	1.000	$r^{\prime}; h_1=q$	$r'; h_1 = q$	$r' \circ h_1 = q$. This gives the existence of an r with the required property, $r' \circ f = r$ and $r' \circ f \circ g = r \circ g = p$.
fong-spivak-seven-sketches F03815	324	1.000	$r^{\prime}; f; g=r; g=p$	$r'; f; g = r; g = p$	$r' \circ h_1 = q$. This gives the existence of an r with the required property, $r' \circ f = r$ and $r' \circ f \circ g = r \circ g = p$.
fong-spivak-seven-sketches F03816	324	0.626	$r_{\{0\}}$	r_0	To see uniqueness, suppose given another morphisms r_0 such that $r_0 \circ f \circ g = p$ and $r_0 \circ h_1 = q$:
fong-spivak-seven-sketches F03817	324	0.626	$f; g=p$	$f; g = p$	To see uniqueness, suppose given another morphisms r_0 such that $r_0 \circ f \circ g = p$ and $r_0 \circ h_1 = q$:
fong-spivak-seven-sketches F03818	324	0.626	$r_{\{0\}}; h_1=q$	$r_0; h_1 = q$	To see uniqueness, suppose given another morphisms r_0 such that $r_0 \circ f \circ g = p$ and $r_0 \circ h_1 = q$:
fong-spivak-seven-sketches F03819	324	0.741	$r_{\{0\}}: f=r$	$r_0: f = r$	Then by the uniqueness of r , we must have $r_0 \circ f = r$, and then by the uniqueness of r' , we must have
fong-spivak-seven-sketches F03820	324	0.741	$r^{\prime}$	r'	Then by the uniqueness of r , we must have $r_0 \circ f = r$, and then by the uniqueness of r' , we must have
fong-spivak-seven-sketches F03821	324	1.000	$r_{\{0\}}=r^{\prime}$	$r_0 = r'$	$r_0 = r'$. This proves the first result.
fong-spivak-seven-sketches F03822	324	0.994	$r; h_2=q; f^{\prime}$	$r; h_2 = q; f'$	i.e. where $r \circ h_2 = q \circ f'$. Then letting $p := r \circ g$, we have
fong-spivak-seven-sketches F03823	324	0.994	$p:=r; g$	$p := r; g$	i.e. where $r \circ h_2 = q \circ f'$. Then letting $p := r \circ g$, we have
fong-spivak-seven-sketches F03824	325	0.999	$r^{\prime}; f; g=p$	$r'; f; g = p$	that $r' \circ f \circ g = p$ and $r_0 \circ h_1 = q$, as shown:
fong-spivak-seven-sketches F03825	325	0.422	$r_{\{0\}}:=r^{\prime}; f$	$r_0 := r'; f$	But now let $r_0 := r' \circ f$. It satisfies $r_0 \circ g = p$ and $r_0 \circ h_2 = q \circ f'$, and r satisfies the same equations:
fong-spivak-seven-sketches F03826	325	0.422	$r_{\{0\}}; g=p$	$r_0; g = p$	But now let $r_0 := r' \circ f$. It satisfies $r_0 \circ g = p$ and $r_0 \circ h_2 = q \circ f'$, and r satisfies the same equations:
fong-spivak-seven-sketches F03827	325	0.422	$r_{\{0\}}; h_2=q; f^{\prime}$	$r_0; h_2 = q; f'$	But now let $r_0 := r' \circ f$. It satisfies $r_0 \circ g = p$ and $r_0 \circ h_2 = q \circ f'$, and r satisfies the same equations:
fong-spivak-seven-sketches F03828	325	0.554	$A, B, A^{\prime}, B^{\prime}$	A, B, A', B'	follows that r' is a pullback of the (A, B, A', B') square, as desired.
fong-spivak-seven-sketches F03829	325	1.000	$a_{\{1\}}, a_{\{2\}} \in A$	$a_1, a_2 \in A$	A function $f: A \rightarrow B$ is injective iff for all $a_1, a_2 \in A$, if $f(a_1) = f(a_2)$ then $a_1 = a_2$. It is a monomorphism
fong-spivak-seven-sketches F03830	325	1.000	$f\left(a_{\{1\}}\right)=f\left(a_{\{2\}}\right)$	$f(a_1) = f(a_2)$	A function $f: A \rightarrow B$ is injective iff for all $a_1, a_2 \in A$, if $f(a_1) = f(a_2)$ then $a_1 = a_2$. It is a monomorphism
fong-spivak-seven-sketches F03831	325	1.000	$a_{\{1\}}=a_{\{2\}}$	$a_1 = a_2$	A function $f: A \rightarrow B$ is injective iff for all $a_1, a_2 \in A$, if $f(a_1) = f(a_2)$ then $a_1 = a_2$. It is a monomorphism
fong-spivak-seven-sketches F03832	325	0.402	$g_{\{1\}}, g_{\{2\}}: X \rightarrow A$	$g_1, g_2: X \rightarrow A$	iff for all sets X and functions $g_1, g_2: X \rightarrow A$, if $g_1 \circ f = g_2 \circ f$ then $g_1 = g_2$. Indeed, this comes directly

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03833	325	■ 0.402	$f=g_{\{2\}}$	$f = g_2$	iff for all sets X and functions $g_1, g_2: X \rightarrow A$, if $g_1 \circ f = g_2 \circ f$ then $g_1 = g_2$. Indeed, this comes directly
fong-spivak-seven-sketches F03834	325	■ 0.402	$g_{\{1\}}=g_{\{2\}}$	$g_1 = g_2$	iff for all sets X and functions $g_1, g_2: X \rightarrow A$, if $g_1 \circ f = g_2 \circ f$ then $g_1 = g_2$. Indeed, this comes directly
fong-spivak-seven-sketches F03835	325	■ 1.000	$X=\{*\}$	$X = \{*\}$	$X = \{*\}$ be a one element set, and let $g_1, g_2: X \rightarrow A$ be given by $g_1(*) := a_1$ and $g_2(*) := a_2$. Then
fong-spivak-seven-sketches F03836	325	■ 1.000	$g_{\{1\}}(*) := a_{\{1\}}$	$g_1(*) := a_1$	$X = \{*\}$ be a one element set, and let $g_1, g_2: X \rightarrow A$ be given by $g_1(*) := a_1$ and $g_2(*) := a_2$. Then
fong-spivak-seven-sketches F03837	325	■ 1.000	$g_{\{2\}}(*) := a_{\{2\}}$	$g_2(*) := a_2$	$X = \{*\}$ be a one element set, and let $g_1, g_2: X \rightarrow A$ be given by $g_1(*) := a_1$ and $g_2(*) := a_2$. Then
fong-spivak-seven-sketches F03838	325	■ 0.626	$=g_{\{2\}}$	$= g_2$	$g_1 \circ f = g_2 \circ f$, so $g_1 = g_2$, so $a_1 = a_2$.
fong-spivak-seven-sketches F03839	325	■ 1.000	$g_{\{1\}}(x)=g_{\{2\}}(x)$	$g_1(x) = g_2(x)$	We will have $g_1 = g_2$ if we can show that $g_1(x) = g_2(x)$ for every $x \in X$. So take any $x \in X$; since
fong-spivak-seven-sketches F03840	325	■ 1.000	$f\left(g_{\{1\}}(x)\right)=f\left(g_{\{2\}}(x)\right)$	$f(g_1(x)) = f(g_2(x))$	$f(g_1(x)) = f(g_2(x))$ and f is injective, we have $g_1(x) = g_2(x)$ as desired.
fong-spivak-seven-sketches F03841	325	■ 0.974	$j:=i^{-1}$	$j := i^{-1}$	Let $j := i^{-1}$ be the inverse of i , and consider $g := (f \circ j): A \rightarrow B'$. Then $g \circ i = f$, so by the
fong-spivak-seven-sketches F03842	325	■ 0.974	$g:=(f \circ j): A \rightarrowtail B^{\prime}$	$g := (f \circ j): A \rightarrow B'$	Let $j := i^{-1}$ be the inverse of i , and consider $g := (f \circ j): A \rightarrow B'$. Then $g \circ i = f$, so by the
fong-spivak-seven-sketches F03843	325	■ 0.974	$g \circ i=f$	$g \circ i = f$	Let $j := i^{-1}$ be the inverse of i , and consider $g := (f \circ j): A \rightarrow B'$. Then $g \circ i = f$, so by the
fong-spivak-seven-sketches F03844	325	■ 0.553	$j^{\prime}: A \rightarrowtail A^{\prime}$	$j': A \rightarrow A'$	existence part of the universal property, there is a map $j': A \rightarrow A'$ such that $j' \circ i' = \text{id}_A$ and
fong-spivak-seven-sketches F03845	325	■ 0.553	$j^{\prime} \ ; \ i^{\prime}=\operatorname{id}_{A}$	$j'; i' = \text{id}_A$	existence part of the universal property, there is a map $j': A \rightarrow A'$ such that $j' \circ i' = \text{id}_A$ and
fong-spivak-seven-sketches F03846	325	■ 0.392	$j^{\prime} \ ; \ f^{\prime}=f \circ j$	$j'; f' = f \circ j$	$j' \circ f' = f \circ j$. We will be done if we can show $i' \circ j' = \text{id}_{A'}$. One checks that $(i' \circ j') \circ i' = i'$ and
fong-spivak-seven-sketches F03847	325	■ 0.392	$i^{\prime} \ ; \ j^{\prime}=\operatorname{id}_{A^{\prime}}$	$i'; j' = \text{id}_{A'}$	$j' \circ f' = f \circ j$. We will be done if we can show $i' \circ j' = \text{id}_{A'}$. One checks that $(i' \circ j') \circ i' = i'$ and
fong-spivak-seven-sketches F03848	325	■ 0.392	$\left(i^{\prime} \ ; \ j^{\prime}\right) \ ; \ i^{\prime}=i^{\prime}$	$(i'; j'); i' = i'$	$j' \circ f' = f \circ j$. We will be done if we can show $i' \circ j' = \text{id}_{A'}$. One checks that $(i' \circ j') \circ i' = i'$ and
fong-spivak-seven-sketches F03849 [conf 0.024]	325	■ 0.024	$\left(i^{\prime} \ ; \ j^{\prime}\right) \ ; \ j^{\prime}=i^{\prime} \ ; \ j^{\prime}=f^{\prime} \ ; \ i, j=f^{\prime}$	$(i' \circ j'), f' = i'; j = f'; i, j = f'$	that $(i' \circ j') \circ f' = i' \circ f \circ j = f' \circ i \circ j = f'$. But $\text{id}_{A'}$ also satisfies those properties: $\text{id}_{A'} \circ i' = i'$ and
fong-spivak-seven-sketches F03850 [conf 0.024]	325	■ 0.024	$\operatorname{id}_{A^{\prime}}$	$\text{id}_{A'}$	that $(i' \circ j') \circ f' = i' \circ f \circ j = f' \circ i \circ j = f'$. But $\text{id}_{A'}$ also satisfies those properties: $\text{id}_{A'} \circ i' = i'$ and
fong-spivak-seven-sketches F03851 [conf 0.024]	325	■ 0.024	$\operatorname{id}_{A^{\prime}} \circ i^{\prime}=i^{\prime}$	$\text{id}_{A'} \circ i' = i'$	that $(i' \circ j') \circ f' = i' \circ f \circ j = f' \circ i \circ j = f'$. But $\text{id}_{A'}$ also satisfies those properties: $\text{id}_{A'} \circ i' = i'$ and
fong-spivak-seven-sketches F03852	325	■ 0.511	$f^{\prime}=f^{\prime}$	$f' = f'$	$\text{id}_{A'} \circ f' = f'$, so by the uniqueness part of the universal property, $(i' \circ j') = \text{id}_{A'}$.
fong-spivak-seven-sketches F03853	325	■ 0.511	$\left(i^{\prime} \ ; \ j^{\prime}\right)=\operatorname{id}_{A^{\prime}}$	$(i'; j') = \text{id}_{A'}$	$\text{id}_{A'} \circ f' = f'$, so by the uniqueness part of the universal property, $(i' \circ j') = \text{id}_{A'}$.
fong-spivak-seven-sketches F03854	325	■ 0.549	$g: X \rightarrowtail A$	$g: X \rightarrow A$	So take any object X and morphisms $g: X \rightarrow A$ and $h: X \rightarrow B$ such that $g \circ f = h \circ \text{id}_B$. We need
fong-spivak-seven-sketches F03855	325	■ 0.549	$h: X \rightarrowtail B$	$h: X \rightarrow B$	So take any object X and morphisms $g: X \rightarrow A$ and $h: X \rightarrow B$ such that $g \circ f = h \circ \text{id}_B$. We need
fong-spivak-seven-sketches F03856	325	■ 0.549	$g \circ f=h \circ \operatorname{id}_B$	$g \circ f = h \circ \text{id}_B$	So take any object X and morphisms $g: X \rightarrow A$ and $h: X \rightarrow B$ such that $g \circ f = h \circ \text{id}_B$. We need
fong-spivak-seven-sketches F03857	325	■ 0.554	$r: X \rightarrowtail A$	$r: X \rightarrow A$	to show there is a unique morphism $r: X \rightarrow A$ such that $r \circ \text{id}_A = g$ and $r \circ f = h$. That's easy:
fong-spivak-seven-sketches F03858	325	■ 0.554	$r \ ; \ \operatorname{id}_A=g$	$r; \text{id}_A = g$	to show there is a unique morphism $r: X \rightarrow A$ such that $r \circ \text{id}_A = g$ and $r \circ f = h$. That's easy:
fong-spivak-seven-sketches F03859	325	■ 0.554	$r \ ; \ f=h$	$r; f = h$	to show there is a unique morphism $r: X \rightarrow A$ such that $r \circ \text{id}_A = g$ and $r \circ f = h$. That's easy:
fong-spivak-seven-sketches F03860	325	■ 1.000	$r=g$	$r = g$	the first requirement forces $r = g$ and the second requirement is then fulfilled.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03861	326	0.666	$v \otimes w$	$v \otimes w$	$v \otimes w$ serves as a meet operation, i.e. it satisfies the same universal property as $\wedge$, namely
fong-spivak-seven-sketches F03862	326	0.988	$v \leq (w \multimap x)$	$v \leq (w \multimap x)$	namely $v \leq (w \multimap x)$ iff $v \otimes W \leq x$. So $\mathcal{V}$ is a cartesian closed category, and of course it is a
fong-spivak-seven-sketches F03863	326	0.988	$v \otimes W \leq x$	$v \otimes W \leq x$	namely $v \leq (w \multimap x)$ iff $v \otimes W \leq x$. So $\mathcal{V}$ is a cartesian closed category, and of course it is a
fong-spivak-seven-sketches F03864	326	0.998	$\mathbb{N}^{\mathrm{op}} \times \mathbb{N}^{\mathrm{op}}$	$\mathbb{N}^{\mathrm{op}} \times \mathbb{N}^{\mathrm{op}}$	This is the product preorder $\mathbb{N}^{\mathrm{op}} \times \mathbb{N}^{\mathrm{op}}$: its objects are pairs $(a, b) \in \mathbb{N} \times \mathbb{N}$ with $(a, b) \leq (a', b')$
fong-spivak-seven-sketches F03865	326	0.998	$(a, b) \in \mathbb{N} \times \mathbb{N}$	$(a, b) \in \mathbb{N} \times \mathbb{N}$	This is the product preorder $\mathbb{N}^{\mathrm{op}} \times \mathbb{N}^{\mathrm{op}}$: its objects are pairs $(a, b) \in \mathbb{N} \times \mathbb{N}$ with $(a, b) \leq (a', b')$
fong-spivak-seven-sketches F03866	326	0.998	$(a, b) \leq (a', b')$	$(a, b) \leq (a', b')$	This is the product preorder $\mathbb{N}^{\mathrm{op}} \times \mathbb{N}^{\mathrm{op}}$: its objects are pairs $(a, b) \in \mathbb{N} \times \mathbb{N}$ with $(a, b) \leq (a', b')$
fong-spivak-seven-sketches F03867	326	1.000	$a' \leq a$	$a' \leq a$	iff, in the usual ordering on $\mathbb{N}$ we have $a' \leq a$ and $b' \leq b$. But you can just look at the diagram.
fong-spivak-seven-sketches F03868	326	1.000	$b' \leq b$	$b' \leq b$	iff, in the usual ordering on $\mathbb{N}$ we have $a' \leq a$ and $b' \leq b$. But you can just look at the diagram.
fong-spivak-seven-sketches F03869	326	0.658	$(0, 0)$	$(0, 0)$	It has a top element, $(0, 0)$, and it has binary meets, $(a, b) \wedge (a', b') = (\max(a, a'), \max(b, b'))$. But
fong-spivak-seven-sketches F03870	326	0.658	$(a, b) \wedge (a', b') = (\max(a, a'), \max(b, b'))$	$(a, b) \wedge (a', b') = (\max(a, a'), \max(b, b'))$	It has a top element, $(0, 0)$, and it has binary meets, $(a, b) \wedge (a', b') = (\max(a, a'), \max(b, b'))$. But
fong-spivak-seven-sketches F03871	327	0.958	$x \multimap y$	$x \multimap y$	each x, y , the hom-object $x \multimap y$ exists. The formula for it is $x \multimap y = \bigvee \{w \mid w \wedge x \leq y\}$, i.e. we
fong-spivak-seven-sketches F03872	327	0.958	$x \multimap y = \bigvee \{w \mid w \wedge x \leq y\}$	$x \multimap y = \bigvee \{w \mid w \wedge x \leq y\}$	each x, y , the hom-object $x \multimap y$ exists. The formula for it is $x \multimap y = \bigvee \{w \mid w \wedge x \leq y\}$, i.e. we
fong-spivak-seven-sketches F03873	327	0.997	$y \wedge x \leq y$	$y \wedge x \leq y$	need these particular joins to exist. Since $y \wedge x \leq y$, we have $y \leq x \multimap y$. So we can replace the
fong-spivak-seven-sketches F03874	327	0.997	$y \leq x \multimap y$	$y \leq x \multimap y$	need these particular joins to exist. Since $y \wedge x \leq y$, we have $y \leq x \multimap y$. So we can replace the
fong-spivak-seven-sketches F03875	327	0.431	$x \multimap y = \bigvee \{w \mid y \leq w$	$x \multimap y = \bigvee \{w \mid y \leq w$	formula with $x \multimap y = \bigvee \{w \mid y \leq w$ and $w \wedge x \leq y\}$. But the set of elements in $\mathbb{N}^{\mathrm{op}} \times \mathbb{N}^{\mathrm{op}}$ that
fong-spivak-seven-sketches F03876	327	0.431	$w \wedge x \leq y\}$	$w \wedge x \leq y\}$	formula with $x \multimap y = \bigvee \{w \mid y \leq w$ and $w \wedge x \leq y\}$. But the set of elements in $\mathbb{N}^{\mathrm{op}} \times \mathbb{N}^{\mathrm{op}}$ that
fong-spivak-seven-sketches F03877	327	0.431	$\mathbb{N}^{\mathrm{op}} \times \mathbb{N}^{\mathrm{op}}$	$\mathbb{N}^{\mathrm{op}} \times \mathbb{N}^{\mathrm{op}}$	formula with $x \multimap y = \bigvee \{w \mid y \leq w$ and $w \wedge x \leq y\}$. But the set of elements in $\mathbb{N}^{\mathrm{op}} \times \mathbb{N}^{\mathrm{op}}$ that
fong-spivak-seven-sketches F03878	327	1.000	$m: \mathbb{Z} \rightarrow \mathbb{B}$	$m: \mathbb{Z} \rightarrow \mathbb{B}$	Let $m: \mathbb{Z} \rightarrow \mathbb{B}$ be the characteristic function of the inclusion $\mathbb{N} \subseteq \mathbb{Z}$.
fong-spivak-seven-sketches F03879	327	1.000	$\mathbb{N} \subseteq \mathbb{Z}$	$\mathbb{N} \subseteq \mathbb{Z}$	Let $m: \mathbb{Z} \rightarrow \mathbb{B}$ be the characteristic function of the inclusion $\mathbb{N} \subseteq \mathbb{Z}$.
fong-spivak-seven-sketches F03880	327	1.000	$\ulcorner m \urcorner(-5) =$	$\ulcorner m \urcorner(-5) =$	1. $\ulcorner m \urcorner(-5) = \text{false}$.
fong-spivak-seven-sketches F03881	327	1.000	$\ulcorner m \urcorner(0) =$	$\ulcorner m \urcorner(0) =$	2. $\ulcorner m \urcorner(0) = \text{true}$.
fong-spivak-seven-sketches F03882	327	1.000	$A \subseteq \mathbb{B}$	$A \subseteq \mathbb{B}$	1. The sort of thing (**) we're looking for is a subobject of $\mathbb{B}$, say $A \subseteq \mathbb{B}$. This would have a
fong-spivak-seven-sketches F03883	327	0.993	$\ulcorner \Rightarrow \urcorner: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$	$\ulcorner \Rightarrow \urcorner: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$	3. The characteristic function for $P \Rightarrow Q$ is the function $\ulcorner \Rightarrow \urcorner: \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{B}$ given by the first, second,
fong-spivak-seven-sketches F03884	327	0.991	$\subseteq \mathbb{B} \times \mathbb{B}$	$\subseteq \mathbb{B} \times \mathbb{B}$	4. It classifies the subset $\{(\text{true}, \text{true}), (\text{false}, \text{true}), (\text{false}, \text{false})\}$.
fong-spivak-seven-sketches F03885	327	1.000	$\ulcorner E \urcorner, \ulcorner P \urcorner, \ulcorner T \urcorner: \mathbb{N} \rightarrow \mathbb{B}$	$\ulcorner E \urcorner, \ulcorner P \urcorner, \ulcorner T \urcorner: \mathbb{N} \rightarrow \mathbb{B}$	Say that $\ulcorner E \urcorner, \ulcorner P \urcorner, \ulcorner T \urcorner: \mathbb{N} \rightarrow \mathbb{B}$ classify respectively the subsets $E := \{n \in \mathbb{N} \mid n \text{ is even}\}$, $P := \{n \in \mathbb{N} \mid$
fong-spivak-seven-sketches F03886	327	1.000	$\}$, $P := \{n \in \mathbb{N} \mid n \text{ is prime}\}$	$\}, P := \{n \in \mathbb{N} \mid$	Say that $\ulcorner E \urcorner, \ulcorner P \urcorner, \ulcorner T \urcorner: \mathbb{N} \rightarrow \mathbb{B}$ classify respectively the subsets $E := \{n \in \mathbb{N} \mid n \text{ is even}\}$, $P := \{n \in \mathbb{N} \mid$
fong-spivak-seven-sketches F03887	327	0.994	$\ulcorner E \urcorner(17) =$	$\ulcorner E \urcorner(17) =$	1. $\ulcorner E \urcorner(17) = \text{false}$ because 17 is not even.
fong-spivak-seven-sketches F03888	327	1.000	$\ulcorner P \urcorner(17) =$	$\ulcorner P \urcorner(17) =$	2. $\ulcorner P \urcorner(17) = \text{true}$ because 17 is prime.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03889	327	0.857	$\ulcorner T \urcorner(17) =$	$\ulcorner T \urcorner(17) =$	3. $\ulcorner T \urcorner(17) = \text{true}$ because $17 \geq 10$.
fong-spivak-seven-sketches F03890	327	0.857	$17 \geq 10$	$17 \geq 10$	3. $\ulcorner T \urcorner(17) = \text{true}$ because $17 \geq 10$.
fong-spivak-seven-sketches F03891	327	0.679	$B(x, \epsilon) := \{x' \in \mathbb{R} \mid x - x' < \epsilon\}$	$B(x, \epsilon) := \{x' \in \mathbb{R} \mid x - x' < \epsilon\}$	1. The 1-dimensional analogue of an ϵ -ball around a point $x \in \mathbb{R}$ is $B(x, \epsilon) := \{x' \in \mathbb{R} \mid x - x' < \epsilon\}$,
fong-spivak-seven-sketches F03892	327	1.000	$x \in U$	$x \in U$	2. A subset $U \subseteq \mathbb{R}$ is open if, for every $x \in U$ there is some $\epsilon > 0$ such that $B(x, \epsilon) \subseteq U$.
fong-spivak-seven-sketches F03893	327	1.000	$B(x, \epsilon) \subseteq U$	$B(x, \epsilon) \subseteq U$	2. A subset $U \subseteq \mathbb{R}$ is open if, for every $x \in U$ there is some $\epsilon > 0$ such that $B(x, \epsilon) \subseteq U$.
fong-spivak-seven-sketches F03894	327	1.000	$U_1 := \{x \in \mathbb{R} \mid 0 < x < 2\}$	$U_1 := \{x \in \mathbb{R} \mid 0 < x < 2\}$	3. Let $U_1 := \{x \in \mathbb{R} \mid 0 < x < 2\}$ and $U_2 := \{x \in \mathbb{R} \mid 1 < x < 3\}$. Then $U := U_1 \cup U_2 = \{x \in \mathbb{R} \mid 0 <$
fong-spivak-seven-sketches F03895	327	1.000	$U_2 := \{x \in \mathbb{R} \mid 1 < x < 3\}$	$U_2 := \{x \in \mathbb{R} \mid 1 < x < 3\}$	3. Let $U_1 := \{x \in \mathbb{R} \mid 0 < x < 2\}$ and $U_2 := \{x \in \mathbb{R} \mid 1 < x < 3\}$. Then $U := U_1 \cup U_2 = \{x \in \mathbb{R} \mid 0 <$
fong-spivak-seven-sketches F03896	327	1.000	$U := U_1 \cup U_2 = \{x \in \mathbb{R} \mid 0 <$	$U := U_1 \cup U_2 = \{x \in \mathbb{R} \mid 0 <$	3. Let $U_1 := \{x \in \mathbb{R} \mid 0 < x < 2\}$ and $U_2 := \{x \in \mathbb{R} \mid 1 < x < 3\}$. Then $U := U_1 \cup U_2 = \{x \in \mathbb{R} \mid 0 <$
fong-spivak-seven-sketches F03897	327	1.000	$x < 3\}$	$x < 3\}$	$x < 3\}$.
fong-spivak-seven-sketches F03898	327	1.000	$y = (a, b)$	$y = (a, b)$	² If $y = (a, b)$ then there are exactly $(a + 1) * (b + 1)$ elements y' for which $y \leq y'$.
fong-spivak-seven-sketches F03899	327	1.000	$(a + 1) * (b + 1)$	$(a + 1) * (b + 1)$	² If $y = (a, b)$ then there are exactly $(a + 1) * (b + 1)$ elements y' for which $y \leq y'$.
fong-spivak-seven-sketches F03900	328	1.000	$I = \{1, 2, 3, 4, \ldots\}$	$I = \{1, 2, 3, 4, \ldots\}$	4. Let $I = \{1, 2, 3, 4, \ldots\}$ and for each $i \in I$ let $U_i := \{x \in \mathbb{R} \mid \frac{1}{i} < x < 1\}$, so we have $U_1 \subseteq U_2 \subseteq$
fong-spivak-seven-sketches F03901	328	1.000	$U_i := \{x \in \mathbb{R} \mid \frac{1}{i} < x < 1\}$	$U_i := \left\{x \in \mathbb{R} \mid \frac{1}{i} < x < 1\right\}$	4. Let $I = \{1, 2, 3, 4, \ldots\}$ and for each $i \in I$ let $U_i := \{x \in \mathbb{R} \mid \frac{1}{i} < x < 1\}$ so we have $U_1 \subseteq U_2 \subseteq$
fong-spivak-seven-sketches F03902	328	1.000	$U_1 \subseteq U_2 \subseteq$	$U_1 \subseteq U_2 \subseteq$	4. Let $I = \{1, 2, 3, 4, \ldots\}$ and for each $i \in I$ let $U_i := \{x \in \mathbb{R} \mid \frac{1}{i} < x < 1\}$, so we have $U_1 \subseteq U_2 \subseteq$
fong-spivak-seven-sketches F03903	328	1.000	$U_3 \subseteq \cdots$	$U_3 \subseteq \cdots$	$U_3 \subseteq \cdots$. Their union is $U := \bigcup_{i \in I} U_i = \{x \in \mathbb{R} \mid 0 < x < 1\}$.
fong-spivak-seven-sketches F03904	328	1.000	$U := \bigcup_{i \in I} U_i = \{x \in \mathbb{R} \mid 0 < x < 1\}$	$U := \bigcup_{i \in I} U_i = \{x \in \mathbb{R} \mid 0 < x < 1\}$	$U_3 \subseteq \cdots$. Their union is $U := \bigcup_{i \in I} U_i = \{x \in \mathbb{R} \mid 0 < x < 1\}$.
fong-spivak-seven-sketches F03905	328	1.000	$\varnothing \subseteq X$	$\varnothing \subseteq X$	1. The coarse topology on X is the one whose only open sets are $X \subseteq X$ and $\varnothing \subseteq X$. This is a
fong-spivak-seven-sketches F03906	328	0.999	$\bigcup_{i \in I} A_i$	$\bigcup_{i \in I} A_i$	$\bigcup_{i \in I} A_i$ is X iff $A_i = X$ for some $i \in I$).
fong-spivak-seven-sketches F03907	328	0.999	$A_i = X$	$A_i = X$	$\bigcup_{i \in I} A_i$ is X iff $A_i = X$ for some $i \in I$).
fong-spivak-seven-sketches F03908	328	1.000	$A \subseteq X$	$A \subseteq X$	2. The fine topology on X is the one where every subset $A \subseteq X$ is considered open. All the
fong-spivak-seven-sketches F03909	328	0.405	$(X, \mathbf{P}(X))$	$(X, \mathbf{P}(X))$	3. If $(X, \mathbf{P}(X))$ is discrete, $(Y, \mathbf{Op}_Y)$ is any topological space, and $f: X \rightarrow Y$ is any function then it
fong-spivak-seven-sketches F03910	328	1.000	$f^{-1}(U) \subseteq X$	$f^{-1}(U) \subseteq X$	is continuous. Indeed, this just means that for any open set $U \subseteq Y$ the preimage $f^{-1}(U) \subseteq X$ is
fong-spivak-seven-sketches F03911	328	1.000	$\varnothing \rightarrow \{1\} \rightarrow \{1, 2\}$	$\varnothing \rightarrow \{1\} \rightarrow \{1, 2\}$	1. The Hasse diagram for the Sierpinsky topology is $\varnothing \rightarrow \{1\} \rightarrow \{1, 2\}$.
fong-spivak-seven-sketches F03912	328	0.994	$U_i = U$	$U_i = U$	• $U_i = U$ for some $i \in I$.
fong-spivak-seven-sketches F03913	328	1.000	$Y = B \cap Y$	$Y = B \cap Y$	1. We want to show that $Y \in \mathbf{Op}_{\mathcal{?} \cap Y}$. We need to find $B \in \mathbf{Op}$ such that $Y = B \cap Y$; this is easy, you
fong-spivak-seven-sketches F03914	328	1.000	$B = Y$	$B = Y$	could take $B = Y$ or $B = X$, or anything in between.
fong-spivak-seven-sketches F03915	328	1.000	$B = X$	$B = X$	could take $B = Y$ or $B = X$, or anything in between.
fong-spivak-seven-sketches F03916	328	1.000	$\varnothing = \varnothing \cap Y$	$\varnothing = \varnothing \cap Y$	union. $\varnothing = \varnothing \cap Y$, so according to the formula, $\varnothing \in \mathbf{Op}_{\mathcal{?} \cap Y}$. Suppose that $A_1, A_2 \in \mathbf{Op}_{\mathcal{?} \cap Y}$. Then

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03917	328	■ 1.000	$\varnothing \in \mathbf{Op}_{? \cap Y}$	$\emptyset \in \mathbf{Op}_{? \cap Y}$	union. $\emptyset = \emptyset \cap Y$, so according to the formula, $\emptyset \in \mathbf{Op}_{? \cap Y}$. Suppose that $A_1, A_2 \in \mathbf{Op}_{? \cap Y}$. Then
fong-spivak-seven-sketches F03918	328	■ 1.000	$A_{\{1\}}, A_{\{2\}} \in \mathbf{Op}_{? \cap Y}$	$A_1, A_2 \in \mathbf{Op}_{? \cap Y}$	union. $\emptyset = \emptyset \cap Y$, so according to the formula, $\emptyset \in \mathbf{Op}_{? \cap Y}$. Suppose that $A_1, A_2 \in \mathbf{Op}_{? \cap Y}$. Then
fong-spivak-seven-sketches F03919	328	■ 0.999	$B_{\{1\}}, B_{\{2\}} \in \mathbf{Op}$	$B_1, B_2 \in \mathbf{Op}$	there exist $B_1, B_2 \in \mathbf{Op}$ with $A_1 = B_1 \cap Y$ and $A_2 = B_2 \cap Y$. But then $A_1 \cap A_2 = (B_1 \cap Y) \cap (B_2 \cap Y) =$
fong-spivak-seven-sketches F03920	328	■ 0.999	$A_{\{1\}} = B_{\{1\}} \cap Y$	$A_1 = B_1 \cap Y$	there exist $B_1, B_2 \in \mathbf{Op}$ with $A_1 = B_1 \cap Y$ and $A_2 = B_2 \cap Y$. But then $A_1 \cap A_2 = (B_1 \cap Y) \cap (B_2 \cap Y) =$
fong-spivak-seven-sketches F03921	328	■ 0.999	$A_{\{2\}} = B_{\{2\}} \cap Y$	$A_2 = B_2 \cap Y$	there exist $B_1, B_2 \in \mathbf{Op}$ with $A_1 = B_1 \cap Y$ and $A_2 = B_2 \cap Y$. But then $A_1 \cap A_2 = (B_1 \cap Y) \cap (B_2 \cap Y) =$
fong-spivak-seven-sketches F03922	328	■ 0.999	$A_{\{1\}} \cap A_{\{2\}} = \left(B_{\{1\}} \cap Y \right) \cap \left(B_{\{2\}} \cap Y \right) =$	$A_1 \cap A_2 = (B_1 \cap Y) \cap (B_2 \cap Y) =$	there exist $B_1, B_2 \in \mathbf{Op}$ with $A_1 = B_1 \cap Y$ and $A_2 = B_2 \cap Y$. But then $A_1 \cap A_2 = (B_1 \cap Y) \cap (B_2 \cap Y) =$
fong-spivak-seven-sketches F03923	328	■ 0.995	$\left(B_{\{1\}} \cap B_{\{2\}} \right) \cap Y$	$(B_1 \cap B_2) \cap Y$	$(B_1 \cap B_2) \cap Y$, so it is in $\mathbf{Op}_{? \cap Y}$ since $B_1 \cap B_2 \in \mathbf{Op}$. The same idea works for arbitrary unions:
fong-spivak-seven-sketches F03924	328	■ 0.995	$B_{\{1\}} \cap B_{\{2\}} \in \mathbf{Op}$	$B_1 \cap B_2 \in \mathbf{Op}$	$(B_1 \cap B_2) \cap Y$, so it is in $\mathbf{Op}_{? \cap Y}$ since $B_1 \cap B_2 \in \mathbf{Op}$. The same idea works for arbitrary unions:
fong-spivak-seven-sketches F03925	328	■ 1.000	$A_{\{i\}}$	A_i	given a set I and A_i for each $i \in I$, we have $A_i = B_i \cap Y$ for some $B_i \in \mathbf{Op}$, and
fong-spivak-seven-sketches F03926	328	■ 1.000	$A_{\{i\}} = B_{\{i\}} \cap Y$	$A_i = B_i \cap Y$	given a set I and A_i for each $i \in I$, we have $A_i = B_i \cap Y$ for some $B_i \in \mathbf{Op}$, and
fong-spivak-seven-sketches F03927	328	■ 1.000	$B_{\{i\}} \in \mathbf{Op}$	$B_i \in \mathbf{Op}$	given a set I and A_i for each $i \in I$, we have $A_i = B_i \cap Y$ for some $B_i \in \mathbf{Op}$, and
fong-spivak-seven-sketches F03928	328	■ 1.000	$U \subseteq \mathbf{Op}$	$U \subseteq \mathbf{Op}$	by an open set $U \subseteq \mathbf{Op}$. It might look something like this:
fong-spivak-seven-sketches F03929	328	■ 1.000	$\mathcal{C}(B, C) =$	$\mathcal{C}(B, C) =$	paths between them, of the monoidal product of distances along that path. For example, $\mathcal{C}(B, C) =$
fong-spivak-seven-sketches F03930	329	■ 1.000	$\left(U_3 \wedge U_1 \right) \vee \left(U_4 \wedge U_2 \right)$	$(U_3 \wedge U_1) \vee (U_4 \wedge U_2)$	$(U_3 \wedge U_1) \vee (U_4 \wedge U_2)$, because $\wedge$ is the monoidal product in $\mathcal{V}$.
fong-spivak-seven-sketches F03931	329	■ 1.000	$\mathcal{C}(a, b)$	$\mathcal{C}(a, b)$	In general, we can thus imagine the open set $\mathcal{C}(a, b)$ as a kind of ‘size restriction’ for getting from a to b ,
fong-spivak-seven-sketches F03932	329	■ 1.000	$\{a_1, a_2\}$	$\{a_1, a_2\}$	1. The fiber of f over a is $\{a_1, a_2\}$.
fong-spivak-seven-sketches F03933	329	■ 1.000	$\{c_1\}$	$\{c_1\}$	2. The fiber of f over c is $\{c_1\}$.
fong-spivak-seven-sketches F03934	329	■ 1.000	$\mathrm{Sec}_f(V_2) = \varnothing$	$\mathrm{Sec}_f(V_2) = \emptyset$	2. When $V_2 = \{a, b, c, d\}$, there are no sections: $\mathrm{Sec}_f(V_2) = \emptyset$.
fong-spivak-seven-sketches F03935	329	■ 0.949	$\mathrm{Sec}_f(V_3)$	$\mathrm{Sec}_f(V_3)$	3. When $V_3 = \{a, b, d, e\}$, the set $\mathrm{Sec}_f(V_3)$ has $2 * 3 * 1 * 2 = 12$ elements.
fong-spivak-seven-sketches F03936	329	■ 0.949	$2 * 3 * 1 * 2 = 12$	$2 * 3 * 1 * 2 = 12$	3. When $V_3 = \{a, b, d, e\}$, the set $\mathrm{Sec}_f(V_3)$ has $2 * 3 * 1 * 2 = 12$ elements.
fong-spivak-seven-sketches F03937	330	■ 1.000	$g_{\{1\}} := \left(a_{\{1\}}, b_{\{1\}} \right)$	$g_1 := (a_1, b_1)$	1. Let $g_1 := (a_1, b_1)$ and $g_2 := (b_2, e_1)$; these do not agree on the overlap.
fong-spivak-seven-sketches F03938	330	■ 1.000	$g_{\{2\}} := \left(b_{\{2\}}, e_{\{1\}} \right)$	$g_2 := (b_2, e_1)$	1. Let $g_1 := (a_1, b_1)$ and $g_2 := (b_2, e_1)$; these do not agree on the overlap.
fong-spivak-seven-sketches F03939	330	■ 1.000	$g \in \mathrm{Sec}_f(U_1 \cup U_2)$	$g \in \mathrm{Sec}_f(U_1 \cup U_2)$	2. No, there’s no section $g \in \mathrm{Sec}_f(U_1 \cup U_2)$ for which $g _{U_1} = g_1$ and $g _{U_2} = g_2$.
fong-spivak-seven-sketches F03940	330	■ 1.000	$g _{U_1} = g_1$	$g _{U_1} = g_1$	2. No, there’s no section $g \in \mathrm{Sec}_f(U_1 \cup U_2)$ for which $g _{U_1} = g_1$ and $g _{U_2} = g_2$.
fong-spivak-seven-sketches F03941	330	■ 1.000	$g _{U_2} = g_2$	$g _{U_2} = g_2$	2. No, there’s no section $g \in \mathrm{Sec}_f(U_1 \cup U_2)$ for which $g _{U_1} = g_1$ and $g _{U_2} = g_2$.
fong-spivak-seven-sketches F03942	330	■ 0.997	$F(\{1, 2\}) \rightarrow F(\{1\}) \rightarrow F(\varnothing)$	$F(\{1, 2\}) \rightarrow F(\{1\}) \rightarrow F(\emptyset)$	2. A presheaf F on $\mathbf{Op}$ consists of any three sets and any two functions $F(\{1, 2\}) \rightarrow F(\{1\}) \rightarrow F(\emptyset)$
fong-spivak-seven-sketches F03943	330	■ 1.000	$F(\varnothing) \cong \{1\}$	$F(\emptyset) \cong \{1\}$	4. As explained in Example 7.36, F will be a sheaf iff $F(\emptyset) \cong \{1\}$. Thus we the category of sheaves
fong-spivak-seven-sketches F03944	330	■ 1.000	$F(\{1, 2\}) \rightarrow F(\{1\})$	$F(\{1, 2\}) \rightarrow F(\{1\})$	is equivalent to that of just two sets and one function $F(\{1, 2\}) \rightarrow F(\{1\})$.

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F03945	330	■ 1.000	$S(\backslash\mathrm{varnothing})=\backslash\{()\backslash$	$S(\emptyset) = \{()\}$	$S(\emptyset) = \{()\}$ by the sheaf condition (see Example 7.36). So the only data in a sheaf $S \in \mathbf{Shv}(X)$ is the set
fong-spivak-seven-sketches F03946	330	■ 1.000	$S(\backslash\{1\}\backslash)$	$S(\{1\})$	$S(\{1\})$. This is how we get the correspondence between sets and sheaves on the one point space.
fong-spivak-seven-sketches F03947	330	■ 0.751	$\backslash\mathrm{0}\mathrm{omega}:\backslash\mathrm{mathbf{0}\ p\}(X)^{\backslash\mathrm{mathrm{op}}}\backslash\rightarrow$	$\Omega : \mathbf{Op}(X)^{\mathrm{op}} \rightarrow$	According to Eq. (7.50), the subobject classifier $\Omega : \mathbf{Op}(X)^{\mathrm{op}} \rightarrow \mathbf{Set}$ in $\mathbf{Shv}(X)$ should be the functor
fong-spivak-seven-sketches F03948	330	■ 0.947	$\backslash\mathrm{0}\mathrm{omega}(\backslash\{1\}\backslash)$	$\Omega(\{1\})$	where $\Omega(\{1\})$ is the set of open sets of $\{1\}$. So we're hoping to see that there is a one-to-one correspon-
fong-spivak-seven-sketches F03949	330	■ 1.000	$\backslash\mathrm{mathbf{0}\ p\}(\backslash\{1\}\backslash)$	$\mathbf{Op}(\{1\})$	dence between the set $\mathbf{Op}(\{1\})$ and the set $\mathbb{B} = \{\mathrm{true}, \mathrm{false}\}$ of booleans. Indeed there is: there are
fong-spivak-seven-sketches F03950	330	■ 0.600	$\backslash\mathrm{0}\mathrm{omega}(U):=\backslash\mathrm{left}\{U^{\backslash\mathrm{prime}}\}\backslash\mathrm{in}\backslash\mathrm{mathbf{0}\ p\}\backslash\mathrm{mid}\backslash U^{\backslash\mathrm{prime}}\}\backslash\mathrm{subse}\mathrm{teq}\backslash U\backslash\mathrm{right}\backslash$	$\Omega(U) := \{U' \in \mathbf{Op} \mid U' \subseteq U\}$	By Eqs. (7.50) and (7.51) the definition of $\Omega(U)$ is $\Omega(U) := \{U' \in \mathbf{Op} \mid U' \subseteq U\}$, and the definition of
fong-spivak-seven-sketches F03951	330	■ 1.000	$U^{\backslash\mathrm{prime}}\}\backslash\mathrm{mapsto}\backslash U^{\backslash\mathrm{prime}}\}\backslash\mathrm{cap}\backslash V$	$U' \mapsto U' \cap V$	the restriction map for $V \subseteq U$ is $U' \mapsto U' \cap V$.
fong-spivak-seven-sketches F03952	331	■ 0.998	$U^{\backslash\mathrm{prime}}\}\backslash\mathrm{subse}\mathrm{teq}\backslash U$	$U' \subseteq U$	1. It is functorial: given $W \subseteq V \subseteq U$ and $U' \subseteq U$, we indeed have $(U' \cap V) \cap W = U' \cap W$, since
fong-spivak-seven-sketches F03953	331	■ 0.998	$\backslash\mathrm{left}(U^{\backslash\mathrm{prime}}\}\backslash\mathrm{cap}\backslash V\backslash\mathrm{right})\backslash\mathrm{cap}\backslash W=U^{\backslash\mathrm{prime}}\}\backslash\mathrm{cap}\backslash W$	$(U' \cap V) \cap W = U' \cap W$	1. It is functorial: given $W \subseteq V \subseteq U$ and $U' \subseteq U$, we indeed have $(U' \cap V) \cap W = U' \cap W$, since
fong-spivak-seven-sketches F03954	331	■ 1.000	$W\backslash\mathrm{subse}\mathrm{teq}\backslash V$	$W \subseteq V$	$W \subseteq V$. For functoriality, we also need preservation of identities, and this amounts to $U' \cap U = U'$
fong-spivak-seven-sketches F03955	331	■ 1.000	$U^{\backslash\mathrm{prime}}\}\backslash\mathrm{cap}\backslash U=U^{\backslash\mathrm{prime}}\}$	$U' \cap U = U'$	$W \subseteq V$. For functoriality, we also need preservation of identities, and this amounts to $U' \cap U = U'$
fong-spivak-seven-sketches F03956	331	■ 0.947	$G^{\backslash\mathrm{prime}}\}$	G'	There is only one that classifies G' , and here it is. Let's write $\gamma := \ulcorner G' \urcorner$.
fong-spivak-seven-sketches F03957	331	■ 0.947	$\backslash\mathrm{gamma}:=\backslash\mathrm{left}\ulcorner G^{\backslash\mathrm{prime}}\}\backslash\mathrm{right}\backslash.$	$\gamma := \ulcorner G' \urcorner$	There is only one that classifies G' , and here it is. Let's write $\gamma := \ulcorner G' \urcorner$.
fong-spivak-seven-sketches F03958	331	■ 1.000	$\backslash\mathrm{gamma}(D)=\emptyset$	$\gamma(D) = 0$	• Since D is missing from G' , we have $\gamma(D) = 0$ (vertex: missing).
fong-spivak-seven-sketches F03959	331	■ 1.000	$A,\ B,\ C$	A, B, C	• Since vertices A, B, C are present in G' we have $\gamma(A) = \gamma(B) = \gamma(C) = V$ (vertex: present).
fong-spivak-seven-sketches F03960	331	■ 1.000	$\backslash\mathrm{gamma}(A)=\backslash\mathrm{gamma}(B)=\backslash\mathrm{gamma}(C)=V$	$\gamma(A) = \gamma(B) = \gamma(C) = V$	• Since vertices A, B, C are present in G' we have $\gamma(A) = \gamma(B) = \gamma(C) = V$ (vertex: present).
fong-spivak-seven-sketches F03961	331	■ 1.000	$\backslash\mathrm{gamma}(i)=(V,\ 0\ ;\ 0)$	$\gamma(i) = (V, 0; 0)$	• The above forces $\gamma(i) = (V, 0; 0)$ (arrow from present vertex to missing vertex: missing).
fong-spivak-seven-sketches F03962	331	■ 1.000	$\backslash\mathrm{gamma}(f)=(V,\ V\ ;\ A)$	$\gamma(f) = (V, V; A)$	• Since the arrow f is in G' , we have $\gamma(f) = (V, V; A)$ (arrow from present vertex to present vertex:
fong-spivak-seven-sketches F03963	331	■ 1.000	$\backslash\mathrm{gamma}(g)=\backslash\mathrm{gamma}(h)=(V,\ V\ ;\ 0)$	$\gamma(g) = \gamma(h) = (V, V; 0)$	• Since the arrows g and h are missing in G' , we have $\gamma(g) = \gamma(h) = (V, V; 0)$ (arrow from present
fong-spivak-seven-sketches F03964	331	■ 1.000	$U=\backslash\mathrm{mathbb{R}}\backslash-\backslash\{0\}\backslash\backslash\mathrm{subse}\mathrm{teq}\backslash\mathrm{mathbb{R}}\backslash$	$U = \mathbb{R} - \{0\} \subseteq \mathbb{R}$	With $U := \mathbb{R} - \{0\} \subseteq \mathbb{R}$, we have:
fong-spivak-seven-sketches F03965	331	■ 1.000	$\backslash\mathrm{mathbb{R}}\backslash-U=\backslash\{0\}\backslash$	$\mathbb{R} - U = \{0\}$	1. The complement of U is $\mathbb{R} - U = \{0\}$ and $\neg U$ is its interior, which is $\neg U = \emptyset$.
fong-spivak-seven-sketches F03966	331	■ 1.000	$\backslash\mathrm{neg}\backslash U=\backslash\mathrm{varnothing}\backslash$	$\neg U = \emptyset$	1. The complement of U is $\mathbb{R} - U = \{0\}$ and $\neg U$ is its interior, which is $\neg U = \emptyset$.
fong-spivak-seven-sketches F03967	331	■ 1.000	$\backslash\mathrm{mathbb{R}}\backslash-\backslash\mathrm{varnothing}\backslash=\backslash\mathrm{mathbb{R}}\backslash$	$\mathbb{R} - \emptyset = \mathbb{R}$	2. The complement of $\neg U$ is $\mathbb{R} - \emptyset = \mathbb{R}$, and this is open, so $\neg\neg U = \mathbb{R}$.
fong-spivak-seven-sketches F03968	331	■ 1.000	$\backslash\mathrm{neg}\backslash\mathrm{neg}\backslash U=\backslash\mathrm{mathbb{R}}\backslash$	$\neg\neg U = \mathbb{R}$	2. The complement of $\neg U$ is $\mathbb{R} - \emptyset = \mathbb{R}$, and this is open, so $\neg\neg U = \mathbb{R}$.
fong-spivak-seven-sketches F03969	331	■ 0.943	$\backslash\mathrm{neg}\backslash\mathrm{neg}\backslash U\backslash\mathrm{subse}\mathrm{teq}$	$\neg\neg U \subseteq$	4. It is false that $\neg\neg U \supseteq U$.
fong-spivak-seven-sketches F03970	331	■ 0.995	$V\backslash\mathrm{in}\backslash\mathrm{mathbf{0}\ p\}$	$V \in \mathbf{Op}$	1. If for any $V \in \mathbf{Op}$ we have $\top \wedge V = V$ then when $V = X$ we have $\top \wedge X := \top \cap X = X$, but
fong-spivak-seven-sketches F03971	331	■ 0.995	$\backslash\mathrm{top}\backslash\mathrm{wedge}\backslash V=V$	$\top \wedge V = V$	1. If for any $V \in \mathbf{Op}$ we have $\top \wedge V = V$ then when $V = X$ we have $\top \wedge X := \top \cap X = X$, but
fong-spivak-seven-sketches F03972	331	■ 0.995	$V=X$	$V = X$	1. If for any $V \in \mathbf{Op}$ we have $\top \wedge V = V$ then when $V = X$ we have $\top \wedge X := \top \cap X = X$, but
fong-spivak-seven-sketches F03973	331	■ 0.995	$\backslash\mathrm{top}\backslash\mathrm{wedge}\backslash X:=\backslash\mathrm{top}\backslash\mathrm{cap}\backslash X=X$	$\top \wedge X := \top \cap X = X$	1. If for any $V \in \mathbf{Op}$ we have $\top \wedge V = V$ then when $V = X$ we have $\top \wedge X := \top \cap X = X$, but

Identifier	Page	Conf.	LaTeX source	Rendered	Image
fong-spivak-seven-sketches F04002	333	■ 1.000	$s \in S(U), p(s)$	$s \in S(U), p(s)$	2. Let U be the interval 2019/01/01 – 2019/02/01. For an arbitrary person $s \in S(U)$, $p(s)$ is a subset
fong-spivak-seven-sketches F04003	333	■ 1.000	$j(j(p(s)))=j(p(s))$	$j(j(p(s))) = j(p(s))$	5. It is true that $j(j(p(s))) = j(p(s))$, because suppose given a time during which “if Bob is in San
fong-spivak-seven-sketches F04004	333	■ 1.000	$0 \leq 2 \leq 6 \leq 8$	$0 \leq 2 \leq 6 \leq 8$	1. Since $0 \leq 2 \leq 6 \leq 8$, we have $[2, 6] \in o_{[0,8]}$ by the above formula.
fong-spivak-seven-sketches F04005	333	■ 0.977	$[2,6] \in^? o_{\{[0,5]\} \cup o_{\{[4,8]\}}}$	$[2, 6] \in^? o_{[0,5]} \cup o_{[4,8]}$	2. In order to have $[2, 6] \in^? o_{[0,5]} \cup o_{[4,8]}$, we would need to have either $[2, 6] \in^? o_{[0,5]}$ or $[2, 6] \in^? o_{[4,8]}$.
fong-spivak-seven-sketches F04006	333	■ 0.977	$[2,6] \in^? o_{\{[0,5]\}}$	$[2, 6] \in^? o_{[0,5]}$	2. In order to have $[2, 6] \in^? o_{[0,5]} \cup o_{[4,8]}$, we would need to have either $[2, 6] \in^? o_{[0,5]}$ or $[2, 6] \in^? o_{[4,8]}$.
fong-spivak-seven-sketches F04007	333	■ 0.977	$[2,6] \in^? o_{\{[4,8]\}}$	$[2, 6] \in^? o_{[4,8]}$	2. In order to have $[2, 6] \in^? o_{[0,5]} \cup o_{[4,8]}$, we would need to have either $[2, 6] \in^? o_{[0,5]}$ or $[2, 6] \in^? o_{[4,8]}$.
fong-spivak-seven-sketches F04008	333	■ 0.510	$U^{\prime} \subseteq \mathbb{R}$	$U' \subseteq \mathbb{R}$	A subset $U \subseteq \mathbb{R}$ is open in the subspace topology of $\mathbb{R} \subseteq \mathbb{IR}$ iff there is an open set $U' \subseteq \mathbb{IR}$ with
fong-spivak-seven-sketches F04009	333	■ 1.000	$U=U^{\prime} \cap \mathbb{R}$	$U = U' \cap \mathbb{R}$	$U = U' \cap \mathbb{R}$. We want to show that this is the case iff U is open in the usual topology.
fong-spivak-seven-sketches F04010	333	■ 0.999	$U^{\prime}=\bigcup_{i \in I} o_{\left[a_{i}, b_{i}\right]}\backslash\right\}$	$U' = \bigcup_{i \in I} o_{[a_i, b_i]}$	basic opens, $U' = \bigcup_{i \in I} o_{[a_i, b_i]}$, where $o_{[a_i, b_i]} = \{[d, u] \in \mathbb{IR} \mid a_i < d < u < b_i\}$. Since $\mathbb{R} = \{[x, x] \in \mathbb{IR}\}$,
fong-spivak-seven-sketches F04011	333	■ 0.999	$o_{\left[a_{i}, b_{i}\right]}\backslash\right\}=\left\{[d, u] \in \mathbb{R} \mid a_{i} < d < u < b_{i}\right\}$	$o_{[a_i, b_i]} = \{[d, u] \in \mathbb{R} \mid a_i < d < u < b_i\}$	basic opens, $U' = \bigcup_{i \in I} o_{[a_i, b_i]}$, where $o_{[a_i, b_i]} = \{[d, u] \in \mathbb{IR} \mid a_i < d < u < b_i\}$. Since $\mathbb{R} = \{[x, x] \in \mathbb{IR}\}$,
fong-spivak-seven-sketches F04012	333	■ 0.999	$\mathbb{R}=\{[x, x] \in \mathbb{R}\}$	$\mathbb{R} = \{[x, x] \in \mathbb{R}\}$	basic opens, $U' = \bigcup_{i \in I} o_{[a_i, b_i]}$, where $o_{[a_i, b_i]} = \{[d, u] \in \mathbb{IR} \mid a_i < d < u < b_i\}$. Since $\mathbb{R} = \{[x, x] \in \mathbb{IR}\}$,
fong-spivak-seven-sketches F04013	333	■ 1.000	$U^{\prime} \cap \mathbb{R}$	$U' \cap \mathbb{R}$	the intersection $U' \cap \mathbb{R}$ will then be
fong-spivak-seven-sketches F04014	333	■ 1.000	$B\left(m_{i}, r_{i}\right)$	$B\left(m_i, r_i\right)$	and this is just the union of open balls $B(m_i, r_i)$ where $m_i := \frac{a_i+b_i}{2}$ is the midpoint and $r_i := \frac{b_i-a_i}{2}$ is
fong-spivak-seven-sketches F04015	333	■ 1.000	$m_i:=\frac{a_i+b_i}{2}$	$m_i := \frac{a_i + b_i}{2}$	and this is just the union of open balls $B(m_i, r_i)$ where $m_i := \frac{a_i+b_i}{2}$ is the midpoint and $r_i := \frac{b_i-a_i}{2}$ is
fong-spivak-seven-sketches F04016	333	■ 1.000	$r_i:=\frac{b_i-a_i}{2}$	$r_i := \frac{b_i - a_i}{2}$	and this is just the union of open balls $B(m_i, r_i)$ where $m_i := \frac{a_i+b_i}{2}$ is the midpoint and $r_i := \frac{b_i-a_i}{2}$ is
fong-spivak-seven-sketches F04017	333	■ 0.877	$\left(a_{i}, b_{i}\right)$	(a_i, b_i)	the radius of the interval (a_i, b_i) . The open balls $B(m_i, r_i)$ are open in the usual topology on $\mathbb{R}$ and the
fong-spivak-seven-sketches F04018	333	■ 1.000	$U=\bigcup_{j \in J} B\left(m_{j}, \epsilon_{j}\right)$	$U = \bigcup_{j \in J} B\left(m_j, \epsilon_j\right)$	Suppose that U is open in the usual topology. Then $U = \bigcup_{j \in J} B(m_j, \epsilon_j)$ for some set J . Let $a_j := m_j - \epsilon_j$
fong-spivak-seven-sketches F04019	333	■ 1.000	$a_j:=m_j-\epsilon_j$	$a_j := m_j - \epsilon_j$	Suppose that U is open in the usual topology. Then $U = \bigcup_{j \in J} B(m_j, \epsilon_j)$ for some set J . Let $a_j := m_j - \epsilon_j$
fong-spivak-seven-sketches F04020	333	■ 1.000	$b_j:=m_j+\epsilon_j$	$b_j := m_j + \epsilon_j$	and $b_j := m_j + \epsilon_j$. Then
fong-spivak-seven-sketches F04021	333	■ 0.950	$H_X(U):=$	$H_X(U) :=$	Fix any topological space $(X, \mathbf{Op}_X)$ and any subset $R \subseteq \mathbb{IR}$ of the interval domain. Define $H_X(U) :=$
fong-spivak-seven-sketches F04022	333	■ 1.000	$\{f: U \cap R \rightarrow X \mid f$	$\{f: U \cap R \rightarrow X \mid f$	$\{f: U \cap R \rightarrow X \mid f$ is continuous}.
fong-spivak-seven-sketches F04023	333	■ 1.000	$f: U \cap R \rightarrow X$	$f: U \cap R \rightarrow X$	1. H_X is a presheaf: given $V \subseteq U$ the restriction map sends the continuous function $f: U \cap R \rightarrow X$
fong-spivak-seven-sketches F04024	333	■ 1.000	$V \cap R \subseteq U \cap R$	$V \cap R \subseteq U \cap R$	to its restriction along the subset $V \cap R \subseteq U \cap R$.
fong-spivak-seven-sketches F04025	333	■ 1.000	$U=\bigcup_i U_i$	$U = \bigcup_i U_i$	2. It is a sheaf: given any family U_i of open sets with $U = \bigcup_i U_i$ and a continuous function
fong-spivak-seven-sketches F04026	333	■ 1.000	$f_i: U_i \cap R \rightarrow X$	$f_i: U_i \cap R \rightarrow X$	$f_i: U_i \cap R \rightarrow X$ for each i , agreeing on overlaps, they can be glued together to give a continuous
fong-spivak-seven-sketches F04027	333	■ 0.913	$U \cap R$	$U \cap R$	function on all of $U \cap R$, since $U \cap R = (\bigcup_i U_i) \cap R = \bigcup_i (U_i \cap R)$.
fong-spivak-seven-sketches F04028	333	■ 0.913	$U \cap R=\left(\bigcup_i U_i\right) \cap R=\bigcup_i\left(U_i \cap R\right)$	$U \cap R = \left(\bigcup_i U_i\right) \cap R = \bigcup_i (U_i \cap R)$	function on all of $U \cap R$, since $U \cap R = (\bigcup_i U_i) \cap R = \bigcup_i (U_i \cap R)$.