

bradley_spring22: equation evidence

Display equations

Identifier	Page	Conf.	LaTeX source	Rendered	Image
bradley_spring22_ EQ0005	13	■ 0.965	$\begin{array}{l} \left(\frac{1}{2}, \frac{1}{2}\right) \text{ and } (1,0) \text{ are elements of the set } \Delta_2, \\ \left(\frac{4}{10}, \frac{5}{10}, \frac{1}{10}\right) \text{ and } \left(\frac{1}{5}, \frac{3}{5}, \frac{1}{5}\right) \text{ are elements of the set } \Delta_3, \\ \left(\frac{2}{7}, \frac{1}{7}, 0, \frac{4}{7}\right) \text{ and } \left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}\right) \text{ are elements of the set } \Delta_4, \end{array}$	$\left(\frac{1}{2}, \frac{1}{2}\right) \text{ and } (1,0) \text{ are elements of the set } \Delta_2, \\ \left(\frac{4}{10}, \frac{5}{10}, \frac{1}{10}\right) \text{ and } \left(\frac{1}{5}, \frac{3}{5}, \frac{1}{5}\right) \text{ are elements of the set } \Delta_3, \\ \left(\frac{2}{7}, \frac{1}{7}, 0, \frac{4}{7}\right) \text{ and } \left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}\right) \text{ are elements of the set } \Delta_4,$	$\left(\frac{1}{2}, \frac{1}{2}\right) \text{ and } (1,0) \text{ are elements of the set } \Delta_2, \\ \left(\frac{4}{10}, \frac{5}{10}, \frac{1}{10}\right) \text{ and } \left(\frac{1}{5}, \frac{3}{5}, \frac{1}{5}\right) \text{ are elements of the set } \Delta_3, \\ \left(\frac{2}{7}, \frac{1}{7}, 0, \frac{4}{7}\right) \text{ and } \left(\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}\right) \text{ are elements of the set } \Delta_4,$
bradley_spring22_ EQ0016	28	■ 0.999	$d\left(p \circ (q^1, q^2, \dots, q^n)\right) \stackrel{\text{sort}}{=} d(p) + p_1 d(q^1) + p_2 d(q^2) + \dots + p_n d(q^n),$	$d\left(p \circ (q^1, q^2, \dots, q^n)\right) \stackrel{\text{sort}}{=} d(p) + p_1 d(q^1) + p_2 d(q^2) + \dots + p_n d(q^n),$	$d\left(p \circ (q^1, q^2, \dots, q^n)\right) \stackrel{\text{sort}}{=} d(p) + p_1 d(q^1) + p_2 d(q^2) + \dots + p_n d(q^n),$
bradley_spring22_ EQ0006 (3)	14	■ 0.999	$\text{Entropy defines a collection of functions } \{H_n: \Delta_n \rightarrow \mathbb{R}\}.$	$\text{Entropy defines a collection of functions } \{H_n: \Delta_n \rightarrow \mathbb{R}\}.$	$\text{Entropy defines a collection of functions } \{H_n: \Delta_n \rightarrow \mathbb{R}\}.$
bradley_spring22_ EQ0001 (1)	12	■ 1.000	$H(p) = -p_1 \log(p_1) - p_2 \log(p_2) - \dots - p_n \log(p_n).$	$H(p) = -p_1 \log(p_1) - p_2 \log(p_2) - \dots - p_n \log(p_n).$	$H(p) = -p_1 \log(p_1) - p_2 \log(p_2) - \dots - p_n \log(p_n).$
bradley_spring22_ EQ0002	12	■ 1.000	$H(p) = -1 \log(1) - 0 \log(0) = -\log(1) = 0.$	$H(p) = -1 \log(1) - 0 \log(0) = -\log(1) = 0.$	$H(p) = -1 \log(1) - 0 \log(0) = -\log(1) = 0.$
bradley_spring22_ EQ0003	12	■ 1.000	$H(p) = -\frac{1}{2} \log\left(\frac{1}{2}\right) - \frac{1}{2} \log\left(\frac{1}{2}\right) = -\log\left(\frac{1}{2}\right) = -\log(1) - (-\log(2)) = \log(2).$	$H(p) = -\frac{1}{2} \log\left(\frac{1}{2}\right) - \frac{1}{2} \log\left(\frac{1}{2}\right) = -\log\left(\frac{1}{2}\right) = -\log(1) - (-\log(2)) = \log(2).$	$H(p) = -\frac{1}{2} \log\left(\frac{1}{2}\right) - \frac{1}{2} \log\left(\frac{1}{2}\right) = -\log\left(\frac{1}{2}\right) = -\log(1) - (-\log(2)) = \log(2).$
bradley_spring22_ EQ0004 (2)	13	■ 1.000	$\text{For each natural number } n, \text{ there is a function } H: \Delta_n \rightarrow \mathbb{R}.$	$\text{For each natural number } n, \text{ there is a function } H: \Delta_n \rightarrow \mathbb{R}.$	$\text{For each natural number } n, \text{ there is a function } H: \Delta_n \rightarrow \mathbb{R}.$
bradley_spring22_ EQ0007	18	■ 1.000	$p \circ (q, r) = \left(\frac{1}{5}, \frac{1}{4}, \frac{1}{20}, \frac{3}{20}, \frac{7}{20}\right).$	$p \circ (q, r) = \left(\frac{1}{5}, \frac{1}{4}, \frac{1}{20}, \frac{3}{20}, \frac{7}{20}\right).$	$p \circ (q, r) = \left(\frac{1}{5}, \frac{1}{4}, \frac{1}{20}, \frac{3}{20}, \frac{7}{20}\right).$
bradley_spring22_ EQ0008 (4)	21	■ 1.000	$H(p \circ (q, r)) = H(p) + \frac{1}{2} H(q) + \frac{1}{2} H(r).$	$H(p \circ (q, r)) = H(p) + \frac{1}{2} H(q) + \frac{1}{2} H(r).$	$H(p \circ (q, r)) = H(p) + \frac{1}{2} H(q) + \frac{1}{2} H(r).$
bradley_spring22_ EQ0009 (5)	22	■ 1.000	$H(p \circ (q^1, q^2, \dots, q^n)) = H(p) + p_1 H(q^1) + p_2 H(q^2) + \dots + p_n H(q^n).$	$H(p \circ (q^1, q^2, \dots, q^n)) = H(p) + p_1 H(q^1) + p_2 H(q^2) + \dots + p_n H(q^n).$	$H(p \circ (q^1, q^2, \dots, q^n)) = H(p) + p_1 H(q^1) + p_2 H(q^2) + \dots + p_n H(q^n).$
bradley_spring22_ EQ0010 (6)	24	■ 1.000	$H(p \circ (q^1, q^2)) = H(p) + p_1 H(q^1) + p_2 H(q^2).$	$H(p \circ (q^1, q^2)) = H(p) + p_1 H(q^1) + p_2 H(q^2).$	$H(p \circ (q^1, q^2)) = H(p) + p_1 H(q^1) + p_2 H(q^2).$
bradley_spring22_ EQ0011 (7)	24	■ 1.000	$H(\bullet \bullet) = H(\bullet) + \bullet H(\bullet).$	$H(\bullet \bullet) = H(\bullet) + \bullet H(\bullet).$	$H(\bullet \bullet) = H(\bullet) + \bullet H(\bullet).$
bradley_spring22_ EQ0012 (8)	24	■ 1.000	$H(\bullet \bullet) = H(\bullet) \bullet + \bullet H(\bullet).$	$H(\bullet \bullet) = H(\bullet) \bullet + \bullet H(\bullet).$	$H(\bullet \bullet) = H(\bullet) \bullet + \bullet H(\bullet).$
bradley_spring22_ EQ0013 (9)	26	■ 1.000	entropy:	entropy:	entropy:

Identifier	Page	Conf.	LaTeX source	Rendered	Image
bradley_spring22_ EQ0014	26	■ 1.000	$d(\text{\texttt{\textbackslash bullet \textbackslash bullet}})=d(\text{\texttt{\textbackslash bullet}}) \text{\texttt{\textbackslash bullet}}+\text{\texttt{\textbackslash bullet}} d(\text{\texttt{\textbackslash bullet}})$	$d(\bullet\bullet) = d(\bullet)\bullet + \bullet d(\bullet)$	$d(\textcolor{violet}{\bullet}\textcolor{teal}{\bullet}) = d(\textcolor{teal}{\bullet})\textcolor{violet}{\bullet} + \textcolor{teal}{\bullet}d(\textcolor{violet}{\bullet})$
bradley_spring22_ EQ0015	26	■ 1.000	$H(\text{\texttt{\textbackslash bullet \textbackslash bullet}})=H(\text{\texttt{\textbackslash bullet}})+\text{\texttt{\textbackslash bullet}} H(\text{\texttt{\textbackslash bullet}})$	$H(\bullet\bullet) = H(\bullet) + \bullet H(\bullet)$	$H(\textcolor{violet}{\bullet}\textcolor{teal}{\bullet}) = H(\textcolor{teal}{\bullet}) + \textcolor{teal}{\bullet}H(\textcolor{violet}{\bullet})$