

mielke-geometroynamics: equation evidence

3 rows show a REFINEMENT below, not the OCR reading: 3 verified against the inferred. Each is marked [refined] beside its identifier. The model is unchanged — the original reading is still what ‘latex’ holds; this report chose the refinement for them. Rows: mielke-geometroynamics_EQ0378, mielke-geometroynamics_EQ0402, mielke-geometroynamics_EQ0972

Display equations

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometroynamics_313 EQ0966 (14.2.7) [conf 0.024]		■ 0.024	$\begin{aligned} d\hat{s}^2 &= \frac{1}{4+K} \operatorname{Tr} \left(\hat{\vartheta} \otimes_s \hat{\vartheta}^T \right) = \hat{g}_{AB} \hat{\vartheta}^A \otimes_s \hat{\vartheta}^B; \quad A, B, = 0, \dots, 3+K \\ &= \frac{1}{4+K} \operatorname{Tr} \left[\begin{array}{c c} \vartheta \otimes \vartheta + \hat{\ell}^2 \omega \otimes \omega & \hat{\ell} \omega \otimes \overset{\circ}{\vartheta} \\ \hline \hat{\ell} \vartheta \otimes \omega & \begin{smallmatrix} \cdot & \cdot \\ \vartheta \otimes \end{smallmatrix} \end{array} \right]. \end{aligned}$	$d\hat{s}^2 = \frac{1}{4+K} \operatorname{Tr} \left(\hat{\vartheta} \otimes_s \hat{\vartheta}^T \right) = \hat{g}_{AB} \hat{\vartheta}^A \otimes_s \hat{\vartheta}^B; \quad A, B, = 0, \dots, 3+K$ $= \frac{1}{4+K} \operatorname{Tr} \left[\begin{array}{c c} \vartheta \otimes \vartheta + \hat{\ell}^2 \omega \otimes \omega & \hat{\ell} \omega \otimes \overset{\circ}{\vartheta} \\ \hline \hat{\ell} \vartheta \otimes \omega & \begin{smallmatrix} \cdot & \cdot \\ \vartheta \otimes \vartheta \end{smallmatrix} \end{array} \right]. \quad (1.$	$d\hat{s}^2 = \frac{1}{4+K} \operatorname{Tr}(\hat{\vartheta} \otimes_s \hat{\vartheta}^T) = \hat{g}_{AB} \hat{\vartheta}^A \otimes_s \hat{\vartheta}^B; \quad A, B, = 0, \dots, 3+K$ $= \frac{1}{4+K} \operatorname{Tr} \left[\frac{\vartheta \otimes \vartheta + \hat{\ell}^2 \omega \otimes \omega}{\hat{\ell} \overset{\circ}{\vartheta} \otimes \omega} \middle \frac{\hat{\ell} \omega \otimes \overset{\circ}{\vartheta}}{\overset{\circ}{\vartheta} \otimes \overset{\circ}{\vartheta}} \right]. \quad (1.$
mielke-geometroynamics_134 EQ0403 (6.4.23) [conf 0.028]		■ 0.028	$\Psi_{\parallel}(\vartheta) = \exp \left(- \int_{M_3} \overset{(\pm)}{C}_{\text{TT}} \right) = \exp \left(\frac{-1}{2\ell^2} \int_{M_3} \left[\overset{*}{+}_{\vartheta_B} \wedge \overset{*}{T}_{\bar{T}}^B \right] \right)$	$\Psi_{\parallel}(\vartheta) = \exp \left(- \int_{M_3} \overset{(\pm)}{C}_{\text{TT}} \right) = \exp \left(\frac{-1}{2\ell^2} \int_{M_3} \left[\overset{*}{+}_{\vartheta_B} \wedge \overset{*}{T}_{\bar{T}}^B \right] \right)$	$\Psi_{\parallel}(\vartheta) = \exp \left(- \int_{M_3} \overset{(\pm)}{C}_{\text{TT}} \right) = \exp \left(\frac{-1}{2\ell^2} \int_{M_3} \left[\overset{*}{\vartheta}_B \wedge \overset{*}{T}^B \right] \right)$
mielke-geometroynamics_275 EQ0850 (12.3.3) [conf 0.032]		■ 0.032	$\begin{aligned} \not{D}^2 &= -\frac{1}{2} \gamma^\alpha \gamma^\beta \left(\{ D_\alpha^{\{\}}, D_\beta^{\{\}} \} + [D_\alpha^{\{\}}, D_\beta^{\{\}}] \right) - 2im \not{D}^{\}} \\ &\cong -\square - \frac{1}{8} \sigma^{\alpha\beta} R_{\alpha\beta\mu\nu}^{\{\}} \sigma^{\mu\nu} \\ &\quad - \frac{i}{4} \gamma_5 \left(\not{D}^{\{\}} \not{b} \right) + \frac{1}{2} \gamma_5 \sigma^{\alpha\beta} \mathscr{A}_\alpha D_\beta^{\{\}} - \frac{1}{16} \mathscr{A}_\alpha \mathscr{A}^\alpha - m^2 \end{aligned}$	$D^2 = -\frac{1}{2} \gamma^\alpha \gamma^\beta \left(\{ D_\alpha^{\{\}}, D_\beta^{\{\}} \} + [D_\alpha^{\{\}}, D_\beta^{\{\}}] \right) - 2im \not{D}^{\}} \\ - \frac{i}{4} \gamma_5 \left(\not{D}^{\{\}} \not{\mathscr{A}} \right) + \frac{1}{2} \gamma_5 \sigma^{\alpha\beta} \mathscr{A}_\alpha D_\beta^{\{\}} + m^2 - \frac{1}{2} m \gamma_5 \not{\mathscr{A}} - \frac{1}{16} \not{\mathscr{A}} \not{\mathscr{A}} \\ \cong -\square - \frac{1}{8} \sigma^{\alpha\beta} R_{\alpha\beta\mu\nu}^{\{\}} \sigma^{\mu\nu} \\ - \frac{i}{4} \gamma_5 \left(\not{D}^{\{\}} \not{b} \right) + \frac{1}{2} \gamma_5 \sigma^{\alpha\beta} \mathscr{A}_\alpha D_\beta^{\{\}} - \frac{1}{16} \mathscr{A}_\alpha \mathscr{A}^\alpha - m^2$	$\not{D}^2 = -\frac{1}{2} \gamma^\alpha \gamma^\beta \left(\{ D_\alpha^{\{\}}, D_\beta^{\{\}} \} + [D_\alpha^{\{\}}, D_\beta^{\{\}}] \right) - 2im \not{D}^{\{\}}$ $- \frac{i}{4} \gamma_5 (\not{D}^{\{\}} \not{\mathscr{A}}) + \frac{1}{2} \gamma_5 \sigma^{\alpha\beta} \mathscr{A}_\alpha D_\beta^{\{\}} + m^2 - \frac{1}{2} m \gamma_5 \not{\mathscr{A}} - \frac{1}{16} \not{\mathscr{A}} \not{\mathscr{A}}$ $\cong -\square - \frac{1}{8} \sigma^{\alpha\beta} R_{\alpha\beta\mu\nu}^{\{\}} \sigma^{\mu\nu}$ $- \frac{i}{4} \gamma_5 (\not{D}^{\{\}} \not{\mathscr{A}}) + \frac{1}{2} \gamma_5 \sigma^{\alpha\beta} \mathscr{A}_\alpha D_\beta^{\{\}} - \frac{1}{16} \mathscr{A}_\alpha \mathscr{A}^\alpha - m^2$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_134 EQ0402 (6.4.22) [conf 0.042] [refined: inferred]	134	■ 0.042	$\begin{array}{ll} \text{q: } & \stackrel{*}{\operatorname{\underline{\vartheta}}}\{\stackrel{*}{\operatorname{\underline{\vartheta}}}\}(\vartheta) = \stackrel{*}{\operatorname{\underline{\vartheta}}}\{\stackrel{*}{\operatorname{\underline{\vartheta}}}\}(\vartheta), \\ \text{p: } & \stackrel{*}{\Pi}_{\alpha} \Psi_{\parallel}(\vartheta) = -i\ell^2 \frac{\delta}{\delta \stackrel{*}{\vartheta}_{\alpha}} \Psi_{\parallel}(\vartheta), \end{array}$	$q : \quad \stackrel{*}{\vartheta}_{\parallel} \Psi_{\parallel}(\vartheta) = \stackrel{*}{\vartheta}_{\parallel} \Psi_{\parallel}(\vartheta),$ $\stackrel{\pm}{p}: \quad \stackrel{*}{\Pi}_{\alpha} \Psi_{\parallel}(\vartheta) = -i\ell^2 \frac{\delta}{\delta \stackrel{*}{\vartheta}_{\alpha}} \Psi_{\parallel}(\vartheta),$	$q : \quad \stackrel{*}{\vartheta}_{\parallel} \Psi_{\parallel}(\vartheta) = \stackrel{*}{\vartheta}_{\parallel} \Psi_{\parallel}(\vartheta),$ $\stackrel{\pm}{p}: \quad \stackrel{*}{\Pi}_{\alpha} \Psi_{\parallel}(\vartheta) = -i\ell^2 \frac{\delta}{\delta \stackrel{*}{\vartheta}_{\alpha}} \Psi_{\parallel}(\vartheta),$
mielke-geometrodynamics_101 EQ0293 (4.5.4) [conf 0.052]	101	■ 0.052	$\xi \downarrow L \equiv (\xi \downarrow \vartheta^{\alpha}) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + (\xi \downarrow \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\xi \downarrow D\Psi) \wedge \frac{\partial L}{\partial D\Psi}.$	$\xi \downarrow L \equiv (\xi \downarrow \vartheta^{\alpha}) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + (\xi \downarrow \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\xi \downarrow D\Psi) \wedge \frac{\partial L}{\partial D\Psi}.$	$\xi \downarrow L \equiv (\xi \downarrow \vartheta^{\alpha}) \wedge \frac{\partial L}{\partial \vartheta^{\alpha}} + (\xi \downarrow \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\xi \downarrow D\Psi) \wedge \frac{\partial L}{\partial D\Psi}.$
mielke-geometrodynamics_47 EQ0064 (2.8.6) [conf 0.054]	47	■ 0.054	$\begin{aligned} & \stackrel{*}{\operatorname{\underline{\vartheta}}}\{\stackrel{*}{\operatorname{\underline{\vartheta}}}\}(\vartheta) = \stackrel{*}{\operatorname{\underline{\vartheta}}}\{\stackrel{*}{\operatorname{\underline{\vartheta}}}\}(\vartheta), \\ & \stackrel{*}{\operatorname{\underline{\vartheta}}}\{\stackrel{*}{\operatorname{\underline{\vartheta}}}\}(\vartheta) = -i\ell^2 \frac{\delta}{\delta \stackrel{*}{\operatorname{\underline{\vartheta}}}\{\stackrel{*}{\operatorname{\underline{\vartheta}}}\}(\vartheta)} \Psi_{\parallel}(\vartheta), \end{aligned}$	$\stackrel{*}{\pi}(\gamma_1) = -\frac{1}{2\pi i} \operatorname{Tr} \Omega \quad (=0)$ $\stackrel{*}{\pi}(\gamma_2) = -\frac{1}{8\pi^2} \{ \operatorname{Tr}(\Omega \wedge \Omega) - \operatorname{Tr}(\Omega) \wedge \operatorname{Tr}(\Omega) \}$ $\left(= -\frac{1}{32\pi^2} \epsilon^{\alpha\beta\mu\nu} F_{\alpha\beta}^j F_{\mu\nu j} \sqrt{ g } d^4x \right)$ $\stackrel{*}{\pi}(\gamma_k) = 0 \text{ for } k \geq 3.$	$\stackrel{*}{\pi}(\gamma_1) = -\frac{1}{2\pi i} \operatorname{Tr} \Omega \quad (=0)$ $\stackrel{*}{\pi}(\gamma_2) = -\frac{1}{8\pi^2} \{ \operatorname{Tr}(\Omega \wedge \Omega) - \operatorname{Tr}(\Omega) \wedge \operatorname{Tr}(\Omega) \}$ $\left(= -\frac{1}{32\pi^2} \epsilon^{\alpha\beta\mu\nu} F_{\alpha\beta}^j F_{\mu\nu j} \sqrt{ g } d^4x \right)$ $\stackrel{*}{\pi}(\gamma_k) = 0 \text{ for } k \geq 3.$
mielke-geometrodynamics_133 EQ0400 (6.4.19) [conf 0.060]	133	■ 0.060	$\stackrel{(\pm)}{\mathcal{G}}_{\alpha} := \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} = \stackrel{(\pm)}{D} \stackrel{(\pm)}{\underline{A}_{\alpha}} = 0,$	$\stackrel{(\pm)}{\mathcal{G}}_{\alpha} := \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} = \stackrel{(\pm)}{D} \stackrel{(\pm)}{\underline{A}_{\alpha}} = 0,$	$\stackrel{(\pm)}{\mathcal{G}}_{\alpha} := \stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} = \stackrel{(\pm)}{D} \stackrel{(\pm)}{\underline{A}_{\alpha}} = 0,$
mielke-geometrodynamics_229 EQ0718 (10.9.4) [conf 0.066]	229	■ 0.066	$\begin{aligned} & \Sigma_{\alpha} := e_{\alpha} \downarrow L_{\Psi} - (e_{\alpha} \downarrow \Psi) \wedge \frac{\partial L_{\Psi}}{\partial \Psi} - (e_{\alpha} \downarrow D\Psi) \wedge \frac{\partial L_{\Psi}}{\partial D\Psi} \\ & \quad - (e_{\alpha} \downarrow \bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial \bar{\Psi}} - (e_{\alpha} \downarrow D\bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial D\bar{\Psi}}; \end{aligned}$	$\Sigma_{\alpha} := e_{\alpha} \downarrow L_{\Psi} - (e_{\alpha} \downarrow \Psi) \wedge \frac{\partial L_{\Psi}}{\partial \Psi} - (e_{\alpha} \downarrow \bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial \bar{\Psi}}$ $- (e_{\alpha} \downarrow D\Psi) \wedge \frac{\partial L_{\Psi}}{\partial D\Psi} - (e_{\alpha} \downarrow D\bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial D\bar{\Psi}};$	$\Sigma_{\alpha} := e_{\alpha} \downarrow L_{\Psi} - (e_{\alpha} \downarrow \Psi) \wedge \frac{\partial L_{\Psi}}{\partial \Psi} - (e_{\alpha} \downarrow \bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial \bar{\Psi}}$ $- (e_{\alpha} \downarrow D\Psi) \wedge \frac{\partial L_{\Psi}}{\partial D\Psi} - (e_{\alpha} \downarrow D\bar{\Psi}) \wedge \frac{\partial L_{\Psi}}{\partial D\bar{\Psi}};$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_168 EQ0529 (7.7.49) [conf 0.099]	168	■ 0.099	$\begin{aligned} L_{\text{Euler}} &:= (-1)^{\text{sig}+1} \text{Tr} \left\{ \Omega \wedge \Omega^{(*)} \right\} = \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= dC_{\text{RR}^{(*)}} = \frac{(-1)^{\text{sig}}}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta^{(*)}} - \frac{1}{3} \Gamma_{\alpha}^{\beta^{(*)}} \wedge \Gamma_{\beta}^{\gamma} \wedge \Gamma_{\gamma}^{\alpha} \right) \\ &\equiv \frac{1}{2} R_{\alpha\beta} \wedge {}^*R^{\alpha\beta} - 2 \text{Ric}_{\alpha\beta} \wedge {}^*\text{Ric}^{\alpha\beta} + \frac{1}{2} \text{Ric}_{\alpha}^{\alpha} \wedge {}^*\text{Ric}_{\beta}^{\beta} \\ &\equiv -L_{\text{SKY}} \\ &\quad - 2 \left(\text{Ric}_{\alpha\beta} \wedge {}^*\text{Ric}^{\alpha\beta} - \frac{1}{4} \text{Ric}_{\alpha}^{\alpha} \wedge {}^*\text{Ric}_{\beta}^{\beta} \right) \end{aligned}$	$\begin{aligned} L_{\text{Euler}} &:= (-1)^{\text{sig}+1} \text{Tr} \left\{ \Omega \wedge \Omega^{(*)} \right\} = \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= dC_{\text{RR}^{(*)}} = \frac{(-1)^{\text{sig}}}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta^{(*)}} - \frac{1}{3} \Gamma_{\alpha}^{\beta^{(*)}} \wedge \Gamma_{\beta}^{\gamma} \wedge \Gamma_{\gamma}^{\alpha} \right) \\ &\equiv \frac{1}{2} R_{\alpha\beta} \wedge {}^*R^{\alpha\beta} - 2 \text{Ric}_{\alpha\beta} \wedge {}^*\text{Ric}^{\alpha\beta} + \frac{1}{2} \text{Ric}_{\alpha}^{\alpha} \wedge {}^*\text{Ric}_{\beta}^{\beta} \\ &\equiv -L_{\text{SKY}} \\ &\quad - 2 \left(\text{Ric}_{\alpha\beta} \wedge {}^*\text{Ric}^{\alpha\beta} - \frac{1}{4} \text{Ric}_{\alpha}^{\alpha} \wedge {}^*\text{Ric}_{\beta}^{\beta} \right) \end{aligned}$	$\begin{aligned} L_{\text{Euler}} &:= (-1)^{\text{sig}+1} \text{Tr} \left\{ \Omega \wedge \Omega^{(*)} \right\} = \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= dC_{\text{RR}^{(*)}} = \frac{(-1)^{\text{sig}}}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta^{(*)}} - \frac{1}{3} \Gamma_{\alpha}^{\beta^{(*)}} \wedge \Gamma_{\beta}^{\gamma} \wedge \Gamma_{\gamma}^{\alpha} \right) \\ &\equiv \frac{1}{2} R_{\alpha\beta} \wedge {}^*R^{\alpha\beta} - 2 \text{Ric}_{\alpha\beta} \wedge {}^*\text{Ric}^{\alpha\beta} + \frac{1}{2} \text{Ric}_{\alpha}^{\alpha} \wedge {}^*\text{Ric}_{\beta}^{\beta} \\ &\equiv -L_{\text{SKY}} \\ &\quad - 2 \left(\text{Ric}_{\alpha\beta} \wedge {}^*\text{Ric}^{\alpha\beta} - \frac{1}{4} \text{Ric}_{\alpha}^{\alpha} \wedge {}^*\text{Ric}_{\beta}^{\beta} \right) \end{aligned}$
mielke-geometrodynamics_315 EQ0974 (14.2.15) [conf 0.099]	315	■ 0.099	$\Theta^{\perp} := \pi(\hat{\Theta} - [\hat{\alpha}, \hat{\vartheta}]) = \left[\frac{\Theta}{0} \middle \frac{\hat{\ell}\Omega'}{\pi\hat{\Theta} = 0} = 0 \right],$	$\Theta^{\perp} := \pi(\hat{\Theta} - [\hat{\alpha}, \hat{\vartheta}]) = \left[\frac{\Theta}{0} \middle \frac{\hat{\ell}\Omega'}{\pi\hat{\Theta} = 0} = 0 \right],$	$\Theta^{\perp} := \pi(\hat{\Theta} - [\hat{\alpha}, \hat{\vartheta}]) = \left[\frac{\Theta}{0} \middle \frac{\hat{\ell}\Omega'}{\pi\hat{\Theta} = 0} = 0 \right],$
mielke-geometrodynamics_180 EQ0553 (8.4.7)	180	■ 0.100	$\begin{aligned} L_{\text{SKY}}^{(*)} &= \frac{1}{2} \text{Tr} \left(\overset{\pm}{\Omega}^g \wedge {}^* \overset{\pm}{\Omega}^g \right) \\ &= -\frac{1}{2} R^{\alpha\beta} \wedge {}^*R_{\alpha\beta} \mp \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= -\frac{1}{4} \left(R_{\alpha\beta} \pm {}^*R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} \pm {}^*R^{\alpha\beta^{(*)}} \right). \end{aligned}$	$\begin{aligned} L_{\text{SKY}}^{(*)} &= \frac{1}{2} \text{Tr} \left(\overset{\pm}{\Omega}^g \wedge {}^* \overset{\pm}{\Omega}^g \right) \\ &= -\frac{1}{2} R^{\alpha\beta} \wedge {}^*R_{\alpha\beta} \mp \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= -\frac{1}{4} \left(R_{\alpha\beta} \pm {}^*R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} \pm {}^*R^{\alpha\beta^{(*)}} \right). \end{aligned}$	$\begin{aligned} L_{\text{SKY}}^{(*)} &= \frac{1}{2} \text{Tr} \left(\overset{\pm}{\Omega}^g \wedge {}^* \overset{\pm}{\Omega}^g \right) \\ &= -\frac{1}{2} R^{\alpha\beta} \wedge {}^*R_{\alpha\beta} \mp \frac{(-1)^{\text{sig}}}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}^{(*)} \\ &= -\frac{1}{4} \left(R_{\alpha\beta} \pm {}^*R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} \pm {}^*R^{\alpha\beta^{(*)}} \right). \end{aligned}$
mielke-geometrodynamics_104 EQ0311 (4.6.3)	104	■ 0.111	$\begin{aligned} \widetilde{\leftarrow}_{\widetilde{y}} \widetilde{D} \widetilde{\xi}^{\alpha} &= \widetilde{y} \rfloor \left(\widetilde{D} \widetilde{D} \widetilde{\xi}^{\alpha} \right) + \widetilde{D}(\widetilde{y} \rfloor \widetilde{D} \widetilde{\xi}^{\alpha}) \\ &= y \rfloor \left(D D \xi^{\alpha} + R^{(T)\alpha} \right) = 0, \end{aligned}$	$\begin{aligned} \widetilde{\leftarrow}_{\widetilde{y}} \widetilde{D} \widetilde{\xi}^{\alpha} &= \widetilde{y} \rfloor \left(\widetilde{D} \widetilde{D} \widetilde{\xi}^{\alpha} \right) + \widetilde{D}(\widetilde{y} \rfloor \widetilde{D} \widetilde{\xi}^{\alpha}) \\ &= y \rfloor (D D \xi^{\alpha} + R^{(T)\alpha}) = 0, \end{aligned}$	$\begin{aligned} \widetilde{\leftarrow}_{\widetilde{y}} \widetilde{D} \widetilde{\xi}^{\alpha} &= \widetilde{y} \rfloor (\widetilde{D} \widetilde{D} \widetilde{\xi}^{\alpha}) + \widetilde{D}(\widetilde{y} \rfloor \widetilde{D} \widetilde{\xi}^{\alpha}) \\ &= y \rfloor (D D \xi^{\alpha} + R^{(T)\alpha}) = 0, \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometroynamics_128 EQ0378 (6.2.29) [re-fined: inferred]	128	■ 0.136	preserve the term Pontryagin index for the classification of real associated vector bundles (\textsc{Mayer} \& \textsc{Drechsler} 1977). Nevertheless, the denotation of \textsc{Belavin}	preserve the term Pontryagin index for the classification of real associated vector bundles (Mayer&Drechsler1977). Nevertheless, the denotation of Belavin	$DD\vartheta^\alpha = R_\beta{}^\alpha \wedge \vartheta^\beta = 0, \ D \ D \overline{\Pi}_\alpha = -R_\alpha{}^\beta \wedge \overline{\Pi}_\beta = \pm \frac{i}{2} \ell^2 (e_\alpha \rfloor \overline{\Pi}^\beta) \wedge \Sigma_\beta \cong 0.$
mielke-geometroynamics_346 EQ1059 (15.2.16)	346	■ 0.147	\begin{aligned} \psi = & \left\{ \begin{array}{lll l} \mathcal{P}_a & \mathcal{P}_b & \mathcal{P}_c & \nu_e \\ \eta_a & \eta_b & \eta_c & e^- \\ \lambda_a & \lambda_b & \lambda_c & \mu^- \\ \hline c_a & c_b & c_c & \nu_\mu \end{array} \right. \begin{array}{l} \text{up} \\ \text{down} \\ \text{strange} \\ \text{charm} \end{array} \begin{array}{l} \uparrow \\ \downarrow \end{array} \text{flavors} \\ & \text{red} \quad \text{yellow} \quad \text{blue} \quad \text{lilac} \end{aligned}	$\psi = \left[\begin{array}{ccc c} \mathcal{P}_a & \mathcal{P}_b & \mathcal{P}_c & \nu_e \\ \eta_a & \eta_b & \eta_c & e^- \\ \lambda_a & \lambda_b & \lambda_c & \mu^- \\ \hline c_a & c_b & c_c & \nu_\mu \end{array} \right] \begin{array}{l} \text{up} \\ \text{down} \\ \text{strange} \\ \text{charm} \end{array} \begin{array}{l} \uparrow \\ \downarrow \end{array} \text{flavors}$ red yellow blue lilac	$\psi = \left[\begin{array}{ccc c} \mathcal{P}_a & \mathcal{P}_b & \mathcal{P}_c & \nu_e \\ \eta_a & \eta_b & \eta_c & e^- \\ \lambda_a & \lambda_b & \lambda_c & \mu^- \\ \hline c_a & c_b & c_c & \nu_\mu \end{array} \right] \begin{array}{l} \text{up} \\ \text{down} \\ \text{strange} \\ \text{charm} \end{array} \begin{array}{l} \uparrow \\ \downarrow \end{array} \text{flavors}$ red yellow blue lilac
mielke-geometroynamics_314 EQ0972 (14.2.13) [re-fined: inferred]	314	■ 0.153	\overset{(1)}{D}\alpha := \overset{(0)}{d}\alpha - \omega^g \wedge \alpha - \alpha \wedge \omega^g	$\overset{(1)}{D}\alpha := \overset{(0)}{d}\alpha - \omega^g \wedge \alpha - \alpha \wedge \omega^g$	$\overset{(\circ)}{D}\alpha := \overset{(\circ)}{d}\alpha - \omega^g \wedge \alpha - \alpha \wedge \overset{\circ}{\omega}$
mielke-geometroynamics_95 EQ0264 (4.3.29)	95	■ 0.154	\Gamma_{ij} = \left\{ \begin{array}{c} \kappa \\ ij \end{array} \right\} - K_{ij}^\kappa,	$\Gamma_{ij}' = \left\{ \begin{array}{c} \kappa \\ ij \end{array} \right\} - K_{ij}^\kappa,$	$\Gamma_{ij}^{\cdot,\kappa} = \left\{ \begin{array}{c} \kappa \\ ij \end{array} \right\} - K_{ij}^{\cdot,\kappa},$
mielke-geometroynamics_100 EQ0292 (4.5.3)	100	■ 0.165	\begin{aligned} \mathfrak{f}_\xi L = & \left(\mathfrak{f}_\xi g_{\alpha\beta} \right) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \left(\mathfrak{f}_\xi \vartheta^\alpha \right) \wedge \frac{\partial L}{\partial \vartheta^\alpha} \\ & + \left(\mathfrak{f}_\xi \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\mathfrak{f}_\xi D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \\ & = D \left[\left(\xi \rfloor \vartheta^\alpha \right) \wedge \frac{\partial L}{\partial \vartheta^\alpha} + \left(\xi \rfloor \Psi \right) \wedge \frac{\partial L}{\partial \Psi} + \left(\xi \rfloor D\Psi \right) \wedge \frac{\partial L}{\partial D\Psi} \right] \\ & - \left(\xi \rfloor \vartheta^\alpha \right) D \frac{\partial L}{\partial \vartheta^\alpha} + \left(\xi \rfloor T^\alpha \right) \wedge \frac{\partial L}{\partial \vartheta^\alpha} + \left(\xi \rfloor R_\beta^\gamma \right) \wedge I^\beta{}_\gamma \Psi \wedge \frac{\partial L}{\partial D\Psi} \\ & + \left(\xi \rfloor D\Psi \right) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p (\xi \rfloor \Psi) \wedge D \frac{\delta L}{\delta \Psi} . \end{aligned}	$\mathfrak{f}_\xi L = (\mathfrak{f}_\xi g_{\alpha\beta}) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + (\mathfrak{f}_\xi \vartheta^\alpha) \wedge \frac{\partial L}{\partial \vartheta^\alpha} + (\mathfrak{f}_\xi \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\mathfrak{f}_\xi D\Psi) \wedge \frac{\partial L}{\partial D\Psi} = D \left[(\xi \rfloor \vartheta^\alpha) \wedge \frac{\partial L}{\partial \vartheta^\alpha} + (\xi \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\xi \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} \right] - (\xi \rfloor \vartheta^\alpha) D \frac{\partial L}{\partial \vartheta^\alpha} + (\xi \rfloor T^\alpha) \wedge \frac{\partial L}{\partial \vartheta^\alpha} + (\xi \rfloor R_\beta^\gamma) \wedge I^\beta{}_\gamma \Psi \wedge \frac{\partial L}{\partial D\Psi} + (\xi \rfloor D\Psi) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p (\xi \rfloor \Psi) \wedge D \frac{\delta L}{\delta \Psi} .$	$\mathfrak{f}_\xi L = (\mathfrak{f}_\xi g_{\alpha\beta}) \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + (\mathfrak{f}_\xi \vartheta^\alpha) \wedge \frac{\partial L}{\partial \vartheta^\alpha} + (\mathfrak{f}_\xi \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\mathfrak{f}_\xi D\Psi) \wedge \frac{\partial L}{\partial D\Psi} = D \left[(\xi \rfloor \vartheta^\alpha) \wedge \frac{\partial L}{\partial \vartheta^\alpha} + (\xi \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi} + (\xi \rfloor D\Psi) \wedge \frac{\partial L}{\partial D\Psi} \right] - (\xi \rfloor \vartheta^\alpha) D \frac{\partial L}{\partial \vartheta^\alpha} + (\xi \rfloor T^\alpha) \wedge \frac{\partial L}{\partial \vartheta^\alpha} + (\xi \rfloor R_\beta^\gamma) \wedge I^\beta{}_\gamma \Psi \wedge \frac{\partial L}{\partial D\Psi} + (\xi \rfloor D\Psi) \wedge \frac{\delta L}{\delta \Psi} + (-1)^p (\xi \rfloor \Psi) \wedge D \frac{\delta L}{\delta \Psi} .$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00734 (11.1.1)	237	■ 0.220	$\pi_1\left(SO_0(s, n-s), e\right)=\begin{cases}\mathbb{Z}_2 & n=2k \\ \mathbb{Z}_2\oplus\mathbb{Z}_2 & n=4k \\ \mathbb{Z}_4 & n=2(2k+1)\end{cases}$	$\pi_1(SO_0(s, n-s), e) = \begin{cases} \mathbb{Z}_2 & n=2k \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2 & n=4k \\ \mathbb{Z}_4 & n=2(2k+1) \end{cases}$	$\pi_1(SO_0(s, n-s), e) = \begin{cases} \mathbb{Z}_2 & n=2k \\ \mathbb{Z}_2 \oplus \mathbb{Z}_2 & n=4k \\ \mathbb{Z}_4 & n=2(2k+1). \end{cases}$
mielke-geometrodynamics_E00961 (14.2.1)	312	■ 0.220	$\hat{\vartheta}=\hat{\vartheta}(m, g)=\underbrace{\left[\begin{array}{c c} \vartheta & \hat{\ell}\omega \\ \hline 0 & \vartheta \end{array}\right]}_4\}^4 K.$	$\hat{\vartheta}=\hat{\vartheta}(m, g)=\underbrace{\left[\begin{array}{c c} \vartheta & \hat{\ell}\omega \\ \hline 0 & \vartheta \end{array}\right]}_4\}^4 K.$	$\hat{\vartheta}=\hat{\vartheta}(m, g)=\underbrace{\left[\begin{array}{c c} \vartheta & \hat{\ell}\omega \\ \hline 0 & \vartheta \end{array}\right]}_4\}^4 K.$
mielke-geometrodynamics_E00211 (4.2.3)	85	■ 0.223	$\begin{aligned} &*\widetilde{\Omega}=\chi\left(d\widetilde{\omega}-\frac{1}{2}[\widetilde{\omega},\widetilde{\omega}]\right) \\ &=d\omega^L-\omega^L\wedge\omega^L+d\omega^T-\omega^T\wedge\omega^L-\omega^L\wedge\omega^T+\frac{1}{2}[\omega^T,\omega^T] \\ &=\Omega^L+\Omega^T \end{aligned}$	$\begin{aligned} *\widetilde{\Omega} &= \chi \left(d\widetilde{\omega} - \frac{1}{2}[\widetilde{\omega}, \widetilde{\omega}] \right) \\ &= d\omega^L - \omega^L \wedge \omega^L + d\omega^T - \omega^T \wedge \omega^L - \omega^L \wedge \omega^T + \frac{1}{2}[\omega^T, \omega^T] \\ &= \Omega^L + \Omega^T \end{aligned}$	$\begin{aligned} *\widetilde{\Omega} &= \chi(d\widetilde{\omega} - \frac{1}{2}[\widetilde{\omega}, \widetilde{\omega}]) \\ &= d\omega^L - \omega^L \wedge \omega^L + d\omega^T - \omega^T \wedge \omega^L - \omega^L \wedge \omega^T + \frac{1}{2}[\omega^T, \omega^T] \\ &= \Omega^L + \Omega^T \end{aligned}$
mielke-geometrodynamics_E00266 (4.3.31)	95	■ 0.227	$R_{ijc}^{:d}::}=R_{ijc}^{\{!\dots d\}-2D_{[i}K_{j]c}^{:d}::}+2K_{[i}^{:h}::}K_{j]h}^{:d}::}.$	$R_{ijc}^{:d}::}=R_{ijc}^{\{!\dots d\}-2D_{[i}K_{j]c}^{:d}::}+2K_{[i}^{:h}::}K_{j]h}^{:d}::}.$	$R_{ijc}^{:d}::}=R_{ijc}^{\{!\dots d\}-2D_{[i}K_{j]c}^{:d}::}+2K_{[i}^{:h}::}K_{j]h}^{:d}::}.$
mielke-geometrodynamics_E00286 (4.4.15)	99	■ 0.229	$D\Pi_{\square}^T-E_{\square}=\Sigma_{\square}$	$D\Pi_{\square}^T-E_{\square}=\Sigma_{\square}$	$D\Pi_{\square}^T-E_{\square}=\Sigma_{\square}$
mielke-geometrodynamics_E00337 (5.3.14)	114	■ 0.239	$\begin{aligned} &R^{\{\alpha\beta\}}\wedge\eta_{\alpha\beta}\equiv R^{\alpha\beta}\wedge\eta_{\alpha\beta}-K^{\alpha\mu}\wedge K_{\mu}^{\beta}\wedge\eta_{\alpha\beta}+K^{\alpha\beta}\wedge T^{\gamma}\wedge\eta_{\alpha\beta\gamma} \\ &\quad +d\left(K^{\alpha\beta}\wedge\eta_{\alpha\beta}\right) \\ &=R^{\alpha\beta}\wedge\eta_{\alpha\beta}+T^{\alpha}\wedge\left(-^{(1)}T_{\alpha}+2^{(2)}T_{\alpha}+\frac{1}{2}^{(3)}T_{\alpha}\right)+2dC_{TT}^* \end{aligned}$	$\begin{aligned} R^{\{\alpha\beta\}}\wedge\eta_{\alpha\beta} &\equiv R^{\alpha\beta}\wedge\eta_{\alpha\beta}-K^{\alpha\mu}\wedge K_{\mu}^{\beta}\wedge\eta_{\alpha\beta}+K^{\alpha\beta}\wedge T^{\gamma}\wedge\eta_{\alpha\beta\gamma} \\ &\quad +d\left(K^{\alpha\beta}\wedge\eta_{\alpha\beta}\right) \\ &=R^{\alpha\beta}\wedge\eta_{\alpha\beta}+T^{\alpha}\wedge\left(-^{(1)}T_{\alpha}+2^{(2)}T_{\alpha}+\frac{1}{2}^{(3)}T_{\alpha}\right)+2dC_{TT}^* \end{aligned}$	$\begin{aligned} R^{\{\alpha\beta\}}\wedge\eta_{\alpha\beta} &\equiv R^{\alpha\beta}\wedge\eta_{\alpha\beta}-K^{\alpha\mu}\wedge K_{\mu}^{\beta}\wedge\eta_{\alpha\beta}+K^{\alpha\beta}\wedge T^{\gamma}\wedge\eta_{\alpha\beta\gamma} \\ &\quad +d(K^{\alpha\beta}\wedge\eta_{\alpha\beta}) \\ &= R^{\alpha\beta}\wedge\eta_{\alpha\beta}+T^{\alpha}\wedge\left(-^{(1)}T_{\alpha}+2^{(2)}T_{\alpha}+\frac{1}{2}^{(3)}T_{\alpha}\right)+2dC_{TT}^* \end{aligned} \tag{5.3.}$
mielke-geometrodynamics_E00835 (12.2.3)	271	■ 0.247	$\begin{aligned} &L_{\mathrm{D}}=L\left(\gamma,\psi,D^{(f)}\psi\right)-\frac{i}{2}\overline{\psi}\left(*\gamma\wedge K-K\wedge*\gamma\right)\psi+A\wedge j \\ &=L\left(\gamma,\psi,D^{(\dagger)}\psi\right)+\frac{1}{4}\mathscr{A}\wedge j_5+A\wedge j \\ &=L\left(\gamma,\psi,D^{(i)}\psi\right)-T^{\alpha}\wedge\mu_{\alpha}+A\wedge j. \end{aligned}$	$\begin{aligned} L_{\mathrm{D}} &= L\left(\gamma,\psi,D^{(f)}\psi\right)-\frac{i}{2}\overline{\psi}\left(*\gamma\wedge K-K\wedge*\gamma\right)\psi+A\wedge j \\ &= L\left(\gamma,\psi,D^{(\dagger)}\psi\right)+\frac{1}{4}\mathscr{A}\wedge j_5+A\wedge j \\ &= L\left(\gamma,\psi,D^{(i)}\psi\right)-T^{\alpha}\wedge\mu_{\alpha}+A\wedge j. \end{aligned}$	$\begin{aligned} L_{\mathrm{D}} &= L(\gamma,\psi,D^{\dagger}\psi)-\frac{i}{2}\overline{\psi}\left(*\gamma\wedge K-K\wedge*\gamma\right)\psi+A\wedge j \\ &= L(\gamma,\psi,D^{\dagger}\psi)+\frac{1}{4}\mathscr{A}\wedge j_5+A\wedge j \\ &= L(\gamma,\psi,D^{\dagger}\psi)-T^{\alpha}\wedge\mu_{\alpha}+A\wedge j. \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_009527 (7.7.47)	168	■ 0.282	$\begin{aligned} L_{\mathrm{Pontr}} := & \, d C_{\mathrm{RR}} = \frac{1}{2} R_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha} \wedge R_{\alpha\beta}^{\{\}} \wedge R^{\{\}}_{\alpha\beta} \\ & - \frac{1}{12} d \left[{}^* \mathcal{A} \wedge R^{\{\}} - \frac{1}{3} \mathcal{A} \wedge d \mathcal{A} + \frac{1}{9} {}^* \mathcal{A} \wedge {}^* (\mathcal{A} \wedge {}^* \mathcal{A}) \right]. \end{aligned}$	$L_{\mathrm{Pontr}} := d C_{\mathrm{RR}} = \frac{1}{2} R_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha} = -\frac{1}{2} R_{\alpha\beta}^{\{\}} \wedge R^{\{j\alpha\beta}$ $- \frac{1}{12} d \left[{}^* \mathcal{A} \wedge R^{\{\}} - \frac{1}{3} \mathcal{A} \wedge d \mathcal{A} + \frac{1}{9} {}^* \mathcal{A} \wedge {}^* (\mathcal{A} \wedge {}^* \mathcal{A}) \right].$	$L_{\mathrm{Pontr}} := d C_{\mathrm{RR}} = \frac{1}{2} R_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha} = -\frac{1}{2} R_{\alpha\beta}^{\{\}} \wedge R^{\{\}}_{\alpha\beta} \\ - \frac{1}{12} d \left[{}^* \mathcal{A} \wedge R^{\{\}} - \frac{1}{3} \mathcal{A} \wedge d \mathcal{A} + \frac{1}{9} {}^* \mathcal{A} \wedge {}^* (\mathcal{A} \wedge {}^* \mathcal{A}) \right].$
mielke-geometrodynamics_00942 (14.1.23)	306	■ 0.287	$\xi \wedge {}^* \xi = \pm \frac{\Omega \wedge {}^* \Omega \operatorname{Tr}^* (\Omega \wedge {}^* \Omega) - (-1)^{2z} \Omega \wedge \Omega \operatorname{Tr}^* (\Omega \wedge \Omega)}{\sqrt{(\operatorname{Tr}^* (\Omega \wedge {}^* \Omega))^2 + (-1)^{2z} (\operatorname{Tr}^* (\Omega \wedge \Omega))^2}}.$	$\xi \wedge {}^* \xi = \pm \frac{\Omega \wedge {}^* \Omega \operatorname{Tr}^* (\Omega \wedge {}^* \Omega) - (-1)^{2z} \Omega \wedge \Omega \operatorname{Tr}^* (\Omega \wedge \Omega)}{\sqrt{(\operatorname{Tr}^* (\Omega \wedge {}^* \Omega))^2 + (-1)^{2z} (\operatorname{Tr}^* (\Omega \wedge \Omega))^2}}.$	$\xi \wedge {}^* \xi = \pm \frac{\Omega \wedge {}^* \Omega \operatorname{Tr}^* (\Omega \wedge {}^* \Omega) - (-1)^{2z} \Omega \wedge \Omega \operatorname{Tr}^* (\Omega \wedge \Omega)}{\sqrt{(\operatorname{Tr}^* (\Omega \wedge {}^* \Omega))^2 + (-1)^{2z} (\operatorname{Tr}^* (\Omega \wedge \Omega))^2}}.$
mielke-geometrodynamics_00440 (6.8.11)	143	■ 0.288	$\left. D \Sigma_{\alpha} \cong (e_{\alpha} \rfloor T^{\gamma}) \wedge \Sigma_{\gamma} + (e_{\alpha} \rfloor R^{\gamma\delta}) \wedge \tau_{\gamma\delta} \right.$	$D\Sigma_{\alpha} \cong (e_{\alpha} \rfloor T^{\gamma}) \wedge \Sigma_{\gamma} + (e_{\alpha} \rfloor R^{\gamma\delta}) \wedge \tau_{\gamma\delta}$	$D\Sigma_{\alpha} \cong (e_{\alpha} \rfloor T^{\gamma}) \wedge \Sigma_{\gamma} + (e_{\alpha} \rfloor R^{\gamma\delta}) \wedge \tau_{\gamma\delta}$
mielke-geometrodynamics_00315 (4.6.7)	105	■ 0.292	$\Delta \Phi(\tilde{\Gamma}, \gamma) \simeq \frac{i}{\hbar} \int_S \left(R^{(T)\alpha} P_{\alpha} + R_{\alpha}^{(L)\beta} L^{\alpha}_{\beta} \right)$	$\Delta \Phi(\tilde{\Gamma}, \gamma) \simeq \frac{i}{\hbar} \int_S \left(R^{(T)\alpha} P_{\alpha} + R_{\alpha}^{(L)\beta} L^{\alpha}_{\beta} \right)$	$\Delta \Phi(\tilde{\Gamma}, \gamma) \simeq \frac{i}{\hbar} \int_S (R^{(T)\alpha} P_{\alpha} + R_{\alpha}^{(L)\beta} L^{\alpha}_{\beta}).$
mielke-geometrodynamics_00965 (14.2.5)	313	■ 0.299	$\begin{aligned} & \stackrel{\circ}{\Theta} := \dot{\stackrel{\circ}{\vartheta}} - [\stackrel{\circ}{\omega}, \stackrel{\circ}{\vartheta}] = (1 - 2b) \stackrel{\circ}{\vartheta} \wedge \stackrel{\circ}{\vartheta}, \\ & \stackrel{\circ}{\Omega} := \dot{\stackrel{\circ}{\omega}} - \stackrel{\circ}{\omega} \wedge \stackrel{\circ}{\omega} = b(1 - b) \stackrel{\circ}{\vartheta} \wedge \stackrel{\circ}{\vartheta}. \end{aligned}$	$\begin{aligned} \stackrel{\circ}{\Theta} &:= \dot{\stackrel{\circ}{\vartheta}} - [\stackrel{\circ}{\omega}, \stackrel{\circ}{\vartheta}] = (1 - 2b) \stackrel{\circ}{\vartheta} \wedge \stackrel{\circ}{\vartheta}, \\ \stackrel{\circ}{\Omega} &:= \dot{\stackrel{\circ}{\omega}} - \stackrel{\circ}{\omega} \wedge \stackrel{\circ}{\omega} = b(1 - b) \stackrel{\circ}{\vartheta} \wedge \stackrel{\circ}{\vartheta}. \end{aligned}$	$\begin{aligned} \stackrel{\circ}{\Theta} &:= \dot{\stackrel{\circ}{\vartheta}} - [\stackrel{\circ}{\omega}, \stackrel{\circ}{\vartheta}] = (1 - 2b) \stackrel{\circ}{\vartheta} \wedge \stackrel{\circ}{\vartheta}, \\ \stackrel{\circ}{\Omega} &:= \dot{\stackrel{\circ}{\omega}} - \stackrel{\circ}{\omega} \wedge \stackrel{\circ}{\omega} = b(1 - b) \stackrel{\circ}{\vartheta} \wedge \stackrel{\circ}{\vartheta}. \end{aligned}$
mielke-geometrodynamics_00226 (4.2.19)	87	■ 0.299	$\begin{aligned} A^{-1} \omega^{\mathrm{T}} &= A^{-1} \tilde{\vartheta} - G^{-1} D G A^{-1} \tilde{\varphi} \\ &= G^{-1} \tilde{\vartheta} - G^{-1} D \tilde{\varphi}, \end{aligned}$	$\begin{aligned} A^{-1} \omega^{\mathrm{T}} &= A^{-1} \tilde{\vartheta} - G^{-1} D G A^{-1} \tilde{\varphi} \\ &= G^{-1} \tilde{\vartheta} - G^{-1} D \tilde{\varphi}, \end{aligned}$	$\begin{aligned} A^{-1} \omega^{\mathrm{T}} &= A^{-1} \tilde{\vartheta} - G^{-1} D G A^{-1} \tilde{\varphi} \\ &= G^{-1} \tilde{\vartheta} - G^{-1} D \tilde{\varphi}, \end{aligned}$
mielke-geometrodynamics_00407 (6.4.27)	135	■ 0.315	$\Psi_{\Lambda}(A) = \exp \left(\frac{3}{\Lambda} \int_{M_3} {}^{(\pm)} \underline{C}_{\mathrm{RR}}^{\{\}} \right),$	$\Psi_{\Lambda}(A) = \exp \left(\frac{3}{\Lambda} \int_{M_3} {}^{(\pm)} \underline{C}_{\mathrm{RR}}^{\{\}} \right),$	$\Psi_{\Lambda}(A) = \exp \left(\frac{3}{\Lambda} \int_{M_3} {}^{(\pm)} \underline{C}_{\mathrm{RR}}^{\{\}} \right),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometroynamics_EQ0196 (4.1.8)	82	■ 0.322	$\begin{aligned} & \left[P_{\alpha}, P_{\beta} = 0 \right] \quad (P_{\alpha} := \partial / \partial x^{\alpha}), \\ & \left[P_{\alpha}, L_{\gamma \delta} \right] = g_{\alpha \delta} P_{\gamma} - g_{\alpha \gamma} P_{\delta}, \\ & \left[L_{\alpha \beta}, L_{\gamma \delta} \right] = c_{\alpha \beta \gamma \delta}^{\varepsilon} L_{\varepsilon \gamma}, \\ & c_{\alpha \beta \delta \gamma}^{\varepsilon \zeta} := \delta_{(\alpha}^{\varepsilon} \delta_{\delta}^{\zeta]} \eta_{\beta \gamma)}. \end{aligned}$	$\begin{aligned} & [P_{\alpha}, P_{\beta} = 0] \quad (P_{\alpha} := \partial / \partial x^{\alpha}), \\ & [P_{\alpha}, L_{\gamma \delta}] = g_{\alpha \delta} P_{\gamma} - g_{\alpha \gamma} P_{\delta}, \\ & [L_{\alpha \beta}, L_{\gamma \delta}] = c_{\alpha \beta \gamma \delta}^{\varepsilon} L_{\varepsilon \gamma}, \\ & c_{\alpha \beta \delta \gamma}^{\varepsilon \zeta} := \delta_{(\alpha}^{\varepsilon} \delta_{\delta}^{\zeta]} \eta_{\beta \gamma)}. \end{aligned}$	$\begin{aligned} & [P_{\alpha}, P_{\beta} = 0] \quad (P_{\alpha} := \partial / \partial x^{\alpha}), \\ & [P_{\alpha}, L_{\gamma \delta}] = g_{\alpha \delta} P_{\gamma} - g_{\alpha \gamma} P_{\delta}, \\ & [L_{\alpha \beta}, L_{\gamma \delta}] = c_{\alpha \beta \gamma \delta}^{\varepsilon} L_{\varepsilon \gamma}, \\ & c_{\alpha \beta \delta \gamma}^{\varepsilon \zeta} := \delta_{(\alpha}^{\varepsilon} \delta_{\delta}^{\zeta]} \eta_{\beta \gamma)}. \end{aligned}$
mielke-geometroynamics_EQ0597 (9.1.15)	196	■ 0.323	$\left[\mathrm{T}_{\alpha \beta \gamma} \right] = \left[\begin{array}{cccc} f & -h & \cdot & \cdot \\ \cdot & \cdot & -k & \cdot \\ \cdot & \cdot & \cdot & -k \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & -g \\ \cdot & \cdot & g & \cdot \\ 0 & 1 & 2 & 3 \end{array} \right] \begin{array}{l} 01 \\ 02 \\ 03 \\ 23 \\ 31 \\ 12 \end{array}$	$[T_{\alpha \beta}{}^{\gamma}] = \begin{bmatrix} f & -h & \cdot & \cdot & 01 \\ \cdot & \cdot & -k & \cdot & 02 \\ \cdot & \cdot & \cdot & -k & 03 \\ \cdot & \cdot & \cdot & \cdot & 23 \\ \cdot & \cdot & \cdot & -g & 31 \\ \cdot & \cdot & g & \cdot & 12 \\ 0 & 1 & 2 & 3 & \end{bmatrix}$	$[T_{\alpha \beta}{}^{\gamma}] = \begin{bmatrix} f & -h & \cdot & \cdot \\ \cdot & \cdot & -k & \cdot \\ \cdot & \cdot & \cdot & -k \\ \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & -g \\ \cdot & \cdot & g & \cdot \\ 0 & 1 & 2 & 3 \end{bmatrix} \begin{array}{l} 01 \\ 02 \\ 03 \\ 23 \\ 31 \\ 12 \end{array}$
mielke-geometroynamics_EQ0803 (11.4.15)	255	■ 0.324	$\psi = \frac{4}{\ell} \left(\frac{2\pi m}{3} \right)^{1/2} e^{-\mu/2 - \nu/4 - i\omega m t} \left[\begin{array}{cc} iH(\rho) & x_{\kappa}^m \\ F(\rho) & \frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } x_{\kappa}^m \end{array} \right].$	$\psi = \frac{4}{\ell} \left(\frac{2\pi m}{3} \right)^{1/2} e^{-\mu/2 - \nu/4 - i\omega m t} \left[\begin{array}{cc} iH(\rho) & x_{\kappa}^m \\ F(\rho) & \frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } x_{\kappa}^m \end{array} \right].$	$\psi = \frac{4}{\ell} \left(\frac{2\pi m}{3} \right)^{1/2} e^{-\mu/2 - \nu/4 - i\omega m t} \left[\begin{array}{cc} iH(\rho) & x_{\kappa}^m \\ F(\rho) & \frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } x_{\kappa}^m \end{array} \right].$
mielke-geometroynamics_EQ1082 (A.7)	365	■ 0.326	$\begin{aligned} & \gamma^0 := \begin{pmatrix} \sigma^0 & 0 \\ 0 & \sigma^0 \end{pmatrix}, \quad \gamma^k := \begin{pmatrix} 0 & \sigma^k \\ -\sigma^k & 0 \end{pmatrix}; \\ & \gamma^5 := i\gamma^0 \gamma^1 \gamma^2 \gamma^3 = \begin{pmatrix} 0 & \sigma^0 \\ \sigma^0 & 0 \end{pmatrix}. \end{aligned}$	$\gamma^0 := \begin{pmatrix} \sigma^0 & 0 \\ 0 & \sigma^0 \end{pmatrix}, \quad \gamma^k := \begin{pmatrix} 0 & \sigma^k \\ -\sigma^k & 0 \end{pmatrix};$ $\gamma^5 := i\gamma^0 \gamma^1 \gamma^2 \gamma^3 = \begin{pmatrix} 0 & \sigma^0 \\ \sigma^0 & 0 \end{pmatrix}.$	$\gamma^0 := \begin{pmatrix} \sigma^0 & 0 \\ 0 & \sigma^0 \end{pmatrix}, \quad \gamma^k := \begin{pmatrix} 0 & \sigma^k \\ -\sigma^k & 0 \end{pmatrix};$ $\gamma^5 := i\gamma^0 \gamma^1 \gamma^2 \gamma^3 = \begin{pmatrix} 0 & \sigma^0 \\ \sigma^0 & 0 \end{pmatrix}.$
mielke-geometroynamics_EQ0978 (14.2.19)	316	■ 0.331	$\begin{aligned} & \pi \hat{L}_{ } = -\frac{\hat{\ell}^2}{\ell^{*2}} \mathrm{Tr}(\Omega' \wedge {}^* \Omega' \wedge \underbrace{\vartheta \wedge \cdots \wedge \vartheta}_{\text{K terms}}) \\ & = -\frac{1}{\alpha_g} \mathrm{Tr}(\Omega' \wedge {}^* \Omega') \sqrt{\det \overset{\circ}{g}} = L'_{\mathrm{YM}} d_{\mu}(G) \end{aligned}$	$\pi \hat{L}_{ } = -\frac{\hat{\ell}^2}{\ell^{*2}} \mathrm{Tr}(\Omega' \wedge {}^* \Omega' \wedge \underbrace{\vartheta \wedge \cdots \wedge \vartheta}_{\text{K terms}})$ $= -\frac{1}{\alpha_g} \mathrm{Tr}(\Omega' \wedge {}^* \Omega') \sqrt{\det \overset{\circ}{g}} = L'_{\mathrm{YM}} d_{\mu}(G)$	$\pi \hat{L}_{ } = -\frac{\hat{\ell}^2}{\ell^{*2}} \mathrm{Tr}(\Omega' \wedge {}^* \Omega' \wedge \underbrace{\vartheta \wedge \cdots \wedge \vartheta}_{\text{K terms}})$ $= -\frac{1}{\alpha_g} \mathrm{Tr}(\Omega' \wedge {}^* \Omega') \sqrt{\det \overset{\circ}{g}} = L'_{\mathrm{YM}} d_{\mu}(G)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_128 EQ0377 (6.2.28)		■ 0.396	$\stackrel{\scriptscriptstyle{(\pm)}}{\rm L}_{\parallel} = -\frac{1}{2}\left[\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{D}_{\alpha} + d\left(\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}\right)\right] = -\frac{1}{2}\left(\stackrel{\scriptscriptstyle{(\pm)}}{D}\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},\right.$	$\stackrel{\scriptscriptstyle{(\pm)}}{L}_{\parallel} = -\frac{1}{2}\left[\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{D}_{\alpha} + d\left(\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}\right)\right] = -\frac{1}{2}\left(\stackrel{\scriptscriptstyle{(\pm)}}{D}\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},\right.$	$\stackrel{\scriptscriptstyle{(\pm)}}{L}_{\parallel} = -\frac{1}{2}\left[\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{D}\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha} + d(\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha})\right] = -\frac{1}{2}\stackrel{\scriptscriptstyle{(\pm)}}{D}\vartheta^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},$
mielke-geometrodynamics_94 EQ0262 (4.3.27)		■ 0.399	$\begin{aligned} \Omega^g &= \Omega^{\{\}} - dK + K\wedge\omega^{\{\}} + \omega^{\{\}}\wedge K - K\wedge K \\ &= \Omega^{\{\}} - D^{\{\}}K - K\wedge K = \Omega^{\{\}} - DK + K\wedge K \end{aligned}$	$\Omega^g = \Omega^{\{\}} - dK + K\wedge\omega^{\{\}} + \omega^{\{\}}\wedge K - K\wedge K$ $= \Omega^{\{\}} - D^{\{\}}K - K\wedge K = \Omega^{\{\}} - DK + K\wedge K$	$\Omega^g = \Omega^{\{\}} - dK + K\wedge\omega^{\{\}} + \omega^{\{\}}\wedge K - K\wedge K$ $= \Omega^{\{\}} - D^{\{\}}K - K\wedge K = \Omega^{\{\}} - DK + K\wedge K$
mielke-geometrodynamics_184 EQ0570 (8.6.8)		■ 0.403	$\left.E_{\alpha}:=\partial L/\partial\vartheta^{\alpha}=e_{\alpha}\rfloor L+\left(e_{\alpha}\rfloor T^{\beta}\right)\wedge H_{\beta}+\left(e_{\alpha}\rfloor R^{\beta\gamma}\right)\wedge H_{\beta\gamma}\right.$	$E_{\alpha}:=\partial L/\partial\vartheta^{\alpha}=e_{\alpha}\rfloor L+\left(e_{\alpha}\rfloor T^{\beta}\right)\wedge H_{\beta}+\left(e_{\alpha}\rfloor R^{\beta\gamma}\right)\wedge H_{\beta\gamma}$	$E_{\alpha}:=\partial L/\partial\vartheta^{\alpha}=e_{\alpha}\rfloor L+\left(e_{\alpha}\rfloor T^{\beta}\right)\wedge H_{\beta}+\left(e_{\alpha}\rfloor R^{\beta\gamma}\right)\wedge H_{\beta\gamma}$
mielke-geometrodynamics_103 EQ0306 (4.5.16)		■ 0.406	$\left.\widetilde{\varepsilon}_{\mathrm{MA}}:=\varepsilon_{\mathrm{RC}}+\xi\rfloor L\cong-\left(\mathcal{L}_{\xi}\Gamma_{\beta}^{\gamma}\right)\wedge H^{\beta}_{\gamma}+dH_{\infty}\right.$	$\widetilde{\varepsilon}_{\mathrm{MA}}:=\varepsilon_{\mathrm{RC}}+\xi\rfloor L\cong-\left(\mathcal{L}_{\xi}\Gamma_{\beta}^{\gamma}\right)\wedge H^{\beta}_{\gamma}+dH_{\infty}$	$\widetilde{\varepsilon}_{\mathrm{MA}}:=\varepsilon_{\mathrm{RC}}+\xi\rfloor L\cong-\left(\mathcal{L}_{\xi}\Gamma_{\beta}^{\gamma}\right)\wedge H^{\beta}_{\gamma}+dH_{\infty}$
mielke-geometrodynamics_114 EQ0336 (5.3.13)		■ 0.407	$R^{\{\alpha\beta\}}=R^{\{\{\jmath\alpha\beta\}-D^{\{\{\}\}\}}K^{\{\alpha\beta\}}-K^{\{\{\alpha\beta\}}\}_{\mu}}\wedge K^{\{\mu\beta\}}$	$R^{\alpha\beta}=R^{\{j\alpha\beta}-D^{\{\}}K^{\alpha\beta}-K^{\alpha}_{\mu}\wedge K^{\mu\beta}$	$R^{\alpha\beta}=R^{\{\}\alpha\beta}-D^{\{\}}K^{\alpha\beta}-K^{\alpha}_{\mu}\wedge K^{\mu\beta}$
mielke-geometrodynamics_375 EQ1136 (C.4)		■ 0.409	$c_{ij}^k=-c_{ji}^k$	$c_{ij}^k=-c_{ji}^k$	$c_{ij}^k=-c_{ji}^k$
mielke-geometrodynamics_279 EQ0861 (12.4.8)		■ 0.420	$\begin{aligned} dC_{\mathrm{RR}} &= -\operatorname{Tr}(\Omega\wedge\Omega) \\ &= \frac{1}{2}R_{\alpha\beta}^{\{\}}\wedge R^{\{\beta\alpha\beta}} \\ &+ \frac{1}{12}d\left[{}^*\mathcal{A}\wedge R^{\{\}}-\frac{1}{3}\mathcal{A}\wedge d\mathcal{A}+\frac{1}{9}{}^*\mathcal{A}\wedge{}^*(\mathcal{A}\wedge{}^*\mathcal{A})\right]. \end{aligned}$	$dC_{\mathrm{RR}} = -\operatorname{Tr}(\Omega\wedge\Omega)$ $= \frac{1}{2}R_{\alpha\beta}^{\{\}}\wedge R^{\{\beta\alpha\beta}}$ $+ \frac{1}{12}d\left[{}^*\mathcal{A}\wedge R^{\{\}}-\frac{1}{3}\mathcal{A}\wedge d\mathcal{A}+\frac{1}{9}{}^*\mathcal{A}\wedge{}^*(\mathcal{A}\wedge{}^*\mathcal{A})\right].$	$dC_{\mathrm{RR}} = -\operatorname{Tr}(\Omega\wedge\Omega)$ $= \frac{1}{2}R_{\alpha\beta}^{\{\}}\wedge R^{\{\}\alpha\beta}$ $+ \frac{1}{12}d\left[{}^*\mathcal{A}\wedge R^{\{\}}-\frac{1}{3}\mathcal{A}\wedge d\mathcal{A}+\frac{1}{9}{}^*\mathcal{A}\wedge{}^*(\mathcal{A}\wedge{}^*\mathcal{A})\right].$
mielke-geometrodynamics_151 EQ0451 (7.2.4)		■ 0.424	$\stackrel{\scriptscriptstyle{+g}}{\Omega}:=\frac{1}{2}\left(\Omega^g\pm{}^*\Omega^{g(*)}\right),$	$\stackrel{\scriptscriptstyle{+g}}{\Omega}:=\frac{1}{2}\left(\Omega^g\pm{}^*\Omega^{g(*)}\right),$	$\stackrel{\scriptscriptstyle{\pm}}{\Omega}^g:=\frac{1}{2}\left(\Omega^g\pm{}^*\Omega^{g(*)}\right),$
mielke-geometrodynamics_127 EQ0372 (6.2.23)		■ 0.437	$\left.\stackrel{\scriptscriptstyle{(\pm)}}{D}\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}\mp\frac{i}{4}\ell^2e_{\alpha}\right]\left(\stackrel{\scriptscriptstyle{(+)}}{\Pi}^{\beta}\wedge\stackrel{\scriptscriptstyle{(-)}}{\Pi}_{\beta}\right)=\Sigma_{\alpha},$	$\stackrel{\scriptscriptstyle{(\pm)}}{D}\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}\mp\frac{i}{4}\ell^2e_{\alpha}\right]\left(\stackrel{\scriptscriptstyle{(+)}}{\Pi}^{\beta}\wedge\stackrel{\scriptscriptstyle{(-)}}{\Pi}_{\beta}\right)=\Sigma_{\alpha},$	$\stackrel{\scriptscriptstyle{(\pm)}}{D}\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}\mp\frac{i}{4}\ell^2e_{\alpha}\rfloor\left(\stackrel{\scriptscriptstyle{(+)}}{\Pi}^{\beta}\wedge\stackrel{\scriptscriptstyle{(-)}}{\Pi}_{\beta}\right)=\Sigma_{\alpha},$
mielke-geometrodynamics_322 EQ0997 (14.3.10)		■ 0.442	$D^{\{(\text{int})\}}=\stackrel{\scriptscriptstyle{\circ}}{\gamma}\wedge{}^*\stackrel{\scriptscriptstyle{\circ}}{D}=\stackrel{\scriptscriptstyle{\circ}}{\gamma}\wedge{}^*(\stackrel{\scriptscriptstyle{\circ}}{d}+i\stackrel{\scriptscriptstyle{\circ}}{\tilde{\omega}})$	$D^{(\text{int})}=\stackrel{\scriptscriptstyle{\circ}}{\gamma}\wedge{}^*\stackrel{\scriptscriptstyle{\circ}}{D}=\stackrel{\scriptscriptstyle{\circ}}{\gamma}\wedge{}^*(\stackrel{\scriptscriptstyle{\circ}}{d}+i\stackrel{\scriptscriptstyle{\circ}}{\tilde{\omega}})$	$\not{D}^{(int)}=\stackrel{\scriptscriptstyle{\circ}}{\gamma}\wedge{}^*\stackrel{\scriptscriptstyle{\circ}}{D}=\stackrel{\scriptscriptstyle{\circ}}{\gamma}\wedge{}^*(\stackrel{\scriptscriptstyle{\circ}}{d}+i\stackrel{\scriptscriptstyle{\circ}}{\tilde{\omega}})$
mielke-geometrodynamics_156 EQ0474 (7.4.2)		■ 0.444	$R_{\alpha\beta}\equiv C_{\alpha\beta}+\frac{1}{2}Ri^{\tau}_{\alpha\beta}-\frac{R}{12}\vartheta_{\alpha}\wedge\vartheta_{\beta},$	$R_{\alpha\beta}\equiv C_{\alpha\beta}+\frac{1}{2}Ri^{\tau}_{\alpha\beta}-\frac{R}{12}\vartheta_{\alpha}\wedge\vartheta_{\beta},$	$R_{\alpha\beta}\equiv C_{\alpha\beta}+\frac{1}{2}Ri^{\tau}_{\alpha\beta}-\frac{R}{12}\vartheta_{\alpha}\wedge\vartheta_{\beta},$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_ E00681 (10.3.38)	221	■ 0.452	$\begin{gathered} \chi=\theta_{\mathrm{L}}\,\varepsilon-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon},\quad \quad \quad \Lambda=\frac{(-1)^s}{4\ell^2}\left(\theta_{\mathrm{L}}\varepsilon^3-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^3\right),\quad \quad \theta_{\mathrm{T}}=\frac{(-1)^s}{2}\left(\theta_{\mathrm{L}}\varepsilon^2-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^2\right), \\ \Delta\theta_{\mathrm{L}}=\theta_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}},\quad \Delta\theta_{\mathrm{TL}}=\frac{1}{2}\left(\theta_{\mathrm{L}}\varepsilon-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}\right). \end{gathered}$	$\chi=\theta_{\mathrm{L}}\varepsilon-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon},\quad \Lambda=\frac{(-1)^s}{4\ell^2}\left(\theta_{\mathrm{L}}\varepsilon^3-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^3\right),\quad \theta_{\mathrm{T}}=\frac{(-1)^s}{2}\left(\theta_{\mathrm{L}}\varepsilon^2-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^2\right),$ $\Delta\theta_{\mathrm{L}}=\theta_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}},\quad \Delta\theta_{\mathrm{TL}}=\frac{1}{2}\left(\theta_{\mathrm{L}}\varepsilon-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}\right).$	$\chi=\theta_{\mathrm{L}}\varepsilon-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon},\quad \Lambda=\frac{(-1)^s}{4\ell^2}\left(\theta_{\mathrm{L}}\varepsilon^3-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^3\right),\quad \theta_{\mathrm{T}}=\frac{(-1)^s}{2}\left(\theta_{\mathrm{L}}\varepsilon^2-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}^2\right),$ $\Delta\theta_{\mathrm{L}}=\theta_{\mathrm{L}}-\tilde{\theta}_{\mathrm{L}},\quad \Delta\theta_{\mathrm{TL}}=\frac{1}{2}\left(\theta_{\mathrm{L}}\varepsilon-\tilde{\theta}_{\mathrm{L}}\tilde{\varepsilon}\right).$
mielke-geometrodynamics_ E00845 (12.2.13)	273	■ 0.457	$C_{\mathrm{TT}}\cong\frac{1}{4}j_5,\quad \mathscr{A}=2\ell^2{}^*C_{\mathrm{TT}}=(\ell^2/2)\,\bar{\psi}\gamma_5\gamma\psi.$	$C_{\mathrm{TT}}\cong\frac{1}{4}j_5,\quad \mathscr{A}=2\ell^2{}^*C_{\mathrm{TT}}=(\ell^2/2)\,\bar{\psi}\gamma_5\gamma\psi.$	$C_{\mathrm{TT}}\cong\frac{1}{4}j_5,\quad \mathscr{A}=2\ell^2{}^*C_{\mathrm{TT}}=(\ell^2/2)\,\bar{\psi}\gamma_5\gamma\psi.$
mielke-geometrodynamics_ E00375 (6.2.26)	128	■ 0.466	$\stackrel{\scriptscriptstyle{(\pm)}}{\stackrel{\scriptscriptstyle{(\pm)}}{L}}_{\parallel}=\mp\frac{i}{4}\ell^2\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},\quad \quad \overline{\stackrel{\scriptscriptstyle{(\pm)}}{L}}_{\parallel}=\stackrel{\scriptscriptstyle{(-)}}{L}_{\parallel},$	$\stackrel{\scriptscriptstyle{(\pm)}}{L}_{\parallel}=\mp\frac{i}{4}\ell^2\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},\quad \quad \overline{\stackrel{\scriptscriptstyle{(\pm)}}{L}}_{\parallel}=\stackrel{\scriptscriptstyle{(-)}}{L}_{\parallel},$	$\stackrel{\scriptscriptstyle{(\pm)}}{L}_{\parallel}=\mp\frac{i}{4}\ell^2\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}^{\alpha}\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},\quad \quad \overline{\stackrel{\scriptscriptstyle{(\pm)}}{L}}_{\parallel}=\stackrel{\scriptscriptstyle{(-)}}{L}_{\parallel},$
mielke-geometrodynamics_ E00222 (4.2.15)	87	■ 0.466	$\widetilde{\varphi}:A(M)\rightarrow C^{\infty}(\overset{\circ}{V})$	$\widetilde{\varphi}:A(M)\rightarrow C^{\infty}(\overset{\circ}{V})$	$\widetilde{\varphi}:A(M)\rightarrow C^{\infty}(\overset{\circ}{V})$
mielke-geometrodynamics_ E00604 (9.2.3)	199	■ 0.474	$\begin{aligned} L_{\mathrm{Euler}}&,\;=\;{}^*(\gamma_2^g)=\frac{1}{(4\pi)^2}\operatorname{Tr}\left(\Omega^g\wedge\Omega^{g(*)}\right) \\ &=\frac{(-1)^{s/4}}{32\pi^2}R^{\alpha\beta\gamma\delta}{}^*R^*_{\alpha\beta\gamma\delta}\sqrt{ g }d^4x. \end{aligned}$	$L_{\mathrm{Euler}}\,,\;=\;{}^*(\gamma_2^g)=\frac{1}{(4\pi)^2}\operatorname{Tr}\left(\Omega^g\wedge\Omega^{g(*)}\right)$ $=\frac{(-1)^{s/4}}{32\pi^2}R^{\alpha\beta\gamma\delta}{}^*R^*_{\alpha\beta\gamma\delta}\sqrt{ g }d^4x.$	$L_{\mathrm{Euler}}\,,\;=\;{}^*(\gamma_2^g)=\frac{1}{(4\pi)^2}\operatorname{Tr}(\Omega^g\wedge\Omega^{g(*)})$ $=\frac{(-1)^{s/4}}{32\pi^2}R^{\alpha\beta\gamma\delta}{}^*R^*_{\alpha\beta\gamma\delta}\sqrt{ g }d^4x.$
mielke-geometrodynamics_ E00401 (6.4.20)	133	■ 0.482	$\mathscr{H}_{\parallel}=\underline{D}-\mathscr{A}_{\hat{0}}^{\scriptscriptstyle{(\pm)}}=0,$	$\mathscr{H}_{\parallel}=\underline{D}-\mathscr{A}_{\hat{0}}^{\scriptscriptstyle{(\pm)}}=0,$	$\mathscr{H}_{\parallel}=\underline{D}-\mathscr{A}_{\hat{0}}^{\scriptscriptstyle{(\pm)}}=0,$
mielke-geometrodynamics_ E00750 (11.1.17)	241	■ 0.484	$D_{\left[\left.\left.\right _{\overline{\operatorname{S}0}(1,3)}\right]^{\operatorname{F}}\right]^{\operatorname{F}}}=\underbrace{D^{\frac{1}{2}}\times\cdots\times D^{\frac{1}{2}}}_{(n-4)/4}.$	$D_{\left[\left.\left.\right _{\overline{\operatorname{S}0}(1,3)}\right]^{\operatorname{F}}\right]^{\operatorname{F}}}=\underbrace{D^{\frac{1}{2}}\times\cdots\times D^{\frac{1}{2}}}_{(n-4)/4}.$	$D_{\left[\left.\left.\right _{\overline{\operatorname{S}0}(1,3)}\right]^{\operatorname{F}}\right]^{\operatorname{F}}}=\underbrace{D^{\frac{1}{2}}\times\cdots\times D^{\frac{1}{2}}}_{(n-4)/4}.$
mielke-geometrodynamics_ E00686 (10.3.44)	222	■ 0.497	$C_l=E_l^{\alpha}\ast\left\{D^{\{\}}\left[2\left(e^{\beta}\lrcorner D^{\{\}}H_{[\alpha\beta]}\right)+\frac{1}{2}\vartheta_{\alpha}\wedge\left(e^{\beta}\lrcorner e^{\gamma}\lrcorner D^{\{\}}H_{[\beta\gamma]}\right)\right]\right\}$	$C_l=E_l^{\alpha}\ast\left\{D^{\{\}}\left[2\left(e^{\beta}\lrcorner D^{\{\}}H_{[\alpha\beta]}\right)+\frac{1}{2}\vartheta_{\alpha}\wedge\left(e^{\beta}\lrcorner e^{\gamma}\lrcorner D^{\{\}}H_{[\beta\gamma]}\right)\right]\right\}$	$C_l=E_l^{\alpha}\ast\left\{D^{\{\}}\left[2\left(e^{\beta}\lrcorner D^{\{\}}H_{[\alpha\beta]}\right)+\frac{1}{2}\vartheta_{\alpha}\wedge\left(e^{\beta}\lrcorner e^{\gamma}\lrcorner D^{\{\}}H_{[\beta\gamma]}\right)\right]\right\}$
mielke-geometrodynamics_ E00515 (7.7.33)	166	■ 0.497	$T:=\frac{1}{4}\operatorname{Tr}(\tilde{\gamma}\lrcorner\Theta)=e_{\alpha}\rfloor T^{\alpha},\quad \mathscr{A}:=\frac{1}{4}{}^*\operatorname{Tr}(\gamma\wedge\Theta)={}^*(\vartheta_{\alpha}\wedge T^{\alpha}).$	$T:=\frac{1}{4}\operatorname{Tr}(\tilde{\gamma}\lrcorner\Theta)=e_{\alpha}\rfloor T^{\alpha},\quad \mathscr{A}:=\frac{1}{4}{}^*\operatorname{Tr}(\gamma\wedge\Theta)={}^*(\vartheta_{\alpha}\wedge T^{\alpha}).$	$T:=\frac{1}{4}\operatorname{Tr}(\check{\gamma}\lrcorner\Theta)=e_{\alpha}\rfloor T^{\alpha},\quad \mathscr{A}:=\frac{1}{4}{}^*\operatorname{Tr}(\gamma\wedge\Theta)={}^*(\vartheta_{\alpha}\wedge T^{\alpha}).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_00799 (11.4.11)	254	■ 0.505	$ds^2=\frac{1}{\lambda^2}\left(R_{\circ}^2d\lambda^2-\mathrm{o}_{ij}dy^i\otimes_sdy^j\right).$	$ds^2=\frac{1}{\lambda^2}\left(R_{\circ}^2d\lambda^2-\mathrm{o}_{ij}dy^i\otimes_sdy^j\right).$	$ds^2=\frac{1}{\lambda^2}\left(R_{\circ}^2d\lambda^2-\mathrm{o}_{ij}dy^i\otimes_sdy^j\right).$
mielke-geometrodynamics_00960 (14.1.42)	309	■ 0.508	$\xi_{\mu\nu}\xi_{\rho\sigma}=-\frac{1}{2}\frac{\bar{R}_{\mu\nu\rho\sigma}}{R_{\alpha\beta}R^{j\alpha\beta}}-\frac{1}{2}\frac{\bar{R}_{\mu\nu\gamma\delta}\bar{R}_{\rho\sigma}{}^{\gamma\delta}}{R_{\alpha\beta}R^{j\alpha\beta}}.$	$\xi_{\mu\nu}\xi_{\rho\sigma}=-\frac{1}{2}\bar{R}_{\mu\nu\rho\sigma}-\frac{1}{2}\frac{\bar{R}_{\mu\nu\gamma\delta}\bar{R}_{\rho\sigma}{}^{\gamma\delta}}{R_{\alpha\beta}R^{j\alpha\beta}}.$	$\xi_{\mu\nu}\xi_{\rho\sigma}=-\frac{1}{2}\bar{R}_{\mu\nu\rho\sigma}-\frac{1}{2}\frac{\bar{R}_{\mu\nu\gamma\delta}\bar{R}_{\rho\sigma}{}^{\gamma\delta}}{R_{\alpha\beta}R^{j\alpha\beta}}.$
mielke-geometrodynamics_01131 (B.43)	373	■ 0.509	$\langle\cdot,\cdot\rangle_M:\begin{cases}\bigwedge^pT^*(M)\times C_p(M,\mathbb{R})\rightarrow\mathbb{R}\\\langle\alpha^{(p)},c_p\rangle_M\rightarrow\int_{c_p}\alpha^{(p)}\end{cases}$	$\langle\cdot,\cdot\rangle_M:\begin{cases}\bigwedge^pT^*(M)\times C_p(M,\mathbb{R})\rightarrow\mathbb{R}\\\langle\alpha^{(p)},c_p\rangle_M\rightarrow\int_{c_p}\alpha^{(p)}\end{cases}$	$\langle\cdot,\cdot\rangle_M:\begin{cases}\bigwedge^pT^*(M)\times C_p(M,\mathbb{R})\rightarrow\mathbb{R}\\\langle\alpha^{(p)},c_p\rangle_M\rightarrow\int_{c_p}\alpha^{(p)}\end{cases}$
mielke-geometrodynamics_00294 (4.5.5)	101	■ 0.510	$\left.\Sigma_{\alpha}\right L-(e_{\alpha}\rfloor D\Psi)\wedge\frac{\partial L}{\partial D\Psi}-(e_{\alpha}\rfloor\Psi)\wedge\frac{\partial L}{\partial\Psi}.$	$\Sigma_{\alpha}=e_{\alpha}\rfloor L-(e_{\alpha}\rfloor D\Psi)\wedge\frac{\partial L}{\partial D\Psi}-(e_{\alpha}\rfloor\Psi)\wedge\frac{\partial L}{\partial\Psi}.$	$\Sigma_{\alpha}=e_{\alpha}\rfloor L-(e_{\alpha}\rfloor D\Psi)\wedge\frac{\partial L}{\partial D\Psi}-(e_{\alpha}\rfloor\Psi)\wedge\frac{\partial L}{\partial\Psi}.$
mielke-geometrodynamics_00118 (3.3.19)	59	■ 0.513	$L_{\mathrm{Pontr}}:=-\frac{1}{2}F\wedge F=\mathbf{E}\cdot\mathbf{B}\eta.$	$L_{\mathrm{Pontr}}:=-\frac{1}{2}F\wedge F=\mathbf{E}\cdot\mathbf{B}\eta.$	$L_{\mathrm{Pontr}}:=-\frac{1}{2}F\wedge F=\mathbf{E}\cdot\mathbf{B}\eta.$
mielke-geometrodynamics_00533 (7.7.53)	169	■ 0.513	$\begin{aligned}G&:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma}\gamma^{\alpha}\\&=G^{(\beta)}+\frac{(-1)^{\mathrm{sig}}}{12}(e_{\alpha}\rfloor\mathscr{A}\wedge{}^*\mathscr{A}-\frac{1}{3}\mathscr{A}\wedge e_{\alpha}\rfloor{}^*\mathscr{A})\gamma^{\alpha}+\frac{(-1)^{\mathrm{sig}}}{6}\gamma\wedge d\mathscr{A}\end{aligned}$	$G:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma}\gamma^{\alpha}=G^{(\beta)}+\frac{(-1)^{\mathrm{sig}}}{12}(e_{\alpha}\rfloor\mathscr{A}\wedge{}^*\mathscr{A}-\frac{1}{3}\mathscr{A}\wedge e_{\alpha}\rfloor{}^*\mathscr{A})\gamma^{\alpha}+\frac{(-1)^{\mathrm{sig}}}{6}\gamma\wedge d\mathscr{A}$	$G:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma}\gamma^{\alpha}=G^{(\beta)}+\frac{(-1)^{\mathrm{sig}}}{12}\left(e_{\alpha}\rfloor\mathscr{A}\wedge{}^*\mathscr{A}-\frac{1}{3}\mathscr{A}\wedge e_{\alpha}\rfloor{}^*\mathscr{A}\right)\gamma^{\alpha}+\frac{(-1)^{\mathrm{sig}}}{6}\gamma\wedge d\mathscr{A}\tag{7.7.5}$
mielke-geometrodynamics_00898 (13.5.3)	288	■ 0.515	$\hat{B}:=B_A{}^B{}^A{}_B=(b_{\alpha}P_{*}^{\alpha}+\hat{b}^{\alpha}P_{\alpha})\ell^2+B_{\alpha}{}^{\beta}L^{\alpha}{}_{\beta}$	$\hat{B}:=B_A{}^B{}^A{}_B=(b_{\alpha}P_{*}^{\alpha}+\hat{b}^{\alpha}P_{\alpha})\ell^2+B_{\alpha}{}^{\beta}L^{\alpha}{}_{\beta}$	$\hat{B}:=B_A{}^B{}^A{}_B=(b_{\alpha}P_{*}^{\alpha}+\hat{b}^{\alpha}P_{\alpha})\ell^2+B_{\alpha}{}^{\beta}L^{\alpha}{}_{\beta}$
mielke-geometrodynamics_00845 (2.6.8)	43	■ 0.516	$(db)_{\mathrm{hor}}=Db=\omega\otimes b$	$(db)_{\mathrm{hor}}=Db=\omega\otimes b$	$(db)_{\mathrm{hor}}=Db=\omega\otimes b$
mielke-geometrodynamics_00876 (13.3.5)	285	■ 0.518	$\begin{aligned}&\left\{ \left[P_{\alpha}, P_{\beta} \right] = 0, \left[P_{*}^{\alpha}, P_{*}^{\beta} \right] = 0, \right. \\&\left. \left[L_{\beta}^{\alpha}, P_{\gamma} \right] = \delta_{\gamma}^{\alpha} P_{\beta}, \left[L_{\beta}^{\alpha}, P_{*}^{\gamma} \right] = -\delta_{\beta}^{\gamma} P_{*}^{\alpha} . \right. \\&\left. \left[L_{\beta}^{\alpha}, P_{\gamma} \right] = \delta_{\gamma}^{\alpha} P_{\beta}, \left[L_{\beta}^{\alpha}, P_{*}^{\gamma} \right] = -\delta_{\beta}^{\gamma} P_{*}^{\alpha} . \right.\end{aligned}$	$\begin{aligned}&\left[P_{\alpha}, P_{\beta} \right] = 0, \left[P_{*}^{\alpha}, P_{*}^{\beta} \right] = 0, \\&\left[L_{\beta}^{\alpha}, P_{\gamma} \right] = \delta_{\gamma}^{\alpha} P_{\beta}, \left[L_{\beta}^{\alpha}, P_{*}^{\gamma} \right] = -\delta_{\beta}^{\gamma} P_{*}^{\alpha} .\end{aligned}$	$\begin{aligned}&\left[P_{\alpha}, P_{\beta} \right] = 0, \quad \left[P_{*}^{\alpha}, P_{*}^{\beta} \right] = 0, \\&\left[L^{\alpha}{}_{\beta}, P_{\gamma} \right] = \delta_{\gamma}^{\alpha} P_{\beta}, \quad \left[L^{\alpha}{}_{\beta}, P_{*}^{\gamma} \right] = -\delta_{\beta}^{\gamma} P_{*}^{\alpha} .\end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_320 E08998 (14.3.3)		■ 0.526	$\begin{aligned} \mathrm{d} \left(\overline{\hat{\psi}} \wedge \hat{\gamma} \right) &= \frac{1}{2} \left(\overline{\hat{\psi}} \wedge \hat{\gamma} \right) \wedge \hat{\gamma} \\ &+ \frac{1}{2} \overline{\hat{\psi}} [\hat{\alpha}, \hat{\gamma}] \hat{\psi} . \end{aligned}$	$\hat{L}_D = \frac{i}{2} \left(\overline{\hat{\psi}} \wedge \hat{\gamma} \right) \wedge \hat{\gamma} + \frac{1}{2} \overline{\hat{\psi}} [\hat{\alpha}, \hat{\gamma}] \hat{\psi} .$	$\hat{L}_D = \frac{i}{2} \left(\overline{\hat{\psi}} \wedge \hat{\gamma} \right) \wedge \hat{\gamma} + \frac{1}{2} \overline{\hat{\psi}} [\hat{\alpha}, \hat{\gamma}] \hat{\psi} .$
mielke-geometrodynamics_152 E08456 (7.2.8)		■ 0.527	$\bar{R}_{..cd} = 2R_{[c}^{[a} \delta_{d]}^{b]} ; \quad \bar{R}_{..cd} = R_{..cd} - \frac{1}{4} \delta_{cd} R ,$	$\bar{R}_{..cd} = 2R_{[c}^{[a} \delta_{d]}^{b]} ; \quad \bar{R}_{..cd} = R_{..cd} - \frac{1}{4} \delta_{cd} R ,$	$\bar{R}_{..cd} = 2R_{[c}^{[a} \delta_{d]}^{b]} ; \quad \bar{R}_{..cd} = R_{..cd} - \frac{1}{4} \delta_{cd} R ,$
mielke-geometrodynamics_125 E08360 (6.2.11)		■ 0.531	$L_{\parallel} \equiv L_{\mathrm{HE}} + \frac{1}{2\kappa} R_{\alpha\beta} \wedge^* (\vartheta^\alpha \wedge \vartheta^\beta) + \frac{2\ell^2}{\kappa} dC_{\mathrm{TT}}^* ,$	$L_{\parallel} \equiv L_{\mathrm{HE}} + \frac{1}{2\kappa} R_{\alpha\beta} \wedge^* (\vartheta^\alpha \wedge \vartheta^\beta) + \frac{2\ell^2}{\kappa} dC_{\mathrm{TT}}^* ,$	$L_{\parallel} \equiv L_{\mathrm{HE}} + \frac{1}{2\kappa} R_{\alpha\beta} \wedge^* (\vartheta^\alpha \wedge \vartheta^\beta) + \frac{2\ell^2}{\kappa} dC_{\mathrm{TT}}^* ,$
mielke-geometrodynamics_142 E08437 (6.8.8)		■ 0.532	$E_{\alpha} := \frac{\partial L_g}{\partial \vartheta^\alpha} = e_{\alpha} \lrcorner L_g + (e_{\alpha} \lrcorner T^{\beta}) \wedge H_{\beta} + (e_{\alpha} \lrcorner R^{\beta\gamma}) \wedge H_{\beta\gamma} ,$	$E_{\alpha} := \frac{\partial L_g}{\partial \vartheta^\alpha} = e_{\alpha} \lrcorner L_g + (e_{\alpha} \lrcorner T^{\beta}) \wedge H_{\beta} + (e_{\alpha} \lrcorner R^{\beta\gamma}) \wedge H_{\beta\gamma} ,$	$E_{\alpha} := \frac{\partial L_g}{\partial \vartheta^\alpha} = e_{\alpha} \lrcorner L_g + (e_{\alpha} \lrcorner T^{\beta}) \wedge H_{\beta} + (e_{\alpha} \lrcorner R^{\beta\gamma}) \wedge H_{\beta\gamma} ,$
mielke-geometrodynamics_82 E08198 (4.1.11)		■ 0.533	$\begin{aligned} e_{\alpha}(m) &\rightarrow e_{\alpha}'(m) = g_{\alpha}^{\beta} e_{\beta}(m) , \\ g &= [g_{\alpha}^{\beta}] \in GL(n, \mathbb{R}) . \end{aligned}$	$e_{\alpha}(m) \rightarrow e_{\alpha}'(m) = g_{\alpha}^{\beta} e_{\beta}(m) , \\ g = [g_{\alpha}^{\beta}] \in GL(n, \mathbb{R}) .$	$e_{\alpha}(m) \rightarrow e_{\alpha}'(m) = g_{\alpha}^{\beta} e_{\beta}(m) , \\ g = [g_{\alpha}^{\beta}] \in GL(n, \mathbb{R}) .$
mielke-geometrodynamics_196 E08599 (9.1.17)		■ 0.534	$\begin{aligned} f &= -\frac{1}{r} \left(1 + \frac{\kappa}{2\ell^2} r^2 \right) e^{-\nu/2} , \quad g = \frac{1}{r} e^{\nu/2} \\ h &= -\frac{1}{(+)\ell^2} \frac{\sqrt{-\kappa}}{3} , \quad k = +\frac{1}{(-)2\ell^2} \sqrt{-3\kappa} . \end{aligned}$	$f = -\frac{1}{r} \left(1 + \frac{\kappa}{2\ell^2} r^2 \right) e^{-\nu/2} , \quad g = \frac{1}{r} e^{\nu/2} \\ h = -\frac{1}{(+)\ell^2} \frac{\sqrt{-\kappa}}{3} , \quad k = +\frac{1}{(-)2\ell^2} \sqrt{-3\kappa} .$	$f = -\frac{1}{r} \left(1 + \frac{\kappa}{2\ell^2} r^2 \right) e^{-\nu/2} , \quad g = \frac{1}{r} e^{\nu/2} \\ h = -\frac{1}{(+)\ell^2} \frac{\sqrt{-\kappa}}{3} , \quad k = +\frac{1}{(-)2\ell^2} \sqrt{-3\kappa} .$
mielke-geometrodynamics_241 E08752 (11.1.19)		■ 0.535	$\Psi = \Psi \otimes \tilde{b} \in C^{\infty} \left(V^{\frac{1}{2}} \right) .$	$\Psi = \Psi \otimes \tilde{b} \in C^{\infty} \left(V^{\frac{1}{2}} \right) .$	$\Psi = \psi \otimes \tilde{b} \in C^{\infty} (V^{\frac{1}{2}}) .$
mielke-geometrodynamics_344 E01847 (15.2.4)		■ 0.536	$\begin{aligned} \bar{E}_i &= \frac{1}{2} \left\{ E_{i\alpha}^{(f)j} \gamma^{\alpha} + i^* E_{i\alpha}^{(f)j} \gamma^{\alpha} \gamma^5 \right\} \lambda_j^{(f)} \\ &\oplus \frac{1}{2} \left\{ E_{i\alpha}^{(c)j} \gamma^{\alpha} + i^* E_{i\alpha}^{(c)j} \gamma^{\alpha} \gamma^5 \right\} \lambda_j^{(c)} \end{aligned}$	$\bar{E}_i = \frac{1}{2} \left\{ E_{i\alpha}^{(f)j} \gamma^{\alpha} + i^* E_{i\alpha}^{(f)j} \gamma^{\alpha} \gamma^5 \right\} \lambda_j^{(f)} \\ \oplus \frac{1}{2} \left\{ E_{i\alpha}^{(c)j} \gamma^{\alpha} + i^* E_{i\alpha}^{(c)j} \gamma^{\alpha} \gamma^5 \right\} \lambda_j^{(c)}$	$\bar{E}_i = \frac{1}{2} \left\{ E_{i\alpha}^{(f)j} \gamma^{\alpha} + i^* E_{i\alpha}^{(f)j} \gamma^{\alpha} \gamma^5 \right\} \lambda_j^{(f)} \\ \oplus \frac{1}{2} \left\{ E_{i\alpha}^{(c)j} \gamma^{\alpha} + i^* E_{i\alpha}^{(c)j} \gamma^{\alpha} \gamma^5 \right\} \lambda_j^{(c)}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_317 EQ9982 (14.2.23)		■ 0.537	$\int_G \pi \hat{L}_E = \frac{1}{\ell^2} L_W + L'_{YM} + \frac{1}{2\ell^2} \vartheta \wedge \vartheta \wedge \vartheta \wedge \vartheta \int_G \mathring{R} d\mu(G).$	$\int_G \pi \hat{L}_E = \frac{1}{\ell^2} L_W + L'_{YM} + \frac{1}{2\ell^2} \vartheta \wedge \vartheta \wedge \vartheta \wedge \vartheta \int_G \mathring{R} d\mu(G).$	$\int_G \pi \hat{L}_E = \frac{1}{\ell^2} L_W + L'_{YM} + \frac{1}{2\ell^2} \vartheta \wedge \vartheta \wedge \vartheta \wedge \vartheta \int_G \mathring{R} d\mu(G).$
mielke-geometrodynamics_375 EQ1134 (C.2)		■ 0.539	$\exp : \mathfrak{g} \rightarrow G \ni g = \exp \left(\iota_{\xi^j} I_j \right)$	$\exp : \mathfrak{g} \rightarrow G \ni g = \exp \left(\iota_{\xi^j} I_j \right)$	$\exp : \mathfrak{g} \rightarrow G \ni g = \exp (\iota \xi^j I_j)$
mielke-geometrodynamics_323 EQ1002 (14.3.15)		■ 0.541	$(i \not{D})^2 = (-1)^{3n-1+s} \operatorname{Tr} \{ \gamma \wedge \gamma \wedge^* (DD) - \gamma \wedge (D\gamma) \wedge^* D \}$	$(i \not{D})^2 = (-1)^{3n-1+s} \operatorname{Tr} \{ \gamma \wedge \gamma \wedge^* (DD) - \gamma \wedge (D\gamma) \wedge^* D \}$	$(i \not{D})^2 = (-1)^{3n-1+s} \operatorname{Tr} \{ \gamma \wedge \gamma \wedge^* (D D) - \gamma \wedge (D \gamma) \wedge^* D \}$
mielke-geometrodynamics_365 EQ1085 (A.11)		■ 0.541	$C(0,1) = \mathbb{C}$ (field of complex numbers), $C(1,0) = \mathbb{R} \oplus \mathbb{R},$	$C(0,1) = \mathbb{C}$ (field of complex numbers), $C(1,0) = \mathbb{R} \oplus \mathbb{R},$	$C(0,1) = \mathbb{C}$ (field of complex numbers), $C(1,0) = \mathbb{R} \oplus \mathbb{R},$
mielke-geometrodynamics_33 EQ0004 (2.1.3)		■ 0.543	$e_{\alpha}(m) = e_{,\alpha}^i(m) \partial_i.$	$e_{\alpha}(m) = e_{,\alpha}^i(m) \partial_i.$	$e_{\alpha}(m) = e_{,\alpha}^i(m) \partial_i.$
mielke-geometrodynamics_181 EQ0558 (8.5.1)		■ 0.544	$\Delta L_{\mathrm{FP}} = s \operatorname{Tr} \left\{ \frac{\theta_{\mathrm{T}}^* i}{4\ell^2} \bar{\chi} \wedge^* \sigma - \bar{c} d^* \gamma \right\}$	$\Delta L_{\mathrm{FP}} = s \operatorname{Tr} \left\{ \frac{\theta_{\mathrm{T}}^* i}{4\ell^2} \bar{\chi} \wedge^* \sigma - \bar{c} d^* \gamma \right\}$	$\Delta L_{\mathrm{FP}} = s \operatorname{Tr} \left\{ \frac{\theta_{\mathrm{T}}^* i}{4\ell^2} \bar{\chi} \wedge^* \sigma - \bar{c} d^* \gamma \right\}$
mielke-geometrodynamics_102 EQ0304 (4.5.14)		■ 0.545	$D\Sigma_{\alpha} \simeq 0$ $D(\tau_{\alpha\beta} + x_{[\alpha} \wedge \Sigma_{\beta]}) \simeq 0.$	$D\Sigma_{\alpha} \simeq 0$ $D(\tau_{\alpha\beta} + x_{[\alpha} \wedge \Sigma_{\beta]}) \simeq 0.$	$D\Sigma_{\alpha} \simeq 0,$ $D(\tau_{\alpha\beta} + x_{[\alpha} \wedge \Sigma_{\beta]}) \simeq 0.$
mielke-geometrodynamics_95 EQ0270 (4.3.35)		■ 0.546	$R := R_{\mu}^{\mu} = R_{\alpha\mu}^{\cdot\mu\alpha} = * (\Omega^g \wedge \vartheta \wedge \vartheta).$	$R := R_{\mu}^{\mu} = R_{\alpha\mu}^{\cdot\mu\alpha} = * (\Omega^g \wedge \vartheta \wedge \vartheta).$	$R := R_{\mu}^{\mu} = R_{\alpha\mu}^{\cdot\mu\alpha} = * (\Omega^g \wedge \vartheta \wedge \vartheta).$
mielke-geometrodynamics_71 EQ0174 (3.7.2)		■ 0.550	$D^* \Theta^{(1)} = \frac{\lambda}{2} \left(\frac{\mu^2}{\lambda} - \bar{\varphi} \cdot \varphi \right) \varphi \eta$ $\Theta^{(1)} = D\phi^{(0)}$	$D^* \Theta^{(1)} = \frac{\lambda}{2} \left(\frac{\mu^2}{\lambda} - \bar{\varphi} \cdot \varphi \right) \varphi \eta$ $\Theta^{(1)} = D\phi^{(0)}$	$D^* \Theta^{(1)} = \frac{\lambda}{2} \left(\frac{\mu^2}{\lambda} - \bar{\varphi} \cdot \varphi \right) \varphi \eta,$ $\Theta^{(1)} = D\phi^{(0)}.$
mielke-geometrodynamics_134 EQ0404 (6.4.24)		■ 0.551	$-\ell^2 \frac{\delta}{\delta \vartheta_B} \Psi_{\parallel}(\vartheta) = \mathring{T}^{(\pm) B} \Psi_{\parallel}(\vartheta).$	$-\ell^2 \frac{\delta}{\delta \vartheta_B} \Psi_{\parallel}(\vartheta) = \mathring{T}^{(\pm) B} \Psi_{\parallel}(\vartheta).$	$-\ell^2 \frac{\delta}{\delta \vartheta_B} \Psi_{\parallel}(\vartheta) = \mathring{T}^{(\pm) B} \Psi_{\parallel}(\vartheta).$
mielke-geometrodynamics_132 EQ0392 (6.4.7)		■ 0.552	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - (n \rfloor D \Psi) \wedge \frac{\partial L}{\partial D \Psi} - (n \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - (n \rfloor D \Psi) \wedge \frac{\partial L}{\partial D \Psi} - (n \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$	$n^{\alpha} \Sigma_{\alpha} = n \rfloor L - (n \rfloor D \Psi) \wedge \frac{\partial L}{\partial D \Psi} - (n \rfloor \Psi) \wedge \frac{\partial L}{\partial \Psi}.$
mielke-geometrodynamics_285 EQ0879 (13.3.4)		■ 0.553	$\mathring{L}_{AB} := \hat{g}_{[A C} C^C{}_{ B]} = -\mathring{L}_{BA}$	$\mathring{L}_{AB} := \hat{g}_{[A C} C^C{}_{ B]} = -\mathring{L}_{BA}$	$\mathring{L}_{AB} := \hat{g}_{[A C} \mathring{E}^C{}_{ B]} = -\mathring{L}_{BA}$
mielke-geometrodynamics_34 EQ0013 (2.2.8)		■ 0.554	$ds^2 = g_{ij} dx^i \otimes_s dx^j \stackrel{(*)}{=} - \sum_{i=1}^s \left(\mathring{\vartheta}^i \right)^2 + \sum_{j=s+1}^n \left(\mathring{\vartheta}^j \right)^2.$	$ds^2 = g_{ij} dx^i \otimes_s dx^j \stackrel{(*)}{=} - \sum_{i=1}^s \left(\mathring{\vartheta}^i \right)^2 + \sum_{j=s+1}^n \left(\mathring{\vartheta}^j \right)^2.$	$ds^2 = g_{ij} dx^i \otimes_s dx^j \stackrel{(*)}{=} - \sum_{i=1}^s (\mathring{\vartheta}^i)^2 + \sum_{j=s+1}^n (\mathring{\vartheta}^j)^2.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_325 EQ1011 (14.4.6)		■ 0.554	$r_{(-)}^{+}=\frac{\hbar}{M^{*2}c}\left(M_{(-)}^{+}\sqrt{M^2-J(J+1)\frac{M^{*4}}{M^2}-\alpha_eM^{*2}\mathrm{Tr}\bar{Q}^2}\right)$	$r_{(-)}^{+}=\frac{\hbar}{M^{*2}c}\left(M_{(-)}^{+}\sqrt{M^2-J(J+1)\frac{M^{*4}}{M^2}-\alpha_eM^{*2}\mathrm{Tr}\bar{Q}^2}\right)$	$r_{(-)}^{+}=\frac{\hbar}{M^{*2}c}\left(M_{(-)}^{+}\sqrt{M^2-J(J+1)\frac{M^{*4}}{M^2}-\alpha_eM^{*2}\mathrm{Tr}\bar{Q}^2}\right)$
mielke-geometrodynamics_130 EQ0385 (6.3.3)		■ 0.561	$\stackrel{\scriptscriptstyle{(\pm)}}{\Gamma}_{\alpha}^{\scriptscriptstyle{(\pm)}}\beta\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\beta}+\Sigma_{\alpha}\cong d\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},\quad\stackrel{\scriptscriptstyle{(\pm)}}{S}_{\alpha}=\stackrel{\scriptscriptstyle{(\mp)}}{S}_{\alpha}$	$\stackrel{\scriptscriptstyle{(\pm)}}{S}_{\alpha}:=\stackrel{\scriptscriptstyle{(\pm)}}{\Gamma}_{\alpha}^{\scriptscriptstyle{(\pm)}}\beta\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\beta}+\Sigma_{\alpha}\cong d\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},\quad\stackrel{\scriptscriptstyle{(\pm)}}{S}_{\alpha}=\stackrel{\scriptscriptstyle{(\mp)}}{S}_{\alpha}$	$\stackrel{\scriptscriptstyle{(\pm)}}{S}_{\alpha}:=\stackrel{\scriptscriptstyle{(\pm)}}{\Gamma}_{\alpha}^{\scriptscriptstyle{(\pm)}}\beta\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\beta}+\Sigma_{\alpha}\cong d\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha},\quad\stackrel{\scriptscriptstyle{(\pm)}}{S}_{\alpha}=\stackrel{\scriptscriptstyle{(\mp)}}{S}_{\alpha}$
mielke-geometrodynamics_195 EQ0596 (9.1.14)		■ 0.564	$(\stackrel{\scriptscriptstyle{+}}{D}K)-K\wedge K=\stackrel{\scriptscriptstyle{+}}{\Omega}_C\{\}$	$(\stackrel{\scriptscriptstyle{+}}{D}K)-K\wedge K=\stackrel{\scriptscriptstyle{+}}{\Omega}_C\{\}.$	$(\stackrel{\scriptscriptstyle{+}}{D}K)-K\wedge K=\stackrel{\scriptscriptstyle{+}}{\Omega}_C\{\}.$
mielke-geometrodynamics_70 EQ0169 (3.6.4)		■ 0.570	$D_{\pm}\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)=\mp\frac{i}{2}\bar{\psi}^*D_{\pm}\psi=\mp\frac{1}{2}\bar{\psi}^*\gamma\wedge^*(i^*\gamma\wedge D_{\pm}\psi).$	$D_{\pm}\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)=\mp\frac{i}{2}\bar{\psi}^*D_{\pm}\psi=\mp\frac{1}{2}\bar{\psi}^*\gamma\wedge^*(i^*\gamma\wedge D_{\pm}\psi).$	$D_{\pm}\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)=\mp\frac{i}{2}\bar{\psi}^*D_{\pm}\psi=\mp\frac{1}{2}\bar{\psi}^*\gamma\wedge^*(i^*\gamma\wedge D_{\pm}\psi).$
mielke-geometrodynamics_286 EQ0889 (13.4.8)		■ 0.576	$\begin{aligned} & \Gamma^{\scriptscriptstyle{(T)}}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}},\quad \Gamma^{\scriptscriptstyle{(T)}}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}}=\left[\Gamma^{\scriptscriptstyle{(T)}}+D\tau(x)\right]\Lambda(x); \\ & \Gamma^{\scriptscriptstyle{(T)}}_{*}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}}_{*}=\left[\Gamma^{\scriptscriptstyle{(T)}}_{*}+D\tau_{*}(x)\right]\Lambda(x); \end{aligned}$	$\Gamma^{\scriptscriptstyle{(T)}}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}}=\Lambda^{-1}(x)\left[\Gamma^{\scriptscriptstyle{(T)}}+D\tau(x)\right],$ $\Gamma^{\scriptscriptstyle{(T)}}_{*}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}}_{*}=\left[\Gamma^{\scriptscriptstyle{(T)}}_{*}+D\tau_{*}(x)\right]\Lambda(x);$	$\Gamma^{\scriptscriptstyle{(T)}}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}}=\Lambda^{-1}(x)\left[\Gamma^{\scriptscriptstyle{(T)}}+D\tau(x)\right],$ $\Gamma^{\scriptscriptstyle{(T)}}_{*}\xrightarrow{A^{-1}(x)}\Gamma^{\scriptscriptstyle{(T)'}}_{*}=\left[\Gamma^{\scriptscriptstyle{(T)}}_{*}+D\tau_{*}(x)\right]\Lambda(x);$
mielke-geometrodynamics_59 EQ0116 (3.3.17)		■ 0.585	$dH=j\left[\begin{array}{l}\boldsymbol{\nabla}\times\boldsymbol{B}-\frac{\partial\boldsymbol{E}}{c^2\partial t}=\boldsymbol{\mu J}\\\boldsymbol{\nabla}\cdot\boldsymbol{E}=\rho/\varepsilon.\end{array}\right].$	$dH=j\left\{\begin{array}{l}\boldsymbol{\nabla}\times\boldsymbol{B}-\frac{\partial\boldsymbol{E}}{c^2\partial t}=\boldsymbol{\mu J}\\\boldsymbol{\nabla}\cdot\boldsymbol{E}=\rho/\varepsilon.\end{array}\right.$	$dH=j\left\{\begin{array}{l}\boldsymbol{\nabla}\times\boldsymbol{B}-\frac{\partial\boldsymbol{E}}{c^2\partial t}=\boldsymbol{\mu J}\\\boldsymbol{\nabla}\cdot\boldsymbol{E}=\rho/\varepsilon.\end{array}\right.$
mielke-geometrodynamics_129 EQ0381 (6.2.32)		■ 0.588	$\stackrel{\scriptscriptstyle{(\pm)}}{D}\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}:=D\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}\pm\frac{i}{2}\ell^2\left(e_{\alpha}\lrcorner\stackrel{\scriptscriptstyle{(\mp)}}{\Pi}\right)\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\beta},$	$\stackrel{\scriptscriptstyle{(\pm)}}{D}\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}:=D\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}\pm\frac{i}{2}\ell^2\left(e_{\alpha}\lrcorner\stackrel{\scriptscriptstyle{(\mp)}}{\Pi}\right)\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\beta},$	$\stackrel{\scriptscriptstyle{(\pm)}}{D}\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}:=D\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\alpha}\pm\frac{i}{2}\ell^2\left(e_{\alpha}\lrcorner\stackrel{\scriptscriptstyle{(\mp)}}{\Pi}\right)\wedge\stackrel{\scriptscriptstyle{(\pm)}}{\Pi}_{\beta},$
mielke-geometrodynamics_45 EQ0056 (2.7.6)		■ 0.592	$\phi=\varphi\otimes b=G^{-1}\varphi\otimes^Gb\in C^\infty(V^\rho).$	$\phi=\varphi\otimes b=G^{-1}\varphi\otimes^Gb\in C^\infty(V^\rho).$	$\phi=\varphi\otimes b=G^{-1}\varphi\otimes^Gb\in C^\infty(V^\rho).$
mielke-geometrodynamics_74 EQ0187 (3.7.17)		■ 0.593	$\begin{aligned} & \left(D^*\Omega\right)=\vec{D}^*\vec{\Omega}+\left[\omega_0,\Omega_0\right] \\ & =\vec{D}^*\vec{\Omega}+\left[\omega_0,\Theta^\perp\right]=0. \end{aligned}$	$\left(D^*\Omega\right)=\vec{D}^*\vec{\Omega}+\left[\omega_0,\Omega_0\right]$ $=\vec{D}^*\vec{\Omega}+\left[\omega_0,\Theta^\perp\right]=0.$	$\left(D^*\Omega\right)=\vec{D}^*\vec{\Omega}+\left[\omega_0,\Omega_0\right]$ $=\vec{D}^*\vec{\Omega}+\left[\omega_0,\Theta^\perp\right]=0.$
mielke-geometrodynamics_186 EQ0581 (8.6.19)		■ 0.593	$\widehat{\Sigma}_{\alpha}:=\Sigma_{\alpha}-D^{\{\}}\mu_{\alpha},$	$\widehat{\Sigma}_{\alpha}:=\Sigma_{\alpha}-D^{\{\}}\mu_{\alpha},$	$\widehat{\Sigma}_{\alpha}:=\Sigma_{\alpha}-D^{\{\}}\mu_{\alpha},$
mielke-geometrodynamics_307 EQ0948 (14.1.29)		■ 0.593	$\begin{aligned} & D^*\Omega=(D^*\xi-(-1)^z\xi\wedge D\delta)\cos\delta \\ & \quad -(-1)^z(D\xi+(-1)^{z*}\xi\wedge D\delta)\sin\delta=\tau_e \end{aligned}$	$D^*\Omega=(D^*\xi-(-1)^z\xi\wedge D\delta)\cos\delta$ $-(-1)^z(D\xi+(-1)^{z*}\xi\wedge D\delta)\sin\delta=\tau_e$	$D^*\Omega=(D^*\xi-(-1)^z\xi\wedge D\delta)\cos\delta$ $-(-1)^z(D\xi+(-1)^{z*}\xi\wedge D\delta)\sin\delta=\tau_e$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_125 EQ0357 (6.2.8)	125	■ 0.600	$L_{\mathrm{HE}}=-\frac{1}{2\kappa}R_{\alpha\beta}^{\{\}}\wedge^*(\vartheta^{\alpha}\wedge\vartheta^{\beta})$	$L_{\mathrm{HE}}=-\frac{1}{2\kappa}R_{\alpha\beta}^{\{\}}\wedge^*(\vartheta^{\alpha}\wedge\vartheta^{\beta})$	$L_{\mathrm{HE}}=-\frac{1}{2\kappa}R_{\alpha\beta}^{\{\}}\wedge^*(\vartheta^{\alpha}\wedge\vartheta^{\beta})$
mielke-geometrodynamics_225 EQ0704 (10.6.3)	225	■ 0.601	$\square F-m_{\mathrm{photon}}^2F=0,$	$\square F-m_{\mathrm{photon}}^2F=0,$	$\square F-m_{\mathrm{photon}}^2F=0,$
mielke-geometrodynamics_303 EQ0927 (14.1.8)	303	■ 0.604	$\begin{aligned} \Sigma_{\mathrm{YM}}&=\frac{(-1)^s}{2\alpha_g}\mathrm{Tr}\left\{\Omega\wedge^*(\vartheta\wedge^*\Omega)+^*\vartheta\wedge^*(\Omega\wedge^*\Omega)\right\} \\ &=\frac{(-1)^s}{4\alpha_g}\mathrm{Tr}\left\{\Omega\wedge^*(\vartheta\wedge^*\Omega)-^*\Omega\wedge^*(\vartheta\wedge\Omega)\right\}. \end{aligned}$	$\Sigma_{\mathrm{YM}}=\frac{(-1)^s}{2\alpha_g}\mathrm{Tr}\left\{\Omega\wedge^*(\vartheta\wedge^*\Omega)+^*\vartheta\wedge^*(\Omega\wedge^*\Omega)\right\} \\ =\frac{(-1)^s}{4\alpha_g}\mathrm{Tr}\left\{\Omega\wedge^*(\vartheta\wedge^*\Omega)-^*\Omega\wedge^*(\vartheta\wedge\Omega)\right\}.$	$\begin{aligned} \Sigma_{\mathrm{YM}}&=\frac{(-1)^s}{2\alpha_g}\mathrm{Tr}\left\{\Omega\wedge^*(\vartheta\wedge^*\Omega)+^*\vartheta\wedge^*(\Omega\wedge^*\Omega)\right\} \\ &=\frac{(-1)^s}{4\alpha_g}\mathrm{Tr}\left\{\Omega\wedge^*(\vartheta\wedge^*\Omega)-^*\Omega\wedge^*(\vartheta\wedge\Omega)\right\}. \end{aligned}$
mielke-geometrodynamics_112 EQ0326 (5.3.3)	112	■ 0.613	$\left.\widehat{D}G_{\alpha}\right _{\mathrm{e}_{\alpha}}\equiv\frac{1}{2}\left(e_{\alpha}\rfloor R^{\beta\gamma}\right)\wedge\eta_{\beta\gamma\mu}\wedge T^{\mu}$	$\widehat{D}G_{\alpha}\equiv\frac{1}{2}(e_{\alpha}\rfloor R^{\beta\gamma})\wedge\eta_{\beta\gamma\mu}\wedge T^{\mu}$	$\widehat{D}G_{\alpha}\equiv\frac{1}{2}(e_{\alpha}\rfloor R^{\beta\gamma})\wedge\eta_{\beta\gamma\mu}\wedge T^{\mu}$
mielke-geometrodynamics_242 EQ0758 (11.1.25)	242	■ 0.615	$\gamma=E_{\cdot j}^{\alpha}(m)\gamma_{\alpha}\otimes dx^j=\gamma_jdx^j\in C^{\infty}(T_{\mathbb{C}}^*(M)\times CV)$	$\gamma=E_{\cdot j}^{\alpha}(m)\gamma_{\alpha}\otimes dx^j=\gamma_jdx^j\in C^{\infty}(T_{\mathbb{C}}^*(M)\times CV)$	$\gamma=E_{\cdot j}^{\alpha}(m)\gamma_{\alpha}\otimes dx^j=\gamma_jdx^j\in C^{\infty}(T_{\mathbb{C}}^*(M)\times CV)$
mielke-geometrodynamics_308 EQ0953 (14.1.35)	308	■ 0.616	$\begin{aligned} &(-1)^z\mathrm{Tr}\{\vartheta\wedge^*\xi\wedge^*\xi-\vartheta\wedge\xi\wedge\xi\}\wedge d\delta=\mathrm{Tr}\{\xi\wedge\vartheta\wedge D^*\xi+^*\xi\wedge\vartheta\wedge D\xi\} \\ &=\frac{1}{2}D\mathrm{Tr}\left\{^*(\xi\wedge^*(\vartheta\wedge^*\xi))-^*\xi\wedge^*(\vartheta\wedge\xi)\right\} \\ &=(-1)^s2\alpha_gD\Sigma_{\mathrm{YM}}=\frac{\alpha_g}{\ell^{*2}}D(\vartheta\wedge^*\Omega^g)\equiv2\frac{\alpha_g}{\ell^{*2}}\Theta\wedge^*\Omega^g. \end{aligned}$	$\begin{aligned} &(-1)^z\mathrm{Tr}\{\vartheta\wedge^*\xi\wedge^*\xi-\vartheta\wedge\xi\wedge\xi\}\wedge d\delta=\mathrm{Tr}\{\xi\wedge\vartheta\wedge D^*\xi+^*\xi\wedge\vartheta\wedge D\xi\} \\ &=\frac{1}{2}D\mathrm{Tr}\left\{^*(\xi\wedge^*(\vartheta\wedge^*\xi))-^*\xi\wedge^*(\vartheta\wedge\xi)\right\} \\ &=(-1)^s2\alpha_gD\Sigma_{\mathrm{YM}}=\frac{\alpha_g}{\ell^{*2}}D(\vartheta\wedge^*\Omega^g)\equiv2\frac{\alpha_g}{\ell^{*2}}\Theta\wedge^*\Omega^g. \end{aligned}$	$\begin{aligned} &(-1)^z\mathrm{Tr}\{\vartheta\wedge^*\xi\wedge^*\xi-\vartheta\wedge\xi\wedge\xi\}\wedge d\delta=\mathrm{Tr}\{\xi\wedge\vartheta\wedge D^*\xi+^*\xi\wedge\vartheta\wedge D\xi\} \\ &=\frac{1}{2}D\mathrm{Tr}\left\{^*(\xi\wedge^*(\vartheta\wedge^*\xi))-^*\xi\wedge^*(\vartheta\wedge\xi)\right\} \\ &=(-1)^s2\alpha_gD\Sigma_{\mathrm{YM}}=\frac{\alpha_g}{\ell^{*2}}D(\vartheta\wedge^*\Omega^g)\equiv2\frac{\alpha_g}{\ell^{*2}}\Theta\wedge^*\Omega^g. \end{aligned}\tag{14.1.3}$
mielke-geometrodynamics_322 EQ0998 (14.3.11)	322	■ 0.619	$\begin{array}{l} \psi_{\mathrm{L}}:\quad\gamma^5=+1,\quad\overset{\circ}{\gamma}^{K+1}=+\widehat{\gamma}^{n+1} \\ \psi_{\mathrm{R}}:\quad\gamma^5=-1,\quad\overset{\circ}{\gamma}^{K+1}=-\widehat{\gamma}^{n+1} \end{array}$	$\begin{array}{l} \psi_{\mathrm{L}}:\quad\gamma^5=+1,\quad\overset{\circ}{\gamma}^{K+1}=+\widehat{\gamma}^{n+1} \\ \psi_{\mathrm{R}}:\quad\gamma^5=-1,\quad\overset{\circ}{\gamma}^{K+1}=-\widehat{\gamma}^{n+1} \end{array}$	$\begin{array}{l} \psi_{\mathrm{L}}:\quad\gamma^5=+1,\quad\overset{\circ}{\gamma}^{K+1}=+\widehat{\gamma}^{n+1} \\ \psi_{\mathrm{R}}:\quad\gamma^5=-1,\quad\overset{\circ}{\gamma}^{K+1}=-\widehat{\gamma}^{n+1} \end{array}$
mielke-geometrodynamics_167 EQ0525 (7.7.44)	167	■ 0.623	$\begin{aligned} C_{\mathrm{TT}}&:=\frac{1}{8\ell^2}\mathrm{Tr}(\gamma\wedge\Theta)=\frac{1}{2\ell^2}\vartheta^{\alpha}\wedge T_{\alpha}=-\frac{(-1)^{\mathrm{sig}}}{2\ell^2}*{\mathscr{A}}, \\ C_{\mathrm{RR}}&:=\mathrm{Tr}\left(\Gamma\wedge\Omega^g-\frac{1}{3}\Gamma\wedge\Gamma\wedge\Gamma\right), \end{aligned}$	$\begin{aligned} C_{\mathrm{TT}}&:=\frac{1}{8\ell^2}\mathrm{Tr}(\gamma\wedge\Theta)=\frac{1}{2\ell^2}\vartheta^{\alpha}\wedge T_{\alpha}=-\frac{(-1)^{\mathrm{sig}}}{2\ell^2}*{\mathscr{A}}, \\ C_{\mathrm{RR}}&:=\mathrm{Tr}\left(\Gamma\wedge\Omega^g-\frac{1}{3}\Gamma\wedge\Gamma\wedge\Gamma\right), \end{aligned}$	$\begin{aligned} C_{\mathrm{TT}}&:=\frac{1}{8\ell^2}\mathrm{Tr}(\gamma\wedge\Theta)=\frac{1}{2\ell^2}\vartheta^{\alpha}\wedge T_{\alpha}=-\frac{(-1)^{\mathrm{sig}}}{2\ell^2}*{\mathscr{A}}, \\ C_{\mathrm{RR}}&:=\mathrm{Tr}\left(\Gamma\wedge\Omega^g-\frac{1}{3}\Gamma\wedge\Gamma\wedge\Gamma\right), \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_ EQ0050 (2.6.13)	44	■ 0.626	$\begin{aligned} D D D b = & D(\Omega \otimes b) = (d\Omega + [\Omega, \omega]) \otimes b \\ & = (d\Omega + \Omega \wedge \omega - \omega \wedge \Omega) \otimes b \\ & = (dd\omega - d\omega \wedge \omega + \omega \wedge d\omega \\ & \quad + d\omega \wedge \omega - \omega \wedge \omega \wedge \omega \\ & \quad + \omega \wedge \omega \wedge \omega - \omega \wedge d\omega) \otimes b \equiv 0. \end{aligned}$	$DDDb = D(\Omega \otimes b) = (d\Omega + [\Omega, \omega]) \otimes b \\ = (d\Omega + \Omega \wedge \omega - \omega \wedge \Omega) \otimes b \\ = (dd\omega - d\omega \wedge \omega + \omega \wedge d\omega \\ + d\omega \wedge \omega - \omega \wedge \omega \wedge \omega \\ + \omega \wedge \omega \wedge \omega - \omega \wedge d\omega) \otimes b \equiv 0.$	$DDDb = D(\Omega \otimes b) = (d\Omega + [\Omega, \omega]) \otimes b \\ = (d\Omega + \Omega \wedge \omega - \omega \wedge \Omega) \otimes b \\ = (dd\omega - d\omega \wedge \omega + \omega \wedge d\omega \\ + d\omega \wedge \omega - \omega \wedge \omega \wedge \omega \\ + \omega \wedge \omega \wedge \omega - \omega \wedge d\omega) \otimes b \equiv 0.$
mielke-geometrodynamics_ EQ0115 (3.3.16)	59	■ 0.630	$d F \equiv 0 \quad \text{quad} \left\{ \begin{array}{l} \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} \equiv 0 \\ \nabla \cdot \mathbf{B} \equiv 0 \end{array} \right.$	$dF \equiv 0 \quad \left\{ \begin{array}{l} \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} \equiv 0 \\ \nabla \cdot \mathbf{B} \equiv 0 \end{array} \right.$	$dF \equiv 0 \quad \left\{ \begin{array}{l} \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} \equiv 0 \\ \nabla \cdot \mathbf{B} \equiv 0. \end{array} \right.$
mielke-geometrodynamics_ EQ0171 (3.6.6)	70	■ 0.630	$i^{\gamma} \wedge \gamma \wedge D_{\pm} \psi = 0 \quad \text{and} \quad F^{\pm} = \frac{1}{2} \bar{\psi} \sigma_{\pm} \psi = m \mathcal{P}_{\pm},$	$i^* \gamma \wedge D_{\pm} \psi = 0 \quad \text{and} \quad F^{\pm} = \frac{1}{2} \bar{\psi} \sigma_{\pm} \psi = m \mathcal{P}_{\pm},$	$i^* \gamma \wedge D_{\pm} \psi = 0 \quad \text{and} \quad F^{\pm} = \frac{1}{2} \bar{\psi} \sigma_{\pm} \psi = m \mathcal{P}_{\pm},$
mielke-geometrodynamics_ EQ0189 (3.7.19)	74	■ 0.631	$\{ \}^{\gamma} \wedge \Omega = (\stackrel{+}{-}) \Omega \quad \text{text } \{. \}$	$^* \Omega = (\stackrel{+}{-}) \Omega.$	$^* \Omega = (\stackrel{+}{-}) \Omega.$
mielke-geometrodynamics_ EQ0788 (11.3.7)	249	■ 0.635	$\left\{ i \gamma^{\kappa} \nabla_{\kappa}^{(l)} - \frac{6}{n^2} \ell^{*2} \gamma^{n+1} \gamma^{\mu} \bar{\psi} \gamma^{n+1} \gamma_{\mu} \psi - m \right\} \psi = 0.$	$\left\{ i \gamma^{\kappa} \nabla_{\kappa}^{(l)} - \frac{6}{n^2} \ell^{*2} \gamma^{n+1} \gamma^{\mu} \bar{\psi} \gamma^{n+1} \gamma_{\mu} \psi - m \right\} \psi = 0.$	$\left\{ i \gamma^{\kappa} \nabla_{\kappa}^{(l)} - \frac{6}{n^2} \ell^{*2} \gamma^{n+1} \gamma^{\mu} \bar{\psi} \gamma^{n+1} \gamma_{\mu} \psi - m \right\} \psi = 0.$
mielke-geometrodynamics_ EQ0164 (3.5.12)	68	■ 0.636	$L_{\mathrm{YM}} + L_{\infty} = (\stackrel{+}{-}) \frac{1}{2\alpha_g} \mathrm{Tr} \left(^* \Omega \stackrel{-}{+} \Omega \right)^2 \geq 0$	$L_{\mathrm{YM}} + L_{\infty} = (\stackrel{+}{-}) \frac{1}{2\alpha_g} \mathrm{Tr} \left(^* \Omega \stackrel{-}{+} \Omega \right)^2 \geq 0$	$L_{\mathrm{YM}} + L_{\infty} = (\stackrel{+}{-}) \frac{1}{2\alpha_g} \mathrm{Tr} (^* \Omega \stackrel{-}{+} \Omega)^2 \geq 0$
mielke-geometrodynamics_ EQ0549 (8.4.3)	179	■ 0.637	$\begin{aligned} L_{\mathrm{FP}} &:= -s \mathrm{Tr} \left[\bar{\chi} \wedge ^* \Omega^g + \bar{\Phi} D^* \Psi + \frac{1}{2} \rho \bar{\chi} \wedge ^* \beta + \bar{C} d^* \Gamma + \frac{1}{2} \bar{C} \wedge ^* \mathbb{B} \right] \\ &= \frac{1}{2} s \left[\bar{\chi}^{\alpha\beta} \wedge ^* R_{\alpha\beta}^{(\pm)} + \bar{\Phi}_{\alpha\beta} D^* \Psi^{\alpha\beta} + \frac{1}{2} \rho \bar{\chi}^{\alpha\beta} \wedge ^* b_{\alpha\beta} + \tilde{\Psi}_{\mathrm{F}} \right] \end{aligned}$	$L_{\mathrm{FP}} := -s \mathrm{Tr} \left\{ \bar{\chi} \wedge ^* \Omega^g + \bar{\Phi} D^* \Psi + \frac{1}{2} \rho \bar{\chi} \wedge ^* \beta + \bar{C} d^* \Gamma + \frac{1}{2} \bar{C} \wedge ^* \mathbb{B} \right\} \\ = \frac{1}{2} s \left[\bar{\chi}^{\alpha\beta} \wedge ^* R_{\alpha\beta}^{(\pm)} + \bar{\Phi}_{\alpha\beta} D^* \Psi^{\alpha\beta} + \frac{1}{2} \rho \bar{\chi}^{\alpha\beta} \wedge ^* b_{\alpha\beta} + \tilde{\Psi}_{\mathrm{F}} \right]$	$L_{\mathrm{FP}} := -s \mathrm{Tr} \left\{ \bar{\chi} \wedge ^* \Omega^g + \bar{\Phi} D^* \Psi + \frac{1}{2} \rho \bar{\chi} \wedge ^* \beta + \bar{C} d^* \Gamma + \frac{1}{2} \bar{C} \wedge ^* \mathbb{B} \right\} \\ = \frac{1}{2} s \left[\bar{\chi}^{\alpha\beta} \wedge ^* R_{\alpha\beta}^{(\pm)} + \bar{\Phi}_{\alpha\beta} D^* \Psi^{\alpha\beta} + \frac{1}{2} \rho \bar{\chi}^{\alpha\beta} \wedge ^* b_{\alpha\beta} + \tilde{\Psi}_{\mathrm{F}} \right]. \quad (8)$
mielke-geometrodynamics_ EQ1058 (15.2.15)	346	■ 0.638	$\bar{D}^{\{ \} \} := d + i \bar{\omega}^{\{ \} \}$	$\bar{D}^{\{ \} \} := d + i \bar{\omega}^{\{ \} \}$	$\bar{D}^{\{ \} \} := d + i \bar{\omega}^{\{ \} \}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_901008 (14.4.3)	325	■ 0.640	$\begin{aligned} d s^2 &= -\frac{\Delta}{\rho^2} \left[c d t - J_3 \frac{\hbar}{M} \sin^2 \vartheta d \phi \right]^2 + \frac{\rho^2}{\Delta} d r + \rho^2 d \vartheta^2 \\ &+ \frac{\sin^2 \vartheta}{\rho^2} \left\{ \left[r^2 + J(J+1) \frac{\hbar^2}{M^2 c^2} \right] d \phi - J_3 \frac{\hbar}{M} d t \right\}^2 \end{aligned}$	$ds^2 = -\frac{\Delta}{\rho^2} \left[cdt - J_3 \frac{\hbar}{M} \sin^2 \vartheta d\phi \right]^2 + \frac{\rho^2}{\Delta} dr + \rho^2 d\vartheta^2 + \frac{\sin^2 \vartheta}{\rho^2} \left\{ \left[r^2 + J(J+1) \frac{\hbar^2}{M^2 c^2} \right] d\phi - J_3 \frac{\hbar}{M} dt \right\}^2$	$ds^2 = -\frac{\Delta}{\rho^2} \left[cdt - J_3 \frac{\hbar}{M} \sin^2 \vartheta d\phi \right]^2 + \frac{\rho^2}{\Delta} dr + \rho^2 d\vartheta^2 + \frac{\sin^2 \vartheta}{\rho^2} \left\{ \left[r^2 + J(J+1) \frac{\hbar^2}{M^2 c^2} \right] d\phi - J_3 \frac{\hbar}{M} dt \right\}^2.$
mielke-geometrodynamics_900255 (4.3.20)	93	■ 0.642	$\stackrel{*}{\Omega}^g = \frac{1}{4} R_{\alpha\beta}^{..cd} L_{cd} \otimes \vartheta^\alpha \wedge \vartheta^\beta.$	$\bar{\sigma}^* \Omega^g = \frac{1}{4} R_{\alpha\beta}^{..cd} L_{cd} \otimes \vartheta^\alpha \wedge \vartheta^\beta.$	$\bar{\sigma}^* \Omega^g = \frac{1}{4} R_{\alpha\beta}^{..cd} L_{cd} \otimes \vartheta^\alpha \wedge \vartheta^\beta.$
mielke-geometrodynamics_901015 (14.4.10)	327	■ 0.646	$A = \{ \}^* \omega = A^{(\circ)} \lambda_3.$	$A = \bar{\sigma} \omega = A^{(\circ)} \lambda_3.$	$A = \bar{\sigma} \omega = A^{(\circ)} \lambda_3.$
mielke-geometrodynamics_900119 (3.3.20)	59	■ 0.646	$\begin{aligned} \Sigma_\alpha &:= e_\alpha \rfloor L - (e_\alpha \rfloor dA) \wedge \frac{\partial L}{\partial(dA)} \\ &= \frac{1}{2} [(e_\alpha \rfloor F) \wedge H - F \wedge (e_\alpha \rfloor H)] \end{aligned}$	$\Sigma_\alpha := e_\alpha \rfloor L - (e_\alpha \rfloor dA) \wedge \frac{\partial L}{\partial(dA)} = \frac{1}{2} [(e_\alpha \rfloor F) \wedge H - F \wedge (e_\alpha \rfloor H)]$	$\Sigma_\alpha := e_\alpha \rfloor L - (e_\alpha \rfloor dA) \wedge \frac{\partial L}{\partial(dA)} = \frac{1}{2} [(e_\alpha \rfloor F) \wedge H - F \wedge (e_\alpha \rfloor H)]$
mielke-geometrodynamics_900120 (3.3.21)	60	■ 0.651	$d\Sigma_\alpha \simeq (e_\alpha \rfloor F) \wedge j$	$d\Sigma_\alpha \simeq (e_\alpha \rfloor F) \wedge j$	$d\Sigma_\alpha \simeq (e_\alpha \rfloor F) \wedge j$
mielke-geometrodynamics_900497 (7.7.15)	163	■ 0.653	$ds^2 = o_{\alpha\beta} \vartheta^\alpha \otimes \vartheta^\beta = g_{ij} dx^i \otimes dx^j, \quad o_{\alpha\beta} := \text{diag}\{1, \underbrace{-1, \dots, -1}_{\text{sig}}, \dots\},$	$ds^2 = o_{\alpha\beta} \vartheta^\alpha \otimes \vartheta^\beta = g_{ij} dx^i \otimes dx^j, \quad o_{\alpha\beta} := \text{diag}\{1, \underbrace{-1, \dots, -1}_{\text{sig}}, \dots\},$	$ds^2 = o_{\alpha\beta} \vartheta^\alpha \otimes \vartheta^\beta = g_{ij} dx^i \otimes dx^j, \quad o_{\alpha\beta} := \text{diag}\{1, \underbrace{-1, \dots, -1}_{\text{sig}}, \dots\},$
mielke-geometrodynamics_900800 (11.4.12)	254	■ 0.658	$\left\{ i\gamma^k \partial_k - \frac{3i}{2R_o \lambda} \gamma^0 - \frac{m}{\lambda} \right\} \psi = 0,$	$\left\{ i\gamma^k \partial_k - \frac{3i}{2R_o \lambda} \gamma^0 - \frac{m}{\lambda} \right\} \psi = 0,$	$\left\{ i\gamma^k \partial_k - \frac{3i}{2R_o \lambda} \gamma^0 - \frac{m}{\lambda} \right\} \psi = 0,$
mielke-geometrodynamics_900316 (4.6.8)	105	■ 0.664	$\Phi(\widetilde{\Gamma}, \gamma) = (2\pi \hbar \mathcal{M}_n G / \hbar \ell c^2) = 2\pi n$	$\Phi(\widetilde{\Gamma}, \gamma) = (2\pi \hbar \mathcal{M}_n G / \hbar \ell c^2) = 2\pi n$	$\Phi(\widetilde{\Gamma}, \gamma) = (2\pi \hbar \mathcal{M}_n G / \hbar \ell c^2) = 2\pi n,$
mielke-geometrodynamics_900335 (5.3.12)	113	■ 0.665	$\begin{aligned} \widetilde{T}_{\mu\nu} &= T_{\mu\nu} + \ell^{*2} \left\{ -4\tau_{\mu\cdot[\lambda]}^{\cdot\kappa} \tau_{\nu\cdot[\kappa]}^{\cdot\lambda} - 2\tau_{\mu}^{\cdot\kappa\lambda} \tau_{\nu\kappa\lambda} + \tau_{\cdot,\mu}^{\kappa\lambda} \tau_{\kappa\lambda\nu} \right. \\ &\quad \left. + \frac{1}{2} g_{\mu\nu} \left(4\tau_{\delta\cdot[\lambda]}^{\cdot\kappa} \tau_{\cdot,\kappa]}^{\delta\lambda} + \tau^{\delta\kappa\lambda} \tau_{\delta\kappa\lambda} \right) \right\}. \end{aligned}$	$\widetilde{T}_{\mu\nu} = T_{\mu\nu} + \ell^{*2} \left\{ -4\tau_{\mu\cdot[\lambda]}^{\cdot\kappa} \tau_{\nu\cdot[\kappa]}^{\cdot\lambda} - 2\tau_{\mu}^{\cdot\kappa\lambda} \tau_{\nu\kappa\lambda} + \tau_{\cdot,\mu}^{\kappa\lambda} \tau_{\kappa\lambda\nu} + \frac{1}{2} g_{\mu\nu} \left(4\tau_{\delta\cdot[\lambda]}^{\cdot\kappa} \tau_{\cdot,\kappa]}^{\delta\lambda} + \tau^{\delta\kappa\lambda} \tau_{\delta\kappa\lambda} \right) \right\}.$	$\widetilde{T}_{\mu\nu} = T_{\mu\nu} + \ell^{*2} \left\{ -4\tau_{\mu\cdot[\lambda]}^{\cdot\kappa} \tau_{\nu\cdot[\kappa]}^{\cdot\lambda} - 2\tau_{\mu}^{\cdot\kappa\lambda} \tau_{\nu\kappa\lambda} + \tau_{\cdot,\mu}^{\kappa\lambda} \tau_{\kappa\lambda\nu} + \frac{1}{2} g_{\mu\nu} (4\tau_{\delta\cdot[\lambda]}^{\cdot\kappa} \tau_{\cdot,\kappa]}^{\delta\lambda} + \tau^{\delta\kappa\lambda} \tau_{\delta\kappa\lambda}) \right\}.$
mielke-geometrodynamics_901055 (15.2.12)	345	■ 0.665	$\bar{L}_{\mathrm{w.}}(f) = \frac{i}{8} \text{Tr}(\bar{\gamma} \wedge \bar{\gamma} \wedge {}^* \bar{\Omega}).$	$\bar{L}_{\mathrm{w.}}(f) = \frac{i}{8} \text{Tr}(\bar{\gamma} \wedge \bar{\gamma} \wedge {}^* \bar{\Omega}).$	$\bar{L}_{\mathrm{w.}}(f) = \frac{i}{8} \text{Tr}(\bar{\gamma} \wedge \bar{\gamma} \wedge {}^* \bar{\Omega}).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_377 EQ1145 (C.13)		■ 0.667	$\stackrel{\circ}{g}_{lm}:=c_{lk}{}^i c_{mi}{}^k$	$\stackrel{\circ}{g}_{lm}:=c_{lk}{}^i c_{mi}{}^k$	$\stackrel{\circ}{g}_{lm}:=c_{lk}{}^i c_{mi}{}^k$
mielke-geometrodynamics_151 EQ0450 (7.2.3)		■ 0.676	$\Omega^g=\stackrel{+}{\Omega}^g+\bar{\Omega}^g$	$\Omega^g=\stackrel{+}{\Omega}^g+\bar{\Omega}^g$	$\Omega^g=\stackrel{+}{\Omega}^g+\bar{\Omega}^g$
mielke-geometrodynamics_64 EQ0143 (3.4.14)		■ 0.679	$\operatorname{div} \iota := (-1)^{1+s} d^* \iota = 0$	$\operatorname{div} \iota := (-1)^{1+s} d^* \iota = 0$	$\operatorname{div} \iota := (-1)^{1+s} d^* \iota = 0$
mielke-geometrodynamics_155 EQ0472 (7.3.11)		■ 0.682	$\stackrel{+}{\Omega}{}^{\{\}}=0\Longleftarrowtail R_{ij}{}^{\{\}}=0\Longleftarrowtail R_{ij}{}^{\{\}}=\Lambda'g_{ij}.$	<i>(not rendered)</i>	$\stackrel{+}{\Omega}{}^{\{\}}=0\Longleftrightarrow \mathcal{R}_{ij}{}^{\{\}}=0\Longleftrightarrow R_{ij}{}^{\{\}}=\Lambda'g_{ij}.$
mielke-geometrodynamics_256 EQ0812 (11.4.24)		■ 0.686	$\begin{aligned} & \partial_{\rho^*} H + \frac{1+\kappa}{\rho} e^{(\nu-\mu)/2} H = \frac{1}{\beta} \left[\omega + e^{\nu/2} - \varepsilon e^{-\mu} \left(H^2 - F^2 - \frac{2}{3} a H^2 \right) \right] F, \\ & \partial_{\rho^*} F + \frac{1-\kappa}{\rho} e^{(\nu-\mu)/2} F = \frac{1}{\beta} \left[-\omega + e^{\nu/2} - \varepsilon e^{-\mu} \left(H^2 - F^2 + \frac{2}{3} a F^2 \right) \right] H. \end{aligned}$	$\begin{aligned} & \partial_{\rho^*} H + \frac{1+\kappa}{\rho} e^{(\nu-\mu)/2} H = \frac{1}{\beta} \left[\omega + e^{\nu/2} - \varepsilon e^{-\mu} \left(H^2 - F^2 - \frac{2}{3} a H^2 \right) \right] F, \\ & \partial_{\rho^*} F + \frac{1-\kappa}{\rho} e^{(\nu-\mu)/2} F = \frac{1}{\beta} \left[-\omega + e^{\nu/2} - \varepsilon e^{-\mu} \left(H^2 - F^2 + \frac{2}{3} a F^2 \right) \right] H. \end{aligned}$	$\begin{aligned} & \partial_{\rho^*} H + \frac{1+\kappa}{\rho} e^{(\nu-\mu)/2} H = \frac{1}{\beta} \left[\omega + e^{\nu/2} - \varepsilon e^{-\mu} \left(H^2 - F^2 - \frac{2}{3} a H^2 \right) \right] F, \\ & \partial_{\rho^*} F + \frac{1-\kappa}{\rho} e^{(\nu-\mu)/2} F = \frac{1}{\beta} \left[-\omega + e^{\nu/2} - \varepsilon e^{-\mu} \left(H^2 - F^2 + \frac{2}{3} a F^2 \right) \right] H. \end{aligned}$
mielke-geometrodynamics_111 EQ0321 (5.2.5)		■ 0.686	$\tau_{\alpha\beta}:=\frac{\partial L_{\mathrm{D}}}{\partial\Gamma^{\alpha\beta}}=\frac{1}{8}\bar{\Psi}\left({}^*\gamma\sigma_{\alpha\beta}+\sigma_{\alpha\beta}{}^*\gamma\right)\Psi=\frac{1}{4}\eta_{\alpha\beta\gamma\delta}\bar{\Psi}\gamma^{\delta}\gamma_5\Psi\eta^{\gamma}=\tau_{\alpha\beta\gamma}\eta^{\gamma}$	$\tau_{\alpha\beta}:=\frac{\partial L_{\mathrm{D}}}{\partial\Gamma^{\alpha\beta}}=\frac{1}{8}\bar{\Psi}\left({}^*\gamma\sigma_{\alpha\beta}+\sigma_{\alpha\beta}{}^*\gamma\right)\Psi=\frac{1}{4}\eta_{\alpha\beta\gamma\delta}\bar{\Psi}\gamma^{\delta}\gamma_5\Psi\eta^{\gamma}=\tau_{\alpha\beta\gamma}\eta^{\gamma}$	$\tau_{\alpha\beta}:=\frac{\partial L_{\mathrm{D}}}{\partial\Gamma^{\alpha\beta}}=\frac{1}{8}\bar{\Psi}\left({}^*\gamma\sigma_{\alpha\beta}+\sigma_{\alpha\beta}{}^*\gamma\right)\Psi=\frac{1}{4}\eta_{\alpha\beta\gamma\delta}\bar{\Psi}\gamma^{\delta}\gamma_5\Psi\eta^{\gamma}=\tau_{\alpha\beta\gamma}\eta^{\gamma}$
mielke-geometrodynamics_131 EQ0389 (6.4.4)		■ 0.688	$\mathscr{H}=\mathscr{K}-\dot{\Psi}\wedge\left(\frac{\partial\mathscr{K}}{\partial\dot{\Psi}}\right),$	$\mathscr{H}=\mathscr{K}-\dot{\Psi}\wedge\left(\frac{\partial\mathscr{K}}{\partial\dot{\Psi}}\right),$	$\mathscr{H}=\mathscr{K}-\dot{\Psi}\wedge\left(\frac{\partial\mathscr{K}}{\partial\dot{\Psi}}\right),$
mielke-geometrodynamics_307 EQ0949 (14.1.30)		■ 0.688	$D\Omega=(D\xi+(-1)^z*\xi\wedge D\delta)\cos\delta+(-1)^z(D^*\xi-(-1)^z\xi\wedge D\delta)\sin\delta=0.$	$D\Omega=(D\xi+(-1)^z*\xi\wedge D\delta)\cos\delta+(-1)^z(D^*\xi-(-1)^z\xi\wedge D\delta)\sin\delta=0.$	$D\Omega=(D\xi+(-1)^z*\xi\wedge D\delta)\cos\delta+(-1)^z(D^*\xi-(-1)^z\xi\wedge D\delta)\sin\delta=0.$
mielke-geometrodynamics_224 EQ0696 (10.5.7)		■ 0.694	$D^*\gamma={}^*\gamma\wedge T,\quad T:=e_{\alpha}{}_{\lceil}T^{\alpha},$	$D^*\gamma={}^*\gamma\wedge T,\quad T:=e_{\alpha}{}_{\lceil}T^{\alpha},$	$D^*\gamma={}^*\gamma\wedge T,\quad T:=e_{\alpha}{}_{\lceil}T^{\alpha},$
mielke-geometrodynamics_175 EQ0534 (8.2.1)		■ 0.699	$s\Gamma=\Psi-DC,$ $sC=\Phi-\frac{1}{2}[C,C],$ $s\Omega^g=-D\Psi-[C,\Omega^g],$ $s\Psi=-D\Phi-[C,\Psi],$ $s\Phi=-[C,\Phi].$	$s\Gamma=\Psi-DC,$ $sC=\Phi-\frac{1}{2}[C,C],$ $s\Omega^g=-D\Psi-[C,\Omega^g],$ $s\Psi=-D\Phi-[C,\Psi],$ $s\Phi=-[C,\Phi].$	$s\Gamma=\Psi-DC,$ $sC=\Phi-\frac{1}{2}[C,C],$ $s\Omega^g=-D\Psi-[C,\Omega^g],$ $s\Psi=-D\Phi-[C,\Psi],$ $s\Phi=-[C,\Phi].$
mielke-geometrodynamics_112 EQ0329 (5.3.6)		■ 0.699	$D\Theta\equiv[\Omega^g,\gamma],\quad D\Omega^g\equiv 0,$	$D\Theta\equiv[\Omega^g,\gamma],\quad D\Omega^g\equiv 0,$	$D\Theta\equiv[\Omega^g,\gamma],\quad D\Omega^g\equiv 0,$
mielke-geometrodynamics_155 EQ0471 (7.3.10)		■ 0.706	$\bar{\Omega}{}^{\{\}}=0\Longleftarrowtail\Omega_{\mathrm{C}}{}^{\{\}}=0\text{ and }R^{\{\}}=0.$	$\bar{\Omega}{}^{\{\}}=0\Longleftrightarrow\Omega_{\mathrm{C}}{}^{\{\}}=0\text{ and }R^{\{\}}=0.$	$\overline{\Omega}{}^{\{\}}=0\Longleftrightarrow\Omega_{\mathrm{C}}{}^{\{\}}=0\text{ and }R^{\{\}}=0.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_140EQ0747 (11.1.14)	240	0.706	$M=\mathbb{R} \times S^3 \times W^3$.	$M = \mathbb{R} \times S^3 \times W^3$.	$M = \mathbb{R} \times S^3 \times W^3$.
mielke-geometrodynamics_140EQ0314 (4.6.6)	105	0.710	$\Phi(\stackrel{\sim}{\Gamma},\gamma)=P\exp\left[(i/\hbar)\oint\left(\Gamma^{(T)\alpha}P_\alpha+\Gamma_\alpha^{(L)\beta}L^\alpha_\beta\right)\right]$	$\Phi(\stackrel{\sim}{\Gamma},\gamma)=P\exp\left[(i/\hbar)\oint\left(\Gamma^{(T)\alpha}P_\alpha+\Gamma_\alpha^{(L)\beta}L^\alpha_\beta\right)\right]$	$\Phi(\stackrel{\sim}{\Gamma},\gamma)=P\exp\left[(i/\hbar)\oint\left(\Gamma^{(T)\alpha}P_\alpha+\Gamma_\alpha^{(L)\beta}L^\alpha_\beta\right)\right]$
mielke-geometrodynamics_140EQ0108 (3.3.9)	58	0.711	$L_{\mathrm{Pontr}}=-\frac{1}{2}F\wedge F=-\frac{1}{2}dC,$	$L_{\mathrm{Pontr}}=-\frac{1}{2}F\wedge F=-\frac{1}{2}dC,$	$L_{\mathrm{Pontr}}=-\frac{1}{2}F\wedge F=-\frac{1}{2}dC,$
mielke-geometrodynamics_140EQ0545 (8.3.6)	177	0.711	$d\rightarrow\exp(\zeta)d\exp(-\zeta).$	$d\rightarrow\exp(\zeta)d\exp(-\zeta).$	$d\rightarrow\exp(\zeta)d\exp(-\zeta).$
mielke-geometrodynamics_140EQ0583 (9.1.1)	192	0.712	$ds^2=-e^vdt^2+e^\lambda dr^2+r^2d\Omega^{n-2}$	$ds^2=-e^vdt^2+e^\lambda dr^2+r^2d\Omega^{n-2}$	$ds^2=-e^vdt^2+e^\lambda dr^2+r^2d\Omega^{n-2}$
mielke-geometrodynamics_140EQ0260 (4.3.25)	94	0.718	$\stackrel{*}{K}=\frac{1}{2}K_{\alpha}^{\beta c}L_{\beta c}\otimes\vartheta^\alpha.$	$\stackrel{*}{K}=\frac{1}{2}K_{\alpha}^{\beta c}L_{\beta c}\otimes\vartheta^\alpha.$	$\stackrel{*}{K}=\frac{1}{2}K_{\alpha}^{\beta c}L_{\beta c}\otimes\vartheta^\alpha.$
mielke-geometrodynamics_140EQ0878 (13.3.7)	285	0.722	$\mathrm{SL}(n+1,\mathbb{R})\approx\mathrm{A}_*(n,\mathbb{R})\otimes\mathrm{GL}(n,\mathbb{R})\otimes\mathbb{R}_*^n.$	$\mathrm{SL}(n+1,\mathbb{R})\approx\mathrm{A}_*(n,\mathbb{R})\otimes\mathrm{GL}(n,\mathbb{R})\otimes\mathbb{R}_*^n.$	$\mathrm{SL}(n+1,\mathbb{R})\approx\mathrm{A}_*(n,\mathbb{R})\otimes\mathrm{GL}(n,\mathbb{R})\otimes\mathbb{R}_*^n.$
mielke-geometrodynamics_140EQ0041 (2.6.4)	42	0.725	$\stackrel{*}{\Omega}=\frac{1}{2}F_{\alpha\beta}L_j\otimes\vartheta^\alpha\wedge\vartheta^\beta$	$\stackrel{*}{\Omega}=\frac{1}{2}F_{\alpha\beta}L_j\otimes\vartheta^\alpha\wedge\vartheta^\beta$	$\stackrel{*}{\Omega}=\frac{1}{2}F_{\alpha\beta}L_j\otimes\vartheta^\alpha\wedge\vartheta^\beta$
mielke-geometrodynamics_140EQ0574 (8.6.12)	185	0.725	$L_{\mathrm{eff}}:=L_{\mathrm{qPG}}(**)-dC_\theta=-\frac{\theta_{\mathrm{T}}^*}{2\ell^2}R^{\{\alpha\beta\}}\wedge\eta_{\alpha\beta}+\frac{\theta_{\mathrm{T}}^*\Lambda_{\mathrm{eff}}}{\ell^2}\eta.$	$L_{\mathrm{eff}}:=L_{\mathrm{qPG}}(**)-dC_\theta=-\frac{\theta_{\mathrm{T}}^*}{2\ell^2}R^{\{\alpha\beta\}}\wedge\eta_{\alpha\beta}+\frac{\theta_{\mathrm{T}}^*\Lambda_{\mathrm{eff}}}{\ell^2}\eta.$	$L_{\mathrm{eff}}:=L_{\mathrm{qPG}}(**)-dC_\theta=-\frac{\theta_{\mathrm{T}}^*}{2\ell^2}R^{\{\alpha\beta\}}\wedge\eta_{\alpha\beta}+\frac{\theta_{\mathrm{T}}^*\Lambda_{\mathrm{eff}}}{\ell^2}\eta.$
mielke-geometrodynamics_140EQ0366 (6.2.13)	126	0.725	$\stackrel{(\pm)}{L}_{\mathrm{EC}}:=L_{\mathrm{EC}}\pm idC_{\mathrm{TT}}=-\frac{1}{2\ell^2}\stackrel{(\pm)}{R}_{\alpha\beta}\wedge*(\vartheta^\alpha\wedge\vartheta^\beta)+\frac{\Lambda}{\ell^2}\eta.$	$\stackrel{(\pm)}{L}_{\mathrm{EC}}:=L_{\mathrm{EC}}\pm idC_{\mathrm{TT}}=-\frac{1}{2\ell^2}\stackrel{(\pm)}{R}_{\alpha\beta}\wedge*(\vartheta^\alpha\wedge\vartheta^\beta)+\frac{\Lambda}{\ell^2}\eta.$	$\stackrel{(\pm)}{L}_{\mathrm{EC}}:=L_{\mathrm{EC}}\pm idC_{\mathrm{TT}}=-\frac{1}{2\ell^2}\stackrel{(\pm)}{R}_{\alpha\beta}\wedge*(\vartheta^\alpha\wedge\vartheta^\beta)+\frac{\Lambda}{\ell^2}\eta.$
mielke-geometrodynamics_140EQ0585 (9.1.3)	193	0.725	$ds^2=\frac{n-2}{2\Lambda_{\mathrm{eff}}t^2}(-dt^2+dr^2)+\frac{(n-2)(n-3)}{2\Lambda_{\mathrm{eff}}}d\Omega^{n-2}$	$ds^2=\frac{n-2}{2\Lambda_{\mathrm{eff}}t^2}(-dt^2+dr^2)+\frac{(n-2)(n-3)}{2\Lambda_{\mathrm{eff}}}d\Omega^{n-2}$	$ds^2=\frac{n-2}{2\Lambda_{\mathrm{eff}}t^2}(-dt^2+dr^2)+\frac{(n-2)(n-3)}{2\Lambda_{\mathrm{eff}}}d\Omega^{n-2}$
mielke-geometrodynamics_140EQ0021 (2.4.3)	38	0.726	$\phi^{(p)}:=\varphi^p\otimes b\in C^\infty(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^\rho),\quad b\in C^\infty(V^\rho).$	$\phi^{(p)}:=\varphi^p\otimes b\in C^\infty(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^\rho),\quad b\in C^\infty(V^\rho).$	$\phi^{(p)}:=\varphi^p\otimes b\in C^\infty(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^\rho),\quad b\in C^\infty(V^\rho).$
mielke-geometrodynamics_140EQ0209 (4.2.1)	85	0.728	$\stackrel{*}{\chi}\stackrel{\sim}{\omega}=\omega^{\mathrm{L}}\otimes\omega^{\mathrm{T}},$	$\stackrel{*}{\chi}\stackrel{\sim}{\omega}=\omega^{\mathrm{L}}\otimes\omega^{\mathrm{T}},$	$\stackrel{*}{\chi}\stackrel{\sim}{\omega}=\omega^{\mathrm{L}}\otimes\omega^{\mathrm{T}},$
mielke-geometrodynamics_140EQ0309 (4.6.1)	104	0.728	$\stackrel{\sim}{\Gamma}:=\Gamma^{(T)\alpha}P_\alpha+\Gamma_\alpha^{(L)\beta}L^\alpha_\beta$	$\stackrel{\sim}{\Gamma}:=\Gamma^{(T)\alpha}P_\alpha+\Gamma_\alpha^{(L)\beta}L^\alpha_\beta$	$\stackrel{\sim}{\Gamma}:=\Gamma^{(T)\alpha}P_\alpha+\Gamma_\alpha^{(L)\beta}L^\alpha_\beta$
mielke-geometrodynamics_140EQ0009 (2.2.4)	34	0.729	$E_i^\alpha e^i_\beta=\delta^\alpha_\beta.$	$E_i^\alpha e^i_\beta=\delta^\alpha_\beta.$	$E_i^\alpha e^i_\beta=\delta^\alpha_\beta.$
mielke-geometrodynamics_140EQ0384 (6.3.2)	130	0.730	$H_\alpha^\parallel=\frac{1}{\ell^2}\left[\vartheta^\beta\wedge*(d\vartheta_\beta\wedge\vartheta_\alpha)-\frac{1}{2}\vartheta_\alpha\wedge*(d\vartheta^\beta\wedge\vartheta_\beta)\right]=\frac{1}{2\ell^2}\Gamma^{\{\beta\gamma\}}\wedge\eta_{\alpha\beta\gamma}$	$H_\alpha^\parallel=\frac{1}{\ell^2}\left[\vartheta^\beta\wedge*(d\vartheta_\beta\wedge\vartheta_\alpha)-\frac{1}{2}\vartheta_\alpha\wedge*(d\vartheta^\beta\wedge\vartheta_\beta)\right]=\frac{1}{2\ell^2}\Gamma^{\{\beta\gamma\}}\wedge\eta_{\alpha\beta\gamma}$	$H_\alpha^\parallel=\frac{1}{\ell^2}\left[\vartheta^\beta\wedge*(d\vartheta_\beta\wedge\vartheta_\alpha)-\frac{1}{2}\vartheta_\alpha\wedge*(d\vartheta^\beta\wedge\vartheta_\beta)\right]=\frac{1}{2\ell^2}\Gamma^{\{\beta\gamma\}}\wedge\eta_{\alpha\beta\gamma}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E08686 (9.2.5)	199	■ 0.753	$\chi(M)=\int_M\left(\frac{1}{4\pi}\left \frac{\partial}{\partial t}\right ^2+\frac{1}{3}\left \frac{\partial}{\partial t}\right ^2\right)\mathrm{d}V$	$\chi(M)=\int_M\left(\gamma_2^g-\frac{1}{(4\pi)^2}\int_{\partial M}\left\{\alpha\wedge\Omega^{g(*)}+\frac{2}{3}\alpha\wedge(\alpha\wedge\alpha)^{(*)}\right\}\right)$	$\chi(M)=\int_M\left(\pi(\gamma_2^g)-\frac{1}{(4\pi)^2}\int_{\partial M}\left\{\alpha\wedge\Omega^{g(*)}+\frac{2}{3}\alpha\wedge(\alpha\wedge\alpha)^{(*)}\right\}\right)$
mielke-geometrodynamics_E0992 (14.3.5)	320	■ 0.754	$\stackrel{\circ}{\chi}^{\pm}_{\rho F}(\xi)=\pm\chi^{\pm}_{\rho F}(\xi)$	$\stackrel{\circ}{\gamma}^{K+1}\chi^{\pm}_{\rho F}(\xi)=\pm\chi^{\pm}_{\rho F}(\xi)$	$\stackrel{\circ}{\gamma}^{K+1}\chi^{\pm}_{\rho F}(\xi)=\pm\chi^{\pm}_{\rho F}(\xi)$
mielke-geometrodynamics_E08249 (4.3.14)	92	■ 0.755	$D_k g_{ij}=E_k^\alpha \nabla_\alpha g_{ij}=0$	$D_k g_{ij}=E_k^\alpha \nabla_\alpha g_{ij}=0$	$D_k g_{ij}=E_k^\alpha \nabla_\alpha g_{ij}=0$
mielke-geometrodynamics_E08399 (6.4.17)	133	■ 0.755	$\begin{aligned} & \left[D_\alpha, D_\beta\right] = -T_{\alpha\beta}{}^\gamma(x) D_\gamma + R_{\alpha\beta\gamma}{}^\delta(x) L^\gamma{}_\delta, \\ & \left[L^\alpha{}_\beta, D_\gamma\right] = \delta_\gamma^\alpha D_\beta, \\ & \left[L^\alpha{}_\beta, L^\gamma{}_\delta\right] = \delta_\delta^\alpha L^\gamma{}_\beta - \delta_\beta^\gamma L^\alpha{}_\delta, \end{aligned}$	$\begin{aligned} & [D_\alpha, D_\beta] = -T_{\alpha\beta}{}^\gamma(x) D_\gamma + R_{\alpha\beta\gamma}{}^\delta(x) L^\gamma{}_\delta, \\ & [L^\alpha{}_\beta, D_\gamma] = \delta_\gamma^\alpha D_\beta, \\ & [L^\alpha{}_\beta, L^\gamma{}_\delta] = \delta_\delta^\alpha L^\gamma{}_\beta - \delta_\beta^\gamma L^\alpha{}_\delta, \end{aligned}$	$\begin{aligned} & [D_\alpha, D_\beta] = -T_{\alpha\beta}{}^\gamma(x) D_\gamma + R_{\alpha\beta\gamma}{}^\delta(x) L^\gamma{}_\delta, \\ & [L^\alpha{}_\beta, D_\gamma] = \delta_\gamma^\alpha D_\beta, \\ & [L^\alpha{}_\beta, L^\gamma{}_\delta] = \delta_\delta^\alpha L^\gamma{}_\beta - \delta_\beta^\gamma L^\alpha{}_\delta, \end{aligned}$
mielke-geometrodynamics_E09957 (14.1.39)	309	■ 0.756	$\partial_\kappa\delta=\sqrt{ g }\varepsilon_{\kappa\lambda\rho\sigma}R^{\{\sigma}{}_{\cdot\nu}{}^{\rho}R^{\}\nu\lambda}/\left(R^{\{\sigma}{}_{\cdot\nu}{}^{\rho}R^{\}\nu\lambda}\right)$	$\partial_\kappa\delta=\sqrt{ g }\varepsilon_{\kappa\lambda\rho\sigma}R^{\{\sigma}{}_{\cdot\nu}{}^{\rho}R^{\}\nu\lambda}/\left(R^{\{\sigma}{}_{\cdot\nu}{}^{\rho}R^{\}\nu\lambda}\right)$	$\partial_\kappa\delta=\sqrt{ g }\varepsilon_{\kappa\lambda\rho\sigma}R^{\{\sigma}{}_{\cdot\nu}{}^{\rho}R^{\}\nu\lambda}/\left(R^{\{\sigma}{}_{\cdot\nu}{}^{\rho}R^{\}\nu\lambda}\right)$
mielke-geometrodynamics_E09978 (14.2.11)	314	■ 0.758	$\hat{\Omega}=\hat{d}\hat{\omega}-\hat{\omega}\wedge\hat{\omega}=\left[\begin{array}{c c}\Omega^g+\alpha\wedge\alpha^T&-D\alpha\\\hline\stackrel{\circ}{\Omega}+\alpha^T\wedge\alpha&\end{array}\right]$	$\hat{\Omega}=\hat{d}\hat{\omega}-\hat{\omega}\wedge\hat{\omega}=\left[\begin{array}{c c}\Omega^g+\alpha\wedge\alpha^T&-D\alpha\\\hline\stackrel{\circ}{\Omega}+\alpha^T\wedge\alpha&\end{array}\right]$	$\hat{\Omega}=\hat{d}\hat{\omega}-\hat{\omega}\wedge\hat{\omega}=\left[\begin{array}{c c}\Omega^g+\alpha\wedge\alpha^T&-D\alpha\\\hline\stackrel{\circ}{\Omega}+\alpha^T\wedge\alpha&\end{array}\right]$
mielke-geometrodynamics_E01816 (14.4.11)	327	■ 0.758	$ds^2=-\left(1-\left(\frac{r_\circ}{r}\right)^4\right)\frac{\Delta}{r^2}dt^2+\frac{r^2}{\Delta}dr^2+r^2d\Omega^2$	$ds^2=-\left(1-\left(\frac{r_\circ}{r}\right)^4\right)\frac{\Delta}{r^2}dt^2+\frac{r^2}{\Delta}dr^2+r^2d\Omega^2$	$ds^2=-\left(1-\left(\frac{r_\circ}{r}\right)^4\right)\frac{\Delta}{r^2}dt^2+\frac{r^2}{\Delta}dr^2+r^2d\Omega^2$
mielke-geometrodynamics_E08318 (4.6.2)	104	■ 0.760	$\widetilde{D}\widetilde{\xi}^\alpha=d\xi^\alpha+\Gamma_\beta^{(L)\alpha}\wedge\xi^\beta+\Gamma^{(T)\alpha}$	$\widetilde{D}\widetilde{\xi}^\alpha=d\xi^\alpha+\Gamma_\beta^{(L)\alpha}\wedge\xi^\beta+\Gamma^{(T)\alpha}$	$\widetilde{D}\widetilde{\xi}^\alpha=d\xi^\alpha+\Gamma_\beta^{(L)\alpha}\wedge\xi^\beta+\Gamma^{(T)\alpha}$
mielke-geometrodynamics_E08832 (12.1.4)	270	■ 0.763	$\langle d\widetilde{j}_5\rangle=0,$	$\langle d\widetilde{j}_5\rangle=0,$	$\langle d\widetilde{j}_5\rangle=0,$
mielke-geometrodynamics_E08231 (4.2.24)	88	■ 0.766	$T_{\alpha\beta}^{\cdot\cdot c}=2D_{[\alpha}E_{\cdot \beta]}^c=2\left(\partial_{[\alpha}E_{\cdot \beta]}^c+\Gamma_{[\alpha \cdot}^cE_{\cdot \beta]}^d\right)$	$T_{\alpha\beta}^{\cdot\cdot c}=2D_{[\alpha}E_{\cdot \beta]}^c=2\left(\partial_{[\alpha}E_{\cdot \beta]}^c+\Gamma_{[\alpha \cdot}^cE_{\cdot \beta]}^d\right)$	$T_{\alpha\beta}^{\cdot\cdot c}=2D_{[\alpha}E_{\cdot \beta]}^c=2\left(\partial_{[\alpha}E_{\cdot \beta]}^c+\Gamma_{[\alpha \cdot}^cE_{\cdot \beta]}^d\right)$
mielke-geometrodynamics_E01879 (A.4)	365	■ 0.766	$[\sigma^i,\sigma^j]=\varepsilon^{ij}{}_k\sigma^k$	$[\sigma^i,\sigma^j]=\varepsilon^{ij}{}_k\sigma^k$	$[\sigma^i,\sigma^j]=\varepsilon^{ij}{}_k\sigma^k$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_97 EQ0276 (4.4.4)		■ 0.767	$\begin{aligned} & \delta L_g = \delta \vartheta \wedge E - \delta \vartheta \wedge D \Pi^T + d(\delta \vartheta \wedge \Pi^T) \\ & - [\delta \omega^g, \vartheta] \wedge [\Pi_{[]}^T, \vartheta] - \delta \omega^g \wedge D \Pi^L + d(\delta \omega^g \wedge \Pi^L), \end{aligned}$	$\delta L_g = \delta \vartheta \wedge E - \delta \vartheta \wedge D \Pi^T + d(\delta \vartheta \wedge \Pi^T) - [\delta \omega^g, \vartheta] \wedge [\Pi_{[]}^T, \vartheta] - \delta \omega^g \wedge D \Pi^L + d(\delta \omega^g \wedge \Pi^L),$	$\delta L_g = \delta \vartheta \wedge E - \delta \vartheta \wedge D \Pi^T + d(\delta \vartheta \wedge \Pi^T) - [\delta \omega^g, \vartheta] \wedge [\Pi_{[]}^T, \vartheta] - \delta \omega^g \wedge D \Pi^L + d(\delta \omega^g \wedge \Pi^L),$
mielke-geometrodynamics_209 EQ0613 (10.1.1)		■ 0.769	$\vartheta^\alpha = E_i^\alpha dx^i \quad \text{and} \quad \Gamma_\alpha^\star := \frac{1}{2} \eta_{\alpha\beta\gamma} \Gamma^{\beta\gamma}.$	$\vartheta^\alpha = E_i^\alpha dx^i \quad \text{and} \quad \Gamma_\alpha^\star := \frac{1}{2} \eta_{\alpha\beta\gamma} \Gamma^{\beta\gamma}.$	$\vartheta^\alpha = E_i^\alpha dx^i \quad \text{and} \quad \Gamma_\alpha^\star := \frac{1}{2} \eta_{\alpha\beta\gamma} \Gamma^{\beta\gamma}.$
mielke-geometrodynamics_355 EQ1066 (16.2.5)		■ 0.770	$\rho^* = \int_o^\rho \frac{dx}{1 \pm x^2} = \begin{cases} \operatorname{artan} \rho, \kappa > 0 \\ \operatorname{Arth} \rho, \kappa < 0 \end{cases}$	$\rho^* = \int_o^\rho \frac{dx}{1 \pm x^2} = \begin{cases} \operatorname{artan} \rho, \kappa > 0 \\ \operatorname{Arth} \rho, \kappa < 0 \end{cases}$	$\rho^* = \int_o^\rho \frac{dx}{1 \pm x^2} = \begin{cases} \operatorname{artan} \rho, \kappa > 0 \\ \operatorname{Arth} \rho, \kappa < 0 \end{cases}.$
mielke-geometrodynamics_337 EQ1030 (15.1.6)		■ 0.771	$\begin{aligned} & \frac{1}{2} \vartheta^g \wedge \Omega^g + \Lambda^* \vartheta_g = \ell^{*2} \{ \Sigma_{\text{m}}(\text{leptons}) + \Sigma^g \}, \\ & \frac{1}{2} \vartheta^f \wedge \Omega^f = \ell^2 \{ \Sigma_{\text{m}}(\text{hadrons}) + \Sigma^f \}. \end{aligned}$	$\frac{1}{2} \vartheta^g \wedge \Omega^g + \Lambda^* \vartheta_g = \ell^{*2} \{ \Sigma_{\text{m}}(\text{leptons}) + \Sigma^g \},$ $\frac{1}{2} \vartheta^f \wedge \Omega^f = \ell^2 \{ \Sigma_{\text{m}}(\text{hadrons}) + \Sigma^f \}.$	$\frac{1}{2} \vartheta^g \wedge \Omega^g + \Lambda^* \vartheta_g = \ell^{*2} \{ \Sigma_{\text{m}}(\text{leptons}) + \Sigma^g \},$ $\frac{1}{2} \vartheta^f \wedge \Omega^f = \ell^2 \{ \Sigma_{\text{m}}(\text{hadrons}) + \Sigma^f \}.$
mielke-geometrodynamics_59 EQ0112 (3.3.13)		■ 0.772	$\begin{aligned} j := & \mathscr{J} \wedge dt + \hat{\rho} \\ = & -(J_x dy \wedge dz + J_y dz \wedge dx + J_z dx \wedge dy) \wedge dt \\ & + \rho dx \wedge dy \wedge dz. \end{aligned}$	$j := -\mathscr{J} \wedge dt + \hat{\rho} = -(J_x dy \wedge dz + J_y dz \wedge dx + J_z dx \wedge dy) \wedge dt + \rho dx \wedge dy \wedge dz$	$j := -\mathscr{J} \wedge dt + \hat{\rho} = -(J_x dy \wedge dz + J_y dz \wedge dx + J_z dx \wedge dy) \wedge dt + \rho dx \wedge dy \wedge dz.$
mielke-geometrodynamics_62 EQ0130 (3.4.1)		■ 0.780	$\mathscr{F} := \frac{1}{\alpha_g} \operatorname{Tr} \left\{ \Omega \wedge \Omega \right\} = \frac{1}{4\alpha_g} F_{\alpha\beta}^j F^{\alpha\beta}_j$	$\mathscr{F} := \frac{1}{\alpha_g} \operatorname{Tr} \{ \Omega \wedge \Omega \} = \frac{1}{4\alpha_g} F_{\alpha\beta}^j F^{\alpha\beta}_j$	$\mathscr{F} := \frac{1}{\alpha_g} \operatorname{Tr} \{ \Omega \wedge \Omega \} = \frac{1}{4\alpha_g} F_{\alpha\beta}^j F^{\alpha\beta}_j$
mielke-geometrodynamics_155 EQ0470 (7.3.9)		■ 0.782	$\Omega^{g(*)} = \zeta \Omega^g,$	$*\Omega^{g(*)} = \zeta \Omega^g,$	$*\Omega^{g(*)} = \zeta \Omega^g,$
mielke-geometrodynamics_43 EQ0046 (2.6.9)		■ 0.783	$\Delta b = \int_{\partial U} \omega \otimes b = \int_U d(\omega \otimes b)_{\text{hor}} = \int_U D(\omega \otimes b)$	$\Delta b = \int_{\partial U} \omega \otimes b = \int_U d(\omega \otimes b)_{\text{hor}} = \int_U D(\omega \otimes b) = \int_U \Omega \otimes b \simeq \Omega \otimes b$	$\Delta b = \int_{\partial U} \omega \otimes b = \int_U d(\omega \otimes b)_{\text{hor}} = \int_U D(\omega \otimes b) = \int_U \Omega \otimes b \simeq \Omega \otimes b$
mielke-geometrodynamics_344 EQ1049 (15.2.6)		■ 0.785	$\bar{\omega} = \bar{\omega}_i^j \bar{L}_j \otimes dx^i = \bar{\omega}_i dx^i$	$*\bar{\omega} = \bar{\omega}_i^j \bar{L}_j \otimes dx^i = \bar{\omega}_i dx^i$	$*\bar{\omega} = \bar{\omega}_i^j \bar{L}_j \otimes dx^i = \bar{\omega}_i dx^i$
mielke-geometrodynamics_63 EQ0133 (3.4.4)		■ 0.785	$L_\omega = L(\mathscr{F}, \mathscr{F}) \eta.$	$L_\omega = L(\mathscr{F}, \mathscr{F}) \eta.$	$L_\omega = L(\mathscr{F}, \mathscr{F}) \eta.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00312 (4.6.4)	104	■ 0.786	$\begin{aligned} \Delta \xi^\alpha &= - \oint_C y \rfloor (DD\xi^\alpha) = \oint_C y \rfloor R^{(T)\alpha} = \int_S R^{(T)\alpha} \\ &\simeq \frac{1}{2} (T_{ij}^\alpha - R_{ij\beta}^\alpha \xi^\beta) \int_S dy^i \wedge dy^j . \end{aligned}$	$\Delta \xi^\alpha = - \oint_C y \rfloor (DD\xi^\alpha) = \oint_C y \rfloor R^{(T)\alpha} = \int_S R^{(T)\alpha} \\ \simeq \frac{1}{2} (T_{ij}^\alpha - R_{ij\beta}^\alpha \xi^\beta) \int_S dy^i \wedge dy^j$	$\Delta \xi^\alpha = - \oint_C y \rfloor (DD\xi^\alpha) = \oint_C y \rfloor R^{(T)\alpha} = \int_S R^{(T)\alpha} \\ \simeq \frac{1}{2} (T_{ij}^\alpha - R_{ij\beta}^\alpha \xi^\beta) \int_S dy^i \wedge dy^j .$
mielke-geometrodynamics_E00224 (4.2.17)	87	■ 0.787	$\widetilde{\varphi} \rightarrow A^{-1} \widetilde{\varphi} = G^{-1} (\widetilde{\varphi} - T) .$	$\widetilde{\varphi} \rightarrow A^{-1} \widetilde{\varphi} = G^{-1} (\widetilde{\varphi} - T) .$	$\widetilde{\varphi} \rightarrow A^{-1} \widetilde{\varphi} = G^{-1} (\widetilde{\varphi} - T) .$
mielke-geometrodynamics_E00941 (14.1.22)	306	■ 0.788	$\tan 2\delta = (-1)^z \frac{\mathrm{Tr}^*(\Omega \wedge \Omega)}{\mathrm{Tr}^*(\Omega \wedge *\Omega)} .$	$\tan 2\delta = (-1)^z \frac{\mathrm{Tr}^*(\Omega \wedge \Omega)}{\mathrm{Tr}^*(\Omega \wedge *\Omega)} .$	$\tan 2\delta = (-1)^z \frac{\mathrm{Tr}^*(\Omega \wedge \Omega)}{\mathrm{Tr}^*(\Omega \wedge *\Omega)} .$
mielke-geometrodynamics_E00760 (11.1.27)	242	■ 0.791	$\left(\gamma^{n+1}\right)^2=\mp \mathbb{1} \quad .$	$\left(\gamma^{n+1}\right)^2=\mp \mathbb{1} .$	$(\gamma^{n+1})^2=\mp \mathbb{1} .$
mielke-geometrodynamics_E00725 (10.10.5)	230	■ 0.793	$\delta_{\mathrm{susy}} L \cong \bar{\sigma} \left(i \gamma^\alpha \Psi \wedge \frac{\delta L}{\delta \vartheta^\alpha} - 2 D \frac{\delta L}{\delta \bar{\Psi}} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} \right) + 2 d \left(\bar{\sigma} \wedge \frac{\delta L}{\delta \bar{\Psi}} \right) .$	$\delta_{\mathrm{susy}} L \cong \bar{\sigma} \left(i \gamma^\alpha \Psi \wedge \frac{\delta L}{\delta \vartheta^\alpha} - 2 D \frac{\delta L}{\delta \bar{\Psi}} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} \right) + 2 d \left(\bar{\sigma} \wedge \frac{\delta L}{\delta \bar{\Psi}} \right) .$	$\delta_{\mathrm{susy}} L \cong \bar{\sigma} \left(i \gamma^\alpha \Psi \wedge \frac{\delta L}{\delta \vartheta^\alpha} - 2 D \frac{\delta L}{\delta \bar{\Psi}} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} \right) + 2 d \left(\bar{\sigma} \wedge \frac{\delta L}{\delta \bar{\Psi}} \right) .$
mielke-geometrodynamics_E00810 (11.4.22)	256	■ 0.793	$\frac{3}{8} \ell^2 \bar{\psi} \psi = 4 \pi m e^{-\mu - \nu / 2} (H^2 - F^2) \left Y_{ \kappa - 1}^{ \kappa - 1} (\vartheta, \phi) \right ^2$	$\frac{3}{8} \ell^2 \bar{\psi} \psi = 4 \pi m e^{-\mu - \nu / 2} (H^2 - F^2) \left Y_{ \kappa - 1}^{ \kappa - 1} (\vartheta, \phi) \right ^2$	$\frac{3}{8} \ell^2 \bar{\psi} \psi = 4 \pi m e^{-\mu - \nu / 2} (H^2 - F^2) \left Y_{ \kappa - 1}^{ \kappa - 1} (\vartheta, \phi) \right ^2$
mielke-geometrodynamics_E01032 (15.1.9)	338	■ 0.793	$\frac{1}{2} * (\vartheta^g \wedge *\Omega^g) + \Lambda \vartheta^g = 0,$	$\frac{1}{2} * (\vartheta^g \wedge *\Omega^g) + \Lambda \vartheta^g = 0,$	$\frac{1}{2} * (\vartheta^g \wedge *\Omega^g) + \Lambda \vartheta^g = 0,$
mielke-geometrodynamics_E00305 (4.5.15)	103	■ 0.793	$\varepsilon_{\mathrm{RC}} := \xi^\alpha \Sigma_\alpha + \left(e_\beta \lrcorner \widehat{D} \xi^\gamma \right) \tau^\beta{}_\gamma ,$	$\varepsilon_{\mathrm{RC}} := \xi^\alpha \Sigma_\alpha + \left(e_\beta \lrcorner \widehat{D} \xi^\gamma \right) \tau^\beta{}_\gamma ,$	$\varepsilon_{\mathrm{RC}} := \xi^\alpha \Sigma_\alpha + (e_\beta \lrcorner \widehat{D} \xi^\gamma) \tau^\beta{}_\gamma ,$
mielke-geometrodynamics_E00848 (12.3.1)	274	■ 0.794	$\langle dj_5 \rangle = 2 i m \langle \bar{\psi} \gamma_5 \psi \rangle \eta - \frac{1}{4 \pi^2} \mathrm{Tr} (G \wedge G) - \frac{1}{96 \pi^2} \left[2 R_{\alpha \beta}^{(\downarrow)} \wedge R^{(\uparrow) \alpha \beta} + d \mathscr{A} \wedge d \mathscr{A} \right]$	$\langle dj_5 \rangle = 2 i m \langle \bar{\psi} \gamma_5 \psi \rangle \eta - \frac{1}{4 \pi^2} \mathrm{Tr} (G \wedge G) - \frac{1}{96 \pi^2} \left[2 R_{\alpha \beta}^{(\downarrow)} \wedge R^{(\uparrow) \alpha \beta} + d \mathscr{A} \wedge d \mathscr{A} \right]$	$\langle dj_5 \rangle = 2 i m \langle \bar{\psi} \gamma_5 \psi \rangle \eta - \frac{1}{4 \pi^2} \mathrm{Tr} (G \wedge G) - \frac{1}{96 \pi^2} \left[2 R_{\alpha \beta}^{(\downarrow)} \wedge R^{(\uparrow) \alpha \beta} + d \mathscr{A} \wedge d \mathscr{A} \right]$
mielke-geometrodynamics_E00319 (5.2.3)	110	■ 0.795	$\begin{aligned} L_{\mathrm{D}} &= L(\gamma, \psi, D^{\downarrow} \psi) - \frac{i}{2} \bar{\psi} (*\gamma \wedge K - K \wedge *\gamma) \psi \\ &= L(\gamma, \psi, D^{\downarrow} \psi) + \frac{1}{4} \mathscr{A} \wedge j_5 . \end{aligned}$	$\begin{aligned} L_{\mathrm{D}} &= L(\gamma, \psi, D^{\downarrow} \psi) - \frac{i}{2} \bar{\psi} (*\gamma \wedge K - K \wedge *\gamma) \psi \\ &= L(\gamma, \psi, D^{\downarrow} \psi) + \frac{1}{4} \mathscr{A} \wedge j_5 . \end{aligned}$	$\begin{aligned} L_{\mathrm{D}} &= L(\gamma, \psi, D^{\downarrow} \psi) - \frac{i}{2} \bar{\psi} (*\gamma \wedge K - K \wedge *\gamma) \psi \\ &= L(\gamma, \psi, D^{\downarrow} \psi) + \frac{1}{4} \mathscr{A} \wedge j_5 . \end{aligned}$
mielke-geometrodynamics_E00552 (8.4.6)	179	■ 0.797	$\stackrel{+}{\Omega} := \frac{1}{2} \left(\Omega^g \pm *\Omega^{g(*)} \right) = 0$	$\stackrel{+}{\Omega} := \frac{1}{2} \left(\Omega^g \pm *\Omega^{g(*)} \right) = 0$	$\stackrel{\pm}{\Omega}{}^g := \frac{1}{2} \left(\Omega^g \pm *\Omega^{g(*)} \right) = 0$
mielke-geometrodynamics_E01001 (14.3.14)	323	■ 0.798	$(i \not{D})^2 := - \mathrm{Tr} \{ \gamma \wedge * D^* (\gamma \wedge * D) \} .$	$(i \not{D})^2 := - \mathrm{Tr} \{ \gamma \wedge * D^* (\gamma \wedge * D) \} .$	$(i \not{D})^2 := - \mathrm{Tr} \{ \gamma \wedge * D^* (\gamma \wedge * D) \} .$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00652 (2.8.8)	47	■ 0.803	$c_{\{2\}}(M)=-\int_M^*\pi\left(\gamma_2\right)=\frac{1}{8\pi^2}\int_M\mathrm{Tr}(\Omega\wedge\Omega)\wedge\Omega\,.$	$c_2(M)=-\int_M^*\pi\left(\gamma_2\right)=\frac{1}{8\pi^2}\int_M\mathrm{Tr}(\Omega\wedge\Omega).$	$c_2(M)=-\int_M^*\pi(\gamma_2)=\frac{1}{8\pi^2}\int_M\mathrm{Tr}(\Omega\wedge\Omega).$
mielke-geometrodynamics_E00652 (10.3.9)	216	■ 0.803	$H_{\{\alpha\}}=-\frac{\gamma\varepsilon}{\ell^2(1+\delta\varepsilon)}\vartheta_\alpha\quad\Leftrightarrow\quad T_\alpha=\frac{\varepsilon}{\ell}\eta_\alpha,$	$H_\alpha=-\frac{\gamma\varepsilon}{\ell^2(1+\delta\varepsilon)}\vartheta_\alpha\quad\Leftrightarrow\quad T_\alpha=\frac{\varepsilon}{\ell}\eta_\alpha,$	$H_\alpha=-\frac{\gamma\varepsilon}{\ell^2(1+\delta\varepsilon)}\vartheta_\alpha\quad\Leftrightarrow\quad T_\alpha=\frac{\varepsilon}{\ell}\eta_\alpha,$
mielke-geometrodynamics_E01065 (16.2.4)	355	■ 0.803	$\rho:=\sqrt{\frac{1}{3}\Lambda_{\mathrm{eff}}}\,r,\quad r= \mathbf{x} $	$\rho:=\sqrt{\frac{1}{3}\Lambda_{\mathrm{eff}}}\,r,\quad r= \mathbf{x} $	$\rho:=\sqrt{\frac{1}{3}\Lambda_{\mathrm{eff}}}\,r,\quad r= \mathbf{x} $
mielke-geometrodynamics_E00166 (3.6.1)	69	■ 0.806	$\begin{aligned} L_{\mathrm{SW}}=&\frac{1}{2}\overline{D_{\pm}\psi}\wedge^*D_{\pm}\psi\mp i\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)^2\\ &=\mp i\mathrm{Tr}\left(F^{\pm}\wedge F^{\pm}\right)+\frac{1}{2}\overline{D_{\pm}\psi}\wedge^*D_{\pm}\psi\\ &\quad\pm i\bar{\psi}\sigma_{\pm}\psi\wedge F^{\pm}\mp\frac{i}{4}\bar{\psi}\sigma_{\pm}\psi\wedge\bar{\psi}\sigma_{\pm}\psi; \end{aligned}$	$\begin{aligned} L_{\mathrm{SW}}=&\frac{1}{2}\overline{D_{\pm}\psi}\wedge^*D_{\pm}\psi\mp i\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)^2\\ &=\mp i\mathrm{Tr}\left(F^{\pm}\wedge F^{\pm}\right)+\frac{1}{2}\overline{D_{\pm}\psi}\wedge^*D_{\pm}\psi\\ &\quad\pm i\bar{\psi}\sigma_{\pm}\psi\wedge F^{\pm}\mp\frac{i}{4}\bar{\psi}\sigma_{\pm}\psi\wedge\bar{\psi}\sigma_{\pm}\psi; \end{aligned}$	$\begin{aligned} L_{\mathrm{SW}}=&\frac{1}{2}\overline{D_{\pm}\psi}\wedge^*D_{\pm}\psi\mp i\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)^2\\ &=\mp i\mathrm{Tr}\left(F^{\pm}\wedge F^{\pm}\right)+\frac{1}{2}\overline{D_{\pm}\psi}\wedge^*D_{\pm}\psi\\ &\quad\pm i\bar{\psi}\sigma_{\pm}\psi\wedge F^{\pm}\mp\frac{i}{4}\bar{\psi}\sigma_{\pm}\psi\wedge\bar{\psi}\sigma_{\pm}\psi; \end{aligned}$
mielke-geometrodynamics_E01070 (16.2.9)	356	■ 0.806	$V_{\mathrm{eff}}^o(\rho^*)=\left(\frac{1}{\beta^2}-2\right)\left[1+\tan^2\rho^*\right]$	$V_{\mathrm{eff}}^o(\rho^*)=\left(\frac{1}{\beta^2}-2\right)\left[1+\tan^2\rho^*\right]$	$V_{\mathrm{eff}}^o(\rho^*)=\left(\frac{1}{\beta^2}-2\right)\left[1+\tan^2\rho^*\right]$
mielke-geometrodynamics_E00446 (6.9.2)	144	■ 0.806	$\begin{aligned} \stackrel{\theta}{L}&=\frac{1}{2\ell}\left[T^\alpha\wedge^*T_\alpha+\theta_{\mathrm{T}}\left(T^\alpha\wedge T_\alpha+R_{\alpha\beta}\wedge\vartheta^\alpha\wedge\vartheta^\beta\right)\right.\\ &\quad\left.+\frac{1}{2}\left[R^{\alpha\beta}\wedge^*R_{\alpha\beta}+\theta_{\mathrm{L}}R^{\alpha\beta}\wedge R_{\alpha\beta}\right]. \end{aligned}$	$\begin{aligned} \stackrel{\theta}{L}&=\frac{1}{2\ell}\left[T^\alpha\wedge^*T_\alpha+\theta_{\mathrm{T}}\left(T^\alpha\wedge T_\alpha+R_{\alpha\beta}\wedge\vartheta^\alpha\wedge\vartheta^\beta\right)\right.\\ &\quad\left.+\frac{1}{2}\left[R^{\alpha\beta}\wedge^*R_{\alpha\beta}+\theta_{\mathrm{L}}R^{\alpha\beta}\wedge R_{\alpha\beta}\right]. \end{aligned}$	$\begin{aligned} \stackrel{\theta}{L}&=\frac{1}{2\ell}\left[T^\alpha\wedge^*T_\alpha+\theta_{\mathrm{T}}\left(T^\alpha\wedge T_\alpha+R_{\alpha\beta}\wedge\vartheta^\alpha\wedge\vartheta^\beta\right)\right.\\ &\quad\left.+\frac{1}{2}\left[R^{\alpha\beta}\wedge^*R_{\alpha\beta}+\theta_{\mathrm{L}}R^{\alpha\beta}\wedge R_{\alpha\beta}\right]. \end{aligned}$
mielke-geometrodynamics_E00610 (9.3.1)	202	■ 0.806	$\begin{aligned} dC_{\mathrm{RR}^{(\star)}}&:=\frac{1}{2}d\left(\Gamma_{\alpha\beta}\wedge R^{\alpha\beta(\star)}-\frac{1}{3}\Gamma_\alpha{}^{\beta(\star)}\wedge\Gamma_\beta{}^\gamma\wedge\Gamma_\gamma{}^\alpha\right)\\ &\equiv-L_{\mathrm{SKY}}-2\mathrm{Ric}_{\alpha\beta}\wedge^*\mathrm{Ric}^{\alpha\beta}+\frac{1}{2}\mathrm{Ric}_\alpha{}^\alpha\wedge^*\mathrm{Ric}_\beta{}^\beta \end{aligned}$	$\begin{aligned} dC_{\mathrm{RR}^{(\star)}}&:=\frac{1}{2}d\left(\Gamma_{\alpha\beta}\wedge R^{\alpha\beta(\star)}-\frac{1}{3}\Gamma_\alpha{}^{\beta(\star)}\wedge\Gamma_\beta{}^\gamma\wedge\Gamma_\gamma{}^\alpha\right)\\ &\equiv-L_{\mathrm{SKY}}-2\mathrm{Ric}_{\alpha\beta}\wedge^*\mathrm{Ric}^{\alpha\beta}+\frac{1}{2}\mathrm{Ric}_\alpha{}^\alpha\wedge^*\mathrm{Ric}_\beta{}^\beta \end{aligned}$	$\begin{aligned} dC_{\mathrm{RR}^{(\star)}}&:=\frac{1}{2}d\left(\Gamma_{\alpha\beta}\wedge R^{\alpha\beta(\star)}-\frac{1}{3}\Gamma_\alpha{}^{\beta(\star)}\wedge\Gamma_\beta{}^\gamma\wedge\Gamma_\gamma{}^\alpha\right)\\ &\equiv-L_{\mathrm{SKY}}-2\mathrm{Ric}_{\alpha\beta}\wedge^*\mathrm{Ric}^{\alpha\beta}+\frac{1}{2}\mathrm{Ric}_\alpha{}^\alpha\wedge^*\mathrm{Ric}_\beta{}^\beta \end{aligned}$
mielke-geometrodynamics_E00767 (11.1.34)	244	■ 0.807	$\psi\longrightarrow S^{-1}\psi=S^{-1}\psi$	$\psi\longrightarrow S^{-1}\psi=S^{-1}\psi$	$\psi\longrightarrow S^{-1}\psi=S^{-1}\psi$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_95 EQ0267 (4.3.32)	95	■ 0.808	$\begin{aligned} & R_{(i\ j)\ \kappa l}^{\{\}} = R_{ij(\kappa l)}^{\{\}} = 0 \\ & R_{ij\kappa l}^{\}} = R_{\kappa lij}^{\}}. \end{aligned}$	$R_{(ij)\kappa l}^{\{\}} = R_{ij(\kappa l)}^{\{\}} = 0$ $R_{ij\kappa l}^{\}} = R_{\kappa lij}^{\}}.$	$R_{(ij)\kappa l}^{\{\}} = R_{ij(\kappa l)}^{\{\}} = 0$ $R_{ij\kappa l}^{\{\}} = R_{\kappa lij}^{\{\}}.$
mielke-geometrodynamics_101 EQ0298 (4.5.8)	101	■ 0.811	$\begin{aligned} & \delta L \wedge = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha} \\ & + \delta \Gamma_\alpha{}^\beta \wedge I^\alpha{}_\beta \Psi \wedge \frac{\partial L}{\partial D\Psi} + \delta \Psi \wedge \frac{\delta L}{\delta \Psi} + D \left(\delta \Psi \wedge \frac{\partial L}{\partial D\Psi} \right). \end{aligned}$	$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha}$ $+ \delta \Gamma_\alpha{}^\beta \wedge I^\alpha{}_\beta \Psi \wedge \frac{\partial L}{\partial D\Psi} + \delta \Psi \wedge \frac{\delta L}{\delta \Psi} + D \left(\delta \Psi \wedge \frac{\partial L}{\partial D\Psi} \right)$	$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta \vartheta^\alpha \wedge \frac{\partial L}{\partial \vartheta^\alpha}$ $+ \delta \Gamma_\alpha{}^\beta \wedge I^\alpha{}_\beta \Psi \wedge \frac{\partial L}{\partial D\Psi} + \delta \Psi \wedge \frac{\delta L}{\delta \Psi} + D \left(\delta \Psi \wedge \frac{\partial L}{\partial D\Psi} \right).$
mielke-geometrodynamics_295 EQ0919 (13.8.5)	295	■ 0.811	$D\ T^{\{\alpha\}} \equiv R_{\{\beta\}\{\}}^{\{\}} \wedge \vartheta^{\beta}$	$DT^\alpha \equiv R_\beta{}^\alpha \wedge \vartheta^\beta$	$DT^\alpha \equiv R_\beta{}^\alpha \wedge \vartheta^\beta$
mielke-geometrodynamics_254 EQ0802 (11.4.14)	254	■ 0.812	$\begin{aligned} & \left[i\gamma^0 \partial_0 - e^{(\nu-\mu)/2} \left[i\vec{\gamma} \cdot \vec{\partial} + i\gamma^1 \partial_r \left(\frac{\mu}{2} + \frac{\nu}{4} \right) \right] \right. \\ & \left. + \frac{3\varepsilon}{8} \ell^2 e^{\nu/2} (\bar{\psi} \psi - a \bar{\psi} \gamma^5 \psi \gamma^5) - e^{\nu/2} m \right] \psi = 0 \end{aligned}$	$\left\{ i\gamma^0 \partial_0 - e^{(\nu-\mu)/2} \left[i\vec{\gamma} \cdot \vec{\partial} + i\gamma^1 \partial_r \left(\frac{\mu}{2} + \frac{\nu}{4} \right) \right] \right.$ $\left. + \frac{3\varepsilon}{8} \ell^2 e^{\nu/2} (\bar{\psi} \psi - a \bar{\psi} \gamma^5 \psi \gamma^5) - e^{\nu/2} m \right\} \psi = 0$	$\left\{ i\gamma^0 \partial_0 - e^{(\nu-\mu)/2} \left[i\vec{\gamma} \cdot \vec{\partial} + i\gamma^1 \partial_r \left(\frac{\mu}{2} + \frac{\nu}{4} \right) \right] \right.$ $\left. + \frac{3\varepsilon}{8} \ell^2 e^{\nu/2} (\bar{\psi} \psi - a \bar{\psi} \gamma^5 \psi \gamma^5) - e^{\nu/2} m \right\} \psi = 0$
mielke-geometrodynamics_255 EQ0807 (11.4.19)	255	■ 0.812	$\vec{\sigma} \cdot \vec{L} \chi_\kappa^m = -(\kappa + 1) \chi_\kappa^m.$	$\vec{\sigma} \cdot \vec{L} \chi_\kappa^m = -(\kappa + 1) \chi_\kappa^m.$	$\vec{\sigma} \cdot \vec{L} \chi_\kappa^m = -(\kappa + 1) \chi_\kappa^m.$
mielke-geometrodynamics_132 EQ0397 (6.4.12)	132	■ 0.813	$\mathcal{H} \cong n^\alpha \mathcal{G}_\alpha + \Gamma_\perp^{\alpha\beta} \mathcal{G}_{\alpha\beta},$	$\mathcal{H} \cong n^\alpha \mathcal{G}_\alpha + \Gamma_\perp^{\alpha\beta} \mathcal{G}_{\alpha\beta},$	$\mathcal{H} \cong n^\alpha \mathcal{G}_\alpha + \Gamma_\perp^{\alpha\beta} \mathcal{G}_{\alpha\beta},$
mielke-geometrodynamics_227 EQ0709 (10.7.5)	227	■ 0.813	$R_{\{\beta\}\{\}}^{\{\}} \wedge \vartheta^{\beta} = \frac{\ell^2}{2} \varepsilon (x^\alpha Y \wedge X \wedge Y - y^\alpha X \wedge X \wedge Y) = 0.$	$R_\beta{}^\alpha \wedge \vartheta^\beta = \frac{\ell^2}{2} \varepsilon (x^\alpha Y \wedge X \wedge Y - y^\alpha X \wedge X \wedge Y) = 0.$	$R_\beta{}^\alpha \wedge \vartheta^\beta = \frac{\ell^2}{2} \varepsilon (x^\alpha Y \wedge X \wedge Y - y^\alpha X \wedge X \wedge Y) = 0.$
mielke-geometrodynamics_85 EQ0210 (4.2.2)	85	■ 0.814	$\stackrel{*}{\omega} \approx \left[\begin{array}{c c} \omega^L & \omega^T \\ \hline 0 & 0 \end{array} \right].$	$\stackrel{*}{\omega} \approx \left[\begin{array}{c c} \omega^L & \omega^T \\ \hline 0 & 0 \end{array} \right].$	$\stackrel{*}{\omega} \approx \left[\begin{array}{c c} \omega^L & \omega^T \\ \hline 0 & 0 \end{array} \right].$
mielke-geometrodynamics_244 EQ0769 (11.1.36)	244	■ 0.817	$\widetilde{\omega} \longrightarrow {}^S \widetilde{\omega} = S \widetilde{\omega} S^{-1} + (dS) S^{-1}$	$\widetilde{\omega} \longrightarrow {}^S \widetilde{\omega} = S \widetilde{\omega} S^{-1} + (dS) S^{-1}$	$\widetilde{\omega} \longrightarrow {}^S \widetilde{\omega} = S \widetilde{\omega} S^{-1} + (dS) S^{-1}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_008398 (6.4.14)	133	■ 0.820	$\begin{aligned} & \left(\mathbf{G}_\alpha(t, \mathbf{x}), \mathbf{G}_\beta(t, \mathbf{y}) \right) = \left(-T_{\alpha\beta}{}^\gamma \mathcal{G}_\gamma + R_{\alpha\beta\gamma}{}^\delta \mathcal{G}^\gamma{}_\delta \right) \cdot \delta(\mathbf{x} - \mathbf{y}), \\ & \left(\mathcal{G}^\alpha{}_\beta(t, \mathbf{x}), \mathcal{G}_\gamma(t, \mathbf{y}) \right) = \delta_\beta^\gamma \mathcal{G}_\beta \delta(\mathbf{x} - \mathbf{y}), \\ & \left(\mathcal{G}^\alpha{}_\beta(t, \mathbf{x}), \mathcal{G}^\gamma{}_\delta(t, \mathbf{y}) \right) = \left(\delta_\delta^\alpha \mathcal{G}^\gamma{}_\beta - \delta_\beta^\gamma \mathcal{G}^\alpha{}_\delta \right) \cdot \delta(\mathbf{x} - \mathbf{y}). \end{aligned}$	$\begin{aligned} & \left(\mathcal{G}_\alpha(t, \mathbf{x}), \mathcal{G}_\beta(t, \mathbf{y}) \right) = \left(-T_{\alpha\beta}{}^\gamma \mathcal{G}_\gamma + R_{\alpha\beta\gamma}{}^\delta \mathcal{G}^\gamma{}_\delta \right) \cdot \delta(\mathbf{x} - \mathbf{y}), \\ & \left(\mathcal{G}^\alpha{}_\beta(t, \mathbf{x}), \mathcal{G}_\gamma(t, \mathbf{y}) \right) = \delta_\beta^\gamma \mathcal{G}_\beta \delta(\mathbf{x} - \mathbf{y}), \\ & \left(\mathcal{G}^\alpha{}_\beta(t, \mathbf{x}), \mathcal{G}^\gamma{}_\delta(t, \mathbf{y}) \right) = \left(\delta_\delta^\alpha \mathcal{G}^\gamma{}_\beta - \delta_\beta^\gamma \mathcal{G}^\alpha{}_\delta \right) \cdot \delta(\mathbf{x} - \mathbf{y}). \end{aligned}$	$\begin{aligned} & \left(\mathcal{G}_\alpha(t, \mathbf{x}), \mathcal{G}_\beta(t, \mathbf{y}) \right) = \left(-T_{\alpha\beta}{}^\gamma \mathcal{G}_\gamma + R_{\alpha\beta\gamma}{}^\delta \mathcal{G}^\gamma{}_\delta \right) \cdot \delta(\mathbf{x} - \mathbf{y}), \\ & \left(\mathcal{G}^\alpha{}_\beta(t, \mathbf{x}), \mathcal{G}_\gamma(t, \mathbf{y}) \right) = \delta_\beta^\gamma \mathcal{G}_\beta \delta(\mathbf{x} - \mathbf{y}), \\ & \left(\mathcal{G}^\alpha{}_\beta(t, \mathbf{x}), \mathcal{G}^\gamma{}_\delta(t, \mathbf{y}) \right) = \left(\delta_\delta^\alpha \mathcal{G}^\gamma{}_\beta - \delta_\beta^\gamma \mathcal{G}^\alpha{}_\delta \right) \cdot \delta(\mathbf{x} - \mathbf{y}). \end{aligned}$
mielke-geometrodynamics_008059 (2.7.9)	45	■ 0.820	$\bar{\varphi} \rightarrow \bar{\varphi} G \quad \in C^\infty(V^{\bar{s}}).$	$\bar{\varphi} \rightarrow \bar{\varphi} G \quad \in C^\infty(V^{\bar{s}}).$	$\bar{\varphi} \rightarrow \bar{\varphi} G \quad \in C^\infty(V^{\bar{s}}).$
mielke-geometrodynamics_009993 (14.3.6)	320	■ 0.822	$\int_G \pi \hat{L}_D = \frac{i}{2} \left(\bar{\psi} \gamma \wedge {}^* D^{g+\omega} \psi + (\overline{D^{g+\omega} \psi}) \wedge {}^* \gamma \psi \right) + \frac{\hat{\ell}}{2} \bar{\psi} \gamma \wedge \gamma \wedge {}^* \Omega \psi - \hat{m} \bar{\psi} \psi \eta,$	$\int_G \pi \hat{L}_D = \frac{i}{2} \left(\bar{\psi} \gamma \wedge {}^* D^{g+\omega} \psi + (\overline{D^{g+\omega} \psi}) \wedge {}^* \gamma \psi \right) + \frac{\hat{\ell}}{2} \bar{\psi} \gamma \wedge \gamma \wedge {}^* \Omega \psi - \hat{m} \bar{\psi} \psi \eta,$	$\int_G \pi \hat{L}_D = \frac{i}{2} \left(\bar{\psi} \gamma \wedge {}^* D^{g+\omega} \psi + (\overline{D^{g+\omega} \psi}) \wedge {}^* \gamma \psi \right) + \frac{\hat{\ell}}{2} \bar{\psi} \gamma \wedge \gamma \wedge {}^* \Omega \psi - \hat{m} \bar{\psi} \psi \eta,$
mielke-geometrodynamics_00844 (12.2.12)	273	■ 0.831	$\mathrm{L} = \frac{1}{2\ell^2} \mathrm{Tr}(\Omega \wedge {}^* \sigma) + \mathrm{L}_D = \frac{1}{2\ell^2} \mathrm{R}^{\alpha\beta} \wedge \eta_{\alpha\beta} + \mathrm{L}_D,$	$\mathrm{L} = \frac{1}{2\ell^2} \mathrm{Tr}(\Omega \wedge {}^* \sigma) + \mathrm{L}_D = \frac{1}{2\ell^2} \mathrm{R}^{\alpha\beta} \wedge \eta_{\alpha\beta} + \mathrm{L}_D,$	$\mathrm{L} = \frac{1}{2\ell^2} \mathrm{Tr}(\Omega \wedge {}^* \sigma) + \mathrm{L}_D = \frac{1}{2\ell^2} \mathrm{R}^{\alpha\beta} \wedge \eta_{\alpha\beta} + \mathrm{L}_D,$
mielke-geometrodynamics_008218 (4.2.10)	86	■ 0.836	$\omega^L \rightarrow A^{-1} \omega^L = G^{-1} \omega^L G - G^{-1} dG,$ $\omega^T \rightarrow A^{-1} \omega^T = G^{-1} \omega^T - G^{-1} DT.$	$\omega^L \rightarrow A^{-1} \omega^L = G^{-1} \omega^L G - G^{-1} dG,$ $\omega^T \rightarrow A^{-1} \omega^T = G^{-1} \omega^T - G^{-1} DT.$	$\omega^L \rightarrow A^{-1} \omega^L = G^{-1} \omega^L G - G^{-1} dG,$ $\omega^T \rightarrow A^{-1} \omega^T = G^{-1} \omega^T - G^{-1} DT.$
mielke-geometrodynamics_008248 (4.3.5)	90	■ 0.836	$F(m, f\vartheta) = f F(m, \vartheta),$ $F(m, \vartheta) > 0 \quad \text{if} \quad \vartheta \neq 0.$	$F(m, f\vartheta) = f F(m, \vartheta),$ $F(m, \vartheta) > 0 \quad \text{if} \quad \vartheta \neq 0.$	$F(m, f\vartheta) = f F(m, \vartheta),$ $F(m, \vartheta) > 0 \quad \text{if} \quad \vartheta \neq 0.$
mielke-geometrodynamics_008445 (6.9.1)	144	■ 0.837	$ds^2 = - \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right)^{-1} dr^2 + r^2 d\Omega^2,$	$ds^2 = - \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right)^{-1} dr^2 + r^2 d\Omega^2,$	$ds^2 = - \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right) dt^2 + \left(1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right)^{-1} dr^2 + r^2 d\Omega^2,$
mielke-geometrodynamics_008227 (4.2.20)	88	■ 0.839	$\widetilde{\vartheta} \rightarrow A^{-1} \widetilde{\vartheta} = G^{-1} \widetilde{\vartheta}.$	$\widetilde{\vartheta} \rightarrow A^{-1} \widetilde{\vartheta} = G^{-1} \widetilde{\vartheta}.$	$\widetilde{\vartheta} \rightarrow A^{-1} \widetilde{\vartheta} = G^{-1} \widetilde{\vartheta}.$
mielke-geometrodynamics_008773 (11.2.2)	245	■ 0.840	$i\gamma \wedge {}^* D\psi - \frac{i}{2} (D \wedge {}^* \gamma) \psi - m\psi\eta = 0$	$i\gamma \wedge {}^* D\psi - \frac{i}{2} (D \wedge {}^* \gamma) \psi - m\psi\eta = 0$	$i\gamma \wedge {}^* D\psi - \frac{i}{2} (D \wedge {}^* \gamma) \psi - m\psi\eta = 0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_1E08811 (11.4.23)	256	0.840	$\frac{3}{8} \ell^2 \bar{\psi} \gamma^5 \psi = 4 \pi m e^{-\mu - v/2} \frac{2}{3} H F \left Y_{ \kappa -1}^{ \kappa -1}(\vartheta, \phi) \right ^2$		
mielke-geometrodynamics_1E09584 (9.1.2)	192	0.843	$\begin{aligned} e^{\nu} &= e^{-\lambda} = 1 - \left(\frac{2\mu}{r} \right)^{n-3} - \frac{2\Lambda_{\text{eff.}}}{(n-1)(n-2)} r^2 \\ \mu &= \frac{M}{m^{*2}}; \quad d\Omega^2 := d\vartheta^2 + \sin^2 \vartheta d\phi^2 \end{aligned}$		
mielke-geometrodynamics_1E09349 (5.4.7)	117	0.845	$\left\langle dj_5 \right\rangle = 2im \left\langle \bar{\psi} \gamma_5 \psi \right\rangle \eta - \frac{1}{96\pi^2} \left[2R_{\alpha\beta}^{\{\}} \wedge R^{\{\beta\alpha\beta} + d\mathscr{A} \wedge d\mathscr{A} \right]$		
mielke-geometrodynamics_1E09792 (11.4.4)	253	0.845	$\begin{aligned} ds^2 &= g_{kl} dx^k \otimes_s dx^l \\ &= e^v dt^2 - e^\mu \left(dr^2 + r^2 d\vartheta^2 + r^2 \sin^2 \vartheta d\phi^2 \right). \end{aligned}$		
mielke-geometrodynamics_1E09742 (11.1.9)	238	0.846	$\Lambda : \begin{cases} \widetilde{SO}_\circ(s, n-s) & \longrightarrow SO_\circ(s, n-s), \\ \widetilde{\Psi} & \longrightarrow g^\alpha{}_\beta = -\frac{i}{N} \text{Tr} \left(\gamma^\alpha \text{Si} \gamma_\beta S^{-1} \right), \end{cases}$		
mielke-geometrodynamics_1E08897 (13.5.2)	288	0.846	$\begin{aligned} \hat{R} &:= R_A{}^B E^A{}_B := \left[d\Gamma_A{}^B - \Gamma_A{}^C \wedge \Gamma_C{}^B \right] E^A{}_B \\ &= \left[R_\alpha{}^\beta - \frac{\Lambda}{3} \theta_\alpha \wedge \vartheta^\beta \right] L^\alpha{}_\beta + T^\beta P_\beta + D\theta_\alpha P_*^\alpha \end{aligned}$		
mielke-geometrodynamics_1E09296 (4.5.7)	101	0.847	$F_{\alpha} = (e_{\alpha} \lrcorner D\Psi) \frac{\delta L}{\delta \Psi} + (-1)^p (e_{\alpha} \lrcorner \Psi) \wedge D \frac{\delta L}{\delta \Psi}$		
mielke-geometrodynamics_1E09241 (4.3.6)	91	0.848	$g_{\alpha\beta}(m, \vartheta) := \frac{1}{2} \frac{\partial^2 F^2(m, \vartheta)}{\partial \vartheta^\alpha \otimes_s \partial \vartheta^\beta}.$		
mielke-geometrodynamics_1E09478 (7.4.6)	157	0.849	$\operatorname{Ric}_{\alpha\beta} := \operatorname{Ric}_{\alpha\beta} - \frac{1}{4} \operatorname{Rg}_{\alpha\beta} = 0.$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_101 EQ0295 (4.5.6)	101	0.850	$\begin{aligned} D \Sigma_{\alpha} &\wedge \left(\left(e_{\alpha} \rfloor T^{\beta} \right) \wedge \left(\left(e_{\alpha} \rfloor R^{\beta\gamma} \right) \wedge \tau_{\beta\gamma} + F_{\alpha} \right) \right) \\ &\simeq (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma}. \end{aligned}$	$D\Sigma_{\alpha} = (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma} + F_{\alpha}$ $\simeq (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma}.$	$D\Sigma_{\alpha} = (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma} + F_{\alpha}$ $\simeq (e_{\alpha} \rfloor T^{\beta}) \wedge \Sigma_{\beta} + (e_{\alpha} \rfloor R^{\beta\gamma}) \wedge \tau_{\beta\gamma}.$
mielke-geometrodynamics_272 EQ0842 (12.2.10)	272	0.852	$\mu := e_{\alpha} \downarrow \mu^{\alpha} = \frac{3}{4} * j_5$	$\mu := e_{\alpha} \downarrow \mu^{\alpha} = \frac{3}{4} * j_5$	$\mu := e_{\alpha} \rfloor \mu^{\alpha} = \frac{3}{4} * j_5$
mielke-geometrodynamics_164 EQ0506 (7.7.24)	164	0.852	$\gamma_5 := (i/4!)^* (\gamma \wedge \gamma \wedge \gamma \wedge \gamma)$	$\gamma_5 := (i/4!)^* (\gamma \wedge \gamma \wedge \gamma \wedge \gamma)$	$\gamma_5 := (i/4!)^* (\gamma \wedge \gamma \wedge \gamma \wedge \gamma)$
mielke-geometrodynamics_326 EQ1014 (14.4.9)	326	0.856	$M^2 = \left(M_{\text{ir}} + \frac{\alpha_e M^{*2}}{4M_{\text{ir}}} \text{Tr} \bar{Q}^2 \right)^2 + \frac{1}{4} J(J+1) \frac{M^{*4}}{M_{\text{ir}}^2},$	$M^2 = \left(M_{\text{ir}} + \frac{\alpha_e M^{*2}}{4M_{\text{ir}}} \text{Tr} \bar{Q}^2 \right)^2 + \frac{1}{4} J(J+1) \frac{M^{*4}}{M_{\text{ir}}^2},$	$M^2 = \left(M_{\text{ir}} + \frac{\alpha_e M^{*2}}{4M_{\text{ir}}} \text{Tr} \bar{Q}^2 \right)^2 + \frac{1}{4} J(J+1) \frac{M^{*4}}{M_{\text{ir}}^2},$
mielke-geometrodynamics_241 EQ0753 (11.1.20)	241	0.859	$\hat{\Psi}_{ \widehat{SO}(1,3)} = \begin{bmatrix} \Psi_1 \\ \Psi_2 \\ \vdots \\ \Psi_{[\frac{n}{4}]} \end{bmatrix}.$	$\hat{\Psi}_{ \widehat{SO}(1,3)} = \begin{bmatrix} \Psi_1 \\ \Psi_2 \\ \vdots \\ \Psi_{[\frac{n}{4}]} \end{bmatrix}.$	$\hat{\Psi}_{ \widehat{SO}(1,3)} = \begin{bmatrix} \Psi_1 \\ \Psi_2 \\ \vdots \\ \Psi_{[\frac{n}{4}]} \end{bmatrix}.$
mielke-geometrodynamics_247 EQ0783 (11.3.2)	247	0.862	$\begin{aligned} L_{\mathrm{D}} &= \frac{i}{2} \left(\bar{\psi} \gamma \wedge * D^{\dagger} \psi + \overline{(D^{\dagger} \psi)} \wedge * \gamma \psi \right) - m \bar{\psi} \psi \\ &+ \frac{1}{2} \bar{\psi} [\tilde{K}, * \gamma] \psi =: L_{\mathrm{D}}^{\dagger} + L_{\mathrm{NL}}. \end{aligned}$	$L_{\mathrm{D}} = \frac{i}{2} \left(\bar{\psi} \gamma \wedge * D^{\dagger} \psi + \overline{(D^{\dagger} \psi)} \wedge * \gamma \psi \right) - m \bar{\psi} \psi$ $+ \frac{1}{2} \bar{\psi} [\tilde{K}, * \gamma] \psi =: L_{\mathrm{D}}^{\dagger} + L_{\mathrm{NL}}.$	$L_{\mathrm{D}} = \frac{i}{2} \left(\bar{\psi} \gamma \wedge * D^{\dagger} \psi + \overline{(D^{\dagger} \psi)} \wedge * \gamma \psi \right) - m \bar{\psi} \psi$ $+ \frac{1}{2} \bar{\psi} [\tilde{K}, * \gamma] \psi =: L_{\mathrm{D}}^{\dagger} + L_{\mathrm{NL}}.$
mielke-geometrodynamics_369 EQ1100 (B.12)	369	0.864	$\{ \}^* \alpha^{\{p\}} := \frac{1}{(n-p)! \cdot (p!)} \sqrt{ \det g_{ij} } \varepsilon_{\alpha_1 \dots \alpha_p \beta_1 \dots \beta_{n-p}} \times A^{\alpha_1 \dots \alpha_p} \vartheta^{\beta_1} \wedge \dots \wedge \vartheta^{\beta_{n-p}}.$	$* \alpha^{\{p\}} := \frac{1}{(n-p)! \cdot (p!)} \sqrt{ \det g_{ij} } \varepsilon_{\alpha_1 \dots \alpha_p \beta_1 \dots \beta_{n-p}} \times A^{\alpha_1 \dots \alpha_p} \vartheta^{\beta_1} \wedge \dots \wedge \vartheta^{\beta_{n-p}}.$	$* \alpha^{\{p\}} := \frac{1}{(n-p)! \cdot (p!)} \sqrt{ \det g_{ij} } \varepsilon_{\alpha_1 \dots \alpha_p \beta_1 \dots \beta_{n-p}} \times A^{\alpha_1 \dots \alpha_p} \vartheta^{\beta_1} \wedge \dots \wedge \vartheta^{\beta_{n-p}}.$
mielke-geometrodynamics_74 EQ0184 (3.7.14)	74	0.864	$\vec{\Omega} = d\vec{\omega} - (\overrightarrow{\omega} \wedge)$	$\vec{\Omega} = d\vec{\omega} - (\overrightarrow{\omega} \wedge)$	$\vec{\Omega} = d\vec{\omega} - (\overrightarrow{\omega} \wedge \vec{\omega}) = d\vec{\omega} - \vec{\omega} \wedge \vec{\omega}.$
mielke-geometrodynamics_69 EQ0165 (3.5.13)	69	0.865	$\{ \}^* \Omega = (\stackrel{+}{-}) \Omega ;$	$* \Omega = (\stackrel{+}{-}) \Omega ;$	$* \Omega = (\stackrel{+}{-}) \Omega ;$
mielke-geometrodynamics_238 EQ0737 (11.1.4)	238	0.867	$x^{\alpha} \rightarrow x'^{\alpha} = g_{\beta}^{\alpha} x^{\beta}, [g_{\beta}^{\alpha}] \in SO(s, n-s), x^{\alpha} \in \mathbf{E}^n.$	$x^{\alpha} \rightarrow x'^{\alpha} = g_{\beta}^{\alpha} x^{\beta}, [g_{\beta}^{\alpha}] \in SO(s, n-s), x^{\alpha} \in \mathbf{E}^n.$	$x^{\alpha} \rightarrow x'^{\alpha} = g_{\beta}^{\alpha} x^{\beta}, [g_{\beta}^{\alpha}] \in SO(s, n-s), x^{\alpha} \in \mathbf{E}^n.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0853 (12.3.6)	276	■ 0.868	$\left\langle \vartheta^\alpha \wedge \sigma_\alpha \right\rangle = -\frac{1}{3\pi^2} \left[\text{Tr} (G \wedge {}^* G) + \frac{1}{24} \left(2R^{\alpha\beta} \wedge R_{\alpha\beta}^{(j\star)} + d\mathcal{A} \wedge {}^* d\mathcal{A} \right) \right],$	$\langle \vartheta^\alpha \wedge \sigma_\alpha \rangle = -\frac{1}{3\pi^2} \left[\text{Tr} (G \wedge {}^* G) + \frac{1}{24} \left(2R^{\alpha\beta} \wedge R_{\alpha\beta}^{(j\star)} + d\mathcal{A} \wedge {}^* d\mathcal{A} \right) \right],$	$\langle \vartheta^\alpha \wedge \sigma_\alpha \rangle = -\frac{1}{3\pi^2} \left[\text{Tr} (G \wedge {}^* G) + \frac{1}{24} \left(2R^{\alpha\beta} \wedge R_{\alpha\beta}^{(j\star)} + d\mathcal{A} \wedge {}^* d\mathcal{A} \right) \right],$
mielke-geometrodynamics_EQ0774 (11.2.3)	245	■ 0.868	$i(\overline{D\psi}) \wedge {}^* \gamma + \frac{i}{2} \bar{\psi} D \wedge {}^* \gamma + m \bar{\psi} \eta = 0,$	$i(\overline{D\psi}) \wedge {}^* \gamma + \frac{i}{2} \bar{\psi} D \wedge {}^* \gamma + m \bar{\psi} \eta = 0,$	$i(\overline{D\psi}) \wedge {}^* \gamma + \frac{i}{2} \bar{\psi} D \wedge {}^* \gamma + m \bar{\psi} \eta = 0,$
mielke-geometrodynamics_EQ0548 (8.4.2)	178	■ 0.871	$s L_{\text{Pontr}} = -2 d \text{Tr} \{ \Psi \wedge \Omega^g \} = -d (\Psi_\alpha{}^\beta \wedge R_\beta{}^\alpha).$	$s L_{\text{Pontr}} = -2 d \text{Tr} \{ \Psi \wedge \Omega^g \} = -d (\Psi_\alpha{}^\beta \wedge R_\beta{}^\alpha).$	$s L_{\text{Pontr}} = -2 d \text{Tr} \{ \Psi \wedge \Omega^g \} = -d (\Psi_\alpha{}^\beta \wedge R_\beta{}^\alpha).$
mielke-geometrodynamics_EQ0244 (4.3.9)	91	■ 0.873	$g_{ij} e_\alpha^i e_\beta^j = g_{\alpha\beta}.$	$g_{ij} e_\alpha^i e_\beta^j = g_{\alpha\beta}.$	$g_{ij} e_\alpha^i e_\beta^j = g_{\alpha\beta}.$
mielke-geometrodynamics_EQ0890 (13.4.9)	287	■ 0.876	$\hat{\xi} = \begin{pmatrix} \xi \\ 1 \end{pmatrix} = \begin{pmatrix} \xi^\alpha P_\alpha \\ 1 \end{pmatrix}, \quad \hat{\xi}_* = (\xi_*, 1) = (\xi_\beta^* P_\beta^*, 1)$	$\hat{\xi} = \begin{pmatrix} \xi \\ 1 \end{pmatrix} = \begin{pmatrix} \xi^\alpha P_\alpha \\ 1 \end{pmatrix}, \quad \hat{\xi}_* = (\xi_*, 1) = (\xi_\beta^* P_\beta^*, 1)$	$\hat{\xi} = \begin{pmatrix} \xi \\ 1 \end{pmatrix} = \begin{pmatrix} \xi^\alpha P_\alpha \\ 1 \end{pmatrix}, \quad \hat{\xi}_* = (\xi_*, 1) = (\xi_\beta^* P_\beta^*, 1)$
mielke-geometrodynamics_EQ1007 (14.4.2)	325	■ 0.877	$\begin{aligned} A^{(\circ)} &= {}^* \omega^{(\circ)} = -\frac{r^2}{\rho^2} \left[dt - J_3 \frac{\hbar}{Mc} \sin^2 \vartheta d\phi \right] Q^{(\circ)} \cos \delta \\ &\quad - \frac{\cos \vartheta}{\rho^2} \left[\left(r^2 + J(J+1) \frac{\hbar^2}{M^2 c^2} \right) d\phi - J_3 \frac{\hbar}{M} dt \right] P^{(\circ)} \sin \delta \end{aligned}$	$\begin{aligned} A^{(\circ)} &= {}^* \omega^{(\circ)} = -\frac{r^2}{\rho^2} \left[dt - J_3 \frac{\hbar}{Mc} \sin^2 \vartheta d\phi \right] Q^{(\circ)} \cos \delta \\ &\quad - \frac{\cos \vartheta}{\rho^2} \left[\left(r^2 + J(J+1) \frac{\hbar^2}{M^2 c^2} \right) d\phi - J_3 \frac{\hbar}{M} dt \right] P^{(\circ)} \sin \delta \end{aligned}$	$\begin{aligned} A^{(\circ)} &= {}^* \omega^{(\circ)} = -\frac{r^2}{\rho^2} \left[dt - J_3 \frac{\hbar}{Mc} \sin^2 \vartheta d\phi \right] Q^{(\circ)} \cos \delta \\ &\quad - \frac{\cos \vartheta}{\rho^2} \left[\left(r^2 + J(J+1) \frac{\hbar^2}{M^2 c^2} \right) d\phi - J_3 \frac{\hbar}{M} dt \right] P^{(\circ)} \sin \delta. \end{aligned}$
mielke-geometrodynamics_EQ1062 (16.2.1)	355	■ 0.877	$[\square - (mc/\hbar)^2] \varphi = 0$	$[\square - (mc/\hbar)^2] \varphi = 0$	$[\square - (mc/\hbar)^2] \varphi = 0$
mielke-geometrodynamics_EQ0051 (2.7.1)	44	■ 0.881	$G(p): P \rightarrow P, \quad G(p) \in \mathcal{G}_p$	$G(p): P \rightarrow P, \quad G(p) \in \mathcal{G}_p$	$G(p): P \rightarrow P, \quad G(p) \in \mathcal{G}_p$
mielke-geometrodynamics_EQ0178 (3.7.8)	72	■ 0.881	$\langle O ^\lambda \varphi O \rangle = \mu / \sqrt{\lambda}$	$\langle O ^\lambda \varphi O \rangle = \mu / \sqrt{\lambda}$	$\langle O ^\lambda \varphi O \rangle = \mu / \sqrt{\lambda}$
mielke-geometrodynamics_EQ0028 (2.5.1)	40	■ 0.885	$V_p^* = T_p^* (F_m(M)),$	$V_p^* = T_p^* (F_m(M)),$	$V_p^* = T_p^* (F_m(M)),$
mielke-geometrodynamics_EQ0920 (13.8.6)	295	■ 0.888	$\begin{aligned} dC_{\text{RR}(\star)} &:= \frac{1}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta(\star)} - \frac{1}{3} \Gamma_\alpha{}^{\beta(\star)} \wedge \Gamma_{\beta}{}^\gamma \wedge \Gamma_\gamma{}^\alpha \right) \\ &\equiv -L_{\text{SKY}} - 2 \text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} + \frac{1}{2} \text{Ric}_\alpha{}^\alpha \wedge {}^* \text{Ric}_\beta{}^\beta \end{aligned}$	$\begin{aligned} dC_{\text{RR}(\star)} &:= \frac{1}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta(\star)} - \frac{1}{3} \Gamma_\alpha{}^{\beta(\star)} \wedge \Gamma_{\beta}{}^\gamma \wedge \Gamma_\gamma{}^\alpha \right) \\ &\equiv -L_{\text{SKY}} - 2 \text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} + \frac{1}{2} \text{Ric}_\alpha{}^\alpha \wedge {}^* \text{Ric}_\beta{}^\beta \end{aligned}$	$\begin{aligned} dC_{\text{RR}(\star)} &:= \frac{1}{2} d \left(\Gamma_{\alpha\beta} \wedge R^{\alpha\beta(\star)} - \frac{1}{3} \Gamma_\alpha{}^{\beta(\star)} \wedge \Gamma_{\beta}{}^\gamma \wedge \Gamma_\gamma{}^\alpha \right) \\ &\equiv -L_{\text{SKY}} - 2 \text{Ric}_{\alpha\beta} \wedge {}^* \text{Ric}^{\alpha\beta} + \frac{1}{2} \text{Ric}_\alpha{}^\alpha \wedge {}^* \text{Ric}_\beta{}^\beta \end{aligned}$
mielke-geometrodynamics_EQ0674 (10.3.31)	220	■ 0.890	$\check{\mathcal{E}}_\alpha = -\frac{\theta_{\text{T}} \theta_{\text{TL}}}{2\ell^2} \left(\frac{1}{\theta_{\text{TL}}} d\vartheta_\alpha - \frac{4\ell}{\theta_{\text{T}} \theta_{\text{L}}} \check{\mathcal{J}}_\alpha^* \right),$	$\check{\mathcal{E}}_\alpha = -\frac{\theta_{\text{T}} \theta_{\text{TL}}}{2\ell^2} \left(\frac{1}{\theta_{\text{TL}}} d\vartheta_\alpha - \frac{4\ell}{\theta_{\text{T}} \theta_{\text{L}}} \check{\mathcal{J}}_\alpha^* \right),$	$\check{\mathcal{E}}_\alpha = -\frac{\theta_{\text{T}} \theta_{\text{TL}}}{2\ell^2} \left(\frac{1}{\theta_{\text{TL}}} d\vartheta_\alpha - \frac{4\ell}{\theta_{\text{T}} \theta_{\text{L}}} \check{\mathcal{J}}_\alpha^* \right),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_169 EQ0531 (7.7.51)		■ 0.890	$\left.\left.R:=e_{\beta}\right\rfloor e_{\alpha}\right\rfloor R^{\alpha\beta},\quad\left.R^{\alpha\beta}\wedge\vartheta_{\alpha}\wedge\vartheta_{\beta}\equiv R^{\alpha\beta}\wedge\eta_{\alpha\beta}\equiv-R\eta,\right.\quad\left.R^{\alpha\beta}\wedge\vartheta_{\alpha}\wedge\vartheta_{\beta}\equiv R^{\alpha\beta}\wedge\eta_{\alpha\beta}\equiv-R\eta,\right.$		
mielke-geometrodynamics_176 EQ0540 (8.3.1)		■ 0.891	$\begin{aligned} (d\oplus s)\gamma + [\widetilde{\Gamma}, \gamma] &= \widetilde{\Theta} \\ (d\oplus s)\widetilde{\Theta} + [\widetilde{\Gamma}, \widetilde{\Theta}] &\equiv [\widetilde{\Omega}^g, \gamma] \end{aligned}$		$(d\oplus s)\gamma + [\widetilde{\Gamma}, \gamma] = \widetilde{\Theta},$ $(d\oplus s)\widetilde{\Theta} + [\widetilde{\Gamma}, \widetilde{\Theta}] \equiv [\widetilde{\Omega}^g, \gamma],$
mielke-geometrodynamics_303 EQ0925 (14.1.5)		■ 0.892	$\begin{aligned} &D^*\Omega=\tau_e, \quad D\Omega\equiv 0, \\ &\frac{1}{2}*(\vartheta\wedge*\Omega^g)=\ell^{*2}(\Sigma_{\mathrm{YM}}+\Sigma_{\mathrm{mat}}), \end{aligned}$	$D^*\Omega=\tau_e,$ $D\Omega\equiv 0,$ $\frac{1}{2}*(\vartheta\wedge*\Omega^g)=\ell^{*2}(\Sigma_{\mathrm{YM}}+\Sigma_{\mathrm{mat}}),$	$D^*\Omega=\tau_e,$ $D\Omega\equiv 0,$ $\frac{1}{2}*(\vartheta\wedge*\Omega^g)=\ell^{*2}(\Sigma_{\mathrm{YM}}+\Sigma_{\mathrm{mat}}),$
mielke-geometrodynamics_283 EQ0867 (13.2.3)		■ 0.893	$B\cong dA,\quad dB\cong dF\equiv 0,$	$B\cong dA,\quad dB\cong dF\equiv 0,$	$B\cong dA,\quad dB\cong dF\equiv 0,$
mielke-geometrodynamics_201 EQ0609 (9.2.11)		■ 0.894	$\begin{aligned} &\left(1+\zeta^2\right)\int_{M^4}L_{\mathrm{SKY}}-(-1)^s\zeta(4\pi)^2\cdot\chi\left(M^4\right) \\ &=\frac{1}{4}\int_{M^4}\left(\Omega^g-\zeta^*\Omega^{g(*)}\right)\wedge^*(\Omega^g-\zeta^*\Omega^{g(*)})\geq 0, \end{aligned}$	$\left(1+\zeta^2\right)\int_{M^4}L_{\mathrm{SKY}}-(-1)^s\zeta(4\pi)^2\cdot\chi\left(M^4\right)$ $=\frac{1}{4}\int_{M^4}\left(\Omega^g-\zeta^*\Omega^{g(*)}\right)\wedge^*(\Omega^g-\zeta^*\Omega^{g(*)})\geq 0,$	$(1+\zeta^2)\int_{M^4}L_{\mathrm{SKY}}-(-1)^s\zeta(4\pi)^2\cdot\chi(M^4)$ $=\frac{1}{4}\int_{M^4}(\Omega^g-\zeta^*\Omega^{g(*)})\wedge^*(\Omega^g-\zeta^*\Omega^{g(*)})\geq 0,$
mielke-geometrodynamics_309 EQ0959 (14.1.41)		■ 0.894	$\oint d\delta=2\pi b_1=2\pi v$	$\oint d\delta=2\pi b_1=2\pi v$	$\oint d\delta=2\pi b_1=2\pi v$
mielke-geometrodynamics_102 EQ0299 (4.5.10)		■ 0.896	$\begin{aligned} &\delta g_{\alpha\beta}=2\varepsilon_{(\alpha}{}^\gamma g_{\beta)\gamma},\quad\delta\vartheta^\alpha=-\vartheta^\beta\varepsilon_\beta{}^\alpha, \\ &\delta\Gamma_\alpha{}^\beta=D\varepsilon_\alpha{}^\beta,\quad\delta\Psi=-\varepsilon_\alpha{}^\beta I^\alpha{}_\beta\Psi. \end{aligned}$	$\delta g_{\alpha\beta}=2\varepsilon_{(\alpha}{}^\gamma g_{\beta)\gamma},\quad\delta\vartheta^\alpha=-\vartheta^\beta\varepsilon_\beta{}^\alpha,$ $\delta\Gamma_\alpha{}^\beta=D\varepsilon_\alpha{}^\beta,\quad\delta\Psi=-\varepsilon_\alpha{}^\beta I^\alpha{}_\beta\Psi.$	$\delta g_{\alpha\beta}=2\varepsilon_{(\alpha}{}^\gamma g_{\beta)\gamma},\quad\delta\vartheta^\alpha=-\vartheta^\beta\varepsilon_\beta{}^\alpha,$ $\delta\Gamma_\alpha{}^\beta=D\varepsilon_\alpha{}^\beta,\quad\delta\Psi=-\varepsilon_\alpha{}^\beta I^\alpha{}_\beta\Psi.$
mielke-geometrodynamics_90 EQ0236 (4.3.1)		■ 0.896	$\begin{aligned} (\Delta s)^2 &= -c^2(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 \\ &= g_{ij}\Delta x^i\Delta x^j. \end{aligned}$	$(\Delta s)^2 = -c^2(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2$ $= g_{ij}\Delta x^i\Delta x^j.$	$(\Delta s)^2 = -c^2(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2$ $= g_{ij}\Delta x^i\Delta x^j.$
mielke-geometrodynamics_323 EQ1005 (14.3.18)		■ 0.896	$\begin{aligned} &\not{D}^2\gamma\wedge\gamma\wedge^*\{D,D\}+\frac{1}{2}\gamma\wedge\gamma\wedge^*\tilde{\Omega}-\gamma\wedge\tilde{\Theta}\wedge^*D \\ &= \left(\Box^g+\frac{1}{4}R\right)\eta-\vartheta\wedge\Theta\wedge^*D \end{aligned}$	$D^2=\mathrm{Tr}\left(\frac{1}{2}\gamma\wedge\gamma\wedge^*\{D,D\}+\frac{1}{2}\gamma\wedge\gamma\wedge^*\tilde{\Omega}-\gamma\wedge\tilde{\Theta}\wedge^*D\right)$ $=\left(\Box^g+\frac{1}{4}R\right)\eta-\vartheta\wedge\Theta\wedge^*D$	$\not{D}^2=\mathrm{Tr}\left(\frac{1}{2}\gamma\wedge\gamma\wedge^*\{D,D\}+\frac{1}{2}\gamma\wedge\gamma\wedge^*\tilde{\Omega}-\gamma\wedge\tilde{\Theta}\wedge^*D\right)$ $=(\Box^g+\frac{1}{4}R)\eta-\vartheta\wedge\Theta\wedge^*D$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E08238 (4.2.23)	88	0.899	$\stackrel{*}{\sigma} \Theta = \frac{1}{2} T_{\alpha\beta}^{\cdot c} P_c \otimes \vartheta^\alpha \wedge \vartheta^\beta = \frac{1}{2} T_{ij}^{\cdot c} P_c \otimes dx^i \wedge dx^j,$		
mielke-geometrodynamics_E08723 (10.10.3)	230	0.900	$\begin{aligned} \delta_{\text{susy}} \vartheta^\alpha &= i\bar{\sigma} \Psi \gamma^\alpha, & \delta_{\text{susy}} \Gamma_\alpha^* &= i\bar{\sigma} \gamma_\alpha^* D\Psi + ic\bar{\sigma} (\gamma_\alpha \Psi + e_\alpha]^* \Psi), \\ \delta_{\text{susy}} \Psi &= 2D\sigma + c\gamma\sigma, \end{aligned}$		$\delta_{\text{susy}} \vartheta^\alpha = i\bar{\sigma} \Psi \gamma^\alpha, \quad \delta_{\text{susy}} \Gamma_\alpha^* = i\bar{\sigma} \gamma_\alpha^* D\Psi + ic\bar{\sigma} (\gamma_\alpha \Psi + e_\alpha]^* \Psi),$ $\delta_{\text{susy}} \Psi = 2D\sigma + c\gamma\sigma, \tag{1}$
mielke-geometrodynamics_E08216 (4.2.8)	86	0.902	$\widetilde{\omega} \rightarrow^{A^{-1}} \widetilde{\omega} = A^{-1} \widetilde{\omega} A - A^{-1} dA, \quad A \in \mathscr{A}_p.$		$\widetilde{\omega} \rightarrow^{A^{-1}} \widetilde{\omega} = A^{-1} \widetilde{\omega} A - A^{-1} dA, \quad A \in \mathscr{A}_p.$
mielke-geometrodynamics_E01842 (15.1.19)	340	0.902	$\Gamma_{\mu\alpha\beta} : (2^+) \oplus 2(1^-) \oplus (0^+) \oplus (2^-) \oplus 2(1^+) \oplus (0^-),$		$\Gamma_{\mu\alpha\beta} : (2^+) \oplus 2(1^-) \oplus (0^+) \oplus (2^-) \oplus 2(1^+) \oplus (0^-),$
mielke-geometrodynamics_E08481 (7.5.2)	157	0.903	$\square^c g_{ij} = \square^c \left(\Omega^2 o_{ij}^{\text{Mink}} \right) = \Omega^6 \square^c o_{ij}^{\text{Mink}} = 0$		$\square^c g_{ij} = \square^c (\Omega^2 o_{ij}^{\text{Mink}}) = \Omega^6 \square^c o_{ij}^{\text{Mink}} = 0$
mielke-geometrodynamics_E08117 (3.3.18)	59	0.905	$L_{\text{Max}} := -\frac{1}{2} F \wedge^* F = \frac{1}{2} (\mathbf{E}^2 - \mathbf{B}^2) \eta,$		$L_{\text{Max}} := -\frac{1}{2} F \wedge^* F = \frac{1}{2} (\mathbf{E}^2 - \mathbf{B}^2) \eta,$
mielke-geometrodynamics_E01118 (B.22)	371	0.905	$\square \alpha = 0,$		$\square \alpha = 0,$
mielke-geometrodynamics_E08831 (12.1.3)	270	0.907	$\tilde{j}_5 := j_5 + (1/96\pi^2) A \wedge dA$		$\tilde{j}_5 := j_5 + (1/96\pi^2) A \wedge dA$
mielke-geometrodynamics_E08489 (6.4.29)	136	0.907	$T^A = \frac{1}{f} (df \wedge \vartheta^A - 2\eta^{0A\beta\gamma} \vartheta_\beta \wedge \vartheta_\gamma) = -\frac{2}{f} \eta^A{}_{\mu\nu} \vartheta^\mu \wedge \vartheta^\nu = \pm^* T^A, \quad T^{\hat{0}} = 0,$		$T^A = \frac{1}{f} (df \wedge \vartheta^A - 2\eta^{0A\beta\gamma} \vartheta_\beta \wedge \vartheta_\gamma) = -\frac{2}{f} \eta^A{}_{\mu\nu} \vartheta^\mu \wedge \vartheta^\nu = \pm^* T^A, \quad T^{\hat{0}} = 0,$
mielke-geometrodynamics_E08657 (10.3.14)	217	0.907	$E_\alpha = e_\alpha] L(\star) - \frac{2\gamma}{\delta \ell^2} T_\alpha,$		$E_\alpha = e_\alpha] L(\star) - \frac{2\gamma}{\delta \ell^2} T_\alpha,$
mielke-geometrodynamics_E08396 (6.4.11)	132	0.909	$\begin{aligned} \mathcal{H} \cong & n^\alpha \frac{\delta L}{\delta \vartheta^\alpha} + \Gamma_\perp^{\alpha\beta} \frac{\delta L}{\delta \Gamma_{\alpha\beta}} \\ & - \underline{d}[(n]\Psi) \wedge \frac{\partial L}{\partial D\Psi} + n^\alpha \frac{\partial L}{\partial T^\alpha} + \Gamma_\perp^{\alpha\beta} \frac{\partial L}{\partial R^{\alpha\beta}} \Big]. \end{aligned}$		$\mathcal{H} \cong n^\alpha \frac{\delta L}{\delta \vartheta^\alpha} + \Gamma_\perp^{\alpha\beta} \frac{\delta L}{\delta \Gamma_{\alpha\beta}} - \underline{d}[(n]\Psi) \wedge \frac{\partial L}{\partial D\Psi} + n^\alpha \frac{\partial L}{\partial T^\alpha} + \Gamma_\perp^{\alpha\beta} \frac{\partial L}{\partial R^{\alpha\beta}} \Big].$
mielke-geometrodynamics_E08641 (10.2.12)	213	0.909	$ds^2 = -(1 - \Lambda_{\text{eff}} r^2) dt^2 + (1 - \Lambda_{\text{eff}} r^2)^{-1} dr^2 + r^2 d\phi^2$		$ds^2 = -(1 - \Lambda_{\text{eff}} r^2) dt^2 + (1 - \Lambda_{\text{eff}} r^2)^{-1} dr^2 + r^2 d\phi^2$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0263 (4.3.28)	94	■ 0.912	$\Gamma_{ij}^{\cdot k} = e_b^{\cdot \kappa} \Gamma_{i\alpha}^{\cdot b} E_{\cdot j}^\alpha + e_\beta^{\cdot \kappa} \partial_i E_{\cdot j}^\beta$	$\Gamma_{ij}^{\cdot k} = e_b^{\cdot \kappa} \Gamma_{i\alpha}^{\cdot b} E_{\cdot j}^\alpha + e_\beta^{\cdot \kappa} \partial_i E_{\cdot j}^\beta$	$\Gamma_{ij}^{\cdot \kappa} = e_b^{\cdot \kappa} \Gamma_{i\alpha}^{\cdot b} E_{\cdot j}^\alpha + e_\beta^{\cdot \kappa} \partial_i E_{\cdot j}^\beta$
mielke-geometrodynamics_EQ0364 (6.2.16)	126	■ 0.912	$D \, D \, \wedge_{\lambda_{\alpha \, \beta}} = -2 \, R_{[\lambda_{\alpha \, \mid \beta}]} \wedge_{\gamma} = 0$	$DD\lambda_{\alpha\beta} = -2R_{[\alpha]}{}^\gamma \wedge \lambda_{\gamma \beta]} = 0$	$DD\lambda_{\alpha\beta} = -2R_{[\alpha]}{}^\gamma \wedge \lambda_{\gamma \beta]} = 0$
mielke-geometrodynamics_EQ0840 (12.2.8)	272	■ 0.916	$\begin{aligned} \tau_{\alpha\beta} &:= \frac{\partial L_D}{\partial \Gamma^{\alpha\beta}} = \frac{1}{8} \bar{\psi} \left(\gamma^\alpha \sigma_{\alpha\beta} + \sigma_{\alpha\beta} \gamma^\beta \right) \psi \\ &= \frac{1}{4} \eta_{\alpha\beta\gamma\delta} \bar{\psi} \gamma^\delta \gamma_5 \psi \eta^\gamma = \tau_{\alpha\beta\gamma} \eta^\gamma = \vartheta_{[\alpha} \wedge \mu_{\beta]} . \end{aligned}$	$\tau_{\alpha\beta} := \frac{\partial L_D}{\partial \Gamma^{\alpha\beta}} = \frac{1}{8} \bar{\psi} \left(\gamma^\alpha \sigma_{\alpha\beta} + \sigma_{\alpha\beta} \gamma^\beta \right) \psi \\ = \frac{1}{4} \eta_{\alpha\beta\gamma\delta} \bar{\psi} \gamma^\delta \gamma_5 \psi \eta^\gamma = \tau_{\alpha\beta\gamma} \eta^\gamma = \vartheta_{[\alpha} \wedge \mu_{\beta]}$	$\tau_{\alpha\beta} := \frac{\partial L_D}{\partial \Gamma^{\alpha\beta}} = \frac{1}{8} \bar{\psi} \left(\gamma^\alpha \sigma_{\alpha\beta} + \sigma_{\alpha\beta} \gamma^\beta \right) \psi \\ = \frac{1}{4} \eta_{\alpha\beta\gamma\delta} \bar{\psi} \gamma^\delta \gamma_5 \psi \eta^\gamma = \tau_{\alpha\beta\gamma} \eta^\gamma = \vartheta_{[\alpha} \wedge \mu_{\beta]} .$
mielke-geometrodynamics_EQ0232 (4.2.25)	88	■ 0.917	$\Theta \rightarrow \Theta = G^{-1} \Theta$	$\Theta \rightarrow {}^{A^{-1}}\Theta = G^{-1}\Theta,$	$\Theta \rightarrow {}^{A^{-1}}\Theta = G^{-1}\Theta,$
mielke-geometrodynamics_EQ0577 (8.6.15)	185	■ 0.920	$-E_\alpha = \frac{\theta_T}{2} R^{\{\beta\gamma\}} \wedge \eta_{\alpha\beta\gamma} - \theta_T^* \Lambda_{\text{eff}} \eta_\alpha = \ell^2 \widehat{\Sigma}_\alpha .$	$-E_\alpha = \frac{\theta_T^*}{2} R^{\{\beta\gamma\}} \wedge \eta_{\alpha\beta\gamma} - \theta_T^* \Lambda_{\text{eff}} \eta_\alpha = \ell^2 \widehat{\Sigma}_\alpha .$	$-E_\alpha = \frac{\theta_T^*}{2} R^{\{\beta\gamma\}} \wedge \eta_{\alpha\beta\gamma} - \theta_T^* \Lambda_{\text{eff}} \eta_\alpha = \ell^2 \widehat{\Sigma}_\alpha .$
mielke-geometrodynamics_EQ1031 (15.1.8)	337	■ 0.920	$\begin{aligned} &\frac{1}{2} \left(\vartheta^f \wedge \Omega^f \right) - a \ell^2 \vartheta^f \wedge \left(2 L_{fg} - \frac{1}{2 b \ell^2} \left(\Omega^g \wedge \vartheta^g \wedge \vartheta^g + 2 \Lambda \eta \right) \right) \\ &= \ell^2 \left(\Sigma_{\text{m}}(\text{hadrons}) + \frac{a}{b} \left(\vartheta^g \wedge \Sigma_{\text{m}}(\text{leptons}) \right) \wedge \vartheta^f \right) \end{aligned}$	$\frac{1}{2} \left(\vartheta^f \wedge \Omega^f \right) - a \ell^2 \vartheta^f \wedge \left\{ 2 L_{fg} - \frac{1}{2 b \ell^2} \left(\Omega^g \wedge \vartheta^g \wedge \vartheta^g + 2 \Lambda \eta \right) \right\} \\ = \ell^2 \left\{ \Sigma_{\text{m}}(\text{ hadrons }) + \frac{a}{b} \left(\vartheta^g \wedge \Sigma_{\text{m}}(\text{ leptons }) \right) \wedge \vartheta^f \right\}$	$\frac{1}{2} \left(\vartheta^f \wedge \Omega^f \right) - a \ell^2 \vartheta^f \wedge \left\{ 2 L_{fg} - \frac{1}{2 b \ell^2} \left(\Omega^g \wedge \vartheta^g \wedge \vartheta^g + 2 \Lambda \eta \right) \right\} \\ = \ell^2 \left\{ \Sigma_{\text{m}}(\text{hadrons}) + \frac{a}{b} \left(\vartheta^g \wedge \Sigma_{\text{m}}(\text{leptons}) \right) \wedge \vartheta^f \right\}$
mielke-geometrodynamics_EQ0473 (7.4.1)	156	■ 0.921	$\begin{aligned} &\int \mathcal{D}\Gamma \exp \left[\int L_{\text{SKY}}^{(*)} / \hbar \right] \\ &= \int \mathcal{D}\Gamma \exp \left[- \int \left(R_{\alpha\beta} + {}^* R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} + {}^* R^{\alpha\beta(*)} \right) / 4 \hbar \right] \end{aligned}$	$\int \mathcal{D}\Gamma \exp \left[\int L_{\text{SKY}}^{(*)} / \hbar \right] \\ = \int \mathcal{D}\Gamma \exp \left[- \int \left(R_{\alpha\beta} + {}^* R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} + {}^* R^{\alpha\beta(*)} \right) / 4 \hbar \right]$	$\int \mathcal{D}\Gamma \exp \left[\int L_{\text{SKY}}^{(*)} / \hbar \right] \\ = \int \mathcal{D}\Gamma \exp \left[- \int \left(R_{\alpha\beta} + {}^* R_{\alpha\beta}^{(*)} \right) \wedge {}^* \left(R^{\alpha\beta} + {}^* R^{\alpha\beta(*)} \right) / 4 \hbar \right]$
mielke-geometrodynamics_EQ0343 (5.4.1)	115	■ 0.922	$k \, \partial_k \Gamma_k = \frac{1}{2} \, \text{S Tr} \left\{ \left[\Gamma_k^{(2)} + R_k \right]^{-1} \left(k \, \partial_k R_k \right) \right\} ,$	$k\partial_k\Gamma_k = \frac{1}{2} \, \text{S Tr} \left\{ \left[\Gamma_k^{(2)} + R_k \right]^{-1} \left(k \, \partial_k R_k \right) \right\} ,$	$k\partial_k\Gamma_k = \frac{1}{2} \text{STr} \left\{ \left[\Gamma_k^{(2)} + R_k \right]^{-1} \left(k \, \partial_k R_k \right) \right\} ,$
mielke-geometrodynamics_EQ0829 (12.1.1)	269	■ 0.922	$\bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \rightarrow \bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \mathbf{P} \exp \left\{ i \int_x^{x+\varepsilon} A \right\} .$	$\bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \rightarrow \bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \mathbf{P} \exp \left\{ i \int_x^{x+\varepsilon} A \right\} .$	$\bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \rightarrow \bar{\psi}(x) \gamma_5^* \gamma \psi(x + \varepsilon) \mathbf{P} \exp \left\{ i \int_x^{x+\varepsilon} A \right\} .$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0736 (11.1.3)	238	■ 0.922	$\gamma^0 := \left(\begin{array}{rr} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{array} \right), \quad \gamma^k := \left(\begin{array}{rr} 0 & \sigma^k \\ -\sigma^k & 0 \end{array} \right), \quad \sigma^k : \text{Pauli matrices.}$	$\gamma^0 := \left(\begin{array}{rr} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{array} \right), \quad \gamma^k := \left(\begin{array}{rr} 0 & \sigma^k \\ -\sigma^k & 0 \end{array} \right), \quad \sigma^k : \text{Pauli matrices.}$	$\gamma^0 := \left(\begin{array}{rr} \mathbb{1} & 0 \\ 0 & -\mathbb{1} \end{array} \right), \quad \gamma^k := \left(\begin{array}{rr} 0 & \sigma^k \\ -\sigma^k & 0 \end{array} \right), \quad \sigma^k : \text{Pauli matrices.}$
mielke-geometrodynamics_EQ0661 (10.3.18)	217	■ 0.923	$\square H - (-1)^{s*} D^* D H = dt + \frac{\gamma}{\delta \ell^2} d\mathcal{A} + \frac{1}{\delta \ell} dh,$	$\square H - (-1)^{s*} D^* D H = dt + \frac{\gamma}{\delta \ell^2} d\mathcal{A} + \frac{1}{\delta \ell} dh,$	$\square H - (-1)^s D^* D H = dt + \frac{\gamma}{\delta \ell^2} d\mathcal{A} + \frac{1}{\delta \ell} dh,$
mielke-geometrodynamics_EQ0685 (10.3.43)	222	■ 0.923	$C_{\alpha} - E_{\alpha} = \Sigma_{\alpha} - D^{\{ \}} \mu_{\alpha},$	$C_{\alpha} - E_{\alpha} = \Sigma_{\alpha} - D^{\{ \}} \mu_{\alpha},$	$C_{\alpha} - E_{\alpha} = \Sigma_{\alpha} - D^{\{ \}} \mu_{\alpha},$
mielke-geometrodynamics_EQ0702 (10.6.1)	225	■ 0.924	$L_{\{A\}} = \frac{1}{2} \left(F \wedge {}^* F + m_{\text{photon}} F \wedge A \right).$	$L_A = \frac{1}{2} \left(F \wedge {}^* F + m_{\text{photon}} F \wedge A \right).$	$L_A = \frac{1}{2} (F \wedge {}^* F + m_{\text{photon}} F \wedge A).$
mielke-geometrodynamics_EQ0221 (4.2.14)	87	■ 0.924	$\Omega^{\mathrm{T}} \rightarrow {}^{A^{-1}} \Omega^{\mathrm{T}} = G^{-1} \left(\Omega^{\mathrm{T}} - D D T \right) = G^{-1} \Omega^{\mathrm{T}} - G^{-1} \Omega^{\mathrm{L}} \otimes T.$	$\Omega^{\mathrm{T}} \rightarrow {}^{A^{-1}} \Omega^{\mathrm{T}} = G^{-1} \left(\Omega^{\mathrm{T}} - D D T \right) = G^{-1} \Omega^{\mathrm{T}} - G^{-1} \Omega^{\mathrm{L}} \otimes T.$	$\Omega^{\mathrm{T}} \rightarrow {}^{A^{-1}} \Omega^{\mathrm{T}} = G^{-1} (\Omega^{\mathrm{T}} - D D T) = G^{-1} \Omega^{\mathrm{T}} - G^{-1} \Omega^{\mathrm{L}} \otimes T.$
mielke-geometrodynamics_EQ0443 (6.8.14)	143	■ 0.924	$H_{\alpha} = -\frac{1}{\ell^2} * \left(\sum_{i=1}^3 a_i^{(i)} T_{\alpha} \right),$	$H_{\alpha} = -\frac{1}{\ell^2} * \left(\sum_{i=1}^3 a_i^{(i)} T_{\alpha} \right),$	$H_{\alpha} = -\frac{1}{\ell^2} * \left(\sum_{i=1}^3 a_i^{(i)} T_{\alpha} \right),$
mielke-geometrodynamics_EQ0192 (4.1.2)	81	■ 0.924	$a := \left[\begin{array}{c c} g & a \\ \hline 0 & 1 \end{array} \right] = \left[\begin{array}{c c} \mathbb{1}_n & a \\ \hline 0 & 1 \end{array} \right] \left[\begin{array}{c c} g & 0 \\ \hline 0 & 1 \end{array} \right] = a^{\mathrm{T}} \cdot a^{\mathrm{L}} \in A(n, \mathbb{R}).$	$a := \left[\begin{array}{c c} g & a \\ \hline 0 & 1 \end{array} \right] = \left[\begin{array}{c c} \mathbb{1}_n & a \\ \hline 0 & 1 \end{array} \right] \left[\begin{array}{c c} g & 0 \\ \hline 0 & 1 \end{array} \right] = a^{\mathrm{T}} \cdot a^{\mathrm{L}} \in A(n, \mathbb{R}).$	$a := \left[\begin{array}{c c} g & a \\ \hline 0 & 1 \end{array} \right] = \left[\begin{array}{c c} \mathbb{1}_n & a \\ \hline 0 & 1 \end{array} \right] \left[\begin{array}{c c} g & 0 \\ \hline 0 & 1 \end{array} \right] = a^{\mathrm{T}} \cdot a^{\mathrm{L}} \in A(n, \mathbb{R}).$
mielke-geometrodynamics_EQ0857 (12.4.4)	278	■ 0.925	$\lfloor \mathbf{b}_{\{n\}}^{\mathscr{G}^A} \rfloor \cong n \right\rfloor \left(D \underline{\tau}^A \right),$	<i>(not rendered)</i>	$\mathbb{L}_n \mathscr{G}^A \cong n \rfloor (D \underline{\tau}^A),$
mielke-geometrodynamics_EQ1025 (15.1.1)	336	■ 0.926	$\begin{aligned} L_{f-g} = & \frac{1}{\ell^2} L_{\mathrm{W}}(f) + \frac{1}{\ell^{*2}} L_{\mathrm{W}}(g) + L_{fg} + \frac{\Lambda}{\ell^{*2}} \vartheta^g \wedge \vartheta^g \wedge \vartheta^g \wedge \vartheta^g \\ & + L_{\mathrm{m}}(\text{hadrons}, f) + L_{\mathrm{m}}(\text{leptons}, g). \end{aligned}$	$L_{f-g} = \frac{1}{\ell^2} L_{\mathrm{W}}(f) + \frac{1}{\ell^{*2}} L_{\mathrm{W}}(g) + L_{fg} + \frac{\Lambda}{\ell^{*2}} \vartheta^g \wedge \vartheta^g \wedge \vartheta^g \wedge \vartheta^g + L_{\mathrm{m}}(\text{hadrons}, f) + L_{\mathrm{m}}(\text{leptons}, g).$	$L_{f-g} = \frac{1}{\ell^2} L_{\mathrm{W}}(f) + \frac{1}{\ell^{*2}} L_{\mathrm{W}}(g) + L_{fg} + \frac{\Lambda}{\ell^{*2}} \vartheta^g \wedge \vartheta^g \wedge \vartheta^g \wedge \vartheta^g + L_{\mathrm{m}}(\text{hadrons}, f) + L_{\mathrm{m}}(\text{leptons}, g).$
mielke-geometrodynamics_EQ0109 (3.3.10)	58	■ 0.927	$L_{\{\mathrm{Max}\}} = -\frac{1}{2} F \wedge {}^* F,$	$L_{\mathrm{Max}} = -\frac{1}{2} F \wedge {}^* F,$	$L_{\mathrm{Max}} = -\frac{1}{2} F \wedge {}^* F,$
mielke-geometrodynamics_EQ0587 (9.1.5)	194	■ 0.928	$dr^* := e^{(\lambda-v)/2} dr.$	$dr^* := e^{(\lambda-v)/2} dr.$	$dr^* := e^{(\lambda-v)/2} dr.$
mielke-geometrodynamics_EQ1101 (B.13)	370	■ 0.929	$*(\underbrace{\vartheta \wedge \cdots \wedge \vartheta}_{\mathrm{p~products}}) = \frac{p!}{(n-p)!} \underbrace{\vartheta \wedge \cdots \wedge \vartheta}_{\mathrm{n-p~products}}(\star),$	$*(\underbrace{\vartheta \wedge \cdots \wedge \vartheta}_{\mathrm{p~products}}) = \frac{p!}{(n-p)!} \underbrace{\vartheta \wedge \cdots \wedge \vartheta}_{\mathrm{n-p~products}}(\star),$	$*(\underbrace{\vartheta \wedge \cdots \wedge \vartheta}_{\mathrm{p~products}}) = \frac{p!}{(n-p)!} \underbrace{\vartheta \wedge \cdots \wedge \vartheta}_{\mathrm{n-p~products}}(\star),$
mielke-geometrodynamics_EQ0413 (6.5.1)	137	■ 0.929	$W(A, \gamma) = \operatorname{Tr} \mathbf{P} \exp \oint A = \operatorname{Tr} \mathbf{P} \exp \oint ds \dot{\gamma}^i(s) A_i^J(\gamma(s)) \lambda_J,$	$W(A, \gamma) = \operatorname{Tr} \mathbf{P} \exp \oint A = \operatorname{Tr} \mathbf{P} \exp \oint ds \dot{\gamma}^i(s) A_i^J(\gamma(s)) \lambda_J,$	$W(A, \gamma) = \operatorname{Tr} \mathbf{P} \exp \oint A = \operatorname{Tr} \mathbf{P} \exp \oint ds \dot{\gamma}^i(s) A_i^J(\gamma(s)) \lambda_J,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_153 EQ0465 (7.3.4)	153	■ 0.929	$\backslash\mathrm{left}.\backslash\mathrm{left}.\mathrm{E}_{\{\alpha\}}:=\partial_{\mathrm{L}_{\mathrm{SKY}}} / \backslash\mathrm{partial} \backslash\mathrm{vartheta}^{\{\alpha\}}=\frac{1}{2}\backslash\mathrm{left}(\mathrm{e}_{\{\alpha\}} \backslash\mathrm{right}\backslash\mathrm{rfloor} \mathrm{R}^{\{\mu} \backslash\mathrm{nu}\} \backslash\mathrm{wedge}\{ \}^{\{*\}} \mathrm{R}_{\{\mu} \backslash\mathrm{nu}\}}-\mathrm{R}^{\{\mu} \backslash\mathrm{nu}\} \backslash\mathrm{wedge} \mathrm{e}_{\{\alpha\}}\backslash\mathrm{right}\backslash\mathrm{rfloor}\{ \}^{\{*\}} \mathrm{R}_{\{\mu} \backslash\mathrm{nu}\}}\backslash\mathrm{right})=0$	$E_{\alpha}:=\partial L_{\mathrm{SKY}}/\partial\vartheta^{\alpha}=\frac{1}{2}\left(e_{\alpha}\rfloor R^{\mu\nu}\wedge{}^*R_{\mu\nu}-R^{\mu\nu}\wedge e_{\alpha}\rfloor{}^*R_{\mu\nu}\right)=0$	$E_{\alpha}:=\partial L_{\mathrm{SKY}}/\partial\vartheta^{\alpha}=\frac{1}{2}\left(e_{\alpha}\rfloor R^{\mu\nu}\wedge{}^*R_{\mu\nu}-R^{\mu\nu}\wedge e_{\alpha}\rfloor{}^*R_{\mu\nu}\right)=0$
mielke-geometrodynamics_338 EQ1034 (15.1.11)	338	■ 0.930	$\backslash\mathrm{Lambda}_{\{\mathrm{text}\{eff.\}\}^{\{f\}}}=\frac{2}{\kappa b}a \backslash\mathrm{Lambda}-2a\ell^{2*}L_{fg}$	$\Lambda_{\mathrm{eff.}}^f=\frac{2a}{\kappa b}\Lambda-2a\ell^{2*}L_{fg}$	$\Lambda_{\mathrm{eff.}}^f=\frac{2a}{\kappa b}\Lambda-2a\ell^{2*}L_{fg}$
mielke-geometrodynamics_165 EQ0512 (7.7.30)	165	■ 0.931	$\backslash\mathrm{begin}\{\mathrm{aligned}\} \backslash\mathrm{Omega}^{\{g\}}:=\mathrm{d} \backslash\mathrm{Gamma}+\backslash\mathrm{Gamma} \backslash\mathrm{wedge} \backslash\mathrm{Gamma} \&=\frac{i}{4}\backslash\mathrm{left}(\mathrm{d} \backslash\mathrm{Gamma}_{\{\alpha\}}\{ \}^{\{\beta\}}-\backslash\mathrm{Gamma}_{\{\alpha\}}\{ \}^{\{\gamma\}} \backslash\mathrm{wedge} \backslash\mathrm{Gamma}_{\{\gamma\}}\{ \}^{\{\beta\}} \backslash\mathrm{right}) \backslash\mathrm{sigma}^{\{\alpha\}}\{ \}_{\{\beta\}} \backslash\backslash \&=\frac{i}{4} \mathrm{R}^{\{\alpha\} \backslash\beta\} \backslash\mathrm{sigma}_{\{\alpha\} \backslash\beta\}=\frac{i}{8} \mathrm{R}_{ij}^{\alpha\beta} \sigma_{\alpha\beta} \mathrm{d} x^i \wedge \mathrm{d} x^j, \backslash\mathrm{end}\{\mathrm{aligned}\}$	$\Omega^g:=d\Gamma+\Gamma\wedge\Gamma=\frac{i}{4}\left(d\Gamma_{\alpha}{}^{\beta}-\Gamma_{\alpha}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\beta}\right)\sigma^{\alpha}{}_{\beta}=\frac{i}{4}R^{\alpha\beta}\sigma_{\alpha\beta}=\frac{i}{8}R_{ij}{}^{\alpha\beta}\sigma_{\alpha\beta}dx^i\wedge dx^j,$	$\Omega^g:=d\Gamma+\Gamma\wedge\Gamma=\frac{i}{4}\left(d\Gamma_{\alpha}{}^{\beta}-\Gamma_{\alpha}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\beta}\right)\sigma^{\alpha}{}_{\beta}=\frac{i}{4}R^{\alpha\beta}\sigma_{\alpha\beta}=\frac{i}{8}R_{ij}{}^{\alpha\beta}\sigma_{\alpha\beta}dx^i\wedge dx^j,$
mielke-geometrodynamics_277 EQ0854 (12.4.1)	277	■ 0.931	$\backslash\mathrm{stackrel}\{(\pm)\}\{V\}_{\{\mathrm{EC}\}}:=V_{\{\mathrm{EC}\}} \backslash\mathrm{p m} \mathrm{i} \mathrm{d} \mathrm{C}_{\{\mathrm{TT}\}}=\backslash\mathrm{p m} \backslash\mathrm{frac}\{1\}\{2\} \ell^{2*} \backslash\mathrm{operatorname}\{\mathrm{Tr}\}\backslash\mathrm{left}\{\backslash\mathrm{left}(1 \backslash\mathrm{mp} \backslash\mathrm{gamma}_{\{5\}}\backslash\mathrm{right}) \backslash\mathrm{Omega} \backslash\mathrm{wedge} \backslash\mathrm{sigma}\backslash\mathrm{right}\}=-\backslash\mathrm{frac}\{1\}\{2\} \ell^{2*} \backslash\mathrm{stackrel}\{(\pm)\}\{R\}\{ \}^{\{\alpha\} \backslash\beta\} \backslash\mathrm{wedge} \backslash\mathrm{eta}_{\{\alpha\} \backslash\beta\}}+\backslash\mathrm{frac}\{\backslash\mathrm{Lambda}\}\{\ell^{2*}\} \backslash\mathrm{eta}$	$\stackrel{(\pm)}{V}_{\mathrm{EC}}:=V_{\mathrm{EC}}\pm idC_{\mathrm{TT}}=\pm\frac{1}{2\ell^2}\mathrm{Tr}\{(1\mp\gamma_5)\Omega\wedge\sigma\}=-\frac{1}{2\ell^2}\stackrel{(\pm)}{R}{}^{\alpha\beta}\wedge\eta_{\alpha\beta}+\frac{\Lambda}{\ell^2}\eta$	$\stackrel{(\pm)}{V}_{\mathrm{EC}}:=V_{\mathrm{EC}}\pm idC_{\mathrm{TT}}=\pm\frac{1}{2\ell^2}Tr\{(1\mp\gamma_5)\Omega\wedge\sigma\}=-\frac{1}{2\ell^2}\stackrel{(\pm)}{R}{}^{\alpha\beta}\wedge\eta_{\alpha\beta}+\frac{\Lambda}{\ell^2}\eta$
mielke-geometrodynamics_169 EQ0530 (7.7.50)	169	■ 0.931	$\backslash\mathrm{left}.\{ \}^{\{*\}} \mathrm{R}_{\{\alpha\} \backslash\beta\}}^{\{(\star)\}} \backslash\mathrm{equiv}(-1)^{\{\mathrm{sig}\}} \mathrm{R}_{\{\alpha\} \backslash\beta\}}+\mathrm{e}_{[\{\alpha\}}\backslash\mathrm{right}\backslash\mathrm{rfloor} \mathrm{G}_{\{\beta\}}]+\backslash\mathrm{frac}\{1\}\{4\} \mathrm{R} \backslash\mathrm{eta}_{\{\alpha\} \backslash\beta\}}+(-1)^{\{\mathrm{sig}\}} \mathrm{D}_{[\{\alpha\}} \mathrm{T}_{\{\beta\}}]$	${}^*R_{\alpha\beta}^{(\star)}\equiv(-1)^{\mathrm{sig}}R_{\alpha\beta}+e_{[\alpha}\rfloor G_{\beta]}+\frac{1}{4}R\eta_{\alpha\beta}+(-1)^{\mathrm{sig}+1}D_{[\alpha}T_{\beta]}$	${}^*R_{\alpha\beta}^{(\star)}\equiv(-1)^{\mathrm{sig}}R_{\alpha\beta}+e_{[\alpha}\rfloor G_{\beta]}+\frac{1}{4}R\eta_{\alpha\beta}+(-1)^{\mathrm{sig}+1}D_{[\alpha}T_{\beta]}$
mielke-geometrodynamics_180 EQ0556 (8.4.10)	180	■ 0.933	$\mathrm{R}_{\{\alpha\} \backslash\beta\}}=\backslash\mathrm{mp}\{ \}^{\{*\}} \mathrm{R}_{\{\alpha\} \backslash\beta\}}^{\{(\star)\}}$	$R_{\alpha\beta}=\mp{}^*R_{\alpha\beta}^{(\star)}$	$R_{\alpha\beta}=\mp{}^*R_{\alpha\beta}^{(\star)}$
mielke-geometrodynamics_178 EQ0547 (8.4.1)	178	■ 0.935	$\backslash\mathrm{begin}\{\mathrm{aligned}\} \mathrm{L}_{\{\mathrm{Pontr}\}} \&:=\mathrm{d} \mathrm{C}_{\{\mathrm{RR}\}}=\backslash\mathrm{operatorname}\{\mathrm{Tr}\}\backslash\mathrm{left}\{\backslash\mathrm{Omega}^{\{g\}} \backslash\mathrm{wedge} \backslash\mathrm{Omega}^{\{g\}} \backslash\mathrm{right}\}=\backslash\mathrm{frac}\{1\}\{2\} \mathrm{R}_{\{\alpha\}}\{ \}^{\{\beta\}} \backslash\mathrm{wedge} \mathrm{R}_{\{\beta\}}\{ \}^{\{\alpha\}} \backslash\backslash \&=\backslash\mathrm{frac}\{1\}\{2\} \mathrm{d}\backslash\mathrm{left}(\backslash\mathrm{Gamma}_{\{\alpha\}}\{ \}^{\{\beta\}} \backslash\mathrm{wedge} \mathrm{R}_{\{\beta\}}\{ \}^{\{\alpha\}}-\backslash\mathrm{frac}\{1\}\{3\} \backslash\mathrm{Gamma}_{\{\alpha\}}\{ \}^{\{\beta\}} \backslash\mathrm{wedge} \backslash\mathrm{Gamma}_{\{\beta\}}\{ \}^{\{\gamma\}} \backslash\mathrm{wedge} \backslash\mathrm{Gamma}_{\{\gamma\}}\{ \}^{\{\beta\}} \backslash\mathrm{right}) \backslash\mathrm{end}\{\mathrm{aligned}\}$	$L_{\mathrm{Pontr}}:=dC_{\mathrm{RR}}=\mathrm{Tr}\{\Omega^g\wedge\Omega^g\}=\frac{1}{2}R_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}=\frac{1}{2}d\left(\Gamma_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}-\frac{1}{3}\Gamma_{\alpha}{}^{\beta}\wedge\Gamma_{\beta}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\alpha}\right)$	$L_{\mathrm{Pontr}}:=dC_{\mathrm{RR}}=\mathrm{Tr}\{\Omega^g\wedge\Omega^g\}=\frac{1}{2}R_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}=\frac{1}{2}d\left(\Gamma_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}-\frac{1}{3}\Gamma_{\alpha}{}^{\beta}\wedge\Gamma_{\beta}{}^{\gamma}\wedge\Gamma_{\gamma}{}^{\alpha}\right),$
mielke-geometrodynamics_259 EQ0824 (11.4.38)	259	■ 0.935	$\backslash\mathrm{left}.\backslash\mathrm{left}.\mathrm{r}_{\{s\}} \backslash\mathrm{equiv}\backslash\mathrm{angle} \mid \backslash\mathrm{vec}\{x\}\backslash\mathrm{right}\ ^{\{2\}} \backslash\mathrm{right}\backslash\mathrm{angle}_{\{\psi\}}^{\{1 / 2\}}:=\backslash\mathrm{left}(\backslash\mathrm{frac}\{\mathrm{int}_{\{H^{\{3\}}\}} \backslash\mathrm{vec}\{x\} ^{\{2\}} \backslash\mathrm{bar}\{\psi\}^{\{*\}} \backslash\mathrm{gamma} \backslash\psi\}\{\mathrm{int}_{\{H^{\{3\}}\}} \backslash\mathrm{bar}\{\psi\}^{\{*\}} \backslash\mathrm{gamma} \backslash\psi\}\backslash\mathrm{right})^{\{1 / 2\}}.$	$r_s\equiv\langle \vec{x} ^2\rangle_{\psi}^{1/2}:=\left(\frac{\int_{H^3} \vec{x} ^2\bar{\psi}^*\gamma\psi}{\int_{H^3}\bar{\psi}^*\gamma\psi}\right)^{1/2}.$	$r_s\equiv\langle \vec{x} ^2\rangle_{\psi}^{1/2}:=\left(\frac{\int_{H^3} \vec{x} ^2\bar{\psi}^*\gamma\psi}{\int_{H^3}\bar{\psi}^*\gamma\psi}\right)^{1/2}.$
mielke-geometrodynamics_99 EQ0288 (4.4.17)	99	■ 0.936	$\backslash\mathrm{left}[\backslash\mathrm{Omega}^{\{g\}}, \backslash\mathrm{Pi}^{\{\mathrm{L}\}}\backslash\mathrm{right}]+\backslash\mathrm{mathrm}\{\mathrm{E}_{\{[]\}}\}=\mathrm{D} \backslash\mathrm{tau}_{\{s\}}-\backslash\mathrm{Sigma}_{\{[]\}}.$	$[\Omega^g,\Pi^{\mathrm{L}}]+E_{[]} = D\tau_s - \Sigma_{[]}.$	$[\Omega^g,\Pi^{\mathrm{L}}]+E_{[]} = D\tau_s - \Sigma_{[]}.$
mielke-geometrodynamics_196 EQ0598 (9.1.16)	196	■ 0.936	$\mathrm{f}=-\mathrm{h}=\mathrm{g}=\mathrm{k}=-\backslash\mathrm{frac}\{\mu\}\{r^{\{2\}}\} \mathrm{e}^{\{-\backslash\mathrm{nu} / 2\}}.$	$f=-h=g=k=-\frac{\mu}{r^2}e^{-\nu/2}.$	$f=-h=g=k=-\frac{\mu}{r^2}e^{-\nu/2}.$
mielke-geometrodynamics_83 EQ0203 (4.1.17)	83	■ 0.937	$\backslash\mathrm{stackrel}\{*\}\{\backslash\mathrm{sigma}\}^{\{*\}} \backslash\mathrm{vartheta}^{\{\beta\}}=\mathrm{E}_{\{j\}}^{\{\beta\}} \mathrm{d} x^{\{j\}}$	$\stackrel{*}{\sigma}{}^*\vartheta^{\beta}=E_j^{\beta}dx^j$	$\stackrel{*}{\sigma}{}^*\vartheta^{\beta}=E_j^{\beta}dx^j$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_00815 (11.4.29)	257	■ 0.946	$\begin{aligned} & H^{\prime} \simeq \frac{1+\omega}{\beta} F \\ & F^{\prime}+\frac{2}{\rho} F \simeq \frac{1-\omega}{\beta} H \end{aligned}$	$H^{\prime} \simeq \frac{1+\omega}{\beta} F$ $F^{\prime}+\frac{2}{\rho} F \simeq \frac{1-\omega}{\beta} H$	$H^{\prime} \simeq \frac{1+\omega}{\beta} F,$ $F^{\prime}+\frac{2}{\rho} F \simeq \frac{1-\omega}{\beta} H.$
mielke-geometrodynamics_01021 (14.4.16)	328	■ 0.947	$S_{\mathrm{A}}:=4\pi\,r_{+}^2=16\pi\,M_{\mathrm{ir}}^2\frac{\hbar^2}{M^*4c^2},$	$S_{\mathrm{A}}:=4\pi r_{+}^2=16\pi M_{\mathrm{ir}}^2\frac{\hbar^2}{M^*4c^2},$	$S_{\mathrm{A}}:=4\pi r_{+}^2=16\pi\,M_{\mathrm{ir}}^2\frac{\hbar^2}{M^*4c^2},$
mielke-geometrodynamics_00855 (12.4.2)	277	■ 0.947	$\mathrm{\mathscr{H}}_{\Lambda}(\underline{\Psi}(\underline{\Gamma}))=0$	$\mathcal{H}_{\Lambda}\Psi^{(1)}(\underline{\Gamma})=0$	$\mathcal{H}_{\Lambda}\Psi^{(\pm)}(\underline{\Gamma})=0$
mielke-geometrodynamics_00452 (7.2.5)	151	■ 0.948	$C_{\{.\,\,c\,d\}^{\{a\,b\}}}=R_{\{.\,\,c\,d\}^{\{a\,b\}}}-2\,R_{\{.\,[c\,]^{\{[a\,\,\cdot\,]\,\,\delta_{d\,]\}^{\{b\,]\}}+\frac{R}{6}\,\,\delta_{[c\,]^{\{a\,]\,\,\delta_{d\,]\}^{\{b\,]\}}\,}.$	$C_{..cd}^{ab}=R_{..cd}^{ab}-2R_{.[c}^{[a\,\cdot}\delta_{d]}^{b]}+\frac{R}{6}\delta_{[c}^a\delta_{d]}^b.$	$C_{..cd}^{ab}=R_{..cd}^{ab}-2R_{.[c}^{[a\,\cdot}\delta_{d]}^{b]}+\frac{R}{6}\delta_{[c}^a\delta_{d]}^b.$
mielke-geometrodynamics_00061 (2.8.2)	46	■ 0.949	$\{ \}^{*}\backslash\left(\gamma_k\backslash\right)=\frac{(-1)^{k}}{(2\pi i)^k k!}\operatorname{Tr}(\backslash\Omega\wedge\cdots\wedge\backslash\omega\wedge\backslash\Omega)\,.$	$^*(\gamma_k)=\frac{(-1)^k}{(2\pi i)^k k!}\operatorname{Tr}(\Omega\wedge\cdots\wedge\Omega).$	$^*\pi(\gamma_k)=\frac{(-1)^k}{(2\pi i)^k k!}\operatorname{Tr}(\Omega\wedge\cdots\wedge\Omega).$
mielke-geometrodynamics_00756 (11.1.23)	242	■ 0.950	$\begin{aligned} S(p) &= \exp \imath \theta_{\alpha\beta}(m) \Lambda^{-1}(L^{\alpha\beta}) = \exp \imath \theta_{\alpha\beta}(m) \sigma^{\alpha\beta} \\ &= \exp \left(-\frac{1}{4} \theta_{\alpha\beta}(m) \left[\gamma^{\alpha}, \gamma^{\beta} \right] \right). \end{aligned}$	$S(p)=\exp \imath \theta_{\alpha\beta}(m) \Lambda^{-1}\left(L^{\alpha\beta}\right)=\exp \imath \theta_{\alpha\beta}(m) \sigma^{\alpha\beta}$ $=\exp \left(-\frac{1}{4} \theta_{\alpha\beta}(m)\left[\gamma^{\alpha}, \gamma^{\beta}\right]\right).$	$S(p)=\exp \imath \theta_{\alpha\beta}(m) \Lambda^{-1}(L^{\alpha\beta})=\exp \imath \theta_{\alpha\beta}(m) \sigma^{\alpha\beta}$ $=\exp \left(-\frac{1}{4} \theta_{\alpha\beta}(m)\left[\gamma^{\alpha}, \gamma^{\beta}\right]\right).$
mielke-geometrodynamics_00939 (14.1.20)	305	■ 0.950	$\boldsymbol{\xi} \wedge \boldsymbol{\xi} = 0 \quad (= 2 \, \vec{E} \cdot \vec{B} \text{ in 3-dimensional vector notation })$	$\xi \wedge \xi = 0 \quad (= 2 \vec{E} \cdot \vec{B} \text{ in 3-dimensional vector notation })$	$\xi \wedge \xi = 0 \quad (= 2 \, \vec{E} \cdot \vec{B} \text{ in 3-dimensional vector notation })$
mielke-geometrodynamics_00568 (8.6.5)	183	■ 0.950	$L_{\{\theta\}}:=\theta_{\mathrm{T}}dC_{\mathrm{TT}}+\theta_{\mathrm{T}}^*dC_{\mathrm{TT}^*}+\theta_{\mathrm{L}}dC_{\mathrm{RR}}+\theta_{\mathrm{L}}^*(-1)^{\mathrm{sig}+1}dC_{\mathrm{RR}^*}=dC_{\theta}.$	$L_{\theta}:=\theta_{\mathrm{T}}dC_{\mathrm{TT}}+\theta_{\mathrm{T}}^*dC_{\mathrm{TT}^*}+\theta_{\mathrm{L}}dC_{\mathrm{RR}}+\theta_{\mathrm{L}}^*(-1)^{\mathrm{sig}+1}dC_{\mathrm{RR}^*}=dC_{\theta}.$	$L_{\theta}:=\theta_{\mathrm{T}}dC_{\mathrm{TT}}+\theta_{\mathrm{T}}^*dC_{\mathrm{TT}^*}+\theta_{\mathrm{L}}dC_{\mathrm{RR}}+\theta_{\mathrm{L}}^*(-1)^{\mathrm{sig}+1}dC_{\mathrm{RR}^*}=dC_{\theta}.$
mielke-geometrodynamics_00234 (4.2.27)	88	■ 0.950	$\widetilde{D}\widetilde{\varphi}=\widetilde{D}\widetilde{\varphi}=0$	$\widetilde{\widetilde{D}}\widetilde{\widetilde{\varphi}}=D\widetilde{\widetilde{\varphi}}=0$	$\widetilde{\widetilde{D}}\widetilde{\widetilde{\varphi}}=D\widetilde{\widetilde{\varphi}}=0$
mielke-geometrodynamics_00302 (4.5.9)	102	■ 0.950	$\begin{aligned} & \widehat{D}\Sigma_{\alpha} \cong (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \Delta^{\beta} \gamma \\ & \quad - \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta\gamma}) \wedge \sigma^{\beta\gamma} \end{aligned}$	$\widehat{D}\Sigma_{\alpha} \cong (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \Delta^{\beta} \gamma$ $- \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta\gamma}) \wedge \sigma^{\beta\gamma}$	$\widehat{D}\Sigma_{\alpha} \cong (e_{\alpha} \rfloor R_{\beta}^{\gamma}) \wedge \Delta^{\beta} \gamma$ $- \frac{1}{2} (e_{\alpha} \rfloor Q_{\beta\gamma}) \wedge \sigma^{\beta\gamma}$
mielke-geometrodynamics_00151 (3.4.22)	66	■ 0.951	$L_{\{\text{Mills}\}}\backslash\left(\mathscr{F},\{ \}^{*}\backslash\mathscr{F}\right)=\mathscr{F}\backslash\left[1-\backslash\left(1+\frac{1}{b^2}\mathscr{F}^2\right)\backslash\right]^{\{2\}}\backslash\right)^{-1}\backslash\right]$	$L_{\mathrm{Mills}}\left(\mathscr{F},*\mathscr{F}\right)=\mathscr{F}\left[1-\left(1+\frac{1}{b^2}\mathscr{F}^2\right)^{-1}\right]$	$L_{\mathrm{Mills}}(\mathscr{F},*\mathscr{F})=\mathscr{F}\left[1-\left(1+\frac{1}{b^2}\mathscr{F}^2\right)^{-1}\right]$
mielke-geometrodynamics_00560 (8.5.3)	181	■ 0.953	$\stackrel{\pm}{\Omega}^g=i\frac{\theta_{\mathrm{T}}^*}{4\ell^2}\sigma$	$\stackrel{\pm}{\Omega}^g=i\frac{\theta_{\mathrm{T}}^*}{4\ell^2}\sigma$	$\stackrel{\pm}{\Omega}^g=i\frac{\theta_{\mathrm{T}}^*}{4\ell^2}\sigma$
mielke-geometrodynamics_01106 (B.18)	370	■ 0.953	$\delta\alpha^{\{p\}}:=(-1)^{pn+n+1+s*}d^*\alpha^{\{p\}}.$	$\delta\alpha^{(p)}:=(-1)^{pn+n+1+s*}d^*\alpha^{(p)}.$	$\delta\alpha^{(p)}:=(-1)^{pn+n+1+s*}d^*\alpha^{(p)}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_248 EQ0786 (11.3.5)	248	■ 0.954	$\begin{aligned} L_{\mathrm{NL}} &= -3 \ell^{*2} \tau_s \wedge^* \tau_s = 3 \ell^{*2} \bar{\psi}^* \gamma \psi \wedge^* (\bar{\psi}^* \gamma \psi) \\ &= 3 \ell^{*2} \bar{\psi} \gamma^{n+1} \gamma \psi \wedge^* (\bar{\psi} \gamma^{n+1} \gamma \psi) . \end{aligned}$	$L_{\mathrm{NL}} = -3 \ell^{*2} \tau_s \wedge^* \tau_s = 3 \ell^{*2} \bar{\psi}^* \gamma \psi \wedge^* (\bar{\psi}^* \gamma \psi) \\ = 3 \ell^{*2} \bar{\psi} \gamma^{n+1} \gamma \psi \wedge^* (\bar{\psi} \gamma^{n+1} \gamma \psi) .$	$L_{\mathrm{NL}} = -3 \ell^{*2} \tau_s \wedge^* \tau_s = 3 \ell^{*2} \bar{\psi}^* \gamma \psi \wedge^* (\bar{\psi}^* \gamma \psi) \\ = 3 \ell^{*2} \bar{\psi} \gamma^{n+1} \gamma \psi \wedge^* (\bar{\psi} \gamma^{n+1} \gamma \psi) .$
mielke-geometrodynamics_231 EQ0727 (10.10.7)	231	■ 0.955	$\begin{aligned} & i \gamma^\alpha \Psi \wedge \frac{\delta L}{\delta \vartheta^\alpha} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} - 2 D \frac{\delta L}{\delta \bar{\Psi}} \\ & \cong i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\mathrm{TL}}}{\ell} R_\alpha^* + \frac{\theta_{\mathrm{T}}}{\ell^2} T_\alpha + \Sigma_\alpha \right) \\ & + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - D \left(i D \Psi + \frac{i}{2} m \gamma \wedge \Psi \right) \\ & = i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\mathrm{TL}}}{\ell} R_\alpha^* + \frac{\theta_{\mathrm{T}}}{\ell^2} T_\alpha \right) + \gamma^\alpha \Psi \wedge \left(\frac{1}{4} m \bar{\Psi} \gamma_\alpha \Psi \right) \\ & + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - i R_\alpha^* \gamma^\alpha \Psi - \frac{i}{2} m T_\alpha \gamma^\alpha \Psi \\ & + \frac{i}{2} m \gamma \wedge D \Psi . \end{aligned}$	$i \gamma^\alpha \Psi \wedge \frac{\delta L}{\delta \vartheta^\alpha} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} - 2 D \frac{\delta L}{\delta \bar{\Psi}} \\ \cong i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\mathrm{TL}}}{\ell} R_\alpha^* + \frac{\theta_{\mathrm{T}}}{\ell^2} T_\alpha + \Sigma_\alpha \right) \\ + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - D \left(i D \Psi + \frac{i}{2} m \gamma \wedge \Psi \right) \\ = i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\mathrm{TL}}}{\ell} R_\alpha^* + \frac{\theta_{\mathrm{T}}}{\ell^2} T_\alpha \right) + \gamma^\alpha \Psi \wedge \left(\frac{1}{4} m \bar{\Psi} \gamma_\alpha \Psi \right) \\ + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - i R_\alpha^* \gamma^\alpha \Psi - \frac{i}{2} m T_\alpha \gamma^\alpha \Psi \\ + \frac{i}{2} m \gamma \wedge D \Psi .$	$i \gamma^\alpha \Psi \wedge \frac{\delta L}{\delta \vartheta^\alpha} + c \gamma \wedge \frac{\delta L}{\delta \bar{\Psi}} - 2 D \frac{\delta L}{\delta \bar{\Psi}} \\ \cong i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\mathrm{TL}}}{\ell} R_\alpha^* + \frac{\theta_{\mathrm{T}}}{\ell^2} T_\alpha + \Sigma_\alpha \right) \\ + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - D \left(i D \Psi + \frac{i}{2} m \gamma \wedge \Psi \right) \\ = i \gamma^\alpha \Psi \wedge \left(\frac{\theta_{\mathrm{TL}}}{\ell} R_\alpha^* + \frac{\theta_{\mathrm{T}}}{\ell^2} T_\alpha \right) + \gamma^\alpha \Psi \wedge \left(\frac{1}{4} m \bar{\Psi} \gamma_\alpha \Psi \right) \\ + c \gamma \wedge \left(\frac{i}{2} D \Psi + \frac{i}{4} m \gamma \wedge \Psi \right) - i R_\alpha^* \gamma^\alpha \Psi - \frac{i}{2} m T_\alpha \gamma^\alpha \Psi \\ + \frac{i}{2} m \gamma \wedge D \Psi .$
mielke-geometrodynamics_244 EQ0768 (11.1.35)	244	■ 0.956	$\bar{\psi} \longrightarrow{ }^{\wedge \{S^{-1}\}} \bar{\psi} = \bar{\psi} S ,$	$\bar{\psi} \longrightarrow S^{-1} \bar{\psi} = \bar{\psi} S ,$	$\bar{\psi} \longrightarrow^{S^{-1}} \bar{\psi} = \bar{\psi} S ,$
mielke-geometrodynamics_315 EQ0973 (14.2.14)	315	■ 0.956	$\begin{aligned} \hat{\Theta} &:= \hat{D} \hat{\vartheta} = \hat{d} \hat{\vartheta} - [\hat{\omega}, \hat{\vartheta}] \\ &= \left[\begin{array}{c c} \Theta - \hat{\ell} \omega \wedge \alpha^{\mathrm{T}} & \hat{\ell} \left(d \omega - \omega^g \wedge \omega - \omega \wedge \overset{\circ}{\omega} \right) + \vartheta \wedge \alpha + \alpha \wedge \overset{\circ}{\vartheta} \\ - \alpha^{\mathrm{T}} \wedge \vartheta - \vartheta \wedge \alpha^{\mathrm{T}} & \overset{\circ}{\Theta} - \hat{\ell} \alpha^{\mathrm{T}} \wedge \omega \end{array} \right] . \end{aligned}$	$\hat{\Theta} := \hat{D} \hat{\vartheta} = \hat{d} \hat{\vartheta} - [\hat{\omega}, \hat{\vartheta}] \\ = \left[\begin{array}{c c} \Theta - \hat{\ell} \omega \wedge \alpha^{\mathrm{T}} & \hat{\ell} \left(d \omega - \omega^g \wedge \omega - \omega \wedge \overset{\circ}{\omega} \right) + \vartheta \wedge \alpha + \alpha \wedge \overset{\circ}{\vartheta} \\ - \alpha^{\mathrm{T}} \wedge \vartheta - \vartheta \wedge \alpha^{\mathrm{T}} & \overset{\circ}{\Theta} - \hat{\ell} \alpha^{\mathrm{T}} \wedge \omega \end{array} \right] .$	$\hat{\Theta} := \hat{D} \hat{\vartheta} = \hat{d} \hat{\vartheta} - [\hat{\omega}, \hat{\vartheta}] \\ = \left[\begin{array}{c c} \Theta - \hat{\ell} \omega \wedge \alpha^{\mathrm{T}} & \hat{\ell} (d \omega - \omega^g \wedge \omega - \omega \wedge \overset{\circ}{\omega}) + \vartheta \wedge \alpha + \alpha \wedge \overset{\circ}{\vartheta} \\ - \alpha^{\mathrm{T}} \wedge \vartheta - \vartheta \wedge \alpha^{\mathrm{T}} & \overset{\circ}{\Theta} - \hat{\ell} \alpha^{\mathrm{T}} \wedge \omega \end{array} \right] .$
mielke-geometrodynamics_243 EQ0761 (11.1.28)	243	■ 0.957	$\frac{1}{N} \operatorname{Tr} \left(\gamma \otimes_{\mathrm{s}} \gamma \right) = d s^2 = g_{ij}(m) d x^i \otimes_{\mathrm{s}} d x^j .$	$\frac{1}{N} \operatorname{Tr} (\gamma \otimes_{\mathrm{s}} \gamma) = d s^2 = g_{ij}(m) d x^i \otimes_{\mathrm{s}} d x^j .$	$\frac{1}{N} \operatorname{Tr} (\gamma \otimes_{\mathrm{s}} \gamma) = d s^2 = g_{ij}(m) d x^i \otimes_{\mathrm{s}} d x^j .$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_160 EQ0486 (7.7.2)	160	■ 0.957	$\begin{aligned} & \vartheta^\alpha \rightarrow \widetilde{\vartheta}^\alpha := \vartheta^\alpha + \ell^2 e_\beta \Big]{}^* H^{\alpha\beta}, \\ & \Gamma_\alpha{}^\beta \rightarrow \widetilde{\Gamma}_\alpha{}^\beta := \Gamma_\alpha{}^\beta + \ell^2 e_\alpha \Big]{}^* H^\beta. \end{aligned}$	$\vartheta^\alpha \rightarrow \widetilde{\vartheta}^\alpha := \vartheta^\alpha + \ell^2 e_\beta \Big]{}^* H^{\alpha\beta},$ $\Gamma_\alpha{}^\beta \rightarrow \widetilde{\Gamma}_\alpha{}^\beta := \Gamma_\alpha{}^\beta + \ell^2 e_\alpha \Big]{}^* H^\beta.$	$\vartheta^\alpha \rightarrow \widetilde{\vartheta}^\alpha := \vartheta^\alpha + \ell^2 e_\beta \Big]{}^* H^{\alpha\beta},$ $\Gamma_\alpha{}^\beta \rightarrow \widetilde{\Gamma}_\alpha{}^\beta := \Gamma_\alpha{}^\beta + \ell^2 e_\alpha \Big]{}^* H^\beta.$
mielke-geometrodynamics_370 E01104 (B.16)	370	■ 0.957	$\begin{aligned} & \left\langle \alpha^{(p)}, \beta^{(p)} \right\rangle := \alpha^{(p)} \wedge^* \beta^{(p)} = \beta^{(p)} \wedge^* \alpha^{(p)} \\ & = (-1)^{s*} (\alpha^{(p)} \wedge^* \beta^{(p)}) \eta \\ & = \frac{(-1)^s}{p!} A^{\alpha_1 \cdots \alpha_p} B_{\alpha_1 \cdots \alpha_p} d\mu(M) \in C^\infty(\wedge^n T^*(M)) \end{aligned}$	$\left\langle \alpha^{(p)}, \beta^{(p)} \right\rangle := \alpha^{(p)} \wedge^* \beta^{(p)} = \beta^{(p)} \wedge^* \alpha^{(p)}$ $= (-1)^{s*} (\alpha^{(p)} \wedge^* \beta^{(p)}) \eta$ $= \frac{(-1)^s}{p!} A^{\alpha_1 \cdots \alpha_p} B_{\alpha_1 \cdots \alpha_p} d\mu(M) \in C^\infty(\wedge^n T^*(M))$	$\langle \alpha^{(p)}, \beta^{(p)} \rangle := \alpha^{(p)} \wedge^* \beta^{(p)} = \beta^{(p)} \wedge^* \alpha^{(p)}$ $= (-1)^{s*} (\alpha^{(p)} \wedge^* \beta^{(p)}) \eta$ $= \frac{(-1)^s}{p!} A^{\alpha_1 \cdots \alpha_p} B_{\alpha_1 \cdots \alpha_p} d\mu(M) \in C^\infty(\wedge^n T^*(M))$
mielke-geometrodynamics_176 E00537 (8.2.4)	176	■ 0.958	$\begin{array}{l} s\,\bar{C}=\mathbb{B},\quad s\,\mathbb{B}=0, \\ s\,\bar{\chi}=\beta,\quad s\,\beta=0, \\ s\,\bar{\Phi}=\bar{\eta},\quad s\,\bar{\eta}=0. \end{array}$	$s\bar{C}=\mathbb{B},\quad s\mathbb{B}=0,$ $s\bar{\chi}=\beta,\quad s\beta=0,$ $s\bar{\Phi}=\bar{\eta},\quad s\bar{\eta}=0.$	$s\bar{C}=\mathbb{B},\quad s\mathbb{B}=0,$ $s\bar{\chi}=\beta,\quad s\beta=0,$ $s\bar{\Phi}=\bar{\eta},\quad s\bar{\eta}=0.$
mielke-geometrodynamics_95 E00269 (4.3.34)	95	■ 0.958	$R_{\{i\}j}:=R_{[\alpha}i\}j]^{\cdots\alpha}$	$R_{ij}:=R_{\alpha ij}^{\cdots\alpha}$	$R_{ij}:=R_{\alpha ij}^{\cdots\alpha}$
mielke-geometrodynamics_132 EQ0394 (6.4.9)	132	■ 0.958	$\left.\mathscr{H}\right.\cong n^{\alpha}\underline{\Sigma}_{\alpha}+\Gamma_{\perp}^{\alpha\beta}\rho\left(I_{\alpha\beta}\right)\underline{\Psi}\wedge\frac{\partial L}{\partial D\Psi}-\underline{d}\left[\left(n\right]\Psi\right)\wedge\frac{\partial L}{\partial D\Psi}\Big].$	$\mathcal{H}\cong n^{\alpha}\underline{\Sigma}_{\alpha}+\Gamma_{\perp}^{\alpha\beta}\rho\left(I_{\alpha\beta}\right)\underline{\Psi}\wedge\frac{\partial L}{\partial D\Psi}-\underline{d}\left[\left(n\right]\Psi\right)\wedge\frac{\partial L}{\partial D\Psi}\Big].$	$\mathcal{H}\cong n^{\alpha}\underline{\Sigma}_{\alpha}+\Gamma_{\perp}^{\alpha\beta}\rho\left(I_{\alpha\beta}\right)\underline{\Psi}\wedge\frac{\partial L}{\partial D\Psi}-\underline{d}\left[\left(n\right]\Psi\right)\wedge\frac{\partial L}{\partial D\Psi}\Big].$
mielke-geometrodynamics_135 EQ0406 (6.4.26)	135	■ 0.959	$\stackrel{\scriptscriptstyle\pm)}{\underline{D}}\stackrel{\scriptscriptstyle\pm)}{\underline{T}}^B=0$	$\stackrel{(\pm)}{\underline{D}}\stackrel{(\pm)}{\underline{T}}^B=0$	$\stackrel{(\pm)}{\underline{D}}\stackrel{(\pm)}{\underline{T}}^B=0$
mielke-geometrodynamics_152 EQ0458 (7.2.10)	152	■ 0.959	$\mathfrak{s}\mathfrak{o}(4)\approx\mathfrak{s}\mathfrak{o}(3)\times\mathfrak{s}\mathfrak{o}(3)\approx\mathfrak{s}\mathfrak{u}(2)\times\mathfrak{s}\mathfrak{u}(2).$	$\mathfrak{so}(4)\approx\mathfrak{so}(3)\times\mathfrak{so}(3)\approx\mathfrak{su}(2)\times\mathfrak{su}(2).$	$\mathfrak{so}(4)\approx\mathfrak{so}(3)\times\mathfrak{so}(3)\approx\mathfrak{su}(2)\times\mathfrak{su}(2).$
mielke-geometrodynamics_308 EQ0954 (14.1.36)	308	■ 0.959	$\begin{aligned} & \left\langle \operatorname{Tr}_{\{ \} \{ \} }^* \xi \wedge \vartheta \wedge \xi - \xi \wedge \vartheta \wedge^* \xi \right\rangle \wedge d\delta \\ & = \operatorname{Tr}_{\{ \} \{ \} }^* \xi \wedge \vartheta \wedge D^* \xi + \xi \wedge \vartheta \wedge D\xi \\ & = -\frac{1}{2} D \operatorname{Tr}_{\{ \} \{ \} }^* \xi \wedge \vartheta \wedge^* \xi - \xi \wedge \vartheta \wedge \xi \Big\} . \end{aligned}$	$(-1)^z \operatorname{Tr}_{\{ \} \{ \} }^* \xi \wedge \vartheta \wedge \xi - \xi \wedge \vartheta \wedge^* \xi \Big\} \wedge d\delta$ $= \operatorname{Tr}_{\{ \} \{ \} }^* \xi \wedge \vartheta \wedge D^* \xi + \xi \wedge \vartheta \wedge D\xi$ $= -\frac{1}{2} D \operatorname{Tr}_{\{ \} \{ \} }^* \xi \wedge \vartheta \wedge^* \xi - \xi \wedge \vartheta \wedge \xi \Big\} .$	$(-1)^z \operatorname{Tr}_{\{ \} \{ \} }^* \xi \wedge \vartheta \wedge \xi - \xi \wedge \vartheta \wedge^* \xi \Big\} \wedge d\delta$ $= \operatorname{Tr}_{\{ \} \{ \} }^* \xi \wedge \vartheta \wedge D^* \xi + \xi \wedge \vartheta \wedge D\xi$ $= -\frac{1}{2} D \operatorname{Tr}_{\{ \} \{ \} }^* \xi \wedge \vartheta \wedge^* \xi - \xi \wedge \vartheta \wedge \xi \Big\} .$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_112 E08338 (5.3.7)	112	■ 0.960	$G:=G_{\alpha}\gamma^{\alpha}\wedge\gamma^{\alpha}:=\frac{1}{2}R^{\mu\nu}\wedge\eta_{\mu\nu\lambda}\gamma^{\lambda}=-i\gamma_5\left(\Omega^g\wedge\gamma+\gamma\wedge\Omega^g\right)=i\left[\gamma,\gamma_5\Omega^g\right].$	$G:=G_{\alpha}\gamma^{\alpha}:=\frac{1}{2}R^{\mu\nu}\wedge\eta_{\mu\nu\lambda}\gamma^{\lambda}=-i\gamma_5\left(\Omega^g\wedge\gamma+\gamma\wedge\Omega^g\right)=i\left[\gamma,\gamma_5\Omega^g\right].$	$G:=G_{\alpha}\gamma^{\alpha}:=\frac{1}{2}R^{\mu\nu}\wedge\eta_{\mu\nu\lambda}\gamma^{\lambda}=-i\gamma_5(\Omega^g\wedge\gamma+\gamma\wedge\Omega^g)=i[\gamma,\gamma_5\Omega^g].$
mielke-geometrodynamics_303 E0926 (14.1.7)	303	■ 0.960	$D\left(\vartheta\wedge\Omega^g\right)\equiv\Theta\wedge\Omega^g.$	$D\left(\vartheta\wedge\Omega^g\right)\equiv\Theta\wedge\Omega^g.$	$D(\vartheta\wedge\Omega^g)\equiv\Theta\wedge\Omega^g.$
mielke-geometrodynamics_92 E08248 (4.3.13)	92	■ 0.961	$D\phi^{(\circ)}=\vartheta^{\alpha}\nabla_{\alpha}\phi^{(\circ)}.$	$D\phi^{(\circ)}=\vartheta^{\alpha}\nabla_{\alpha}\phi^{(\circ)}.$	$D\phi^{(\circ)}=\vartheta^{\alpha}\nabla_{\alpha}\phi^{(\circ)}.$
mielke-geometrodynamics_98 E08282 (4.4.11)	98	■ 0.961	$\begin{aligned} d\Theta&=dd\vartheta-d[\omega^L,\vartheta] \\ &=-[d\omega^L,\vartheta]+[\omega^L,d\vartheta] \\ &=-[\Omega^L,\vartheta]-[\omega^L\wedge\omega^L,\vartheta]+[\omega^L,\Theta]+[\omega^L,[\omega^L,\vartheta]] \\ &=[\omega^L,\Theta]+[\vartheta,\Omega^L] \end{aligned}$	$\begin{aligned} d\Theta&=dd\vartheta-d[\omega^L,\vartheta] \\ &=-[d\omega^L,\vartheta]+[\omega^L,d\vartheta] \\ &=-[\Omega^L,\vartheta]-[\omega^L\wedge\omega^L,\vartheta]+[\omega^L,\Theta]+[\omega^L,[\omega^L,\vartheta]] \\ &=[\omega^L,\Theta]+[\vartheta,\Omega^L] \end{aligned}$	$\begin{aligned} d\Theta&=dd\vartheta-d[\omega^L,\vartheta] \\ &=-[d\omega^L,\vartheta]+[\omega^L,d\vartheta] \\ &=-[\Omega^L,\vartheta]-[\omega^L\wedge\omega^L,\vartheta]+[\omega^L,\Theta]+[\omega^L,[\omega^L,\vartheta]] \\ &=[\omega^L,\Theta]+[\vartheta,\Omega^L]. \end{aligned}$
mielke-geometrodynamics_34 E09016 (2.2.5)	34	■ 0.962	$T_{\{p\}^q}(M):=\bigcup_{m\in M}T_m^*(M)\otimes^qT_m(M).$	$T_p^q(M):=\bigcup_{m\in M}\otimes^pT_m^*(M)\otimes^qT_m(M).$	$T_p^q(M):=\bigcup_{m\in M}\otimes^pT_m^*(M)\otimes^qT_m(M).$
mielke-geometrodynamics_115 E08348 (5.3.17)	115	■ 0.962	$\begin{aligned} G_{\alpha}&:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma} \\ &=G_{\alpha}^{\{}}+\frac{(-1)^s}{12}\left(e_{\alpha}\rfloor\mathcal{A}\wedge{}^*\mathcal{A}-\frac{1}{3}\mathcal{A}\wedge e_{\alpha}\rfloor{}^*\mathcal{A}\right)+\frac{(-1)^s}{6}\vartheta_{\alpha}\wedge d\mathcal{A} \end{aligned}$	$\begin{aligned} G_{\alpha}&:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma} \\ &=G_{\alpha}^{\{}}+\frac{(-1)^s}{12}\left(e_{\alpha}\rfloor\mathcal{A}\wedge{}^*\mathcal{A}-\frac{1}{3}\mathcal{A}\wedge e_{\alpha}\rfloor{}^*\mathcal{A}\right)+\frac{(-1)^s}{6}\vartheta_{\alpha}\wedge d\mathcal{A} \end{aligned}$	$\begin{aligned} G_{\alpha}&:=\frac{1}{2}R^{\beta\gamma}\wedge\eta_{\alpha\beta\gamma} \\ &=G_{\alpha}^{\{}}+\frac{(-1)^s}{12}\left(e_{\alpha}\rfloor\mathcal{A}\wedge{}^*\mathcal{A}-\frac{1}{3}\mathcal{A}\wedge e_{\alpha}\rfloor{}^*\mathcal{A}\right)+\frac{(-1)^s}{6}\vartheta_{\alpha}\wedge d\mathcal{A} \end{aligned} \tag{5.3}$
mielke-geometrodynamics_140 E08424 (6.7.6)	140	■ 0.962	$\begin{aligned} \frac{1}{2\kappa}T^{\gamma}\wedge\eta_{\alpha\beta\gamma}-\frac{D\theta_{\mathrm{T}}}{2\ell^2}\wedge\vartheta_{\alpha}\wedge\vartheta_{\beta}+D\theta_{\mathrm{L}}\wedge R_{\alpha\beta}&=\vartheta_{[\alpha}\wedge\mu_{\beta]} \\ &=\frac{1}{4}\vartheta_{\alpha}\wedge\vartheta_{\beta}\wedge{}^*j_5. \end{aligned}$	$\begin{aligned} \frac{1}{2\kappa}T^{\gamma}\wedge\eta_{\alpha\beta\gamma}-\frac{D\theta_{\mathrm{T}}}{2\ell^2}\wedge\vartheta_{\alpha}\wedge\vartheta_{\beta}+D\theta_{\mathrm{L}}\wedge R_{\alpha\beta}&=\vartheta_{[\alpha}\wedge\mu_{\beta]} \\ &=\frac{1}{4}\vartheta_{\alpha}\wedge\vartheta_{\beta}\wedge{}^*j_5. \end{aligned}$	$\begin{aligned} \frac{1}{2\kappa}T^{\gamma}\wedge\eta_{\alpha\beta\gamma}-\frac{D\theta_{\mathrm{T}}}{2\ell^2}\wedge\vartheta_{\alpha}\wedge\vartheta_{\beta}+D\theta_{\mathrm{L}}\wedge R_{\alpha\beta}&=\vartheta_{[\alpha}\wedge\mu_{\beta]} \\ &=\frac{1}{4}\vartheta_{\alpha}\wedge\vartheta_{\beta}\wedge{}^*j_5. \end{aligned}$
mielke-geometrodynamics_255 E08889 (11.4.21)	255	■ 0.962	$i\gamma^0\vec{\gamma}\cdot\vec{\partial}\left[\begin{array}{cc} iH(\rho) & \chi_{\kappa}^m \\ -F(\rho) & \chi_{-\kappa}^m \end{array}\right]=\frac{\vec{\sigma}\cdot\vec{x}}{ \vec{x} }\left(\partial_r-\frac{\vec{\sigma}\cdot\vec{L}}{r}\right)\left[\begin{array}{cc} iF(\rho) & \chi_{-\kappa}^m \\ H(\rho) & \chi_{\kappa}^m \end{array}\right].$	$i\gamma^0\vec{\gamma}\cdot\vec{\partial}\left[\begin{array}{cc} iH(\rho) & \chi_{\kappa}^m \\ -F(\rho) & \chi_{-\kappa}^m \end{array}\right]=\frac{\vec{\sigma}\cdot\vec{x}}{ \vec{x} }\left(\partial_r-\frac{\vec{\sigma}\cdot\vec{L}}{r}\right)\left[\begin{array}{cc} iF(\rho) & \chi_{-\kappa}^m \\ H(\rho) & \chi_{\kappa}^m \end{array}\right].$	$i\gamma^0\vec{\gamma}\cdot\vec{\partial}\left[\begin{array}{cc} iH(\rho) & \chi_{\kappa}^m \\ -F(\rho) & \chi_{-\kappa}^m \end{array}\right]=\frac{\vec{\sigma}\cdot\vec{x}}{ \vec{x} }\left(\partial_r-\frac{\vec{\sigma}\cdot\vec{L}}{r}\right)\left[\begin{array}{cc} iF(\rho) & \chi_{-\kappa}^m \\ H(\rho) & \chi_{\kappa}^m \end{array}\right].$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00475 (7.4.3)	156	■ 0.963	$\{ \}^{\{ * \}} R_{\{\alpha \ \beta \}}^{\{ (\star) \}} = - C_{\{\alpha \ \beta \}} + \frac{1}{2} R \ i^{\{\tau \} \{ \}}_{\{\alpha \ \beta \}} + \frac{R}{12} \vartheta_{\alpha} \wedge \vartheta_{\beta} .$	${}^*R_{\alpha\beta}^{(*)} = -C_{\alpha\beta} + \frac{1}{2} Ri^{\tau}_{\alpha\beta} + \frac{R}{12} \vartheta_{\alpha} \wedge \vartheta_{\beta}.$	${}^*R_{\alpha\beta}^{(*)} = -C_{\alpha\beta} + \frac{1}{2} Ri\check{z}_{\alpha\beta} + \frac{R}{12} \vartheta_{\alpha} \wedge \vartheta_{\beta}.$
mielke-geometrodynamics_E00485 (7.7.1)	159	■ 0.963	$g_{\{i \ j\}} \rightarrow \widetilde{g}_{\{i \ j\}} = g_{\{i \ j\}} + a \operatorname{Ric}_{\{i \ j\}} + b \ g_{\{i \ j\}} \operatorname{Ric}_{\{k\}}^{\{k\}}$	$g_{ij} \rightarrow \widetilde{g}_{ij} = g_{ij} + a \operatorname{Ric}_{ij} + b g_{ij} \operatorname{Ric}_k^k$	$g_{ij} \rightarrow \widetilde{g}_{ij} = g_{ij} + a \operatorname{Ric}_{ij} + b g_{ij} \operatorname{Ric}_k^k$
mielke-geometrodynamics_E01143 (C.11)	376	■ 0.963	$(A, \ B) := \operatorname{Tr}(\operatorname{Ad} \ A \operatorname{Ad} \ B) = c_{\iota k} c_{s i}^k A^{\iota} B^s =: \overset{\circ}{g}_{\iota s} A^{\iota} B^s$	$(A, B) := \operatorname{Tr}(\operatorname{Ad} A \operatorname{Ad} B) = c_{\iota k}^i c_{s i}^k A^{\iota} B^s =: \overset{\circ}{g}_{\iota s} A^{\iota} B^s$	$(A, B) := \operatorname{Tr}(\operatorname{Ad} A \operatorname{Ad} B) = c_{\iota k}^i c_{s i}^k A^{\iota} B^s =: \overset{\circ}{g}_{\iota s} A^{\iota} B^s$
mielke-geometrodynamics_E01122 (B.34)	372	■ 0.963	$H^{\{p\}} \backslash \left(S^{\{n\}} \backslash \right) = 0 \quad \text{for} \quad 1 \leq p \leq n-1 \ .$	$H^p(S^n) = 0 \quad \text{for} \quad 1 \leq p \leq n-1.$	$H^p(S^n) = 0 \quad \text{for} \quad 1 \leq p \leq n-1.$
mielke-geometrodynamics_E00791 (11.4.3)	252	■ 0.964	$\backslash \left\{ i \ \gamma^{\{k\}} \ \nabla_{\{k\}}^{\{ (t) \}} + \frac{3}{8} \backslash \operatorname{varepsilon}_{\{8\}} \ \ell^{\{2\}} \backslash \left(\backslash \operatorname{bar} \{ \psi \} \ \psi - a \ \operatorname{bar} \{ \psi \} \ \gamma^{\{5\}} \ \psi \ \gamma^{\{5\}} \right) - m \right\} \ \psi = 0$	$\left\{ i \gamma^k \nabla_k^{(t)} + \frac{3\varepsilon}{8} \ell^2 \left(\bar{\psi} \psi - a \bar{\psi} \gamma^5 \psi \gamma^5 \right) - m \right\} \psi = 0$	$\left\{ i \gamma^k \nabla_k^{(l)} + \frac{3\varepsilon}{8} \ell^2 (\bar{\psi} \psi - a \bar{\psi} \gamma^5 \psi \gamma^5) - m \right\} \psi = 0$
mielke-geometrodynamics_E00575 (8.6.13)	185	■ 0.964	$\backslash \left. \backslash \left. R := e_{\{\beta \}} \right\rfloor e_{\{\alpha \}} \right\rfloor R^{\{\alpha \ \beta \}} = \frac{6 \left(\theta_{\mathrm{T}}^{\star} - a_0 \right)}{\ell^2 \left(\theta_{\mathrm{L}}^{\star} - b_6 \right)}$	$R := e_{\beta} \rfloor e_{\alpha} \rfloor R^{\alpha\beta} = \frac{6 \left(\theta_{\mathrm{T}}^{\star} - a_0 \right)}{\ell^2 \left(\theta_{\mathrm{L}}^{\star} - b_6 \right)}$	$R := e_{\beta} \rfloor e_{\alpha} \rfloor R^{\alpha\beta} = \frac{6(\theta_{\mathrm{T}}^{\star} - a_0)}{\ell^2(\theta_{\mathrm{L}}^{\star} - b_6)}$
mielke-geometrodynamics_E00996 (14.3.9)	321	■ 0.965	$i \ \widehat{D} \ \widehat{\psi} := i \ \widehat{\gamma} \wedge^* \widehat{D} \widehat{\psi} = i \gamma \wedge^* D \widehat{\psi} + i \ \mathcal{D}^{(int)} \widehat{\psi} = 0,$	$i \widehat{D} \widehat{\psi} := i \widehat{\gamma} \wedge^* \widehat{D} \widehat{\psi} = i \gamma \wedge^* D \widehat{\psi} + i \ \mathcal{D}^{(int)} \widehat{\psi} = 0,$	$i \widehat{\mathcal{D}} \widehat{\psi} := i \widehat{\gamma} \wedge^* \widehat{D} \widehat{\psi} = i \gamma \wedge^* D \widehat{\psi} + i \mathcal{D}^{(int)} \widehat{\psi} = 0,$
mielke-geometrodynamics_E00131 (3.4.2)	62	■ 0.966	$\{ \}^{\{ * \}} \ \operatorname{mathscr{F}} := \frac{1}{\alpha_g} \operatorname{Tr} \{ \{ \}^{\{ * \}} (\backslash 0 \omega \ \wedge \ \backslash 0 \omega) \ \right\} = \frac{1}{4 \alpha_g} F_{\alpha \beta}^{\{ \}} F^{\alpha \beta}_{\{ \}}^{\{ * \}} \ .$	${}^*\mathcal{F} := \frac{1}{\alpha_g} \operatorname{Tr} \{ {}^*(\Omega \wedge \Omega) \} = \frac{1}{4\alpha_g} F_{\alpha\beta}{}^j {}^*F^{\alpha\beta}{}_j.$	${}^*\mathcal{F} := \frac{1}{\alpha_g} \operatorname{Tr} \{ {}^*(\varOmega \wedge \varOmega) \} = \frac{1}{4\alpha_g} F_{\alpha\beta}{}^j {}^*F^{\alpha\beta}{}_j.$
mielke-geometrodynamics_E01089 (A.16)	366	■ 0.967	$\begin{aligned} & \ \backslash \left(\widehat{\gamma}^{\{A\}} \right)^{\{ \} \operatorname{Tr} \{ \}} := \backslash \left(\widehat{\gamma}^{\{ \mu \}}, \ \widehat{\gamma}^{\{ 2 \ p+1 \}}, \ \widehat{\gamma}^{\{ 2 \ p+2 \}} \right), \ \backslash \backslash \\ & \ \& \ \widehat{\gamma}^{\{ \mu \}} := \gamma^{\{ \mu \}} \ \otimes \left[\bigotimes_{i=1}^{\left[\frac{n-2}{2} \right]} \sigma^1 \right], \\ & \ \widehat{\gamma}^{2p+1} := i \mathbb{1} \otimes \left(\bigotimes_{i=1}^{p-2} \sigma^0 \right) \otimes \sigma^3 \otimes \left(\bigotimes_{j=1}^{l-p+1} \sigma^1 \right), \\ & \ \widehat{\gamma}^{2p+2} := i \mathbb{1} \otimes \left(\bigotimes_{i=1}^{p-2} \sigma^0 \right) \otimes \sigma^2 \otimes \left(\bigotimes_{j=1}^{l-p+2} \sigma^1 \right), \end{aligned}$	$\begin{aligned} (\widehat{\gamma}^A)^{\mathrm{T}} &:= (\widehat{\gamma}^{\mu}, \widehat{\gamma}^{2p+1}, \widehat{\gamma}^{2p+2}), \\ \widehat{\gamma}^{\mu} &:= \gamma^{\mu} \otimes \left[\bigotimes_{i=1}^{\left[\frac{n-2}{2} \right]} \sigma^1 \right], \\ \widehat{\gamma}^{2p+1} &:= i \mathbb{1} \otimes \left(\bigotimes_{i=1}^{p-2} \sigma^0 \right) \otimes \sigma^3 \otimes \left(\bigotimes_{j=1}^{l-p+1} \sigma^1 \right), \\ \widehat{\gamma}^{2p+2} &:= i \mathbb{1} \otimes \left(\bigotimes_{i=1}^{p-2} \sigma^0 \right) \otimes \sigma^2 \otimes \left(\bigotimes_{j=1}^{l-p+2} \sigma^1 \right), \end{aligned}$	$\begin{aligned} (\widehat{\gamma}^A)^{\mathrm{T}} &:= (\widehat{\gamma}^{\mu}, \widehat{\gamma}^{2p+1}, \widehat{\gamma}^{2p+2}), \\ \widehat{\gamma}^{\mu} &:= \gamma^{\mu} \otimes \left[\bigotimes_{i=1}^{\left[\frac{n-2}{2} \right]} \sigma^1 \right], \\ \widehat{\gamma}^{2p+1} &:= i \mathbb{1} \otimes \left(\bigotimes_{i=1}^{p-2} \sigma^0 \right) \otimes \sigma^3 \otimes \left(\bigotimes_{j=1}^{l-p+1} \sigma^1 \right), \\ \widehat{\gamma}^{2p+2} &:= i \mathbb{1} \otimes \left(\bigotimes_{i=1}^{p-2} \sigma^0 \right) \otimes \sigma^2 \otimes \left(\bigotimes_{j=1}^{l-p+2} \sigma^1 \right), \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_ EQ1090 (A.17)	367	■ 0.967	$\stackrel{*}{\alpha}\left(e_{\alpha_1},\ldots,e_{\alpha_p}\right):=\alpha\left(\sigma_*e_{\alpha_1},\ldots,\sigma_*e_{\alpha_p}\right).$	$\stackrel{*}{\sigma}\alpha\left(e_{\alpha_1},\ldots,e_{\alpha_p}\right):=\alpha\left(\sigma_*e_{\alpha_1},\ldots,\sigma_*e_{\alpha_p}\right).$	
mielke-geometrodynamics_ EQ0592 (9.1.10)	195	■ 0.968	$\stackrel{+}{\Omega}=\frac{\gamma\kappa}{\ell^{*2}}\vartheta\wedge\vartheta.$	$\stackrel{+}{\Omega}=\frac{\gamma\kappa}{\ell^{*2}}\vartheta\wedge\vartheta.$	
mielke-geometrodynamics_ EQ0611 (9.3.2)	202	■ 0.968	$L_{\mathrm{SKY}}:=-\frac{1}{2}R_{\alpha\beta}\wedge{}^*R^{\alpha\beta},$	$L_{\mathrm{SKY}}:=-\frac{1}{2}R_{\alpha\beta}\wedge{}^*R^{\alpha\beta},$	
mielke-geometrodynamics_ EQ0841 (12.2.9)	272	■ 0.968	$\mu_{\alpha}=\frac{1}{4}\vartheta_{\alpha}\wedge{}^*j_5.$	$\mu_{\alpha}=\frac{1}{4}\vartheta_{\alpha}\wedge{}^*j_5.$	
mielke-geometrodynamics_ EQ0636 (10.2.7)	212	■ 0.970	$T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha},$	$T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha},$	
mielke-geometrodynamics_ EQ1137 (C.5)	376	■ 0.970	$3\,c_{[i s}{}^pc_{ jk]}{}^s:=c_{is}{}^pc_{jk}{}^s+c_{js}{}^pc_{ki}{}^s+c_{ks}{}^pc_{ij}{}^s\equiv 0.$	$3c_{[i s}{}^pc_{ jk]}{}^s:=c_{is}{}^pc_{jk}{}^s+c_{js}{}^pc_{ki}{}^s+c_{ks}{}^pc_{ij}{}^s\equiv 0.$	
mielke-geometrodynamics_ EQ0415 (6.5.3)	137	■ 0.970	$\begin{aligned} \langle\!\langle 0 W(\gamma) 0\rangle\!\rangle^{(1)} &= \int_{\mathcal{C} \mathcal{G}} \mathcal{D}A e^{iC} W(A,\gamma) \\ &= \frac{1}{4\pi} \oint ds \oint dt \dot{\gamma}^{\mu}(s) \dot{\gamma}^{\nu}(t) \eta_{0\mu\nu\kappa} \frac{\gamma^{\kappa}(s) - \gamma^{\kappa}(t)}{ \gamma(s) - \gamma(t) ^3} \end{aligned}$	$\langle\!\langle 0 W(\gamma) 0\rangle\!\rangle^{(1)} = \int_{\mathcal{C} \mathcal{G}} \mathcal{D}A e^{iC} W(A,\gamma) = \frac{1}{4\pi} \oint ds \oint dt \dot{\gamma}^{\mu}(s) \dot{\gamma}^{\nu}(t) \eta_{0\mu\nu\kappa} \frac{\gamma^{\kappa}(s) - \gamma^{\kappa}(t)}{ \gamma(s) - \gamma(t) ^3}$	$\begin{aligned} \langle 0 W(\gamma) 0\rangle^{(1)} &= \int_{\mathcal{C} \mathcal{G}} \mathcal{D}A e^{iC} W(A,\gamma) \\ &= \frac{1}{4\pi} \oint ds \oint dt \dot{\gamma}^{\mu}(s) \dot{\gamma}^{\nu}(t) \eta_{0\mu\nu\kappa} \frac{\gamma^{\kappa}(s) - \gamma^{\kappa}(t)}{ \gamma(s) - \gamma(t) ^3} \end{aligned}$
mielke-geometrodynamics_ EQ0495 (7.7.13)	162	■ 0.971	$\{ \}^*\!\!\!\left(\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta}\right)=\eta^{\alpha\beta\gamma\delta},\text{ where }\eta_{\alpha\beta\gamma\delta}:=+\delta_{\alpha\beta\gamma\delta}^{0123},$	$\left(\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta}\right)=\eta^{\alpha\beta\gamma\delta},\text{ where }\eta_{\alpha\beta\gamma\delta}:=+\delta_{\alpha\beta\gamma\delta}^{0123},$	$\left(\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta}\right)=\eta^{\alpha\beta\gamma\delta},\text{ where }\eta_{\alpha\beta\gamma\delta}:=+\delta_{\alpha\beta\gamma\delta}^{0123},$
mielke-geometrodynamics_ EQ1147 (C.16)	377	■ 0.971	$\begin{aligned} \left[\lambda_i,\lambda_j\right] &= 2if_{ij}^k\lambda_k\quad (c_{ij}^k=2if_{ig}^k), \\ \left\{\lambda_i,\lambda_j\right\}_+ &= \frac{4}{N}\delta_{ij}+2d_{ij}^k\lambda_k. \end{aligned}$	$\begin{aligned} [\lambda_i,\lambda_j] &= 2if_{ij}^k\lambda_k\quad (c_{ij}^k=2if_{ig}^k), \\ \{\lambda_i,\lambda_j\}_+ &= \frac{4}{N}\delta_{ij}+2d_{ij}^k\lambda_k. \end{aligned}$	$\begin{aligned} [\lambda_i,\lambda_j] &= 2if_{ij}^k\lambda_k\quad (c_{ij}^k=2if_{ig}^k), \\ \{\lambda_i,\lambda_j\}_+ &= \frac{4}{N}\delta_{ij}+2d_{ij}^k\lambda_k. \end{aligned}$
mielke-geometrodynamics_ EQ0129 (3.3.30)	61	■ 0.972	$\begin{aligned} j:=\frac{\delta L_{\mathrm{matter}}}{\delta A} &= \frac{\partial L_{\mathrm{matter}}}{\partial A} + d\frac{\partial L_{\mathrm{matter}}}{\partial dA} \\ &= \Psi\wedge\frac{\partial L_{\mathrm{matter}}}{\partial D\Psi} + D\frac{\partial L_{\mathrm{matter}}}{\partial B}, \end{aligned}$	$\begin{aligned} j:=\frac{\delta L_{\mathrm{matter}}}{\delta A} &= \frac{\partial L_{\mathrm{matter}}}{\partial A} + d\frac{\partial L_{\mathrm{matter}}}{\partial dA} \\ &= \Psi\wedge\frac{\partial L_{\mathrm{matter}}}{\partial D\Psi} + D\frac{\partial L_{\mathrm{matter}}}{\partial B}, \end{aligned}$	$\begin{aligned} j:=\frac{\delta L_{\mathrm{matter}}}{\delta A} &= \frac{\partial L_{\mathrm{matter}}}{\partial A} + d\frac{\partial L_{\mathrm{matter}}}{\partial dA} \\ &= \Psi\wedge\frac{\partial L_{\mathrm{matter}}}{\partial D\Psi} + D\frac{\partial L_{\mathrm{matter}}}{\partial B}, \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_248 EQ0784 (11.3.3)		■ 0.972	$\begin{aligned} L_{\mathrm{NL}} &= \frac{1}{2} \bar{\psi} [\tilde{K}, {}^*\gamma] \psi = -3i^*[\Theta, \vartheta] \wedge (\bar{\psi}^* \gamma \psi) \\ &= -3\tau_s \wedge [\Theta, \vartheta]. \end{aligned}$	$L_{\mathrm{NL}} = \frac{1}{2} \bar{\psi} [\tilde{K}, {}^*\gamma] \psi = -3i^*[\Theta, \vartheta] \wedge (\bar{\psi}^* \gamma \psi) = -3\tau_s \wedge [\Theta, \vartheta].$	$L_{\mathrm{NL}} = \frac{1}{2} \bar{\psi} [\tilde{K}, {}^*\gamma] \psi = -3i^*[\Theta, \vartheta] \wedge (\bar{\psi}^* \gamma \psi) = -3\tau_s \wedge [\Theta, \vartheta].$
mielke-geometrodynamics_218 EQ0666 (10.3.23)		■ 0.972	$\square \vartheta_\alpha + \frac{\ell}{\gamma} {}^*D^* (\delta \ell E_\alpha + S_\alpha) = 0,$	$\square \vartheta_\alpha + \frac{\ell}{\gamma} {}^*D^* (\delta \ell E_\alpha + S_\alpha) = 0,$	$\square \vartheta_\alpha + \frac{\ell}{\gamma} {}^*D^* (\delta \ell E_\alpha + S_\alpha) = 0,$
mielke-geometrodynamics_306 EQ0945 (14.1.26)		■ 0.972	$\begin{aligned} \Sigma_{\mathrm{YM}}(\Omega) \wedge \Sigma_{\mathrm{YM}}(\Omega) &= \frac{1}{4\alpha_g} \mathrm{Tr} \{ -\xi \wedge \xi \wedge (\vartheta \wedge {}^*\xi) \wedge (\vartheta \wedge {}^*\xi) \\ &+ \vartheta \wedge \vartheta \wedge {}^*(\xi \wedge {}^*\xi)^2 - \vartheta \wedge {}^*\xi \wedge {}^*(\vartheta \wedge {}^*\xi) \wedge {}^*(\xi \wedge {}^*\xi) \} \\ &= \vartheta \wedge \vartheta \wedge {}^*(\Sigma_{\mathrm{YM}}(\xi) \wedge {}^*\Sigma_{\mathrm{YM}}(\xi)) \end{aligned}$	$\Sigma_{\mathrm{YM}}(\Omega) \wedge \Sigma_{\mathrm{YM}}(\Omega) = \frac{1}{4\alpha_g} \mathrm{Tr} \{ -\xi \wedge \xi \wedge (\vartheta \wedge {}^*\xi) \wedge (\vartheta \wedge {}^*\xi) \\ + \vartheta \wedge \vartheta \wedge {}^*(\xi \wedge {}^*\xi)^2 - \vartheta \wedge {}^*\xi \wedge {}^*(\vartheta \wedge {}^*\xi) \wedge {}^*(\xi \wedge {}^*\xi) \} \\ = \vartheta \wedge \vartheta \wedge {}^*(\Sigma_{\mathrm{YM}}(\xi) \wedge {}^*\Sigma_{\mathrm{YM}}(\xi))$	$\Sigma_{\mathrm{YM}}(\Omega) \wedge \Sigma_{\mathrm{YM}}(\Omega) = \frac{1}{4\alpha_g} \mathrm{Tr} \{ -\xi \wedge \xi \wedge (\vartheta \wedge {}^*\xi) \wedge (\vartheta \wedge {}^*\xi) \\ + \vartheta \wedge \vartheta \wedge {}^*(\xi \wedge {}^*\xi)^2 - \vartheta \wedge {}^*\xi \wedge {}^*(\vartheta \wedge {}^*\xi) \wedge {}^*(\xi \wedge {}^*\xi) \} \\ = \vartheta \wedge \vartheta \wedge {}^*(\Sigma_{\mathrm{YM}}(\xi) \wedge {}^*\Sigma_{\mathrm{YM}}(\xi))$
mielke-geometrodynamics_153 EQ0462 (7.3.1)		■ 0.973	$L_{\mathrm{SKY}} = \mathrm{Tr}(\Omega^g \wedge {}^*\Omega^g) = -\frac{1}{2} R^{\alpha\beta} \wedge {}^*R_{\alpha\beta},$	$L_{\mathrm{SKY}} = \mathrm{Tr}(\Omega^g \wedge {}^*\Omega^g) = -\frac{1}{2} R^{\alpha\beta} \wedge {}^*R_{\alpha\beta},$	$L_{\mathrm{SKY}} = \mathrm{Tr}(\Omega^g \wedge {}^*\Omega^g) = -\frac{1}{2} R^{\alpha\beta} \wedge {}^*R_{\alpha\beta},$
mielke-geometrodynamics_124 EQ0356 (6.2.7)		■ 0.974	$D T^{\alpha} \equiv R_{\beta}^{\alpha} \wedge \vartheta^{\beta}$	$DT^\alpha \equiv R_\beta^\alpha \wedge \vartheta^\beta$	$DT^\alpha \equiv R_\beta^\alpha \wedge \vartheta^\beta$
mielke-geometrodynamics_228 EQ0712 (10.8.1)		■ 0.975	$\Psi = \Psi_i dx^i = \Psi_\alpha \vartheta^\alpha$	$\Psi = \Psi_i dx^i = \Psi_\alpha \vartheta^\alpha$	$\Psi = \Psi_i dx^i = \Psi_\alpha \vartheta^\alpha$
mielke-geometrodynamics_314 EQ0971 (14.2.12)		■ 0.975	$\hat{d} := \left[\begin{array}{c c} d & 0 \\ 0 & \overset{\circ}{d} \end{array} \right],$	$\hat{d} := \left[\begin{array}{c c} d & 0 \\ 0 & \overset{\circ}{d} \end{array} \right],$	$\hat{d} := \left[\begin{array}{c c} d & 0 \\ 0 & \overset{\circ}{d} \end{array} \right],$
mielke-geometrodynamics_181 EQ0557 (8.4.11)		■ 0.975	$\left\langle dj_5 \right\rangle = 2i^*m \left\langle \bar{\psi} \gamma_5 \psi \right\rangle - \frac{1}{48\pi^2} \left(R_{\alpha\beta}^{\{ \} } \wedge R^{\{\beta\alpha\beta} + \frac{1}{2} d\mathscr{A} \wedge d\mathscr{A} \right),$	$\langle dj_5 \rangle = 2i^*m \langle \bar{\psi} \gamma_5 \psi \rangle - \frac{1}{48\pi^2} \left(R_{\alpha\beta}^{\{ \} } \wedge R^{\{\beta\alpha\beta} + \frac{1}{2} d\mathscr{A} \wedge d\mathscr{A} \right),$	$\langle dj_5 \rangle = 2i^*m \langle \bar{\psi} \gamma_5 \psi \rangle - \frac{1}{48\pi^2} \left(R_{\alpha\beta}^{\{ \} } \wedge R^{\{ \} \alpha\beta} + \frac{1}{2} d\mathscr{A} \wedge d\mathscr{A} \right),$
mielke-geometrodynamics_160 EQ0487 (7.7.4)		■ 0.977	$\left. \left. \left. \eta_{\alpha\beta} \quad \rightarrow \quad \tilde{\eta}_{\alpha\beta} = \eta_{\alpha\beta} + \ell^2 \eta_{\alpha\beta\gamma} \wedge e_\mu \right]^* H^{\mu\gamma} + \frac{\ell^4}{2} \eta_{\alpha\beta\gamma\delta} (e_\mu]^* H^{\gamma\mu} \right) \wedge (e_\nu]^* H^{\delta\nu} \right).$	$\eta_{\alpha\beta} \quad \rightarrow \quad \tilde{\eta}_{\alpha\beta} = \eta_{\alpha\beta} + \ell^2 \eta_{\alpha\beta\gamma} \wedge e_\mu]^* H^{\mu\gamma} + \frac{\ell^4}{2} \eta_{\alpha\beta\gamma\delta} (e_\mu]^* H^{\gamma\mu} \right) \wedge (e_\nu]^* H^{\delta\nu} \right).$	$\eta_{\alpha\beta} \quad \rightarrow \quad \tilde{\eta}_{\alpha\beta} = \eta_{\alpha\beta} + \ell^2 \eta_{\alpha\beta\gamma} \wedge e_\mu]^* H^{\mu\gamma} + \frac{\ell^4}{2} \eta_{\alpha\beta\gamma\delta} (e_\mu]^* H^{\gamma\mu} \right) \wedge (e_\nu]^* H^{\delta\nu} \right).$
mielke-geometrodynamics_355 EQ1068 (16.2.7)		■ 0.978	$\left[\partial_{\rho^*}^2 - V_{\mathrm{eff}}^{\iota}(\rho^*) + \frac{\omega^2}{\beta^2} \right] F = 0$	$\left[\partial_{\rho^*}^2 - V_{\mathrm{eff}}^{\iota}(\rho^*) + \frac{\omega^2}{\beta^2} \right] F = 0$	$\left[\partial_{\rho^*}^2 - V_{\mathrm{eff}}^{\iota}(\rho^*) + \frac{\omega^2}{\beta^2} \right] F = 0$
mielke-geometrodynamics_371 EQ1114 (B.26)		■ 0.979	$H^{\{p\}}(M) = 0 \text{ for } p > n,$	$H^p(M) = 0 \text{ for } p > n,$	$H^p(M) = 0 \text{ for } p > n,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_291EQ0007 (2.2.2)	33	■ 0.980	$\vartheta=\vartheta^{\alpha}\,P_{\alpha}\in C^{\infty}\left(T^{\ast}(M)\right),$	$\vartheta=\vartheta^{\alpha}P_{\alpha}\in C^{\infty}(T^{\ast}(M)),$	$\vartheta=\vartheta^{\alpha}P_{\alpha}\in C^{\infty}(T^{\ast}(M)),$
mielke-geometrodynamics_291EQ0908 (13.5.13)	291	■ 0.980	$D\,g_{\alpha\beta}=0,$	$Dg_{\alpha\beta}=0,$	$Dg_{\alpha\beta}=0,$
mielke-geometrodynamics_291EQ0536 (8.2.3)	175	■ 0.980	$\begin{aligned} (d\oplus s)\,\widetilde{\Gamma}+\frac{1}{2}[\widetilde{\Gamma},\widetilde{\Gamma}]&=\widetilde{\Omega}^g\\ (d\oplus s)\widetilde{\Omega}^g+\left[\widetilde{\Gamma},\widetilde{\Omega}^g\right]&\equiv 0 \end{aligned}$	$(d\oplus s)\widetilde{\Gamma}+\frac{1}{2}[\widetilde{\Gamma},\widetilde{\Gamma}]=\widetilde{\Omega}^g$ $(d\oplus s)\widetilde{\Omega}^g+\left[\widetilde{\Gamma},\widetilde{\Omega}^g\right]\equiv 0$	$(d\oplus s)\widetilde{\Gamma}+\frac{1}{2}[\widetilde{\Gamma},\widetilde{\Gamma}]=\widetilde{\Omega}^g,$ $(d\oplus s)\widetilde{\Omega}^g+[\widetilde{\Gamma},\widetilde{\Omega}^g]\equiv 0,$
mielke-geometrodynamics_291EQ0911 (13.6.1)	292	■ 0.980	$\begin{aligned} ds^2&\,=\frac{\ell^2}{\mu^2}\left\langle\hat{D}\Phi^A\otimes_s\hat{D}\Phi_A\right\rangle=\frac{\ell^2}{\mu^2}\hat{D}\Phi_0^A\otimes_s\hat{D}\Phi_{A0}\\&=\ell^2\Gamma_4^\alpha\otimes_s\Gamma_\alpha^4=(\vartheta^\alpha-D\xi^\alpha)\otimes_s(\theta_\alpha-D\xi_\alpha^*)\\&\stackrel{*}{=} \vartheta^\alpha\otimes_s\theta_\alpha=g_{ij}dx^i\otimes_sdx^j \end{aligned}$	$ds^2:=\frac{\ell^2}{\mu^2}\left\langle\hat{D}\Phi^A\otimes_s\hat{D}\Phi_A\right\rangle=\frac{\ell^2}{\mu^2}\hat{D}\Phi_0^A\otimes_s\hat{D}\Phi_{A0}$ $=\ell^2\Gamma_4^\alpha\otimes_s\Gamma_\alpha^4=(\vartheta^\alpha-D\xi^\alpha)\otimes_s(\theta_\alpha-D\xi_\alpha^*)$ $\stackrel{*}{=} \vartheta^\alpha\otimes_s\theta_\alpha=g_{ij}dx^i\otimes_sdx^j$	$ds^2:=\frac{\ell^2}{\mu^2}\langle\hat{D}\Phi^A\otimes_s\hat{D}\Phi_A\rangle=\frac{\ell^2}{\mu^2}\hat{D}\Phi_0^A\otimes_s\hat{D}\Phi_{A0}$ $=\ell^2\Gamma_4^\alpha\otimes_s\Gamma_\alpha^4=(\vartheta^\alpha-D\xi^\alpha)\otimes_s(\theta_\alpha-D\xi_\alpha^*)$ $\stackrel{*}{=} \vartheta^\alpha\otimes_s\theta_\alpha=g_{ij}dx^i\otimes_sdx^j.$
mielke-geometrodynamics_291EQ0931 (14.1.12)	304	■ 0.981	$\Omega\rightarrowtail e^{*\delta}\Omega:=\Omega\cos\delta+(-1)^z*\Omega\sin\delta.$	$\Omega\rightarrow e^{*\delta}\Omega:=\Omega\cos\delta+(-1)^z*\Omega\sin\delta.$	$\Omega\rightarrow e^{*\delta}\Omega:=\Omega\cos\delta+(-1)^z*\Omega\sin\delta.$
mielke-geometrodynamics_291EQ0671 (10.3.28)	219	■ 0.982	$\check{\mathscr{H}}_{\alpha}^{\star}:=dH_{\alpha}^{\star}\cong\tau_{\alpha}^{\star}+E_{\alpha}^{\star}-(-1)^s\eta_{\alpha\beta\gamma}\Gamma^{\star\beta}\wedge H^{\star\gamma},$	$\check{\mathscr{H}}_{\alpha}^{\star}:=dH_{\alpha}^{\star}\cong\tau_{\alpha}^{\star}+E_{\alpha}^{\star}-(-1)^s\eta_{\alpha\beta\gamma}\Gamma^{\star\beta}\wedge H^{\star\gamma},$	$\check{\mathscr{H}}_{\alpha}^{\star}:=dH_{\alpha}^{\star}\cong\tau_{\alpha}^{\star}+E_{\alpha}^{\star}-(-1)^s\eta_{\alpha\beta\gamma}\Gamma^{\star\beta}\wedge H^{\star\gamma},$
mielke-geometrodynamics_291EQ1086 (A.12)	366	■ 0.982	$\begin{aligned} &\mathrm{C}(1,1)=\mathrm{C}(2,0)=\mathrm{GL}(2,\mathbb{R}),\\ &\mathrm{C}(0,2)=\mathbb{H}\quad\text{(field of quaternions)}, \end{aligned}$	$\begin{aligned} \mathrm{C}(1,1)&=\mathrm{C}(2,0)=\mathrm{GL}(2,\mathbb{R}),\\ \mathrm{C}(0,2)&=\mathbb{H}\quad\text{(field of quaternions)}, \end{aligned}$	$\begin{aligned} \mathrm{C}(1,1)&=\mathrm{C}(2,0)=\mathrm{GL}(2,\mathbb{R}),\\ \mathrm{C}(0,2)&=\mathbb{H}\quad\text{(field of quaternions)}, \end{aligned}$
mielke-geometrodynamics_291EQ0229 (4.2.22)	88	■ 0.982	$\Theta:=D\vartheta=d\vartheta-\left[\omega^{\mathrm{L}},\vartheta\right]$	$\Theta:=D\vartheta=d\vartheta-\left[\omega^{\mathrm{L}},\vartheta\right]$	$\Theta:=D\vartheta=d\vartheta-\left[\omega^{\mathrm{L}},\vartheta\right]$
mielke-geometrodynamics_291EQ0836 (12.2.4)	271	■ 0.983	$\mathscr{A}:=\frac{1}{4}{}^*\mathrm{Tr}(\gamma\wedge D\gamma)={}^*(\vartheta^\alpha\wedge T_\alpha)=\frac{1}{2}T^{[\alpha\beta\gamma]}{}_{\eta_{\alpha\beta\gamma}}=\mathscr{A}_idx^i,$	$\mathscr{A}:=\frac{1}{4}{}^*\mathrm{Tr}(\gamma\wedge D\gamma)={}^*(\vartheta^\alpha\wedge T_\alpha)=\frac{1}{2}T^{[\alpha\beta\gamma]}{}_{\eta_{\alpha\beta\gamma}}=\mathscr{A}_idx^i,$	$\mathscr{A}:=\frac{1}{4}{}^*\mathrm{Tr}(\gamma\wedge D\gamma)={}^*(\vartheta^\alpha\wedge T_\alpha)=\frac{1}{2}T^{[\alpha\beta\gamma]}{}_{\eta_{\alpha\beta\gamma}}=\mathscr{A}_idx^i,$
mielke-geometrodynamics_291EQ0511 (7.7.29)	165	■ 0.983	$\begin{aligned} \Theta&:=D\gamma=d\gamma+[\Gamma,\gamma]=\left(d\vartheta^\alpha+\Gamma_\beta^\alpha\wedge\vartheta^\beta\right)\gamma_\alpha\\&=T^\alpha\gamma_\alpha=\frac{1}{2}T_{ij}^\alpha\gamma_\alpha dx^i\wedge dx^j \end{aligned}$	$\begin{aligned} \Theta:=D\gamma&=d\gamma+[\Gamma,\gamma]=\left(d\vartheta^\alpha+\Gamma_\beta^\alpha\wedge\vartheta^\beta\right)\gamma_\alpha\\&=T^\alpha\gamma_\alpha=\frac{1}{2}T_{ij}^\alpha\gamma_\alpha dx^i\wedge dx^j \end{aligned}$	$\begin{aligned} \Theta:=D\gamma&=d\gamma+[\Gamma,\gamma]=\left(d\vartheta^\alpha+\Gamma_\beta^\alpha\wedge\vartheta^\beta\right)\gamma_\alpha\\&=T^\alpha\gamma_\alpha=\frac{1}{2}T_{ij}^\alpha\gamma_\alpha dx^i\wedge dx^j, \end{aligned}$
mielke-geometrodynamics_291EQ0637 (10.2.8)	212	■ 0.983	$\Gamma_{\alpha}^{\star}=\Gamma_{\alpha}^{\{\}\star}-K_{\alpha}^{\star}$	$\Gamma_{\alpha}^{\star}=\Gamma_{\alpha}^{\{\}\star}-K_{\alpha}^{\star}$	$\Gamma_{\alpha}^{\star}=\Gamma_{\alpha}^{\{\}\star}-K_{\alpha}^{\star}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_ E00572 (8.6.10)	184	■ 0.990	$\begin{aligned} H_{\alpha}(* *) &= \frac{\theta_{\mathrm{T}}^*}{2\ell^2} K^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha} = \frac{1}{\ell^2} \left[\theta_{\mathrm{T}}^* K_{\alpha}^{(*)} - \theta_{\mathrm{T}} T_{\alpha} \right] \\ &= \frac{\theta_{\mathrm{T}}^*}{\ell^2} * \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta} \rfloor T^{\beta}) - \frac{1}{2} e_{\alpha} \rfloor (T^{\beta} \wedge \vartheta_{\beta}) \right] - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha} \\ &\simeq -\partial L_{\theta} / \partial T^{\alpha}. \end{aligned}$	$H_{\alpha}(**) = \frac{\theta_{\mathrm{T}}^*}{2\ell^2} K^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha} = \frac{1}{\ell^2} \left[\theta_{\mathrm{T}}^* K_{\alpha}^{(*)} - \theta_{\mathrm{T}} T_{\alpha} \right] \\ = \frac{\theta_{\mathrm{T}}^*}{\ell^2} * \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta} \rfloor T^{\beta}) - \frac{1}{2} e_{\alpha} \rfloor (T^{\beta} \wedge \vartheta_{\beta}) \right] - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha} \\ \simeq -\partial L_{\theta} / \partial T^{\alpha}.$	$H_{\alpha}(**) = \frac{\theta_{\mathrm{T}}^*}{2\ell^2} K^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha} = \frac{1}{\ell^2} \left[\theta_{\mathrm{T}}^* K_{\alpha}^{(*)} - \theta_{\mathrm{T}} T_{\alpha} \right] \\ = \frac{\theta_{\mathrm{T}}^*}{\ell^2} * \left[T_{\alpha} - \vartheta_{\alpha} \wedge (e_{\beta} \rfloor T^{\beta}) - \frac{1}{2} e_{\alpha} \rfloor (T^{\beta} \wedge \vartheta_{\beta}) \right] - \frac{\theta_{\mathrm{T}}}{\ell^2} T_{\alpha} \\ \simeq -\partial L_{\theta} / \partial T^{\alpha}.$
mielke-geometrodynamics_ E00110 (3.3.11)	58	■ 0.990	$\begin{aligned} F := & E \wedge dt + B \\ &= (E_x dx + E_y dy + E_z dz) \wedge dt \\ &+ B_x dy \wedge dz + B_y dz \wedge dx + B_z dx \wedge dy. \end{aligned}$	$F := E \wedge dt + B \\ = (E_x dx + E_y dy + E_z dz) \wedge dt \\ + B_x dy \wedge dz + B_y dz \wedge dx + B_z dx \wedge dy.$	$F := E \wedge dt + B \\ = (E_x dx + E_y dy + E_z dz) \wedge dt \\ + B_x dy \wedge dz + B_y dz \wedge dx + B_z dx \wedge dy.$
mielke-geometrodynamics_ E00142 (3.4.13)	64	■ 0.990	$\begin{aligned} D \tau &= DD\Pi^{\Omega} = [\Omega, \Pi^{\Omega}] \\ &= \frac{2}{\alpha_g} \left\{ [\Omega, * \Omega] \frac{\partial L}{\partial \mathcal{F}} + (-1)^s [\Omega, \Omega] \frac{\partial L}{\partial * \mathcal{F}} \right\} = 0. \end{aligned}$	$D\tau = DD\Pi^{\Omega} = [\Omega, \Pi^{\Omega}] \\ = \frac{2}{\alpha_g} \left\{ [\Omega, * \Omega] \frac{\partial L}{\partial \mathcal{F}} + (-1)^s [\Omega, \Omega] \frac{\partial L}{\partial * \mathcal{F}} \right\} = 0.$	$D\tau = DD\Pi^{\Omega} = [\Omega, \Pi^{\Omega}] \\ = \frac{2}{\alpha_g} \left\{ [\Omega, * \Omega] \frac{\partial L}{\partial \mathcal{F}} + (-1)^s [\Omega, \Omega] \frac{\partial L}{\partial * \mathcal{F}} \right\} = 0.$
mielke-geometrodynamics_ E00155 (3.5.3)	67	■ 0.990	$\stackrel{*}{\sigma} \Omega = \frac{1}{4} F_{\alpha\beta\mu\nu} \sigma^{\mu\nu} \otimes \vartheta^{\alpha} \wedge \vartheta^{\beta}$	$\stackrel{*}{\sigma} \Omega = \frac{1}{4} F_{\alpha\beta\mu\nu} \sigma^{\mu\nu} \otimes \vartheta^{\alpha} \wedge \vartheta^{\beta}$	$\stackrel{*}{\sigma} \Omega = \frac{1}{4} F_{\alpha\beta\mu\nu} \sigma^{\mu\nu} \otimes \vartheta^{\alpha} \wedge \vartheta^{\beta}$
mielke-geometrodynamics_ E00682 (10.3.40)	221	■ 0.990	$\theta_{\mathrm{L}} C_{\mathrm{L}} - \tilde{\theta}_{\mathrm{L}} \tilde{C}_{\mathrm{L}} = L_{\mathrm{MB}} + \frac{\Delta\theta_{\mathrm{TL}}}{\ell} d(\Gamma^{*\alpha} \wedge \vartheta_{\alpha}),$	$\theta_{\mathrm{L}} C_{\mathrm{L}} - \tilde{\theta}_{\mathrm{L}} \tilde{C}_{\mathrm{L}} = L_{\mathrm{MB}} + \frac{\Delta\theta_{\mathrm{TL}}}{\ell} d(\Gamma^{*\alpha} \wedge \vartheta_{\alpha}),$	$\theta_{\mathrm{L}} C_{\mathrm{L}} - \tilde{\theta}_{\mathrm{L}} \tilde{C}_{\mathrm{L}} = L_{\mathrm{MB}} + \frac{\Delta\theta_{\mathrm{TL}}}{\ell} d(\Gamma^{*\alpha} \wedge \vartheta_{\alpha}),$
mielke-geometrodynamics_ E00874 (13.3.2)	284	■ 0.990	$[Z^A{}_B, Z^C{}_D] = \delta_D^A Z^C{}_B - \delta_B^C Z^A{}_D$	$[Z^A{}_B, Z^C{}_D] = \delta_D^A Z^C{}_B - \delta_B^C Z^A{}_D$	$[\mathcal{E}^A{}_B, \mathcal{E}^C{}_D] = \delta_D^A \mathcal{E}^C{}_B - \delta_B^C \mathcal{E}^A{}_D$
mielke-geometrodynamics_ E00042 (2.6.5)	42	■ 0.991	$F_{\alpha\beta}{}^j = \partial_{\alpha} \Gamma_{\beta}{}^j - \partial_{\beta} \Gamma_{\alpha}{}^j - c_{kl}{}^j \Gamma_{\alpha}{}^k \Gamma_{\beta}{}^l,$	$F_{\alpha\beta}{}^j = \partial_{\alpha} \Gamma_{\beta}{}^j - \partial_{\beta} \Gamma_{\alpha}{}^j - c_{kl}{}^j \Gamma_{\alpha}{}^k \Gamma_{\beta}{}^l,$	$F_{\alpha\beta}{}^j = \partial_{\alpha} \Gamma_{\beta}{}^j - \partial_{\beta} \Gamma_{\alpha}{}^j - c_{kl}{}^j \Gamma_{\alpha}{}^k \Gamma_{\beta}{}^l,$
mielke-geometrodynamics_ E00233 (4.2.26)	88	■ 0.991	$\Omega^{\mathrm{T}} = f\Theta + df \wedge \vartheta - \Omega^{\mathrm{L}} \otimes \widetilde{\varphi}$	$\Omega^{\mathrm{T}} = f\Theta + df \wedge \vartheta - \Omega^{\mathrm{L}} \otimes \widetilde{\varphi}$	$\Omega^{\mathrm{T}} = f\Theta + df \wedge \vartheta - \Omega^{\mathrm{L}} \otimes \widetilde{\varphi}$
mielke-geometrodynamics_ E00778	246	■ 0.991	$\langle -\infty D\tau_A \infty \rangle^{\scriptscriptstyle (+)} = \frac{1}{(4.)8\pi^2} \mathrm{Tr} \left(\Omega^g \wedge \Omega^{g(*)} \right).$	$\langle -\infty D\tau_A \infty \rangle^{\scriptscriptstyle (+)} = \frac{1}{(4.)8\pi^2} \mathrm{Tr} (\Omega^g \wedge \Omega^{g(*)}).$	$\langle -\infty D\tau_A \infty \rangle^{\scriptscriptstyle (+)} = \frac{1}{(4.)8\pi^2} \mathrm{Tr} (\Omega^g \wedge \Omega^{g(*)}).$
mielke-geometrodynamics_ E00644 (10.3.1)	214	■ 0.991	$\widetilde{\Gamma}_{\alpha}^{*} = \Gamma_{\alpha}^{*} - (-1)^s \frac{\varepsilon}{2\ell} \vartheta_{\alpha},$	$\widetilde{\Gamma}_{\alpha}^{*} = \Gamma_{\alpha}^{*} - (-1)^s \frac{\varepsilon}{2\ell} \vartheta_{\alpha},$	$\widetilde{\Gamma}_{\alpha}^{*} = \Gamma_{\alpha}^{*} - (-1)^s \frac{\varepsilon}{2\ell} \vartheta_{\alpha},$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_370 EQ1102 (B.14)	370	■ 0.994	$\begin{aligned} d_{\mu}(M) \ \& \ :=\{ }^{\{ * \}} 1\!=\!\sqrt{\left \operatorname{det} g_{\alpha\beta}\right }\,\vartheta^1\wedge\cdots\wedge\vartheta^n=\eta \\ & \ =\sqrt{\left \operatorname{det} g_{ij}\right }dx^1\wedge\cdots\wedge dx^n=\frac{1}{n}\vartheta\wedge^*\vartheta. \end{aligned}$	$d_{\mu}(M) \ :=\ ^*1=\sqrt{ \det g_{\alpha\beta} }\vartheta^1\wedge\cdots\wedge\vartheta^n=\eta$ $=\sqrt{ \det g_{ij} }dx^1\wedge\cdots\wedge dx^n=\frac{1}{n}\vartheta\wedge^*\vartheta.$	$d_{\mu}(M) \ :=\ ^*1=\sqrt{ \det g_{\alpha\beta} }\vartheta^1\wedge\cdots\wedge\vartheta^n=\eta$ $=\sqrt{ \det g_{ij} }dx^1\wedge\cdots\wedge dx^n=\frac{1}{n}\vartheta\wedge^*\vartheta.$
mielke-geometrodynamics_154 EQ0467 (7.3.6)	154	■ 0.994	$\begin{aligned} L_{\mathrm{SKY}}^{\left(\star\right)}\& \ =-\frac{1}{2}R^{\alpha\beta}\wedge^*R_{\alpha\beta}-\frac{\left(-1\right)^s}{2}R^{\alpha\beta}\wedge R_{\alpha\beta}^{\left(\star\right)} \\ & \ =-\frac{1}{4}\left(R_{\alpha\beta}+^*R_{\alpha\beta}^{\left(\star\right)}\right)\wedge^*\left(R^{\alpha\beta}+^*R^{\alpha\beta\left(\star\right)}\right) \\ & \ =\frac{1}{2}\mathrm{Tr}\left(\bar{\Omega}^g\wedge^*\bar{\Omega}^g\right). \end{aligned}$	$L_{\mathrm{SKY}}^{\left(\star\right)}=-\frac{1}{2}R^{\alpha\beta}\wedge^*R_{\alpha\beta}-\frac{\left(-1\right)^s}{2}R^{\alpha\beta}\wedge R_{\alpha\beta}^{\left(\star\right)}$ $=-\frac{1}{4}\left(R_{\alpha\beta}+^*R_{\alpha\beta}^{\left(\star\right)}\right)\wedge^*\left(R^{\alpha\beta}+^*R^{\alpha\beta\left(\star\right)}\right)$ $=\frac{1}{2}\mathrm{Tr}\left(\bar{\Omega}^g\wedge^*\bar{\Omega}^g\right).$	$L_{\mathrm{SKY}}^{\left(\star\right)}=-\frac{1}{2}R^{\alpha\beta}\wedge^*R_{\alpha\beta}-\frac{\left(-1\right)^s}{2}R^{\alpha\beta}\wedge R_{\alpha\beta}^{\left(\star\right)}$ $=-\frac{1}{4}\left(R_{\alpha\beta}+^*R_{\alpha\beta}^{\left(\star\right)}\right)\wedge^*\left(R^{\alpha\beta}+^*R^{\alpha\beta\left(\star\right)}\right)$ $=\frac{1}{2}\mathrm{Tr}(\bar{\Omega}^g\wedge^*\bar{\Omega}^g).$
mielke-geometrodynamics_163 EQ0498 (7.7.16)	163	■ 0.994	$\{ }^{\{ * \}} \ \Phi\!=\!\frac{1}{(4-p!)\ p!}\ \sqrt{\operatorname{det} g_{ij}}\varepsilon^{i_1\cdots i_p}_{j_1\cdots j_{4-p}}\ \Phi_{i_1\cdots i_p}dx^{j_1}\wedge\cdots\wedge dx^{j_{4-p}}$	$^*\Phi:=\frac{1}{(4-p!)p!}\sqrt{\det g_{ij}}\varepsilon^{i_1\cdots i_p}_{j_1\cdots j_{4-p}}\Phi_{i_1\cdots i_p}dx^{j_1}\wedge\cdots\wedge dx^{j_{4-p}}$	$^*\Phi:=\frac{1}{(4-p!)p!}\sqrt{\det g_{ij}}\varepsilon^{i_1\cdots i_p}_{j_1\cdots j_{4-p}}\Phi_{i_1\cdots i_p}dx^{j_1}\wedge\cdots\wedge dx^{j_{4-p}},$
mielke-geometrodynamics_230 EQ0724 (10.10.4)	230	■ 0.994	$\delta_{\mathrm{susy}}L\!=\!i\overline{\sigma}\,\Psi\gamma^{\alpha}\wedge\frac{\delta L}{\delta\vartheta^{\alpha}}+\delta_{\mathrm{susy}}\Gamma_{\alpha}^{\star}\wedge\frac{\delta L}{\delta\Gamma_{\alpha}^{\star}}+\left(2\overline{D\sigma}+c\overline{\sigma}\gamma\right)\wedge\frac{\delta L}{\delta\overline{\Psi}},$	$\delta_{\mathrm{susy}}L=i\overline{\sigma}\Psi\gamma^{\alpha}\wedge\frac{\delta L}{\delta\vartheta^{\alpha}}+\delta_{\mathrm{susy}}\Gamma_{\alpha}^{\star}\wedge\frac{\delta L}{\delta\Gamma_{\alpha}^{\star}}+(2\overline{D\sigma}+c\overline{\sigma}\gamma)\wedge\frac{\delta L}{\delta\overline{\Psi}},$	$\delta_{\mathrm{susy}}L=i\overline{\sigma}\,\Psi\gamma^{\alpha}\wedge\frac{\delta L}{\delta\vartheta^{\alpha}}+\delta_{\mathrm{susy}}\Gamma_{\alpha}^{\star}\wedge\frac{\delta L}{\delta\Gamma_{\alpha}^{\star}}+\left(2\overline{D\sigma}+c\overline{\sigma}\gamma\right)\wedge\frac{\delta L}{\delta\overline{\Psi}},$
mielke-geometrodynamics_53 EQ0081 (3.1.6)	53	■ 0.994	$\begin{aligned} \delta L \ \& \ =\delta\!L\ \varphi\wedge^*\!L_{\varphi}+\delta\!L\ \varphi\wedge^*\!L_{\varphi}+d\!L\ \varphi\wedge^*\!L_{\varphi}+d\!L\ \varphi\wedge^*\!L_{\varphi} \\ & \ =\delta\!L\ \varphi\wedge^*\!L_{\varphi}+d\!L\ \varphi\wedge^*\!L_{\varphi}+d\!L\ \varphi\wedge^*\!L_{\varphi}+d\!L\ \varphi\wedge^*\!L_{\varphi}, \end{aligned}$	$\delta L=\delta\varphi\wedge^*L_{\varphi}+\delta(d\varphi)\wedge\pi$ $=\delta\varphi\wedge^*L_{\varphi}+d(\delta\varphi)\wedge\pi$ $=\delta\varphi\wedge(^*L_{\varphi}+d\pi)+d(\delta\varphi\wedge\pi),$	$\delta L=\delta\varphi\wedge^*L_{\varphi}+\delta(d\varphi)\wedge\pi$ $=\delta\varphi\wedge^*L_{\varphi}+d(\delta\varphi)\wedge\pi$ $=\delta\varphi\wedge(^*L_{\varphi}+d\pi)+d(\delta\varphi\wedge\pi),$
mielke-geometrodynamics_165 EQ0509 (7.7.27)	165	■ 0.994	$D\!=\!d+[\Gamma,\quad]$	$D=d+[\Gamma,\quad]$	$D=d+[\Gamma,\quad]$
mielke-geometrodynamics_130 EQ0387 (6.4.2)	130	■ 0.994	$\left.\{ }^{\{ \perp \}} \Psi\!=\!d\ t\wedge\Psi_{\{ \perp \}}\right\} ;\quad\Psi_{\{ \perp \}}\!=\!n\right\rfloor\Psi,$	$^{\perp}\Psi:=dt\wedge\Psi_{\perp};\quad\Psi_{\perp}:=n\rfloor\Psi,$	$^{\perp}\Psi:=dt\wedge\Psi_{\perp};\quad\Psi_{\perp}:=n\rfloor\Psi,$
mielke-geometrodynamics_214 EQ0646 (10.3.3)	214	■ 0.995	$\Gamma_{\alpha}^{\left(\star\right)}\!=\!\left(-1\right)^s\frac{\varepsilon}{2\ell}\vartheta_{\alpha},$	$\Gamma_{\alpha}^{\star}=(-1)^s\frac{\varepsilon}{2\ell}\vartheta_{\alpha},$	$\Gamma_{\alpha}^{\star}=(-1)^s\frac{\varepsilon}{2\ell}\vartheta_{\alpha},$
mielke-geometrodynamics_255 EQ0806 (11.4.18)	255	■ 0.995	$\overrightarrow{J^2}\chi_{\kappa}^m=j(j+1)\chi_{\kappa}^m$	$\overrightarrow{J^2}\chi_{\kappa}^m=j(j+1)\chi_{\kappa}^m$	$\overrightarrow{J^2}\chi_{\kappa}^m=j(j+1)\chi_{\kappa}^m$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_00995 (14.3.8)	321	■ 0.995	$L_{\text{anom.}}=\frac{\widehat{\ell}}{2}\bar{\psi}\gamma\wedge\gamma\wedge{}^*\Omega\psi=\frac{\ell^*}{4\sqrt{\alpha_g}}\bar{\psi}F_{\mu\nu}\gamma^5[\gamma^\mu,\gamma^\nu]\psi\eta.$	$L_{\text{anom.}}=\frac{\widehat{\ell}}{2}\bar{\psi}\gamma\wedge\gamma\wedge{}^*\Omega\psi=\frac{\ell^*}{4\sqrt{\alpha_g}}\bar{\psi}F_{\mu\nu}\gamma^5[\gamma^\mu,\gamma^\nu]\psi\eta.$	
mielke-geometrodynamics_00998 (3.2.8)	56	■ 0.995	$\Omega\rightarrow{}^{G^{-1}}\Omega=G^{-1}\Omega G.$	$\Omega\rightarrow{}^{G^{-1}}\Omega=G^{-1}\Omega G.$	
mielke-geometrodynamics_00930 (14.1.11)	304	■ 0.995	$\mathrm{R}=\{ \}^*\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)=0$	$\mathrm{R}=\left(\Omega^g\wedge\vartheta\wedge\vartheta\right)=0$	
mielke-geometrodynamics_01033 (15.1.10)	338	■ 0.995	$\frac{1}{2}\{ \}^*\left(\vartheta^f\wedge{}^*\Omega^f\right)+\Lambda_{\mathrm{eff}}^f\vartheta^f=0$	$\frac{1}{2}\left(\vartheta^f\wedge{}^*\Omega^f\right)+\Lambda_{\mathrm{eff}}^f\vartheta^f=0$	
mielke-geometrodynamics_00658 (10.3.15)	217	■ 0.995	$\{ \}^*\mathrm{D}\left(\vartheta^{\alpha}\wedge H_{\alpha}\right)-\{ \}^*\left(T^{\alpha}\wedge H_{\alpha}\right)=\frac{\gamma}{\delta\ell^2}\{ \}^*\left(\vartheta^{\alpha}\wedge T_{\alpha}\right)+\frac{1}{\delta\ell}\{ \}^*\left(\eta^{\alpha}\wedge H_{\alpha}\right).$	${}^*D(\vartheta^\alpha\wedge H_\alpha)-{}^*(T^\alpha\wedge H_\alpha)=\frac{\gamma}{\delta\ell^2}{}^*(\vartheta^\alpha\wedge T_\alpha)+\frac{1}{\delta\ell}{}^*(\eta^\alpha\wedge H_\alpha).$	${}^*D(\vartheta^\alpha\wedge H_\alpha)-{}^*(T^\alpha\wedge H_\alpha)=\frac{\gamma}{\delta\ell^2}{}^*(\vartheta^\alpha\wedge T_\alpha)+\frac{1}{\delta\ell}{}^*(\eta^\alpha\wedge H_\alpha).$
mielke-geometrodynamics_00207 (4.1.21)	83	■ 0.995	$\delta:\left\{\begin{array}{lll}A(n,\mathbb{R})&\times A(M)&\longrightarrow\\ \downarrow & \downarrow & \\(g,a)&(e(m),x)&=(\mathbf{g.e}(m),g^{-1}(x-a)).\end{array}\right.$	$\delta:\left\{\begin{array}{lll}A(n,\mathbb{R})&\times A(M)&\longrightarrow\\ \downarrow & \downarrow & \\(g,a)&(e(m),x)&=(\mathbf{g.e}(m),g^{-1}(x-a)).\end{array}\right.$	$\delta:\left\{\begin{array}{lll}A(n,\mathbb{R})&\times A(M)&\longrightarrow\\ \downarrow & \downarrow & \\(g,a)&(e(m),x)&=(g.e(m),g^{-1}(x-a)).\end{array}\right.$
mielke-geometrodynamics_00154 (3.5.2)	66	■ 0.995	$\stackrel{\scriptscriptstyle (-)}{\sigma}_{\mu\nu}:=\left\{\begin{array}{l} \sigma^{ij}=\frac{1}{2}\varepsilon^{ijk}\sigma_k\\ \sigma^{i0}=\left(-\right)\frac{1}{2}\sigma^i\end{array}\left i,j,k=1,2,3\right.\right.$	$\stackrel{\scriptscriptstyle (-)}{\sigma}_{\mu\nu}:=\left\{\begin{array}{l} \sigma^{ij}=\frac{1}{2}\varepsilon^{ijk}\sigma_k\\ \sigma^{i0}=\left(-\right)\frac{1}{2}\sigma^i\end{array}\left i,j,k=1,2,3\right.\right.$	$\stackrel{\scriptscriptstyle (-)}{\sigma}_{\mu\nu}:=\left\{\begin{array}{l} \sigma^{ij}=\frac{1}{2}\varepsilon^{ijk}\sigma_k\\ \sigma^{i0}=\left(-\right)\frac{1}{2}\sigma^i\end{array}\left i,j,k=1,2,3\right.\right.$
mielke-geometrodynamics_00928 (14.1.9)	304	■ 0.995	$\begin{aligned} \Sigma_{\mu\nu}^{\mathrm{YM}}&=\frac{1}{\alpha_g}\mathrm{Tr}\left(F_{\mu\alpha}F_{\nu}^{\alpha}-\frac{1}{4}g_{\mu\nu}F_{\alpha\beta}F^{\alpha\beta}\right)\\ &=\frac{1}{2\alpha_g}\mathrm{Tr}\left(F_{\mu\alpha}F_{\nu}^{\alpha}-(-1)^s{}^*F_{\mu\alpha}{}^*F_{\nu}^{\alpha}\right), \end{aligned}$	$\Sigma_{\mu\nu}^{\mathrm{YM}}=\frac{1}{\alpha_g}\mathrm{Tr}\left(F_{\mu\alpha}F_{\nu}^{\alpha}-\frac{1}{4}g_{\mu\nu}F_{\alpha\beta}F^{\alpha\beta}\right)\\ =\frac{1}{2\alpha_g}\mathrm{Tr}\left(F_{\mu\alpha}F_{\nu}^{\alpha}-(-1)^s{}^*F_{\mu\alpha}{}^*F_{\nu}^{\alpha}\right),$	$\Sigma_{\mu\nu}^{\mathrm{YM}}=\frac{1}{\alpha_g}\mathrm{Tr}(F_{\mu\alpha}F_{\nu}^{\alpha}-\frac{1}{4}g_{\mu\nu}F_{\alpha\beta}F^{\alpha\beta})\\ =\frac{1}{2\alpha_g}\mathrm{Tr}(F_{\mu\alpha}F_{\nu}^{\alpha}-(-1)^s{}^*F_{\mu\alpha}{}^*F_{\nu}^{\alpha}),$
mielke-geometrodynamics_00806 (2.2.1)	33	■ 0.996	$\left.e_{\beta}\right]\vartheta^{\alpha}=\delta^{\alpha}_{\beta}.$	$e_{\beta}\rfloor\vartheta^{\alpha}=\delta^{\alpha}_{\beta}.$	$e_{\beta}\rfloor\vartheta^{\alpha}=\delta^{\alpha}_{\beta}.$
mielke-geometrodynamics_00284 (4.4.13)	99	■ 0.996	$D[\Theta,\vartheta]=[D\Theta,\vartheta]+[\Theta,D\vartheta]$ $=[[\vartheta,\Omega^g],\vartheta]+[\Theta,\Theta]$ $=\vartheta\wedge\Omega^g\wedge\vartheta-\Omega^g\wedge\vartheta\wedge\vartheta+\vartheta\wedge\vartheta\wedge\Omega^g-\vartheta\wedge\Omega^g\wedge\vartheta$ $=[\vartheta\wedge\vartheta,\Omega^g].$	$D[\Theta,\vartheta]=[D\Theta,\vartheta]+[\Theta,D\vartheta]$ $=[[\vartheta,\Omega^g],\vartheta]+[\Theta,\Theta]$ $=\vartheta\wedge\Omega^g\wedge\vartheta-\Omega^g\wedge\vartheta\wedge\vartheta+\vartheta\wedge\vartheta\wedge\Omega^g-\vartheta\wedge\Omega^g\wedge\vartheta$ $=[\vartheta\wedge\vartheta,\Omega^g].$	$D[\Theta,\vartheta]=[D\Theta,\vartheta]+[\Theta,D\vartheta]$ $=[[\vartheta,\Omega^g],\vartheta]+[\Theta,\Theta]$ $=\vartheta\wedge\Omega^g\wedge\vartheta-\Omega^g\wedge\vartheta\wedge\vartheta+\vartheta\wedge\vartheta\wedge\Omega^g-\vartheta\wedge\Omega^g\wedge\vartheta$ $=[\vartheta\wedge\vartheta,\Omega^g].$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_ EQ0602 (9.2.1)	198	■ 0.997	$\begin{aligned} p_1(M^4) &= \frac{1}{2(2\pi)^2} \int_{M^4} \mathrm{Tr}(\Omega^L \wedge \Omega^L) \\ &= \frac{(-1)^{s/2}}{(4\pi)^2} \int_{M^4} R_{\alpha\beta\gamma\delta}^* R^{\alpha\beta\gamma\delta} \sqrt{ g } d^4x \end{aligned}$	$p_1(M^4) = \frac{1}{2(2\pi)^2} \int_{M^4} \mathrm{Tr}(\Omega^L \wedge \Omega^L) = \frac{(-1)^{s/2}}{(4\pi)^2} \int_{M^4} R_{\alpha\beta\gamma\delta}^* R^{\alpha\beta\gamma\delta} \sqrt{ g } d^4x$	$p_1(M^4) = \frac{1}{2(2\pi)^2} \int_{M^4} \mathrm{Tr}(\Omega^L \wedge \Omega^L) = \frac{(-1)^{s/2}}{(4\pi)^2} \int_{M^4} R_{\alpha\beta\gamma\delta}^* R^{\alpha\beta\gamma\delta} \sqrt{ g } d^4x$
mielke-geometrodynamics_ EQ0717 (10.9.3)	229	■ 0.997	$\Sigma_{\alpha} := \frac{\delta L_{\Psi}}{\delta \vartheta^{\alpha}} = \frac{\partial L_{\Psi}}{\partial \vartheta_{\alpha}} + D \frac{\partial L_{\Psi}}{\partial T^{\alpha}},$	$\Sigma_{\alpha} := \frac{\delta L_{\Psi}}{\delta \vartheta^{\alpha}} = \frac{\partial L_{\Psi}}{\partial \vartheta_{\alpha}} + D \frac{\partial L_{\Psi}}{\partial T^{\alpha}},$	$\Sigma_{\alpha} := \frac{\delta L_{\Psi}}{\delta \vartheta^{\alpha}} = \frac{\partial L_{\Psi}}{\partial \vartheta_{\alpha}} + D \frac{\partial L_{\Psi}}{\partial T^{\alpha}},$
mielke-geometrodynamics_ EQ0766 (11.1.33)	243	■ 0.997	$\stackrel{*}{\sigma} \tilde{\omega} = \frac{1}{2} \omega^{\alpha\beta} \sigma_{\alpha\beta} = \frac{1}{2} \Gamma_i^{\alpha\beta} \sigma_{\alpha\beta} \otimes dx^i =: \Gamma_i dx^i.$	$\stackrel{*}{\sigma} \tilde{\omega} = \frac{1}{2} \omega^{\alpha\beta} \sigma_{\alpha\beta} = \frac{1}{2} \Gamma_i^{\alpha\beta} \sigma_{\alpha\beta} \otimes dx^i =: \Gamma_i dx^i.$	$\stackrel{*}{\sigma} \tilde{\omega} = \frac{1}{2} \omega^{\alpha\beta} \sigma_{\alpha\beta} = \frac{1}{2} \Gamma_i^{\alpha\beta} \sigma_{\alpha\beta} \otimes dx^i =: \Gamma_i dx^i.$
mielke-geometrodynamics_ EQ1003 (14.3.16)	323	■ 0.997	$D D = \frac{1}{2} \{D, D\} + \frac{1}{2} [D, D] = \frac{1}{2} \{D, D\} + \frac{1}{2} \tilde{\Omega}$	$D D = \frac{1}{2} \{D, D\} + \frac{1}{2} [D, D] = \frac{1}{2} \{D, D\} + \frac{1}{2} \tilde{\Omega}$	$D D = \frac{1}{2} \{D, D\} + \frac{1}{2} [D, D] = \frac{1}{2} \{D, D\} + \frac{1}{2} \tilde{\Omega}$
mielke-geometrodynamics_ EQ0225 (4.2.18)	87	■ 0.997	$\omega^T = \tilde{\vartheta} - D \tilde{\varphi}$	$\omega^T = \tilde{\vartheta} - D \tilde{\varphi}$	$\omega^T = \tilde{\vartheta} - D \tilde{\varphi}$
mielke-geometrodynamics_ EQ0048 (2.6.11)	43	■ 0.997	$D D D b \equiv 0.$	$D D D b \equiv 0.$	$D D D b \equiv 0.$
mielke-geometrodynamics_ EQ0099 (3.2.9)	56	■ 0.997	$\begin{aligned} \{ \}^{\Delta} \omega &:= \{ \}^{\Delta} \omega - \omega = G^{-1} [\omega, G] - G^{-1} dG \\ &= -G^{-1} DG = -G^{-1} D. \end{aligned}$	$\Delta \omega := G^{-1} \omega - \omega = G^{-1} [\omega, G] - G^{-1} dG = -G^{-1} DG = -G^{-1} D.$	$\Delta \omega := G^{-1} \omega - \omega = G^{-1} [\omega, G] - G^{-1} dG = -G^{-1} DG = -G^{-1} D.$
mielke-geometrodynamics_ EQ0219 (4.2.12)	86	■ 0.997	$\tilde{\tilde{\Omega}} \rightarrow^{A^{-1}} \tilde{\tilde{\Omega}} = A^{-1} \tilde{\tilde{\Omega}} A$	$\tilde{\tilde{\Omega}} \rightarrow^{A^{-1}} \tilde{\tilde{\Omega}} = A^{-1} \tilde{\tilde{\Omega}} A$	$\tilde{\tilde{\Omega}} \rightarrow^{A^{-1}} \tilde{\tilde{\Omega}} = A^{-1} \tilde{\tilde{\Omega}} A$
mielke-geometrodynamics_ EQ0411 (6.4.31)	136	■ 0.997	$n_{\mathrm{NY}} := \int_{R^4} dC_{\mathrm{TT}} = \int_{S^3_{\infty}} \underline{C}_{\mathrm{TT}} = 3 \mathrm{Vol}(S^3) k = 6\pi^2 k.$	$n_{\mathrm{NY}} := \int_{R^4} dC_{\mathrm{TT}} = \int_{S^3_{\infty}} \underline{C}_{\mathrm{TT}} = 3 \mathrm{Vol}(S^3) k = 6\pi^2 k.$	$n_{\mathrm{NY}} := \int_{R^4} dC_{\mathrm{TT}} = \int_{S^3_{\infty}} \underline{C}_{\mathrm{TT}} = 3 \mathrm{Vol}(S^3) k = 6\pi^2 k.$
mielke-geometrodynamics_ EQ0492 (7.7.10)	161	■ 0.997	$H_{\alpha\beta} := -\frac{\partial L}{\partial R^{\alpha\beta}} = \tilde{H}_{\alpha\beta} + \frac{1}{2} R^{\mu\nu} \wedge (\partial \tilde{H}_{\mu\nu} / \partial R^{\alpha\beta})$	$H_{\alpha\beta} := -\frac{\partial L}{\partial R^{\alpha\beta}} = \tilde{H}_{\alpha\beta} + \frac{1}{2} R^{\mu\nu} \wedge (\partial \tilde{H}_{\mu\nu} / \partial R^{\alpha\beta})$	$H_{\alpha\beta} := -\frac{\partial L}{\partial R^{\alpha\beta}} = \tilde{H}_{\alpha\beta} + \frac{1}{2} R^{\mu\nu} \wedge (\partial \tilde{H}_{\mu\nu} / \partial R^{\alpha\beta})$
mielke-geometrodynamics_ EQ0877 (13.3.6)	285	■ 0.997	$\left[P_{\alpha}, P_{*}^{\beta} \right] = \frac{1}{\ell^2} L^{\beta}{}_{\alpha}.$	$\left[P_{\alpha}, P_{*}^{\beta} \right] = \frac{1}{\ell^2} L^{\beta}{}_{\alpha}.$	$\left[P_{\alpha}, P_{*}^{\beta} \right] = \frac{1}{\ell^2} L^{\beta}{}_{\alpha}.$
mielke-geometrodynamics_ EQ0962 (14.2.2)	312	■ 0.997	$\stackrel{\circ}{\vartheta} = \varphi_j d\xi^j,$	$\stackrel{\circ}{\vartheta} = \varphi_j d\xi^j,$	$\stackrel{\circ}{\vartheta} = \varphi_j d\xi^j,$
mielke-geometrodynamics_ EQ1060 (16.1.1)	354	■ 0.997	$R_{ij}^f - \frac{1}{2} f_{ij} (R^f - 2\Lambda_{\mathrm{eff}}) = \ell^2 T_{ij}$	$R_{ij}^f - \frac{1}{2} f_{ij} (R^f - 2\Lambda_{\mathrm{eff}}) = \ell^2 T_{ij}$	$R_{ij}^f - \frac{1}{2} f_{ij} (R^f - 2\Lambda_{\mathrm{eff}}) = \ell^2 T_{ij}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_315 EQ0976 (14.2.17)		■ 0.997	$\begin{aligned} \hat{L}_{\mathrm{W}.} &= \frac{1}{2} \operatorname{Tr}(\hat{\Omega} \wedge \underbrace{\hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}}_{2+K \text{ termms}}) \\ &= \hat{L}_{\mathrm{W}.}^g + \hat{L}_{\mathrm{W}.}^G + \ell^{*2} \hat{L}_{\parallel} - \frac{1}{2} d \operatorname{Tr}(\hat{\alpha} \wedge \hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}) \end{aligned}$	$\hat{L}_{\mathrm{W}.} = \frac{1}{2} \operatorname{Tr}(\hat{\Omega} \wedge \underbrace{\hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}}_{2+K \text{ termms}})$ $= \hat{L}_{\mathrm{W}.}^g + \hat{L}_{\mathrm{W}.}^G + \ell^{*2} \hat{L}_{\parallel} - \frac{1}{2} d \operatorname{Tr}(\hat{\alpha} \wedge \hat{\vartheta} \wedge \dots \wedge \hat{\vartheta})$	$\hat{L}_{\mathrm{W}.} = \frac{1}{2} \operatorname{Tr}(\hat{\Omega} \wedge \underbrace{\hat{\vartheta} \wedge \dots \wedge \hat{\vartheta}}_{2+K \text{ termms}})$ $= \hat{L}_{\mathrm{W}.}^g + \hat{L}_{\mathrm{W}.}^G + \ell^{*2} \hat{L}_{\parallel} - \frac{1}{2} d \operatorname{Tr}(\hat{\alpha} \wedge \hat{\vartheta} \wedge \dots \wedge \hat{\vartheta})$
mielke-geometrodynamics_153 EQ0466 (7.3.5)		■ 0.997	$G_{\alpha}^{\{\}} := \frac{1}{2} R^{\{\} \beta \gamma} \wedge \eta_{\alpha \beta \gamma}, \quad G_{ij} := \operatorname{Ric}_{ij}^{\{\}} - \frac{1}{2} g_{ij},$	$G_{\alpha}^{\{\}} := \frac{1}{2} R^{\{\} \beta \gamma} \wedge \eta_{\alpha \beta \gamma}, \quad G_{ij} := \operatorname{Ric}_{ij}^{\{\}} - \frac{1}{2} g_{ij},$	$G_{\alpha}^{\{\}} := \frac{1}{2} R^{\{\} \beta \gamma} \wedge \eta_{\alpha \beta \gamma}, \quad G_{ij} := \operatorname{Ric}_{ij}^{\{\}} - \frac{1}{2} g_{ij},$
mielke-geometrodynamics_58 EQ0105 (3.3.6)		■ 0.998	$H := -\frac{\partial L}{\partial dA} = -\frac{\partial L}{\partial F} \quad \text{and} \quad j := \frac{\delta L_{\mathrm{mat}}}{\delta A},$	$H := -\frac{\partial L}{\partial dA} = -\frac{\partial L}{\partial F} \quad \text{and} \quad j := \frac{\delta L_{\mathrm{mat}}}{\delta A},$	$H := -\frac{\partial L}{\partial dA} = -\frac{\partial L}{\partial F} \quad \text{and} \quad j := \frac{\delta L_{\mathrm{mat}}}{\delta A},$
mielke-geometrodynamics_63 EQ0136 (3.4.7)		■ 0.998	$\begin{aligned} \delta \Omega &= \delta(d\omega - \omega \wedge \omega) = \delta d\omega - \delta\omega \wedge \omega - \omega \wedge \delta\omega \\ &= d\delta\omega - [\omega, \delta\omega] = D\delta\omega. \end{aligned}$	$\delta \Omega = \delta(d\omega - \omega \wedge \omega) = \delta d\omega - \delta\omega \wedge \omega - \omega \wedge \delta\omega$ $= d\delta\omega - [\omega, \delta\omega] = D\delta\omega.$	$\delta \Omega = \delta(d\omega - \omega \wedge \omega) = \delta d\omega - \delta\omega \wedge \omega - \omega \wedge \delta\omega$ $= d\delta\omega - [\omega, \delta\omega] = D\delta\omega.$
mielke-geometrodynamics_131 EQ0390 (6.4.5)		■ 0.998	$\mathscr{H} = n \rfloor L - \ell_n \underline{\Psi} \wedge \left(\frac{\partial L}{\partial d\Psi} \right).$	$\mathscr{H} = n \rfloor L - \ell_n \underline{\Psi} \wedge \left(\frac{\partial L}{\partial d\Psi} \right).$	$\mathscr{H} = n \rfloor L - \ell_n \underline{\Psi} \wedge \left(\frac{\partial L}{\partial d\Psi} \right).$
mielke-geometrodynamics_195 EQ0594 (9.1.12)		■ 0.998	$\bar{\omega}^{\{\}} = 0.$	$\bar{\Omega}^{\{\}} = 0.$	$\bar{\Omega}^{\{\}} = 0.$
mielke-geometrodynamics_326 EQ1012 (14.4.7)		■ 0.998	$\begin{aligned} S_{\mathrm{A}} &:= 4\pi \left(r_+^2 + \frac{J(J+1)\hbar^2}{M^2 c^2} \right) \\ &= \frac{8\pi\hbar^2}{M^{*4} c^2} \left(M^2 + \frac{\alpha_e}{2} M^{*2} \operatorname{Tr} \bar{Q}^2 + M \sqrt{M^2 - J(J+1) \frac{M^{*4}}{M^2} - \alpha_e M^{*2} \operatorname{Tr} \bar{Q}^2} \right) \end{aligned}$	$S_{\mathrm{A}} := 4\pi \left(r_+^2 + \frac{J(J+1)\hbar^2}{M^2 c^2} \right)$ $= \frac{8\pi\hbar^2}{M^{*4} c^2} \left(M^2 + \frac{\alpha_e}{2} M^{*2} \operatorname{Tr} \bar{Q}^2 + M \sqrt{M^2 - J(J+1) \frac{M^{*4}}{M^2} - \alpha_e M^{*2} \operatorname{Tr} \bar{Q}^2} \right)$	$S_{\mathrm{A}} := 4\pi \left(r_+^2 + \frac{J(J+1)\hbar^2}{M^2 c^2} \right)$ $= \frac{8\pi\hbar^2}{M^{*4} c^2} \left(M^2 + \frac{\alpha_e}{2} M^{*2} \operatorname{Tr} \bar{Q}^2 + M \sqrt{M^2 - J(J+1) \frac{M^{*4}}{M^2} - \alpha_e M^{*2} \operatorname{Tr} \bar{Q}^2} \right)$
mielke-geometrodynamics_256 EQ0813 (11.4.26)		■ 0.998	$d \rho^{*} := e^{(\mu - \nu)/2} d \rho.$	$d \rho^{*} := e^{(\mu - \nu)/2} d \rho.$	$d \rho^{*} := e^{(\mu - \nu)/2} d \rho.$
mielke-geometrodynamics_49 EQ0073 (2.8.16)		■ 0.998	$\begin{aligned} &\delta \stackrel{\infty}{\omega} - \stackrel{\infty}{\omega} \wedge \stackrel{\infty}{\omega} \\ &= -d(G^{-1}) \wedge dG - G^{-1} ddG - (G^{-1} dG) \wedge (G^{-1} dG) \\ &= -d(G^{-1}) \wedge dG + G^{-1} G d(G^{-1}) \wedge dG = 0 \end{aligned}$	$d \stackrel{\infty}{\omega} - \stackrel{\infty}{\omega} \wedge \stackrel{\infty}{\omega}$ $= -d(G^{-1}) \wedge dG - G^{-1} ddG - (G^{-1} dG) \wedge (G^{-1} dG)$ $= -d(G^{-1}) \wedge dG + G^{-1} G d(G^{-1}) \wedge dG = 0$	$d \stackrel{\infty}{\omega} - \stackrel{\infty}{\omega} \wedge \stackrel{\infty}{\omega}$ $= -d(G^{-1}) \wedge dG - G^{-1} ddG - (G^{-1} dG) \wedge (G^{-1} dG)$ $= -d(G^{-1}) \wedge dG + G^{-1} G d(G^{-1}) \wedge dG = 0$
mielke-geometrodynamics_86 EQ0214 (4.2.6)		■ 0.998	$\mathscr{G}(n, \mathbb{R}) := C^{\infty}(A(M) \times_{\mathrm{Ad}} GL(n, \mathbb{R}))$	$\mathscr{G}(n, \mathbb{R}) := C^{\infty}(A(M) \times_{\mathrm{Ad}} GL(n, \mathbb{R}))$	$\mathscr{G}(n, \mathbb{R}) := C^{\infty}(A(M) \times_{\mathrm{Ad}} GL(n, \mathbb{R}))$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_373 E01126 (B.38)	373	■ 0.998	$\partial: C_p(M, \mathbb{R}) \rightarrow C_{p-1}(M, \mathbb{R}),$	$\partial: C_p(M, \mathbb{R}) \rightarrow C_{p-1}(M, \mathbb{R}),$	
mielke-geometrodynamics_166 E00516 (7.7.34)	166	■ 0.998	$\widetilde{T}=T-3\,d\varphi,\quad \mathscr{A}^\theta=\mathscr{A}-id\theta\,.$	$\widetilde{T}=T-3d\varphi,\quad \mathscr{A}^\theta=\mathscr{A}-id\theta\,.$	
mielke-geometrodynamics_313 E00967 (14.2.8)	313	■ 0.998	$d\stackrel{\circ}{s}^2=\stackrel{\circ}{g}_{ij}\,d\xi^i\otimes_s d\xi^j$	$d\stackrel{\circ}{s}^2=\stackrel{\circ}{g}_{ij}\,d\xi^i\otimes_s d\xi^j,$	
mielke-geometrodynamics_46 E00060 (2.8.1)	46	■ 0.999	$\operatorname{det}\!\left(1+\frac{i}{2\pi}\,\Omega\right)=\stackrel{*}{\operatorname{det}}\!\left(1+\gamma_1+\cdots+\gamma_m\right)$	$\det\left(1+\frac{i}{2\pi}\Omega\right)=\overset{*}{\pi}\left(1+\gamma_1+\cdots+\gamma_m\right)$	
mielke-geometrodynamics_48 E00071 (2.8.14)	48	■ 0.999	$\stackrel{\infty}{\omega}:= -G^{-1}dG,$	$\overset{\infty}{\omega}:= -G^{-1}dG,$	
mielke-geometrodynamics_89 E00235 (4.2.28)	89	■ 0.999	$Q_{\kappa\mu\nu}:= -D_{\kappa}\,g_{\mu\nu}$	$Q_{\kappa\mu\nu}:= -D_{\kappa}\,g_{\mu\nu}$	
mielke-geometrodynamics_97 E00278 (4.4.6)	97	■ 0.999	$\begin{aligned} &\& D\,\Pi^{\mathrm{T}}-E=\Sigma \\ &\& \Pi^{\mathrm{L}}+\Pi_{\square}^{\mathrm{T}}=\tau_{\mathrm{s}} \end{aligned}$	$\begin{aligned} D\Pi^{\mathrm{T}}-E&=\Sigma \\ D\Pi^{\mathrm{L}}+\Pi_{\square}^{\mathrm{T}}&=\tau_{\mathrm{s}} \end{aligned}$	
mielke-geometrodynamics_105 E00313 (4.6.5)	105	■ 0.999	$\Phi(A,\gamma)=P\,\exp\left[(i/\hbar)\oint A\right],$	$\Phi(A,\gamma)=P\,\exp[(i/\hbar)\oint A],$	
mielke-geometrodynamics_125 E00358 (6.2.9)	125	■ 0.999	$L_{\parallel}:= -\frac{1}{2\kappa}T^{\alpha}\wedge^{*}\left({}^{(1)}T_{\alpha}-2{}^{(2)}T_{\alpha}-\frac{1}{2}{}^{(3)}T_{\alpha}\right),$	$L_{\parallel}:= -\frac{1}{2\kappa}T^{\alpha}\wedge^{*}\left({}^{(1)}T_{\alpha}-2{}^{(2)}T_{\alpha}-\frac{1}{2}{}^{(3)}T_{\alpha}\right),$	
mielke-geometrodynamics_152 E00461 (7.2.13)	152	■ 0.999	$\stackrel{(\mp)}{\Omega}_{\mathrm{C}}=0$	$\overset{(\mp)}{\Omega}_{\mathrm{C}}=0$	
mielke-geometrodynamics_209 E00620 (10.1.8)	209	■ 0.999	$\frac{\delta C_{\mathrm{T}}}{\delta \vartheta^{\alpha}}=\frac{1}{\ell^2}T_{\alpha}\quad\text{and}\quad\frac{\delta C_{\mathrm{T}}}{\delta \Gamma^{\star\alpha}}=\frac{(-1)^s}{\ell^2}\eta_{\alpha},$	$\frac{\delta C_{\mathrm{T}}}{\delta \vartheta^{\alpha}}=\frac{1}{\ell^2}T_{\alpha}\quad\text{and}\quad\frac{\delta C_{\mathrm{T}}}{\delta \Gamma^{\star\alpha}}=\frac{(-1)^s}{\ell^2}\eta_{\alpha},$	
mielke-geometrodynamics_218 E00667 (10.3.24)	218	■ 0.999	$\left[\square+(-1)^s m^2\right]\vartheta^{\alpha}\cong 0,$	$\left[\square+(-1)^s m^2\right]\vartheta^{\alpha}\cong 0,$	
mielke-geometrodynamics_220 E00678 (10.3.35)	220	■ 0.999	$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}}-\check{\mathcal{E}}_{\alpha}=\frac{\theta_{\mathrm{TL}}}{\ell}d\Gamma_{\alpha}^{\star},$	$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}}-\check{\mathcal{E}}_{\alpha}=\frac{\theta_{\mathrm{TL}}}{\ell}d\Gamma_{\alpha}^{\star},$	
mielke-geometrodynamics_238 E00740 (11.1.7)	238	■ 0.999	$XX=-\mathrm{o}_{\alpha\beta}x^{\alpha}x^{\beta}\mathbb{1}$	$XX=-\mathrm{o}_{\alpha\beta}x^{\alpha}x^{\beta}\mathbb{1}$	
mielke-geometrodynamics_246 E00776 (11.2.5)	246	■ 0.999	$L_{\mathrm{D}}=\frac{i}{2}\left(\bar{\psi}\gamma\wedge^{*}d\psi+d\bar{\psi}\wedge^{*}\gamma\psi\right)-m\bar{\psi}\psi\eta-\frac{1}{2}\bar{\psi}\left[\widetilde{\omega},*\gamma\right]\psi,$	$L_{\mathrm{D}}=\frac{i}{2}\left(\bar{\psi}\gamma\wedge^{*}d\psi+d\bar{\psi}\wedge^{*}\gamma\psi\right)-m\bar{\psi}\psi\eta-\frac{1}{2}\bar{\psi}\left[\widetilde{\omega},*\gamma\right]\psi,$	
mielke-geometrodynamics_255 E00805 (11.4.17)	255	■ 0.999	$\kappa=\mp\left(j+\frac{1}{2}\right)\quad\text{for}\quad j=l\pm\frac{1}{2}.$	$\kappa=\mp\left(j+\frac{1}{2}\right)\quad\text{for}\quad j=l\pm\frac{1}{2}.$	
mielke-geometrodynamics_320 E00989 (14.3.2)	320	■ 0.999	$\hat{L}_{\mathrm{D}}.=\frac{i}{2}\left(\overline{\hat{\psi}}\hat{\gamma}\wedge^{*}\hat{D}\hat{\psi}+\overline{\left(\hat{D}\hat{\psi}\right)}\wedge^{*}\hat{\gamma}\hat{\psi}\right)-\hat{m}\bar{\hat{\psi}}\hat{\psi}\eta$	$\hat{L}_{\mathrm{D}}.=\frac{i}{2}\left(\bar{\hat{\psi}}\hat{\gamma}\wedge^{*}\hat{D}\hat{\psi}+\overline{\left(\hat{D}\hat{\psi}\right)}\wedge^{*}\hat{\gamma}\hat{\psi}\right)-\hat{m}\bar{\hat{\psi}}\hat{\psi}\eta$	

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ1057 (15.2.14)	346	■ 0.999	$i\,\bar{\gamma}\wedge^*\bar{D}^{\bar{\mathfrak{U}}}\psi-6\ell^2\bar{\gamma}^5\bar{\gamma}\wedge^*(\bar{\psi}\bar{\gamma}^5\bar{\gamma}\psi)\,\psi-m\psi\eta=0$	$i\,\bar{\gamma}\wedge^*\bar{D}^{\bar{\mathfrak{U}}}\psi-6\ell^2\bar{\gamma}^5\bar{\gamma}\wedge^*(\bar{\psi}\bar{\gamma}^5\bar{\gamma}\psi)\psi-m\psi\eta=0$	
mielke-geometrodynamics_EQ0669 (10.3.26)	219	■ 0.999	$m=\frac{\theta_{\mathrm{T}}\theta_{\mathrm{TL}}}{2(\theta_{\mathrm{TL}}^2+\theta_{\mathrm{T}}\theta_{\mathrm{L}})}\ell,\quad\Lambda_{\mathrm{ind}}=\frac{\theta_{\mathrm{T}}^2\left[-(-1)^s3\theta_{\mathrm{TL}}^2+8\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right]}{4\left[(-1)^s\theta_{\mathrm{TL}}^2-2\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right]^2\ell^2},$	$m=\frac{\theta_{\mathrm{T}}\theta_{\mathrm{TL}}}{2(\theta_{\mathrm{TL}}^2+\theta_{\mathrm{T}}\theta_{\mathrm{L}})\ell},\quad\Lambda_{\mathrm{ind}}=\frac{\theta_{\mathrm{T}}^2[-(-1)^s3\theta_{\mathrm{TL}}^2+8\theta_{\mathrm{T}}\theta_{\mathrm{L}}]}{4[(-1)^s\theta_{\mathrm{TL}}^2-2\theta_{\mathrm{T}}\theta_{\mathrm{L}}]^2\ell^2},$	
mielke-geometrodynamics_EQ0929 (14.1.10)	304	■ 0.999	$\begin{aligned} &\operatorname{Tr}\left(\Sigma_{\mathrm{YM}}\wedge^*\vartheta\right)=\\ &\operatorname{Tr}\left(\vartheta\wedge^*\Sigma_{\mathrm{YM}}\right)\quad\&=\frac{1}{2\alpha_g}\operatorname{Tr}\left\{\vartheta\wedge\Omega\wedge^*(\vartheta\wedge^*\Omega)+\vartheta\wedge^*\vartheta\wedge^*(\Omega\wedge^*\Omega)\right\}=0 \end{aligned}$	$\begin{aligned} &\operatorname{Tr}\left(\Sigma_{\mathrm{YM}}\wedge^*\vartheta\right)=\operatorname{Tr}\left(\vartheta\wedge^*\Sigma_{\mathrm{YM}}\right)\\ &=\frac{1}{2\alpha_g}\operatorname{Tr}\left\{\vartheta\wedge\Omega\wedge^*(\vartheta\wedge^*\Omega)+\vartheta\wedge^*\vartheta\wedge^*(\Omega\wedge^*\Omega)\right\}=0 \end{aligned}$	$\begin{aligned} &\operatorname{Tr}(\Sigma_{\mathrm{YM}}\wedge^*\vartheta)=\operatorname{Tr}(\vartheta\wedge^*\Sigma_{\mathrm{YM}})\\ &=\frac{1}{2\alpha_g}\operatorname{Tr}\left\{\vartheta\wedge\Omega\wedge^*(\vartheta\wedge^*\Omega)+\vartheta\wedge^*\vartheta\wedge^*(\Omega\wedge^*\Omega)\right\}=0 \end{aligned}$
mielke-geometrodynamics_EQ0093 (3.2.3)	55	■ 0.999	$\omega^{\prime}\rightarrow^{G^{-1}}\omega^{\prime}=G^{-1}\omega^{\prime}G+\left(dG^{-1}\right)G$	$\omega^{\prime}\rightarrow^{G^{-1}}\omega^{\prime}=G^{-1}\omega^{\prime}G+\left(dG^{-1}\right)G$	
mielke-geometrodynamics_EQ0113 (3.3.14)	59	■ 0.999	$H=\left\{\right\}^*\!F\;.$	$H=\left\{\right\}^*\!F.$	$H=\left\{\right\}^*\!F\;.$
mielke-geometrodynamics_EQ0173 (3.7.1)	71	■ 0.999	$L_{\mathrm{H}}=-\bar{\Theta}^{(1)}\wedge^*D\phi^0+\bar{\Theta}^{(1)}\wedge^*\Theta^{(1)}+\frac{\lambda}{4}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)^2\eta.$	$L_{\mathrm{H}}=-\bar{\Theta}^{(1)}\wedge^*D\phi^0+\bar{\Theta}^{(1)}\wedge^*\Theta^{(1)}+\frac{\lambda}{4}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)^2\eta.$	$L_{\mathrm{H}}=-\bar{\Theta}^{(1)}\wedge^*D\phi^0+\bar{\Theta}^{(1)}\wedge^*\Theta^{(1)}+\frac{\lambda}{4}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)^2\eta.$
mielke-geometrodynamics_EQ0201 (4.1.15)	83	■ 0.999	$\delta_g^*\vartheta=g^{-1}\vartheta,\quad g\in GL(n,\mathbb{R}),$	$\delta_g^*\vartheta=g^{-1}\vartheta,\quad g\in GL(n,\mathbb{R}),$	
mielke-geometrodynamics_EQ0275 (4.4.3)	97	■ 0.999	$\begin{aligned} &d(\delta\vartheta)-[\omega^{\mathrm{L}},\delta\vartheta]-[\delta\omega^{\mathrm{L}},\vartheta] \\ &=D(\delta\vartheta)-[\delta\omega^{\mathrm{L}},\vartheta]. \end{aligned}$	$\begin{aligned} &=d(\delta\vartheta)-[\omega^{\mathrm{L}},\delta\vartheta]-[\delta\omega^{\mathrm{L}},\vartheta] \\ &=D(\delta\vartheta)-[\delta\omega^{\mathrm{L}},\vartheta]. \end{aligned}$	$\begin{aligned} &=d(\delta\vartheta)-[\omega^{\mathrm{L}},\delta\vartheta]-[\delta\omega^{\mathrm{L}},\vartheta] \\ &=D(\delta\vartheta)-[\delta\omega^{\mathrm{L}},\vartheta]. \end{aligned}$
mielke-geometrodynamics_EQ0341 (5.3.18)	115	■ 0.999	$\left\{\right\}^*\!\mathscr{A}=\frac{\kappa}{2}j_5,$	$\left\{\right\}^*\!\mathscr{A}=\frac{\kappa}{2}j_5,$	$\left\{\right\}^*\!\mathscr{A}=\frac{\kappa}{2}j_5,$
mielke-geometrodynamics_EQ0441 (6.8.12)	143	■ 0.999	$D\tau_{\alpha\beta}+\vartheta_{[\alpha}\wedge\Sigma_{\beta]}\cong0,$	$D\tau_{\alpha\beta}+\vartheta_{[\alpha}\wedge\Sigma_{\beta]}\cong0,$	$D\tau_{\alpha\beta}+\vartheta_{[\alpha}\wedge\Sigma_{\beta]}\cong0,$
mielke-geometrodynamics_EQ0476 (7.4.4)	156	■ 0.999	$C_{\alpha\beta}=0\quad\&\text{ and }\quad R=0,$	$C_{\alpha\beta}=0\quad\text{ and }\quad R=0,$	$C_{\alpha\beta}=0\quad\text{ and }\quad R=0,$
mielke-geometrodynamics_EQ0519 (7.7.37)	166	■ 0.999	$D\Theta\equiv[\Omega^g,\gamma],\quad D\Omega^g\equiv0,$	$D\Theta\equiv[\Omega^g,\gamma],\quad D\Omega^g\equiv0,$	$D\Theta\equiv[\Omega^g,\gamma],\quad D\Omega^g\equiv0,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_183 EQ0567 (8.6.4)	183	■ 0.999	$\begin{aligned} H_{\alpha\beta}^{(*)} &= \theta_L R_{\alpha\beta} + \theta_L^* R_{\alpha\beta}^{(*)} + \frac{\theta_T}{2\ell^2} \eta_{\alpha\beta} - \frac{\theta_T}{2\ell^2} \vartheta_\alpha \wedge \vartheta_\beta \\ &\simeq -\partial L_\theta / \partial R^{\alpha\beta}, \end{aligned}$	$H_{\alpha\beta}^{(*)} = \theta_L R_{\alpha\beta} + \theta_L^* R_{\alpha\beta}^{(*)} + \frac{\theta_T}{2\ell^2} \eta_{\alpha\beta} - \frac{\theta_T}{2\ell^2} \vartheta_\alpha \wedge \vartheta_\beta$ $\simeq -\partial L_\theta / \partial R^{\alpha\beta}$	$H_{\alpha\beta}^{(*)} = \theta_L R_{\alpha\beta} + \theta_L^* R_{\alpha\beta}^{(*)} + \frac{\theta_T}{2\ell^2} \eta_{\alpha\beta} - \frac{\theta_T}{2\ell^2} \vartheta_\alpha \wedge \vartheta_\beta$ $\simeq -\partial L_\theta / \partial R^{\alpha\beta},$
mielke-geometrodynamics_210 EQ0624 (10.1.12)	210	■ 0.999	$D H_{\alpha} - E_{\alpha} = \Sigma_{\alpha},$	$D H_{\alpha} - E_{\alpha} = \Sigma_{\alpha},$	$D H_{\alpha} - E_{\alpha} = \Sigma_{\alpha},$
mielke-geometrodynamics_220 EQ0673 (10.3.30)	220	■ 0.999	$H_{\alpha} = -\frac{\theta_T}{2\ell^2} \vartheta_{\alpha} - \frac{\theta_{TL}}{\ell} \Gamma_{\alpha}^*, \quad H_{\alpha}^* = -\frac{\theta_L}{2} \Gamma_{\alpha}^*,$	$H_{\alpha} = -\frac{\theta_T}{2\ell^2} \vartheta_{\alpha} - \frac{\theta_{TL}}{\ell} \Gamma_{\alpha}^*, \quad H_{\alpha}^* = -\frac{\theta_L}{2} \Gamma_{\alpha}^*,$	$H_{\alpha} = -\frac{\theta_T}{2\ell^2} \vartheta_{\alpha} - \frac{\theta_{TL}}{\ell} \Gamma_{\alpha}^*, \quad H_{\alpha}^* = -\frac{\theta_L}{2} \Gamma_{\alpha}^*,$
mielke-geometrodynamics_230 EQ0722 (10.10.2)	230	■ 0.999	$\delta L = \delta \vartheta^{\alpha} \wedge \frac{\delta L}{\delta \vartheta^{\alpha}} + \delta \Gamma_{\alpha}^* \wedge \frac{\delta L}{\delta \Gamma_{\alpha}^*} + \delta \bar{\Psi} \wedge \frac{\delta L}{\delta \bar{\Psi}},$	$\delta L = \delta \vartheta^{\alpha} \wedge \frac{\delta L}{\delta \vartheta^{\alpha}} + \delta \Gamma_{\alpha}^* \wedge \frac{\delta L}{\delta \Gamma_{\alpha}^*} + \delta \bar{\Psi} \wedge \frac{\delta L}{\delta \bar{\Psi}},$	$\delta L = \delta \vartheta^{\alpha} \wedge \frac{\delta L}{\delta \vartheta^{\alpha}} + \delta \Gamma_{\alpha}^* \wedge \frac{\delta L}{\delta \Gamma_{\alpha}^*} + \delta \bar{\Psi} \wedge \frac{\delta L}{\delta \bar{\Psi}},$
mielke-geometrodynamics_242 EQ0759 (11.1.26)	242	■ 0.999	$\{ \}^* (\underbrace{\gamma \wedge \cdots \wedge \gamma}_{p}) = \frac{p!}{(n-p)!} \gamma^{\wedge n+1} \underbrace{\gamma \wedge \cdots \wedge \gamma}_{n-p}.$	$*(\underbrace{\gamma \wedge \cdots \wedge \gamma}_p) = \frac{p!}{(n-p)!} \gamma^{\wedge n+1} \underbrace{\gamma \wedge \cdots \wedge \gamma}_{n-p}.$	$*(\underbrace{\gamma \wedge \cdots \wedge \gamma}_p) = \frac{p!}{(n-p)!} \gamma^{\wedge n+1} \underbrace{\gamma \wedge \cdots \wedge \gamma}_{n-p}.$
mielke-geometrodynamics_247 EQ0780 (11.2.8)	247	■ 0.999	$(\psi, \chi) := \int_{H^{n-1}} \tau_s(\psi, \chi) = i \int_{H^{n-1}} \psi^* \gamma \chi$	$(\psi, \chi) := \int_{H^{n-1}} \tau_s(\psi, \chi) = i \int_{H^{n-1}} \psi^* \gamma \chi$	$(\psi, \chi) := \int_{H^{n-1}} \tau_s(\psi, \chi) = i \int_{H^{n-1}} \psi^* \gamma \chi$
mielke-geometrodynamics_251 EQ0789 (11.4.1)	251	■ 0.999	$E_r = \sqrt{\alpha} r_{\circ}^2 \left[1 + \left(\frac{r}{r_{\circ}} \right)^4 \right]^{-1/2} \sim \sqrt{\alpha} \frac{1}{r^2}$	$E_r = \sqrt{\alpha} r_{\circ}^2 \left[1 + \left(\frac{r}{r_{\circ}} \right)^4 \right]^{-1/2} \sim \sqrt{\alpha} \frac{1}{r^2}$	$E_r = \sqrt{\alpha} r_{\circ}^2 \left[1 + \left(\frac{r}{r_{\circ}} \right)^4 \right]^{-1/2} \sim \sqrt{\alpha} \frac{1}{r^2}$
mielke-geometrodynamics_259 EQ0825 (11.4.39)	259	■ 0.999	$m_{\pi} = \sqrt{1 - \omega^2} m$	$m_{\pi} = \sqrt{1 - \omega^2} m$	$m_{\pi} = \sqrt{1 - \omega^2} m$
mielke-geometrodynamics_287 EQ0892 (13.4.11)	287	■ 0.999	$\vartheta \xrightarrow{A^{-1}(x)} \vartheta' = \Lambda^{-1}(x) \vartheta, \quad \theta \xrightarrow{A^{-1}(x)} \theta' = \theta \Lambda^{-1}(x).$	$\vartheta \xrightarrow{A^{-1}(x)} \vartheta' = \Lambda^{-1}(x) \vartheta, \quad \theta \xrightarrow{A^{-1}(x)} \theta' = \theta \Lambda^{-1}(x).$	$\vartheta \xrightarrow{A^{-1}(x)} \vartheta' = \Lambda^{-1}(x) \vartheta, \quad \theta \xrightarrow{A^{-1}(x)} \theta' = \theta \Lambda^{-1}(x).$
mielke-geometrodynamics_304 EQ0932 (14.1.13)	304	■ 0.999	$e^{*\delta} \circ e^{*\varepsilon} = e^{*\varepsilon} \circ e^{*\delta} = e^{*(\delta+\varepsilon)},$	$e^{*\delta} \circ e^{*\varepsilon} = e^{*\varepsilon} \circ e^{*\delta} = e^{*(\delta+\varepsilon)},$	$e^{*\delta} \circ e^{*\varepsilon} = e^{*\varepsilon} \circ e^{*\delta} = e^{*(\delta+\varepsilon)},$
mielke-geometrodynamics_305 EQ0940 (14.1.21)	305	■ 0.999	$\xi \wedge \xi = \xi \wedge^* \xi = 0$	$\xi \wedge \xi = \xi \wedge^* \xi = 0$	$\xi \wedge \xi = \xi \wedge^* \xi = 0$
mielke-geometrodynamics_307 EQ0951 (14.1.33)	307	■ 0.999	$\begin{aligned} D^* \xi - (-1)^z \xi \wedge D \delta &= \bar{\tau}_e \\ D \xi + (-1)^z \xi \wedge D \delta &= \bar{\tau}_p \end{aligned}$	$\begin{aligned} D^* \xi - (-1)^z \xi \wedge D \delta &= \bar{\tau}_e \\ D \xi + (-1)^z \xi \wedge D \delta &= \bar{\tau}_p \end{aligned}$	$\begin{aligned} D^* \xi - (-1)^z \xi \wedge D \delta &= \bar{\tau}_e, \\ D \xi + (-1)^z \xi \wedge D \delta &= \bar{\tau}_p. \end{aligned}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_314 EQ9968 (14.2.9)		■ 0.999	$\hat{\omega}=\left[\begin{array}{c c} \omega^g & -\alpha \\ \hline \alpha^T & \hat{\omega} \end{array}\right]=:\omega^g\oplus\hat{\omega}-\hat{\alpha};\quad \hat{\alpha}=\left[\begin{array}{c c} 0 & \alpha \\ \hline -\alpha^T & 0 \end{array}\right]$		
mielke-geometrodynamics_339 EQ1039 (15.1.16)		■ 0.999	$\begin{aligned} ds_{ij}^2 &= f_{ij} dx^i \otimes_s dx^j \\ &= -\frac{3\gamma}{2} e^{-\lambda^f} dt^2 + 2 \left[\gamma \left\{ 1 - \left(1 + \frac{9\gamma}{4} \right) e^{\lambda^f - \lambda^g} + \frac{9\gamma}{4} e^{2(\lambda^f - \lambda^g)} \right\} \right]^{1/2} dt dr \\ &\quad + e^{\lambda^g} \left\{ \frac{2}{3} + \frac{3\gamma}{2} (1 - e^{\lambda^g - \lambda^f}) \right\} dr^2 + \frac{2}{3} r^2 d\Omega^2 \end{aligned}$		
mielke-geometrodynamics_366 EQ1088 (A.15)		■ 0.999	$C(1,n-1)=\left[\bigotimes_{i=1}^{[\frac{n-2}{2}]}C(0,2)\right]\otimes C(1,1).$		
mielke-geometrodynamics_142 EQ0434 (6.8.5)		■ 0.999	$H_{\alpha} := -\frac{\partial L_g}{\partial d\vartheta^{\alpha}} = -\frac{\partial L_g}{\partial T^{\alpha}}, \quad \text{and} \quad H_{\alpha\beta} := -\frac{\partial L_g}{\partial d\Gamma^{\alpha\beta}} = -\frac{\partial L_g}{\partial R^{\alpha\beta}}.$		
mielke-geometrodynamics_183 EQ0566 (8.6.3)		■ 0.999	$H_{\alpha\beta} := -\partial L_{\text{qPG}}/\partial R^{\alpha\beta} = \frac{a_0}{2\ell^2} \eta_{\alpha\beta} + \left(\sum_{N=1}^6 b_{(N)} {}^{(N)}R_{\alpha\beta} \right)$		
mielke-geometrodynamics_254 EQ0798 (11.4.10)		■ 0.999	$\begin{aligned} &\left\{ i\bar{\gamma}^k \left(\partial_k + i\bar{\Gamma}_k + \frac{3}{4}\partial_k\mu \right) \right. \\ &\quad \left. + \frac{3\varepsilon}{8}\ell^2 e^{\mu/2} \left(\bar{\psi}\psi - \text{a}\bar{\psi}\gamma^5\psi\gamma^5 \right) - e^{\mu/2}m \right\} \psi = 0. \end{aligned}$		
mielke-geometrodynamics_270 EQ0833 (12.2.1)		■ 0.999	$L_{\mathrm{D}}=L(\gamma,\psi,\mathscr{D}\psi)=\frac{i}{2}\left\{\bar{\psi}^*\gamma\wedge\mathscr{D}\psi+\overline{\mathscr{D}\psi}\wedge^*\gamma\psi\right\}-m\bar{\psi}\psi\eta,$		
mielke-geometrodynamics_38 EQ0020 (2.4.2)		■ 1.000	$V^{\rho}:=V^{\rho}(M,\mathbb{C}^N,\rho(G)\subseteq GL(N,\mathbb{C}),P),$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_55 EQ0091 (3.2.1)	55	■ 1.000	$\mathrm{0} \rightarrow \mathrm{G}^{-1} \mathrm{0} \quad \mathrm{0} \rightarrow \mathrm{G}^{-1} \mathrm{0}$	$\mathcal{O} \rightarrow G^{-1} \mathcal{O} = G^{-1} \mathcal{O} G.$	$\mathcal{O} \rightarrow G^{-1} \mathcal{O} = G^{-1} \mathcal{O} G.$
mielke-geometrodynamics_57 EQ0101 (3.3.2)	57	■ 1.000	$A \rightarrow A + d \theta(x) \quad \Rightarrow \quad F \rightarrow d A + d d \theta = d A = F$	$A \rightarrow A + d \theta(x) \quad \Rightarrow \quad F \rightarrow d A + d d \theta = d A = F.$	$A \rightarrow A + d \theta(x) \quad \Rightarrow \quad F \rightarrow d A + d d \theta = d A = F .$
mielke-geometrodynamics_63 EQ0137 (3.4.8)	63	■ 1.000	$\begin{aligned} \delta L_{\omega} &= \delta \Omega \wedge \Pi^{\Omega} = D(\delta \omega) \wedge \Pi^{\Omega} \\ &= -\delta \omega \wedge D \Pi^{\Omega} + d(\delta \omega \wedge \Pi^{\Omega}). \end{aligned}$	$\delta L_{\omega} = \delta \Omega \wedge \Pi^{\Omega} = D(\delta \omega) \wedge \Pi^{\Omega} \\ = -\delta \omega \wedge D \Pi^{\Omega} + d(\delta \omega \wedge \Pi^{\Omega}).$	$\delta L_{\omega} = \delta \Omega \wedge \Pi^{\Omega} = D(\delta \omega) \wedge \Pi^{\Omega} \\ = -\delta \omega \wedge D \Pi^{\Omega} + d(\delta \omega \wedge \Pi^{\Omega}).$
mielke-geometrodynamics_65 EQ0146 (3.4.17)	65	■ 1.000	$Q = Q^{(i)} I_i := \int_{H^3} \iota = \int_{\partial H^3} \Pi^{\Omega},$	$Q = Q^{(i)} I_i := \int_{H^3} \iota = \int_{\partial H^3} \Pi^{\Omega},$	$Q = Q^{(i)} I_i := \int_{H^3} \iota = \int_{\partial H^3} \Pi^{\Omega},$
mielke-geometrodynamics_72 EQ0175 (3.7.4)	72	■ 1.000	$\tau_{\mathrm{H}} = - \left[\phi^{(0)}, * \Theta^{(1)} \right]$	$\tau_{\mathrm{H}} = - \left[\phi^{(0)}, * \Theta^{(1)} \right]$	$\tau_{\mathrm{H}} = - [\phi^{(0)}, * \Theta^{(1)}]$
mielke-geometrodynamics_82 EQ0195 (4.1.5)	82	■ 1.000	$\mathfrak{a}(n, \mathbb{R}) = \mathbb{R}^n \otimes \mathfrak{gl}(n, \mathbb{R})$	$\mathfrak{a}(n, \mathbb{R}) = \mathbb{R}^n \otimes \mathfrak{gl}(n, \mathbb{R})$	$\mathfrak{a}(n, \mathbb{R}) = \mathbb{R}^n \otimes \mathfrak{gl}(n, \mathbb{R})$
mielke-geometrodynamics_84 EQ0208 (4.1.22)	84	■ 1.000	$T_{\mathrm{A}}(M) = \stackrel{\circ}{V}(M, \mathbb{R}^n, A(n, \mathbb{R}), A(M)) \subset V^{\mathrm{id}}$	$T_{\mathrm{A}}(M) = \stackrel{\circ}{V}(M, \mathbb{R}^n, A(n, \mathbb{R}), A(M)) \subset V^{\mathrm{id}}$	$T_{\mathrm{A}}(M) = \stackrel{\circ}{V}(M, \mathbb{R}^n, A(n, \mathbb{R}), A(M)) \subset V^{\mathrm{id}}$
mielke-geometrodynamics_85 EQ0212 (4.2.4)	85	■ 1.000	$\Omega^{\mathrm{T}} := D \omega^{\mathrm{T}} = d \omega^{\mathrm{T}} - [\omega^{\mathrm{T}}, \omega^{\mathrm{L}}],$	$\Omega^{\mathrm{T}} := D \omega^{\mathrm{T}} = d \omega^{\mathrm{T}} - [\omega^{\mathrm{T}}, \omega^{\mathrm{L}}],$	$\Omega^{\mathrm{T}} := D \omega^{\mathrm{T}} = d \omega^{\mathrm{T}} - [\omega^{\mathrm{T}}, \omega^{\mathrm{L}}],$
mielke-geometrodynamics_86 EQ0215 (4.2.7)	86	■ 1.000	$\mathcal{T}(n, \mathbb{R}) := C^{\infty}(A(M) \times_{\mathrm{Ad}} \mathbb{R}^n)$	$\mathcal{T}(n, \mathbb{R}) := C^{\infty}(A(M) \times_{\mathrm{Ad}} \mathbb{R}^n)$	$\mathcal{T}(n, \mathbb{R}) := C^{\infty}(A(M) \times_{\mathrm{Ad}} \mathbb{R}^n)$
mielke-geometrodynamics_86 EQ0217 (4.2.9)	86	■ 1.000	$A = \left[\begin{array}{c c} G & T \\ \hline 0 & 1 \end{array} \right] \in \mathcal{A}_p, \quad G \in \mathcal{G}_p, T \in \mathcal{T}_p$	$A = \left[\begin{array}{c c} G & T \\ \hline 0 & 1 \end{array} \right] \in \mathcal{A}_p, \quad G \in \mathcal{G}_p, T \in \mathcal{T}_p$	$A = \left[\begin{array}{c c} G & T \\ \hline 0 & 1 \end{array} \right] \in \mathcal{A}_p, \quad G \in \mathcal{G}_p, T \in \mathcal{T}_p$
mielke-geometrodynamics_91 EQ0243 (4.3.8)	91	■ 1.000	$\cos \phi = \frac{g_{\mu \nu} d V^{\mu} d W^{\nu}}{\sqrt{g_{\alpha \beta} d V^{\alpha} d V^{\beta}} \sqrt{g_{\alpha \beta} d W^{\alpha} d W^{\beta}}}.$	$\cos \phi = \frac{g_{\mu \nu} d V^{\mu} d W^{\nu}}{\sqrt{g_{\alpha \beta} d V^{\alpha} d V^{\beta}} \sqrt{g_{\alpha \beta} d W^{\alpha} d W^{\beta}}}.$	$\cos \phi = \frac{g_{\mu \nu} d V^{\mu} d W^{\nu}}{\sqrt{g_{\alpha \beta} d V^{\alpha} d V^{\beta}} \sqrt{g_{\alpha \beta} d W^{\alpha} d W^{\beta}}}.$
mielke-geometrodynamics_111 EQ0323 (5.2.7)	111	■ 1.000	$\begin{aligned} \Sigma_{\alpha} &\rightarrow \sigma_{\alpha} := \Sigma_{\alpha} - D \mu_{\alpha} + e_{\beta} \rfloor (T^{\beta} \wedge \mu_{\alpha}), \\ \tau_{\alpha \beta} &\rightarrow \hat{\tau}_{\alpha \beta} := \tau_{\alpha \beta} - \vartheta_{[\alpha} \wedge \mu_{\beta]} = 0 \end{aligned}$	$\Sigma_{\alpha} \rightarrow \sigma_{\alpha} := \Sigma_{\alpha} - D \mu_{\alpha} + e_{\beta} \rfloor (T^{\beta} \wedge \mu_{\alpha}), \\ \tau_{\alpha \beta} \rightarrow \hat{\tau}_{\alpha \beta} := \tau_{\alpha \beta} - \vartheta_{[\alpha} \wedge \mu_{\beta]} = 0$	$\Sigma_{\alpha} \rightarrow \sigma_{\alpha} := \Sigma_{\alpha} - D \mu_{\alpha} + e_{\beta} \rfloor (T^{\beta} \wedge \mu_{\alpha}), \\ \tau_{\alpha \beta} \rightarrow \hat{\tau}_{\alpha \beta} := \tau_{\alpha \beta} - \vartheta_{[\alpha} \wedge \mu_{\beta]} = 0$
mielke-geometrodynamics_112 EQ0328 (5.3.5)	112	■ 1.000	$\Theta := D \gamma \quad \text{and} \quad \Omega^g := d \Gamma + \Gamma \wedge \Gamma.$	$\Theta := D \gamma \quad \text{and} \quad \Omega^g := d \Gamma + \Gamma \wedge \Gamma.$	$\Theta := D \gamma \quad \text{and} \quad \Omega^g := d \Gamma + \Gamma \wedge \Gamma.$
mielke-geometrodynamics_127 EQ0370 (6.2.21)	127	■ 1.000	$\stackrel{(\pm)}{\Pi}_{\alpha} = -\partial \stackrel{(\pm)}{L} / \partial T^{\alpha} = H_{\alpha}^{\parallel} \mp (i / \kappa) T_{\alpha}$	$\stackrel{(\pm)}{\Pi}_{\alpha} = -\partial \stackrel{(\pm)}{L} / \partial T^{\alpha} = H_{\alpha}^{\parallel} \mp (i / \kappa) T_{\alpha}$	$\stackrel{(\pm)}{\Pi}_{\alpha} = -\partial \stackrel{(\pm)}{L} / \partial T^{\alpha} = H_{\alpha}^{\parallel} \mp (i / \kappa) T_{\alpha}$
mielke-geometrodynamics_129 EQ0380 (6.2.31)	129	■ 1.000	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} = 0,$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} = 0,$	$\stackrel{(\pm)}{D} \stackrel{(\pm)}{\Pi}_{\alpha} = 0,$
mielke-geometrodynamics_141 EQ0428 (6.7.10)	141	■ 1.000	$\kappa D \theta_{\mathrm{L}} \wedge D T_{\beta} = -T \wedge \eta_{\beta} = D \eta_{\beta},$	$\kappa D \theta_{\mathrm{L}} \wedge D T_{\beta} = -T \wedge \eta_{\beta} = D \eta_{\beta},$	$\kappa D \theta_{\mathrm{L}} \wedge D T_{\beta} = -T \wedge \eta_{\beta} = D \eta_{\beta},$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_167 E08522 (7.7.41)	167	■ 1.000	$\begin{gathered} K^{\{(\star)\}}:=\eta_{\alpha\beta\gamma}\wedge K^{\beta\gamma}\gamma^{\alpha}\wedge K^{\{\beta\gamma\}}\gamma^{\alpha},\\ \Omega^{g(\star)}:=\frac{i}{8}R_{\alpha\beta}\eta^{\alpha\beta\gamma\delta}\sigma_{\gamma\delta}=-\frac{1}{4}R^{\alpha\beta}\gamma_5\sigma_{\alpha\beta}=i\gamma_5\Omega^g. \end{gathered}$	$K^{(\star)}:=\eta_{\alpha\beta\gamma}\wedge K^{\beta\gamma}\gamma^{\alpha},$ $\Omega^{g(\star)}:=\frac{i}{8}R_{\alpha\beta}\eta^{\alpha\beta\gamma\delta}\sigma_{\gamma\delta}=-\frac{1}{4}R^{\alpha\beta}\gamma_5\sigma_{\alpha\beta}=i\gamma_5\Omega^g.$	$K^{(\star)}:=\eta_{\alpha\beta\gamma}\wedge K^{\beta\gamma}\gamma^{\alpha},$ $\Omega^{g(\star)}:=\frac{i}{8}R_{\alpha\beta}\eta^{\alpha\beta\gamma\delta}\sigma_{\gamma\delta}=-\frac{1}{4}R^{\alpha\beta}\gamma_5\sigma_{\alpha\beta}=i\gamma_5\Omega^g.$
mielke-geometrodynamics_168 E08526 (7.7.46)	168	■ 1.000	$L_{\mathrm{NY}}:=dC_{\mathrm{TT}}=\frac{1}{2\ell^2}\left(T^{\alpha}\wedge T_{\alpha}+R_{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}\right),$	$L_{\mathrm{NY}}:=dC_{\mathrm{TT}}=\frac{1}{2\ell^2}\left(T^{\alpha}\wedge T_{\alpha}+R_{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}\right),$	$L_{\mathrm{NY}}:=dC_{\mathrm{TT}}=\frac{1}{2\ell^2}\left(T^{\alpha}\wedge T_{\alpha}+R_{\alpha\beta}\wedge\vartheta^{\alpha}\wedge\vartheta^{\beta}\right),$
mielke-geometrodynamics_175 E08535 (8.2.2)	175	■ 1.000	$\widetilde{\Gamma}:=\Gamma\oplus C,\quad\widetilde{\Omega}^g:=\Omega^g\oplus\Psi\oplus\Phi.$	$\widetilde{\Gamma}:=\Gamma\oplus C,\quad\widetilde{\Omega}^g:=\Omega^g\oplus\Psi\oplus\Phi.$	$\widetilde{F}:=F\oplus C,\quad\widetilde{\Omega}^g:=\Omega^g\oplus\Psi\oplus\Phi.$
mielke-geometrodynamics_176 E08538 (8.2.5)	176	■ 1.000	$s_{\mathrm{loc}}(\alpha+\lambda)=-d\lambda,\quad s_{\mathrm{loc}}\lambda=0$	$s_{\mathrm{loc}}(\alpha+\lambda)=-d\lambda,\quad s_{\mathrm{loc}}\lambda=0$	$s_{\mathrm{loc}}(\alpha+\lambda)=-d\lambda,\quad s_{\mathrm{loc}}\lambda=0$
mielke-geometrodynamics_176 E08539 (8.2.6)	176	■ 1.000	$(d\oplus s_{\mathrm{loc}})(\alpha+\lambda)=d\alpha.$	$(d\oplus s_{\mathrm{loc}})(\alpha+\lambda)=d\alpha.$	$(d\oplus s_{\mathrm{loc}})(\alpha+\lambda)=d\alpha.$
mielke-geometrodynamics_185 E08573 (8.6.11)	185	■ 1.000	$\begin{aligned} L_{\mathrm{qPG}}^{(*)}&=\frac{\Lambda}{\ell^2}\eta-\frac{a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2}T^{\alpha}\wedge H_{\alpha}^{(*)}-\frac{1}{2}R^{\alpha\beta}\wedge H_{\alpha\beta}^{(*)}\\ &=\frac{\Lambda}{\ell^2}\eta-\frac{\theta_{\mathrm{T}}^{*}+a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2\ell^2}T^{\alpha}\wedge\left[\theta_{\mathrm{T}}^{*}K_{\alpha}^{(*)}-\theta_{\mathrm{T}}T_{\alpha}\right]\\ &\quad -\frac{1}{2}R^{\alpha\beta}\wedge\left[\theta_{\mathrm{L}}R_{\alpha\beta}+\theta_{\mathrm{L}}^{*}R_{\alpha\beta}^{(*)}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}\right]. \end{aligned}$	$L_{\mathrm{qPG}}^{(*)}=\frac{\Lambda}{\ell^2}\eta-\frac{a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2}T^{\alpha}\wedge H_{\alpha}^{(*)}-\frac{1}{2}R^{\alpha\beta}\wedge H_{\alpha\beta}^{(*)}$ $=\frac{\Lambda}{\ell^2}\eta-\frac{\theta_{\mathrm{T}}^{*}+a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2\ell^2}T^{\alpha}\wedge\left[\theta_{\mathrm{T}}^{*}K_{\alpha}^{(*)}-\theta_{\mathrm{T}}T_{\alpha}\right]$ $-\frac{1}{2}R^{\alpha\beta}\wedge\left[\theta_{\mathrm{L}}R_{\alpha\beta}+\theta_{\mathrm{L}}^{*}R_{\alpha\beta}^{(*)}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}\right].$	$L_{\mathrm{qPG}}^{(**)}=\frac{\Lambda}{\ell^2}\eta-\frac{a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2}T^{\alpha}\wedge H_{\alpha}^{(**)}-\frac{1}{2}R^{\alpha\beta}\wedge H_{\alpha\beta}^{(**)}$ $=\frac{\Lambda}{\ell^2}\eta-\frac{\theta_{\mathrm{T}}^{*}+a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2\ell^2}T^{\alpha}\wedge\left[\theta_{\mathrm{T}}^{*}K_{\alpha}^{(*)}-\theta_{\mathrm{T}}T_{\alpha}\right]$ $-\frac{1}{2}R^{\alpha\beta}\wedge\left[\theta_{\mathrm{L}}R_{\alpha\beta}+\theta_{\mathrm{L}}^{*}R_{\alpha\beta}^{(*)}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}\right]. \quad (8.6)$
mielke-geometrodynamics_186 E08580 (8.6.18)	186	■ 1.000	$G_{\alpha}^{\{\}}-\Lambda_{\mathrm{eff}}\eta_{\alpha}=\kappa_{\mathrm{eff}}\widehat{\Sigma}_{\alpha}.$	$G_{\alpha}^{\{\}}-\Lambda_{\mathrm{eff}}\eta_{\alpha}=\kappa_{\mathrm{eff}}\widehat{\Sigma}_{\alpha}.$	$G_{\alpha}^{\{\}}-\Lambda_{\mathrm{eff}}\eta_{\alpha}=\kappa_{\mathrm{eff}}\widehat{\Sigma}_{\alpha}.$
mielke-geometrodynamics_195 E08595 (9.1.13)	195	■ 1.000	$R^{\{\}}=-\frac{6\kappa\gamma}{\ell^{*2}}.$	$R^{\{\}}=-\frac{6\kappa\gamma}{\ell^{*2}}.$	$R^{\{\}}=-\frac{6\kappa\gamma}{\ell^{*2}}.$
mielke-geometrodynamics_214 E08645 (10.3.2)	214	■ 1.000	$\widetilde{T}_{\alpha}=T_{\alpha}-\frac{\varepsilon}{\ell}\eta_{\alpha},\quad\widetilde{R}_{\alpha}^{*}=R_{\alpha}^{*}-(-1)^s\frac{\varepsilon}{2\ell}T_{\alpha}+(-1)^s\frac{\varepsilon^2}{4\ell^2}\eta_{\alpha}$	$\widetilde{T}_{\alpha}=T_{\alpha}-\frac{\varepsilon}{\ell}\eta_{\alpha},\quad\widetilde{R}_{\alpha}^{*}=R_{\alpha}^{*}-(-1)^s\frac{\varepsilon}{2\ell}T_{\alpha}+(-1)^s\frac{\varepsilon^2}{4\ell^2}\eta_{\alpha}$	$\widetilde{T}_{\alpha}=T_{\alpha}-\frac{\varepsilon}{\ell}\eta_{\alpha},\quad\widetilde{R}_{\alpha}^{*}=R_{\alpha}^{*}-(-1)^s\frac{\varepsilon}{2\ell}T_{\alpha}+(-1)^s\frac{\varepsilon^2}{4\ell^2}\eta_{\alpha}$
mielke-geometrodynamics_220 E08676 (10.3.33)	220	■ 1.000	$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}}=-(-1)^s\frac{\theta_{\mathrm{T}}\theta_{\mathrm{L}}}{2\chi\ell}\left(d\Gamma_{\alpha}^{*}+\frac{2}{\theta_{\mathrm{L}}}\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}*}\right),$	$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}}=-(-1)^s\frac{\theta_{\mathrm{T}}\theta_{\mathrm{L}}}{2\chi\ell}\left(d\Gamma_{\alpha}^{*}+\frac{2}{\theta_{\mathrm{L}}}\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}*}\right),$	$\check{\mathcal{E}}_{\alpha}^{\mathrm{MB}}=-(-1)^s\frac{\theta_{\mathrm{T}}\theta_{\mathrm{L}}}{2\chi\ell}\left(d\Gamma_{\alpha}^{*}+\frac{2}{\theta_{\mathrm{L}}}\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}*}\right),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_222 EQ0688 (10.3.46)		■ 1.000	$G_{\alpha}^{\{\}} - \tilde{A}\eta_{\alpha} = \ell \tilde{\Sigma}_{\alpha},$	$G_{\alpha}^{\{\}} - \tilde{A}\eta_{\alpha} = \ell \tilde{\Sigma}_{\alpha},$	$G_{\alpha}^{\{\}} - \tilde{A}\eta_{\alpha} = \ell \tilde{\Sigma}_{\alpha},$
mielke-geometrodynamics_223 EQ0690 (10.5.1)		■ 1.000	$\sigma^{\wedge\{1\}}=\left(\begin{array}{ll} 0 & 1 \\ 0 & 1 \end{array}\right), \quad \sigma^{\wedge\{2\}}=\left(\begin{array}{cc} 0 & -i \\ 0 & 0 \end{array}\right), \quad \sigma^{\wedge\{3\}}=\left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right).$	$\sigma^1 = \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right), \quad \sigma^2 = \left(\begin{array}{cc} 0 & -i \\ i & 0 \end{array}\right), \quad \sigma^3 = \left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right).$	$\sigma^1 = \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right), \quad \sigma^2 = \left(\begin{array}{cc} 0 & -i \\ i & 0 \end{array}\right), \quad \sigma^3 = \left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right).$
mielke-geometrodynamics_227 EQ0710 (10.7.6)		■ 1.000	$\begin{aligned} \vartheta^{\hat{0}} &= dt + \ell^2 \sigma \rho^{*2} [1 - \cos(\rho/\rho^*)] d\phi \\ \vartheta^{\hat{1}} &= d\rho, \quad \vartheta^{\hat{2}} = \rho^* \sin(\rho/\rho^*) d\phi, \\ \Gamma^{\hat{1}\hat{2}} &= \cos(\rho/\rho^*) d\phi = -\Gamma^{\hat{2}\hat{1}}. \end{aligned}$	$\vartheta^{\hat{0}} = dt + \ell^2 \sigma \rho^{*2} [1 - \cos(\rho/\rho^*)] d\phi$ $\vartheta^{\hat{1}} = d\rho, \quad \vartheta^{\hat{2}} = \rho^* \sin(\rho/\rho^*) d\phi$ $\Gamma^{\hat{1}\hat{2}} = \cos(\rho/\rho^*) d\phi = -\Gamma^{\hat{2}\hat{1}}$	$\vartheta^{\hat{0}} = dt + \ell^2 \sigma \rho^{*2} [1 - \cos(\rho/\rho^*)] d\phi$ $\vartheta^{\hat{1}} = d\rho, \quad \vartheta^{\hat{2}} = \rho^* \sin(\rho/\rho^*) d\phi,$ $\Gamma^{\hat{1}\hat{2}} = \cos(\rho/\rho^*) d\phi = -\Gamma^{\hat{2}\hat{1}}.$
mielke-geometrodynamics_229 EQ0716 (10.9.2)		■ 1.000	$D \Psi = d \Psi - \frac{1}{2} \gamma_{\alpha} \Gamma^{*\alpha} \wedge \Psi,$	$D\Psi = d\Psi - \frac{1}{2} \gamma_{\alpha} \Gamma^{*\alpha} \wedge \Psi,$	$D\Psi = d\Psi - \frac{1}{2} \gamma_{\alpha} \Gamma^{*\alpha} \wedge \Psi,$
mielke-geometrodynamics_229 EQ0720 (10.9.6)		■ 1.000	$\tau_{\alpha}^{\star} := \frac{1}{2} \eta_{\alpha\beta\gamma} \tau^{\beta\gamma} = \frac{(-1)^s}{2} \frac{\delta L_{\Psi}}{\delta \Gamma_{\alpha}^{\star}} = \frac{(-1)^s}{2} \frac{i}{4} \bar{\Psi} \wedge \gamma_{\alpha} \Psi.$	$\tau_{\alpha}^{\star} := \frac{1}{2} \eta_{\alpha\beta\gamma} \tau^{\beta\gamma} = \frac{(-1)^s}{2} \frac{\delta L_{\Psi}}{\delta \Gamma_{\alpha}^{\star}} = \frac{(-1)^s}{2} \frac{i}{4} \bar{\Psi} \wedge \gamma_{\alpha} \Psi.$	$\tau_{\alpha}^{\star} := \frac{1}{2} \eta_{\alpha\beta\gamma} \tau^{\beta\gamma} = \frac{(-1)^s}{2} \frac{\delta L_{\Psi}}{\delta \Gamma_{\alpha}^{\star}} = \frac{(-1)^s}{2} \frac{i}{4} \bar{\Psi} \wedge \gamma_{\alpha} \Psi.$
mielke-geometrodynamics_231 EQ0728 (10.10.8)		■ 1.000	$\gamma^{\alpha} \Psi \wedge \bar{\Psi} \gamma_{\alpha} \Psi = 0,$	$\gamma^{\alpha} \Psi \wedge \bar{\Psi} \gamma_{\alpha} \Psi = 0,$	$\gamma^{\alpha} \Psi \wedge \bar{\Psi} \gamma_{\alpha} \Psi = 0,$
mielke-geometrodynamics_244 EQ0770 (11.1.37)		■ 1.000	$D \longrightarrow^{S^{-1}} D = S^{-1} D S.$	$D \longrightarrow^{S^{-1}} D = S^{-1} D S.$	$D \longrightarrow^{S^{-1}} D = S^{-1} D S.$
mielke-geometrodynamics_272 EQ0839 (12.2.7)		■ 1.000	$j_{\pm} := \frac{1}{2} \bar{\psi} (1 \pm \gamma_5)^* \gamma \psi = \bar{\psi}_{\text{L,R}}^* \gamma \psi_{\text{L,R}}.$	$j_{\pm} := \frac{1}{2} \bar{\psi} (1 \pm \gamma_5)^* \gamma \psi = \bar{\psi}_{\text{L,R}}^* \gamma \psi_{\text{L,R}}.$	$j_{\pm} := \frac{1}{2} \bar{\psi} (1 \pm \gamma_5)^* \gamma \psi = \bar{\psi}_{\text{L,R}}^* \gamma \psi_{\text{L,R}}.$
mielke-geometrodynamics_284 EQ0873 (13.3.1)		■ 1.000	$\not{L}^A{}_B := L^A{}_B - \frac{1}{n} \delta_B^A L^C{}_C.$	$\not{L}^A{}_B := L^A{}_B - \frac{1}{n} \delta_B^A L^C{}_C.$	$\not{L}^A{}_B := L^A{}_B - \frac{1}{n} \delta_B^A L^C{}_C.$
mielke-geometrodynamics_287 EQ0893 (13.4.12)		■ 1.000	$A_*(n, \mathbb{R})/\mathrm{GL}(n, \mathbb{R}) \approx \mathbb{R}^n \otimes \mathbb{R}_*^n,$	$A_*(n, \mathbb{R})/\mathrm{GL}(n, \mathbb{R}) \approx \mathbb{R}^n \otimes \mathbb{R}_*^n,$	$A_*(n, \mathbb{R})/\mathrm{GL}(n, \mathbb{R}) \approx \mathbb{R}^n \otimes \mathbb{R}_*^n,$
mielke-geometrodynamics_290 EQ0907 (13.5.12)		■ 1.000	$\tilde{L}_{\mathrm{SSB}} + d\hat{C} \cong \frac{1}{2\mu} \eta_{\alpha\beta\gamma\delta} \left[R^{\alpha\beta} - \frac{\Lambda}{3} \vartheta^{\alpha} \wedge \vartheta^{\beta} \right] \wedge \left[R^{\gamma\delta} - \frac{\Lambda}{3} \vartheta^{\gamma} \wedge \vartheta^{\delta} \right].$	$\tilde{L}_{\mathrm{SSB}} + d\hat{C} \cong \frac{1}{2\mu} \eta_{\alpha\beta\gamma\delta} \left[R^{\alpha\beta} - \frac{\Lambda}{3} \vartheta^{\alpha} \wedge \vartheta^{\beta} \right] \wedge \left[R^{\gamma\delta} - \frac{\Lambda}{3} \vartheta^{\gamma} \wedge \vartheta^{\delta} \right].$	$\tilde{L}_{\mathrm{SSB}} + d\hat{C} \cong \frac{1}{2\mu} \eta_{\alpha\beta\gamma\delta} \left[R^{\alpha\beta} - \frac{\Lambda}{3} \vartheta^{\alpha} \wedge \vartheta^{\beta} \right] \wedge \left[R^{\gamma\delta} - \frac{\Lambda}{3} \vartheta^{\gamma} \wedge \vartheta^{\delta} \right].$
mielke-geometrodynamics_307 EQ0947 (14.1.28)		■ 1.000	$R_{i\alpha}^{\{\}} R_{\cdot j}^{\{\}\alpha} = \frac{1}{4} g_{ij} R_{\alpha\beta}^{\{\}} R^{\{\}\alpha\beta}$	$R_{i\alpha}^{\{\}} R_{\cdot j}^{\{\}\alpha} = \frac{1}{4} g_{ij} R_{\alpha\beta}^{\{\}} R^{\{\}\alpha\beta}$	$R_{i\alpha}^{\{\}} R_{\cdot j}^{\{\}\alpha} = \frac{1}{4} g_{ij} R_{\alpha\beta}^{\{\}} R^{\{\}\alpha\beta}$
mielke-geometrodynamics_323 EQ1004 (14.3.17)		■ 1.000	$\widetilde{\Theta} := D\gamma = \frac{1}{2} T_{\alpha\beta}^{\cdot\cdot c} \gamma_c \otimes \vartheta^{\alpha} \wedge \vartheta^{\beta}$	$\widetilde{\Theta} := D\gamma = \frac{1}{2} T_{\alpha\beta}^{\cdot\cdot c} \gamma_c \otimes \vartheta^{\alpha} \wedge \vartheta^{\beta}$	$\widetilde{\Theta} := D\gamma = \frac{1}{2} T_{\alpha\beta}^{\cdot\cdot c} \gamma_c \otimes \vartheta^{\alpha} \wedge \vartheta^{\beta}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ1013 (14.4.8)	326	■ 1.000	$M_{\mathrm{ir}}^2:=\frac{S_{\mathrm{A}}}{16\pi}\frac{M^{*4}c^2}{\hbar^2}$	$M_{\mathrm{ir}}^2:=\frac{S_{\mathrm{A}}}{16\pi}\frac{M^{*4}c^2}{\hbar^2}$	$M_{\mathrm{ir}}^2:=\frac{S_{\mathrm{A}}}{16\pi}\frac{M^{*4}c^2}{\hbar^2}$
mielke-geometrodynamics_EQ1135 (C.3)	375	■ 1.000	$\left[I_i,I_j\right]=c_{ij}^{k}I_k\text{ .}$	$[I_i,I_j]=c_{ij}^{k}I_k.$	$[I_i,I_j]=c_{ij}^{k}I_k.$
mielke-geometrodynamics_EQ0001 (1.1)	24	■ 1.000	$\ell^*:=\sqrt{8\pi\hbar G_{\mathrm{N}}/c^3}\simeq 8\times 10^{-33}\text{ cm},$	$\ell^*:=\sqrt{8\pi\hbar G_N/c^3}\simeq 8\times 10^{-33}\text{ cm},$	$\ell^*:=\sqrt{8\pi\hbar G_N/c^3}\simeq 8\times 10^{-33}\text{ cm},$
mielke-geometrodynamics_EQ0003 (2.1.2)	32	■ 1.000	$e^{\mathrm{f}}:=\left.\frac{df(s(t))}{dt}\right _{t_o=s^{-1}(m)}=\frac{\partial f}{\partial x^i}\cdot\frac{dx^i(s(t))}{dt}$	$e^{\mathrm{f}}:=\left.\frac{df(s(t))}{dt}\right _{t_o=s^{-1}(m)}=\frac{\partial f}{\partial x^i}\cdot\frac{dx^i(s(t))}{dt}$	$e^{\mathrm{f}}:=\left.\frac{df(s(t))}{dt}\right _{t_o=\bar{s}^{-1}(m)}=\frac{\partial f}{\partial x^i}\cdot\frac{dx^i(s(t))}{dt}$
mielke-geometrodynamics_EQ0005 (2.1.4)	33	■ 1.000	$T(M)=\bigcup_{m\in M}T_m(M)$	$T(M)=\bigcup_{m\in M}T_m(M)$	$T(M)=\bigcup_{m\in M}T_m(M).$
mielke-geometrodynamics_EQ0008 (2.2.3)	34	■ 1.000	$\vartheta^{\alpha}=\mathrm{E}_{\!j}^{\alpha}(m)\,\mathrm{d}\,x^{\!j}\text{ .}$	$\vartheta^{\alpha}=\mathrm{E}_{\!j}^{\alpha}(m)dx^j.$	$\vartheta^{\alpha}=\mathrm{E}_{\!j}^{\alpha}(m)\,dx^j.$
mielke-geometrodynamics_EQ0011 (2.2.6)	34	■ 1.000	$T=T_{\alpha_1\cdots\alpha_p}^{\beta_1\cdots\beta_q}(m)\vartheta^{\alpha_1}\otimes\cdots\otimes\vartheta^{\alpha_p}\otimes e_{\beta_1}\otimes\cdots\otimes e_{\beta_q}\in C^{\infty}\left(T_p^q(M)\right).$	$T=T_{\alpha_1\cdots\alpha_p}^{\beta_1\cdots\beta_q}(m)\vartheta^{\alpha_1}\otimes\cdots\otimes\vartheta^{\alpha_p}\otimes e_{\beta_1}\otimes\cdots\otimes e_{\beta_q}\in C^{\infty}\left(T_p^q(M)\right).$	$T=T_{\alpha_1\cdots\alpha_p}^{\beta_1\cdots\beta_q}(m)\vartheta^{\alpha_1}\otimes\cdots\otimes\vartheta^{\alpha_p}\otimes e_{\beta_1}\otimes\cdots\otimes e_{\beta_q}\in C^{\infty}(T_p^q(M)).$
mielke-geometrodynamics_EQ0012 (2.2.7)	34	■ 1.000	$ds^2=g_{\alpha\beta}\vartheta^{\alpha}\otimes_s\vartheta^{\beta}=:g_{ij}dx^i\otimes_sdx^j.$	$ds^2=g_{\alpha\beta}\vartheta^{\alpha}\otimes_s\vartheta^{\beta}=:g_{ij}dx^i\otimes_sdx^j.$	$ds^2=g_{\alpha\beta}\vartheta^{\alpha}\otimes_s\vartheta^{\beta}=:g_{ij}dx^i\otimes_sdx^j.$
mielke-geometrodynamics_EQ0014 (2.2.9)	35	■ 1.000	$\alpha^{\{p\}}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^{\infty}\left(\wedge^pT^*(M)\right).$	$\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^{\infty}\left(\wedge^pT^*(M)\right).$	$\alpha^{(p)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^{\infty}(\wedge^pT^*(M)).$
mielke-geometrodynamics_EQ0015 (2.3.1)	36	■ 1.000	$\pi\colon F\rightarrowtail M\text{ .}$	$\pi:F\rightarrowtail M.$	$\pi:F\rightarrowtail M.$
mielke-geometrodynamics_EQ0016 (2.3.2)	37	■ 1.000	$\sigma\colon U\rightarrowtail F\quad\text{where}\quad\pi\circ\sigma=\mathrm{id}$	$\sigma:U\rightarrowtail F\quad\text{where}\quad\pi\circ\sigma=\mathrm{id}$	$\sigma:U\rightarrowtail F\quad\text{where}\quad\pi\circ\sigma=\mathrm{id}$
mielke-geometrodynamics_EQ0017 (2.3.3)	37	■ 1.000	$\iota\!\left(m,g_1g_2\right)=\iota\!\left(m,g_1\right)\cdot g_2,\quad m\in U,g\in G.$	$\iota(m,g_1g_2)=\iota(m,g_1)\cdot g_2,\quad m\in U,g\in G.$	$\iota(m,g_1g_2)=\iota(m,g_1)\cdot g_2,\quad m\in U,g\in G.$
mielke-geometrodynamics_EQ0019 (2.4.1)	38	■ 1.000	$\delta_g(p,\zeta):=\left(pg,g^{-1}\zeta\right)\subset P\times F,\quad g\in G.$	$\delta_g(p,\zeta):=(pg,g^{-1}\zeta)\subset P\times F,\quad g\in G.$	$\delta_g(p,\zeta):=(pg,g^{-1}\zeta)\subset P\times F,\quad g\in G.$
mielke-geometrodynamics_EQ0022 (2.4.4)	38	■ 1.000	$\varphi^{\{p\}}=\left\{\varphi^{\left(p\right)A}\mid A=1,\ldots,N\right\}.$	$\varphi^{(p)}=\left\{\varphi^{(p)A}\mid A=1,\ldots,N\right\}.$	$\varphi^{(p)}=\left\{\varphi^{(p)A} A=1,\ldots,N\right\}.$
mielke-geometrodynamics_EQ0023 (2.4.5)	39	■ 1.000	$L_j:=\left[\left.\frac{\partial\rho_A^{B}(g(\xi))}{\partial\xi^j}\right _{g=e}\right]\in T_e(\rho(G))$	$L_j:=\left[\left.\frac{\partial\rho_A^{B}(g(\xi))}{\partial\xi^j}\right _{g=e}\right]\in T_e(\rho(G))$	$L_j:=\left[\left.\frac{\partial\rho_A^{B}(g(\xi))}{\partial\xi^j}\right _{g=e}\right]\in T_e(\rho(G))$
mielke-geometrodynamics_EQ0025 (2.4.7)	39	■ 1.000	$\phi^{\{p\}}=\frac{1}{p!}\phi_{\alpha_1\cdots\alpha_p}^jL_j\otimes\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^{\infty}\left(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^{Ad}\right).$	$\phi^{(p)}=\frac{1}{p!}\phi_{\alpha_1\cdots\alpha_p}^jL_j\otimes\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^{\infty}\left(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^{Ad}\right).$	$\phi^{(p)}=\frac{1}{p!}\phi_{\alpha_1\cdots\alpha_p}^jL_j\otimes\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^{\infty}\left(\wedge^pT_{\mathbb{C}}^*(M)\otimes V^{Ad}\right).$
mielke-geometrodynamics_EQ0026 (2.4.8)	39	■ 1.000	$\left[\phi^{\{p\}},\psi^{\{q\}}\right]:=\phi^{\{p\}}\wedge\psi^{\{q\}}-(-1)^{pq}\psi^{\{q\}}\wedge\phi^{\{p\}}=(-1)^{pq+1}\left[\psi^{\{q\}},\phi^{\{p\}}\right]$	$\left[\phi^{(p)},\psi^{(q)}\right]:=\phi^{(p)}\wedge\psi^{(q)}-(-1)^{pq}\psi^{(q)}\wedge\phi^{(p)}=(-1)^{pq+1}\left[\psi^{(q)},\phi^{(p)}\right]$	$\left[\phi^{(p)},\psi^{(q)}\right]:=\phi^{(p)}\wedge\psi^{(q)}-(-1)^{pq}\psi^{(q)}\wedge\phi^{(p)}=(-1)^{pq+1}\left[\psi^{(q)},\phi^{(p)}\right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00027 (2.4.9)	39	■ 1.000	$\left[\varphi^{\{2\,k\}},\varphi^{\{2\,k\}}\right]=-\left[\varphi^{\{2\,k\}},\varphi^{\{2\,k\}}\right]=0\;.$	$\left[\phi^{(2k)},\phi^{(2k)}\right]=-\left[\phi^{(2k)},\phi^{(2k)}\right]=0.$	$\left[\phi^{(2k)},\phi^{(2k)}\right]=-\left[\phi^{(2k)},\phi^{(2k)}\right]=0.$
mielke-geometrodynamics_E00029 (2.5.2)	40	■ 1.000	$\omega(e\,g)=g^{-1}\,\omega(e)\,g,\quad e\in T(P)\;.$	$\omega(eg)=g^{-1}\omega(e)g,\quad e\in T(P).$	$\omega(eg)=g^{-1}\omega(e)g,\quad e\in T(P).$
mielke-geometrodynamics_E00030 (2.5.3)	40	■ 1.000	$\omega(p\,d\,g)=g^{-1}\,d\,g\quad\text{in}\quad T_e(G)\approx\mathfrak{g}$	$\omega(pdg)=g^{-1}dg\quad\in\quad T_e(G)\approx\mathfrak{g}$	$\omega(p\,d\,g)=g^{-1}dg\quad\in\quad T_e(G)\approx\mathfrak{g}$
mielke-geometrodynamics_E00031 (2.5.4)	40	■ 1.000	$D:C^{\infty}\left(V^{\rho}\right)\longrightarrow C^{\infty}\left(T_{\mathbb{C}}^{\ast}(M)\otimes V^{\rho}\right)$	$D:C^{\infty}(V^{\rho})\longrightarrow C^{\infty}(T_{\mathbb{C}}^{\ast}(M)\otimes V^{\rho})$	$D:C^{\infty}(V^{\rho})\longrightarrow C^{\infty}(T_{\mathbb{C}}^{\ast}(M)\otimes V^{\rho})$
mielke-geometrodynamics_E00032 (2.5.6)	41	■ 1.000	$D\,b=\omega\otimes b,\quad b\in C^{\infty}\left(V^{\rho}\right)\;.$	$Db=\omega\otimes b,\quad b\in C^{\infty}(V^{\rho}).$	$Db=\omega\otimes b,\quad b\in C^{\infty}(V^{\rho}).$
mielke-geometrodynamics_E00033 (2.5.7)	41	■ 1.000	$D\left(\varphi^{\left(p\right)}\otimes b\right)=d\varphi^{\left(p\right)}\otimes b+(-1)^p\varphi^{\left(p\right)}\otimes Db.$	$D\left(\varphi^{(p)}\otimes b\right)=d\varphi^{(p)}\otimes b+(-1)^p\varphi^{(p)}\otimes Db.$	$D(\varphi^{(p)}\otimes b)=d\varphi^{(p)}\otimes b+(-1)^p\varphi^{(p)}\otimes Db.$
mielke-geometrodynamics_E00034 (2.5.8)	41	■ 1.000	$D\left(\varphi^{\left(p\right)}\otimes b\right)=\left\{d\varphi^p+(-1)^p\left[\varphi^{\left(p\right)},\omega\right]\right\}\otimes b.$	$D\left(\varphi^{(p)}\otimes b\right)=\left\{d\varphi^p+(-1)^p\left[\varphi^{(p)},\omega\right]\right\}\otimes b.$	$D(\varphi^{(p)}\otimes b)=\left\{d\varphi^p+(-1)^p[\varphi^{(p)},\omega]\right\}\otimes b.$
mielke-geometrodynamics_E00035 (2.5.9)	41	■ 1.000	$D\,b_{\{A\}}=\omega_{\{A\}}\{\,\}^{\{B\}}\,b_{\{B\}},\,A,B=1,\,\ldots,\,N\;.$	$Db_A=\omega_A{}^Bb_B,A,B=1,\ldots,N.$	$Db_A=\omega_A{}^Bb_B,A,B=1,\ldots,N.$
mielke-geometrodynamics_E00036 (2.5.10)	41	■ 1.000	$\Gamma:=\stackrel{\ast}{\sigma}\,\omega=\Gamma_{\alpha}{}^jL_j\otimes\vartheta^{\alpha},\quad j=1,\ldots,\dim G,$	$\Gamma:=\overset{\ast}{\sigma}\,\omega=\Gamma_{\alpha}{}^jL_j\otimes\vartheta^{\alpha},\quad j=1,\ldots,\dim G,$	$\Gamma:=\overset{\ast}{\sigma}\,\omega=\Gamma_{\alpha}{}^jL_j\otimes\vartheta^{\alpha},\quad j=1,\ldots,\dim G,$
mielke-geometrodynamics_E00037 (2.5.11)	41	■ 1.000	$A_{\alpha}{}^j{}_{\{m\}}:=\Gamma_{\alpha}{}^j{}_{\{m\}}(m)_{\mid G=U(f)}$	$A_{\alpha}{}^j(m):=\Gamma_{\alpha}{}^j(m)_{ G=U(f)}$	$A_{\alpha}{}^j(m):=\Gamma_{\alpha}{}^j(m)_{ G=U(f)}$
mielke-geometrodynamics_E00038 (2.6.1)	42	■ 1.000	$D\,D\,b=: \,\Omega\otimes b\;.$	$DDb=:\Omega\otimes b.$	$D\,D\,b=:\,\Omega\otimes b.$
mielke-geometrodynamics_E00039 (2.6.2)	42	■ 1.000	$\begin{aligned} D\,D\,b &= D(\omega\otimes b) \,\&= d\,\omega\otimes b - \omega\wedge Db \\ &\otimes b - \omega\wedge D\,b \,\&= (d\,\omega - \omega\wedge\omega)\otimes b \end{aligned}$	$\begin{aligned} DDb &= D(\omega\otimes b) = d\omega\otimes b - \omega\wedge Db \\ &= (d\omega - \omega\wedge\omega)\otimes b \end{aligned}$	$\begin{aligned} DDb &= D(\omega\otimes b) = d\omega\otimes b - \omega\wedge Db \\ &= (d\omega - \omega\wedge\omega)\otimes b \end{aligned}$
mielke-geometrodynamics_E00040 (2.6.3)	42	■ 1.000	$\Omega=d\,\omega-\omega\wedge\omega\quad\Omega=d\,\omega-\frac{1}{2}[\omega,\omega]$	$\Omega=d\omega-\omega\wedge\omega=d\omega-\frac{1}{2}[\omega,\omega]$	$\Omega=d\omega-\omega\wedge\omega=d\omega-\frac{1}{2}[\omega,\omega]\;.$
mielke-geometrodynamics_E00043 (2.6.6)	42	■ 1.000	$F_{\{i\,j\}}=\partial_{\{i\}}\,A_{\{j\}}-\partial_{\{j\}}\,A_{\{i\}}$	$F_{ij}=\partial_iA_j-\partial_jA_i$	$F_{ij}=\partial_iA_j-\partial_jA_i$
mielke-geometrodynamics_E00044 (2.6.7)	42	■ 1.000	$\Delta b=b-b(\circ)=\oint(d\,b)_{\mathrm{hor}}$	$\Delta b=b-b(\circ)=\oint (db)_{\mathrm{hor}}$	$\Delta b=b-b(\circ)=\oint (db)_{\mathrm{hor}},$
mielke-geometrodynamics_E00047 (2.6.10)	43	■ 1.000	$K=\lim_{U\rightarrow 0}\left(\int_U\Omega\otimes b\right)/\left(\int_U1\otimes b\right)$	$K=\lim_{U\rightarrow 0}\left(\int_U\Omega\otimes b\right)/\left(\int_U1\otimes b\right)$	$K=\lim_{U\rightarrow 0}\left(\int_U\Omega\otimes b\right)/\left(\int_U1\otimes b\right)$
mielke-geometrodynamics_E00049 (2.6.12)	43	■ 1.000	$D\,\Omega\equiv 0$	$D\Omega\equiv 0$	$D\Omega\equiv 0$
mielke-geometrodynamics_E00052 (2.7.2)	44	■ 1.000	$G(g\,p)=g\,G(p),\quad g\in G,\quad p\in P\;;$	$G(gp)=gG(p),\quad g\in G,\quad p\in P;$	$G(g\,p)=g\,G(p),\quad g\in G,\quad p\in P;$
mielke-geometrodynamics_E00053 (2.7.3)	44	■ 1.000	$\pi\circ G=\pi\;.$	$\pi\circ G=\pi.$	$\pi\circ G=\pi.$
mielke-geometrodynamics_E00054 (2.7.4)	45	■ 1.000	$\mathscr{G}_p\approx C^{\infty}\left(P\times_{\mathrm{Ad}}G\right)\;.$	$\mathscr{G}_p\approx C^{\infty}(P\times_{\mathrm{Ad}}G).$	$\mathscr{G}_p\approx C^{\infty}(P\times_{\mathrm{Ad}}G).$
mielke-geometrodynamics_E00055 (2.7.5)	45	■ 1.000	$G(p)=\exp i\,\theta^k(m)\,L_{\{k\}}\in\mathscr{G}_p\;.$	$G(p)=\exp i\theta^k(m)L_k\in\mathscr{G}_p.$	$G(p)=\exp i\,\theta^k(m)L_k\in\mathscr{G}_p.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_45 EQ0057 (2.7.7)	45	■ 1.000	$\varphi \rightarrow^{G^{-1}} \varphi := G^{-1} \varphi \quad \in C^\infty(M, \mathbb{C}), G \in \mathcal{G}_V,$ $\varphi \in C^\infty(M, \mathbb{C}), G \in \mathcal{G}_V,$	$\varphi \rightarrow^{G^{-1}} \varphi := G^{-1} \varphi \quad \in C^\infty(M, \mathbb{C}), G \in \mathcal{G}_V,$	$\varphi \rightarrow^{G^{-1}} \varphi := G^{-1} \varphi \quad \in C^\infty(M, \mathbb{C}), G \in \mathcal{G}_V,$
mielke-geometrodynamics_46 EQ0063 (2.8.4)	46	■ 1.000	$p_k(V) = (-1)^k c_{2k}(V).$	$p_k(V) = (-1)^k c_{2k}(V).$	$p_k(V) = (-1)^k c_{2k}(V).$
mielke-geometrodynamics_47 EQ0066 (2.8.9)	47	■ 1.000	$\text{Tr}(\Omega \wedge \Omega) = d \text{Tr} \left(\omega \wedge \Omega + \frac{1}{3} \omega \wedge \omega \wedge \omega \right)$	$\text{Tr}(\Omega \wedge \Omega) = d \text{Tr} \left(\omega \wedge \Omega + \frac{1}{3} \omega \wedge \omega \wedge \omega \right)$	$\text{Tr}(\Omega \wedge \Omega) = d \text{Tr} \left(\omega \wedge \Omega + \frac{1}{3} \omega \wedge \omega \wedge \omega \right)$
mielke-geometrodynamics_48 EQ0067 (2.8.10)	48	■ 1.000	$\begin{aligned} & \text{Tr} \left(d \left(\omega \wedge d\omega - \frac{2}{3} \omega \wedge \omega \wedge \omega \right) \right) = \text{Tr} \{ d\omega \wedge d\omega - \omega \wedge dd\omega \\ & - \frac{2}{3} d\omega \wedge \omega \wedge \omega + \frac{2}{3} \omega \wedge d\omega \wedge \omega - \frac{2}{3} \omega \wedge \omega \wedge d\omega \} \\ & = \text{Tr} \{ d\omega \wedge d\omega - d\omega \wedge \omega \wedge \omega - \omega \wedge \omega \wedge d\omega \} \\ & = \text{Tr} \{ (d\omega - \omega \wedge \omega) \wedge (d\omega - \omega \wedge \omega) \} = \text{Tr}(\Omega \wedge \Omega) \end{aligned}$	$\begin{aligned} & \text{Tr} \left\{ d \left(\omega \wedge d\omega - \frac{2}{3} \omega \wedge \omega \wedge \omega \right) \right\} = \text{Tr} \{ d\omega \wedge d\omega - \omega \wedge dd\omega \\ & - \frac{2}{3} d\omega \wedge \omega \wedge \omega + \frac{2}{3} \omega \wedge d\omega \wedge \omega - \frac{2}{3} \omega \wedge \omega \wedge d\omega \} \\ & = \text{Tr} \{ d\omega \wedge d\omega - d\omega \wedge \omega \wedge \omega - \omega \wedge \omega \wedge d\omega \} \\ & = \text{Tr} \{ (d\omega - \omega \wedge \omega) \wedge (d\omega - \omega \wedge \omega) \} = \text{Tr}(\Omega \wedge \Omega) \end{aligned}$	$\begin{aligned} & \text{Tr} \left\{ d \left(\omega \wedge d\omega - \frac{2}{3} \omega \wedge \omega \wedge \omega \right) \right\} = \text{Tr} \{ d\omega \wedge d\omega - \omega \wedge dd\omega \\ & - \frac{2}{3} d\omega \wedge \omega \wedge \omega + \frac{2}{3} \omega \wedge d\omega \wedge \omega - \frac{2}{3} \omega \wedge \omega \wedge d\omega \} \\ & = \text{Tr} \{ d\omega \wedge d\omega - d\omega \wedge \omega \wedge \omega - \omega \wedge \omega \wedge d\omega \} \\ & = \text{Tr} \{ (d\omega - \omega \wedge \omega) \wedge (d\omega - \omega \wedge \omega) \} = \text{Tr}(\Omega \wedge \Omega). \end{aligned}$
mielke-geometrodynamics_48 EQ0068 (2.8.11)	48	■ 1.000	$\text{Tr}(\Omega \wedge \Omega) = 0.$	$\text{Tr}(\Omega \wedge \Omega) = 0.$	$\text{Tr}(\Omega \wedge \Omega) = 0.$
mielke-geometrodynamics_48 EQ0070 (2.8.13)	48	■ 1.000	$S_{\text{YM}} = \int_M L_{\text{YM}}$	$S_{\text{YM}} = \int_M L_{\text{YM}}$	$S_{\text{YM}} = \int_M L_{\text{YM}}$
mielke-geometrodynamics_48 EQ0072 (2.8.15)	48	■ 1.000	$\omega \sim \tilde{\omega}$	$\omega \sim \tilde{\omega}$	$\omega \sim \tilde{\omega}$
mielke-geometrodynamics_49 EQ0074 (2.8.17)	49	■ 1.000	$\Omega(\tilde{\omega}) = 0.$	$\Omega(\tilde{\omega}) = 0.$	$\Omega(\tilde{\omega}) = 0.$
mielke-geometrodynamics_49 EQ0075 (2.8.18)	49	■ 1.000	$c_2(M) = \frac{1}{24\pi^2} \int_{\partial M} \text{Tr}(\tilde{\omega} \wedge \tilde{\omega} \wedge \tilde{\omega}),$	$c_2(M) = \frac{1}{24\pi^2} \int_{\partial M} \text{Tr}(\tilde{\omega} \wedge \tilde{\omega} \wedge \tilde{\omega}),$	$c_2(M) = \frac{1}{24\pi^2} \int_{\partial M} \text{Tr}(\tilde{\omega} \wedge \tilde{\omega} \wedge \tilde{\omega}),$
mielke-geometrodynamics_52 EQ0076 (3.1.1)	52	■ 1.000	$L = L(\varphi, d\varphi)\eta$	$L = L(\varphi, d\varphi)\eta$	$L = L(\varphi, d\varphi)\eta$
mielke-geometrodynamics_52 EQ0077 (3.1.2)	52	■ 1.000	$\delta \int_M L = 0$	$\delta \int_M L = 0$	$\delta \int_M L = 0$
mielke-geometrodynamics_52 EQ0078 (3.1.3)	52	■ 1.000	$\delta\varphi = 0 \quad \text{at the boundary } \partial M \text{ of } M$	$\delta\varphi = 0 \quad \text{at the boundary } \partial M \text{ of } M$	$\delta\varphi = 0 \quad \text{at the boundary } \partial M \text{ of } M$
mielke-geometrodynamics_52 EQ0079 (3.1.4)	52	■ 1.000	$\delta(d\varphi) := d(\varphi + \delta\varphi) - d\varphi = d\delta\varphi$	$\delta(d\varphi) := d(\varphi + \delta\varphi) - d\varphi = d\delta\varphi$	$\delta(d\varphi) := d(\varphi + \delta\varphi) - d\varphi = d\delta\varphi$
mielke-geometrodynamics_52 EQ0080 (3.1.5)	52	■ 1.000	$\pi := \frac{\partial L}{\partial(\partial_\alpha \varphi)} \vartheta^\alpha$	$\pi := \frac{\partial L}{\partial(\partial_\alpha \varphi)} \vartheta^\alpha$	$\pi := \frac{\partial L}{\partial(\partial_\alpha \varphi)} \vartheta^\alpha$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00082 (3.1.7)	53	■ 1.000	$\int_M d(\delta\varphi\wedge\pi)=\int_{\partial M}\delta\varphi\wedge\pi$	$\int_M d(\delta\varphi\wedge\pi)=\int_{\partial M}\delta\varphi\wedge\pi$	$\int_M d(\delta\varphi\wedge\pi)=\int_{\partial M}\delta\varphi\wedge\pi.$
mielke-geometrodynamics_E00083 (3.1.9)	53	■ 1.000	$\lambda^\varphi:=(-1)^sL_\varphi+{}^*d\pi=0,$	$\lambda^\varphi:=(-1)^sL_\varphi+{}^*d\pi=0,$	$\lambda^\varphi:=(-1)^sL_\varphi+{}^*d\pi=0,$
mielke-geometrodynamics_E00084 (3.1.10)	53	■ 1.000	$\frac{\partial L}{\partial\varphi}-\partial_\alpha\frac{\partial L}{\partial(\partial_\alpha\varphi)}=0.$	$\frac{\partial L}{\partial\varphi}-\partial_\alpha\frac{\partial L}{\partial(\partial_\alpha\varphi)}=0.$	$\frac{\partial L}{\partial\varphi}-\partial_\alpha\frac{\partial L}{\partial(\partial_\alpha\varphi)}=0.$
mielke-geometrodynamics_E00085 (3.1.11)	53	■ 1.000	$\{ \}^{\ast}H:=L-d\,\varphi\wedge\pi$	${}^*H:=L-d\varphi\wedge\pi$	${}^*H:=L-d\varphi\wedge\pi$
mielke-geometrodynamics_E00086 (3.1.12)	53	■ 1.000	$d^{\ast}H=dL+d\varphi\wedge{}^*L_\varphi+d\varphi\wedge d\pi-dd\varphi\wedge\pi$	$d^*H=dL+d\varphi\wedge{}^*L_\varphi+d\varphi\wedge d\pi-dd\varphi\wedge\pi$	$d^*H=dL+d\varphi\wedge{}^*L_\varphi+d\varphi\wedge d\pi-dd\varphi\wedge\pi$
mielke-geometrodynamics_E00087 (3.1.13)	53	■ 1.000	$d^{\ast}H\,\simeq\,0$	$d^*H\simeq 0$	$d^*H\simeq 0$
mielke-geometrodynamics_E00088 (3.1.14)	54	■ 1.000	$\beta\wedge{}^*D\psi=m\psi\eta.$	$\beta\wedge{}^*D\psi=m\psi\eta.$	$\beta\wedge{}^*D\psi=m\psi\eta.$
mielke-geometrodynamics_E00089 (3.1.15)	54	■ 1.000	$\square\varphi+m^2\varphi=0$	$\square\varphi+m^2\varphi=0$	$\square\varphi+m^2\varphi=0$
mielke-geometrodynamics_E00090 (3.1.16)	54	■ 1.000	$\overline{L}_{\mathrm{mat}}(\overline{\psi}\psi,\overline{\psi}\beta\cdot d\psi)\eta=\overline{\psi}\beta\wedge{}^*d\psi-m\overline{\psi}\psi\eta.$	$\overline{L}_{\mathrm{mat}}=\overline{L}_{\mathrm{mat}}(\overline{\psi}\psi,\overline{\psi}\beta\cdot d\psi)\eta=\overline{\psi}\beta\wedge{}^*d\psi-m\overline{\psi}\psi\eta.$	$\overline{L}_{\mathrm{mat}}=\overline{L}_{\mathrm{mat}}(\overline{\psi}\psi,\overline{\psi}\beta\cdot d\psi)\eta=\overline{\psi}\beta\wedge{}^*d\psi-m\overline{\psi}\psi\eta.$
mielke-geometrodynamics_E00092 (3.2.2)	55	■ 1.000	$L_{\mathrm{mat}}=L(\overline{\varphi}\varphi,\overline{\varphi}\beta\cdot(d+\omega')\varphi)\eta=L(\overline{\varphi}\varphi,\overline{\varphi}\beta\cdot D'\varphi)\eta$	$L_{\mathrm{mat}}=L(\overline{\varphi}\varphi,\overline{\varphi}\beta\cdot(d+\omega')\varphi)\eta=L(\overline{\varphi}\varphi,\overline{\varphi}\beta\cdot D'\varphi)\eta$	$L_{\mathrm{mat}}=L(\overline{\varphi}\varphi,\overline{\varphi}\beta\cdot(d+\omega')\varphi)\eta=L(\overline{\varphi}\varphi,\overline{\varphi}\beta\cdot D'\varphi)\eta$
mielke-geometrodynamics_E00094 (3.2.4)	55	■ 1.000	$D\left(\{ \}^{\ast}G\right)b=\{ \}^{\ast}G\,\omega\otimes\{ \}^{\ast}G\,b\,.$	$D\left({}^G b\right)={}^G\omega\otimes{}^G b.$	$D({}^G b)={}^G\omega\otimes{}^G b.$
mielke-geometrodynamics_E00095 (3.2.5)	55	■ 1.000	$\begin{aligned} D\left(\{ \}^{\ast}G\right)b&\&=d\,G\,\otimes b+G\,\omega\otimes b\\ &=\left\{d\,G\,G^{-1}+G\,\omega\,G^{-1}\right\}\otimes{}^G b. \end{aligned}$	$D\left({}^G b\right)=dG\otimes b+G\omega\otimes b\\ =\left\{dGG^{-1}+G\omega G^{-1}\right\}\otimes{}^G b.$	$D({}^G b)=dG\otimes b+G\omega\otimes b\\ =\left\{dGG^{-1}+G\omega G^{-1}\right\}\otimes{}^G b.$
mielke-geometrodynamics_E00096 (3.2.6)	55	■ 1.000	$\omega\rightarrow{}^G\omega=G\omega G^{-1}+dG\,G^{-1}.$	$\omega\rightarrow{}^G\omega=G\omega G^{-1}+dG\,G^{-1}.$	$\omega\rightarrow{}^G\omega=G\omega G^{-1}+dG\,G^{-1}.$
mielke-geometrodynamics_E00102 (3.3.3)	57	■ 1.000	$\theta=-\frac{e}{\hbar}\int\mathbf{A}\cdot d\mathbf{l}$	$\theta=-\frac{e}{\hbar}\int\mathbf{A}\cdot d\mathbf{l}$	$\theta=-\frac{e}{\hbar}\int\mathbf{A}\cdot d\mathbf{l}.$
mielke-geometrodynamics_E00103 (3.3.4)	57	■ 1.000	$L=L(A,d\,A)+L_{\mathrm{mat}}(A,d\,A,\Psi,d\,\Psi)\,.$	$L=L(A,dA)+L_{\mathrm{mat}}(A,dA,\Psi,d\Psi).$	$L=L(A,dA)+L_{\mathrm{mat}}(A,dA,\Psi,d\Psi).$
mielke-geometrodynamics_E00104 (3.3.5)	57	■ 1.000	$\frac{\delta L}{\delta A}:=\frac{\partial L}{\partial A}+d\frac{\partial L}{\partial dA}=0,$	$\frac{\delta L}{\delta A}:=\frac{\partial L}{\partial A}+d\frac{\partial L}{\partial dA}=0,$	$\frac{\delta L}{\delta A}:=\frac{\partial L}{\partial A}+d\frac{\partial L}{\partial dA}=0,$
mielke-geometrodynamics_E00106 (3.3.7)	58	■ 1.000	$dF\equiv 0,\quad dH=j$	$dF\equiv 0,\quad dH=j$	$dF\equiv 0,\quad dH=j\,.$
mielke-geometrodynamics_E00107 (3.3.8)	58	■ 1.000	$dj\,\simeq\,0\,.$	$dj\simeq 0.$	$dj\simeq 0\,.$
mielke-geometrodynamics_E00111 (3.3.12)	58	■ 1.000	$H:=-\mathfrak{H}\wedge dt+\mathfrak{D}=\frac{1}{2}H_{ij}dx^i\wedge dx^j$	$H:=-\mathfrak{H}\wedge dt+\mathfrak{D}=\frac{1}{2}H_{ij}dx^i\wedge dx^j$	$H:=-\mathfrak{H}\wedge dt+\mathfrak{D}=\frac{1}{2}H_{ij}dx^i\wedge dx^j\,,$
mielke-geometrodynamics_E00114 (3.3.15)	59	■ 1.000	$H_{mn}=\frac{\sqrt{ g }}{2}\varepsilon_{mnkl}g^{ki}g^{lj}F_{ij}\,.$	$H_{mn}=\frac{\sqrt{ g }}{2}\varepsilon_{mnkl}g^{ki}g^{lj}F_{ij}.$	$H_{mn}=\frac{\sqrt{ g }}{2}\varepsilon_{mnkl}g^{ki}g^{lj}F_{ij}\,.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_60 EQ0121 (3.3.22)	60	■ 1.000	$L_{\mathrm{BF}}=-B \wedge F=-B \wedge d A$.	$L_{\mathrm{BF}}=-B \wedge F=-B \wedge d A$.	$L_{\mathrm{BF}}=-B \wedge F=-B \wedge d A$.
mielke-geometrodynamics_60 EQ0122 (3.3.23)	60	■ 1.000	$\begin{aligned} \tilde{L}_{\mathrm{BF}} &:= -B \wedge d A + \frac{1}{2} B \wedge B \\ &\cong -B \wedge d A + d C \end{aligned}$	$\tilde{L}_{\mathrm{BF}}:= -B \wedge d A + \frac{1}{2} B \wedge B \\ \cong -B \wedge d A + d C$	$\tilde{L}_{\mathrm{BF}}:= -B \wedge d A + \frac{1}{2} B \wedge B \\ \cong -B \wedge d A + d C$.
mielke-geometrodynamics_60 EQ0123 (3.3.24)	60	■ 1.000	$B \cong d A, \quad d B \cong d F \equiv 0$	$B \cong d A, \quad d B \cong d F \equiv 0$	$B \cong d A, \quad d B \cong d F \equiv 0$,
mielke-geometrodynamics_60 EQ0124 (3.3.25)	60	■ 1.000	$C=\frac{1}{2} A \wedge F, \quad d C=\frac{1}{2} F \wedge F$,	$C=\frac{1}{2} A \wedge F, \quad d C=\frac{1}{2} F \wedge F$,	$C=\frac{1}{2} A \wedge F, \quad d C=\frac{1}{2} F \wedge F$,
mielke-geometrodynamics_61 EQ0125 (3.3.26)	61	■ 1.000	$A \rightarrow A^{\prime}=A+d \theta(x)$.	$A \rightarrow A^{\prime}=A+d \theta(x)$.	$A \rightarrow A^{\prime}=A+d \theta(x)$.
mielke-geometrodynamics_61 EQ0126 (3.3.27)	61	■ 1.000	$A \rightarrow \widetilde{A}=A+\psi, \quad B \rightarrow \widetilde{B}=B+d \psi$,	$A \rightarrow \widetilde{A}=A+\psi, \quad B \rightarrow \widetilde{B}=B+d \psi$,	$A \rightarrow \widetilde{A}=A+\psi, \quad B \rightarrow \widetilde{B}=B+d \psi$,
mielke-geometrodynamics_61 EQ0127 (3.3.28)	61	■ 1.000	$L_{\mathrm{Max}}=-B \wedge d A+\frac{1}{2} B \wedge \wedge^{*} B+L_{\mathrm{matter}}$,	$L_{\mathrm{Max}}=-B \wedge d A+\frac{1}{2} B \wedge \wedge^{*} B+L_{\mathrm{matter}}$,	$L_{\mathrm{Max}}=-B \wedge d A+\frac{1}{2} B \wedge \wedge^{*} B+L_{\mathrm{matter}}$,
mielke-geometrodynamics_61 EQ0128 (3.3.29)	61	■ 1.000	$-d B=d^{*} F \cong j$,	$-d B=d^{*} F \cong j$,	$-d B=d^{*} F \cong j$,
mielke-geometrodynamics_63 EQ0132 (3.4.3)	63	■ 1.000	$\alpha_g:=g^2 / \hbar c$	$\alpha_g:=g^2 / \hbar c$	$\alpha_g:=g^2 / \hbar c$
mielke-geometrodynamics_63 EQ0134 (3.4.5)	63	■ 1.000	$\delta L_{\omega}:=\delta \Omega \wedge \Pi^{\Omega}$.	$\delta L_{\omega}:=\delta \Omega \wedge \Pi^{\Omega}$.	$\delta L_{\omega}:=\delta \Omega \wedge \Pi^{\Omega}$.
mielke-geometrodynamics_63 EQ0138 (3.4.9)	63	■ 1.000	$L=L_{\mathrm{mat}}+L_{\omega}$,	$L=L_{\mathrm{mat}}+L_{\omega}$,	$L=L_{\mathrm{mat}}+L_{\omega}$,
mielke-geometrodynamics_63 EQ0139 (3.4.10)	63	■ 1.000	$\delta L_{\mathrm{mat}}:=\delta \omega \wedge \tau$.	$\delta L_{\mathrm{mat}}:=\delta \omega \wedge \tau$.	$\delta L_{\mathrm{mat}}:=\delta \omega \wedge \tau$.
mielke-geometrodynamics_64 EQ0140 (3.4.11)	64	■ 1.000	$D \Pi^{\Omega}=\tau$	$D \Pi^{\Omega}=\tau$	$\underline{D \Pi^{\Omega}=\tau}$.
mielke-geometrodynamics_64 EQ0141 (3.4.12)	64	■ 1.000	$D \Omega \equiv 0$.	$D \Omega \equiv 0$.	$D \Omega \equiv 0$.
mielke-geometrodynamics_64 EQ0145 (3.4.16)	64	■ 1.000	$d \Pi^{\Omega}+\left[\omega, \Pi^{\Omega}\right]=\tau$.	$d \Pi^{\Omega}+\left[\omega, \Pi^{\Omega}\right]=\tau$.	$d \Pi^{\Omega}+\left[\omega, \Pi^{\Omega}\right]=\tau$.
mielke-geometrodynamics_65 EQ0147 (3.4.18)	65	■ 1.000	$Q_e=\frac{1}{2} Y+I_3$.	$Q_e=\frac{1}{2} Y+I_3$.	$Q_e=\frac{1}{2} Y+I_3$.
mielke-geometrodynamics_65 EQ0148 (3.4.19)	65	■ 1.000	$L_{\mathrm{YM}}=-\frac{1}{\alpha_g} \operatorname{Tr}\left(\Omega \wedge \wedge^{*} \Omega\right)$.	$L_{\mathrm{YM}}=-\frac{1}{\alpha_g} \operatorname{Tr}\left(\Omega \wedge \wedge^{*} \Omega\right)$.	$L_{\mathrm{YM}}=-\frac{1}{\alpha_g} \operatorname{Tr}\left(\Omega \wedge \wedge^{*} \Omega\right)$.
mielke-geometrodynamics_65 EQ0150 (3.4.21)	65	■ 1.000	$\Pi^{\Omega}=-\frac{2}{\alpha_g} \wedge^{*} \Omega$	$\Pi^{\Omega}=-\frac{2}{\alpha_g} \wedge^{*} \Omega$	$\Pi^{\Omega}=-\frac{2}{\alpha_g} \wedge^{*} \Omega$.
mielke-geometrodynamics_66 EQ0153 (3.5.1)	66	■ 1.000	$\Pi^{\Omega}=\zeta \Omega$.	$\Pi^{\Omega}=\zeta \Omega$.	$\Pi^{\Omega}=\zeta \Omega$.
mielke-geometrodynamics_67 EQ0156 (3.5.4)	67	■ 1.000	$\wedge^{*} \sigma^{\mu \nu}=\sigma^{\mu \nu}, \quad \wedge^{*} \bar{\sigma}^{\mu \nu}=-\bar{\sigma}^{\mu \nu},$ $\bar{\sigma}^{\mu \nu}=-\bar{\sigma}^{\mu \nu}$,	$\wedge^{*} \sigma^{\mu \nu}=\sigma^{\mu \nu}, \quad \wedge^{*} \bar{\sigma}^{\mu \nu}=-\bar{\sigma}^{\mu \nu},$	$\wedge^{*} \sigma^{\mu \nu}=\sigma^{\mu \nu}, \quad \wedge^{*} \bar{\sigma}^{\mu \nu}=-\bar{\sigma}^{\mu \nu},$
mielke-geometrodynamics_67 EQ0158 (3.5.6)	67	■ 1.000	$A_{\mu}=i \sigma_{\mu i} \partial^i \ln h$.	$A_{\mu}=i \sigma_{\mu i} \partial^i \ln h$.	$A_{\mu}=i \sigma_{\mu i} \partial^i \ln h$.
mielke-geometrodynamics_67 EQ0159 (3.5.7)	67	■ 1.000	$\square h-h^{-1} \partial_i h \partial^i h=h^3$	$\square h-h^{-1} \partial_i h \partial^i h=h^3$	$\square h-h^{-1} \partial_i h \partial^i h=h^3$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_67 EQ0160 (3.5.8)	67	■ 1.000	$h=\sum_{i=1}^{k+1} \frac{l_{(i)}^2}{\left(x-x_{(i)}\right)^2},$	$h=\sum_{i=1}^{k+1} \frac{l_{(i)}^2}{\left(x-x_{(i)}\right)^2},$	$h=\sum_{i=1}^{k+1} \frac{l_{(i)}^2}{\left(x-x_{(i)}\right)^2},$
mielke-geometrodynamics_67 EQ0161 (3.5.9)	67	■ 1.000	$p_{\{1\}}(M)=k \enspace .$	$p_1(M)=k.$	$p_1(M)=k.$
mielke-geometrodynamics_68 EQ0162 (3.5.10)	68	■ 1.000	$\left\langle \infty \mid -\infty \right\rangle =Z \int_{\mathcal{M}_\omega} d\omega \mu[\omega] \exp i \int_M (L_\omega + L_\infty).$	$\langle \infty \mid -\infty \rangle =Z \int_{\mathcal{M}_\omega} d\omega \mu[\omega] \exp i \int_M (L_\omega + L_\infty)$	$\langle \infty \mid -\infty \rangle =Z \int_{\mathcal{M}_\omega} d\omega \mu[\omega] \exp i \int_M (L_\omega + L_\infty).$
mielke-geometrodynamics_68 EQ0163 (3.5.11)	68	■ 1.000	$L_{\{\infty\}}=(\stackrel{+}{-})\frac{1}{\alpha_g}\operatorname{Tr}(\Omega\wedge\Omega)$	$L_\infty=(\stackrel{+}{-})\frac{1}{\alpha_g}\operatorname{Tr}(\Omega\wedge\Omega)$	$L_\infty=(\stackrel{+}{-})\frac{1}{\alpha_g}\operatorname{Tr}(\Omega\wedge\Omega)$
mielke-geometrodynamics_70 EQ0167 (3.6.2)	70	■ 1.000	$\mathscr{P}_{\{\pm\}}:=\bar{\psi}\sigma_{\pm}\psi/\sigma_{\pm}\psi/2m,$	$\mathscr{P}_{\pm}:=\bar{\psi}\sigma_{\pm}\psi/2m,$	$\mathscr{P}_{\pm}:=\overline{\psi}\sigma_{\pm}\psi/2m,$
mielke-geometrodynamics_70 EQ0168 (3.6.3)	70	■ 1.000	$D_{\{\pm\}}\psi\mp i\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)\wedge\sigma_{\pm}\psi=0$	$D_{\pm}^*D_{\pm}\psi\mp i\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)\wedge\sigma_{\pm}\psi=0$	$D_{\pm}^*D_{\pm}\psi\mp i\left(F^{\pm}-\frac{1}{2}\bar{\psi}\sigma_{\pm}\psi\right)\wedge\sigma_{\pm}\psi=0$
mielke-geometrodynamics_70 EQ0170 (3.6.5)	70	■ 1.000	$D^*\psi=\left(\nabla^*\nabla+\frac{1}{4}R+\frac{1}{2}F^+\right)\psi.$	$D^*D\psi=\left(\nabla^*\nabla+\frac{1}{4}R+\frac{1}{2}F^+\right)\psi.$	$D^*D\psi=\left(\nabla^*\nabla+\frac{1}{4}R+\frac{1}{2}F^+\right)\psi.$
mielke-geometrodynamics_70 EQ0172 (3.6.7)	70	■ 1.000	$\Omega_{\{\pm\}}^g=\frac{1}{2\ell^2}\sigma_{\pm},$	$\Omega_{\pm}^g=\frac{1}{2\ell^2}\sigma_{\pm},$	$\Omega_{\pm}^g=\frac{1}{2\ell^2}\sigma_{\pm},$
mielke-geometrodynamics_72 EQ0176 (3.7.5)	72	■ 1.000	$\begin{aligned} & D^*\Theta^{\{1\}}-\frac{\lambda}{2}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)\varphi\eta=0, \\ & D\Pi^{\Omega}+\left[\phi^{(0)},*\Theta^{\{1\}}\right]=0. \end{aligned}$	$D^*\Theta^{\{1\}}-\frac{\lambda}{2}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)\varphi\eta=0,$ $D\Pi^{\Omega}+\left[\phi^{(0)},*\Theta^{\{1\}}\right]=0.$	$D^*\Theta^{\{1\}}-\frac{\lambda}{2}\left(\frac{\mu^2}{\lambda}-\bar{\varphi}\cdot\varphi\right)\varphi\eta=0,$ $D\Pi^{\Omega}+\left[\phi^{(0)},*\Theta^{\{1\}}\right]=0.$
mielke-geometrodynamics_72 EQ0177 (3.7.7)	72	■ 1.000	$\bar{\varphi}\cdot\varphi:=\varphi^{*j}\varphi_j=\frac{\mu^2}{\lambda}.$	$\bar{\varphi}\cdot\varphi:=\varphi^{*j}\varphi_j=\frac{\mu^2}{\lambda}.$	$\bar{\varphi}\cdot\varphi:=\varphi^{*j}\varphi_j=\frac{\mu^2}{\lambda}.$
mielke-geometrodynamics_72 EQ0179 (3.7.9)	72	■ 1.000	$\varphi_{\min}:P\rightarrow C^\infty\left(\mathring{V}(M,\rho(G/H),\rho(G),P)\subset V^\rho\right).$	$\varphi_{\min}:P\rightarrow C^\infty\left(\mathring{V}(M,\rho(G/H),\rho(G),P)\subset V^\rho\right).$	$\varphi_{\min}:P\rightarrow C^\infty\left(\mathring{V}(M,\rho(G/H),\rho(G),P)\subset V^\rho\right).$
mielke-geometrodynamics_73 EQ0180 (3.7.10)	73	■ 1.000	$H:=\left\{h\in G\mid \rho(h)\mathrm{v}_0=\mathrm{v}_0\right\},$	$H:=\left\{h\in G\mid \rho(h)\mathrm{v}_0=\mathrm{v}_0\right\},$	$H:=\left\{h\in G \rho(h)\mathrm{v}_0=\mathrm{v}_0\right\},$
mielke-geometrodynamics_73 EQ0181 (3.7.11)	73	■ 1.000	$Q=\left\{p\in P\mid \varphi(p)=\sigma(\mathrm{v}_0)\right\}$	$Q=\left\{p\in P\mid \varphi(p)=\sigma(\mathrm{v}_0)\right\}$	$Q=\left\{p\in P \varphi(p)=\sigma(\mathrm{v}_0)\right\}$
mielke-geometrodynamics_73 EQ0182 (3.7.12)	73	■ 1.000	$D\phi^{\{0\}}=0.$	$D\phi^{(0)}=0.$	$D\phi^{\{0\}}=0.$
mielke-geometrodynamics_73 EQ0183 (3.7.13)	73	■ 1.000	$\begin{aligned} \Theta^{\perp} &:= \Omega_0 = d\omega_0 - (\omega \wedge \omega)_0 = d\omega_0 - \omega_0 \wedge \omega - \omega \wedge \omega_0 \\ &= d\omega_0 - [\omega_0, \omega] = D\omega_0 \end{aligned}$	$\Theta^{\perp}:=\Omega_0=d\omega_0-(\omega\wedge\omega)_0=d\omega_0-\omega_0\wedge\omega-\omega\wedge\omega_0$ $=d\omega_0-[\omega_0,\omega]=D\omega_0$	$\Theta^{\perp}:=\Omega_0=d\omega_0-(\omega\wedge\omega)_0=d\omega_0-\omega_0\wedge\omega-\omega\wedge\omega_0$ $=d\omega_0-[\omega_0,\omega]=D\omega_0$
mielke-geometrodynamics_74 EQ0185 (3.7.15)	74	■ 1.000	$d_{\{0\}}\Omega_0=0.$	$d_0^*\Omega_0=0.$	$d_0^*\Omega_0=0.$
mielke-geometrodynamics_74 EQ0186 (3.7.16)	74	■ 1.000	$\left(D^*\Omega\right)_0=\vec{D}^*\Omega_0=\vec{D}^*\Theta^{\perp}=0$	$(D^*\Omega)_0=\vec{D}^*\Omega_0=\vec{D}^*\Theta^{\perp}=0$	$(D^*\Omega)_0=\vec{D}^*\Omega_0=\vec{D}^*\Theta^{\perp}=0$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0188 (3.7.18)	74	■ 1.000	$\omega_{\{0\}}=\varphi^{\{j\}}L_{\{j\}}\otimes\vartheta_{\{0\}}.$	$\omega_0=\varphi^jL_j\otimes\vartheta_0.$	$\omega_0=\varphi^jL_j\otimes\vartheta_0.$
mielke-geometrodynamics_EQ0190	80	■ 1.000	$\left[\begin{array}{l} T_{\{G\}}:\\ \quad T_{\{g\}}\,\psi(m)=D\left(g_{\{m\}}^{-1}\,g_{g_{\{g\}}^{-1}\,m},\,m_{\{0\}}\right)\,\psi\left(g^{-1}m\right)\\ \cdot\,\left[\right]\end{array}\right]$	$T_G:\quad T_g\psi(m)=D\left(g_m^{-1}gg_{g^{-1}m},m_0\right)\psi\left(g^{-1}m\right).$	$T_G:\quad T_g\psi(m)=D\left(g_m^{-1}gg_{g^{-1}m},m_0\right)\psi(g^{-1}m).$
mielke-geometrodynamics_EQ0193 (4.1.3)	81	■ 1.000	$A(n,\,\mathbb{R})=\mathbb{R}^n\otimes GL(n,\mathbb{R})$	$A(n,\mathbb{R})=\mathbb{R}^n\otimes GL(n,\mathbb{R})$	$A(n,\mathbb{R})=\mathbb{R}^n\otimes GL(n,\mathbb{R})$
mielke-geometrodynamics_EQ0194 (4.1.4)	81	■ 1.000	$P:=\mathbb{R}^n\otimes\mathbb{O}(1,\,n-1)\subset A(n,\,\mathbb{R}),$	$P:=\mathbb{R}^n\otimes\mathbb{O}(1,n-1)\subset A(n,\mathbb{R}),$	$P:=\mathbb{R}^n\otimes\mathbb{O}(1,n-1)\subset A(n,\mathbb{R}),$
mielke-geometrodynamics_EQ0197 (4.1.10)	82	■ 1.000	$e_{\{\alpha\}}(m)=e_{\{\alpha\}}^{\{i\}}\,\partial_{\{i\}}$	$e_{\alpha}(m)=e_{\alpha}^i\partial_i$	$e_{\alpha}(m)=e_{\alpha}^i\partial_i$
mielke-geometrodynamics_EQ0199 (4.1.13)	82	■ 1.000	$L(M)=\bigcup_{m\in M}L_{\{m\}}(M)$	$L(M)=\bigcup_{m\in M}L_m(M)$	$L(M)=\bigcup_{m\in M}L_m(M)$
mielke-geometrodynamics_EQ0200 (4.1.14)	83	■ 1.000	$L(M)=P(M,\,GL(n,\,\mathbb{R})),\,\pi,\,\delta)$	$L(M)=P(M, GL(n,\mathbb{R}),\pi,\delta)$	$L(M)=P(M, GL(n,\mathbb{R}),\pi,\delta)$
mielke-geometrodynamics_EQ0202 (4.1.16)	83	■ 1.000	$\vartheta(e)=0\,\Leftrightarrow\pi(e)=0,\,e\in T_{\{p\}}(P),$	$\vartheta(e)=0\,\Leftrightarrow\pi(e)=0,e\in T_p(P),$	$\vartheta(e)=0\,\Leftrightarrow\pi(e)=0,e\in T_p(P),$
mielke-geometrodynamics_EQ0204 (4.1.18)	83	■ 1.000	$T(M)=V\left(M,\,\mathbb{R}^n,\,GL(n,\,\mathbb{R}),\,L(M)\right)$	$T(M)=V\left(M,\mathbb{R}^n, GL(n,\mathbb{R}),L(M)\right)$	$T(M)=V(M,\mathbb{R}^n, GL(n,\mathbb{R}),L(M))$
mielke-geometrodynamics_EQ0205 (4.1.19)	83	■ 1.000	$A(M):=P(M,\,A(n,\,\mathbb{R})),\,\pi,\,\delta)=L(M)\times\mathbb{R}^n.$	$A(M):=P(M,A(n,\mathbb{R}),\pi,\delta)=L(M)\times\mathbb{R}^n.$	$A(M):=P(M,A(n,\mathbb{R}),\pi,\delta)=L(M)\times\mathbb{R}^n.$
mielke-geometrodynamics_EQ0206 (4.1.20)	83	■ 1.000	$\beta:A(n,\,\mathbb{R})\rightarrow GL(n,\mathbb{R})=A(n,\mathbb{R})/\mathbb{R}^n.$	$\beta:A(n,\mathbb{R})\rightarrow GL(n,\mathbb{R})=A(n,\mathbb{R})/\mathbb{R}^n.$	$\beta:A(n,\mathbb{R})\rightarrow GL(n,\mathbb{R})=A(n,\mathbb{R})/\mathbb{R}^n.$
mielke-geometrodynamics_EQ0213 (4.2.5)	86	■ 1.000	$\mathscr{A}(n,\,\mathbb{R}):=C^{\infty}\left(A(M)\times_{\mathrm{Ad}}A(n,\,\mathbb{R})\right).$	$\mathscr{A}(n,\mathbb{R}):=C^{\infty}\left(A(M)\times_{\mathrm{Ad}}A(n,\mathbb{R})\right).$	$\mathscr{A}(n,\,\mathbb{R}):=C^{\infty}\left(A(M)\times_{\mathrm{Ad}}A(n,\,\mathbb{R})\right).$
mielke-geometrodynamics_EQ0223 (4.2.16)	87	■ 1.000	$Q_{\{A\}}=\left\{p\in A(M)\, \,\widetilde{\varphi}(p)=\sigma(0_m)\right\}=L(M)$	$Q_A=\left\{p\in A(M)\, \,\widetilde{\varphi}(p)=\sigma(0_m)\right\}=L(M)$	$Q_A=\{p\in A(M) \widetilde{\varphi}(p)=\sigma(0_m)\}=L(M)$
mielke-geometrodynamics_EQ0237 (4.3.2)	90	■ 1.000	$\ell(\mathscr{C}):=\int_{t_0}^{t_1}ds(t)$	$\ell(\mathscr{C}):=\int_{t_0}^{t_1}ds(t)$	$\ell(\mathscr{C}):=\int_{t_0}^{t_1}ds(t)$
mielke-geometrodynamics_EQ0238 (4.3.3)	90	■ 1.000	$\ell\left(\mathscr{C}_1+\mathscr{C}_2\right)=\ell\left(\mathscr{C}_1\right)+\ell\left(\mathscr{C}_2\right)$	$\ell\left(\mathscr{C}_1+\mathscr{C}_2\right)=\ell\left(\mathscr{C}_1\right)+\ell\left(\mathscr{C}_2\right)$	$\ell(\mathscr{C}_1+\mathscr{C}_2)=\ell(\mathscr{C}_1)+\ell(\mathscr{C}_2)$
mielke-geometrodynamics_EQ0239 (4.3.4)	90	■ 1.000	$ds=F(m,\vartheta)dt,$	$ds=F(m,\vartheta)dt,$	$ds=F(m,\vartheta)dt,$
mielke-geometrodynamics_EQ0242 (4.3.7)	91	■ 1.000	$ds^2=(F(m,\vartheta)dt)^2=g_{\{\alpha\}\{\beta\}}\vartheta^{\{\alpha\}}\otimes\vartheta^{\{\beta\}}=g_{\{i\}\{j\}}(m)\,dx^{\{i\}}\otimes_sdx^{\{j\}};$	$ds^2=(F(m,\vartheta)dt)^2=g_{\alpha\beta}\vartheta^{\alpha}\otimes\vartheta^{\beta}=g_{ij}(m)dx^i\otimes_sdx^j;$	$ds^2=(F(m,\vartheta)dt)^2=g_{\alpha\beta}\vartheta^{\alpha}\otimes\vartheta^{\beta}=g_{ij}(m)dx^i\otimes_sdx^j;$
mielke-geometrodynamics_EQ0246 (4.3.11)	92	■ 1.000	$L^{\{g\}}(M):=P(M,\,\mathbb{O}(s,\,n-s),\,\pi,\,\delta).$	$L^g(M):=P(M,O(s,n-s),\pi,\delta).$	$L^g(M):=P(M,O(s,n-s),\pi,\delta).$
mielke-geometrodynamics_EQ0247 (4.3.12)	92	■ 1.000	$D\,e_{\{\alpha\}}^{\{\cdot\}}=0$	$De_{\alpha}^{\cdot}=0$	$D\,e_{\alpha}^{\cdot}=0$
mielke-geometrodynamics_EQ0251 (4.3.16)	92	■ 1.000	$\Gamma_{\{i\}}^{\cdot\alpha\beta}=-\Gamma_{\{i\}}^{\cdot\beta\alpha}$	$\Gamma_i^{\cdot\alpha\beta}=-\Gamma_i^{\cdot\beta\alpha}$	$\Gamma_i^{\cdot\alpha\beta}=-\Gamma_i^{\cdot\beta\alpha}$
mielke-geometrodynamics_EQ0252 (4.3.17)	93	■ 1.000	$g_{\{i\}\{j\}}\stackrel{(*)}{=}o_{\{i\}\{j\}}$	$g_{ij}\stackrel{(*)}{=}o_{ij}$	$g_{ij}\stackrel{(*)}{=}o_{ij}$
mielke-geometrodynamics_EQ0253 (4.3.18)	93	■ 1.000	$\partial_{\{k\}}g_{\{i\}\{j\}}\stackrel{(*)}{=}0.$	$\partial_kg_{ij}\stackrel{(*)}{=}0.$	$\partial_kg_{ij}\stackrel{(*)}{=}0.$
mielke-geometrodynamics_EQ0254 (4.3.19)	93	■ 1.000	$\Omega^g=d\omega^g-\omega^g\wedge\omega^g$	$\Omega^g=d\omega^g-\omega^g\wedge\omega^g$	$\Omega^g=d\omega^g-\omega^g\wedge\omega^g$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0257 (4.3.22)	94	■ 1.000	$\backslash\omega^{\mathrm{g}}=\backslash\omega^{\{\}\mathrm{}}\cdot K$.	$\omega^g = \omega^{\{\}} - K.$	$\omega^g = \omega^{\{\}} - K.$
mielke-geometrodynamics_EQ0258 (4.3.23)	94	■ 1.000	$\mathrm{d}\backslash\vartheta\mathrm{-}\backslash\left[\backslash\omega^{\{\}\mathrm{}},\backslash\vartheta\mathrm{right}\right]=0$	$d\vartheta - \left[\omega^{\{\}},\vartheta\right] = 0$	$d\vartheta - [\omega^{\{\}},\vartheta] = 0$
mielke-geometrodynamics_EQ0259 (4.3.24)	94	■ 1.000	$\backslash\Theta=[K,\backslash\vartheta\mathrm{a}]$	$\Theta = [K,\vartheta]$	$\Theta = [K,\vartheta]$
mielke-geometrodynamics_EQ0265 (4.3.30)	95	■ 1.000	$\backslash\left\{\backslash\begin{array}{c}\kappa\\i\ j\end{array}\right\}\backslash\mathrm{g}^{\{\kappa\ l\}}\backslash\left(\backslash\partial_{\mathrm{i}}\backslash\mathrm{g}_{\{\mathrm{j}\ l\}}+\backslash\partial_{\mathrm{j}}\backslash\mathrm{g}_{\{\mathrm{i}\ l\}}-\backslash\partial_{\mathrm{l}}\backslash\mathrm{g}_{\{\mathrm{i}\ j\}}\right)$	$\left\{\begin{array}{c}\kappa\\i\ j\end{array}\right\} = \frac{1}{2}g^{\kappa l}(\partial_i g_{jl} + \partial_j g_{il} - \partial_l g_{ij})$	$\left\{\begin{array}{c}\kappa\\i\ j\end{array}\right\} = \frac{1}{2}g^{\kappa l}(\partial_i g_{jl} + \partial_j g_{il} - \partial_l g_{ij})$
mielke-geometrodynamics_EQ0272 (4.4.1)	96	■ 1.000	$\mathrm{L}_{\{\mathrm{g}\}}=\mathrm{L}\backslash\left(\mathrm{g}_{\{\backslash\alpha\ \backslash\beta\}};\backslash\vartheta\mathrm{a},\backslash\left(\backslash\omega^{\mathrm{g}}\backslash\right),\backslash\Theta\mathrm{a},\backslash\Omega^{\mathrm{g}}\backslash\right)$.	$L_g = L(g_{\alpha\beta};\vartheta,(\omega^g),\Theta,\Omega^g).$	$L_g = L(g_{\alpha\beta};\vartheta,(\omega^g),\Theta,\Omega^g).$
mielke-geometrodynamics_EQ0273 (4.4.2)	97	■ 1.000	$\backslash\delta\mathrm{L}_{\{\mathrm{g}\}}=: \backslash\delta\mathrm{a}\backslash\vartheta\mathrm{a}\backslash\wedge\mathrm{E}+\backslash\delta\mathrm{a}\backslash\Theta\backslash\wedge\backslash\mathrm{Pi}^{\{\mathrm{mat}\mathrm{r}\mathrm{m}\{T\}\}}+\backslash\delta\mathrm{a}\backslash\Omega^{\mathrm{g}}\backslash\wedge\backslash\mathrm{Pi}^{\{\mathrm{mat}\mathrm{r}\mathrm{m}\{L\}\}},$	$\delta L_g =: \delta\vartheta \wedge E + \delta\Theta \wedge \Pi^T + \delta\Omega^g \wedge \Pi^L,$	$\delta L_g =: \delta\vartheta \wedge E + \delta\Theta \wedge \Pi^T + \delta\Omega^g \wedge \Pi^L,$
mielke-geometrodynamics_EQ0274 (4.4.7)	97	■ 1.000	$\backslash\delta\mathrm{a}\backslash\Theta\mathrm{a}=\backslash\delta\mathrm{a}\backslash\left(\mathrm{d}\backslash\vartheta\mathrm{a}\backslash\left[\backslash\omega^{\mathrm{L}}\backslash\right),\backslash\vartheta\mathrm{a}\backslash\right)\backslash\right)$	$\delta\Theta = \delta\left(d\vartheta - [\omega^L,\vartheta]\right)$	$\delta\Theta = \delta(d\vartheta - [\omega^L,\vartheta])$
mielke-geometrodynamics_EQ0277 (4.4.5)	97	■ 1.000	$\mathrm{S}=\backslash\mathrm{int}_{\mathrm{M}}\backslash\left(\mathrm{L}_{\{\mathrm{g}\}}+\mathrm{L}_{\{\mathrm{mat}\mathrm{r}\mathrm{m}\{mat\}\}}\backslash\right)$	$S = \int_M (L_g + L_{\mathrm{mat}})$	$S = \int_M (L_g + L_{\mathrm{mat}})$
mielke-geometrodynamics_EQ0279 (4.4.8)	98	■ 1.000	$\backslash\delta\mathrm{L}_{\{\backslash\mathrm{t}\mathrm{e}\mathrm{x}\mathrm{t}\ \{\mathrm{matter}\}\}}=: \backslash\delta\mathrm{a}\backslash\vartheta\mathrm{a}\backslash\wedge\backslash\Sigma+\backslash\delta\mathrm{a}\backslash\Omega^{\mathrm{g}}\backslash\wedge\backslash\tau_{\{\mathrm{s}\}}+\backslash\delta\mathrm{a}\backslash\backslash\varphi\mathrm{i}\backslash\wedge\backslash\lambda\mathrm{b}\mathrm{d}\mathrm{a}^{\{\backslash\varphi\mathrm{i}\}}$	$\delta L_{\mathrm{matter}} =: \delta\vartheta \wedge \Sigma + \delta\omega^g \wedge \tau_s + \delta_\varphi \wedge \lambda^\varphi$	$\delta L_{\mathrm{matter}} =: \delta\vartheta \wedge \Sigma + \delta\omega^g \wedge \tau_s + \delta_\varphi \wedge \lambda^\varphi$
mielke-geometrodynamics_EQ0280 (4.4.9)	98	■ 1.000	$\mathrm{D}\backslash\Theta\mathrm{a}\backslash\equiv\backslash\left[\backslash\vartheta\mathrm{a},\backslash\Omega^{\mathrm{L}}\backslash\right)$	$D\Theta \equiv [\vartheta,\Omega^L]$	$D\Theta \equiv [\vartheta,\Omega^L]$
mielke-geometrodynamics_EQ0281 (4.4.10)	98	■ 1.000	$\mathrm{D}\backslash\Omega^{\mathrm{g}}\backslash\equiv\backslash 0,$	$D\Omega^g \equiv 0,$	$D\Omega^g \equiv 0,$
mielke-geometrodynamics_EQ0283 (4.4.12)	99	■ 1.000	$\mathrm{D}[\backslash\Theta\mathrm{a},\backslash\vartheta\mathrm{a}]\backslash\equiv\backslash\left[\backslash\vartheta\mathrm{a}\backslash\wedge\backslash\vartheta\mathrm{a},\backslash\Omega^{\mathrm{g}}\backslash\right)$	$D[\Theta,\vartheta] \equiv [\vartheta \wedge \vartheta,\Omega^g]$	$D[\Theta,\vartheta] \equiv [\vartheta \wedge \vartheta,\Omega^g]$
mielke-geometrodynamics_EQ0289 (4.4.18)	100	■ 1.000	$\backslash\left[\backslash\Omega^{\mathrm{g}}\backslash,\backslash\mathrm{Pi}^{\{\mathrm{L}\}}\backslash\right)+\backslash\mathrm{mat}\mathrm{r}\mathrm{m}\{\mathrm{E}\}_{\{\{\}\}}=0$	$[\Omega^g,\Pi^L] + \mathrm{E}_{\square} = 0$	$[\Omega^g,\Pi^L] + \mathrm{E}_{\square} = 0$
mielke-geometrodynamics_EQ0290 (4.5.1)	100	■ 1.000	$\mathrm{L}_{\{\mathrm{mat}\mathrm{r}\mathrm{m}\{mat\}\}}=\mathrm{L}\backslash\left(\mathrm{g}_{\{\backslash\alpha\ \backslash\beta\}},\backslash\vartheta\mathrm{a}^{\{\backslash\alpha\}},\backslash\mathrm{Psi},\mathrm{D}\backslash\mathrm{Psi}\backslash\right)$.	$L_{\mathrm{mat}} = L(g_{\alpha\beta},\vartheta^\alpha,\Psi,D\Psi).$	$L_{\mathrm{mat}} = L(g_{\alpha\beta},\vartheta^\alpha,\Psi,D\Psi).$
mielke-geometrodynamics_EQ0291 (4.5.2)	100	■ 1.000	$\backslash\Sigma_{\{\backslash\alpha\}}:=\backslash\frac{\{\partial\}\backslash\mathrm{L}\{\}\{\partial\}\backslash\vartheta\mathrm{a}^{\{\backslash\alpha\}}}{\{\partial\}\backslash\vartheta^{\backslash\alpha}},\backslash\mathrm{quad}\backslash\sigma\mathrm{a}^{\{\backslash\alpha\ \backslash\beta\}}:=2\backslash\frac{\{\partial\}\backslash\mathrm{L}\{\}\{\partial\}\backslash\mathrm{g}_{\{\backslash\alpha\ \backslash\beta\}}}{\{\partial\}\backslash\mathrm{g}_{\backslash\alpha\ \backslash\beta}},\backslash\mathrm{quad}\backslash\mathrm{t}\mathrm{e}\mathrm{x}\mathrm{t}\ \{\mathrm{and}\}\backslash\mathrm{quad}\backslash\tau_{\{\backslash\alpha\ \backslash\beta\}}:=\backslash\frac{\{\partial\}\backslash\mathrm{L}\{\}\{\partial\}\backslash\Gamma^{\{\backslash\alpha\ \backslash\beta\}}}{\{\partial\}\backslash\Gamma^{\backslash\alpha\ \backslash\beta}}=\backslash\mathrm{I}_{\{\backslash\alpha\ \backslash\beta\}}\backslash\mathrm{Psi}\backslash\wedge\backslash\frac{\{\partial\}\backslash\mathrm{L}\{\}\{\partial\}\backslash\mathrm{D}\backslash\mathrm{Psi}\}}{\{\partial\}\backslash\mathrm{D}\backslash\mathrm{Psi}}$	$\Sigma_\alpha := \frac{\partial L}{\partial\vartheta^\alpha}, \quad \sigma^{\alpha\beta} := 2\frac{\partial L}{\partial g_{\alpha\beta}}, \quad \text{and} \quad \tau_{\alpha\beta} := \frac{\partial L}{\partial\Gamma^{\alpha\beta}} = I_{\alpha\beta}\Psi \wedge \frac{\partial L}{\partial D\Psi}$	$\Sigma_\alpha := \frac{\partial L}{\partial\vartheta^\alpha}, \quad \sigma^{\alpha\beta} := 2\frac{\partial L}{\partial g_{\alpha\beta}}, \quad \text{and} \quad \tau_{\alpha\beta} := \frac{\partial L}{\partial\Gamma^{\alpha\beta}} = I_{\alpha\beta}\Psi \wedge \frac{\partial L}{\partial D\Psi}$
mielke-geometrodynamics_EQ0297 (8.10)	101	■ 1.000	$\backslash\delta\mathrm{L}=\backslash\delta\mathrm{g}_{\{\backslash\alpha\ \backslash\beta\}}\backslash\wedge\backslash\frac{\{\partial\}\backslash\mathrm{L}\{\}\{\partial\}\backslash\mathrm{g}_{\{\backslash\alpha\ \backslash\beta\}}}{\{\partial\}\backslash\mathrm{g}_{\backslash\alpha\ \backslash\beta}}+\backslash\delta\mathrm{a}\backslash\vartheta\mathrm{a}^{\{\backslash\alpha\}}\backslash\wedge\backslash\frac{\{\partial\}\backslash\mathrm{L}\{\}\{\partial\}\backslash\vartheta\mathrm{a}^{\{\backslash\alpha\}}}{\{\partial\}\backslash\vartheta^{\backslash\alpha}}+\backslash\delta\mathrm{a}\backslash\mathrm{Psi}\backslash\wedge\backslash\frac{\{\partial\}\backslash\mathrm{L}\{\}\{\partial\}\backslash\mathrm{Psi}\}}{\{\partial\}\backslash\mathrm{Psi}}+\backslash\delta\mathrm{a}\backslash(\mathrm{D}\backslash\mathrm{Psi})\backslash\wedge\backslash\frac{\{\partial\}\backslash\mathrm{L}\{\}\{\partial\}\backslash\mathrm{D}\backslash\mathrm{Psi}\}}{\{\partial\}\backslash\mathrm{D}\backslash\mathrm{Psi}}$.	$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta\vartheta^\alpha \wedge \frac{\partial L}{\partial\vartheta^\alpha} + \delta\Psi \wedge \frac{\partial L}{\partial\Psi} + \delta(D\Psi) \wedge \frac{\partial L}{\partial D\Psi}.$	$\delta L = \delta g_{\alpha\beta} \wedge \frac{\partial L}{\partial g_{\alpha\beta}} + \delta\vartheta^\alpha \wedge \frac{\partial L}{\partial\vartheta^\alpha} + \delta\Psi \wedge \frac{\partial L}{\partial\Psi} + \delta(D\Psi) \wedge \frac{\partial L}{\partial D\Psi}.$
mielke-geometrodynamics_EQ0300 (4.5.11)	102	■ 1.000	$\backslash\delta\mathrm{L}=-\backslash\varepsilon_{\backslash\alpha}^{\backslash\beta}\backslash\left[-\sigma^{\alpha\gamma}\backslash\mathrm{g}_{\gamma\beta}+\vartheta^\alpha\backslash\wedge\backslash\frac{\{\partial\}\backslash\mathrm{L}\{\}\{\partial\}\backslash\mathrm{g}_{\{\backslash\gamma\ \backslash\beta\}}}{\{\partial\}\backslash\mathrm{g}_{\backslash\gamma\ \backslash\beta}}+\mathrm{D}\backslash\frac{\{\partial\}\backslash\mathrm{L}\{\}\{\partial\}\backslash\vartheta\mathrm{a}^{\{\backslash\alpha\}}}{\{\partial\}\backslash\vartheta^{\backslash\alpha}}+\mathrm{I}^{\alpha}_{\backslash\beta}\backslash\mathrm{Psi}\backslash\wedge\backslash\frac{\{\partial\}\backslash\mathrm{L}\{\}\{\partial\}\backslash\mathrm{Psi}\}}{\{\partial\}\backslash\mathrm{Psi}}\backslash\right)$.	$\delta L = -\varepsilon_\alpha{}^\beta \left[-\sigma^{\alpha\gamma} g_{\gamma\beta} + \vartheta^\alpha \wedge \frac{\partial L}{\partial\vartheta^\beta} + D\frac{\partial L}{\partial\Gamma_\alpha{}^\beta} + I^\alpha{}_\beta \Psi \wedge \frac{\delta L}{\delta\Psi} \right].$	$\delta L = -\varepsilon_\alpha{}^\beta \left[-\sigma^{\alpha\gamma} g_{\gamma\beta} + \vartheta^\alpha \wedge \frac{\partial L}{\partial\vartheta^\beta} + D\frac{\partial L}{\partial\Gamma_\alpha{}^\beta} + I^\alpha{}_\beta \Psi \wedge \frac{\delta L}{\delta\Psi} \right].$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_102 EQ0301 (4.5.12)	102	■ 1.000	$D \tau_{\alpha \beta} + \vartheta_{[\alpha} \wedge \Sigma_{\beta]} = -I_{\alpha \beta} \Psi \wedge \frac{\delta L}{\delta \Psi} \simeq 0.$	$D \tau_{\alpha \beta} + \vartheta_{[\alpha} \wedge \Sigma_{\beta]} = -I_{\alpha \beta} \Psi \wedge \frac{\delta L}{\delta \Psi} \simeq 0.$	
mielke-geometrodynamics_102 EQ0303 (4.5.13)	102	■ 1.000	$\begin{aligned} & R_{\alpha}{}^{\beta} = d\Gamma_{\alpha}{}^{\beta} \\ & - \Gamma_{\alpha}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\beta} \end{aligned}$	$R_{\alpha}{}^{\beta} = d\Gamma_{\alpha}{}^{\beta} - \Gamma_{\alpha}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\beta}$	$R_{\alpha}{}^{\beta} = d\Gamma_{\alpha}{}^{\beta} - \Gamma_{\alpha}{}^{\gamma} \wedge \Gamma_{\gamma}{}^{\beta}$
mielke-geometrodynamics_103 EQ0307 (4.5.17)	103	■ 1.000	$\hat{M} := M^3 / \left[M^2 + \frac{\Lambda}{3} J^2 \right], \quad \hat{J} := JM^2 / \left[M^2 + \frac{\Lambda}{3} J^2 \right],$	$\hat{M} := M^3 / \left[M^2 + \frac{\Lambda}{3} J^2 \right], \quad \hat{J} := JM^2 / \left[M^2 + \frac{\Lambda}{3} J^2 \right],$	$\hat{M} := M^3 / [M^2 + \frac{\Lambda}{3} J^2], \quad \hat{J} := JM^2 / [M^2 + \frac{\Lambda}{3} J^2],$
mielke-geometrodynamics_110 EQ0317 (5.2.1)	110	■ 1.000	$L_{\mathrm{D}} = L(\gamma, \psi, D\psi) = \frac{i}{2} \left\{ \bar{\psi}^* \gamma \wedge D\psi + \overline{D\psi} \wedge^* \gamma \psi \right\} - m \bar{\psi} \psi \eta,$	$L_{\mathrm{D}} = L(\gamma, \psi, D\psi) = \frac{i}{2} \left\{ \bar{\psi}^* \gamma \wedge D\psi + \overline{D\psi} \wedge^* \gamma \psi \right\} - m \bar{\psi} \psi \eta,$	$L_{\mathrm{D}} = L(\gamma, \psi, D\psi) = \frac{i}{2} \left\{ \bar{\psi}^* \gamma \wedge D\psi + \overline{D\psi} \wedge^* \gamma \psi \right\} - m \bar{\psi} \psi \eta,$
mielke-geometrodynamics_111 EQ0322 (5.2.6)	111	■ 1.000	$\left. dU = T^{\alpha} \wedge \mu_{\alpha} - \vartheta^{\alpha} \wedge D\mu_{\alpha} = \vartheta^{\alpha} \wedge [e_{\beta}] (T^{\beta} \wedge \mu_{\alpha}) - D\mu_{\alpha} \right $	$dU = T^{\alpha} \wedge \mu_{\alpha} - \vartheta^{\alpha} \wedge D\mu_{\alpha} = \vartheta^{\alpha} \wedge [e_{\beta}] (T^{\beta} \wedge \mu_{\alpha}) - D\mu_{\alpha}$	$dU = T^{\alpha} \wedge \mu_{\alpha} - \vartheta^{\alpha} \wedge D\mu_{\alpha} = \vartheta^{\alpha} \wedge [e_{\beta}] (T^{\beta} \wedge \mu_{\alpha}) - D\mu_{\alpha}$
mielke-geometrodynamics_111 EQ0324 (5.3.1)	111	■ 1.000	$L_{\mathrm{EC}} := -\frac{1}{2\kappa} R^{\alpha\beta} \wedge \eta_{\alpha\beta},$	$L_{\mathrm{EC}} := -\frac{1}{2\kappa} R^{\alpha\beta} \wedge \eta_{\alpha\beta},$	$L_{\mathrm{EC}} := -\frac{1}{2\kappa} R^{\alpha\beta} \wedge \eta_{\alpha\beta},$
mielke-geometrodynamics_112 EQ0325 (5.3.2)	112	■ 1.000	$G_{\alpha} := \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} = \kappa \Sigma_{\alpha},$	$G_{\alpha} := \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} = \kappa \Sigma_{\alpha},$	$G_{\alpha} := \frac{1}{2} R^{\beta\gamma} \wedge \eta_{\alpha\beta\gamma} = \kappa \Sigma_{\alpha},$
mielke-geometrodynamics_112 EQ0327 (5.3.4)	112	■ 1.000	$D^{\{\} \{ \}} G_{\alpha}^{\{\} \{ \}} \equiv 0$	$D^{\{\} \{ \}} G_{\alpha}^{\{\} \{ \}} \equiv 0$	$D^{\{\} \{ \}} G_{\alpha}^{\{\} \{ \}} \equiv 0$
mielke-geometrodynamics_112 EQ0331 (5.3.8)	112	■ 1.000	$D[\gamma, \Theta] \equiv 2i[\sigma, \Omega^g], \quad D[\gamma, \Omega^g] \equiv [\Theta, \Omega^g],$	$D[\gamma, \Theta] \equiv 2i[\sigma, \Omega^g], \quad D[\gamma, \Omega^g] \equiv [\Theta, \Omega^g],$	$D[\gamma, \Theta] \equiv 2i[\sigma, \Omega^g], \quad D[\gamma, \Omega^g] \equiv [\Theta, \Omega^g],$
mielke-geometrodynamics_113 EQ0332 (5.3.9)	113	■ 1.000	$DG \equiv i[\Theta, \gamma_5 \Omega^g]$	$DG \equiv i[\Theta, \gamma_5 \Omega^g]$	$DG \equiv i[\Theta, \gamma_5 \Omega^g]$
mielke-geometrodynamics_113 EQ0333 (5.3.10)	113	■ 1.000	$\eta_{\alpha\beta\gamma} \wedge T^{\gamma} = 2\kappa \tau_{\alpha\beta}$	$\eta_{\alpha\beta\gamma} \wedge T^{\gamma} = 2\kappa \tau_{\alpha\beta}$	$\eta_{\alpha\beta\gamma} \wedge T^{\gamma} = 2\kappa \tau_{\alpha\beta}$
mielke-geometrodynamics_113 EQ0334 (5.3.11)	113	■ 1.000	$T_{\mu\nu} := \frac{2}{\sqrt{g}} \frac{\delta L}{\delta g_{\mu\nu}}$	$T_{\mu\nu} := \frac{2}{\sqrt{g}} \frac{\delta L}{\delta g_{\mu\nu}}$	$T_{\mu\nu} := \frac{2}{\sqrt{g}} \frac{\delta L}{\delta g_{\mu\nu}}$
mielke-geometrodynamics_114 EQ0338 (5.3.15)	114	■ 1.000	$\{ \}^{\{3\}} T_{\alpha} = \frac{(-1)^s}{3} * (\vartheta_{\alpha} \wedge \mathscr{A})$	$^{(3)}T_{\alpha} = \frac{(-1)^s}{3} * (\vartheta_{\alpha} \wedge \mathscr{A})$	$^{(3)}T_{\alpha} = \frac{(-1)^s}{3} * (\vartheta_{\alpha} \wedge \mathscr{A})$
mielke-geometrodynamics_115 EQ0342 (5.3.19)	115	■ 1.000	$\Delta \tilde{L} := \kappa (L_{\mathrm{EC}} - L_{\mathrm{HE}}) \simeq \frac{\kappa^2}{48} j_5 \wedge^* j_5 = 4f^2 j_5 \wedge^* j_5,$	$\Delta \tilde{L} := \kappa (L_{\mathrm{EC}} - L_{\mathrm{HE}}) \simeq \frac{\kappa^2}{48} j_5 \wedge^* j_5 = 4f^2 j_5 \wedge^* j_5,$	$\Delta \tilde{L} := \kappa (L_{\mathrm{EC}} - L_{\mathrm{HE}}) \simeq \frac{\kappa^2}{48} j_5 \wedge^* j_5 = 4f^2 j_5 \wedge^* j_5,$
mielke-geometrodynamics_116 EQ0344 (5.4.2)	116	■ 1.000	$L_{\mathrm{HE}} := L_{\mathrm{HE}} + \frac{\Lambda}{\kappa} \eta$	$L_{\mathrm{HE}} := L_{\mathrm{HE}} + \frac{\Lambda}{\kappa} \eta$	$L_{\mathrm{HE}} := L_{\mathrm{HE}} + \frac{\Lambda}{\kappa} \eta$
mielke-geometrodynamics_116 EQ0345 (5.4.3)	116	■ 1.000	$g_{\mathrm{N}} := \kappa k^2, \quad \lambda := \Lambda / k^2,$	$g_{\mathrm{N}} := \kappa k^2, \quad \lambda := \Lambda / k^2,$	$g_{\mathrm{N}} := \kappa k^2, \quad \lambda := \Lambda / k^2,$
mielke-geometrodynamics_116 EQ0346 (5.4.4)	116	■ 1.000	$k \frac{\partial}{\partial k} g_{\mathrm{N}} = \beta_1(g_{\mathrm{N}}, \lambda) = (2 + d_{\mathrm{N}}) g_{\mathrm{N}}, \quad k \frac{\partial}{\partial k} \lambda = \beta_2(g_{\mathrm{N}}, \lambda),$	$k \frac{\partial}{\partial k} g_{\mathrm{N}} = \beta_1(g_{\mathrm{N}}, \lambda) = (2 + d_{\mathrm{N}}) g_{\mathrm{N}}, \quad k \frac{\partial}{\partial k} \lambda = \beta_2(g_{\mathrm{N}}, \lambda),$	$k \frac{\partial}{\partial k} g_{\mathrm{N}} = \beta_1(g_{\mathrm{N}}, \lambda) = (2 + d_{\mathrm{N}}) g_{\mathrm{N}}, \quad k \frac{\partial}{\partial k} \lambda = \beta_2(g_{\mathrm{N}}, \lambda),$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_117 EQ0348 (5.4.6)		■ 1.000	$f^2=\frac{1}{192} \kappa^2=3 \times 2^{-10}\backslash\left(\frac{\mu}{\Lambda}\right)^2=2^{-8} \frac{\mu}{\rho_{\Lambda}}$	$f^2 = \frac{1}{192} \kappa^2 = 3 \times 2^{-10} \left(\frac{\mu}{\Lambda} \right)^2 = 2^{-8} \frac{\mu}{\rho_{\Lambda}}$	$f^2 = \frac{1}{192} \kappa^2 = 3 \times 2^{-10} \left(\frac{\mu}{\Lambda} \right)^2 = 2^{-8} \frac{\mu}{\rho_{\Lambda}}$
mielke-geometrodynamics_123 EQ0350 (6.2.1)		■ 1.000	$C_{\mathrm{TT}}:=\frac{1}{2} \ell^2 \backslash\left(\vartheta^{\alpha} \wedge T_{\alpha}\right)=-\frac{(-1)^{\mathrm{sig}}}{2 \ell^2} * \mathscr{A},$	$C_{\mathrm{TT}} := \frac{1}{2\ell^2} (\vartheta^{\alpha} \wedge T_{\alpha}) = -\frac{(-1)^{\mathrm{sig}}}{2\ell^2} * \mathscr{A},$	$C_{\mathrm{TT}} := \frac{1}{2\ell^2} \left(\vartheta^{\alpha} \wedge T_{\alpha} \right) = -\frac{(-1)^{\mathrm{sig}}}{2\ell^2} * \mathscr{A},$
mielke-geometrodynamics_124 EQ0351 (6.2.2)		■ 1.000	$\begin{aligned} C_{\mathrm{RR}} \& \; :=-\operatorname{Tr} \backslash\left(\Gamma \wedge \Omega-\frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma\right) \backslash\& \; =-\frac{1}{2} \backslash\left(\Gamma_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha}+\frac{1}{3} \Gamma_{\alpha}^{\beta} \wedge \Gamma_{\beta}^{\gamma} \wedge \Gamma_{\gamma}^{\alpha}\right) . \end{aligned}$	$C_{\mathrm{RR}} := -\operatorname{Tr} \left(\Gamma \wedge \Omega - \frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma \right) \\ = -\frac{1}{2} \left(\Gamma_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha} + \frac{1}{3} \Gamma_{\alpha}^{\beta} \wedge \Gamma_{\beta}^{\gamma} \wedge \Gamma_{\gamma}^{\alpha} \right) .$	$C_{\mathrm{RR}} := -\operatorname{Tr} \left(\Gamma \wedge \Omega - \frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma \right) \\ = -\frac{1}{2} \left(\Gamma_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha} + \frac{1}{3} \Gamma_{\alpha}^{\beta} \wedge \Gamma_{\beta}^{\gamma} \wedge \Gamma_{\gamma}^{\alpha} \right) .$
mielke-geometrodynamics_124 EQ0352 (6.2.3)		■ 1.000	$\hat{C}=C_{\mathrm{RR}}-2 C_{\mathrm{TT}}$	$\hat{C} = C_{\mathrm{RR}} - 2C_{\mathrm{TT}}$	$\hat{C} = C_{\mathrm{RR}} - 2C_{\mathrm{TT}}$
mielke-geometrodynamics_124 EQ0353 (6.2.4)		■ 1.000	$d C_{\mathrm{TT}} \equiv \frac{1}{2 \ell^2} \left(T^{\alpha} \wedge T_{\alpha}+R_{\alpha \beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta}\right),$	$dC_{\mathrm{TT}} \equiv \frac{1}{2\ell^2} \left(T^{\alpha} \wedge T_{\alpha} + R_{\alpha\beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta} \right),$	$dC_{\mathrm{TT}} \equiv \frac{1}{2\ell^2} \left(T^{\alpha} \wedge T_{\alpha} + R_{\alpha\beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta} \right),$
mielke-geometrodynamics_124 EQ0354 (6.2.5)		■ 1.000	$d C_{\mathrm{RR}}=-\operatorname{Tr}(\Omega \wedge \Omega)=\frac{1}{2} R^{\alpha \beta} \wedge R_{\alpha \beta}$	$dC_{\mathrm{RR}} = -\operatorname{Tr}(\Omega \wedge \Omega) = \frac{1}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}$	$dC_{\mathrm{RR}} = -\operatorname{Tr}(\Omega \wedge \Omega) = \frac{1}{2} R^{\alpha\beta} \wedge R_{\alpha\beta}$
mielke-geometrodynamics_124 EQ0355 (6.2.6)		■ 1.000	$D R_{\alpha \beta} \equiv 0,$	$DR_{\alpha\beta} \equiv 0,$	$DR_{\alpha\beta} \equiv 0,$
mielke-geometrodynamics_125 EQ0359 (6.2.10)		■ 1.000	$H_{\alpha}^{\parallel}:=-\partial L_{\parallel} / \partial T^{\alpha}=(1 / \kappa) \eta_{\alpha \beta \gamma} \wedge K^{\beta \gamma}$	$H_{\alpha}^{\parallel} := -\partial L_{\parallel} / \partial T^{\alpha} = (1/\kappa) \eta_{\alpha\beta\gamma} \wedge K^{\beta\gamma}$	$H_{\alpha}^{\parallel} := -\partial L_{\parallel} / \partial T^{\alpha} = (1/\kappa) \eta_{\alpha\beta\gamma} \wedge K^{\beta\gamma}$
mielke-geometrodynamics_126 EQ0361 (6.2.12)		■ 1.000	$\tilde{L}_{\parallel}=L_{\parallel}-R^{\alpha \beta} \wedge \lambda_{\alpha \beta} .$	$\tilde{L}_{\parallel} = L_{\parallel} - R^{\alpha\beta} \wedge \lambda_{\alpha\beta}.$	$\tilde{L}_{\parallel} = L_{\parallel} - R^{\alpha\beta} \wedge \lambda_{\alpha\beta}.$
mielke-geometrodynamics_126 EQ0362 (6.2.14)		■ 1.000	$\begin{gathered} D H_{\alpha}^{\parallel}-E_{\alpha}^{\parallel}=\Sigma_{\alpha}, \backslash D \lambda_{\alpha \beta}+\vartheta_{[\alpha} \wedge H_{\beta]}^{\parallel}=\tau_{\alpha \beta}, \end{gathered}$	$DH_{\alpha}^{\parallel} - E_{\alpha}^{\parallel} = \Sigma_{\alpha}, \\ D\lambda_{\alpha\beta} + \vartheta_{[\alpha} \wedge H_{\beta]}^{\parallel} = \tau_{\alpha\beta},$	$DH_{\alpha}^{\parallel} - E_{\alpha}^{\parallel} = \Sigma_{\alpha}, \\ D\lambda_{\alpha\beta} + \vartheta_{[\alpha} \wedge H_{\beta]}^{\parallel} = \tau_{\alpha\beta},$
mielke-geometrodynamics_126 EQ0363 (6.2.15)		■ 1.000	$R^{\alpha \beta}=0 .$	$R^{\alpha\beta} = 0.$	$R^{\alpha\beta} = 0.$
mielke-geometrodynamics_126 EQ0365 (6.2.17)		■ 1.000	$D\left(\tau_{\alpha \beta}-\vartheta_{[\alpha} \wedge H_{\beta]}^{\parallel}\right)=0$	$D \left(\tau_{\alpha\beta} - \vartheta_{[\alpha} \wedge H_{\beta]}^{\parallel} \right) = 0$	$D(\tau_{\alpha\beta} - \vartheta_{[\alpha} \wedge H_{\beta]}^{\parallel}) = 0$
mielke-geometrodynamics_127 EQ0369 (6.2.20)		■ 1.000	$\stackrel{(\pm)}{C}_{\mathrm{TT}}:=\frac{1}{2 \ell^2} \left(\vartheta^{\alpha} \wedge \stackrel{(\pm)}{T}_{\alpha}\right),$	$\stackrel{(\pm)}{C}_{\mathrm{TT}} := \frac{1}{2\ell^2} \left(\vartheta^{\alpha} \wedge \stackrel{(\pm)}{T}_{\alpha} \right),$	$\stackrel{(\pm)}{C}_{\mathrm{TT}} := \frac{1}{2\ell^2} \left(\vartheta^{\alpha} \wedge \stackrel{(\pm)}{T}_{\alpha} \right),$
mielke-geometrodynamics_127 EQ0373 (6.2.24)		■ 1.000	$\stackrel{(\pm)}{D} \Pi_{\alpha}=\Sigma_{\alpha} .$	$\stackrel{(\pm)}{D} \Pi_{\alpha} = \Sigma_{\alpha}.$	$\stackrel{(\pm)}{D} \Pi_{\alpha} = \Sigma_{\alpha}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_128 EQ0379 (6.2.30)	128	■ 1.000	$\stackrel{\scriptscriptstyle\text{L}}{\operatorname{Tr}}(\stackrel{\scriptscriptstyle\text{L}}{\mathrm{YM}})=\mathrm{mp}(1/4)$ $\operatorname{Tr}(\stackrel{\scriptscriptstyle\text{L}}{\mathrm{F}}\wedge\stackrel{\scriptscriptstyle\text{L}}{\mathrm{F}})$	$\stackrel{(\pm)}{L}_{\mathrm{YM}}=\mp(1/4)\operatorname{Tr}(\stackrel{(\pm)}{F}\wedge\stackrel{(\pm)}{F})$	$\stackrel{(\pm)}{L}_{\mathrm{YM}}=\mp(1/4)\operatorname{Tr}\left(\stackrel{(\pm)}{F}\wedge\stackrel{(\pm)}{F}\right)$
mielke-geometrodynamics_129 EQ0383 (6.3.1)	129	■ 1.000	$\check{E}_{\alpha}:=E_{\alpha}+\Gamma_{\alpha}^{\beta}\wedge H_{\beta}+\Sigma_{\alpha}\cong dH_{\alpha},$ $\wedge H_{\beta}+\Sigma_{\alpha}\cong dH_{\alpha},$	$\check{E}_{\alpha}:=E_{\alpha}+\Gamma_{\alpha}^{\beta}\wedge H_{\beta}+\Sigma_{\alpha}\cong dH_{\alpha},$	$\check{E}_{\alpha}:=E_{\alpha}+\Gamma_{\alpha}^{\beta}\wedge H_{\beta}+\Sigma_{\alpha}\cong dH_{\alpha},$
mielke-geometrodynamics_130 EQ0386 (6.4.1)	130	■ 1.000	$n:=\partial_t-N^A\partial/\partial x^A,\quad A=1,2,3,$	$n:=\partial_t-N^A\partial/\partial x^A,\quad A=1,2,3,$	$n:=\partial_t-N^A\partial/\partial x^A,\quad A=1,2,3,$
mielke-geometrodynamics_132 EQ0395 (6.4.10)	132	■ 1.000	$\Sigma_{\alpha}:=\frac{\partial L}{\partial \vartheta^{\alpha}},\quad \tau_{\alpha\beta}:=\frac{\partial L}{\partial \Gamma^{\alpha\beta}}=\rho(I_{\alpha\beta})\Psi\wedge\frac{\partial L}{\partial D\Psi}$	$\Sigma_{\alpha}:=\frac{\partial L}{\partial \vartheta^{\alpha}},\quad \tau_{\alpha\beta}:=\frac{\partial L}{\partial \Gamma^{\alpha\beta}}=\rho(I_{\alpha\beta})\Psi\wedge\frac{\partial L}{\partial D\Psi}$	$\Sigma_{\alpha}:=\frac{\partial L}{\partial \vartheta^{\alpha}},\quad \tau_{\alpha\beta}:=\frac{\partial L}{\partial \Gamma^{\alpha\beta}}=\rho(I_{\alpha\beta})\Psi\wedge\frac{\partial L}{\partial D\Psi}$
mielke-geometrodynamics_136 EQ0408 (6.4.28)	136	■ 1.000	$d s^2=h^2 d r^2+f^2\left[d \psi^2+\sin ^2 \psi\left(d \theta^2+\sin ^2 \theta d \phi^2\right)\right],$ $\psi\left(d \theta^2+\sin ^2 \theta d \phi^2\right)\right],$	$d s^2=h^2 d r^2+f^2\left[d \psi^2+\sin ^2 \psi\left(d \theta^2+\sin ^2 \theta d \phi^2\right)\right],$	$d s^2=h^2 d r^2+f^2\left[d \psi^2+\sin ^2 \psi\left(d \theta^2+\sin ^2 \theta d \phi^2\right)\right],$
mielke-geometrodynamics_136 EQ0410 (6.4.30)	136	■ 1.000	$\ell^2\operatorname{\underline{C}}_{\mathrm{TT}}=\frac{1}{2}\vartheta^{\alpha}\wedge d\underline{\vartheta}_{\alpha}=\frac{1}{2}\underline{\vartheta}^A\wedge d\underline{\vartheta}_A=3\underline{\vartheta}^{\hat{1}}\wedge\underline{\vartheta}^{\hat{2}}\wedge\underline{\vartheta}^{\hat{3}}$	$\ell^2\operatorname{\underline{C}}_{\mathrm{TT}}=\frac{1}{2}\vartheta^{\alpha}\wedge d\underline{\vartheta}_{\alpha}=\frac{1}{2}\underline{\vartheta}^A\wedge d\underline{\vartheta}_A=3\underline{\vartheta}^{\hat{1}}\wedge\underline{\vartheta}^{\hat{2}}\wedge\underline{\vartheta}^{\hat{3}}$	$\ell^2\operatorname{\underline{C}}_{\mathrm{TT}}=\frac{1}{2}\underline{\vartheta}^{\alpha}\wedge d\underline{\vartheta}_{\alpha}=\frac{1}{2}\underline{\vartheta}^A\wedge d\underline{\vartheta}_A=3\underline{\vartheta}^{\hat{1}}\wedge\underline{\vartheta}^{\hat{2}}\wedge\underline{\vartheta}^{\hat{3}}$
mielke-geometrodynamics_136 EQ0412 (6.4.32)	136	■ 1.000	$f=\frac{a}{r^2+c^2}r^2$	$f=\frac{ar^2}{r^2+c^2}$	$f=\frac{ar^2}{r^2+c^2}$
mielke-geometrodynamics_137 EQ0414 (6.5.2)	137	■ 1.000	$\Psi(\gamma)=\int_{\mathscr{C}/\mathscr{G}}\mathscr{D}A\,W(A,\gamma)\Psi(A).$ $A\,W(A,\gamma)\,\Psi(A).$	$\Psi(\gamma)=\int_{\mathscr{C}/\mathscr{G}}\mathscr{D}A\,W(A,\gamma)\Psi(A).$	$\Psi(\gamma)=\int_{\mathscr{C}/\mathscr{G}}\mathscr{D}A\,W(A,\gamma)\Psi(A).$
mielke-geometrodynamics_137 EQ0416 (6.5.4)	137	■ 1.000	$\langle 0 W(\gamma) 0\rangle e^{\Lambda G(\gamma)}V(\gamma),$	$\langle 0 W(\gamma) 0\rangle e^{\Lambda G(\gamma)}V(\gamma),$	$\langle 0 W(\gamma) 0\rangle e^{\Lambda G(\gamma)}V(\gamma),$
mielke-geometrodynamics_138 EQ0417 (6.6.1)	138	■ 1.000	$\int T=\int d\sigma=2\pi N$	$\int T=\int d\sigma=2\pi N$	$\int T=\int d\sigma=2\pi N,$
mielke-geometrodynamics_139 EQ0418 (6.6.2)	139	■ 1.000	$W^N:=S^3\#^N\left(S^1\times S^2\right).$	$W^N:=S^3\#^N\left(S^1\times S^2\right).$	$W^N:=S^3\#^N(S^1\times S^2).$
mielke-geometrodynamics_140 EQ0419 (6.7.1)	140	■ 1.000	$L:=L_{\mathrm{EC}}+L_{\theta}+L_{\mathrm{D}},$	$L:=L_{\mathrm{EC}}+L_{\theta}+L_{\mathrm{D}},$	$L:=L_{\mathrm{EC}}+L_{\theta}+L_{\mathrm{D}},$
mielke-geometrodynamics_140 EQ0420 (6.7.2)	140	■ 1.000	$L_{\theta}=\theta_{\mathrm{T}}dC_{\mathrm{TT}}+\theta_{\mathrm{L}}dC_{\mathrm{RR}}$	$L_{\theta}=\theta_{\mathrm{T}}dC_{\mathrm{TT}}+\theta_{\mathrm{L}}dC_{\mathrm{RR}}$	$L_{\theta}=\theta_{\mathrm{T}}dC_{\mathrm{TT}}+\theta_{\mathrm{L}}dC_{\mathrm{RR}}$
mielke-geometrodynamics_140 EQ0421 (6.7.3)	140	■ 1.000	$H_{\alpha}:= -\frac{\partial L}{\partial T^{\alpha}}= -\frac{\theta_{\mathrm{T}}}{\ell^2}T_{\alpha},$	$H_{\alpha}:= -\frac{\partial L}{\partial T^{\alpha}}= -\frac{\theta_{\mathrm{T}}}{\ell^2}T_{\alpha},$	$H_{\alpha}:= -\frac{\partial L}{\partial T^{\alpha}}= -\frac{\theta_{\mathrm{T}}}{\ell^2}T_{\alpha},$
mielke-geometrodynamics_140 EQ0422 (6.7.4)	140	■ 1.000	$H_{\alpha\beta}:= -\frac{\partial L}{\partial R^{\alpha\beta}}= \frac{1}{2\kappa}\eta_{\alpha\beta}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\theta_{\mathrm{L}}R_{\alpha\beta}$	$H_{\alpha\beta}:= -\frac{\partial L}{\partial R^{\alpha\beta}}= \frac{1}{2\kappa}\eta_{\alpha\beta}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\theta_{\mathrm{L}}R_{\alpha\beta}$	$H_{\alpha\beta}:= -\frac{\partial L}{\partial R^{\alpha\beta}}= \frac{1}{2\kappa}\eta_{\alpha\beta}-\frac{\theta_{\mathrm{T}}}{2\ell^2}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\theta_{\mathrm{L}}R_{\alpha\beta}$
mielke-geometrodynamics_140 EQ0423 (6.7.5)	140	■ 1.000	$D\theta_{\mathrm{T}}\wedge T_{\alpha}+\theta_{\mathrm{T}}DT_{\alpha}+\ell^2E_{\alpha}=-\ell^2\Sigma_{\alpha}.$	$D\theta_{\mathrm{T}}\wedge T_{\alpha}+\theta_{\mathrm{T}}DT_{\alpha}+\ell^2E_{\alpha}=-\ell^2\Sigma_{\alpha}.$	$D\theta_{\mathrm{T}}\wedge T_{\alpha}+\theta_{\mathrm{T}}DT_{\alpha}+\ell^2E_{\alpha}=-\ell^2\Sigma_{\alpha}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_140 EQ0425 (6.7.7)	140	■ 1.000	$C_{\mathrm{TT}} \cong \frac{\kappa}{4\ell^2} j_5$	$C_{\mathrm{TT}} \cong \frac{\kappa}{4\ell^2} j_5$	$C_{\mathrm{TT}} \cong \frac{\kappa}{4\ell^2} j_5$
mielke-geometrodynamics_141 EQ0427 (6.7.9)	141	■ 1.000	$T^\gamma \wedge \eta_{\gamma\beta} + \kappa D\theta_{\mathrm{L}} \wedge DT_\beta = 0.$	$T^\gamma \wedge \eta_{\gamma\beta} + \kappa D\theta_{\mathrm{L}} \wedge DT_\beta = 0.$	$T^\gamma \wedge \eta_{\gamma\beta} + \kappa D\theta_{\mathrm{L}} \wedge DT_\beta = 0.$
mielke-geometrodynamics_141 EQ0429 (6.7.11)	141	■ 1.000	$\kappa d\theta_{\mathrm{L}} \wedge T_\beta = -\eta_\beta$	$\kappa d\theta_{\mathrm{L}} \wedge T_\beta = -\eta_\beta$	$\kappa d\theta_{\mathrm{L}} \wedge T_\beta = -\eta_\beta$
mielke-geometrodynamics_141 EQ0430 (6.7.12)	141	■ 1.000	$\kappa \ell^2 d\theta_{\mathrm{L}} \wedge C_{\mathrm{TT}} = 2\eta.$	$\kappa \ell^2 d\theta_{\mathrm{L}} \wedge C_{\mathrm{TT}} = 2\eta.$	$\kappa \ell^2 d\theta_{\mathrm{L}} \wedge C_{\mathrm{TT}} = 2\eta.$
mielke-geometrodynamics_141 EQ0431 (6.7.13)	141	■ 1.000	$D T^\gamma \wedge \eta_{\alpha\beta\gamma} + \kappa D\theta_{\mathrm{T}} \wedge T_{[\alpha} \wedge \vartheta_{\beta]}/\ell^2 = 2\kappa (T_{[\alpha} \wedge \mu_{\beta]} - \vartheta_{[\alpha} \wedge D\mu_{\beta]}).$	$DT^\gamma \wedge \eta_{\alpha\beta\gamma} + \kappa D\theta_{\mathrm{T}} \wedge T_{[\alpha} \wedge \vartheta_{\beta]}/\ell^2 = 2\kappa (T_{[\alpha} \wedge \mu_{\beta]} - \vartheta_{[\alpha} \wedge D\mu_{\beta]}).$	$DT^\gamma \wedge \eta_{\alpha\beta\gamma} + \kappa D\theta_{\mathrm{T}} \wedge T_{[\alpha} \wedge \vartheta_{\beta]}/\ell^2 = 2\kappa (T_{[\alpha} \wedge \mu_{\beta]} - \vartheta_{[\alpha} \wedge D\mu_{\beta]}).$
mielke-geometrodynamics_142 EQ0432 (6.8.1)	142	■ 1.000	$W = \int \left[L(\vartheta^\alpha, T^\alpha, R^{\alpha\beta}, \Psi, D\Psi) + L_g(\vartheta^\alpha, T^\alpha, R^{\alpha\beta}) \right].$	$W = \int \left[L(\vartheta^\alpha, T^\alpha, R^{\alpha\beta}, \Psi, D\Psi) + L_g(\vartheta^\alpha, T^\alpha, R^{\alpha\beta}) \right].$	$W = \int \left[L(\vartheta^\alpha, T^\alpha, R^{\alpha\beta}, \Psi, D\Psi) + L_g(\vartheta^\alpha, T^\alpha, R^{\alpha\beta}) \right].$
mielke-geometrodynamics_142 EQ0433 (6.8.3)	142	■ 1.000	$\begin{gathered} \frac{\delta L}{\delta \Psi} = \frac{\partial L}{\partial \Psi} - (-1)^p D \frac{\partial L}{\partial D\Psi} = 0, \quad (\text{MATTER}) \\ DH_\alpha - E_\alpha = \Sigma_\alpha, \quad (\text{FIRST}) \\ DH_{\alpha\beta} - E_{\alpha\beta} = \tau_{\alpha\beta}. \quad (\text{SECOND}) \end{gathered}$	$\frac{\delta L}{\delta \Psi} = \frac{\partial L}{\partial \Psi} - (-1)^p D \frac{\partial L}{\partial D\Psi} = 0, \quad (\text{MATTER})$ $DH_\alpha - E_\alpha = \Sigma_\alpha, \quad (\text{FIRST})$ $DH_{\alpha\beta} - E_{\alpha\beta} = \tau_{\alpha\beta}. \quad (\text{SECOND})$	$\frac{\delta L}{\delta \Psi} = \frac{\partial L}{\partial \Psi} - (-1)^p D \frac{\partial L}{\partial D\Psi} = 0, \quad (\text{MATTER})$ $DH_\alpha - E_\alpha = \Sigma_\alpha, \quad (\text{FIRST})$ $DH_{\alpha\beta} - E_{\alpha\beta} = \tau_{\alpha\beta}. \quad (\text{SECOND})$
mielke-geometrodynamics_142 EQ0435 (6.8.6)	142	■ 1.000	$\Sigma_\alpha := \frac{\delta L}{\delta \vartheta^\alpha} = \frac{\partial L}{\partial \vartheta^\alpha} + D \frac{\partial L}{\partial T^\alpha}$	$\Sigma_\alpha := \frac{\delta L}{\delta \vartheta^\alpha} = \frac{\partial L}{\partial \vartheta^\alpha} + D \frac{\partial L}{\partial T^\alpha}$	$\Sigma_\alpha := \frac{\delta L}{\delta \vartheta^\alpha} = \frac{\partial L}{\partial \vartheta^\alpha} + D \frac{\partial L}{\partial T^\alpha}$
mielke-geometrodynamics_142 EQ0436 (6.8.7)	142	■ 1.000	$\tau_{\alpha\beta} := \frac{\delta L}{\delta \Gamma^{\alpha\beta}} = \rho(I_{\alpha\beta}) \Psi \wedge \frac{\partial L}{\partial (D\Psi)} + \vartheta_{[\alpha} \wedge \frac{\partial L}{\partial T^{\beta]}} + D \frac{\partial L}{\partial R^{\alpha\beta}},$	$\tau_{\alpha\beta} := \frac{\delta L}{\delta \Gamma^{\alpha\beta}} = \rho(I_{\alpha\beta}) \Psi \wedge \frac{\partial L}{\partial (D\Psi)} + \vartheta_{[\alpha} \wedge \frac{\partial L}{\partial T^{\beta]}} + D \frac{\partial L}{\partial R^{\alpha\beta}},$	$\tau_{\alpha\beta} := \frac{\delta L}{\delta \Gamma^{\alpha\beta}} = \rho(I_{\alpha\beta}) \Psi \wedge \frac{\partial L}{\partial (D\Psi)} + \vartheta_{[\alpha} \wedge \frac{\partial L}{\partial T^{\beta]}} + D \frac{\partial L}{\partial R^{\alpha\beta}},$
mielke-geometrodynamics_142 EQ0438 (6.8.9)	142	■ 1.000	$E_{\alpha\beta} := -\vartheta_{[\alpha} \wedge H_{\beta]}$	$E_{\alpha\beta} := -\vartheta_{[\alpha} \wedge H_{\beta]}$	$E_{\alpha\beta} := -\vartheta_{[\alpha} \wedge H_{\beta]}$
mielke-geometrodynamics_143 EQ0439 (6.8.10)	143	■ 1.000	$\vartheta^\alpha \wedge E_\alpha = 4L_g + 2T^\alpha \wedge H_\alpha + 2R^{\alpha\beta} \wedge H_{\alpha\beta}.$	$\vartheta^\alpha \wedge E_\alpha = 4L_g + 2T^\alpha \wedge H_\alpha + 2R^{\alpha\beta} \wedge H_{\alpha\beta}.$	$\vartheta^\alpha \wedge E_\alpha = 4L_g + 2T^\alpha \wedge H_\alpha + 2R^{\alpha\beta} \wedge H_{\alpha\beta}.$
mielke-geometrodynamics_143 EQ0442 (6.8.13)	143	■ 1.000	$L_{\mathrm{qPG}} = -\frac{1}{2} T^\alpha \wedge H_\alpha - \frac{1}{2} R^{\alpha\beta} \wedge \left(H_{\alpha\beta} - \frac{a_0}{2\ell^2} \eta_{\alpha\beta} \right),$	$L_{\mathrm{qPG}} = -\frac{1}{2} T^\alpha \wedge H_\alpha - \frac{1}{2} R^{\alpha\beta} \wedge \left(H_{\alpha\beta} - \frac{a_0}{2\ell^2} \eta_{\alpha\beta} \right),$	$L_{\mathrm{qPG}} = -\frac{1}{2} T^\alpha \wedge H_\alpha - \frac{1}{2} R^{\alpha\beta} \wedge \left(H_{\alpha\beta} - \frac{a_0}{2\ell^2} \eta_{\alpha\beta} \right),$
mielke-geometrodynamics_143 EQ0444 (6.8.15)	143	■ 1.000	$H_{\alpha\beta} = -\frac{a_0}{2\ell^2} \eta_{\alpha\beta} - \frac{1}{\kappa} * \left(\sum_{j=1}^6 b_j^{(j)} R_{\alpha\beta} \right).$	$H_{\alpha\beta} = -\frac{a_0}{2\ell^2} \eta_{\alpha\beta} - \frac{1}{\kappa} * \left(\sum_{j=1}^6 b_j^{(j)} R_{\alpha\beta} \right).$	$H_{\alpha\beta} = -\frac{a_0}{2\ell^2} \eta_{\alpha\beta} - \frac{1}{\kappa} * \left(\sum_{j=1}^6 b_j^{(j)} R_{\alpha\beta} \right).$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0447 (6.9.3)	144	■ 1.000	$\eta=\{ \}^{\{*\}} 1=\frac{1}{4!}\eta_{\alpha\beta\gamma\delta}\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta},$	$\eta = {}^*1 = \frac{1}{4!}\eta_{\alpha\beta\gamma\delta}\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta},$	$\eta = {}^*1 = \frac{1}{4!}\eta_{\alpha\beta\gamma\delta}\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta},$
mielke-geometrodynamics_EQ0448 (6.9.4)	145	■ 1.000	$p=\frac{1}{8}\pi^{\{2\}}\int R_{\{\alpha\beta\}}^{\{\}}\wedge R_{\{\beta\alpha\}}^{\{\}}.$	$p = \frac{1}{8\pi^2}\int R_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}.$	$p = \frac{1}{8\pi^2}\int R_{\alpha}{}^{\beta}\wedge R_{\beta}{}^{\alpha}.$
mielke-geometrodynamics_EQ0453 (7.2.1)	151	■ 1.000	$[\ L_{\{a\ b\}}^{\{(\star)\}}:=\frac{1}{2}\varepsilon_{abcd}L^{\{c\ d\}}_{\{a\ b\}}\ L^{\{c\ d\}}=L_{\{a\ b\}}\ .\]$	$L_{ab}^{(\star)}:=\frac{1}{2}\varepsilon_{abcd}L^{cd}=L_{ab}.$	$L_{ab}^{(\star)}:=\frac{1}{2}\varepsilon_{abcd}L^{cd}=L_{ab}.$
mielke-geometrodynamics_EQ0454 (7.2.6)	152	■ 1.000	$\Omega^g=\Omega_{\mathrm{C}}+\bar{\Omega}^g-\frac{1}{6}\vartheta\wedge\vartheta R.$	$\Omega^g = \Omega_C + \bar{\Omega}^g - \frac{1}{6}\vartheta\wedge\vartheta R.$	$\Omega^g = \Omega_C + \bar{\Omega}^g - \frac{1}{6}\vartheta\wedge\vartheta R.$
mielke-geometrodynamics_EQ0455 (7.2.7)	152	■ 1.000	$\bar{\Omega}^g=\vartheta\wedge^*(\vartheta\wedge\Omega^g)-\frac{1}{4}\vartheta\wedge\vartheta R,$	$\bar{\Omega}^g = \vartheta\wedge^*(\vartheta\wedge\Omega^g) - \frac{1}{4}\vartheta\wedge\vartheta R,$	$\bar{\Omega}^g = \vartheta\wedge^*(\vartheta\wedge\Omega^g) - \frac{1}{4}\vartheta\wedge\vartheta R,$
mielke-geometrodynamics_EQ0457 (7.2.9)	152	■ 1.000	$\operatorname{Tr}\left(\vartheta\wedge\vartheta\wedge\Omega_{\mathrm{C}}\right)=\operatorname{Tr}\left(\vartheta\wedge\vartheta\wedge\bar{\Omega}^g\right)=0$	$\operatorname{Tr}\left(\vartheta\wedge\vartheta\wedge\Omega_{\mathrm{C}}\right)=\operatorname{Tr}\left(\vartheta\wedge\vartheta\wedge\bar{\Omega}^g\right)=0$	$\operatorname{Tr}(\vartheta\wedge\vartheta\wedge\Omega_{\mathrm{C}})=\operatorname{Tr}(\vartheta\wedge\vartheta\wedge\bar{\Omega}^g)=0$
mielke-geometrodynamics_EQ0463 (7.3.2)	153	■ 1.000	$D^{\{*\}}\Omega^g=0,$	$D^*\Omega^g=0,$	$D^*\Omega^g=0,$
mielke-geometrodynamics_EQ0464 (7.3.3)	153	■ 1.000	$D^{\{*\}}R_{\{\alpha\beta\}}=0,$	$D^*R_{\alpha\beta}=0,$	$D^*R_{\alpha\beta}=0,$
mielke-geometrodynamics_EQ0468 (7.3.7)	154	■ 1.000	$R_{\{\alpha\beta\}}=-\{ \}^{\{*\}}R_{\{\alpha\beta\}}^{\{(\star)\}},$	$R_{\alpha\beta}=-{}^*R_{\alpha\beta}^{(\star)},$	$R_{\alpha\beta}=-{}^*R_{\alpha\beta}^{(\star)},$
mielke-geometrodynamics_EQ0469 (7.3.8)	154	■ 1.000	$R_{\{\alpha\beta\}}=\{ \}^{\{*\}}R_{\{\alpha\beta\}}^{\{(\star)\}},$	$R_{\alpha\beta}={}^*R_{\alpha\beta}^{(\star)},$	$R_{\alpha\beta}={}^*R_{\alpha\beta}^{(\star)},$
mielke-geometrodynamics_EQ0479 (7.4.7)	157	■ 1.000	$G_{\alpha}-\Lambda\eta_{\alpha}=0$	$G_{\alpha}-\Lambda\eta_{\alpha}=0$	$G_{\alpha}-\Lambda\eta_{\alpha}=0$
mielke-geometrodynamics_EQ0480 (7.5.1)	157	■ 1.000	$\square^c:=\square-\frac{1}{6}R,\quad\square:=(1/\sqrt{g})\partial_i(\sqrt{g}g^{ij}\partial_j).$	$\square^c:=\square-\frac{1}{6}R,\quad\square:=(1/\sqrt{g})\partial_i(\sqrt{g}g^{ij}\partial_j).$	$\square^c:=\square-\frac{1}{6}R,\quad\square:=(1/\sqrt{g})\partial_i(\sqrt{g}g^{ij}\partial_j).$
mielke-geometrodynamics_EQ0482 (7.5.3)	157	■ 1.000	$\left(\square-\frac{2}{3}\Lambda\right)h_{ij}=0$	$\left(\square-\frac{2}{3}\Lambda\right)h_{ij}=0$	$\left(\square-\frac{2}{3}\Lambda\right)h_{ij}=0$
mielke-geometrodynamics_EQ0483 (7.5.4)	158	■ 1.000	$\partial^ih_{ij}-\partial_jh=0$	$\partial^ih_{ij}-\partial_jh=0$	$\partial^ih_{ij}-\partial_jh=0$
mielke-geometrodynamics_EQ0484 (7.5.5)	158	■ 1.000	$\Delta m=\sqrt{2\Lambda/3}>0$	$\Delta m = \sqrt{2\Lambda/3} > 0$	$\Delta m = \sqrt{2\Lambda/3} > 0$
mielke-geometrodynamics_EQ0488 (7.7.5)	160	■ 1.000	$L_{\{\mathrm{EC}\}}=-\frac{1}{2}\kappa^{\{2\}}R^{\{\alpha\beta\}}\wedge\eta_{\alpha\beta}$	$L_{\mathrm{EC}}=-\frac{1}{2\kappa^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}$	$L_{\mathrm{EC}}=-\frac{1}{2\kappa^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}$
mielke-geometrodynamics_EQ0489 (7.7.6)	160	■ 1.000	$\widetilde{L}=-\sum_{k=0}^K(1/2k)R^{\alpha\beta}\wedge\stackrel{(2k)}{H}_{\alpha\beta}+L_{\theta},$	$\tilde{L}=-\sum_{k=0}^K(1/2k)R^{\alpha\beta}\wedge\stackrel{(2k)}{H}_{\alpha\beta}+L_{\theta},$	$\tilde{L}=-\sum_{k=0}^K(1/2k)R^{\alpha\beta}\wedge\stackrel{(2k)}{H}_{\alpha\beta}+L_{\theta},$
mielke-geometrodynamics_EQ0491 (7.7.9)	161	■ 1.000	$\widetilde{L}\rightarrow L=-\frac{1}{2}\left(R^{\alpha\beta}\wedge\widetilde{H}_{\alpha\beta}-\widetilde{L}\right)$	$\tilde{L}\rightarrow L=-\frac{1}{2}\left(R^{\alpha\beta}\wedge\tilde{H}_{\alpha\beta}-\tilde{L}\right)$	$\tilde{L}\rightarrow L=-\frac{1}{2}\left(R^{\alpha\beta}\wedge\tilde{H}_{\alpha\beta}-\tilde{L}\right)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_161 EQ0493 (7.7.11)		■ 1.000	$\widetilde{H}_{\alpha\beta\mu\nu}:=\frac{\partial^2\widetilde{L}}{\partial R^{\mu\nu}\partial R^{\alpha\beta}}=-\frac{\partial\widetilde{H}_{\mu\nu}}{\partial R^{\alpha\beta}}$		
mielke-geometrodynamics_161 EQ0494 (7.7.12)		■ 1.000	$\widetilde{\eta}_{\alpha\beta}:=\frac{\theta_{\mathrm{T}}^*}{2}\eta_{\alpha\beta}-\frac{\theta_{\mathrm{T}}}{2}\vartheta_{\alpha}\wedge\vartheta_{\beta}-\ell^2\widetilde{H}_{\alpha\beta}=0$		
mielke-geometrodynamics_162 EQ0496 (7.7.14)		■ 1.000	$\eta:=\eta_{\alpha\beta\gamma\delta}\vartheta^{\alpha}\wedge\vartheta^{\beta}\wedge\vartheta^{\gamma}\wedge\vartheta^{\delta}/4!$		
mielke-geometrodynamics_163 EQ0499 (7.7.17)		■ 1.000	$\gamma_{\alpha}\gamma_{\beta}+\gamma_{\beta}\gamma_{\alpha}=2o_{\alpha\beta}\mathbf{1}_4.$		
mielke-geometrodynamics_163 EQ0501 (7.7.19)		■ 1.000	$\gamma:=\gamma^{\alpha}\vartheta_{\alpha}$		
mielke-geometrodynamics_163 EQ0502 (7.7.20)		■ 1.000	${}^*\gamma=\gamma^{\alpha}\eta_{\alpha}=\frac{i}{6}\gamma_5\gamma\wedge\gamma\wedge\gamma,$		
mielke-geometrodynamics_164 EQ0503 (7.7.21)		■ 1.000	$\sigma:=\frac{1}{2}\sigma_{\alpha\beta}\vartheta^{\alpha}\wedge\vartheta^{\beta}=\frac{i}{2}\gamma\wedge\gamma,\quad{}^*\sigma=\frac{1}{2}\sigma_{\alpha\beta}\eta^{\alpha\beta}=: \sigma^{(*)}=i\gamma_5\sigma.$		
mielke-geometrodynamics_164 EQ0504 (7.7.22)		■ 1.000	$\sigma^{(*)}:=i\gamma_5\sigma,\quad\sigma^{(**)}=i^2\gamma_5^2\sigma=-\sigma.$		
mielke-geometrodynamics_164 EQ0505 (7.7.23)		■ 1.000	$[\gamma,\sigma^{(*)}]=\gamma\wedge i\gamma_5\sigma-i\gamma_5\sigma\wedge\gamma=-i\gamma_5(\gamma\wedge\sigma+\sigma\wedge\gamma)=2\gamma_5\gamma\wedge\gamma\wedge\gamma$		
mielke-geometrodynamics_165 EQ0508 (7.7.26)		■ 1.000	$\gamma\rightarrow\gamma^{\theta}=e^{i\gamma^5\theta}\gamma e^{-i\gamma^5\theta}$		
mielke-geometrodynamics_165 EQ0510 (7.7.28)		■ 1.000	$\gamma:=\vartheta^{\alpha}\gamma_{\alpha},\quad\Gamma:=\frac{i}{4}\Gamma^{\alpha\beta}\sigma_{\alpha\beta}=\Gamma^{\{\}}-K,$		
mielke-geometrodynamics_166 EQ0513 (7.7.31)		■ 1.000	$\Omega^{(*)}=\frac{1}{4}R_{\alpha\beta}^{(*)}\sigma^{\alpha\beta},\quad R_{\alpha\beta}^{(*)}:=\frac{1}{2}\eta_{\alpha\beta\gamma\delta}R^{\gamma\delta}.$		
mielke-geometrodynamics_166 EQ0514 (7.7.32)		■ 1.000	$D\sigma=\frac{i}{2}[\Theta,\gamma]$		
mielke-geometrodynamics_166 EQ0517 (7.7.35)		■ 1.000	$d^*\gamma=0,\quad d^*\Gamma=0$		
mielke-geometrodynamics_166 EQ0518 (7.7.36)		■ 1.000	$DD\Phi=[\Omega^g,\Phi],$		

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E08528 (7.7.38)	166	■ 1.000	$G:=G_{\alpha}\gamma^{\alpha}\wedge\eta_{\mu\nu\lambda}\gamma^{\lambda}=-i\gamma_5(\Omega^g\wedge\gamma+\gamma\wedge\Omega^g)=i[\gamma,\gamma_5\Omega^g].$	$G:=G_{\alpha}\gamma^{\alpha}:=\frac{1}{2}R^{\mu\nu}\wedge\eta_{\mu\nu\lambda}\gamma^{\lambda}=-i\gamma_5(\Omega^g\wedge\gamma+\gamma\wedge\Omega^g)=i[\gamma,\gamma_5\Omega^g].$	$G:=G_{\alpha}\gamma^{\alpha}:=\frac{1}{2}R^{\mu\nu}\wedge\eta_{\mu\nu\lambda}\gamma^{\lambda}=-i\gamma_5(\Omega^g\wedge\gamma+\gamma\wedge\Omega^g)=i[\gamma,\gamma_5\Omega^g].$
mielke-geometrodynamics_E08521 (7.7.39)	167	■ 1.000	$D[\gamma,\Theta]\equiv 2i[\sigma,\Omega^g],\quad D[\gamma,\Omega^g]\equiv [\Theta,\Omega^g],\quad DG\equiv i[\Theta,\gamma_5\Omega^g],$	$D[\gamma,\Theta]\equiv 2i[\sigma,\Omega^g],\quad D[\gamma,\Omega^g]\equiv [\Theta,\Omega^g],\quad DG\equiv i[\Theta,\gamma_5\Omega^g],$	$D[\gamma,\Theta]\equiv 2i[\sigma,\Omega^g],\quad D[\gamma,\Omega^g]\equiv [\Theta,\Omega^g],\quad DG\equiv i[\Theta,\gamma_5\Omega^g],$
mielke-geometrodynamics_E08523 (7.7.42)	167	■ 1.000	$D R_{\alpha\beta}^{\star}\equiv 0,$	$DR_{\alpha\beta}^{(\star)}\equiv 0,$	$DR_{\alpha\beta}^{(\star)}\equiv 0,$
mielke-geometrodynamics_E08532 (7.7.52)	169	■ 1.000	$\{ \}^{\{6\}}R^{\alpha\beta}=-\frac{1}{12}R\vartheta^{\alpha}\wedge\vartheta^{\beta}$	$^{(6)}R^{\alpha\beta}=-\frac{1}{12}R\vartheta^{\alpha}\wedge\vartheta^{\beta}$	$^{(6)}R^{\alpha\beta}=-\frac{1}{12}R\vartheta^{\alpha}\wedge\vartheta^{\beta}$
mielke-geometrodynamics_E08541 (8.3.2)	177	■ 1.000	$s\gamma=\psi-[C,\gamma]$	$s\gamma=\psi-[C,\gamma]$	$s\gamma=\psi-[C,\gamma]$
mielke-geometrodynamics_E08542 (8.3.3)	177	■ 1.000	$s\sigma:=\frac{i}{2}s(\gamma\wedge\gamma)=\frac{i}{2}([\psi,\gamma]-[[C,\gamma],\gamma])=\frac{i}{2}[\psi,\gamma]-[C,\sigma].$	$s\sigma:=\frac{i}{2}s(\gamma\wedge\gamma)=\frac{i}{2}([\psi,\gamma]-[[C,\gamma],\gamma])=\frac{i}{2}[\psi,\gamma]-[C,\sigma].$	$s\sigma:=\frac{i}{2}s(\gamma\wedge\gamma)=\frac{i}{2}([\psi,\gamma]-[[C,\gamma],\gamma])=\frac{i}{2}[\psi,\gamma]-[C,\sigma].$
mielke-geometrodynamics_E08543 (8.3.4)	177	■ 1.000	$s\bar{c}=b,\quad sb=0.$	$s\bar{c}=b,\quad sb=0.$	$s\bar{c}=b,\quad sb=0.$
mielke-geometrodynamics_E08546 (8.3.7)	178	■ 1.000	$ds^2=\frac{1}{4}\mathrm{Tr}\{\gamma\otimes\gamma\}^*o_{\alpha\beta}D\xi^{\alpha}\otimes D\xi^{\beta}$	$ds^2=\frac{1}{4}\mathrm{Tr}\{\gamma\otimes\gamma\}^*o_{\alpha\beta}D\xi^{\alpha}\otimes D\xi^{\beta}$	$ds^2=\frac{1}{4}\mathrm{Tr}\{\gamma\otimes\gamma\}^*o_{\alpha\beta}D\xi^{\alpha}\otimes D\xi^{\beta}$
mielke-geometrodynamics_E08558 (8.4.4)	179	■ 1.000	$L_{\mathrm{TG}}=dC_{\mathrm{RR}}+L_{\mathrm{FP}}=dC_{\mathrm{RR}}+s\{\cdots\}$	$L_{\mathrm{TG}}=dC_{\mathrm{RR}}+L_{\mathrm{FP}}=dC_{\mathrm{RR}}+s\{\cdots\}$	$L_{\mathrm{TG}}=dC_{\mathrm{RR}}+L_{\mathrm{FP}}=dC_{\mathrm{RR}}+s\{\cdots\}$
mielke-geometrodynamics_E08559 (8.5.2)	181	■ 1.000	$\sigma:=\frac{i}{2}\gamma\wedge\gamma.$	$\sigma:=\frac{i}{2}\gamma\wedge\gamma.$	$\sigma:=\frac{i}{2}\gamma\wedge\gamma.$
mielke-geometrodynamics_E08563 (8.5.6)	182	■ 1.000	$\Omega^{g(\theta)}:=\sqrt{2}\left(\Omega^g\sin\theta+^*\Omega^{g(\star)}\cos\theta\right)$	$\Omega^{g(\theta)}:=\sqrt{2}\left(\Omega^g\sin\theta+^*\Omega^{g(\star)}\cos\theta\right)$	$\Omega^{g(\theta)}:=\sqrt{2}\left(\Omega^g\sin\theta+^*\Omega^{g(\star)}\cos\theta\right)$
mielke-geometrodynamics_E08564 (8.6.1)	183	■ 1.000	$L_{\mathrm{qPG}}=\frac{\Lambda}{\ell^2}\eta-\frac{a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2}T^{\alpha}\wedge H_{\alpha}-\frac{1}{2}R^{\alpha\beta}\wedge H_{\alpha\beta}.$	$L_{\mathrm{qPG}}=\frac{\Lambda}{\ell^2}\eta-\frac{a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2}T^{\alpha}\wedge H_{\alpha}-\frac{1}{2}R^{\alpha\beta}\wedge H_{\alpha\beta}.$	$L_{\mathrm{qPG}}=\frac{\Lambda}{\ell^2}\eta-\frac{a_0}{4\ell^2}R^{\alpha\beta}\wedge\eta_{\alpha\beta}-\frac{1}{2}T^{\alpha}\wedge H_{\alpha}-\frac{1}{2}R^{\alpha\beta}\wedge H_{\alpha\beta}.$
mielke-geometrodynamics_E08565 (8.6.2)	183	■ 1.000	$H_{\alpha}:=-\partial L_{\mathrm{qPG}}/\partial T^{\alpha}=\frac{1}{\ell^2}^*\left(\sum_{M=1}^3a_{(M)}^{(M)}T_{\alpha}\right),$	$H_{\alpha}:=-\partial L_{\mathrm{qPG}}/\partial T^{\alpha}=\frac{1}{\ell^2}^*\left(\sum_{M=1}^3a_{(M)}^{(M)}T_{\alpha}\right),$	$H_{\alpha}:=-\partial L_{\mathrm{qPG}}/\partial T^{\alpha}=\frac{1}{\ell^2}^*\left(\sum_{M=1}^3a_{(M)}^{(M)}T_{\alpha}\right),$
mielke-geometrodynamics_E08569 (8.6.7)	184	■ 1.000	$\begin{aligned} D H_{\alpha}-E_{\alpha} &= \Sigma_{\alpha}, \\ D H_{\alpha\beta}+\vartheta_{[\alpha}\wedge H_{\beta]} &= \tau_{\alpha\beta}, \end{aligned}$	$\begin{aligned} D H_{\alpha}-E_{\alpha} &= \Sigma_{\alpha}, \\ D H_{\alpha\beta}+\vartheta_{[\alpha}\wedge H_{\beta]} &= \tau_{\alpha\beta}, \end{aligned}$	$\begin{aligned} D H_{\alpha}-E_{\alpha} &= \Sigma_{\alpha}, \\ D H_{\alpha\beta}+\vartheta_{[\alpha}\wedge H_{\beta]} &= \tau_{\alpha\beta}, \end{aligned}$
mielke-geometrodynamics_E08571 (8.6.9)	184	■ 1.000	$\frac{\theta_{\mathrm{T}}^*}{2\ell^2}\eta_{\alpha\beta\gamma}T^{\gamma}+\frac{\theta_{\mathrm{T}}}{\ell^2}\vartheta_{[\alpha}\wedge T_{\beta]}+\vartheta_{[\alpha}\wedge H_{\beta]}(**)=\tau_{\alpha\beta}.$	$\frac{\theta_{\mathrm{T}}^*}{2\ell^2}\eta_{\alpha\beta\gamma}T^{\gamma}+\frac{\theta_{\mathrm{T}}}{\ell^2}\vartheta_{[\alpha}\wedge T_{\beta]}+\vartheta_{[\alpha}\wedge H_{\beta]}(**)=\tau_{\alpha\beta}.$	$\frac{\theta_{\mathrm{T}}^*}{2\ell^2}\eta_{\alpha\beta\gamma}T^{\gamma}+\frac{\theta_{\mathrm{T}}}{\ell^2}\vartheta_{[\alpha}\wedge T_{\beta]}+\vartheta_{[\alpha}\wedge H_{\beta]}(**)=\tau_{\alpha\beta}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_185 EQ0576 (8.6.14)	185	■ 1.000	$\Lambda_{\mathrm{eff}}=\frac{1}{\theta_{\mathrm{T}}}\left[\Lambda-\frac{1}{4}(\theta_{\mathrm{T}}^*-a_0)R\right]=\frac{\Lambda}{\theta_{\mathrm{T}}} - \Lambda_{\theta}$	$\Lambda_{\mathrm{eff}} = \frac{1}{\theta_{\mathrm{T}}^*} \left[\Lambda - \frac{1}{4} (\theta_{\mathrm{T}}^* - a_0) R \right] = \frac{\Lambda}{\theta_{\mathrm{T}}^*} - \Lambda_{\theta}$	$\Lambda_{\mathrm{eff}} = \frac{1}{\theta_{\mathrm{T}}^*} \left[\Lambda - \frac{1}{4} (\theta_{\mathrm{T}}^* - a_0) R \right] = \frac{\Lambda}{\theta_{\mathrm{T}}^*} - \Lambda_{\theta}$
mielke-geometrodynamics_186 EQ0578 (8.6.16)	186	■ 1.000	$\widetilde{L}_{\mathrm{qPG}}=L_{\mathrm{qPG}}+\lambda_{\alpha}\wedge T^{\alpha}.$	$\widetilde{L}_{\mathrm{qPG}} = L_{\mathrm{qPG}} + \lambda_{\alpha} \wedge T^{\alpha}.$	$\widetilde{L}_{\mathrm{qPG}} = L_{\mathrm{qPG}} + \lambda_{\alpha} \wedge T^{\alpha}.$
mielke-geometrodynamics_188 EQ0582 (8.6.20)	188	■ 1.000	$\Lambda_{\theta}=\frac{3\left(\theta_{\mathrm{T}}^*-a_0\right)^2}{2\ell^2\left(\theta_{\mathrm{L}}^*-b_6\right)\theta_{\mathrm{T}}^*}$	$\Lambda_{\theta} = \frac{3(\theta_{\mathrm{T}}^* - a_0)^2}{2\ell^2(\theta_{\mathrm{L}}^* - b_6)\theta_{\mathrm{T}}^*}$	$\Lambda_{\theta} = \frac{3(\theta_{\mathrm{T}}^* - a_0)^2}{2\ell^2(\theta_{\mathrm{L}}^* - b_6)\theta_{\mathrm{T}}^*}$
mielke-geometrodynamics_193 EQ0586 (9.1.4)	193	■ 1.000	$d\,s^2=\left(\frac{N(r\pm t)}{r}\right)^2(-dt^2+dr^2+r^2d\Omega^2),$	$ds^2 = \left(\frac{N(r\pm t)}{r}\right)^2(-dt^2 + dr^2 + r^2d\Omega^2),$	$ds^2 = \left(\frac{N(r\pm t)}{r}\right)^2(-dt^2 + dr^2 + r^2d\Omega^2),$
mielke-geometrodynamics_194 EQ0588 (9.1.6)	194	■ 1.000	$u^2-v^2=e^{r\,*/\,4\,\mu}$	$u^2 - v^2 = e^{r^{*/4\mu}}$	$u^2 - v^2 = e^{r^{*/4\mu}}$
mielke-geometrodynamics_194 EQ0589 (9.1.7)	194	■ 1.000	$(u+v)/(u-v)=e^{t\,/\,2\,\mu},$	$(u+v)/(u-v) = e^{t/2\mu},$	$(u+v)/(u-v) = e^{t/2\mu},$
mielke-geometrodynamics_194 EQ0590 (9.1.8)	194	■ 1.000	$d\,s^2=16\,\mu^2\,e^{\nu-r^{*/2\mu}}(-dv^2+du^2)+r^2d\Omega^2,$	$ds^2 = 16\mu^2e^{\nu-r^{*/2\mu}}(-dv^2 + du^2) + r^2d\Omega^2,$	$ds^2 = 16\mu^2e^{\nu-r^{*/2\mu}}(-dv^2 + du^2) + r^2d\Omega^2,$
mielke-geometrodynamics_195 EQ0591 (9.1.9)	195	■ 1.000	$\Omega^{\{\}}=\Omega_{\mathrm{C}}^{\{\}}+\bar{\Omega}^{\{\}}-\frac{1}{6}R^{\{\}}\vartheta\wedge\vartheta,$	$\Omega^{\{\}} = \Omega_{\mathrm{C}}^{\{\}} + \bar{\Omega}^{\{\}} - \frac{1}{6}R^{\{\}}\vartheta \wedge \vartheta,$	$\Omega^{\{\}} = \Omega_{\mathrm{C}}^{\{\}} + \bar{\Omega}^{\{\}} - \frac{1}{6}R^{\{\}}\vartheta \wedge \vartheta,$
mielke-geometrodynamics_197 EQ0600 (9.1.18)	197	■ 1.000	$\begin{aligned} d\,s_{\mathrm{Sds}}^2&\,\&=\left(1-\frac{2\,\mu}{r}+\frac{\kappa}{4\ell^{*2}}\right)d\tau^2\\ &\,\&+\left(1-\frac{2\,\mu}{r}+\frac{\kappa}{4\ell^{*2}}\right)^{-1}dr^2+r^2d\Omega^2\end{aligned}$	$ds_{\mathrm{Sds}}^2 = \left(1 - \frac{2\mu}{r} + \frac{\kappa}{4\ell^{*2}}\right)d\tau^2 + \left(1 - \frac{2\mu}{r} + \frac{\kappa}{4\ell^{*2}}\right)^{-1}dr^2 + r^2d\Omega^2$	$ds_{\mathrm{Sds}}^2 = \left(1 - \frac{2\mu}{r} + \frac{\kappa}{4\ell^{*2}}\right)d\tau^2 + \left(1 - \frac{2\mu}{r} + \frac{\kappa}{4\ell^{*2}}\right)^{-1}dr^2 + r^2d\Omega^2,$
mielke-geometrodynamics_197 EQ0601 (9.1.19)	197	■ 1.000	$\begin{aligned} f&=\frac{2\,\mu}{r^2}\left(1-\frac{2\,\mu}{r}+\frac{\kappa}{4\ell^{*2}}\right)^{-1/2}\\ g&=h=k=0\end{aligned}$	$f = \frac{2\mu}{r^2} \left(1 - \frac{2\mu}{r} + \frac{\kappa}{4\ell^{*2}}\right)^{-1/2}$ $g = h = k = 0$	$f = \frac{2\mu}{r^2} \left(1 - \frac{2\mu}{r} + \frac{\kappa}{4\ell^{*2}}\right)^{-1/2}$ $g = h = k = 0$
mielke-geometrodynamics_198 EQ0603 (9.2.2)	198	■ 1.000	$p_1\left(M^4\right)=-3\,\tau$	$p_1\left(M^4\right) = -3\tau$	$p_1(M^4) = -3\tau$
mielke-geometrodynamics_200 EQ0607 (9.2.8)	200	■ 1.000	$\begin{aligned} \chi(M\,\#\,N)\,\&=\chi(M)+\chi(N)-\chi\left(S^{\{n\}}\right),\\ \chi(M\times N)\,\&=\chi(M)\cdot\chi(N),\\ \chi\left(M^2\right)&=2(g-1),\\ \chi\left(\mathbb{C}P^2\right)&=3\end{aligned}$	$\chi(M\#N) = \chi(M) + \chi(N) - \chi(S^n),$ $\chi(M\times N) = \chi(M) \cdot \chi(N),$ $\chi(M^2) = 2(g-1),$ $\chi(\mathbb{C}P^2) = 3$	$\chi(M\#N) = \chi(M) + \chi(N) - \chi(S^n),$ $\chi(M\times N) = \chi(M) \cdot \chi(N),$ $\chi(M^2) = 2(g-1),$ $\chi(\mathbb{C}P^2) = 3$
mielke-geometrodynamics_201 EQ0608 (9.2.10)	201	■ 1.000	$\chi\left(M^{2\,k+1}\right)=0$	$\chi\left(M^{2k+1}\right) = 0$	$\chi(M^{2k+1}) = 0$
mielke-geometrodynamics_208 EQ0612 (10.0.1)	208	■ 1.000	$L_{\mathrm{EC}}=-\frac{\chi}{\ell}\vartheta^{\alpha}\wedge R_{\alpha}^{\star}\wedge R_{\alpha}^{\star}$	$L_{\mathrm{EC}} = -\frac{\chi}{\ell}\vartheta^{\alpha} \wedge R_{\alpha}^{\star}$	$L_{\mathrm{EC}} = -\frac{\chi}{\ell}\vartheta^{\alpha} \wedge R_{\alpha}^{\star}$
mielke-geometrodynamics_209 EQ0614 (10.1.2)	209	■ 1.000	$T^{\alpha}:=d\vartheta^{\alpha}-(-1)^s\eta^{\alpha\beta}\wedge\Gamma_{\beta}^{\star}$	$T^{\alpha} := d\vartheta^{\alpha} - (-1)^s\eta^{\alpha\beta} \wedge \Gamma_{\beta}^{\star}$	$T^{\alpha} := d\vartheta^{\alpha} - (-1)^s\eta^{\alpha\beta} \wedge \Gamma_{\beta}^{\star}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_E00615 (10.1.3)	209	■ 1.000	$R_{\alpha}^{\star}=\frac{1}{2} \eta_{\alpha} \beta \gamma \wedge \frac{1}{2} \eta_{\alpha} \beta \gamma \wedge \Gamma_{\beta}^{\star} \wedge \Gamma_{\gamma}^{\star}+\frac{(-1)^s}{2} \eta_{\alpha} \beta \gamma \wedge \Gamma_{\beta}^{\star} \wedge \Gamma_{\gamma}^{\star},$	$R_{\alpha}^{\star}=\frac{1}{2} \eta_{\alpha \beta \gamma} R^{\beta \gamma}:=d \Gamma_{\alpha}^{\star}+\frac{(-1)^s}{2} \eta_{\alpha}^{\beta \gamma} \Gamma_{\beta}^{\star} \wedge \Gamma_{\gamma}^{\star},$	$R_{\alpha}^{\star}=\frac{1}{2} \eta_{\alpha \beta \gamma} R^{\beta \gamma}:=d \Gamma_{\alpha}^{\star}+\frac{(-1)^s}{2} \eta_{\alpha}^{\beta \gamma} \Gamma_{\beta}^{\star} \wedge \Gamma_{\gamma}^{\star},$
mielke-geometrodynamics_E00616 (10.1.4)	209	■ 1.000	$D T^{\alpha} \equiv(-1)^s \eta^{\alpha \beta} \wedge R_{\beta}^{\star}$	$D T^{\alpha} \equiv(-1)^s \eta^{\alpha \beta} \wedge R_{\beta}^{\star}$	$D T^{\alpha} \equiv(-1)^s \eta^{\alpha \beta} \wedge R_{\beta}^{\star}$
mielke-geometrodynamics_E00617 (10.1.5)	209	■ 1.000	$D R_{\alpha}^{\star} \equiv 0$	$D R_{\alpha}^{\star} \equiv 0$	$D R_{\alpha}^{\star} \equiv 0$
mielke-geometrodynamics_E00618 (10.1.6)	209	■ 1.000	$C_{\mathrm{T}}:=\frac{1}{2} \ell \wedge \vartheta^{\alpha} \wedge T_{\alpha}=-\frac{(-1)^s}{\ell^2} \eta^{\alpha} \wedge K_{\alpha}^{\star},$	$C_{\mathrm{T}}:=\frac{1}{2 \ell^2} \vartheta^{\alpha} \wedge T_{\alpha}=-\frac{(-1)^s}{\ell^2} \eta^{\alpha} \wedge K_{\alpha}^{\star},$	$C_{\mathrm{T}}:=\frac{1}{2 \ell^2} \vartheta^{\alpha} \wedge T_{\alpha}=-\frac{(-1)^s}{\ell^2} \eta^{\alpha} \wedge K_{\alpha}^{\star},$
mielke-geometrodynamics_E00619 (10.1.7)	209	■ 1.000	$C_{\mathrm{L}}:=(-1)^s \Gamma^{\star \alpha} \wedge R_{\alpha}^{\star}-\frac{1}{3 !} \eta_{\alpha \beta \gamma} \Gamma^{\star \alpha} \wedge \Gamma^{\star \beta} \wedge \Gamma^{\star \gamma}.$	$C_{\mathrm{L}}:=(-1)^s \Gamma^{\star \alpha} \wedge R_{\alpha}^{\star}-\frac{1}{3 !} \eta_{\alpha \beta \gamma} \Gamma^{\star \alpha} \wedge \Gamma^{\star \beta} \wedge \Gamma^{\star \gamma}.$	$C_{\mathrm{L}}:=(-1)^s \Gamma^{\star \alpha} \wedge R_{\alpha}^{\star}-\frac{1}{3 !} \eta_{\alpha \beta \gamma} \Gamma^{\star \alpha} \wedge \Gamma^{\star \beta} \wedge \Gamma^{\star \gamma}.$
mielke-geometrodynamics_E00621 (10.1.9)	209	■ 1.000	$\frac{\delta C_{\mathrm{L}}}{\delta \vartheta^{\alpha}}=0 \quad \text { and } \quad \frac{\delta C_{\mathrm{L}}}{\delta \Gamma^{\star \alpha}}=(-1)^s 2 R_{\alpha}^{\star},$	$\frac{\delta C_{\mathrm{L}}}{\delta \vartheta^{\alpha}}=0 \quad \text { and } \quad \frac{\delta C_{\mathrm{L}}}{\delta \Gamma^{\star \alpha}}=(-1)^s 2 R_{\alpha}^{\star},$	$\frac{\delta C_{\mathrm{L}}}{\delta \vartheta^{\alpha}}=0 \quad \text { and } \quad \frac{\delta C_{\mathrm{L}}}{\delta \Gamma^{\star \alpha}}=(-1)^s 2 R_{\alpha}^{\star},$
mielke-geometrodynamics_E00622 (10.1.10)	210	■ 1.000	$C_{\mathrm{TL}}:=\frac{1}{\ell}\left(\Gamma^{\star \alpha} \wedge T_{\alpha}-\frac{(-1)^s}{2} \eta_{\alpha \beta \gamma} \Gamma^{\star \alpha} \wedge \Gamma^{\star \beta} \wedge \vartheta^{\gamma}\right).$	$C_{\mathrm{TL}}:=\frac{1}{\ell}\left(\Gamma^{\star \alpha} \wedge T_{\alpha}-\frac{(-1)^s}{2} \eta_{\alpha \beta \gamma} \Gamma^{\star \alpha} \wedge \Gamma^{\star \beta} \wedge \vartheta^{\gamma}\right).$	$C_{\mathrm{TL}}:=\frac{1}{\ell}\left(\Gamma^{\star \alpha} \wedge T_{\alpha}-\frac{(-1)^s}{2} \eta_{\alpha \beta \gamma} \Gamma^{\star \alpha} \wedge \Gamma^{\star \beta} \wedge \vartheta^{\gamma}\right).$
mielke-geometrodynamics_E00625 (10.1.13)	210	■ 1.000	$D H_{\alpha}^{\star}-S_{\alpha}=\tau_{\alpha}^{\star},$	$D H_{\alpha}^{\star}-S_{\alpha}=\tau_{\alpha}^{\star},$	$D H_{\alpha}^{\star}-S_{\alpha}=\tau_{\alpha}^{\star},$
mielke-geometrodynamics_E00626 (10.1.14)	211	■ 1.000	$H_{\alpha}:=\frac{\partial L}{\partial T^{\alpha}}, \quad H_{\alpha}^{\star}:=\frac{(-1)^s}{2} \frac{\partial L}{\partial R^{\star \alpha}},$	$H_{\alpha}:=\frac{\partial L}{\partial T^{\alpha}}, \quad H_{\alpha}^{\star}:=\frac{(-1)^s}{2} \frac{\partial L}{\partial R^{\star \alpha}},$	$H_{\alpha}:=\frac{\partial L}{\partial T^{\alpha}}, \quad H_{\alpha}^{\star}:=\frac{(-1)^s}{2} \frac{\partial L}{\partial R^{\star \alpha}},$
mielke-geometrodynamics_E00628 (10.1.16)	211	■ 1.000	$S_{\alpha}:=\frac{(-1)^s}{2} \frac{\partial L}{\partial \Gamma^{\star \alpha}}=\frac{1}{2} \eta_{\alpha \beta} \wedge H^{\beta}.$	$S_{\alpha}:=\frac{(-1)^s}{2} \frac{\partial L}{\partial \Gamma^{\star \alpha}}=\frac{1}{2} \eta_{\alpha \beta} \wedge H^{\beta}.$	$S_{\alpha}:=\frac{(-1)^s}{2} \frac{\partial L}{\partial \Gamma^{\star \alpha}}=\frac{1}{2} \eta_{\alpha \beta} \wedge H^{\beta}.$
mielke-geometrodynamics_E00629 (10.1.17)	211	■ 1.000	$\Sigma_{\alpha}:=\frac{\delta L_{\mathrm{MAT}}}{\delta \vartheta^{\alpha}}, \quad \tau_{\alpha}^{\star}:=\frac{(-1)^s}{2} \frac{\delta L_{\mathrm{MAT}}}{\delta \Gamma^{\star \alpha}},$	$\Sigma_{\alpha}:=\frac{\delta L_{\mathrm{MAT}}}{\delta \vartheta^{\alpha}}, \quad \tau_{\alpha}^{\star}:=\frac{(-1)^s}{2} \frac{\delta L_{\mathrm{MAT}}}{\delta \Gamma^{\star \alpha}},$	$\Sigma_{\alpha}:=\frac{\delta L_{\mathrm{MAT}}}{\delta \vartheta^{\alpha}}, \quad \tau_{\alpha}^{\star}:=\frac{(-1)^s}{2} \frac{\delta L_{\mathrm{MAT}}}{\delta \Gamma^{\star \alpha}},$
mielke-geometrodynamics_E00630 (10.2.1)	211	■ 1.000	$L_{\mathrm{EC}}:=\frac{\chi}{\ell} \vartheta^{\alpha} \wedge R_{\alpha}^{\star} \equiv-\chi C_{\mathrm{TL}}-\frac{\chi}{\ell} d\left(\Gamma_{\alpha}^{\star} \wedge \vartheta^{\alpha}\right)$	$L_{\mathrm{EC}}:=\frac{\chi}{\ell} \vartheta^{\alpha} \wedge R_{\alpha}^{\star} \equiv-\chi C_{\mathrm{TL}}-\frac{\chi}{\ell} d\left(\Gamma_{\alpha}^{\star} \wedge \vartheta^{\alpha}\right)$	$L_{\mathrm{EC}}:=\frac{\chi}{\ell} \vartheta^{\alpha} \wedge R_{\alpha}^{\star} \equiv-\chi C_{\mathrm{TL}}-\frac{\chi}{\ell} d\left(\Gamma_{\alpha}^{\star} \wedge \vartheta^{\alpha}\right)$
mielke-geometrodynamics_E00631 (10.2.2)	211	■ 1.000	$L_{\mathrm{MB}}\left(\vartheta^{\alpha}, \Gamma_{\alpha}^{\star}\right)=\theta_{\mathrm{T}} C_{\mathrm{T}}+\theta_{\mathrm{L}} C_{\mathrm{L}}+\theta_{\mathrm{TL}} C_{\mathrm{TL}},$	$L_{\mathrm{MB}}\left(\vartheta^{\alpha}, \Gamma_{\alpha}^{\star}\right)=\theta_{\mathrm{T}} C_{\mathrm{T}}+\theta_{\mathrm{L}} C_{\mathrm{L}}+\theta_{\mathrm{TL}} C_{\mathrm{TL}},$	$L_{\mathrm{MB}}\left(\vartheta^{\alpha}, \Gamma_{\alpha}^{\star}\right)=\theta_{\mathrm{T}} C_{\mathrm{T}}+\theta_{\mathrm{L}} C_{\mathrm{L}}+\theta_{\mathrm{TL}} C_{\mathrm{TL}},$
mielke-geometrodynamics_E00632 (10.2.3)	212	■ 1.000	$-\theta_{\mathrm{TL}} R_{\alpha}^{\star}-\frac{1}{\ell} \theta_{\mathrm{T}} T_{\alpha}=\ell \Sigma_{\alpha}$	$-\theta_{\mathrm{TL}} R_{\alpha}^{\star}-\frac{1}{\ell} \theta_{\mathrm{T}} T_{\alpha}=\ell \Sigma_{\alpha}$	$-\theta_{\mathrm{TL}} R_{\alpha}^{\star}-\frac{1}{\ell} \theta_{\mathrm{T}} T_{\alpha}=\ell \Sigma_{\alpha}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_212 E00634 (10.2.5)	212	■ 1.000	$T_{\{\alpha\}}-\frac{\varepsilon}{\ell}\eta_{\alpha}=\frac{2}{A}\ell\left(\theta_{\mathrm{TL}}\tau_{\alpha}^{\star}-\theta_{\mathrm{L}}\ell\Sigma_{\alpha}\right)$	$T_{\alpha}-\frac{\varepsilon}{\ell}\eta_{\alpha}=\frac{2}{A}\ell\left(\theta_{\mathrm{TL}}\tau_{\alpha}^{\star}-\theta_{\mathrm{L}}\ell\Sigma_{\alpha}\right)$	$T_{\alpha}-\frac{\varepsilon}{\ell}\eta_{\alpha}=\frac{2}{A}\ell\left(\theta_{\mathrm{TL}}\tau_{\alpha}^{\star}-\theta_{\mathrm{L}}\ell\Sigma_{\alpha}\right)$
mielke-geometrodynamics_212 E00635 (10.2.6)	212	■ 1.000	$R_{\{\alpha\}}^{\star}-\frac{\rho}{\ell^2}\eta_{\alpha}=\frac{2}{A}\left(\theta_{\mathrm{TL}}\ell\Sigma_{\alpha}-\theta_{\mathrm{T}}\tau_{\alpha}^{\star}\right)$	$R_{\alpha}^{\star}-\frac{\rho}{\ell^2}\eta_{\alpha}=\frac{2}{A}\left(\theta_{\mathrm{TL}}\ell\Sigma_{\alpha}-\theta_{\mathrm{T}}\tau_{\alpha}^{\star}\right)$	$R_{\alpha}^{\star}-\frac{\rho}{\ell^2}\eta_{\alpha}=\frac{2}{A}\left(\theta_{\mathrm{TL}}\ell\Sigma_{\alpha}-\theta_{\mathrm{T}}\tau_{\alpha}^{\star}\right)$
mielke-geometrodynamics_213 E00638 (10.2.9)	213	■ 1.000	$R_{\{\alpha\}}^{\star}\equiv R_{\{\alpha\}}^{\{\}\star}-DK_{\alpha}^{\star}+\frac{1}{2}\eta_{\alpha\beta\gamma}K^{\star\beta}\wedge K^{\star\gamma}$	$R_{\alpha}^{\star}\equiv R_{\alpha}^{\{\}\star}-DK_{\alpha}^{\star}+\frac{1}{2}\eta_{\alpha\beta\gamma}K^{\star\beta}\wedge K^{\star\gamma}$	$R_{\alpha}^{\star}\equiv R_{\alpha}^{\{\}\star}-DK_{\alpha}^{\star}+\frac{1}{2}\eta_{\alpha\beta\gamma}K^{\star\beta}\wedge K^{\star\gamma}$
mielke-geometrodynamics_213 E00640 (10.2.11)	213	■ 1.000	$\Lambda_{\mathrm{eff}}=-\left[4\left(\rho-\left(1+2(-1)^s\right)\varepsilon^2\right)/4\ell^2=\theta_{\mathrm{T}}^2\left[\left(9+2(-1)^s\right)\theta_{\mathrm{TL}}^2+8\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right]/(2A\ell)^2\right]$	$\Lambda_{\mathrm{eff}}=-\left[4\rho-\left(1+2(-1)^s\right)\varepsilon^2\right]/4\ell^2=\theta_{\mathrm{T}}^2\left[\left(9+2(-1)^s\right)\theta_{\mathrm{TL}}^2+8\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right]/(2A\ell)^2$	$\Lambda_{\mathrm{eff}}=-\left[4\rho-\left(1+2(-1)^s\right)\varepsilon^2\right]/4\ell^2=\theta_{\mathrm{T}}^2\left[\left(9+2(-1)^s\right)\theta_{\mathrm{TL}}^2+8\theta_{\mathrm{T}}\theta_{\mathrm{L}}\right]/(2A\ell)^2$
mielke-geometrodynamics_213 E00642 (10.2.13)	213	■ 1.000	$ds^2=-N^2(r)dt^2+N^{-2}(r)dr^2+r^2\left[d\phi+N^{\phi}(r)dt\right]^2$	$ds^2=-N^2(r)dt^2+N^{-2}(r)dr^2+r^2\left[d\phi+N^{\phi}(r)dt\right]^2$	$ds^2=-N^2(r)dt^2+N^{-2}(r)dr^2+r^2\left[d\phi+N^{\phi}(r)dt\right]^2$
mielke-geometrodynamics_213 E00643 (10.2.14)	213	■ 1.000	$N^2(r)=-M-\Lambda_{\mathrm{eff}}r^2+\frac{J^2}{4r^2},\quad N^{\phi}(r)=-\frac{J}{2r^2},$	$N^2(r)=-M-\Lambda_{\mathrm{eff}}r^2+\frac{J^2}{4r^2},\quad N^{\phi}(r)=-\frac{J}{2r^2},$	$N^2(r)=-M-\Lambda_{\mathrm{eff}}r^2+\frac{J^2}{4r^2},\quad N^{\phi}(r)=-\frac{J}{2r^2},$
mielke-geometrodynamics_214 E00647 (10.3.4)	214	■ 1.000	$S_{\{\alpha\}}=\frac{\ell}{\varepsilon}E_{\alpha}$	$S_{\alpha}=\frac{\ell}{\varepsilon}E_{\alpha}$	$S_{\alpha}=\frac{\ell}{\varepsilon}E_{\alpha}$
mielke-geometrodynamics_215 E00648 (10.3.5)	215	■ 1.000	$H_{\{\alpha\}}^{\star}=\frac{\ell}{\varepsilon}H_{\alpha}$	$H_{\alpha}^{\star}=\frac{\ell}{\varepsilon}H_{\alpha}$	$H_{\alpha}^{\star}=\frac{\ell}{\varepsilon}H_{\alpha}$
mielke-geometrodynamics_215 E00649 (10.3.6)	215	■ 1.000	$T_{\{\alpha\}}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\{\alpha\}}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha}$	$T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha}$	$T_{\alpha}=\frac{\varepsilon}{\ell}\eta_{\alpha},\quad R_{\alpha}^{\star}=\frac{\rho}{\ell^2}\eta_{\alpha},$
mielke-geometrodynamics_215 E00650 (10.3.7)	215	■ 1.000	$H_{\{\alpha\}}^{\star}=\delta\ell H_{\alpha}+\frac{\gamma}{\ell}\vartheta_{\alpha}$	$H_{\alpha}^{\star}=\delta\ell H_{\alpha}+\frac{\gamma}{\ell}\vartheta_{\alpha}$	$H_{\alpha}^{\star}=\delta\ell H_{\alpha}+\frac{\gamma}{\ell}\vartheta_{\alpha}$
mielke-geometrodynamics_215 E00651 (10.3.8)	215	■ 1.000	$\delta\ell DH_{\alpha}+\frac{\gamma}{\ell}T_{\alpha}-\frac{1}{2}\eta_{\alpha\beta}\wedge H^{\beta}=\tau_{\alpha}^{\star}$	$\delta\ell DH_{\alpha}+\frac{\gamma}{\ell}T_{\alpha}-\frac{1}{2}\eta_{\alpha\beta}\wedge H^{\beta}=\tau_{\alpha}^{\star}$	$\delta\ell DH_{\alpha}+\frac{\gamma}{\ell}T_{\alpha}-\frac{1}{2}\eta_{\alpha\beta}\wedge H^{\beta}=\tau_{\alpha}^{\star}$
mielke-geometrodynamics_216 E00653 (10.3.10)	216	■ 1.000	$H_{\{\alpha\}}^{\star}=\frac{\gamma}{\ell(1+\delta\varepsilon)}\vartheta_{\alpha}=-\frac{\ell}{\varepsilon}H_{\alpha},$	$H_{\alpha}^{\star}=\frac{\gamma}{\ell(1+\delta\varepsilon)}\vartheta_{\alpha}=-\frac{\ell}{\varepsilon}H_{\alpha},$	$H_{\alpha}^{\star}=\frac{\gamma}{\ell(1+\delta\varepsilon)}\vartheta_{\alpha}=-\frac{\ell}{\varepsilon}H_{\alpha},$
mielke-geometrodynamics_217 E00659 (10.3.16)	217	■ 1.000	$H=\left(\vartheta^{\alpha}\wedge H_{\alpha}\right),$	$H=\left(\vartheta^{\alpha}\wedge H_{\alpha}\right),$	$H=\left(\vartheta^{\alpha}\wedge H_{\alpha}\right),$
mielke-geometrodynamics_217 E00660 (10.3.17)	217	■ 1.000	$t=\left(T^{\alpha}\wedge H_{\alpha}\right),\quad \mathscr{A}:=\left(\vartheta^{\alpha}\wedge T_{\alpha}\right),\quad h=\left(\eta^{\alpha}\wedge H_{\alpha}\right),$	$t=\left(T^{\alpha}\wedge H_{\alpha}\right),\quad \mathscr{A}:=\left(\vartheta^{\alpha}\wedge T_{\alpha}\right),\quad h=\left(\eta^{\alpha}\wedge H_{\alpha}\right),$	$t=\left(T^{\alpha}\wedge H_{\alpha}\right),\quad \mathscr{A}:=\left(\vartheta^{\alpha}\wedge T_{\alpha}\right),\quad h=\left(\eta^{\alpha}\wedge H_{\alpha}\right),$
mielke-geometrodynamics_217 E00663 (10.3.20)	217	■ 1.000	$ds^2=(-1)^sdt^2+d\mathscr{A}^2+dh^2$	$ds^2=(-1)^sdt^2+d\mathscr{A}^2+dh^2$	$ds^2=(-1)^sdt^2+d\mathscr{A}^2+dh^2$
mielke-geometrodynamics_218 E00665 (10.3.22)	218	■ 1.000	$D^{\star}\vartheta_{\alpha}:=\frac{\ell}{\gamma}\left[D^{\star}H_{\alpha}^{\star}-\delta\ell D^{\star}H_{\alpha}\right]$	$D^{\star}\vartheta_{\alpha}:=\frac{\ell}{\gamma}\left[D^{\star}H_{\alpha}^{\star}-\delta\ell D^{\star}H_{\alpha}\right]$	$D^{\star}\vartheta_{\alpha}:=\frac{\ell}{\gamma}\left[D^{\star}H_{\alpha}^{\star}-\delta\ell D^{\star}H_{\alpha}\right]$
mielke-geometrodynamics_219 E00668 (10.3.25)	219	■ 1.000	$D^{\star}\vartheta^{\alpha}\cong 0,$	$D^{\star}\vartheta^{\alpha}\cong 0,$	$D^{\star}\vartheta^{\alpha}\cong 0,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_220 EQ0672 (10.3.29)	220	■ 1.000	$P_{\alpha}=\int \check{\mathrm{P}}_{\alpha}, \quad \mathrm{J}_{\alpha}=\int \check{\mathrm{J}}_{\alpha}^{\star}.$	$P_{\alpha}=\int \check{\mathcal{E}}_{\alpha}, \quad J_{\alpha}=\int \check{\mathcal{J}}_{\alpha}^{\star}.$	$P_{\alpha}=\int \check{\mathcal{E}}_{\alpha}, \quad J_{\alpha}=\int \check{\mathcal{J}}_{\alpha}^{\star}.$
mielke-geometrodynamics_220 EQ0675 (10.3.32)	220	■ 1.000	$\check{\mathrm{J}}_{\alpha}^{\star}=-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$	$\check{\mathcal{J}}_{\alpha}^{\star}=-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$	$\check{\mathcal{J}}_{\alpha}^{\star}=-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$
mielke-geometrodynamics_220 EQ0677 (10.3.34)	220	■ 1.000	$\check{\mathrm{J}}_{\alpha}^{\mathrm{MB}\star}=(-1)^s\frac{\chi}{2\ell}d\vartheta_{\alpha}-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$	$\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}\star}=(-1)^s\frac{\chi}{2\ell}d\vartheta_{\alpha}-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$	$\check{\mathcal{J}}_{\alpha}^{\mathrm{MB}\star}=(-1)^s\frac{\chi}{2\ell}d\vartheta_{\alpha}-\frac{\theta_{\mathrm{L}}}{2}d\Gamma_{\alpha}^{\star}$
mielke-geometrodynamics_221 EQ0679 (10.3.36)	221	■ 1.000	$\widetilde{\mathrm{C}}_{\mathrm{L}}=(-1)^s\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{R}_{\alpha}^{\star}-\frac{1}{3!}\eta_{\alpha\beta\gamma}\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{\Gamma}^{\star\beta}\wedge\widetilde{\Gamma}^{\star\gamma},$	$\widetilde{C}_{\mathrm{L}}=(-1)^s\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{R}_{\alpha}^{\star}-\frac{1}{3!}\eta_{\alpha\beta\gamma}\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{\Gamma}^{\star\beta}\wedge\widetilde{\Gamma}^{\star\gamma},$	$\widetilde{C}_{\mathrm{L}}=(-1)^s\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{R}_{\alpha}^{\star}-\frac{1}{3!}\eta_{\alpha\beta\gamma}\widetilde{\Gamma}^{\star\alpha}\wedge\widetilde{\Gamma}^{\star\beta}\wedge\widetilde{\Gamma}^{\star\gamma},$
mielke-geometrodynamics_221 EQ0683 (10.3.41)	221	■ 1.000	$c=12\cdot 4\pi\theta_{\mathrm{L}}, \quad \tilde{c}=12\cdot 4\pi\tilde{\theta}_{\mathrm{L}},$	$c=12\cdot 4\pi\theta_{\mathrm{L}}, \quad \tilde{c}=12\cdot 4\pi\tilde{\theta}_{\mathrm{L}},$	$c=12\cdot 4\pi\theta_{\mathrm{L}}, \quad \tilde{c}=12\cdot 4\pi\tilde{\theta}_{\mathrm{L}},$
mielke-geometrodynamics_222 EQ0684 (10.3.42)	222	■ 1.000	$\widetilde{L}=L+\lambda_{\alpha}\wedge T^{\alpha}.$	$\widetilde{L}=L+\lambda_{\alpha}\wedge T^{\alpha}.$	$\widetilde{L}=L+\lambda_{\alpha}\wedge T^{\alpha}.$
mielke-geometrodynamics_223 EQ0691 (10.5.2)	223	■ 1.000	$[\sigma^{\alpha},\sigma^{\beta}]=2i\eta^{\alpha\beta\gamma}\sigma_{\gamma}.$	$[\sigma^{\alpha},\sigma^{\beta}]=2i\eta^{\alpha\beta\gamma}\sigma_{\gamma}.$	$[\sigma^{\alpha},\sigma^{\beta}]=2i\eta^{\alpha\beta\gamma}\sigma_{\gamma}.$
mielke-geometrodynamics_224 EQ0692 (10.5.3)	224	■ 1.000	$\gamma_0=i\sigma^2, \quad \gamma_1=\sigma^1, \quad \gamma_2=\sigma^3.$	$\gamma_0=i\sigma^2, \quad \gamma_1=\sigma^1, \quad \gamma_2=\sigma^3.$	$\gamma_0=i\sigma^2, \quad \gamma_1=\sigma^1, \quad \gamma_2=\sigma^3.$
mielke-geometrodynamics_224 EQ0693 (10.5.4)	224	■ 1.000	$\gamma_{\alpha}\gamma_{\beta}=g_{\alpha\beta}\mathbf{1}+\eta_{\alpha\beta\nu}\gamma^{\nu},$	$\gamma_{\alpha}\gamma_{\beta}=g_{\alpha\beta}\mathbf{1}+\eta_{\alpha\beta\nu}\gamma^{\nu},$	$\gamma_{\alpha}\gamma_{\beta}=g_{\alpha\beta}\mathbf{1}+\eta_{\alpha\beta\nu}\gamma^{\nu},$
mielke-geometrodynamics_224 EQ0695 (10.5.6)	224	■ 1.000	$i^{\star}\gamma\wedge D\psi-m\psi\eta-\frac{i}{2}(D^{\star}\gamma)\psi=0$	$i^{\star}\gamma\wedge D\psi-m\psi\eta-\frac{i}{2}(D^{\star}\gamma)\psi=0$	$i^{\star}\gamma\wedge D\psi-m\psi\eta-\frac{i}{2}(D^{\star}\gamma)\psi=0$
mielke-geometrodynamics_224 EQ0697 (10.5.8)	224	■ 1.000	$\hat{D}:=D-\frac{1}{2}T,$	$\hat{D}:=D-\frac{1}{2}T,$	$\hat{D}:=D-\frac{1}{2}T,$
mielke-geometrodynamics_224 EQ0698 (10.5.9)	224	■ 1.000	$i^{\star}\gamma\wedge\hat{D}\psi-m\eta\psi=0,$	$i^{\star}\gamma\wedge\hat{D}\psi-m\eta\psi=0,$	$i^{\star}\gamma\wedge\hat{D}\psi-m\eta\psi=0,$
mielke-geometrodynamics_224 EQ0699 (10.5.10)	224	■ 1.000	$\hat{R}=R-\frac{1}{2}dT=\frac{i}{4}R^{\alpha\beta}\sigma_{\alpha\beta}-\frac{1}{2}dT$	$\hat{R}=R-\frac{1}{2}dT=\frac{i}{4}R^{\alpha\beta}\sigma_{\alpha\beta}-\frac{1}{2}dT$	$\hat{R}=R-\frac{1}{2}dT=\frac{i}{4}R^{\alpha\beta}\sigma_{\alpha\beta}-\frac{1}{2}dT$
mielke-geometrodynamics_225 EQ0703 (10.6.2)	225	■ 1.000	$d^{\star}F-m_{\mathrm{photon}}F=0$	$d^{\star}F-m_{\mathrm{photon}}F=0$	$d^{\star}F-m_{\mathrm{photon}}F=0$
mielke-geometrodynamics_226 EQ0705 (10.7.1)	226	■ 1.000	$X:=x_{\alpha}\vartheta^{\alpha}, \quad Y:=y_{\alpha}\vartheta^{\alpha}.$	$X:=x_{\alpha}\vartheta^{\alpha}, \quad Y:=y_{\alpha}\vartheta^{\alpha}.$	$X:=x_{\alpha}\vartheta^{\alpha}, \quad Y:=y_{\alpha}\vartheta^{\alpha}.$
mielke-geometrodynamics_226 EQ0706 (10.7.2)	226	■ 1.000	$\Sigma_{\alpha}=\varepsilon x_{\alpha}X\wedge Y, \quad \tau_{\alpha}^{\star}=\sigma y_{\alpha}X\wedge Y,$	$\Sigma_{\alpha}=\varepsilon x_{\alpha}X\wedge Y, \quad \tau_{\alpha}^{\star}=\sigma y_{\alpha}X\wedge Y,$	$\Sigma_{\alpha}=\varepsilon x_{\alpha}X\wedge Y, \quad \tau_{\alpha}^{\star}=\sigma y_{\alpha}X\wedge Y,$
mielke-geometrodynamics_226 EQ0707 (10.7.3)	226	■ 1.000	$\vartheta^{\alpha}\wedge\Sigma_{\alpha}=0, \quad \vartheta^{\alpha}\wedge\tau_{\alpha}^{\star}=0$	$\vartheta^{\alpha}\wedge\Sigma_{\alpha}=0, \quad \vartheta^{\alpha}\wedge\tau_{\alpha}^{\star}=0$	$\vartheta^{\alpha}\wedge\Sigma_{\alpha}=0, \quad \vartheta^{\alpha}\wedge\tau_{\alpha}^{\star}=0$
mielke-geometrodynamics_227 EQ0708 (10.7.4)	227	■ 1.000	$R^{\alpha\beta}=\varepsilon\ell^2x^{[\alpha}y^{\beta]}X\wedge Y,$	$R^{\alpha\beta}=\varepsilon\ell^2x^{[\alpha}y^{\beta]}X\wedge Y,$	$R^{\alpha\beta}=\varepsilon\ell^2x^{[\alpha}y^{\beta]}X\wedge Y,$
mielke-geometrodynamics_228 EQ0711 (10.7.7)	228	■ 1.000	$\mathscr{A}:=\ast(\vartheta^{\alpha}\wedge T_{\alpha})=-(-1)^s\frac{\varepsilon}{\ell^2}$	$\mathscr{A}:=\ast(\vartheta^{\alpha}\wedge T_{\alpha})=-(-1)^s\frac{\varepsilon}{\ell^2}$	$\mathscr{A}:=\ast(\vartheta^{\alpha}\wedge T_{\alpha})=-(-1)^s\frac{\varepsilon}{\ell^2}$
mielke-geometrodynamics_228 EQ0713 (10.8.2)	228	■ 1.000	$\gamma=\gamma_{\alpha}\vartheta^{\alpha}.$	$\gamma=\gamma_{\alpha}\vartheta^{\alpha}.$	$\gamma=\gamma_{\alpha}\vartheta^{\alpha}.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0714	228	■ 1.000	$\backslash\mathrm{bar}\{\Psi\} \wedge \Psi=0, \quad \backslash\mathrm{quad} \backslash\mathrm{bar}\{\Psi\} \wedge \gamma_5 \backslash\mathrm{gamma}_{\{5\}} \backslash\mathrm{gamma}^{\{\alpha\}} \Psi=0, \quad \backslash\mathrm{quad} \backslash\mathrm{bar}\{\Psi\} \wedge \gamma_5 \backslash\mathrm{gamma}_{\{5\}} \Psi=0$.	$\bar{\Psi} \wedge \Psi = 0, \quad \bar{\Psi} \wedge \gamma_5 \gamma^\alpha \Psi = 0, \quad \bar{\Psi} \wedge \gamma_5 \Psi = 0.$	$\bar{\Psi} \wedge \Psi = 0, \quad \bar{\Psi} \wedge \gamma_5 \gamma^\alpha \Psi = 0, \quad \bar{\Psi} \wedge \gamma_5 \Psi = 0.$
mielke-geometrodynamics_EQ0715 (10.9.1)	229	■ 1.000	$L_{\mathrm{RS}}=\frac{i}{4}(\backslash\mathrm{bar}\{\Psi\} \wedge D \Psi-\Psi \wedge \overline{D \Psi})+\frac{i}{4} m \backslash\mathrm{bar}\{\Psi\} \wedge \gamma \wedge \Psi,$	$L_{\mathrm{RS}}=\frac{i}{4}(\bar{\Psi} \wedge D \Psi - \Psi \wedge \overline{D \Psi}) + \frac{i}{4} m \bar{\Psi} \wedge \gamma \wedge \Psi,$	$L_{\mathrm{RS}}=\frac{i}{4}\left(\bar{\Psi} \wedge D \Psi - \Psi \wedge \overline{D \Psi}\right) + \frac{i}{4} m \bar{\Psi} \wedge \gamma \wedge \Psi,$
mielke-geometrodynamics_EQ0719 (10.9.5)	229	■ 1.000	$\Sigma_{\alpha}=-\frac{i}{4} m \backslash\mathrm{bar}\{\Psi\} \wedge \gamma_{\alpha} \Psi=-(-1)^s 2 m \tau_{\alpha}^{\star}$	$\Sigma_{\alpha}=-\frac{i}{4} m \bar{\Psi} \wedge \gamma_{\alpha} \Psi = -(-1)^s 2 m \tau_{\alpha}^{\star}$	$\Sigma_{\alpha}=-\frac{i}{4} m \bar{\Psi} \wedge \gamma_{\alpha} \Psi = -(-1)^s 2 m \tau_{\alpha}^{\star},$
mielke-geometrodynamics_EQ0721 (10.10.1)	230	■ 1.000	$L=L_{\mathrm{infy}}\left(\vartheta^{\alpha}, \Gamma_{\alpha}^{\star}, \Psi\right)=L_{\mathrm{MB}}+L_{\Psi}$	$L=L_{\infty}\left(\vartheta^{\alpha}, \Gamma_{\alpha}^{\star}, \Psi\right)=L_{\mathrm{MB}}+L_{\Psi}$	$L=L_{\infty}\left(\vartheta^{\alpha}, \Gamma_{\alpha}^{\star}, \Psi\right)=L_{\mathrm{MB}}+L_{\Psi}$
mielke-geometrodynamics_EQ0726 (10.10.6)	230	■ 1.000	$\frac{2}{\delta L} \frac{\delta L}{\delta \Psi}=D \Psi+\frac{1}{2} m \gamma \wedge \Psi \cong 0$	$\frac{2}{i} \frac{\delta L}{\delta \Psi}=D \Psi+\frac{1}{2} m \gamma \wedge \Psi \cong 0$	$\frac{2}{i} \frac{\delta L}{\delta \Psi}=D \Psi+\frac{1}{2} m \gamma \wedge \Psi \cong 0$
mielke-geometrodynamics_EQ0729 (10.10.9)	231	■ 1.000	$\gamma \wedge \gamma=-2 \gamma^{\alpha} \eta_{\alpha}$	$\gamma \wedge \gamma=-2 \gamma^{\alpha} \eta_{\alpha}$	$\gamma \wedge \gamma=-2 \gamma^{\alpha} \eta_{\alpha}$
mielke-geometrodynamics_EQ0730 (10.10.10)	231	■ 1.000	$i\left[\left(\frac{\theta_{\mathrm{TL}}}{\ell}-1\right) R_{\alpha}^{\star}+\left(\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right) T_{\alpha}+\frac{m^2}{2} \eta_{\alpha}\right] \wedge \gamma^{\alpha} \Psi=0,$	$i\left[\left(\frac{\theta_{\mathrm{TL}}}{\ell}-1\right) R_{\alpha}^{\star}+\left(\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right) T_{\alpha}+\frac{m^2}{2} \eta_{\alpha}\right] \wedge \gamma^{\alpha} \Psi=0,$	$i\left[\left(\frac{\theta_{\mathrm{TL}}}{\ell}-1\right) R_{\alpha}^{\star}+\left(\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right) T_{\alpha}+\frac{m^2}{2} \eta_{\alpha}\right] \wedge \gamma^{\alpha} \Psi=0,$
mielke-geometrodynamics_EQ0731 (10.10.11)	231	■ 1.000	$\begin{aligned} & i\left[\left(\frac{\theta_{\mathrm{TL}}}{\ell}-1\right) R_{\alpha}^{\star}+\left(\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right) T_{\alpha}+\frac{m^2}{2} \eta_{\alpha}\right] \wedge \gamma^{\alpha} \Psi \simeq 0 \\ & +\tau_{\alpha}^{\star}] \wedge \gamma^{\alpha} \Psi \simeq 0 \end{aligned}$	$i\left[\left(\theta_{\mathrm{L}}+\frac{\theta_{\mathrm{TL}}}{\ell}-1\right) R_{\alpha}^{\star}+\left((-1)^s \frac{\theta_{\mathrm{TL}}}{\ell}+\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right) T_{\alpha}+\frac{1}{2}\left(\frac{\theta_{\mathrm{T}}}{\ell^2}+m^2\right) \eta_{\alpha}+\tau_{\alpha}^{\star}\right] \wedge \gamma^{\alpha} \Psi \simeq 0$	$i\left[\left(\theta_{\mathrm{L}}+\frac{\theta_{\mathrm{TL}}}{\ell}-1\right) R_{\alpha}^{\star}+\left((-1)^s \frac{\theta_{\mathrm{TL}}}{\ell}+\frac{\theta_{\mathrm{T}}}{\ell^2}-\frac{m}{2}\right) T_{\alpha}+\frac{1}{2}\left(\frac{\theta_{\mathrm{T}}}{\ell^2}+m^2\right) \eta_{\alpha}+\tau_{\alpha}^{\star}\right] \wedge \gamma^{\alpha} \Psi \simeq 0, \quad (10.10)$
mielke-geometrodynamics_EQ0732 (10.10.12)	232	■ 1.000	$\theta_{\mathrm{T}} \simeq-m^2 \ell^2, \theta_{\mathrm{TL}} \simeq \frac{(-1)^s}{2} m(2 m+1) \ell, \theta_{\mathrm{L}} \simeq 1-\frac{\theta_{\mathrm{TL}}}{\ell}=1-\frac{(-1)^s}{2} m(2 m+1)$	$\theta_{\mathrm{T}} \simeq-m^2 \ell^2, \theta_{\mathrm{TL}} \simeq \frac{(-1)^s}{2} m(2 m+1) \ell, \theta_{\mathrm{L}} \simeq 1-\frac{\theta_{\mathrm{TL}}}{\ell}=1-\frac{(-1)^s}{2} m(2 m+1)$	$\theta_{\mathrm{T}} \simeq-m^2 \ell^2, \theta_{\mathrm{TL}} \simeq \frac{(-1)^s}{2} m(2 m+1) \ell, \theta_{\mathrm{L}} \simeq 1-\frac{\theta_{\mathrm{TL}}}{\ell}=1-\frac{(-1)^s}{2} m(2 m+1)$
mielke-geometrodynamics_EQ0733 (10.10.13)	232	■ 1.000	$\vartheta_{\alpha}=(-1)^s \ell \Gamma_{\alpha}^{\star}+\bar{\sigma} \gamma_{\alpha} \Psi.$	$\vartheta_{\alpha}=(-1)^s \ell \Gamma_{\alpha}^{\star}+\bar{\sigma} \gamma_{\alpha} \Psi.$	$\vartheta_{\alpha}=(-1)^s \ell \Gamma_{\alpha}^{\star}+\bar{\sigma} \gamma_{\alpha} \Psi .$
mielke-geometrodynamics_EQ0735 (11.1.2)	238	■ 1.000	$\gamma_{\alpha} \gamma_{\beta}+\gamma_{\beta} \gamma_{\alpha}=2 \mathrm{o}_{\alpha \beta} \mathbb{1}$	$\gamma_{\alpha} \gamma_{\beta}+\gamma_{\beta} \gamma_{\alpha}=2 \mathrm{o}_{\alpha \beta} \mathbb{1}$	$\gamma_{\alpha} \gamma_{\beta}+\gamma_{\beta} \gamma_{\alpha}=2 \mathrm{o}_{\alpha \beta} \mathbb{1}$
mielke-geometrodynamics_EQ0738 (11.1.5)	238	■ 1.000	$X:=\iota x^{\alpha} \gamma_{\alpha} \longleftrightarrow x^{\alpha}=-\frac{l}{N} \mathrm{Tr}\left(\gamma^{\alpha} X\right).$	$X:=\iota x^{\alpha} \gamma_{\alpha} \longleftrightarrow x^{\alpha}=-\frac{l}{N} \mathrm{Tr}\left(\gamma^{\alpha} X\right).$	$X:=\iota x^{\alpha} \gamma_{\alpha} \longleftrightarrow x^{\alpha}=-\frac{l}{N} \mathrm{Tr}\left(\gamma^{\alpha} X\right).$
mielke-geometrodynamics_EQ0739 (11.1.6)	238	■ 1.000	$\det X=-\mathrm{o}_{\alpha \beta} x^{\alpha} x^{\beta},$	$\det X=-\mathrm{o}_{\alpha \beta} x^{\alpha} x^{\beta},$	$\det X=-\mathrm{o}_{\alpha \beta} x^{\alpha} x^{\beta},$
mielke-geometrodynamics_EQ0741 (11.1.8)	238	■ 1.000	$X-X^{\prime}=S X S^{-1}, \quad S \in \widetilde{S O}_{\circ}(s, n-s).$	$X-X^{\prime}=S X S^{-1}, \quad S \in \widetilde{S O}_{\circ}(s, n-s).$	$X-X^{\prime}=S X S^{-1}, \quad S \in \widetilde{S O}_{\circ}(s, n-s).$
mielke-geometrodynamics_EQ0743 (11.1.10)	239	■ 1.000	$\sigma^{\alpha \beta}:=\frac{i}{4}\left[\gamma^{\alpha}, \gamma^{\beta}\right], \quad \Lambda\left(\sigma^{\alpha \beta}\right)=L^{\alpha \beta},$	$\sigma^{\alpha \beta}:=\frac{i}{4}\left[\gamma^{\alpha}, \gamma^{\beta}\right], \quad \Lambda\left(\sigma^{\alpha \beta}\right)=L^{\alpha \beta},$	$\sigma^{\alpha \beta}:=\frac{i}{4}\left[\gamma^{\alpha}, \gamma^{\beta}\right], \quad \Lambda\left(\sigma^{\alpha \beta}\right)=L^{\alpha \beta},$
mielke-geometrodynamics_EQ0744 (11.1.11)	239	■ 1.000	$L_{\mathrm{circ}}^g(M):=P\left(M, S O_{\circ}(s, n-s), \pi, \delta\right)$	$L_{\mathrm{circ}}^g(M):=P\left(M, S O_{\circ}(s, n-s), \pi, \delta\right)$	$L_{\mathrm{circ}}^g(M):=P\left(M, S O_{\circ}(s, n-s), \pi, \delta\right)$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0745 (11.1.12)	239	■ 1.000	$\widetilde{L}(M):=P\left(M,\widetilde{SO}_{\circ}(s,n-2),\tilde{\pi},\tilde{\delta}\right)\ni\tilde{p}$	$\tilde{L}(M):=P\left(M,\widetilde{SO}_{\circ}(s,n-2),\tilde{\pi},\tilde{\delta}\right)\ni\tilde{p}$	$\tilde{L}(M):=P(M,\widetilde{SO}_{\circ}(s,n-2),\tilde{\pi},\tilde{\delta})\ni\tilde{p}$
mielke-geometrodynamics_EQ0746 (11.1.13)	239	■ 1.000	$f(\tilde{p}S)=f(\tilde{p})\Lambda(S),S\in\widetilde{SO}_{\circ}(s,n-2).$	$f(\tilde{p}S)=f(\tilde{p})\Lambda(S),S\in\widetilde{SO}_{\circ}(s,n-2).$	$f(\tilde{p}S)=f(\tilde{p})A(S),S\in\widetilde{SO}_{\circ}(s,n-2).$
mielke-geometrodynamics_EQ0748 (11.1.15)	240	■ 1.000	$\tilde{L}(M^4)=M^4\times SL(2,\mathbb{C})$	$\tilde{L}(M^4)=M^4\times SL(2,\mathbb{C})$	$\tilde{L}(M^4)=M^4\times SL(2,\mathbb{C})$
mielke-geometrodynamics_EQ0749 (11.1.16)	241	■ 1.000	$V^{\bar{s}}:=V\left(M,\mathbb{C}^N,\rho^{\bar{s}}\left(\widetilde{SO}_{\circ}(s,n-s)\right)\subset GL(N,\mathbb{C}),\tilde{L}(M)\right)$	$V^{\bar{s}}:=V\left(M,\mathbb{C}^N,\rho^{\bar{s}}\left(\widetilde{SO}_{\circ}(s,n-s)\right)\subset GL(N,\mathbb{C}),\tilde{L}(M)\right)$	$V^{\bar{s}}:=V(M,\mathbb{C}^N,\rho^{\bar{s}}(\widetilde{SO}_{\circ}(s,n-s))\subset GL(N,\mathbb{C}),\tilde{L}(M))$
mielke-geometrodynamics_EQ0751 (11.1.18)	241	■ 1.000	$D^{\frac{1}{2}}:=D^{\left(-\frac{1}{2},\frac{3}{2}\right)}\oplus D^{\left(\frac{1}{2},\frac{3}{2}\right)}\left(\widetilde{SO}_{\circ}(1,3)\right)$	$D^{\frac{1}{2}}:=D^{\left(-\frac{1}{2},\frac{3}{2}\right)}\oplus D^{\left(\frac{1}{2},\frac{3}{2}\right)}\left(\widetilde{SO}_{\circ}(1,3)\right)$	$D^{\frac{1}{2}}:=D^{\left(-\frac{1}{2},\frac{3}{2}\right)}\oplus D^{\left(\frac{1}{2},\frac{3}{2}\right)}\left(\widetilde{SO}_{\circ}(1,3)\right)$
mielke-geometrodynamics_EQ0754 (11.1.21)	241	■ 1.000	$\psi\rightarrow^{S^{-1}}\psi:=S^{-1}\psi$	$\psi\rightarrow^{S^{-1}}\psi:=S^{-1}\psi$	$\psi\rightarrow^{S^{-1}}\psi:=S^{-1}\psi$
mielke-geometrodynamics_EQ0755 (11.1.22)	242	■ 1.000	$\begin{gathered} S(p)\in\mathscr{G}_{\{S\}}\approx C^{\infty}\left(\tilde{L}(M)\times_{\mathrm{Ad}}\widetilde{SO}_{\circ}(s,n-s)\right)\\ \Lambda\left(S_{(p)}\right)=G(p)\in\mathscr{G}_p \end{gathered}$	$S(p)\in\mathscr{G}_S\approx C^{\infty}\left(\tilde{L}(M)\times_{\mathrm{Ad}}\widetilde{SO}_{\circ}(s,n-s)\right)\\ \Lambda\left(S_{(p)}\right)=G(p)\in\mathscr{G}_p$	$S(p)\in\mathscr{G}_S\approx C^{\infty}(\tilde{L}(M)\times_{\mathrm{Ad}}\widetilde{SO}_{\circ}(s,n-s))\\ \Lambda(S_{(p)})=G(p)\in\mathscr{G}_p$
mielke-geometrodynamics_EQ0757 (11.1.24)	242	■ 1.000	$S^{-1}(p)\gamma_{\alpha}S(p)=\gamma_{\beta}G_{\alpha}^{\beta}(p),\quad[G_{\alpha}^{\beta}(p)]\in\mathscr{G}_p.$	$S^{-1}(p)\gamma_{\alpha}S(p)=\gamma_{\beta}G_{\alpha}^{\beta}(p),\quad[G_{\alpha}^{\beta}(p)]\in\mathscr{G}_p.$	$S^{-1}(p)\gamma_{\alpha}S(p)=\gamma_{\beta}G_{\alpha}^{\beta}(p),\quad[G_{\alpha}^{\beta}(p)]\in\mathscr{G}_p.$
mielke-geometrodynamics_EQ0762 (11.1.29)	243	■ 1.000	$\bar{\psi}:=\psi^{+}\gamma^0,$	$\bar{\psi}:=\psi^{+}\gamma^0,$	$\bar{\psi}:=\psi^{+}\gamma^0,$
mielke-geometrodynamics_EQ0763 (11.1.30)	243	■ 1.000	$\bar{\sigma}^{\alpha\beta}:=\gamma^0\sigma^{\alpha\beta}+\gamma^0=\sigma^{\alpha\beta},\quad\bar{\gamma}^{\alpha}:=\gamma^0\gamma^{\alpha}+\gamma^0=\gamma^{\alpha}.$	$\bar{\sigma}^{\alpha\beta}:=\gamma^0\sigma^{\alpha\beta}+\gamma^0=\sigma^{\alpha\beta},\quad\bar{\gamma}^{\alpha}:=\gamma^0\gamma^{\alpha}+\gamma^0=\gamma^{\alpha}.$	$\bar{\sigma}^{\alpha\beta}:=\gamma^0\sigma^{\alpha\beta}+\gamma^0=\sigma^{\alpha\beta},\quad\bar{\gamma}^{\alpha}:=\gamma^0\gamma^{\alpha}+\gamma^0=\gamma^{\alpha}.$
mielke-geometrodynamics_EQ0764 (11.1.31)	243	■ 1.000	$D\psi=d\psi+i\widetilde{\omega}\psi,$	$D\psi=d\psi+i\widetilde{\omega}\psi,$	$D\psi=d\psi+i\widetilde{\omega}\psi,$
mielke-geometrodynamics_EQ0765 (11.1.32)	243	■ 1.000	$\Lambda(i\widetilde{\omega})=\omega^{\mathfrak{g}}$	$\Lambda(i\widetilde{\omega})=\omega^{\mathfrak{g}}$	$\Lambda(i\widetilde{\omega})=\omega^{\mathfrak{g}}$
mielke-geometrodynamics_EQ0771 (11.1.38)	244	■ 1.000	$\gamma\longrightarrow^{S^{-1}}\gamma=S^{-1}\gamma S$	$\gamma\longrightarrow^{S^{-1}}\gamma=S^{-1}\gamma S$	$\gamma\longrightarrow^{S^{-1}}\gamma=S^{-1}\gamma S$
mielke-geometrodynamics_EQ0775 (11.2.4)	246	■ 1.000	$\overline{\widetilde{\omega}}=\gamma^0\widetilde{\omega}^{+}\gamma^0=\widetilde{\omega}$	$\overline{\widetilde{\omega}}=\gamma^0\widetilde{\omega}^{+}\gamma^0=\widetilde{\omega}$	$\overline{\widetilde{\omega}}=\gamma^0\widetilde{\omega}^{+}\gamma^0=\widetilde{\omega}$
mielke-geometrodynamics_EQ0777 (11.2.6)	246	■ 1.000	$\tau_s=i\left(\bar{\psi}^{*}\gamma\psi\right).$	$\tau_s=i\left(\bar{\psi}^{*}\gamma\psi\right).$	$\tau_s=i(\bar{\psi}^{*}\gamma\psi).$
mielke-geometrodynamics_EQ0779 (11.2.7)	247	■ 1.000	$d\tau_s=0$	$d\tau_s=0$	$d\tau_s=0$
mielke-geometrodynamics_EQ0781 (11.2.9)	247	■ 1.000	$(\psi,\chi)=i\int_{H^{n-1}}\psi^{+}\chi d^{n-1}x$	$(\psi,\chi)=i\int_{H^{n-1}}\psi^{+}\chi d^{n-1}x$	$(\psi,\chi)=i\int_{H^{n-1}}\psi^{+}\chi d^{n-1}x.$
mielke-geometrodynamics_EQ0782 (11.3.1)	247	■ 1.000	$\widetilde{\omega}=\widetilde{\omega}^{\{\}}-\widetilde{K},\quad\Lambda(i\widetilde{K})=K$	$\widetilde{\omega}=\widetilde{\omega}^{\{\}}-\widetilde{K},\quad\Lambda(i\widetilde{K})=K$	$\widetilde{\omega}=\widetilde{\omega}^{\{\}}-\widetilde{K},\quad\Lambda(i\widetilde{K})=K$
mielke-geometrodynamics_EQ0785 (11.3.4)	248	■ 1.000	$[\Theta,\vartheta]=\ell^{*2}\tau_s.$	$[\Theta,\vartheta]=\ell^{*2}\tau_s.$	$[\Theta,\vartheta]=\ell^{*2}\tau_s.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0790 (11.4.2)	252	■ 1.000	$\begin{aligned} & \left(\bar{\psi} i \gamma^5 \gamma_{\mu} \psi\right) \left(\bar{\psi} i \gamma^5 \gamma^{\mu} \psi\right) = \left(\bar{\psi} \gamma_{\mu} \psi\right) \left(\bar{\psi} \gamma^{\mu} \psi\right) \\ & = \left(\bar{\psi} \psi\right)^2 - \left(\bar{\psi} \gamma^5 \psi\right)^2 . \end{aligned}$	$\left(\bar{\psi} i \gamma^5 \gamma_{\mu} \psi\right) \left(\bar{\psi} i \gamma^5 \gamma^{\mu} \psi\right) = \left(\bar{\psi} \gamma_{\mu} \psi\right) \left(\bar{\psi} \gamma^{\mu} \psi\right) = \left(\bar{\psi} \psi\right)^2 - \left(\bar{\psi} \gamma^5 \psi\right)^2 .$	
mielke-geometrodynamics_EQ0793 (11.4.5)	253	■ 1.000	$\rho := \frac{\sqrt{2\pi}}{\ell^*} r,$	$\rho := \frac{\sqrt{2\pi}}{\ell^*} r,$	$\rho := \frac{\sqrt{2\pi}}{\ell^*} r,$
mielke-geometrodynamics_EQ0794 (11.4.6)	253	■ 1.000	$\bar{g}_{kl} \longleftarrow g_{kl} = e^{\mu} \bar{g}_{kl}.$	$\bar{g}_{kl} \longleftarrow g_{kl} = e^{\mu} \bar{g}_{kl}.$	$\bar{g}_{kl} \longleftarrow g_{kl} = e^{\mu} \bar{g}_{kl}.$
mielke-geometrodynamics_EQ0795 (11.4.7)	253	■ 1.000	$\bar{E}_{\cdot k}^{\alpha} \longleftarrow E_{\cdot k}^{\alpha} = e^{\mu/2} \bar{E}_{\cdot k}^{\alpha}$	$\bar{E}_{\cdot k}^{\alpha} \longleftarrow E_{\cdot k}^{\alpha} = e^{\mu/2} \bar{E}_{\cdot k}^{\alpha}$	$\bar{E}_{\cdot k}^{\alpha} \longleftarrow E_{\cdot k}^{\alpha} = e^{\mu/2} \bar{E}_{\cdot k}^{\alpha}$
mielke-geometrodynamics_EQ0796 (11.4.8)	253	■ 1.000	$\overline{\left\{ \begin{array}{c} \alpha \\ kl \end{array} \right\}} = \overline{\left\{ \begin{array}{c} \alpha \\ kl \end{array} \right\}} + \frac{1}{2} \left(\delta_k^{\alpha} \partial_l + \delta_l^{\alpha} \partial_k - g_{kl} \partial^{\alpha} \right) \mu$	$\left\{ \begin{array}{c} \alpha \\ kl \end{array} \right\} = \overline{\left\{ \begin{array}{c} \alpha \\ kl \end{array} \right\}} + \frac{1}{2} \left(\delta_k^{\alpha} \partial_l + \delta_l^{\alpha} \partial_k - g_{kl} \partial^{\alpha} \right) \mu$	$\left\{ \begin{array}{c} \alpha \\ kl \end{array} \right\} = \overline{\left\{ \begin{array}{c} \alpha \\ kl \end{array} \right\}} + \frac{1}{2} \left(\delta_k^{\alpha} \partial_l + \delta_l^{\alpha} \partial_k - g_{kl} \partial^{\alpha} \right) \mu$
mielke-geometrodynamics_EQ0797 (11.4.9)	253	■ 1.000	$\Gamma_k = \bar{\Gamma}_k - \frac{1}{4} \delta_k^{[\alpha} \partial^{\beta]} \mu \sigma_{\alpha\beta}.$	$\Gamma_k = \bar{\Gamma}_k - \frac{1}{4} \delta_k^{[\alpha} \partial^{\beta]} \mu \sigma_{\alpha\beta}.$	$\Gamma_k = \bar{\Gamma}_k - \frac{1}{4} \delta_k^{[\alpha} \partial^{\beta]} \mu \sigma_{\alpha\beta}.$
mielke-geometrodynamics_EQ0801 (11.4.13)	254	■ 1.000	$\begin{aligned} & \bar{\Gamma}_0 = \frac{i}{4} e^{(\nu-\mu)/2} \partial_r (\nu - \mu) \gamma_0 \gamma_1 \\ & \bar{\Gamma}_1 = 0, \quad \bar{\Gamma}_2 = \frac{i}{2} \gamma_2 \gamma_1 \\ & \bar{\Gamma}_3 = \frac{i}{2} (\sin \vartheta \gamma_3 \gamma_1 + \cos \vartheta \gamma_3 \gamma_2) \end{aligned}$	$\begin{aligned} \bar{\Gamma}_0 &= \frac{i}{4} e^{(\nu-\mu)/2} \partial_r (\nu - \mu) \gamma_0 \gamma_1 \\ \bar{\Gamma}_1 &= 0, \quad \bar{\Gamma}_2 = \frac{i}{2} \gamma_2 \gamma_1 \\ \bar{\Gamma}_3 &= \frac{i}{2} (\sin \vartheta \gamma_3 \gamma_1 + \cos \vartheta \gamma_3 \gamma_2) \end{aligned}$	$\begin{aligned} \bar{T}_0 &= \frac{i}{4} e^{(v-\mu)/2} \partial_r (v - \mu) \gamma_0 \gamma_1 \\ \bar{T}_1 &= 0, \quad \bar{T}_2 = \frac{i}{2} \gamma_2 \gamma_1 \\ \bar{T}_3 &= \frac{i}{2} (\sin \vartheta \gamma_3 \gamma_1 + \cos \vartheta \gamma_3 \gamma_2) \end{aligned}$
mielke-geometrodynamics_EQ0804 (11.4.16)	255	■ 1.000	$\chi_{\kappa}^m = \sum_{\bar{m}=\pm 1/2} C \left(l \frac{1}{2} j; m - \bar{m}, \bar{m} \right) Y_l^{m-\bar{m}}(\vartheta, \phi) \chi^{\bar{m}}.$	$\chi_{\kappa}^m = \sum_{\bar{m}=\pm 1/2} C \left(l \frac{1}{2} j; m - \bar{m}, \bar{m} \right) Y_l^{m-\bar{m}}(\vartheta, \phi) \chi^{\bar{m}}.$	$\chi_{\kappa}^m = \sum_{\bar{m}=\pm 1/2} C \left(l \frac{1}{2} j; m - \bar{m}, \bar{m} \right) Y_l^{m-\bar{m}}(\vartheta, \phi) \chi^{\bar{m}}.$
mielke-geometrodynamics_EQ0808 (11.4.20)	255	■ 1.000	$\frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } \chi_{\kappa}^m = -\chi_{-\kappa}^m$	$\frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } \chi_{\kappa}^m = -\chi_{-\kappa}^m$	$\frac{\vec{\sigma} \cdot \vec{x}}{ \vec{x} } \chi_{\kappa}^m = -\chi_{-\kappa}^m$
mielke-geometrodynamics_EQ0814 (11.4.27)	257	■ 1.000	$\beta = \frac{M^*}{2m}, \quad M^* := \frac{\sqrt{8\pi} \hbar}{c \ell^*}.$	$\beta = \frac{M^*}{2m}, \quad M^* := \frac{\sqrt{8\pi} \hbar}{c \ell^*}.$	$\beta = \frac{M^*}{2m}, \quad M^* := \frac{\sqrt{8\pi} \hbar}{c \ell^*}.$
mielke-geometrodynamics_EQ0816 (11.4.30)	257	■ 1.000	$F'' + \frac{2}{\rho} F' - \frac{2}{\rho^2} F \simeq \frac{1-\omega^2}{\beta^2} F$	$F'' + \frac{2}{\rho} F' - \frac{2}{\rho^2} F \simeq \frac{1-\omega^2}{\beta^2} F$	$F'' + \frac{2}{\rho} F' - \frac{2}{\rho^2} F \simeq \frac{1-\omega^2}{\beta^2} F$
mielke-geometrodynamics_EQ0817 (11.4.31)	257	■ 1.000	$F \sim C_{\infty} \left(\frac{1}{\bar{\rho}} + \frac{1}{\bar{\rho}^2} \right) e^{-\bar{\rho}}$	$F \sim C_{\infty} \left(\frac{1}{\bar{\rho}} + \frac{1}{\bar{\rho}^2} \right) e^{-\bar{\rho}}$	$F \sim C_{\infty} \left(\frac{1}{\bar{\rho}} + \frac{1}{\bar{\rho}^2} \right) e^{-\bar{\rho}}$
mielke-geometrodynamics_EQ0818 (11.4.32)	257	■ 1.000	$H \sim C_{\infty} \sqrt{\frac{1+\omega}{1-\omega}} \frac{1}{\bar{\rho}} e^{-\bar{\rho}}$	$H \sim C_{\infty} \sqrt{\frac{1+\omega}{1-\omega}} \frac{1}{\bar{\rho}} e^{-\bar{\rho}}$	$H \sim C_{\infty} \sqrt{\frac{1+\omega}{1-\omega}} \frac{1}{\bar{\rho}} e^{-\bar{\rho}}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometroynamics_E08819 (11.4.33)	257	■ 1.000	$\bar{\rho}:=\frac{1}{\beta}\sqrt{1-\omega^2}\rho$	$\bar{\rho}:=\frac{1}{\beta}\sqrt{1-\omega^2}\rho$	$\bar{\rho}:=\frac{1}{\beta}\sqrt{1-\omega^2}\rho$
mielke-geometroynamics_E08821 (11.4.35)	258	■ 1.000	$F(\rho)\simeq C_0f^1(\bar{\rho})\frac{1}{\cosh\bar{\rho}}$	$F(\rho)\simeq C_0f^1(\bar{\rho})\frac{1}{\cosh\bar{\rho}}$	$F(\rho)\simeq C_0f^1(\bar{\rho})\frac{1}{\cosh\bar{\rho}}$
mielke-geometroynamics_E08822 (11.4.36)	258	■ 1.000	$H(\rho)\simeq C_0\sqrt{\frac{1+\omega}{1-\omega}}f^0(\bar{\rho})\frac{1}{\cosh\bar{\rho}}$	$H(\rho)\simeq C_0\sqrt{\frac{1+\omega}{1-\omega}}f^0(\bar{\rho})\frac{1}{\cosh\bar{\rho}}$	$H(\rho)\simeq C_0\sqrt{\frac{1+\omega}{1-\omega}}f^0(\bar{\rho})\frac{1}{\cosh\bar{\rho}}$
mielke-geometroynamics_E08823 (11.4.37)	258	■ 1.000	$C_0=(1-\omega)\sqrt{\frac{-2}{\varepsilon(\omega+1)}}$	$C_0=(1-\omega)\sqrt{\frac{-2}{\varepsilon(\omega+1)}}$	$C_0=(1-\omega)\sqrt{\frac{-2}{\varepsilon(\omega+1)}}$
mielke-geometroynamics_E08826 (11.4.40)	259	■ 1.000	$m_s=\frac{1}{c}\int_{H^3}\Sigma_{\mathrm{D}}^0$	$m_s=\frac{1}{c}\int_{H^3}\Sigma_{\mathrm{D}}^0$	$m_s=\frac{1}{c}\int_{H^3}\Sigma_{\mathrm{D}}^0$
mielke-geometroynamics_E08827 (11.5.1)	262	■ 1.000	$Q:=ie\int_{H^3}\bar{\psi}^*\gamma\psi=ke,\quad k=0,1,2,\dots$	$Q:=ie\int_{H^3}\bar{\psi}^*\gamma\psi=ke,\quad k=0,1,2,\dots$	$Q:=ie\int_{H^3}\bar{\psi}^*\gamma\psi=ke,\quad k=0,1,2,\dots$
mielke-geometroynamics_E08828 (11.5.2)	263	■ 1.000	$i\gamma\wedge^*d\psi(x)+e^2\int_{y\in M}\bar{\psi}(y)\gamma\psi(y)D_{\mathrm{F}}(y-x)\wedge^*\gamma\psi(x)=m\psi(x)\eta$	$i\gamma\wedge^*d\psi(x)+e^2\int_{y\in M}\bar{\psi}(y)\gamma\psi(y)D_{\mathrm{F}}(y-x)\wedge^*\gamma\psi(x)=m\psi(x)\eta$	$i\gamma\wedge^*d\psi(x)+e^2\int_{y\in M}\bar{\psi}(y)\gamma\psi(y)D_{\mathrm{F}}(y-x)\wedge^*\gamma\psi(x)=m\psi(x)\eta$
mielke-geometroynamics_E08830 (12.1.2)	269	■ 1.000	$\langle dj_5(x)\rangle=2im\langle P\rangle-(1/96\pi^2)F\wedge F$	$\langle dj_5(x)\rangle=2im\langle P\rangle-(1/96\pi^2)F\wedge F$	$\langle dj_5(x)\rangle=2im\langle P\rangle-(1/96\pi^2)F\wedge F$
mielke-geometroynamics_E08838 (12.2.6)	271	■ 1.000	$dj_5=2imP=2im\bar{\psi}\gamma_5\psi\eta$	$dj_5=2imP=2im\bar{\psi}\gamma_5\psi\eta$	$dj_5=2imP=2im\bar{\psi}\gamma_5\psi\eta$
mielke-geometroynamics_E08843 (12.2.11)	272	■ 1.000	$L_{a\psi\psi}=\frac{1}{2}d\theta\wedge j_5=\frac{1}{2f_a}da\wedge\bar{\psi}^*\gamma\gamma_5\psi,$	$L_{a\psi\psi}=\frac{1}{2}d\theta\wedge j_5=\frac{1}{2f_a}da\wedge\bar{\psi}^*\gamma\gamma_5\psi,$	$L_{a\psi\psi}=\frac{1}{2}d\theta\wedge j_5=\frac{1}{2f_a}da\wedge\bar{\psi}^*\gamma\gamma_5\psi,$
mielke-geometroynamics_E08846 (12.2.14)	273	■ 1.000	$dj_5\cong 4dC_{\mathrm{TT}}=\frac{2}{\ell^2}(T^\alpha\wedge T_\alpha+R_{\alpha\beta}\wedge\vartheta^\alpha\wedge\vartheta^\beta)$	$dj_5\cong 4dC_{\mathrm{TT}}=\frac{2}{\ell^2}(T^\alpha\wedge T_\alpha+R_{\alpha\beta}\wedge\vartheta^\alpha\wedge\vartheta^\beta)$	$dj_5\cong 4dC_{\mathrm{TT}}=\frac{2}{\ell^2}(T^\alpha\wedge T_\alpha+R_{\alpha\beta}\wedge\vartheta^\alpha\wedge\vartheta^\beta)$
mielke-geometroynamics_E08849 (12.3.2)	275	■ 1.000	$K(t,x,\mathcal{D}^2)=(4\pi)^{-n/2}\sum_{k=0}^\infty t^{(k-n)/2}K_k(x,\mathcal{D}^2)$	$K(t,x,\mathcal{D}^2)=(4\pi)^{-n/2}\sum_{k=0}^\infty t^{(k-n)/2}K_k(x,\mathcal{D}^2)$	$K(t,x,\mathcal{D}^2)=(4\pi)^{-n/2}\sum_{k=0}^\infty t^{(k-n)/2}K_k(x,\mathcal{D}^2)$
mielke-geometroynamics_E08852 (12.3.5)	276	■ 1.000	$-K_2/t=(2\ell^2/t)dC_{\mathrm{TT}}\cong(\ell^2/2t)dj_5=(im\ell^2/t)\bar{\psi}\gamma_5\psi\sim\ell^2M^2m^4\rightarrow 0.$	$-K_2/t=(2\ell^2/t)dC_{\mathrm{TT}}\cong(\ell^2/2t)dj_5=(im\ell^2/t)\bar{\psi}\gamma_5\psi\sim\ell^2M^2m^4\rightarrow 0.$	$-K_2/t=(2\ell^2/t)dC_{\mathrm{TT}}\cong(\ell^2/2t)dj_5=(im\ell^2/t)\bar{\psi}\gamma_5\psi\sim\ell^2M^2m^4\rightarrow 0.$
mielke-geometroynamics_E08856 (12.4.3)	277	■ 1.000	$\mathscr{G}^A\cong 0,\quad A=1,2,3$	$\mathscr{G}^A\cong 0,\quad A=1,2,3$	$\mathscr{G}^A\cong 0,\quad A=1,2,3$
mielke-geometroynamics_E08858 (12.4.5)	278	■ 1.000	$\gamma:=\gamma_\alpha\vartheta^\alpha,\quad{}^*\gamma=\gamma^\alpha\eta_\alpha.$	$\gamma:=\gamma_\alpha\vartheta^\alpha,\quad{}^*\gamma=\gamma^\alpha\eta_\alpha.$	$\gamma:=\gamma_\alpha\vartheta^\alpha,\quad{}^*\gamma=\gamma^\alpha\eta_\alpha.$
mielke-geometroynamics_E08859 (12.4.6)	278	■ 1.000	$\Theta:=D\gamma=T^\alpha\gamma_\alpha,\quad\Omega:=d\Gamma+\Gamma\wedge\Gamma=\frac{i}{4}R^{\alpha\beta}\sigma_{\alpha\beta}$	$\Theta:=D\gamma=T^\alpha\gamma_\alpha,\quad\Omega:=d\Gamma+\Gamma\wedge\Gamma=\frac{i}{4}R^{\alpha\beta}\sigma_{\alpha\beta}$	$\Theta:=D\gamma=T^\alpha\gamma_\alpha,\quad\Omega:=d\Gamma+\Gamma\wedge\Gamma=\frac{i}{4}R^{\alpha\beta}\sigma_{\alpha\beta}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_278 EQ0860 (12.4.7)	278	■ 1.000	$C_{\mathrm{RR}} := -\operatorname{Tr}(\Gamma \wedge \Omega - \frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma)$.	$C_{\mathrm{RR}} := -\operatorname{Tr} \left(\Gamma \wedge \Omega - \frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma \right).$	$C_{\mathrm{RR}} := -\operatorname{Tr} \left(\Gamma \wedge \Omega - \frac{1}{3} \Gamma \wedge \Gamma \wedge \Gamma \right).$
mielke-geometrodynamics_279 EQ0862 (12.4.9)	279	■ 1.000	$C_{\mathrm{TT}} := \frac{1}{8\ell^2} \operatorname{Tr}(\gamma \wedge \Theta) = \frac{1}{2\ell^2} \vartheta^\alpha \wedge T_\alpha.$	$C_{\mathrm{TT}} := \frac{1}{8\ell^2} \operatorname{Tr}(\gamma \wedge \Theta) = \frac{1}{2\ell^2} \vartheta^\alpha \wedge T_\alpha.$	$C_{\mathrm{TT}} := \frac{1}{8\ell^2} \operatorname{Tr}(\gamma \wedge \Theta) = \frac{1}{2\ell^2} \vartheta^\alpha \wedge T_\alpha.$
mielke-geometrodynamics_279 EQ0863 (12.4.10)	279	■ 1.000	$dC_{\mathrm{TT}} = \frac{1}{8\ell^2} \operatorname{Tr}(\Theta \wedge \Theta - 4i\Omega \wedge \sigma) = \frac{1}{2\ell^2} (T^\alpha \wedge T_\alpha + R_{\alpha\beta} \wedge \vartheta^\alpha \wedge \vartheta^\beta).$	$dC_{\mathrm{TT}} = \frac{1}{8\ell^2} \operatorname{Tr}(\Theta \wedge \Theta - 4i\Omega \wedge \sigma) = \frac{1}{2\ell^2} (T^\alpha \wedge T_\alpha + R_{\alpha\beta} \wedge \vartheta^\alpha \wedge \vartheta^\beta).$	$dC_{\mathrm{TT}} = \frac{1}{8\ell^2} \operatorname{Tr}(\Theta \wedge \Theta - 4i\Omega \wedge \sigma) = \frac{1}{2\ell^2} (T^\alpha \wedge T_\alpha + R_{\alpha\beta} \wedge \vartheta^\alpha \wedge \vartheta^\beta).$
mielke-geometrodynamics_279 EQ0864 (12.4.11)	279	■ 1.000	$\hat{C}_{\mathrm{RR}} = C_{\mathrm{RR}} - 2C_{\mathrm{TT}}$	$\hat{C}_{\mathrm{RR}} = C_{\mathrm{RR}} - 2C_{\mathrm{TT}}$	$\hat{C}_{\mathrm{RR}} = C_{\mathrm{RR}} - 2C_{\mathrm{TT}}$
mielke-geometrodynamics_283 EQ0865 (13.2.1)	283	■ 1.000	$L_{\mathrm{BF}} = -B \wedge F = -B \wedge dA.$	$L_{\mathrm{BF}} = -B \wedge F = -B \wedge dA.$	$L_{\mathrm{BF}} = -B \wedge F = -B \wedge dA.$
mielke-geometrodynamics_283 EQ0866 (13.2.2)	283	■ 1.000	$\tilde{L}_{\mathrm{BF}} := -B \wedge dA + \frac{1}{2} B \wedge B \cong -B \wedge dA + dC$	$\tilde{L}_{\mathrm{BF}} := -B \wedge dA + \frac{1}{2} B \wedge B \cong -B \wedge dA + dC$	$\tilde{L}_{\mathrm{BF}} := -B \wedge dA + \frac{1}{2} B \wedge B \cong -B \wedge dA + dC.$
mielke-geometrodynamics_283 EQ0868 (13.2.4)	283	■ 1.000	$C = \frac{1}{2} A \wedge F, \quad dC = \frac{1}{2} F \wedge F,$	$C = \frac{1}{2} A \wedge F, \quad dC = \frac{1}{2} F \wedge F,$	$C = \frac{1}{2} A \wedge F, \quad dC = \frac{1}{2} F \wedge F,$
mielke-geometrodynamics_283 EQ0869 (13.2.5)	283	■ 1.000	$A \rightarrow A' = A + d\theta(x),$	$A \rightarrow A' = A + d\theta(x),$	$A \rightarrow A' = A + d\theta(x),$
mielke-geometrodynamics_284 EQ0870 (13.2.6)	284	■ 1.000	$A \rightarrow \tilde{A} = A + \sigma, \quad B \rightarrow \tilde{B} = B + d\sigma,$	$A \rightarrow \tilde{A} = A + \sigma, \quad B \rightarrow \tilde{B} = B + d\sigma,$	$A \rightarrow \tilde{A} = A + \sigma, \quad B \rightarrow \tilde{B} = B + d\sigma,$
mielke-geometrodynamics_284 EQ0871 (13.2.7)	284	■ 1.000	$L_{\mathrm{Max}} = -B \wedge dA + \frac{1}{2} B \wedge {}^*B + L_{\mathrm{matter}},$	$L_{\mathrm{Max}} = -B \wedge dA + \frac{1}{2} B \wedge {}^*B + L_{\mathrm{matter}},$	$L_{\mathrm{Max}} = -B \wedge dA + \frac{1}{2} B \wedge {}^*B + L_{\mathrm{matter}},$
mielke-geometrodynamics_284 EQ0872 (13.2.8)	284	■ 1.000	$-dB = d{}^*F \cong j,$	$-dB = d{}^*F \cong j,$	$-dB = d{}^*F \cong j,$
mielke-geometrodynamics_284 EQ0875 (13.3.3)	284	■ 1.000	$[L^\alpha{}_\beta, L^\gamma{}_\delta] = \delta^\alpha_\delta L^\gamma{}_\beta - \delta^\gamma_\beta L^\alpha{}_\delta,$	$[L^\alpha{}_\beta, L^\gamma{}_\delta] = \delta^\alpha_\delta L^\gamma{}_\beta - \delta^\gamma_\beta L^\alpha{}_\delta,$	$[L^\alpha{}_\beta, L^\gamma{}_\delta] = \delta^\alpha_\delta L^\gamma{}_\beta - \delta^\gamma_\beta L^\alpha{}_\delta,$
mielke-geometrodynamics_285 EQ0880 (13.4.1)	285	■ 1.000	$A(x) := \left(\begin{array}{cc} \Lambda(x) & \tau(x) \\ \tau_*(x) & 1 \end{array} \right),$	$A(x) := \left\{ \begin{array}{cc} \Lambda(x) & \tau(x) \\ \tau_*(x) & 1 \end{array} \right\},$	$A(x) := \left\{ \begin{array}{cc} \Lambda(x) & \tau(x) \\ \tau_*(x) & 1 \end{array} \right\},$
mielke-geometrodynamics_285 EQ0881 (13.4.2)	285	■ 1.000	$\hat{\Gamma} = \left(\begin{array}{cc} \Gamma^{(L)} & \Gamma^{(T)} \\ \Gamma_*^{(T)} & 0 \end{array} \right) = \left(\begin{array}{cc} \Gamma_\alpha^{(L)\beta} L_\beta^\alpha & \Gamma^{(T)\alpha} \ell P_\alpha \\ \Gamma_\beta^{*(T)} & \ell P_*^\beta \end{array} \right)$	$\hat{\Gamma} = \left(\begin{array}{cc} \Gamma^{(L)} & \Gamma^{(T)} \\ \Gamma_*^{(T)} & 0 \end{array} \right) = \left(\begin{array}{cc} \Gamma_\alpha^{(L)\beta} L_\beta^\alpha & \Gamma^{(T)\alpha} \ell P_\alpha \\ \Gamma_\beta^{*(T)} & \ell P_*^\beta \end{array} \right)$	$\hat{\Gamma} = \left(\begin{array}{cc} \Gamma^{(L)} & \Gamma^{(T)} \\ \Gamma_*^{(T)} & 0 \end{array} \right) = \left(\begin{array}{cc} \Gamma_\alpha^{(L)\beta} L_\beta^\alpha & \Gamma^{(T)\alpha} \ell P_\alpha \\ \Gamma_\beta^{*(T)} & \ell P_*^\beta \end{array} \right)$
mielke-geometrodynamics_286 EQ0882 (13.4.3)	286	■ 1.000	$\hat{\Gamma} \xrightarrow{A^{-1}(x)} \hat{\Gamma}' = A^{-1}(x) \hat{\Gamma} A(x) + A^{-1}(x) dA(x).$	$\hat{\Gamma} \xrightarrow{A^{-1}(x)} \hat{\Gamma}' = A^{-1}(x) \hat{\Gamma} A(x) + A^{-1}(x) dA(x).$	$\hat{\Gamma} \xrightarrow{A^{-1}(x)} \hat{\Gamma}' = A^{-1}(x) \hat{\Gamma} A(x) + A^{-1}(x) dA(x).$
mielke-geometrodynamics_286 EQ0883	286	■ 1.000	$\tau \tau_* = -\Lambda \tau_* \Lambda^{-1} \tau.$	$\tau \tau_* = -\Lambda \tau_* \Lambda^{-1} \tau.$	$\tau \tau_* = -\Lambda \tau_* \Lambda^{-1} \tau.$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometroynamics_286 EQ0884 (13.4.4)		■ 1.000	$\hat{\mathrm{R}}:=\mathrm{d}\,\hat{\Gamma}+\hat{\Gamma}\wedge\hat{\Gamma}=\left(\begin{array}{cc}\mathrm{R}^{\left(L\right)}+\Gamma^{\left(L\right)}\wedge\Gamma_{*}^{\left(L\right)} & d\Gamma^{\left(T\right)}+\Gamma^{\left(L\right)}\wedge\Gamma^{\left(T\right)}\\d\Gamma_{*}^{\left(T\right)}+\Gamma_{*}^{\left(T\right)}\wedge\Gamma^{\left(L\right)} & 0\end{array}\right)$	$\hat{R} := d\hat{\Gamma} + \hat{\Gamma} \wedge \hat{\Gamma} = \left(\begin{array}{cc} R^{(L)} + \Gamma^{(L)} \wedge \Gamma_*^{(L)} & d\Gamma^{(T)} + \Gamma^{(L)} \wedge \Gamma^{(T)} \\ d\Gamma_*^{(T)} + \Gamma_*^{(T)} \wedge \Gamma^{(L)} & 0 \end{array} \right)$	$\hat{R} := d\hat{\Gamma} + \hat{\Gamma} \wedge \hat{\Gamma} = \left(\begin{array}{cc} R^{(L)} + \Gamma^{(L)} \wedge \Gamma_*^{(L)} & d\Gamma^{(T)} + \Gamma^{(L)} \wedge \Gamma^{(T)} \\ d\Gamma_*^{(T)} + \Gamma_*^{(T)} \wedge \Gamma^{(L)} & 0 \end{array} \right),$
mielke-geometroynamics_286 EQ0885		■ 1.000	$\mathrm{R}^{\left(L\right)}:=\mathrm{d}\,\Gamma^{\left(L\right)}+\Gamma^{\left(L\right)}\wedge\Gamma^{\left(L\right)}$	$R^{(L)} := d\Gamma^{(L)} + \Gamma^{(L)} \wedge \Gamma^{(L)}$	$R^{(L)} := d\Gamma^{(L)} + \Gamma^{(L)} \wedge \Gamma^{(L)}$
mielke-geometroynamics_286 EQ0886 (13.4.5)		■ 1.000	$\mathrm{R}^{\left(T\right)}:=\mathrm{d}\,\Gamma^{\left(T\right)}+\Gamma^{\left(L\right)}\wedge\Gamma^{\left(T\right)}=\left(\mathrm{T}^{\left(\beta\right)}-\mathrm{R}_{\alpha}^{\beta}\xi^{\alpha}\right)\ell\mathrm{P}_{\beta}$	$R^{(T)} := d\Gamma^{(T)} + \Gamma^{(L)} \wedge \Gamma^{(T)} = \left(T^{\beta} - R_{\alpha}^{\beta}\xi^{\alpha}\right)\ell P_{\beta}$	$R^{(T)} := d\Gamma^{(T)} + \Gamma^{(L)} \wedge \Gamma^{(T)} = \left(T^{\beta} - R_{\alpha}^{\beta}\xi^{\alpha}\right)\ell P_{\beta}$
mielke-geometroynamics_286 EQ0887 (13.4.6)		■ 1.000	$\hat{\mathrm{R}}\,\mathrm{xrightarrow{A^{-1}(x)}}\,\hat{\mathrm{R}}^{\prime}=\mathrm{A}^{-1}(x)\hat{\mathrm{R}}\mathrm{A}(x).$	$\hat{R} \xrightarrow{A^{-1}(x)} \hat{R}' = A^{-1}(x)\hat{R}A(x).$	$\hat{R} \xrightarrow{A^{-1}(x)} \hat{R}' = A^{-1}(x)\hat{R}A(x).$
mielke-geometroynamics_286 EQ0888 (13.4.7)		■ 1.000	$\Gamma^{\left(L\right)}\,\mathrm{xrightarrow{A^{-1}(x)}}\,\Gamma^{\left(L\right)\prime}=\Lambda^{-1}(x)\Gamma^{\left(L\right)}\Lambda(x)+\Lambda^{-1}(x)d\Lambda(x)$	$\Gamma^{(L)} \xrightarrow{A^{-1}(x)} \Gamma^{(L)'} = \Lambda^{-1}(x)\Gamma^{(L)}\Lambda(x) + \Lambda^{-1}(x)d\Lambda(x)$	$\Gamma^{(L)} \xrightarrow{A^{-1}(x)} \Gamma^{(L)'} = \Lambda^{-1}(x)\Gamma^{(L)}\Lambda(x) + \Lambda^{-1}(x)d\Lambda(x)$
mielke-geometroynamics_287 EQ0891 (13.4.10)		■ 1.000	$\vartheta:=\Gamma^{\left(T\right)}+D\xi:=\vartheta^{\alpha}P_{\alpha},\quad\theta:=\Gamma_{*}^{\left(T\right)}+D\xi_{*}:=\theta_{\beta}P_{*}^{\beta},$	$\vartheta := \Gamma^{(T)} + D\xi := \vartheta^{\alpha}P_{\alpha}, \quad \theta := \Gamma_*^{(T)} + D\xi_* := \theta_{\beta}P_{*}^{\beta},$	$\vartheta := \Gamma^{(T)} + D\xi := \vartheta^{\alpha}P_{\alpha}, \quad \theta := \Gamma_*^{(T)} + D\xi_* := \theta_{\beta}P_{*}^{\beta},$
mielke-geometroynamics_287 EQ0894 (13.4.13)		■ 1.000	$\mathrm{D}\,\mathrm{x}\stackrel{*}{\mathrm{rel}}\mathrm{=}\,0,\quad\mathrm{D}\,\mathrm{x}_*\stackrel{*}{\mathrm{rel}}\mathrm{=}\,0,$	$D\xi \stackrel{*}{=} 0, \quad D\xi_* \stackrel{*}{=} 0,$	$D\xi \stackrel{*}{=} 0, \quad D\xi_* \stackrel{*}{=} 0,$
mielke-geometroynamics_287 EQ0895 (13.4.14)		■ 1.000	$\overline{\Gamma}=\left(\begin{array}{cc}\Gamma^{\left(L\right)} & \vartheta\\ \theta & 0\end{array}\right),$	$\bar{\Gamma} = \left(\begin{array}{cc} \Gamma^{(L)} & \vartheta \\ \theta & 0 \end{array} \right),$	$\bar{\Gamma} = \left(\begin{array}{cc} \Gamma^{(L)} & \vartheta \\ \theta & 0 \end{array} \right),$
mielke-geometroynamics_288 EQ0896 (13.5.1)		■ 1.000	$\hat{\Gamma}=\Gamma_{\alpha}^{\beta}L^{\alpha}_{\beta}+\vartheta^{\alpha}P_{\alpha}+\theta_{\beta}P_{*}^{\beta},$	$\hat{\Gamma} = \Gamma_{\alpha}^{\beta}L^{\alpha}_{\beta} + \vartheta^{\alpha}P_{\alpha} + \theta_{\beta}P_{*}^{\beta},$	$\hat{\Gamma} = \Gamma_{\alpha}^{\beta}L^{\alpha}_{\beta} + \vartheta^{\alpha}P_{\alpha} + \theta_{\beta}P_{*}^{\beta},$
mielke-geometroynamics_288 EQ0899 (13.5.4)		■ 1.000	$\tilde{L}_{\mathrm{SL}(5,\mathrm{R})}=-\mathrm{Tr}\{\hat{B}\wedge\hat{R}\}-d\hat{C}.$	$\tilde{L}_{\mathrm{SL}(5,\mathbb{R})} = -\mathrm{Tr}\{\hat{B} \wedge \hat{R}\} - d\hat{C}.$	$\tilde{L}_{\mathrm{SL}(5,\mathbb{R})} = -\mathrm{Tr}\{\hat{B} \wedge \hat{R}\} - d\hat{C}.$
mielke-geometroynamics_289 EQ0900 (13.5.5)		■ 1.000	$\hat{\mathrm{D}}\,\mathrm{B}_{\mathrm{A}}^{\mathrm{B}}\cong\hat{\mathrm{D}}\,\mathrm{B}_{\mathrm{A}}^{\mathrm{B}}\equiv 0.$	$\hat{D}B_A^B \cong \hat{D}\hat{R}_A^B \equiv 0.$	$\hat{D}B_A^B \cong \hat{D}\hat{R}_A^B \equiv 0.$
mielke-geometroynamics_289 EQ9901 (13.5.6)		■ 1.000	$\tilde{L}_{\mathrm{SSB}}=\tilde{L}_{\mathrm{SL}(5,\mathrm{R})}+\frac{1}{2}\eta_{\mathrm{A}\,\mathrm{B}\,\mathrm{C}\,\mathrm{D}\,\mathrm{E}}\,\mathrm{B}^{\mathrm{A}\,\mathrm{B}}\wedge\mathrm{B}^{\mathrm{C}\,\mathrm{D}}\Phi^{\mathrm{E}}.$	$\tilde{L}_{\mathrm{SSB}} = \tilde{L}_{\mathrm{SL}(5,\mathbb{R})} + \frac{1}{2}\eta_{ABCDE}B^{AB} \wedge B^{CD}\Phi^E.$	$\tilde{L}_{\mathrm{SSB}} = \tilde{L}_{\mathrm{SL}(5,\mathbb{R})} + \frac{1}{2}\eta_{ABCDE}B^{AB} \wedge B^{CD}\Phi^E.$
mielke-geometroynamics_289 EQ9902 (13.5.7)		■ 1.000	$\hat{g}_{\mathrm{A}\,\mathrm{B}}\Phi^{\mathrm{A}}\Phi^{\mathrm{B}}=\mu^2,\quad\Phi_{\mathrm{E}}\hat{\mathrm{D}}\Phi^{\mathrm{E}}=0$	$\hat{g}_{AB}\Phi^A\Phi^B = \mu^2, \quad \Phi_E\hat{D}\Phi^E = 0$	$\hat{g}_{AB}\Phi^A\Phi^B = \mu^2, \quad \Phi_E\hat{D}\Phi^E = 0,$
mielke-geometroynamics_289 EQ9903 (13.5.8)		■ 1.000	$\left\langle\Phi^E\right\rangle=\Phi_0^E=(0,0,0,0,\mu)^T$	$\langle\Phi^E\rangle = \Phi_0^E = (0,0,0,0,\mu)^T$	$\langle\Phi^E\rangle = \Phi_0^E = (0,0,0,0,\mu)^T$
mielke-geometroynamics_289 EQ9904 (13.5.9)		■ 1.000	$\tilde{L}_{\mathrm{SSB}}=-\mathrm{Tr}\{\hat{B}\wedge\hat{R}\}+\frac{\mu}{2}\eta_{\alpha\beta\gamma\delta}B^{\alpha\beta}\wedge B^{\gamma\delta}-dC_{\mathrm{RR}}+2dC_{\mathrm{TT}}$	$\tilde{L}_{\mathrm{SSB}} = -\mathrm{Tr}\{\hat{B} \wedge \hat{R}\} + \frac{\mu}{2}\eta_{\alpha\beta\gamma\delta}B^{\alpha\beta} \wedge B^{\gamma\delta} - dC_{\mathrm{RR}} + 2dC_{\mathrm{TT}}$	$\tilde{L}_{\mathrm{SSB}} = -\mathrm{Tr}\{\hat{B} \wedge \hat{R}\} + \frac{\mu}{2}\eta_{\alpha\beta\gamma\delta}B^{\alpha\beta} \wedge B^{\gamma\delta} - dC_{\mathrm{RR}} + 2dC_{\mathrm{TT}}$
mielke-geometroynamics_290 EQ9905 (13.5.10)		■ 1.000	$\mathrm{d}\,\mathrm{C}_{\mathrm{C}}=\mathrm{d}\,\mathrm{C}_{\mathrm{RR}}-2\,\mathrm{d}\,\mathrm{C}_{\mathrm{TT}},$	$d\hat{C} = dC_{\mathrm{RR}} - 2dC_{\mathrm{TT}},$	$d\hat{C} = dC_{\mathrm{RR}} - 2dC_{\mathrm{TT}},$
mielke-geometroynamics_290 EQ9906 (13.5.11)		■ 1.000	$\mathrm{B}_{\alpha\beta}\cong\frac{1}{\mu}\eta_{\alpha\beta\gamma\delta}\left[R^{\gamma\delta}-\frac{\Lambda}{3}\theta^{\gamma}\wedge\vartheta^{\delta}\right]$	$B_{\alpha\beta} \cong \frac{1}{\mu}\eta_{\alpha\beta\gamma\delta} \left[R^{\gamma\delta} - \frac{\Lambda}{3}\theta^{\gamma} \wedge \vartheta^{\delta} \right]$	$B_{\alpha\beta} \cong \frac{1}{\mu}\eta_{\alpha\beta\gamma\delta} \left[R^{\gamma\delta} - \frac{\Lambda}{3}\theta^{\gamma} \wedge \vartheta^{\delta} \right]$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0909 (13.5.14)	291	■ 1.000	$\mathrm{\tilde{L}}_{\mathrm{SSB}} \cong \frac{1}{2} \kappa R^{\alpha \beta} \wedge \eta_{\alpha \beta} + \frac{\Lambda}{\kappa} \eta$	$\tilde{L}_{\mathrm{SSB}} \cong -\frac{1}{2\kappa} R^{\alpha\beta} \wedge \eta_{\alpha\beta} + \frac{\Lambda}{\kappa} \eta$	$\tilde{L}_{\mathrm{SSB}} \cong -\frac{1}{2\kappa} R^{\alpha\beta} \wedge \eta_{\alpha\beta} + \frac{\Lambda}{\kappa} \eta$
mielke-geometrodynamics_EQ0910 (13.5.15)	291	■ 1.000	$\mu=\frac{4}{3} \kappa \Lambda=\frac{1}{3}(2 \kappa)^2 \rho_{\Lambda} .$	$\mu=\frac{4}{3} \kappa \Lambda=\frac{1}{3}(2 \kappa)^2 \rho_{\Lambda} .$	$\mu=\frac{4}{3} \kappa \Lambda=\frac{1}{3}(2 \kappa)^2 \rho_{\Lambda} .$
mielke-geometrodynamics_EQ0912 (13.7.1)	293	■ 1.000	$g_{\mathrm{N}}:=\kappa k^2, \quad \lambda:=\Lambda / k^2,$	$g_{\mathrm{N}}:=\kappa k^2, \quad \lambda:=\Lambda / k^2,$	$g_{\mathrm{N}}:=\kappa k^2, \quad \lambda:=\Lambda / k^2,$
mielke-geometrodynamics_EQ0913 (13.7.2)	293	■ 1.000	$k \frac{\partial}{\partial k} g_{\mathrm{N}}=\beta_1\left(g_{\mathrm{N}}, \lambda\right)=\left(2+d_{\mathrm{N}}\right) g_{\mathrm{N}}, \quad k \frac{\partial}{\partial k} \lambda=\beta_2\left(g_{\mathrm{N}}, \lambda\right),$	$k \frac{\partial}{\partial k} g_{\mathrm{N}}=\beta_1\left(g_{\mathrm{N}}, \lambda\right)=\left(2+d_{\mathrm{N}}\right) g_{\mathrm{N}}, \quad k \frac{\partial}{\partial k} \lambda=\beta_2\left(g_{\mathrm{N}}, \lambda\right),$	$k \frac{\partial}{\partial k} g_{\mathrm{N}}=\beta_1\left(g_{\mathrm{N}}, \lambda\right)=\left(2+d_{\mathrm{N}}\right) g_{\mathrm{N}}, \quad k \frac{\partial}{\partial k} \lambda=\beta_2\left(g_{\mathrm{N}}, \lambda\right),$
mielke-geometrodynamics_EQ0914 (13.7.3)	293	■ 1.000	$\mu_* \leq \frac{4}{3} g_{\mathrm{N}*} \lambda_* \simeq 0.2$	$\mu_* \leq \frac{4}{3} g_{\mathrm{N}*} \lambda_* \simeq 0.2$	$\mu_* \leq \frac{4}{3} g_{\mathrm{N}*} \lambda_* \simeq 0.2$
mielke-geometrodynamics_EQ0916 (13.8.2)	295	■ 1.000	$d C_{\mathrm{TT}}=\frac{1}{2 \ell^2}\left(T^{\alpha} \wedge T_{\alpha}+R_{\alpha \beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta}\right)$	$d C_{\mathrm{TT}}=\frac{1}{2 \ell^2}\left(T^{\alpha} \wedge T_{\alpha}+R_{\alpha \beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta}\right)$	$d C_{\mathrm{TT}}=\frac{1}{2 \ell^2}\left(T^{\alpha} \wedge T_{\alpha}+R_{\alpha \beta} \wedge \vartheta^{\alpha} \wedge \vartheta^{\beta}\right)$
mielke-geometrodynamics_EQ0917 (13.8.3)	295	■ 1.000	$d C_{\mathrm{RR}}=-\frac{1}{2} R_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha}$	$d C_{\mathrm{RR}}=-\frac{1}{2} R_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha}$	$d C_{\mathrm{RR}}=-\frac{1}{2} R_{\alpha}^{\beta} \wedge R_{\beta}^{\alpha}$
mielke-geometrodynamics_EQ0918 (13.8.4)	295	■ 1.000	$D R_{\alpha}^{\beta} \equiv 0,$	$D R_{\alpha}^{\beta} \equiv 0,$	$D R_{\alpha}^{\beta} \equiv 0,$
mielke-geometrodynamics_EQ0921 (13.8.7)	296	■ 1.000	$L_{\mathrm{SKY}}:=-\frac{1}{2} R_{\alpha \beta} \wedge * R^{\alpha \beta},$	$L_{\mathrm{SKY}}:=-\frac{1}{2} R_{\alpha \beta} \wedge * R^{\alpha \beta},$	$L_{\mathrm{SKY}}:=-\frac{1}{2} R_{\alpha \beta} \wedge * R^{\alpha \beta},$
mielke-geometrodynamics_EQ0922 (14.1.1)	302	■ 1.000	$L=L_g+L_{\omega}+L_{\mathrm{mat}}$	$L=L_g+L_{\omega}+L_{\mathrm{mat}}$	$L=L_g+L_{\omega}+L_{\mathrm{mat}}$
mielke-geometrodynamics_EQ0923 (14.1.2)	302	■ 1.000	$L_{\omega-g}=\mathrm{Tr}\left\{\wedge^*\left(\vartheta \wedge \Omega^g\right) \wedge^*\left(\vartheta \wedge \Omega\right)\right\}$	$L_{\omega-g}=\mathrm{Tr}\left\{\wedge^*\left(\vartheta \wedge \Omega^g\right) \wedge^*\left(\vartheta \wedge \Omega\right)\right\}$	$L_{\omega-g}=\mathrm{Tr}\left\{\wedge^*\left(\vartheta \wedge \Omega^g\right) \wedge^*\left(\vartheta \wedge \Omega\right)\right\}$
mielke-geometrodynamics_EQ0924 (14.1.3)	303	■ 1.000	$L_{\mathrm{GMD}}.=\frac{1}{\ell^2} L_{\mathrm{W}}.+L_{\mathrm{YM}}+L_{\mathrm{mat}}$	$L_{\mathrm{GMD}}.=\frac{1}{\ell^2} L_{\mathrm{W}}.+L_{\mathrm{YM}}+L_{\mathrm{mat}}$	$L_{\mathrm{GMD}}.=\frac{1}{\ell^2} L_{\mathrm{W}}.+L_{\mathrm{YM}}+L_{\mathrm{mat}}$
mielke-geometrodynamics_EQ0933 (14.1.14)	305	■ 1.000	$e^{*\pi / 2}=(-1)^z *, \quad z:=\frac{s-1}{2} .$	$e^{*\pi / 2}=(-1)^z *, \quad z:=\frac{s-1}{2} .$	$e^{*\pi / 2}=(-1)^z *, \quad z:=\frac{s-1}{2} .$
mielke-geometrodynamics_EQ0934 (14.1.15)	305	■ 1.000	$\xi:=e^{*(-\delta)} \Omega$	$\xi:=e^{*(-\delta)} \Omega$	$\xi:=e^{*(-\delta)} \Omega$
mielke-geometrodynamics_EQ0935 (14.1.16)	305	■ 1.000	$\Omega \wedge^* \Omega \rightarrow \xi \wedge^* \xi=\Omega \wedge^* \Omega \cos 2 \delta-(-1)^z \Omega \wedge \Omega \sin 2 \delta$	$\Omega \wedge^* \Omega \rightarrow \xi \wedge^* \xi=\Omega \wedge^* \Omega \cos 2 \delta-(-1)^z \Omega \wedge \Omega \sin 2 \delta$	$\Omega \wedge^* \Omega \rightarrow \xi \wedge^* \xi=\Omega \wedge^* \Omega \cos 2 \delta-(-1)^z \Omega \wedge \Omega \sin 2 \delta$
mielke-geometrodynamics_EQ0936 (14.1.17)	305	■ 1.000	$\Omega \wedge \Omega \rightarrow \xi \wedge \xi=\Omega \wedge \Omega \cos 2 \delta-(-1)^z \Omega \wedge^* \Omega \sin 2 \delta .$	$\Omega \wedge \Omega \rightarrow \xi \wedge \xi=\Omega \wedge \Omega \cos 2 \delta-(-1)^z \Omega \wedge^* \Omega \sin 2 \delta .$	$\Omega \wedge \Omega \rightarrow \xi \wedge \xi=\Omega \wedge \Omega \cos 2 \delta-(-1)^z \Omega \wedge^* \Omega \sin 2 \delta .$
mielke-geometrodynamics_EQ0937 (14.1.18)	305	■ 1.000	$\vec{E}=\left\{E_j:=F_{j 0} \mid j=1,2,3\right\}$	$\vec{E}=\left\{E_j:=F_{j 0} \mid j=1,2,3\right\}$	$\vec{E}=\left\{E_j:=F_{j 0} \mid j=1,2,3\right\}$
mielke-geometrodynamics_EQ0938 (14.1.19)	305	■ 1.000	$\vec{B}=\left\{B_j:=\frac{1}{2} \epsilon_{i j k} F^{j k} \mid i, j, k=1,2,3\right\}$	$\vec{B}=\left\{B_j:=\frac{1}{2} \epsilon_{i j k} F^{j k} \mid i, j, k=1,2,3\right\}$	$\vec{B}=\left\{B_j:=\frac{1}{2} \epsilon_{i j k} F^{j k} \mid i, j, k=1,2,3\right\}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_EQ0943 (14.1.24)	306	■ 1.000	$\Omega = e^{*\delta} \xi$.	$\Omega = e^{*\delta} \xi$.	$\Omega = e^{*\delta} \xi$.
mielke-geometrodynamics_EQ0944 (14.1.25)	306	■ 1.000	$\Sigma_{\mathrm{YM}}(\Omega) = \Sigma_{\mathrm{YM}}(\xi)$	$\Sigma_{\mathrm{YM}}(\Omega) = \Sigma_{\mathrm{YM}}(\xi)$	$\Sigma_{\mathrm{YM}}(\Omega) = \Sigma_{\mathrm{YM}}(\xi)$
mielke-geometrodynamics_EQ0950 (14.1.31)	307	■ 1.000	$\tau_e = \bar{\tau}_e \cos \delta - (-1)^z \bar{\tau}_p \sin \delta$	$\tau_e = \bar{\tau}_e \cos \delta - (-1)^z \bar{\tau}_p \sin \delta$	$\tau_e = \bar{\tau}_e \cos \delta - (-1)^z \bar{\tau}_p \sin \delta$
mielke-geometrodynamics_EQ0952 (14.1.34)	307	■ 1.000	$D \delta = d \delta = 0,$	$D \delta = d \delta = 0,$	$D \delta = d \delta = 0,$
mielke-geometrodynamics_EQ0955 (14.1.37)	308	■ 1.000	$(-1)^z \Sigma_{\mathrm{YM}}(\xi) \wedge d \delta = D \Sigma_{\mathrm{YM}}(\xi).$	$(-1)^z \Sigma_{\mathrm{YM}}(\xi) \wedge d \delta = D \Sigma_{\mathrm{YM}}(\xi).$	$(-1)^z \Sigma_{\mathrm{YM}}(\xi) \wedge d \delta = D \Sigma_{\mathrm{YM}}(\xi).$
mielke-geometrodynamics_EQ0958 (14.1.40)	309	■ 1.000	$\delta = \int_{c_1} d \delta + \delta_0$	$\delta = \int_{c_1} d \delta + \delta_0$	$\delta = \int_{c_1} d \delta + \delta_0.$
mielke-geometrodynamics_EQ0969 (14.2.10)	314	■ 1.000	$\hat{\omega}^{AB} = -\hat{\omega}^{BA}, \quad A, B = 0, \dots, 3 + K$	$\hat{\omega}^{AB} = -\hat{\omega}^{BA}, \quad A, B = 0, \dots, 3 + K$	$\hat{\omega}^{AB} = -\hat{\omega}^{BA}, \quad A, B = 0, \dots, 3 + K$
mielke-geometrodynamics_EQ0975 (14.2.16)	315	■ 1.000	$\Omega^{\prime} := \pi \left(d \omega - \omega \wedge \overset{\circ}{\omega} - \omega^g \wedge \omega \right) = \Omega - \omega^g \wedge \omega.$	$\Omega^{\prime} := \pi \left(d \omega - \omega \wedge \overset{\circ}{\omega} - \omega^g \wedge \omega \right) = \Omega - \omega^g \wedge \omega.$	$\Omega^{\prime} := \pi (d \omega - \omega \wedge \overset{\circ}{\omega} - \omega^g \wedge \omega) = \Omega - \omega^g \wedge \omega.$
mielke-geometrodynamics_EQ0980 (14.2.21)	316	■ 1.000	$\hat{g}_{AB}(x, \xi^j) = \sum_{n=-\infty}^{+\infty} g_{AB}^{(n)}(x) Y_n(\xi^j)$	$\hat{g}_{AB}(x, \xi^j) = \sum_{n=-\infty}^{+\infty} g_{AB}^{(n)}(x) Y_n(\xi^j)$	$\hat{g}_{AB}(x, \xi^j) = \sum_{n=-\infty}^{+\infty} g_{AB}^{(n)}(x) Y_n(\xi^j)$
mielke-geometrodynamics_EQ0981 (14.2.22)	317	■ 1.000	$M^* = \sqrt{\hbar c / G_{\mathrm{N}}} \simeq 10^{19} \mathrm{GeV} / c^2.$	$M^* = \sqrt{\hbar c / G_{\mathrm{N}}} \simeq 10^{19} \mathrm{GeV} / c^2.$	$M^* = \sqrt{\hbar c / G_{\mathrm{N}}} \simeq 10^{19} \mathrm{GeV} / c^2.$
mielke-geometrodynamics_EQ0983 (14.2.24)	318	■ 1.000	$G = SU(3) \times SU(2) \times U(1)$	$G = SU(3) \times SU(2) \times U(1)$	$G = SU(3) \times SU(2) \times U(1)$
mielke-geometrodynamics_EQ0984 (14.2.25)	318	■ 1.000	$B = \mathbb{C} P^2 \times S^2 \times S^1,$	$B = \mathbb{C} P^2 \times S^2 \times S^1,$	$B = \mathbb{C} P^2 \times S^2 \times S^1,$
mielke-geometrodynamics_EQ0985 (14.2.26)	318	■ 1.000	$G / H = \frac{SU(3) \times SU(2) \times U(1)}{SU(2) \times U(1) \times U(1)},$	$G / H = \frac{SU(3) \times SU(2) \times U(1)}{SU(2) \times U(1) \times U(1)},$	$G / H = \frac{SU(3) \times SU(2) \times U(1)}{SU(2) \times U(1) \times U(1)},$
mielke-geometrodynamics_EQ0986 (14.2.27)	318	■ 1.000	$\widetilde{S}^7 := \frac{SO(5) \times SO(3)}{SO(3) \times SO(3)}.$	$\widetilde{S}^7 := \frac{SO(5) \times SO(3)}{SO(3) \times SO(3)}.$	$\widetilde{S}^7 := \frac{SO(5) \times SO(3)}{SO(3) \times SO(3)}.$
mielke-geometrodynamics_EQ0988 (14.3.1)	319	■ 1.000	$\varphi \rightarrow \varphi^{\mathrm{C}} := \bar{\varphi}$	$\varphi \rightarrow \varphi^{\mathrm{C}} := \bar{\varphi}$	$\varphi \rightarrow \varphi^{\mathrm{C}} := \bar{\varphi}$
mielke-geometrodynamics_EQ0991 (14.3.4)	320	■ 1.000	$\hat{\psi} = \sum_{F=1}^{\dim \rho(G)} \left(\psi_{-}^F(x) \chi_{\rho F}^{-}(\xi) + \psi_{+}^F(x) \chi_{\rho F}^{+}(\xi) \right).$	$\hat{\psi} = \sum_{F=1}^{\dim \rho(G)} \left(\psi_{-}^F(x) \chi_{\rho F}^{-}(\xi) + \psi_{+}^F(x) \chi_{\rho F}^{+}(\xi) \right).$	$\hat{\psi} = \sum_{F=1}^{\dim \rho(G)} (\psi_{-}^F(x) \chi_{\rho F}^{-}(\xi) + \psi_{+}^F(x) \chi_{\rho F}^{+}(\xi)).$
mielke-geometrodynamics_EQ0994 (14.3.7)	321	■ 1.000	$D^{g+\omega} := d + i \widetilde{\omega}^g \otimes i \omega$	$D^{g+\omega} := d + i \widetilde{\omega}^g \otimes i \omega$	$D^{g+\omega} := d + i \widetilde{\omega}^g \otimes i \omega$
mielke-geometrodynamics_EQ0999 (14.3.12)	322	■ 1.000	$(\widehat{\gamma}^{n+1})^2 = -1 \Rightarrow \widehat{\gamma}^{n+1} \widehat{\psi} = \mp i \widehat{\psi}.$	$(\widehat{\gamma}^{n+1})^2 = -1 \Rightarrow \widehat{\gamma}^{n+1} \widehat{\psi} = \mp i \widehat{\psi}.$	$(\widehat{\gamma}^{n+1})^2 = -1 \Rightarrow \widehat{\gamma}^{n+1} \widehat{\psi} = \mp i \widehat{\psi}.$
mielke-geometrodynamics_EQ1000 (14.3.13)	322	■ 1.000	$(\widehat{\gamma}^{n+1})^2 = +1 \Rightarrow \widehat{\gamma}^{n+1} \widehat{\psi} = \mp \widehat{\psi}.$	$(\widehat{\gamma}^{n+1})^2 = +1 \Rightarrow \widehat{\gamma}^{n+1} \widehat{\psi} = \mp \widehat{\psi}.$	$(\widehat{\gamma}^{n+1})^2 = +1 \Rightarrow \widehat{\gamma}^{n+1} \widehat{\psi} = \mp \widehat{\psi}.$
mielke-geometrodynamics_EQ1006 (14.4.1)	324	■ 1.000	$\bar{Q}^{(i)} = Q^{(i)} \cos \delta + P^{(i)} \sin \delta, \quad Q^{(\circ)} \equiv Q_e.$	$\bar{Q}^{(i)} = Q^{(i)} \cos \delta + P^{(i)} \sin \delta, \quad Q^{(\circ)} \equiv Q_e.$	$\bar{Q}^{(i)} = Q^{(i)} \cos \delta + P^{(i)} \sin \delta, \quad Q^{(\circ)} \equiv Q_e.$
mielke-geometrodynamics_EQ1009 (14.4.4)	325	■ 1.000	$\rho^2 := r^2 + \frac{J(J+1) \hbar^2}{M^2 c^2} \cos^2 \vartheta$	$\rho^2 := r^2 + \frac{J(J+1) \hbar^2}{M^2 c^2} \cos^2 \vartheta$	$\rho^2 := r^2 + \frac{J(J+1) \hbar^2}{M^2 c^2} \cos^2 \vartheta$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_325 EQ1010 (14.4.5)		■ 1.000	$\Delta:=r^{\wedge\{2\}}-\frac{2\text{ M }\hbar}{M^{\wedge\{2\}}c}\text{ r}+\frac{J(J+1)}{\hbar^{\wedge\{2\}}\{M^{\wedge\{2\}}c^{\wedge\{2\}}+\frac{\alpha_e}{\hbar^{\wedge\{2\}}\{M^{\wedge\{2\}}c^{\wedge\{2\}}}\operatorname{Tr}\bar{Q}^{\wedge\{2\}}$	$\Delta := r^2 - \frac{2M\hbar}{M^{*2}c}r + \frac{J(J+1)\hbar^2}{M^2c^2} + \frac{\alpha_e\hbar^2}{M^{*2}c^2}\operatorname{Tr}\bar{Q}^2$	$\Delta := r^2 - \frac{2M\hbar}{M^{*2}c}r + \frac{J(J+1)\hbar^2}{M^2c^2} + \frac{\alpha_e\hbar^2}{M^{*2}c^2}\operatorname{Tr}\bar{Q}^2$
mielke-geometrodynamics_327 EQ1017 (14.4.12)		■ 1.000	$d\text{ s}^{\wedge\{2\}}=-\frac{\bar{\Delta}}{\bar{\rho}^2}\left(\frac{4N\hbar}{Mc}\sin^2\left(\frac{\vartheta}{2}\right)d\phi+c\,dt\right)^2+\frac{\bar{\rho}^2}{\bar{\Delta}}\left(dr^2+\bar{\Delta}d\Omega^2\right),$	$ds^2 = -\frac{\bar{\Delta}}{\bar{\rho}^2}\left(\frac{4N\hbar}{Mc}\sin^2\left(\frac{\vartheta}{2}\right)d\phi+c\,dt\right)^2 + \frac{\bar{\rho}^2}{\bar{\Delta}}\left(dr^2+\bar{\Delta}d\Omega^2\right),$	$ds^2 = -\frac{\bar{\Delta}}{\bar{\rho}^2}\left(\frac{4N\hbar}{Mc}\sin^2\left(\frac{\vartheta}{2}\right)d\phi+c\,dt\right)^2 + \frac{\bar{\rho}^2}{\bar{\Delta}}(dr^2+\bar{\Delta}d\Omega^2),$
mielke-geometrodynamics_328 EQ1018 (14.4.13)		■ 1.000	$\bar{\Delta}:=r^{\wedge\{2\}}-\frac{2\text{ M }\hbar}{M^{\wedge\{2\}}c}\text{ r}-\frac{N^{\wedge\{2\}}\hbar^{\wedge\{2\}}\{M^{\wedge\{2\}}c^{\wedge\{2\}}+\frac{\alpha_e}{\hbar^{\wedge\{2\}}\{M^{\wedge\{2\}}c^{\wedge\{2\}}}\operatorname{Tr}\bar{Q}^{\wedge\{2\}}$	$\bar{\Delta} := r^2 - \frac{2M\hbar}{M^{*2}c}r - \frac{N^2\hbar^2}{M^2c^2} + \frac{\alpha_e\hbar^2}{M^{*2}c^2}\operatorname{Tr}\bar{Q}^2$	$\bar{\Delta} := r^2 - \frac{2M\hbar}{M^{*2}c}r - \frac{N^2\hbar^2}{M^2c^2} + \frac{\alpha_e\hbar^2}{M^{*2}c^2}\operatorname{Tr}\bar{Q}^2$
mielke-geometrodynamics_328 EQ1019 (14.4.14)		■ 1.000	$\bar{\rho}^{\wedge\{2\}}:=r^{\wedge\{2\}}+\frac{N^{\wedge\{2\}}\hbar^{\wedge\{2\}}\{M^{\wedge\{2\}}c^{\wedge\{2\}}$	$\bar{\rho}^2 := r^2 + \frac{N^2\hbar^2}{M^2c^2}$	$\bar{\rho}^2 := r^2 + \frac{N^2\hbar^2}{M^2c^2}$
mielke-geometrodynamics_328 EQ1020 (14.4.15)		■ 1.000	$\begin{aligned} A = & \; -\frac{1}{\bar{\rho}^2}\left(\bar{Q}\cos\delta+N\bar{Q}\sin\delta\right)\left(\bar{Q}\cos\delta+N\bar{Q}\sin\delta\right)c\,dt \\ & +\frac{1-\cos\vartheta}{\bar{\rho}^2}\bar{Q}\left[-\bar{\rho}^2\sin\delta+2r\frac{N\hbar}{Mc}\cos\delta\right]d\phi. \end{aligned}$	$A = -\frac{1}{\bar{\rho}^2}(\bar{Q}\cos\delta+N\bar{Q}\sin\delta)c\,dt + \frac{1-\cos\vartheta}{\bar{\rho}^2}\bar{Q}\left[-\bar{\rho}^2\sin\delta+2r\frac{N\hbar}{Mc}\cos\delta\right]d\phi$	$A = -\frac{1}{\bar{\rho}^2}(\bar{Q}\cos\delta+N\bar{Q}\sin\delta)c\,dt + \frac{1-\cos\vartheta}{\bar{\rho}^2}\bar{Q}\left[-\bar{\rho}^2\sin\delta+2r\frac{N\hbar}{Mc}\cos\delta\right]d\phi.$
mielke-geometrodynamics_328 EQ1022 (14.4.17)		■ 1.000	$M^{\wedge\{2\}}/M_{\mathrm{ir}}^{\wedge\{2\}}=\frac{\left(\left(1+\frac{1}{4}\left(\alpha_e^2\operatorname{Tr}\bar{Q}^2-2N^2\right)\left(M^*/M_{\mathrm{ir}}\right)^4\right)\right)^2}{1-\frac{1}{4}N^2\left(M^*/M_{\mathrm{ir}}\right)^4},$	$M^2/M_{\mathrm{ir}}^2 = \frac{\left[1+\frac{1}{4}\left(\alpha_e^2\operatorname{Tr}\bar{Q}^2-2N^2\right)\left(M^*/M_{\mathrm{ir}}\right)^4\right]^2}{1-\frac{1}{4}N^2\left(M^*/M_{\mathrm{ir}}\right)^4},$	$M^2/M_{\mathrm{ir}}^2 = \frac{\left[1+\frac{1}{4}(\alpha_e^2\operatorname{Tr}\bar{Q}^2-2N^2)(M^*/M_{\mathrm{ir}})^4\right]^2}{1-\frac{1}{4}N^2(M^*/M_{\mathrm{ir}})^4},$
mielke-geometrodynamics_335 EQ1023 (15.0.1)		■ 1.000	$\ell:=2\pi\hbar/\left(M_Pc\right)=\ell^*/\sqrt{\kappa}\simeq1.3\times10^{-13}\,\mathrm{cm}$	$\ell := 2\pi\hbar/(M_Pc) = \ell^*/\sqrt{\kappa} \simeq 1.3 \times 10^{-13} \operatorname{cm}$	$\ell := 2\pi\hbar/(M_Pc) = \ell^*/\sqrt{\kappa} \simeq 1.3 \times 10^{-13} \operatorname{cm}$
mielke-geometrodynamics_335 EQ1024 (15.0.2)		■ 1.000	$V_{\mathrm{Yu}}(r)\sim\frac{1}{r}e^{-rm_{\pi}c/\hbar},$	$V_{\mathrm{Yu}}(r) \sim \frac{1}{r}e^{-rm_{\pi}c/\hbar},$	$V_{\mathrm{Yu}}(r) \sim \frac{1}{r}e^{-rm_{\pi}c/\hbar},$
mielke-geometrodynamics_336 EQ1026 (15.1.2)		■ 1.000	$L_{\{f\,g\}}=P\left(\vartheta^{\wedge\{f\}},\,\vartheta^{\wedge\{g\}}\right)\eta,$	$L_{fg} = P\left(\vartheta^f,\vartheta^g\right)\eta,$	$L_{fg} = P(\vartheta^f,\vartheta^g)\eta,$
mielke-geometrodynamics_337 EQ1027 (15.1.3)		■ 1.000	$\delta L_{\{f\,g\}}=: \delta\vartheta^{\wedge\{f\}}\wedge\Sigma^{\wedge\{f\}}+\delta\vartheta^{\wedge\{g\}}\wedge\Sigma^{\wedge\{g\}}.$	$\delta L_{fg} =: \delta\vartheta^f\wedge\Sigma^f+\delta\vartheta^g\wedge\Sigma^g.$	$\delta L_{fg} =: \delta\vartheta^f\wedge\Sigma^f+\delta\vartheta^g\wedge\Sigma^g.$
mielke-geometrodynamics_337 EQ1028 (15.1.4)		■ 1.000	$\frac{1}{a}\vartheta^{\wedge\{a\}}\wedge\Sigma^{\wedge\{f\}}+\frac{1}{b}\vartheta^{\wedge\{b\}}\wedge\Sigma^{\wedge\{g\}}=2\,L_{\{f\,g\}}.$	$\frac{1}{a}\vartheta^f\wedge\Sigma^f+\frac{1}{b}\vartheta^g\wedge\Sigma^g=2L_{fg}.$	$\frac{1}{a}\vartheta^f\wedge\Sigma^f+\frac{1}{b}\vartheta^g\wedge\Sigma^g=2\,L_{fg}.$
mielke-geometrodynamics_337 EQ1029 (15.1.5)		■ 1.000	$L_{\{f\,g\}}^{\wedge\{\prime\}}=\frac{m_f^2}{\ell^2}\left \det g_{ij}\right ^u\left \det f_{ij}\right ^{1/2-u}\left(\vartheta^f-\vartheta^g\right)\wedge\left(\vartheta^f-\vartheta^g\right)\wedge\vartheta^g\wedge\vartheta^g$	$L'_{fg} = \frac{m_f^2}{\ell^2} \det g_{ij} ^u \det f_{ij} ^{1/2-u}\left(\vartheta^f-\vartheta^g\right)\wedge\left(\vartheta^f-\vartheta^g\right)\wedge\vartheta^g\wedge\vartheta^g$	$L'_{fg} = \frac{m_f^2}{\ell^2} \det g_{ij} ^u \det f_{ij} ^{1/2-u}(\vartheta^f-\vartheta^g)\wedge(\vartheta^f-\vartheta^g)\wedge\vartheta^g\wedge\vartheta^g$
mielke-geometrodynamics_338 EQ1035 (15.1.12)		■ 1.000	$\vartheta^{\wedge\{m_f\}}:=\vartheta^{\wedge\{f\}}-\vartheta^{\wedge\{g\}}$	$\vartheta^{m_f} := \vartheta^f - \vartheta^g$	$\vartheta^{m_f} := \vartheta^f - \vartheta^g$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_338 EQ1036 (15.1.13)		■ 1.000	$\begin{aligned} ds^2 &= \frac{1}{1+\kappa} g^{\alpha\beta} \left(\vartheta_\alpha^g \otimes_s \vartheta_\beta^g + \kappa \vartheta_\alpha^f \otimes \vartheta_\beta^f \right) \\ &= \frac{1}{1+\kappa} (g_{ij} + \kappa f_{ij}) dx^i \otimes_s dx^j \end{aligned}$	$ds^2 = \frac{1}{1+\kappa} g^{\alpha\beta} \left(\vartheta_\alpha^g \otimes_s \vartheta_\beta^g + \kappa \vartheta_\alpha^f \otimes \vartheta_\beta^f \right) \\ = \frac{1}{1+\kappa} (g_{ij} + \kappa f_{ij}) dx^i \otimes_s dx^j$	$ds^2 = \frac{1}{1+\kappa} g^{\alpha\beta} \left(\vartheta_\alpha^g \otimes_s \vartheta_\beta^g + \kappa \vartheta_\alpha^f \otimes \vartheta_\beta^f \right) \\ = \frac{1}{1+\kappa} (g_{ij} + \kappa f_{ij}) dx^i \otimes_s dx^j.$
mielke-geometrodynamics_338 EQ1037 (15.1.14)		■ 1.000	$\kappa = \ell^{*2} / \ell^2 = G_{\mathrm{N}} / G_{\mathrm{S}} = 0.38 \times 10^{-38}$	$\kappa = \ell^{*2} / \ell^2 = G_{\mathrm{N}} / G_{\mathrm{S}} = 0.38 \times 10^{-38}$	$\kappa = \ell^{*2} / \ell^2 = G_{\mathrm{N}} / G_{\mathrm{S}} = 0.38 \times 10^{-38}$
mielke-geometrodynamics_339 EQ1038 (15.1.15)		■ 1.000	$e^{-\lambda^{g,f}} := 1 - \frac{2\mu^{g,f}}{r} - \frac{1}{3} \Lambda^{g,f} r^2,$	$e^{-\lambda^{g,f}} := 1 - \frac{2\mu^{g,f}}{r} - \frac{1}{3} \Lambda^{g,f} r^2,$	$e^{-\lambda^{g,f}} := 1 - \frac{2\mu^{g,f}}{r} - \frac{1}{3} \Lambda^{g,f} r^2,$
mielke-geometrodynamics_339 EQ1040 (15.1.17)		■ 1.000	$e^{-\lambda^f} \simeq 1 - \frac{2\mu^f}{r} \exp(-r m_f c / \hbar).$	$e^{-\lambda^f} \simeq 1 - \frac{2\mu^f}{r} \exp(-r m_f c / \hbar).$	$e^{-\lambda^f} \simeq 1 - \frac{2\mu^f}{r} \exp(-r m_f c / \hbar).$
mielke-geometrodynamics_340 EQ1041 (15.1.18)		■ 1.000	$E_{\alpha\beta} : (2^+) \oplus 2(1^-) \oplus 2(0^+) \oplus (1^+) \\ \oplus 2(0^+) \oplus 2(1^-) \oplus 2(0^+) \oplus (1^+)$	$E_{\alpha\beta} : (2^+) \oplus 2(1^-) \oplus 2(0^+) \oplus (1^+)$	$E_{\alpha\beta} : (2^+) \oplus 2(1^-) \oplus 2(0^+) \oplus (1^+)$
mielke-geometrodynamics_342 EQ1043 (15.1.20)		■ 1.000	$\frac{M_{\mathrm{U}}}{M_{\mathrm{P}}} = \frac{\ell_{\mathrm{H}}}{\ell^*} \cdot \frac{\ell}{\ell^*} = \frac{1}{\kappa} \frac{\ell_{\mathrm{H}}}{\ell} \simeq 10^{80}$	$\frac{M_{\mathrm{U}}}{M_{\mathrm{P}}} = \frac{\ell_{\mathrm{H}}}{\ell^*} \cdot \frac{\ell}{\ell^*} = \frac{1}{\kappa} \frac{\ell_{\mathrm{H}}}{\ell} \simeq 10^{80}$	$\frac{M_{\mathrm{U}}}{M_{\mathrm{P}}} = \frac{\ell_{\mathrm{H}}}{\ell^*} \cdot \frac{\ell}{\ell^*} = \frac{1}{\kappa} \frac{\ell_{\mathrm{H}}}{\ell} \simeq 10^{80}$
mielke-geometrodynamics_343 EQ1044 (15.2.1)		■ 1.000	$\begin{aligned} \bar{G} &= SL(2f, \mathbb{C}) \otimes SL(2c, \mathbb{C}) \\ &\supset SU(f)_{\mathrm{L}} \otimes SU(f)_{\mathrm{R}} \otimes SU(c)_{\mathrm{L}} \otimes SU(c)_{\mathrm{R}} \end{aligned}$	$\bar{G} = SL(2f, \mathbb{C}) \otimes SL(2c, \mathbb{C}) \\ \supset SU(f)_{\mathrm{L}} \otimes SU(f)_{\mathrm{R}} \otimes SU(c)_{\mathrm{L}} \otimes SU(c)_{\mathrm{R}}$	$\bar{G} = SL(2f, \mathbb{C}) \otimes SL(2c, \mathbb{C}) \\ \supset SU(f)_{\mathrm{L}} \otimes SU(f)_{\mathrm{R}} \otimes SU(c)_{\mathrm{L}} \otimes SU(c)_{\mathrm{R}}$
mielke-geometrodynamics_343 EQ1045 (15.2.2)		■ 1.000	$\bar{L}(M) := M^4 \times \bar{G} \supset M^4 \times SL(2, \mathbb{C})$	$\bar{L}(M) := M^4 \times \bar{G} \supset M^4 \times SL(2, \mathbb{C})$	$\bar{L}(M) := M^4 \times \bar{G} \supset M^4 \times SL(2, \mathbb{C})$
mielke-geometrodynamics_344 EQ1046 (15.2.3)		■ 1.000	$\bar{\gamma} = \bar{E}_j^\alpha(m) \gamma_\alpha \otimes dx^j = \bar{E}_j dx^j$	$\bar{\gamma} = \bar{E}_j^\alpha(m) \gamma_\alpha \otimes dx^j = \bar{E}_j dx^j$	$\bar{\gamma} = \bar{E}_j^\alpha(m) \gamma_\alpha \otimes dx^j = \bar{E}_j dx^j$
mielke-geometrodynamics_344 EQ1048 (15.2.5)		■ 1.000	$\frac{1}{4} \mathrm{Tr}(\bar{\gamma} \otimes_s \bar{\gamma}) = f_{ij} dx^i \otimes_s dx^j$	$\frac{1}{4} \mathrm{Tr}(\bar{\gamma} \otimes_s \bar{\gamma}) = f_{ij} dx^i \otimes_s dx^j$	$\frac{1}{4} \mathrm{Tr}(\bar{\gamma} \otimes_s \bar{\gamma}) = f_{ij} dx^i \otimes_s dx^j$
mielke-geometrodynamics_345 EQ1052 (15.2.9)		■ 1.000	$\bar{\omega}_i = \frac{1}{2} \Gamma_i^{(f)\alpha\beta j} \sigma_{\alpha\beta} \lambda_j^{(f)} \oplus \frac{1}{2} \Gamma_i^{(c)\alpha\beta j} \sigma_{\alpha\beta} \lambda_j^{(c)} \oplus A_i.$	$\bar{\omega}_i = \frac{1}{2} \Gamma_i^{(f)\alpha\beta j} \sigma_{\alpha\beta} \lambda_j^{(f)} \oplus \frac{1}{2} \Gamma_i^{(c)\alpha\beta j} \sigma_{\alpha\beta} \lambda_j^{(c)} \oplus A_i.$	$\bar{\omega}_i = \frac{1}{2} \Gamma_i^{(f)\alpha\beta j} \sigma_{\alpha\beta} \lambda_j^{(f)} \oplus \frac{1}{2} \Gamma_i^{(c)\alpha\beta j} \sigma_{\alpha\beta} \lambda_j^{(c)} \oplus A_i.$
mielke-geometrodynamics_345 EQ1053 (15.2.10)		■ 1.000	$\bar{\Theta} := \bar{D}\bar{\gamma} = d\bar{\gamma} - [\bar{\omega}, \bar{\gamma}]$	$\bar{\Theta} := \bar{D}\bar{\gamma} = d\bar{\gamma} - [\bar{\omega}, \bar{\gamma}]$	$\bar{\Theta} := \bar{D}\bar{\gamma} = d\bar{\gamma} - [\bar{\omega}, \bar{\gamma}]$
mielke-geometrodynamics_345 EQ1054 (15.2.11)		■ 1.000	$\bar{\Omega} = d\bar{\omega} - \bar{\omega} \wedge \bar{\omega}$	$\bar{\Omega} = d\bar{\omega} - \bar{\omega} \wedge \bar{\omega}$	$\bar{\Omega} = d\bar{\omega} - \bar{\omega} \wedge \bar{\omega}$
mielke-geometrodynamics_346 EQ1056 (15.2.13)		■ 1.000	$\psi = \left\{ \psi^{(q_f, q_c)} \mid q_f = 1, \dots, f; q_c = 1, \dots, c \right\}.$	$\psi = \left\{ \psi^{(q_f, q_c)} \mid q_f = 1, \dots, f; q_c = 1, \dots, c \right\}.$	$\psi = \{ \psi^{(q_f, q_c)} \mid q_f = 1, \dots, f; q_c = 1, \dots, c \}.$
mielke-geometrodynamics_354 EQ1061 (16.1.2)		■ 1.000	$B = \frac{2\hbar c}{\ell^{*2}} \Lambda_{\mathrm{eff}} = -\frac{3\kappa}{2} \frac{\hbar c}{\ell^{*4}}$	$B = \frac{2\hbar c}{\ell^{*2}} \Lambda_{\mathrm{eff}} = -\frac{3\kappa}{2} \frac{\hbar c}{\ell^{*4}}$	$B = \frac{2\hbar c}{\ell^{*2}} \Lambda_{\mathrm{eff}} = -\frac{3\kappa}{2} \frac{\hbar c}{\ell^{*4}}$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_355 EQ1063 (16.2.2)	355	■ 1.000	$\square=\frac{1}{\sqrt{ f }}\partial_i\left(f^{ij}\sqrt{ f }\partial_j\right)$	$\square = \frac{1}{\sqrt{ f }}\partial_i\left(f^{ij}\sqrt{ f }\partial_j\right)$	$\square = \frac{1}{\sqrt{ f }}\partial_i(f^{ij}\sqrt{ f }\partial_j)$
mielke-geometrodynamics_355 EQ1064 (16.2.3)	355	■ 1.000	$\varphi=m\frac{c}{\hbar}\frac{1}{\rho}F_{nm}^{\iota}(\rho^*)Y_{\iota}^m(\vartheta,\phi)e^{-it\omega mc^2/\hbar}.$	$\varphi = m\frac{c}{\hbar}\frac{1}{\rho}F_{nm}^{\iota}(\rho^*)Y_{\iota}^m(\vartheta,\phi)e^{-it\omega mc^2/\hbar}.$	$\varphi = m\frac{c}{\hbar}\frac{1}{\rho}F_{nm}^{\iota}(\rho^*)Y_{\iota}^m(\vartheta,\phi)e^{-it\omega mc^2/\hbar}.$
mielke-geometrodynamics_355 EQ1067 (16.2.6)	355	■ 1.000	$\beta=\frac{\sqrt{2\pi\kappa}}{\ell^*}/\frac{mc}{\hbar}=\frac{\ell_m}{\ell}.$	$\beta = \frac{\sqrt{2\pi\kappa}}{\ell^*}/\frac{mc}{\hbar} = \frac{\ell_m}{\ell}.$	$\beta = \frac{\sqrt{2\pi\kappa}}{\ell^*}\Big/\frac{mc}{\hbar} = \frac{\ell_m}{\ell}.$
mielke-geometrodynamics_358 EQ1072 (16.3.1)	358	■ 1.000	$\operatorname{Tr}\bar{Q}^2=-\frac{4}{\alpha_e}b\{\alpha_e\}Y+I(I+1)-\frac{1}{4}Y^2.$	$\operatorname{Tr}\bar{Q}^2 = -\frac{4b}{\alpha_e}Y + I(I+1) - \frac{1}{4}Y^2.$	$\operatorname{Tr}\bar{Q}^2 = -\frac{4b}{\alpha_e}Y + I(I+1) - \frac{1}{4}Y^2.$
mielke-geometrodynamics_358 EQ1073 (16.3.2)	358	■ 1.000	$Q_e=I_3+\frac{1}{2}Y-\frac{2}{3}C.$	$Q_e = I_3 + \frac{1}{2}Y - \frac{2}{3}C.$	$Q_e = I_3 + \frac{1}{2}Y - \frac{2}{3}C.$
mielke-geometrodynamics_359 EQ1074 (16.3.3)	359	■ 1.000	$\frac{M^2}{M^{*2}}=\left\{1-bY+\frac{\alpha_e}{4}\left[I(I+1)-\frac{1}{4}Y^2\right]\right\}^2+\frac{1}{4}J(J+1)$	$\frac{M^2}{M^{*2}} = \left\{1-bY + \frac{\alpha_e}{4}\left[I(I+1) - \frac{1}{4}Y^2\right]\right\}^2 + \frac{1}{4}J(J+1)$	$\frac{M^2}{M^{*2}} = \left\{1-bY + \frac{\alpha_e}{4}[I(I+1) - \frac{1}{4}Y^2]\right\}^2 + \frac{1}{4}J(J+1)$
mielke-geometrodynamics_360 EQ1075 (16.3.4)	360	■ 1.000	$T_{\bullet}=\frac{2\hbar c}{S_Ak_B}\left(r_+-\frac{M}{M^{*2}}\frac{\hbar}{c}\right).$	$T_{\bullet} = \frac{2\hbar c}{S_Ak_B}\left(r_+ - \frac{M}{M^{*2}}\frac{\hbar}{c}\right).$	$T_{\bullet} = \frac{2\hbar c}{S_Ak_B}\left(r_+ - \frac{M}{M^{*2}}\frac{\hbar}{c}\right).$
mielke-geometrodynamics_364 EQ1076 (A.1)	364	■ 1.000	$x^Ay_A:=\sum_{A=0}^n\sum_{B=0}^nx^Ay^Bg_{AB}.$	$x^Ay_A := \sum_{A=0}^n\sum_{B=0}^nx^Ay^Bg_{AB}.$	$x^Ay_A := \sum_{A=0}^n\sum_{B=0}^nx^Ay^Bg_{AB}.$
mielke-geometrodynamics_364 EQ1077 (A.2)	364	■ 1.000	$c=\hbar=1.$	$c = \hbar = 1.$	$c = \hbar = 1.$
mielke-geometrodynamics_365 EQ1080 (A.5)	365	■ 1.000	$\left\{\sigma^i,\sigma^j\right\}=g^{ij}$	$\left\{\sigma^i,\sigma^j\right\} = g^{ij}$	$\{\sigma^i,\sigma^j\} = g^{ij}$
mielke-geometrodynamics_365 EQ1081 (A.6)	365	■ 1.000	$\left(\gamma^{\alpha}\right)^{\mathrm{T}}:=\left(\gamma^0,\gamma^k\right),$	$(\gamma^{\alpha})^{\mathrm{T}} := (\gamma^0,\gamma^k),$	$(\gamma^{\alpha})^{\mathrm{T}} := (\gamma^0,\gamma^k),$
mielke-geometrodynamics_365 EQ1083 (A.8)	365	■ 1.000	$\gamma^{\alpha}\gamma^{\beta}+\gamma^{\beta}\gamma^{\alpha}=2g^{\alpha\beta}$	$\gamma^{\alpha}\gamma^{\beta} + \gamma^{\beta}\gamma^{\alpha} = 2g^{\alpha\beta}$	$\gamma^{\alpha}\gamma^{\beta} + \gamma^{\beta}\gamma^{\alpha} = 2g^{\alpha\beta}$
mielke-geometrodynamics_365 EQ1084 (A.9)	365	■ 1.000	$\gamma^{0+}=\gamma^0,\quad\gamma^{k+}=-\gamma^k,\quad\gamma^{5+}=\gamma^5.$	$\gamma^{0+} = \gamma^0,\quad \gamma^{k+} = -\gamma^k,\quad \gamma^{5+} = \gamma^5.$	$\gamma^{0+} = \gamma^0,\quad \gamma^{k+} = -\gamma^k,\quad \gamma^{5+} = \gamma^5.$
mielke-geometrodynamics_366 EQ1087 (A.14)	366	■ 1.000	$C(1,3)=C(0,2)\otimes C(1,1).$	$C(1,3) = C(0,2) \otimes C(1,1).$	$C(1,3) = C(0,2) \otimes C(1,1).$
mielke-geometrodynamics_368 EQ1091 (B.1)	368	■ 1.000	$\alpha^{\left(p\right)}=\frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\in C^{\infty}\left(\wedge^pT^{\ast}(M)\right).$	$\alpha^{(p)} = \frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1} \wedge \cdots \wedge \vartheta^{\alpha_p} \in C^{\infty}\left(\wedge^pT^{\ast}(M)\right).$	$\alpha^{(p)} = \frac{1}{p!}A_{\alpha_1\cdots\alpha_p}\vartheta^{\alpha_1} \wedge \cdots \wedge \vartheta^{\alpha_p} \in C^{\infty}(\wedge^pT^{\ast}(M)).$
mielke-geometrodynamics_368 EQ1092 (B.2)	368	■ 1.000	$\begin{gathered}\alpha^{\{p\}}\wedge\beta^{\{q\}}:=\frac{1}{(p!)\cdot(q!)}A_{\alpha_1\cdots\alpha_p}\cdot B_{\beta_1\cdots\beta_q}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\\\wedge\vartheta^{\beta_1}\wedge\cdots\wedge\vartheta^{\beta_q}\\\alpha^{\{p\}}\wedge\beta^{\{q\}}:=\frac{1}{(p!)\cdot(q!)}A_{\alpha_1\cdots\alpha_p}\cdot B_{\beta_1\cdots\beta_q}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\\\wedge\vartheta^{\beta_1}\wedge\cdots\wedge\vartheta^{\beta_q}\end{gathered}$	$\alpha^{(p)}\wedge\beta^{(q)}:=\frac{1}{(p!)\cdot(q!)}A_{\alpha_1\cdots\alpha_p}\cdot B_{\beta_1\cdots\beta_q}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\\\wedge\vartheta^{\beta_1}\wedge\cdots\wedge\vartheta^{\beta_q}$	$\alpha^{(p)}\wedge\beta^{(q)}:=\frac{1}{(p!)\cdot(q!)}A_{\alpha_1\cdots\alpha_p}\cdot B_{\beta_1\cdots\beta_q}\vartheta^{\alpha_1}\wedge\cdots\wedge\vartheta^{\alpha_p}\\\wedge\vartheta^{\beta_1}\wedge\cdots\wedge\vartheta^{\beta_q}$
mielke-geometrodynamics_368 EQ1093 (B.3)	368	■ 1.000	$\alpha^{\{p\}}\wedge\beta^{\{q\}}=(-1)^{pq}\beta^{\{q\}}\wedge\alpha^{\{p\}},$	$\alpha^{(p)}\wedge\beta^{(q)} = (-1)^{pq}\beta^{(q)}\wedge\alpha^{(p)},$	$\alpha^{(p)}\wedge\beta^{(q)} = (-1)^{pq}\beta^{(q)}\wedge\alpha^{(p)},$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_368 EQ1094 (B.4)	368	■ 1.000	$\mathrm{D}(M):=\sum_{p=0}^n\wedge^pT^*(M)$	$D(M):=\sum_{p=0}^n\wedge^pT^*(M)$	$D(M):=\sum_{p=0}^n\wedge^pT^*(M)$
mielke-geometrodynamics_369 EQ1095 (B.5)	369	■ 1.000	$d\left(\alpha^{\left(p\right)}\wedge\beta^{\left(q\right)}\right)=d\alpha^{\left(p\right)}\wedge\beta^{\left(q\right)}+(-1)^p\alpha^{\left(p\right)}\wedge d\beta^{\left(q\right)}$	$d\left(\alpha^{\left(p\right)}\wedge\beta^{\left(q\right)}\right)=d\alpha^{\left(p\right)}\wedge\beta^{\left(q\right)}+(-1)^p\alpha^{\left(p\right)}\wedge d\beta^{\left(q\right)}$	$d(\alpha^{(p)}\wedge\beta^{(q)})=d\alpha^{(p)}\wedge\beta^{(q)}+(-1)^p\alpha^{(p)}\wedge d\beta^{(q)}$
mielke-geometrodynamics_369 EQ1097 (B.7)	369	■ 1.000	$\sigma:M\rightarrow N,\quad\sigma\in\mathscr{D}(M)$	$\sigma:M\rightarrow N,\quad\sigma\in\mathscr{D}(M)$	$\sigma:M\rightarrow N,\quad\sigma\in\mathscr{D}(M)$
mielke-geometrodynamics_369 EQ1098 (B.8)	369	■ 1.000	$\sigma_*:\wedge^pT^*(N)\rightarrow\wedge^pT^*(M)$	$\sigma_*:\wedge^pT^*(N)\rightarrow\wedge^pT^*(M)$	$\sigma_*:\wedge^pT^*(N)\rightarrow\wedge^pT^*(M)$
mielke-geometrodynamics_370 EQ1103 (B.15)	370	■ 1.000	$\{ \}^{* *} \alpha^{\left(p\right)}=(-1)^{p\left(n+1\right)+s}\alpha^{\left(p\right)}\text{ .}$	$**\alpha^{(p)}=(-1)^{p(n+1)+s}\alpha^{(p)}.$	$**\alpha^{(p)}=(-1)^{p(n+1)+s}\alpha^{(p)}.$
mielke-geometrodynamics_370 EQ1105 (B.17)	370	■ 1.000	$(\alpha,\beta):=\int_M\alpha^{\left(p\right)}\wedge^*\beta^{\left(p\right)}=(\beta,\alpha)$	$(\alpha,\beta):=\int_M\alpha^{(p)}\wedge^*\beta^{(p)}=(\beta,\alpha)$	$(\alpha,\beta):=\int_M\alpha^{(p)}\wedge^*\beta^{(p)}=(\beta,\alpha)$
mielke-geometrodynamics_370 EQ1107 (B.19)	370	■ 1.000	$\square\alpha^{\left(p\right)}:=d\delta\alpha^{\left(p\right)}+\delta d\alpha^{\left(p\right)}\text{ .}$	$\square\alpha^{(p)}:=d\delta\alpha^{(p)}+\delta d\alpha^{(p)}.$	$\square\alpha^{(p)}:=d\delta\alpha^{(p)}+\delta d\alpha^{(p)}.$
mielke-geometrodynamics_371 EQ1108 (B.20)	371	■ 1.000	$\begin{aligned}(\alpha,\square\alpha)&=(\alpha,d\delta\alpha)+(\alpha,\delta d\alpha)\\&=(\delta\alpha,\delta\alpha)+(d\alpha,d\alpha).\end{aligned}$	$\begin{aligned}(\alpha,\square\alpha)&=(\alpha,d\delta\alpha)+(\alpha,\delta d\alpha)\\&=(\delta\alpha,\delta\alpha)+(d\alpha,d\alpha).\end{aligned}$	$\begin{aligned}(\alpha,\square\alpha)&=(\alpha,d\delta\alpha)+(\alpha,\delta d\alpha)\\&=(\delta\alpha,\delta\alpha)+(d\alpha,d\alpha).\end{aligned}$
mielke-geometrodynamics_371 EQ1109 (B.21)	371	■ 1.000	$d\alpha=0\text{ .}$	$d\alpha=0.$	$d\alpha=0.$
mielke-geometrodynamics_371 EQ1111 (B.23)	371	■ 1.000	$\beta^{\left(p\right)}=d\alpha^{\left(p-1\right)}\in C^\infty(B(M))$	$\beta^{(p)}=d\alpha^{(p-1)}\in C^\infty(B(M))$	$\beta^{(p)}=d\alpha^{(p-1)}\in C^\infty(B(M))$
mielke-geometrodynamics_371 EQ1112 (B.24)	371	■ 1.000	$H(M):=Z(M)/B(M)$	$H(M):=Z(M)/B(M)$	$H(M):=Z(M)/B(M)$
mielke-geometrodynamics_371 EQ1113 (B.25)	371	■ 1.000	$H(M)=\sum_{p=0}^nH^p(M):=\sum_{p=0}^nZ^p(M)/B^p(M)\text{ .}$	$H(M)=\sum_{p=0}^nH^p(M):=\sum_{p=0}^nZ^p(M)/B^p(M).$	$H(M)=\sum_{p=0}^nH^p(M):=\sum_{p=0}^nZ^p(M)/B^p(M).$
mielke-geometrodynamics_371 EQ1115 (B.27)	371	■ 1.000	$H^{\left\{n\right\}}(M)=\wedge^{\left\{n\right\}}T^*(M)/B^{\left\{n\right\}}(M)\text{ .}$	$H^n(M)=\wedge^nT^*(M)/B^n(M).$	$H^n(M)=\wedge^nT^*(M)/B^n(M).$
mielke-geometrodynamics_371 EQ1116 (B.28)	371	■ 1.000	$H^{\left\{\emptyset\right\}}(M)=Z^{\left\{\emptyset\right\}}(M)$	$H^0(M)=Z^0(M)$	$H^0(M)=Z^0(M)$
mielke-geometrodynamics_372 EQ1117 (B.29)	372	■ 1.000	$H^{\left\{\emptyset\right\}}(M)\approx\mathbb{R}$	$H^0(M)\approx\mathbb{R}$	$H^0(M)\approx\mathbb{R}$
mielke-geometrodynamics_372 EQ1118 (B.30)	372	■ 1.000	$b_{\left\{p\right\}}:=\operatorname{dim}H^{\left\{p\right\}}(M)$	$b_p:=\dim H^p(M)$	$b_p:=\dim H^p(M)$
mielke-geometrodynamics_372 EQ1119 (B.31)	372	■ 1.000	$\mathrm{f}_{\mathrm{M}}(t)=\sum_{p=0}^nb_{\left\{p\right\}}t^{\left\{p\right\}}$	$\mathrm{f}_{\mathrm{M}}(t)=\sum_{p=0}^nb_pt^p$	$\mathrm{f}_{\mathrm{M}}(t)=\sum_{p=0}^nb_pt^p,$
mielke-geometrodynamics_372 EQ1120 (B.32)	372	■ 1.000	$\chi(M):=\mathrm{f}_{\mathrm{M}}(-1)=\sum_{p=0}^n(-1)^{\left\{p\right\}}b_{\left\{p\right\}},$	$\chi(M):=\mathrm{f}_{\mathrm{M}}(-1)=\sum_{p=0}^n(-1)^pb_p,$	$\chi(M):=\mathrm{f}_{\mathrm{M}}(-1)=\sum_{p=0}^n(-1)^pb_p,$
mielke-geometrodynamics_372 EQ1121 (B.33)	372	■ 1.000	$\mathrm{H}^{\left\{\emptyset\right\}}\left(S^{\left\{n\right\}}\right)=H^{\left\{n\right\}}\left(S^{\left\{n\right\}}\right)\approx\mathbb{R}$	$H^0(S^n)=H^n(S^n)\approx\mathbb{R}$	$H^0(S^n)=H^n(S^n)\approx\mathbb{R}$
mielke-geometrodynamics_372 EQ1123 (B.35)	372	■ 1.000	$\mathrm{f}_{S^{\left\{n\right\}}}(t)=1+t^{\left\{n\right\}},$	$\mathrm{f}_{S^n}(t)=1+t^n,$	$\mathrm{f}_{S^n}(t)=1+t^n,$

Identifier	Page	Conf.	LaTeX source	Rendered	Image
mielke-geometrodynamics_372 EQ1124 (B.36)		■ 1.000	$\chi\left(S^{\{n\}}\right)=\left\{\begin{array}{l} 0 \text { for } n=2 k+1, \\ 2 \text { for } n=2 k, \end{array}\right.$	$\chi\left(S^n\right)=\left\{\begin{array}{l} 0 \text { for } n=2 k+1, \\ 2 \text { for } n=2 k, \end{array}\right.$	$\chi\left(S^n\right)=\left\{\begin{array}{l} 0 \text { for } n=2 k+1, \\ 2 \text { for } n=2 k, \end{array}\right.$
mielke-geometrodynamics_372 EQ1125 (B.37)		■ 1.000	$\Delta_p:=\left\{x \in E^{\{n\}} \mid x=\sum_{i=0}^p \lambda^i \eta_i, \lambda^i \geq 0, \sum_{i=0}^p \lambda^i=1\right\}.$	$\Delta_p:=\left\{x \in E^n \mid x=\sum_{i=0}^p \lambda^i \eta_i, \lambda^i \geq 0, \sum_{i=0}^p \lambda^i=1\right\}.$	$\Delta_p:=\left\{x \in E^n \mid x=\sum_{i=0}^p \lambda^i \eta_i, \lambda^i \geq 0, \sum_{i=0}^p \lambda^i=1\right\}.$
mielke-geometrodynamics_373 EQ1127 (B.39)		■ 1.000	$\partial c=\sum_{i=0}^p(-1)^i c\left(\sum_{j=0}^{i-1} \lambda^j \eta_j+\sum_{j=i}^{p-1} \lambda^j \eta_{j+1}\right).$	$\partial c=\sum_{i=0}^p(-1)^i c\left(\sum_{j=0}^{i-1} \lambda^j \eta_j+\sum_{j=i}^{p-1} \lambda^j \eta_{j+1}\right).$	$\partial c=\sum_{i=0}^p(-1)^i c\left(\sum_{j=0}^{i-1} \lambda^j \eta_j+\sum_{j=i}^{p-1} \lambda^j \eta_{j+1}\right).$
mielke-geometrodynamics_373 EQ1129 (B.41)		■ 1.000	$H_{\{\mathrm{p}\}}(M, \mathbb{R}):=Z_{\{\mathrm{p}\}}(M, \mathbb{R}) / B_{\{\mathrm{p}\}}(M, \mathbb{R})$	$H_p(M, \mathbb{R}):=Z_p(M, \mathbb{R}) / B_p(M, \mathbb{R})$	$H_p(M, \mathbb{R}):=Z_p(M, \mathbb{R}) / B_p(M, \mathbb{R})$
mielke-geometrodynamics_373 EQ1130 (B.42)		■ 1.000	$\int_{c_{p+1}} d \alpha^{\{p\}}=\int_{\partial c_{p+1}} \alpha^{\{p\}}$	$\int_{c_{p+1}} d \alpha^{(p)}=\int_{\partial c_{p+1}} \alpha^{(p)}$	$\int_{c_{p+1}} d \alpha^{(p)}=\int_{\partial c_{p+1}} \alpha^{(p)}.$
mielke-geometrodynamics_373 EQ1132 (B.44)		■ 1.000	$\langle d \alpha, c\rangle_M=\langle \alpha, \partial c\rangle_M$	$\langle d \alpha, c\rangle_M=\langle \alpha, \partial c\rangle_M$	$\langle d \alpha, c\rangle_M=\langle \alpha, \partial c\rangle_M$
mielke-geometrodynamics_375 EQ1133 (C.1)		■ 1.000	$I_{\{j\}}:=\left.\frac{\partial g(\xi)}{\partial \xi^j}\right _{g=e} \in T_e(G) \approx \mathfrak{g}$	$I_j:=\left.\frac{\partial g(\xi)}{\partial \xi^j}\right _{g=e} \in T_e(G) \approx \mathfrak{g}$	$I_j:=\left.\frac{\partial g(\xi)}{\partial \xi^j}\right _{g=e} \in T_e(G) \approx \mathfrak{g}$
mielke-geometrodynamics_376 EQ1138 (C.6)		■ 1.000	$[\mathrm{AB}]:=\mathrm{A}^i \mathrm{B}^j \mathrm{c}_{ij}^k \mathrm{I}_k.$	$[\mathrm{AB}]:=\mathrm{A}^i \mathrm{B}^j \mathrm{c}_{ij}^k \mathrm{I}_k.$	$[\mathrm{AB}]:=\mathrm{A}^i \mathrm{B}^j \mathrm{c}_{ij}^k \mathrm{I}_k.$
mielke-geometrodynamics_376 EQ1139 (C.7)		■ 1.000	$\rho_A^C\left(g_1\right) \rho_C^B\left(g_2\right)=\rho_A^B\left(g_1 g_2\right)$	$\rho_A^C\left(g_1\right) \rho_C^B\left(g_2\right)=\rho_A^B\left(g_1 g_2\right)$	$\rho_A^C\left(g_1\right) \rho_C^B\left(g_2\right)=\rho_A^B\left(g_1 g_2\right)$
mielke-geometrodynamics_376 EQ1140 (C.8)		■ 1.000	$[A, B] \longrightarrow \rho([A, B])=[\rho(A), \rho(B)].$	$[A, B] \longrightarrow \rho([A, B])=[\rho(A), \rho(B)].$	$[A, B] \longrightarrow \rho([A, B])=[\rho(A), \rho(B)].$
mielke-geometrodynamics_376 EQ1141 (C.9)		■ 1.000	$\operatorname{Ad} A(B):=[A, B] ; \quad A, B \in \mathfrak{g},$	$\operatorname{Ad} A(B):=[A, B]; \quad A, B \in \mathfrak{g},$	$\operatorname{Ad} A(B):=[A, B]; \quad A, B \in \mathfrak{g},$
mielke-geometrodynamics_376 EQ1142 (C.10)		■ 1.000	$(\operatorname{Ad} A)_j^k=c_{ij}^k A^i.$	$(\operatorname{Ad} A)_j^k=c_{ij}^k A^i.$	$(\operatorname{Ad} A)_j^k=c_{ij}^k A^i.$
mielke-geometrodynamics_376 EQ1144 (C.12)		■ 1.000	$(A, B)=2 N \operatorname{Tr}(A B)-2 \operatorname{Tr}(A) \operatorname{Tr}(B)$	$(A, B)=2 N \operatorname{Tr}(A B)-2 \operatorname{Tr}(A) \operatorname{Tr}(B)$	$(A, B)=2 N \operatorname{Tr}(A B)-2 \operatorname{Tr}(A) \operatorname{Tr}(B)$
mielke-geometrodynamics_377 EQ1146 (C.14)		■ 1.000	$g=\exp ^{\left(-\frac{i}{2} \xi^j \lambda_j\right)} \in S U(N), j=1, \ldots, N^2-1 .$	$g=\exp ^{\left(-\frac{i}{2} \xi^j \lambda_j\right)} \in S U(N), j=1, \ldots, N^2-1 .$	$g=\exp ^{\left(-\frac{i}{2} \xi^j \lambda_j\right)} \in S U(N), j=1, \ldots, N^2-1 .$
mielke-geometrodynamics_377 EQ1148 (C.17)		■ 1.000	$\lambda_i \lambda_j=\frac{2}{N} \delta_{ij}+\left(d_{ij}^k+i f_{ij}^k\right) \lambda_k.$	$\lambda_i \lambda_j=\frac{2}{N} \delta_{ij}+\left(d_{ij}^k+i f_{ij}^k\right) \lambda_k.$	$\lambda_i \lambda_j=\frac{2}{N} \delta_{ij}+\left(d_{ij}^k+i f_{ij}^k\right) \lambda_k.$
mielke-geometrodynamics_377 EQ1149 (C.18)		■ 1.000	$\operatorname{Tr}\left(\lambda_i \lambda_j\right)=2 \delta_{ij}.$	$\operatorname{Tr}\left(\lambda_i \lambda_j\right)=2 \delta_{ij}.$	$\operatorname{Tr}\left(\lambda_i \lambda_j\right)=2 \delta_{ij}.$