


so far ...


- In the process of training a neural network, we look for **weights(network parameters)** that minimize the loss function.
- How to find weights?
 - an optimization problem → **SGD**
 - at each step, need to calculate gradients w.r.t every parameter.

Chain Rule



$$\frac{\partial y}{\partial x} = \frac{\partial y}{\partial C} \times \frac{\partial C}{\partial B} \times \frac{\partial B}{\partial A} \times \frac{\partial A}{\partial x}$$

Chain Rule - Multiple Path



$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial y_1} \frac{\partial y_1}{\partial x} + \frac{\partial z}{\partial y_2} \frac{\partial y_2}{\partial x}$$


Backpropagation is just repeated application of the chain rule.

A Simple Example

Consider following computational graph:

$$f(x, y, z) = (x + y)z$$

e.g. $x = -2, y = 5, z = -4$



$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$


$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want: $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$


A Simple Example

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$



$$\frac{\partial f}{\partial f}$$




$$\frac{\partial f}{\partial z}$$

A Simple Example

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$




$$\frac{\partial f}{\partial q}$$

A Simple Example

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$



$$\frac{\partial f}{\partial y}$$


Chain rule:

$$\frac{\partial f}{\partial y} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial y}$$

A Simple Example

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$




$$\frac{\partial f}{\partial x}$$


Chain rule:

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$


Gradient Backward Propagation



Forward-Backward Passes




Example (2)



Assume loss is as follows:

$$J = y^* \log y + (1 - y^*) \log(1 - y)$$



Example (2)

Forward

$$J = y^* \log y + (1 - y^*) \log(1 - y)$$

$$y = \frac{1}{1 + \exp(-b)}$$

$$b = \sum_{j=0}^D \beta_j z_j$$

$$z_j = \frac{1}{1 + \exp(-a_j)}$$

$$a_j = \sum_{i=0}^M \alpha_{ji} x_i$$

Backward

$$\frac{dJ}{dy} = \frac{y^*}{y} + \frac{(1 - y^*)}{y - 1}$$

$$\frac{dJ}{db} = \frac{dJ}{dy} \frac{dy}{db}, \quad \frac{dy}{db} = \frac{\exp(-b)}{(\exp(-b) + 1)^2}$$

$$\frac{dJ}{d\beta_j} = \frac{dJ}{db} \frac{db}{d\beta_j}, \quad \frac{db}{d\beta_j} = z_j$$

$$\frac{dJ}{dz_j} = \frac{dJ}{db} \frac{db}{dz_j}, \quad \frac{db}{dz_j} = \beta_j$$

$$\frac{dJ}{da_j} = \frac{dJ}{dz_j} \frac{dz_j}{da_j}, \quad \frac{dz_j}{da_j} = \frac{\exp(-a_j)}{(\exp(-a_j) + 1)^2}$$

$$\frac{dJ}{d\alpha_{ji}} = \frac{dJ}{da_j} \frac{da_j}{d\alpha_{ji}}, \quad \frac{da_j}{d\alpha_{ji}} = x_i$$

$$\frac{dJ}{dx_i} = \frac{dJ}{da_j} \frac{da_j}{dx_i}, \quad \frac{da_j}{dx_i} = \sum_{j=0}^D \alpha_{ji}$$