Adamax 10mg

QUICK GUIDE

No evidence of any kind ♥



WHAT IS ADAMAX?

Adamax is sold as a modified version of the Russian nootropic Semax. No published research of any kind was found: no human, animal or chemistry data. Its catalog page has a dosing table and nothing else.

Strongest evidence: N/E · Not enough evidence

RECONSTITUTION

Add 3 mL bacteriostatic water to the 10 mg vial.

This yields 3.3 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 3 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog 8-week ladder (grade E)

WEEK	DOSE (Daily)	UNITS (U-100 SYRINGE)
Weeks 1–2	300 mcg	9 units
Weeks 3–4	500 mcg	15 units
Weeks 5–6	750 mcg	22.5 units
Weeks 7–8	1 mg	30 units

Cycle: catalog: 8–12 weeks on, washout equal to the cycle length.

POTENTIAL BENEFITS

- **Evidence:** None found. N/E · Not enough evidence
- **Identity:** Structure not verified. N/E · Not enough evidence

POTENTIAL SIDE EFFECTS

- Unknown

Vial size isn't stated; 10 mg is implied.
Assuming it acts like Semax

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade N/E: Not enough evidence

For educational use only.


PEPPERSDVO