ARA-290 (Cibinetide) 16mg
QUICK GUIDE

Tissue repair without the EPO effects ♥



WHAT IS IT?

ARA-290 (cibinetide) is a small piece of erythropoietin that keeps EPO's tissue-protective signaling without stimulating red blood cells. In 28-day trials for nerve damage in sarcoidosis, it increased small nerve fibers in the cornea and skin. Development stopped before phase 3.

Strongest evidence: B · Human trials

RECONSTITUTION

Add 1.5 mL bacteriostatic water to the 16 mg vial.

This yields 10.7 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 1.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Trial course: 4 mg daily × 28 days (grade B)

WEEK	DOSE (Once daily (catalog and trials))	UNITS (U-100 SYRINGE)
Weeks 1–4	4 mg	37.5 units

Cycle: catalog: 4/8/12 weeks on, 2/4/6 weeks off.

POTENTIAL BENEFITS

- **Nerve fibers:** Corneal and skin small-fiber counts rose (phase 2). B · Human trials
- **No EPO effect:** Doesn't raise red-cell production. B · Human trials
- **Pain:** Placebo-corrected pain benefit not significant. B · Human trials

POTENTIAL SIDE EFFECTS

- Well tolerated in 28-day trials
- Injection-site reactions

2 mg = 18.75 units exactly; the catalog prints 18 (rounding).

Unit math otherwise matches (10.7 mg/mL).

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade B: Human trials

For educational use only.


PEPPERSDVO