BPC-157 / TB-500 / Cartalax 5mg
QUICK GUIDE

Catalog blend: untested as a combination ♥



WHAT IS IT?

BPC-157 / TB-500 / Cartalax is a pre-made catalog blend of BPC-157, TB-500, Cartalax. Each ingredient keeps its own evidence grade (BPC-157 C, TB-500 D, Cartalax D). No study of this combination was found.

Strongest evidence: **E · Community / vendor**

RECONSTITUTION

Add 2 mL bacteriostatic water to the 5 mg vial.

This yields 2.5 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog protocol (grade E)

WEEK	DOSE (Per catalog protocol)	UNITS (U-100 SYRINGE)
Weeks 1–2	150 mcg	6 units
Weeks 3–8	200 mcg	8 units

Cycle: Catalog cycle.

POTENTIAL BENEFITS

- **BPC-157:** 5 mg per vial · grade C **C · Human observational**
- **TB-500:** 5 mg per vial · grade D **D · Animal / lab**
- **Cartalax:** 10 mg per vial · grade D **D · Animal / lab**

POTENTIAL SIDE EFFECTS

- See each ingredient's sheet
- Fixed-ratio dosing

The catalog gives units only. Per 6 units: 150 mcg BPC-157, 150 mcg TB-500, 300 mcg Cartalax (computed).

Stub page: no description.

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade E: Community / vendor

For educational use only.


PEPPERSDVO