BPC-157 10mg

QUICK GUIDE

The most popular repair peptide, built on animal data ♥



WHAT IS BPC-157?

BPC-157 is a 15-amino-acid peptide based on a stomach protein. Hundreds of rodent studies report faster healing of tendons, muscle and the gut. In people there are only three tiny uncontrolled pilots and no randomized trial. The FDA restricts compounding it.

Strongest evidence: C · Human observational

RECONSTITUTION

Add 2.5 mL bacteriostatic water to the 10 mg vial.

This yields 4 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog ladder, Mon–Fri (grade E)

WEEK	DOSE (1–3× daily, 5 days a week)	UNITS (U-100 SYRINGE)
Week 1	250 mcg	6.3 units
Weeks 2–4	500 mcg	12.5 units
Weeks 5–8	750 mcg	18.8 units

The catalog allows 1–3 injections a day: set 'Injections/day'.

Cycle: catalog: 8–10 weeks on, 4–6 weeks off.

POTENTIAL BENEFITS

- **Rodent repair:** Faster tendon, muscle and gut healing in animals. D · Animal / lab
- **Human data:** 3 small uncontrolled pilots; no RCTs. C · Human observational
- **Catalog math:** 20 mg vial: '4 units = 250 mcg' is really 320 mcg. E · Community / vendor

POTENTIAL SIDE EFFECTS

- Few adverse events reported in tiny pilots
- Injection-site reactions
- Unknown long-term risk

20 mg vial (8 mg/mL, 80 mcg/unit): '4 units = 250 mcg' is really 320 mcg (28% over); '10 units = 750 mcg' is really 800 mcg.

10 mg vial matches (40 mcg/unit).

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade C: Human observational

For educational use only.

PEPPERSDVO