Cagrilintide 10mg

QUICK GUIDE

Amylin pathway satiety ♥



WHAT IS CAGRILINTIDE?

Cagrilintide is a once-weekly long-acting amylin analog. Amylin is a satiety hormone released with insulin, and it acts on a different pathway from GLP-1. Alone it produced about 11% weight loss in phase 2. Paired with semaglutide (CagriSema), it reached about 20% in phase 3.

Strongest evidence: B · Human trials

RECONSTITUTION

Add 2 mL bacteriostatic water to the 10 mg vial.

This yields 5 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog weekly ladder to 2.7 mg (grade E)

WEEK	DOSE (Once weekly, same day each week)	UNITS (U-100 SYRINGE)
Week 1	200 mcg	4 units
Week 2	400 mcg	8 units
Week 3	750 mcg	15 units
Week 4	1.2 mg	24 units
Week 5	1.7 mg	34 units
Week 6	2.2 mg	44 units
Weeks 7–8	2.7 mg	54 units

Much faster than the trials (16 weeks to 2.4 mg in REDEFINE 1).

Cycle: catalog: 8 weeks on, 4–8 weeks off (56 / 28–56 days). No trial basis.

POTENTIAL BENEFITS

- **Monotherapy:** Up to −10.8% at 4.5 mg over 26 weeks (phase 2). B · Human trials
- **With semaglutide:** −20.4% vs −3.0% at 68 weeks (REDEFINE 1). B · Human trials
- **Satiety:** Amylin receptors in the area postrema and hypothalamus. B · Human trials
- **Fast catalog ladder:** Catalog escalates weekly; trials took 6–16 weeks. E · Community / vendor

POTENTIAL SIDE EFFECTS

- Nausea: 20–47% vs 18% placebo (phase 2)
- Other GI effects, dose-related
- Injection-site reactions
- Combined with semaglutide: more GI effects than either alone

Escalation: the catalog steps up every week to 2.7 mg in 7 weeks. REDEFINE 1 took 16 weeks to reach 2.4 mg, and phase 2 took up to 6 weeks.

Top dose: the catalog's 2.7 mg is above the 2.4 mg phase 3 dose.

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade B: Human trials

For educational use only.

PEPPERSDVO