Cerebrolysin 64mg

QUICK GUIDE

IV brain-peptide drug abroad; the catalog powder isn't it



WHAT IS CEREBROLYSIN?

Cerebrolysin is a peptide mixture made from pig brain. It's sold as an IV solution in many countries outside the US for dementia, stroke and brain injury. A Cochrane review found no survival benefit in stroke and more non-fatal serious adverse events. The catalog's powder, dosed in milligrams, isn't the same product.

Strongest evidence: A · Approved use

RECONSTITUTION

Add 8 mL bacteriostatic water to the 64 mg vial.

This yields 8 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 8 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog: 8 mg daily, rising to 3×/day (grade E)

WEEK	DOSE (1–3× daily)	UNITS (U-100 SYRINGE)
Weeks 1–4	8 mg	100 units

The catalog goes 1×/day (wk 1), 2×/day (wk 2), 3×/day (wks 3–4): adjust 'Injections/day'.

Cycle: catalog: 4 weeks on, 2–4 weeks off.

POTENTIAL BENEFITS

- **Approved abroad:** IV solution for dementia, stroke, brain trauma. A · Approved use
- **Stroke review:** No mortality benefit; more non-fatal serious AEs. A · Approved use
- **Catalog form:** Powder at mg doses vs a 215 mg/mL IV solution. E · Community / vendor

POTENTIAL SIDE EFFECTS

- Non-fatal serious adverse events increased in stroke trials
- Contraindicated in epilepsy

Form: the approved Cerebrolysin is a 215.2 mg/mL solution. The catalog sells a powder at about 8 mg/mL after mixing.

Dose: clinical use is milliliters IV (e.g. 30 mL/day); the catalog's 8–24 mg/day SC is a tiny fraction.

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade A: Approved use

For educational use only.


PEPPERSDVO