CJC-1295 (no DAC) 10mg
QUICK GUIDE

Short-acting GHRH analog, no human trials ♥



WHAT IS CJC-1295 (NO DAC)?

CJC-1295 without DAC (also called modified GRF 1–29) is a breakdown-resistant version of sermorelin that gives a short pulse of growth hormone. The long-acting DAC version reached human trials. This no-DAC form never did.

Strongest evidence: E · Community / vendor

RECONSTITUTION

Add 2.5 mL bacteriostatic water to the 10 mg vial.

This yields 4 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog 1×/day: 100 → 150 mcg (grade E)

WEEK	DOSE (1–3 times daily)	UNITS (U-100 SYRINGE)
Weeks 1–4	100 mcg	2.5 units
Weeks 5–12	150 mcg	3.8 units

Cycle: catalog: 12 weeks on, 6–12 weeks off.

POTENTIAL BENEFITS

- **Mechanism:** GHRH receptor: a short natural GH pulse. E · Community / vendor
- **DAC cousin:** The DAC form raised GH/IGF-1 for 9–11 days in humans. B · Human trials
- **Catalog math:** '5 units = 150 mcg' is really 200 mcg. E · Community / vendor

POTENTIAL SIDE EFFECTS

- No human data for this form
- Flushing, head rush (GHRH class, anecdotal)
- Water retention at higher GH

Catalog: '5 units (150 mcg)'. At 4 mg/mL (40 mcg/unit), 5 units is 200 mcg, so it's 33% over the stated dose. The catalog's own 3×/day table correctly prints 150 mcg as 4 units.

100 mcg = 2.5 units; the catalog prints 3 (rounding).

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade E: Community / vendor

For educational use only.


PEPPERSDVO