GHRP-2 10mg

QUICK GUIDE

Potent GH secretagogue, a diagnostic in Japan ♥



WHAT IS GHRP-2?

GHRP-2 (pralmorelin) is a six-amino-acid peptide that activates the ghrelin receptor and causes a strong growth-hormone release. Japan approved it as a one-time diagnostic test for GH deficiency. Repeated use for body composition is untested.

Strongest evidence: A · Approved use

RECONSTITUTION

Add 3 mL bacteriostatic water to the 10 mg vial.

This yields 3.3 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 3 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog: 150 mcg at bedtime (grade E)

WEEK	DOSE (1–3 times daily)	UNITS (U-100 SYRINGE)
Weeks 1–12	150 mcg	4.5 units

Cycle: catalog: 8–12 weeks on, 4 weeks off.

POTENTIAL BENEFITS

- **Approved use:** GH-deficiency diagnostic in Japan. A · Approved use
- **Strong GH release:** Marked GH rise in healthy people. A · Approved use
- **Catalog math:** '6 units = 150 mcg' is really 200 mcg. E · Community / vendor

POTENTIAL SIDE EFFECTS

- Hunger
- Slight cortisol / prolactin rise
- Water retention, flushing (class)

Catalog: '6 units = 150 mcg'. At 10 mg in 3 mL (33.3 mcg/unit), 6 units is 200 mcg, 33% over. 150 mcg is 4.5 units.

300 mcg = 9 units matches.

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade A: Approved use

For educational use only.


PEPPERSDVO