Semaglutide / Cagrilintide 5/5 5mg
QUICK GUIDE

A studied pairing, as a research powder ♥



WHAT IS IT?

Semaglutide / Cagrilintide 5/5 is a pre-made catalog blend of Semaglutide, Cagrilintide. Each ingredient keeps its own evidence grade (Semaglutide A, Cagrilintide B). This pairing was tested in phase 3 as CagriSema, though not as this research powder.

Strongest evidence: B · Human trials

RECONSTITUTION

Add 1.5 mL bacteriostatic water to the 5 mg vial.

This yields 3.3 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 1.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog protocol (grade E)

WEEK	DOSE (Once weekly)	UNITS (U-100 SYRINGE)
Weeks 1–4	250 mcg	7.5 units
Weeks 5–8	500 mcg	15 units
Weeks 9–12	750 mcg	22.5 units
Weeks 13–16	1 mg	30 units

POTENTIAL BENEFITS

- **Semaglutide:** 5 mg per vial · grade A A · Approved use
- **Cagrilintide:** 5 mg per vial · grade B B · Human trials

POTENTIAL SIDE EFFECTS

- See each ingredient's sheet
- Fixed-ratio dosing

UNIT MATH ERROR: the catalog says to mix with 1.5 mL, but its table (5 units = 0.25 mg … 50 units = 2.5 mg) only works with 1 mL. At 1.5 mL, 5 units = 0.167 mg of each drug: one-third less than stated at every step.

The catalog's 2.5 mg maximum is above the CagriSema trial dose (2.4/2.4 mg).

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade B: Human trials

For educational use only.

PEPPERSDVO