Retatrutide / Cagrilintide 12.5/2.5 12.5mg
QUICK GUIDE

Catalog blend: untested as a combination ♥



WHAT IS IT?

Retatrutide / Cagrilintide 12.5/2.5 is a pre-made catalog blend of Retatrutide, Cagrilintide. Each ingredient keeps its own evidence grade (Retatrutide B, Cagrilintide B). No study of this combination was found.

Strongest evidence: E · Community / vendor

RECONSTITUTION

Add 1.5 mL bacteriostatic water to the 12.5 mg vial.

This yields 8.3 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 1.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog protocol (grade E)

WEEK	DOSE (Once weekly)	UNITS (U-100 SYRINGE)
Weeks 1–4	1 mg	12 units
Weeks 5–8	3 mg	36 units

POTENTIAL BENEFITS

- **Retatrutide:** 12.5 mg per vial · grade B — B · Human trials
- **Cagrilintide:** 2.5 mg per vial · grade B — B · Human trials

POTENTIAL SIDE EFFECTS

- See each ingredient's sheet
- Fixed-ratio dosing

The 1 mg retatrutide start is below the 2 mg trial start.

The catalog's anti-cagrilintide antibody claim wasn't verified.

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade E: Community / vendor

For educational use only.


PEPPERSDVO