Retatrutide 30mg

QUICK GUIDE

Next-generation metabolic peptide ♥



WHAT IS RETATRUTIDE?

Retatrutide is a single peptide that activates three gut-hormone receptors: GIP, GLP-1 and glucagon. GLP-1 and GIP reduce appetite and improve blood-sugar control. The glucagon arm is thought to raise energy use and liver fat burning. In trials it has produced larger average weight loss than any approved GLP-1 drug.

Strongest evidence: B · Human trials

RECONSTITUTION

Add 2.5 mL bacteriostatic water to the 30 mg vial.

This yields 12 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Trial ladder 2→4→8→12 mg (phase 2, grade B)

WEEK	DOSE (Once weekly, same day each week)	UNITS (U-100 SYRINGE)
Weeks 1–4	2 mg	16.7 units
Weeks 5–8	4 mg	33.3 units
Weeks 9–12	8 mg	66.7 units
Week 13 onward	12 mg	100 units

Phase 2 12-mg arm: 4-week steps. Phase 3 maintenance doses were 4, 9 or 12 mg; edit the last step to match.

POTENTIAL BENEFITS

- **Significant weight loss:** −28.3% mean at 12 mg over 80 weeks (TRIUMPH-1). B · Human trials
- **Glycemic control:** HbA1c −2.02% at 24 wk vs −1.41% on dulaglutide (T2D). B · Human trials
- **Liver fat:** ≈ −82% relative liver fat at 8–12 mg by week 24. B · Human trials
- **Appetite suppression:** Central GLP-1R satiety plus slower gastric emptying. D · Animal / lab
- **Energy expenditure:** Glucagon-receptor arm raises energy use in rodents; not yet quantified in humans. D · Animal / lab
- **Knee OA:** Pain and function improved alongside weight loss (TRIUMPH-4). B · Human trials

POTENTIAL SIDE EFFECTS

- Nausea: 42% at 12 mg (15% placebo)
- Diarrhea: 32%
- Constipation: 26%
- Vomiting: 25%

Start dose: the catalog uses 4 mg for GLP-experienced users. Phase 2 found more GI events with a 4 mg start than a 2 mg start.

The catalog gives no escalation schedule. The trials used 4-week steps to 12 mg (phase 2) and 4 / 9 / 12 mg maintenance (phase 3).

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade B: Human trials

For educational use only.

PEPPERSDVO