CJC-1295 (no DAC) / Ipamorelin 10/10 20mg
QUICK GUIDE

Catalog blend: untested as a combination ♥



WHAT IS IT?

CJC-1295 (no DAC) / Ipamorelin 10/10 is a pre-made catalog blend of CJC-1295 (no DAC), Ipamorelin. Each ingredient keeps its own evidence grade (CJC-1295 (no DAC) E, Ipamorelin B). No study of this combination was found.

Strongest evidence: E · Community / vendor

RECONSTITUTION

Add 2.5 mL bacteriostatic water to the 20 mg vial.

This yields 8 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog protocol (grade E)

WEEK	DOSE (Per catalog protocol)	UNITS (U-100 SYRINGE)
Weeks 1–2	100 mcg	1.3 units
Weeks 3–4	150 mcg	1.9 units
Weeks 5–6	200 mcg	2.5 units
Weeks 7–12	250 mcg	3.1 units

Cycle: Catalog cycle.

POTENTIAL BENEFITS

- **CJC-1295 (no DAC):** 10 mg per vial · grade E E · Community / vendor
- **Ipamorelin:** 10 mg per vial · grade B B · Human trials

POTENTIAL SIDE EFFECTS

- See each ingredient's sheet
- Fixed-ratio dosing

The dose table is internally inconsistent: 1 → 4 units can't equal 100 → 250 mcg. At 8 mg/mL combined (80 mcg/unit), 4 units is 320 mcg combined (stated 250) and 5 units is 400 (stated 300). The catalog admits it rounds partial units.

The stated mcg don't say whether they mean per peptide or combined; this sheet uses combined, which fits the first rows best.

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade E: Community / vendor

For educational use only.

PEPPERSDVO