IGF-1 LR3 1mg

QUICK GUIDE

Long-acting IGF-1 analog, no human data ♥



WHAT IS IGF-1 LR3?

IGF-1 LR3 is an engineered, longer-lasting form of insulin-like growth factor 1, made mainly as a cell-culture reagent. It acts directly on IGF-1 receptors and bypasses growth hormone. It has never been studied in humans. Its approved cousin, mecasermin, carries a serious hypoglycemia warning.

Strongest evidence: D · Animal / lab

RECONSTITUTION

Add 2.5 mL 0.6% acetic acid to the 1 mg vial.

This yields 0.4 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog 8-week ladder 33→100 mcg (grade E)

WEEK	DOSE (Once daily)	UNITS (U-100 SYRINGE)
Week 1	33 mcg	8.3 units
Week 2	50 mcg	12.5 units
Weeks 3–4	75 mcg	18.7 units
Weeks 5–8	100 mcg	25 units

Cycle: catalog: 8 weeks on, 4–8 weeks off.

POTENTIAL BENEFITS

- **Direct IGF-1:** Bypasses GH; weak binding to IGF-binding proteins. D · Animal / lab
- **Hypoglycemia:** The approved IGF-1 drug warns of severe hypoglycemia. A · Approved use
- **Human data:** None. N/E · Not enough evidence

POTENTIAL SIDE EFFECTS

- Hypoglycemia (class)
- No human data for LR3

Catalog advises fasted dosing. The approved IGF-1 drug's label says to dose with a meal to avoid hypoglycemia.

Unit math: every unit value the catalog prints matches (0.4 mg/mL = 4 mcg/unit).

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade D: Animal / lab

For educational use only.

PEPPERSDVO