Ipamorelin 10mg

QUICK GUIDE

Ghrelin-receptor GH secretagogue



WHAT IS IPAMORELIN?

Ipamorelin is a small peptide that activates the ghrelin receptor and triggers a pulse of growth hormone. It was tested in humans only for bowel recovery after surgery, and that trial missed its goal. Its body-composition use is untested.

Strongest evidence: B · Human trials

RECONSTITUTION

Add 2.5 mL bacteriostatic water to the 10 mg vial.

This yields 4 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog: 100 mcg at bedtime (grade E)

WEEK	DOSE (1–3 times daily)	UNITS (U-100 SYRINGE)
Weeks 1–12	100 mcg	2.5 units

Cycle: catalog: 8–12 weeks on, 4–6 weeks off.

POTENTIAL BENEFITS

- **GH pulse:** Ghrelin-receptor agonist: pituitary GH release. D · Animal / lab
- **Phase 2:** Postoperative ileus trial not significant (p=0.15). B · Human trials
- **SC use:** No body-composition trials. N/E · Not enough evidence

POTENTIAL SIDE EFFECTS

- Adverse events similar to placebo in the ileus trial
- Hunger, water retention (class, anecdotal)
- Injection-site reactions

100 mcg = 2.5 units and 300 mcg = 7.5 units; the catalog prints 3 and 8 (rounding).

Active cancer (raises GH/IGF-1)

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade B: Human trials

For educational use only.


PEPPERSDVO