LL-37 5mg

QUICK GUIDE

Human antimicrobial peptide: studied on wounds ♥



WHAT IS LL-37?

LL-37 is the only human cathelicidin, an antimicrobial peptide that also helps wounds heal. Applied to chronic leg ulcers, the lower doses sped healing several-fold in a small trial. Injected LL-37 has never been studied in humans.

Strongest evidence: **B · Human trials**

RECONSTITUTION

Add 2.5 mL bacteriostatic water to the 5 mg vial.

This yields 2 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog 12-week ladder (grade E)

WEEK	DOSE (Daily, then 5 days/week)	UNITS (U-100 SYRINGE)
Week 1	50 mcg	2.5 units
Week 2	100 mcg	5 units
Week 3	150 mcg	7.5 units
Week 4	200 mcg	10 units
Week 5	300 mcg	15 units
Week 6	350 mcg	17.5 units
Week 7	400 mcg	20 units
Weeks 8–12	450 mcg	22.5 units

Units are computed from the stated mcg, not the catalog's printed units.

Cycle: catalog: 8–12 weeks on, 4–8 weeks off.

POTENTIAL BENEFITS

- **Antimicrobial:** Disrupts bacterial membranes and biofilms. B · Human trials
- **Leg ulcers:** Healing rate ~6× placebo at 0.5 mg/mL topical. B · Human trials
- **Catalog math:** Weeks 5–12 units run ≈13% short. E · Community / vendor

POTENTIAL SIDE EFFECTS

- Topical: no safety concerns in the wound trial
- Injected: no data
- Injection-site irritation (cationic peptide)

Weeks 5–12: the catalog's units are short of its stated mcg. 300 mcg = 15 u (catalog 13), 350 = 17.5 (15), 400 = 20 (18), 450 = 22.5 (20). Following the units gives about 13% less than stated.

Typo: 'Week 4: 10 units (200cg)' means 200 mcg.

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade B: Human trials

For educational use only.

PEPPERSDVO