Melanotan II 10mg

QUICK GUIDE

Tanning peptide with a case-report trail ♥



WHAT IS MELANOTAN II?

Melanotan II is a synthetic hormone analog that activates several melanocortin receptors at once: it tans the skin and causes erections and nausea. It has never been approved, and published case reports include melanoma, priapism and kidney injury.

Strongest evidence: C · Human observational

RECONSTITUTION

Add 2.5 mL bacteriostatic water to the 10 mg vial.

This yields 4 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog loading ladder (grade E)

WEEK	DOSE (Daily loading, then 1–2×/week)	UNITS (U-100 SYRINGE)
Week 1	250 mcg	6.3 units
Week 2	500 mcg	12.5 units
Week 3	750 mcg	18.8 units
Weeks 4–8	1 mg	25 units

Units are computed from the stated mcg.

Cycle: catalog: 8–16 weeks on, 8–16 weeks off.

POTENTIAL BENEFITS

- **Tanning:** MC1R-driven pigmentation without UV. C · Human observational
- **Moles:** New and darkening moles; melanoma case reports. C · Human observational
- **Priapism:** Ischemic priapism requiring surgery reported. C · Human observational
- **Quality:** Gray-market vials held 43–88% of the labeled amount. C · Human observational

POTENTIAL SIDE EFFECTS

- Nausea, flushing (dose-limiting)
- Spontaneous erections; priapism reported
- New or darkening moles
- Appetite suppression

Catalog: '28 units (1,000 mcg)'. At 4 mg/mL, 1,000 mcg is 25 units; 28 units is 1,120 mcg (12% over).

500 mcg = 12.5 units; the catalog prints 13 (rounding).

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade C: Human observational

For educational use only.

PEPPERSDVO