Methylene Blue (oral)
QUICK GUIDE

Oral · grade A ♥



WHAT IS IT?

Methylene Blue (oral): redox dye; MAO-A inhibitor. The approved use is IV methylene blue (ProvayBlue) for acquired methemoglobinemia (accelerated approval). Oral methylene blue isn't FDA-approved for anything.

Strongest evidence: A · Approved use

HOW IT'S TAKEN

25 mg pills

1 pill = 25 mg

The approved use is IV methylene blue (ProvayBlue) for acquired methemoglobinemia (accelerated approval) A · Approved use

Oral LMTM (a methylene-blue derivative) failed its Alzheimer's phase 3 primary endpoints C · Human observational

DOSING PROTOCOL

Catalog: 1 pill (grade E)

WEEK	DOSE (1 pill daily)
Weeks 1–8	1 pill

Cycle: Catalog cycle.

POTENTIAL BENEFITS

- ♥ **Evidence:** The approved use is IV methylene blue (ProvayBlue) for acquired methemoglobinemia (accelerate... A · Approved use
- ♥ **Evidence:** Oral LMTM (a methylene-blue derivative) failed its Alzheimer's phase 3 primary endpoints. C · Human observational
- ♥ **Evidence:** No adequate trials support cognitive or anti-aging use in healthy adults. C · Human observational
- ♥ **Caution:** BOXED WARNING (label): serious or fatal serotonin syndrome with SSRIs, SNRIs, MAOIs and other... A · Approved use

POTENTIAL SIDE EFFECTS

- BOXED WARNING
- Contraindicated in G6PD deficiency
- Turns urine blue-green

BOXED WARNING (label): serious or fatal serotonin syndrome with SSRIs, SNRIs, MAOIs and other serotonergic drugs (opioids, dextromethorphan); none within 72 h

Contraindicated in G6PD deficiency (hemolysis)

Oral use

Store as the label directs.

Evidence grade A: Approved use

For educational use only.


PEPPERSDVO