NAD+ 500mg

QUICK GUIDE

Energy coenzyme, thin injection data ♥



WHAT IS NAD+?

NAD+ is a coenzyme every cell uses to make energy and repair DNA, and levels fall with age. IV infusions are popular in clinics, but plasma levels barely move in the first hours, and the only human data are a small pharmacokinetic pilot. Under-the-skin NAD+ hasn't been studied.

Strongest evidence: C · Human observational

RECONSTITUTION

Add 2.5 mL bacteriostatic water to the 500 mg vial.

This yields 200 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog ladder (grade E)

WEEK	DOSE (2–7 times weekly)	UNITS (U-100 SYRINGE)
Week 1	25 mg	12.5 units
Weeks 2–6	50 mg	25 units
Weeks 7–12	100 mg	50 units

The catalog moves to 4×/week from week 3: change the days then.

Cycle: catalog: up to 12 weeks on, 4–6 weeks off.

POTENTIAL BENEFITS

- **Energy metabolism:** Core redox cofactor; feeds sirtuins and PARPs. C · Human observational
- **Fast clearance:** No plasma rise in the first 2 h of IV infusion. C · Human observational
- **Side effects:** Flushing, nausea, chest tightness. C · Human observational

POTENTIAL SIDE EFFECTS

- Flushing, nausea, cramping
- Chest tightness (IV, rate-related)
- Injection-site discomfort

25 mg = 12.5 units exactly; the catalog prints 12 (rounding).

Unit math otherwise matches (200 mg/mL).

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 30 days.

Evidence grade C: Human observational

For educational use only.

PEPPERSDVO