Oxytocin 10mg

QUICK GUIDE

The bonding hormone: proven in labor, unproven daily ♥



WHAT IS OXYTOCIN?

Oxytocin is the pituitary hormone behind labor contractions and milk let-down, and it also acts on the brain's social circuits. IV oxytocin (Pitocin) is a standard obstetric drug. A large trial of intranasal oxytocin for autism found no benefit, and daily injections haven't been studied.

Strongest evidence: A · Approved use

RECONSTITUTION

Add 2.5 mL bacteriostatic water to the 10 mg vial.

This yields 4 mg/mL.

1. Swab both vial tops with an alcohol pad.
2. Draw 2.5 mL of water and inject it slowly down the vial wall.
3. Roll gently until dissolved; don't shake.
4. Label and refrigerate.

DOSING PROTOCOL

Catalog: 100 mcg daily (grade E)

WEEK	DOSE (Once daily)	UNITS (U-100 SYRINGE)
Weeks 1–8	100 mcg	2.5 units

Cycle: catalog: 8 weeks on, 4 weeks off.

POTENTIAL BENEFITS

- **Approved:** IV/IM for labor (Pitocin). A · Approved use
- **Social effects:** Large autism trial: no benefit. B · Human trials
- **Water balance:** Antidiuretic: water intoxication risk. A · Approved use
- **Catalog math:** '6 units = 200 mcg' is really 240 mcg. E · Community / vendor

POTENTIAL SIDE EFFECTS

- Water retention / low sodium (antidiuretic effect)
- Uterine contractions
- Nausea, headache (intranasal studies)

Catalog: '6 units (200 mcg)'. At 4 mg/mL, 6 units is 240 mcg (20% over); 200 mcg is 5 units.

100 mcg = 2.5 units; the catalog prints 3 (rounding).

For subcutaneous injection

Refrigerate after mixing. Catalog: use within 60 days.

Evidence grade A: Approved use

For educational use only.


PEPPERSDVO