Bremelanotide / PT-141 (nasal)
QUICK GUIDE

Intranasal form: does it get in? ♥



WHAT IS IT?

Bremelanotide / PT-141 (nasal) is the catalog's intranasal version of PT-141. Intranasal bremelanotide WAS tested: the FDA halted the phase II nasal trials in 2007 because of blood-pressure increases, and the nasal form was abandoned in 2008. The developer blamed erratic nasal uptake causing high blood levels in some users; it was reformulated as the SC injection now approved.

Strongest evidence: B · Human trials

HOW IT'S TAKEN

Nasal spray

1 spray = 100 mcg

Intranasal bremelanotide WAS tested: the FDA halted the phase II nasal trials in 2007 because of blood-pressure increases, and the nasal form was abandoned in 2008 B · Human trials

DOSING PROTOCOL

- 3–4 sprays (300–400 mcg) 30–60 min before sex
- Catalog: "trials used 10–20 mg without adverse effects... we are not recommending a maximum"

POTENTIAL BENEFITS

- ♥ **Route evidence:** Intranasal bremelanotide WAS tested: the FDA halted the phase II nasal trials in 2007 be... B · Human trials

POTENTIAL SIDE EFFECTS

- See the molecule's own sheet

DANGEROUS CLAIM: the catalog says nasal trials used 10–20 mg "without adverse effects" and sets no maximum. In fact the nasal trials were halted by the FDA over blood-pressure rises. The approved injection is 1.75 mg, at most once in 24 hours and 8 times a month.

Everything on the molecule's own sheet

Intranasal use

Store as the label directs.

Evidence grade B: Human trials

For educational use only.

PEPPERSDVO